

# Lecture 7: Hypothesis testing part I

## Statistical Methods for Data Science

**Yinan Yu**

Department of Computer Science and Engineering

November 25, 2021

# Today

## 1 Terminology

- Experiment and parameter of interest
- Null hypothesis and alternative hypothesis
- Test statistic
- Null distribution  $f(s \mid H_0)$
- Significance level  $\alpha$ , power and *p*-value

## 2 Example

## 3 *p*-hacking

## 4 Summary



## Learning outcome

- Be able to explain the following terminology
  - Null hypothesis  $H_0$  and alternative hypothesis  $H_A$
  - Test statistic  $s$
  - Null distribution  $f(s | H_0)$
  - Significance level  $\alpha$  and power
  - $p$ -value
- Be able to design and interpret the one-sample z-test
- Be able to explain the concept of  $p$ -hacking

# Today

## 1 Terminology

- Experiment and parameter of interest
- Null hypothesis and alternative hypothesis
- Test statistic
- Null distribution  $f(s \mid H_0)$
- Significance level  $\alpha$ , power and  $p$ -value

## 2 Example

## 3 $p$ -hacking

## 4 Summary



# Example

If you control the diet of your ducks, they lose 2.1 kg after one month on average

## Example

If you control the diet of your ducks, they lose 2.1 kg after one month on average

- Company A has developed a drug D to help ducks lose weight. They claim that **on average** the drug works better than diet control

## Example

If you control the diet of your ducks, they lose 2.1 kg after one month on average

- Company A has developed a drug D to help ducks lose weight. They claim that **on average** the drug works better than diet control
- Company B has developed a drug E and they claim that drug E is more effective than drug D **on average**

## Example

If you control the diet of your ducks, they lose 2.1 kg after one month on average

- Company A has developed a drug D to help ducks lose weight. They claim that **on average** the drug works better than diet control
- Company B has developed a drug E and they claim that drug E is more effective than drug D **on average**

You need to help your chonker ducks lose weight. Which drug do you buy? Or should you just control their diet?



## Example

If you control the diet of your ducks, they lose 2.1 kg after one month on average

- Company A has developed a drug D to help ducks lose weight. They claim that **on average** the drug works better than diet control
- Company B has developed a drug E and they claim that drug E is more effective than drug D **on average**

You need to help your chonker ducks lose weight. Which drug do you buy? Or should you just control their diet?

- If company A tested drug D on 30 ducks and the average weight loss after one month is 2.2 kg, would you buy drug D instead of regular diet control?

## Example

If you control the diet of your ducks, they lose 2.1 kg after one month on average

- Company A has developed a drug D to help ducks lose weight. They claim that **on average** the drug works better than diet control
- Company B has developed a drug E and they claim that drug E is more effective than drug D **on average**

You need to help your chonker ducks lose weight. Which drug do you buy? Or should you just control their diet?

- If company A tested drug D on 30 ducks and the average weight loss after one month is 2.2 kg, would you buy drug D instead of regular diet control?
- What if company A tested drug D on 30 ducks and the average weight loss after one month is 2.3 kg? Would you buy drug D instead of regular diet control in this case?

## Example

If you control the diet of your ducks, they lose 2.1 kg after one month on average

- Company A has developed a drug D to help ducks lose weight. They claim that **on average** the drug works better than diet control
- Company B has developed a drug E and they claim that drug E is more effective than drug D **on average**

You need to help your chonker ducks lose weight. Which drug do you buy? Or should you just control their diet?

- If company A tested drug D on 30 ducks and the average weight loss after one month is 2.2 kg, would you buy drug D instead of regular diet control?
- What if company A tested drug D on 30 ducks and the average weight loss after one month is 2.3 kg? Would you buy drug D instead of regular diet control in this case?
- What if company A tested drug D on 100 ducks and the average weight loss after one month is 2.3 kg?

## Example

If you control the diet of your ducks, they lose 2.1 kg after one month on average

- Company A has developed a drug D to help ducks lose weight. They claim that **on average** the drug works better than diet control
- Company B has developed a drug E and they claim that drug E is more effective than drug D **on average**

You need to help your chonker ducks lose weight. Which drug do you buy? Or should you just control their diet?

- If company A tested drug D on 30 ducks and the average weight loss after one month is 2.2 kg, would you buy drug D instead of regular diet control?
- What if company A tested drug D on 30 ducks and the average weight loss after one month is 2.3 kg? Would you buy drug D instead of regular diet control in this case?
- What if company A tested drug D on 100 ducks and the average weight loss after one month is 2.3 kg?
- Now company B tested drug E on 30 ducks and the average weight loss after one month is 2.5 kg, while drug D results in 2.3 kg weight loss with the same setup, would you buy drug E instead of drug D?

## Example

If you control the diet of your ducks, they lose 2.1 kg after one month on average

- Company A has developed a drug D to help ducks lose weight. They claim that **on average** the drug works better than diet control
- Company B has developed a drug E and they claim that drug E is more effective than drug D **on average**

You need to help your chonker ducks lose weight. Which drug do you buy? Or should you just control their diet?

- If company A tested drug D on 30 ducks and the average weight loss after one month is 2.2 kg, would you buy drug D instead of regular diet control?
- What if company A tested drug D on 30 ducks and the average weight loss after one month is 2.3 kg? Would you buy drug D instead of regular diet control in this case?
- What if company A tested drug D on 100 ducks and the average weight loss after one month is 2.3 kg?
- Now company B tested drug E on 30 ducks and the average weight loss after one month is 2.5 kg, while drug D results in 2.3 kg weight loss with the same setup, would you buy drug E instead of drug D?

How would you make your decision?

# Hypothesis

- **Hypothesis:** a hypothesis is a proposed explanation for a phenomenon (wikipedia)

# Hypothesis

- **Hypothesis:** a hypothesis is a proposed explanation for a phenomenon (wikipedia)
- **Statistical hypothesis:** a proposed distribution that explains a set of random variables

# Hypothesis

- **Hypothesis:** a hypothesis is a proposed explanation for a phenomenon (wikipedia)
- **Statistical hypothesis:** a proposed distribution that explains a set of random variables
- **Hypothesis testing in statistics:** we want to decide if it is likely that a random variable follows the distribution proposed by the statistical hypothesis



# Hypothesis

- **Hypothesis:** a hypothesis is a proposed explanation for a phenomenon (wikipedia)
- **Statistical hypothesis:** a proposed distribution that explains a set of random variables
- **Hypothesis testing in statistics:** we want to decide if it is likely that a random variable follows the distribution proposed by the statistical hypothesis
  - The test is based on sample statistics, which are computed from data

# Hypothesis

- **Hypothesis:** a hypothesis is a proposed explanation for a phenomenon (wikipedia)
- **Statistical hypothesis:** a proposed distribution that explains a set of random variables
- **Hypothesis testing in statistics:** we want to decide if it is likely that a random variable follows the distribution proposed by the statistical hypothesis
  - The test is based on sample statistics, which are computed from data
  - Hypothesis + data  $\rightarrow$  decision on rejecting/not rejecting the hypothesis

# Hypothesis testing: a list to go through

- A “boring” statement
- Experiment
- Data  $x$ , random variable  $X$
- Parameter of interest  $\theta$
- Parameter estimate  $\hat{\theta}$
- Null hypothesis  $H_0$
- Alternative hypothesis  $H_A$
- Test statistic  $s$
- Null distribution  $f(s \mid H_0)$
- Significance level  $\alpha$
- $p$ -value

## Experiment and parameter of interest

## Experiment design

- Before formulating the statistical hypothesis, we need a **“boring” statement**: a claim that we would like to test against, e.g. drug D is not more effective than regular diet on average; drug E works the same as drug D on average

## Experiment design

- Before formulating the statistical hypothesis, we need a **“boring” statement**: a claim that we would like to test against, e.g. drug D is not more effective than regular diet on average; drug E works the same as drug D on average
- How do we test the “boring” statement?

## Experiment design

- Before formulating the statistical hypothesis, we need a **“boring” statement**: a claim that we would like to test against, e.g. drug D is not more effective than regular diet on average; drug E works the same as drug D on average
- How do we test the “boring” statement? We design and run **experiments** to collect evidence (**data**)

## Experiment design

- Before formulating the statistical hypothesis, we need a **“boring” statement**: a claim that we would like to test against, e.g. drug D is not more effective than regular diet on average; drug E works the same as drug D on average
- How do we test the “boring” statement? We design and run **experiments** to collect evidence (**data**)
- Example 1: recall if you control the diet of your ducks, they lose 2.1 kg after one month on average
  - A **“boring” statement**: drug D is not more effective than regular diet on average



## Experiment design

- Before formulating the statistical hypothesis, we need a **“boring” statement**: a claim that we would like to test against, e.g. drug D is not more effective than regular diet on average; drug E works the same as drug D on average
- How do we test the “boring” statement? We design and run **experiments** to collect evidence (**data**)
- Example 1: recall if you control the diet of your ducks, they lose 2.1 kg after one month on average
  - A **“boring” statement**: drug D is not more effective than regular diet on average
  - **Experiment** (5 sec):

# Experiment design

- Before formulating the statistical hypothesis, we need a **“boring” statement**: a claim that we would like to test against, e.g. drug D is not more effective than regular diet on average; drug E works the same as drug D on average
- How do we test the “boring” statement? We design and run **experiments** to collect evidence (**data**)
- Example 1: recall if you control the diet of your ducks, they lose 2.1 kg after one month on average
  - A **“boring” statement**: drug D is not more effective than regular diet on average
  - **Experiment** (5 sec): test drug D on  $N$  chonker ducks and record the average weight loss after one month

# Experiment design

- Before formulating the statistical hypothesis, we need a **“boring” statement**: a claim that we would like to test against, e.g. drug D is not more effective than regular diet on average; drug E works the same as drug D on average
- How do we test the “boring” statement? We design and run **experiments** to collect evidence (**data**)
- Example 1: recall if you control the diet of your ducks, they lose 2.1 kg after one month on average
  - A **“boring” statement**: drug D is not more effective than regular diet on average
  - **Experiment** (5 sec): test drug D on  $N$  chonker ducks and record the average weight loss after one month
  - **Data** and **random variable** (5 sec):

# Experiment design

- Before formulating the statistical hypothesis, we need a **“boring” statement**: a claim that we would like to test against, e.g. drug D is not more effective than regular diet on average; drug E works the same as drug D on average
- How do we test the “boring” statement? We design and run **experiments** to collect evidence (**data**)
- Example 1: recall if you control the diet of your ducks, they lose 2.1 kg after one month on average
  - **A “boring” statement**: drug D is not more effective than regular diet on average
  - **Experiment** (5 sec): test drug D on  $N$  chonker ducks and record the average weight loss after one month
  - **Data** and **random variable** (5 sec): data -  $x_i$ ; weight loss after one month for  $i = 1, \dots, N$ ; random variable -  $X_i$  i.i.d.

# Experiment design

- Before formulating the statistical hypothesis, we need a **“boring” statement**: a claim that we would like to test against, e.g. drug D is not more effective than regular diet on average; drug E works the same as drug D on average
- How do we test the “boring” statement? We design and run **experiments** to collect evidence (**data**)
- Example 1: recall if you control the diet of your ducks, they lose 2.1 kg after one month on average
  - **A “boring” statement**: drug D is not more effective than regular diet on average
  - **Experiment** (5 sec): test drug D on  $N$  chonker ducks and record the average weight loss after one month
  - **Data** and **random variable** (5 sec): data -  $x_i$  weight loss after one month for  $i = 1, \dots, N$ ; random variable -  $X_i$  i.i.d.
  - **Parameter of interest** (5 sec):

# Experiment design

- Before formulating the statistical hypothesis, we need a **“boring” statement**: a claim that we would like to test against, e.g. drug D is not more effective than regular diet on average; drug E works the same as drug D on average
- How do we test the “boring” statement? We design and run **experiments** to collect evidence (**data**)
- Example 1: recall if you control the diet of your ducks, they lose 2.1 kg after one month on average
  - **A “boring” statement**: drug D is not more effective than regular diet on average
  - **Experiment** (5 sec): test drug D on  $N$  chonker ducks and record the average weight loss after one month
  - **Data** and **random variable** (5 sec): data -  $x_i$  weight loss after one month for  $i = 1, \dots, N$ ; random variable -  $X_i$  i.i.d.
  - **Parameter of interest** (5 sec): the average weight loss  $\mu_D$

# Experiment design

- Before formulating the statistical hypothesis, we need a **“boring” statement**: a claim that we would like to test against, e.g. drug D is not more effective than regular diet on average; drug E works the same as drug D on average
- How do we test the “boring” statement? We design and run **experiments** to collect evidence (**data**)
- Example 1: recall if you control the diet of your ducks, they lose 2.1 kg after one month on average
  - A **“boring” statement**: drug D is not more effective than regular diet on average
  - **Experiment** (5 sec): test drug D on  $N$  chonker ducks and record the average weight loss after one month
  - **Data** and **random variable** (5 sec): data -  $x_i$  weight loss after one month for  $i = 1, \dots, N$ ; random variable -  $X_i$  i.i.d.
  - **Parameter of interest** (5 sec): the average weight loss  $\mu_D$
  - **Parameter estimate** (5 sec):

# Experiment design

- Before formulating the statistical hypothesis, we need a **“boring” statement**: a claim that we would like to test against, e.g. drug D is not more effective than regular diet on average; drug E works the same as drug D on average
- How do we test the “boring” statement? We design and run **experiments** to collect evidence (**data**)
- Example 1: recall if you control the diet of your ducks, they lose 2.1 kg after one month on average
  - **A “boring” statement**: drug D is not more effective than regular diet on average
  - **Experiment** (5 sec): test drug D on  $N$  chonker ducks and record the average weight loss after one month
  - **Data** and **random variable** (5 sec): data -  $x_i$  weight loss after one month for  $i = 1, \dots, N$ ; random variable -  $X_i$  i.i.d.
  - **Parameter of interest** (5 sec): the average weight loss  $\mu_D$
  - **Parameter estimate** (5 sec):  $\hat{\mu}_D = \bar{x} = \frac{1}{N} \sum_{i=1}^N x_i$



# Experiment design

- Before formulating the statistical hypothesis, we need a **“boring” statement**: a claim that we would like to test against, e.g. drug D is not more effective than regular diet on average; drug E works the same as drug D on average
- How do we test the “boring” statement? We design and run **experiments** to collect evidence (**data**)
- Example 1: recall if you control the diet of your ducks, they lose 2.1 kg after one month on average
  - **A “boring” statement**: drug D is not more effective than regular diet on average
  - **Experiment** (5 sec): test drug D on  $N$  chonker ducks and record the average weight loss after one month
  - **Data** and **random variable** (5 sec): data -  $x_i$  weight loss after one month for  $i = 1, \dots, N$ ; random variable -  $X_i$  i.i.d.
  - **Parameter of interest** (5 sec): the average weight loss  $\mu_D$
  - **Parameter estimate** (5 sec):  $\hat{\mu}_D = \bar{x} = \frac{1}{N} \sum_{i=1}^N x_i$

Then we can use  $\bar{x}$  to approximate  $\mu_D$  and check if it is greater than diet control (2.1 kg)

## Experiment design (cont.)

- Example 2:
  - A **“boring” statement**: drug E and drug D work the same on average

## Experiment design (cont.)

- Example 2:
  - A “boring” statement: drug E and drug D work the same on average
  - Experiment (5 sec):

## Experiment design (cont.)

- Example 2:
  - **A “boring” statement**: drug E and drug D work the same on average
  - **Experiment** (5 sec): test drug D on  $N_D$  chonker ducks and record the average weight loss after one month; test drug E on another  $N_E$  chonker ducks and record the average weight loss after one month

## Experiment design (cont.)

- Example 2:
  - **A “boring” statement**: drug E and drug D work the same on average
  - **Experiment** (5 sec): test drug D on  $N_D$  chonker ducks and record the average weight loss after one month; test drug E on another  $N_E$  chonker ducks and record the average weight loss after one month
  - **Data** and **random variable** (5 sec):

## Experiment design (cont.)

- Example 2:
  - **A “boring” statement**: drug E and drug D work the same on average
  - **Experiment** (5 sec): test drug D on  $N_D$  chonker ducks and record the average weight loss after one month; test drug E on another  $N_E$  chonker ducks and record the average weight loss after one month
  - **Data** and **random variable** (5 sec): data -  $x_i$  weight loss using drug D after one month for  $i = 1, \dots, N_D$ ; random variable -  $X_i$  i.i.d.; likewise, we have data  $y_j$  and random variable  $Y_j$  for drug E

## Experiment design (cont.)

- Example 2:
  - **A “boring” statement**: drug E and drug D work the same on average
  - **Experiment** (5 sec): test drug D on  $N_D$  chonker ducks and record the average weight loss after one month; test drug E on another  $N_E$  chonker ducks and record the average weight loss after one month
  - **Data** and **random variable** (5 sec): data -  $x_i$  weight loss using drug D after one month for  $i = 1, \dots, N_D$ ; random variable -  $X_i$  i.i.d.; likewise, we have data  $y_j$  and random variable  $Y_j$  for drug E
  - **Parameter of interest** (5 secs):

## Experiment design (cont.)

- Example 2:
  - **A “boring” statement**: drug E and drug D work the same on average
  - **Experiment** (5 sec): test drug D on  $N_D$  chonker ducks and record the average weight loss after one month; test drug E on another  $N_E$  chonker ducks and record the average weight loss after one month
  - **Data** and **random variable** (5 sec): data -  $x_i$  weight loss using drug D after one month for  $i = 1, \dots, N_D$ ; random variable -  $X_i$  i.i.d.; likewise, we have data  $y_j$  and random variable  $Y_j$  for drug E
  - **Parameter of interest** (5 secs): the average weight loss  $\mu_D$  and  $\mu_E$  for drug  $D$  and  $E$ , respectively



## Experiment design (cont.)

- Example 2:
  - **A “boring” statement**: drug E and drug D work the same on average
  - **Experiment** (5 sec): test drug D on  $N_D$  chonker ducks and record the average weight loss after one month; test drug E on another  $N_E$  chonker ducks and record the average weight loss after one month
  - **Data** and **random variable** (5 sec): data -  $x_i$  weight loss using drug D after one month for  $i = 1, \dots, N_D$ ; random variable -  $X_i$  i.i.d.; likewise, we have data  $y_j$  and random variable  $Y_j$  for drug E
  - **Parameter of interest** (5 secs): the average weight loss  $\mu_D$  and  $\mu_E$  for drug  $D$  and  $E$ , respectively
  - **Parameter estimate** (5 secs):

## Experiment design (cont.)

- Example 2:

- **A “boring” statement**: drug E and drug D work the same on average
- **Experiment** (5 sec): test drug D on  $N_D$  chonker ducks and record the average weight loss after one month; test drug E on another  $N_E$  chonker ducks and record the average weight loss after one month
- **Data** and **random variable** (5 sec): data -  $x_i$  weight loss using drug D after one month for  $i = 1, \dots, N_D$ ; random variable -  $X_i$  i.i.d.; likewise, we have data  $y_j$  and random variable  $Y_j$  for drug E
- **Parameter of interest** (5 secs): the average weight loss  $\mu_D$  and  $\mu_E$  for drug  $D$  and  $E$ , respectively
- **Parameter estimate** (5 secs):  $\hat{\mu}_D = \bar{x} = \frac{1}{N_D} \sum_{i=1}^{N_D} x_i$  and  $\hat{\mu}_E = \bar{y} = \frac{1}{N_E} \sum_{j=1}^{N_E} y_j$

## Experiment design (cont.)

- Example 2:
  - **A “boring” statement**: drug E and drug D work the same on average
  - **Experiment** (5 sec): test drug D on  $N_D$  chonker ducks and record the average weight loss after one month; test drug E on another  $N_E$  chonker ducks and record the average weight loss after one month
  - **Data** and **random variable** (5 sec): data -  $x_i$  weight loss using drug D after one month for  $i = 1, \dots, N_D$ ; random variable -  $X_i$  i.i.d.; likewise, we have data  $y_j$  and random variable  $Y_j$  for drug E
  - **Parameter of interest** (5 secs): the average weight loss  $\mu_D$  and  $\mu_E$  for drug D and E, respectively
  - **Parameter estimate** (5 secs):  $\hat{\mu}_D = \bar{x} = \frac{1}{N_D} \sum_{i=1}^{N_D} x_i$  and  $\hat{\mu}_E = \bar{y} = \frac{1}{N_E} \sum_{j=1}^{N_E} y_j$

Then we use  $\bar{x}$  and  $\bar{y}$  to approximate  $\mu_D$  and  $\mu_E$  to see if they are the same

## Experiment design (cont.)

- We make our decision by observing data; if the evidence does not support the “boring” statement, we **reject the statement**; otherwise, we **do not reject the statement**

## Experiment design (cont.)

- We make our decision by observing data; if the evidence does not support the “boring” statement, we **reject the statement**; otherwise, we **do not reject the statement**
- But we can never prove or accept the statement - we can only **reject** a statement by showing counterexamples

## Experiment design (cont.)

- We make our decision by observing data; if the evidence does not support the “boring” statement, we **reject the statement**; otherwise, we **do not reject the statement**
- But we can never prove or accept the statement - we can only **reject** a statement by showing counterexamples
- The logic here is: if a statement is true, then the evidence should support the statement

## Experiment design (cont.)

- We make our decision by observing data; if the evidence does not support the “boring” statement, we **reject the statement**; otherwise, we **do not reject the statement**
- But we can never prove or accept the statement - we can only **reject** a statement by showing counterexamples
- The logic here is: if a statement is true, then the evidence should support the statement  $\iff$  if the evidence does not support the statement, the statement is considered false

## Experiment design (cont.)

- We make our decision by observing data; if the evidence does not support the “boring” statement, we **reject the statement**; otherwise, we **do not reject the statement**
- But we can never prove or accept the statement - we can only **reject** a statement by showing counterexamples
- The logic here is: if a statement is true, then the evidence should support the statement  $\iff$  if the evidence does not support the statement, the statement is considered false  $\niff$  if the evidence supports the statement, the statement must be true



# Null hypothesis and alternative hypothesis

# Hypotheses $H_0$ and $H_A$

- **Statistical hypothesis:** a proposed distribution

# Hypotheses $H_0$ and $H_A$

- **Statistical hypothesis**: a proposed distribution - a statement about the **parameter of interest**

## Hypotheses $H_0$ and $H_A$

- **Statistical hypothesis**: a proposed distribution - a statement about the **parameter of interest**
- **Null hypothesis  $H_0$** : the “boring” statement translated into a mathematical expression

# Hypotheses $H_0$ and $H_A$

- **Statistical hypothesis**: a proposed distribution - a statement about the **parameter of interest**
- **Null hypothesis  $H_0$** : the “boring” statement translated into a mathematical expression
  - Example 1: drug D is not more effective than regular diet on average

$$H_0 : \mu_D = 2.1$$

# Hypotheses $H_0$ and $H_A$

- **Statistical hypothesis**: a proposed distribution - a statement about the **parameter of interest**
- **Null hypothesis  $H_0$** : the “boring” statement translated into a mathematical expression
  - Example 1: drug D is not more effective than regular diet on average

$$H_0 : \mu_D = 2.1$$

- Example 2: drug E and drug D work the same on average (5 sec)

# Hypotheses $H_0$ and $H_A$

- **Statistical hypothesis**: a proposed distribution - a statement about the **parameter of interest**
- **Null hypothesis  $H_0$** : the “boring” statement translated into a mathematical expression
  - Example 1: drug D is not more effective than regular diet on average

$$H_0 : \mu_D = 2.1$$

- Example 2: drug E and drug D work the same on average (5 sec)

$$H_0 : \mu_D = \mu_E$$

# Hypotheses $H_0$ and $H_A$

- **Statistical hypothesis**: a proposed distribution - a statement about the **parameter of interest**
- **Null hypothesis  $H_0$** : the “boring” statement translated into a mathematical expression
  - Example 1: drug D is not more effective than regular diet on average

$$H_0 : \mu_D = 2.1$$

- Example 2: drug E and drug D work the same on average (5 sec)

$$H_0 : \mu_D = \mu_E$$

- **Alternative hypothesis  $H_A$** : a complementary alternative explanation to the “boring” statement



# Hypotheses $H_0$ and $H_A$

- **Statistical hypothesis**: a proposed distribution - a statement about the **parameter of interest**
- **Null hypothesis  $H_0$** : the “boring” statement translated into a mathematical expression
  - Example 1: drug D is not more effective than regular diet on average

$$H_0 : \mu_D = 2.1$$

- Example 2: drug E and drug D work the same on average (5 sec)

$$H_0 : \mu_D = \mu_E$$

- **Alternative hypothesis  $H_A$** : a complementary alternative explanation to the “boring” statement
  - Example 1: drug D is more effective than regular diet on average (5 sec)

# Hypotheses $H_0$ and $H_A$

- **Statistical hypothesis**: a proposed distribution - a statement about the **parameter of interest**
- **Null hypothesis  $H_0$** : the “boring” statement translated into a mathematical expression
  - Example 1: drug D is not more effective than regular diet on average

$$H_0 : \mu_D = 2.1$$

- Example 2: drug E and drug D work the same on average (5 sec)

$$H_0 : \mu_D = \mu_E$$

- **Alternative hypothesis  $H_A$** : a complementary alternative explanation to the “boring” statement
  - Example 1: drug D is more effective than regular diet on average (5 sec)

$$H_A : \mu_D > 2.1$$

# Hypotheses $H_0$ and $H_A$

- **Statistical hypothesis**: a proposed distribution - a statement about the **parameter of interest**
- **Null hypothesis  $H_0$** : the “boring” statement translated into a mathematical expression
  - Example 1: drug D is not more effective than regular diet on average

$$H_0 : \mu_D = 2.1$$

- Example 2: drug E and drug D work the same on average (5 sec)

$$H_0 : \mu_D = \mu_E$$

- **Alternative hypothesis  $H_A$** : a complementary alternative explanation to the “boring” statement
  - Example 1: drug D is more effective than regular diet on average (5 sec)

$$H_A : \mu_D > 2.1$$

- Example 2: drug E and drug D do not work the same on average (5 sec)

# Hypotheses $H_0$ and $H_A$

- **Statistical hypothesis**: a proposed distribution - a statement about the **parameter of interest**
- **Null hypothesis  $H_0$** : the “boring” statement translated into a mathematical expression
  - Example 1: drug D is not more effective than regular diet on average

$$H_0 : \mu_D = 2.1$$

- Example 2: drug E and drug D work the same on average (5 sec)

$$H_0 : \mu_D = \mu_E$$

- **Alternative hypothesis  $H_A$** : a complementary alternative explanation to the “boring” statement
  - Example 1: drug D is more effective than regular diet on average (5 sec)

$$H_A : \mu_D > 2.1$$

- Example 2: drug E and drug D do not work the same on average (5 sec)

$$H_A : \mu_D \neq \mu_E$$

## Hypotheses $H_0$ and $H_A$ (cont.)

Questions:

- Question 1: why are  $H_A : \mu_D > 2.1$  and  $H_0 : \mu_D = 2.1$  complementary to each other? What about  $H_A : \mu_D < 2.1$ ?

## Hypotheses $H_0$ and $H_A$ (cont.)

Questions:

- Question 1: why are  $H_A : \mu_D > 2.1$  and  $H_0 : \mu_D = 2.1$  complementary to each other? What about  $H_A : \mu_D < 2.1$ ?

Answer: an implicit assumption here is that  $\mu_D$  will not be smaller than 2.1

## Hypotheses $H_0$ and $H_A$ (cont.)

Questions:

- Question 1: why are  $H_A : \mu_D > 2.1$  and  $H_0 : \mu_D = 2.1$  complementary to each other? What about  $H_A : \mu_D < 2.1$ ?  
Answer: an implicit assumption here is that  $\mu_D$  will not be smaller than 2.1
- Question 2: can  $H_0$  and  $H_A$  be ANYTHING I want? Like a magic mirror!?

## Hypotheses $H_0$ and $H_A$ (cont.)

Questions:

- Question 1: why are  $H_A : \mu_D > 2.1$  and  $H_0 : \mu_D = 2.1$  complementary to each other? What about  $H_A : \mu_D < 2.1$ ?

Answer: an implicit assumption here is that  $\mu_D$  will not be smaller than 2.1

- Question 2: can  $H_0$  and  $H_A$  be ANYTHING I want? Like a magic mirror!?

Answer: no



## Hypotheses $H_0$ and $H_A$ (cont.)

Questions:

- Question 1: why are  $H_A : \mu_D > 2.1$  and  $H_0 : \mu_D = 2.1$  complementary to each other? What about  $H_A : \mu_D < 2.1$ ?

Answer: an implicit assumption here is that  $\mu_D$  will not be smaller than 2.1

- Question 2: can  $H_0$  and  $H_A$  be ANYTHING I want? Like a magic mirror!?

Answer: no

- Follow up question: what are the choices for  $H_0$  and  $H_A$ ?

## Choices for $H_0$

- In this course, we only deal with null hypotheses **with an equal sign** in them - only one fixed choice for the distribution proposed by  $H_0$

## Choices for $H_0$

- In this course, we only deal with null hypotheses **with an equal sign** in them - only one fixed choice for the distribution proposed by  $H_0$
- **Null hypothesis  $H_0$** : two cases

## Choices for $H_0$

- In this course, we only deal with null hypotheses **with an equal sign** in them - only one fixed choice for the distribution proposed by  $H_0$
- **Null hypothesis  $H_0$** : two cases
  - **One-sample test**: to test a data distribution against a theoretical probability distribution, i.e. for a given constant  $c$

$$H_0 : \theta = c$$

For example, is a binary classifier more accurate than random?

## Choices for $H_0$

- In this course, we only deal with null hypotheses **with an equal sign** in them - only one fixed choice for the distribution proposed by  $H_0$
- **Null hypothesis  $H_0$** : two cases
  - **One-sample test**: to test a data distribution against a theoretical probability distribution, i.e. for a given constant  $c$

$$H_0 : \theta = c$$

For example, is a binary classifier more accurate than random?  $H_0 : p = 50\%$

## Choices for $H_0$

- In this course, we only deal with null hypotheses **with an equal sign** in them - only one fixed choice for the distribution proposed by  $H_0$
- **Null hypothesis  $H_0$** : two cases
  - **One-sample test**: to test a data distribution against a theoretical probability distribution, i.e. for a given constant  $c$

$$H_0 : \theta = c$$

For example, is a binary classifier more accurate than random?  $H_0 : p = 50\%$

- **Two-sample test**: to test a data distribution against another data distribution, i.e.

$$H_0 : \theta_1 = \theta_2$$

For example, is classifier A better than classifier B?

# Choices for $H_0$

- In this course, we only deal with null hypotheses **with an equal sign** in them - only one fixed choice for the distribution proposed by  $H_0$
- **Null hypothesis  $H_0$** : two cases
  - **One-sample test**: to test a data distribution against a theoretical probability distribution, i.e. for a given constant  $c$

$$H_0 : \theta = c$$

For example, is a binary classifier more accurate than random?  $H_0 : p = 50\%$

- **Two-sample test**: to test a data distribution against another data distribution, i.e.

$$H_0 : \theta_1 = \theta_2$$

For example, is classifier A better than classifier B?  $H_0 : p_A = p_B$

## Choices for $H_0$

- In this course, we only deal with null hypotheses **with an equal sign** in them - only one fixed choice for the distribution proposed by  $H_0$
- **Null hypothesis  $H_0$** : two cases
  - **One-sample test**: to test a data distribution against a theoretical probability distribution, i.e. for a given constant  $c$

$$H_0 : \theta = c$$

For example, is a binary classifier more accurate than random?  $H_0 : p = 50\%$

- **Two-sample test**: to test a data distribution against another data distribution, i.e.

$$H_0 : \theta_1 = \theta_2$$

For example, is classifier A better than classifier B?  $H_0 : p_A = p_B$

- We have seen one-sample test and two-sample test in the Q-Q plot lecture
- In practice, you can narrow down your choice of hypotheses by making a Q-Q plot



## Choices for $H_A$

### Given

$$H_0 : \theta = \beta$$

where  $\beta$  can be either a constant (one-sample test) or a parameter from another data distribution (two-sample test)

- **Alternative hypothesis**  $H_A$ :  $H_A$  can be **one-tailed** or **two-tailed**
  - **One-tailed:**

$$H_A : \theta > \beta$$

or

$$H_A : \theta < \beta$$

- **Two-tailed:**

$$H_A : \theta \neq \beta \iff \theta < \beta \text{ or } \theta > \beta$$

## Summary: choices for $H_0$ and $H_A$

Putting everything together,

	One-sample test	Two-sample test
Two-tailed	$H_0 : \theta = c, H_A : \theta \neq c$	$H_0 : \theta_1 = \theta_2, H_A : \theta_1 \neq \theta_2$
One-tailed	$H_0 : \theta = c, H_A : \theta > c$	$H_0 : \theta_1 = \theta_2, H_A : \theta_1 > \theta_2$
	$H_0 : \theta = c, H_A : \theta < c$	$H_0 : \theta_1 = \theta_2, H_A : \theta_1 < \theta_2$

where  $\theta, \theta_1, \theta_2$  are the parameters of interest and  $c$  is a constant

## Test statistic

# Test statistic

- **Test statistic  $s$ , random variable  $S$** : the statistic used for testing the hypothesis

# Test statistic

- **Test statistic  $s$ , random variable  $S$** : the statistic used for testing the hypothesis
  - $s$  is the **observation**

# Test statistic

- **Test statistic  $s$ , random variable  $S$** : the statistic used for testing the hypothesis
  - $s$  is the **observation**
  - Given a set of parameters of interest and a set of estimates,  $s$  is typically a **standardized statistic** computed from the estimates

# Test statistic

- **Test statistic  $s$ , random variable  $S$** : the statistic used for testing the hypothesis
  - $s$  is the **observation**
  - Given a set of parameters of interest and a set of estimates,  $s$  is typically a **standardized statistic** computed from the estimates
  - **Purpose**: to compare  $s$  with a **standard distribution**, e.g. the standard Gaussian distribution  $\mathcal{N}(0, 1)$ , to see if it is likely that the standard distribution is the underlying distribution of  $S$ , i.e. if the null hypothesis is plausible

# Test statistic

- **Test statistic  $s$ , random variable  $S$** : the statistic used for testing the hypothesis
  - $s$  is the **observation**
  - Given a set of parameters of interest and a set of estimates,  $s$  is typically a **standardized statistic** computed from the estimates
  - **Purpose**: to compare  $s$  with a **standard distribution**, e.g. the standard Gaussian distribution  $\mathcal{N}(0, 1)$ , to see if it is likely that the standard distribution is the underlying distribution of  $S$ , i.e. if the null hypothesis is plausible
- What is needed for computing the test statistic?
  - Assumptions on random variables  $X_i$
  - We only need the null hypothesis  $H_0$  (not  $H_A$ ) to choose the test statistic



# Test statistic

- **Test statistic  $s$ , random variable  $S$** : the statistic used for testing the hypothesis
  - $s$  is the **observation**
  - Given a set of parameters of interest and a set of estimates,  $s$  is typically a **standardized statistic** computed from the estimates
  - **Purpose**: to compare  $s$  with a **standard distribution**, e.g. the standard Gaussian distribution  $\mathcal{N}(0, 1)$ , to see if it is likely that the standard distribution is the underlying distribution of  $S$ , i.e. if the null hypothesis is plausible
- What is needed for computing the test statistic?
  - Assumptions on random variables  $X_i$
  - We only need the null hypothesis  $H_0$  (not  $H_A$ ) to choose the test statistic

Note: in this course, we only deal with null hypothesis where we are able to express the PDF/PMF  $f(s \mid H_0)$ , i.e.  $H_0$  with an equal sign in them

## Test statistic (cont.)

Example 1. one-sample test

- **Data:**  $x_1, \dots, x_N$
- **Random variable:**  $X_1, \dots, X_N$  i.i.d. **Gaussian with known  $\sigma$**
- **Parameter of interest:**  $\mu_D$
- **Parameter estimate:**  $\bar{x}$
- **Null hypothesis:**  $H_0 : \mu_D = 2.1$
- **Test statistic:** **standardized  $\bar{x}$  assuming the null hypothesis**

## Test statistic (cont.)

Example 1. one-sample test

- **Data:**  $x_1, \dots, x_N$
- **Random variable:**  $X_1, \dots, X_N$  i.i.d. **Gaussian with known  $\sigma$**
- **Parameter of interest:**  $\mu_D$
- **Parameter estimate:**  $\bar{x}$
- **Null hypothesis:**  $H_0 : \mu_D = 2.1$
- **Test statistic:** **standardized  $\bar{x}$  assuming the null hypothesis**
  - Recall: what is **standardization**?

# Test statistic (cont.)

## Example 1. one-sample test

- **Data:**  $x_1, \dots, x_N$
- **Random variable:**  $X_1, \dots, X_N$  i.i.d. **Gaussian with known  $\sigma$**
- **Parameter of interest:**  $\mu_D$
- **Parameter estimate:**  $\bar{x}$
- **Null hypothesis:**  $H_0 : \mu_D = 2.1$
- **Test statistic:** **standardized  $\bar{x}$  assuming the null hypothesis**
  - Recall: what is **standardization**?
    - Random variable  $X$ :  $Y = \frac{X - \mu_X}{\sigma_X}$

## Test statistic (cont.)

### Example 1. one-sample test

- **Data:**  $x_1, \dots, x_N$
- **Random variable:**  $X_1, \dots, X_N$  i.i.d. **Gaussian with known  $\sigma$**
- **Parameter of interest:**  $\mu_D$
- **Parameter estimate:**  $\bar{x}$
- **Null hypothesis:**  $H_0 : \mu_D = 2.1$
- **Test statistic:** **standardized  $\bar{x}$  assuming the null hypothesis**
  - Recall: what is **standardization**?
    - Random variable  $X$ :  $Y = \frac{X - \mu_X}{\sigma_X}$
    - Data  $x$ :  $y = \frac{x - \mu_X}{\sigma_X}$

## Test statistic (cont.)

### Example 1. one-sample test

- **Data:**  $x_1, \dots, x_N$
- **Random variable:**  $X_1, \dots, X_N$  i.i.d. **Gaussian with known  $\sigma$**
- **Parameter of interest:**  $\mu_D$
- **Parameter estimate:**  $\bar{x}$
- **Null hypothesis:**  $H_0 : \mu_D = 2.1$
- **Test statistic:** **standardized  $\bar{x}$  assuming the null hypothesis**
  - Recall: what is **standardization**?
    - Random variable  $X$ :  $Y = \frac{X - \mu_X}{\sigma_X}$
    - Data  $x$ :  $y = \frac{x - \mu_X}{\sigma_X}$
  - Recall: what are we trying to do?

## Test statistic (cont.)

### Example 1. one-sample test

- **Data:**  $x_1, \dots, x_N$
- **Random variable:**  $X_1, \dots, X_N$  i.i.d. **Gaussian with known  $\sigma$**
- **Parameter of interest:**  $\mu_D$
- **Parameter estimate:**  $\bar{x}$
- **Null hypothesis:**  $H_0 : \mu_D = 2.1$
- **Test statistic:** **standardized  $\bar{x}$  assuming the null hypothesis**
  - Recall: what is **standardization**?
    - Random variable  $X$ :  $Y = \frac{X - \mu_X}{\sigma_X}$
    - Data  $x$ :  $y = \frac{x - \mu_X}{\sigma_X}$
  - Recall: what are we trying to do? - Decide how likely data follows **the distribution described by the null hypothesis**

## Test statistic (cont.)

### Example 1. one-sample test

- **Data:**  $x_1, \dots, x_N$
- **Random variable:**  $X_1, \dots, X_N$  i.i.d. **Gaussian with known  $\sigma$**
- **Parameter of interest:**  $\mu_D$
- **Parameter estimate:**  $\bar{x}$
- **Null hypothesis:**  $H_0 : \mu_D = 2.1$
- **Test statistic:** **standardized  $\bar{x}$  assuming the null hypothesis**
  - Recall: what is **standardization**?
    - Random variable  $X$ :  $Y = \frac{X - \mu_X}{\sigma_X}$
    - Data  $x$ :  $y = \frac{x - \mu_X}{\sigma_X}$
  - Recall: what are we trying to do? - Decide how likely data follows **the distribution described by the null hypothesis?**
  - What is the **distribution described by the null hypothesis?**



## Test statistic (cont.)

### Example 1. one-sample test

- **Data:**  $x_1, \dots, x_N$
- **Random variable:**  $X_1, \dots, X_N$  i.i.d. **Gaussian with known  $\sigma$**
- **Parameter of interest:**  $\mu_D$
- **Parameter estimate:**  $\bar{x}$
- **Null hypothesis:**  $H_0 : \mu_D = 2.1$
- **Test statistic:** **standardized  $\bar{x}$  assuming the null hypothesis**
  - Recall: what is **standardization**?
    - Random variable  $X$ :  $Y = \frac{X - \mu_X}{\sigma_X}$
    - Data  $x$ :  $y = \frac{x - \mu_X}{\sigma_X}$
  - Recall: what are we trying to do? - Decide how likely data follows **the distribution described by the null hypothesis**
  - What is the **distribution described by the null hypothesis**?
    - Gaussian distribution with standard deviation  $\sigma$  and mean  $\mu_D = 2.1$

## Test statistic (cont.)

### Example 1. one-sample test

- **Data:**  $x_1, \dots, x_N$
- **Random variable:**  $X_1, \dots, X_N$  i.i.d. **Gaussian with known  $\sigma$**
- **Parameter of interest:**  $\mu_D$
- **Parameter estimate:**  $\bar{x}$
- **Null hypothesis:**  $H_0 : \mu_D = 2.1$
- **Test statistic:** **standardized  $\bar{x}$  assuming the null hypothesis**
  - Recall: what is **standardization**?
    - Random variable  $X$ :  $Y = \frac{X - \mu_X}{\sigma_X}$
    - Data  $x$ :  $y = \frac{x - \mu_X}{\sigma_X}$
  - Recall: what are we trying to do? - Decide how likely data follows **the distribution described by the null hypothesis**?
  - What is the **distribution described by the null hypothesis**?
    - Gaussian distribution with standard deviation  $\sigma$  and mean  $\mu_D = 2.1$
  - **Assuming the null hypothesis:** data are assumed to be generated from the distribution described by the null hypothesis

## Test statistic (cont.)

### Example 1. one-sample test

- **Data:**  $x_1, \dots, x_N$
- **Random variable:**  $X_1, \dots, X_N$  i.i.d. **Gaussian with known  $\sigma$**
- **Parameter of interest:**  $\mu_D$
- **Parameter estimate:**  $\bar{x}$
- **Null hypothesis:**  $H_0 : \mu_D = 2.1$
- **Test statistic:** **standardized  $\bar{x}$  assuming the null hypothesis**
  - Recall: what is **standardization**?
    - Random variable  $X$ :  $Y = \frac{X - \mu_X}{\sigma_X}$
    - Data  $x$ :  $y = \frac{x - \mu_X}{\sigma_X}$
  - Recall: what are we trying to do? - Decide how likely data follows **the distribution described by the null hypothesis?**
  - What is the **distribution described by the null hypothesis?**
    - Gaussian distribution with standard deviation  $\sigma$  and mean  $\mu_D = 2.1$
  - **Assuming the null hypothesis:** data are assumed to be generated from the distribution described by the null hypothesis -  $X_i \sim \mathcal{N}(\mu_D, \sigma^2)$

## Test statistic (cont.)

### Example 1. one-sample test

- **Data:**  $x_1, \dots, x_N$
- **Random variable:**  $X_1, \dots, X_N$  i.i.d. **Gaussian with known  $\sigma$**
- **Parameter of interest:**  $\mu_D$
- **Parameter estimate:**  $\bar{x}$
- **Null hypothesis:**  $H_0 : \mu_D = 2.1$
- **Test statistic:** **standardized  $\bar{x}$  assuming the null hypothesis**
  - Recall: what is **standardization**?
    - Random variable  $X$ :  $Y = \frac{X - \mu_X}{\sigma_X}$
    - Data  $x$ :  $y = \frac{x - \mu_X}{\sigma_X}$
  - Recall: what are we trying to do? - Decide how likely data follows **the distribution described by the null hypothesis?**
  - What is the **distribution described by the null hypothesis?**
    - Gaussian distribution with standard deviation  $\sigma$  and mean  $\mu_D = 2.1$
  - **Assuming the null hypothesis:** data are assumed to be generated from the distribution described by the null hypothesis -  $X_i \sim \mathcal{N}(\mu_D, \sigma^2)$

**Standardize  $\bar{x}$  (15 sec)**

## Test statistic (cont.)

### Example 1. one-sample test

- **Data:**  $x_1, \dots, x_N$
- **Random variable:**  $X_1, \dots, X_N$  i.i.d. **Gaussian with known  $\sigma$**
- **Parameter of interest:**  $\mu_D$
- **Parameter estimate:**  $\bar{x}$
- **Null hypothesis:**  $H_0 : \mu_D = 2.1$
- **Test statistic:** **standardized  $\bar{x}$  assuming the null hypothesis**
  - Recall: what is **standardization**?
    - Random variable  $X$ :  $Y = \frac{X - \mu_X}{\sigma_X}$
    - Data  $x$ :  $y = \frac{x - \mu_X}{\sigma_X}$
  - Recall: what are we trying to do? - Decide how likely data follows **the distribution described by the null hypothesis**?
  - What is the **distribution described by the null hypothesis**?
    - Gaussian distribution with standard deviation  $\sigma$  and mean  $\mu_D = 2.1$
  - **Assuming the null hypothesis:** data are assumed to be generated from the distribution described by the null hypothesis -  $X_i \sim \mathcal{N}(\mu_D, \sigma^2)$

**Standardize  $\bar{x}$  (15 sec)**

$$z = \frac{\bar{x} - 2.1}{\sigma / \sqrt{N}}$$

## Test statistic (cont.)

### Example 2. two-sample test

- **Data:**  $x_1, \dots, x_{N_D}$  and  $y_1, \dots, y_{N_E}$
- **Random variable:**  $X_1, \dots, X_{N_D}$  i.i.d. **Gaussian with known  $\sigma_D$** ;  $Y_1, \dots, Y_{N_E}$  i.i.d. **Gaussian with known  $\sigma_E$** ;  $X_i$  and  $Y_j$  independent
- **Parameter of interest:**  $\mu_D, \mu_E$
- **Parameter estimate:**  $\bar{x}, \bar{y}$
- **Null hypothesis:**  $H_0 : \mu_D = \mu_E \iff H_0 : \mu_D - \mu_E = 0$
- **Test statistic:** standardized  $\bar{x} - \bar{y}$  assuming the null hypothesis

$$z = \frac{\bar{x} - \bar{y}}{\sqrt{\sigma_D^2/N_D + \sigma_E^2/N_E}} \quad (\text{explained later})$$


Null distribution  $f(s \mid H_0)$

# Null distribution

- **Null distribution  $f(s | H_0)$ :** the distribution of the test statistic given the null hypothesis
- Example:
  - **Data:**  $x_1, \dots, x_N$
  - **Random variable:**  $X_1, \dots, X_N$  i.i.d. Gaussian with known  $\sigma$
  - **Parameter of interest:**  $\mu$
  - **Parameter estimate:**  $\bar{x}$
  - **Null hypothesis:**  $H_0 : \mu = \mu_0$
  - **Test statistic:**

$$z = \frac{\bar{x} - \mu_0}{\sigma/\sqrt{N}}$$

- **Null distribution:** standard Gaussian distribution






## Significance level $\alpha$ , power and $p$ -value

# Significance level

Given a null hypothesis  $H_0 : \mu = 2.1$  and the null distribution  $f(s | H_0)$ , we decide if we reject the hypothesis or not by observing data

- Run some experiments and collect data  $x_1, \dots, x_N$
- Estimate the parameter of interest  $\hat{\theta}$ , e.g.  $\hat{\mu} = \bar{x} = \frac{1}{N} \sum_{i=1}^N x_i$
- Standardize  $\hat{\theta}$  assuming  $H_0$  to compute the test statistic, e.g.

$$z = \frac{\bar{x} - 2.1}{\sigma/\sqrt{N}} = 4.0$$




- Does this evidence support the hypothesis  $H_0$ ?

# Significance level

Given a null hypothesis  $H_0 : \mu = 2.1$  and the null distribution  $f(s | H_0)$ , we decide if we reject the hypothesis or not by observing data

- Run some experiments and collect data  $x_1, \dots, x_N$
- Estimate the parameter of interest  $\hat{\theta}$ , e.g.  $\hat{\mu} = \bar{x} = \frac{1}{N} \sum_{i=1}^N x_i$
- Standardize  $\hat{\theta}$  assuming  $H_0$  to compute the test statistic, e.g.


$$z = \frac{\bar{x} - 2.1}{\sigma/\sqrt{N}} = 4.0$$



- Does this evidence support the hypothesis  $H_0$ ? Probably not since it's so far away from the center?


## Significance level (cont.)

- What about this observation?



## Significance level (cont.)


- What about this observation?



- To be able to answer the question, you need to decide where you draw the line

## Significance level (cont.)


- What about this observation?




- To be able to answer the question, you need to decide where you draw the line - define a **rejection region** by choosing a significance level

## Significance level (cont.)

- What about this observation?




- To be able to answer the question, you need to decide where you draw the line - define a **rejection region** by choosing a significance level
- Significance level  $\alpha$** : red area under the curve




In these three images,  $\alpha = 0.05$

## Significance level (cont.)

- What about this observation?



- To be able to answer the question, you need to decide where you draw the line - define a **rejection region** by choosing a significance level
- Significance level  $\alpha$** : red area under the curve




In these three images,  $\alpha = 0.05$

More conservative  $\Rightarrow$  less probable to reject  $H_0$ , which indicates a smaller rejection region




## Significance level (cont.)

- What about this observation?



- To be able to answer the question, you need to decide where you draw the line - define a **rejection region** by choosing a significance level
- Significance level  $\alpha$** : red area under the curve



In these three images,  $\alpha = 0.05$

More conservative  $\Rightarrow$  less probable to reject  $H_0$ , which indicates a smaller rejection region

Two-tailed  $H_A$  is more conservative


## Significance level (cont.)

What is needed for choosing a meaningful  $\alpha$ ?

- Null distribution
- $H_A$  one-tailed or two-tailed

# Interpretation of $\alpha$


- $\alpha = P(\text{reject } H_0 \mid H_0 \text{ is true})$  - the probability of making such a mistake



- The rejection region indicates that  $H_0$  is **unlikely**, but the probability is not zero
- It is possible that  $H_0$  is true, but our observation happens to fall in the rejection region
- If  $H_0$  is true and our observation falls in the rejection region, we will **mistakenly** reject  $H_0$
- The probability of making this type of mistakes is  $\alpha$
- Similar to the confidence interval,  $1 - \alpha$  is called the **confidence level**

# Interpretation of $\alpha$


- $\alpha = P(\text{reject } H_0 \mid H_0 \text{ is true})$  - the probability of making such a mistake



- The rejection region indicates that  $H_0$  is **unlikely**, but the probability is not zero
- It is possible that  $H_0$  is true, but our observation happens to fall in the rejection region
- If  $H_0$  is true and our observation falls in the rejection region, we will **mistakenly** reject  $H_0$
- The probability of making this type of mistakes is  $\alpha$
- Similar to the confidence interval,  $1 - \alpha$  is called the **confidence level** - “with 95% confidence, rejecting  $H_0$  is the right thing to do”

# Interpretation of $\alpha$

- $\alpha = P(\text{reject } H_0 \mid H_0 \text{ is true})$  - the probability of making such a mistake



- The rejection region indicates that  $H_0$  is **unlikely**, but the probability is not zero
- It is possible that  $H_0$  is true, but our observation happens to fall in the rejection region
- If  $H_0$  is true and our observation falls in the rejection region, we will **mistakenly** reject  $H_0$
- The probability of making this type of mistakes is  $\alpha$
- Similar to the confidence interval,  $1 - \alpha$  is called the **confidence level** - “with 95% confidence, rejecting  $H_0$  is the right thing to do”
- Define the significance level **before you run the experiments** so that you can't cheat!

# Significance level and power

- Contingency table:

	$y = H_A$	$y = H_0$
$\hat{y} = \text{reject } H_0$	TP	FP (Type I error)
$\hat{y} = \text{do not reject } H_0$	FN (Type II error)	TN

# Significance level and power

- Contingency table:

	$y = H_A$	$y = H_0$
$\hat{y} = \text{reject } H_0$	TP	FP (Type I error)
$\hat{y} = \text{do not reject } H_0$	FN (Type II error)	TN

- Significance level  $\alpha$ : incorrectly rejecting  $H_0$

$$\alpha = P(\text{type I error})$$

# Significance level and power

- Contingency table:

	$y = H_A$	$y = H_0$
$\hat{y} = \text{reject } H_0$	TP	FP (Type I error)
$\hat{y} = \text{do not reject } H_0$	FN (Type II error)	TN

- Significance level**  $\alpha$ : incorrectly rejecting  $H_0$

$$\alpha = P(\text{type I error})$$

- Power**: correctly rejecting  $H_0$

$$\text{power} = P(\text{reject } H_0 \mid H_A) = 1 - P(\text{type II error})$$



# Significance level and power

- Contingency table:



	$y = H_A$	$y = H_0$
$\hat{y} = \text{reject } H_0$	TP	FP (Type I error)
$\hat{y} = \text{do not reject } H_0$	FN (Type II error)	TN

- Significance level  $\alpha$ : incorrectly rejecting  $H_0$

$$\alpha = P(\text{type I error})$$

- Power: correctly rejecting  $H_0$

$$\text{power} = P(\text{reject } H_0 | H_A) = 1 - P(\text{type II error})$$



# Significance level and power

- Contingency table:



	$y = H_A$	$y = H_0$
$\hat{y} = \text{reject } H_0$	TP	FP (Type I error)
$\hat{y} = \text{do not reject } H_0$	FN (Type II error)	TN

- Significance level  $\alpha$ : incorrectly rejecting  $H_0$

$$\alpha = P(\text{type I error})$$

- Power: correctly rejecting  $H_0$

$$\text{power} = P(\text{reject } H_0 | H_A) = 1 - P(\text{type II error})$$



- What is needed for computing the power?

# Significance level and power

- Contingency table:



	$y = H_A$	$y = H_0$
$\hat{y} = \text{reject } H_0$	TP	FP (Type I error)
$\hat{y} = \text{do not reject } H_0$	FN (Type II error)	TN

- Significance level**  $\alpha$ : incorrectly rejecting  $H_0$

$$\alpha = P(\text{type I error})$$

- Power**: correctly rejecting  $H_0$

$$\text{power} = P(\text{reject } H_0 | H_A) = 1 - P(\text{type II error})$$



- What is needed for computing the power?  $f(s | H_0)$ ,  $f(s | H_A)$

# $p$ -value

- $p$ -value:

- One-tailed:




- Right tail:  $p = P(S \geq s | H_0)$ , e.g.  $1 - \text{stats.norm.cdf}(s, 0, 1)$
- Left tail:  $p = P(S \leq s | H_0)$ , e.g.  $\text{stats.norm.cdf}(s, 0, 1)$

- Two-tailed:

- $p = 2 \min(P(S \leq s | H_0), P(S \geq s | H_0))$ , e.g.  $2 * \min(\text{stats.norm.cdf}(s, 0, 1), 1 - \text{stats.norm.cdf}(s, 0, 1))$

Note: for example, if  $f(s | H_0)$  is symmetric around zero and  $s < 0$ ,

$$p = 2P(S \leq s | H_0)$$



- What is needed for computing the  $p$ -value? (10 sec)

# $p$ -value

- $p$ -value:

- One-tailed:




- Right tail:  $p = P(S \geq s | H_0)$ , e.g.  $1 - \text{stats.norm.cdf}(s, 0, 1)$
- Left tail:  $p = P(S \leq s | H_0)$ , e.g.  $\text{stats.norm.cdf}(s, 0, 1)$

- Two-tailed:

- $p = 2 \min(P(S \leq s | H_0), P(S \geq s | H_0))$ , e.g.  $2 * \min(\text{stats.norm.cdf}(s, 0, 1), 1 - \text{stats.norm.cdf}(s, 0, 1))$

Note: for example, if  $f(s | H_0)$  is symmetric around zero and  $s < 0$ ,

$$p = 2P(S \leq s | H_0)$$



- What is needed for computing the  $p$ -value? (10 sec)

- Null distribution
- Alternative hypothesis  $H_A$  to know one-tailed or two-tailed
- Observation - test statistic computed from data

## Summary: steps for hypothesis testing

- Step 1 Make a “boring” statement
- Step 2 Design an **experiment**
- Step 3 Describe the **data** generated from the experiment and the corresponding random variables
- Step 4 Describe the parameter of interest and their estimates
- Step 5 Translate the “boring” statement into a statistical hypothesis and call it the **null hypothesis  $H_0$**
- Step 6 Find the expression for the **test statistic  $s$**
- Step 7 Find the expression for the **null distribution**
- Step 8 Define **an alternative hypothesis  $H_A$** : one-tailed or two-tailed
- Step 9 Choose a **significance level  $\alpha$**  (the tail), which defines the **rejection region**
- Step 10 Collect **data**
- Step 11 Compute the test statistic from data
- Step 12 Compute the  $p$ -value
- Step 13 If  $p\text{-value} < \alpha$ , i.e. the test statistic falls in the rejection region of the null distribution, then we reject the hypothesis  $H_0$ ; otherwise, we fail to reject  $H_0$ .

# Today

- 1 Terminology
- 2 Example
- 3  $p$ -hacking
- 4 Summary

## Example

Recall example: if you control the diet of your ducks, they lose 2.1 kg after one month on average. Company A has developed a drug D to help ducks lose weight. They claim that on average the drug works better than diet control. Here is the set up for the experiment.



# Example

Recall example: if you control the diet of your ducks, they lose 2.1 kg after one month on average. Company A has developed a drug D to help ducks lose weight. They claim that on average the drug works better than diet control. Here is the set up for the experiment.

Step 1 Make a “boring” statement (5 secs):

## Example

Recall example: if you control the diet of your ducks, they lose 2.1 kg after one month on average. Company A has developed a drug D to help ducks lose weight. They claim that on average the drug works better than diet control. Here is the set up for the experiment.

**Step 1** Make a “boring” statement (5 secs): **drug D works the same as diet**

# Example

Recall example: if you control the diet of your ducks, they lose 2.1 kg after one month on average. Company A has developed a drug D to help ducks lose weight. They claim that on average the drug works better than diet control. Here is the set up for the experiment.

Step 1 Make a “boring” statement (5 secs): **drug D works the same as diet**

Step 2 Design an **experiment** (choose  $N = 30$ ) (10 secs):

## Example

Recall example: if you control the diet of your ducks, they lose 2.1 kg after one month on average. Company A has developed a drug D to help ducks lose weight. They claim that on average the drug works better than diet control. Here is the set up for the experiment.

- Step 1 Make a “boring” statement (5 secs): **drug D works the same as diet**
- Step 2 Design an **experiment** (choose  $N = 30$ ) (10 secs): **let 30 chonker ducks take drug D and measure their weight loss after one month**

## Example

Recall example: if you control the diet of your ducks, they lose 2.1 kg after one month on average. Company A has developed a drug D to help ducks lose weight. They claim that on average the drug works better than diet control. Here is the set up for the experiment.

- Step 1 Make a “boring” statement (5 secs): **drug D works the same as diet**
- Step 2 Design an **experiment** (choose  $N = 30$ ) (10 secs): **let 30 chonker ducks take drug D and measure their weight loss after one month**
- Step 3 Describe the **data** and **random variables** with assumptions about their distributions (5 secs):

# Example

Recall example: if you control the diet of your ducks, they lose 2.1 kg after one month on average. Company A has developed a drug D to help ducks lose weight. They claim that on average the drug works better than diet control. Here is the set up for the experiment.

- Step 1 Make a “boring” statement (5 secs): **drug D works the same as diet**
- Step 2 Design an **experiment** (choose  $N = 30$ ) (10 secs): **let 30 chonker ducks take drug D and measure their weight loss after one month**
- Step 3 Describe the **data** and **random variables** with assumptions about their distributions (5 secs): **weight loss  $x_1, \dots, x_{30}$ ;  $X_1, \dots, X_{30}$  i.i.d. Gaussian random variables**

# Example

Recall example: if you control the diet of your ducks, they lose 2.1 kg after one month on average. Company A has developed a drug D to help ducks lose weight. They claim that on average the drug works better than diet control. Here is the set up for the experiment.

- Step 1 Make a “boring” statement (5 secs): **drug D works the same as diet**
- Step 2 Design an **experiment** (choose  $N = 30$ ) (10 secs): **let 30 chonker ducks take drug D and measure their weight loss after one month**
- Step 3 Describe the **data** and **random variables** with assumptions about their distributions (5 secs): **weight loss  $x_1, \dots, x_{30}$ ;  $X_1, \dots, X_{30}$  i.i.d. Gaussian random variables** - let's make an additional assumption to simplify the problem - the standard deviation of  $X_i$   $\sigma = 0.6$  is known

# Example

Recall example: if you control the diet of your ducks, they lose 2.1 kg after one month on average. Company A has developed a drug D to help ducks lose weight. They claim that on average the drug works better than diet control. Here is the set up for the experiment.

- Step 1 Make a “boring” statement (5 secs): **drug D works the same as diet**
- Step 2 Design an **experiment** (choose  $N = 30$ ) (10 secs): **let 30 chonker ducks take drug D and measure their weight loss after one month**
- Step 3 Describe the **data** and **random variables** with assumptions about their distributions (5 secs): **weight loss  $x_1, \dots, x_{30}$ ;  $X_1, \dots, X_{30}$  i.i.d. Gaussian random variables** - let's make an additional assumption to simplify the problem - the standard deviation of  $X_i$   $\sigma = 0.6$  is known
- Step 4 Describe the parameter of interest and their estimates (10 secs):



# Example

Recall example: if you control the diet of your ducks, they lose 2.1 kg after one month on average. Company A has developed a drug D to help ducks lose weight. They claim that on average the drug works better than diet control. Here is the set up for the experiment.

- Step 1 Make a “boring” statement (5 secs): **drug D works the same as diet**
- Step 2 Design an **experiment** (choose  $N = 30$ ) (10 secs): **let 30 chonker ducks take drug D and measure their weight loss after one month**
- Step 3 Describe the **data** and **random variables** with assumptions about their distributions (5 secs): **weight loss  $x_1, \dots, x_{30}$ ;  $X_1, \dots, X_{30}$  i.i.d. Gaussian random variables** - let's make an additional assumption to simplify the problem - the standard deviation of  $X_i$   $\sigma = 0.6$  is known
- Step 4 Describe the parameter of interest and their estimates (10 secs): **the mean value  $\mu_D$  and  $\hat{\mu}_D = \bar{x}$**

# Example

Recall example: if you control the diet of your ducks, they lose 2.1 kg after one month on average. Company A has developed a drug D to help ducks lose weight. They claim that on average the drug works better than diet control. Here is the set up for the experiment.

- Step 1 Make a “boring” statement (5 secs): **drug D works the same as diet**
- Step 2 Design an **experiment** (choose  $N = 30$ ) (10 secs): **let 30 chonker ducks take drug D and measure their weight loss after one month**
- Step 3 Describe the **data** and **random variables** with assumptions about their distributions (5 secs): **weight loss  $x_1, \dots, x_{30}$ ;  $X_1, \dots, X_{30}$  i.i.d. Gaussian random variables** - let's make an additional assumption to simplify the problem - the standard deviation of  $X_i$   $\sigma = 0.6$  is known
- Step 4 Describe the parameter of interest and their estimates (10 secs): **the mean value  $\mu_D$  and  $\hat{\mu}_D = \bar{x}$**
- Step 5 Translate the “boring” statement into a statistical hypothesis and call it the **null hypothesis  $H_0$**  (10 secs):

# Example

Recall example: if you control the diet of your ducks, they lose 2.1 kg after one month on average. Company A has developed a drug D to help ducks lose weight. They claim that on average the drug works better than diet control. Here is the set up for the experiment.

- Step 1 Make a “boring” statement (5 secs): **drug D works the same as diet**
- Step 2 Design an **experiment** (choose  $N = 30$ ) (10 secs): **let 30 chonker ducks take drug D and measure their weight loss after one month**
- Step 3 Describe the **data** and **random variables** with assumptions about their distributions (5 secs): **weight loss  $x_1, \dots, x_{30}$ ;  $X_1, \dots, X_{30}$  i.i.d. Gaussian random variables** - let's make an additional assumption to simplify the problem - the standard deviation of  $X_i$   $\sigma = 0.6$  is known
- Step 4 Describe the parameter of interest and their estimates (10 secs): **the mean value  $\mu_D$  and  $\hat{\mu}_D = \bar{x}$**
- Step 5 Translate the “boring” statement into a statistical hypothesis and call it the **null hypothesis  $H_0$**  (10 secs):  **$H_0 : \mu_D = 2.1$**

# Example

Recall example: if you control the diet of your ducks, they lose 2.1 kg after one month on average. Company A has developed a drug D to help ducks lose weight. They claim that on average the drug works better than diet control. Here is the set up for the experiment.

- Step 1 Make a “boring” statement (5 secs): **drug D works the same as diet**
- Step 2 Design an **experiment** (choose  $N = 30$ ) (10 secs): **let 30 chonker ducks take drug D and measure their weight loss after one month**
- Step 3 Describe the **data** and **random variables** with assumptions about their distributions (5 secs): **weight loss  $x_1, \dots, x_{30}$ ;  $X_1, \dots, X_{30}$  i.i.d. Gaussian random variables** - let's make an additional assumption to simplify the problem - the standard deviation of  $X_i$   $\sigma = 0.6$  is known
- Step 4 Describe the parameter of interest and their estimates (10 secs): **the mean value  $\mu_D$  and  $\hat{\mu}_D = \bar{x}$**
- Step 5 Translate the “boring” statement into a statistical hypothesis and call it the **null hypothesis  $H_0$**  (10 secs):  **$H_0 : \mu_D = 2.1$**
- Step 6 Find the expression for the **test statistic  $s$**  (60 secs):

# Example

Recall example: if you control the diet of your ducks, they lose 2.1 kg after one month on average. Company A has developed a drug D to help ducks lose weight. They claim that on average the drug works better than diet control. Here is the set up for the experiment.

- Step 1 Make a “boring” statement (5 secs): **drug D works the same as diet**
- Step 2 Design an **experiment** (choose  $N = 30$ ) (10 secs): **let 30 chonker ducks take drug D and measure their weight loss after one month**
- Step 3 Describe the **data** and **random variables** with assumptions about their distributions (5 secs): **weight loss  $x_1, \dots, x_{30}$ ;  $X_1, \dots, X_{30}$  i.i.d. Gaussian random variables** - let's make an additional assumption to simplify the problem - the standard deviation of  $X_i$   $\sigma = 0.6$  is known
- Step 4 Describe the parameter of interest and their estimates (10 secs): **the mean value  $\mu_D$  and  $\hat{\mu}_D = \bar{x}$**
- Step 5 Translate the “boring” statement into a statistical hypothesis and call it the **null hypothesis  $H_0$**  (10 secs):  **$H_0 : \mu_D = 2.1$**
- Step 6 Find the expression for the **test statistic  $s$**  (60 secs):

$$s = z = \frac{\bar{x} - 2.1}{\sigma/\sqrt{30}}$$

# Example

Recall example: if you control the diet of your ducks, they lose 2.1 kg after one month on average. Company A has developed a drug D to help ducks lose weight. They claim that on average the drug works better than diet control. Here is the set up for the experiment.

- Step 1 Make a “boring” statement (5 secs): **drug D works the same as diet**
- Step 2 Design an **experiment** (choose  $N = 30$ ) (10 secs): **let 30 chonker ducks take drug D and measure their weight loss after one month**
- Step 3 Describe the **data** and **random variables** with assumptions about their distributions (5 secs): **weight loss  $x_1, \dots, x_{30}$ ;  $X_1, \dots, X_{30}$  i.i.d. Gaussian random variables** - let's make an additional assumption to simplify the problem - the standard deviation of  $X_i$   $\sigma = 0.6$  is known
- Step 4 Describe the parameter of interest and their estimates (10 secs): **the mean value  $\mu_D$  and  $\hat{\mu}_D = \bar{x}$**
- Step 5 Translate the “boring” statement into a statistical hypothesis and call it the **null hypothesis  $H_0$**  (10 secs):  **$H_0 : \mu_D = 2.1$**
- Step 6 Find the expression for the **test statistic  $s$**  (60 secs):

$$s = z = \frac{\bar{x} - 2.1}{\sigma / \sqrt{30}}$$

- Step 7 Find the expression for the **null distribution  $f(s | H_0)$**  (10 secs):

# Example

Recall example: if you control the diet of your ducks, they lose 2.1 kg after one month on average. Company A has developed a drug D to help ducks lose weight. They claim that on average the drug works better than diet control. Here is the set up for the experiment.

- Step 1 Make a “boring” statement (5 secs): **drug D works the same as diet**
- Step 2 Design an **experiment** (choose  $N = 30$ ) (10 secs): **let 30 chonker ducks take drug D and measure their weight loss after one month**
- Step 3 Describe the **data** and **random variables** with assumptions about their distributions (5 secs): **weight loss  $x_1, \dots, x_{30}$ ;  $X_1, \dots, X_{30}$  i.i.d. Gaussian random variables** - let's make an additional assumption to simplify the problem - the standard deviation of  $X_i$   $\sigma = 0.6$  is known
- Step 4 Describe the parameter of interest and their estimates (10 secs): **the mean value  $\mu_D$  and  $\hat{\mu}_D = \bar{x}$**
- Step 5 Translate the “boring” statement into a statistical hypothesis and call it the **null hypothesis  $H_0$**  (10 secs):  **$H_0 : \mu_D = 2.1$**
- Step 6 Find the expression for the **test statistic  $s$**  (60 secs):

$$s = z = \frac{\bar{x} - 2.1}{\sigma / \sqrt{30}}$$

- Step 7 Find the expression for the **null distribution  $f(s | H_0)$**  (10 secs):

$$f(z | H_0) = \frac{1}{\sqrt{2\pi}} e^{-\frac{z^2}{2}}$$

## Example (cont.)

Step 8 Define **an alternative hypothesis**  $H_A$  (10 secs):



## Example (cont.)

Step 8 Define **an alternative hypothesis**  $H_A$  (10 secs):

$$H_A : \mu_D \neq 2.1 \text{ or } H_A : \mu_D > 2.1$$

## Example (cont.)

Step 8 Define **an alternative hypothesis**  $H_A$  (10 secs):

$$H_A : \mu_D \neq 2.1 \text{ or } H_A : \mu_D > 2.1$$

One-tailed or two-tailed

## Example (cont.)

Step 8 Define **an alternative hypothesis**  $H_A$  (10 secs):

$$H_A : \mu_D \neq 2.1 \text{ or } H_A : \mu_D > 2.1$$

One-tailed or two-tailed

- **Two-tailed (5 secs):**

## Example (cont.)

Step 8 Define **an alternative hypothesis**  $H_A$  (10 secs):

$$H_A : \mu_D \neq 2.1 \text{ or } H_A : \mu_D > 2.1$$

One-tailed or two-tailed

- **Two-tailed (5 secs):**  $H_A : \mu_D \neq 2.1$

## Example (cont.)

Step 8 Define **an alternative hypothesis**  $H_A$  (10 secs):

$$H_A : \mu_D \neq 2.1 \text{ or } H_A : \mu_D > 2.1$$

One-tailed or two-tailed

- **Two-tailed (5 secs):**  $H_A : \mu_D \neq 2.1$
- **One-tailed (5 secs):**

## Example (cont.)

Step 8 Define **an alternative hypothesis**  $H_A$  (10 secs):

$$H_A : \mu_D \neq 2.1 \text{ or } H_A : \mu_D > 2.1$$

One-tailed or two-tailed

- **Two-tailed (5 secs):**  $H_A : \mu_D \neq 2.1$
- **One-tailed (5 secs):**  $H_A : \mu_D > 2.1$

## Example (cont.)

Step 8 Define **an alternative hypothesis**  $H_A$  (10 secs):

$$H_A : \mu_D \neq 2.1 \text{ or } H_A : \mu_D > 2.1$$

One-tailed or two-tailed

- **Two-tailed (5 secs):**  $H_A : \mu_D \neq 2.1$
- **One-tailed (5 secs):**  $H_A : \mu_D > 2.1$

Step 9 Choose a **significance level**  $\alpha$  (the tail), which defines the rejection region (5 secs):

## Example (cont.)

Step 8 Define **an alternative hypothesis**  $H_A$  (10 secs):

$$H_A : \mu_D \neq 2.1 \text{ or } H_A : \mu_D > 2.1$$

One-tailed or two-tailed

- **Two-tailed (5 secs):**  $H_A : \mu_D \neq 2.1$
- **One-tailed (5 secs):**  $H_A : \mu_D > 2.1$

Step 9 Choose a **significance level**  $\alpha$  (the tail), which defines the rejection region (5 secs): e.g.  $\alpha = 0.05$



## Example (cont.)

Step 8 Define **an alternative hypothesis**  $H_A$  (10 secs):

$$H_A : \mu_D \neq 2.1 \text{ or } H_A : \mu_D > 2.1$$

One-tailed or two-tailed

- **Two-tailed (5 secs):**  $H_A : \mu_D \neq 2.1$
- **One-tailed (5 secs):**  $H_A : \mu_D > 2.1$

Step 9 Choose a **significance level**  $\alpha$  (the tail), which defines the rejection region (5 secs): e.g.  $\alpha = 0.05$

Step 10 Collect 30 ducks in 20 secs and feed them drugs

## Example (cont.)

Step 8 Define **an alternative hypothesis**  $H_A$  (10 secs):

$$H_A : \mu_D \neq 2.1 \text{ or } H_A : \mu_D > 2.1$$

One-tailed or two-tailed

- **Two-tailed (5 secs):**  $H_A : \mu_D \neq 2.1$
- **One-tailed (5 secs):**  $H_A : \mu_D > 2.1$

Step 9 Choose a **significance level**  $\alpha$  (the tail), which defines the rejection region (5 secs): e.g.  $\alpha = 0.05$

Step 10 Collect 30 ducks in 20 secs and feed them drugs - great job! Weights measured after one month  $x_1, \dots, x_{30}$

## Example (cont.)

Step 8 Define **an alternative hypothesis**  $H_A$  (10 secs):

$$H_A : \mu_D \neq 2.1 \text{ or } H_A : \mu_D > 2.1$$

One-tailed or two-tailed

- **Two-tailed (5 secs):**  $H_A : \mu_D \neq 2.1$
- **One-tailed (5 secs):**  $H_A : \mu_D > 2.1$

Step 9 Choose a **significance level**  $\alpha$  (the tail), which defines the rejection region (5 secs): e.g.  $\alpha = 0.05$

Step 10 Collect 30 ducks in 20 secs and feed them drugs - great job! Weights measured after one month  $x_1, \dots, x_{30}$

Say  $\frac{1}{30} \sum_{i=1}^{30} x_i = 2.2$

Step 11 Compute the test statistic from data (5 secs):

## Example (cont.)

Step 8 Define **an alternative hypothesis**  $H_A$  (10 secs):

$$H_A : \mu_D \neq 2.1 \text{ or } H_A : \mu_D > 2.1$$

One-tailed or two-tailed

- **Two-tailed (5 secs):**  $H_A : \mu_D \neq 2.1$
- **One-tailed (5 secs):**  $H_A : \mu_D > 2.1$

Step 9 Choose a **significance level**  $\alpha$  (the tail), which defines the rejection region (5 secs): e.g.  $\alpha = 0.05$

Step 10 Collect 30 ducks in 20 secs and feed them drugs - great job! Weights measured after one month  $x_1, \dots, x_{30}$

$$\text{Say } \frac{1}{30} \sum_{i=1}^{30} x_i = 2.2$$

Step 11 Compute the test statistic from data (5 secs):

$$z_0 = \frac{2.2 - 2.1}{0.6/\sqrt{30}} = 0.91$$

## Example (cont.)

Step 12 Compute the  $p$ -value (20 secs):

## Example (cont.)

Step 12 Compute the  $p$ -value (20 secs):

- For  $H_A : \mu_D > 2.1$  (one-tailed):  $p = P(Z \geq z_0 \mid H_0) = 0.1807 > \alpha$

## Example (cont.)

Step 12 Compute the *p*-value (20 secs):



- For  $H_A : \mu_D > 2.1$  (**one-tailed**):  $p = P(Z \geq z_0 \mid H_0) = 0.1807 > \alpha$
- For  $H_A : \mu_D \neq 2.1$  (**two-tailed**):  $p = 2P(Z \geq z_0 \mid H_0) = 0.3613 > \alpha$

## Example (cont.)

Step 12 Compute the  $p$ -value (20 secs):

- For  $H_A : \mu_D > 2.1$  (**one-tailed**):  $p = P(Z \geq z_0 | H_0) = 0.1807 > \alpha$
- For  $H_A : \mu_D \neq 2.1$  (**two-tailed**):  $p = 2P(Z \geq z_0 | H_0) = 0.3613 > \alpha$

Step 13 If  $p\text{-value} < \alpha$ , i.e. the test statistic falls in the rejection region of the null distribution, then we reject the hypothesis  $H_0$



Do not reject  $H_0$  for both one-tailed and two-tailed  $H_A$





## Example (cont.)

Step 12 Compute the  $p$ -value (20 secs):

- For  $H_A : \mu_D > 2.1$  (**one-tailed**):  $p = P(Z \geq z_0 | H_0) = 0.1807 > \alpha$
- For  $H_A : \mu_D \neq 2.1$  (**two-tailed**):  $p = 2P(Z \geq z_0 | H_0) = 0.3613 > \alpha$

Step 13 If  $p\text{-value} < \alpha$ , i.e. the test statistic falls in the rejection region of the null distribution, then we reject the hypothesis  $H_0$



Do not reject  $H_0$  for both one-tailed and two-tailed  $H_A$



What does it mean?

## Example (cont.)

Step 12 Compute the  $p$ -value (20 secs):

- For  $H_A : \mu_D > 2.1$  (one-tailed):  $p = P(Z \geq z_0 | H_0) = 0.1807 > \alpha$
- For  $H_A : \mu_D \neq 2.1$  (two-tailed):  $p = 2P(Z \geq z_0 | H_0) = 0.3613 > \alpha$

Step 13 If  $p\text{-value} < \alpha$ , i.e. the test statistic falls in the rejection region of the null distribution, then we reject the hypothesis  $H_0$



**Do not reject  $H_0$  for both one-tailed and two-tailed  $H_A$**

What does it mean? - Based on this test, you will stick to diet control instead of buying drug D.

## Example (cont.)

What if  $\bar{x} = 2.3$ ?

## Example (cont.)

What if  $\bar{x} = 2.3$ ?

Step 11 Compute the test statistic from data (5 secs):

## Example (cont.)

What if  $\bar{x} = 2.3$ ?

Step 11 Compute the test statistic from data (5 secs):

$$z_0 = \frac{2.3 - 2.1}{0.6/\sqrt{30}} = 1.826$$

## Example (cont.)

What if  $\bar{x} = 2.3$ ?

**Step 11** Compute the test statistic from data (5 secs):

$$z_0 = \frac{2.3 - 2.1}{0.6/\sqrt{30}} = 1.826$$

**Step 12** Compute the *p*-value (20 secs):

## Example (cont.)

What if  $\bar{x} = 2.3$ ?

Step 11 Compute the test statistic from data (5 secs):

$$z_0 = \frac{2.3 - 2.1}{0.6/\sqrt{30}} = 1.826$$

Step 12 Compute the *p*-value (20 secs):

- **For  $H_A : \mu_D > 2.1$  (one-tailed):**  $p = P(Z \geq z_0 \mid H_0) = 0.0339 < \alpha$
- **For  $H_A : \mu_D \neq 2.1$  (two-tailed):**  $p = 2P(Z \geq z_0 \mid H_0) = 0.0679 > \alpha$

## Example (cont.)

What if  $\bar{x} = 2.3$ ?

Step 11 Compute the test statistic from data (5 secs):

$$z_0 = \frac{2.3 - 2.1}{0.6/\sqrt{30}} = 1.826$$



Step 12 Compute the *p*-value (20 secs):

- **For  $H_A : \mu_D > 2.1$  (one-tailed):**  $p = P(Z \geq z_0 \mid H_0) = 0.0339 < \alpha$
- **For  $H_A : \mu_D \neq 2.1$  (two-tailed):**  $p = 2P(Z \geq z_0 \mid H_0) = 0.0679 > \alpha$



## Example (cont.)



**Step 13** If  $p\text{-value} < \alpha$ , i.e. the test statistic falls in the rejection region of the null distribution, then we reject the hypothesis  $H_0$



Reject  $H_0$  for one-tailed  $H_A$ ; do not reject  $H_0$  for two-tailed  $H_A$   
for the same confidence level  $1 - \alpha = 95\%$

## Example (cont.)

**Step 13** If  $p\text{-value} < \alpha$ , i.e. the test statistic falls in the rejection region of the null distribution, then we reject the hypothesis  $H_0$



**Reject  $H_0$  for one-tailed  $H_A$ ; do not reject  $H_0$  for two-tailed  $H_A$  for the same confidence level  $1 - \alpha = 95\%$**

Note: the two-tailed test is more conservative - if the data passes a two-tailed test, it is more conclusive than one-tailed test for the same confidence level

## Example (cont.)

What if  $\bar{x} = 2.3$  with  $N = 100$ ?

## Example (cont.)

What if  $\bar{x} = 2.3$  with  $N = 100$ ?

Step 11 Compute the test statistic from data (5 secs):

## Example (cont.)

What if  $\bar{x} = 2.3$  with  $N = 100$ ?

Step 11 Compute the test statistic from data (5 secs):

$$z_0 = \frac{2.3 - 2.1}{0.6/\sqrt{100}} = 3.33$$

## Example (cont.)

What if  $\bar{x} = 2.3$  with  $N = 100$ ?

Step 11 Compute the test statistic from data (5 secs):

$$z_0 = \frac{2.3 - 2.1}{0.6/\sqrt{100}} = 3.33$$

Step 12 Compute the *p*-value (20 secs):

## Example (cont.)

What if  $\bar{x} = 2.3$  with  $N = 100$ ?

**Step 11** Compute the test statistic from data (5 secs):



$$z_0 = \frac{2.3 - 2.1}{0.6/\sqrt{100}} = 3.33$$

**Step 12** Compute the *p*-value (20 secs):

- **For  $H_A : \mu_D > 2.1$  (one-tailed):**  $p = P(Z \geq z_0 \mid H_0) = 0.0004 < \alpha$
- **For  $H_A : \mu_D \neq 2.1$  (two-tailed):**  $p = 2P(Z \geq z_0 \mid H_0) = 0.0009 < \alpha$

## Example (cont.)

**Step 13** If  $p\text{-value} < \alpha$ , i.e. the test statistic falls in the rejection region of the null distribution, then we reject the hypothesis  $H_0$



**Reject  $H_0$  for both one-tailed and two-tailed  $H_A$**



Note:

- With more data, it becomes more certain that we should reject  $H_0$  in favor of  $H_A$  given the observation  $\bar{x} = 2.3$



## Example (cont.)

**Step 13** If  $p\text{-value} < \alpha$ , i.e. the test statistic falls in the rejection region of the null distribution, then we reject the hypothesis  $H_0$



**Reject  $H_0$  for both one-tailed and two-tailed  $H_A$**

Note:

- With more data, it becomes more certain that we should reject  $H_0$  in favor of  $H_A$  given the observation  $\bar{x} = 2.3$

This test is called **one-sample z-test**

# Today

- 1 Terminology
- 2 Example
- 3  $p$ -hacking**
- 4 Summary

## Recall: one-tailed vs two-tailed tests



- *p*-value indicates how “surprising” the observation is

## Recall: one-tailed vs two-tailed tests

- $p$ -value indicates how “surprising” the observation is
- “Surprising” observation usually means potential novelty



## Recall: one-tailed vs two-tailed tests

- $p$ -value indicates how “surprising” the observation is
- “Surprising” observation usually means potential novelty
- In one of the examples, we have shown that we reject the null hypothesis for the one-tailed test and we fail to reject the null hypothesis for the two-tailed test given the same significance level



## Recall: one-tailed vs two-tailed tests



- $p$ -value indicates how “surprising” the observation is
- “Surprising” observation usually means potential novelty
- In one of the examples, we have shown that we reject the null hypothesis for the one-tailed test and we fail to reject the null hypothesis for the two-tailed test given the same significance level



- In this example, if we use the two-tailed test, we will not claim that we have observed potential novelty with the experiment, whereas if we use the one-tailed test, we claim that we do observe potential novelty

## Recall: one-tailed vs two-tailed tests

- $p$ -value indicates how “surprising” the observation is
- “Surprising” observation usually means potential novelty
- In one of the examples, we have shown that we reject the null hypothesis for the one-tailed test and we fail to reject the null hypothesis for the two-tailed test given the same significance level



- In this example, if we use the two-tailed test, we will not claim that we have observed potential novelty with the experiment, whereas if we use the one-tailed test, we claim that we do observe potential novelty
- The conclusion we draw depends on which test we conduct

## Variation of the *p*-value

- *p*-value is computed from data



## Variation of the $p$ -value

- $p$ -value is computed from data
- Data is random -  **$p$ -value is random**

## Variation of the *p*-value

- *p*-value is computed from data
- Data is random - ***p*-value is random**
- With the same experiment set up, if we switch to a different sample, *p*-value will be different

# *p*-hacking

- Many factors can result in a different *p*-value
- *p*-hacking refers to situations where researchers are **trying multiple things until they get the desired result**
- This action can be a conscious decision, a subconscious decision or even an unconscious action
- *p*-hacking can be tricky to identify
- Suggestions to avoid *p*-hacking, e.g. one should always **report effect sizes and confidence intervals**
- Reference:
  - <https://www.nature.com/news/scientific-method-statistical-errors-1.14700>
  - Why Most Published Research Findings Are False?

## *p*-hacking (cont.)

What should I do!?



## p-hacking (cont.)



What should I do!?

- Be honest and explicit about your assumptions

## p-hacking (cont.)



What should I do!?

- Be honest and explicit about your assumptions
- Be “conservative”


## p-hacking (cont.)



What should I do!?

- Be honest and explicit about your assumptions
- Be “conservative”
- Be skeptical about your result - **don't let go of any doubt!**

## p-hacking (cont.)



What should I do!?

- Be honest and explicit about your assumptions
- Be “conservative”
- Be skeptical about your result - **don't let go of any doubt!**
- Assume the first success is always **too good to be true**



## p-hacking (cont.)



What should I do!?

- Be honest and explicit about your assumptions
- Be “conservative”
- Be skeptical about your result - **don't let go of any doubt!**
- Assume the first success is always **too good to be true** - **try to prove yourself wrong**

## p-hacking (cont.)



What should I do!?

- Be honest and explicit about your assumptions
- Be “conservative”
- Be skeptical about your result - **don't let go of any doubt!**
- Assume the first success is always **too good to be true** - **try to prove yourself wrong** - be a proper scientist

# Today

- 1 Terminology
- 2 Example
- 3  $p$ -hacking
- 4 Summary

# Summary

So far:

- Data types and data containers
- Descriptive data analysis: descriptive statistics, visualization
- Probability distributions, events, random variables, PMF, PDF, parameters
- CDF, Q-Q plot, how to compare two distributions (data vs theoretical, data vs data)
- Modeling
- Parameter estimation: maximum likelihood estimation (MLE) and maximum a posteriori estimation (MAP)
- Classification, multinomial naive Bayes classifier, Gaussian naive Bayes classifier
- Central limit theorem, interval estimation
- Hypothesis test

# Summary

So far:

- Data types and data containers
- Descriptive data analysis: descriptive statistics, visualization
- Probability distributions, events, random variables, PMF, PDF, parameters
- CDF, Q-Q plot, how to compare two distributions (data vs theoretical, data vs data)
- Modeling
- Parameter estimation: maximum likelihood estimation (MLE) and maximum a posteriori estimation (MAP)
- Classification, multinomial naive Bayes classifier, Gaussian naive Bayes classifier
- Central limit theorem, interval estimation
- Hypothesis test

Next:

- More examples, test statistics; comparison of two classifiers

# Summary

So far:

- Data types and data containers
- Descriptive data analysis: descriptive statistics, visualization
- Probability distributions, events, random variables, PMF, PDF, parameters
- CDF, Q-Q plot, how to compare two distributions (data vs theoretical, data vs data)
- Modeling
- Parameter estimation: maximum likelihood estimation (MLE) and maximum a posteriori estimation (MAP)
- Classification, multinomial naive Bayes classifier, Gaussian naive Bayes classifier
- Central limit theorem, interval estimation
- Hypothesis test

Next:

- More examples, test statistics; comparison of two classifiers

Before next lecture:

- Steps for hypothesis testing

