Lecture 2: Probability Distribution Statistical Methods for Data Science

Yinan Yu

Department of Computer Science and Engineering

November 4, 2021

Today

- Probability distribution
 - Why probability distributions?
 - Terminology
 - Some probability distributions that you should know by heart
- 2 Demo
- Summary





Learning outcome

- Be able to explain the following concepts: experiment, event, random variable, probability distribution
- Given the PDF/PMF, be able to describe the probability distribution of a continuous/discrete random variable
- Be able to compute conditional probability
- Understand Gaussian distribution and Bernoulli distribution: 1)
 PDF/PMF; 2) what are the parameters? 3) what happens to the shape of the distribution if we change the parameters?
- Be able to apply the learning routine to study a new probability distribution yourself





Today

- Probability distribution
 - Why probability distributions?
 - Terminology
 - Some probability distributions that you should know by heart
- 2 Demo
- Summary





Why probability distributions?
Terminology

Why probability distributions?





Terminology

Histogram vs probability distribution

You need to estimate the weight distribution of your 1000 ducks without weighing all of them





You need to estimate the weight distribution of your 1000 ducks without weighing all of them, because, well, data collection is expensive.






You need to estimate the weight distribution of your 1000 ducks without weighing all of them, because, well, data collection is expensive. You weighed 20 ducks and you plotted the histogram of the weights.






You need to estimate the weight distribution of your 1000 ducks without weighing all of them, because, well, data collection is expensive. You weighed 20 ducks and you plotted the histogram of the weights. Your best friend Jack looked at the histogram and suggested that you should use a Gaussian distribution to make a better estimation of the distribution.







You need to estimate the weight distribution of your 1000 ducks without weighing all of them, because, well, data collection is expensive. You weighed 20 ducks and you plotted the histogram of the weights. Your best friend Jack looked at the histogram and suggested that you should use a Gaussian distribution to make a better estimation of the distribution.









 Question 1: Why can't I just use descriptive analysis, like the histogram, to describe the data distribution? Why should I use probability distributions?





 Question 1: Why can't I just use descriptive analysis, like the histogram, to describe the data distribution? Why should I use probability distributions?

To address this question, let's describe the data distribution using a **histogram** and a **Gaussian distribution** to see the difference.





Here are the weights of the $20\ ducks$ in kg

duck id	1	2	3	4	 19	20
weight	6.98	5.43	2.97	7.07	 4.63	7.27





Here are the weights of the 20 ducks in kg

d	uck id	1	2	3	4	 19	20
٧	veight	6.98	5.43	2.97	7.07	 4.63	7.27

Let's try to describe these ducks using a histogram with 5 bins.


• Divide the range of the data into 5 bins



Here are the weights of the 20 ducks in kg

duck id	1	2	3	4	 19	20
weight	6.98	5.43	2.97	7.07	 4.63	7.27

- Divide the range of the data into 5 bins
- There are 3 ducks within the first bin [2.965, 4.596]; there are 5 ducks within second bin [4.596, 6.226], etc.








Here are the weights of the 20 ducks in kg

duck id	1	2	3	4	 19	20
weight	6.98	5.43	2.97	7.07	 4.63	7.27

Let's try to describe these ducks using a histogram with 5 bins.

- Divide the range of the data into 5 bins
- There are 3 ducks within the first bin [2.965, 4.596]; there are 5 ducks within second bin [4.596, 6.226], etc.



Now you want to use the histogram to describe the distribution of 1000 ducks






Here are the weights of the 20 ducks in kg

duck id	1	2	3	4	 19	20
weight	6.98	5.43	2.97	7.07	 4.63	7.27

- Divide the range of the data into 5 bins
- There are 3 ducks within the first bin [2.965, 4.596]; there are 5 ducks within second bin [4.596, 6.226], etc.



- Now you want to use the histogram to describe the distribution of 1000 ducks
- What is the chance of a duck weighing between 2.965 kg and 4.596 kg?






Here are the weights of the 20 ducks in kg

duck	d	1	2	3	4	 19	20
weigh	t	6.98	5.43	2.97	7.07	 4.63	7.27

- Divide the range of the data into 5 bins
- There are 3 ducks within the first bin [2.965, 4.596]; there are 5 ducks within second bin [4.596, 6.226], etc.



- Now you want to use the histogram to describe the distribution of 1000 ducks
- What is the chance of a duck weighing between 2.965 kg and 4.596 kg? $\frac{3}{20}$






Here are the weights of the 20 ducks in kg

duck id	1	2	3	4	 19	20
weight	6.98	5.43	2.97	7.07	 4.63	7.27

- Divide the range of the data into 5 bins
- There are 3 ducks within the first bin [2.965, 4.596]; there are 5 ducks within second bin [4.596, 6.226], etc.



- Now you want to use the histogram to describe the distribution of 1000 ducks
- What is the chance of a duck weighing between 2.965 kg and 4.596 kg? $\frac{3}{20}$ How about between 3.1 kg and 3.4 kg?





Resolution: the number of bins per kilogram

$$\frac{\text{number of bins}}{\textit{range}} = \frac{\text{number of bins}}{\text{max}(\textit{weights}) - \text{min}(\textit{weights})} = \frac{5}{11.118 - 2.965} = 0.61$$







Resolution: the number of bins per kilogram

$$\frac{\text{number of bins}}{\textit{range}} = \frac{\text{number of bins}}{\text{max}(\textit{weights}) - \text{min}(\textit{weights})} = \frac{5}{11.118 - 2.965} = 0.61$$

- \bullet The chance of weighing between 3.1 and 3.4: (3.4 3.1) \times resolution \times $\frac{3}{20}=0.028$
- The chance of weighing between 4.1 and 4.4: $(4.4-4.1) \times resolution \times \frac{20}{20} = 0.028$









Resolution: the number of bins per kilogram

$$\frac{\text{number of bins}}{\text{range}} = \frac{\text{number of bins}}{\text{max}(\text{weights}) - \text{min}(\text{weights})} = \frac{5}{11.118 - 2.965} = 0.61$$

- \bullet The chance of weighing between 3.1 and 3.4: (3.4 3.1) \times resolution \times $\frac{3}{20}=0.028$
- The chance of weighing between 4.1 and 4.4: $(4.4-4.1) \times resolution \times \frac{3}{20} = 0.028$






· We lose the resolution within each bin due to quantization




Resolution: the number of bins per kilogram

$$\frac{\text{number of bins}}{\text{range}} = \frac{\text{number of bins}}{\text{max}(\text{weights}) - \text{min}(\text{weights})} = \frac{5}{11.118 - 2.965} = 0.61$$

- \bullet The chance of weighing between 3.1 and 3.4: (3.4 3.1) \times resolution \times $\frac{3}{20}=0.028$
- The chance of weighing between 4.1 and 4.4: $(4.4-4.1) \times resolution \times \frac{3}{20} = 0.028$






 We lose the resolution within each bin due to quantization - you don't know what's going on within each bin




Resolution: the number of bins per kilogram

$$\frac{\text{number of bins}}{\text{range}} = \frac{\text{number of bins}}{\text{max}(\text{weights}) - \text{min}(\text{weights})} = \frac{5}{11.118 - 2.965} = 0.61$$

- \bullet The chance of weighing between 3.1 and 3.4: (3.4 3.1) \times resolution \times $\frac{3}{20}=0.028$
- The chance of weighing between 4.1 and 4.4: $(4.4-4.1) \times resolution \times \frac{3}{20} = 0.028$






- We lose the resolution within each bin due to quantization you don't know what's going on within each bin
- And if we increase the number of bins? 5 -> 200




Resolution: the number of bins per kilogram


$$\frac{\text{number of bins}}{\text{range}} = \frac{\text{number of bins}}{\text{max}(\text{weights}) - \text{min}(\text{weights})} = \frac{5}{11.118 - 2.965} = 0.61$$

- The chance of weighing between 3.1 and 3.4: $(3.4 3.1) \times resolution \times \frac{3}{20} = 0.028$
- The chance of weighing between 4.1 and 4.4: $(4.4-4.1) \times resolution \times \frac{3}{20} = 0.028$





- We lose the resolution within each bin due to quantization you don't know what's going on within each bin
- And if we increase the number of bins? 5 -> 200







 Descriptive analysis (e.g. histogram) does not generalize well to unseen data






- Descriptive analysis (e.g. histogram) does not generalize well to unseen data
- Now let's try to use a Gaussian distribution to describe the data





- Descriptive analysis (e.g. histogram) does not generalize well to unseen data
- Now let's try to use a Gaussian distribution to describe the data
- First, we assume that data is generated from a Gaussian distribution





 A Gaussian distribution is described by a function that looks similar to this histogram

$$f(x) = \frac{1}{\sqrt{2\pi\sigma^2}}e^{-\frac{1}{2}\left(\frac{x-\mu}{\sigma}\right)^2} = a \text{ scalar}$$



 A Gaussian distribution is described by a function that looks similar to this histogram

$$\mathbf{f(x)} = rac{1}{\sqrt{2\pi\sigma^2}} \mathrm{e}^{-rac{1}{2}\left(rac{x-\mu}{\sigma}
ight)^2} = \mathsf{a} \; \mathsf{scalar}$$


• This function is sufficiently defined by two parameters μ and σ .



 A Gaussian distribution is described by a function that looks similar to this histogram

$$f(x) = \frac{1}{\sqrt{2\pi\sigma^2}}e^{-\frac{1}{2}\left(\frac{x-\mu}{\sigma}\right)^2} = a \text{ scalar}$$

- This function is sufficiently defined by two parameters μ and σ .
- The shape of the function (e.g. given μ =7 and σ =2):








 A Gaussian distribution is described by a function that looks similar to this histogram

$$f(x) = rac{1}{\sqrt{2\pi\sigma^2}}e^{-rac{1}{2}\left(rac{x-\mu}{\sigma}
ight)^2} = a$$
 scalar


- This function is sufficiently defined by two parameters μ and σ .
- The shape of the function (e.g. given μ =7 and σ =2):




• We will try to use this function instead of the histogram to describe the data.




- Describe the distribution:
 - Histogram (using 0.61 bins to describe 1 kg):
 - The chance of $weight \in [3.1, 3.4]$: $(3.4 3.1) \times resolution \times \frac{3}{20} = 0.028$ The chance of $weight \in [4.1, 4.4]$: $(4.4 4.1) \times resolution \times \frac{3}{20} = 0.028$









- Describe the distribution:
 - Histogram (using 0.61 bins to describe 1 kg):
 - The chance of $weight \in [3.1, 3.4]$: $(3.4 3.1) \times resolution \times \frac{3}{20} = 0.028$
 - The chance of weight $\in [4.1, 4.4]$: $(4.4 4.1) \times resolution \times \frac{3}{20} = 0.028$





- Gaussian distribution (using infinite bins to describe 1 kg):
 - The chance of $weight \in [3.1, 3.4]$: $\int_{3.1}^{3.4} f(t) dt = 0.010$
 - The chance of weight $\in [4.1, 4.4]$: $\int_{4.1}^{4.4} f(t) dt = 0.023$





• Descriptive analysis: a histogram with M bins (e.g. M=5)





- Descriptive analysis: a *histogram* with M bins (e.g. M=5)
- Gaussian distribution: a mathematical function $f(x) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{1}{2}\left(\frac{x-\mu}{\sigma}\right)^2}$





- ullet Descriptive analysis: a *histogram* with M bins (e.g. M=5)
- Gaussian distribution: a mathematical function $f(x) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{1}{2}\left(\frac{x-\mu}{\sigma}\right)^2}$

Comparison

	Histogram		Gaussian distribution		
Representation	M values		mathematical function		
Number of parameters	М -		2 (μ and σ)	+	
Resolution	$\frac{M}{\max(x)-\min(x)}$		infinity	+	
Analytical properties	No	-	Yes	+	
Assumptions	No +		Yes	-	
Can be directly computed from data	Yes +		Parameters unknown	-	





- Descriptive analysis: a histogram with M bins (e.g. M = 5)
- Gaussian distribution: a mathematical function $f(x) = \frac{1}{\sqrt{2-2}} e^{-\frac{1}{2} \left(\frac{x-\mu}{\sigma}\right)^2}$

Comparison

	Histogram		Gaussian distribution		
Representation	M values		mathematical function f		
Number of parameters	М	-	2 (μ and σ)	+	
Resolution	$\frac{M}{\max(x)-\min(x)}$	-	infinity	+	
Analytical properties	No	-	Yes	+	
Assumptions	No	+	Yes	-	
Can be directly computed from data	Yes	+	Parameters unknown	-	

• For your use case, you want to estimate the distribution of your 1000 ducks without weighing all of them. It is hard to do that from the histogram.





- Descriptive analysis: a histogram with M bins (e.g. M = 5)
- Gaussian distribution: a mathematical function $f(x) = \frac{1}{\sqrt{2-\sigma^2}} e^{-\frac{1}{2}\left(\frac{x-\mu}{\sigma}\right)^2}$

Comparison

	Histogram		Gaussian distribution	
Representation	M values		mathematical function	
Number of parameters	М	-	2 (μ and σ)	+
Resolution	$\frac{M}{\max(x)-\min(x)}$		infinity	+
Analytical properties	No	-	Yes	+
Assumptions	No	+	Yes	-
Can be directly computed from data	Yes	+	Parameters unknown	-

• For your use case, you want to estimate the distribution of your 1000 ducks without weighing all of them. It is hard to do that from the histogram. The histogram describes the data you have seen



- Descriptive analysis: a *histogram* with M bins (e.g. M=5)
- Gaussian distribution: a mathematical function $f(x) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{1}{2}\left(\frac{x-\mu}{\sigma}\right)^2}$

Comparison

Histogram		Gaussian distributio	n
M values		mathematical functio	n f
М	-	2 (μ and σ)	+
$\frac{M}{\max(x)-\min(x)}$	-	infinity	+
No	-	Yes	+
No	+	Yes	-
Yes	+	Parameters unknown	-
	M values M max(x)-min(x) No No	$\begin{array}{c c} M & - \\ \hline M \\ \hline max(x) - min(x) \\ \hline No & - \\ \hline No & + \\ \end{array}$	

 For your use case, you want to estimate the distribution of your 1000 ducks without weighing all of them. It is hard to do that from the histogram. The histogram describes the data you have seen, but it does not generalize well to unseen data.



- Descriptive analysis: a *histogram* with M bins (e.g. M=5)
- Gaussian distribution: a mathematical function $f(x) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{1}{2}\left(\frac{x-\mu}{\sigma}\right)^2}$
- Comparison

	Histogram	Histogram Gaussian distributio		
Representation	M values		mathematical functio	n f
Number of parameters	М	-	2 (μ and σ)	+
Resolution	$\frac{M}{\max(x)-\min(x)}$	-	infinity	+
Analytical properties	No	-	Yes	+
Assumptions	No	+	Yes	-
Can be directly computed from data	Yes +		Parameters unknown	-

- For your use case, you want to estimate the distribution of your 1000 ducks
 without weighing all of them. It is hard to do that from the histogram. The
 histogram describes the data you have seen, but it does not generalize well to
 unseen data.
- Statistical modelling using probability distributions (e.g. a Gaussian distribution) can help you with that!





- Descriptive analysis: a *histogram* with M bins (e.g. M=5)
- Gaussian distribution: a mathematical function $f(x) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{1}{2}\left(\frac{x-\mu}{\sigma}\right)^2}$
- Comparison

	Histogram		Gaussian distribution		
Representation	M values		mathematical functio	n f	
Number of parameters	М	-	2 (μ and σ)	+	
Resolution	$\frac{M}{\max(x)-\min(x)}$	-	infinity	+	
Analytical properties	No	-	Yes	+	
Assumptions	No	+	Yes	-	
Can be directly computed from data	Yes +		Parameters unknown	-	

- For your use case, you want to estimate the distribution of your 1000 ducks
 without weighing all of them. It is hard to do that from the histogram. The
 histogram describes the data you have seen, but it does not generalize well to
 unseen data.
- Statistical modelling using probability distributions (e.g. a Gaussian distribution) can help you with that!
- This concludes the question why we use probability distributions instead of histograms to describe your ducks.





- Descriptive analysis: a *histogram* with M bins (e.g. M=5)
- Gaussian distribution: a mathematical function $f(x) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{1}{2}\left(\frac{x-\mu}{\sigma}\right)^2}$

Comparison

	Histogram		Gaussian distribution		
Representation	M values		mathematical functio	n f	
Number of parameters	М	-	2 (μ and σ)	+	
Resolution	$\frac{M}{\max(x)-\min(x)}$	-	infinity	+	
Analytical properties	No	-	Yes	+	
Assumptions	No	+	Yes	-	
Can be directly computed from data	Yes +		Parameters unknown	-	

- For your use case, you want to estimate the distribution of your 1000 ducks
 without weighing all of them. It is hard to do that from the histogram. The
 histogram describes the data you have seen, but it does not generalize well to
 unseen data.
- Statistical modelling using probability distributions (e.g. a Gaussian distribution) can help you with that!
- This concludes the question why we use probability distributions instead of histograms to describe your ducks.

Disclaimer: we used a continuous distribution to illustrate the comparison between a histogram and a probability distribution. A discrete



Choosing a probability distribution


 Question 2: How do I know which probability distribution I should use to describe the data? How do I know that it should be a Gaussian distribution?





Choosing a probability distribution

- Question 2: How do I know which probability distribution I should use to describe the data? How do I know that it should be a Gaussian distribution?
 - Short answer: if the probability distribution looks like the histogram, then go for it!








Choosing a probability distribution

- Question 2: How do I know which probability distribution I should use to describe the data? How do I know that it should be a Gaussian distribution?
 - Short answer: if the probability distribution looks like the histogram, then go for it!




• Long answer will be given in lecture 3.





Parameter estimation and evaluation


• Question 3: Okay fine, let's say we describe the data with a Gaussian distribution. How do I know what the parameters μ and σ are?





Parameter estimation and evaluation

• Question 3: Okay fine, let's say we describe the data with a Gaussian distribution. How do I know what the parameters μ and σ are?



 This is done by parameter estimation. In lecture 3 & 4, we will talk about the maximum likelihood estimation (MLE) and the maximum a posteriori estimation (MAP).



Why probability distributions? Terminology Some probability distributions that you should know by heart

Terminology





• Experiment: an action that leads to one outcome. For example, we weigh a duck and look at its weight. The outcome is weight = 2 kg.





- Experiment: an action that leads to one outcome. For example, we weigh a duck and look at its weight. The outcome is weight = 2 kg.
- Sample space: the set of all possible outcomes from the experiment. The sample space of the previous example is any real value between 0 and ∞ .





- Experiment: an action that leads to one outcome. For example, we weigh a duck and look at its weight. The outcome is weight = 2 kg.
- Sample space: the set of all possible outcomes from the experiment. The sample space of the previous example is any real value between 0 and ∞ .
- Event: a subset of the sample space, for example, a duck weighs between 5kg and 6kg.



- Experiment: an action that leads to one outcome. For example, we weigh a duck and look at its weight. The outcome is weight = 2 kg.
- Sample space: the set of all possible outcomes from the experiment. The sample space of the previous example is any real value between 0 and ∞ .
- Event: a subset of the sample space, for example, a duck weighs between 5kg and 6kg.
- Probability distribution: the probability of the occurrence of *any* event in the sample space, e.g. P(a duck weighs between a kg and b kg) for any $0 < a < b < \infty$ (not only for a = 5 and b = 6).





- Experiment: an action that leads to one outcome. For example, we weigh a duck and look at its weight. The outcome is weight = 2 kg.
- Sample space: the set of all possible outcomes from the experiment. The sample space of the previous example is any real value between 0 and ∞ .
- Event: a subset of the sample space, for example, a duck weighs between 5kg and 6kg.
- Probability distribution: the probability of the occurrence of any event in the sample space, e.g. P(a duck weighs between a kg and b kg) for any $0 < a < b < \infty$ (not only for a = 5 and b = 6).
- Random variable X:
 - Heuristically, X assigns a numerical value to each outcome of the experiment:

$$X: \mathsf{weight} o \mathbb{R}$$

- X follows some underlying probability distribution.
- Discrete random variable and continuous random variable: depends on the sample space of the
 experiment; the underlying distributions are called discrete distribution and continuous
 distribution, respectively. For example, weights are continuous so X from this example is a
 continuous random variable.





- Experiment: an action that leads to one outcome. For example, we weigh a duck and look at its weight. The outcome is weight = 2 kg.
- Sample space: the set of all possible outcomes from the experiment. The sample space of the previous example is any real value between 0 and ∞ .
- Event: a subset of the sample space, for example, a duck weighs between 5kg and 6kg.
- Probability distribution: the probability of the occurrence of any event in the sample space, e.g. P(a duck weighs between a kg and b kg) for any $0 < a < b < \infty$ (not only for a = 5 and b = 6).
- Random variable X:
 - Heuristically, X assigns a numerical value to each outcome of the experiment:

$$X: \mathsf{weight} o \mathbb{R}$$

- X follows some underlying probability distribution.
- Discrete random variable and continuous random variable: depends on the sample space of the
 experiment; the underlying distributions are called discrete distribution and continuous
 distribution, respectively. For example, weights are continuous so X from this example is a
 continuous random variable.
- Data x: a value drawn from the underlying distribution of X.
 - We use a capital letter (e.g. X) to denote a random variable and the corresponding lower case letter (e.g. x) to denote the data generated from the underlying distribution of X.
 - Discrete random variable: categorical data or discrete numerical data
 - Continuous random variable: continuous numerical data





A probability distribution describes the probabilities of occurrence of all possible events.





A probability distribution describes the probabilities of occurrence of all possible events.

More precisely, a probability distribution is defined by a function f_X (also denoted as f if neglecting X does not cause confusion), where





A probability distribution describes the probabilities of occurrence of all possible events.

More precisely, a probability distribution is defined by a function f_X (also denoted as f if neglecting X does not cause confusion), where

• for discrete distribution, the probability mass function (PMF) is used, where

$$f_X(x_i) = P(X = x_i)$$

where $0 \le f_X(x_i) \le 1$ for all x_i (discrete).



A probability distribution describes the probabilities of occurrence of all possible events.

More precisely, a probability distribution is defined by a function f_X (also denoted as f if neglecting X does not cause confusion), where

• for discrete distribution, the probability mass function (PMF) is used, where

$$f_X(x_i) = P(X = x_i)$$

where $0 \le f_X(x_i) \le 1$ for all x_i (discrete).

 for continuous distribution, the probability density function (PDF) is used, where

$$P(a \le X \le b) = \int_a^b f_X(x) dx, \ \forall a, b \in \mathbb{R}, a \le b$$

where $f_X(x) \ge 0$ for all x (continuous).





A probability distribution describes the probabilities of occurrence of all possible events.

More precisely, a probability distribution is defined by a function f_X (also denoted as f if neglecting X does not cause confusion), where

• for discrete distribution, the probability mass function (PMF) is used, where

$$f_X(x_i) = P(X = x_i)$$

where $0 \le f_X(x_i) \le 1$ for all x_i (discrete).

 for continuous distribution, the probability density function (PDF) is used, where

$$P(a \le X \le b) = \int_a^b f_X(x) dx, \ \forall a, b \in \mathbb{R}, a \le b$$

where $f_X(x) \ge 0$ for all x (continuous).

P(event) is the probability of the **event** occurring.






Example: continuous random variables and PDF

- Experiment: you weigh a duck and look at its weight
- Sample space: $0 < weight < \infty$
- Random variable X : weight $\to \mathbb{R}$
 - $\bullet \ \ X = x \ \text{if the duck weighs} \ x \ \text{kg for} \ 0 < x < \infty$
 - X follows a Gaussian distribution with parameters μ and σ ; denoted as $X \sim \mathcal{N}(\mu, \sigma^2)$
- PDF: f_X(x)


$$P(a \le X \le b) = \underbrace{\int_a^b f_X(x) dx}_{\text{Integral} = \text{area under the PDF curve}} \forall a, b \in \mathbb{R}, a \le b$$





Example: discrete random variables and PMF

- Experiment: you measure the color of the duck.
- Sample space: the color can be either red or blue
- Random variable $X : color \rightarrow \mathbb{Z}$
 - $X = \begin{cases} 0, & \text{if a duck is red} \\ 1, & \text{if a duck is blue} \end{cases}$
 - X follows a Bernoulli distribution with parameter p; denoted as $X \sim Bernoulli(p)$
- PMF: $f_X(x_i) = P(X = x_i)$








Differences between PMF and PDF

- Discrete distribution $f_X(x_i) = P(X = x_i)$:
 - · y-axis represents the probability itself
- Continuous distribution:
 - $P(a \le X \le b) = \int_a^b f_X(x) dx$: **y-axis** f(x) DOES NOT represent the probability itself.
 - For continuous distributions, the probability at any given value is always 0, i.e.
 P(X = a) = P(a ≤ X ≤ a) = ∫_a^a f_X(x)dx ≡ 0. Example: what is the probability of a duck weighing exactly 4.32028374... kg?







Conditional probability

Given events A and B,

$$P(A \mid B) = \frac{P(A \text{ and } B)}{P(B)} = \frac{P(A \cap B)}{P(B)}$$

The probability of event A given event B.





- Experiment: You ask your ducks to stand in a row again and look at their colors and sizes.
- Sample space: The color can be either red or blue; the size can be either slim or chonker.
- Data:

duck id	1	2	3	4	5	6
color	red	red	blue	blue	blue	red
size	chonker	slim	slim	chonker	chonker	slim

- Event:
 - · A: a duck is blue
 - B: a duck is a chonker





- Experiment: You ask your ducks to stand in a row again and look at their colors and sizes.
- Sample space: The color can be either red or blue; the size can be either slim or chonker.
- Data:

duck id	1	2	3	4	5	6
color	red	red	blue	blue	blue	red
size	chonker	slim	slim	chonker	chonker	slim

- Event:
 - · A: a duck is blue
 - B: a duck is a chonker
- Estimate the conditional probability $P(A \mid B) = \frac{P(A \text{ and } B)}{P(B)}$ from this data



- Experiment: You ask your ducks to stand in a row again and look at their colors and sizes.
- Sample space: The color can be either red or blue; the size can be either slim or chonker.
- Data:

duck id	1	2	3	4	5	6
color	red	red	blue	blue	blue	red
size	chonker	slim	slim	chonker	chonker	slim

- Event:
 - · A: a duck is blue
 - B: a duck is a chonker
- Estimate the conditional probability $P(A \mid B) = \frac{P(A \text{ and } B)}{P(B)}$ from this data
 - \bullet P(A and B): the probability that a duck is both blue and a chonker is (10 secs)



- Experiment: You ask your ducks to stand in a row again and look at their colors and sizes.
- Sample space: The color can be either red or blue; the size can be either slim or chonker.
- Data:

duck id	1	2	3	4	5	6
color	red	red	blue	blue	blue	red
size	chonker	slim	slim	chonker	chonker	slim

- Event:
 - · A: a duck is blue
 - B: a duck is a chonker
- Estimate the conditional probability $P(A \mid B) = \frac{P(A \text{ and } B)}{P(B)}$ from this data
 - P(A and B): the probability that a duck is both blue and a chonker is (10 secs)

$$\frac{count(blue and chonker)}{total} = \frac{1}{2}$$





Example:

- Experiment: You ask your ducks to stand in a row again and look at their colors and sizes.
- Sample space: The color can be either red or blue; the size can be either slim or chonker.
- Data:

duck id	1	2	3	4	5	6
color	red	red	blue	blue	blue	red
size	chonker	slim	slim	chonker	chonker	slim

- Event:
 - · A: a duck is blue
 - B: a duck is a chonker
- Estimate the conditional probability $P(A \mid B) = \frac{P(A \text{ and } B)}{P(B)}$ from this data
 - \bullet P(A and B): the probability that a duck is both blue and a chonker is (10 secs)

$$\frac{count(blue and chonker)}{total} = \frac{2}{6}$$

• P(B): the probability that a duck is a chonker is (10 secs)



Example:

- Experiment: You ask your ducks to stand in a row again and look at their colors and sizes.
- Sample space: The color can be either red or blue; the size can be either slim or chonker.
- Data:

duck id	1	2	3	4	5	6
color	red	red	blue	blue	blue	red
size	chonker	slim	slim	chonker	chonker	slim

- Event:
 - A: a duck is blue
 - B: a duck is a chonker
- Estimate the conditional probability $P(A \mid B) = \frac{P(A \text{ and } B)}{P(B)}$ from this data
 - P(A and B): the probability that a duck is both blue and a chonker is (10 secs)

$$\frac{\textit{count}(\textit{blue and chonker})}{\textit{total}} = \frac{2}{6}$$

P(B): the probability that a duck is a chonker is (10 secs)

$$\frac{count(\mathsf{chonker})}{\mathsf{total}} = \frac{3}{6}$$





Example:

- Experiment: You ask your ducks to stand in a row again and look at their colors and sizes.
- Sample space: The color can be either red or blue; the size can be either slim or chonker.
- Data:

duck id	1	2	3	4	5	6
color	red	red	blue	blue	blue	red
size	chonker	slim	slim	chonker	chonker	slim

- Event:
 - · A: a duck is blue
 - B: a duck is a chonker
- Estimate the conditional probability $P(A \mid B) = \frac{P(A \text{ and } B)}{P(B)}$ from this data
 - P(A and B): the probability that a duck is both blue and a chonker is (10 secs)

$$\frac{count(blue and chonker)}{total} = \frac{2}{6}$$

P(B): the probability that a duck is a chonker is (10 secs)

$$\frac{count(chonker)}{total} = \frac{3}{6}$$

· Conditional probability:





Example:

- Experiment: You ask your ducks to stand in a row again and look at their colors and sizes.
- Sample space: The color can be either red or blue; the size can be either slim or chonker.
- Data:

duck id	1	2	3	4	5	6
color	red	red	blue	blue	blue	red
size	chonker	slim	slim	chonker	chonker	slim

- Event:
 - · A: a duck is blue
 - B: a duck is a chonker
- Estimate the conditional probability $P(A \mid B) = \frac{P(A \text{ and } B)}{P(B)}$ from this data
 - P(A and B): the probability that a duck is both blue and a chonker is (10 secs)

$$\frac{count(blue and chonker)}{total} = \frac{2}{6}$$

P(B): the probability that a duck is a chonker is (10 secs)

$$\frac{count(chonker)}{total} = \frac{3}{6}$$

Conditional probability:

$$P(A \mid B) = \frac{P(A \text{ and } B)}{P(B)} = \frac{P(A \cap B)}{P(B)} = \frac{2/6}{3/6} = \frac{2}{3}$$





An alternative way to estimate $P(A \mid B)$:

- Count the chonkers: 3
- Count blue ducks within the chonker gang: 2
- $P(A \mid B) = \frac{2}{3}$



An alternative way to estimate $P(A \mid B)$:

- Count the chonkers: 3
- Count blue ducks within the chonker gang: 2
- $P(A \mid B) = \frac{2}{3}$

Side note: from this calculation, can you make a bold statement of the probability distribution of all the ducks in the world? No. It is only an estimation based on the data you got.



As an exercise, let's define the random variables.

- Experiment: You ask your ducks to stand in a row again and look at their colors and sizes.
- Sample space: The color can be either red or blue; the size can be either slim or chonker.
- Data:

duck id	1	2	3	4	5	6
color	red	red	blue	blue	blue	red
size	chonker	slim	slim	chonker	chonker	slim

Event:

- A: a duck is blue
- B: a duck is a chonker





As an exercise, let's define the random variables.

- Experiment: You ask your ducks to stand in a row again and look at their colors and sizes.
- Sample space: The color can be either red or blue; the size can be either slim or chonker.
- Data:

duck id	1	2	3	4	5	6
color	red	red	blue	blue	blue	red
size	chonker	slim	slim	chonker	chonker	slim

- Event:
 - · A: a duck is blue
 - B: a duck is a chonker
- Random variables: X, Y

Hint: $X : \mathsf{color} \to \mathbb{Z}, \ Y : \mathsf{size} \to \mathbb{Z} \ (10 \ \mathsf{secs})$





As an exercise, let's define the random variables.

- Experiment: You ask your ducks to stand in a row again and look at their colors and sizes.
- Sample space: The color can be either red or blue; the size can be either slim or chonker.
- Data:

duck id	1	2	3	4	5	6
color	red	red	blue	blue	blue	red
size	chonker	slim	slim	chonker	chonker	slim

- Event:
 - A: a duck is blue
 - B: a duck is a chonker
- Random variables: X, Y

Hint:
$$X : \operatorname{color} \to \mathbb{Z}$$
, $Y : \operatorname{size} \to \mathbb{Z}$ (10 secs)

$$X = \begin{cases} 0, \text{ duck is red} \\ 1, \text{ duck is blue} \end{cases} \text{ and } Y = \begin{cases} 0, \text{ duck is slim} \\ 1, \text{ duck is a chonker} \end{cases}$$



As an exercise, let's define the random variables.

- Experiment: You ask your ducks to stand in a row again and look at their colors and sizes.
- Sample space: The color can be either red or blue; the size can be either slim or chonker.
- Data:

duck id	1	2	3	4	5	6
color	red	red	blue	blue	blue	red
size	chonker	slim	slim	chonker	chonker	slim

- Event:
 - A: a duck is blue
 - B: a duck is a chonker
- Random variables: X, Y

Hint:
$$X : \operatorname{color} \to \mathbb{Z}, Y : \operatorname{size} \to \mathbb{Z}$$
 (10 secs)

$$X = \begin{cases} 0, \text{ duck is red} \\ 1, \text{ duck is blue} \end{cases} \text{ and } Y = \begin{cases} 0, \text{ duck is slim} \\ 1, \text{ duck is a chonker} \end{cases}$$

Are they continuous or discrete? (2 secs)





As an exercise, let's define the random variables.

- Experiment: You ask your ducks to stand in a row again and look at their colors and sizes.
- Sample space: The color can be either red or blue; the size can be either slim or chonker.
- Data:

duck id	1	2	3	4	5	6
color	red	red	blue	blue	blue	red
size	chonker	slim	slim	chonker	chonker	slim

- Event:
 - · A: a duck is blue
 - B: a duck is a chonker
- Random variables: X, Y

Hint:
$$X : \operatorname{color} \to \mathbb{Z}$$
, $Y : \operatorname{size} \to \mathbb{Z}$ (10 secs)

$$X = \begin{cases} 0, \text{ duck is red} \\ 1, \text{ duck is blue} \end{cases} \text{ and } Y = \begin{cases} 0, \text{ duck is slim} \\ 1, \text{ duck is a chonker} \end{cases}$$

Are they continuous or discrete? (2 secs) Discrete





As an exercise, let's define the random variables.

- Experiment: You ask your ducks to stand in a row again and look at their colors and sizes.
- Sample space: The color can be either red or blue; the size can be either slim or chonker.
- Data:

duck id	1	2	3	4	5	6
color	red	red	blue	blue	blue	red
size	chonker	slim	slim	chonker	chonker	slim

- Event:
 - · A: a duck is blue
 - B: a duck is a chonker
- Random variables: X, Y

Hint: $X : \operatorname{color} \to \mathbb{Z}, Y : \operatorname{size} \to \mathbb{Z}$ (10 secs)

$$X = \begin{cases} 0, \text{ duck is red} \\ 1, \text{ duck is blue} \end{cases} \text{ and } Y = \begin{cases} 0, \text{ duck is slim} \\ 1, \text{ duck is a chonker} \end{cases}$$

Are they continuous or discrete? (2 secs) Discrete

Write the conditional probability in terms of the random variables X and Y (10 secs), i.e.

$$P(A \mid B) = \frac{P(A \text{ and } B)}{P(B)} =$$





As an exercise, let's define the random variables.

- Experiment: You ask your ducks to stand in a row again and look at their colors and sizes.
- Sample space: The color can be either red or blue; the size can be either slim or chonker.
- Data:

duck id	1	2	3	4	5	6
color	red	red	blue	blue	blue	red
size	chonker	slim	slim	chonker	chonker	slim

- Event:
 - · A: a duck is blue
 - B: a duck is a chonker
- Random variables: X, Y

Hint: $X : \operatorname{color} \to \mathbb{Z}, Y : \operatorname{size} \to \mathbb{Z}$ (10 secs)

$$X = \begin{cases} 0, \text{ duck is red} \\ 1, \text{ duck is blue} \end{cases} \text{ and } Y = \begin{cases} 0, \text{ duck is slim} \\ 1, \text{ duck is a chonker} \end{cases}$$

Are they continuous or discrete? (2 secs) Discrete

Write the conditional probability in terms of the random variables X and Y (10 secs), i.e.

$$P(A \mid B) = \frac{P(A \text{ and } B)}{P(B)} = P(X = 1 \mid Y = 1) = \frac{P(X = 1 \text{ and } Y = 1)}{P(Y = 1)}$$





Independent events

Two events A and B are independent if and only if

$$P(A \text{ and } B) := P(A \cap B) = P(A)P(B)$$

$$\iff$$
 $P(A \mid B) = P(A), P(B \mid A) = P(B)$ (conditional probability)

$$\iff$$
 log $(P(A \text{ and } B) = \log(P(A \cap B)) = \log(P(A)) + \log(P(B))$





Bayes' rule

Given events A and B,

$$P(A \mid B) = \frac{P(B \mid A)P(A)}{P(B)}$$

Just a heads-up!





Summary: Terminology

- Experiment
- Sample space
- Event
- Random variable:
 - Discrete random variable
 - Continuous random variable
- Data
- Probability distribution:
 - Discrete distribution: P(event) is described by the probability mass function (PMF)
 - Continuous distribution: P(event) is described by the probability density function (PDF)

What are their differences?

- Conditional probability of events
- Independent events
- Bayes' rule





Terminology
Some probability distributions that you should know by heart

Some probability distributions that you should know by heart





Probability distribution	Continuous/discrete	Apply to data type
Bernoulli distribution	Discrete	Categorical (nominal)
Categorical distribution	Discrete	Categorical (nominal)
Discrete uniform	Discrete	Numerical (discrete)
Gaussian distribution	Continuous	Numerical (continuous)





Probability distribution	Continuous/discrete	Apply to data type
Bernoulli distribution	Discrete	Categorical (nominal)
Categorical distribution	Discrete	Categorical (nominal)
Discrete uniform	Discrete	Numerical (discrete)
Gaussian distribution	Continuous	Numerical (continuous)

Recall

- Discrete distribution is described by the probability mass function (PMF)
- Continuous distribution is described by the probability density function (PDF)





Probability distribution	Continuous/discrete	Apply to data type
Bernoulli distribution	Discrete	Categorical (nominal)
Categorical distribution	Discrete	Categorical (nominal)
Discrete uniform	Discrete	Numerical (discrete)
Gaussian distribution	Continuous	Numerical (continuous)

Recall

- Discrete distribution is described by the probability mass function (PMF)
- Continuous distribution is described by the probability density function (PDF)

For each distribution, you need to know:

- its PMF or PDF: the equation and the shape
- its parameters
- its applications





Probability distribution	Continuous/discrete	Apply to data type
Bernoulli distribution	Discrete	Categorical (nominal)
Categorical distribution	Discrete	Categorical (nominal)
Discrete uniform	Discrete	Numerical (discrete)
Gaussian distribution	Continuous	Numerical (continuous)

Recall

- Discrete distribution is described by the probability mass function (PMF)
- Continuous distribution is described by the probability density function (PDF)

For each distribution, you need to know:

- its PMF or PDF: the equation and the shape
- its parameters
- its applications
- how to estimate the parameters (next lecture)





Probability distribution	Continuous/discrete	Apply to data type
Bernoulli distribution	Discrete	Categorical (nominal)
Categorical distribution	Discrete	Categorical (nominal)
Discrete uniform	Discrete	Numerical (discrete)
Gaussian distribution	Continuous	Numerical (continuous)





In your town everybody has ducks (of course). Ducks in this town ONLY have TWO colors: blue and red. What is the probability distribution we use to describe the duck colors in your town?





In your town everybody has ducks (of course). Ducks in this town ONLY have TWO colors: blue and red. What is the probability distribution we use to describe the duck colors in your town?

• Let X be a discrete random variable $X = \begin{cases} 0 & \text{a duck is red} \\ 1 & \text{a duck is blue} \end{cases}$



In your town everybody has ducks (of course). Ducks in this town ONLY have TWO colors: blue and red. What is the probability distribution we use to describe the duck colors in your town?

- Let X be a discrete random variable $X = \begin{cases} 0 & \text{a duck is red} \\ 1 & \text{a duck is blue} \end{cases}$
- Given *p* the probability of a duck being blue, we can express the probability distribution as follows:

$$P(a \text{ duck is red}) = P(X = 0) = 1 - p$$

 $P(a \text{ duck is blue}) = P(X = 1) = p$





In your town everybody has ducks (of course). Ducks in this town ONLY have TWO colors: blue and red. What is the probability distribution we use to describe the duck colors in your town?

- Let X be a discrete random variable $X = \begin{cases} 0 & \text{a duck is red} \\ 1 & \text{a duck is blue} \end{cases}$
- Given p the probability of a duck being blue, we can express the probability distribution as follows:

$$P(a \text{ duck is red}) = P(X = 0) = 1 - p$$

 $P(a \text{ duck is blue}) = P(X = 1) = p$

• What is the PMF?



In your town everybody has ducks (of course). Ducks in this town ONLY have TWO colors: blue and red. What is the probability distribution we use to describe the duck colors in your town?

- Let X be a discrete random variable $X = \begin{cases} 0 & \text{a duck is red} \\ 1 & \text{a duck is blue} \end{cases}$
- Given *p* the probability of a duck being blue, we can express the probability distribution as follows:

$$P(a \text{ duck is red}) = P(X = 0) = 1 - p$$

 $P(a \text{ duck is blue}) = P(X = 1) = p$

What is the PMF?Merge these two equations:

$$P(X=k) = f_X(k) \equiv f_X(k \mid p) = pk + (1-p)(1-k), \ k \in \{0,1\}, p \in [0,1]$$

Note: here we use a \mid to indicate that the parameter p is given.






- Discrete distribution
- Applies to nominal data with 2 categories
- PMF:
 - Equation

$$f_X(k \mid p) = pk + (1-p)(1-k), k \in \{0,1\}, p \in [0,1]$$

Shape



Parameters: p





Probability distribution	Continuous/discrete	Apply to data type
Bernoulli distribution	Discrete	Categorical (nominal)
Categorical distribution	Discrete	Categorical (nominal)
Discrete uniform	Discrete	Numerical (discrete)
Gaussian distribution	Continuous	Numerical (continuous)





Why probability distributions?
Terminology
Some probability distributions that you should know by heart

Categorical distribution

In Jack's town, ducks have FOUR colors: blue, red, green and gray. What is the probability distribution of duck colors in Jack's town?





In Jack's town, ducks have FOUR colors: blue, red, green and gray. What is the probability distribution of duck colors in Jack's town?

Given p₁ the probability of a duck being blue, p₂ the probability of a duck being red, p₃ the probability of a duck being green and p₄ the probability of a duck being gray. Note that p₁ + p₂ + p₃ + p₄ = 1.



In Jack's town, ducks have FOUR colors: blue, red, green and gray. What is the probability distribution of duck colors in Jack's town?

• Given p_1 the probability of a duck being blue, p_2 the probability of a duck being red, p_3 the probability of a duck being green and p_4 the probability of a duck being gray. Note that $p_1 + p_2 + p_3 + p_4 = 1$.

• Let
$$X$$
 be a discrete random variable $X = \begin{cases} 1 & \text{a duck is blue} \\ 2 & \text{a duck is red} \\ 3 & \text{a duck is green} \\ 4 & \text{a duck is gray} \end{cases}$





In Jack's town, ducks have FOUR colors: blue, red, green and gray. What is the probability distribution of duck colors in Jack's town?

- Given p_1 the probability of a duck being blue, p_2 the probability of a duck being red, p_3 the probability of a duck being green and p_4 the probability of a duck being gray. Note that $p_1 + p_2 + p_3 + p_4 = 1$.
- Let X be a discrete random variable $X = \begin{cases} 1 & \text{a duck is blue} \\ 2 & \text{a duck is red} \\ 3 & \text{a duck is green} \\ 4 & \text{a duck is gray} \end{cases}$
- Now we can express the probability distribution as follows:

$$P(a \text{ duck is blue}) = P(X = 1) = p_1$$

$$P(a \text{ duck is red}) = P(X = 2) = p_2$$

$$P(a \text{ duck is green}) = P(X = 3) = p_3$$

$$P(a \text{ duck is gray}) = P(X = 4) = p_4$$



Some probability distributions that you should know by heart

In Jack's town, ducks have FOUR colors: blue, red, green and gray. What is the probability distribution of duck colors in Jack's town?

- Given p_1 the probability of a duck being blue, p_2 the probability of a duck being red, p_3 the probability of a duck being green and p_4 the probability of a duck being gray. Note that $p_1 + p_2 + p_3 + p_4 = 1$.
- Let X be a discrete random variable $X = \begin{cases} 1 & \text{a duck is blue} \\ 2 & \text{a duck is red} \\ 3 & \text{a duck is green} \\ 4 & \text{a duck is gray} \end{cases}$
- Now we can express the probability distribution as follows:

$$P(\text{a duck is blue}) = P(X = 1) = p_1$$

 $P(\text{a duck is red}) = P(X = 2) = p_2$

$$P(a \text{ duck is green}) = P(X = 3) = p_3$$

$$P(a \text{ duck is gray}) = P(X = 4) = p_4$$

What is the PMF?



In Jack's town, ducks have FOUR colors: blue, red, green and gray. What is the probability distribution of duck colors in Jack's town?

- Given p_1 the probability of a duck being blue, p_2 the probability of a duck being red, p_3 the probability of a duck being green and p_4 the probability of a duck being gray. Note that $p_1 + p_2 + p_3 + p_4 = 1$.
- Let X be a discrete random variable $X = \begin{cases} 1 & \text{a duck is blue} \\ 2 & \text{a duck is red} \\ 3 & \text{a duck is green} \\ 4 & \text{a duck is gray} \end{cases}$
- Now we can express the probability distribution as follows:

$$P(a \text{ duck is blue}) = P(X = 1) = p_1$$

 $P(a \text{ duck is red}) = P(X = 2) = p_2$

$$P(\text{a duck is red}) = P(X = 2) = p_2$$

$$P(a \text{ duck is green}) = P(X = 3) = p_3$$

$$P(a \text{ duck is gray}) = P(X = 4) = p_4$$

What is the PMF?

$$P(X = k) = f_X(k) \equiv f_X(k \mid p_1, p_2, p_3, p_4) = p_k, \sum_{i=1}^4 p_i = 1, p_i \ge 0, \ k \in \{1, \dots, 4\}$$





In Jack's town, ducks have FOUR colors: blue, red, green and gray. What is the probability distribution of duck colors in Jack's town?

- Given p₁ the probability of a duck being blue, p₂ the probability of a duck being red, p₃ the probability of a duck being green and p₄ the probability of a duck being gray. Note that p₁ + p₂ + p₃ + p₄ = 1.
- Let X be a discrete random variable $X = \begin{cases} 1 & \text{a duck is blue} \\ 2 & \text{a duck is red} \\ 3 & \text{a duck is green} \\ 4 & \text{a duck is gray} \end{cases}$
- Now we can express the probability distribution as follows:
 - $P(a ext{ duck is blue}) = P(X = 1) = p_1$ $P(a ext{ duck is red}) = P(X = 2) = p_2$ $P(a ext{ duck is green}) = P(X = 3) = p_3$ $P(a ext{ duck is gray}) = P(X = 4) = p_4$
- What is the PMF?

$$P(X = k) = f_X(k) \equiv f_X(k \mid p_1, p_2, p_3, p_4) = p_k, \sum_{i=1}^4 p_i = 1, p_i \ge 0, \ k \in \{1, \dots, 4\}$$

Note: categorical distribution is also called multinoulli distribution. It is a generalization of the Bernoulli distribution.






- Discrete distribution
- Applies to nominal data with n > 0 categories
- PMF:
 - Equation

$$f_X(k \mid p_1, p_2, \dots, p_n) = p_k, \sum_{i=1}^n p_i = 1, p_i \ge 0, k \in \{1, \dots, n\}$$

Shape



• Parameters: p_k , $k \in \{1, \dots, n\}$ for given n; n-1 parameters $(\sum_{i=1}^n p_i = 1)$.





Probability distribution	Continuous/discrete	Apply to data type
Bernoulli distribution	Discrete	Categorical (nominal)
Categorical distribution	Discrete	Categorical (nominal)
Discrete uniform	Discrete	Numerical (discrete)
Gaussian distribution	Continuous	Numerical (continuous)



Discrete uniform distribution

Meanwhile, back to your town, a team of scientists crunched some numbers and they stated that the number of ducks that each person has follows a uniform distribution between 1 and 1000. What does that mean?





Discrete uniform distribution

Meanwhile, back to your town, a team of scientists crunched some numbers and they stated that the number of ducks that each person has follows a uniform distribution between 1 and 1000. What does that mean?

ullet Given integers a and b with $a \leq b$ (here we have a=1 and b=1000)



Meanwhile, back to your town, a team of scientists crunched some numbers and they stated that the number of ducks that each person has follows a uniform distribution between 1 and 1000. What does that mean?

- Given integers a and b with $a \le b$ (here we have a=1 and b=1000)
- Let X be a discrete random variable: X = k if a person has k ducks



Meanwhile, back to your town, a team of scientists crunched some numbers and they stated that the number of ducks that each person has follows a uniform distribution between 1 and 1000. What does that mean?

- Given integers a and b with $a \le b$ (here we have a = 1 and b = 1000)
- Let X be a discrete random variable: X = k if a person has k ducks
- We can express the probability distribution as follows:

$$P(a person has k ducks) = P(X = k) = \frac{1}{b-a+1}$$





Meanwhile, back to your town, a team of scientists crunched some numbers and they stated that the number of ducks that each person has follows a uniform distribution between 1 and 1000. What does that mean?

- ullet Given integers a and b with $a \leq b$ (here we have a=1 and b=1000)
- Let X be a discrete random variable: X = k if a person has k ducks
- We can express the probability distribution as follows: $P(a \text{ person has } k \text{ ducks}) = P(X = k) = \frac{1}{b-a+1}$
- What is the PMF of a discrete uniform distribution?



Meanwhile, back to your town, a team of scientists crunched some numbers and they stated that the number of ducks that each person has follows a uniform distribution between 1 and 1000. What does that mean?

- Given integers a and b with $a \le b$ (here we have a = 1 and b = 1000)
- Let X be a discrete random variable: X = k if a person has k ducks
- We can express the probability distribution as follows: $P(a \text{ person has } k \text{ ducks}) = P(X = k) = \frac{1}{b-a+1}$
- What is the PMF of a discrete uniform distribution?

$$P(X = k) = f_X(k) \equiv f_X(k \mid a, b) = \frac{1}{b - a + 1}$$






- Discrete distribution
- Applies to discrete numerical data
- PMF:
 - Equation

$$f_X(k \mid a, b) = \frac{1}{b-a+1}, \ a \le b, \ a, b \text{ integers}$$

Shape



• Parameters: integers a, b





Probability distributions

Probability distribution	Continuous/discrete	Apply to data type
Bernoulli distribution	Discrete	Categorical (nominal)
Categorical distribution	Discrete	Categorical (nominal)
Discrete uniform	Discrete	Numerical (discrete)
Gaussian distribution	Continuous	Numerical (continuous)





Still at your town, a scientist told you that the weights of your ducks follow a Gaussian distribution with mean $\mu=7$ and standard deviation $\sigma=2$. What does that mean?





Still at your town, a scientist told you that the weights of your ducks follow a Gaussian distribution with mean $\mu=7$ and standard deviation $\sigma=2$. What does that mean?

• Let X be a continuous random variable: X = x if a duck weighs x kg





Still at your town, a scientist told you that the weights of your ducks follow a Gaussian distribution with mean $\mu=7$ and standard deviation $\sigma=2$. What does that mean?

- Let X be a continuous random variable: X = x if a duck weighs x kg
- Let $f_X(x)$ be the PDF, the probability distribution can be expressed as (10 secs):



Still at your town, a scientist told you that the weights of your ducks follow a Gaussian distribution with mean $\mu=7$ and standard deviation $\sigma=2$. What does that mean?

- Let X be a continuous random variable: X = x if a duck weighs x kg
- Let $f_X(x)$ be the PDF, the probability distribution can be expressed as (10 secs):

$$P(a \le X \le b) = \int_a^b f_X(x) dx$$
 for all $a < b$



Still at your town, a scientist told you that the weights of your ducks follow a Gaussian distribution with mean $\mu=7$ and standard deviation $\sigma=2$. What does that mean?

- Let X be a continuous random variable: X = x if a duck weighs x kg
- Let $f_X(x)$ be the PDF, the probability distribution can be expressed as (10 secs):

$$P(a \le X \le b) = \int_a^b f_X(x) dx$$
 for all $a < b$

• What is the PDF of a Gaussian distribution with $\mu = 7$ and $\sigma = 2$?



Still at your town, a scientist told you that the weights of your ducks follow a Gaussian distribution with mean $\mu = 7$ and standard deviation $\sigma = 2$. What does that mean?

- Let X be a continuous random variable: X = x if a duck weighs x kg
- Let $f_X(x)$ be the PDF, the probability distribution can be expressed as (10 secs):

$$P(a \le X \le b) = \int_a^b f_X(x) dx$$
 for all $a < b$

• What is the PDF of a Gaussian distribution with $\mu = 7$ and $\sigma = 2$?


$$f_X(x \mid \mu, \sigma) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{1}{2}(\frac{x-\mu}{\sigma})^2} = \frac{1}{\sqrt{8\pi}} e^{-\frac{1}{2}(\frac{x-7}{2})^2}$$



- Continuous distribution
- Applies to continuous numerical data
- PDF:
- Equation

$$f_X(x) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{1}{2}\left(\frac{x-\mu}{\sigma}\right)^2}, \ \mu \in \mathbb{R}, \ \sigma \in \mathbb{R}^+$$

Shape



• Parameters: $\mu, \ \sigma$





Probability distributions

Probability distribution	Continuous/discrete	Apply to data type
Bernoulli distribution	Discrete	Categorical (nominal)
Categorical distribution	Discrete	Categorical (nominal)
Discrete uniform	Discrete	Numerical (discrete)
Gaussian distribution	Continuous	Numerical (continuous)

Hooray!



An important note





An important note

These probability distributions DO NOT ONLY apply to duck related applications!

Check out what data types they apply to!

We are going to talk about more applications in the future





An important note

These probability distributions DO NOT ONLY apply to duck related applications!

Check out what data types they apply to!

We are going to talk about more applications in the future (even though they won't be as important as ducks)





Today

- Probability distribution
- 2 Demo
- Summary





Demo

Code demo

- Image processing
- Natural language processing
- Table with numerical data
- Table with categorical data





Today

- Probability distribution
- 2 Demo
- Summary





- Data types, data containers, descriptive statistics (e.g. sample mean, sample variance, data quantile), visualization (e.g. histogram)
- Probability distributions, sample space, events, random variables, PMF, PDF, parameters



- Data types, data containers, descriptive statistics (e.g. sample mean, sample variance, data quantile), visualization (e.g. histogram)
- Probability distributions, sample space, events, random variables, PMF, PDF, parameters

Not yet:

- How to estimate parameters, such as μ and σ in a Gaussian distribution?
- How to choose the probability distribution for a given data set?



- Data types, data containers, descriptive statistics (e.g. sample mean, sample variance, data quantile), visualization (e.g. histogram)
- Probability distributions, sample space, events, random variables, PMF, PDF, parameters

Not yet:

- \bullet How to estimate parameters, such as μ and σ in a Gaussian distribution?
- How to choose the probability distribution for a given data set?

Next:

Q-Q plot and mathematical modeling





- Data types, data containers, descriptive statistics (e.g. sample mean, sample variance, data quantile), visualization (e.g. histogram)
- Probability distributions, sample space, events, random variables, PMF, PDF, parameters

Not yet:

- How to estimate parameters, such as μ and σ in a Gaussian distribution?
- How to choose the probability distribution for a given data set?

Next:

Q-Q plot and mathematical modeling

Before next lecture:

- Quantile
- Probability distributions from today and their parameters
- PMF and PDF





Stay safe!





