


Gaussian Naïve Bayes

Nazerfard, Ehsan
nazerfard@aut.ac.ir



Gaussian/Normal Distribution

- A lot of things in nature follow the normal distribution, such as people's height, etc.



Gaussian Distribution

- A type of continuous distribution
- (Univariate) probability density function:

$$f(x) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{1}{2}\left(\frac{x-\mu}{\sigma}\right)^2}$$

- where μ and σ^2 are the mean and variance of the distribution.



Gaussian Distribution (cont.)

- (Multivariate) probability density function:

$$f_{\mathbf{X}}(x_1, \dots, x_k) = \frac{1}{\sqrt{(2\pi)^k |\Sigma|}} e^{-\frac{1}{2}(\mathbf{X}-\boldsymbol{\mu})^T \Sigma^{-1} (\mathbf{X}-\boldsymbol{\mu})}$$

$$\mathbf{X} = (X_1, \dots, X_k)^T$$

- where $\mu_{k \times 1}$ and $\Sigma_{k \times k}$ are the mean vector and covariance matrix of the distribution:
- $\boldsymbol{\mu} = \mathbb{E}[\mathbf{X}] = [\mathbb{E}[X_1], \dots, \mathbb{E}[X_k]]^T$
- $\Sigma = \mathbb{E}[(\mathbf{X} - \boldsymbol{\mu})(\mathbf{X} - \boldsymbol{\mu})^T]$

Gaussian naïve Bayes (GNB)

- ❑ How to distinguish children from adults, using a Gaussian naïve Bayes classifier?
 - Classes: {a,c}
 - Features: height and weight (continuous inputs)

- ❑ Training data $\{h_i, w_i, y_i\}$
 - 12 children (black)
 - 4 adults (red)




Figure © Victor Lavrenko, Lecture on GNB, University of Edinburgh

Prior Probabilities

□ Prior class probabilities:

- $p(Y = a) = 4/16 = 0.25$
- $p(Y = c) = 12/16 = 0.75$




Figure © Victor Lavrenko, Lecture on GNB, University of Edinburgh

Children's Features


- Assuming that children's heights and weights follow the Gaussian distribution:

- $\mu_{h,c} = \frac{1}{12} \sum_{i:y_i=c} h^i$

- $\sigma_{h,c}^2 = \frac{1}{12-1} \sum_{i:y_i=c} (h^i - \mu_{h,c})^2$

- $\mu_{w,c} = \frac{1}{12} \sum_{i:y_i=c} w^i$


- $\sigma_{w,c}^2 = \frac{1}{12-1} \sum_{i:y_i=c} (w^i - \mu_{w,c})^2$



2-Dimensional Gaussian Distribution


$$\boldsymbol{\mu} = \begin{bmatrix} \mu_X \\ \mu_Y \end{bmatrix}, \quad \boldsymbol{\Sigma} = \begin{bmatrix} \sigma_X^2 & \rho\sigma_X\sigma_Y \\ \rho\sigma_X\sigma_Y & \sigma_Y^2 \end{bmatrix}$$

where $\rho = \frac{\text{Cov}(X,Y)}{\sigma_X\sigma_Y}$ is the correlation coefficient between X and Y .



Independence Assumption

- Multiplying two Gaussian distributions



- The same thing for the adult's features

Test Phase

- $X = \langle h_X, w_X \rangle$ (test data)
 - $p(h_X|Y = a) = \frac{1}{\sqrt{2\pi\sigma_{h,a}^2}} \exp\left(-\frac{(h_X - \mu_{h,a})^2}{2\sigma_{h,a}^2}\right)$
 - $p(w_X|Y = a) = \frac{1}{\sqrt{2\pi\sigma_{w,a}^2}} \exp\left(-\frac{(w_X - \mu_{w,a})^2}{2\sigma_{w,a}^2}\right)$
- Similarly compute $p(h_X|Y = c)$ and $p(w_X|Y = c)$




Figure © Victor Lavrenko, Lecture on GNB, University of Edinburgh

Classification

□ Classification Rule:

$$y_{MLE} = \operatorname{argmax}_{y \in \{a, c\}} p(Y = y | X) = \operatorname{argmax}_{y \in \{a, c\}} p(X | Y = y) p(Y = y)$$

□ Test data: $X = \langle h_X, w_X \rangle$

If $p(\langle h_X, w_X \rangle | Y = a) p(Y = a) > p(\langle h_X, w_X \rangle | Y = c) p(Y = c)$

In dependent assumption
→

If $p(h_X | Y = a) p(w_X | Y = a) p(Y = a) > p(h_X | Y = c) p(w_X | Y = c) p(Y = c)$

Then Class is addult

Else Class is children

A Gentle Correction

- Population mean vs. Sample mean

$$\mu_{h,c} = \frac{1}{12} \sum_{i:y_i=c} h^i \rightarrow \bar{X}_{h,c} = \frac{1}{12} \sum_{i:y_i=c} h^i$$

- Population variance vs. Sample variance

$$\sigma_{h,c}^2 = \frac{1}{12-1} \sum_{i:y_i=c} (h^i - \mu_{h,c})^2 \rightarrow S_{h,c}^2 = \frac{1}{12-1} \sum_{i:y_i=c} (h^i - \bar{X}_{h,c})^2$$



Parametric Form of $P(Y|X)$ for GNB Classifier

□ Assumptions

- Y is boolean, governed by a Bernoulli distribution
- $\mathbf{X} = \langle X_1, X_2, \dots, X_n \rangle$, where X_i is a continuous random variable
- For each X_i , $P(X_i|Y = y_k)$ is a Gaussian distribution of the form $N(\mu_{ik}, \sigma_i)$
- For all i and $j \neq i$, X_i and X_j are conditionally independent given Y

GNB Parametric Form of $P(Y|X)$

- We derive the parametric form of $p(Y|X)$ that follows from the set of GNB assumptions:

$$\begin{aligned} p(Y = 1|\mathbf{X}) &= \frac{p(\mathbf{X}|Y = 1)p(Y = 1)}{p(\mathbf{X})} = \frac{p(\mathbf{X}|Y = 1)p(Y = 1)}{p(\mathbf{X}|Y = 1)p(Y = 1) + p(\mathbf{X}|Y = 0)p(Y = 0)} \\ &= \frac{1}{1 + \frac{p(\mathbf{X}|Y = 0)p(Y = 0)}{p(\mathbf{X}|Y = 1)p(Y = 1)}} \end{aligned}$$

GNB Parametric Form of $P(Y|X)$

- We derive the parametric form of $p(Y|X)$ that follows from the set of GNB assumptions:

$$\begin{aligned}
 p(Y = 1|\mathbf{X}) &= \frac{p(\mathbf{X}|Y = 1)p(Y = 1)}{p(\mathbf{X})} = \frac{p(\mathbf{X}|Y = 1)p(Y = 1)}{p(\mathbf{X}|Y = 1)p(Y = 1) + p(\mathbf{X}|Y = 0)p(Y = 0)} \\
 &= \frac{1}{1 + \frac{p(\mathbf{X}|Y = 0)p(Y = 0)}{p(\mathbf{X}|Y = 1)p(Y = 1)}} = \frac{1}{1 + \exp\left[\ln \frac{p(\mathbf{X}|Y = 0)p(Y = 0)}{p(\mathbf{X}|Y = 1)p(Y = 1)}\right]} \\
 &= \frac{1}{1 + \exp\left[\ln \frac{p(\mathbf{X}|Y = 0)}{p(\mathbf{X}|Y = 1)} + \ln \frac{p(Y=0)}{p(Y=1)}\right]} = \frac{1}{1 + \exp\left[\sum_{i=1}^n \ln \frac{p(x_i|Y = 0)}{p(x_i|Y = 1)} + \ln \frac{p(Y=0)}{p(Y=1)}\right]} \quad (\text{Eq. I})
 \end{aligned}$$

GNB Parametric Form of $P(Y|X)$

- The sigma term in the denominator:

$$\begin{aligned}
 \sum_{i=1}^n \ln \frac{p(x_i|Y=0)}{p(x_i|Y=1)} &= \sum_i \ln \frac{\frac{1}{\sqrt{2\pi\sigma_i^2}} \exp\left(\frac{-(X_i - \mu_{i0})^2}{2\sigma_i^2}\right)}{\frac{1}{\sqrt{2\pi\sigma_i^2}} \exp\left(\frac{-(X_i - \mu_{i1})^2}{2\sigma_i^2}\right)} \\
 &= \sum_i \ln \left(\exp \left(\frac{(X_i - \mu_{i1})^2 - (X_i - \mu_{i0})^2}{2\sigma_i^2} \right) \right) = \sum_i \frac{(X_i^2 - 2X_i\mu_{i1} + \mu_{i1}^2) - (X_i^2 - 2X_i\mu_{i0} + \mu_{i0}^2)}{2\sigma_i^2} \\
 &= \sum_i \frac{2X_i(\mu_{i0} - \mu_{i1}) + \mu_{i1}^2 - \mu_{i0}^2}{2\sigma_i^2}
 \end{aligned}$$

GNB Parametric Form of $P(Y|X)$

- The sigma term in the denominator:

$$\begin{aligned}
 \sum_{i=1}^n \ln \frac{p(x_i|Y=0)}{p(x_i|Y=1)} &= \sum_i \ln \frac{\frac{1}{\sqrt{2\pi\sigma_i^2}} \exp\left(\frac{-(X_i - \mu_{i0})^2}{2\sigma_i^2}\right)}{\frac{1}{\sqrt{2\pi\sigma_i^2}} \exp\left(\frac{-(X_i - \mu_{i1})^2}{2\sigma_i^2}\right)} \\
 &= \sum_i \ln \left(\exp \left(\frac{(X_i - \mu_{i1})^2 - (X_i - \mu_{i0})^2}{2\sigma_i^2} \right) \right) = \sum_i \frac{(X_i^2 - 2X_i\mu_{i1} + \mu_{i1}^2) - (X_i^2 - 2X_i\mu_{i0} + \mu_{i0}^2)}{2\sigma_i^2} \\
 &= \sum_i \frac{2X_i(\mu_{i0} - \mu_{i1}) + \mu_{i1}^2 - \mu_{i0}^2}{2\sigma_i^2} = \sum_i \left(\frac{\mu_{i0} - \mu_{i1}}{\sigma_i^2} X_i + \frac{\mu_{i1}^2 - \mu_{i0}^2}{2\sigma_i^2} \right) \quad (\text{Eq. II})
 \end{aligned}$$

GNB Parametric Form of $P(Y|X)$

$$I, II \Rightarrow p(Y = 1|\mathbf{X}) = \frac{1}{1 + \exp[\sum_{i=1}^n \left(\frac{\mu_{i0} - \mu_{i1}}{\sigma_i^2} X_i + \frac{\mu_{i1}^2 - \mu_{i0}^2}{2\sigma_i^2} \right) + \ln \frac{p(Y = 0)}{p(Y = 1)}]}$$

GNB Parametric Form of $P(Y|X)$

$$I, II \Rightarrow p(Y = 1|X) = \frac{1}{1 + \exp[\sum_{i=1}^n \left(\frac{\mu_{i0} - \mu_{i1}}{\sigma_i^2} X_i + \frac{\mu_{i1}^2 - \mu_{i0}^2}{2\sigma_i^2} \right) + \ln \frac{p(Y = 0)}{p(Y = 1)}]}$$

Or equivalently:

$$p(Y = 1|X) = \frac{1}{1 + \exp(\sum_{i=1}^n w_i X_i + w_0)}$$

$$\text{where } w_i = \frac{\mu_{i0} - \mu_{i1}}{\sigma_i^2}, w_0 = \ln \frac{1 - P(Y=1)}{P(Y=1)} + \sum_{i=1}^n \frac{\mu_{i1}^2 - \mu_{i0}^2}{2\sigma_i^2}$$

GNB Parametric Form of $P(Y|X)$

$$I, II \Rightarrow p(Y = 1|X) = \frac{1}{1 + \exp[\sum_{i=1}^n \left(\frac{\mu_{i0} - \mu_{i1}}{\sigma_i^2} X_i + \frac{\mu_{i1}^2 - \mu_{i0}^2}{2\sigma_i^2} \right) + \ln \frac{p(Y = 0)}{p(Y = 1)}]}$$

Or equivalently:

$$p(Y = 1|X) = \frac{1}{1 + \exp(\sum_{i=1}^n w_i X_i + w_0)}$$

$$\text{where } w_i = \frac{\mu_{i0} - \mu_{i1}}{\sigma_i^2}, w_0 = \ln \frac{1 - p(Y=1)}{p(Y=1)} + \sum_{i=1}^n \frac{\mu_{i1}^2 - \mu_{i0}^2}{2\sigma_i^2}$$

$$p(Y = 0|X) = 1 - p(Y = 1|X) = \frac{\exp(\sum_{i=1}^n w_i X_i + w_0)}{1 + \exp(\sum_{i=1}^n w_i X_i + w_0)}$$

GNB Parametric Form of $P(Y|X)$

If $p(Y = 1|X) \geq p(Y = 0 |X)$ Then Class is +

If $\frac{p(Y=1 |X)}{p(Y = 0 |X)} = \frac{1}{\exp(\sum_{i=1}^n w_i X_i + w_0)} \geq 1$ Then Class is +

GNB Parametric Form of $P(Y|X)$

If $p(Y = 1|X) \geq p(Y = 0 |X)$ Then Class is +

If $\frac{p(Y=1 |X)}{p(Y = 0 |X)} = \frac{1}{\exp(\sum_{i=1}^n w_i X_i + w_0)} \geq 1$ Then Class is +

If $\exp(\sum_{i=1}^n w_i X_i + w_0) \leq 1$ Then Class is +

If $\sum_{i=1}^n w_i X_i + w_0 \leq 0$ Then Class is +; Else Class is -

GNB Parametric Form of $P(Y|X)$

If $p(Y = 1|X) \geq p(Y = 0 |X)$ Then Class is +

If $\frac{p(Y=1 |X)}{p(Y = 0 |X)} = \frac{1}{\exp(\sum_{i=1}^n w_i X_i + w_0)} \geq 1$ Then Class is +

If $\exp(\sum_{i=1}^n w_i X_i + w_0) \leq 1$ Then Class is +

If $\sum_{i=1}^n w_i X_i + w_0 \leq 0$ Then Class is +; Else Class is -

Therefore, naïve Bayes learning algorithm can be viewed as a linear classifier, under the assumption that variance of each feature is independent of the class (i.e if $\sigma_{ik} = \sigma_i$).

References

- ❑ Victor Lavrenko, Lecture on Gaussian Naïve Bayes, University of Edinburgh
- ❑ Tom Mitchell, Generative and Discriminative Classifiers: Naïve Bayes and Logistic Regression, Machine Learning (2nd ed.), Chapter 3, McGraw Hill, 2015

