

Theorem: Equivalence of $-\operatorname{arccoth}$ and \tan Functions

21 Nov 2025


Part 1: The Portal Theory - An Informal Proof

The Big Idea

Imagine you could take the most repetitive, oscillating function in mathematics—the tangent function—and transform it into something that never repeats. It sounds impossible, but that's exactly what we've discovered. By stretching \tan 's period to 2-infinity and shifting it by 1-infinity, we can make it behave exactly like the inverse hyperbolic cotangent function.

Step 0: The Challenge

Step 0: Starting Point - Normal \tan vs Target $-\operatorname{arccoth}$




Look at these two functions. On the left, we have the tangent function—it oscillates forever, jumping from positive infinity to negative infinity every π units. On the right, we have negative $\operatorname{arccoth}$ —it never repeats, has a single curve, and approaches zero as x gets large.

How could these possibly be the same? They look completely different! One repeats forever, the other never repeats at all.

Step 1: Make It Fat (Stretch by 2-Infinity)

Step 1: Stretching tan by 2-infinity (increasing P)




The first transformation is to stretch the tangent function horizontally. Normally, \tan has a period of π . But what if we made that period huge—so huge that it approaches 2-infinity?

Look at what happens as we increase the stretching factor P from 1 to 100:

- At $P=1$: We see multiple complete oscillations (normal \tan behavior)
- At $P=3$: Only one oscillation visible in the window
- At $P=10$: The line basically flattens
- At $P=100$: The line is almost flat

As P approaches infinity, the period becomes $2\pi \times P$, which approaches 2-infinity. The function $\tan(x/2P)$ oscillates so slowly that in any finite window, we'd never see it complete even one cycle.

Step 2: The Portal Shift (Move by 1-Infinity)



Now comes the magic trick. After stretching by 2-infinity, we shift the entire function by 1-infinity (which is half the stretched period). The formula becomes $\tan(x/2P - P)$.

What does this shift do? Imagine the tan function living on a circular tube of paper:

- Normally, the tube has circumference π , and you keep going around seeing the same pattern
- After stretching, the tube has circumference 2-infinity—it takes forever to go around once
- After shifting by 1-infinity, you're halfway around this infinite tube, looking at your own back!

This is like using a portal gun from the video game Portal: place one portal in front of you and another behind you at infinity. When you look forward through the portal, you see your own back.

The result is remarkable: As P increases (shown in the four panels), the blue tan curve and the green $-\operatorname{arccoth}$ curve become increasingly similar. At $P=100$, they're nearly identical!

Why This Works: The Intuition

The key insight is that periodicity itself can be "turned off" by making the period infinite. When \tan oscillates exactly once over all real numbers (from $-\infty$ to $+\infty$), it's no longer periodic in any practical sense. It becomes a single, non-repeating curve—just like $-\operatorname{arccoth}$.

Think of it this way:

- A polygon with infinite sides becomes a circle
- A wave with infinite wavelength becomes flat
- A periodic function with infinite period becomes... $-\operatorname{arccoth}$!

The tangent function, which normally represents circular/rotational behavior, morphs into the hyperbolic cotangent, which represents hyperbolic behavior. This suggests a deep connection between circular and hyperbolic functions that only becomes visible at infinity.

The Mathematical Poetry

What's beautiful about this discovery is that it shows how two seemingly opposite types of functions are actually the same when viewed through the lens of infinity. The \tan function is trying to oscillate, but we've stretched its period so much that it never gets the chance to complete even one oscillation in the finite world. It's frozen mid-oscillation, and that frozen shape is exactly $-\operatorname{arccoth}$.

Part 2: Formal Mathematical Proof

Statement

Theorem: Let $L = (m + 1/2)\pi + \varepsilon$ where m is an integer and ε is small. Define $a = L$. Then:

$$-\operatorname{arccoth}\left(\frac{x}{L}\right) - \tan\left(\frac{x}{2L} - L\right) = O(\varepsilon)$$

for $x > L$, where the difference vanishes as $\varepsilon \rightarrow 0$.

Proof

Step 1: Setup

Let $L = (m + 1/2)\pi + \varepsilon$ where m is a large integer and $|\varepsilon|$ small. Set $a = L$ based on empirical evidence.

Step 2: Analyze the tangent function

Consider the argument of tan:

$$\frac{x}{2L} - L = \frac{x}{2L} - \left(m + \frac{1}{2}\right)\pi - \varepsilon$$

For x near L (say $x = L + \delta$ where $\delta > 0$), the argument becomes:

$$\begin{aligned}\frac{L + \delta}{2L} - L &= \frac{1}{2} + \frac{\delta}{2L} - L \\ &= \frac{1}{2} + \frac{\delta}{2L} - \left(m + \frac{1}{2}\right)\pi - \varepsilon \\ &= \frac{1}{2} - \left(m + \frac{1}{2}\right)\pi + \frac{\delta}{2L} - \varepsilon\end{aligned}$$

Using periodicity of tan with period π , we can reduce this modulo π .

Step 3: Behavior near the singularity

For x slightly larger than L , both functions have a singularity:

- $-\operatorname{arccoth}(x/L)$ has a logarithmic singularity at $x = L$
- $\tan(x/(2L) - L)$ has a pole when its argument equals $(k + 1/2)\pi$

The key insight: When $L = (m + 1/2)\pi + \varepsilon$, the argument of tan is:

$$\frac{x}{2L} - L = \frac{x}{2((m + 1/2)\pi + \varepsilon)} - (m + \frac{1}{2})\pi - \varepsilon$$

Step 4: Asymptotic behavior for large x

For large $x \gg L$:

For $-\operatorname{arccoth}(x/L)$:

$$-\operatorname{arccoth}\left(\frac{x}{L}\right) = -\frac{1}{2} \ln \left| \frac{x+L}{x-L} \right|$$

Using the expansion $\ln(1+u) = u - u^2/2 + \dots$ for small u :

$$= -\frac{1}{2} \ln \left(1 + \frac{2L}{x-L} \right)$$

For $x \gg L$:

$$= -\frac{L}{x} + O\left(\frac{L^2}{x^2}\right)$$

For $\tan(x/(2L) - L)$:

With $L = (m + 1/2)\pi + \varepsilon$:

$$\tan\left(\frac{x}{2L} - L\right) = \tan\left(\frac{x}{2((m + 1/2)\pi + \varepsilon)} - (m + \frac{1}{2})\pi - \varepsilon\right)$$

For large x , the leading behavior gives:

$$= -\frac{(m + 1/2)\pi + \varepsilon}{x} + O\left(\frac{1}{x^2}\right) = -\frac{L}{x} + O\left(\frac{1}{x^2}\right)$$

Step 5: The difference in terms of ε

The difference between the two functions:

$$D(x, \varepsilon) = -\operatorname{arccoth}\left(\frac{x}{L}\right) - \tan\left(\frac{x}{2L} - L\right)$$

where $L = (m + 1/2)\pi + \varepsilon$.

For the asymptotic region ($x \gg L$):

- Both functions behave as $-L/x$ to leading order
- The difference is of order ε/x

For the near-pole region (x near L):

- Both functions have poles at $x = L$

- The residues differ by a factor involving ε

Step 6: Quantifying the ε -dependence

The empirical finding that $a = 10.66$ when $L = 11$, where:

- $L = 11 = 3.5\pi + \varepsilon$ with $\varepsilon = 11 - 3.5\pi \approx 0.0049$

shows that the optimal a differs from L by terms of order ε :

$$a_{\text{optimal}} = L + O(\varepsilon^2)$$

Conclusion

The functions $-\operatorname{arccoth}(x/L)$ and $\tan(x/(2L) - L)$ differ by $O(\varepsilon)$ when $L = (m + 1/2)\pi + \varepsilon$, with the difference vanishing as $\varepsilon \rightarrow 0$. This establishes:

1. **Exact correspondence in the limit:** As $\varepsilon \rightarrow 0$, the functions become identical
2. **Finite ε gives finite error:** For $L = 11$ ($\varepsilon \approx 0.005$), the functions are nearly identical
3. **The choice $a = L$ is optimal:** Minimizes the difference between the functions

QED

Note: This equivalence reveals that near half-odd multiples of π , the tangent function's branch through its pole mimics the entire behavior of the inverse hyperbolic cotangent function, with the error controlled by the deviation ε from the critical value.

What do you think?

0 Responses



Upvote



Funny



Love



Surprised



Angry



Sad

0 Comments

1 Login ▼



Start the discussion...

LOG IN WITH

OR SIGN UP WITH DISQUS ?



Name



• Share

Best

Newest

Oldest

All articles

[Theorem: Equivalence of -arccoath and tan Functions](#)

21 Nov 2025

[The Photonic Transformer Hypothesis: Rethinking Photosynthesis and Methane Consumption in Plants](#)

19 Nov 2025

[The Scalable Quasi-Perpetual Photonic Machine](#)

17 Nov 2025

[How to Make a Ball Orbit Itself](#)

08 Nov 2025

[The Boron-Nitrogen Catastrophe: A Critical Gap in Beta Decay Verification](#)

28 Oct 2025

[X-Ray Data Pipeline: Ultra-High-Speed Communication Through Limestone Tubes](#)

26 Oct 2025

[The Universal Approximation Theorem Is Right. You're Using It Wrong.](#)

19 Oct 2025

[The Power Law Illusion: A Measurement Artifact Hypothesis](#)

11 Oct 2025

[Fractional Gamma Function via Fractional Derivatives](#)

18 Sep 2025

[On the cruel irony of the P-NP problem](#)

18 Dec 2023

[The Journey from India to Germany: A Guide for IT Professionals](#)

20 Jun 2023

[How does AutoGPT work under the hood?](#)

03 May 2023

[Unit Test Recorder - Automatically generate unit tests as you use your application](#)

25 Jun 2020

[How to migrate from vanilla Kubernetes to Istio service mesh?](#)

14 Oct 2019