

MA 211
Notes

Patrick Rall

Contents

Chapter 1	Module 01: Limits	Page 2
1.1	The Idea of Limits	2
1.2	Definitions of Limits	4
1.3	Techniques for Computing Limits	5
Chapter 2	Module 02: Derivatives	Page 7
2.1	Introducing the Derivative	7
2.2	Derivative as a Function	9

Chapter 1

Module 01: Limits


1.1 The Idea of Limits

Average Velocity

The average velocity over time interval $[t_0, t]$ is the change in position divided by the time elapsed.

$$v_{avg} = \frac{s(t) - s(t_0)}{t - t_0}$$

The average velocity is simply the slope of the line joining points $(t_0, s(t_0))$ and $(t, s(t))$ on the graph.




Instantaneous Velocity

Instantaneous velocity is determined by computing average velocities over time interval $[t_0, t]$ that decrease in length. As t approaches t_0 , the v_{avg} approaches the instantaneous velocity $v(t_0)$.

$$v(t_0) = \lim_{t \rightarrow t_0} \frac{s(t) - s(t_0)}{t - t_0}$$

Slope of the Tangent Line

As we take the limit of v_{avg} as t approaches t_0 , we get a tangent line.



We can define find the slope of the tangent line by finding the limit of the slope of the secant line as t approaches t_0 .

$$m(t_0) = \lim_{t \rightarrow t_0} \frac{s(t) - s(t_0)}{t - t_0}.$$

The limit defines the instantaneous velocity, therefore the instantaneous velocity at point P of the graph is the slope of the tangent line at point P of the position curve.

There are parallels to the idea of limits. As $t \rightarrow 1$, the average velocity approaches the instantaneous velocity, just as the secant line approaches the tangent line.

1.2 Definitions of Limits

Defining the Limit of a Function (Preliminary)

The function f is defined for all x except possibly at a . If $f(x)$ is arbitrarily close to L for all x sufficiently close to a , then $f(x)$ is said to be **tending to L as x approaches a** . The limit of $f(x)$ as x approaches a is denoted by

$$\lim_{x \rightarrow a} f(x) = L.$$

Often times the $\lim_{x \rightarrow a} f(x) = f(a)$, but this is not always the case and other methods for finding limits are available.

Finding Limits from a Graph

Finding Limits from a Table

We create a table of values of a function corresponding to values of x that are close to a .

Example 1.2.1

Make a conjecture about $\lim_{x \rightarrow 1} \frac{\sqrt{x}-1}{x-1}$ using the tabular method.

x	0.9	0.99	0.999	0.9999	1.0001	1.001	1.01	1.1
$f(x)$	0.5132	0.5013	0.5001	0.5000	0.5000	0.4999	0.4988	0.4881

We can make a mathematical conjecture that the $\lim_{x \rightarrow 1} \frac{\sqrt{x}-1}{x-1} = 0.5$.

One-Sided Limits

The $\lim_{x \rightarrow a} f(x) = L$ is referred to as two-sided limit because $f(x)$ approaches L as x approaches a from both sides. However, there are times when we are only interested in the limit as x approaches a from one side. In this case, we use the notation $\lim_{x \rightarrow a^+} f(x) = L$ or $\lim_{x \rightarrow a^-} f(x) = L$ to denote the limit as x approaches a from the positive side or the negative side, respectively.

Definition of One-Sided Limits

1. Right Sided Limits Suppose f is defined for all x for all x near a with $x > a$. If $f(x)$ is arbitrarily close to L for all x sufficiently close to a with $x > a$, we write,

$$\lim_{x \rightarrow a^+} f(x) = L.$$

2. Left Sided Limits Analogously, suppose f is defined for all x for all x near a with $x < a$. If $f(x)$ is arbitrarily close to L for all x sufficiently close to a with $x < a$, we write,

$$\lim_{x \rightarrow a^-} f(x) = L.$$

Theorem 1.2.1

The $\lim_{x \rightarrow a} f(x) = L$ if and only if $\lim_{x \rightarrow a^-} f(x) = \lim_{x \rightarrow a^+} f(x) = L$.

1.3 Techniques for Computing Limits

This section of the notes will cover the techniques for precisely computing the limits of functions without having the need to use mathematical evidence to create a conjecture.

Limits of Linear Functions

If $f(x) = mx + b$, then the $\lim_{x \rightarrow a} f(x) = f(a)$. Linear functions allow for direct substitution of x with a to find the limit.

Theorem 1.3.1

Let a, b and m be real numbers. For linear functions $f(x) = mx + b$, we have

$$\lim_{x \rightarrow a} f(x) = ma + b.$$

Limit Laws

Theorem 1.3.2 Limit Laws

1. Sum Law $\lim_{x \rightarrow a} (f(x) + g(x)) = \lim_{x \rightarrow a} f(x) + \lim_{x \rightarrow a} g(x)$
2. Difference Law $\lim_{x \rightarrow a} (f(x) - g(x)) = \lim_{x \rightarrow a} f(x) - \lim_{x \rightarrow a} g(x)$
3. Constant Multiple Law $\lim_{x \rightarrow a} (cf(x)) = c \lim_{x \rightarrow a} f(x)$
4. Product Law $\lim_{x \rightarrow a} (f(x)g(x)) = (\lim_{x \rightarrow a} f(x))(\lim_{x \rightarrow a} g(x))$
5. Quotient Law $\lim_{x \rightarrow a} \left(\frac{f(x)}{g(x)} \right) = \frac{\lim_{x \rightarrow a} f(x)}{\lim_{x \rightarrow a} g(x)}$, provided $\lim_{x \rightarrow a} g(x) \neq 0$
6. Power Law $\lim_{x \rightarrow a} (f(x))^n = (\lim_{x \rightarrow a} f(x))^n$
7. Root Law $\lim_{x \rightarrow a} (f(x))^{1/n} = (\lim_{x \rightarrow a} f(x))^{1/n}$, provided $f(x) > 0$, for x near a , if n is even

Limits of Polynomial Functions and Rational Functions

The limit laws are used to find the limits of polynomial functions and rational functions. The limit of a polynomial function is found by direct substitution, $\lim_{x \rightarrow a} p(x) = p(a)$. And using the quotient limit law, we can find the limit of rational functions,

$$\lim_{x \rightarrow a} \frac{p(x)}{q(x)} = \frac{\lim_{x \rightarrow a} p(x)}{\lim_{x \rightarrow a} q(x)} = \frac{p(a)}{q(a)}, \text{ provided } q(a) \neq 0.$$

Theorem 1.3.3 Limits of Polynomial and Rational Functions

1. Polynomial functions: $\lim_{x \rightarrow a} p(x) = p(a)$
2. Rational functions: $\lim_{x \rightarrow a} \frac{p(x)}{q(x)} = \frac{p(a)}{q(a)}$, provided $q(a) \neq 0$

One Sided Limits

Theorem 1.3.1, 1.3.2, and 1.3.3 can be used to find the one-sided limits of functions. The one-sided limits are found by using the one-sided limit laws.

Theorem 1.3.4 One Sided Limits

Limit Laws 1-6 hold then the $\lim_{x \rightarrow a}$ replaced with $\lim_{x \rightarrow a^+}$ or $\lim_{x \rightarrow a^-}$. Law 7 is modified as follows. Assume $n > 0$ is an integer.

7. Root a. $\lim_{x \rightarrow a^+} (f(x))^{1/n} = (\lim_{x \rightarrow a^+} f(x))^{1/n}$, provided $f(x) \geq 0$, for x near a with $x > a$, if n is even
- b. $\lim_{x \rightarrow a^-} (f(x))^{1/n} = (\lim_{x \rightarrow a^-} f(x))^{1/n}$, provided $f(x) \geq 0$, for x near a with $x < a$, if n is even

Other Techniques for Computing Limits

1. Factor and cancel
2. Use Conjugates

Squeeze Theorem

Chapter 2

Module 02: Derivatives

2.1 Introducing the Derivative

The problem that was introduced in Chapter 2 about finding the slope of the tangent is important for several reasons.

- We identify the slope of the tangent line with the *instantaneous rate of change* of a function.
- The slopes of the tangent line as they change along a curve are the values of a new function called the *derivative*.
- If a curve represents the trajectory of a moving object, the tangent line at a point on the curve indicates the direction of the motion at that point.


The instantaneous velocity at time $t = a$ is the limit of the average velocity as $t \rightarrow a$:

$$v_{inst} = \lim_{t \rightarrow a} \frac{s(t) - s(a)}{t - a}$$

We also learned that there are special geometric meanings to these quantities. The average velocity is the slope of the secant line and the instantaneous velocity is the slope of the tangent line.

Tangent lines and Rates of Change

Consider the curve $y = f(x)$ and a secant line intersecting the curve at points $P(a, f(a))$ and $Q(x, f(x))$. The difference $f(x) - f(a)$ is the change in the value of f on the interval $[a, x]$, while $x - a$ is the change in x . As discussed before the slope of the secant line \overleftrightarrow{PQ} , is the $\lim_{x \rightarrow a} \frac{f(x) - f(a)}{x - a}$



Definition Rate of Change and the Slope of the Tangent Line

The **average rate of change** in f on the interval $[a, x]$ is the corresponding secant line:

$$m_{sec} = \frac{f(x) - f(a)}{x - a}$$

The **instantaneous rate of change** in f at a is

$$m_{tan} = \lim_{x \rightarrow a} \frac{f(x) - f(a)}{x - a}$$

which is also the slope of the tangent line at $(a, f(a))$ provided that the limit exists. Tangent line through $(a, f(a))$ with slope m_{tan} . Its equation is

$$y - f(a) = m_{tan}(x - a)$$

Alternative Definition Rate of Change and the Slope of the Tangent Line

The **average rate of change** in f on the interval $[a, a + h]$ is the slope of the corresponding secant line:

$$m_{sec} = \frac{f(a + h) - f(a)}{h}$$

The **instantaneous rate of change** in f at a is

$$m_{tan} = \lim_{h \rightarrow 0} \frac{f(a + h) - f(a)}{h}$$

The Derivative

Computing the slope of the line tangent to the graph of a function f at a point a gives us the instantaneous rate of change in f at a . This is called the derivative.

The **derivative of f at a** written,


$$f'(a) = \lim_{h \rightarrow 0} \frac{f(a + h) - f(a)}{h}$$

2.2 Derivative as a Function

In this section we now extend our understandings of derivatives to all points in the domain f to create new function called the derivative of f .

The Derivative Function

Due to the fact that the tangent of any given function moves, the slope of the tangent line will also change. For this reason, the slope on the tangent line for the function f is itself a function, call the *derivative* of f .



Definition The Derivative Function

The **derivative** f' is the function

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

provided the limit exists and x is in the domain f .

Example 2.2.1 (Find the derivative of $f(x) = -x^2 + 6x$)

$$\begin{aligned} f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{-(x+h)^2 + 6(x+h) - (-x^2 + 6x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{-(x^2 + 2xh + h^2) + 6x + 6h + x^2 - 6x}{h} \\ &= \lim_{h \rightarrow 0} \frac{h(-2x - h + 6)}{h} \\ &= \lim_{h \rightarrow 0} (-2x - h + 6) \\ &= \boxed{-2x + 6} \end{aligned}$$

Derivative Notation

Several notations for the derivatives are used. Recall the equation for the slope of the secant line,

$$m_{sec} = \frac{f(x + \Delta x) - f(x)}{\Delta x} = \frac{\Delta y}{\Delta x}$$

By letting $\Delta x \rightarrow 0$, the slope of the tangent line at $(x, f(x))$ is

$$f'(x) = \lim_{\Delta x \rightarrow 0} \frac{(f(x + \Delta x) - f(x))}{\Delta x} = \lim_{\Delta x \rightarrow 0} \frac{\Delta y}{\Delta x} = \frac{dy}{dx}$$

Continuity

If a function is differentiable at a point, then it is also continuous at that point.

Theorem 2.2.1 Differentiable Implies Continuous

If f is differentiable at a , then f is continuous at a .