Master thesis notes

Giacomo Galliani

Academic year 2026-2027

capitolo 1 **NLO computations in the subtraction scheme** Page 2

1.1 The singularities of the NLO real corrections for DY processes 2
The soft limit — 3 • The collinear limit — 4

1.2 The subtraction scheme for $q\bar{q} \to V + g$ 4

1 NLO computations in the subtraction scheme

1.1 The singularities of the NLO real corrections for DY processes

Consider the production of a lepton pair, by annichilation of a quark and a antiquark. These process are known as DY process. At LO (in α_s), there are two diagrams, when the exchange is mediated by a photon or a Z boson. The amplitude is known. We label p_1 the momentum of the quark, and p_2 of the anti-quark. The two diagrams are in figure 1.1, and we write their amplitudes as

$$\mathcal{M}(1,2) = \delta_{ij}\bar{v}(p_2)\gamma^\mu u(p_1)L_\mu(p_\ell, p_{\bar{\ell}}) \tag{1.1}$$

The function L_μ contains the propagator of the photon/Z boson, as well as the lepton vertex. So we just used the electromagnetic vertex of the quarks.

As is well known, if we have to compute the cross section at NLO, there are two type of corrections we need to compute. Naturally there are the 1 loop diagrams, that are called virtual, because in the sub-process, an off-shell virtual particle is created. Then there are the real diagrams: in those, an extra gluon appears in the final space. This processes are real because in those a real on-shell gluon is created, and in particular the final phase space is different. Real diagrams are tree level: they should be easy, but the integration over the final state gluon gives some trouble.

Let us discuss in detail. In figure 1.2 we've drawn the relevant diagrams, both real and virtual. Let's focus on the real. Their amplitude is obtained by eq. 1.1, by adding the multiplicative factor of the extra color vertex, times the quark propagator, that can take values of $p_1 - p_4$, or $-(p_2 - p_4)$, if p_4 is the momentum of the emitted gluon.

Thus, we write

Figura 1.1: the LO diagrams for DY process




Figura 1.2: NLO diagrams for DY process



$$\mathcal{M}(1,2,4) = \epsilon^*_\nu(p_4) g_{s,b} (T^a)_{ij} \bar{v}(p_2) \left(\gamma^\mu \frac{\not{p}_1 - \not{p}_4}{(p_1 - p_4)^2} \gamma^\nu - \gamma^\nu \frac{\not{p}_2 - \not{p}_4}{(p_2 - p_4)^2} \gamma^\mu \right) u(p_1) L_\mu(p_\ell, p_{\bar{\ell}}) \tag{1.2}$$

1.1.1 The soft limit

To simplify things, we'd like to use Dirac's equation, that tells us $\not{p}_1 u(p_1) = 0 = \bar{v}(p_2)\not{p}_2$ (we will always consider massless quarks). In both cases, we anti-commute with the γ^ν Dirac gamma,

$$\begin{aligned} \not{p}_1 \gamma^\nu u(p_1) &= \left\{ \not{p}_1, \gamma^\nu \right\} u(p_1) = 2p_1^\nu u(p_1) \\ \bar{v}(p_2)\gamma^\nu \not{p}_2 &= 2\bar{v}(p_2) p_2^\nu \end{aligned} \tag{1.3}$$

Now consider the "soft" limit, where $E_4 \to 0$: then we can take $\not{p}_4 = 0$ in the numerator. For the denominator instead we use $(p_1 - p_4)^2 = -2p_1 \cdot p_4 = -2E_1 E_4 (1 - \cos\theta_{14}) = -2E_1 E_4 \rho_{14}$. In general, we indicate $\rho_{ij} = (1 - n_i \cdot n_j)$, if n_i are unit vector that point in the same direction as p_i. Therefore $\rho_{ij} \in [0, 2]$, $\forall i, j$.

We define $S_i A := \lim_{E_i \to 0} A$. We find

$$S_4 \mathcal{M}(1,2,4) = g_{s,b} \epsilon^*_\nu(p_4) (T^a)_{ij} \bar{v}(p_2) \left(-\frac{\gamma^\mu p_1^\nu}{E_1 E_4 \rho_{14}} + \frac{\gamma^\mu p_2^\nu}{E_2 E_4 \rho_{24}} \right) u(p_1) L_\mu(p_\ell, p_{\bar{\ell}}) \tag{1.4}$$

squaring this, we can write

$$\begin{aligned} S_4 |\mathcal{M}(1,2,4)|^2 &= -\frac{g_{s,b}^2 (N_c^2 - 1)}{2N_c E_4^2} \left(-\frac{p_1^\nu}{E_1 \rho_{14}} + \frac{p_2^\nu}{E_2 \rho_{24}} \right)^2 |\mathcal{M}(1,2)|^2 \\ &= 2\frac{g_{s,b}^2 C_F}{E_4^2} \left(\frac{p_1 \cdot p_2}{E_1 E_2 \rho_{14} \rho_{24}} \right) |\mathcal{M}(1,2)|^2 = 2\frac{g_{s,b}^2 C_F}{E_4^2} \left(\frac{\rho_{12}}{\rho_{14}\rho_{24}} \right) |\mathcal{M}(1,2)|^2 \end{aligned} \tag{1.5}$$

In the first line, the factor $N_c^2 - 1$ comes from $\mathrm{Tr}(T^a T^a) = \frac{1}{2}\delta^{aa} = \frac{N_c^2 - 1}{2}$, and the factor $N_c^- 1$ is the color factor of $|\mathcal{M}(1,2)|^2$. Thus we see $C_F = \frac{N_c^2 - 1}{2N_c}$ appearing naturally. This is the known factorization of the matrix element in the soft limit, and the factor before $|\mathcal{M}(1,2)|^2$ is called *eikonal* factor. In can be shown that this is happens universally, not only for DY process.

Now that we have properly obtained the eikonal factor, we see that its integral over the gluon finale phase space is not finite. Of course this is a problem: we cannot, for example, put the integral in a MC program and get a sensibile result.

This is not the only singularity of the amplitude squared. In the soft limit, the denominator cointains the factors $\rho_{14}\rho_{24} \propto (1-\cos\theta_{14})(1-\cos\theta_{24})$ that go to zero as θ^2 as $\theta \to 0$. Thus also the angular integral is divergent.

1.1.2 The collinear limit

That the angular integral is divergent in the collinear limit (i.e. when one of the θs goes to zero) is true also when the gluon emmitted is *hard*. To see this we go back to the amplitude, eq. 1.2.

1.2 The subtraction scheme for $q\bar{q} \to V + g$

As mentioned before, if an integral is divergent, we cannot integrate it numerically. A possibile strategy for dealing with this problem is defining some operators that extract the singularities. Then one can try to compute analitically the integral in the particular limit, and use a numerical algorithm (e.g. Monte Carlo integrator) to compute the regular integral.

To see how we can do this in practice, first we introduce some useful notation. We also switch to the more general case where the process is $q\bar{q} \to V + g$, with V colorless particles. We denote the contribution of the real diagrams of $d\sigma^{NLO}$ as

$$d\sigma^R = \frac{1}{2s}\int [dg_4] F_{LM}(1,2,4) \tag{1.6}$$

where $(2s)^{-1}$ is the flux factor, $[dg_4] = \frac{d^{d-1}p_4\theta(E_{max}-E_4)}{(2\pi)^d 2E_4}$ and

$$F_{LM}(1,2,4) = d\text{Lips}_V |\mathcal{M}(1,2,4)|^2 \mathcal{F}_{kin}(1,2,4) \tag{1.7}$$

where $d\text{Lips}_V$ is the lorentz invariant phase space of the particles V, and $\mathcal{F}_{kin}(1,2,4)$ is a IR safe observable.

Now notice that the soft limit is not the only point in the phase space that is not integrable. There is also the so called *collinear* singularity, that arises if θ_{14} or θ_{24} vanishes (or, equivalently, ρ_{12} or ρ_{14} go to 0). To be able to integrate the matrix element numerically we must therefor subtract both the soft and the collinear divergences. To this aim we also define the collinear operator,

$$C_{ij}A = \lim_{\rho_{ij}\to 0} A \tag{1.8}$$

The isoleted divergences of $F_{LM}(1,2,4)$ are given by $C_{41}F_{LM}(1,2,4)$, $C_{42}F_{LM}(1,2,4)$, $S_4F_{LM}(1,2,4)$. We thus re-write $d\sigma^R$ as (using the convention where $\int[dg_4]\ldots = <\ldots>$)

$$\begin{aligned} d\sigma^R &= \frac{1}{2s} < (C_{41}+C_{42})F_{LM}(1,2,4) > + \frac{1}{2s} < (1-(C_{41}+C_{42}))F_{LM}(1,2,4) > \\ &= \frac{1}{2s}(< (C_{41}+C_{42})F_{LM}(1,2,4) > + < (I-C_{41}-C_{42})S_4F_{LM}(1,2,4) > + < \hat{O}_{NLO}F_{LM}(1,2,4) >) \end{aligned} \tag{1.9}$$

where we defined $\hat{O}_{NLO} = (I - C41 - C_{42})(I - S_4)$. The term $\propto \hat{O}_{NLO}$ is now free from any singularity by costruction, and can be numerically computed safely. Hence we must compute only the two extra terms.

Claim 1.2.1

The term $< (I - C_{41} - C_{42}) S_4 F_{LM}(1,2,4) >$ is zero

To see this, we use the expresion for $S_4 F_{LM}(1,2,4)$ that we derived earlier (eq. 1.5). In the general case, the formula can be written as

$$S_4 F_{LM}(1,2,4) = \frac{2 g_{s,b}^2 C_F}{E_4^2} \frac{\rho_{12}}{\rho_{14}\rho_{24}} F_{LM}(1,2) \tag{1.10}$$

We have to compute the action of C_{14} as well as C_{24} on this. Of course the limit affects only the fraction $\frac{\rho_{12}}{\rho_{14}\rho_{24}}$. In the CM frame, $\rho_{12} = 2$. Also, if p_4 is allined with p_1, it must be anti-parallel to p_2. We conclude that

$$C_{14} \frac{\rho_{12}}{\rho_{14}\rho_{24}} = \frac{1}{\rho_{14}}, \quad C_{24} \frac{\rho_{12}}{\rho_{14}\rho_{24}} = \frac{1}{\rho_{24}} \implies (C_{14} + C_{24}) \frac{\rho_{12}}{\rho_{14}\rho_{24}} = \frac{1}{\rho_{14}} + \frac{1}{\rho_{24}} = \frac{\rho_{14} + \rho_{24}}{\rho_{14}\rho_{24}} \tag{1.11}$$

Again since we're in the CM frame, $\rho_{14} + \rho_{24} = 2 = \rho_{12}$. We conclude that $C_{14} + C_{42}$ leaves $SF_{LM}(1,2,4)$ unchanged, and so $(I - C_{41} - C_{42}) S_4 F_{LM}(1,2,4) = 0$.

The only term that we have to deal with is therefore

$$< (C_{41} + C_{42}) F_{LM}(1,2,4) > \tag{1.12}$$

When we derived the collinear limit factorization for the DY process (eq.), we mentioned that it was completely universal. We can safely write than

$$C_{41} F_{LM}(1,2,4) = \frac{g_{s,b}^2}{E_4^2 \rho_{14}} \frac{1-z}{z} P_{qq}(z) F_{LM}(z \cdot 1, 2) \tag{1.13}$$

The splitting function is defined as before, and $z = 1 - E_4/E_1$. We also simplified the denominator using $(p_1 - p_4)^2 = -2 p_1 \cdot p_4 = -2 E_1 E_4 \rho_{14} = -2 E_4^2 \rho_{14} \frac{E_1}{E_4} = -2 \frac{E_4^2 \rho_{14}}{1-z}$.

Problems start to arise when we consider this term integrated over the gluon final phase space. We need to evaluate the integral in dimensional regularization, so we can cancel it with the UV divergences of the virutual NLO corrections. We write then

$$[dg_4] = \frac{E_4^{d-3} dE_4 d\cos\theta (1 - \cos^2\theta)^{d/2-2} d\Omega_{d-2}}{2(2\pi)^{d-1}} \tag{1.14}$$

since $E_4 = |p_4|$. Our integrand (eq. 1.13) depends on E_4 through z, and the angular dependence is only in ρ_{14}^{-1}. Our next step is therefore to compute the angular integral, and to write the energy integral in terms of z.

We obtain

$$< C_{41} F_{LM}(1,2,4) >= -\frac{[\alpha_s]}{\epsilon} \frac{\Gamma^2(1-\epsilon)}{\Gamma(1-2\epsilon)} (2E_1)^{-2\epsilon} \int_{z_{\min}}^1 \frac{dz}{(1-z)^{2\epsilon}} P_{qq}(z) \frac{F_{LM}(z \cdot 1, 2)}{z} \tag{1.15}$$

Now we need to write to write the z integral in such a way that the combination with the virtual corrections is straightforward.

Notice that when $z \to 1$ the integral is divergent (or at least would be divergent in the limit $\epsilon \to 0$): this is of course the soft limit. We write $F_{LM}(1,2,4)/z = G(z)$, and we split P_{qq} in a regular function plus a singular, when $z \to 1$:

$$P_{qq}(z) = \frac{2C_F}{1-z} + P_{qq}^{reg}(z) \tag{1.16}$$