

# Fundamentals of Statistical Learning - Unit 1

Master's Degree in Data Science, Sapienza Università di Roma  
Faculty of Information Engineering, Informatics and Statistics

Department of Computer Science

**Author:** Gianmaria Romano

## Reference Bibliography

- F. M. Dekking, C. Kraaikamp, H. P. Lopuhaa, L. E. Meester: *A Modern Introduction to Probability and Statistics*  
L. Wasserman: *All of Statistics: A Concise Course in Statistical Inference*  
N. Matloff: *The Art of R Programming*

Academic Year 2026/2027

### Disclaimer

This document contains notes for the **Fundamentals of Statistical Learning - Unit 1** course, given by **Professor Brutti** as part of the Master's Degree in Data Science at **Sapienza Università di Roma** during the **first** semester of Academic Year **2026/2027**.

**Note:** These notes may not cover all details of each lecture, so it is highly suggested to consult the Professor's official resources and materials.

**Important:** These notes are free to share and use, but please remember to credit me as the author and do not try to hide/remove this page.

### Contacts

For further information or requests, feel free to contact me privately or reach out through my repository: <https://github.com/GianmariaRomano/Data-Science-Notes>.

### License

These documents are distributed under the GNU General Public License 3.0, a form of copyleft intended for use on a manual, textbook or other documents. Material licensed under the current version of the license can be used for any purpose, as long as the use meets certain conditions:

- All previous authors of the work must be attributed.
- All changes to the work must be logged.
- All derivative works must be licensed under the same license.
- The full text of the license, unmodified invariant sections as defined by the author if any, and any other added warranty disclaimers (such as a general disclaimer alerting readers that the document may not be accurate for example) and copyright notices from previous versions must be maintained.
- Technical measures such as DRM may not be used to control or obstruct distribution or editing of the document.

# Contents

|          |                                                         |           |
|----------|---------------------------------------------------------|-----------|
| <b>1</b> | <b>Defining probability</b>                             | <b>3</b>  |
| 1.1      | Introducing a qualitative idea of probability . . . . . | 3         |
| 1.2      | Formalizing probability using set theory . . . . .      | 3         |
| 1.2.1    | Set operations on probability spaces . . . . .          | 3         |
| 1.2.2    | Sample space partitions . . . . .                       | 4         |
| 1.3      | Probability measure functions . . . . .                 | 4         |
| 1.3.1    | Probability bounds . . . . .                            | 6         |
| <b>2</b> | <b>Conditional probability</b>                          | <b>7</b>  |
| 2.1      | Understanding conditional probability . . . . .         | 7         |
| 2.1.1    | The (conditional) odds of an event . . . . .            | 8         |
| 2.1.2    | The multiplication rule . . . . .                       | 8         |
| 2.2      | Defining event independence . . . . .                   | 8         |
| 2.3      | Generalizing event independence . . . . .               | 9         |
| 2.3.1    | Pairwise and mutual independence . . . . .              | 9         |
| 2.3.2    | Conditional independence . . . . .                      | 10        |
| 2.4      | The law of total probability . . . . .                  | 10        |
| 2.5      | Bayes' theorem . . . . .                                | 10        |
| 2.5.1    | The posterior odds of a hypothesis . . . . .            | 11        |
| <b>3</b> | <b>Random variables</b>                                 | <b>12</b> |
| 3.1      | Defining random variables . . . . .                     | 12        |
| 3.1.1    | Discrete random variables . . . . .                     | 13        |
| 3.1.2    | Continuous random variables . . . . .                   | 13        |
| 3.2      | The quantile function of a random variable . . . . .    | 15        |

# Chapter 1

## Defining probability

### 1.1 Introducing a qualitative idea of probability

Probability is the field of mathematics that studies events and how likely they are to occur in order to describe relationships or patterns in existing models.

Starting from a model's data generation process, probability aims to investigate the properties of the observed outcomes, whereas inference makes use of the observed outcomes to study the underlying data generation process.

#### Example: Diffusion models

Diffusion models are a type of generative model that create new data by gradually reversing a noising process.

From a mathematical perspective, a diffusion model can be described using a *Markovian Direct Acyclic Graph* whose nodes represent the observable or latent variables involved in the process and whose edges denote the transition probabilities of the model.

### 1.2 Formalizing probability using set theory

It is possible to formalize probabilistic experiments using set theory.

In fact, starting from a sample space  $\Omega$ , which represents the set of all possible outcomes, an event  $A$  on  $\Omega$  can be defined as the set of all favourable outcomes in which the event does occur.

#### Example: Tossing a coin

Suppose you toss a coin: it is possible to construct a random experiment by defining the sample space  $\Omega = \{Heads, Tails\}$  and considering the event  $A = \{Heads\}$ .

#### 1.2.1 Set operations on probability spaces

Given two events  $A$  and  $B$  on the same sample space  $\Omega$ , it is possible to define three main operations:

- **Union:** The set  $A \cup B = \{\omega \in \Omega : \omega \in A \text{ or } \omega \in B\}$  corresponds to the event in which at least one of  $A$  or  $B$ , or both, occurs.  
In particular, it is always the case that  $A \cup \Omega = \Omega$  and  $A \cup \emptyset = A$ .
- **Intersection:** The set  $A \cap B = \{\omega \in \Omega : \omega \in A \text{ and } \omega \in B\}$  corresponds to the event in which both  $A$  and  $B$  occur.  
In particular, it is always the case that  $A \cap \Omega = A$  and  $A \cap \emptyset = \emptyset$ .
- **Complement:** The set  $A^C = \Omega \setminus A = \{\omega \in \Omega : \omega \notin A\}$  corresponds to the event in which  $A$  does not occur.  
In particular, it is always the case that  $A \cup A^C = \Omega$  and  $A \cap A^C = \emptyset$ .

Additionally, given a sequence of events  $\{A_1, \dots, A_n\}$  on the same sample space  $\Omega$ , De Morgan's laws allow to switch between the union and intersection operations through the following identities:

$$\left( \bigcup_{i=1}^n A_i \right)^C = \bigcap_{i=1}^n A_i^C \text{ and } \left( \bigcap_{i=1}^n A_i \right)^C = \bigcup_{i=1}^n A_i^C$$

### 1.2.2 Sample space partitions

A sequence of events  $\{A_1, \dots, A_n\}$  is said to form a partition on a sample space  $\Omega$  if the following conditions are satisfied:

1.  $A_1 \cup \dots \cup A_n = \Omega$ , typically requiring  $A_i \neq \emptyset \forall i = 1, \dots, n$  as well.
2. The sets in the sequence are pairwise disjoint, meaning that  $A_i \cap A_j = \emptyset \forall i \neq j$ .

Generally speaking, partitions come in handy to explain the probability of an event by decomposing it as a union of disjoint sets with respect to a sequence of mutually exclusive hypotheses.

## 1.3 Probability measure functions

While several definitions of probability exist, Kolmogorov's axiomatic definition is the most commonly used when defining probability measures.

According to this definition, a probability measure function  $\mathbb{P} : \mathcal{P}(\Omega) \rightarrow [0, 1]$  assigns to each event  $A \subseteq \Omega$  a probability  $\mathbb{P}(A)$  such that:

1.  $\mathbb{P}(A) \in [0, 1] \forall A \subseteq \Omega$ .
2.  $\mathbb{P}(\emptyset) = 0$  and  $\mathbb{P}(\Omega) = 1$ .
3.  $\mathbb{P}(A \cup B) = \mathbb{P}(A) + \mathbb{P}(B) - \mathbb{P}(A \cap B)$ , with  $\mathbb{P}(A \cup B) = \mathbb{P}(A) + \mathbb{P}(B)$  if and only if  $A \cap B = \emptyset$ .  
This last axiom can be generalized for a sequence of  $n$  events through the inclusion-exclusion principle, which states that

$$\mathbb{P}\left(\bigcup_{i=1}^n A_i\right) = \sum_{k=1}^n (-1)^{k+1} \left( \sum_{1 \leq i_1 < \dots < i_k \leq n} \mathbb{P}(A_{i_1} \cap \dots \cap A_{i_k}) \right)$$

### Example: Checking polynomial identities using a randomized algorithm

Consider two polynomials of degree  $d$ :

$$F(x) = \prod_{i=1}^d (x - a_i) \text{ and } G(x) = \sum_{i=1}^d c_i x^i$$

A simple way of checking whether  $F(x) \equiv G(x)$  involves expanding  $F(x)$  into its canonical form and compare the resulting coefficient with those of  $G(x)$ , although this algorithm would require performing  $\Theta(d^2)$  operations.

Knowing that, for any randomly chosen point  $x_0$ , evaluating  $F(x_0)$  and  $G(x_0)$  requires  $O(d)$  time, it is possible to cut down complexity using a randomized algorithm:

---

**Algorithm 1:** Randomized Algorithm for Polynomial Identities.

---

**Data:**  $F(x), G(x)$

**Result:** Whether  $F(x) \equiv G(x)$ .

Generate  $r$  uniformly at random from a set  $\mathcal{I} = \{1, \dots, 100d\}$ ;

Compute  $F(r)$  and  $G(r)$ ;

**if**  $F(r) = G(r)$  **then**

$F(x) \equiv G(x)$ ;

**else**

$F(x) \not\equiv G(x)$ ;

---

However, the randomized nature of this algorithm can result in a wrong evaluation.

In fact, while  $F(x) \equiv G(x)$  results in  $F(r) = G(r) \forall r$ , it might be the case that, for some  $r$ ,  $F(r) = G(r)$  even if  $F(x) \not\equiv G(x)$ : this implies that the algorithm has one-sided error as it can make mistakes only when deciding that  $F(x) \equiv G(x)$ .

Therefore, define the event

$$E = \{F(r) = G(r) \text{ but } F(x) \not\equiv G(x)\}$$

This event occurs if and only if  $r$  is a root of the polynomial  $H(x) = F(x) - G(x)$ : since  $H(x)$  will always have degree at most  $d$ , it will have at most  $d$  roots, allowing to bound the error probability as follows:

$$\mathbb{P}(E) = \frac{|\{r \in \mathcal{I} : r \text{ is a root of } H(x)\}|}{|\mathcal{I}|} \leq \frac{d}{100d} = \frac{1}{100}$$

As a consequence to Kolmogorov's axiomatic definition of probability, every probability space  $(\Omega, \mathbb{P})$  satisfies the property that  $\mathbb{P}(A^C) = 1 - \mathbb{P}(A) \forall A \subseteq \Omega$ .

### Proof:

Consider the partition  $\Omega = A \cup A^C$  and apply Kolmogorov's axioms.

$$1 = \mathbb{P}(\Omega) = \mathbb{P}(A \cup A^C) = \mathbb{P}(A) + \mathbb{P}(A^C) \Rightarrow \mathbb{P}(A^C) = 1 - \mathbb{P}(A)$$

### 1.3.1 Probability bounds

Boole's inequality states that, for every sequence of events  $\{A_1, \dots, A_n\}$ , it holds that

$$\mathbb{P}\left(\bigcup_{i=1}^n A_i\right) \leq \sum_{i=1}^n \mathbb{P}(A_i)$$

**Proof:**

This inequality can be proved by induction on the length of the sequence.

- For the base case, give by  $n = 1$ , the inequality is trivially true as  $\mathbb{P}(A_1) \leq \mathbb{P}(A_1)$ .
- Assuming the inequality is true for some  $n$ , the goal is to show that it is true for  $(n + 1)$  as well.

Reformulate  $A_1 \cup \dots \cup A_{n+1} = (A_1 \cup \dots \cup A_n) \cup A_{n+1}$  and let

$$\begin{aligned} \mathbb{P}\left(\bigcup_{i=1}^{n+1} A_i\right) &= \mathbb{P}\left(\left(\bigcup_{i=1}^n A_i\right) \cup A_{n+1}\right) = \mathbb{P}\left(\bigcup_{i=1}^n A_i\right) + \mathbb{P}(A_{n+1}) - \mathbb{P}\left(\left(\bigcup_{i=1}^n A_i\right) \cap A_{n+1}\right) \\ &\stackrel{\mathbb{P}(\cdot) \geq 0}{\leq} \mathbb{P}\left(\bigcup_{i=1}^n A_i\right) + \mathbb{P}(A_{n+1}) \stackrel{\text{hp. ind.}}{\leq} \sum_{i=1}^n \mathbb{P}(A_i) + \mathbb{P}(A_{n+1}) = \sum_{i=1}^{n+1} \mathbb{P}(A_i) \end{aligned}$$

Having proved the base case and the inductive case, this inequality must be true  $\forall n \in \mathbb{N}$ , which implies that probability measure functions are always sub-additive.

As a consequence to Boole's inequality, Bonferroni's inequality states that, for every sequence of events  $\{A_1, \dots, A_n\}$ , it must be the case that

$$\mathbb{P}\left(\bigcap_{i=1}^n A_i\right) \geq \sum_{i=1}^n \mathbb{P}(A_i) - (n - 1)$$

**Proof:**

Reformulate the joint probability of the events using De Morgan's laws.

$$\begin{aligned} \mathbb{P}\left(\bigcap_{i=1}^n A_i\right) &= 1 - \mathbb{P}\left(\left(\bigcap_{i=1}^n A_i\right)^C\right) \stackrel{\text{De Morgan}}{=} 1 - \mathbb{P}\left(\bigcup_{i=1}^n A_i^C\right) \\ &\stackrel{\text{Boole}}{\geq} 1 - \sum_{i=1}^n \mathbb{P}(A_i^C) = 1 - \sum_{i=1}^n (1 - \mathbb{P}(A_i)) \\ &= 1 - \left(\sum_{i=1}^n 1 - \sum_{i=1}^n \mathbb{P}(A_i)\right) = \sum_{i=1}^n \mathbb{P}(A_i) - (n - 1) \end{aligned}$$

## Chapter 2

# Conditional probability

### 2.1 Understanding conditional probability

Given a sample space  $\Omega$ , the conditional probability that event  $A$  occurs knowing that event  $B$  was observed can be defined as follows:

$$\mathbb{P}(A|B) = \frac{\mathbb{P}(A \cap B)}{\mathbb{P}(B)}, \text{ with } \mathbb{P}(B) > 0.$$

#### Example: Measles and red spots

Consider the two events

$$A = \{\text{I have measles}\} \text{ and } B = \{\text{I have red spots on my face}\}$$

Knowing that red spots are a common symptom of measles, it will be the case that  $\mathbb{P}(B|A) = 1$ . However, the same thing cannot be said about  $\mathbb{P}(A|B)$  as red spots might be caused by other external factors, such as eating hot food or getting a scratch.

From a broader perspective, a conditional probability distribution can be described using a probability measure function that satisfies the following properties:

1.  $\mathbb{P}(A|B) \in [0, 1] \forall A \subseteq \Omega$ .
2.  $\mathbb{P}(\Omega|B) = 1$ .
3. For any sequence of disjoint events  $\{A_1, \dots, A_n\}$ , it is always the case that

$$\mathbb{P}\left(\bigcup_{i=1}^n A_i \middle| B\right) = \sum_{i=1}^n \mathbb{P}(A_i|B)$$

As a consequence, it is always the case that  $\mathbb{P}(A^C|B) = 1 - \mathbb{P}(A|B) \forall A \subseteq \Omega$ .

**Proof:**

Let  $B = B \cap \Omega$  on the partition  $\Omega = A \cup A^C$ .

$$\begin{aligned}\mathbb{P}(B) &= \mathbb{P}(B \cap \Omega) = \mathbb{P}(B \cap (A \cup A^C)) = \mathbb{P}((B \cap A) \cup (B \cap A^C)) \\ &= \mathbb{P}(A \cap B) + \mathbb{P}(A^C \cap B) \Rightarrow \mathbb{P}(A^C \cap B) = \mathbb{P}(B) - \mathbb{P}(A \cap B)\end{aligned}$$

Therefore, applying the definition of conditional probability will result in

$$\mathbb{P}(A^C|B) = \frac{\mathbb{P}(A^C \cap B)}{\mathbb{P}(B)} = \frac{\mathbb{P}(B) - \mathbb{P}(A \cap B)}{\mathbb{P}(B)} = \frac{\mathbb{P}(B)}{\mathbb{P}(B)} - \frac{\mathbb{P}(A \cap B)}{\mathbb{P}(B)} = 1 - \mathbb{P}(A|B)$$

**2.1.1 The (conditional) odds of an event**

Generally speaking, the odds of an event can be seen as the likelihood that the event will occur relative to it not occurring.

For this reason, the odds of an event  $A \subseteq \Omega$  with probability  $\mathbb{P}(A) \in [0, 1)$  can be computed as

$$\mathcal{O}(A) = \frac{\mathbb{P}(A)}{\mathbb{P}(A^C)} = \frac{\mathbb{P}(A)}{1 - \mathbb{P}(A)}$$

More specifically, the conditional odds of event  $A$  given that a second event  $B$  occurred can be computed as

$$\mathcal{O}(A|B) = \frac{\mathbb{P}(A|B)}{\mathbb{P}(A^C|B)} = \frac{\mathbb{P}(A \cap B)}{\mathbb{P}(A^C \cap B)}$$

**2.1.2 The multiplication rule**

The multiplication rule allows to reformulate the joint probability of two events  $A$  and  $B$  according to the following equation:

$$\mathbb{P}(A \cap B) = \mathbb{P}(A|B)\mathbb{P}(B) = \mathbb{P}(B|A)\mathbb{P}(A)$$

In particular, repeatedly applying the multiplication rule to a sequence of events  $\{A_1, \dots, A_n\}$  results in the chain rule, which can be formulated as

$$\mathbb{P}\left(\bigcap_{i=1}^n A_i\right) = \mathbb{P}(A_1) \cdot \mathbb{P}(A_2|A_1) \cdot \dots \cdot \mathbb{P}(A_n|A_1 \cap \dots \cap A_{n-1})$$

**2.2 Defining event independence**

Two events  $A$  and  $B$  on a sample space  $\Omega$  are said to be independent if knowing that  $B$  occurred does not add information about the probability of  $A$  occurring, resulting in

$$A \perp B \Leftrightarrow \mathbb{P}(A|B) = \mathbb{P}(A) \text{ and } \mathbb{P}(B|A) = \mathbb{P}(B), \text{ resulting in } \mathbb{P}(A \cap B) = \mathbb{P}(A)\mathbb{P}(B).$$

Depending on the context, event independence can be assumed or verified, although it is always possible to show that, if  $A$  and  $B$  are events such that  $\mathbb{P}(A), \mathbb{P}(B) > 0$  and  $A \cap B = \emptyset$ , then the events cannot be independent.

**Proof:**

It is possible to notice that having  $A \cap B = \emptyset$  and knowing that one of the two events occurred would violate the definition of event independence because

$$\mathbb{P}(A)\mathbb{P}(B) > 0, \text{ but } \mathbb{P}(A \cap B) = \mathbb{P}(\emptyset) = 0.$$

## 2.3 Generalizing event independence

For  $n > 2$  events, independence can be defined at several distinct levels.

### 2.3.1 Pairwise and mutual independence

A (countable) sequence of events  $\{A_1, \dots, A_n\}$  is said to be pairwise independent if

$$A_i \perp A_j \quad \forall i \neq j \Leftrightarrow \mathbb{P}(A_i \cap A_j) = \mathbb{P}(A_i)\mathbb{P}(A_j) \quad \forall i \neq j$$

Additionally, a (countable) sequence of events  $\{A_1, \dots, A_n\}$  is said to be mutually independent if, for any group of  $k$  events in the sequence, it is the case that

$$\mathbb{P}\left(\bigcap_{i=1}^k A_{n_i}\right) = \prod_{i=1}^k \mathbb{P}(A_{n_i}), \text{ for } 1 \leq k \leq n \text{ and } k \in \mathbb{N}.$$

It should be noticed that, while mutual independence always implies pairwise independence, the converse statement is not true because a sequence of events can be pairwise independent but not mutually independent.

**Example: A special die**

Given a balanced four-faced die, which has a red face, a green face, a blue face and a striped face with all three colours, define the events

$$R = \{\text{I see red}\}, G = \{\text{I see green}\} \text{ and } B = \{\text{I see blue}\}$$

It is possible to show that these events are pairwise independent as

$$\mathbb{P}(R \cap G) = 0.25 = 0.5 \cdot 0.5 = \mathbb{P}(R)\mathbb{P}(G)$$

$$\mathbb{P}(R \cap B) = 0.25 = 0.5 \cdot 0.5 = \mathbb{P}(R)\mathbb{P}(B)$$

$$\mathbb{P}(G \cap B) = 0.25 = 0.5 \cdot 0.5 = \mathbb{P}(G)\mathbb{P}(B)$$

However, the events are not mutually independent because

$$\mathbb{P}(R \cap G \cap B) = 0.25 \neq 0.125 = 0.5 \cdot 0.5 \cdot 0.5 = \mathbb{P}(R)\mathbb{P}(G)\mathbb{P}(B)$$

### 2.3.2 Conditional independence

Two events  $A$  and  $B$  are said to be conditionally independent given an event  $C$  if, knowing that  $C$  occurred, observing event  $B$  does not add information about the probability of  $A$  occurring, resulting in

$$\mathbb{P}(A \cap B|C) = \mathbb{P}(A|C)\mathbb{P}(B|C), \text{ with } \mathbb{P}(C) > 0.$$

#### Example: University admissions

For a student applying to a Master's Degree course, define the events

$$DS = \{\text{Accepted to Data Science}\} \text{ and } ST = \{\text{Accepted to Statistics}\}$$

Knowing that event  $DS$  occurred can provide additional information about the likelihood of event  $ST$  occurring, meaning that the two events are not independent.

However, if the admission procedure is solely based on the student's GPA  $C$ , knowing that one event occurred does not provide additional information about the other event occurring as well, making  $DS$  and  $ST$  conditionally independent given event  $C$ .

## 2.4 The law of total probability

Given a sample space  $\Omega$ , the law of total probability allows to explain the probability of an event  $A \subseteq \Omega$  by decomposing it over the partition  $\Omega = B \cup B^C$ , resulting in

$$\mathbb{P}(A) = \mathbb{P}(A \cap B) + \mathbb{P}(A \cap B^C) = \mathbb{P}(A|B)\mathbb{P}(B) + \mathbb{P}(A|B^C)\mathbb{P}(B^C)$$

#### Proof:

Let  $A = A \cap \Omega$  on the partition  $\Omega = B \cup B^C$ .

$$\begin{aligned} \mathbb{P}(A) &= \mathbb{P}(A \cap \Omega) = \mathbb{P}(A \cap (B \cup B^C)) = \mathbb{P}((A \cap B) \cup (A \cap B^C)) \\ &= \mathbb{P}(A \cap B) + \mathbb{P}(A \cap B^C) \stackrel{\text{Multiplication rule}}{=} \mathbb{P}(A|B)\mathbb{P}(B) + \mathbb{P}(A|B^C)\mathbb{P}(B^C) \end{aligned}$$

Additionally, it is possible to generalize the law of total probability to explain the probability of an event  $E$  with respect to a sequence of mutually exclusive hypotheses  $\{H_1, \dots, H_n\}$  as

$$\mathbb{P}(E) = \sum_{i=1}^n \mathbb{P}(E \cap H_i) = \sum_{i=1}^n \mathbb{P}(E|H_i)\mathbb{P}(H_i)$$

## 2.5 Bayes' theorem

Bayes' theorem allows to switch between conditional probabilities by setting

$$\mathbb{P}(A|B) = \frac{\mathbb{P}(A \cap B)}{\mathbb{P}(B)} = \frac{\mathbb{P}(B|A)\mathbb{P}(A)}{\mathbb{P}(B)} \text{ with } \mathbb{P}(B) > 0.$$

In particular, when considering a sequence of mutually exclusive hypotheses  $\{H_1, \dots, H_n\}$  with known prior probabilities, Bayes' theorem allows to compute the posterior probability of hypothesis  $H_i$  given the observed evidence  $E$  as

$$\mathbb{P}(H_i|E) = \frac{\mathbb{P}(E|H_i)\mathbb{P}(H_i)}{\mathbb{P}(E)} = \frac{\mathbb{P}(E|H_i)\mathbb{P}(H_i)}{\sum_{j=1}^n \mathbb{P}(E|H_j)\mathbb{P}(H_j)}$$

In fact, the posterior probability of each hypothesis given the observed evidence will be computed in terms of the likelihood  $\mathbb{P}(E|H_i)$ , which describes the probability of seeing the evidence given that the hypothesis  $H_i$  is true, the prior probability  $\mathbb{P}(H_i)$ , which describes the probability of  $H_i$  being true, and the marginal probability  $\mathbb{P}(E)$ , which denotes the total probability of seeing the evidence under all possible hypotheses.

#### Example: Iterated Bayes for medical diagnosis

Consider the events

$$P = \{\text{I test positive to a disease}\} \text{ and } D = \{\text{I actually have the disease}\}$$

In particular, the disease affects 0.1% of the population and it is known that a dedicated test correctly detects 99% of disease case but produces a false positive result on 1% of occasions, which means that

$$\mathbb{P}(D|P) = \frac{\mathbb{P}(P|D)\mathbb{P}(D)}{\mathbb{P}(P|D)\mathbb{P}(D) + \mathbb{P}(P|D^C)\mathbb{P}(D^C)} = \frac{0.99 \cdot 0.001}{(0.99 \cdot 0.001) + (0.01 \cdot 0.999)} = 9\%$$

Since the posterior probability appears to be quite low, Bayes' rule is applied once again, using the posterior probability as the new prior probability, resulting in

$$\mathbb{P}(D|P) = \frac{\mathbb{P}(P|D)\mathbb{P}(D)}{\mathbb{P}(P|D)\mathbb{P}(D) + \mathbb{P}(P|D^C)\mathbb{P}(D^C)} = \frac{0.99 \cdot 0.09}{(0.99 \cdot 0.09) + (0.01 \cdot 0.91)} = 91\%$$

### 2.5.1 The posterior odds of a hypothesis

When considering an event  $E$  on a sequence of mutually exclusive hypotheses  $\{H_1, \dots, H_n\}$ , it is possible to compute the posterior odds of hypothesis  $H_i$  given evidence  $E$  as

$$\mathcal{O}(H_i|E) = \frac{\mathbb{P}(H_i|E)}{\mathbb{P}(H_i^C|E)} = \frac{\frac{\mathbb{P}(E|H_i)\mathbb{P}(H_i)}{\mathbb{P}(E)}}{\frac{\mathbb{P}(E|H_i^C)\mathbb{P}(H_i^C)}{\mathbb{P}(E)}} = \frac{\mathbb{P}(E|H_i)\mathcal{O}(H_i)}{\mathbb{P}(E|H_i)}$$

# Chapter 3

## Random variables

### 3.1 Defining random variables

Given a probability space  $(\Omega, \mathbb{P})$ , a random variable is a (measurable) mapping  $X : \Omega \rightarrow S_X \subseteq \mathbb{R}$  that maps every outcome  $\omega \in \Omega$  onto a value  $X(\omega) \in \mathbb{R}$ , with the set  $S_X = \{X(\omega) : \omega \in \Omega\}$  forming the “support” of  $X$ .

This concept can be extended to define a random vector, also known as a  $p$ -variate random variable, as a (measurable) mapping  $\mathbf{X} : \Omega \rightarrow \mathbb{R}^p$  that assigns a vector  $\mathbf{X}(\omega) \in \mathbb{R}^p$  to each outcome  $\omega \in \Omega$ .




Figure 3.1: A visual idea of what is a random variable.

Since this mapping describes the probability of the random variable  $X$  falling within a subset of  $\mathbb{R}$ , it is possible to define the pre-image of a set  $E \subseteq \mathbb{R}$  as the set  $X^{-1}(E) = \{\omega \in \Omega : X(\omega) \in E\} \subseteq \Omega$ . This definition allows to describe the probability distribution of a random variable  $X$  as

$$\mathbb{P}(X \in E) = \mathbb{P}(\{\omega \in \Omega : X(\omega) \in E\}) = \mathbb{P}(X^{-1}(E))$$




Figure 3.2: A visual idea of the pre-image of a random variable.

When defining the probability distribution of a random variable  $X$ , it should be noticed that  $\mathbb{P}(X \in E)$  represents an induced distribution supported on  $\mathbb{R}$  that is constructed based on the assumed distribution  $\mathbb{P}(X^{-1}(E))$  supported on  $\Omega$ .

### 3.1.1 Discrete random variables

A random variable is said to be “discrete“ if its support is a countable set, which means that the random variable can take at most a finite, or at least countable, number of values.

For this reason, a discrete random variable  $X : \Omega \rightarrow S_X$  can be described using a probability mass function  $p_X(x) = \mathbb{P}(X = x)$  that satisfies the following conditions:

- $p_X(x) \geq 0 \ \forall x \in \mathbb{R}$ , with  $p_X(x) > 0 \ \forall x \in S_X$  specifically.
- $\sum_{x \in S_X} p_X(x) = 1$ .

#### Example: Repeated coin flips

Consider 5 independent flips of a fair coin: since each flip can be seen as a repetition of a binary experiment with sample space  $\Omega = \{Heads, Tails\}$ , it is possible to define the overall outcome space as the Cartesian product  $\Omega^{\otimes 5}$ .

Assuming symmetry and independence between coin flips, each outcome  $\omega \in \Omega^{\otimes 5}$  is equally likely and has probability  $\mathbb{P}(\omega) = \frac{1}{|\Omega^{\otimes 5}|} = \frac{1}{32}$ .

Define now the random variable  $X = \{\text{Number of tails in 5 independent coin flips}\}$ : on its support  $S_X = \{0, 1, 2, 3, 4, 5\}$ , the probability definition of  $X$  can be defined as

$$p_X(x) = \mathbb{P}(X = x) = \binom{5}{x} \left(\frac{1}{2}\right)^5, \text{ for } x \in S_X.$$

### 3.1.2 Continuous random variables

A random variable is said to be “continuous“ if its support is an uncountable set.

Since, in this setting,  $\mathbb{P}(X = x) \rightarrow 0 \ \forall x \in \mathbb{R}$ , a continuous random variable  $X : \Omega \rightarrow S_X$  is described using a probability density function  $f_X(x)$  that satisfies the following conditions:

- $\mathbb{P}(a \leq X \leq b) = \mathbb{P}(X \in [a, b]) = \int_a^b f_X(x) \, dx$ .
- $f_X(x) \geq 0 \ \forall x \in \mathbb{R}$ , with  $f_X(x) > 0 \ \forall x \in S_X$  specifically.
- $\int_{\mathbb{R}} f_X(x) \, dx = 1$ .

In particular, all information about a continuous random variable  $X$  can be retrieved from its cumulative distribution function  $F_X : \mathbb{R} \rightarrow [0, 1]$ , which can be defined as

$$F_X(x) = \mathbb{P}(X \leq x) = \mathbb{P}(X \in (-\infty, x]) \ \forall x \in \mathbb{R}$$

More specifically, a function  $F_X : \mathbb{R} \rightarrow [0, 1]$  acts as a cumulative distribution function on some probability measure  $\mathbb{P}$  if and only if the following conditions are met:

- $F_X(x)$  is non-decreasing, which means that  $F_X(x_1) \leq F_X(x_2) \forall x_1 < x_2$ .
- $F_X(x)$  is normalized, meaning that

$$\lim_{x \rightarrow -\infty} F_X(x) = 0 \quad \lim_{x \rightarrow +\infty} F_X(x) = 1$$

- $F_X(x)$  is right-continuous, which means that

$$F_X(x) = F_X(x^+) = \lim_{t \rightarrow x^+} F_X(t)$$

Furthermore, the cumulative distribution function of a random variable  $X$  always satisfies several properties:

- $\mathbb{P}(x_1 < X \leq x_2) = \mathbb{P}(X \in (x_1, x_2]) = F(x_2) - F(x_1)$ .
- $\mathbb{P}(X > x) = \mathbb{P}(X \in (x, +\infty)) = 1 - F_X(x)$ .
- If  $X$  is a continuous random variable with probability density function  $f_X(x)$ , then

$$F_X(x) = \int_{-\infty}^x f_X(t) dt \text{ and } f_X(x) = F'_X(x) \text{ wherever } F_X \text{ is differentiable.}$$

It should be noticed that this last property can also be formulated for discrete random variables. In fact, if  $X$  is a discrete random variable with probability mass function  $p_X(x)$ , then its cumulative function is given by

$$F_X(x) = \sum_{x_i \leq x} p_X(x_i) = \mathbb{P}(X < x) + p_X(x), \text{ with } p_X(x) = F_X(x) - F_X(x^-).$$

### Example: Playing darts

Suppose you throw a dart on a disc of radius  $r$  at random. Define the random variable  $X = \{\text{Distance of the hitting point from the center}\}$  and note that its support  $[0, r]$  is an uncountable set.

$$\mathbb{P}(X \leq x) = 0 \quad \forall x < 0 \text{ as distances are not negative.}$$

$$\mathbb{P}(X \leq x) = 1 \quad \forall x > r \text{ as the dart did hit the disc.}$$

$$\mathbb{P}(X \leq x) = \frac{\pi x^2}{\pi r^2} = \frac{x^2}{r^2} \quad \forall x \in [0, r]$$

Therefore, it is possible to define the cumulative distribution function of  $X$  as

$$F_X(x) = \begin{cases} 0 & \text{if } x < 0 \\ \frac{x^2}{r^2} & \text{if } x \in [0, r] \\ 1 & \text{if } x > r \end{cases}$$

This implies that the probability density function will be

$$f_X(x) = \begin{cases} \frac{2x}{r^2} & \text{if } x \in [0, r] \\ 0 & \text{otherwise} \end{cases}, \text{ or } f_X(x) = \frac{2x}{r^2} \mathbb{I}_{[0, r]}(x), \text{ with } \mathbb{I}_{[0, r]}(x) = \begin{cases} 1 & \text{if } x \in [0, r] \\ 0 & \text{otherwise} \end{cases}$$

### 3.2 The quantile function of a random variable

Let  $X : \Omega \rightarrow S_X$  be a continuous random variable with cumulative distribution function  $F_X(x)$ : for a given level  $p \in [0, 1]$ , the quantile function is defined as

$$q_p = \mathbb{Q}(p) = \inf\{x \in S_X : F_X(x) = p\}$$

In particular, if  $F_X(x)$  is strictly increasing in  $\mathbb{R}$ , then it is also invertible and it is possible to set

$$q_p = \mathbb{Q}(p) = F_X^{-1}(p), \text{ with } p = F_X(q_p) \int_{-\infty}^{q_p} f_X(t) dt \text{ representing the quantile at level } p.$$

However, since discrete random variables tend to have weakly monotonic cumulative distribution functions, a more general definition of quantile function is typically used.

In fact, given any random variable  $X : \Omega \rightarrow S_X$  with cumulative distribution function  $F_X(x)$ , the quantile function for a given level  $p \in [0, 1]$  is formulated as

$$q_p = \mathbb{Q}(p) = \inf\{x \in S_X : F_X(x) \geq p\}$$