

Due Date: 11:59 pm on 2/17/2026

Instructor: Dr. Tutku Karacolak

ECE 370 Electromagnetic Fields and Waves

HOMEWORK #3

Name: Gianni Avilan

Problem	Course outcome	Grade
1	1-c, 1-d	/30
2	1-a, 1-d	/30
3	1-c, 1-d	/40
TOTAL:		/100

1- Solve Problem 4.32 from the textbook.

4.32 A circular ring of charge of radius a lies in the x - y plane and is centered at the origin.

Assume also that the ring is in air and carries a uniform density ρ_ℓ .

a. Show that the electrical potential at $(0, 0, z)$ is given by

$$V = \rho_\ell a / [2\epsilon_0 (a^2 + z^2)^{1/2}].$$

b. Find the corresponding electric field $\mathbf{E}$.

2- A ceramic sphere of radius $\frac{1}{2}$ m is charged with a surface charge density $\rho_s = 1.5$ C/m². The relative permittivity of the ceramic is $\epsilon_r = 4$. The region outside of the sphere has $\epsilon_r = 2$. The electric flux density on the outside surface of the ceramic is

$$\vec{D} = 0.7\hat{a}_R + 2\hat{a}_\theta - 3\hat{a}_\phi \quad (\text{C/m}^2)$$

Use electric field boundary conditions to find the electric flux density on the inside surface of the sphere.

3- The potential in free space is given by:

$$V(x, y, z) = -x^2y(z^2 + 1) \text{ (V)}$$

Determine

- (a) the vector electric field. (12 pts)
- (b) the vector electric flux density. (8 pts)
- (c) the potential difference V_{AB} between $A=(1, 2, 5)$ m and $B=(1, 2, 1)$ m by integrating the electric field along the straight line path between these two points. (20 pts)

1. a) We may consider the given ring of charge to be a collection of infinitesimal point charges, and find the electric potential at some point on the z-axis by summing the total effect of each point charge. Here, we refer to **electric potential** as the difference in the potential energy of a unit charge (1C) at some point P , with the implication that there is some quantifiable electric field intensity at P , versus that same unit charge's potential energy infinitely far away from P , with the implication that there is no electric field at infinity.

For a point charge with charge q , this electric potential is given by

$$V = \frac{q}{4\pi\epsilon|R - R_1|},$$

with $|R - R_1|$ representing the shortest distance between the point charge and P . Since there is no directional component to electric potential, and it reduces to a scalar, we may use the superposition principle to find the total influence of any number of point charges on the electric potential at P as the sum of their influences. Factoring out constant terms,

$$V = \frac{1}{4\pi\epsilon} \sum_{i=1}^N \frac{q_i}{|R - R_1|}.$$

As mentioned earlier, we may reduce the given ring of charge of radius a to a collection of point charges q_i , where each point charge carries a charge of ρ_ℓ , we will note that, by centering the ring of charge at the origin of the x-y axis, each of these infinitesimal point charges lies at an equal distance to any given point along the z-axis, so that for any one point charge, $|R - R_1| = \sqrt{z^2 + a^2}$. Since this term is constant for any point $(0, 0, z)$, we may now express the electric potential at z as

$$\frac{1}{4\pi\epsilon\sqrt{z^2 + a^2}} \sum_{i=1}^N \rho_\ell dl.$$

We will also assume a permittivity of ϵ_0 , since the permittivity of air is virtually identical to free space permittivity. Summing all point charges around the entire ring, we could derive the corresponding integral using cylindrical coordinates, or simply use the well-known formula for the circumference of a circle, $2\pi R$, and reduce the formula for electric potential at $(0, 0, z)$ to

$$V_z = \frac{\rho_\ell 2\pi a}{4\pi\epsilon_0\sqrt{z^2 + a^2}} = \frac{\rho_\ell a}{2\epsilon_0\sqrt{z^2 + a^2}} V.$$

- b) Though we could use the methods from the previous homework, such as Gauss's Law, to derive the electric field $\vec{E}$ at some point $(0, 0, z)$, it is easier to use the electric potential itself to derive the electric field. We will use the relationship

$$\vec{E} = -\nabla V;$$

in other words, electric field is the gradient of electric potential.

Finding electric field at $(0, 0, z)$,

$$\vec{E}_z = -\left(\frac{\partial V_z}{\partial x} \hat{\alpha}_x + \frac{\partial V_z}{\partial y} \hat{\alpha}_y + \frac{\partial V_z}{\partial z} \hat{\alpha}_z\right) = -\frac{\partial V_z}{\partial z} \hat{\alpha}_z = -\frac{\rho_\ell a}{2\epsilon_0} \frac{d}{dz} \frac{1}{\sqrt{z^2 + a^2}} = \frac{\rho_\ell a z}{2\epsilon_0(z^2 + a^2)^{\frac{3}{2}}} \hat{\alpha}_z \frac{V}{m}$$

2. Using boundary conditions, we know that

$$D_{1n} - D_{2n} = \rho_s \Rightarrow D_{1n} = \rho_s + D_{2n}.$$

Given surface charge density $1.5 \frac{C}{m^2}$ and flux density normal to the outer boundary of the ceramic sphere of 0.7,

$$D_{1n} = 1.5 + 0.7 = 2.2 \hat{\alpha}_R \frac{C}{m^2}$$

Then, finding flux density tangent to the outer surface, we may use

$$\frac{D_{1t}}{\varepsilon_1} = \frac{D_{2t}}{\varepsilon_2} \Rightarrow D_{1t} = \frac{\varepsilon_1}{\varepsilon_2} D_{2t}.$$

Given the components of $\vec{D}_2$ tangent to the boundary, $2\hat{\alpha}_\theta - 3\hat{\alpha}_\phi$, and permittivities of $\varepsilon_1 = 4$, $\varepsilon_2 = 2$,

$$D_{2t} = \frac{4}{2}(2\hat{\alpha}_\theta - 3\hat{\alpha}_\phi) = 4\hat{\alpha}_\theta - 6\hat{\alpha}_\phi.$$

Combining the two components,

$$\vec{D}_1 = 2.2\hat{\alpha}_R + 4\hat{\alpha}_\theta - 6\hat{\alpha}_\phi.$$

3. a) To find the vector electric field, we may use

$$\vec{E} = -\nabla V.$$

Taking the partial derivative for each direction in the x-y-z plane,

$$-\left(\frac{\partial V}{\partial dx}\hat{\alpha}_x + \frac{\partial V}{\partial dy}\hat{\alpha}_y + \frac{\partial V}{\partial dz}\hat{\alpha}_z\right) = 2xy(z^2 + 1)\hat{\alpha}_x + x^2(z^2 + 1)\hat{\alpha}_y + 2xyz\hat{\alpha}_z \frac{\text{V}}{\text{C}}.$$

- b) In free space, the electric flux density $\vec{D} = \varepsilon_0 \vec{E}$, and so

$$\vec{D} = \varepsilon_0 2xy(z^2 + 1)\hat{\alpha}_x + \varepsilon_0 x^2(z^2 + 1)\hat{\alpha}_y + \varepsilon_0 2xyz\hat{\alpha}_z \frac{\text{C}}{\text{m}^2}$$

- c) Since the straight line path from A to B represents a change in z but no change in $x = 1$ or $y = 2$, we can simply evaluate the z-component at each point,

$$\int_{z=1}^{z=5} 4z \, dz = 48 \text{ V}.$$