

AMPAH PRAH MORTUARIES

STANDARD OPERATING PROCEDURES (SOPs)A

Post-Exposure Prophylaxis (PEP)

Infectious Exposures

Blood, visually bloody body fluids, semen, vaginal secretions, cerebrospinal fluid, peritoneal fluid, pleural fluid, pericardial fluid, synovial fluid, and amniotic fluid are all potentially infectious.

Non-Infectious Exposures

Feces, urine, vomitus, nasal secretions, saliva, sputum, sweat, tears, are NOT considered to be infectious unless they are **visibly bloody**.

A. Needle sticks, lacerations, or exposure of non-intact skin (i.e., open wounds, abrasions, chapped skin, or areas of dermatitis)

1. Allow wound to bleed but do not squeeze enough to bruise and do not suck wound
2. Wash the affected area gently with soap and water but do not scrub strongly or use nail brush

B. Mucous Membrane Exposure:

1. Irrigate the affected area (eye, mouth, etc.) with clean water
2. Inform QA team leader / supervisor to be evaluated for PEP as soon as exposure occurs.

Exposed Person

1. Exposed individuals should undergo rapid HIV testing immediately after exposure.

2. A brief medical history, medication list, and allergies should be obtained from the exposed to ensure no contraindications to ARVs and to tailor the regimen appropriately should there be any potential interactions
3. A pregnancy test should be offered to all exposed females.
4. ARVs should be started as soon as possible, ideally within 2 hours of exposure, up to 72 hours post-exposure for any of the following exposures listed above.