



WARNING!
READ
CAREFULLY!



The following document has not been checked for mistakes, and although every effort possible has been made to ensure that the information in this document is correct, it might still contain some mistakes. The following document is given “as is”, without any warranty whatsoever.

1 Integration

Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a continuous function, $P = \{a = x_0 < x_1 < \dots < x_n < b\}$ a partition of the interval $[a, b]$ and $\xi_k \in [x_k, x_{k+1}]$ points in each subinterval, then the sum

$$S(f, P, \xi) = \sum_{k=0}^{n-1} \left(\inf_{I_k} f \right) (x_{k+1} - x_k)$$

is called the Riemann sum to f and to

$$U(f, P) := \sum_{k=0}^{n-1} \left(\inf_{I_k} f \right) (x_{k+1} - x_k)$$

$$O(f, P) := \sum_{k=0}^{n-1} f(\sup_{I_k} f) (x_{k+1} - x_k)$$

(where $I_k = [x_k, x_{k+1}]$) are called the lower & upper Riemann sums. Similarly

$$\int_a^b f dx = \sup \{ U(f, P), P \in P \}$$

and

$$\int_a^b f dx = \inf \{ O(f, P), P \in P \}$$

are called lower and upper Integrals of f . f is called Riemann integrable if

$$\int_a^b f dx = \int_a^{\bar{b}} f dx$$

Fact

1. Each continuous function is Riemann integrable
2. Each monotone function is Riemann integrable

■

Properties

Let f, g Riemann integrable on I , $\alpha, \beta \in \mathbb{R}$ for

1.

$$\int_a^b (\alpha f + \beta g) dx = \alpha \int_a^b f dx + \beta \int_a^b g dx$$

2. if $f(x) \leq g(x), \forall x \in [a, b]$ then

$$\int_a^b f dx < \int_a^b g dx$$

?? attached ??
page 1
middle

Chopped
content,
page 1
middle

Chopped
content,
page 1
middle to
bottom

1 INTEGRATION

3.

$$\left| \int_a^b f(x) dx \right| \leq \int_a^b |f(x)| dx$$

4.

$$\left(\inf_I f \right) (b-a) \leq \int_a^b f(x) dx \leq \left(\sup_I f \right) (b-a)$$

5.

$$\int_a^b f dx = - \int_b^a f dx$$

6.

$$\int_a^b f dx = \int_a^c f dx + \int_c^b f dx, \quad \forall a, b, c \in \mathbb{R}$$

Fact (MWS der Integralrechnung)

$f : [a, b] \rightarrow \mathbb{R}$ continuous, then $\exists \rho \in (a, b)$

$$\int_a^b f(x) dx = f(\rho)(b-a)$$

■

Fact (Fundamental of calculus)

1. Let $f : [a, b] \rightarrow \mathbb{R}$ continuous. Define

$$F(x) := \int_a^x f(t) dt \quad \forall x \in [a, b]$$

then F is differentiable and $F' = f$. F is called primitive (Stammfunktion) of f .

2. If G is primitive of f , then $G = F + c$ for some constant

3. Let F be primitive of f then

$$\int_a^b f(x) dx = F(b) - F(a)$$

■

Can't re-read, page 3 middle to top

Can't re-read, page 3 middle to bottom

?? any ?? page 3 middle to bottom

How do we calculate integrals?

1. **Partial Integration:** Follows from product rule for differentiation

$$\int f(x)g'(x)dx = f(x)g(x) - \int f'(x)g(x)dx$$

$$\int_a^b f(x)g'(x)dx = f(x)g(x)|_a^b - \int_a^b f'(x)g(x)dx$$

2. **Substitution:** It follows from chain rule for differentiation

$$\int f(x)dx = \int f(\varphi(y))\varphi'(y)dy$$

$$\int_{\tilde{a}}^{\tilde{b}} f(x)dx = \int_a^b f(\varphi(y))\varphi'(y)dy$$

where $\varphi(a) = \tilde{a}$, $\varphi(b) = \tilde{b}$.

3. **Partialfractions:** To integrate rational functions of the form $\frac{P(x)}{Q(x)}$ with P, Q polynomials.

- If $\deg P < \deg Q$

We expand the rational function in simpler rational functions by finding the roots of $Q(x)$. Say

$$Q(x) = (x-a_1)\dots(x-a_n)\cdot(A_1x^2 + B_1x + C_1)\dots(A_nx^2 + B_nx + C_n)$$

Then for each linear factor $(x - a_i)$ of multiplicity r_i we need

$$\frac{\alpha_{i1}}{(x - a_i)} + \frac{\alpha_{i2}}{(x - a_i)^2} + \dots + \frac{\alpha_{ir}}{(x - a_i)^r}$$

with constants $\alpha_{i1}, \dots, \alpha_{ir}$. For each quadratic factor $A_ix^2 + B_ix + C_i$ of multiplicity S we have

$$\frac{\beta_{i1}x + \gamma_{i1}}{A_ix^2 + B_ix + C_i} + \dots + \frac{\beta_{iS}x + \gamma_{iS}}{A_ix^2 + B_ix + C_i}$$

with constants β_{ij}, γ_{ij} , $j = 1, \dots, S$.

- If $\deg P \geq \deg Q$

We first do long division to unite

$$\frac{P}{Q} \cdot R(x) + \frac{\tilde{P}(x)}{Q(x)}$$

with polynomials $R(X)$ and $\tilde{P}(x)$ with $\deg \tilde{P} < \deg Q$. Proceed as in the first case

Can't understand 1 word out of two, last paragraph in page 5

Chopped content
page 5
middle to
bottom

1 INTEGRATION

Beispiel

Berechne

1.

$$\int_1^2 \frac{\sqrt{1+\ln(x)}}{x} dx$$

2.

$$\int \cos(x) \cosh(x) dx$$

3.

$$\int \frac{x^2 - x + 2}{x^3 - x^2 + x - 1} dx$$

Solution

1. Note $\frac{1}{x}$ is the derivative of $(\ln(x) + 1)$. Hence using substitution $u = (1 + \ln(x)) \Rightarrow du = \frac{1}{x} dx$

$$\int_1^2 \frac{(1 + \ln(x))^{\frac{1}{2}}}{x} dx = \int_1^{1+\ln(2)} u^{\frac{1}{2}} du = \left. \frac{u^{\frac{3}{2}}}{\frac{3}{2}} \right|_1^{1+\ln(2)} = \frac{2}{3} \left[(1 + \ln(2))^{\frac{3}{2}} - 1 \right]$$

2. This one is suitable for Integral by parts:

$$\begin{aligned} I &= \int \underbrace{\cos(x)}_u \underbrace{\cosh(x)}_{v'} dx & \begin{array}{ll} u = \cos(x) & v' = \cosh(x) \\ u' = -\sin(x) & v = \sinh(x) \end{array} \\ I &= \cos(x) \sinh(x) + \int \underbrace{\sin(x)}_u \underbrace{\sinh(x)}_{v'} dx & \begin{array}{ll} u = \sin(x) & v' = \sinh(x) \\ u' = \cos(x) & v = \cosh(x) \end{array} \\ I &= \cos(x) \sinh(x) + \left[\sin(x) \cosh(x) - \int \cos(x) \cosh(x) dx \right] \\ I &= \cos(x) \sinh(x) + \sin(x) \cosh(x) - I \\ 2I &= \cos(x) \sinh(x) + \sin(x) \cosh(x) \\ x &= \frac{1}{2} (\cos(x) \sinh(x) + \sin(x) \cosh(x)) + C \end{aligned}$$

3.

$$\int \frac{x^2 - x + 2}{x^3 - x^2 + x - 1} dx$$

Note: $x^3 - x^2 + x - 1 = x^2(x - 1) + (x - 1) = (x^2 + 1)(x - 1)$

$$\begin{aligned} \frac{x^2 - x + 2}{x^3 - x^2 + x - 1} &= \frac{A}{x - 1} + \frac{Bx + C}{x^2 + 1} = \frac{1}{x - 1} - \frac{1}{x^2 + 1} \\ A(x^2 + 1) + (x - 1)(Bx + C) &= x^2 - x + 2 \end{aligned}$$

1 INTEGRATION

$$\begin{aligned}x = 1 &\Rightarrow 2A = 2 \Rightarrow A = 1 \\x = 2 &\Rightarrow 5 + 2B + C = 4 \Rightarrow 2B + C = -1 \\x = 0 &\Rightarrow 1 - C = 2 \Rightarrow C = -1 \\&\Rightarrow B = 0\end{aligned}$$

$$\Rightarrow \int \frac{x^2 - x + 2}{x^3 - x^2 + x - 1} dx = \ln|x - 1| - \tan^{-1}(x) + C$$

Improper Integrals

The improper integral of an integrable function f on (a, b) which is integrable on any subinterval $[a', b']$ we define the improper integral as

$$\int_a^b f(x) dx := \lim_{a' \searrow a} \lim_{b' \nearrow b} \int_{a'}^{b'} f(x) dx$$

Example (Basisprüfung Frühling 2010)

Untersuche ob das uneigentliche Integral $\int_1^{\infty} \frac{1}{x^2+x} dx$ konvergiert. Falls ja, berechne den Wert.

Note: since $\frac{1}{x^2+x} < \frac{1}{x^2}$ and $\int_1^{\infty} \frac{1}{x^2} dx$ is convergent, so is $\int \frac{1}{x^2+x}$

$$\begin{aligned}\int \frac{1}{x^2+x} &= \int \left(\frac{1}{x} - \frac{1}{x+1} \right) dx \\&= \ln|x| - \ln|x+1| = \ln \left| \frac{x}{x+1} \right|\end{aligned}$$

$$\begin{aligned}\int_1^{\infty} \frac{1}{x^2+x} &= \lim_{b \rightarrow \infty} \int_1^b \frac{1}{x^2+x} dx \\&= \lim_{b \rightarrow \infty} \ln \left| \frac{x}{x+1} \right|_1^b \\&= \lim_{b \rightarrow \infty} \ln \left(\frac{b}{b+1} \right) - \ln \left(\frac{1}{2} \right) \\&= \ln(1) + \ln(2) = \ln(2)\end{aligned}$$

limit exists and is equal to $\ln(2)$

Fact

1.

$$\forall s \in \mathbb{R}, a > 0, \int_a^{\infty} \frac{dx}{x^s} = \begin{cases} \frac{a^{1-s}}{s-1} & s > 1 \\ \infty & s \leq 1 \end{cases}$$

1 INTEGRATION

2. If f is in $[a, \infty)$ continuous, and $\exists c$ and $s > 1$ so that $|f(x)| \leq \frac{c}{x^s}, \forall x \geq a$, then $\int_a^\infty f(x)dx$ converges (Majorantenkriterium)

3. If f is in $[a, \infty)$ continuous, and $\exists c > 0$ $f(x) \geq \frac{c}{x}, \forall x \geq a$, then $\int_a^\infty f(x)dx$ diverges to ∞ .

Can't understand,
page 9
bottom

Example

$$\int_1^\infty \frac{t^2 + 4}{(1 + 4t^2)^{\frac{3}{2}}} dt \text{ diverges}$$

$$\int_1^\infty \frac{t^2 + 4}{(1 + 4t^2)^{\frac{5}{2}}} dt \text{ converges}$$

Because

$$\text{if } t \text{ is large enough, say } t > 2 \begin{cases} t^2 < t^2 + 4 < 2t^2 \\ 4t^2 < 1 + 4t^2 < 5t^2 \end{cases}$$

Hence

$$\frac{t^2}{(5t)^{2\alpha}} < \frac{t^2 + 4}{(1 + 4t^2)^\alpha} < \frac{2t^2}{(4t)^{2\alpha}}$$

$$\frac{C_1}{(t)^{2\alpha-2}} < \frac{t^2 + 4}{(1 + 4t^2)^\alpha} < \frac{C_2}{(4t)^{2\alpha-2}}$$

$$\Rightarrow \text{if } \alpha = \frac{3}{2}$$

$$\frac{t^2 + 4}{(1 + 4t^2)^{\frac{3}{2}}} > \frac{C_1}{t}$$

And since $\int_1^\infty \frac{dt}{t}$ diverges, so does

$$\int_1^\infty \frac{t^2 + 4}{(1 + 4t^2)^{\frac{3}{2}}} dt$$

Similarly since for $\alpha = \frac{5}{3} : 2\alpha - 2 = \frac{10}{3} - 2 > 1$ and

$$\int \frac{C_2}{t^{\frac{4}{3}}} dt$$

converges, so does

$$\int \frac{t^2 + 4}{(1 + 4t^2)^{\frac{5}{3}}} dt < \int \frac{C_2}{t^{\frac{4}{3}}} dt$$

2 Differential Equations

2.1 Linear Differential equations with constant coefficients

To solve a linear differential equation of the form $Ly = b(x)$, where

$$L := \frac{d^n}{dx^n} + a_{n-1} \frac{d^{n-1}}{dx^{n-1}} + \cdots + a_1 \frac{d}{dx} + a_0$$

$b(x)$ a function, and $a_i \in \mathbb{R}$

1. Find a homogeneous solution y_H , namely a solution of $Ly = 0$
2. Find a special solution y_S of $Ly = b(x)$ using the method “Ansatz von Typ der rechten Seite”
3. Then the general solution is given by

$$y = y_H + y_S$$

How to find the homogeneous solution y_H of $Ly = 0$

1. Find the characteristic polynom of L , namely

$$P_l(\lambda) = \lambda^n + a_{n-1}\lambda^{n-1} + \cdots + a_1\lambda + a_0$$

2. Fact

If $\lambda_1, \dots, \lambda_r \in \mathbb{C}$ are the Pairwise distinct roots of $P(\lambda) = 0$ with associated multiplicates m_1, \dots, m_r , then the functions

$$x \rightarrow x^k e^{\lambda_j x}, 1 \leq k \leq m_j, 0 \leq k \leq m_j$$

Form a system of fundamental solutions of the homogeneous equations $Ly = 0$

Variant

If L has real coefficients, every pair of complex, non-real roots $\lambda_j = \mu_j \pm i v_j$ of multiply m_j give a fundamental solution

$$x^k e^{(\mu_j \pm i v_j)x} = x^k e^{\mu_j x} (\cos(v_j x) \pm i \sin(v_j x))$$

for $0 \leq k < m_j$. So one can, as a basis, take

$$x^k e^{\mu_j x} \cos v_j x \text{ and } x^k e^{\mu_j x} \sin v_j x$$

instead of

$$x^k e^{(\mu_j + i v_j)x} \text{ and } x^k e^{(\mu_j - i v_j)x}$$

Then the general homogeneous solution is of the form

$$y_H(x) = \sum_{j=1}^r \sum_{k=0}^{m_j} c_{jk} x^k e^{\lambda_j x}$$

with constant c_{jk}

Not really sure if it says variant, page 13 middle to bottom

■

2 DIFFERENTIAL EQUATIONS

How to find the special solution of Inhomogeneous $Ly = b(x)$ using the method of “Ansatz”

Fact

1. Let $\lambda \in \mathbb{C}$. if λ is not a solution of $P_L(\lambda)$, then the inhomogeneous DGL $Ly = e^{\lambda x}$ has particular solutions $y = \frac{1}{P_L(\lambda)} e^{\lambda x}$
2. Let $\lambda \in \mathbb{C}$, m its multiplicity as a solution of $P_L(\lambda) = 0$ (m can be zero, which means λ is not a solution of $P_L(\lambda) = 0$).

Let $Q(x)$ a polynomial of degree k . Then a particular solution of $Ly(x) = Q(x)e^{\lambda x}$ is of the form

$$y(x) = R(x)e^{\lambda x}$$

for a polynomial $R(x)$ of degree $k + m$

3. Let l has a real coefficient. Let $\mu, v \in \mathbb{R}$, m the multiplicity of $\mu \pm iv$ as a solution of $P_L(\lambda) = 0$ ($m = 0$ means $\mu \pm iv$ is not a root of P_L). Let $Q(x)$ and $R(x)$ be polynomials of degree $\leq k$. Then the particular solution of the inhomogeneous Differential equation

$$Ly = Q(x)e^{\mu x} \cos vx + R(x)e^{\mu x} \sin x$$

is of the form

$$y(x) = S(x)e^{\mu x} \cos vx + T(x)e^{\mu x} \sin x$$

for polynomials S, T of degree $\leq k + m$

■

Chopped
content,
page 14
bottom

Chopped
content,
page 15
top

Boundary (Rand) or Initial value Problems (Anfangswert)

When we are given a differential equation $Ly = b(x)$ together with either

Boundary values	or	Initial values
$y(a_1) = A_1$		$y(a) = A_1$
$y(a_2) = A_2$		$y'(a) = A_2$
\vdots		\vdots
$y(a_n) = A_n$		$y^{(n-1)}(a) = A_n$

We first find the general solution $y = y_H + y_S$. Then we determine the constants C_1, \dots, C_n in the homogeneous solution using the given boundary or initial values.

Example

1. Basisprüfung Frühling 2011

- (a) Bestimme alle Lösungen $y = y(x)$ der DGL $y^{(4)} - y = 0$ welche für $|x| \rightarrow \infty$ beschränkt bleiben
- (b) Bestimme eine Lösung $y = y(x)$ der DGL $y^{(4)} - y = e^{-x} + x$

2 DIFFERENTIAL EQUATIONS

Solution

- (a) We first find the characteristic polynomial

$$x^4 - 1 = 0 \Rightarrow (x^2 - 1)(x^2 + 1) = 0$$

$$\lambda = \pm 1, \lambda = \pm i$$

The homogeneous solution is of the form

$$y_H = C_1 e^x + C_2 e^{-x} + C_3 \cos x + C_4 \sin x$$

Since

$$e^x \rightarrow \infty \text{ as } x \rightarrow \infty$$

$$e^{-x} \rightarrow \infty \text{ as } x \rightarrow -\infty$$

The solutions that satisfy are $y_H = C_3 \cos x + C_4 \sin x$ with $C_3, C_4 \in \mathbb{R}$

- (b) We use the superposition principle: Let $b_1(x) = e^{-x}$, $b_2(x) = x$

i. y_{1S} is a solution of $Ly = b_1(x)$

ii. y_{2S} is a solution of $Ly = b_2(x)$

Then $y_{1S} + y_{2S}$ is a solution of $Ly = e^{-x} + x$. Since $\lambda = -1$ is a root of $P_L(\lambda) = 0$ of multiplicity, i.e. since e^{-x} is also a solution of $Ly = 0$

For y_{1S} we take as “ansatz” $y_{1S} = Cxe^{-x}$. For y_{2S} we take as “ansatz” $y_{2S} = Dx + E$ a polynomial of degree 1. So a special solution is of the form

$$y_S = Cxe^{-x} + Dx + E$$

To find the constants C, D, E, F we put this solution into the differential equation $y^{(4)} - y = e^{-x} + x$.

$$y'(x) = C[e^{-x} - xe^{-x}] + D$$

$$y''(x) = C[-e^{-x} - [e^{-x} - xe^{-x}]] = C[-2e^{-x} + xe^{-x}]$$

$$y^{(3)}(x) = C[2e^{-x} + [e^{-x} - xe^{-x}]] = C[3e^{-x} - xe^{-x}]$$

$$y^{(4)}(x) = C[-3e^{-x} + [e^{-x} - xe^{-x}]] = C[-4e^{-x} + xe^{-x}]$$

Then

$$y_S^{(4)} - y_S(x) = C \cdot 4e^{-x} + Cxe^{-x} - [Cxe^{-x} + Dx + E]$$

$$\Rightarrow 4Ce^{-x} - Dx - E = e^{-x} + x$$

$$\Rightarrow 4C = 1, D = -1$$

Hence a particular solution is

$$y_S = \frac{1}{4}xe^{-x} - x$$

General solution is

$$y = y_H + y_S$$

$$= C_1 e^x + C_2 e^{-x} + C_3 \cos x + C_4 \sin x + \frac{1}{4}xe^{-x} - x$$

Can't read, page 16 bottom

2 DIFFERENTIAL EQUATIONS

2. Basisprüfung Sommer 2013

- (a) Für welche Werte des Parameters $a \in \mathbb{R}$ strebt die allgemeine Lösung $y(x)$ der DGL $y'' + 2y' + ay = 0$ unabhängig von den Anfangsbedingungen gegen 0 für $x =$
- (b) Finden Sie eine homogene DGL 2. Ordnung mit konstanten Koeffizienten, deren allgemeine Lösung $y(x) = e^{-x} + 2xe^{-x}$ ist. Was sind dann die Anfangsbedingungen bei $x = 0$?

Chopped content, page 18 bottom

Solution

- (a) The characteristic polynomial of $y'' + 2y' + ay = 0$ is $\lambda^2 + 2\lambda + a = 0$. The roots are

$$\lambda_{1,2} = \frac{-2 \pm \sqrt{4 - 4a}}{2} = -1 \pm \sqrt{1 - a}$$

- If $1 - a < 0$ then there are 2 complex conjugate roots. Let $|1 - a| = b^2$ then the general solution is of the form

$$y(x) = C_1 e^{-x} \cos(bx) + C_2 e^{-x} \sin(bx)$$

and this goes to zero as $b \rightarrow 0$, hence a since e^{-x} decays to zero.

- If $1 - a = 0$ then -1 is a double root. Then the solutions are of the form $C_1 e^{-x} + C_2 x e^{-x}$ and this will not decay to zero as $x \rightarrow \infty$

Similarly if $1 - a > 0$ then one of the roots will be positive if $\sqrt{1 - a} > 1$ and that root can lead to a growing solution.

Hence when $\sqrt{1 - a} > 1$, or $1 - a > 1$, or when $a < 0$, then $-1 + \sqrt{1 - a} > 0$. If on the other hand $0 < a < 1$ then again $-1 \pm \sqrt{1 - a} < 0$ and both solutions are decaying. Hence if $0 < a < 1$ or $a > 1$ then the solutions are decaying of initial values, and if $a = 1$ or $a < 0$ then the solution can be growing.

- (b) Let $Ly = 0$ be of the form $y'' + a_1 y' + a_0 y = 0$ then the characteristic polynomial is $\lambda^2 + a_1 \lambda + a_0 = 0$. Since we want e^{-x} and $x e^{-x}$ to be the solutions, we should have $\lambda = -1$ with multiplicity 2, that means

$$P_L(\lambda) = \lambda^2 + 2\lambda + 1$$

and the differential equation is $y'' + 2y' + y = 0$. The general solution of this equation is

$$y_H = C_1 e^{-x} + C_2 x e^{-x}$$

Since the solution is $e^{-x} + 2x e^{-x}$, $C_1 = 1$, $C_2 = 2$ we must have $y(0) = e^{-0} + 2 \cdot 0 = 1$

$$y'(0) = -e^{-x} + 2[e^{-x} - x e^{-x}] \Big|_{x=0} = -1 + 2 = 1$$

So we can choose $y(0) = 1$, $y'(0) = 1$

Can't read, page 19 middle

Can't understand, page 20 middle to bottom

2.2 Solving Differential Equations by separation of variables

Example

1. (Frühling 2011) Bestimme die Lösung $y = y(x)$ der DGL $y' = e^{x-y}$ mit $y(0) = 0$

Solution

$$\begin{aligned}\frac{dy}{dx} &= e^x \cdot e^{-y} \\ \Rightarrow e^y dy &= e^x dx \\ \Rightarrow \int e^y dx &= \int e^x dx \\ \Rightarrow e^y &= e^x + C\end{aligned}$$

Since $y(0) = 0 \Rightarrow e^0 = e^0 + C \Rightarrow C = 0$ and $y = x$

2. (Herbst 2010) Bestimme eine Lösung der DGL

$$y + xy' = \frac{1}{y} \left(x + \frac{1}{x} \right), x > 0$$

mit $y(1) = 2, y(x) > 0, \forall x$. (Hinweise: substituieren $u(x) = xy(x)$).

$$u'(x) = y(x) + xy'(x)$$

Left hand side is

$$\begin{aligned}u'(x) &= \frac{x}{y} + \frac{1}{xy} = \frac{x^2}{u} + \frac{1}{u} = \frac{x^2 + 1}{u} \\ \Rightarrow \frac{u'}{u} &= x^2 + 1 \Rightarrow \ln u = \frac{x^3}{3} + x + C \\ y(1) = 2 &\Rightarrow u(1) = 1 \cdot y(1) = 2 \\ \Rightarrow \ln 2 &= \frac{1}{3} + 1 + C \Rightarrow C = (\ln 2) - \frac{4}{3} \\ u &= C e^{\frac{x^3}{3} + x}\end{aligned}$$

In my opinion there is an error, my solution (the same as Wolfram Mathematica) is $y(x) = \pm \sqrt{\frac{c+2x^3+6x}{3}} \frac{1}{x}$

2.3 Differentiation in many variables

A function $f : \Omega \subset \mathbb{R}^n \rightarrow \mathbb{R}$ is (total) differentiable in x_0 if it exists a linear map $A : \mathbb{R}^n \rightarrow \mathbb{R}$ such that

$$f(x) = f(x_0) + A(x - x_0) + R(x_0, x)$$

Where $\lim_{x \rightarrow x_0} \frac{R(x_0, x)}{|x - x_0|} = 0$. In this case A is called the differential of f at x_0 and it's denoted as $(df)(x_0)$.

Let (A_1, A_2, \dots, A_n) be a matrix representation of the linear map $A : \mathbb{R}^n \rightarrow \mathbb{R}$ (wrt to the standard Basis). Then f differentiable at x_0 means:

$$f(x) = f(x_0) + A_1(x^1 - x_0^1) + A_2(x^2 - x_0^2) + \dots + A_n(x^n - x_0^n) + R(x, x_0)$$

2 DIFFERENTIAL EQUATIONS

$$P(x, x_0) = f(x_0) + [A_1 \quad \dots \quad A_n] \begin{bmatrix} x^1 - x_0^1 \\ \dots \\ x^n - x_0^n \end{bmatrix}$$

is the equation of the tangent plane at the point $f(x_0)$ on the surface formed by the graph of f .

Fact

if $f : \Omega \rightarrow \mathbb{R}$ is differentiable in $x_0 \in \Omega$ then the partial derivative exist and the differential $df(x_0)$ has the matrix representation

$$\left(\frac{\partial f}{\partial x}(x_0) \quad \dots \quad \frac{\partial f}{\partial x^n}(x_0) \right) = \nabla f$$

the gradient of f .

■

Fact

f differentiable in $x_0 \Rightarrow f$ is continuous in x_0 .

■

Fact

If all partial derivatives of f are continuous then f is differentiable

■

Using these last two facts and the definition of differentiability one can study if a given function is differentiable or not.

2.3.1 Differentiation rules

Let $f, g : \Omega \rightarrow \mathbb{R}$ be differentiable in x_0 . Then:

1. $d(f \pm g)(x_0) = df(x_0) \pm dg(x_0)$
2. $d(fg)(x_0) = g(x_0)df(x_0) + f(x_0)dg(x_0)$
3. $d(f/g)(x_0) = \frac{g(x_0)df(x_0) - f(x_0)dg(x_0)}{(g(x_0))^2}$
4. Let $h : \mathbb{R} \rightarrow \mathbb{R}$ be differentiable in $g(x_0)$, then

$$d(hog)(x_0) = h'(g(x_0))dg(x_0)$$

5. Let $H : I \subset \mathbb{R} \rightarrow \Omega \subset \mathbb{R}^n$ be differentiable in $t_0 \in \mathbb{R}$ and f differentiable in $H(t_0)$. Then

$$\frac{d}{dt}(f \circ H)(t_0) = df(H(t_0))H'(t_0)$$

where $H(t) = (H_1(t), \dots, H_n(t))$ and $H'(t) = (H'_1(t), \dots, H'_n(t))$

2 DIFFERENTIAL EQUATIONS

2.3.2 Directional derivative

The directional derivative of f in the direction of a unit vector $e \in \mathbb{R}^n - \{0\}$ is given by $d_e f(x_0) = \nabla f(x_0) \cdot e$.

2.3.3 Particular higher derivatives

One can similarly define higher derivatives order partial derivatives for functions $f \in C^m(\Omega)$.

Fact (Schwarz)

if $f \in C^2(\Omega)$ then $\frac{\partial^2 f}{\partial x^i \partial x^j} = \frac{\partial^2 f}{\partial x^j \partial x^i}$ and in general for $f \in C^m(\Omega)$ all n partial derivatives of f of order $\leq m$ are independent of the order of differentiation. Using higher order derivatives one can analogous to the 1-dimensional case define a Taylor approximation of f .

■

Fact

Let $f \in C^m(\Omega)$, $f : \Omega \rightarrow \mathbb{R}$, $\Omega \in \mathbb{R}$ and $x_1, x_0 \in \Omega$. Then

$$f(x_1) = f(x_0) + \nabla f(x_0)(x_1 - x_0) + \frac{1}{2} \sum_{i,j=1}^2 \frac{\partial^2 f}{\partial x^i \partial x^j}(x_0)(x_1^i - x_0^i)(x_1^j - x_0^j) + R_3(f, x_1, x_0)$$

Where $\lim_{x_1 \rightarrow x_0} \frac{R(f, x_1, x_0)}{\|x_1 - x_0\|} = 0$

■

The analogue of the second derivative is given by the matrix of partial derivatives of order 2. This matrix is called the Hesse-matrix of f .

$$\text{Hess}(f) = \nabla^2 f = \left(\frac{\partial^2 f}{\partial x^i \partial x^j} \right)_{i,j=1,\dots,n}$$

2.3.4 The extrema of a function $f : \Omega \rightarrow \mathbb{R}$

Definition

A point $x \in \Omega$ is called a critical point if $\nabla f(x) = 0$.

Fact

f differentiable, x_0 is called local extrema of f then x_0 is a critical point

■

2 DIFFERENTIAL EQUATIONS

Fact

Let x_0 be a critical point of f . Then we have three different cases:

1. x_0 is a local minima if $\nabla^2 f(x_0)$ is positive definite.
2. x_0 is a local maxima if $\nabla^2 f(x_0)$ is negative definite.
3. Otherwise it is a saddle point (Sattelpunkt).

■

Fact

Let $f : \Omega \rightarrow \mathbb{R}$ be continuous and differentiable on an open set $\Omega \subset \mathbb{R}^n$. Let $\partial\Omega$ be the boundary of Ω . Then every global extrema of f is either a critical point of f in Ω or a global extremal point of $f|_{\partial\Omega}$ (f restricted to the boundary)

■

Example (FS 2011)

Sei $f(x, y) = 4x^2y^2 - x^2 - 4y^2 + 1$. Bestimme die globalen Extrema von f auf dem Gebiet $\Omega = \{(x, y) = \frac{x^2}{4} + y^2 \leq 1, y \geq 0\}$.

Solution

We first find the critical points:

$$\nabla f = \begin{pmatrix} 8xy^2 - 2x \\ 8x^2y - 8y \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

$$2x(4y^2 - 1) = 0 \Rightarrow x = 0 \text{ or } y = \pm \frac{1}{2}$$

$$8y(x^2 - 1) = 0 \Rightarrow x = \pm 1 \text{ or } y = 0$$

$(0, 0), (\pm 1, \pm \frac{1}{2})$ are the critical points of f . Since $y \geq 0$ we only take $P_1 = (0, 0), P_{2,3} = (\pm 1, \frac{1}{2})$. Then we need to compute $\text{Hess}(f)$.

$$\text{Hess}(f) = \begin{pmatrix} 8y^2 - 2 & 16xy \\ 16xy & 8x^2 - 8 \end{pmatrix}$$

$$\text{Hess}(f)(0, 0) = \begin{pmatrix} -2 & 0 \\ 0 & -8 \end{pmatrix} \Rightarrow \text{negative definite} \Rightarrow \text{local maxima}$$

$$\text{Hess}(f)(1, \frac{1}{2}) = \begin{pmatrix} 0 & 8 \\ 8 & 0 \end{pmatrix} \Rightarrow \text{indefinite}$$

$$\text{Hess}(f)(-1, \frac{1}{2}) = \begin{pmatrix} 0 & -8 \\ -8 & 0 \end{pmatrix} \Rightarrow \text{indefinite}$$

To find global extrema we need to look at f on the boundary of Ω , which is the curve $\frac{x^2}{4} + y^2 = 1$ and the line $y = 0$. First on the line $y = 0$ let $g = f|_{y=0} = -x^2 + 1$.

$$g'(x) = -2x \Rightarrow x = 0, P_1(0, 0) \text{ is a point we need to check}$$

2 DIFFERENTIAL EQUATIONS

We also need to check the corners $P_{4,5} = (\pm 2, 0)$. On the ellipse: let $h = f|_{\frac{x^2}{4} + y^2 = 1} = -x^4 + 4x^2 - 3$.

$$h'(x) = -4x^3 + 8x = 0 \Rightarrow P_6 = (0, 1), P_{7,8} = (\pm\sqrt{2}, \frac{1}{\sqrt{2}})$$

Now we look at the values of f at these points. $f(P_1) = f(0, 0) = 1, f(P_{2,3}) = 0, f(P_{4,5}) = -3, f(P_6) = -3, f(P_{7,8}) = 1$. f has also a minima at $(\pm 2, 0), (0, 1)$ and a maxima at $(0, 0), (\pm\sqrt{2}, \frac{1}{\sqrt{2}})$.

Example (FS 2010)

1. Bestimme das Taylorpolynom erster Ordnung der Funktion $f(x, y) = e^{x^2}(x + y)$ um dem Punkt $(1, 1)$.
2. Bestimme $c \in \mathbb{R}$ so dass der Vektor $\begin{pmatrix} 1 \\ -1 \\ c \end{pmatrix}$ tangential an den Graphen $g(f) = \{(x, y, f(x, y)) : (x, y) \in \mathbb{R}^2 \text{ im Punkt } (1, 1, z) \text{ liegt.}\}$

Solution

1.

$$\nabla f = \begin{pmatrix} 2xe^{x^2}(x+y) + e^{x^2} \\ e^{x^2} \end{pmatrix}, \nabla f(1, 1) = \begin{pmatrix} 5e \\ e \end{pmatrix}, f(1, 1) = 2e$$

$$\begin{aligned} f(x, y) &= 2e + \begin{pmatrix} 5e \\ e \end{pmatrix} \cdot (x-1, y-1) + r_2(x, y) \\ &= 2e + 5e(x-1) + e(y-1) + (x-1)(\nabla^2 f)(t) \begin{pmatrix} x-1 \\ y-1 \end{pmatrix} \end{aligned}$$

The Taylor polynomial of order 1 is $z = 2e + 5e(x-1) + e(y-1)$

2. The vector $(1, -1, c)^T$ must be perpendicular to the normal vector of the plane, that is (from part 1) $n = (5e, e, -1)^T$. Hence

$$(1, -1, c) \cdot (5e, e, -1) = 0 \Rightarrow 4e - c = 0 \Rightarrow c = 4e$$

2.4 Line (Weg) integral

Let $v : \Omega \rightarrow \mathbb{R}^n$ be a vector field and γ a curve with parametrization $\gamma : [a, b] \rightarrow \Omega, t \rightarrow \gamma(t)$. Then the line integral of v along γ is defined as

$$\int_{\gamma} v ds = \int_a^b \langle v(\gamma(t)), \gamma'(t) \rangle dt$$

2 DIFFERENTIAL EQUATIONS

Facts

1. $\int_{\gamma} v ds$ is independent of the parametrization of the path.
2. $\int_{\gamma_1 + \gamma_2} v ds = \int_{\gamma_1} v ds + \int_{\gamma_2} v ds$
3. $\int_{\gamma} v ds = - \int_{-\gamma} v ds$
4. If v is the gradient vector field associated to a function f i.e. $v = df$ then $\int_{\gamma} v ds = f(\gamma(b)) - f(\gamma(a))$, where $\gamma : [a, b] \rightarrow \Omega$.

Equivalent one can change everything in terms of 1-forms $\lambda = \lambda_1 dx^1 + \lambda_2 dx^2 + \dots + \lambda_n dx^n$. Then

$$\int_{\gamma} \lambda = \int_a^b \lambda(\gamma(t)) \gamma'(t) dt$$

Fact

$\lambda : \Omega \rightarrow L(\mathbb{R}^n, \mathbb{R})$ a constant 1-form, then the following are equivalent:

1. $\exists f \in C^1(\Omega) : df = \lambda$
2. For every 2 continuous C'-paths γ_1, γ_2 with $\gamma_i : [a_i, b_i] \rightarrow \Omega$ with the same beginning and ending points:

$$\int_{\gamma_1} \lambda = \int_{\gamma_2} \lambda$$

3. For every closed curve γ , $\int_{\gamma} \lambda = 0$

■

Definition

A vector field $V : \Omega \rightarrow \mathbb{R}^n$ is called conservative if $\int_{\gamma} V ds = 0$ for every closed curve γ .

Fact

For a simply connected region Ω , we have


$$V \text{ is conservative} \iff v = \nabla f \text{ for some function } f$$

■

Example (FS 2010)

Berechne das Wegintegral $\int_{\gamma} v ds$ entlang des eingezeichneten Weg für das Vektorfeld $v(x, y) = \begin{pmatrix} xy^2 \\ -y \end{pmatrix}$

2 DIFFERENTIAL EQUATIONS



Solution

We first parametrize the curves:

1. $\gamma_1 = (1, t), t \in [0, e] \Rightarrow \gamma'_1 = (0, 1)$
2. $-\gamma_2 = (t, e^t), t \in [0, 1] \Rightarrow -\gamma'_2 = (1, e^t)$
3. $\gamma_3 = (0, 1 - t), t \in [0, 1] \Rightarrow \gamma'_3 = (0, -1)$
4. $\gamma_4 = (t, -t), t \in [0, 1] \Rightarrow \gamma'_4 = (1, -1)$

$$\int_{\gamma} v ds = \int_0^e (t^2, -t) \cdot (0, 1) dt + \int_1^0 (te^{2t}, -e^t) \cdot (1, e^t) dt + \dots$$

See the last example of Green's theorem for the rest.

2.5 Integration in \mathbb{R}^n

The Riemann integral in \mathbb{R}^n is constructed in analogue way to the case in $n=1$. With Riemann sums over subintervals is replaced with sums over "subrectangles". dx is replaced with an n -dimensional volume element $dvol_n$ which we denote also by $d\mu(x)$.

Fact

For a normal-region

$$D = \{(x, y) \in \mathbb{R}^2 : a \leq x \leq b, \phi(x) < y < \psi(x)\}$$

Where ϕ, ψ are continuous functions: $\mathbb{R} \rightarrow \mathbb{R}$ we have

$$\int_D f d\mu(x) = \int_a^b \int_{\phi(x)}^{\psi(x)} f(x, y) dy dx$$

■

2 DIFFERENTIAL EQUATIONS

Fact

For a rectangle $Q = [a, b] \times [c, d] \in \mathbb{R}^2$

$$\int_Q f d\mu = \int_a^b \int_c^d f dy dx = \int_c^d \int_a^b f dx dy$$

■

2.5.1 Substitution in \mathbb{R}^n

Let $u, v \in \mathbb{R}^n$ open intervals, $\Phi : u \rightarrow v$ bijective continuous differentiable with $\det(d\Phi(y)) \neq 0 \forall y \in u$. Then for $f : v \rightarrow \mathbb{R}$ continuous we have


$$\int_v f(x) d\mu(x) = \int_u f(\Phi(y)) |\det(d\Phi(y))| d\mu(y)$$

Example (FS 2010)

Bestimme das Integral

$$\int_{\Omega} x \sqrt{x^2 + y^2} \log(\sqrt{x^2 + y^2}) d\mu(x, y)$$

wobei



Solution

Given the symmetry of the region it's better to use polar coordinates. Let

$$\begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} r \cos(\sigma) \\ r \sin(\sigma) \end{pmatrix} \Rightarrow dx dy = r dr d\sigma$$

2 DIFFERENTIAL EQUATIONS

The region in (r, σ) domain is simply $1 \leq r \leq 2, \frac{\pi}{4} \leq \sigma \leq \frac{7\pi}{4}$.

$$\begin{aligned} \int_{\Omega} x \sqrt{x^2 + y^2} \log \sqrt{x^2 + y^2} d\mu(x, y) &= \int_{\frac{\pi}{4}}^{\frac{7\pi}{4}} \int_1^2 r \cos(\sigma) r \log(r) r dr d\sigma \\ &= \int_{\frac{\pi}{4}}^{\frac{7\pi}{4}} \cos(\sigma) d\sigma \int_1^2 r^3 \log(r) dr \\ &= \left[-\sqrt{2} + \frac{1}{4} x^4 \log(x) - \frac{x^4}{16} \right]_1^2 \\ &= \log(16) - \frac{15}{16} \end{aligned}$$

2.6 Green's theorem

Let $\Omega \subset \mathbb{R}^2$ whose boundary $\partial\Omega$ has a C' parametrization. Let $U \subset \Omega$ and $f = \frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y}$ where $P, Q \in C'(U)$. Then

$$\int \int_{\Omega} \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) d\mu = \int_{\partial\Omega} P dx + Q dy$$

Let $V = (P, Q)$ be a vectorfield then

$$\int_{\partial\Omega} V ds = \int \int_{\Omega} (\text{rot}(V)) d\mu$$

Where $\text{rot}(V) = \frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y}$ and the line integral is taken around the boundary of Ω in positive sense.

add last
example
(it's either
in my no-
tes nor in
her)