



# Support Session

Wednesdays from 13:00 to 14:00.

Venue: TB 318+

The support session is another opportunity (in addition to the office hours) for you to get your questions answered by tutors.



# Mid-Semester Examination

Wednesday 3<sup>rd</sup> November from 13:30 to 14:30.

Confirm your venue from CPSO

Topics covered would be those topics treated in Seminars 1 to 4


Revision Materials would be made available on Moodle, and you would be  
informed when they are live



Please make sure you bring a permissible calculator to the examination

Permissible Calculator  
CASIO fx-82E family

*fx-82ES PLUS*



*fx-82ES PLUS 2nd edition*



*fx-82ES PLUS A*





# Lecture 4


Topics covered in this lecture session

1. Inverse Trigonometric functions.
2. Expressing  $a \cos x + b \sin x$  in the form  $r \cos(\theta - x)$ .



# Inverse Trigonometric Functions

The graph of the sine function over  $\mathbb{R}$ , indicates that it is not one-one however, if we restrict the domain to  $\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$ , then sine function is one-one and its inverse exists.





It is denoted by  $\sin^{-1}$  **or**  $\arcsin$  and is defined by

$$y = \sin x \quad \Leftrightarrow \quad x = \sin^{-1} y \quad ; \quad -\frac{\pi}{2} \leq x \leq \frac{\pi}{2}$$



# Inverse Trigonometric Functions

Inverse Function	Domain of Inverse function $\equiv$ Range of Trigonometric function	Range of Inverse function <i>i.e.</i> <u>Restricted Domain</u> for Trigonometric function	Graph of Inverse Trigonometric function
$\cos^{-1} x$ or arccos	$[-1, 1]$	$[0, \pi]$	
$\sin^{-1} x$ or arcsin	$[-1, 1]$	$\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$	





# Inverse Trigonometric Functions

Inverse Function	Domain of Inverse function $\equiv$ Range of Trigonometric function	Range of Inverse function <i>i.e.</i> <u>Restricted Domain</u> for Trigonometric function	Graph of Inverse Trigonometric function
$\tan^{-1} x$ or arctan	$\mathbb{R}$	$\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$	
$\sec^{-1} x$ or arcsec	$\mathbb{R} - (-1, 1)$	$[0, \pi] - \left\{\frac{\pi}{2}\right\}$	




# Inverse Trigonometric Functions

Inverse Function	Domain of Inverse function $\equiv$ Range of Trigonometric function	Range of Inverse function <i>i.e.</i> <u>Restricted Domain</u> for Trigonometric function	Graph of Inverse Trigonometric function
$\operatorname{cosec}^{-1} x$ or $\operatorname{arccosec}$	$\mathbb{R} - (-1, 1)$	$\left[-\frac{\pi}{2}, \frac{\pi}{2}\right] - \{0\}$	
$\cot^{-1} x$ or $\operatorname{arccot}$	$\mathbb{R}$	$(0, \pi)$	






# Inverse Trigonometric Functions




Reference Angle  $\theta \in (0, 2\pi)$

Restricted domain for Inverse Trigonometric functions

$[0, \pi]$ , used for  $\cos^{-1}$



$[-\frac{\pi}{2}, \frac{\pi}{2}]$ , used for  $\sin^{-1}$



$(-\frac{\pi}{2}, \frac{\pi}{2})$ , used for  $\tan^{-1}$




# Inverse Trigonometric Functions

Find the values of:

(i)  $\cos^{-1} \left( \frac{-1}{\sqrt{2}} \right)$

(ii)  $\tan^{-1} \left( \frac{-1}{\sqrt{3}} \right)$

(iii)  $\sin^{-1} \left( \sin \left( \frac{7\pi}{4} \right) \right)$





## Inverse Trigonometric Functions

### Question 4-N36-Q1

Solve  $\cos^{-1} \left( \cos \left( \frac{14\pi}{3} \right) \right) - \sin^{-1} \left( -\frac{\sqrt{3}}{2} \right)$

A  $\frac{\pi}{2}$

B  $\pi$

C  $2\pi$



# Expressing $a \cos x + b \sin x$ in the form $r \cos(\theta - x)$

Sometimes it is important to express

$f(x) = a \cos x + b \sin x$  in the form  $r \cos(\theta - x)$ ,

so as to

- determine the range of  $f$ ;
- find the period of  $f$ ;
- sketch the graph of the function  $f$ .



# Expressing $a \cos x + b \sin x$ in the form $r \cos(\theta - x)$

**The Method:** Let  $a = r \cos \theta$   
 $b = r \sin \theta$

where  $\theta$  and  $r$  are to be determined.

- Squaring and adding  $\Rightarrow a^2 + b^2 = r^2 \Rightarrow r = \sqrt{a^2 + b^2}$
- Dividing the second equation by the first equation, gives

$$\frac{r \sin \theta}{r \cos \theta} = \frac{b}{a} \Rightarrow \tan \theta = \frac{b}{a} \quad (\text{from which } \theta \text{ can be found})$$



# Expressing $a \cos x + b \sin x$ in the form $r \cos(\theta - x)$

Thus,  $f(x) = a \cos x + b \sin x$

$$= r \cos \theta \cos x + r \sin \theta \sin x$$

$$= r [\cos \theta \cos x + \sin \theta \sin x]$$

$$= r \cos(\theta - x) \quad \text{or} \quad r \cos(x - \theta)$$

$$\text{because, } \cos(-\theta) = \cos \theta$$



# Expressing $a \cos x + b \sin x$ in the form $r \cos(\theta - x)$

1. Prove that  $\cos 2x - \sqrt{3} \sin 2x = 2 \cos \left( 2x + \frac{\pi}{3} \right)$ .

2. Express  $\sin x - \sqrt{3} \cos x$  in the form  $R \sin(x - \alpha)$ ,  
where  $R > 0$  and  $0 < \alpha < \pi/2$ .

Hence (i) sketch the graph of  $y = f(x) = \sin x - \sqrt{3} \cos x$ ;

(ii) find the range of  $f$ ;

(iii) find the period of  $f$ .



# Expressing $a \cos x + b \sin x$ in the form $r \cos(\theta - x)$

1. Prove that  $\cos 2x - \sqrt{3} \sin 2x = 2 \cos \left( 2x + \frac{\pi}{3} \right)$ .

2. Express  $\sin x - \sqrt{3} \cos x$  in the form  $R \sin(x - \alpha)$ ,  
where  $R > 0$  and  $0 < \alpha < \pi/2$ .

Hence (i) sketch the graph of  $y = f(x) = \sin x - \sqrt{3} \cos x$ ;

(ii) find the range of  $f$ ;

(iii) find the period of  $f$ .





# Expressing $a \cos x + b \sin x$ in the form $r \cos(\theta - x)$

1. Prove that  $\cos 2x - \sqrt{3} \sin 2x = 2 \cos \left( 2x + \frac{\pi}{3} \right)$ .

2. Express  $\sin x - \sqrt{3} \cos x$  in the form  $R \sin(x - \alpha)$ ,  
where  $R > 0$  and  $0 < \alpha < \pi/2$ .

Hence (i) sketch the graph of  $y = f(x) = \sin x - \sqrt{3} \cos x$ ;

(ii) find the range of  $f$ ;

(iii) find the period of  $f$ .



# Inverse Trigonometric Functions

## Question 4-N36-Q2

Express  $4 \sin x - 3 \cos x$  in the form  $r \sin(x - \alpha)$

A  $5 \sin(x + 45.00^\circ)$

B  $5 \sin(x - 30.00^\circ)$

C  $5 \sin(x + 36.87^\circ)$



# Suggested Reading

[Foundation Algebra](#) by P. Gajjar.

(Chapter 6)