

CELEN037 Seminar 2



University of
Nottingham
UK | CHINA | MALAYSIA



- Chain Rule for Differentiation
- The Fast-Track Chain Rule Method
- Logarithmic Differentiation
- Implicit Differentiation
- Derivatives of Inverse Functions



Chain Rule

$$\frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dv} \cdot \frac{dv}{dw} \cdot \frac{dw}{dx}$$

Chain Rule

$$\frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dv} \cdot \frac{dv}{dw} \cdot \frac{dw}{dx}$$

Example: Given $y = \ln(\sec x)$, find $\frac{dy}{dx}$ using Chain Rule

Chain Rule

$$\frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dv} \cdot \frac{dv}{dw} \cdot \frac{dw}{dx}$$

Example: Given $y = \ln(\sec x)$, find $\frac{dy}{dx}$ using Chain Rule

Solution:

Chain Rule

$$\frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dv} \cdot \frac{dv}{dw} \cdot \frac{dw}{dx}$$

Example: Given $y = \ln(\sec x)$, find $\frac{dy}{dx}$ using Chain Rule

Solution:

Let $u = \sec x$, so that $y = \ln u$.

Chain Rule

$$\frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dv} \cdot \frac{dv}{dw} \cdot \frac{dw}{dx}$$

Example: Given $y = \ln(\sec x)$, find $\frac{dy}{dx}$ using Chain Rule

Solution:

Let $u = \sec x$, so that $y = \ln u$.

Then $\frac{du}{dx} = \sec x \cdot \tan x$, and $\frac{dy}{du} = \frac{1}{u}$.

Chain Rule

$$\frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dv} \cdot \frac{dv}{dw} \cdot \frac{dw}{dx}$$

Example: Given $y = \ln(\sec x)$, find $\frac{dy}{dx}$ using Chain Rule

Solution:

Let $u = \sec x$, so that $y = \ln u$.

Then $\frac{du}{dx} = \sec x \cdot \tan x$, and $\frac{dy}{du} = \frac{1}{u}$.

Hence $\frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dx} = \frac{1}{u} \cdot \sec x \cdot \tan x$

Chain Rule

$$\frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dv} \cdot \frac{dv}{dw} \cdot \frac{dw}{dx}$$

Example: Given $y = \ln(\sec x)$, find $\frac{dy}{dx}$ using Chain Rule

Solution:

Let $u = \sec x$, so that $y = \ln u$.

Then $\frac{du}{dx} = \sec x \cdot \tan x$, and $\frac{dy}{du} = \frac{1}{u}$.

Hence
$$\begin{aligned}\frac{dy}{dx} &= \frac{dy}{du} \cdot \frac{du}{dx} = \frac{1}{u} \cdot \sec x \cdot \tan x \\ &= \frac{1}{\sec x} \cdot \sec x \cdot \tan x\end{aligned}$$

Chain Rule

$$\frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dv} \cdot \frac{dv}{dw} \cdot \frac{dw}{dx}$$

Example: Given $y = \ln(\sec x)$, find $\frac{dy}{dx}$ using Chain Rule

Solution:

Let $u = \sec x$, so that $y = \ln u$.

Then $\frac{du}{dx} = \sec x \cdot \tan x$, and $\frac{dy}{du} = \frac{1}{u}$.

$$\begin{aligned}\text{Hence } \frac{dy}{dx} &= \frac{dy}{du} \cdot \frac{du}{dx} = \frac{1}{u} \cdot \sec x \cdot \tan x \\ &= \frac{1}{\sec x} \cdot \sec x \cdot \tan x \\ &= \tan x\end{aligned}$$



Practice Problems on Worksheet:

- 1: Q1(ii)
- 2: Q1(vii)
- 3: Q1(viii)
- 4: Q1(ix)

Practice Problems on Worksheet:

- 1: Q1(ii)
- 2: Q1(vii)
- 3: Q1(viii)
- 4: Q1(ix)

Answers:

- 1: $-\frac{\cos(\cos(\ln x)) \cdot \sin(\ln x)}{x}$
- 2: $-\tan x \cdot \cos(\ln(\cos x))$
- 3: $-\frac{\sec^2(\cos(\sqrt{x})) \cdot \sin(\sqrt{x})}{2\sqrt{x}}$
- 4: $e^x \cdot \cot(e^x)$



Chain Rule

$$\frac{d}{dx}[f(u)] = f'(u) \cdot \frac{du}{dx}$$

Chain Rule

$$\frac{d}{dx}[f(u)] = f'(u) \cdot \frac{du}{dx}$$

Example: Given $y = \sqrt{\sin(e^{\cos x})}$, find $\frac{dy}{dx}$ using Chain Rule

Chain Rule

$$\frac{d}{dx}[f(u)] = f'(u) \cdot \frac{du}{dx}$$

Example: Given $y = \sqrt{\sin(e^{\cos x})}$, find $\frac{dy}{dx}$ using Chain Rule

Solution:

Chain Rule

$$\frac{d}{dx}[f(u)] = f'(u) \cdot \frac{du}{dx}$$

Example: Given $y = \sqrt{\sin(e^{\cos x})}$, find $\frac{dy}{dx}$ using Chain Rule

Solution:

$$\frac{dy}{dx} = \frac{1}{2} \cdot \frac{1}{\sqrt{\sin(e^{\cos x})}} \cdot \frac{d}{dx} (\sin(e^{\cos x}))$$

Chain Rule

$$\frac{d}{dx}[f(u)] = f'(u) \cdot \frac{du}{dx}$$

Example: Given $y = \sqrt{\sin(e^{\cos x})}$, find $\frac{dy}{dx}$ using Chain Rule

Solution:

$$\begin{aligned}\frac{dy}{dx} &= \frac{1}{2} \cdot \frac{1}{\sqrt{\sin(e^{\cos x})}} \cdot \frac{d}{dx} (\sin(e^{\cos x})) \\ &= \frac{1}{2\sqrt{\sin(e^{\cos x})}} \cdot \cos(e^{\cos x}) \cdot \frac{d}{dx} (e^{\cos x})\end{aligned}$$

Chain Rule

$$\frac{d}{dx}[f(u)] = f'(u) \cdot \frac{du}{dx}$$

Example: Given $y = \sqrt{\sin(e^{\cos x})}$, find $\frac{dy}{dx}$ using Chain Rule

Solution:

$$\begin{aligned}\frac{dy}{dx} &= \frac{1}{2} \cdot \frac{1}{\sqrt{\sin(e^{\cos x})}} \cdot \frac{d}{dx} (\sin(e^{\cos x})) \\ &= \frac{1}{2\sqrt{\sin(e^{\cos x})}} \cdot \cos(e^{\cos x}) \cdot \frac{d}{dx} (e^{\cos x}) \\ &= \frac{\cos(e^{\cos x})}{2\sqrt{\sin(e^{\cos x})}} \cdot e^{\cos x} \cdot \frac{d}{dx} (\cos x)\end{aligned}$$

Chain Rule

$$\frac{d}{dx}[f(u)] = f'(u) \cdot \frac{du}{dx}$$

Example: Given $y = \sqrt{\sin(e^{\cos x})}$, find $\frac{dy}{dx}$ using Chain Rule

Solution:

$$\begin{aligned}\frac{dy}{dx} &= \frac{1}{2} \cdot \frac{1}{\sqrt{\sin(e^{\cos x})}} \cdot \frac{d}{dx}(\sin(e^{\cos x})) \\&= \frac{1}{2\sqrt{\sin(e^{\cos x})}} \cdot \cos(e^{\cos x}) \cdot \frac{d}{dx}(e^{\cos x}) \\&= \frac{\cos(e^{\cos x})}{2\sqrt{\sin(e^{\cos x})}} \cdot e^{\cos x} \cdot \frac{d}{dx}(\cos x) \\&= -\frac{\cos(e^{\cos x}) \cdot e^{\cos x} \cdot \sin x}{2\sqrt{\sin(e^{\cos x})}}\end{aligned}$$



Practice Problems on Worksheet:

1: Q2(ii)

2: Q2(iii)

3: Q2(iv)

4: Q2(v)

Practice Problems on Worksheet:

1: Q2(ii)

2: Q2(iii)

3: Q2(iv)

4: Q2(v)

Answers:

$$1: -\frac{\cos(\cos(\ln x)) \cdot \sin(\ln x)}{x}$$

$$2: -\tan x \cdot \cos(\ln(\cos x))$$

$$3: -\frac{\sec^2(\cos(\sqrt{x})) \cdot \sin(\sqrt{x})}{2\sqrt{x}}$$

$$4: e^x \cdot \cot(e^x)$$



Example: Given $y = (\tan x)^{\sin x}$, find $\frac{dy}{dx}$



Example: Given $y = (\tan x)^{\sin x}$, find $\frac{dy}{dx}$

Solution:



Example: Given $y = (\tan x)^{\sin x}$, find $\frac{dy}{dx}$

Solution:

$$\ln y = \ln ((\tan x)^{\sin x})$$



Example: Given $y = (\tan x)^{\sin x}$, find $\frac{dy}{dx}$

Solution:

$$\begin{aligned}\ln y &= \ln ((\tan x)^{\sin x}) \\ &= \sin x \cdot \ln(\tan x)\end{aligned}$$

Example: Given $y = (\tan x)^{\sin x}$, find $\frac{dy}{dx}$

Solution:

$$\begin{aligned}\ln y &= \ln ((\tan x)^{\sin x}) \\ &= \sin x \cdot \ln(\tan x)\end{aligned}$$

Then $\frac{d}{dx}(\ln y) = \frac{d}{dx}(\sin x \cdot \ln(\tan x))$

Example: Given $y = (\tan x)^{\sin x}$, find $\frac{dy}{dx}$

Solution:

$$\begin{aligned}\ln y &= \ln ((\tan x)^{\sin x}) \\ &= \sin x \cdot \ln(\tan x)\end{aligned}$$

Then $\frac{d}{dx}(\ln y) = \frac{d}{dx}(\sin x \cdot \ln(\tan x))$

$$\frac{1}{y} \cdot \frac{dy}{dx} = \sin x \cdot \frac{1}{\tan x} \cdot \sec^2 x + \ln(\tan x) \cdot \cos x$$

Example: Given $y = (\tan x)^{\sin x}$, find $\frac{dy}{dx}$

Solution:

$$\begin{aligned}\ln y &= \ln ((\tan x)^{\sin x}) \\ &= \sin x \cdot \ln(\tan x)\end{aligned}$$

Then $\frac{d}{dx}(\ln y) = \frac{d}{dx}(\sin x \cdot \ln(\tan x))$

$$\begin{aligned}\frac{1}{y} \cdot \frac{dy}{dx} &= \sin x \cdot \frac{1}{\tan x} \cdot \sec^2 x + \ln(\tan x) \cdot \cos x \\ &= \sec x + \ln(\tan x) \cdot \cos x\end{aligned}$$

Example: Given $y = (\tan x)^{\sin x}$, find $\frac{dy}{dx}$

Solution:

$$\begin{aligned}\ln y &= \ln ((\tan x)^{\sin x}) \\ &= \sin x \cdot \ln(\tan x)\end{aligned}$$

Then $\frac{d}{dx}(\ln y) = \frac{d}{dx}(\sin x \cdot \ln(\tan x))$

$$\begin{aligned}\frac{1}{y} \cdot \frac{dy}{dx} &= \sin x \cdot \frac{1}{\tan x} \cdot \sec^2 x + \ln(\tan x) \cdot \cos x \\ &= \sec x + \ln(\tan x) \cdot \cos x\end{aligned}$$

Hence $\frac{dy}{dx} = y \cdot (\sec x + \ln(\tan x) \cdot \cos x)$

Example: Given $y = (\tan x)^{\sin x}$, find $\frac{dy}{dx}$

Solution:

$$\begin{aligned}\ln y &= \ln ((\tan x)^{\sin x}) \\ &= \sin x \cdot \ln(\tan x)\end{aligned}$$

Then $\frac{d}{dx}(\ln y) = \frac{d}{dx}(\sin x \cdot \ln(\tan x))$

$$\begin{aligned}\frac{1}{y} \cdot \frac{dy}{dx} &= \sin x \cdot \frac{1}{\tan x} \cdot \sec^2 x + \ln(\tan x) \cdot \cos x \\ &= \sec x + \ln(\tan x) \cdot \cos x\end{aligned}$$

Hence $\frac{dy}{dx} = y \cdot (\sec x + \ln(\tan x) \cdot \cos x)$

$$= (\tan x)^{\sin x} \cdot (\sec x + \ln(\tan x) \cdot \cos x)$$

Example: Given $y = (\tan x)^{\sin x}$, find $\frac{dy}{dx}$

Solution:

$$\begin{aligned}\ln y &= \ln ((\tan x)^{\sin x}) \\ &= \sin x \cdot \ln(\tan x)\end{aligned}$$

Then $\frac{d}{dx}(\ln y) = \frac{d}{dx}(\sin x \cdot \ln(\tan x))$

$$\begin{aligned}\frac{1}{y} \cdot \frac{dy}{dx} &= \sin x \cdot \frac{1}{\tan x} \cdot \sec^2 x + \ln(\tan x) \cdot \cos x \\ &= \sec x + \ln(\tan x) \cdot \cos x\end{aligned}$$

Hence $\frac{dy}{dx} = y \cdot (\sec x + \ln(\tan x) \cdot \cos x)$

$$= (\tan x)^{\sin x} \cdot (\sec x + \ln(\tan x) \cdot \cos x)$$

Steps

⇐ 1. Take Logarithm on both sides

⇐ 2. Apply rules of logarithms

⇐ 3. Differentiate both sides w.r.t x



Practice Problems on Worksheet:

- 1: Q3(vii)
- 2: Q3(viii)
- 3: Q3(ix)
- 4: Q3(x)

Practice Problems on Worksheet:

1: Q3(vii)

2: Q3(viii)

3: Q3(ix)

4: Q3(x)

Answers:

1: $x^x(\ln x + 1)$

2: $\frac{\sqrt[x]{x}(1 - \ln x)}{x^2}$

3: $\cos(x^x) x^x(\ln x + 1)$

4: $\frac{\sqrt[3]{x} \tan^4 x}{\cos(e^x)} \left(\frac{1}{3x} + \frac{4 \sec^2 x}{\tan x} + \tan(e^x) \cdot e^x \right)$



Example: Given $\ln(x + y) = \ln(xy) + 1$, find $\frac{dy}{dx}$



Example: Given $\ln(x + y) = \ln(xy) + 1$, find $\frac{dy}{dx}$

Solution: Differentiate both sides w.r.t. x :

Example: Given $\ln(x + y) = \ln(xy) + 1$, find $\frac{dy}{dx}$

Solution: Differentiate both sides w.r.t. x :

$$\frac{1}{x + y} \left(1 + \frac{dy}{dx} \right) = \frac{1}{xy} \left(y + x \cdot \frac{dy}{dx} \right) + 0$$

Example: Given $\ln(x + y) = \ln(xy) + 1$, find $\frac{dy}{dx}$

Solution: Differentiate both sides w.r.t. x :

$$\frac{1}{x + y} \left(1 + \frac{dy}{dx} \right) = \frac{1}{xy} \left(y + x \cdot \frac{dy}{dx} \right) + 0$$

$$\frac{dy}{dx} \left(\frac{1}{x + y} - \frac{1}{y} \right) = \frac{1}{x} - \frac{1}{x + y}$$

Example: Given $\ln(x + y) = \ln(xy) + 1$, find $\frac{dy}{dx}$

Solution: Differentiate both sides w.r.t. x :

$$\frac{1}{x + y} \left(1 + \frac{dy}{dx} \right) = \frac{1}{xy} \left(y + x \cdot \frac{dy}{dx} \right) + 0$$

$$\frac{dy}{dx} \left(\frac{1}{x + y} - \frac{1}{y} \right) = \frac{1}{x} - \frac{1}{x + y}$$

$$\frac{dy}{dx} \cdot \frac{-x}{y(x + y)} = \frac{y}{x(x + y)}$$

Example: Given $\ln(x + y) = \ln(xy) + 1$, find $\frac{dy}{dx}$

Solution: Differentiate both sides w.r.t. x :

$$\frac{1}{x + y} \left(1 + \frac{dy}{dx} \right) = \frac{1}{xy} \left(y + x \cdot \frac{dy}{dx} \right) + 0$$

$$\frac{dy}{dx} \left(\frac{1}{x + y} - \frac{1}{y} \right) = \frac{1}{x} - \frac{1}{x + y}$$

$$\frac{dy}{dx} \cdot \frac{-x}{y(x + y)} = \frac{y}{x(x + y)}$$

$$\Rightarrow \frac{dy}{dx} = -\frac{y^2}{x^2}$$



Practice Problems on Worksheet:

- 1: Q4(vii)
- 2: Q4(viii)
- 3: Q4(ix)
- 4: Q4(x)

Practice Problems on Worksheet:

1: Q4(vii)

2: Q4(viii)

3: Q4(ix)

4: Q4(x)

Answers:

$$1: \frac{dy}{dx} = \frac{-x-2xy+2y^2}{x^2-4xy+y}, \quad \left. \frac{dy}{dx} \right|_{(1,1)} = \frac{1}{2}$$

$$2: \frac{dy}{dx} = 1, \quad \left. \frac{dy}{dx} \right|_{(0,0)} = 1$$

$$3: \frac{dy}{dx} = \frac{3x^2-2xy-y^2}{x^2+2xy-3y^2}, \quad \left. \frac{dy}{dx} \right|_{(1,-1)} = -1$$

$$4: \frac{dy}{dx} = \frac{-2xy^3+5y^2+3y-2}{3x^2y^2-3x-10xy}, \quad \left. \frac{dy}{dx} \right|_{(2,1)} = -\frac{1}{7}$$



Derivatives of Inverse Trigonometric Functions

$$\frac{d}{dx} (\sin^{-1} x) = \frac{1}{\sqrt{1-x^2}}, \quad \frac{d}{dx} (\cos^{-1} x) = \frac{-1}{\sqrt{1-x^2}}, \quad \frac{d}{dx} (\tan^{-1} x) = \frac{1}{1+x^2}$$

Derivatives of Inverse Trigonometric Functions

$$\frac{d}{dx} (\sin^{-1} x) = \frac{1}{\sqrt{1-x^2}}, \quad \frac{d}{dx} (\cos^{-1} x) = \frac{-1}{\sqrt{1-x^2}}, \quad \frac{d}{dx} (\tan^{-1} x) = \frac{1}{1+x^2}$$

Example: Given $y = \sec^{-1}(\sqrt{1+x^2}) = \cos^{-1}\left(\frac{1}{\sqrt{1+x^2}}\right)$, $x > 0$, find $\frac{dy}{dx}$

Derivatives of Inverse Trigonometric Functions

$$\frac{d}{dx} (\sin^{-1} x) = \frac{1}{\sqrt{1-x^2}}, \quad \frac{d}{dx} (\cos^{-1} x) = \frac{-1}{\sqrt{1-x^2}}, \quad \frac{d}{dx} (\tan^{-1} x) = \frac{1}{1+x^2}$$

Example: Given $y = \sec^{-1}(\sqrt{1+x^2}) = \cos^{-1}\left(\frac{1}{\sqrt{1+x^2}}\right)$, $x > 0$, find $\frac{dy}{dx}$

Solution: Using $\frac{d}{dx} (\cos^{-1} x) = \frac{-1}{\sqrt{1-x^2}}$, we can get

Derivatives of Inverse Trigonometric Functions

$$\frac{d}{dx} (\sin^{-1} x) = \frac{1}{\sqrt{1-x^2}}, \quad \frac{d}{dx} (\cos^{-1} x) = \frac{-1}{\sqrt{1-x^2}}, \quad \frac{d}{dx} (\tan^{-1} x) = \frac{1}{1+x^2}$$

Example: Given $y = \sec^{-1}(\sqrt{1+x^2}) = \cos^{-1}\left(\frac{1}{\sqrt{1+x^2}}\right)$, $x > 0$, find $\frac{dy}{dx}$

Solution: Using $\frac{d}{dx} (\cos^{-1} x) = \frac{-1}{\sqrt{1-x^2}}$, we can get

$$\frac{dy}{dx} = \frac{-1}{\sqrt{1 - \left(\frac{1}{\sqrt{1+x^2}}\right)^2}} \cdot \frac{d}{dx} \left(\frac{1}{\sqrt{1+x^2}}\right)$$

Derivatives of Inverse Trigonometric Functions

$$\frac{d}{dx} (\sin^{-1} x) = \frac{1}{\sqrt{1-x^2}}, \quad \frac{d}{dx} (\cos^{-1} x) = \frac{-1}{\sqrt{1-x^2}}, \quad \frac{d}{dx} (\tan^{-1} x) = \frac{1}{1+x^2}$$

Example: Given $y = \sec^{-1}(\sqrt{1+x^2}) = \cos^{-1}\left(\frac{1}{\sqrt{1+x^2}}\right)$, $x > 0$, find $\frac{dy}{dx}$

Solution: Using $\frac{d}{dx} (\cos^{-1} x) = \frac{-1}{\sqrt{1-x^2}}$, we can get

$$\begin{aligned} \frac{dy}{dx} &= \frac{-1}{\sqrt{1 - \left(\frac{1}{\sqrt{1+x^2}}\right)^2}} \cdot \frac{d}{dx} \left(\frac{1}{\sqrt{1+x^2}} \right) \\ &= \frac{-1}{\sqrt{1 - \frac{1}{1+x^2}}} \cdot \left(-\frac{1}{2} \right) \frac{1}{\sqrt{(1+x^2)^3}} \cdot 2x \end{aligned}$$

Derivatives of Inverse Trigonometric Functions

$$\frac{d}{dx} (\sin^{-1} x) = \frac{1}{\sqrt{1-x^2}}, \quad \frac{d}{dx} (\cos^{-1} x) = \frac{-1}{\sqrt{1-x^2}}, \quad \frac{d}{dx} (\tan^{-1} x) = \frac{1}{1+x^2}$$

Example: Given $y = \sec^{-1}(\sqrt{1+x^2}) = \cos^{-1}\left(\frac{1}{\sqrt{1+x^2}}\right)$, $x > 0$, find $\frac{dy}{dx}$

Solution: Using $\frac{d}{dx} (\cos^{-1} x) = \frac{-1}{\sqrt{1-x^2}}$, we can get

$$\begin{aligned} \frac{dy}{dx} &= \frac{-1}{\sqrt{1 - \left(\frac{1}{\sqrt{1+x^2}}\right)^2}} \cdot \frac{d}{dx} \left(\frac{1}{\sqrt{1+x^2}} \right) \\ &= \frac{-1}{\sqrt{1 - \frac{1}{1+x^2}}} \cdot \left(-\frac{1}{2} \right) \frac{1}{\sqrt{(1+x^2)^3}} \cdot 2x \\ &= \frac{x}{\sqrt{\frac{x^2}{1+x^2}} \cdot \sqrt{(1+x^2)^3}} \end{aligned}$$

Derivatives of Inverse Trigonometric Functions

$$\frac{d}{dx} (\sin^{-1} x) = \frac{1}{\sqrt{1-x^2}}, \quad \frac{d}{dx} (\cos^{-1} x) = \frac{-1}{\sqrt{1-x^2}}, \quad \frac{d}{dx} (\tan^{-1} x) = \frac{1}{1+x^2}$$

Example: Given $y = \sec^{-1}(\sqrt{1+x^2}) = \cos^{-1}\left(\frac{1}{\sqrt{1+x^2}}\right)$, $x > 0$, find $\frac{dy}{dx}$

Solution: Using $\frac{d}{dx} (\cos^{-1} x) = \frac{-1}{\sqrt{1-x^2}}$, we can get

$$\begin{aligned} \frac{dy}{dx} &= \frac{-1}{\sqrt{1 - \left(\frac{1}{\sqrt{1+x^2}}\right)^2}} \cdot \frac{d}{dx} \left(\frac{1}{\sqrt{1+x^2}} \right) \\ &= \frac{-1}{\sqrt{1 - \frac{1}{1+x^2}}} \cdot \left(-\frac{1}{2} \right) \frac{1}{\sqrt{(1+x^2)^3}} \cdot 2x \end{aligned}$$

$$\begin{aligned} &= \frac{x}{\sqrt{\frac{x^2}{1+x^2}} \cdot \sqrt{(1+x^2)^3}} \\ &= \frac{x}{\underbrace{\sqrt{x^2(1+x^2)^2}}_{x>0}} = \frac{x}{x(1+x^2)} \end{aligned}$$

Derivatives of Inverse Trigonometric Functions

$$\frac{d}{dx} (\sin^{-1} x) = \frac{1}{\sqrt{1-x^2}}, \quad \frac{d}{dx} (\cos^{-1} x) = \frac{-1}{\sqrt{1-x^2}}, \quad \frac{d}{dx} (\tan^{-1} x) = \frac{1}{1+x^2}$$

Example: Given $y = \sec^{-1}(\sqrt{1+x^2}) = \cos^{-1}\left(\frac{1}{\sqrt{1+x^2}}\right)$, $x > 0$, find $\frac{dy}{dx}$

Solution: Using $\frac{d}{dx} (\cos^{-1} x) = \frac{-1}{\sqrt{1-x^2}}$, we can get

$$\begin{aligned} \frac{dy}{dx} &= \frac{-1}{\sqrt{1 - \left(\frac{1}{\sqrt{1+x^2}}\right)^2}} \cdot \frac{d}{dx} \left(\frac{1}{\sqrt{1+x^2}} \right) \\ &= \frac{-1}{\sqrt{1 - \frac{1}{1+x^2}}} \cdot \left(-\frac{1}{2} \right) \frac{1}{\sqrt{(1+x^2)^3}} \cdot 2x \end{aligned}$$

$$\begin{aligned} &= \frac{x}{\sqrt{\frac{x^2}{1+x^2}} \cdot \sqrt{(1+x^2)^3}} \\ &= \frac{x}{\underbrace{\sqrt{x^2(1+x^2)^2}}_{x>0}} = \frac{x}{x(1+x^2)} \\ &= \frac{1}{1+x^2} \end{aligned}$$



Practice Problems on Worksheet:

1: Q5(iii)

2: Q5(iv)

3: Q5(v)

4: Q5(vi)

Practice Problems on Worksheet:

1: Q5(iii)

2: Q5(iv)

3: Q5(v)

4: Q5(vi)

Answers:

1: $\frac{1}{2(1+x^2)}$

2: $\frac{1}{2}$

3: $\frac{2}{1+x^2}$

4: $\frac{2(1-x^2)}{x^4+6x^2+1}$



Day	Time	Venue
Tuesday	12:00 to 14:00	Trent 314a
	13:00 to 14:00	Trent 322
Thursday	16:30 to 17:30	TB 417
	17:00 to 18:00	IAMET 315
Friday	14:00 to 15:00	PB 330
	17:00 to 18:00	TB 417