

CELEN037 Seminar 3



University of
Nottingham
UK | CHINA | MALAYSIA



- Parametric Differentiation
- Maclaurin's Series
- Equations of Tangent and Normal Lines
- Newton-Raphson Method
- Increasing and Decreasing Functions



Example: Given $x = a(t - \sin t)$ and $y = a(1 - \cos t)$ (a is a constant),
find $\left. \frac{dy}{dx} \right|_{t=\frac{\pi}{2}}$.



Example: Given $x = a(t - \sin t)$ and $y = a(1 - \cos t)$ (a is a constant),
find $\left. \frac{dy}{dx} \right|_{t=\frac{\pi}{2}}$.

Solution:



Example: Given $x = a(t - \sin t)$ and $y = a(1 - \cos t)$ (a is a constant),
find $\left. \frac{dy}{dx} \right|_{t=\frac{\pi}{2}}$.

Solution:

$$\frac{dx}{dt} = a(1 - \cos t), \quad \frac{dy}{dt} = a \sin t$$

Example: Given $x = a(t - \sin t)$ and $y = a(1 - \cos t)$ (a is a constant),
find $\left. \frac{dy}{dx} \right|_{t=\frac{\pi}{2}}$.

Solution:

$$\frac{dx}{dt} = a(1 - \cos t), \quad \frac{dy}{dt} = a \sin t$$

$$\frac{dy}{dx} = \frac{\frac{dy}{dt}}{\frac{dx}{dt}} = \frac{a \sin t}{a(1 - \cos t)}$$

Example: Given $x = a(t - \sin t)$ and $y = a(1 - \cos t)$ (a is a constant),
find $\left. \frac{dy}{dx} \right|_{t=\frac{\pi}{2}}$.

Solution:

$$\frac{dx}{dt} = a(1 - \cos t), \quad \frac{dy}{dt} = a \sin t$$

$$\frac{dy}{dx} = \frac{\frac{dy}{dt}}{\frac{dx}{dt}} = \frac{a \sin t}{a(1 - \cos t)} = \frac{\sin t}{1 - \cos t}$$

Example: Given $x = a(t - \sin t)$ and $y = a(1 - \cos t)$ (a is a constant),
find $\left. \frac{dy}{dx} \right|_{t=\frac{\pi}{2}}$.

Solution:

$$\frac{dx}{dt} = a(1 - \cos t), \quad \frac{dy}{dt} = a \sin t$$

$$\frac{dy}{dx} = \frac{\frac{dy}{dt}}{\frac{dx}{dt}} = \frac{a \sin t}{a(1 - \cos t)} = \frac{\sin t}{1 - \cos t}$$

$$\left. \frac{dy}{dx} \right|_{t=\frac{\pi}{2}} = \frac{\sin \frac{\pi}{2}}{1 - \cos \frac{\pi}{2}}$$

Example: Given $x = a(t - \sin t)$ and $y = a(1 - \cos t)$ (a is a constant),
find $\left. \frac{dy}{dx} \right|_{t=\frac{\pi}{2}}$.

Solution:

$$\frac{dx}{dt} = a(1 - \cos t), \quad \frac{dy}{dt} = a \sin t$$

$$\frac{dy}{dx} = \frac{\frac{dy}{dt}}{\frac{dx}{dt}} = \frac{a \sin t}{a(1 - \cos t)} = \frac{\sin t}{1 - \cos t}$$

$$\begin{aligned} \left. \frac{dy}{dx} \right|_{t=\frac{\pi}{2}} &= \frac{\sin \frac{\pi}{2}}{1 - \cos \frac{\pi}{2}} \\ &= \frac{1}{1 - 0} \\ &= 1 \end{aligned}$$



Practice Problems on Worksheet:

- 1: Q1(ii)
- 2: Q1(iv)
- 3: Q1(v)
- 4: Q1(vi)

Practice Problems on Worksheet:

1: Q1(ii)

2: Q1(iv)

3: Q1(v)

4: Q1(vi)

Answers:

1: -1

2: -1

3: $\frac{v_0 \sin \alpha - gt}{v_0 \cos \alpha}$

4: $-\frac{2t}{1 - t^2}$



Example: Obtain the Maclaurin's Series expansions of the following function:

$$f(x) = \frac{1}{1-x}; \quad -1 < x < 1$$



Example: Obtain the Maclaurin's Series expansions of the following function:

$$f(x) = \frac{1}{1-x}; \quad -1 < x < 1$$

Solution:



Example: Obtain the Maclaurin's Series expansions of the following function:

$$f(x) = \frac{1}{1-x}; \quad -1 < x < 1$$

Solution:

$$f(0) = 1$$

Example: Obtain the Maclaurin's Series expansions of the following function:

$$f(x) = \frac{1}{1-x}; \quad -1 < x < 1$$

Solution:

$$f(0) = 1$$

$$f'(x) = \frac{-1}{(1-x)^2} \cdot (-1) = \frac{1}{(1-x)^2} \implies f'(0) = 1$$

Example: Obtain the Maclaurin's Series expansions of the following function:

$$f(x) = \frac{1}{1-x}; \quad -1 < x < 1$$

Solution:

$$f(0) = 1$$

$$f'(x) = \frac{-1}{(1-x)^2} \cdot (-1) = \frac{1}{(1-x)^2} \implies f'(0) = 1$$

$$f''(x) = \frac{-2}{(1-x)^3} \cdot (-1) = \frac{2}{(1-x)^3} \implies f''(0) = 2$$

Example: Obtain the Maclaurin's Series expansions of the following function:

$$f(x) = \frac{1}{1-x}; \quad -1 < x < 1$$

Solution:

$$f(0) = 1$$

$$f'(x) = \frac{-1}{(1-x)^2} \cdot (-1) = \frac{1}{(1-x)^2} \implies f'(0) = 1$$

$$f''(x) = \frac{-2}{(1-x)^3} \cdot (-1) = \frac{2}{(1-x)^3} \implies f''(0) = 2$$

$$f'''(x) = \frac{-6}{(1-x)^4} \cdot (-1) = \frac{6}{(1-x)^4} \implies f'''(0) = 6$$

Example: Obtain the Maclaurin's Series expansions of the following function:

$$f(x) = \frac{1}{1-x}; \quad -1 < x < 1$$

Solution:

$$f(0) = 1$$

$$f'(x) = \frac{-1}{(1-x)^2} \cdot (-1) = \frac{1}{(1-x)^2} \implies f'(0) = 1$$

$$f''(x) = \frac{-2}{(1-x)^3} \cdot (-1) = \frac{2}{(1-x)^3} \implies f''(0) = 2$$

$$f'''(x) = \frac{-6}{(1-x)^4} \cdot (-1) = \frac{6}{(1-x)^4} \implies f'''(0) = 6$$

$$f^{(4)}(x) = \frac{-24}{(1-x)^5} \cdot (-1) = \frac{24}{(1-x)^5} \implies f^{(4)}(0) = 24$$

Example: Obtain the Maclaurin's Series expansions of the following function:

$$f(x) = \frac{1}{1-x}; \quad -1 < x < 1$$

Solution:

$$f(0) = 1$$

$$f'(x) = \frac{-1}{(1-x)^2} \cdot (-1) = \frac{1}{(1-x)^2} \implies f'(0) = 1$$

$$f''(x) = \frac{-2}{(1-x)^3} \cdot (-1) = \frac{2}{(1-x)^3} \implies f''(0) = 2$$

$$f'''(x) = \frac{-6}{(1-x)^4} \cdot (-1) = \frac{6}{(1-x)^4} \implies f'''(0) = 6$$

$$f^{(4)}(x) = \frac{-24}{(1-x)^5} \cdot (-1) = \frac{24}{(1-x)^5} \implies f^{(4)}(0) = 24$$

$$\implies \frac{1}{1-x} = 1 + x \cdot (1) + \frac{x^2}{2!} \cdot (2) + \frac{x^3}{3!} \cdot (6) + \frac{x^4}{4!} \cdot (24) + \dots$$

Example: Obtain the Maclaurin's Series expansions of the following function:

$$f(x) = \frac{1}{1-x}; \quad -1 < x < 1$$

Solution:

$$f(0) = 1$$

$$f'(x) = \frac{-1}{(1-x)^2} \cdot (-1) = \frac{1}{(1-x)^2} \implies f'(0) = 1$$

$$f''(x) = \frac{-2}{(1-x)^3} \cdot (-1) = \frac{2}{(1-x)^3} \implies f''(0) = 2$$

$$f'''(x) = \frac{-6}{(1-x)^4} \cdot (-1) = \frac{6}{(1-x)^4} \implies f'''(0) = 6$$

$$f^{(4)}(x) = \frac{-24}{(1-x)^5} \cdot (-1) = \frac{24}{(1-x)^5} \implies f^{(4)}(0) = 24$$

$$\begin{aligned} \implies \frac{1}{1-x} &= 1 + x \cdot (1) + \frac{x^2}{2!} \cdot (2) + \frac{x^3}{3!} \cdot (6) + \frac{x^4}{4!} \cdot (24) + \dots \\ &= 1 + x + x^2 + x^3 + x^4 + \dots \end{aligned}$$

Practice Problems on Workbook:

- 1: Q2(i)
- 2: Q2(iii)
- 3: Q2(vi)
- 4: Q2(vii)

Practice Problems on Workbook:

1: Q2(i)

2: Q2(iii)

3: Q2(vi)

4: Q2(vii)

Answers:

$$1: 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \dots$$

$$2: 1 - \frac{x}{2} - \frac{x^2}{8} - \frac{x^3}{16} + \dots$$

$$3: e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots, e^{-x} = 1 - x + \frac{x^2}{2!} - \frac{x^3}{3!} + \dots$$

$$4: \ln(1+x) = x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \dots, \ln(1-x) = -x - \frac{x^2}{2} - \frac{x^3}{3} - \frac{x^4}{4} - \dots$$



Example: The equation of the curve is given by $y = \frac{1}{x}$. Obtain the equation of the tangent line and the normal line to the curve at point $(-1, -1)$.



Example: The equation of the curve is given by $y = \frac{1}{x}$. Obtain the equation of the tangent line and the normal line to the curve at point $(-1, -1)$.

Solution:



Example: The equation of the curve is given by $y = \frac{1}{x}$. Obtain the equation of the tangent line and the normal line to the curve at point $(-1, -1)$.

Solution:

$$\frac{dy}{dx} = -\frac{1}{x^2}$$



Example: The equation of the curve is given by $y = \frac{1}{x}$. Obtain the equation of the tangent line and the normal line to the curve at point $(-1, -1)$.

Solution:

$$\frac{dy}{dx} = -\frac{1}{x^2}$$

$$\left. \frac{dy}{dx} \right|_{x=-1} = -\frac{1}{(-1)^2} = -\frac{1}{1} = -1 = m$$



Example: The equation of the curve is given by $y = \frac{1}{x}$. Obtain the equation of the tangent line and the normal line to the curve at point $(-1, -1)$.

Solution:

$$\frac{dy}{dx} = -\frac{1}{x^2}$$

$$\left. \frac{dy}{dx} \right|_{x=-1} = -\frac{1}{(-1)^2} = -\frac{1}{1} = -1 = m$$

Equation of tangent line: $y - y_1 = m(x - x_1)$

Example: The equation of the curve is given by $y = \frac{1}{x}$. Obtain the equation of the tangent line and the normal line to the curve at point $(-1, -1)$.

Solution:

$$\frac{dy}{dx} = -\frac{1}{x^2}$$

$$\left. \frac{dy}{dx} \right|_{x=-1} = -\frac{1}{(-1)^2} = -\frac{1}{1} = -1 = m$$

Equation of tangent line: $y - y_1 = m(x - x_1)$

$$y + x + 2 = 0$$

Example: The equation of the curve is given by $y = \frac{1}{x}$. Obtain the equation of the tangent line and the normal line to the curve at point $(-1, -1)$.

Solution:

$$\frac{dy}{dx} = -\frac{1}{x^2}$$

$$\left. \frac{dy}{dx} \right|_{x=-1} = -\frac{1}{(-1)^2} = -\frac{1}{1} = -1 = m$$

Equation of tangent line: $y - y_1 = m(x - x_1)$

$$y + x + 2 = 0$$

Equation of normal line: $y - y_1 = -\frac{1}{m}(x - x_1)$

Example: The equation of the curve is given by $y = \frac{1}{x}$. Obtain the equation of the tangent line and the normal line to the curve at point $(-1, -1)$.

Solution:

$$\frac{dy}{dx} = -\frac{1}{x^2}$$

$$\left. \frac{dy}{dx} \right|_{x=-1} = -\frac{1}{(-1)^2} = -\frac{1}{1} = -1 = m$$

Equation of tangent line: $y - y_1 = m(x - x_1)$

$$y + x + 2 = 0$$

Equation of normal line: $y - y_1 = -\frac{1}{m}(x - x_1)$

$$y = x$$



Practice Problems on Workbook:

1: Q3(ii)

2: Q3(iii)

Practice Problems on Workbook:

1: Q3(ii)

2: Q3(iii)

Answers:

1: Tangent line $y - 2x + 3 = 0$

Normal line $2y + x + 1 = 0$

2: Tangent line $2y + 3x - 6 = 0$

Normal line $3y - 2x - 9 = 0$



$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)} \quad (n = 0, 1, 2, 3, \dots)$$



$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)} \quad (n = 0, 1, 2, 3, \dots)$$

Example: Use the Newton-Raphson method to approximate the root of $x^4 - \sin x = 1$, correct to 7 d.p., by starting with $x_0 = 1.5$.



$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)} \quad (n = 0, 1, 2, 3, \dots)$$

Example: Use the Newton-Raphson method to approximate the root of $x^4 - \sin x = 1$, correct to 7 d.p., by starting with $x_0 = 1.5$.

Solution:



$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)} \quad (n = 0, 1, 2, 3, \dots)$$

Example: Use the Newton-Raphson method to approximate the root of $x^4 - \sin x = 1$, correct to 7 d.p., by starting with $x_0 = 1.5$.

Solution:

$$f(x) = x^4 - \sin x - 1; \quad f'(x) = 4x^3 - \cos x;$$



$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)} \quad (n = 0, 1, 2, 3, \dots)$$

Example: Use the Newton-Raphson method to approximate the root of $x^4 - \sin x = 1$, correct to 7 d.p., by starting with $x_0 = 1.5$.

Solution:

$$f(x) = x^4 - \sin x - 1; \quad f'(x) = 4x^3 - \cos x; \quad x_{n+1} = x_n - \frac{x_n^4 - \sin x_n - 1}{4x_n^3 - \cos x_n}$$

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)} \quad (n = 0, 1, 2, 3, \dots)$$

Example: Use the Newton-Raphson method to approximate the root of $x^4 - \sin x = 1$, correct to 7 d.p., by starting with $x_0 = 1.5$.

Solution:

$$f(x) = x^4 - \sin x - 1; \quad f'(x) = 4x^3 - \cos x; \quad x_{n+1} = x_n - \frac{x_n^4 - \sin x_n - 1}{4x_n^3 - \cos x_n}$$

Steps on calculator

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)} \quad (n = 0, 1, 2, 3, \dots)$$

Example: Use the Newton-Raphson method to approximate the root of $x^4 - \sin x = 1$, correct to 7 d.p., by starting with $x_0 = 1.5$.

Solution:

$$f(x) = x^4 - \sin x - 1; \quad f'(x) = 4x^3 - \cos x; \quad x_{n+1} = x_n - \frac{x_n^4 - \sin x_n - 1}{4x_n^3 - \cos x_n}$$

Steps on calculator

- (i) Set calculator to RADIAN mode:

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)} \quad (n = 0, 1, 2, 3, \dots)$$

Example: Use the Newton-Raphson method to approximate the root of $x^4 - \sin x = 1$, correct to 7 d.p., by starting with $x_0 = 1.5$.

Solution:

$$f(x) = x^4 - \sin x - 1; \quad f'(x) = 4x^3 - \cos x; \quad x_{n+1} = x_n - \frac{x_n^4 - \sin x_n - 1}{4x_n^3 - \cos x_n}$$

Steps on calculator

- (i) Set calculator to RADIAN
mode: **Shift** **Mode** **4**

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)} \quad (n = 0, 1, 2, 3, \dots)$$

Example: Use the Newton-Raphson method to approximate the root of $x^4 - \sin x = 1$, correct to 7 d.p., by starting with $x_0 = 1.5$.

Solution:

$$f(x) = x^4 - \sin x - 1; \quad f'(x) = 4x^3 - \cos x; \quad x_{n+1} = x_n - \frac{x_n^4 - \sin x_n - 1}{4x_n^3 - \cos x_n}$$

Steps on calculator

- (i) Set calculator to RADIAN
mode: **Shift Mode 4**
- (ii) Fix calculator to **7** digits:

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)} \quad (n = 0, 1, 2, 3, \dots)$$

Example: Use the Newton-Raphson method to approximate the root of $x^4 - \sin x = 1$, correct to 7 d.p., by starting with $x_0 = 1.5$.

Solution:

$$f(x) = x^4 - \sin x - 1; \quad f'(x) = 4x^3 - \cos x; \quad x_{n+1} = x_n - \frac{x_n^4 - \sin x_n - 1}{4x_n^3 - \cos x_n}$$

Steps on calculator

- (i) Set calculator to RADIAN
mode: **Shift Mode 4**
- (ii) Fix calculator to 7 digits:
Shift Mode 6 7

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)} \quad (n = 0, 1, 2, 3, \dots)$$

Example: Use the Newton-Raphson method to approximate the root of $x^4 - \sin x = 1$, correct to 7 d.p., by starting with $x_0 = 1.5$.

Solution:

$$f(x) = x^4 - \sin x - 1; \quad f'(x) = 4x^3 - \cos x; \quad x_{n+1} = x_n - \frac{x_n^4 - \sin x_n - 1}{4x_n^3 - \cos x_n}$$

Steps on calculator

- (i) Set calculator to RADIAN
mode: **Shift Mode 4**
- (ii) Fix calculator to 7 digits:
Shift Mode 6 7
- (iii) Press **1.5** and **=** on
calculator

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)} \quad (n = 0, 1, 2, 3, \dots)$$

Example: Use the Newton-Raphson method to approximate the root of $x^4 - \sin x = 1$, correct to 7 d.p., by starting with $x_0 = 1.5$.

Solution:

$$f(x) = x^4 - \sin x - 1; \quad f'(x) = 4x^3 - \cos x; \quad x_{n+1} = x_n - \frac{x_n^4 - \sin x_n - 1}{4x_n^3 - \cos x_n}$$

Steps on calculator

- (i) Set calculator to RADIAN
mode: **Shift Mode 4**
- (ii) Fix calculator to 7 digits:
Shift Mode 6 7
- (iii) Press **1.5** and $\boxed{=}$ on
calculator ($x_0 = 1.5$)

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)} \quad (n = 0, 1, 2, 3, \dots)$$

Example: Use the Newton-Raphson method to approximate the root of $x^4 - \sin x = 1$, correct to 7 d.p., by starting with $x_0 = 1.5$.

Solution:

$$f(x) = x^4 - \sin x - 1; \quad f'(x) = 4x^3 - \cos x; \quad x_{n+1} = x_n - \frac{x_n^4 - \sin x_n - 1}{4x_n^3 - \cos x_n}$$

Steps on calculator

- (i) Set calculator to Radian mode: **Shift Mode 4**
- (ii) Fix calculator to 7 digits: **Shift Mode 6 7**
- (iii) Press **1.5** and **=** on calculator ($x_0 = 1.5$)
- (iv) Enter the formula on calculator as:

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)} \quad (n = 0, 1, 2, 3, \dots)$$

Example: Use the Newton-Raphson method to approximate the root of $x^4 - \sin x = 1$, correct to 7 d.p., by starting with $x_0 = 1.5$.

Solution:

$$f(x) = x^4 - \sin x - 1; \quad f'(x) = 4x^3 - \cos x; \quad x_{n+1} = x_n - \frac{x_n^4 - \sin x_n - 1}{4x_n^3 - \cos x_n}$$

Steps on calculator

- (i) Set calculator to RADIAN mode: **Shift Mode 4**
- (ii) Fix calculator to 7 digits: **Shift Mode 6 7**
- (iii) Press **1.5** and **=** on calculator ($x_0 = 1.5$)
- (iv) Enter the formula on calculator as:
$$\mathbf{Ans} - \frac{(\mathbf{Ans}^4 - \sin(\mathbf{Ans}) - 1)}{(4 \times \mathbf{Ans}^3 - \cos(\mathbf{Ans}))}$$

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)} \quad (n = 0, 1, 2, 3, \dots)$$

Example: Use the Newton-Raphson method to approximate the root of $x^4 - \sin x = 1$, correct to 7 d.p., by starting with $x_0 = 1.5$.

Solution:

$$f(x) = x^4 - \sin x - 1; \quad f'(x) = 4x^3 - \cos x; \quad x_{n+1} = x_n - \frac{x_n^4 - \sin x_n - 1}{4x_n^3 - \cos x_n}$$

Steps on calculator

- | | |
|---|--|
| <p>(i) Set calculator to RADIAN mode: Shift Mode 4</p> <p>(ii) Fix calculator to 7 digits: Shift Mode 6 7</p> <p>(iii) Press 1.5 and $\boxed{=}$ on calculator ($x_0 = 1.5$)</p> | <p>(iv) Enter the formula on calculator as:
$\mathbf{Ans} - \frac{(\mathbf{Ans}^4 - \sin(\mathbf{Ans}) - 1)}{(4 \times \mathbf{Ans}^3 - \cos(\mathbf{Ans}))}$</p> <p>(v) Press $\boxed{=}$ button successively, and write down all x_n in a table</p> |
|---|--|

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)} \quad (n = 0, 1, 2, 3, \dots)$$

Calculation results

n	x_n
0	1.5000000
1	1.2717667
2	1.1885303
3	1.1778677
4	1.1777035
5	1.1777035

Example: Use the Newton-Raphson method to approximate the root of $x^4 - \sin x = 1$, correct to 7 decimal places.

Solution:

$$f(x) = x^4 - \sin x - 1;$$

Steps on calculator

- Set calculator to RAN mode: **Shift Mode 7**
- Fix calculator to 7 d.p.: **Shift Mode 6 7**
- Press **1.5** and **=** on calculator ($x_0 = 1.5$)

Approximate the root of $x^4 - \sin x = 1$, correct to 7 decimal places.

$$x_{n+1} = x_n - \frac{x_n^4 - \sin x_n - 1}{4x_n^3 - \cos x_n}$$

the formula on calculator

$$\frac{(\mathbf{Ans}^4 - \sin(\mathbf{Ans}) - 1)}{(4 \times \mathbf{Ans}^3 - \cos(\mathbf{Ans}))}$$

- Press **=** button successively, and write down all x_n in a table

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)} \quad (n = 0, 1, 2, 3, \dots)$$

Calculation results

n	x_n
0	1.5000000
1	1.2717667
2	1.1885303
3	1.1778677
4	1.1777035
5	1.1777035

Example: Use the Newton-Raphson method to approximate the root of $x^4 - \sin x = 1$, correct to 7 decimal places.

Solution:

$$f(x) = x^4 - \sin x - 1;$$

Steps on calculator

- Set calculator to RAN mode: **Shift** **Mode**
- Fix calculator to 7 d.p.: **Shift** **Mode** **6** **7**
- Press **1.5** and **=** on calculator ($x_0 = 1.5$)

Thus, the root is
 $x^* = 1.1777035$

Approximate the root of $x^4 - \sin x = 1$, correct to 7 decimal places.

$$x_{n+1} = x_n - \frac{x_n^4 - \sin x_n - 1}{4x_n^3 - \cos x_n}$$

the formula on calculator

$$\frac{(\mathbf{Ans}^4 - \sin(\mathbf{Ans}) - 1)}{(4 \times \mathbf{Ans}^3 - \cos(\mathbf{Ans}))}$$

- Press **=** button successively, and write down all x_n in a table

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)} \quad (n = 0, 1, 2, 3, \dots)$$

Practice Problems on Workbook:

- 1: Q4(ii)
- 2: Q4(iii)
- 3: Q4(iv)
- 4: Q4(vi)

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)} \quad (n = 0, 1, 2, 3, \dots)$$

Practice Problems on Workbook:

- 1: Q4(ii)
- 2: Q4(iii)
- 3: Q4(iv)
- 4: Q4(vi)

Answers:

- 1: $x = 1.895494$
- 2: $x = 1.021690$
- 3: $x = -1.272020$
- 4: $x = 2.09455148$



Example: Show that $f(x) = e^{-2x} + 1$ is always decreasing.



Example: Show that $f(x) = e^{-2x} + 1$ is always decreasing.

Solution:



Example: Show that $f(x) = e^{-2x} + 1$ is always decreasing.

Solution:

$$f'(x) = -2(e^{-2x}) < 0 \text{ for all } x \in \mathbb{R} \quad \therefore f \text{ is always decreasing.}$$



Example: Show that $f(x) = e^{-2x} + 1$ is always decreasing.

Solution:

$$f'(x) = -2(e^{-2x}) < 0 \text{ for all } x \in \mathbb{R} \quad \therefore f \text{ is always decreasing.}$$

Example: Given $f(x) = x^3 - 5x + 1$. Determine the intervals in which the function f is increasing and the intervals in which it is decreasing.



Example: Show that $f(x) = e^{-2x} + 1$ is always decreasing.

Solution:

$$f'(x) = -2(e^{-2x}) < 0 \text{ for all } x \in \mathbb{R} \quad \therefore f \text{ is always decreasing.}$$

Example: Given $f(x) = x^3 - 5x + 1$. Determine the intervals in which the function f is increasing and the intervals in which it is decreasing.

Solution:

Example: Show that $f(x) = e^{-2x} + 1$ is always decreasing.

Solution:

$$f'(x) = -2(e^{-2x}) < 0 \text{ for all } x \in \mathbb{R} \quad \therefore f \text{ is always decreasing.}$$

Example: Given $f(x) = x^3 - 5x + 1$. Determine the intervals in which the function f is increasing and the intervals in which it is decreasing.

Solution:

$$f'(x) = 3x^2 - 5 = 0 \implies x = \pm\sqrt{\frac{5}{3}}$$

Example: Show that $f(x) = e^{-2x} + 1$ is always decreasing.

Solution:

$$f'(x) = -2(e^{-2x}) < 0 \text{ for all } x \in \mathbb{R} \quad \therefore f \text{ is always decreasing.}$$

Example: Given $f(x) = x^3 - 5x + 1$. Determine the intervals in which the function f is increasing and the intervals in which it is decreasing.

Solution:

$$f'(x) = 3x^2 - 5 = 0 \implies x = \pm\sqrt{\frac{5}{3}}$$

For $x < -\sqrt{\frac{5}{3}}$, $f'(x) > 0 \quad \therefore f$ is increasing.



Example: Show that $f(x) = e^{-2x} + 1$ is always decreasing.

Solution:

$$f'(x) = -2(e^{-2x}) < 0 \text{ for all } x \in \mathbb{R} \quad \therefore f \text{ is always decreasing.}$$

Example: Given $f(x) = x^3 - 5x + 1$. Determine the intervals in which the function f is increasing and the intervals in which it is decreasing.

Solution:

$$f'(x) = 3x^2 - 5 = 0 \implies x = \pm\sqrt{\frac{5}{3}}$$

For $x < -\sqrt{\frac{5}{3}}$, $f'(x) > 0 \quad \therefore f$ is increasing.

For $-\sqrt{\frac{5}{3}} < x < \sqrt{\frac{5}{3}}$, $f'(x) < 0 \quad \therefore f$ is decreasing.

Example: Show that $f(x) = e^{-2x} + 1$ is always decreasing.

Solution:

$$f'(x) = -2(e^{-2x}) < 0 \text{ for all } x \in \mathbb{R} \quad \therefore f \text{ is always decreasing.}$$

Example: Given $f(x) = x^3 - 5x + 1$. Determine the intervals in which the function f is increasing and the intervals in which it is decreasing.

Solution:

$$f'(x) = 3x^2 - 5 = 0 \implies x = \pm\sqrt{\frac{5}{3}}$$

For $x < -\sqrt{\frac{5}{3}}$, $f'(x) > 0 \quad \therefore f$ is increasing.

For $-\sqrt{\frac{5}{3}} < x < \sqrt{\frac{5}{3}}$, $f'(x) < 0 \quad \therefore f$ is decreasing.

For $x > \sqrt{\frac{5}{3}}$, $f'(x) > 0 \quad \therefore f$ is increasing.

Example: Show that $f(x) = e^{-2x} + 1$ is always decreasing.

Solution:

$$f'(x) = -2(e^{-2x}) < 0 \text{ for all } x \in \mathbb{R} \quad \therefore f \text{ is always decreasing.}$$

Example: Given $f(x) = x^3 - 5x + 1$. Determine the intervals in which the function f is increasing and the intervals in which it is decreasing.

Solution:

$$f'(x) = 3x^2 - 5 = 0 \implies x = \pm\sqrt{\frac{5}{3}}$$

For $x < -\sqrt{\frac{5}{3}}$, $f'(x) > 0 \quad \therefore f$ is increasing.

For $-\sqrt{\frac{5}{3}} < x < \sqrt{\frac{5}{3}}$, $f'(x) < 0 \quad \therefore f$ is decreasing.

For $x > \sqrt{\frac{5}{3}}$, $f'(x) > 0 \quad \therefore f$ is increasing.

Therefore, $x \in (-\infty, -\sqrt{\frac{5}{3}})$ and $(\sqrt{\frac{5}{3}}, +\infty)$ f is increasing;

$x \in (-\sqrt{\frac{5}{3}}, \sqrt{\frac{5}{3}})$ f is decreasing.



Practice Problems on Workbook:

- 1: Q5(i)
- 2: Q5(ii)
- 3: Q5(iv)
- 4: Q5(vi)

Practice Problems on Workbook:

- 1: Q5(i)
- 2: Q5(ii)
- 3: Q5(iv)
- 4: Q5(vi)

Answers:

- 1: $f' = \sec^2 x > 0$ for all $x \in \mathbb{R}$
- 2: $f' = -\sin x > 0$ for all $x \in \text{Fourth Quadrant}$
- 3: $2x + 2 < 0$ for all $x \in (-\infty, -1)$
- 4: $x \in (-\infty, \frac{-3\sqrt{73}-9}{16})$, f is decreasing; $x \in (\frac{-3\sqrt{73}-9}{16}, 0)$, f is increasing; $x \in (0, \frac{3\sqrt{73}-9}{16})$, f is decreasing; $x \in (\frac{3\sqrt{73}-9}{16}, +\infty)$, f is increasing;

Day	Time	Venue
Tuesday	12:00 to 14:00	Trent 314a
	13:00 to 14:00	Trent 322
Thursday	16:30 to 17:30	TB 417
	17:00 to 18:00	IAMET 315
Friday	14:00 to 15:00	PB 330
	17:00 to 18:00	TB 417

Weekly drop-in Session: Wednesdays 16:00-17:00 at PB115+.