Septal GABAergic Timing in Entorhinal-Hippocampal Retrieval

May 1, 2026

1 Motivation and Gap

Memory updating in the hippocampal-entorhinal system is increasingly described as a problem of retrieval selection rather than storage replacement. Earlier traces are not eliminated when new learning occurs. Distinct engram populations can persist and compete, with accessibility shaped by context and inhibitory tone alongside network state [100, 8, 2]. This view shifts the central question from how a stored value is overwritten to how one learned context-value association is allowed to dominate behavior during retrieval.

The medial septum is a strong candidate regulator of this selection process because septal GABAergic neurons organize hippocampal and entorhinal rhythmic state. Septal output is associated with theta-promoting circuit dynamics and state transitions between active processing and replay, along with long-range timing control within the hippocampal formation [103]. Related work on hippocampal-entorhinal information-sharing networks shows that ensemble structure is not fixed. Core-periphery organization can change across temporal windows, providing a network-level substrate for switching between active computational states [80].

Recent search results therefore suggest a specific mechanistic gap. The medial septum-to-medial entorhinal cortex pathway is repeatedly implicated in oscillatory control and arousal-sensitive coordination, along with behaviorally relevant switching. However, the intermediate transformation is not yet specified at the level needed for a retrieval model. The unresolved step is how septal GABAergic input is converted within medial entorhinal cortex into a timing signal that can reorganize downstream CA1 output. A circuit explanation requires more than the statement that septal activity supports theta. It must specify how inhibitory microcircuits in medial entorhinal cortex alter the timing of entorhinal output and how that timing shift changes the CA1 ensemble state expressed during retrieval.

The primary circuit study directly motivates this gap by showing that inhibition of MS GABAergic terminals in MEC reverses updated behavior and shifts CA1 population activity back toward the pre-update pattern. This places the gap at the level of timing-based retrieval selection after sequential learning in the same context, rather than broad memory enhancement, generalized arousal, or locomotor facilitation.

2 Background

The hippocampal-entorhinal circuit supports memory retrieval through interactions among sparse coding and inhibitory competition, with temporal coordination providing the organizing frame. In the dentate gyrus and CA3, inhibitory control contributes to pattern separation and winner-take-all dynamics, allowing similar inputs to activate distinct population states [45, 41]. In computational models, context-dependent gating and sparse hippocampal memory architectures reduce interference without requiring erasure of prior representations [2, 97].

At the CA1 output stage, the active memory state is not fully captured by firing rate alone. Several retrieved references treat CA1 as a temporal and predictive structure in which spike timing and co-firing relationships interact with replay and phase organization to carry information about sequence and transition structure [33, 50, 101]. This distinction is important for memory updating. If retrieval selection depends on which context-value association controls CA1 temporal ensemble structure, then disruption of the mechanism should be visible in timing relationships and cross-correlated activity, not only in mean rate changes.

The medial entorhinal cortex provides a major route through which spatial and contextual information reaches CA1. The temporoammonic pathway supplies direct entorhinal input to distal CA1 dendrites, where local inhibition can regulate gain and delay sensitivity while shaping synchrony [24, 25]. In parallel, models of the entorhinal-hippocampal loop describe CA1 as a comparison and readout layer that balances internal hippocampal sequences with current entorhinal context [59, 50].

Medial entorhinal inhibitory populations provide a plausible transformation layer between septal input and CA1 output. Medial entorhinal cortex influences CA1 through both direct temporoammonic input and indirect pathways via CA3, providing multiple routes through which inhibitory timing can shape CA1 ensemble structure. Parvalbumin-expressing interneurons and related inhibitory motifs constrain spike timing and support phase locking while shaping the distribution of neural responses in recurrent circuits [51, 56, 57]. Septal GABAergic projections can therefore be treated as an upstream timing source, with medial entorhinal inhibitory microcircuits providing the local timing transformation and CA1 temporal ensemble organization serving as the retrieval readout that drives behavior.

3 Research Goal and Hypothesis

The research goal is to explain how medial septum GABAergic input to medial entorhinal cortex alters CA1 temporal ensemble structure during retrieval after sequential learning in the same context. The analysis focuses on retrieval selection rather than memory storage. The relevant behavioral problem is the expression of the later learned context-value association when prior and updated context-value associations can both be expressed under different retrieval conditions.

The primary circuit study directly motivates this gap by showing that inhibition of MS GABAergic terminals in MEC reverses updated behavior and shifts CA1 population activity back toward the pre-update pattern.

The central hypothesis is that medial septum GABAergic projections regulate retrieval by changing inhibitory microcircuit timing in medial entorhinal cortex. This local timing change is expected to alter the phase and precision of entorhinal output arriving in CA1. The resulting shift in input timing should reorganize CA1 ensemble structure so that the later learned

context value is expressed during retrieval.

Three assumptions define the hypothesis. First, hippocampal updating preserves competing traces rather than removing earlier ones. This assumption is consistent with retrieved work on engram persistence and context-dependent gating, alongside memory architectures that avoid catastrophic overwriting [100, 29, 97]. Second, inhibitory circuits regulate the accessibility of these traces by adjusting thresholds and synchrony while controlling competition between ensembles [41, 95, 86]. Third, CA1 expresses the selected trace through temporal ensemble organization. Evidence from temporal predictive coding and successor-representation models, along with phase-native coding, supports the idea that CA1 output depends on structured timing as well as rate [101, 50, 1].

The hypothesis is deliberately constrained. A broad claim that septal input improves memory is insufficient. The proposed mechanism requires linked effects in sequence: septal GABAergic control of medial entorhinal timing, medial entorhinal inhibitory control of output precision, then CA1 temporal reorganization during the retrieval decision. If one link is absent, the pathway would not provide a complete account of retrieval selection after sequential learning.

4 Proposed Approach

The proposed analysis synthesizes the retrieved circuit and computational literature around a sequence from MS GABAergic output through MEC inhibitory timing to CA1 temporal ensemble structure, which determines retrieval behavior. The first level treats medial septum GABAergic output as the upstream timing source. Work on theta-promoting septal circuits and optogenetic control, alongside sleep-wake state transitions, establishes that septal activity can regulate hippocampal-entorhinal rhythmic state with sufficient temporal specificity to affect information routing [103, 75, 104]. Because septal manipulation can also influence locomotion and arousal, those effects must be separated from retrieval-specific mechanisms [26, 58].

The second level treats medial entorhinal inhibitory microcircuits as the transformation layer. The key question is whether septal GABAergic input changes the timing of local inhibitory populations that regulate entorhinal output. Parvalbumin-related and recurrent inhibitory motifs are relevant because they can narrow spike windows and normalize response distributions while accelerating adaptation to changing input statistics [51, 56, 57]. This level isolates timing as the controlled variable. Evidence that only firing rate or nonspecific excitability changes would not be sufficient.

The third level treats CA1 ensemble dynamics as the retrieval readout. CA1 should be evaluated through temporal structure, including phase relationships and cross-correlations, along with replay-like sequence organization and ensemble composition. A compact readout for this comparison is

$$S_{\mathrm{CA1}}(a, b) = \mathrm{corr}(C_a, C_b),$$

where C_a and C_b are pairwise CA1 cross-correlation matrices from two retrieval conditions. Retrieved models and analyses support temporal coding and predictive coding, along with calcium-imaging-based ensemble decoding, as relevant readouts [33, 101, 17]. The key test is whether altered entorhinal timing can change which CA1 temporal ensemble structure is expressed when prior and updated context-value associations can both be expressed under different retrieval conditions.

Support for the mechanism would require convergence across these levels. Septal control should alter medial entorhinal timing and medial entorhinal inhibitory control should reshape output precision before CA1 shows retrieval-specific temporal reorganization. A dissociation at any level would constrain the model and redirect attention toward downstream output circuits and direct CA1 inhibition, with behavioral state variables treated as a separate alternative.

5 Outcomes and Significance

Four outcomes would support the proposed mechanism.

First, medial septum GABAergic input to medial entorhinal cortex should be required most strongly when retrieval depends on selecting between sequentially learned context-value associations in the same context. Perturbation should impair retrieval precision without producing a general memory failure. Behavioral controls are therefore necessary because septal and GABAergic manipulations can affect locomotion and arousal, along with anxiety-related state [26, 75].

Second, the CA1 effect should appear as altered temporal ensemble structure. Changes in spike timing relationships, phase organization, or cross-correlated ensemble composition would support a timing-based account. A pure firing-rate effect would support a less specific gain-control model. The relevant evidence base includes temporal coding models and dynamic core-periphery ensemble structure, alongside Bayesian decoding of spatially referenced calcium traces [80, 33, 17].

Third, the mechanism should pass through medial entorhinal inhibitory microcircuits. Septal input should affect local timing control before the CA1 readout changes. Evidence for this step would include altered interneuron phase locking, changed entorhinal output precision, or modified gain and whitening dynamics in medial entorhinal populations [93, 56, 19]. Without this intermediate transformation, the pathway would not explain how septal timing becomes a CA1 retrieval-selection signal.

Fourth, retrieval should follow inhibitory competition rules rather than storage rules. Prior and updated context-value associations should both be expressible under different retrieval conditions, while inhibitory timing determines which association becomes behaviorally expressed. This outcome is consistent with models of engram coexistence and inhibitory plasticity, along with context-dependent gating [100, 95, 86].

The significance is that memory updating can be framed as a timing problem at the level of inhibitory control. The hippocampal-entorhinal system would not need to replace one learned association with another. It would instead regulate which association dominates CA1 temporal ensemble structure during retrieval. A negative result would also be informative. If behavioral switching occurs without CA1 temporal reorganization or without medial entorhinal timing control, then the mechanism should be localized outside the proposed MS GABAergic output to MEC inhibitory timing to CA1 temporal ensemble circuit pathway.

References

[1] *A Unified Phase-native Computational Principle Governs Hippocampal Spike Timing and Neural Coding*. 2026. arXiv: 2603.19690. URL: https://arxiv.org/abs/2603.19690.

[2] *Context-modulation of hippocampal dynamics and deep convolutional networks*. 2017. arXiv: 1711.09876. URL: https://arxiv.org/abs/1711.09876.

[3] *Discretization of continuous input spaces in the hippocampal autoencoder*. 2024. arXiv: 2405.14600. URL: https://arxiv.org/abs/2405.14600.

[4] *Linking In-context Learning in Transformers to Human Episodic Memory*. 2024. arXiv: 2405.14992. URL: https://arxiv.org/abs/2405.14992.

[5] *Automatic Neuron Detection in Calcium Imaging Data Using Convolutional Networks*. 2016. arXiv: 1606.07372. URL: https://arxiv.org/abs/1606.07372.

[6] *Feedforward Architectures Driven by Inhibitory Interactions*. 2017. arXiv: 1701.04905. URL: https://arxiv.org/abs/1701.04905.

[7] *Cross-Frequency Coupling Increases Memory Capacity in Oscillatory Neural Networks*. 2022. arXiv: 2204.07163. URL: https://arxiv.org/abs/2204.07163.

[8] *A bio-inspired implementation of a sparse-learning spike-based hippocampus memory model*. 2022. arXiv: 2206.04924. URL: https://arxiv.org/abs/2206.04924.

[9] *Spike-based computational models of bio-inspired memories in the hippocampal CA3 region on SpiNNaker*. 2022. arXiv: 2205.04782. URL: https://arxiv.org/abs/2205.04782.

[10] *Bio-inspired computational memory model of the Hippocampus: an approach to a neuromorphic spike-based Content-Addressable Memory*. 2023. arXiv: 2310.05868. URL: https://arxiv.org/abs/2310.05868.

[11] *Statistical Mechanics of Learning via Reverberation in Bidirectional Associative Memories*. 2023. arXiv: 2307.08365. URL: https://arxiv.org/abs/2307.08365.

[12] *Transformers Remember First, Forget Last: Dual-Process Interference in LLMs*. 2026. arXiv: 2603.00270. URL: https://arxiv.org/abs/2603.00270.

[13] *Recurrent Auto-Encoder With Multi-Resolution Ensemble and Predictive Coding for Multivariate Time-Series Anomaly Detection*. 2022. arXiv: 2202.10001. URL: https://arxiv.org/abs/2202.10001.

[14] *Neurogenesis and multiple plasticity mechanisms enhance associative memory retrieval in a spiking network model of the hippocampus*. 2017. arXiv: 1704.07526. URL: https://arxiv.org/abs/1704.07526.

[15] *Cortical microcircuits as gated-recurrent neural networks*. 2017. arXiv: 1711.02448. URL: https://arxiv.org/abs/1711.02448.

[16] *Recurrent neural network models for working memory of continuous variables: activity manifolds, connectivity patterns, and dynamic codes*. 2021. arXiv: 2111.01275. URL: https://arxiv.org/abs/2111.01275.

[17] *Decoding Neuronal Ensembles from Spatially-Referenced Calcium Traces: A Bayesian Semiparametric Approach*. 2025. arXiv: 2508.09576. URL: https://arxiv.org/abs/2508.09576.

[18] *Grid cells show field-to-field variability and this explains the aperiodic response of inhibitory interneurons*. 2017. arXiv: 1701.04893. URL: https://arxiv.org/abs/1701.04893.

[19] *Adaptive whitening in neural populations with gain-modulating interneurons*. 2023. arXiv: 2301.11955. URL: https://arxiv.org/abs/2301.11955.

[20] *Temporary changes in large-scale memory neural networks after fear learning and extinction in healthy adults*. 2020. arXiv: 2011.00257. URL: https://arxiv.org/abs/2011.00257.

[21] *Outan: An On-Head System for Driving micro-LED Arrays Implanted in Freely Moving Mice*. 2022. arXiv: 2211.13589. URL: https://arxiv.org/abs/2211.13589.

[22] *Neural correlates of episodic memory in the Memento cohort*. 2018. arXiv: 1812.08031. URL: https://arxiv.org/abs/1812.08031.

[23] *Pattern Separation in a Spiking Neural Network of Hippocampus Robust to Imbalanced Excitation/Inhibition*. 2018. arXiv: 1808.00367. URL: https://arxiv.org/abs/1808.00367.

[24] *Effects of synapse location, delay and background stochastic activity on synchronising hippocampal CA1 neurons*. 2024. arXiv: 2409.10460. URL: https://arxiv.org/abs/2409.10460.

[25] *Hippocampal synchronization in a realistic CA1 neuron model*. 2024. arXiv: 2409.10431. URL: https://arxiv.org/abs/2409.10431.

[26] *State-Dependent Modulation of Locomotion by GABAergic Spinal Sensory Neurons*. 2018. arXiv: 1812.00625. URL: https://arxiv.org/abs/1812.00625.

[27] *Human-inspired Episodic Memory for Infinite Context LLMs*. 2024. arXiv: 2407.09450. URL: https://arxiv.org/abs/2407.09450.

[28] *Robust computation with rhythmic spike patterns*. 2019. arXiv: 1901.07718. URL: https://arxiv.org/abs/1901.07718.

[29] *Key-value memory in the brain*. 2025. arXiv: 2501.02950. URL: https://arxiv.org/abs/2501.02950.

[30] *Palimpsest Memories Stored in Memristive Synapses*. 2021. arXiv: 2109.13198. URL: https://arxiv.org/abs/2109.13198.

[31] *Optimal rate-variance coding due to firing threshold adaptation near criticality*. 2025. arXiv: 2509.04106. URL: https://arxiv.org/abs/2509.04106.

[32] *Neuromodulation-inspired gated associative memory networks:extended memory retrieval and emergent multistability*. 2025. arXiv: 2512.13859. URL: https://arxiv.org/abs/2512.13859.

[33] *An uncertainty principle for neural coding: Conjugate representations of position and velocity are mapped onto firing rates and co-firing rates of neural spike trains*. 2019. arXiv: 1912.11126. URL: https://arxiv.org/abs/1912.11126.

[34] *HippoRAG: Neurobiologically Inspired Long-Term Memory for Large Language Models*. 2024. arXiv: 2405.14831. URL: https://arxiv.org/abs/2405.14831.

[35] *Extended temporal association memory by inhibitory Hebbian learning*. 2018. arXiv: 1809.05254. URL: https://arxiv.org/abs/1809.05254.

[36] *Efficient Neural Network Approximation of Robust PCA for Automated Analysis of Calcium Imaging Data*. 2021. arXiv: 2108.01665. URL: https://arxiv.org/abs/2108.01665.

[37] *Local inhibitory plasticity tunes global brain dynamics and allows the emergence of functional brain networks*. 2014. arXiv: 1412.6325. URL: https://arxiv.org/abs/1412.6325.

[38] *Robust retrieval of dynamic sequences through interaction modulation*. 2022. arXiv: 2211.17152. URL: https://arxiv.org/abs/2211.17152.

[39] *SynapticRAG: Enhancing Temporal Memory Retrieval in Large Language Models through Synaptic Mechanisms*. 2024. arXiv: 2410.13553. URL: https://arxiv.org/abs/2410.13553.

[40] *PhiNets: Brain-inspired Non-contrastive Learning Based on Temporal Prediction Hypothesis*. 2024. arXiv: 2405.14650. URL: https://arxiv.org/abs/2405.14650.

[41] *Feedback inhibition shapes emergent computational properties of cortical microcircuit motifs*. 2017. arXiv: 1705.07614. URL: https://arxiv.org/abs/1705.07614.

[42] *Hippocampal representations emerge when training recurrent neural networks on a memory dependent maze navigation task*. 2020. arXiv: 2012.01328. URL: https://arxiv.org/abs/2012.01328.

[43] *Coupled Neural Associative Memories*. 2013. arXiv: 1301.1555. URL: https://arxiv.org/abs/1301.1555.

[44] *Noise Facilitation in Associative Memories of Exponential Capacity*. 2014. arXiv: 1403.3305. URL: https://arxiv.org/abs/1403.3305.

[45] *Dynamical Origin for Winner-Take-All Competition in A Biological Network of The Hippocampal Dentate Gyrus*. 2021. arXiv: 2105.06057. URL: https://arxiv.org/abs/2105.06057.

[46] *Effect of Adult-Born Immature Granule Cells on Pattern Separation in The Hippocampal Dentate Gyrus*. 2023. arXiv: 2301.05387. URL: https://arxiv.org/abs/2301.05387.

[47] *Structural Correlates Of Spatial Navigation And Memory Formation*. 2022. arXiv: 2203.13434. URL: https://arxiv.org/abs/2203.13434.

[48] *Neuronal mechanisms for sequential activation of memory items: dynamics and reliability*. 2019. arXiv: 1904.12133. URL: https://arxiv.org/abs/1904.12133.

[49] *Binding in hippocampal-entorhinal circuits enables compositionality in cognitive maps*. 2024. arXiv: 2406.18808. URL: https://arxiv.org/abs/2406.18808.

[50] *Toward the biological model of the hippocampus as the successor representation agent*. 2020. arXiv: 2006.11975. URL: https://arxiv.org/abs/2006.11975.

[51] *A Computational Analysis of the Function of Three Inhibitory Cell Types in Contextual Visual Processing*. 2016. arXiv: 1609.03622. URL: https://arxiv.org/abs/1609.03622.

[52] *Hippocampal Spatial Mapping As Fast Graph Learning*. 2021. arXiv: 2107.00567. URL: https://arxiv.org/abs/2107.00567.

[53] *Memory-enriched computation and learning in spiking neural networks through Hebbian plasticity*. 2022. arXiv: 2205.11276. URL: https://arxiv.org/abs/2205.11276.

[54] *Emergence of Spatial Representation in an Actor-Critic Agent with Hippocampus-Inspired Sequence Generator*. 2025. arXiv: 2510.09951. URL: https://arxiv.org/abs/2510.09951.

[55] *HippoMM: Hippocampal-inspired Multimodal Memory for Long Audiovisual Event Understanding*. 2025. arXiv: 2504.10739. URL: https://arxiv.org/abs/2504.10739.

[56] *Interneurons accelerate learning dynamics in recurrent neural networks for statistical adaptation*. 2022. arXiv: 2209.10634. URL: https://arxiv.org/abs/2209.10634.

[57] *Shaping the distribution of neural responses with interneurons in a recurrent circuit model*. 2024. arXiv: 2405.17745. URL: https://arxiv.org/abs/2405.17745.

[58] *A System-on-Chip for Closed-loop Optogenetic Sleep Modulation*. 2021. arXiv: 2108.12261. URL: https://arxiv.org/abs/2108.12261.

[59] *GATE: Adaptive Learning with Working Memory by Information Gating in Multi-lamellar Hippocampal Formation*. 2025. arXiv: 2501.12615. URL: https://arxiv.org/abs/2501.12615.

[60] *Bridging Cognitive Maps: a Hierarchical Active Inference Model of Spatial Alternation Tasks and the Hippocampal-Prefrontal Circuit*. 2023. arXiv: 2308.11463. URL: https://arxiv.org/abs/2308.11463.

[61] *Mechanisms underlying the response of mouse cortical networks to optogenetic manipulation*. 2019. arXiv: 1907.00816. URL: https://arxiv.org/abs/1907.00816.

[62] *Bifurcation of spiking oscillations from a center in resonate-and-fire neurons*. 2025. arXiv: 2510.13156. URL: https://arxiv.org/abs/2510.13156.

[63] *Alleviating catastrophic forgetting using context-dependent gating and synaptic stabilization*. 2018. arXiv: 1802.01569. URL: https://arxiv.org/abs/1802.01569.

[64] *A Hippocampus Model for Online One-Shot Storage of Pattern Sequences*. 2019. arXiv: 1905.12937. URL: https://arxiv.org/abs/1905.12937.

[65] *Grid-like structure is optimal for path integration*. 2016. arXiv: 1606.01239. URL: https://arxiv.org/abs/1606.01239.

[66] *Effect of Neuromodulation of Short-Term Plasticity on Information Processing in Hippocampal Interneuron Synapses*. 2018. arXiv: 1804.02115. URL: https://arxiv.org/abs/1804.02115.

[67] *Cross-talk and transitions between multiple spatial maps in an attractor neural network model of the hippocampus: phase diagram (I)*. 2013. arXiv: 1304.1457. URL: https://arxiv.org/abs/1304.1457.

[68] *Hippocampal Spike-Timing Correlations Lead to Hexagonal Grid Fields*. 2017. arXiv: 1706.06914. URL: https://arxiv.org/abs/1706.06914.

[69] *An Efficient Coding Theory for a Dynamic Trajectory Predicts non-Uniform Allocation of Grid Cells to Modules in the Entorhinal Cortex*. 2016. arXiv: 1601.02948. URL: https://arxiv.org/abs/1601.02948.

[70] *Stable Memory Allocation in the Hippocampus: Fundamental Limits and Neural Realization*. 2016. arXiv: 1612.04659. URL: https://arxiv.org/abs/1612.04659.

[71] *A Coupled Neural Field Model for the Standard Consolidation Theory*. 2024. arXiv: 2404.02938. URL: https://arxiv.org/abs/2404.02938.

[72] *CREIMBO: Cross-Regional Ensemble Interactions in Multi-view Brain Observations*. 2024. arXiv: 2405.17395. URL: https://arxiv.org/abs/2405.17395.

[73] *Spatial and Temporal Correlates of Vesicular Release at Hippocampal Synapses*. 2010. arXiv: 1004.2009. URL: https://arxiv.org/abs/1004.2009.

[74] *High fidelity optogenetic control of individual prefrontal cortical pyramidal neurons in vivo*. 2012. arXiv: 1208.2720. URL: https://arxiv.org/abs/1208.2720.

[75] *Bidirectional optogenetic control of inhibitory neurons in freely-moving mice*. 2022. arXiv: 2211.13591. URL: https://arxiv.org/abs/2211.13591.

[76] *Multisensory learning recruits visual neurons into an olfactory memory engram*. 2026. arXiv: 2604.28007. URL: https://arxiv.org/abs/2604.28007.

[77] *Modulation of metastable ensemble dynamics explains the inverted-U relationship between tone discriminability and arousal in auditory cortex*. 2024. arXiv: 2404.03902. URL: https://arxiv.org/abs/2404.03902.

[78] *The cross-frequency mediation mechanism of intracortical information transactions*. 2017. arXiv: 1703.07654. URL: https://arxiv.org/abs/1703.07654.

[79] *An algebraic approach to spike-time neural codes in the hippocampus*. 2022. arXiv: 2211.06692. URL: https://arxiv.org/abs/2211.06692.

[80] *Dynamic core-periphery structure of information sharing networks in the entorhinal cortex and the hippocampus*. 2020. arXiv: 2001.06371. URL: https://arxiv.org/abs/2001.06371.

[81] *Forgetting leads to chaos in attractor networks*. 2021. arXiv: 2112.00119. URL: https://arxiv.org/abs/2112.00119.

[82] *Querying hippocampal replay with subcortical inputs*. 2022. arXiv: 2205.02665. URL: https://arxiv.org/abs/2205.02665.

[83] *ConFER: A Neurally Constrained Computational Model of Context-Dependent Fear Extinction Recall and Relapse*. 2024. arXiv: 2411.08140. URL: https://arxiv.org/abs/2411.08140.

[84] *A Cognitive Architecture for Machine Consciousness and Artificial Superintelligence: Thought Is Structured by the Iterative Updating of Working Memory*. 2022. arXiv: 2203.17255. URL: https://arxiv.org/abs/2203.17255.

[85] *Memory Recall and Spike Frequency Adaptation*. 2016. arXiv: 1601.02974. URL: https://arxiv.org/abs/1601.02974.

[86] *Dis-inhibitory neuronal circuits can control the sign of synaptic plasticity*. 2023. arXiv: 2310.19614. URL: https://arxiv.org/abs/2310.19614.

[87] *Deep Episodic Memory: Encoding, Recalling, and Predicting Episodic Experiences for Robot Action Execution*. 2018. arXiv: 1801.04134. URL: https://arxiv.org/abs/1801.04134.

[88] *A balanced memory network*. 2007. arXiv: 0704.3005. URL: https://arxiv.org/abs/0704.3005.

[89] *A Neural Network Model of Continual Learning with Cognitive Control*. 2022. arXiv: 2202.04773. URL: https://arxiv.org/abs/2202.04773.

[90] *Competition through selective inhibitory synchrony*. 2012. arXiv: 1201.2845. URL: https://arxiv.org/abs/1201.2845.

[91] *Associative Memories via Predictive Coding*. 2021. arXiv: 2109.08063. URL: https://arxiv.org/abs/2109.08063.

[92] *The role of oscillations in grid cells' toroidal topology*. 2025. arXiv: 2501.19262. URL: https://arxiv.org/abs/2501.19262.

[93] *On the organization of grid and place cells: Neural denoising via subspace learning*. 2017. arXiv: 1712.04602. URL: https://arxiv.org/abs/1712.04602.

[94] *Dynamical Mechanisms for Coordinating Long-term Working Memory Based on the Precision of Spike-timing in Cortical Neurons*. 2025. arXiv: 2512.15891. URL: https://arxiv.org/abs/2512.15891.

[95] *Unsupervised learning by a nonlinear network with Hebbian excitatory and anti-Hebbian inhibitory neurons*. 2018. arXiv: 1812.11581. URL: https://arxiv.org/abs/1812.11581.

[96] *AI-native Memory: A Pathway from LLMs Towards AGI*. 2024. arXiv: 2406.18312. URL: https://arxiv.org/abs/2406.18312.

[97] *Content Addressable Memory Without Catastrophic Forgetting by Heteroassociation with a Fixed Scaffold*. 2022. arXiv: 2202.00159. URL: https://arxiv.org/abs/2202.00159.

[98] *Theta, alpha and gamma traveling waves in a multi-item working memory model*. 2021. arXiv: 2103.15266. URL: https://arxiv.org/abs/2103.15266.

[99] *Fast amortized inference of neural activity from calcium imaging data with variational autoencoders*. 2017. arXiv: 1711.01846. URL: https://arxiv.org/abs/1711.01846.

[100] *Engram Memory Encoding and Retrieval: A Neurocomputational Perspective*. 2025. arXiv: 2506.01659. URL: https://arxiv.org/abs/2506.01659.

[101] *Sequential Memory with Temporal Predictive Coding*. 2023. arXiv: 2305.11982. URL: https://arxiv.org/abs/2305.11982.

[102] *Artificial Neuronal Ensembles with Learned Context Dependent Gating*. 2023. arXiv: 2301.07187. URL: https://arxiv.org/abs/2301.07187.

[103] *SWS promoting MHb-IPN-MRN circuit opposes the theta promoting circuit, active wake and REM sleep*. 2015. arXiv: 1510.01100. URL: https://arxiv.org/abs/1510.01100.

[104] *Mechanisms for anesthesia, unawareness, respiratory depression, memory replay and sleep: MHb > IPN > PAG + DRN + MRN > claustrum > cortical slow waves*. 2025. arXiv: 2509.04454. URL: https://arxiv.org/abs/2509.04454.

[105] *Neuronal Response Impedance Mechanism Implementing Cooperative Networks with Low Firing Rates and Microseconds Precision*. 2014. arXiv: 1411.5258. URL: https://arxiv.org/abs/1411.5258.

[106] *Ripple oscillations in the left temporal neocortex are associated with impaired verbal episodic memory encoding*. 2018. arXiv: 1807.10269. URL: https://arxiv.org/abs/1807.10269.

[107] *Anti-Retroactive Interference for Lifelong Learning*. 2022. arXiv: 2208.12967. URL: https://arxiv.org/abs/2208.12967.

[108] *GAM-RAG: Gain-Adaptive Memory for Evolving Retrieval in Retrieval-Augmented Generation*. 2026. arXiv: 2603.01783. URL: https://arxiv.org/abs/2603.01783.

[109] *Time Makes Space: Emergence of Place Fields in Networks Encoding Temporally Continuous Sensory Experiences*. 2024. arXiv: 2408.05798. URL: https://arxiv.org/abs/2408.05798.

[110] *REMI: Reconstructing Episodic Memory During Internally Driven Path Planning*. 2025. arXiv: 2507.02064. URL: https://arxiv.org/abs/2507.02064.

[111] *A Robotic Model of Hippocampal Reverse Replay for Reinforcement Learning*. 2021. arXiv: 2102.11914. URL: https://arxiv.org/abs/2102.11914.

[112] *Generalisation of structural knowledge in the hippocampal-entorhinal system*. 2018. arXiv: 1805.09042. URL: https://arxiv.org/abs/1805.09042.

[113] *BayesPCN: A Continually Learnable Predictive Coding Associative Memory*. 2022. arXiv: 2205.09930. URL: https://arxiv.org/abs/2205.09930.

[114] *Mapping the Genetic-Imaging-Clinical Pathway with Applications to Alzheimer's Disease*. 2020. arXiv: 2007.04558. URL: https://arxiv.org/abs/2007.04558.

[115] *Learning Sparse Spatial Codes for Cognitive Mapping Inspired by Entorhinal-Hippocampal Neurocircuit*. 2019. arXiv: 1910.04590. URL: https://arxiv.org/abs/1910.04590.

[116] *Theories of synaptic memory consolidation and intelligent plasticity for continual learning*. 2024. arXiv: 2405.16922. URL: https://arxiv.org/abs/2405.16922.

[117] *Episodic memory governs choices: An RNN-based reinforcement learning model for decision-making task*. 2021. arXiv: 2103.03679. URL: https://arxiv.org/abs/2103.03679.

[118] *Efficient and accurate extraction of in vivo calcium signals from microendoscopic video data*. 2016. arXiv: 1605.07266. URL: https://arxiv.org/abs/1605.07266.