Septal GABAergic Timing in Entorhinal-Hippocampal Retrieval

Gil Raitses

1 Motivation and Gap

Memory updating in the hippocampal-entorhinal system is increasingly described as a problem of retrieval selection rather than storage replacement. Earlier traces are not eliminated when new learning occurs. Distinct engram populations can persist, compete, and become differentially accessible according to context, inhibitory tone, and network state [100, 8, 2]. This view shifts the central question from how a stored value is overwritten to how one available trace is allowed to dominate behavior during retrieval.

The medial septum is a strong candidate regulator of this selection process because septal GABAergic neurons organize hippocampal and entorhinal rhythmic state. Septal output is associated with theta-promoting circuit dynamics, state transitions between active processing and replay, and long-range timing control within the hippocampal formation [103]. Related work on hippocampal-entorhinal information-sharing networks shows that ensemble structure is not fixed. Core-periphery organization can change across temporal windows, providing a network-level substrate for switching between active computational states [80].

Recent search results therefore suggest a specific mechanistic gap. The medial septum-to-medial entorhinal cortex pathway is repeatedly implicated in oscillatory control, arousal-sensitive coordination, and behaviorally relevant switching. However, the intermediate transformation is not yet specified at the level needed for a retrieval model. The unresolved step is how septal GABAergic input is converted within medial entorhinal cortex into a timing signal that can reorganize downstream CA1 output. A circuit explanation requires more than the statement that septal activity supports theta. It must specify how inhibitory microcircuits in medial entorhinal cortex alter the timing of entorhinal output, and how that timing shift changes the CA1 ensemble state expressed during retrieval.

The gap addressed here is therefore narrow. The relevant mechanism is not broad memory enhancement, generalized arousal, or overall locomotor facilitation. The target mechanism is timing-based retrieval selection after sequential learning in the same context.

2 Background

The hippocampal-entorhinal circuit supports memory retrieval through interactions among sparse coding, inhibitory competition, and temporal coordination. In the dentate gyrus and CA3, inhibitory control contributes to pattern separation and winner-take-all dynamics, allowing similar inputs to activate distinct population states [45, 41]. In computational models, context-dependent gating and sparse hippocampal memory architectures reduce interference without requiring erasure of prior representations [2, 97].

At the CA1 output stage, the active memory state is not fully captured by firing rate alone. Several retrieved references treat CA1 as a temporal and predictive structure in which spike timing, co-firing relationships, replay, and phase organization carry information about sequence and transition structure [33, 50, 101]. This distinction is important for memory updating. If retrieval selection depends on which trace controls CA1 temporal ensemble structure, then disruption of the mechanism should be visible in timing relationships and cross-correlated activity, not only in mean rate changes.

The medial entorhinal cortex provides a major route through which spatial and contextual information reaches CA1. The temporoammonic pathway supplies direct entorhinal input to distal CA1 dendrites, where local inhibition can regulate gain, delay sensitivity, and synchrony [24, 25]. In parallel, models of the entorhinal-hippocampal loop describe CA1 as a comparison and readout layer that balances internal hippocampal sequences with current entorhinal context [59, 50].

Medial entorhinal inhibitory populations provide a plausible transformation layer between septal input and CA1 output. Parvalbumin-expressing interneurons and related inhibitory motifs constrain spike timing, support phase locking, and shape the distribution of neural responses in recurrent circuits [51, 56, 57]. Septal GABAergic projections can therefore be treated as an upstream timing source, medial entorhinal inhibitory microcircuits as the local timing transformation, and CA1 temporal ensemble organization as the retrieval readout.

3 Research Goal and Hypothesis

The research goal is to explain how medial septum GABAergic input to medial entorhinal cortex alters CA1 temporal ensemble structure during retrieval after sequential learning in the same context. The analysis focuses on retrieval selection rather than memory storage. The relevant behavioral problem is the expression of one learned value when multiple memory traces remain available.

The central hypothesis is that medial septum GABAergic projections regulate retrieval by changing inhibitory microcircuit timing in medial entorhinal cortex. This local timing change is expected to alter the phase and precision of entorhinal output arriving in CA1. The resulting shift in input timing should reorganize CA1 ensemble structure so that the later learned context value is expressed during retrieval.

Three assumptions define the hypothesis. First, hippocampal updating preserves competing traces rather than removing earlier ones. This assumption is consistent with retrieved

work on engram persistence, context-dependent gating, and memory architectures that avoid catastrophic overwriting [100, 29, 97]. Second, inhibitory circuits regulate the accessibility of these traces by adjusting thresholds, synchrony, and competition between ensembles [41, 95, 86]. Third, CA1 expresses the selected trace through temporal ensemble organization. Evidence from temporal predictive coding, successor-representation models, and phase-native coding supports the idea that CA1 output depends on structured timing as well as rate [101, 50, 1].

The hypothesis is deliberately constrained. A broad claim that septal input improves memory is insufficient. The proposed mechanism requires a sequence of linked effects: septal GABAergic control of medial entorhinal timing, medial entorhinal inhibitory control of output precision, and CA1 temporal reorganization during the retrieval decision. If one link is absent, the pathway would not provide a complete account of retrieval selection after sequential learning.

4 Proposed Approach

The proposed analysis synthesizes the retrieved circuit and computational literature around three linked levels. The first level treats medial septum GABAergic output as the upstream timing source. Work on theta-promoting septal circuits, optogenetic control, and sleep-wake state transitions establishes that septal activity can regulate hippocampal-entorhinal rhythmic state with sufficient temporal specificity to affect information routing [103, 75, 104]. Because septal manipulation can also influence locomotion and arousal, those effects must be separated from retrieval-specific mechanisms [26, 58].

The second level treats medial entorhinal inhibitory microcircuits as the transformation layer. The key question is whether septal GABAergic input changes the timing of local inhibitory populations that regulate entorhinal output. Parvalbumin-related and recurrent inhibitory motifs are relevant because they can narrow spike windows, normalize response distributions, and accelerate adaptation to changing input statistics [51, 56, 57]. This level isolates timing as the controlled variable. Evidence that only firing rate or nonspecific excitability changes would not be sufficient.

The third level treats CA1 ensemble dynamics as the retrieval readout. CA1 should be evaluated through temporal structure, including phase relationships, cross-correlations, replay-like sequence organization, and ensemble composition. Retrieved models and analyses support the use of temporal coding, predictive coding, and calcium-imaging-based ensemble decoding as relevant readouts [33, 101, 17]. The key test is whether altered entorhinal timing can change which CA1 population state is expressed when competing traces are available.

Support for the mechanism would require convergence across these levels. Septal control should alter medial entorhinal timing, medial entorhinal inhibitory control should reshape output precision, and CA1 should show retrieval-specific temporal reorganization. A dissociation at any level would constrain the model and redirect attention toward downstream output circuits, direct CA1 inhibition, or behavioral state variables.

5 Outcomes and Significance

Four outcomes would support the proposed mechanism. First, medial septum GABAergic input to medial entorhinal cortex should be required most strongly when retrieval depends on selecting between sequentially learned values in the same context. Perturbation should impair retrieval precision without producing a general memory failure. Behavioral controls are therefore necessary because septal and GABAergic manipulations can affect locomotion, arousal, and anxiety-related state [26, 75].

Second, the CA1 effect should appear as altered temporal ensemble structure. Changes in spike timing relationships, phase organization, or cross-correlated ensemble composition would support a timing-based account. A pure firing-rate effect would support a less specific gain-control model. The relevant evidence base includes temporal coding models, dynamic core-periphery ensemble structure, and Bayesian decoding of spatially referenced calcium traces [80, 33, 17].

Third, the mechanism should pass through medial entorhinal inhibitory microcircuits. Septal input should affect local timing control before the CA1 readout changes. Evidence for this step would include altered interneuron phase locking, changed entorhinal output precision, or modified gain and whitening dynamics in medial entorhinal populations [93, 56, 19]. Without this intermediate transformation, the pathway would not explain how septal timing becomes a CA1 retrieval-selection signal.

Fourth, retrieval should follow competition rules rather than storage rules. Earlier traces should remain available, while inhibitory timing determines which trace becomes behaviorally expressed. This outcome is consistent with models of engram coexistence, inhibitory plasticity, and context-dependent gating [100, 95, 86].

The significance is that memory updating can be framed as a timing problem at the level of inhibitory control. The hippocampal-entorhinal system would not need to replace one trace with another. It would instead regulate which trace is permitted to dominate CA1 temporal structure during retrieval. A negative result would also be informative: if behavioral switching occurs without CA1 temporal reorganization or without medial entorhinal timing control, then the mechanism should be localized outside the proposed septal-entorhinal-CA1 sequence.

References

[1] R. Ahmadvand, S. S. Sharif, and Y. M. Banad. *A Unified Phase-native Computational Principle Governs Hippocampal Spike Timing and Neural Coding*. 2026. arXiv: 2603.19690. URL: https://arxiv.org/abs/2603.19690.

[2] J. B. Aimone and W. M. Severa. *Context-modulation of hippocampal dynamics and deep convolutional networks*. 2017. arXiv: 1711.09876. URL: https://arxiv.org/abs/1711.09876.

[3] A. F. Amil, I. T. Freire, and P. F. M. J. Verschure. *Discretization of continuous input spaces in the hippocampal autoencoder*. 2024. arXiv: 2405.14600. URL: https://arxiv.org/abs/2405.14600.

[4] L. Ji-An, C. Y. Zhou, M. K. Benna, and M. G. Mattar. *Linking In-context Learning in Transformers to Human Episodic Memory*. 2024. arXiv: 2405.14992. URL: https://arxiv.org/abs/2405.14992.

[5] N. J. Apthorpe et al. *Automatic Neuron Detection in Calcium Imaging Data Using Convolutional Networks*. 2016. arXiv: 1606.07372. URL: https://arxiv.org/abs/1606.07372.

[6] Y. N. Billeh and M. T. Schaub. *Feedforward Architectures Driven by Inhibitory Interactions*. 2017. arXiv: 1701.04905. URL: https://arxiv.org/abs/1701.04905.

[7] C. Bybee, A. Belsten, and F. T. Sommer. *Cross-Frequency Coupling Increases Memory Capacity in Oscillatory Neural Networks*. 2022. arXiv: 2204.07163. URL: https://arxiv.org/abs/2204.07163.

[8] D. Casanueva-Morato et al. *A bio-inspired implementation of a sparse-learning spike-based hippocampus memory model*. 2022. arXiv: 2206.04924. URL: https://arxiv.org/abs/2206.04924.

[9] D. Casanueva-Morato et al. *Spike-based computational models of bio-inspired memories in the hippocampal CA3 region on SpiNNaker*. 2022. arXiv: 2205.04782. URL: https://arxiv.org/abs/2205.04782.

[10] D. Casanueva-Morato et al. *Bio-inspired computational memory model of the Hippocampus: an approach to a neuromorphic spike-based Content-Addressable Memory*. 2023. arXiv: 2310.05868. URL: https://arxiv.org/abs/2310.05868.

[11] M. S. Centonze, I. Kanter, and A. Barra. *Statistical Mechanics of Learning via Reverberation in Bidirectional Associative Memories*. 2023. arXiv: 2307.08365. URL: https://arxiv.org/abs/2307.08365.

[12] S. Chattaraj and K. Raj. *Transformers Remember First, Forget Last: Dual-Process Interference in LLMs*. 2026. arXiv: 2603.00270. URL: https://arxiv.org/abs/2603.00270.

[13] H. Choi, S. Kim, and P. Kang. *Recurrent Auto-Encoder With Multi-Resolution Ensemble and Predictive Coding for Multivariate Time-Series Anomaly Detection*. 2022. arXiv: 2202.10001. URL: https://arxiv.org/abs/2202.10001.

[14] Y. Chua and C. Tan. *Neurogenesis and multiple plasticity mechanisms enhance associative memory retrieval in a spiking network model of the hippocampus*. 2017. arXiv: 1704.07526. URL: https://arxiv.org/abs/1704.07526.

[15] R. P. Costa et al. *Cortical microcircuits as gated-recurrent neural networks*. 2017. arXiv: 1711.02448. URL: https://arxiv.org/abs/1711.02448.

[16] C. J. Cueva, A. Ardalan, M. Tsodyks, and N. Qian. *Recurrent neural network models for working memory of continuous variables: activity manifolds, connectivity patterns, and dynamic codes*. 2021. arXiv: 2111.01275. URL: https://arxiv.org/abs/2111.01275.

[17] L. D'Angelo, F. Denti, A. Canale, and M. Guindani. *Decoding Neuronal Ensembles from Spatially-Referenced Calcium Traces: A Bayesian Semiparametric Approach*. 2025. arXiv: 2508.09576. URL: https://arxiv.org/abs/2508.09576.

[18] B. Dunn, D. Wennberg, Z. Huang, and Y. Roudi. *Grid cells show field-to-field variability and this explains the aperiodic response of inhibitory interneurons*. 2017. arXiv: 1701.04893. URL: https://arxiv.org/abs/1701.04893.

[19] L. R. Duong et al. *Adaptive whitening in neural populations with gain-modulating interneurons*. 2023. arXiv: 2301.11955. URL: https://arxiv.org/abs/2301.11955.

[20] K. V. Efimov et al. *Temporary changes in large-scale memory neural networks after fear learning and extinction in healthy adults*. 2020. arXiv: 2011.00257. URL: https://arxiv.org/abs/2011.00257.

[21] A. T. Eitan et al. *Outan: An On-Head System for Driving micro-LED Arrays Implanted in Freely Moving Mice*. 2022. arXiv: 2211.13589. URL: https://arxiv.org/abs/2211.13589.

[22] S. Epelbaum et al. *Neural correlates of episodic memory in the Memento cohort*. 2018. arXiv: 1812.08031. URL: https://arxiv.org/abs/1812.08031.

[23] F. Faghihi, H. Samani, and A. A. Moustafa. *Pattern Separation in a Spiking Neural Network of Hippocampus Robust to Imbalanced Excitation/Inhibition*. 2018. arXiv: 1808.00367. URL: https://arxiv.org/abs/1808.00367.

[24] A. Fiasconaro and M. Migliore. *Effects of synapse location, delay and background stochastic activity on synchronising hippocampal CA1 neurons*. 2024. arXiv: 2409.10460. URL: https://arxiv.org/abs/2409.10460.

[25] A. Fiasconaro and M. Migliore. *Hippocampal synchronization in a realistic CA1 neuron model*. 2024. arXiv: 2409.10431. URL: https://arxiv.org/abs/2409.10431.

[26] K. Fidelin et al. *State-Dependent Modulation of Locomotion by GABAergic Spinal Sensory Neurons*. 2018. arXiv: 1812.00625. URL: https://arxiv.org/abs/1812.00625.

[27] Z. Fountas et al. *Human-inspired Episodic Memory for Infinite Context LLMs*. 2024. arXiv: 2407.09450. URL: https://arxiv.org/abs/2407.09450.

[28] E. P. Frady and F. T. Sommer. *Robust computation with rhythmic spike patterns*. 2019. arXiv: 1901.07718. URL: https://arxiv.org/abs/1901.07718.

[29] S. J. Gershman, I. Fiete, and K. Irie. *Key-value memory in the brain*. 2025. arXiv: 2501.02950. URL: https://arxiv.org/abs/2501.02950.

[30] C. Giotis et al. *Palimpsest Memories Stored in Memristive Synapses*. 2021. arXiv: 2109.13198. URL: https://arxiv.org/abs/2109.13198.

[31] M. Girardi-Schappo, L. Maler, and A. Longtin. *Optimal rate-variance coding due to firing threshold adaptation near criticality*. 2025. arXiv: 2509.04106. URL: https://arxiv.org/abs/2509.04106.

[32] D. Goto, H. M. L. Rios, M. Scholz, and S. Vaikuntanathan. *Neuromodulation-inspired gated associative memory networks:extended memory retrieval and emergent multistability*. 2025. arXiv: 2512.13859. URL: https://arxiv.org/abs/2512.13859.

[33] R. Grgurich and H. T. Blair. *An uncertainty principle for neural coding: Conjugate representations of position and velocity are mapped onto firing rates and co-firing rates of neural spike trains*. 2019. arXiv: 1912.11126. URL: https://arxiv.org/abs/1912.11126.

[34] B. J. Gutiérrez et al. *HippoRAG: Neurobiologically Inspired Long-Term Memory for Large Language Models*. 2024. arXiv: 2405.14831. URL: https://arxiv.org/abs/2405.14831.

[35] T. Haga and T. Fukai. *Extended temporal association memory by inhibitory Hebbian learning*. 2018. arXiv: 1809.05254. URL: https://arxiv.org/abs/1809.05254.

[36] S. Han et al. *Efficient Neural Network Approximation of Robust PCA for Automated Analysis of Calcium Imaging Data*. 2021. arXiv: 2108.01665. URL: https://arxiv.org/abs/2108.01665.

[37] P. J. Hellyer, B. Jachs, R. Leech, and C. Clopath. *Local inhibitory plasticity tunes global brain dynamics and allows the emergence of functional brain networks*. 2014. arXiv: 1412.6325. URL: https://arxiv.org/abs/1412.6325.

[38] L. Herron, P. Sartori, and B. Xue. *Robust retrieval of dynamic sequences through interaction modulation*. 2022. arXiv: 2211.17152. URL: https://arxiv.org/abs/2211.17152.

[39] Y. Hou, H. Tamoto, Q. Zhao, and H. Miyashita. *SynapticRAG: Enhancing Temporal Memory Retrieval in Large Language Models through Synaptic Mechanisms*. 2024. arXiv: 2410.13553. URL: https://arxiv.org/abs/2410.13553.

[40] S. Ishikawa, M. Yamada, H. Bao, and Y. Takezawa. *PhiNets: Brain-inspired Non-contrastive Learning Based on Temporal Prediction Hypothesis*. 2024. arXiv: 2405.14650. URL: https://arxiv.org/abs/2405.14650.

[41] Z. Jonke, R. Legenstein, S. Habenschuss, and W. Maass. *Feedback inhibition shapes emergent computational properties of cortical microcircuit motifs*. 2017. arXiv: 1705.07614. URL: https://arxiv.org/abs/1705.07614.

[42] J. Jude and M. H. Hennig. *Hippocampal representations emerge when training recurrent neural networks on a memory dependent maze navigation task*. 2020. arXiv: 2012.01328. URL: https://arxiv.org/abs/2012.01328.

[43] A. Karbasi, A. H. Salavati, and A. Shokrollahi. *Coupled Neural Associative Memories*. 2013. arXiv: 1301.1555. URL: https://arxiv.org/abs/1301.1555.

[44] A. Karbasi, A. H. Salavati, A. Shokrollahi, and L. R. Varshney. *Noise Facilitation in Associative Memories of Exponential Capacity*. 2014. arXiv: 1403.3305. URL: https://arxiv.org/abs/1403.3305.

[45] S.-Y. Kim and W. Lim. *Dynamical Origin for Winner-Take-All Competition in A Biological Network of The Hippocampal Dentate Gyrus*. 2021. arXiv: 2105.06057. URL: https://arxiv.org/abs/2105.06057.

[46] S.-Y. Kim and W. Lim. *Effect of Adult-Born Immature Granule Cells on Pattern Separation in The Hippocampal Dentate Gyrus*. 2023. arXiv: 2301.05387. URL: https://arxiv.org/abs/2301.05387.

[47] S. Knight. *Structural Correlates Of Spatial Navigation And Memory Formation*. 2022. arXiv: 2203.13434. URL: https://arxiv.org/abs/2203.13434.

[48] E. Köksal-Ersöz et al. *Neuronal mechanisms for sequential activation of memory items: dynamics and reliability*. 2019. arXiv: 1904.12133. URL: https://arxiv.org/abs/1904.12133.

[49] C. J. Kymn et al. *Binding in hippocampal-entorhinal circuits enables compositionality in cognitive maps*. 2024. arXiv: 2406.18808. URL: https://arxiv.org/abs/2406.18808.

[50] H. Lee. *Toward the biological model of the hippocampus as the successor representation agent*. 2020. arXiv: 2006.11975. URL: https://arxiv.org/abs/2006.11975.

[51] J. H. Lee, C. Koch, and S. Mihalas. *A Computational Analysis of the Function of Three Inhibitory Cell Types in Contextual Visual Processing*. 2016. arXiv: 1609.03622. URL: https://arxiv.org/abs/1609.03622.

[52] M. Lewis. *Hippocampal Spatial Mapping As Fast Graph Learning*. 2021. arXiv: 2107.00567. URL: https://arxiv.org/abs/2107.00567.

[53] T. Limbacher, O. Özdenizci, and R. Legenstein. *Memory-enriched computation and learning in spiking neural networks through Hebbian plasticity*. 2022. arXiv: 2205.11276. URL: https://arxiv.org/abs/2205.11276.

[54] X.-X. Lin, Y.-H. Yiu, and C. Leibold. *Emergence of Spatial Representation in an Actor-Critic Agent with Hippocampus-Inspired Sequence Generator*. 2025. arXiv: 2510.09951. URL: https://arxiv.org/abs/2510.09951.

[55] Y. Lin et al. *HippoMM: Hippocampal-inspired Multimodal Memory for Long Audiovisual Event Understanding*. 2025. arXiv: 2504.10739. URL: https://arxiv.org/abs/2504.10739.

[56] D. Lipshutz, C. Pehlevan, and D. B. Chklovskii. *Interneurons accelerate learning dynamics in recurrent neural networks for statistical adaptation*. 2022. arXiv: 2209.10634. URL: https://arxiv.org/abs/2209.10634.

[57] D. Lipshutz and E. P. Simoncelli. *Shaping the distribution of neural responses with interneurons in a recurrent circuit model*. 2024. arXiv: 2405.17745. URL: https://arxiv.org/abs/2405.17745.

[58] X. Liu and A. G. Richardson. *A System-on-Chip for Closed-loop Optogenetic Sleep Modulation*. 2021. arXiv: 2108.12261. URL: https://arxiv.org/abs/2108.12261.

[59] Y. Liu, Z. Wang, C. Qiao, and Z. Xu. *GATE: Adaptive Learning with Working Memory by Information Gating in Multi-lamellar Hippocampal Formation*. 2025. arXiv: 2501.12615. URL: https://arxiv.org/abs/2501.12615.

[60] T. V. de Maele, B. Dhoedt, T. Verbelen, and G. Pezzulo. *Bridging Cognitive Maps: a Hierarchical Active Inference Model of Spatial Alternation Tasks and the Hippocampal-Prefrontal Circuit*. 2023. arXiv: 2308.11463. URL: https://arxiv.org/abs/2308.11463.

[61] A. Mahrach et al. *Mechanisms underlying the response of mouse cortical networks to optogenetic manipulation*. 2019. arXiv: 1907.00816. URL: https://arxiv.org/abs/1907.00816.

[62] O. Makarenkov, M. Bezaire, and M. Hasselmo. *Bifurcation of spiking oscillations from a center in resonate-and-fire neurons*. 2025. arXiv: 2510.13156. URL: https://arxiv.org/abs/2510.13156.

[63] N. Y. Masse, G. D. Grant, and D. J. Freedman. *Alleviating catastrophic forgetting using context-dependent gating and synaptic stabilization*. 2018. arXiv: 1802.01569. URL: https://arxiv.org/abs/1802.01569.

[64] J. Melchior et al. *A Hippocampus Model for Online One-Shot Storage of Pattern Sequences*. 2019. arXiv: 1905.12937. URL: https://arxiv.org/abs/1905.12937.

[65] R. Moazzezi. *Grid-like structure is optimal for path integration*. 2016. arXiv: 1606.01239. URL: https://arxiv.org/abs/1606.01239.

[66] E. B. Mokhtari, J. J. Lawrence, and E. F. Stone. *Effect of Neuromodulation of Short-Term Plasticity on Information Processing in Hippocampal Interneuron Synapses*. 2018. arXiv: 1804.02115. URL: https://arxiv.org/abs/1804.02115.

[67] R. Monasson and S. Rosay. *Cross-talk and transitions between multiple spatial maps in an attractor neural network model of the hippocampus: phase diagram (I)*. 2013. arXiv: 1304.1457. URL: https://arxiv.org/abs/1304.1457.

[68] M. M. Monsalve-Mercado and C. Leibold. *Hippocampal Spike-Timing Correlations Lead to Hexagonal Grid Fields*. 2017. arXiv: 1706.06914. URL: https://arxiv.org/abs/1706.06914.

[69] N. W. Mosheiff, H. Agmon, A. Moriel, and Y. Burak. *An Efficient Coding Theory for a Dynamic Trajectory Predicts non-Uniform Allocation of Grid Cells to Modules in the Entorhinal Cortex*. 2016. arXiv: 1601.02948. URL: https://arxiv.org/abs/1601.02948.

[70] W. Mou, Z. Wang, and L. Wang. *Stable Memory Allocation in the Hippocampus: Fundamental Limits and Neural Realization*. 2016. arXiv: 1612.04659. URL: https://arxiv.org/abs/1612.04659.

[71] L. B. Moyse and H. Berry. *A Coupled Neural Field Model for the Standard Consolidation Theory*. 2024. arXiv: 2404.02938. URL: https://arxiv.org/abs/2404.02938.

[72] N. Mudrik, R. Ly, O. Ruebel, and A. S. Charles. *CREIMBO: Cross-Regional Ensemble Interactions in Multi-view Brain Observations*. 2024. arXiv: 2405.17395. URL: https://arxiv.org/abs/2405.17395.

[73] S. Nadkarni, T. Bartol, T. Sejnowski, and H. Levine. *Spatial and Temporal Correlates of Vesicular Release at Hippocampal Synapses*. 2010. arXiv: 1004.2009. URL: https://arxiv.org/abs/1004.2009.

[74] S. Nakamura et al. *High fidelity optogenetic control of individual prefrontal cortical pyramidal neurons in vivo*. 2012. arXiv: 1208.2720. URL: https://arxiv.org/abs/1208.2720.

[75] O. Noked et al. *Bidirectional optogenetic control of inhibitory neurons in freely-moving mice*. 2022. arXiv: 2211.13591. URL: https://arxiv.org/abs/2211.13591.

[76] Z. Okray et al. *Multisensory learning recruits visual neurons into an olfactory memory engram*. 2026. arXiv: 2604.28007. URL: https://arxiv.org/abs/2604.28007.

[77] L. Papadopoulos et al. *Modulation of metastable ensemble dynamics explains the inverted-U relationship between tone discriminability and arousal in auditory cortex*. 2024. arXiv: 2404.03902. URL: https://arxiv.org/abs/2404.03902.

[78] R. Pascual-Marqui et al. *The cross-frequency mediation mechanism of intracortical information transactions*. 2017. arXiv: 1703.07654. URL: https://arxiv.org/abs/1703.07654.

[79] F. W. Pasini et al. *An algebraic approach to spike-time neural codes in the hippocampus*. 2022. arXiv: 2211.06692. URL: https://arxiv.org/abs/2211.06692.

[80] N. Pedreschi et al. *Dynamic core-periphery structure of information sharing networks in the entorhinal cortex and the hippocampus*. 2020. arXiv: 2001.06371. URL: https://arxiv.org/abs/2001.06371.

[81] U. Pereira-Obilinovic, J. Aljadeff, and N. Brunel. *Forgetting leads to chaos in attractor networks*. 2021. arXiv: 2112.00119. URL: https://arxiv.org/abs/2112.00119.

[82] A. Peyrache. *Querying hippocampal replay with subcortical inputs*. 2022. arXiv: 2205.02665. URL: https://arxiv.org/abs/2205.02665.

[83] S. K. Rajagopal and T. A. Polk. *ConFER: A Neurally Constrained Computational Model of Context-Dependent Fear Extinction Recall and Relapse*. 2024. arXiv: 2411.08140. URL: https://arxiv.org/abs/2411.08140.

[84] J. E. Reser. *A Cognitive Architecture for Machine Consciousness and Artificial Superintelligence: Thought Is Structured by the Iterative Updating of Working Memory*. 2022. arXiv: 2203.17255. URL: https://arxiv.org/abs/2203.17255.

[85] J. P. Roach, L. M. Sander, and M. R. Zochowski. *Memory Recall and Spike Frequency Adaptation*. 2016. arXiv: 1601.02974. URL: https://arxiv.org/abs/1601.02974.

[86] J. Rossbroich and F. Zenke. *Dis-inhibitory neuronal circuits can control the sign of synaptic plasticity*. 2023. arXiv: 2310.19614. URL: https://arxiv.org/abs/2310.19614.

[87] J. Rothfuss et al. *Deep Episodic Memory: Encoding, Recalling, and Predicting Episodic Experiences for Robot Action Execution*. 2018. arXiv: 1801.04134. URL: https://arxiv.org/abs/1801.04134.

[88] Y. Roudi and P. E. Latham. *A balanced memory network*. 2007. arXiv: 0704.3005. URL: https://arxiv.org/abs/0704.3005.

[89] J. Russin et al. *A Neural Network Model of Continual Learning with Cognitive Control*. 2022. arXiv: 2202.04773. URL: https://arxiv.org/abs/2202.04773.

[90] U. Rutishauser, J.-J. Slotine, and R. J. Douglas. *Competition through selective inhibitory synchrony*. 2012. arXiv: 1201.2845. URL: https://arxiv.org/abs/1201.2845.

[91] T. Salvatori et al. *Associative Memories via Predictive Coding*. 2021. arXiv: 2109.08063. URL: https://arxiv.org/abs/2109.08063.

[92] G. di Sarra, S. Jha, and Y. Roudi. *The role of oscillations in grid cells' toroidal topology*. 2025. arXiv: 2501.19262. URL: https://arxiv.org/abs/2501.19262.

[93] D. M. Schwartz and O. O. Koyluoglu. *On the organization of grid and place cells: Neural de-noising via subspace learning*. 2017. arXiv: 1712.04602. URL: https://arxiv.org/abs/1712.04602.

[94] T. J. Sejnowski. *Dynamical Mechanisms for Coordinating Long-term Working Memory Based on the Precision of Spike-timing in Cortical Neurons*. 2025. arXiv: 2512.15891. URL: https://arxiv.org/abs/2512.15891.

[95] H. S. Seung. *Unsupervised learning by a nonlinear network with Hebbian excitatory and anti-Hebbian inhibitory neurons*. 2018. arXiv: 1812.11581. URL: https://arxiv.org/abs/1812.11581.

[96] J. Shang et al. *AI-native Memory: A Pathway from LLMs Towards AGI*. 2024. arXiv: 2406.18312. URL: https://arxiv.org/abs/2406.18312.

[97] S. Sharma, S. Chandra, and I. R. Fiete. *Content Addressable Memory Without Catastrophic Forgetting by Heteroassociation with a Fixed Scaffold*. 2022. arXiv: 2202.00159. URL: https://arxiv.org/abs/2202.00159.

[98] G. Soroka and M. Idiart. *Theta, alpha and gamma traveling waves in a multi-item working memory model*. 2021. arXiv: 2103.15266. URL: https://arxiv.org/abs/2103.15266.

[99] A. Speiser et al. *Fast amortized inference of neural activity from calcium imaging data with variational autoencoders*. 2017. arXiv: 1711.01846. URL: https://arxiv.org/abs/1711.01846.

[100] D. Szelogowski. *Engram Memory Encoding and Retrieval: A Neurocomputational Perspective*. 2025. arXiv: 2506.01659. URL: https://arxiv.org/abs/2506.01659.

[101] M. Tang, H. Barron, and R. Bogacz. *Sequential Memory with Temporal Predictive Coding*. 2023. arXiv: 2305.11982. URL: https://arxiv.org/abs/2305.11982.

[102] M. J. Tilley, M. Miller, and D. J. Freedman. *Artificial Neuronal Ensembles with Learned Context Dependent Gating*. 2023. arXiv: 2301.07187. URL: https://arxiv.org/abs/2301.07187.

[103] K. Vadovičová. *SWS promoting MHb-IPN-MRN circuit opposes the theta promoting circuit, active wake and REM sleep*. 2015. arXiv: 1510.01100. URL: https://arxiv.org/abs/1510.01100.

[104] K. Vadovičová. *Mechanisms for anesthesia, unawareness, respiratory depression, memory replay and sleep: MHb > IPN > PAG + DRN + MRN > claustrum > cortical slow waves*. 2025. arXiv: 2509.04454. URL: https://arxiv.org/abs/2509.04454.

[105] R. Vardi et al. *Neuronal Response Impedance Mechanism Implementing Cooperative Networks with Low Firing Rates and Microseconds Precision*. 2014. arXiv: 1411.5258. URL: https://arxiv.org/abs/1411.5258.

[106] Z. J. Waldman et al. *Ripple oscillations in the left temporal neocortex are associated with impaired verbal episodic memory encoding*. 2018. arXiv: 1807.10269. URL: https://arxiv.org/abs/1807.10269.

[107] R. Wang et al. *Anti-Retroactive Interference for Lifelong Learning*. 2022. arXiv: 2208.12967. URL: https://arxiv.org/abs/2208.12967.

[108] Y. Wang et al. *GAM-RAG: Gain-Adaptive Memory for Evolving Retrieval in Retrieval-Augmented Generation*. 2026. arXiv: 2603.01783. URL: https://arxiv.org/abs/2603.01783.

[109] Z. Wang, R. W. D. Tullio, S. Rooke, and V. Balasubramanian. *Time Makes Space: Emergence of Place Fields in Networks Encoding Temporally Continuous Sensory Experiences*. 2024. arXiv: 2408.05798. URL: https://arxiv.org/abs/2408.05798.

[110] Z. Wang et al. *REMI: Reconstructing Episodic Memory During Internally Driven Path Planning*. 2025. arXiv: 2507.02064. URL: https://arxiv.org/abs/2507.02064.

[111] M. T. Whelan, T. J. Prescott, and E. Vasilaki. *A Robotic Model of Hippocampal Reverse Replay for Reinforcement Learning*. 2021. arXiv: 2102.11914. URL: https://arxiv.org/abs/2102.11914.

[112] J. C. R. Whittington et al. *Generalisation of structural knowledge in the hippocampal-entorhinal system*. 2018. arXiv: 1805.09042. URL: https://arxiv.org/abs/1805.09042.

[113] J. Yoo and F. Wood. *BayesPCN: A Continually Learnable Predictive Coding Associative Memory*. 2022. arXiv: 2205.09930. URL: https://arxiv.org/abs/2205.09930.

[114] D. Yu, L. Wang, D. Kong, and H. Zhu. *Mapping the Genetic-Imaging-Clinical Pathway with Applications to Alzheimer's Disease*. 2020. arXiv: 2007.04558. URL: https://arxiv.org/abs/2007.04558.

[115] T. Zeng, X. Li, and B. Si. *Learning Sparse Spatial Codes for Cognitive Mapping Inspired by Entorhinal-Hippocampal Neurocircuit*. 2019. arXiv: 1910.04590. URL: https://arxiv.org/abs/1910.04590.

[116] F. Zenke and A. Laborieux. *Theories of synaptic memory consolidation and intelligent plasticity for continual learning*. 2024. arXiv: 2405.16922. URL: https://arxiv.org/abs/2405.16922.

[117] X. Zhang et al. *Episodic memory governs choices: An RNN-based reinforcement learning model for decision-making task*. 2021. arXiv: 2103.03679. URL: https://arxiv.org/abs/2103.03679.

[118] P. Zhou et al. *Efficient and accurate extraction of in vivo calcium signals from microendoscopic video data*. 2016. arXiv: 1605.07266. URL: https://arxiv.org/abs/1605.07266.