

v2 dynamic AF — diagnostics + v1 comparison


v2 fits : 235 (subjects: 30, ROIs: 8)
v1 fits : 240 (subjects: 30, ROIs: 8)

v2 shared parameters: σ_{AF} , g_{HP} , g_{LP} , g_{HP_dyn} , g_{LP_dyn}
v1 shared parameters: σ_{AF} , g_{HP} , g_{LP}



Sections:

1. v2 parameter distributions per ROI.
2. v2 HP-vs-LP tests (sustained AND dynamic differential).
3. Within-fit (g_{diff_sus} vs g_{diff_dyn}) scatter — does dynamic suppression covary with sustained suppression?
4. v1 vs v2 per-fit comparison: σ_{AF} , g_{HP} , g_{LP} . The v2 sustained estimates SHOULD be similar to v1 if the dynamic component is doing something independent.


v2 parameter distributions per ROI (y-clipped; ↑/↓ = #fits past limit)





v2 HP-vs-LP tests: sustained vs dynamic differential per ROI



HP-LP differential — sustained, dynamic, and total per ROI.
In V2/V3, sustained (negative) and dynamic (positive) cancel → total ~0. In V3AB/VO, they add → total strongly negative.




Is there a dynamic distractor effect? Wilcoxon test of $g_dyn_avg \neq 0$ per ROI.
 Green = significant CAPTURE (gain > 0), Red = significant SUPPRESSION (gain < 0).




Sustained vs dynamic HP-LP differential per fit
(do they covary? both should track HP-suppression if
they measure the same underlying attentional state.)


V1 $r_s=-0.03$, $p=0.857$




V2 $r_s=-0.16$, $p=0.413$




V3 $r_s=-0.26$, $p=0.161$




V3AB $r_s=-0.23$, $p=0.239$




hV4 $r_s=-0.28$, $p=0.146$




LO $r_s=+0.04$, $p=0.825$



TO $r_s=-0.02$, $p=0.915$



VO $r_s=-0.37$, $p=0.043^*$



v1 (static-AF only) vs v2 (with dynamic) sustained params — diagonal = identical fit

