



PRF shifts by HP-distractor location — V1


HP = upper right




HP = upper left



HP = lower left




HP = lower right




PRF shifts by HP-distractor location — V2


HP = upper right




HP = upper left



HP = lower left




HP = lower right




PRF shifts by HP-distractor location — V3


HP = upper right




HP = upper left



HP = lower left




HP = lower right




PRF shifts by HP-distractor location — V3AB


HP = upper right




HP = upper left



HP = lower left




HP = lower right




PRF shifts by HP-distractor location — hV4


HP = upper right




HP = upper left



HP = lower left




HP = lower right




PRF shifts by HP-distractor location — LO


HP = upper right




HP = upper left



HP = lower left




HP = lower right




PRF shifts by HP-distractor location — TO


HP = upper right




HP = upper left



HP = lower left




HP = lower right




PRF shifts by HP-distractor location — VO


HP = upper right




HP = upper left




HP = lower left




HP = lower right




PRF shifts in HP-aligned frame (HP at top, all conditions pooled)




Rotated-coordinate radial shifts by ROI




Radial PRF shift as a function of distance from HP distractor
(faded = 1° bins $\pm$ 95% CI; bold = LOWESS smooth)




Mean PRF shift for voxels within 1.5° of HP distractor
(per-subject means; error = SEM; FDR-corrected)





Mean PRF shift by ROI, stratified by mean PRF distance from HP distractor
(per-subject means; error = 95% CI)




Radial shift vs PRF size, voxels with mean PRF within 2.0° of HP
(faded = 1° bins $\pm$ 95% CI; bold = LOWESS) — collapse $\rightarrow$ RF-size driven; separation $\rightarrow$ structural




HP-anchored vs non-HP-anchored radial shift
(non-HP = three other ring quadrants; should be ~ 0 if effect is HP-specific)




HP vs non-HP shift as a function of distance from anchor
(non-HP = three other ring quadrants per condition)





Random-rotation permutation test
Gray dots = null (rotated HPs); red = observed; black ticks = 95% null interval




Per-anchor HP-vs-LP comparison (projection_away_from_anchor)
For each ring location, contrast voxel shift when that location is HP (red) vs LP (blue, mean of the three conditions where it's LP).




HP-anchored vs opposite-anchored radial shift
(under real suppression: HP > 0, Opposite < 0)




Per-subject means in V3AB + hV4
(n = 28; voxels with mean PRF within 1.5° of HP)



Random-rotation null in V3AB + hV4 combined
observed vs null ($p = 0.4900$, $n_{\text{perm}} = 499$)




Shift along HP-opposite axis vs signed position along axis
 Negative across whole range = pure suppression; sign flip = local suppression + global attraction




Shift along HP-opposite axis in V3AB + hV4 combined
(voxels with $|y_mean_rotated| < 3.0^\circ$; y-axis clipped to $\pm 0.5^\circ$)

Pure suppression: $y < 0$ everywhere · Mixed local-supp + global-attr: sign flip across x




Axial shift along HP-opposite axis in V3AB + hV4
(near-axis voxels: $|x_mean_rotated| < 1.5^\circ$, $|y_mean_rotated| < 3.0^\circ$; $n = 48,114$ rows)
Pure suppression: $y < 0$ everywhere · Mixed local-supp + global-attr: sign flip across x




Paired HP-vs-Opposite metric as a function of distance from HP
 bias-free; sign-flip at large d would mean local suppression + global attraction

HP shift - Opposite shift (deg)
 HP shift - Opposite shift (deg)
 (>0: suppression; <0: attraction)




Paired metric vs distance from HP in V3AB + hV4 combined


near ($d < 1.5^\circ$): mean = $+0.357^\circ$ ($t = +4.18$, $p = 0.0003$, *** suppression) · far ($d > 4.0^\circ$): mean = -0.022° ($t = -1.98$, $p = 0.0580$, n.s. attraction)




Paired metric vs distance from HP, per ROI
Red band = near ($d < 1.5^\circ$); blue band = far ($d > 4.0^\circ$); p-values from one-sample t against zero




Position-based shift along HP-opposite axis per ROI
 (near-axis voxels: $|x_mean_rotated| < 1.5^\circ$, $|y_mean_rotated| < 3.0^\circ$)
 Negative across whole range = pure suppression · sign flip across x = mixed local supp + global attr



Conditionwise sd and amplitude vs distance from HP, per ROI
 Top: AF predicts flat at 0; Bottom: AF (suppression) predicts < 0 near HP



AF fit AF centers — responders only (27/28) AF+ fit AF centers — responders only (27/28)
 ★ = HP ; × = opposite of HP ; ◇ = group mean ★ = HP ; × = opposite of HP ; ◇ = group mean






Model comparison (V3AB + hV4 combined, n = 28 subjects)

Within-voxel R^2 (cond variation only)
 AF responders ($R^2 > 0.01$): 27/28
 AF+ responders ($R^2 > 0.01$): 27/28

 AF responder σ_{AF} : 4.35° (median)
 AF+ responder σ_{AF} : 5.23° (median)


 AF+ $\log(b/a)$ responder: -3.00
 (smaller → closer to simple AF;
 larger → more local effect)

 AIC: AF+ wins in 7/28 subjects.
 Δ_{AIC} (AF - AF+) responders only:
 mean = -5.08




Per-condition AF center deviation from per-subject across-condition mean
Responders only (within-voxel $R^2 > 0.01$)


AF — condition-specific AF deviation
(faded ★ = direction of HP per condition; arrow = group-mean dev $\pm$ SEM)



AF+ — condition-specific AF deviation
(faded ★ = direction of HP per condition; arrow = group-mean dev $\pm$ SEM)



AF fit AF centers — responders only (28/28) AF+ fit AF centers — responders only (28/28)
 ★ = HP ; × = opposite of HP ; ◇ = group mean ★ = HP ; × = opposite of HP ; ◇ = group mean





Model comparison (V3AB + hV4 combined, n = 28 subjects)

Within-voxel R^2 (cond variation only)
 AF responders ($R^2 > 0.01$): 28/28
 AF+ responders ($R^2 > 0.01$): 28/28


 AF responder σ_{AF} : 4.03° (median)
 AF+ responder σ_{AF} : 4.74° (median)

 AF+ $\log(b/a)$ responder: -3.00
 (smaller → closer to simple AF;
 larger → more local effect)

 AIC: AF+ wins in 7/28 subjects.
 Δ_{AIC} (AF - AF+) responders only:
 mean = -7.78




Per-subject R^2 ; red lines mark responder cutoff (0.01)




Per-condition AF center deviation from per-subject across-condition mean
Responders only (within-voxel $R^2 > 0.01$)

AF — condition-specific AF deviation
(faded ★ = direction of HP per condition; arrow = group-mean dev $\pm$ SEM)




AF+ — condition-specific AF deviation
(faded ★ = direction of HP per condition; arrow = group-mean dev $\pm$ SEM)



Rotated-frame AF fit (free σ_{AF}) — HP at (0, +4°), antipode at (0, -4°)
 Suppression: μ_{AF} should land in lower half ($y < 0$). Red x = σ_{AF} saturated at bound.


V3AB (free σ_{AF}) — single AF center per subject

Group mean: (+0.09, +0.03)° (13/28 subj $y < 0$; 17 saturated) $t(y) = +0.39$, $p = 0.707$




hV4 (free σ_{AF}) — single AF center per subject

Group mean: (+0.01, -0.02)° (15/28 subj $y < 0$; 21 saturated) $t(y) = -0.30$, $p = 0.771$



Rotated-frame AF fit ($\sigma_{AF} = 4^\circ$ fixed) — HP at $(0, +4^\circ)$, antipode at $(0, -4^\circ)$
 Suppression: μ_{AF} should land in lower half ($y < 0$). Red $\times$ = σ_{AF} saturated at bound.



Robustness to R^2 threshold (default = 0.2)
 If V3AB/hV4 effects shift sign or vanish across thresholds $\rightarrow$ fragile

