

a
Target: odd orientation, report the dot, never at the singleton
Bar in 3.17° aperture
LP 10%
LP 10%
LP 10%
HP 50%
4°
1°
Singleton distractor (none on 20% of trials)
b
1 s
1.5 s
1 s
1.3 s
2.9 s
4.5 s
Fixation
Search
Feedback
Interval
The bar keeps sweeping, independent of the trials
HP location moves between blocks of 3 runs (12 runs, 2 sessions); example order
c
Visual areas
Polar angle
Eccentricity (°)
pRF size σ (°)
R²
Ventral
Dorsal
V1 V2 V3 hV4 V3AB
VO LO TO IPS FEF
RVF
LVF
0 2 4
0.0 1.5 3.0
0.0 0.3 0.6
d
pRF size σ (°)
pRF eccentricity (°)
V3AB
hV4
V3
V2
V1
e
Voxels with R² ≥ .1 (%)
V1 V2 V3 hV4 VO LO TO V3AB IPS
f
V1, 96% contralateral
V3, 95% contralateral
V3AB, 99% contralateral
IPS, 96% contralateral
1°
2°
3°
Left hemisphere
Right hemisphere
g
Held-out R²
Flexible HRF
Fixed HRF
Gaussian
DoG
DN