a
$N(t) = \frac{a\langle S \cdot M_c, G_c\rangle + b_c}{c|a|\langle S \cdot M_c, G_s\rangle + d_c} - b_c/d_c$
Input gain (m7): $M_c(g) = 1 + \sum_t w_{ct} A_t(g)$, $w = g_{HP}$ at HP, g_{LP} elsewhere
b-field: $b_c = b \exp(w_b F_c)$
d-field: $d_c = d \exp(w_d F_c)$, F_c = HP/LP field (σ_{AF}, θ) over the pRF
Input: field on the image, reaches the surround
b: baseline of the pRF centre
d: normalization constant
b
Pending: held-out comparison (run3, frozen pRFs)
dS = $g_{HP} - g_{LP}$
0.1
0.0
−0.1
c
Pending: recovery
Pending: recovery
Pending: recovery
d
Input (DN)
Input (DoG)
b-field
d-field
Predicted, bar on pRF centre
Predicted, bar in surround
Change (HP = −1)
0.5
0.0
−0.5
−1.0
HP
Next
Opp.
Items, early areas
Decoded bar (DN pRFs)
Measured
2
1
0
−1
e
b/d
b
d
HP − opposite quadrant (log ratio)
0.10
0.05
0.00
−0.05
V1
V2
V3
V3AB
IPS
hV4
VO
LO
TO
Early
Dorsal
Occipitotemporal