

a
The bar stimulus can be decoded from held-out runs
24 s
34 s
45 s
115 s
130 s
142 s
208 s
Bar
Best run sub-03, V1 r = .62
Mean of 30 subjects (V1)
b
Held-out run
Frames shuffled
Decoded vs true bar (r)
0.6
0.4
0.2
0.0
V1
V2
V3
V3AB
IPS
hV4
VO
LO
TO
Early
Dorsal
Occipitotemporal
c
The bar is decoded at lower intensity in the quadrant facing the HP location
V1–V3 pooled
HP
Less
More
–3%
0
3%
d
Decoded intensity vs mean of quadrants (%)
5
0
–5
V1–V3 pooled
HP
Next to HP
Opposite
e
Pre-specified
Exploratory
Decoded intensity, HP – other quadrants
.005
0
–.005
V1
V2
V3
V3AB
IPS
hV4
VO
LO
TO
Early
Dorsal
Occipitotemporal
f
Not driven by the items outside the aperture, and seen in the measured response
HP item
HP – other quadrants (%)
0–1.5°
1.5–2.5°
2.5°+
V1–V3 pooled
Not larger near the items
Aperture edge (items beyond, at 4°)
2
0
–2
–4
0
1
2
3
Eccentricity (°)
g
pRF
Bar crosses pRF
0
10 s
V1–V3: each location minus the mean of the four
HP
Next to HP
Opposite
Δ response vs mean of 4 locations (% signal)
.05
0
–.05
0
5
10
Time from crossing (s)