

a
The bar stimulus can be decoded from held-out runs
24 s
34 s
45 s
115 s
130 s
142 s
208 s
Bar
Best run
sub-03, V1
r = .62
Mean of 28
subjects (V1)
b
Decoded vs true bar (r)
0.6
0.4
0.2
0.0
Held-out run
Frames shuffled
V1
V2
V3
V3AB
IPS
hV4
VO
LO
TO
Early
Dorsal
Occipitotemporal
c
The bar is decoded at lower intensity in the quadrant facing the HP location
Data
Gain-field model
HP
V1–V3 pooled
Less
More
–3%
0
3%
Segment difference (%)
Gain-field model
Data
Next to HP
Opposite
HP quadrant
–2.7
–1.8
–0.9
0
+0.9
+1.8
+2.7
Bar centre (°)
Next to HP + opposite
HP + next to HP
d
Weaker than the quadrants next to it, and not driven by the items outside the aperture
Decoded intensity vs mean of quadrants (%)
V1–V3 pooled
Gain-field model
HP
Next to HP
Opposite
e
HP – other quadrants (%)
0–1.5°
1.5–2.5°
2.5°+
V1–V3 pooled
Gain-field model
Not larger near the items
Aperture edge (items beyond, at 4°)
Eccentricity (°)
f
Pooled over areas also beyond V1–V3 (occipitotemporal); per area reliable in V2 and V3
Pooled
** p = .003
* p = .010
p = .561
** p = .006
Per area
* q = .033
HP – other (%)
All areas
Early
Dorsal
Occipito-temporal
V1
V2
V3
V3AB
IPS
hV4
VO
LO
TO
Early
Dorsal
Occipitotemporal