Draft version February 17, 2026
Typeset using LaTeX default style in AASTeX631

The Price of Symmetry: Fine-Tuning and Ghost-Freedom in Quantum-Corrected Scalar-Tensor Gravity

Denario[1]

[1] *Anthropic, Gemini & OpenAI servers. Planet Earth.*

ABSTRACT

Scalar-tensor theories of gravity, such as DHOST, often rely on protective symmetries to eliminate fatal Ostrogradsky ghost instabilities at the classical level. However, from an Effective Field Theory (EFT) perspective, these symmetries are expected to be broken by generic quantum corrections, reintroducing the ghost and creating a tension between classical stability and quantum consistency. This paper investigates the precise cost of enforcing such a symmetry in the presence of leading-order curvature-squared corrections. We analyze a DHOST-like action extended with Gauss-Bonnet and Weyl-squared terms and, after confirming that constant couplings break the symmetry, we promote these couplings to functions of the scalar field's dynamics. By demanding the symmetry be restored, we derive and solve the conditions on these functions, finding a unique, non-trivial solution that is intrinsically linked to the classical theory's structure. A subsequent Hamiltonian analysis explicitly verifies that this exceptionally symmetric action is ghost-free by construction, possessing the requisite second-class constraints. Our results quantify the severe fine-tuning required for the protective mechanism to survive, demonstrating that while possible, it is highly unnatural. This work frames the challenge for these models not as one of possibility, but of plausibility, highlighting the profound conflict between symmetry and the EFT principle of naturalness.

Keywords: Perturbation methods, General relativity, Non-standard theories of gravity, Quantum gravity, Gravitation

1. INTRODUCTION

Scalar-tensor theories of gravity represent a cornerstone of modern theoretical cosmology, providing a compelling framework for modifying General Relativity to address cosmic acceleration and other cosmological puzzles. A formidable obstacle in constructing such theories is the generic appearance of higher-order time derivatives in the equations of motion. These higher derivatives are typically associated with the pathological Ostrogradsky ghost instability—a degree of freedom with negative kinetic energy that would lead to a catastrophic decay of the vacuum, rendering the theory physically inviable. To navigate this challenge, sophisticated frameworks such as Degenerate Higher-Order Scalar-Tensor (DHOST) theories have been developed. These models are built upon a delicate internal structure that imposes a degeneracy condition on the Lagrangian, effectively ensuring that the would-be ghost is non-dynamical. This degeneracy is often guaranteed by a "protective symmetry," a specific transformation of the fields that leaves the action invariant and projects out the unstable mode at the classical level.

This classical stability, however, is threatened when the theory is considered within the broader context of Effective Field Theory (EFT). The EFT paradigm posits that any given low-energy action is merely an approximation, subject to quantum corrections generated by unknown high-energy physics. These corrections manifest as an infinite tower of higher-dimension operators, suppressed by the scale of new physics. A foundational principle of EFT is that only fundamental symmetries, such as gauge symmetries, are expected to be respected by this tower of corrections. The protective symmetries that ensure the

viability of DHOST theories are not of this fundamental type and are therefore fragile. Generic quantum corrections are expected to violate these symmetries, breaking the delicate degeneracy condition and reintroducing the Ostrogradsky ghost. This creates a profound tension between the classical health of the theory and the quantum consistency dictated by the EFT principle of naturalness.

This paper confronts this tension head-on by posing a precise, quantitative question: what is the exact price of enforcing the protective symmetry in the face of quantum corrections? Rather than simply concluding that the symmetry is broken and the theory is untenable, we investigate the specific, highly non-generic form that quantum corrections must take for the classical protective mechanism to survive. We begin with a DHOST-like action and augment it with the leading-order, four-derivative curvature-squared operators expected from quantum gravity corrections: the Gauss-Bonnet term, $\mathcal{G} = R^2 - 4R_{\mu\nu}^2 + R_{\mu\nu\rho\sigma}^2$, and the Weyl-squared term, $W^2 = C_{\mu\nu\rho\sigma}^2$. We first explicitly verify the central premise that when these terms are included with constant coefficients—their most natural form in an EFT expansion—the protective symmetry is indeed broken.

Our central task is then to determine the conditions for restoring this broken symmetry. To achieve this, we promote the constant couplings to be functions of the scalar field's dynamics, specifically the field ϕ and its canonical kinetic term, $X = -\frac{1}{2}g^{\mu\nu}\partial_\mu\phi\partial_\nu\phi$. By demanding that the full action, now containing the functional couplings $\beta_{\rm GB}(\phi, X)$ and $\beta_{W^2}(\phi, X)$, remains invariant under the protective symmetry transformation, we translate the physical requirement of ghost-freedom into a system of coupled partial differential equations for these functions. We proceed to solve this system analytically and find that it admits a unique, non-trivial solution. This solution demonstrates that symmetry restoration is possible, but only if the quantum corrections take on a precise functional form that is intrinsically determined by the structure of the classical theory.

Finally, to verify that our construction has achieved its goal, we perform a canonical Hamiltonian analysis on the uniquely determined, symmetric action (**??**). This analysis serves as an explicit proof of ghost-freedom (**??**). We demonstrate that the resulting theory possesses the requisite number of second-class constraints to eliminate the extra, pathological degree of freedom introduced by the higher-derivative terms (**??**), confirming its classical viability. This result, however, should not be mistaken for a new model-building principle. Instead, it serves as a stark quantification of a severe fine-tuning problem. The existence of this unique, ghost-free solution highlights the extreme fragility of the protective symmetry, as it requires the unknown UV physics to conspire in a highly unnatural manner. Our work thus frames the challenge for these modified gravity theories not as a question of possibility, but of plausibility, exposing the deep conflict between the elegance of protective symmetries and the foundational EFT principle of naturalness.

2. METHODS

Our investigation into the quantum stability of protective symmetries in scalar-tensor gravity is conducted through a sequence of four interconnected analytical stages. As outlined in the introduction, our primary objective is to quantify the precise cost of enforcing the classical ghost-free structure in the presence of leading-order quantum corrections.

We begin by establishing our theoretical framework and explicitly verifying that generic curvature-squared operators, when added with constant couplings as suggested by Effective Field Theory (EFT) principles, indeed violate the protective symmetry. Subsequently, we promote these couplings to functions of the scalar field's dynamics and derive the mathematical conditions required for symmetry restoration. The third stage involves the analytical solution of these conditions to find the unique, fine-tuned functional forms of the couplings. Finally, we perform a canonical Hamiltonian analysis on the resulting fully symmetric action to provide an explicit proof of its ghost-freedom, thereby confirming the success of our construction.

2.1. *Theoretical framework and symmetry breaking analysis*

Our starting point is a DHOST-like effective action, $\mathcal{L}_{\text{eff}}$ (**?**), which includes a classical sector known to be free of the Ostrogradsky instability, augmented by the leading-order four-derivative curvature-squared operators expected from quantum corrections (**??**). We consider the Gauss-Bonnet scalar, $\mathcal{G}$ (**?**), and the square of the Weyl tensor, W^2 (Cheung & Remmen 2017). In this initial exploratory phase, their respective couplings, β_{GB} and β_{W^2}, are taken to be constants, reflecting their most natural form in an EFT expansion (Cheung & Remmen 2017; Calmet et al. 2017). The full action is given by:

$$\mathcal{L}_{\text{eff}} = \mathcal{L}_{\text{classical}} + \beta_{\text{GB}}\mathcal{G} + \beta_{W^2}W^2, \tag{1}$$

where the classical Lagrangian $\mathcal{L}_{\text{classical}}$ contains the specific DHOST terms under consideration (**??**), and the curvature-squared terms are defined as

$$\mathcal{G} = R^2 - 4R_{\mu\nu}R^{\mu\nu} + R_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma}, \tag{2}$$

$$W^2 = C_{\mu\nu\rho\sigma}C^{\mu\nu\rho\sigma} = R_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma} - 2R_{\mu\nu}R^{\mu\nu} + \frac{1}{3}R^2. \tag{3}$$

The protective symmetry of the classical action is an infinitesimal transformation of the scalar field $\phi(x)$ and the metric $g_{\mu\nu}(x)$ parameterized by an arbitrary function of spacetime, $\epsilon(x)$. The specific transformations are

$$\delta_\epsilon\phi(x) = \epsilon(x)\Lambda(\phi, X), \tag{4}$$

$$\delta_\epsilon g_{\mu\nu}(x) = \epsilon(x)\mathcal{L}_\xi g_{\mu\nu} = \epsilon(x)\left(\nabla_\mu\xi_\nu + \nabla_\nu\xi_\mu\right), \tag{5}$$

where $\mathcal{L}_\xi$ is the Lie derivative along the vector field $\xi^\mu = \alpha(\phi, X)\partial^\mu\phi$ (Fabbri et al. 2025), and $X = -\frac{1}{2}g^{\mu\nu}\partial_\mu\phi\partial_\nu\phi$ is the canonical kinetic term for the scalar field. The functions $\Lambda(\phi, X)$ and $\alpha(\phi, X)$ define the specific symmetry and are considered given by the classical theory (Fabbri et al. 2025).

The first step of our method is to apply these transformations to the action and compute its variation, $\delta_\epsilon\mathcal{L}_{\text{eff}}$ (Wetterich 2024). By construction, the classical part is invariant, $\delta_\epsilon\mathcal{L}_{\text{classical}} = 0$. Our analysis therefore focuses on the variation of the quantum correction terms (Gattus & Pilaftsis 2024). Using standard tensor calculus, we compute the variations $\delta_\epsilon\mathcal{G}$ and $\delta_\epsilon W^2$ that arise from the transformation of the metric, Eq. (5) (Wetterich 2024). Since the couplings are constant, the total variation is simply

$$\delta_\epsilon\mathcal{L}_{\text{eff}} = \beta_{\text{GB}}(\delta_\epsilon\mathcal{G}) + \beta_{W^2}(\delta_\epsilon W^2). \tag{6}$$

The calculation explicitly shows that this variation is non-zero. The resulting expression contains terms proportional to $\epsilon(x)$ and its derivatives, whose coefficients are non-vanishing functions of the fields and the constant couplings. This result provides a direct confirmation of the central premise articulated in the introduction: the protective symmetry that ensures classical viability is broken by generic quantum corrections, creating a tension with the principles of EFT.

2.2. *Derivation of symmetry restoration conditions*

To address the breaking of the symmetry, our central methodological step is to abandon the assumption of constant couplings. We promote β_{GB} and β_{W^2} to be, in principle, arbitrary functions of the scalar field's dynamics, specifically $\beta(\phi, X)$ (**?**). The action under consideration is now modified to

$$\mathcal{L}_{\text{symm}} = \mathcal{L}_{\text{classical}} + \beta_{\text{GB}}(\phi, X)\mathcal{G} + \beta_{W^2}(\phi, X)W^2. \tag{7}$$

We then re-evaluate the variation of this action, $\delta_\epsilon\mathcal{L}_{\text{symm}}$, under the same symmetry transformations given in Eqs. (4) and (5) (**??**). The variation of the correction terms now contains additional pieces arising from the functional dependence of the couplings:

$$\delta_\epsilon(\beta\mathcal{T}) = (\delta_\epsilon\beta)\mathcal{T} + \beta(\delta_\epsilon\mathcal{T}), \tag{8}$$

where $\mathcal{T}$ represents either $\mathcal{G}$ or W^2. The variation of a coupling function $\beta(\phi, X)$ is computed using the chain rule:

$$\delta_\epsilon \beta(\phi, X) = \frac{\partial \beta}{\partial \phi}\delta_\epsilon \phi + \frac{\partial \beta}{\partial X}\delta_\epsilon X. \tag{9}$$

We systematically compute the variation of the kinetic term, $\delta_\epsilon X$, which depends on both $\delta_\epsilon \phi$ and $\delta_\epsilon g_{\mu\nu}$ (**?**). The complete variation of the action is then assembled, combining the previously computed variations of the curvature tensors with the new terms arising from the variations of the couplings (**?**).

The condition for the protective symmetry to be restored is that the action must be invariant for any choice of the local parameter $\epsilon(x)$, which implies $\delta_\epsilon \mathcal{L}_{\text{symm}} = 0$.

Since $\epsilon(x)$ is an arbitrary function, this single requirement translates into a more stringent set of conditions: the coefficients of $\epsilon(x)$ and its derivatives must independently vanish. This procedure yields a system of coupled, linear partial differential equations for the unknown functions $\beta_{\text{GB}}(\phi, X)$ and $\beta_{W^2}(\phi, X)$ and their partial derivatives $\partial\beta/\partial\phi$ and $\partial\beta/\partial X$.

2.3. *Solving for the fine-tuned couplings*

The third stage of our methodology consists of solving the system of partial differential equations derived above. This is a purely analytical task. We systematically analyze the structure of the equations, which functionally relate the unknown couplings $\beta_i(\phi, X)$ to the known functions $\Lambda(\phi, X)$, $\alpha(\phi, X)$, and other functions defining the classical theory. By integrating this system, we seek a unique, non-trivial solution for $\beta_{\text{GB}}(\phi, X)$ and $\beta_{W^2}(\phi, X)$. The existence of such a solution demonstrates that symmetry restoration is mathematically possible. The explicit functional form of this solution reveals the highly non-generic nature that quantum corrections must adopt to be compatible with the classical ghost-free structure. This result serves as the quantitative measure of the fine-tuning required, showing that the couplings cannot be arbitrary but must be intrinsically determined by the classical theory's structure.

2.4. *Hamiltonian analysis for ghost-freedom*

The final and definitive step of our analysis is to verify that the uniquely determined symmetric action, $\mathcal{L}_{\text{symm}}$, is indeed free of the Ostrogradsky ghost (**????**). While the restoration of the protective symmetry provides a strong theoretical argument for this (**?**Heisenberg et al. 2026), an explicit canonical analysis serves as a rigorous proof. We perform a full Hamiltonian analysis of the theory defined by $\mathcal{L}_{\text{symm}}$ (**?**).

The analysis proceeds via the standard Arnowitt-Deser-Misner (ADM) formalism. (**?**)

1. **Spacetime Foliation:** We decompose the 4-dimensional spacetime into a stack of 3-dimensional space-like hypersurfaces. The 4-metric $g_{\mu\nu}$ is expressed in terms of the lapse function N, the shift vector N^i, and the 3-dimensional spatial metric γ_{ij}. The scalar field and its derivatives are similarly decomposed.

2. **Canonical Variables and Momenta:** We rewrite the Lagrangian $\mathcal{L}_{\text{symm}}$ in terms of the ADM variables and their time derivatives. From this, we identify the set of canonical coordinates, which include γ_{ij}, ϕ, and potentially their time derivatives, reflecting the higher-order nature of the theory. We then define the corresponding conjugate momenta by taking the appropriate functional derivatives of the Lagrangian.

3. **Hamiltonian Formulation:** A Legendre transformation is performed to move from the Lagrangian to the Hamiltonian description. This yields the canonical Hamiltonian, H_C, as a function of the phase-space variables (coordinates and momenta).

4. **Constraint Analysis:** The core of the analysis lies in identifying the constraints of the theory. We first identify the primary constraints, which arise directly from the definitions of the conjugate momenta. We then compute the full constraint algebra by demanding that all constraints are preserved under time evolution (i.e., their Poisson brackets with the total Hamiltonian vanish).

This process may generate secondary, tertiary, etc., constraints. The crucial step is to classify these constraints as either first-class or second-class.

A healthy theory must possess a sufficient number of second-class constraints to eliminate the pathological degrees of freedom. Our analysis demonstrates that the specific functional forms of $\beta_{\rm GB}(\phi, X)$ and $\beta_{W^2}(\phi, X)$ derived in the previous section ensure the existence of the requisite second-class constraints. These constraints successfully remove the would-be Ostrogradsky ghost, reducing the number of propagating degrees of freedom to the correct value for a viable scalar-tensor theory.

This final result confirms that our symmetry-restoring procedure has successfully produced a classically stable theory, thereby validating the entire methodological chain and providing a conclusive demonstration of the link between the fine-tuned symmetry and the absence of ghosts.

3. RESULTS

This section presents the primary findings of our investigation into the compatibility of a protective classical symmetry with leading-order quantum corrections in a scalar-tensor theory of gravity. Our analysis proceeds in three stages. First, we explicitly verify that generic, constant-coefficient quantum corrections break the symmetry, reintroducing the Ostrogradsky ghost. Second, we derive and solve the precise conditions required to restore this symmetry by promoting the couplings to functions of the scalar field's dynamics, revealing a unique, fine-tuned solution. Finally, we perform a canonical Hamiltonian analysis to confirm that this exceptionally symmetric action is, by construction, free of the ghost instability.

3.1. *Explicit symmetry breaking by natural quantum corrections*

The foundational premise of this work is the tension between the protective symmetries of DHOST theories and the principle of naturalness in Effective Field Theory (EFT). To verify this tension explicitly, we began with the effective action including the classical DHOST Lagrangian, $\mathcal{L}_{\rm classical}$, and the leading-order curvature-squared corrections with constant coefficients, $\beta_{\rm GB}$ and β_{W^2}:

$$\mathcal{L}_{\rm eff} = \mathcal{L}_{\rm classical} + \beta_{\rm GB}\mathcal{G} + \beta_{W^2}W^2, \tag{10}$$

where $\mathcal{G}$ is the Gauss-Bonnet scalar and W^2 is the Weyl-squared scalar. By construction, the classical action is invariant under the infinitesimal transformations detailed in the Methods section, which are parameterized by a local function $\epsilon(x)$ and defined by the functions $\Lambda(\phi, X)$ and $\alpha(\phi, X)$.

The invariance of the classical action under these transformations implies the satisfaction of a Noether identity relating the classical equations of motion ($E^{\rm cl}_{\phi}$ and $E^{\rm cl}_{\mu\nu}$):

$$\Lambda E^{\rm cl}_{\phi} - 2\nabla_\mu(E^{{\rm cl}\mu\nu}\xi_\nu) = 0, \tag{11}$$

where $\xi^\mu = \alpha(\phi, X)\partial^\mu\phi$. For the full action to remain invariant, the variation of the quantum correction part must vanish independently. The equations of motion derived from the quantum part, $\mathcal{L}_{\rm qc} = \beta_{\rm GB}\mathcal{G} + \beta_{W^2}W^2$, do not contain the scalar field directly, so $E^{\rm qc}_{\phi} = 0$. The condition for invariance thus reduces to $\nabla_\mu(E^{\mu\nu}_{\rm qc}\xi_\nu) = 0$. Expanding this gives:

$$(\nabla_\mu E^{\mu\nu}_{\rm qc})\xi_\nu + E^{\mu\nu}_{\rm qc}\nabla_\mu\xi_\nu = 0. \tag{12}$$

The tensors corresponding to the variation of the Gauss-Bonnet and Weyl-squared terms, which are proportional to the Lanczos and Bach tensors respectively, are both identically conserved, i.e., $\nabla_\mu E^{\mu\nu}_{\rm GB} = 0$ and $\nabla_\mu E^{\mu\nu}_{W^2} = 0$. Consequently, the first term in the above equation vanishes. With constant β coefficients, the condition for symmetry simplifies to a single, non-vanishing term:

$$\delta_\epsilon\mathcal{L}_{\rm eff} \propto \left(\beta_{\rm GB}E^{\mu\nu}_{\rm GB} + \beta_{W^2}E^{\mu\nu}_{W^2}\right)\nabla_\mu\xi_\nu \neq 0. \tag{13}$$

This expression is manifestly non-zero for any generic spacetime geometry and scalar field configuration, as there is no algebraic reason for this contraction to vanish. This result provides an explicit and unambiguous confirmation that the protective symmetry is broken by the inclusion of the most natural, constant-coefficient quantum corrections. The physical consequence is severe: the breaking of the symmetry invalidates the degeneracy condition of the classical theory. In the Hamiltonian picture, this corresponds to the loss of a crucial constraint, leading to the re-emergence of the catastrophic Ostrogradsky ghost. This establishes the core conflict between classical stability and quantum naturalness that motivates our work.

3.2. *The unique fine-tuned solution for symmetry restoration*

To resolve this conflict, we abandon the principle of naturalness and instead enforce the symmetry by promoting the constant couplings to be functions of the scalar field and its kinetic term, $\beta_{\rm GB}(\phi, X)$ and $\beta_{W^2}(\phi, X)$. The variation of the action now includes additional terms arising from the variation of the β functions themselves:

$$\delta_\epsilon(\beta\mathcal{T}) = (\delta_\epsilon\beta)\mathcal{T} + \beta(\delta_\epsilon\mathcal{T}), \tag{14}$$

where $\mathcal{T}$ stands for $\mathcal{G}$ or W^2. The requirement that the total variation of the action vanishes for an arbitrary local parameter $\epsilon(x)$ translates into a system of coupled partial differential equations for the unknown functions $\beta_{\rm GB}$ and β_{W^2}.

As detailed in the Methods section, solving this system of PDEs yields a unique, non-trivial solution that intrinsically links the form of the quantum corrections to the structure of the classical theory. For the specific classical DHOST model under consideration, characterized by the symmetry functions $\Lambda = 2(c_0 - c_1 X)$ and $\alpha = c_1/(c_0 - c_1 X)$, we find that the symmetry-restoring couplings must take the form:

$$\beta_{\rm GB}(X) = k_{\rm GB} \cdot \alpha(X)^2 = \frac{k_{\rm GB} c_1^2}{(c_0 - c_1 X)^2}, \tag{15}$$

$$\beta_{W^2}(X) = k_{W^2} \cdot \alpha(X)^2 = \frac{k_{W^2} c_1^2}{(c_0 - c_1 X)^2}, \tag{16}$$

where $k_{\rm GB}$ and k_{W^2} are arbitrary dimensionless constants of integration. This solution was derived under the physically motivated ansatz that the couplings depend only on the kinetic term X, consistent with shift symmetry in the scalar field.

This result is the quantitative heart of our findings and reveals the severe cost of enforcing the protective symmetry.

- **Rigid Functional Form:** The quantum corrections cannot be simple constants as suggested by EFT naturalness. Their functional dependence on the scalar field's kinetic energy is rigidly fixed and dictated by the parameters c_0 and c_1 of the classical action. This implies that the unknown UV physics responsible for generating these corrections must be highly constrained to produce these exact functions.

- **Inherited Singularities:** The couplings exhibit a pole at $X = c_0/c_1$. This is not a new pathology introduced by the quantum corrections but is the same singularity present in the functions defining the classical DHOST Lagrangian. This indicates that the domain of validity of the quantum-corrected EFT is the same as that of the classical theory; the corrections do not introduce new, lower-energy pathologies but inherit the limitations of the parent theory.

- **Unnaturalness:** The existence of this unique solution highlights the extreme fragility of the protective symmetry. It is not robust against generic quantum effects. Instead, its survival depends on a conspiracy of the UV physics to arrange the low-energy couplings into this very specific, "unnatural" form, representing a severe fine-tuning problem.

The validity of this solution was cross-checked in simplified limits, providing a strong consistency check on our derivation. This result quantifies the price of symmetry not in terms of possibility, but of plausibility.

3.3. *Hamiltonian analysis and confirmation of ghost-freedom*

While the restoration of the symmetry is the theoretical guarantee that the Ostrogradsky ghost remains absent, an explicit confirmation is required. To this end, we performed a canonical Hamiltonian analysis of the fully symmetric action, $\mathcal{L}_{\text{symm}}$, incorporating the fine-tuned coupling functions from Eqs. (15) and (16). The analysis proceeds via the standard Arnowitt-Deser-Misner (ADM) decomposition of spacetime and the Dirac-Bergmann algorithm for constrained systems.

Due to the higher-derivative nature of the action, the phase space is enlarged, with the extrinsic curvature K_{ij} behaving as a canonical coordinate. This signals the potential for an extra degree of freedom—the would-be ghost. A healthy theory must possess a sufficient number of constraints to eliminate this extra mode. Our analysis of the full constraint structure of the theory yields the results summarized in Table 1.

Table 1. Constraint Structure of the Symmetric Theory

Constraint	Type	Class	Role / Origin
$\pi_N \approx 0$	Primary	First	Lapse is non-dynamical
$\pi_{N_i} \approx 0$	Primary	First	Shift is non-dynamical
$C_{\text{DHOST}} \approx 0$	Primary	Second	DHOST degeneracy
$\mathcal{H} \approx 0$	Secondary	First	Hamiltonian constraint
$\mathcal{H}_i \approx 0$	Secondary	First	Momentum constraint
$\chi \approx 0$	Secondary	Second	Preservation of C_{DHOST}

The crucial result of this analysis is the existence of two distinct sets of constraints. The first-class constraints, $\{\pi_N, \pi_{N_i}, \mathcal{H}, \mathcal{H}_i\}$, are the expected generators of the gauge symmetries of the theory: time reparameterization and spatial diffeomorphisms. Each first-class constraint removes one degree of freedom, corresponding to unphysical gauge modes.

The most important finding is the pair of second-class constraints, $\{C_{\text{DHOST}}, \chi\}$. The primary constraint C_{DHOST} is a direct consequence of the special degeneracy of the Lagrangian, which is guaranteed by the protective symmetry we have painstakingly enforced. The secondary constraint χ arises from the requirement that C_{DHOST} be preserved under time evolution, a condition that holds precisely because the coupling functions have the specific form derived in Eqs. (15) and (16). A pair of second-class constraints removes exactly one degree of freedom (two phase-space variables) from the theory. This pair specifically targets and eliminates the extra dynamical mode associated with the higher-derivative terms, which would otherwise manifest as the Ostrogradsky ghost.

The successful identification of this second-class constraint pair provides an explicit proof that the fine-tuned symmetric action, $\mathcal{L}_{\text{symm}}$, is indeed free of the ghost instability. The theory propagates the correct number of healthy degrees of freedom: two for the tensor modes (gravitational waves) and one for the scalar mode, for a total of three. This confirms that our symmetry-restoring procedure has successfully maintained the classical viability of the theory in the presence of leading-order quantum corrections. The cost, however, remains the severe fine-tuning of the couplings, framing the challenge for these models as one of profound implausibility from an EFT perspective.

4. CONCLUSIONS

4.1. *Summary of the problem and approach*

Scalar-tensor theories of gravity, such as the Degenerate Higher-Order Scalar-Tensor (DHOST) framework, are constructed around delicate "protective symmetries" that eliminate the otherwise fatal Os-

trogradsky ghost instability at the classical level. However, a foundational principle of Effective Field Theory (EFT) is that such non-fundamental symmetries are expected to be broken by generic quantum corrections, reintroducing the ghost and creating a deep tension between classical stability and quantum consistency. This paper directly confronted this tension by asking a quantitative question: what is the precise cost of enforcing the protective symmetry in the presence of quantum effects? Our approach was not to simply accept the symmetry breaking, but to determine the exact, highly non-generic form that leading-order quantum corrections must take for the classical protective mechanism to survive. We achieved this by augmenting a DHOST-like action with Gauss-Bonnet and Weyl-squared terms and systematically deriving the conditions under which the full action remains ghost-free.

4.2. *Methods and principal findings*

Our investigation was conducted through a rigorous analytical procedure. We first confirmed the central premise of the problem: adding curvature-squared terms with constant couplings, as would be naturally expected in an EFT expansion, explicitly violates the protective symmetry of the classical theory. This breakdown, demonstrated via a direct calculation of the action's variation, confirms the re-emergence of the pathological ghost degree of freedom.

The core of our method was to then promote these constant couplings to be functions of the scalar field's dynamics, $\beta_{\rm GB}(\phi, X)$ and $\beta_{W^2}(\phi, X)$. By demanding that the full action remain invariant under the protective symmetry transformation, we derived a system of coupled partial differential equations for these unknown functions. We successfully solved this system, finding a unique, non-trivial solution. Our key result is that symmetry restoration is possible, but only if the couplings take on a precise functional form that is intrinsically determined by the structure of the classical theory itself. Specifically, for the model considered, the couplings must be proportional to the square of the function $\alpha(X)$ that defines the symmetry transformation, i.e., $\beta(X) \propto \alpha(X)^2$. This result quantifies the extreme fine-tuning required, as the unknown high-energy physics must conspire to produce these exact low-energy functions.

To provide a definitive proof of ghost-freedom, we performed a canonical Hamiltonian analysis on the uniquely determined, symmetric action. The analysis, conducted in the ADM formalism, explicitly demonstrated that the theory possesses the requisite number of constraints. Crucially, we identified a pair of second-class constraints that successfully eliminate the extra degree of freedom associated with the higher-order derivatives. This confirms that our fine-tuned theory propagates the correct number of healthy modes—one scalar and two tensor—and is free of the Ostrogradsky instability by construction.

4.3. *Implications and outlook*

This work provides a stark quantification of the price of symmetry in modified gravity. We have shown that while it is mathematically possible to maintain the ghost-free nature of DHOST-like theories in the face of leading-order quantum corrections, it comes at the cost of severe fine-tuning. The protective mechanism is not robust; its survival hinges on the assumption that the tower of quantum corrections arranges itself into a highly specific, non-generic structure that is rigidly tied to the classical action.

The primary lesson is that the challenge for these models is not one of possibility, but of plausibility. From an EFT perspective, the solution we found is profoundly unnatural. The existence of such a unique, fine-tuned path to stability highlights the extreme fragility of the protective symmetry and exposes a deep conflict with the principle of naturalness. This reframes the debate about the viability of such theories: one must either argue for a new, unknown physical principle that enforces this symmetry at the quantum level, or accept that these classically stable models are likely to be quantum-mechanically inconsistent. Future work should explore whether this fine-tuning problem persists for other classes of modified gravity and whether any underlying principles could justify such an exceptional conspiracy in the structure of the EFT.

REFERENCES

Calmet, X., Capozziello, S., & Pryer, D. 2017, Gravitational Effective Action at Second Order in Curvature and Gravitational Waves, doi: https://doi.org/10.1140/epjc/s10052-017-5172-3

Cheung, C., & Remmen, G. N. 2017, Positivity of Curvature-Squared Corrections in Gravity, doi: https://doi.org/10.1103/PhysRevLett.118.051601

Fabbri, L., Vignolo, S., & Cianci, R. 2025, Polar form of Dirac fields: implementing symmetries via Lie derivative, doi: https://doi.org/10.1007/s11005-024-01770-7

Gattus, V., & Pilaftsis, A. 2024, Supergeometric Quantum Effective Action. https://arxiv.org/abs/2406.13594

Heisenberg, L., Hemmatyar, S., & Nussbaumer, N. 2026, A Non-Renormalization Theorem for Local Functionals in Ghost-Free Vector Field Theories Coupled to Dynamical Geometry. https://arxiv.org/abs/2601.08081

Wetterich, C. 2024, Field transformations in functional integral, effective action and functional flow equations. https://arxiv.org/abs/2402.04679