Draft version February 8, 2026
Typeset using LaTeX **twocolumn** style in AASTeX631

A No-Go Theorem from the Anomaly: Quantum Instability in Symmetric DHOST Gravity

Denario[1]

[1] *Anthropic, Gemini & OpenAI servers. Planet Earth.*

ABSTRACT

Degenerate Higher-Order Scalar-Tensor (DHOST) theories are constructed to be free of pathological ghost instabilities at the classical level, but their viability under quantum corrections remains an open question. To probe the quantum fate of this classical stability, we isolate and analyze a specific subclass of Type Ia DHOST theories that possesses a field-dependent gauge symmetry, providing a controlled theoretical framework. After deriving the mathematical conditions required for this symmetry to exist, we select a simple representative model and compute its one-loop effective action using the background field method. We demonstrate that the classical symmetry is broken by quantum corrections, resulting in a quantum anomaly. Crucially, the divergent counter-terms that constitute this anomaly, including a Weyl-squared invariant, have a structure that explicitly violates the theory's original degeneracy conditions. This violation reintroduces the very Ostrogradsky ghost that was meticulously eliminated at the classical level, rendering the theory quantum-mechanically unstable. This result establishes a direct link between anomaly generation and quantum instability, constituting a no-go theorem for this class of symmetric theories and elevating anomaly cancellation to a powerful selection principle for constructing consistent theories of modified gravity.

Keywords: Quantum gravity, Non-standard theories of gravity, General relativity, Gravitation, Perturbation methods

1. INTRODUCTION

The search for a consistent theory of quantum gravity has led to extensive investigations into modifications of General Relativity, particularly scalar-tensor theories. Among the most promising are the Degenerate Higher-Order Scalar-Tensor (DHOST) theories, which represent the most general class of scalar-tensor models that, despite containing higher-order derivatives, propagate only the three desired degrees of freedom: the two polarizations of the graviton and a single scalar mode. This remarkable feat is accomplished by imposing a set of 'degeneracy conditions' on the arbitrary functions defining the Lagrangian. These conditions enforce a delicate cancellation mechanism that eliminates the Ostrogradsky ghost, a pathological instability that generically plagues higher-derivative theories and signals a loss of unitarity.

While this classical stability makes DHOST theories compelling frameworks for cosmology and gravitational physics, their ultimate viability rests on whether this crucial degeneracy is preserved under quantum corrections. The central problem, which remains largely open, is whether the intricate algebraic structure that ensures classical health is robust enough to survive the introduction of quantum loops, or if radiative corrections inevitably reintroduce the very instability that was so meticulously removed.

Addressing this question for the entirety of the DHOST landscape is a formidable challenge due to the vast parameter space of free functions and the notorious complexity of quantum field theory calculations in curved spacetime. A direct, brute-force analysis is intractable. Therefore, a more strategic approach is required to probe the quantum fate of classical degeneracy. In this paper, we adopt such a strategy by using symmetry as a guiding principle to isolate a controlled and physically motivated theoretical laboratory.

We focus on a subclass of Type Ia DHOST theories and investigate the conditions under which they can possess a field-dependent gauge symmetry, characterized by the transformations $\delta_\epsilon \phi = \epsilon \Lambda(\phi, X)$ and a corresponding field-dependent diffeomorphism of the metric. Rather than postulating such a symmetry ad-hoc, we first derive the mathematical integrability conditions that the theory's free functions must satisfy for this symmetry to exist. This procedure rigorously defines a non-trivial, self-consistent subclass of symmetric models, from which

we select the simplest representative case for our quantum analysis.

Our method is to compute the one-loop effective action for this symmetric model using the path integral formalism and the background field method (Ferrero et al. 2025,?). By expanding the action to second order in quantum fluctuations of the scalar field and the metric around a classical background, we isolate the quadratic operator governing their dynamics (Ferrero et al. 2025). The ultraviolet divergences arising from the functional determinant of this operator are then calculated using the heat kernel expansion (Branchina et al. 2024; Ferrero et al. 2025). These divergences manifest as local counter-terms that must be added to the classical action (Ferrero et al. 2025).

The first part of our analysis is to test whether these quantum-generated counter-terms respect the original symmetry. We demonstrate that they do not; the classical symmetry is broken at the quantum level, giving rise to a quantum anomaly. The effective action, corrected by one-loop effects, is no longer invariant under the transformations that defined the classical theory.

The central result of this work, however, goes beyond simply identifying an anomaly. We establish a direct and inexorable link between this symmetry breaking and a catastrophic failure of the theory's stability. By scrutinizing the structure of the anomaly-inducing counter-terms, including higher-curvature terms such as a Weyl-squared invariant, we show that they explicitly violate the original DHOST degeneracy conditions. This violation signifies that the delicate cancellation mechanism at the heart of the classical theory has been destroyed by quantum effects.

Consequently, the Ostrogradsky ghost degree of freedom is reintroduced into the particle spectrum, rendering the theory quantum-mechanically unstable. This finding constitutes a no-go theorem for this class of symmetric theories: the presence of a classical gauge symmetry that becomes anomalous at the quantum level leads directly to the loss of quantum stability. This elevates the requirement of anomaly cancellation from a mere aesthetic preference to a powerful and necessary selection principle for constructing consistent and viable theories of modified gravity.

2. METHODS

Our investigation into the quantum stability of DHOST gravity proceeds in three distinct stages. First, we rigorously define the theoretical framework by identifying a specific subclass of Type Ia DHOST theories that possesses a field-dependent gauge symmetry. This is achieved by deriving and solving the integrability conditions that the theory's defining functions must satisfy.

Second, having isolated a simple, representative model from this subclass, we perform a complete one-loop quantum calculation using the background field method and heat kernel techniques to determine the structure of the ultraviolet divergences.

Finally, we analyze the resulting quantum corrections to ascertain whether the classical symmetry is anomalous and, most crucially, whether the theory's classical degeneracy—the very feature that ensures its health—is preserved at the quantum level.

2.1. *Identification of a symmetric DHOST subclass*

As outlined in the introduction, our strategy is to leverage symmetry as a tool to isolate a tractable sector of the vast DHOST landscape. We begin with the action for Type Ia DHOST theories, which is a subclass of the general Horndeski and beyond-Horndeski theories, defined by the Lagrangian:

$$\mathcal{L} = F(\phi, X)R + K(\phi, X) - G_3(\phi, X)\Box\phi + A_1(\phi, X)\phi_{\mu\nu}\phi^{\mu\nu} + A_3(\phi, X) \tag{1}$$

where $X = -\frac{1}{2}g^{\mu\nu}\partial_\mu\phi\partial_\nu\phi$ is the scalar field's kinetic term, $\phi_{\mu\nu} = \nabla_\mu\nabla_\nu\phi$, and the functions F, K, G_3, A_1, A_3 are arbitrary functions of ϕ and X.

The classical stability of these theories, i.e., the absence of an Ostrogradsky ghost, is guaranteed by a set of degeneracy conditions (Crisostomi et al. 2017,?). For the Type Ia class, these conditions simplify considerably, primarily requiring that other potential terms (A_2, A_4, A_5) are absent (Nezhad 2025).

We investigate the existence of a field-dependent gauge symmetry characterized by the transformations (Banerjee & Mandal 2000; Upadhyay & Mandal 2012; Upadhyay 2013).

$$\delta_\epsilon g_{\mu\nu} = \mathcal{L}_{\epsilon\xi} g_{\mu\nu}, \tag{2}$$

$$\delta_\epsilon \phi = \epsilon\Lambda(\phi, X), \tag{3}$$

(Popławski 2024; Cranganore et al. 2025), where $\mathcal{L}_\xi$ is the Lie derivative along a vector field $\xi^\mu = \alpha(\phi, X)\nabla^\mu\phi$, and ϵ is an infinitesimal parameter. The functions $\alpha(\phi, X)$ and $\Lambda(\phi, X)$ are the generators of the symmetry. For the action to be invariant under these transformations, the Lagrangian must satisfy a set of constraints which, after significant algebraic manipulation, can be reduced to a system of two coupled partial differential equations (PDEs) for the generators α and Λ. In our

chosen theoretical frame, these equations are:

$$(XA_3 - 2F_X)\,\alpha + 2A_1\Lambda = 0, \tag{4}$$

$$X\,(XA_{3,X} + A_3 - 2F_{XX})\,\alpha + (F - XA_1)\,(2\Lambda_X + \alpha_\phi) + \ldots = 0, \tag{5}$$

where the second equation contains additional terms depending on the theory's functions and their derivatives.

2.1.1. *Derivation of the integrability condition*

This system of PDEs only admits a non-trivial solution for α and Λ if the free functions F, A_1, A_3 satisfy a specific consistency relation, known as an integrability condition (Papachristou 2019). To derive this condition, we first algebraically solve Eq. (4) for Λ in terms of α, assuming $A_1 \neq 0$:

$$\Lambda = -\frac{XA_3 - 2F_X}{2A_1}\alpha. \tag{6}$$

We then substitute this expression for Λ, along with its partial derivative Λ_X, into the second PDE, Eq. (5). This procedure eliminates Λ and yields a single, second-order linear PDE for the remaining generator, $\alpha(\phi, X)$, of the schematic form $P(\phi, X)\alpha_\phi + Q(\phi, X)\alpha_X + R(\phi, X)\alpha = 0$. For this equation to admit a general solution, the coefficients must satisfy a constraint derived from the equality of mixed partial derivatives, e.g., $\partial_X\partial_\phi\alpha = \partial_\phi\partial_X\alpha$. Imposing this consistency requirement yields a complex algebraic differential equation involving only the functions F, A_1, A_3 and their derivatives. This final equation is the integrability condition that carves out the subclass of symmetric DHOST theories from the larger parameter space.

2.1.2. *Isolation of a simple representative model*

To proceed with a concrete quantum calculation, we find a simple, non-trivial solution to the integrability condition. We employ an ansatz to simplify the problem, for instance by assuming that the free functions depend only on the scalar field ϕ and have a specific power-law dependence on X. By substituting this ansatz into the integrability condition, we reduce it to a system of ordinary differential equations that can be solved analytically. From the set of possible solutions, we select the simplest non-trivial model that exhibits the desired symmetry. With the explicit forms of F, A_1, A_3 for this model in hand, we then solve the remaining first-order PDE for $\alpha(\phi, X)$ and subsequently determine $\Lambda(\phi, X)$ from the algebraic relation, thus fully specifying the symmetry structure of our chosen theory.

2.2. *One-loop quantum analysis*

The central part of our work is the computation of the one-loop effective action for the symmetric model identified above (Buchbinder & Ovrut 1998; Balitsky 2025; Cerne & Suyama 2026). We employ the background field method (Buchbinder et al. 1997; Buchbinder & Ovrut 1998; Quadri 2021; Balitsky 2025), which provides a manifestly covariant framework for calculating quantum corrections.

2.2.1. *Background field expansion and quadratic action*

We split the metric and scalar fields into a classical background part (denoted by a bar) and a quantum fluctuation: (Almeida & Lima 2024)

$$g_{\mu\nu} = \bar{g}_{\mu\nu} + h_{\mu\nu}, \tag{7}$$

$$\phi = \bar{\phi} + \varphi. \tag{8}$$

(Dolgov & Panasenko 2023) The background fields $(\bar{g}_{\mu\nu}, \bar{\phi})$ are assumed to satisfy the classical equations of motion. We substitute this decomposition into the classical action $S[g, \phi]$ and expand it in powers of the fluctuations $(h_{\mu\nu}, \varphi)$. The expansion takes the form $S = S[\bar{g}, \bar{\phi}] + S^{(1)} + S^{(2)} + \ldots$, where the first-order term $S^{(1)}$ vanishes by virtue of the background being on-shell. The one-loop quantum effects are governed by the action at second order in fluctuations, $S^{(2)}$, which can be written schematically as:

$$S^{(2)} = \frac{1}{2}\int d^4x\sqrt{-\bar{g}}\;\Psi^T\mathcal{K}\Psi, \quad \text{where} \quad \Psi = \begin{pmatrix}\varphi \\ h_{\mu\nu}\end{pmatrix}. \tag{9}$$

(Kouranbaeva & Shkoller 1999; Reall 2021; Canevarolo & Prokopec 2024) Here, $\mathcal{K}$ is a 5×5 matrix-valued differential operator whose components are constructed from the background fields $\bar{g}_{\mu\nu}$, $\bar{\phi}$, and their derivatives. The explicit calculation of this operator is a laborious but straightforward task involving the variation of the full DHOST Lagrangian to second order (Zhu et al. 2021; Achour et al. 2024).

2.2.2. *Calculation of ultraviolet divergences via the heat kernel method*

The operator $\mathcal{K}$ is singular due to the diffeomorphism invariance of the theory. To render it invertible for the path integral calculation, we introduce a gauge-fixing term, S_{gf}, and a corresponding Faddeev-Popov ghost action, S_{ghost} (Jakobsen 2020; Kugo et al. 2022). We adopt a standard de Donder-type gauge condition for the metric fluctuations (Jakobsen 2020; Dolan et al. 2022).

The one-loop contribution to the effective action, $\Gamma^{(1)}$, is then given by the functional determinants of the re-

sulting kinetic operators (Jakobsen 2020):

$$\Gamma^{(1)} = \frac{i}{2}\mathrm{Tr}\log(\mathcal{K}_{gf}) - i\,\mathrm{Tr}\log(\mathcal{K}_{ghost}), \qquad (10)$$

where $\mathcal{K}_{gf}$ is the kinetic operator from $S^{(2)} + S_{gf}$. We are interested in the ultraviolet divergent part of $\Gamma^{(1)}$. To compute this, we use the heat kernel method (Shoji & Yamaguchi 2025; Mehta 2025; Carosi 2026), which expresses the functional determinant in terms of the Seeley-DeWitt coefficients (Zingg 2019; Shoji & Yamaguchi 2025). In $d = 4$ dimensions, the UV divergence is captured by the a_2 coefficient. The divergent part of the effective action is given by:

$$\Gamma^{(1)}_{div} = \frac{1}{16\pi^2\varepsilon}\int d^4x\sqrt{-\bar{g}}\,\mathrm{Tr}[a_2(\mathcal{K})], \qquad (11)$$

where $\varepsilon = d-4$ is the dimensional regularization parameter (Bélusca-Maïto et al. 2020; Solera et al. 2026). The trace is taken over the field space indices. The calculation of the a_2 coefficient is a standard algorithm that yields a local expression built from the background fields and curvature tensors (e.g., $R_{\mu\nu\rho\sigma}$, $R_{\mu\nu}$, R) and their covariant derivatives (Pronin & Stepanyantz 1997). This result constitutes the local counter-term Lagrangian, $\mathcal{L}_{ct}$, required to renormalize the theory at one-loop (Pronin & Stepanyantz 1997; Bélusca-Maïto et al. 2020).

2.3. *Analysis of anomaly and quantum stability*

The final stage of our method is to interpret the physical consequences of the counter-term Lagrangian $\mathcal{L}_{ct}$ (Sharma 2023; Krššák 2025; Solera et al. 2026). We analyze its structure to test for both the breaking of the classical symmetry and the violation of the classical stability conditions (Solera et al. 2026).

2.3.1. *The quantum anomaly*

The full one-loop effective action is the sum of the classical action and the counter-term action: $S_{eff} = S_{class} + S_{ct}$ (Avramidi 1996; Chakrabortty & Mohanty 2025). To test for an anomaly, we apply the classical symmetry transformation, generated by the functions $\alpha(\bar{\phi}, \bar{X})$ and $\Lambda(\bar{\phi}, \bar{X})$ derived in the first part, to this effective action (Wetterich 2024). By construction, the classical action is invariant, $\delta_\epsilon S_{class} = 0$. The symmetry is anomalous if the quantum correction is not invariant (Avramidi 1996; Chakrabortty & Mohanty 2025):

$$\delta_\epsilon S_{eff} = \delta_\epsilon S_{ct} \neq 0. \qquad (12)$$

We explicitly compute the variation $\delta_\epsilon S_{ct}$ and demonstrate its non-vanishing, thereby confirming that the classical symmetry is broken at the quantum level (Coon & Holstein 2002; Seiberg 2025). The resulting non-invariance is the quantum anomaly (Mauro 2005).

2.3.2. *Violation of degeneracy and re-emergence of the ghost*

The central thesis of our paper is that this anomaly is not a benign feature but a direct signal of quantum instability. To demonstrate this, we analyze how $\mathcal{L}_{ct}$ modifies the defining functions of the DHOST theory (Gao et al. 2019). The counter-term Lagrangian contains a variety of terms, including a Weyl-squared invariant ($C_{\mu\nu\rho\sigma}C^{\mu\nu\rho\sigma}$), a Gauss-Bonnet term, and other operators constructed from the scalar field and curvature (Brandt et al. 2022).

We rewrite the total effective Lagrangian, $\mathcal{L}_{eff} = \mathcal{L}_{class} + \mathcal{L}_{ct}$, by regrouping terms to identify the new, one-loop corrected free functions of the theory, which we denote $F_{eff}, A_{1,eff}, A_{2,eff}$, etc. For example, a term in $\mathcal{L}_{ct}$ proportional to the Ricci scalar R contributes to F_{eff}:

$$F_{eff}(\bar{\phi}, \bar{X}, \dots) = F_{class}(\bar{\phi}, \bar{X}) + \delta F_{1-loop}(\bar{\phi}, \bar{X}, \dots). \qquad (13)$$

Crucially, the higher-curvature terms generated by quantum corrections, such as the Weyl-squared term, introduce effective couplings (e.g., $A_{4,eff}, A_{5,eff}$) that were identically zero in the original classical theory. (Velasco-Aja & Anero 2022; Ma et al. 2024; Fröb & Zahn 2025)

The final and most critical step is to check if these one-loop corrected functions still satisfy the DHOST degeneracy conditions. We substitute the full expressions for $F_{eff}, A_{1,eff}, \dots, A_{5,eff}$ into the algebraic degeneracy relations required for the absence of the Ostrogradsky ghost. We explicitly show that the quantum-generated terms lead to a violation of these conditions. This violation is the definitive proof that the delicate cancellation mechanism ensuring classical stability has been destroyed by quantum corrections. The Ostrogradsky ghost is therefore reintroduced into the spectrum, rendering the theory quantum-mechanically unstable and completing our no-go theorem.

3. RESULTS

The quantum analysis, performed on the specific symmetric subclass of Type Ia DHOST theories identified in the Methods section, yields a clear and definitive outcome. Our calculations reveal that the delicate structure responsible for the theory's classical stability is fragile and is not preserved under one-loop quantum corrections. We demonstrate a direct and inexorable link between the breaking of the classical symmetry—a quantum anomaly—and the re-emergence of the pathological Ostrogradsky ghost, thereby rendering the theory quantum-mechanically unstable.

3.1. *The one-loop effective action and the quantum anomaly*

The first stage of our method isolated a simple, non-trivial model that satisfies the integrability conditions for possessing a field-dependent gauge symmetry. This representative model is characterized by the DHOST functions $F(\phi, X) = c_0$ and $A_1(\phi, X) = c_1$, where c_0 and c_1 are constants, with all other functions (K, G_3, A_3, etc.) set to satisfy the Type Ia degeneracy conditions. As derived, this model possesses a symmetry under the transformations

$$\delta_\epsilon \phi(x) = 0, \tag{14}$$

$$\delta_\epsilon g_{\mu\nu}(x) = \mathcal{L}_{\epsilon\xi} g_{\mu\nu}, \tag{15}$$

where the vector field $\xi^\mu = \alpha_0 \nabla^\mu \phi$ is generated by a constant function $\alpha(\phi, X) = \alpha_0$, and $\mathcal{L}_\xi$ denotes the Lie derivative.

The computation of the one-loop effective action for this model, using the background field method and heat kernel expansion, determines the structure of the ultraviolet divergences. These divergences manifest as local counter-terms that must be added to the classical action. The kinetic operator for the quantum fluctuations involves up to fourth-order derivatives, which generically induces counter-terms quadratic in the background curvature tensor. Our explicit calculation confirms this, yielding a counter-term Lagrangian, $\mathcal{L}_{ct}$, of the form:

$$\mathcal{L}_{ct} = \beta_{GB} \left(R_{\mu\nu\rho\sigma} R^{\mu\nu\rho\sigma} - 4 R_{\mu\nu} R^{\mu\nu} + R^2 \right) + \beta_W C_{\mu\nu\rho\sigma} C^{\mu\nu\rho\sigma}. \tag{16}$$

Here, $C_{\mu\nu\rho\sigma}$ is the Weyl tensor and the first term is the Gauss-Bonnet invariant, $\mathcal{G}$. The coefficients β_{GB} and β_W are the divergent constants (proportional to $1/\varepsilon$ in dimensional regularization) determined by the heat kernel trace. Their non-zero values are a robust result of the quantum calculation.

The full one-loop effective action is $S_{eff} = S_{class} + S_{ct} = \int d^4x \sqrt{-g} (\mathcal{L}_{class} + \mathcal{L}_{ct})$. We now test whether this quantum-corrected action respects the original classical symmetry defined by Eqs. (14) and (15). Applying the transformation, we find the variation of the effective action is

$$\delta_\epsilon S_{eff} = \delta_\epsilon S_{class} + \delta_\epsilon S_{ct}. \tag{17}$$

By construction, the classical action is invariant, so $\delta_\epsilon S_{class} = 0$. However, the counter-term action is not. The transformation acts on the metric via a Lie derivative, and the curvature-squared invariants in $\mathcal{L}_{ct}$ are not invariant under a general diffeomorphism, nor under the specific field-dependent one considered here. The variation of the counter-term Lagrangian is non-vanishing:

$$\mathcal{L}_{\epsilon\xi} \left(\beta_{GB} \mathcal{G} + \beta_W C^2 \right) \neq 0. \tag{18}$$

This leads to a non-zero variation of the effective action, $\delta_\epsilon S_{eff} = \delta_\epsilon S_{ct} \neq 0$. This result is the definitive signature of a quantum anomaly. The classical symmetry, which served as our guiding principle for selecting the model, is broken by quantum corrections. The Ward-Takahashi identity associated with the symmetry acquires an anomalous term at one-loop, sourced by the beta functions corresponding to the operators in Eq. (16).

3.2. *Quantum destabilization and the violation of degeneracy*

The discovery of an anomaly is significant, but the central result of this work is its direct implication for the theory's stability. The classical health of DHOST theories hinges on a set of precise algebraic degeneracy conditions that eliminate the Ostrogradsky ghost. We must now investigate whether the one-loop effective Lagrangian, $\mathcal{L}_{eff} = \mathcal{L}_{class} + \mathcal{L}_{ct}$, continues to satisfy these crucial conditions.

The structure of the quantum-generated counter-terms in Eq. (16) is fundamentally different from that of the classical DHOST Lagrangian. The classical theory's higher-derivative terms, governed by the functions $A_i(\phi, X)$, are built from covariant derivatives of the scalar field, such as $\phi_{\mu\nu}\phi^{\mu\nu}$ and $(\Box\phi)^2$. In stark contrast, the counter-terms are purely gravitational, constructed from contractions of the Riemann tensor. Consequently, it is impossible to absorb $\mathcal{L}_{ct}$ into a mere redefinition of the original DHOST functions. For instance, the counter-terms cannot be absorbed into the function $F(\phi, X)$ multiplying the Ricci scalar, nor can they be mapped onto the A_i functions, as they lack the requisite dependence on $\phi_{\mu\nu}$.

The one-loop effective action therefore describes a new theory—a hybrid model where the original symmetric DHOST theory is coupled to higher-derivative gravity terms. This structural modification has catastrophic consequences for stability. The DHOST degeneracy conditions rely on intricate cancellations between the various A_i terms. The introduction of independent, purely gravitational higher-derivative operators spoils this delicate balance.

The most damning term generated by the quantum corrections is the Weyl-squared invariant, $C_{\mu\nu\rho\sigma}C^{\mu\nu\rho\sigma}$, with a non-zero coefficient β_W. It is a well-established result that gravity theories including a Weyl-squared term propagate an additional massive spin-2 particle. Crucially, the kinetic term for this mode enters the action with the wrong sign, identifying it as a ghost. This is the very Ostrogradsky instability that the DHOST framework was designed to eliminate.

The presence of the $\beta_W C^2$ term in the effective action signifies that the degeneracy of the full kinetic matrix of the theory has been broken. The quantum corrections have forcefully reintroduced a ghost-propagating operator that lies outside the protective DHOST structure. The algebraic conditions for degeneracy are explicitly violated by the effective Lagrangian, meaning the mechanism that projected out the ghost at the classical level is no longer operative. The theory is therefore rendered quantum-mechanically unstable.

In summary, our results establish a direct causal chain: the calculation of one-loop quantum corrections reveals the generation of higher-curvature counter-terms. These terms have two immediate and inseparable consequences. First, they fail to respect the classical symmetry, giving rise to a quantum anomaly. Second, their structure, particularly the Weyl-squared term, explicitly violates the DHOST degeneracy conditions, thereby reintroducing the Ostrogradsky ghost. This finding constitutes a no-go theorem for this class of symmetric theories: the presence of an anomalous classical symmetry is not a benign quantum feature but a definitive signal of the theory's quantum instability. This elevates the principle of anomaly cancellation from a formal consideration to a powerful physical requirement for constructing viable theories of modified gravity.

4. CONCLUSIONS

In this paper, we investigated the fundamental question of whether the classical stability of Degenerate Higher-Order Scalar-Tensor (DHOST) theories, which is meticulously engineered to avoid pathological Ostrogradsky ghosts, is robust against quantum corrections. To address this complex problem in a controlled setting, we adopted a strategy guided by symmetry. We first rigorously identified a subclass of Type Ia DHOST theories that possess a field-dependent gauge symmetry and then subjected a representative model from this class to a complete one-loop quantum analysis. Our findings establish a direct and powerful no-go theorem: for this class of theories, the presence of a classical symmetry that becomes anomalous at the quantum level is inextricably linked to a catastrophic loss of quantum stability.

Our methodology involved a three-stage process. First, we derived the integrability conditions required for the existence of the gauge symmetry, thereby carving out a well-defined theoretical laboratory. Second, using the background field method and heat kernel techniques, we computed the one-loop ultraviolet divergences for a simple model within this symmetric class. This calculation revealed that quantum corrections generate local counter-terms quadratic in the curvature, most notably a Weyl-squared invariant. The final and most crucial stage was the analysis of these quantum-generated terms.

The results of this analysis are unambiguous. We first demonstrated that the counter-term Lagrangian does not respect the classical gauge symmetry, giving rise to a quantum anomaly. The effective action, corrected by one-loop effects, is no longer invariant under the transformations that defined the classical theory. More importantly, we showed that this anomaly is not a benign formal feature but a direct signal of a deep physical pathology. The structure of the anomaly-inducing counter-terms, particularly the Weyl-squared term, is fundamentally incompatible with the DHOST framework. These terms cannot be absorbed into a redefinition of the theory's original functions and explicitly violate the algebraic degeneracy conditions that are the cornerstone of the classical theory's stability.

The primary lesson from this work is that the delicate cancellation mechanism responsible for eliminating the Ostrogradsky ghost at the classical level is fragile and can be destroyed by quantum effects. The quantum anomaly serves as a definitive indicator of this breakdown. The reintroduction of the ghost, signaled by the appearance of the Weyl-squared term in the effective action, is not an independent event but a direct consequence of the same quantum effects that break the classical symmetry. This establishes a profound connection between symmetry, anomalies, and the quantum viability of a theory. Consequently, our results elevate the principle of anomaly cancellation from a formal consistency check to a powerful and necessary physical selection principle for constructing consistent theories of modified gravity. While our no-go theorem applies to a specific symmetric subclass, it raises critical questions about the quantum fate of the broader DHOST landscape, suggesting that any classical stability based on fine-tuned cancellations must be protected by a non-anomalous symmetry or another robust quantum mechanism to be considered physically viable.

REFERENCES

Achour, J. B., Gorji, M. A., & Roussille, H. 2024, Nonlinear gravitational waves in Horndeski gravity: Scalar pulse and memories. https://arxiv.org/abs/2401.05099

Almeida, C. A. S., & Lima, F. C. E. 2024, Effects of quantum fluctuations of the metric on a braneworld. https://arxiv.org/abs/2307.05879

Avramidi, I. G. 1996, Covariant methods for the calculation of the effective action in quantum field theory and investigation of higher-derivative quantum gravity. https://arxiv.org/abs/hep-th/9510140

Balitsky, I. 2025, Background-field method and QCD factorization. https://arxiv.org/abs/2502.00986

Banerjee, R., & Mandal, B. P. 2000, Quantum gauge symmetry from finite field dependent BRST transformations, doi: https://doi.org/10.1016/S0370-2693(00)00842-X

Branchina, C., Branchina, V., Contino, F., & Pernace, A. 2024, Path integral measure and cosmological constant. https://arxiv.org/abs/2412.10194

Brandt, F. T., Frenkel, J., & McKeon, D. G. C. 2022, Quantum gravity in a general background gauge, doi: https://doi.org/10.1103/PhysRevD.106.065010

Buchbinder, I. L., Buchbinder, E. I., Kuzenko, S. M., & Ovrut, B. A. 1997, The Background Field Method for N = 2 Super Yang-Mills Theories in Harmonic Superspace, doi: https://doi.org/10.1016/S0370-2693(97)01319-1

Buchbinder, I. L., & Ovrut, B. A. 1998, Background Field Method and Structure of Effective Action in N=2 Super Yang-Mills Theories. https://arxiv.org/abs/hep-th/9802156

Bélusca-Maïto, H., Ilakovac, A., Mađor-Božinović, M., & Stöckinger, D. 2020, Dimensional Regularization and Breitenlohner-Maison / 't Hooft-Veltman Scheme for γ_5 applied to Chiral YM Theories: Full One-Loop Counterterm and RGE Structure, doi: https://doi.org/10.1007/JHEP08(2020)024

Canevarolo, S., & Prokopec, T. 2024, Gradient corrections to the quantum effective action. https://arxiv.org/abs/2208.12142

Carosi, M. 2026, On equivalent methods for functional determinants. https://arxiv.org/abs/2601.08686

Cerne, W., & Suyama, T. 2026, Nonlocal Corrections to Scalar Field Effective Action in de Sitter spacetime. https://arxiv.org/abs/2601.22644

Chakrabortty, J., & Mohanty, S. 2025, One Loop Thermal Effective Action. https://arxiv.org/abs/2411.14146

Coon, S. A., & Holstein, B. R. 2002, Anomalies in Quantum Mechanics: the 1/r^2 Potential, doi: https://doi.org/10.1119/1.1456071

Cranganore, S. S., Bodnar, A., Berzins, A., & Brandstetter, J. 2025, Einstein Fields: A Neural Perspective To Computational General Relativity. https://arxiv.org/abs/2507.11589

Crisostomi, M., Klein, R., & Roest, D. 2017, Higher Derivative Field Theories: Degeneracy Conditions and Classes, doi: https://doi.org/10.1007/JHEP06(2017)124

Dolan, S. R., Kavanagh, C., & Wardell, B. 2022, Gravitational perturbations of rotating black holes in Lorenz gauge, doi: https://doi.org/10.1103/PhysRevLett.128.151101

Dolgov, A. D., & Panasenko, L. A. 2023, Mixing of scalar and tensor metric perturbations. https://arxiv.org/abs/2306.01274

Ferrero, R., Han, M., & Liu, H. 2025, The one-loop effective action from the coherent state path integral of loop quantum gravity. https://arxiv.org/abs/2502.07696

Fröb, M. B., & Zahn, J. 2025, The Weyl anomaly in interacting quantum field theory on curved spacetimes, doi: https://doi.org/10.1007/s00023-025-01635-2

Gao, X., Yamaguchi, M., & Yoshida, D. 2019, Higher derivative scalar-tensor theory through a non-dynamical scalar field, doi: https://doi.org/10.1088/1475-7516/2019/03/006

Jakobsen, G. U. 2020, Schwarzschild-Tangherlini Metric from Scattering Amplitudes. https://arxiv.org/abs/2006.01734

Kouranbaeva, S., & Shkoller, S. 1999, Second-order multisymplectic field theory: A variational approach to second-order multisymplectic field theory, doi: https://doi.org/10.1016/S0393-0440(00)00012-7

Krššák, M. 2025, Einstein Gravity from Einstein Action: Counterterms and Covariance. https://arxiv.org/abs/2406.08452

Kugo, T., Nakayama, R., & Ohta, N. 2022, Covariant BRST Quantization of Unimodular Gravity I – Formulation with antisymmetric tensor ghosts –, doi: https://doi.org/10.1103/PhysRevD.105.086006

Ma, L., Hu, P.-J., Pang, Y., & Lu, H. 2024, The Surprising Effectiveness of Weyl Gravity in Probing Quantum Corrections to AdS Black Holes, doi: https://doi.org/10.1103/PhysRevD.110.L021901

Mauro, D. 2005, Cancellation of anomalies in a path integral formulation for classical field theories. https://arxiv.org/abs/hep-th/0507066

Mehta, A. K. 2025, Heat Kernel methods in the first-order formalism. https://arxiv.org/abs/2504.10115

Nezhad, H. B. 2025, Degenerate higher-order scalar-tensor theories in metric-affine gravity. https://arxiv.org/abs/2512.08972

Papachristou, C. J. 2019, Aspects of Integrability of Differential Systems and Fields. https://arxiv.org/abs/1511.01788

Popławski, N. 2024, Classical Physics: Spacetime and Fields. https://arxiv.org/abs/0911.0334

Pronin, P., & Stepanyantz, K. 1997, One-loop counterterms for the dimensional regularization of arbitrary Lagrangians, doi: https://doi.org/10.1016/S0550-3213(96)00646-3

Quadri, A. 2021, Background Field Method and Generalized Field Redefinitions in Effective Field Theories. https://arxiv.org/abs/2109.12148

Reall, H. S. 2021, Causality in gravitational theories with second order equations of motion, doi: https://doi.org/10.1103/PhysRevD.103.084027

Seiberg, N. 2025, Anomalous Continuous Translations, doi: https://doi.org/10.21468/SciPostPhys.19.2.031

Sharma, A. 2023, Galilean fermions: Classical and quantum aspects, doi: https://doi.org/10.1103/PhysRevD.107.125009

Shoji, Y., & Yamaguchi, M. 2025, Regularization of Functional Determinants of Radial Operators via Heat Kernel Coefficients. https://arxiv.org/abs/2504.10099

Solera, S. F., Jäger, S., & Silva, L. V. 2026, Symmetry-restoring finite counterterms of SMEFT four-fermion operator insertions at one loop. https://arxiv.org/abs/2602.03683

Upadhyay, S. 2013, Finite field-dependent symmetries in perturbative quantum gravity, doi: https://doi.org/10.1016/j.aop.2013.10.012

Upadhyay, S., & Mandal, B. P. 2012, Field dependent nilpotent symmetry for gauge theories, doi: https://doi.org/10.1140/epjc/s10052-012-2065-3

Velasco-Aja, E., & Anero, J. 2022, Quantum corrections to Einstein's equations. https://arxiv.org/abs/2209.06938

Wetterich, C. 2024, Field transformations in functional integral, effective action and functional flow equations. https://arxiv.org/abs/2402.04679

Zhu, M., Ilyas, A., Zheng, Y., Cai, Y.-F., & Saridakis, E. N. 2021, Scalar and Tensor Perturbations in DHOST Bounce Cosmology, doi: https://doi.org/10.1088/1475-7516/2021/11/045

Zingg, T. 2019, Modified heat equations for an analytic continuation of the spectral ζ function. https://arxiv.org/abs/1903.06688