

SNESL formalization

Dandan Xue

September 7, 2017

0 Level-0

Draft version 0.0.3: changed some notations; added some definitions and lemmas; fixed some bugs.

0.1 Source language syntax

(Ignore empty sequence for now)

Expressions:

$$e ::= x \mid \mathbf{let} \ x = e_1 \ \mathbf{in} \ e_2 \mid \phi(x_1, \dots, x_k) \mid \{e : x \ \mathbf{in} \ y \ \mathbf{using} \ \cdot\} \\ \phi = \mathbf{const}_n \mid \mathbf{iota} \mid \mathbf{plus}$$

Values:

$$n \in \mathbf{Z} \\ v ::= n \mid \{v_1, \dots, v_k\}$$

0.2 Type system

$$\tau ::= \mathbf{int} \mid \{\tau_1\}$$

Type environment $\Gamma = [x_1 \mapsto \tau_1, \dots, x_i \mapsto \tau_i]$.

- Judgment $\boxed{\Gamma \vdash e : \tau}$

$$\frac{}{\Gamma \vdash x : \tau} (\Gamma(x) = \tau) \quad \frac{\Gamma \vdash e_1 : \tau_1 \quad \Gamma[x \mapsto \tau_1] \vdash e_2 : \tau}{\Gamma \vdash \mathbf{let} \ x = e_1 \ \mathbf{in} \ e_2 : \tau} \\ \frac{\vdash \phi : (\tau_1, \dots, \tau_k) \rightarrow \tau \quad ((\Gamma(x_i) = \tau_i)_{i=1}^k)}{\Gamma \vdash \phi(x_1, \dots, x_k) : \tau} \quad \frac{[x \mapsto \tau_1] \vdash e : \tau}{\Gamma \vdash \{e : x \ \mathbf{in} \ y \ \mathbf{using} \ \cdot\} : \{\tau\}} (\Gamma(y) = \{\tau_1\})$$

- Auxiliary Judgment $\boxed{\vdash \phi : (\tau_1, \dots, \tau_k) \rightarrow \tau}$

$$\frac{}{\vdash \mathbf{const}_n : \mathbf{int}} \quad \frac{}{\vdash \mathbf{iota} : \mathbf{int} \rightarrow \{\mathbf{int}\}} \quad \frac{}{\vdash \mathbf{plus} : \mathbf{int} \rightarrow \mathbf{int} \rightarrow \mathbf{int}}$$

0.3 Source language semantics

$\rho = [x_1 \mapsto v_1, \dots, x_i \mapsto v_i]$

- Judgment $\boxed{\rho \vdash e \downarrow v}$

$$\frac{}{\rho \vdash x \downarrow v} (\rho(x) = v) \quad \frac{\rho \vdash e_1 \downarrow v_1 \quad \rho[x \mapsto v_1] \vdash e_2 \downarrow v}{\rho \vdash \mathbf{let} \ e_1 = x \ \mathbf{in} \ e_2 \downarrow v}$$

$$\frac{\vdash \phi(v_1, \dots, v_k) \downarrow v}{\rho \vdash \phi(x_1, \dots, x_k) \downarrow v} ((\rho(x_i) = v_i)_{i=1}^k) \quad \frac{([x \mapsto v_i] \vdash e \downarrow v'_i)_{i=1}^k}{\rho \vdash \{e : x \text{ in } y \text{ using } \cdot\} \downarrow \{v'_1, \dots, v'_k\}} (\rho(y) = \{v_1, \dots, v_k\})$$

- Auxiliary Judgment $\boxed{\vdash \phi(v_1, \dots, v_k) \downarrow v}$

$$\frac{}{\vdash \text{const}_n() \downarrow n} \quad \frac{}{\vdash \text{iota}(n) \downarrow \{0, 1, \dots, n-1\}} (n \geq 0) \quad \frac{}{\vdash \text{plus}(n_1, n_2) \downarrow n_3} (n_3 = n_1 + n_2)$$

0.4 SVCODE syntax

Stream id:

$$s \in \mathbf{SId} = \mathbf{N} = \{0, 1, 2, \dots\}$$

Stream tree:

$$\mathbf{STree} \ni st ::= s \mid (st_1, s)$$

SVCODE operations:

$$\psi ::= \text{Ctrl} \mid \text{Const}_a \mid \text{ToFlags} \mid \text{Usum} \mid \text{MapTwo} \mid \text{ScanPlus}$$

SVCODE program:

$$\begin{aligned} p ::= & \epsilon \\ & \mid s := \psi(s_1, \dots, s_i) \\ & \mid st := \text{WithCtrl}(s, p) \\ & \mid p_1; p_2 \end{aligned}$$

Target language values:

$$\begin{aligned} b & \in \{\mathbf{T}, \mathbf{F}\} \\ a & ::= n \mid b \mid () \\ \vec{b} & = \langle b_1, \dots, b_i \rangle \\ \vec{a} & = \langle a_1, \dots, a_i \rangle \\ \mathbf{SVal} \ni w & ::= \vec{a} \mid (w, \vec{b}) \end{aligned}$$

Some notations and operations:

- For some a_0 and $\vec{a} = \langle a_1, \dots, a_i \rangle$, $\langle a_0 | \vec{a} \rangle = \langle a_0, a_1, \dots, a_i \rangle$.
- $++ : \mathbf{SVal} \rightarrow \mathbf{SVal} \rightarrow \mathbf{SVal}$
 $\langle a_1, \dots, a_i \rangle ++ \langle a'_1, \dots, a'_i \rangle = \langle a_1, \dots, a_i, a'_1, \dots, a'_i \rangle$
 $(w_1, \vec{b}_1) ++ (w_2, \vec{b}_2) = (w_1 ++ w_2, \vec{b}_1 ++ \vec{b}_2)$
- **sids** converts a $st \in \mathbf{STree}$ to a set of $s \in \mathbf{SId}$:
 $\mathbf{sids}(s) = \{s\}$
 $\mathbf{sids}((st, s)) = \mathbf{sids}(st) \cup \{s\}$
- For some set set of \mathbf{SId} and some $s \in \mathbf{SId}$, $set < s$ denotes $\forall s' \in set. s' < s$.

0.5 SVCODE semantics

$\sigma = [s_1 \mapsto \vec{a}_1, \dots, s_i \mapsto \vec{a}_i]$.

Notation $\sigma_1 \stackrel{\leq s}{=} \sigma_2$ denotes $\forall s' < s. \sigma_1(s') = \sigma_2(s')$.

\vec{a}_c is the control stream.

- Judgment $\boxed{\langle p, \sigma \rangle \downarrow^{\vec{a}_c} \sigma'}$

$$\frac{}{\langle \epsilon, \sigma \rangle \downarrow^{\vec{a}_c} \sigma} \quad \frac{\psi(\vec{a}_1, \dots, \vec{a}_k) \downarrow^{\vec{a}_c} \vec{a}}{\langle s := \psi(s_1, \dots, s_k), \sigma \rangle \downarrow^{\vec{a}_c} \sigma[s \mapsto \vec{a}]} ((\sigma(s_i) = \vec{a}_i)_{i=1}^k)$$

$$\frac{}{\langle st := \text{WithCtrl}(s, p), \sigma \rangle \downarrow^{\vec{a}_c} \sigma[s_1 \mapsto \langle \rangle, \dots, s_i \mapsto \langle \rangle]} (\sigma(s) = \langle \rangle, \text{sids}(st) = \{s_1, \dots, s_i\})$$

$$\frac{\langle p, \sigma \rangle \downarrow^{\vec{a}_s} \sigma''}{\langle st := \text{WithCtrl}(s, p), \sigma \rangle \downarrow^{\vec{a}_c} \sigma[s_1 \mapsto \sigma''(s_1), \dots, s_i \mapsto \sigma''(s_i)]} \left(\begin{array}{l} \sigma(s) = \vec{a}_s = \langle a_0 | \vec{a} \rangle \\ \text{sids}(st) = \{s_1, \dots, s_i\} \end{array} \right)$$

$$\frac{\langle p_1, \sigma \rangle \downarrow^{\vec{a}_c} \sigma'' \quad \langle p_2, \sigma'' \rangle \downarrow^{\vec{a}_c} \sigma'}{\langle p_1; p_2, \sigma \rangle \downarrow^{\vec{a}_c} \sigma'}$$

- Auxiliary Judgment $\boxed{\psi(\vec{a}_1, \dots, \vec{a}_k) \downarrow^{\vec{a}_c} \vec{a}}$

$$\frac{\psi(\vec{a}_{11}, \dots, \vec{a}_{k1}) \downarrow \vec{a}_1 \quad \psi(\vec{a}_{12}, \dots, \vec{a}_{k2}) \downarrow^{\vec{a}_c} \vec{a}_2}{\psi(\vec{a}_{11} ++ \vec{a}_{12}, \dots, \vec{a}_{k1} ++ \vec{a}_{k2}) \downarrow^{\langle a_0 | \vec{a}_c \rangle} \vec{a}} (\vec{a} = \vec{a}_1 ++ \vec{a}_2)$$

$$\frac{}{\psi(\vec{a}_1, \dots, \vec{a}_k) \downarrow^{\langle \rangle} \langle \rangle}$$

- Auxiliary Judgment $\boxed{\psi(\vec{a}_1, \dots, \vec{a}_k) \Downarrow \vec{a}}$

$$\frac{}{\text{Const}_a \Downarrow \langle a \rangle} \quad \frac{}{\text{ToFlags}(\langle n \rangle) \Downarrow \langle F_1, \dots, F_n, T \rangle} \quad \frac{}{\text{MapTwo}_{\oplus}(\langle n_1 \rangle, \langle n_2 \rangle) \Downarrow \langle n_3 \rangle} (n_3 = n_1 \oplus n_2)$$

$$\frac{\psi(\langle F \rangle, \dots, \vec{a}_{k1}) \Downarrow \vec{a}_1 \quad \psi(\vec{a}_{12}, \dots, \vec{a}_{k2}) \Downarrow \vec{a}_2}{\psi(\langle F \rangle ++ \vec{a}_{12}, \dots, \vec{a}_{k1} ++ \vec{a}_{k2}) \Downarrow \vec{a}} (\vec{a} = \vec{a}_1 ++ \vec{a}_2)$$

$$\frac{\psi(\langle T \rangle, \dots, \vec{a}_k) \Downarrow \vec{a}}{\psi(\langle T \rangle, \dots, \vec{a}_k) \Downarrow \vec{a}}$$

- Auxiliary Judgment $\boxed{\psi(\langle b \rangle, \dots, \vec{a}_k) \Downarrow \vec{a}}$

$$\frac{}{\text{Usum}(\langle F \rangle) \Downarrow \langle () \rangle} \quad \frac{}{\text{Usum}(\langle T \rangle) \Downarrow \langle \rangle}$$

Theorem 0.1 (deterministic ??). *If $\langle p, \sigma \rangle \downarrow^{\vec{a}_c} \sigma'$ and $\langle p, \sigma \rangle \downarrow^{\vec{a}_c} \sigma''$, then $\sigma' = \sigma''$.*

Definition 0.1 (Stream prefix).

\vec{a} is a *prefix* of \vec{a}' :

Judgment $\boxed{\vec{a} \sqsubseteq \vec{a}'}$

$$\frac{}{\langle \rangle \sqsubseteq \vec{a}} \quad \frac{\vec{a} \sqsubseteq \vec{a}'}{\langle a_0 | \vec{a} \rangle \sqsubseteq \langle a_0 | \vec{a}' \rangle}$$

Lemma 0.1. *If*

(i) $(\vec{a}'_i \sqsubseteq \vec{a}_i)_{i=1}^k$ and $\psi(\vec{a}'_1, \dots, \vec{a}'_k) \Downarrow \vec{a}'$,

(ii) $(\vec{a}''_i \sqsubseteq \vec{a}_i)_{i=1}^k$ and $\psi(\vec{a}''_1, \dots, \vec{a}''_k) \Downarrow \vec{a}''$

then $(\vec{a}'_i = \vec{a}''_i)_{i=1}^k$ and $\vec{a}' = \vec{a}''$.

0.6 Translation

$\delta = [x_1 \mapsto st_1, \dots, x_i \mapsto st_i]$

- Judgment $\boxed{\delta \vdash e \xRightarrow[s_1]{s_0} (p, st)}$

$$\frac{}{\delta \vdash x \xRightarrow[s_0]{s_0} (\epsilon, st)} (\delta(x) = st) \quad \frac{\delta \vdash e_1 \xRightarrow[s'_0]{s_0} (p_1, st_1) \quad \delta[x \mapsto st_1] \vdash e_2 \xRightarrow[s_1]{s'_0} (p_2, st)}{\delta \vdash \text{let } x = e_1 \text{ in } e_2 \xRightarrow[s_1]{s_0} (p_1; p_2, st)}$$

$$\frac{\vdash \phi(st_1, \dots, st_k) \xRightarrow[s_1]{s_0} (p, st)}{\delta \vdash \phi(x_1, \dots, x_k) \xRightarrow[s_1]{s_0} (p, st)} ((\delta(x_i) = st_i)_{i=1}^k)$$

$$\frac{[x \mapsto st_1] \vdash e \xRightarrow[s_1]{s_0+1} (p, st)}{\delta \vdash \{e : x \text{ in } y \text{ using } \cdot\} \xRightarrow[s_1]{s_0} (s_0 := \text{Usum}(s_2); st := \text{WithCtrl}(s_0, p), (st, s_2))} (\delta(y) = (st_1, s_2))$$

- Auxiliary Judgment $\boxed{\vdash \phi(st_1, \dots, st_k) \xRightarrow[s_1]{s_0} (p, st)}$

$$\frac{}{\text{const}_a() \xRightarrow[s_0]{s_0+1} (s_0 := \text{Const}_a, s_0)}$$

$$\frac{\text{iota}(s) \xRightarrow[s_0]{s_4} (p, (s_3, s_0))}{\left(\begin{array}{l} s_{i+1} = s_i + 1 \\ p = s_0 := \text{ToFlags}(s); \\ s_1 := \text{Usum}(s_0); \\ s_2 := \text{WithCtrl}(s_1, s_2 := \text{Const}_1); \\ s_3 := \text{ScanPlus}(s_0, s_2) \end{array} \right)}$$

$$\frac{}{\text{plus}(s_1, s_2) \xRightarrow[s_0]{s_0+1} (s_0 := \text{MapTwo}_+(s_1, s_2), s_0)}$$

0.7 Value representation

- Judgment $\boxed{v \triangleright_\tau w}$

$$\frac{}{n \triangleright_{\text{int}} \langle n \rangle} \quad \frac{(v_i \triangleright_\tau w_i)_{i=1}^k}{\{v_1, \dots, v_k\} \triangleright_{\{\tau\}} (w, \langle \mathbf{F}_1, \dots, \mathbf{F}_k, \mathbf{T} \rangle)} (w = w_1 ++ w_2 ++ \dots ++ w_k)$$

Lemma 0.2. *If $v \triangleright_\tau w$, $v' \triangleright_\tau w$, then $v = v'$.*

0.8 Correctness proof

Lemma 0.3. *If*

(i) $\vdash \phi : (\tau_1, \dots, \tau_k) \rightarrow \tau$

(ii) $\vdash \phi(v_1, \dots, v_k) \Downarrow v$

$$(iii) \vdash \phi(st_1, \dots, st_k) \xRightarrow[s_1]{s_0} (p, st)$$

$$(iv) (v_i \triangleright_{\tau_i} st_i)_{i=1}^k$$

$$(v) \bigcup_{i=1}^k \mathbf{sids}(st_i) \leq s_0$$

then

$$(i) \langle p, \sigma \rangle \downarrow^{\vec{a}_c} \sigma' \text{ (by } \mathcal{P})$$

$$(ii) v \triangleright_{\tau} \sigma'(st) \text{ (by } \mathcal{V})$$

$$(iii) \sigma' \xRightarrow[\leq s_0]{} \sigma$$

$$(iv) \mathbf{sids}(st) \leq s_1$$

$$(v) s_0 \leq s_1$$

Theorem 0.2. *If*

$$(i) \Gamma \vdash e : \tau \text{ (by some derivation } \mathcal{T})$$

$$(ii) \rho \vdash e \downarrow v \text{ (by } \mathcal{E})$$

$$(iii) \delta \vdash e \xRightarrow[s_1]{s_0} (p, st) \text{ (by } \mathcal{C})$$

$$(iv) \forall x \in \text{dom}(\Gamma). \rho(x) : \Gamma(x) \wedge \mathbf{sids}(\delta(x)) \leq s_0 \wedge \rho(x) \triangleright_{\Gamma(x)} \sigma(\delta(x))$$

then

$$(i) \langle p, \sigma \rangle \downarrow^{(\cdot)} \sigma' \text{ (by } \mathcal{P})$$

$$(ii) v \triangleright_{\tau} \sigma'(st) \text{ (by } \mathcal{V})$$

$$(iii) \sigma' \xRightarrow[\leq s_0]{} \sigma$$

$$(iv) \mathbf{sids}(st) \leq s_1$$

$$(v) s_0 \leq s_1$$