

More Reductions for **NP** Problems


Nabil Mustafa

Computational Complexity

Reductions From Last Time

So far, we have shown the following problems **NP** complete:

- SAT , 3-CNF SAT, INDSET , CLIQUE , VERTEX-COVER , HITTING-SET , INTEGER-PROGRAMMING



HAMILTONIAN

Claim

HAMILTONIAN : Given a directed graph $G = (V, E)$, does G have a Hamiltonian path?

The Hamiltonian path problem (**HAMILTONIAN**) is **NP** complete.

HAMILTONIAN

Claim

HAMILTONIAN : Given a directed graph $G = (V, E)$, does G have a Hamiltonian path?

The Hamiltonian path problem (**HAMILTONIAN**) is **NP** complete.

- Review: Hamiltonian path visits each vertex in G exactly once

HAMILTONIAN

Claim

HAMILTONIAN : Given a directed graph $G = (V, E)$, does G have a Hamiltonian path?

The Hamiltonian path problem (**HAMILTONIAN**) is **NP** complete.

- Review: Hamiltonian path visits each vertex in G exactly once
- Computationally very different from Eulerian paths

HAMILTONIAN

Claim

HAMILTONIAN : Given a directed graph $G = (V, E)$, does G have a Hamiltonian path?

The Hamiltonian path problem (**HAMILTONIAN**) is **NP** complete.

- Review: Hamiltonian path visits each vertex in G exactly once
- Computationally very different from Eulerian paths
- Note that **HAMILTONIAN** is in **NP**

HAMILTONIAN

Claim

HAMILTONIAN : Given a directed graph $G = (V, E)$, does G have a Hamiltonian path?

The Hamiltonian path problem (**HAMILTONIAN**) is **NP** complete.

- Review: Hamiltonian path visits each vertex in G exactly once
- Computationally very different from Eulerian paths
- Note that **HAMILTONIAN** is in **NP**
- We reduce 3-CNF SAT to **HAMILTONIAN**
 - ▶ For a fixed 3-CNF SAT formula, show that it can be transformed into a graph whose Hamiltonian path will give us the assignments for SAT.

Reduction Sketch

$$\phi = \mathcal{C}_1 \wedge \mathcal{C}_2 \wedge \dots \wedge \mathcal{C}_m, \quad n \text{ variables}, \quad \mathcal{C}_i = (x_1^i \vee x_2^i \vee x_3^i)$$

Reduction Sketch

$$\phi = \mathcal{C}_1 \wedge \mathcal{C}_2 \wedge \dots \wedge \mathcal{C}_m, \quad n \text{ variables}, \quad \mathcal{C}_i = (x_1^i \vee x_2^i \vee x_3^i)$$

- Given ϕ , have to construct a graph G such that G has a Hamiltonian path iff ϕ is satisfiable.

Reduction Sketch

$$\phi = \mathcal{C}_1 \wedge \mathcal{C}_2 \wedge \dots \wedge \mathcal{C}_m, \quad n \text{ variables}, \quad \mathcal{C}_i = (x_1^i \vee x_2^i \vee x_3^i)$$

- Given ϕ , have to construct a graph G such that G has a Hamiltonian path iff ϕ is satisfiable.
- We first define the mapping of variables, and then the clauses.

Reduction Sketch


$$\phi = \mathcal{C}_1 \wedge \mathcal{C}_2 \wedge \dots \wedge \mathcal{C}_m, \quad n \text{ variables}, \quad \mathcal{C}_i = (x_1^i \vee x_2^i \vee x_3^i)$$

- Given ϕ , have to construct a graph G such that G has a Hamiltonian path iff ϕ is satisfiable.
- We first define the mapping of variables, and then the clauses.
- Each x_i will correspond to a path (chain) of $6m$ vertices

Reduction Sketch

$$\phi = \mathcal{C}_1 \wedge \mathcal{C}_2 \wedge \dots \wedge \mathcal{C}_m, \quad n \text{ variables}, \quad \mathcal{C}_i = (x_1^i \vee x_2^i \vee x_3^i)$$


- Given ϕ , have to construct a graph G such that G has a Hamiltonian path iff ϕ is satisfiable.
- We first define the mapping of variables, and then the clauses.
- Each x_i will correspond to a path (chain) of $6m$ vertices



Reduction Sketch

$$\phi = \mathcal{C}_1 \wedge \mathcal{C}_2 \wedge \dots \wedge \mathcal{C}_m, \quad n \text{ variables}, \quad \mathcal{C}_i = (x_1^i \vee x_2^i \vee x_3^i)$$

- Given ϕ , have to construct a graph G such that G has a Hamiltonian path iff ϕ is satisfiable.
- We first define the mapping of variables, and then the clauses.
- Each x_i will correspond to a path (chain) of $6m$ vertices
- If we are at the first (or end) vertex, only one path to follow




- Add a start vertex v_{start} which has

- Add a start vertex v_{start} which has
 - ▶ no incoming edges.

- Add a start vertex v_{start} which has
 - ▶ no incoming edges.
 - ▶ an outgoing edge to the first vertex of x_1 chain.

- Add a start vertex v_{start} which has
 - ▶ no incoming edges.
 - ▶ an outgoing edge to the first vertex of x_1 chain.
 - ▶ an outgoing edge to the last vertex of x_1 chain.

- Add a start vertex v_{start} which has
 - ▶ no incoming edges.
 - ▶ an outgoing edge to the first vertex of x_1 chain.
 - ▶ an outgoing edge to the last vertex of x_1 chain.




- Add an end vertex v_{end} which has

- Add an end vertex v_{end} which has
 - ▶ no outgoing edges.

- Add an end vertex v_{end} which has
 - ▶ no outgoing edges.
 - ▶ an incoming edge from the first vertex of x_n chain.

- Add an end vertex v_{end} which has
 - ▶ no outgoing edges.
 - ▶ an incoming edge from the first vertex of x_n chain.
 - ▶ an incoming edge from the last vertex of x_n chain.

- Add an end vertex v_{end} which has
 - ▶ no outgoing edges.
 - ▶ an incoming edge from the first vertex of x_n chain.
 - ▶ an incoming edge from the last vertex of x_n chain.




- From the first and last vertex of each chain x_i , add


- From the first and last vertex of each chain x_i , add
 - ▶ an outgoing edge to the first vertex of chain x_{i+1}

- From the first and last vertex of each chain x_i , add
 - ▶ an outgoing edge to the first vertex of chain x_{i+1}
 - ▶ an outgoing edge to the last vertex of chain x_{i+1}


- From the first and last vertex of each chain x_i , add
 - an outgoing edge to the first vertex of chain x_{i+1}
 - an outgoing edge to the last vertex of chain x_{i+1}




- From the first and last vertex of each chain x_i , add
 - an outgoing edge to the first vertex of chain x_{i+1}
 - an outgoing edge to the last vertex of chain x_{i+1}




- From the first and last vertex of each chain x_i , add
 - ▶ an outgoing edge to the first vertex of chain x_{i+1}
 - ▶ an outgoing edge to the last vertex of chain x_{i+1}



- From the first and last vertex of each chain x_i , add
 - an outgoing edge to the first vertex of chain x_{i+1}
 - an outgoing edge to the last vertex of chain x_{i+1}




Construction Properties




Construction Properties

- Any Hamiltonian path has to start at v_{start}




Construction Properties

- Any Hamiltonian path has to end at v_{end}




Construction Properties

- Any Hamiltonian path first traverses chain x_1 , then x_2 etc.




Construction Properties

- For each chain, only two ways of traversing it.
 - ▶ Left-to-right means $x_i = 1$, right-to-left means $x_i = 0$




Construction Properties

- Each assignment of variables corresponds to a unique Hamiltonian path.




Construction Properties

- Each Hamiltonian path corresponds to a unique variable assignment.



Construction Properties

- So far, no constraints – they will come from the clauses now.



Clause Constraints

- Each clause \mathcal{C}_j corresponds to a new vertex u_j .

Clause Constraints

- Each clause \mathcal{C}_j corresponds to a new vertex u_j .
- If \mathcal{C}_j contains a non-negated literal x_i , add edges to u_j :

Clause Constraints


- Each clause \mathcal{C}_j corresponds to a new vertex u_j .
- If \mathcal{C}_j contains a non-negated literal x_i , add edges to u_j :
 - ▶ Add an incoming edge from a vertex, say v_k , in the x_i chain

Clause Constraints

- Each clause \mathcal{C}_j corresponds to a new vertex u_j .
- If \mathcal{C}_j contains a non-negated literal x_i , add edges to u_j :
 - ▶ Add an incoming edge from a vertex, say v_k , in the x_i chain
 - ▶ Add an outgoing edge to v_{k+1} in the x_i chain

Clause Constraints

- Each clause \mathcal{C}_j corresponds to a new vertex u_j .
- If \mathcal{C}_j contains a non-negated literal x_i , add edges to u_j :
 - ▶ Add an incoming edge from a vertex, say v_k , in the x_i chain
 - ▶ Add an outgoing edge to v_{k+1} in the x_i chain



Clause Constraints

- If \mathcal{C}_j contains a negated literal, reverse the edge directions

Clause Constraints


- If \mathcal{C}_j contains a negated literal, reverse the edge directions
- If \mathcal{C}_j contains a negated literal \bar{x}_i , add edges to u_j :
 - ▶ Add an incoming edge from a vertex, say v_k , in the x_i chain

Clause Constraints

- If \mathcal{C}_j contains a negated literal, reverse the edge directions
- If \mathcal{C}_j contains a negated literal \bar{x}_i , add edges to u_j :
 - ▶ Add an incoming edge from a vertex, say v_k , in the x_i chain
 - ▶ Add an outgoing edge to v_{k-1} in the x_i chain

Clause Constraints

- If \mathcal{C}_j contains a negated literal, reverse the edge directions
- If \mathcal{C}_j contains a negated literal \bar{x}_i , add edges to u_j :
 - ▶ Add an incoming edge from a vertex, say v_k , in the x_i chain
 - ▶ Add an outgoing edge to v_{k-1} in the x_i chain



An Example


Construction of Hamiltonian path for

$$(a \vee b) \wedge (a \vee \bar{b})$$


Here:

- $C_1 = (a \vee b)$
- $C_2 = (a \vee \bar{b})$

The Graph Construction



The Graph Construction




Claim

Hamiltonian path exists **ONLY** if you go from left to right in **a** and chose any one of the two directions for **b**.


Path Corresponding to Assignment 1

$a = 1; b = 1$




Path Corresponding to Assignment 2

$a = 1; b = 0$



Path Corresponding to Assignment 3


$$a = 0; b = 1$$



Error in finding HAMILTONIAN

No HAMILTONIAN PATH as $(a \vee \bar{b})$ is not accessible

Path Corresponding to Assignment 4



Error in finding HAMILTONIAN

No HAMILTONIAN PATH as $(a \vee b)$ is not accessible

Construction Claim

Claim

*The constructed graph G has a Hamiltonian path **iff** ϕ satisfiable.*

Construction Claim

Claim

*The constructed graph G has a Hamiltonian path **iff** ϕ satisfiable.*

Things to note about the final construction:

Construction Claim

Claim

*The constructed graph G has a Hamiltonian path **iff** ϕ satisfiable.*

Things to note about the final construction:

- Any Hamiltonian path has to start at v_{start}

Construction Claim

Claim

*The constructed graph G has a Hamiltonian path **iff** ϕ satisfiable.*

Things to note about the final construction:

- Any Hamiltonian path has to start at v_{start}
- Any Hamiltonian path has to end at v_{end}

Construction Claim

Claim

The constructed graph G has a Hamiltonian path iff ϕ satisfiable.

Things to note about the final construction:

- Any Hamiltonian path has to start at v_{start}
- Any Hamiltonian path has to end at v_{end}
- Any Hamiltonian path first traverses chain x_1 , then x_2 etc.

Construction Claim

Claim

The constructed graph G has a Hamiltonian path iff ϕ satisfiable.

Things to note about the final construction:

- Any Hamiltonian path has to start at v_{start}
- Any Hamiltonian path has to end at v_{end}
- Any Hamiltonian path first traverses chain x_1 , then x_2 etc.
- For each chain, only two ways of traversing it.

Construction Claim

Claim

The constructed graph G has a Hamiltonian path iff ϕ satisfiable.

Things to note about the final construction:

- Any Hamiltonian path has to start at v_{start}
- Any Hamiltonian path has to end at v_{end}
- Any Hamiltonian path first traverses chain x_1 , then x_2 etc.
- For each chain, only two ways of traversing it.
- Each assignment of variables corresponds to a path.

Construction Claim

Claim

The constructed graph G has a Hamiltonian path iff ϕ satisfiable.

Things to note about the final construction:

- Any Hamiltonian path has to start at v_{start}
- Any Hamiltonian path has to end at v_{end}
- Any Hamiltonian path first traverses chain x_1 , then x_2 etc.
- For each chain, only two ways of traversing it.
- Each assignment of variables corresponds to a path.
- Each path corresponds to a unique variable assignment.

Construction Claim

Claim

The constructed graph G has a Hamiltonian path iff ϕ satisfiable.

Things to note about the final construction:

- Any Hamiltonian path has to start at v_{start}
- Any Hamiltonian path has to end at v_{end}
- Any Hamiltonian path first traverses chain x_1 , then x_2 etc.
- For each chain, only two ways of traversing it.
- Each assignment of variables corresponds to a path.
- Each path corresponds to a unique variable assignment.
- If $x_i \in \mathcal{C}_j$ and $x_i = 1$, can visit u_j along the way.

Construction Claim

Claim

The constructed graph G has a Hamiltonian path iff ϕ satisfiable.

Things to note about the final construction:

- Any Hamiltonian path has to start at v_{start}
- Any Hamiltonian path has to end at v_{end}
- Any Hamiltonian path first traverses chain x_1 , then x_2 etc.
- For each chain, only two ways of traversing it.
- Each assignment of variables corresponds to a path.
- Each path corresponds to a unique variable assignment.
- If $x_i \in C_j$ and $x_i = 1$, can visit u_j along the way.
- If $\bar{x}_i \in C_j$ and $x_i = 0$, can visit u_j along the way.

Construction Claim

Claim

The constructed graph G has a Hamiltonian path iff ϕ satisfiable.

Things to note about the final construction:

- Any Hamiltonian path has to start at v_{start}
- Any Hamiltonian path has to end at v_{end}
- Any Hamiltonian path first traverses chain x_1 , then x_2 etc.
- For each chain, only two ways of traversing it.
- Each assignment of variables corresponds to a path.
- Each path corresponds to a unique variable assignment.
- If $x_i \in C_j$ and $x_i = 1$, can visit u_j along the way.
- If $\bar{x}_i \in C_j$ and $x_i = 0$, can visit u_j along the way.
- The above two *only* ways to visit u_j without getting stuck.

SET-COVER

Claim

SET-COVER : A collection $\mathcal{C} = \{S_1, \dots, S_m\}$ of subsets of a base set X , $|X| = n$, and parameter k , find a set cover $\mathcal{C}' \subseteq \mathcal{C}$ of size k
The set cover problem (**SET-COVER**) is **NP** complete.

SET-COVER

Claim

SET-COVER : A collection $\mathcal{C} = \{S_1, \dots, S_m\}$ of subsets of a base set X , $|X| = n$, and parameter k , find a set cover $\mathcal{C}' \subseteq \mathcal{C}$ of size k
The set cover problem (**SET-COVER**) is **NP** complete.

- $\mathcal{C}' = \{S_{i_1}, \dots, S_{i_k}\}$ is a set-cover iff $\bigcup_j S_{i_j} = X$.

SET-COVER

Claim

SET-COVER : A collection $\mathcal{C} = \{S_1, \dots, S_m\}$ of subsets of a base set X , $|X| = n$, and parameter k , find a set cover $\mathcal{C}' \subseteq \mathcal{C}$ of size k
The set cover problem (**SET-COVER**) is **NP** complete.

- $\mathcal{C}' = \{S_{i_1}, \dots, S_{i_k}\}$ is a set-cover iff $\bigcup_j S_{i_j} = X$.
- Note that SET-COVER is in **NP**

SET-COVER

Claim

SET-COVER : A collection $\mathcal{C} = \{S_1, \dots, S_m\}$ of subsets of a base set X , $|X| = n$, and parameter k , find a set cover $\mathcal{C}' \subseteq \mathcal{C}$ of size k
The set cover problem (**SET-COVER**) is **NP** complete.

- $\mathcal{C}' = \{S_{i_1}, \dots, S_{i_k}\}$ is a set-cover iff $\bigcup_j S_{i_j} = X$.
- Note that SET-COVER is in **NP**
- How to prove **NP** hardness?

SET-COVER

Claim

SET-COVER : A collection $\mathcal{C} = \{S_1, \dots, S_m\}$ of subsets of a base set X , $|X| = n$, and parameter k , find a set cover $\mathcal{C}' \subseteq \mathcal{C}$ of size k
The set cover problem (**SET-COVER**) is **NP** complete.

- $\mathcal{C}' = \{S_{i_1}, \dots, S_{i_k}\}$ is a set-cover iff $\bigcup_j S_{i_j} = X$.
- Note that SET-COVER is in **NP**
- How to prove **NP** hardness?
- Reduce from **VERTEX-COVER**

Reduction from VERTEX-COVER

- Given a graph $G = (V, E)$, the idea is that:

Reduction from VERTEX-COVER

- Given a graph $G = (V, E)$, the idea is that:
 - ▶ Each vertex v_i maps to a set C_i

Reduction from VERTEX-COVER

- Given a graph $G = (V, E)$, the idea is that:
 - ▶ Each vertex v_i maps to a set C_i
 - ▶ $\{v_{i_1}, \dots, v_{i_k}\}$ is a vertex-cover iff $\{C_{i_1}, \dots, C_{i_k}\}$ is a set-cover.

Reduction from VERTEX-COVER

- Given a graph $G = (V, E)$, the idea is that:
 - ▶ Each vertex v_i maps to a set C_i
 - ▶ $\{v_{i_1}, \dots, v_{i_k}\}$ is a vertex-cover iff $\{C_{i_1}, \dots, C_{i_k}\}$ is a set-cover.
- The exact construction is as follows:

Reduction from VERTEX-COVER

- Given a graph $G = (V, E)$, the idea is that:
 - ▶ Each vertex v_i maps to a set \mathcal{C}_i
 - ▶ $\{v_{i_1}, \dots, v_{i_k}\}$ is a vertex-cover iff $\{C_{i_1}, \dots, C_{i_k}\}$ is a set-cover.
- The exact construction is as follows:
 - ▶ Each edge $e_j \in E$ maps to an element $a_j \in X$.

Reduction from VERTEX-COVER

- Given a graph $G = (V, E)$, the idea is that:
 - ▶ Each vertex v_i maps to a set C_i
 - ▶ $\{v_{i_1}, \dots, v_{i_k}\}$ is a vertex-cover iff $\{C_{i_1}, \dots, C_{i_k}\}$ is a set-cover.
- The exact construction is as follows:
 - ▶ Each edge $e_j \in E$ maps to an element $a_j \in X$.
 - ▶ Each vertex $v_i \in V$ maps to set $C_i = \{a_j \text{ s.t. } e_j \text{ incident to } v_i\}$

Reduction from VERTEX-COVER

- Given a graph $G = (V, E)$, the idea is that:
 - ▶ Each vertex v_i maps to a set C_i
 - ▶ $\{v_{i_1}, \dots, v_{i_k}\}$ is a vertex-cover iff $\{C_{i_1}, \dots, C_{i_k}\}$ is a set-cover.
- The exact construction is as follows:
 - ▶ Each edge $e_j \in E$ maps to an element $a_j \in X$.
 - ▶ Each vertex $v_i \in V$ maps to set $C_i = \{a_j \text{ s.t. } e_j \text{ incident to } v_i\}$

Claim

$V' = \{v_{i_1}, \dots, v_{i_k}\}$ is a vertex-cover iff $\{C_{i_1}, \dots, C_{i_k}\}$ is a set cover.

Reduction from VERTEX-COVER

- Given a graph $G = (V, E)$, the idea is that:
 - ▶ Each vertex v_i maps to a set C_i
 - ▶ $\{v_{i_1}, \dots, v_{i_k}\}$ is a vertex-cover iff $\{C_{i_1}, \dots, C_{i_k}\}$ is a set-cover.
- The exact construction is as follows:
 - ▶ Each edge $e_j \in E$ maps to an element $a_j \in X$.
 - ▶ Each vertex $v_i \in V$ maps to set $C_i = \{a_j \text{ s.t. } e_j \text{ incident to } v_i\}$

Claim

$V' = \{v_{i_1}, \dots, v_{i_k}\}$ is a vertex-cover iff $\{C_{i_1}, \dots, C_{i_k}\}$ is a set cover.

Proof.

- If e_j incident to v_i , then set C_i contains a_j

Reduction from VERTEX-COVER

- Given a graph $G = (V, E)$, the idea is that:
 - ▶ Each vertex v_i maps to a set C_i
 - ▶ $\{v_{i_1}, \dots, v_{i_k}\}$ is a vertex-cover iff $\{C_{i_1}, \dots, C_{i_k}\}$ is a set-cover.
- The exact construction is as follows:
 - ▶ Each edge $e_j \in E$ maps to an element $a_j \in X$.
 - ▶ Each vertex $v_i \in V$ maps to set $C_i = \{a_j \text{ s.t. } e_j \text{ incident to } v_i\}$

Claim

$V' = \{v_{i_1}, \dots, v_{i_k}\}$ is a vertex-cover iff $\{C_{i_1}, \dots, C_{i_k}\}$ is a set cover.

Proof.

- If e_j incident to v_i , then set C_i contains a_j
- Picking v_i covers all edges incident to $v_i \iff$ picking C_i covers all corresponding elements a_j

Reduction from VERTEX-COVER

- Given a graph $G = (V, E)$, the idea is that:
 - ▶ Each vertex v_i maps to a set C_i
 - ▶ $\{v_{i_1}, \dots, v_{i_k}\}$ is a vertex-cover iff $\{C_{i_1}, \dots, C_{i_k}\}$ is a set-cover.
- The exact construction is as follows:
 - ▶ Each edge $e_j \in E$ maps to an element $a_j \in X$.
 - ▶ Each vertex $v_i \in V$ maps to set $C_i = \{a_j \text{ s.t. } e_j \text{ incident to } v_i\}$

Claim

$V' = \{v_{i_1}, \dots, v_{i_k}\}$ is a vertex-cover iff $\{C_{i_1}, \dots, C_{i_k}\}$ is a set cover.

Proof.

- If e_j incident to v_i , then set C_i contains a_j
- Picking v_i covers all edges incident to $v_i \iff$ picking C_i covers all corresponding elements a_j
- V' covers all edges $\iff C'$ cover all elements



Another Hardness Proof for INTEGER-PROGRAMMING

Another reduction for INTEGER-PROGRAMMING

SET-COVER can be reduced to INTEGER-PROGRAMMING

Another Hardness Proof for INTEGER-PROGRAMMING

Another reduction for INTEGER-PROGRAMMING

SET-COVER can be reduced to INTEGER-PROGRAMMING

Proof.

- Formulate the set-cover problem as an integer program

Another Hardness Proof for INTEGER-PROGRAMMING

Another reduction for INTEGER-PROGRAMMING

SET-COVER can be reduced to INTEGER-PROGRAMMING

Proof.

- Formulate the set-cover problem as an integer program
- Given $\mathcal{C} = \{S_1, \dots, S_m\}$ over $X = \{a_1, \dots, a_n\}$ and k :

Another Hardness Proof for INTEGER-PROGRAMMING

Another reduction for INTEGER-PROGRAMMING

SET-COVER can be reduced to INTEGER-PROGRAMMING

Proof.

- Formulate the set-cover problem as an integer program
- Given $\mathcal{C} = \{S_1, \dots, S_m\}$ over $X = \{a_1, \dots, a_n\}$ and k :
- Integer program variables: $x_j = 1$ iff S_j picked in set-cover

Another Hardness Proof for INTEGER-PROGRAMMING

Another reduction for INTEGER-PROGRAMMING

SET-COVER can be reduced to INTEGER-PROGRAMMING

Proof.

- Formulate the set-cover problem as an integer program
- Given $\mathcal{C} = \{S_1, \dots, S_m\}$ over $X = \{a_1, \dots, a_n\}$ and k :
- Integer program variables: $x_j = 1$ iff S_j picked in set-cover

At most k sets:

Another Hardness Proof for INTEGER-PROGRAMMING

Another reduction for INTEGER-PROGRAMMING

SET-COVER can be reduced to INTEGER-PROGRAMMING

Proof.

- Formulate the set-cover problem as an integer program
- Given $\mathcal{C} = \{S_1, \dots, S_m\}$ over $X = \{a_1, \dots, a_n\}$ and k :
- Integer program variables: $x_j = 1$ iff S_j picked in set-cover

At most k sets:
$$\sum_j x_j \leq k$$

Another Hardness Proof for INTEGER-PROGRAMMING

Another reduction for INTEGER-PROGRAMMING

SET-COVER can be reduced to INTEGER-PROGRAMMING

Proof.

- Formulate the set-cover problem as an integer program
- Given $\mathcal{C} = \{S_1, \dots, S_m\}$ over $X = \{a_1, \dots, a_n\}$ and k :
- Integer program variables: $x_j = 1$ iff S_j picked in set-cover

At most k sets:
$$\sum_j x_j \leq k$$

All elements covered:

Another Hardness Proof for INTEGER-PROGRAMMING

Another reduction for INTEGER-PROGRAMMING

SET-COVER can be reduced to INTEGER-PROGRAMMING

Proof.

- Formulate the set-cover problem as an integer program
- Given $\mathcal{C} = \{S_1, \dots, S_m\}$ over $X = \{a_1, \dots, a_n\}$ and k :
- Integer program variables: $x_j = 1$ iff S_j picked in set-cover

At most k sets:
$$\sum_j x_j \leq k$$

All elements covered:
$$\sum_{j|a_i \in S_j} x_j \geq 1 \quad \forall i$$

Another Hardness Proof for INTEGER-PROGRAMMING

Another reduction for INTEGER-PROGRAMMING

SET-COVER can be reduced to INTEGER-PROGRAMMING

Proof.

- Formulate the set-cover problem as an integer program
- Given $\mathcal{C} = \{S_1, \dots, S_m\}$ over $X = \{a_1, \dots, a_n\}$ and k :
- Integer program variables: $x_j = 1$ iff S_j picked in set-cover

At most k sets:
$$\sum_j x_j \leq k$$

All elements covered:
$$\sum_{j|a_i \in S_j} x_j \geq 1 \quad \forall i$$

Variable 0 or 1:

Another Hardness Proof for INTEGER-PROGRAMMING

Another reduction for INTEGER-PROGRAMMING

SET-COVER can be reduced to INTEGER-PROGRAMMING

Proof.

- Formulate the set-cover problem as an integer program
- Given $\mathcal{C} = \{S_1, \dots, S_m\}$ over $X = \{a_1, \dots, a_n\}$ and k :
- Integer program variables: $x_j = 1$ iff S_j picked in set-cover

At most k sets:
$$\sum_j x_j \leq k$$

All elements covered:
$$\sum_{j|a_i \in S_j} x_j \geq 1 \quad \forall i$$

Variable 0 or 1:
$$x_i \in \{0, 1\}$$



Yet Another Hardness Proof for INTEGER-PROGRAMMING

Yet another reduction for INTEGER-PROGRAMMING

HITTING-SET can be reduced to INTEGER-PROGRAMMING

Yet Another Hardness Proof for INTEGER-PROGRAMMING

Yet another reduction for INTEGER-PROGRAMMING

HITTING-SET can be reduced to INTEGER-PROGRAMMING

Proof.

- Formulate the hitting-set as an integer program

Yet Another Hardness Proof for INTEGER-PROGRAMMING

Yet another reduction for INTEGER-PROGRAMMING

HITTING-SET can be reduced to INTEGER-PROGRAMMING

Proof.

- Formulate the hitting-set as an integer program
- Given $\mathcal{C} = \{S_1, \dots, S_m\}$ over $X = \{a_1, \dots, a_n\}$ and k :

Yet Another Hardness Proof for INTEGER-PROGRAMMING

Yet another reduction for INTEGER-PROGRAMMING

HITTING-SET can be reduced to INTEGER-PROGRAMMING

Proof.

- Formulate the hitting-set as an integer program
- Given $\mathcal{C} = \{S_1, \dots, S_m\}$ over $X = \{a_1, \dots, a_n\}$ and k :
- Integer program variables: $x_j = 1$ iff x_j picked in hitting-set

Yet Another Hardness Proof for INTEGER-PROGRAMMING

Yet another reduction for INTEGER-PROGRAMMING

HITTING-SET can be reduced to INTEGER-PROGRAMMING

Proof.

- Formulate the hitting-set as an integer program
- Given $\mathcal{C} = \{S_1, \dots, S_m\}$ over $X = \{a_1, \dots, a_n\}$ and k :
- Integer program variables: $x_j = 1$ iff x_j picked in hitting-set

At most k elements:

Yet Another Hardness Proof for INTEGER-PROGRAMMING

Yet another reduction for INTEGER-PROGRAMMING

HITTING-SET can be reduced to INTEGER-PROGRAMMING

Proof.

- Formulate the hitting-set as an integer program
- Given $\mathcal{C} = \{S_1, \dots, S_m\}$ over $X = \{a_1, \dots, a_n\}$ and k :
- Integer program variables: $x_j = 1$ iff x_j picked in hitting-set

At most k elements:
$$\sum_j x_j \leq k$$

Yet Another Hardness Proof for INTEGER-PROGRAMMING

Yet another reduction for INTEGER-PROGRAMMING

HITTING-SET can be reduced to INTEGER-PROGRAMMING

Proof.

- Formulate the hitting-set as an integer program
- Given $\mathcal{C} = \{S_1, \dots, S_m\}$ over $X = \{a_1, \dots, a_n\}$ and k :
- Integer program variables: $x_j = 1$ iff x_j picked in hitting-set

At most k elements:
$$\sum_j x_j \leq k$$

All sets covered:

Yet Another Hardness Proof for INTEGER-PROGRAMMING

Yet another reduction for INTEGER-PROGRAMMING

HITTING-SET can be reduced to INTEGER-PROGRAMMING

Proof.

- Formulate the hitting-set as an integer program
- Given $\mathcal{C} = \{S_1, \dots, S_m\}$ over $X = \{a_1, \dots, a_n\}$ and k :
- Integer program variables: $x_j = 1$ iff x_j picked in hitting-set

At most k elements:
$$\sum_j x_j \leq k$$

All sets covered:
$$\sum_{j|a_j \in S_i} x_j \geq 1 \quad \forall i$$

Yet Another Hardness Proof for INTEGER-PROGRAMMING

Yet another reduction for INTEGER-PROGRAMMING

HITTING-SET can be reduced to INTEGER-PROGRAMMING

Proof.

- Formulate the hitting-set as an integer program
- Given $\mathcal{C} = \{S_1, \dots, S_m\}$ over $X = \{a_1, \dots, a_n\}$ and k :
- Integer program variables: $x_j = 1$ iff x_j picked in hitting-set

At most k elements:
$$\sum_j x_j \leq k$$

All sets covered:
$$\sum_{j|a_j \in S_i} x_j \geq 1 \quad \forall i$$

Variable 0 or 1:

Yet Another Hardness Proof for INTEGER-PROGRAMMING

Yet another reduction for INTEGER-PROGRAMMING

HITTING-SET can be reduced to INTEGER-PROGRAMMING

Proof.

- Formulate the hitting-set as an integer program
- Given $\mathcal{C} = \{S_1, \dots, S_m\}$ over $X = \{a_1, \dots, a_n\}$ and k :
- Integer program variables: $x_j = 1$ iff x_j picked in hitting-set

At most k elements:
$$\sum_j x_j \leq k$$

All sets covered:
$$\sum_{j|a_j \in S_i} x_j \geq 1 \quad \forall i$$

Variable 0 or 1:
$$x_i \in \{0, 1\}$$



EXACT-COVER

Claim

EXACT-COVER : A collection $\mathcal{C} = \{S_1, \dots, S_n\}$, $|S_j| = 3$ over a base set X , $|X| = 3m$, find a **disjoint** set cover $\mathcal{C}' \subseteq \mathcal{C}$ of size m .

The exact cover by 3-sets problem (**EXACT-COVER**) is **NP** complete.

EXACT-COVER

Claim

EXACT-COVER : A collection $\mathcal{C} = \{S_1, \dots, S_n\}$, $|S_j| = 3$ over a base set X , $|X| = 3m$, find a **disjoint** set cover $\mathcal{C}' \subseteq \mathcal{C}$ of size m .

The exact cover by 3-sets problem (**EXACT-COVER**) is **NP** complete.

- Note that EXACT-COVER is in **NP**

EXACT-COVER

Claim

EXACT-COVER : A collection $\mathcal{C} = \{S_1, \dots, S_n\}$, $|S_j| = 3$ over a base set X , $|X| = 3m$, find a **disjoint** set cover $\mathcal{C}' \subseteq \mathcal{C}$ of size m .

The exact cover by 3-sets problem (**EXACT-COVER**) is **NP** complete.

- Note that EXACT-COVER is in **NP**
- How to prove **NP** hardness?

EXACT-COVER

Claim

EXACT-COVER : A collection $\mathcal{C} = \{S_1, \dots, S_n\}$, $|S_j| = 3$ over a base set X , $|X| = 3m$, find a **disjoint** set cover $\mathcal{C}' \subseteq \mathcal{C}$ of size m .

The exact cover by 3-sets problem (**EXACT-COVER**) is **NP** complete.

- Note that EXACT-COVER is in **NP**
- How to prove **NP** hardness?
- Reduce from SAT . Read from the Papadimitriou book.

KNAPSACK

Claim

***KNAPSACK** : Given a set $A = \{a_1, \dots, a_n\}$ of n elements, where each element a_i has a weight w_i and a value v_i , both positive integers. Find a subset $A' \subseteq A$ such that*

KNAPSACK

Claim

KNAPSACK : Given a set $A = \{a_1, \dots, a_n\}$ of n elements, where each element a_i has a weight w_i and a value v_i , both positive integers. Find a subset $A' \subseteq A$ such that

$$\textcircled{1} \quad \sum_{a_j \in A'} w_j \leq W,$$

KNAPSACK

Claim

KNAPSACK : Given a set $A = \{a_1, \dots, a_n\}$ of n elements, where each element a_i has a weight w_i and a value v_i , both positive integers. Find a subset $A' \subseteq A$ such that

① $\sum_{a_j \in A'} w_j \leq W,$

② $\sum_{a_j \in A'} v_j \geq K.$

KNAPSACK

Claim

KNAPSACK : Given a set $A = \{a_1, \dots, a_n\}$ of n elements, where each element a_i has a weight w_i and a value v_i , both positive integers. Find a subset $A' \subseteq A$ such that

① $\sum_{a_j \in A'} w_j \leq W,$

② $\sum_{a_j \in A'} v_j \geq K.$

The Knapsack problem (**KNAPSACK**) is **NP** complete.

KNAPSACK

Claim

KNAPSACK : Given a set $A = \{a_1, \dots, a_n\}$ of n elements, where each element a_i has a weight w_i and a value v_i , both positive integers. Find a subset $A' \subseteq A$ such that

① $\sum_{a_j \in A'} w_j \leq W,$

② $\sum_{a_j \in A'} v_j \geq K.$

The Knapsack problem (**KNAPSACK**) is **NP** complete.

- Note that **KNAPSACK** is in **NP**

KNAPSACK

Claim

KNAPSACK : Given a set $A = \{a_1, \dots, a_n\}$ of n elements, where each element a_i has a weight w_i and a value v_i , both positive integers. Find a subset $A' \subseteq A$ such that

① $\sum_{a_j \in A'} w_j \leq W,$

② $\sum_{a_j \in A'} v_j \geq K.$

The Knapsack problem (**KNAPSACK**) is **NP** complete.

- Note that **KNAPSACK** is in **NP**
- How to prove **NP** hardness?

KNAPSACK

Claim

KNAPSACK : Given a set $A = \{a_1, \dots, a_n\}$ of n elements, where each element a_i has a weight w_i and a value v_i , both positive integers. Find a subset $A' \subseteq A$ such that

① $\sum_{a_j \in A'} w_j \leq W,$

② $\sum_{a_j \in A'} v_j \geq K.$

The Knapsack problem (**KNAPSACK**) is **NP** complete.

- Note that **KNAPSACK** is in **NP**
- How to prove **NP** hardness?
- Special case: $K = W$ and $w_i = v_i$ for all i .

KNAPSACK

Claim

KNAPSACK : Given a set $A = \{a_1, \dots, a_n\}$ of n elements, where each element a_i has a weight w_i and a value v_i , both positive integers. Find a subset $A' \subseteq A$ such that

① $\sum_{a_j \in A'} w_j \leq W,$

② $\sum_{a_j \in A'} v_j \geq K.$

The Knapsack problem (**KNAPSACK**) is **NP** complete.

- Note that **KNAPSACK** is in **NP**
- How to prove **NP** hardness?
- Special case: $K = W$ and $w_i = v_i$ for all i . Then problem?

KNAPSACK

Claim

KNAPSACK : Given a set $A = \{a_1, \dots, a_n\}$ of n elements, where each element a_i has a weight w_i and a value v_i , both positive integers. Find a subset $A' \subseteq A$ such that

① $\sum_{a_j \in A'} w_j \leq W,$

② $\sum_{a_j \in A'} v_j \geq K.$

The Knapsack problem (**KNAPSACK**) is **NP** complete.

- Note that **KNAPSACK** is in **NP**
- How to prove **NP** hardness?
- Special case: $K = W$ and $w_i = v_i$ for all i . Then problem?
- Subset sum! Find a subset with value and weight equal to W .

SUBSET-SUM

Claim

*The subset sum problem (SUBSET-SUM) is **NP** complete.*

SUBSET-SUM

Claim

*The subset sum problem (SUBSET-SUM) is **NP** complete.*

- Reduce EXACT-COVER to SUBSET-SUM

SUBSET-SUM

Claim

*The subset sum problem (SUBSET-SUM) is **NP** complete.*

- Reduce EXACT-COVER to SUBSET-SUM
- $\{S_1, \dots, S_n\}, |S_j| = 3$ over a base set $X = \{a_1, \dots, a_{3m}\}$

SUBSET-SUM

Claim

*The subset sum problem (SUBSET-SUM) is **NP** complete.*

- Reduce EXACT-COVER to SUBSET-SUM
- $\{S_1, \dots, S_n\}, |S_j| = 3$ over a base set $X = \{a_1, \dots, a_{3m}\}$
- Map each set S_j to an integer t_j

SUBSET-SUM

Claim

*The subset sum problem (SUBSET-SUM) is **NP** complete.*

- Reduce EXACT-COVER to SUBSET-SUM
- $\{S_1, \dots, S_n\}, |S_j| = 3$ over a base set $X = \{a_1, \dots, a_{3m}\}$
- Map each set S_j to an integer t_j
- If S_j 's are disjoint and cover X , then t_j 's add up to some T

SUBSET-SUM

Claim

*The subset sum problem (SUBSET-SUM) is **NP** complete.*

- Reduce EXACT-COVER to SUBSET-SUM
- $\{S_1, \dots, S_n\}, |S_j| = 3$ over a base set $X = \{a_1, \dots, a_{3m}\}$
- Map each set S_j to an integer t_j
- If S_j 's are disjoint and cover X , then t_j 's add up to some T
- For mapping sets into integers, use the characteristic vector

SUBSET-SUM

Claim

*The subset sum problem (SUBSET-SUM) is **NP** complete.*

- Reduce EXACT-COVER to SUBSET-SUM
- $\{S_1, \dots, S_n\}$, $|S_j| = 3$ over a base set $X = \{a_1, \dots, a_{3m}\}$
- Map each set S_j to an integer t_j
- If S_j 's are disjoint and cover X , then t_j 's add up to some T
- For mapping sets into integers, use the characteristic vector
 - ▶ Represent set S as a vector $v(S)$ where $v_i(S) = 1$ iff $a_i \in S$

SUBSET-SUM

Claim

*The subset sum problem (SUBSET-SUM) is **NP** complete.*

- Reduce EXACT-COVER to SUBSET-SUM
- $\{S_1, \dots, S_n\}, |S_j| = 3$ over a base set $X = \{a_1, \dots, a_{3m}\}$
- Map each set S_j to an integer t_j
- If S_j 's are disjoint and cover X , then t_j 's add up to some T
- For mapping sets into integers, use the characteristic vector
 - ▶ Represent set S as a vector $v(S)$ where $v_i(S) = 1$ iff $a_i \in S$
 - ▶ Example: $S = \{a_1, a_4, a_5\}$ out of 6 elements

SUBSET-SUM

Claim

*The subset sum problem (SUBSET-SUM) is **NP** complete.*

- Reduce EXACT-COVER to SUBSET-SUM
- $\{S_1, \dots, S_n\}, |S_j| = 3$ over a base set $X = \{a_1, \dots, a_{3m}\}$
- Map each set S_j to an integer t_j
- If S_j 's are disjoint and cover X , then t_j 's add up to some T
- For mapping sets into integers, use the characteristic vector
 - ▶ Represent set S as a vector $v(S)$ where $v_i(S) = 1$ iff $a_i \in S$
 - ▶ Example: $S = \{a_1, a_4, a_5\}$ out of 6 elements
 - ▶ $v(S) =$

SUBSET-SUM

Claim

*The subset sum problem (SUBSET-SUM) is **NP** complete.*

- Reduce EXACT-COVER to SUBSET-SUM
- $\{S_1, \dots, S_n\}, |S_j| = 3$ over a base set $X = \{a_1, \dots, a_{3m}\}$
- Map each set S_j to an integer t_j
- If S_j 's are disjoint and cover X , then t_j 's add up to some T
- For mapping sets into integers, use the characteristic vector
 - ▶ Represent set S as a vector $v(S)$ where $v_i(S) = 1$ iff $a_i \in S$
 - ▶ Example: $S = \{a_1, a_4, a_5\}$ out of 6 elements
 - ▶ $v(S) = 1\ 0\ 0\ 1\ 1\ 0$

SUBSET-SUM

Claim

*The subset sum problem (SUBSET-SUM) is **NP** complete.*

- Reduce EXACT-COVER to SUBSET-SUM
- $\{S_1, \dots, S_n\}, |S_j| = 3$ over a base set $X = \{a_1, \dots, a_{3m}\}$
- Map each set S_j to an integer t_j
- If S_j 's are disjoint and cover X , then t_j 's add up to some T
- For mapping sets into integers, use the characteristic vector
 - ▶ Represent set S as a vector $v(S)$ where $v_i(S) = 1$ iff $a_i \in S$
 - ▶ Example: $S = \{a_1, a_4, a_5\}$ out of 6 elements
 - ▶ $v(S) = 1\ 0\ 0\ 1\ 1\ 0$
 - ▶ Easy to see that its a one-to-one mapping

An Example

$\{1, 4, 6\}, \{6, 8, 9\}, \{2, 4, 7\}, \{1, 2, 8\}, \{2, 4, 6\}, \{1, 3, 5\}$

An Example

$\{1, 4, 6\}, \{6, 8, 9\}, \{2, 4, 7\}, \{1, 2, 8\}, \{2, 4, 6\}, \{1, 3, 5\}$

1 2 3 4 5 6 7 8 9

An Example

$\{1, 4, 6\}, \{6, 8, 9\}, \{2, 4, 7\}, \{1, 2, 8\}, \{2, 4, 6\}, \{1, 3, 5\}$

1 2 3 4 5 6 7 8 9

1 0 0 1 0 1 0 0 0

An Example

$\{1, 4, 6\}, \{6, 8, 9\}, \{2, 4, 7\}, \{1, 2, 8\}, \{2, 4, 6\}, \{1, 3, 5\}$

1 2 3 4 5 6 7 8 9

1 0 0 1 0 1 0 0 0

0 0 0 0 0 1 0 1 1

An Example

$\{1, 4, 6\}, \{6, 8, 9\}, \{2, 4, 7\}, \{1, 2, 8\}, \{2, 4, 6\}, \{1, 3, 5\}$

1 2 3 4 5 6 7 8 9

1 0 0 1 0 1 0 0 0

0 0 0 0 0 1 0 1 1

0 1 0 1 0 0 1 0 0

An Example

$\{1, 4, 6\}, \{6, 8, 9\}, \{2, 4, 7\}, \{1, 2, 8\}, \{2, 4, 6\}, \{1, 3, 5\}$

1 2 3 4 5 6 7 8 9

1 0 0 1 0 1 0 0 0

0 0 0 0 0 1 0 1 1

0 1 0 1 0 0 1 0 0

1 1 0 0 0 0 0 1 0

0 1 0 1 0 1 0 0 0

1 0 1 0 1 0 0 0 0

An Example

- Lets take the two sets $\{6, 8, 9\}, \{2, 4, 7\}$.
- The corresponding vectors are:

0 0 0 0 0 1 0 1 1

An Example

- Lets take the two sets $\{6, 8, 9\}, \{2, 4, 7\}$.
- The corresponding vectors are:

0	0	0	0	0	1	0	1	1
0	1	0	1	0	0	1	0	0

An Example

- Lets take the two sets $\{6, 8, 9\}$, $\{2, 4, 7\}$.
- The corresponding vectors are:

0	0	0	0	0	1	0	1	1
0	1	0	1	0	0	1	0	0

- The union of the two sets is: $\{2, 4, 6, 7, 8, 9\}$

An Example

- Lets take the two sets $\{6, 8, 9\}$, $\{2, 4, 7\}$.
- The corresponding vectors are:

0	0	0	0	0	1	0	1	1
0	1	0	1	0	0	1	0	0

- The union of the two sets is: $\{2, 4, 6, 7, 8, 9\}$
- The characteristic vector of the union is:

0	1	0	1	0	1	1	1	1
---	---	---	---	---	---	---	---	---

An Example

- Lets take the two sets $\{6, 8, 9\}$, $\{2, 4, 7\}$.
- The corresponding vectors are:

0	0	0	0	0	1	0	1	1
0	1	0	1	0	0	1	0	0

- The union of the two sets is: $\{2, 4, 6, 7, 8, 9\}$
- The characteristic vector of the union is:

0	1	0	1	0	1	1	1	1
---	---	---	---	---	---	---	---	---

- Any observations?

An Example

$\{1, 4, 6\}, \{6, 8, 9\}, \{2, 4, 7\}, \{1, 2, 8\}, \{2, 4, 6\}, \{1, 3, 5\}$

An Example

$\{1, 4, 6\}$, $\{6, 8, 9\}$, $\{2, 4, 7\}$, $\{1, 2, 8\}$, $\{2, 4, 6\}$, $\{1, 3, 5\}$

An Example

$\{1, 4, 6\}, \{6, 8, 9\}, \{2, 4, 7\}, \{1, 2, 8\}, \{2, 4, 6\}, \{1, 3, 5\}$

$$0 \ 0 \ 0 \ 0 \ 0 \ 1 \ 0 \ 1 \ 1 : 2^3 + 2^1 + 2^0 = 011$$

$$0 \ 1 \ 0 \ 1 \ 0 \ 0 \ 1 \ 0 \ 0 : 2^7 + 2^5 + 2^2 = 164$$

$$1 \ 0 \ 1 \ 0 \ 1 \ 0 \ 0 \ 0 \ 0 : 2^8 + 2^6 + 2^4 = 336$$

An Example

$\{1, 4, 6\}, \{6, 8, 9\}, \{2, 4, 7\}, \{1, 2, 8\}, \{2, 4, 6\}, \{1, 3, 5\}$

$$0 \ 0 \ 0 \ 0 \ 0 \ 1 \ 0 \ 1 \ 1 : 2^3 + 2^1 + 2^0 = 011$$

$$0 \ 1 \ 0 \ 1 \ 0 \ 0 \ 1 \ 0 \ 0 : 2^7 + 2^5 + 2^2 = 164$$

$$1 \ 0 \ 1 \ 0 \ 1 \ 0 \ 0 \ 0 \ 0 : 2^8 + 2^6 + 2^4 = 336$$

- The union covers X and the sets are pair-wise disjoint:

An Example

$\{1, 4, 6\}, \{6, 8, 9\}, \{2, 4, 7\}, \{1, 2, 8\}, \{2, 4, 6\}, \{1, 3, 5\}$

$$0 \ 0 \ 0 \ 0 \ 0 \ 1 \ 0 \ 1 \ 1 \quad : 2^3 + 2^1 + 2^0 = 011$$

$$0 \ 1 \ 0 \ 1 \ 0 \ 0 \ 1 \ 0 \ 0 \quad : 2^7 + 2^5 + 2^2 = 164$$

$$1 \ 0 \ 1 \ 0 \ 1 \ 0 \ 0 \ 0 \ 0 \quad : 2^8 + 2^6 + 2^4 = 336$$

- The union covers X and the sets are pair-wise disjoint:

$$1 \ 1 \ 1 \ 1 \ 1 \ 1 \ 1 \ 1 \ 1 \quad : 2^9 - 1 = 511$$

An Example

$\{1, 4, 6\}, \{6, 8, 9\}, \{2, 4, 7\}, \{1, 2, 8\}, \{2, 4, 6\}, \{1, 3, 5\}$

$$0 \ 0 \ 0 \ 0 \ 0 \ 1 \ 0 \ 1 \ 1 \quad : 2^3 + 2^1 + 2^0 = 11$$

$$0 \ 1 \ 0 \ 1 \ 0 \ 0 \ 1 \ 0 \ 0 \quad : 2^7 + 2^5 + 2^2 = 164$$

$$1 \ 0 \ 1 \ 0 \ 1 \ 0 \ 0 \ 0 \ 0 \quad : 2^8 + 2^6 + 2^4 = 336$$

- The union covers X and the sets are pair-wise disjoint:

$$1 \ 1 \ 1 \ 1 \ 1 \ 1 \ 1 \ 1 \ 1 \quad : 2^9 - 1 = 511$$

- $11 + 164 + 336 = 511$.

An Example

$\{1, 4, 6\}, \{6, 8, 9\}, \{2, 4, 7\}, \{1, 2, 8\}, \{2, 4, 6\}, \{1, 3, 5\}$

$$0 \ 0 \ 0 \ 0 \ 0 \ 1 \ 0 \ 1 \ 1 \quad : 2^3 + 2^1 + 2^0 = 011$$

$$0 \ 1 \ 0 \ 1 \ 0 \ 0 \ 1 \ 0 \ 0 \quad : 2^7 + 2^5 + 2^2 = 164$$

$$1 \ 0 \ 1 \ 0 \ 1 \ 0 \ 0 \ 0 \ 0 \quad : 2^8 + 2^6 + 2^4 = 336$$

- The union covers X and the sets are pair-wise disjoint:

$$1 \ 1 \ 1 \ 1 \ 1 \ 1 \ 1 \ 1 \ 1 \quad : 2^9 - 1 = 511$$

- $11 + 164 + 336 = 511$. So set $W = 2^n - 1$, and check.

An Example

$$\{1, 4, 6\}, \{6, 8, 9\}, \{2, 4, 7\}, \{1, 2, 8\}, \{2, 4, 6\}, \{1, 3, 5\}$$

$$0 \ 0 \ 0 \ 0 \ 0 \ 1 \ 0 \ 1 \ 1 \quad : 2^3 + 2^1 + 2^0 = 011$$

$$0 \ 1 \ 0 \ 1 \ 0 \ 0 \ 1 \ 0 \ 0 \quad : 2^7 + 2^5 + 2^2 = 164$$

$$1 \ 0 \ 1 \ 0 \ 1 \ 0 \ 0 \ 0 \ 0 \quad : 2^8 + 2^6 + 2^4 = 336$$

- The union covers X and the sets are pair-wise disjoint:

$$1 \ 1 \ 1 \ 1 \ 1 \ 1 \ 1 \ 1 \ 1 \quad : 2^9 - 1 = 511$$

- $11 + 164 + 336 = 511$. So set $W = 2^n - 1$, and check.
- There exists subset equal to $W \implies$ exact-cover possible

An Example

$\{1, 4, 6\}, \{6, 8, 9\}, \{2, 4, 7\}, \{1, 2, 8\}, \{2, 4, 6\}, \{1, 3, 5\}$

$$0 \ 0 \ 0 \ 0 \ 0 \ 1 \ 0 \ 1 \ 1 \quad : 2^3 + 2^1 + 2^0 = 011$$

$$0 \ 1 \ 0 \ 1 \ 0 \ 0 \ 1 \ 0 \ 0 \quad : 2^7 + 2^5 + 2^2 = 164$$

$$1 \ 0 \ 1 \ 0 \ 1 \ 0 \ 0 \ 0 \ 0 \quad : 2^8 + 2^6 + 2^4 = 336$$

- The union covers X and the sets are pair-wise disjoint:

$$1 \ 1 \ 1 \ 1 \ 1 \ 1 \ 1 \ 1 \ 1 \quad : 2^9 - 1 = 511$$

- $11 + 164 + 336 = 511$. So set $W = 2^n - 1$, and check.
- There exists subset equal to $W \implies$ exact-cover possible
- Problem:

An Example

$\{1, 4, 6\}, \{6, 8, 9\}, \{2, 4, 7\}, \{1, 2, 8\}, \{2, 4, 6\}, \{1, 3, 5\}$

$$0 \ 0 \ 0 \ 0 \ 0 \ 1 \ 0 \ 1 \ 1 \quad : 2^3 + 2^1 + 2^0 = 011$$

$$0 \ 1 \ 0 \ 1 \ 0 \ 0 \ 1 \ 0 \ 0 \quad : 2^7 + 2^5 + 2^2 = 164$$

$$1 \ 0 \ 1 \ 0 \ 1 \ 0 \ 0 \ 0 \ 0 \quad : 2^8 + 2^6 + 2^4 = 336$$

- The union covers X and the sets are pair-wise disjoint:

$$1 \ 1 \ 1 \ 1 \ 1 \ 1 \ 1 \ 1 \ 1 \quad : 2^9 - 1 = 511$$

- $11 + 164 + 336 = 511$. So set $W = 2^n - 1$, and check.
- There exists subset equal to $W \implies$ exact-cover possible
- **Problem:** False positives.

A Problem

$$\{3, 4\}, \{2, 4\}, \{2, 3, 4\}$$

A Problem

$$\{3, 4\}, \{2, 4\}, \{2, 3, 4\}$$

$$0 \ 0 \ 1 \ 1 \ : 3$$

A Problem

$$\{3, 4\}, \{2, 4\}, \{2, 3, 4\}$$

$$0 \ 0 \ 1 \ 1 \ : 3$$

$$0 \ 1 \ 0 \ 1 \ : 5$$

A Problem

$$\{3, 4\}, \{2, 4\}, \{2, 3, 4\}$$

$$0 \ 0 \ 1 \ 1 \ : 3$$

$$0 \ 1 \ 0 \ 1 \ : 5$$

$$0 \ 1 \ 1 \ 1 \ : 7$$

A Problem

$$\{3, 4\}, \{2, 4\}, \{2, 3, 4\}$$

0 0 1 1 : 3

0 1 0 1 : 5

0 1 1 1 : 7

- The union does **not** cover X , and **not disjoint**.

A Problem

$$\{3, 4\}, \{2, 4\}, \{2, 3, 4\}$$

$$0 \ 0 \ 1 \ 1 \ : 3$$

$$0 \ 1 \ 0 \ 1 \ : 5$$

$$0 \ 1 \ 1 \ 1 \ : 7$$

- The union does **not** cover X , and **not disjoint**.

$$0 \ 0 \ 1 \ 1 \ + \ 0 \ 1 \ 0 \ 1 \ + \ 0 \ 1 \ 1 \ 1$$

A Problem

$$\{3, 4\}, \{2, 4\}, \{2, 3, 4\}$$

$$0 \ 0 \ 1 \ 1 \ : 3$$

$$0 \ 1 \ 0 \ 1 \ : 5$$

$$0 \ 1 \ 1 \ 1 \ : 7$$

- The union does **not** cover X , and **not disjoint**.

$$0 \ 0 \ 1 \ 1 \ + \ 0 \ 1 \ 0 \ 1 \ + \ 0 \ 1 \ 1 \ 1 \ = \ 1 \ 1 \ 1 \ 1$$

A Problem

$$\{3, 4\}, \{2, 4\}, \{2, 3, 4\}$$

$$0 \ 0 \ 1 \ 1 \ : 3$$

$$0 \ 1 \ 0 \ 1 \ : 5$$

$$0 \ 1 \ 1 \ 1 \ : 7$$

- The union does **not** cover X , and **not disjoint**.

$$0 \ 0 \ 1 \ 1 \ + \ 0 \ 1 \ 0 \ 1 \ + \ 0 \ 1 \ 1 \ 1 \ = \ 1 \ 1 \ 1 \ 1 \ = 15$$

A Problem

$$\{3, 4\}, \{2, 4\}, \{2, 3, 4\}$$

$$0 \ 0 \ 1 \ 1 \ : 3$$

$$0 \ 1 \ 0 \ 1 \ : 5$$

$$0 \ 1 \ 1 \ 1 \ : 7$$

- The union does **not** cover X , and **not disjoint**.

$$0 \ 0 \ 1 \ 1 \ + \ 0 \ 1 \ 0 \ 1 \ + \ 0 \ 1 \ 1 \ 1 \ = \ 1 \ 1 \ 1 \ 1 \ = 15$$

- So, where is the problem coming from?

A Problem

$$\{3, 4\}, \{2, 4\}, \{2, 3, 4\}$$

$$0 \ 0 \ 1 \ 1 \ : 3$$

$$0 \ 1 \ 0 \ 1 \ : 5$$

$$0 \ 1 \ 1 \ 1 \ : 7$$

- The union does **not** cover X , and **not disjoint**.

$$0 \ 0 \ 1 \ 1 \ + \ 0 \ 1 \ 0 \ 1 \ + \ 0 \ 1 \ 1 \ 1 \ = \ 1 \ 1 \ 1 \ 1 \ = 15$$

- So, where is the problem coming from?
- **Carrying!** It messes up the addition.

A Solution

- There are actually two different problems due to carrying:

A Solution

- There are actually two different problems due to carrying:
 - ▶ If a column has all 0's, could get 1 from carry-over

A Solution

- There are actually two different problems due to carrying:
 - ▶ If a column has all 0's, could get 1 from carry-over
 - ▶ If a column has more than one 1, could still get one 1

A Solution

- There are actually two different problems due to carrying:
 - ▶ If a column has all 0's, could get 1 from carry-over
 - ▶ If a column has more than one 1, could still get one 1
- **Solution:**

A Solution

- There are actually two different problems due to carrying:
 - ▶ If a column has all 0's, could get 1 from carry-over
 - ▶ If a column has more than one 1, could still get one 1
- **Solution:** Interpret vector over base $3m$, not base 2!

A Solution

- There are actually two different problems due to carrying:
 - ▶ If a column has all 0's, could get 1 from carry-over
 - ▶ If a column has more than one 1, could still get one 1
- **Solution:** Interpret vector over base $3m$, not base 2!
 - ▶ $\{2, 3, 4\} = 0\ 1\ 1\ 1 =$

A Solution

- There are actually two different problems due to carrying:
 - ▶ If a column has all 0's, could get 1 from carry-over
 - ▶ If a column has more than one 1, could still get one 1
- **Solution:** Interpret vector over base $3m$, not base 2!
 - ▶ $\{2, 3, 4\} = 0 \ 1 \ 1 \ 1 = 2^2 + 2^1 + 2^0,$

A Solution

- There are actually two different problems due to carrying:
 - ▶ If a column has all 0's, could get 1 from carry-over
 - ▶ If a column has more than one 1, could still get one 1
- **Solution:** Interpret vector over base $3m$, not base 2!
 - ▶ $\{2, 3, 4\} = 0 \ 1 \ 1 \ 1 = 2^2 + 2^1 + 2^0, \quad 5^2 + 5^1 + 5^0 = 31$

A Solution

- There are actually two different problems due to carrying:
 - ▶ If a column has all 0's, could get 1 from carry-over
 - ▶ If a column has more than one 1, could still get one 1
- **Solution:** Interpret vector over base $3m$, not base 2!
 - ▶ $\{2, 3, 4\} = 0\ 1\ 1\ 1 = 2^2 + 2^1 + 2^0, \quad 5^2 + 5^1 + 5^0 = 31$
 - ▶ $\{3, 4\} = 0\ 0\ 1\ 1 =$

A Solution

- There are actually two different problems due to carrying:
 - ▶ If a column has all 0's, could get 1 from carry-over
 - ▶ If a column has more than one 1, could still get one 1
- **Solution:** Interpret vector over base $3m$, not base 2!
 - ▶ $\{2, 3, 4\} = 0\ 1\ 1\ 1 = 2^2 + 2^1 + 2^0, \quad 5^2 + 5^1 + 5^0 = 31$
 - ▶ $\{3, 4\} = 0\ 0\ 1\ 1 = 2^1 + 2^0,$

A Solution

- There are actually two different problems due to carrying:
 - ▶ If a column has all 0's, could get 1 from carry-over
 - ▶ If a column has more than one 1, could still get one 1
- **Solution:** Interpret vector over base $3m$, not base 2!
 - ▶ $\{2, 3, 4\} = 0\ 1\ 1\ 1 = 2^2 + 2^1 + 2^0, \quad 5^2 + 5^1 + 5^0 = 31$
 - ▶ $\{3, 4\} = 0\ 0\ 1\ 1 = 2^1 + 2^0, \quad 5^1 + 5^0 = 6$

A Solution

- There are actually two different problems due to carrying:
 - ▶ If a column has all 0's, could get 1 from carry-over
 - ▶ If a column has more than one 1, could still get one 1
- **Solution:** Interpret vector over base $3m$, not base 2!
 - ▶ $\{2, 3, 4\} = 0\ 1\ 1\ 1 = 2^2 + 2^1 + 2^0, \quad 5^2 + 5^1 + 5^0 = 31$
 - ▶ $\{3, 4\} = 0\ 0\ 1\ 1 = 2^1 + 2^0, \quad 5^1 + 5^0 = 6$
 - ▶ $\{2, 4\} = 0\ 1\ 0\ 1 =$

A Solution

- There are actually two different problems due to carrying:
 - ▶ If a column has all 0's, could get 1 from carry-over
 - ▶ If a column has more than one 1, could still get one 1
- **Solution:** Interpret vector over base $3m$, not base 2!
 - ▶ $\{2, 3, 4\} = 0\ 1\ 1\ 1 = 2^2 + 2^1 + 2^0, \quad 5^2 + 5^1 + 5^0 = 31$
 - ▶ $\{3, 4\} = 0\ 0\ 1\ 1 = 2^1 + 2^0, \quad 5^1 + 5^0 = 6$
 - ▶ $\{2, 4\} = 0\ 1\ 0\ 1 = 2^2 + 2^0,$

A Solution

- There are actually two different problems due to carrying:
 - ▶ If a column has all 0's, could get 1 from carry-over
 - ▶ If a column has more than one 1, could still get one 1
- **Solution:** Interpret vector over base $3m$, not base 2!
 - ▶ $\{2, 3, 4\} = 0\ 1\ 1\ 1 = 2^2 + 2^1 + 2^0, \quad 5^2 + 5^1 + 5^0 = 31$
 - ▶ $\{3, 4\} = 0\ 0\ 1\ 1 = 2^1 + 2^0, \quad 5^1 + 5^0 = 6$
 - ▶ $\{2, 4\} = 0\ 1\ 0\ 1 = 2^2 + 2^0, \quad 5^2 + 5^0 = 26$

A Solution

- There are actually two different problems due to carrying:
 - ▶ If a column has all 0's, could get 1 from carry-over
 - ▶ If a column has more than one 1, could still get one 1
- **Solution:** Interpret vector over base $3m$, not base 2!
 - ▶ $\{2, 3, 4\} = 0\ 1\ 1\ 1 = 2^2 + 2^1 + 2^0, \quad 5^2 + 5^1 + 5^0 = 31$
 - ▶ $\{3, 4\} = 0\ 0\ 1\ 1 = 2^1 + 2^0, \quad 5^1 + 5^0 = 6$
 - ▶ $\{2, 4\} = 0\ 1\ 0\ 1 = 2^2 + 2^0, \quad 5^2 + 5^0 = 26$

$$0\ 0\ 1\ 1 + 0\ 1\ 0\ 1 + 0\ 1\ 1\ 1$$

A Solution

- There are actually two different problems due to carrying:
 - ▶ If a column has all 0's, could get 1 from carry-over
 - ▶ If a column has more than one 1, could still get one 1
- **Solution:** Interpret vector over base $3m$, not base 2!
 - ▶ $\{2, 3, 4\} = 0\ 1\ 1\ 1 = 2^2 + 2^1 + 2^0, \quad 5^2 + 5^1 + 5^0 = 31$
 - ▶ $\{3, 4\} = 0\ 0\ 1\ 1 = 2^1 + 2^0, \quad 5^1 + 5^0 = 6$
 - ▶ $\{2, 4\} = 0\ 1\ 0\ 1 = 2^2 + 2^0, \quad 5^2 + 5^0 = 26$

$$0\ 0\ 1\ 1 + 0\ 1\ 0\ 1 + 0\ 1\ 1\ 1 = 0\ 2\ 2\ 3$$

A Solution

- There are actually two different problems due to carrying:
 - ▶ If a column has all 0's, could get 1 from carry-over
 - ▶ If a column has more than one 1, could still get one 1
- **Solution:** Interpret vector over base $3m$, not base 2!
 - ▶ $\{2, 3, 4\} = 0\ 1\ 1\ 1 = 2^2 + 2^1 + 2^0, \quad 5^2 + 5^1 + 5^0 = 31$
 - ▶ $\{3, 4\} = 0\ 0\ 1\ 1 = 2^1 + 2^0, \quad 5^1 + 5^0 = 6$
 - ▶ $\{2, 4\} = 0\ 1\ 0\ 1 = 2^2 + 2^0, \quad 5^2 + 5^0 = 26$

$$0\ 0\ 1\ 1 + 0\ 1\ 0\ 1 + 0\ 1\ 1\ 1 = 0\ 2\ 2\ 3 = 63 \neq 1\ 1\ 1\ 1$$

A Solution

- There are actually two different problems due to carrying:
 - ▶ If a column has all 0's, could get 1 from carry-over
 - ▶ If a column has more than one 1, could still get one 1

- **Solution:** Interpret vector over base $3m$, not base 2!

- ▶ $\{2, 3, 4\} = 0\ 1\ 1\ 1 = 2^2 + 2^1 + 2^0, \quad 5^2 + 5^1 + 5^0 = 31$
- ▶ $\{3, 4\} = 0\ 0\ 1\ 1 = 2^1 + 2^0, \quad 5^1 + 5^0 = 6$
- ▶ $\{2, 4\} = 0\ 1\ 0\ 1 = 2^2 + 2^0, \quad 5^2 + 5^0 = 26$

$$0\ 0\ 1\ 1 + 0\ 1\ 0\ 1 + 0\ 1\ 1\ 1 = 0\ 2\ 2\ 3 = 63 \neq 1\ 1\ 1\ 1$$

- Adding at most n 1's \implies no carry-over at any digit

A Solution

- There are actually two different problems due to carrying:
 - ▶ If a column has all 0's, could get 1 from carry-over
 - ▶ If a column has more than one 1, could still get one 1

- **Solution:** Interpret vector over base $3m$, not base 2!

- ▶ $\{2, 3, 4\} = 0\ 1\ 1\ 1 = 2^2 + 2^1 + 2^0, \quad 5^2 + 5^1 + 5^0 = 31$
- ▶ $\{3, 4\} = 0\ 0\ 1\ 1 = 2^1 + 2^0, \quad 5^1 + 5^0 = 6$
- ▶ $\{2, 4\} = 0\ 1\ 0\ 1 = 2^2 + 2^0, \quad 5^2 + 5^0 = 26$

$$0\ 0\ 1\ 1 + 0\ 1\ 0\ 1 + 0\ 1\ 1\ 1 = 0\ 2\ 2\ 3 = 63 \neq 1\ 1\ 1\ 1$$

- Adding at most n 1's \implies no carry-over at any digit
- This solves both the problems since:

A Solution

- There are actually two different problems due to carrying:
 - ▶ If a column has all 0's, could get 1 from carry-over
 - ▶ If a column has more than one 1, could still get one 1

- **Solution:** Interpret vector over base $3m$, not base 2!

- ▶ $\{2, 3, 4\} = 0\ 1\ 1\ 1 = 2^2 + 2^1 + 2^0, \quad 5^2 + 5^1 + 5^0 = 31$
- ▶ $\{3, 4\} = 0\ 0\ 1\ 1 = 2^1 + 2^0, \quad 5^1 + 5^0 = 6$
- ▶ $\{2, 4\} = 0\ 1\ 0\ 1 = 2^2 + 2^0, \quad 5^2 + 5^0 = 26$

$$0\ 0\ 1\ 1 + 0\ 1\ 0\ 1 + 0\ 1\ 1\ 1 = 0\ 2\ 2\ 3 = 63 \neq 1\ 1\ 1\ 1$$

- Adding at most n 1's \implies no carry-over at any digit
- This solves both the problems since:
 - ▶ If a column has all 0's, no carry-over ensures it gets a 0.

A Solution

- There are actually two different problems due to carrying:
 - ▶ If a column has all 0's, could get 1 from carry-over
 - ▶ If a column has more than one 1, could still get one 1

- **Solution:** Interpret vector over base $3m$, not base 2!

- ▶ $\{2, 3, 4\} = 0\ 1\ 1\ 1 = 2^2 + 2^1 + 2^0, \quad 5^2 + 5^1 + 5^0 = 31$
- ▶ $\{3, 4\} = 0\ 0\ 1\ 1 = 2^1 + 2^0, \quad 5^1 + 5^0 = 6$
- ▶ $\{2, 4\} = 0\ 1\ 0\ 1 = 2^2 + 2^0, \quad 5^2 + 5^0 = 26$

$$0\ 0\ 1\ 1 + 0\ 1\ 0\ 1 + 0\ 1\ 1\ 1 = 0\ 2\ 2\ 3 = 63 \neq 1\ 1\ 1\ 1$$

- Adding at most n 1's \implies no carry-over at any digit
- This solves both the problems since:
 - ▶ If a column has all 0's, no carry-over ensures it gets a 0.
 - ▶ If a column has more than one 1, it will get the sum, not 1.

Reductions So Far

