

UiO : **University of Oslo**

Giovanni Domenico Di Salvo

**Proper Holomorphic Embeddings  
of open Riemann Surfaces into  $\mathbb{C}^2$   
and holomorphic mappings  
between complex manifolds with  
dense images**

**Thesis submitted for the degree of Philosophiae Doctor**

Department of Mathematics

Faculty of Mathematics and Natural Sciences



**2023**

© Giovanni Domenico Di Salvo, 2023

*Series of dissertations submitted to the  
Faculty of Mathematics and Natural Sciences, University of Oslo  
No. 1234*

ISSN 1234-5678

All rights reserved. No part of this publication may be  
reproduced or transmitted, in any form or by any means, without permission.

Cover: Hanne Baadsgaard Utigard.

Print production: Reprosentralen, University of Oslo.

*Alla prima parte della mia vita.*

---

*It's time to hear the call  
of all you are  
of all you are!*

SMKC, Boulevard of Broken Hearts

# Preface

This thesis is submitted in partial fulfillment of the requirements for the degree of *Philosophiae Doctor* at the University of Oslo. The research presented here was conducted at the University of Oslo, under the supervision of professor Erlend Fornæss Wold and professor Tuyen Trung Truong.

The thesis is a collection of four papers, presented in chronological order of writing. The papers are preceded by an introductory chapter that provides background information and motivation for the work.

## Acknowledgements

First of all I wish to warmly thank my advisors Erlend F. Wold, for all the discussions (mathematical and otherwise) all the patience and the wise guidance; and Tuyen Trung Truong for giving me this great occasion and for always being available to talk. Thanks to Erik Løv for all the interesting and constructive discussions.

Then all the other colleagues of section 6, providing such a truly joyful environment; in particular Makoto Yamashita for being so available.

Thanks to my collaborator Tyson Ritter and to Karoline Moe for the kind help with L<sup>A</sup>T<sub>E</sub>X.

Thanks to all the people that contributed to help me achieving such a great goal: my friends, my music, my family.

This is not the way I would write anything personal; the way I feel to express my gratitude does not match with any Ph.D. thesis acknowledgement format; I will do it in a different context.

✶ **Giovanni Domenico Di Salvo**

Oslo, June 2023

# List of Papers

## Paper I

G. D. Di Salvo and E. F. Wold. “Proper Holomorphic Embeddings of complements of large Cantor sets in  $\mathbb{C}^2$ ”. Submitted for publication to *Arkiv för Matematik*.

## Paper II

G. D. Di Salvo. “Extended explanation of Orevkov’s paper on proper holomorphic embeddings of complements of Cantor sets in  $\mathbb{C}^2$  and a discussion of their measure”. To be submitted.

## Paper III

G. D. Di Salvo, T. Ritter and E. F. Wold. “Families of Proper Holomorphic Embeddings and Carleman–type Theorems with parameters”. Submitted for publication to *The Journal of Geometric Analysis*.

## Paper IV

G. D. Di Salvo. “Approximation and accumulation results of holomorphic mappings with dense image”. Submitted for publication to *Mathematica Scandinavica*.





# Contents

Preface	iii
List of Papers	v
Contents	vii
Notation	ix
<b>1 Introduction</b>	<b>1</b>
1.1 Historical background and some of the main results in Complex Analysis . . . . .	1
1.2 Approximation Theory . . . . .	3
1.3 Several Complex Variables . . . . .	6
1.4 Stein Manifolds . . . . .	8
1.5 Proper Holomorphic Embeddings . . . . .	10
<b>Bibliography</b>	<b>15</b>
1.6 Summary of Papers . . . . .	20
<b>Papers</b>	<b>22</b>
<b>I Proper Holomorphic Embeddings of complements of large Cantor sets in <math>\mathbb{C}^2</math></b>	<b>23</b>
I.1 Introduction . . . . .	23
I.2 Preliminaries . . . . .	25
I.3 The Induction Step . . . . .	27
I.4 Proof of Theorem I.1.1 . . . . .	30
<b>Bibliography</b>	<b>33</b>
<b>II Extended explanation of Orevkov’s paper on proper holomorphic embeddings of complements of Cantor sets in <math>\mathbb{C}^2</math> and a discussion of their measure</b>	<b>35</b>
II.1 Introduction . . . . .	35
II.2 Definitions and preliminary results . . . . .	36
II.3 General Properties . . . . .	40
II.4 The Cantor set $C$ and the proper holomorphic embedding $\gamma$	45
<b>Bibliography</b>	<b>51</b>
	vii

<b>III</b>	<b>Families of Proper Holomorphic Embeddings and Carleman–type Theorems with parameters</b>	<b>53</b>
III.1	Introduction . . . . .	53
III.2	The Setup . . . . .	54
III.3	Carleman approximation with parameters . . . . .	56
III.4	Andersén–Lempert Theory . . . . .	65
III.5	Proof of Theorem III.1.1 . . . . .	69
	<b>Bibliography</b>	<b>71</b>
<b>IV</b>	<b>Approximation and accumulation results of holomorphic mappings with dense image</b>	<b>73</b>
IV.1	Introduction . . . . .	73
IV.2	Main Results . . . . .	74
IV.3	Technical Tools . . . . .	75
IV.4	Inductive procedure . . . . .	77
IV.5	Proofs . . . . .	82
	<b>Bibliography</b>	<b>85</b>

# Notation

We will denote by  $\mathbb{C}^n$  the usual  $n$ -dimensional complex euclidean space, endowed with the euclidean topology and distance, whose coordinates are  $z = (z_1, \dots, z_n)$ . We denote by  $\mathbb{B}^n := \{z \in \mathbb{C}^n : |z_1|^2 + \dots + |z_n|^2 < 1\}$  the open unit ball in  $\mathbb{C}^n$ ; correspondingly  $R\mathbb{B}^n := \{z \in \mathbb{C}^n : |z_1|^2 + \dots + |z_n|^2 < R^2\}$  is the  $R$ -ball, for  $R > 0$ . When  $n = 1$  we might write  $\triangle$  and  $\triangle_R$  instead of  $\mathbb{B}$  and  $R\mathbb{B}$  respectively.

Given a set  $A$  inside a complex manifold  $X$ , we will denote its topological interior, closure, boundary and the set of its accumulation points by writing  $A^\circ$  (equiv.  $\text{int } A$ ),  $\overline{A}$  (equiv.  $\text{cl } A$ ),  $\partial A$ , and  $\text{Acc } A$  respectively, with respect to the topology of  $X$  (or to some induced topology, it will be clear by the context). When  $A = \mathbb{C}^n$ , then  $\overline{\mathbb{C}^n} = \mathbb{C}^n \cup \{\infty_n\}$  denotes the one-point compactification; in particular for  $n = 1$  we agree to write  $\infty_1 = \infty$  and we get the Riemann sphere  $\overline{\mathbb{C}}$  which is equivalently denoted as the projective complex line  $\mathbb{P}^1(\mathbb{C})$ .

We will refer to a *domain* or to a *region* in  $\mathbb{C}^n$  when dealing with an open connected set.

Given a domain  $D \subset \mathbb{C}^n$  (a complex manifold  $X$  respectively) we will denote by  $\mathcal{O}(D)$  (by  $\mathcal{O}(X)$  respectively) the  $\mathbb{C}$ -algebra of all complex-valued holomorphic functions on  $D$  (on  $X$  respectively). When the target space is any  $Y$  different than  $\mathbb{C}$ , we will highlight it by writing  $\mathcal{O}(X, Y)$ . If  $K$  is compact,  $\mathcal{O}(K)$  will denote the  $\mathbb{C}$ -algebra of all functions holomorphic on some unspecified open neighborhood of  $K$ .  $\mathcal{C}^k(D)$  will denote the space of the  $k$ -times continuously differentiable functions, with  $k \in \mathbb{N} \cup \{\infty\}$  and  $\mathcal{C}_c(D)$  will denote the algebra of continuous functions with compact support. Denote by  $\text{Aut } X$  the group (with respect to the composition) of all holomorphic automorphisms  $X \rightarrow X$ .

Given a compact  $K$  in a complex manifold  $X$ , we define its  $\mathcal{O}(X)$ -convex hull as

$$\widehat{K}_{\mathcal{O}(X)} := \{p \in X : |f(p)| \leq \|f\|_K \ \forall f \in \mathcal{O}(X)\}.$$

A compact  $K$  is said  $\mathcal{O}(X)$ -convex if  $K = \widehat{K}_{\mathcal{O}(X)}$  (*polynomially convex* when  $X = \mathbb{C}^n$ ) and a complex manifold  $X$  is said *holomorphically convex* when  $\widehat{K}_{\mathcal{O}(X)}$  is relatively compact for every compact  $K \subset X$ .

Given a metric space  $(X, d)$ , a subset  $A \subset X$  and a positive number  $\delta$ , we define the *open  $\delta$ -neighborhood* of  $A$  as  $A(\delta) := \{x \in X : d(x, A) < \delta\}$ .



# Chapter 1

## Introduction

The main concern of the present thesis is the investigation of proper holomorphic embeddings of certain open Riemann Surfaces into  $\mathbb{C}^2$ . To give to the reader the big picture of the work done, we present here a general overview of Complex Analysis, the area in which the subject takes place. We will start from the historical background, making a quick journey throughout the subject, spotting the main milestones from the very beginning to the nowadays state of art in the direction of the proper holomorphic embeddings area. In particular, we will dwell on the Approximation Theory, as some of the concerning results, besides being interesting on their own, turn out to be a fundamental tool for one of the main Theorems presented in the thesis.

### 1.1 Historical background and some of the main results in Complex Analysis

The first jolts of Complex Analysis –historically known as *Function Theory*– can be traced back to the 18th century with L. Euler (1707–1783) who exploited complex numbers to establish, among other results, the well-known connection between circular functions and exponential function:

$$e^{it} = \cos t + i \sin t, \quad t \in \mathbb{R}.$$

C. F. Gauss (1777–1855) was aware of both the importance and beauty of the subject, as some memorable lines he wrote to Bessel on December 18th, 1811 testify (see [50]). In particular, he knew about the Cauchy integral formula back then. However, neither Euler nor Gauss participate in the development of Complex Analysis in a systematic way, which took place in the 19th century by A.L Cauchy (1789–1857), B. Riemann (1826–1866), and K. Weierstrass (1815–1897). The three of them contributed with quite different methodologies.

Cauchy followed an analytical approach as to him a holomorphic function was a complex-differentiable function with continuous derivative; his discussion is based on the residue theorem and on the integral formula named after him:

**Theorem 1.1.1** (Cauchy Integral Formula). Let  $D \subset \mathbb{C}$  be a simply connected region and  $\gamma \subset D$  a Jordan closed curve. Then any function  $f \in \mathcal{O}(D)$  satisfies

$$f(z) = \frac{1}{2\pi i} \int_{\gamma} \frac{f(\zeta)}{\zeta - z} d\zeta \quad (1.1.1)$$

for any  $z$  belonging to the open bounded set of  $\mathbb{C}$  identified by such a curve.

Riemann had a more geometric point of view, seeing holomorphic (=conformal) functions as mappings between domains of the complex plane, trying to

## 1. Introduction

---

understand such objects searching for some “intrinsic characteristic” properties rather with formulas (which anyhow raised as an extrinsic representation for his intuitions): he classified the non-empty simply connected domains of the complex plane, in the following, well-celebrated milestone, one of his main results in the field.

**Theorem 1.1.2** (Riemann Mapping Theorem). Any non-empty and simply connected region  $D \subsetneq \mathbb{C}$  is biholomorphically equivalent to the unit ball  $\mathbb{B}$ .

The nature of Weierstrass’ contribution has been more algebraic as he saw holomorphic functions as those which can be developed into convergent power series. In particular, he systematized the work by Lagrange, building a whole rigorous theoretical apparatus on it.

It soon turned out how much these three apparently different approaches to Complex Analysis are indeed intertwined: in 1831 Cauchy exploited his integral representation to express holomorphic functions as power series, used by Riemann as well. On the other side, in 1841 Weierstrass exploited the integral representation to develop holomorphic function on annular domains  $\{r < |z - z_0| < R\} \subset \mathbb{C}$  (where  $0 \leq r < R \leq +\infty$ ) into the so-called *Laurent series*, that is the double-sided series:

$$f(z) = \sum_{n \geq 0} c_n (z - z_0)^n + \sum_{n > 0} \frac{c_{-n}}{(z - z_0)^n} . \quad (1.1.2)$$

Some of the major achievements in Complex Analysis include the

**Theorem 1.1.3** (Liouville, 1844). Every bounded entire function is constant.

From such a result the power of the subject starts to appear as it allows to prove the Fundamental Theorem of Algebra in one line: if a polynomial  $p$  never vanishes,  $1/p$  is entire and bounded, hence constant, so  $p$  is. Another fundamental result is the

**Theorem 1.1.4** (Identity Principle). Consider  $f \in \mathcal{O}(D)$  and  $g \in \mathcal{O}(D')$ , where  $D, D' \subset \mathbb{C}$  are two domains such that  $D \cap D'$  is non-empty and connected. If  $f = g$  on some  $S \subset D \cap D'$  containing an accumulation point, then  $f$  and  $g$  agree on the whole  $D \cap D'$ .

Hence holomorphic functions are very rigid; for example, they are uniquely identified by their behavior on the real line. In particular, this theorem allows to uniquely define a holomorphic function which extends  $f$  (or equivalently  $g$ ) to the bigger domain  $D \cup D'$ . The phenomenon of holomorphic extension becomes very tricky and interesting in higher dimensions; it will be discussed later. Another strong property of holomorphic functions pointing out deep differences between real and complex world is the regularity: holomorphic functions are infinitely many times complex-differentiable, as one can easily see by differentiating the Cauchy Integral Formula, against every kind of real regularity. Moreover, holomorphic functions are analytic, as one can see by expanding  $\frac{1}{\zeta - z} = \sum_{n \geq 0} \frac{z^n}{\zeta^{n+1}}$  in the Cauchy Integral Formula and interchanging

the series with the integral. This is not true for smooth functions in the real sense:  $f(x) = e^{-1/x^2}$ ,  $x \neq 0$  and  $f(0) = 0$  is  $\mathcal{C}^\infty(\mathbb{R})$  but it is not the sum of its Taylor series around 0 as  $f^{(k)}(0) = 0$  for all  $k$ . Behavior in a punctured neighborhood of isolated singularities may sound weird for those who come from the real world: given  $z_0$  a point inside a domain  $D$ , then  $f \in \mathcal{O}(D \setminus \{z_0\})$  admits a Laurent representation near  $z_0$ . The second series in (1.1.2) is called the *principal part* as it describes the behavior of  $f$  near  $z_0$ . In particular we distinguish three cases:

- $c_{-n} = 0$  for all  $n > 0$ ;  $z_0$  is called *removable singularity*, this happens if and only if  $f$  is bounded and holomorphic near  $z_0$  as proved by Riemann;
- $c_{-n} \neq 0$  for a finite number of  $n$ ;  $z_0$  is then a *pole* of order  $m$  (being  $m$  the biggest of such  $n$ ). This is equivalent to have  $(z - z_0)^m f(z)$  bounded and non zero near  $z_0$ ;
- $c_{-n} \neq 0$  for infinitely many values of  $n$ . In this case  $z_0$  is called *essential singularity*.

In the last case the following Theorem holds:

**Theorem 1.1.5** (Big Picard Theorem). Let  $f \in \mathcal{O}(D \setminus \{z_0\})$  with  $z_0$  essential singularity. Then in every neighborhood of  $z_0$  the function  $f$  takes every complex value, with at most one exception, infinitely many times.

At this point, it is worth mentioning thus

**Theorem 1.1.6** (Little Picard Theorem, strengthened). An entire function that is not a polynomial takes every complex value, with at most one exception, infinitely many times.

A special role is played by the coefficient  $c_{-1}$ . It is called *residue* of  $f$  at  $z_0$ , denoted also as  $\text{Res}(f, z_0)$  and it is the main character of one of the most important theorems in the field, which allows to easily compute a large number of integrals:

**Theorem 1.1.7** (Residue Theorem). Let  $D$  be a simply connected region in  $\mathbb{C}$ ,  $\gamma \subset D$  be a closed Jordan curve and  $S$  a finite set contained into the bounded open region defined by  $\gamma$ . If  $f \in \mathcal{O}(D \setminus S)$ , then

$$\frac{1}{2\pi i} \int_{\gamma} f(\zeta) d\zeta = \sum_{s \in S} \text{Res}(f, s) .$$

## 1.2 Approximation Theory

Approximation theory plays a very important role in Complex Analysis, as it allows to construct holomorphic maps with certain desired properties. The proof of the main Theorem in [15] raised the necessity of the formulation and proof of an approximation theorem; to introduce it, we collect here some of the main

goals achieved in the branch, following the presentation provided in [18]. This area of investigation has its roots in two classical theorems from 1885.

The first one deals with approximating continuous functions with polynomials on compact intervals of  $\mathbb{R}$ , which is quite remarkable as an arbitrary continuous function could behave very wildly, whether holomorphic functions are extremely regular.

**Theorem 1.2.1** (Weierstrass (1885), [57]). Let  $f$  be a continuous function on the compact interval  $[a, b] \subset \mathbb{R}$ . Then for every  $\epsilon > 0$  there exists a holomorphic polynomial  $p \in \mathbb{C}[z]$  such that  $\|f - p\|_{[a,b]} < \epsilon$ .

It is worth sketching the proof, as the technique is exploited both in [38] and in [15]: we consider the family of entire functions defined as

$$f_t(z) := \frac{1}{t\sqrt{\pi}} \int_{\mathbb{R}} f(x) e^{-(x-z)^2/t^2} dx, \quad z \in \mathbb{C}, \quad t > 0 \quad (1.2.1)$$

after having continuously extended  $f$  as a function in  $\mathcal{C}_c(\mathbb{R})$ . Then  $f_t \rightarrow f$  uniformly on compacts of  $\mathbb{R}$  as  $t \rightarrow 0$ . The Taylor polynomial  $T_N(f_t)$  of  $f_t$  of order  $N$  does the job for  $t$  small and  $N$  large enough.

The second theorem we were referring to, concerns approximation of holomorphic functions on a neighborhood of a compact in the complex plane by either holomorphic polynomials or by rational functions, depending on whether  $K$  is polynomially convex or not:

**Theorem 1.2.2** (Runge (1885), [53]). Let  $K \subset \mathbb{C}$  be compact and  $f \in \mathcal{O}(K)$ . Then  $f$  can be uniformly approximated on  $K$  by rational functions without poles in  $K$  and by holomorphic polynomials if  $\mathbb{C} \setminus K$  is connected.

The more delicate case of approximating functions which are continuous on  $K$  and holomorphic only on its interior  $K^\circ$  (let us denote the set of such functions  $\mathcal{C}(K) \cap \mathcal{O}(K^\circ)$  by  $\mathcal{A}(K)$ ) has been solved by Mergelyan in the 1950s, generalizing both Weierstrass' and Runge's Theorems:

**Theorem 1.2.3** (Mergelyan (1951), [39], [40], [41]). Let  $K \subset \mathbb{C}$  be compact polynomially convex. Then any  $f \in \mathcal{A}(K)$  can be uniformly approximated on  $K$  by holomorphic polynomials.

Denoting by  $\overline{\mathcal{O}(K)}$  the uniform closure of  $\{f|_K : f \in \mathcal{O}(K)\}$  in  $\mathcal{C}(K)$ , Mergelyan's Theorem implies the so-called *Mergelyan property*:

$$\mathcal{A}(K) = \overline{\mathcal{O}(K)}. \quad (1.2.2)$$

Both Runge's and Mergelyan's Theorem has been generalized a few years later, achieving the respective results on open Riemann surfaces:

**Theorem 1.2.4** (Runge's Theorem on Riemann surfaces, [4], [35], [52]). If  $K$  is a compact set in a Riemann Surface  $X$ , then any holomorphic function on a neighborhood of  $K$  can be uniformly approximated on  $K$  by meromorphic functions on  $X$  without poles in  $K$  and by holomorphic functions on  $X$  if  $X \setminus K$  has no relatively compact connected components.



**Theorem 1.2.5** (Bishop (1958), [6]). If  $K$  is a compact set without holes in an open Riemann Surface  $X$ , any  $f \in \mathcal{A}(K)$  can be uniformly approximated on  $K$  by functions in  $\mathcal{O}(X)$ .

Theorem 1.2.5 is actually a generalization of Theorem 1.2.3, as not having holes is equivalent to polynomial convexity for a compact  $K \subset \mathbb{C}$ . So far we have only dealt with approximation on compact sets; it is quite natural to investigate the approximation on unbounded sets as well. The first known result in this direction is a generalization of Weierstrass' Theorem 1.2.1:

**Theorem 1.2.6** (Carleman (1927), [10]). Given  $f: \mathbb{R} \rightarrow \mathbb{C}$  and  $\epsilon: \mathbb{R} \rightarrow (0, +\infty)$  continuous functions, there exists an entire function  $g \in \mathcal{O}(\mathbb{C})$  such that

$$|f(x) - g(x)| < \epsilon(x) \quad \forall x \in \mathbb{R}.$$

The proof is obtained inductively applying Mergelyan's Theorem 1.2.3. Many other results have been proved in the field, for which we suggest to consult the surveys [18], [8], [25], [26], [27], [28], [30], [29], [31], [32], [37], [61].

The mentioned approximation theorem in [15] is a Carleman-type Theorem. Given a continuous and positive function  $\epsilon: \mathbb{C} \rightarrow (0, +\infty)$ , a family of continuous functions  $f_r: \mathbb{C} \rightarrow \mathbb{C}$ , holomorphic on some ball  $\Delta_{\rho+a}$ , along with  $n$  families of unbounded Lipschitz curves  $\Gamma_{k,r}$ , we inductively construct a family of entire functions  $g_r: \mathbb{C} \rightarrow \mathbb{C}$  approximating  $f_r$  with a pointwise  $\epsilon$ -precision on a smaller ball  $\Delta_\rho$  and all along the  $n$  families of Lipschitz curves  $\Gamma_r = \cup_k \Gamma_{k,r}$ . We start approximating a family of continuous functions  $\alpha_r$  compactly supported on a single family of curves; the approximating family of functions  $H_r$  is the convolution with the gaussian kernel (1.2.1) defined on the curve  $\Gamma_r$  with  $n = 1$ ; being an integral representation, the approximation is preserved plugging in a whole family of functions, each one moving on a single Lipschitz curve; this generalizes a result in [38], holding for a single Lipschitz curve. Then with a Cousin II argument, we modify  $H_r$  into a function  $\xi_r$  which preserves the approximation already achieved and is arbitrarily small on a Pac-Man bounded region disjoint from  $\text{supp } \alpha_r$ : this is necessary to simultaneously approximate on several families of curves. With  $\xi_r$  we gain control on the behavior of the approximation losing "entireness", which is afterwards recovered exploiting a parametric version of Runge Theorem 1.2.2 we purposely proved as a lemma, getting  $Q_r$ . The simultaneous approximation on  $n$  families of curves is obtained by constructing a  $Q_{k,r}$  for each of them, then setting  $Q_r := \sum_k Q_{k,r}$ , which is an entire approximation of  $\alpha_r$  on compact subsets of the curves and arbitrarily small on  $\Delta_\rho$ . Such a construction makes us in business to prove our Carleman-type theorem: we first consider  $f_{N,r}$  the truncated Taylor polynomial of  $f_r$  on  $\Delta_\rho$ ; then we glue  $f_r$  and  $f_{N,r}$  with suitable cut-off functions obtaining  $\alpha_r$ , which we approximate with  $Q_r$ . Then  $f_{N,r} + Q_r$  is an entire approximation of  $f_r$  on  $\Delta_\rho \cup (\Gamma_r \cap \overline{\Delta_U})$ , for some  $U > \rho + a$ . Reiterating the procedure, we obtain approximations  $Q_r^{(m)}$  on the curves further and further, arbitrarily small on the domain on which we have already approximated, allowing  $g_r := f_{N,r} + \sum_{m \geq 1} Q_r^{(m)}$  to converge to an entire approximation of  $f_r$  on  $\Delta_\rho \cup \Gamma_r$ .

Finally, the last paper of this thesis [16], presents four approximation theorems. Nevertheless, being the setting of these results more general than one complex variable, we will describe them further later.

### 1.3 Several Complex Variables

Switching from one to several complex variables, many substantial differences arise quite soon. While certain results, like Liouville Theorem or the Identity Principle, keep holding, other important results turn out to fail. For instance: Riemann Mapping Theorem does not hold anymore, as the following theorem shows:

**Theorem 1.3.1** (Poincaré (1907), [48]). There exists no biholomorphic map between the polydisc  $P^n(0, 1) = \mathbb{B} \times \cdots \times \mathbb{B}$  and the ball  $\mathbb{B}^n$  if  $n > 1$ .

Holomorphic functions in more than one variable have no isolated zeros (see [49], corollary 3.3), or –really striking– when defined on an open neighborhood of  $\partial D$  where  $D$  is an arbitrary domain with connected boundary, they extend holomorphically to the whole  $D$ ; for example any function holomorphic on  $\{z \in \mathbb{C}^n : 1/2 < |z| < 1\}$ , or on a neighborhood of  $\{z \in \mathbb{C}^n : |z| = 1\}$  extends to the whole ball  $\mathbb{B}^n$ ; this is due to geometric properties of the domain of convergence of Laurent representation in several variables (see [49], Theorem 1.6) and leads to one of the most important phenomena in several variables: domains allowing simultaneous holomorphic extension across them. The first instance in this direction was provided already in 1897 by A. Hurwitz, who showed in his lecture at the first International Congress of Mathematicians that a holomorphic function of more than one variable, has no isolated singularity; anyhow, the most famous (and historically relevant) example of such a domain was found by F. Hartogs in 1906 (see [34]):

$$H = \{(z, w) \in \mathbb{C}^2 : |z| < 1, \ 1/2 < |w| < 1\} \cup \{|z| < 1/2, \ |w| < 1\}.$$

In fact any  $f \in \mathcal{O}(H)$  is holomorphically extended on the whole polydisc  $P^2(0, 1) = \mathbb{B} \times \mathbb{B}$  by

$$F(z, w) = \frac{1}{2\pi i} \int_{|\zeta|=3/4} \frac{f(z, \zeta)}{\zeta - w} d\zeta,$$

as  $F$  matches  $f$  on  $\{|z| < 1/2, \ |w| < 3/4\}$  and it is holomorphic on  $\{|z| < 1, \ |w| < 3/4\}$ . This phenomena does not occur in  $\mathbb{C}$ , as, given any domain  $D \subset \mathbb{C}$ , it is enough to consider  $f(z) = (z - p)^{-1}$  for some  $p \in \partial D$  to obtain a function holomorphic on  $D$  which does not extend across it. To talk precisely about this occurrence, we state the following definition.

**Definition 1.3.1.** A function  $f \in \mathcal{O}(D)$  is called *completely singular* at  $p \in \partial D$  if for every  $U$  connected neighborhood of  $p$ , there exists no  $h \in \mathcal{O}(U)$  such that  $f = h$  on some connected component of  $U \cap D$ .  $D$  is a *domain of holomorphy* when there exists  $f \in \mathcal{O}(D)$  completely singular at each boundary point.

The condition to find for each  $p \in \partial D$  a  $f_p \in \mathcal{O}(D)$  completely singular at  $p$ , seems weaker but it is indeed an equivalent condition of being a domain of holomorphy. This is always the case in  $\mathbb{C}$ , but not always in  $\mathbb{C}^n$  with  $n > 1$  as the above examples show. To characterize these domains has been a central point in complex analysis research. Few years after Hartogs' discovery, E. E. Levi had a first major insight about domains with *differentiable boundaries*, that is domains  $D$  such that for each  $p \in \partial D$  there exists  $U$  neighborhood of  $p$  and a real function  $r \in \mathcal{C}^2(U)$  with  $dr \neq 0$ , such that  $D \cap U = \{z \in U : r(z) < 0\}$ . In this setting, he gave a simple differential condition, remarkably similar to the differential characterization of euclidean convex domains:

**Theorem 1.3.2** (Levi (1910/1911)).

- (i)<sub>L</sub> If there exists a holomorphic function on  $U \cap D$  which does not extend across  $p$  (in particular  $D$  is a domain of holomorphy) then

$$L_p(r; t) := \sum_{j,k=1}^n \frac{\partial^2 r}{\partial z_j \partial \bar{z}_k}(p) t_j \bar{t}_k \geq 0$$

for all  $t \in \mathbb{C}^n$  such that  $\sum_{j=1}^n \frac{\partial r}{\partial z_j}(p) t_j = 0$ .

- (ii)<sub>L</sub> If  $L_p(r; t) > 0$  for all  $t \neq 0$  such that  $\sum_{j=1}^n \frac{\partial r}{\partial z_j}(p) t_j = 0$ , then  $U$  can be chosen so that  $U \cap D$  is a domain of holomorphy.

The quadratic form  $t \mapsto L_p(r; t)$  is the complex hessian of  $r$  at  $p$ , commonly called *Levi form*. When the differential condition in (i)<sub>L</sub> or (ii)<sub>L</sub> in the above theorem holds, we say that  $D$  is *Levi pseudoconvex* and *strongly pseudoconvex* respectively. In particular, it follows from (ii)<sub>L</sub>, that a domain that is strongly pseudoconvex near each boundary point, is locally a domain of holomorphy. Proving that a strongly pseudoconvex domain is actually a domain of holomorphy, has been a major problem for 30 years, known as *Levi problem*. It was solved in the early '50 by K. Oka, H. Bremermann, and F. Norguet. So we have a good description of domains of holomorphy when we restrict to domains with differentiable boundaries. To deal with the general case, we need the following definition.

**Definition 1.3.2.** An arbitrary domain  $D \subset \mathbb{C}^n$  is said to be *pseudoconvex* if the mapping  $z \mapsto -\log \text{dist}(z, \partial D)$  is plurisubharmonic on  $D$ , where a real valued  $r \in \mathcal{C}^2(D)$  is *plurisubharmonic* if  $L_p(r; t) \geq 0$  for all  $t \in \mathbb{C}^n$  and all  $p \in D$ .

The general version of the solution of the Levi Problem reads as follows:

**Theorem 1.3.3** ([43], [44], [45] [9], [46]). A domain  $D$  in  $\mathbb{C}^n$  is a domain of holomorphy if and only if it is pseudoconvex.

We conclude this section by spending a couple of words on  $\text{Aut } \mathbb{C}^n$ , the group of holomorphic automorphisms of  $\mathbb{C}^n$ . When  $n = 1$  it is well known that  $\text{Aut } \mathbb{C}$  is given by the affine functions  $z \mapsto az + b$ , where  $a, b \in \mathbb{C}$  and  $a \neq 0$ . When  $n > 1$

## 1. Introduction

---

the situation is much more complicated. To address the problem of describing  $\text{Aut } \mathbb{C}^n$  we give the following

**Definition 1.3.3.** Let  $n > 1$  and write  $z = (z', z_n) \in \mathbb{C}^n$ . The elements of  $\text{Aut } \mathbb{C}^n$  of the form

$$\Psi(z) = (z', e^{f(z')}z_n + g(z')), \quad z \in \mathbb{C}^n$$

and their  $\text{GL}_n(\mathbb{C}^n)$ -conjugates are called *shears*, where  $f, g \in \mathcal{O}(\mathbb{C}^{n-1})$ . Denote by  $\mathcal{S}(n)$  the group generated by the shears. If  $f = 0$ ,  $\Psi$  and their  $\text{SL}_n(\mathbb{C}^n)$ -conjugates are called *additive shears*, while if  $g = 0$ ,  $\Psi$  and their  $\text{GL}_n(\mathbb{C}^n)$ -conjugates are called *over-shears*.

Even if an explicit description of  $\text{Aut } \mathbb{C}^n$  is not known, in 1992 E. Andersén and L. Lempert made a fundamental discovery that started a new theory.

**Theorem 1.3.4** (Andersén and Lempert (1992), [3]).  $\mathcal{S}(n)$  is dense in  $\text{Aut } \mathbb{C}^n$  and not equal to it.

## 1.4 Stein Manifolds

An analytic theory developed on euclidean spaces is naturally generalized to the setting of manifolds. Complex manifolds, in this case. As the concept of domain of holomorphy turned out to be central in the “standard theory”, we wish to carry this notion to the manifold setting, together with other reasonable requests, which are embodied in the following definition provided by K. Stein:

**Definition 1.4.1** (K. Stein (1951), [56]). A complex manifold  $X$  is said to be a *Stein Manifold* (or a *holomorphically complete manifold*) if the following hold:

- (i)<sub>SM</sub> For every pair of distinct points  $x, y \in X$ , there exists a holomorphic function  $f \in \mathcal{O}(X)$  such that  $f(x) \neq f(y)$ .
- (ii)<sub>SM</sub> For every  $p \in X$  there are  $n$  holomorphic functions (where  $n = \dim X$ )  $f_1, \dots, f_n \in \mathcal{O}(X)$  whose differentials  $df_j$  are  $\mathbb{C}$ -linearly independent at  $p$  (so global functions provide local charts).
- (iii)<sub>SM</sub>  $X$  is holomorphically convex.

If  $X = \mathbb{C}^n$ , properties (i)<sub>SM</sub> and (ii)<sub>SM</sub> are automatically fulfilled, while (iii)<sub>SM</sub> is actually equivalent to being a domain of holomorphy:

**Theorem 1.4.1** (Cartan and Thullen (1932), [11]). A Riemann domain  $X$  over  $\mathbb{C}^n$  is a domain of holomorphy if and only if it is holomorphically convex.

A Riemann domain  $X$  is just a complex manifold endowed with a locally biholomorphic mapping  $\pi: X \rightarrow \mathbb{C}^n$ : as analytic continuation leads to multivalued functions, the choice of such a  $\pi$  avoids any kind of ambiguity. Let us highlight that this theorem generalizes the notion of domain of holomorphy

to an abstract setting in which we cannot refer (a priori) to a surrounding space, so it identifies the property of being a domain of holomorphy as an intrinsic characteristic of a space.

As mentioned in Section 1.2, the last paper of this thesis [16] is about approximation theory dealing with holomorphic mappings with dense images; some results covering this topic can be found in [1] and [19]. In particular, the motivation for [16] was to generalize the main result of [22], in which F. Forstnerič and J. Winkelmann proved that the set of all holomorphic mappings  $f: \Delta \rightarrow Y$  with dense image, is dense in  $\mathcal{O}(\Delta, Y)$ , where  $Y$  is a connected complex manifold. In fact, the first result collected in [16] is about approximating holomorphic embeddings  $g: \Omega \rightarrow Y$  with the same kind of mappings with dense images, where  $Y$  is a connected complex manifold and  $\Omega \subset \mathbb{C}^n$  is open, bounded star-shaped and pseudoconvex. The method exploited to prove this result relies on the technique of exposing boundary points introduced in [12] and it turned out to be valid to prove three furthermore general approximation theorems, likewise presented in [16]. The second theorem, like the first one, claims that the set of all non-constant holomorphic mappings  $g: X \rightarrow Y$  with bounded image can be approximated by holomorphic mappings with dense image, where  $X$  is a complex manifold and  $Y$  is a Stein manifold. The remaining two theorems work the other way around, proving that mappings defined on a given compact  $K \subset Y$  minus certain points (hence not a priori holomorphic) with dense image in  $Y$ , are approximated by holomorphic mappings defined on neighborhoods of  $K$ , which is a Stein compact inside a complex manifold  $Y$  in the third theorem and it is a compact inside a Stein manifold  $Y$  in the fourth one. The methodology exploited to prove these results is as follows: we start from a compact  $K$ , which in the last two theorems is given, whether it is obtained as the closure of the image  $\overline{g(X)}$  of the mapping  $g: X \rightarrow Y$  we wish to approximate in the first two; consider moreover a countable dense set  $Q \subset Y$ .  $K$  contains *locally exposable points* (a generalization of the notion of *peak point*), which actually belong to the boundary; pick one of them  $\zeta$  and connect it to a suitable  $q \in Q \setminus K$  with a smoothly embedded path  $\gamma \subset Y \setminus K \cup \{q\}$ . Then Theorem 1.1 in [12] (recalled in the paper in a convenient version) guarantees the existence of a holomorphic mapping  $f$  on a neighborhood of  $K$ , arbitrarily close to the identity on  $K$  minus a small ball  $B$  around  $\zeta$ , and deforming such a ball  $B$  into a tube arbitrarily close to  $\gamma$  and touching  $q$  (namely  $f(\zeta) = q$ ). We repeat the procedure on the compact  $f(K)$  and iterate it; in the limit, we get a mapping  $H$  with dense image, as we constructed it touching new points of  $Q$  at each step and moving arbitrarily little what we have already hit, that guarantees both convergence and approximation. In the first two theorems,  $H \circ g: X \rightarrow Y$  is the approximating holomorphic mapping with dense image. In the remaining two,  $H$  is a mapping with dense image which is approximated by partial compositions of the holomorphic mappings obtained at each step.

## 1.5 Proper Holomorphic Embeddings

From what we have seen so far, Stein manifolds are the natural source of holomorphic functions in a reasonably general setting. We are therefore interested in studying these objects. To represent them more concretely, we would like to identify them as subsets of some Euclidean space, in a faithful way (one-to-one and avoiding creating singularities), respecting the boundaries (which is crucial as the definition of pseudoconvexity actually relies on the boundaries) and the complex structure of a manifold. Hence, given a Stein manifold  $X$  we are looking for *proper holomorphic embeddings*

$$f: X \hookrightarrow \mathbb{C}^N$$

for some  $N$ . Proper mappings between topological spaces are the ones preserving compactness under inverse image. An equivalent characterization (that takes into account the boundary if defined) is that such mappings, map sequences leaving every compact into sequences leaving every compact, thus if a Stein manifold is a bounded domain inside some bigger space (e.g., any domain of holomorphy  $X \subset \mathbb{C}^n$  relatively compact), every sequence approaching the boundary of  $X$  is sent to an unbounded sequence. The first question is then whether such a mapping always exists. An affirmative answer has been provided by R. Remmert in 1956:

**Theorem 1.5.1** (Remmert (1956), [51]). Given any Stein manifold  $X$ , there exists a proper holomorphic embedding  $f: X \hookrightarrow \mathbb{C}^N$  for  $N$  sufficiently large.

The attention is thus towards  $N$  in function of  $n = \dim X$ . In particular, interest is for spotting a minimal  $N$ : the smaller  $N$ , the tighter the euclidian space, the more faithful the representation. A first result in this direction has been given in 1960/1961 independently by E. Bishop and R. Narasimhan, as they proved that  $N$  can be taken to be  $2n + 1$  (see [7], [42]). In 1970, O. Forster conjectured that, if  $n > 1$ ,  $N$  could be taken to be  $\lfloor \frac{3n}{2} \rfloor + 1$ , proving moreover that this is minimal (see [20]): given an integer  $n \geq 2$ , he defined the  $n$ -dimensional Stein manifold

$$X := \begin{cases} Y^m & n = 2m \\ Y^m \times \mathbb{C} & n = 2m + 1, \end{cases}$$

where

$$Y := \{[x : y : z] \in \mathbb{P}^2 : x^2 + y^2 + z^2 \neq 0\}$$

is a Stein surface. Eliashberg and Gromov first, Schürmann later, turned this conjecture into the following

**Theorem 1.5.2** (Eliashberg, Gromov (1992), [17] and Schürmann (1997), [54]). Every Stein manifold with  $n = \dim X > 1$  admits a proper holomorphic embedding into  $\mathbb{C}^N$ , for  $N = \lfloor \frac{3n}{2} \rfloor + 1$ .

The proof breaks down in the case  $n = 1$ . Since Behnke and Stein proved that one-dimensional Stein manifolds are precisely open connected Riemann

Surfaces (see [4] and [5]), we are led to the following long-standing open problem:

**Conjecture 1.5.1.** Every open connected Riemann Surface admits a proper holomorphic embedding into  $\mathbb{C}^2$ .

The first example validating this conjecture dates back to 1970 when Kasahara and Nishino proved it in the case of the unit disk  $\mathbb{B}$  ([55]). The annulus  $\{1 < |z| < r\}$  and the punctured disk  $\mathbb{B} \setminus \{0\}$  have been proved to verify the conjecture by Laufer (1973, see [36]) and Alexander (1977, [2]) respectively. The first quite general result has been given by J. Globevnik and B. Stensønes in 1995: they proved in [33] that any finitely connected domain in  $\mathbb{C}$  without isolated boundary points verifies the conjecture. A more complete account concerning the topic of proper holomorphic embeddings can be found in [13]. To present more general results, we need to gather some definitions.

**Definition 1.5.1.**

- A *bordered Riemann Surface* is the interior  $D$  of a 1-dimensional compact complex manifold  $\overline{D}$ , not necessarily connected, with smooth boundary  $\partial D$  consisting of finitely many closed Jordan curves.
- Given a closed complex curve  $\Sigma \subset \mathbb{C}^2$  (that is the closure of the holomorphic image of some open subset of  $\mathbb{C}$ , namely  $\Sigma = \overline{F(D)}$ , where  $F: D \rightarrow \mathbb{C}^2$  is holomorphic), possibly with boundary, and a linear projection  $\pi: \mathbb{C}^2 \rightarrow \mathbb{C}$ , a point  $p \in \Sigma$  is an *exposed point* (with respect to  $\pi$ ) if the affine complex plane thru  $p$  defined by  $\pi$ , that is  $\Lambda_p := \pi^{-1}\pi(p)$ , meets  $\Sigma$  exactly in  $p$  ( $\Lambda_p \cap \Sigma = \{p\}$ ) and the intersection is transverse:  $T_p\Lambda_p \cap T_p\Sigma = \{0\}$ .
- A family  $\Gamma = \{\gamma_j(t) : j = 1, \dots, m, t \in (-\infty, +\infty)\}$  of pairwise disjoint, smooth curves in  $\mathbb{C}^2$  without self intersections has the *immediate projection property* with respect to some linear projection  $\pi: \mathbb{C}^2 \rightarrow \mathbb{C}$  if:
  - \*  $|\pi(\gamma_j(t))| \rightarrow +\infty$  as  $|t| \rightarrow +\infty$  for every  $j$ ,
  - \* there exists  $M > 0$  such that  $\mathbb{C} \setminus (\pi(\Gamma) \cup R\overline{\mathbb{B}})$  has no relatively components for every  $R \geq M$ .
- A domain  $\Omega \subset \mathbb{C}^n$ ,  $n > 1$ , is a *Fatou-Bieberbach domain* if it is biholomorphic to  $\mathbb{C}^n$ . A biholomorphism  $\Phi: \Omega \rightarrow \mathbb{C}^n$  is called a *Fatou-Bieberbach map*.

The problem of finding proper holomorphic embeddings of such  $D$  into  $\mathbb{C}^2$  can be split into the following two problems:

- A) find an embedding  $f: \overline{D} \hookrightarrow \mathbb{C}^2$  onto a smooth compact complex curve with boundary  $\overline{\Sigma} := f(\overline{D}) \subset \mathbb{C}^2$ , holomorphic on  $D$ .
- B) obtaining properness by pushing the boundary  $\partial\Sigma = f(\partial D)$  to infinity without introducing any interior double points.

E. F. Wold developed a remarkably general method in [58], [59], that allowed him to generalize Globevnik and Stensønes' result to all finitely connected and some infinitely connected planar domains [58], to every domain in the Riemann sphere  $\mathbb{P}^1$  with at most countably many boundary components, none of which are points [24] (with Forstnerič), and to every subset of the torus  $\mathbb{T}$  with finitely many boundary components, none of which being a point [60]. But most importantly, the works [58] and [59] have been the starting point leading Wold and Forstnerič to formulate and prove the following:

**Theorem 1.5.3** (Theorem 4.2, [23]). A  $\mathcal{C}^1$ -smooth embedding  $f: \overline{D} \hookrightarrow \mathbb{C}^2$ , holomorphic on  $D$ , can be approximated in the  $\mathcal{C}^1$ -topology on  $\overline{D} \setminus \cup_j U_j$ , where  $U_j$  is a small neighborhood of a point  $a_j$  in each connected component of the boundary  $\partial D$ , by embeddings  $F: \overline{D} \hookrightarrow \mathbb{C}^2$ , holomorphic on  $D$ , such that every boundary component of  $F(\overline{D})$  has an exposed point.

This result is needed to give a complete solution to problem B) (Corollary 1.2 in [23]). Another fundamental tool is the following

**Lemma 1.5.1** (Lemma 1, [58]). Given  $K \subset \mathbb{C}^2$  compact polynomially convex and  $\Gamma \subset \mathbb{C}^2 \setminus K$  a set of  $m$  real pairwise disjoint smooth curves without self-intersection enjoying the immediate projection property, we can build automorphisms  $\phi \in \text{Aut } \mathbb{C}^2$  arbitrarily close to the identity on  $K$  and pushing the curves  $\Gamma$  outside a ball of arbitrarily big radius.

This Lemma turns out to be a key step of the main result in [59], in which a description of the construction of proper holomorphic embeddings into  $\mathbb{C}^2$  of bordered Riemann Surfaces  $D$  whose boundary  $\partial D$  enjoys the nice projection property is provided. We report it here as resumed in [21]:

**Theorem 1.5.4** (Theorem 4.14.6, [21]). Let  $X \subset \mathbb{C}^n$  be a complex curve ( $n > 1$ ), possibly with singularities, whose boundary  $\Gamma = \partial X$  is as in definition 1.5.1. Then there exists a Fatou–Bieberbach domain  $\Omega \subset \mathbb{C}^n$  such that  $X \subset \Omega$  and  $\partial X \subset \partial \Omega$ . Moreover, a Fatou–Bieberbach map  $\Phi: \Omega \rightarrow \mathbb{C}^n$  can be chosen to be arbitrarily close to the identity on a fixed compact inside  $X$ . In particular  $\Phi|_X: X \hookrightarrow \mathbb{C}^n$  is a proper holomorphic embedding.

The last-mentioned results are the state of art regarding Conjecture 1.5.1, which remains unsolved as not much is known about A). It, therefore, makes sense to search for a counterexample: there have been discussions suggesting that 1.5.1 could have failed for complements of large Cantor sets inside the Riemann sphere. S. Y. Orevkov, wrote the paper [47] to prove the existence of a proper holomorphic embedding  $\mathbb{P}^1 \setminus C \hookrightarrow \mathbb{C}^2$ , without any discussion on the size of the Cantor set  $C$ . Therefore the second paper included in this thesis [14] is focused on the study of the measure of  $C$ . It turns out natural to take it very small: it is there proved that the construction of  $C$  can be performed so that  $C$  has zero Hausdorff dimension. Such a paper is also justified by the extreme synthesis and lack of details in [47], which I have provided. The construction is as follows: a sequence of continuous mappings  $\gamma_n: \overline{\mathbb{C}} \rightarrow \overline{\mathbb{C}^2}$  is built, such that  $\gamma_n: \overline{\mathbb{C}} \setminus \gamma_n^{-1}(\infty_2) \rightarrow \mathbb{C}^2$  is a holomorphic embedding and  $A_n = \gamma_n(\overline{\mathbb{C}})$  is compact



complex curve with an increasing (with  $n$ ) number of unbounded components in both coordinate directions. Call  $\Delta_n$  the preimage through  $\gamma_n$  of the curves  $A_n$  in smaller and smaller (as  $n$  increases) neighborhoods of  $\infty_2$ , whose size depends on a divergent sequence of parameters  $\{R_n\}_n$  on which the whole construction relies on;  $\Delta_n$  is the disjoint union of open discs in  $\mathbb{C}$ , which are as many as the branches of  $A_n$  touching  $\infty_2$ . It turns out that  $\Delta_n$  is a decreasing sequence and every disc tends to shrink to a point, none of which remains isolated. Therefore  $C = \bigcap_n \Delta_n$  is a Cantor set and the limit mapping  $\gamma = \lim_n \gamma_n: \overline{\mathbb{C}} \setminus C \rightarrow \mathbb{C}^2$  is a holomorphic embedding which turns to be proper. The diameter of the discs can be taken arbitrarily little provided  $R_n \rightarrow +\infty$  fast enough, thus  $C$  can be realized to have zero Hausdorff dimension.

Since this last construction provides a proper holomorphic embedding of the complement of a Cantor set which is actually thin, here is where the first paper of this thesis [13] comes into play. We presented in it a construction of the same topological object using the techniques developed by Wold, which allow a construction leading to properly holomorphically embed a considerably large Cantor set, as follows: let  $Q_1$  be a square in the Riemann Sphere  $\mathbb{P}^1$  of side 2. At each step we remove smaller and smaller strips, alternatively vertical and horizontal, getting  $Q_n$ ; set then  $C_n := \mathbb{P}^1 \setminus \overline{Q_n}$ . We inductively construct the holomorphic embeddings  $f_n: C_n \hookrightarrow \mathbb{C}^2$  as a composition  $\phi_n \circ g_n \circ F_n$ , where  $F_n: C_n \hookrightarrow \mathbb{C}^2$  is a holomorphic embedding having exposed points on each of the  $2^{n-1}$  boundary components of the image,  $g_n$  is a rational shear sending these points to infinity in  $\mathbb{C}^2$ , so that  $X_n := g_n \circ F_n(C_n) \subset \mathbb{C}^2$  is an unbounded complex curve enjoying the immediate projection property 1.5.1. Then setting  $r_n := n-2$  we have that  $L_n := r_n \overline{\mathbb{B}^2} \cup K_n$  is polynomially convex for a sufficiently big polynomially convex  $K_n \subset X_n$  (see the proof of Theorem 4.14.6 in [21]), so we can apply Lemma 1.5.1 and get  $\phi_n \in \text{Aut } \mathbb{C}^2$

1. suitably close to the identity on  $L_n$ , and
2. pushing the boundary away from a bigger ball, say  $\phi_n(\partial X_n) \subset \mathbb{C}^2 \setminus r_{n+1} \overline{\mathbb{B}^2}$ .

Extend  $f_n$  to a smooth embedding  $\tilde{f}_n: C_n \cup l_n \hookrightarrow \mathbb{C}^2$ , where  $l_n$  is the set of vertical or horizontal lines dividing the rectangular components of  $Q_n$  into two equal pieces and use Mergelyan Theorem 1.2.3 to approximate  $\tilde{f}_n$  on  $C_n \cup l_n$  with a holomorphic embedding  $\hat{f}_{n+1}: C_{n+1} \hookrightarrow \mathbb{C}^2$ . Theorem 1.5.3 allows now to approximate  $\hat{f}_{n+1}$  with a holomorphic embedding  $F_{n+1}: C_{n+1} \hookrightarrow \mathbb{C}^2$  with an exposed point on every boundary component of the image and the induction may proceed, creating a sequence of holomorphic embeddings  $\{f_n\}_n$  convergent by 1 to a holomorphic embedding  $f$  on  $\left(\bigcup_{n \geq 1} C_n\right)^\circ$ , which equals  $\mathbb{P}^1 \setminus C$ , where  $C$  is a Cantor set of Lebesgue measure arbitrarily close to 4 (by choosing the strips to remove to be suitably small at each step). Properness of the limit function  $f: \mathbb{P}^1 \setminus C \hookrightarrow \mathbb{C}^2$  follows from 2.

The last result to mention in this thesis, is the main Theorem of the third paper [15]. The concern deals with simultaneously properly holomorphically embedding a whole family of complex curves into  $\mathbb{C}^2$ . Namely: given a

## 1. Introduction

---

continuous family of  $n$ -connected domains  $\Omega_r \subset \mathbb{P}^1$  such that none of the components of  $\mathbb{P}^1 \setminus \Omega_r$  reduces to a point, the goal is to find a continuous mapping  $\Xi: \bigcup_r \{r\} \times \Omega_r \rightarrow \mathbb{C}^2$  which properly holomorphically embeds fiberwise, that is  $\Xi(r, \cdot): \Omega_r \hookrightarrow \mathbb{C}^2$  is a proper holomorphic embedding for every  $r$ . The proof structure is similar to the one of Theorem 1.5.4; the main difference lies in the use of Lemma 1.5.1, in fact, we now need a parametric version of it. This issue is solved with the Andersén–Lempert procedure, for which the parametric version of the Carleman Theorem previously presented is fundamental.

# Bibliography

- [1] Alarcón, A. and Forstnerič, F.; Complete densely embedded complex lines in  $\mathbb{C}^2$ . *Proc. Amer. Math. Soc.*, 146(3):1059–1067, 2018.
- [2] Alexander, H.; *Explicit imbedding of the (punctured) disc into  $\mathbb{C}^2$* . *Comment. Math. Helv.*, **52** (1977), 539–544.
- [3] Andersén, E. and Lempert, L.; *On the group of holomorphic automorphisms of  $\mathbb{C}^n$* . *Invent. Math.*, **110**(2) (1992), 371–388.
- [4] Behnke, H. and Stein, K.; *Entwicklung analytischer Funktionen auf Riemannschen Flächen*. *Math. Ann.*, **120** (1948), 430–461.
- [5] Behnke, H. and Stein, K.; *Elementarfunktionen auf Riemannschen Flächen als Hilfsmittel für die Funktionentheorie mehrer Veränderlichen*. *Can. J. Math.*, **2** (1950), 152–165.
- [6] Bishop, E.; *Subalgebras of functions on a Riemann surface*. *Pacific J. Math.*, **8** (1958), 29–50.
- [7] Bishop, E.; *Mappings of partially analytic spaces*. *Am. J. Math.*, **83** (1961), 209–242.
- [8] Boivin, A. and Gauthier P. M.; *Holomorphic and harmonic approximation on Riemann surfaces*. In *Approximation, complex analysis, and potential theory* (Montreal, QC, 2000), volume 37 of NATO Sci. Ser. II Math. Phys. Chem., pages 107–128. Kluwer Acad. Publ., Dordrecht, (2001).
- [9] Bremermann, H. J.; *Über die Äquivalenz der pseudokonvexen Gebiete und der Holomorphiegebiete in Raum von  $n$  komplexen Veränderlichen*. *Math. Ann.*, **128** (1954), 63–91.
- [10] Carleman, T.; *Sur un théorème de Weierstraß*. *Ark. Mat. Astron. Fys.*, **20** (1927), no. 4, 1–5.
- [11] Cartan, H. and Thullen, P.; *Zur Theorie der Singularitäten der Funktionen mehrer komplexen Veränderlichen*. *Math. Ann.*, **106**, (1932), 617–647.
- [12] Deng, F., Fornæss J. E. and Wold E. F.; *Exposing Boundary Points of Strongly Pseudoconvex Subvarieties in Complex Spaces*. *Proceedings Of The AMS*, Volume **146**, no. 6, June 2018, 2473–2487, <http://dx.doi.org/10.1090/proc/13693>, Article electronically published on March 9, 2018.

- [13] Di Salvo, G. D. and Wold, E. F.; *Proper Holomorphic Embeddings of complements of Large Cantor Sets in  $\mathbb{C}^2$* . First paper in this thesis.
- [14] Di Salvo, G. D.; *Extended explanation of Orevkov's paper on the Proper Holomorphic Embeddings of complements of Cantor Spaces in  $\mathbb{C}^2$  and discussion of their measure*. Second paper in this thesis.
- [15] Di Salvo, G. D., Ritter, T. and Wold, E. F.; *Families of Proper Holomorphic Embeddings and Carleman-type Theorems with parameters*. Third paper in this thesis.
- [16] Di Salvo G. D.; *Approximation and accumulation results of holomorphic mappings with dense image*. Fourth paper in this thesis.
- [17] Eliashberg, Y. and Gromov, M.; *Embeddings of Stein manifolds*. Ann. Math., (2), **136** (1992), 123–135.
- [18] Fornæss, J. E., Forstnerič F. and Wold, E. F.; *Holomorphic approximation: the legacy of Weierstrass, Runge, Oka–Weil, and Mergelyan. Advancements in Complex Analysis – From Theory to Practice*. Breaz, Daniel & Rassias, Michael Th. (Ed.), Springer Nature. ISSN 978–3–030–40119–1. p. 133–192.
- [19] Fornæss, J. E. and Stout, E. L.; *Polydiscs in complex manifolds.. Math. Ann.* 227 (1977), no. 2, 145–153.
- [20] Forster, O.; *Plongements des variétés de Stein*. Comment. Math. Helv, **45** (1970), 170–184.
- [21] Forstnerič, F.; *Stein manifolds and holomorphic mappings. The homotopy principle in complex analysis*. Second edition. Ergebnisse der Mathematik und ihrer Grenzgebiete. **3**. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics], **56**. Springer, Cham, (2017).
- [22] Forstnerič, F. and J. Winkelmann, J.; *Holomorphic Discs with Dense Images*. Mathematical Research Letters 12, 265–268 (2005)
- [23] Forstnerič, F. and Wold, E. F.; *Bordered Riemann Surfaces in  $\mathbb{C}^2$* . J. Math. Pures Appl. **91** (2009), 100–114.
- [24] Forstnerič, F. and Wold, E. F.; *Embeddings of infinitely connected planar domains into  $\mathbb{C}^2$* . Anal. PDE, **6** (2013), no. 2, 499–514.
- [25] Fuchs, W. H. J.; *Théorie de l'approximation des fonctions d'une variable complexe*. Séminaire de Mathématiques Supérieures, no. 26 (Été, 1967). Les Presses de l'Université de Montréal, Montreal, Que., (1968).
- [26] Gaier, D.; *Lectures on complex approximation*. Birkhäuser Boston, Inc., Boston, MA, (1987). Translated from the German by Renate McLaughlin.

- [27] Gamelin, T. W.; *Polynomial approximation on thin sets*. Symposium on several complex variables, Lecture Notes in Math. (Park City, Utah, 1970), Springer, Berlin, **184** (1971), 50–78.
- [28] Gamelin, T. W.; *Uniform algebras*, 2nd ed, Chelsea Press, New York, (1984).
- [29] Gauthier, P. and Paramonov, P. V.; *Approximation by solutions of elliptic equations and extension of subharmonic functions*. Mashregghi J., Manolaki M., Gauthier P. (eds), New Trends in Approximation Theory, Volume 81 of Fields Institute Communications. Springer, New York, NY, (2018).
- [30] Gauthier, P.; *Uniform approximation*. Complex potential theory (Montreal, PQ, 1993), Volume **439** of NATO Adv. Sci. Inst. Ser. C Math. Phys. Sci., pages 235–271. Kluwer Acad. Publ., Dordrecht, (1994).
- [31] Gauthier, P. and Hengartner, W.; *Approximation uniforme qualitative sur des ensembles non borné*, Volume **82** of Séminaire de Mathématiques Supérieures [Seminar on Higher Mathematics], Presses de l'Université de Montréal, Montreal, Que., (1982).
- [32] Gauthier, P. and Sharifi, F.; *Uniform approximation in the spherical distance by functions meromorphic on Riemann surfaces*. J. Math. Anal. Appl., **445** no. 2 (2017), 1328–1353.
- [33] Globevnik, J. and Stensønes, B.; *Holomorphic embeddings of planar domains into  $\mathbb{C}^2$* . Math. Ann., **303** (1995), 579–597.
- [34] Hartogs, F.; *Zur Theorie der analytischen Functionen mehrerer unabhängiger Veränderlichen, insbesondere über die Darstellung derselben durch Reihen, welche nach Potenzen einer Veränderlichen fortschreiten*. Math. Anal., **62** no. 2 (1906), 1–88.
- [35] Köditz, H. and Timmann, S.; *Randschlichte meromorphe Funktionen auf endlichen Riemannschen Flächen*. Math. Ann., **217** no. 2 (1975), 157–159.
- [36] Laufer, H. B.; *Imbedding annuli in  $\mathbb{C}^2$* . J. Anal. Math., **26** (1973), 187–215.
- [37] Levenberg, N.; *Approximation in  $\mathbb{C}^N$* . Surv. Approx. Theory, **2** (2006), 92–140.
- [38] Manne, P.; *Carleman approximation in several complex variables*, Ph.D. Thesis. University of Oslo, (1993).
- [39] Mergelyan, S. N.; *On the representation of functions by series of polynomials on closed sets*. Doklady Akad. Nauk SSSR (N.S.), **78** (1951), 405–408.
- [40] Mergelyan, S. N.; *Uniform approximations of functions of a complex variable*. Uspehi Matem. Nauk (N.S.), **7** no. 2(48) (1952), 31–122.
- [41] Mergelyan, S. N.; *Uniform approximations to functions of a complex variable*. Amer. Math. Soc. Translation 1954, no. 101 (1954), 1–99.

- [42] Narasimhan, R.; *Holomorphic mappings of complex spaces*. Proc. Am. Math. Soc., **11** (1960), 800–804.
- [43] Norguet, F.; *Sur le domaines d'holomorphie des fonctions uniformes de plusieurs variables complexes (Passage du local au global)*. Bull. Soc. Math. Fr., **82** (1954), 137–159.
- [44] Oka, K.; *Sur les fonctions uniformes de plusieurs variables. VI: Domaines pseudoconvexes*. Tôhoku Math. J., **49** (1942), 15–52.
- [45] Oka, K.; *Sur les fonctions uniformes de plusieurs variables. VI: Domaines finis sans point critique intérieur*. Tôhoku Math. J., **49** (1942), 15–52.
- [46] Oka, K.; *Collected Papers*. Translated from the French by R. Narasimhan. With commentaries by H. Cartan. Edited by R. Remmert. Springer–Verlag, Berlin (1984).
- [47] Orevkov, S. Y.; *Proper analytic embedding of  $\mathbb{CP}^1$  minus Cantor set into  $\mathbb{C}^2$* . Uspekhi Mat. Nauk. **63** (2008), no. 1(379), 155–156 (in Russian), English transl.: Russian Math. Surveys, **63** (2008), 168–169.
- [48] Poincaré, H.; *Les fonctions analytiques de deux variables et la représentation conforme*. Rend. Circ. Mat. Palermo, **23** (1907), 185–220.
- [49] Range, R. M.; *Holomorphic Functions and Integral Representations in Several Complex Variables*. Springer, Graduate Text in Mathematics **108** (1986).
- [50] Remmert, R.; *Theory of Complex Functions*. Second Edition. Graduate Texts in Mathematics. **122**. Readings in Mathematics. Springer–Verlag, (1989).
- [51] Remmert, R.; *Sur les espaces analytiques holomorphiquement séparables et holomorphiquement convexes*. C. R. Acad. Sci. Paris, **243** (1956), 211–221.
- [52] Royden, H. L.; *Function theory on compact Riemann surfaces*. J. Analyse Math., (1967) 295–327.
- [53] Runge, C.; *Zur Theorie der Eindeutigen Analytischen Functionen*. Acta Math., **6** no. 1 (1885), 229–244.
- [54] Schürmann, J.; *Embeddings of Stein spaces into affine spaces of minimal dimension*. Math. Ann., **307** (1997), 381–399.
- [55] Stehlé, J. –L.; *Plongements du disque dans  $\mathbb{C}^2$* . In: Séminaire P. Lelong (Analyse). Lect. Notes in Math., Springer–Verlag, Berlin, **275** (1970), 119–130.
- [56] Stein, K.; *Analytische Funktionen mehrerer komplexer Veränderlichen zu vorgegebenen Periodizitätsmoduln und das zweite Cousinsche Problem*. Math. Ann., **123** (1951), 201–222.

- [57] Weierstrass, K.; *Über die analytische darstellbarkeit sogenannter willkürlicher functionen einer reellen veränderlichen*. Berl. Ber., **1885** (1885), 633–640, 789–806.
- [58] Wold, E. F.; *Proper holomorphic embeddings of finitely and some infinitely connected subsets of  $\mathbb{C}$  into  $\mathbb{C}^2$* . Math. Z., **252** (2006), 1–9.
- [59] Wold, E. F.; *Embedding Riemann surfaces properly in  $\mathbb{C}^2$* . Int. J. Math., **17** (2006), 963–974.
- [60] Wold, E. F.; *Embedding subsets of tori properly into  $\mathbb{C}^2$* . Ann. Inst. Fourier, **57** (2007), 1537–1555.
- [61] Zalcman, L.; *Analytic capacity and rational approximation*. Lecture Notes in Mathematics, no. 50. Springer-Verlag, Berlin–New York, (1968).

## 1.6 Summary of Papers

**Paper I** presents a construction of a proper holomorphic embedding  $f: \mathbb{P}^1 \setminus C \hookrightarrow \mathbb{C}^2$ , where  $C$  is a Cantor set obtained by removing smaller and smaller vertical and horizontal strips from a square of side 2, allowing to realize it to have Lebesgue measure arbitrarily close to 4.

**Paper II** clarifies the details of a cryptical paper by Orevkov in which a construction of a proper holomorphic embedding  $\gamma: \mathbb{P}^1 \setminus C \hookrightarrow \mathbb{C}^2$  is performed with techniques completely different from the ones exploited in Paper I; in particular, it is proved that such a construction can be done to get the Cantor set  $C$  to have zero Hausdorff dimension.

**Paper III** solves the problem of simultaneously embedding properly holomorphically into  $\mathbb{C}^2$  a whole family of  $n$ -connected domains  $\Omega_r \subset \mathbb{P}^1$  such that none of the components of  $\mathbb{P}^1 \setminus \Omega_r$  reduces to a point, by constructing a continuous mapping  $\Xi: \bigcup_r \{r\} \times \Omega_r \rightarrow \mathbb{C}^2$  such that  $\Xi(r, \cdot): \Omega_r \hookrightarrow \mathbb{C}^2$  is a proper holomorphic embedding for every  $r$ . To this aim, a parametric version of both the Andersén–Lempert procedure and Carleman’s Theorem is formulated and proved.

**Paper IV** displays four approximation theorems for manifold-valued mappings. The first one approximates holomorphic embeddings on pseudoconvex domains in  $\mathbb{C}^n$  with holomorphic embeddings with dense images. The second theorem approximates holomorphic mappings on complex manifolds with bounded images with holomorphic mappings with dense images. The last two theorems work the other way around, constructing (in different settings) sequences of holomorphic mappings (embeddings in the first one) converging to a mapping with dense image defined on a given compact minus certain points (thus in general not holomorphic).



# Papers



# Proper Holomorphic Embeddings of complements of large Cantor sets in $\mathbb{C}^2$

G. D. Di Salvo and E. F. Wold

Published in *Arkiv för Matematik*, October 2022, volume 60, issue 2, pp. 323–333.

## Abstract

We present a construction of a proper holomorphic embedding  $f: \mathbb{P}^1 \setminus C \hookrightarrow \mathbb{C}^2$ , where  $C$  is a Cantor set obtained by removing smaller and smaller vertical and horizontal strips from a square of side 2, allowing to realize it to have Lebesgue measure arbitrarily close to 4.

## Contents

I.1	Introduction . . . . .	23
I.2	Preliminaries . . . . .	25
I.3	The Induction Step . . . . .	27
I.4	Proof of Theorem I.1.1 . . . . .	30

## I.1 Introduction

### I.1.1 The main result

A major unresolved issue, known as Forster’s Conjecture, is whether or not every open Riemann surface  $X$  admits a holomorphic embedding into  $\mathbb{C}^2$ , and, if it does, whether it admits a *proper* holomorphic embedding. For instance, if  $Y$  is a compact Riemann surface, and if  $X = Y \setminus C$  where  $C$  is a closed set whose connected components are all points, it is unknown whether  $X$  embeds (properly or not) into  $\mathbb{C}^2$ . We may consider two extremal cases: (i) the case where  $C$  is a finite set, and (ii) the case where  $C$  is a Cantor set, and we may further consider the simplest compact Riemann surface in this context, namely  $Y = \mathbb{P}^1$ . Then in

## I. Proper Holomorphic Embeddings of complements of large Cantor sets in $\mathbb{C}^2$

case (i), it is clear that  $X$  admits a proper holomorphic embedding into  $\mathbb{C}^2$ , so we will consider the case (ii).

Let  $Q$  denote the square  $Q = [-1, 1] \times [-1, 1] \subset \mathbb{C}$ , and let  $\mu$  denote the 2-dimensional Lebesgue measure on  $\mathbb{C}$ . A procedure for constructing a (large) Cantor set  $C \subset Q$  is as follows (see Section I.3 for a more detailed description). Let  $l_1$  denote the vertical line dividing  $Q$  into two equal pieces, choose  $\delta_1 > 0$  small, and remove an open  $\delta_1$ -neighbourhood of  $l_1$  to obtain a union  $Q_2$  of two disjoint rectangles. Next, let  $l_2^j, j = 1, 2$ , be horizontal lines dividing each rectangle in  $Q_2$  into equal pieces, choose  $\delta_2$  small, and remove an open  $\delta_2$ -neighbourhood of  $l_2^1 \cup l_2^2$  to obtain a disjoint union  $Q_3$  of four rectangles. Next, switch back to vertical lines, choose  $\delta_3$  small to obtain  $Q_4$  and so forth, to obtain a sequence  $\delta_j \rightarrow 0$  and nested sequence  $Q_j$  of rectangles such that  $C = \bigcap_j Q_j$  is a Cantor set contained in  $Q$ . Our main result is the following.

**Theorem I.1.1.** There are sequences  $\{\delta_j\}_j$  converging to zero arbitrarily fast such that the complement  $\mathbb{P}^1 \setminus C$  of the resulting Cantor set admits a proper holomorphic embedding into  $\mathbb{C}^2$ . In particular, for any  $\epsilon > 0$  we may achieve that  $\mu(C) > 4 - \epsilon$ .

The motivation for proving this result is that there have been speculations that considering complements of “fat” Cantor sets could lead to counterexamples to Forster’s Conjecture.

We note that Orevkov [18] showed the existence of a Cantor set  $C \subset \mathbb{P}^1$  such that  $\mathbb{P}^1 \setminus C$  admits a proper holomorphic embedding into  $\mathbb{C}^2$ . His construction is quite cryptical and it is explained in detail in [4], where it is also proved that  $C$  can be obtained to have zero Hausdorff dimension. From such a construction it seems difficult, or perhaps impossible, to achieve that  $C$  is large.

### I.1.2 History

Dealing with Stein manifolds, one of the most important goals to achieve is to embed them properly holomorphically into  $\mathbb{C}^N$  for some  $N$ . A first important result comes from Remmert [19], who proved in 1956 that every  $n$ -dimensional Stein manifold admits a proper holomorphic embedding into  $\mathbb{C}^N$  for  $N$  big enough. Such a result was made precise by Bishop and Narasimhan who independently proved in 1960–61 that  $N$  can be taken to be  $2n + 1$  (see [3] and [17]). In 1970 Forster [7] improved Bishop–Narasimhan’s result, decreasing  $N$  to  $\lfloor \frac{5n}{3} \rfloor + 2$  and proving that it is not possible for  $N$  to go below  $\lfloor \frac{3n}{2} \rfloor + 1$  and conjecturing that the euclidean dimension could have been improved exactly to  $\lfloor \frac{3n}{2} \rfloor + 1$ . Eliashberg, Gromov (1992) and Shürmann (1997) proved the following

**Theorem I.1.2.** [Eliashberg–Gromov [5] (1992) and Shürmann [21] (1997)] Every  $n$ -dimensional Stein manifold  $X$ , with  $n \geq 2$  embeds properly holomorphically into  $\mathbb{C}^N$  with  $N = \lfloor \frac{3n}{2} \rfloor + 1$ .

The proof of the theorem breaks down when  $n = 1$ ; since 1-dimensional Stein manifolds are precisely open connected Riemann surfaces, Forster’s conjecture reduces to the following

**Conjecture I.1.1.** Every open connected Riemann surface embeds properly holomorphically into  $\mathbb{C}^2$ .

So far, only a few open Riemann surfaces are known to admit a proper holomorphic embedding into  $\mathbb{C}^2$ . The first known examples are the open unit disk in  $\mathbb{C}$  (Kasahara–Nishino, 1970, [22]), open annuli in  $\mathbb{C}$  (Laufer, 1973, [14]) and the punctured disk in  $\mathbb{C}$  (Alexander, 1977, [1]). Later (1995) Stensønes and Globevnik proved in [12] that every finitely connected planar domain without isolated boundary points verifies the conjecture. In 2009, Wold and Forstnerič proved the best result known so far: if  $\overline{D}$  is a Riemann surface with smooth enough boundary which admits a smooth embedding into  $\mathbb{C}^2$ , holomorphic on the interior  $D$ , then  $D$  admits a proper holomorphic embedding into  $\mathbb{C}^2$  (see [10] and next section). Other remarkable results include proper holomorphic embeddings of certain Riemann surfaces into  $\mathbb{C}^2$  with interpolation (see [13]), deformation of continuous mappings  $f: S \rightarrow X$  between Stein manifolds into proper holomorphic embeddings under certain hypothesis on the dimension of the spaces (see [2]), embeddings of infinitely connected planar domains into  $\mathbb{C}^2$  (see [11]), the existence of a homotopy of continuous mappings  $f: D \rightarrow \mathbb{C} \times \mathbb{C}^*$  into proper holomorphic embedding whenever  $D$  is a finitely connected planar domain without punctures (see [20]), existence of proper holomorphic embeddings of the unit disc  $\mathbb{B}$  into connected pseudoconvex Runge domains  $\Omega \subset \mathbb{C}^n$  (when  $n \geq 2$ ) whose image contains arbitrarily fixed discrete subsets of  $\Omega$  (see [9]), approximation of proper embeddings on smooth curves contained in a finitely connected planar domain  $D$  into  $\mathbb{C}^n$  (with  $n \geq 2$ ) by proper holomorphic embeddings  $f: D \hookrightarrow \mathbb{C}^n$  (see [15]), and the existence of proper holomorphic embeddings into  $\mathbb{C}^2$  of certain infinitely connected domains  $\Omega$  lying inside a bordered Riemann surface  $\overline{D}$  knowing to admit a proper holomorphic embedding into  $\mathbb{C}^2$  [16].

## I.2 Preliminaries

### I.2.1 Notation

We will use the following notation.

- Given  $K \subset \mathbb{C}$  and a positive real number  $\delta$ , we define the open subset

$$K(\delta) := \{z \in \mathbb{C} : \text{dist}(z, K) < \delta\}.$$

- For a closed subset  $K \subset \mathbb{P}^1$  we denote by  $\mathcal{O}(K)$  the algebra of continuous functions  $f \in \mathcal{C}(K)$  such that there exists an open set  $U \subset \mathbb{P}^1$  containing  $K$ , and  $F \in \mathcal{O}(U)$  with  $F|_K = f$ .
- We let  $\pi_j: \mathbb{C}^2 \rightarrow \mathbb{C}$  denote the projection onto the  $j$ -th coordinate line, and given a point  $p \in \mathbb{C}^2$  we denote the vertical complex line through  $p$  by

$$\Lambda_p := \pi_1^{-1}(\pi_1(p)) = \{(\pi_1(p), \zeta) : \zeta \in \mathbb{C}\}.$$

## I. Proper Holomorphic Embeddings of complements of large Cantor sets in $\mathbb{C}^2$

- If  $X$  is a domain with piecewise smooth boundary in a Riemann surface  $Y$ ,  $f: \overline{X} \rightarrow \mathbb{C}^2$  is a holomorphic map, and if  $a \in \partial X$  is a smooth boundary point, we say that  $f(a)$  is  $\pi_1$ -exposed for  $f(\overline{X})$  if  $f(\overline{X}) \cap \Lambda_{f(a)} = \{f(a)\}$ , and  $\pi_1 \circ f$  is an embedding sufficiently close to  $a$ . Similarly, for a smooth map  $\gamma: [0, 1] \rightarrow \mathbb{C}^2$ , we say that  $\gamma([0, 1])$  is exposed at  $\gamma(1)$  if  $\gamma([0, 1]) \cap \Lambda_{\gamma(1)} = \{\gamma(1)\}$ , and  $\pi_1 \circ \gamma$  is an embedding sufficiently close to 1.

### I.2.2 Results

In this section, we collect the technical tools needed to prove Theorem I.1.1.

The following result is essentially Theorem 4.2 in [10]. Although (1) and (2) were not stated explicitly in [10] they are evident from the proof therein and were added to the corresponding Theorem 2.8 in [11].

**Theorem I.2.1.** Let  $X$  be a smoothly bounded domain in a Riemann surface  $Y$ ,  $f: \overline{X} \hookrightarrow \mathbb{C}^2$  a holomorphic embedding, and  $a_1, \dots, a_m \in \partial X$ . Let  $\gamma_j: [0, 1] \rightarrow \mathbb{C}^2$  ( $j = 1, \dots, m$ ) be smooth embedded arcs with pairwise disjoint images satisfying the following properties:

- $\gamma_j([0, 1]) \cap f(\overline{X}) = \gamma_j(0) = f(a_j)$  for  $j = 1, \dots, m$ , and
- the image  $E := f(\overline{X}) \cup \bigcup_{j=1}^m \gamma_j([0, 1])$  is  $\pi_1$ -exposed at  $\gamma_j(1)$  for  $j = 1, \dots, m$ .

Then given an open set  $V \subset \mathbb{C}^2$  containing  $\bigcup_{j=1}^m \gamma_j([0, 1])$ , an open set  $U \subset Y$  containing the points  $a_j$  that satisfies  $f(\overline{U \cap X}) \subset V$ , and any  $\epsilon > 0$ , there exists a holomorphic embedding  $F: \overline{X} \hookrightarrow \mathbb{C}^2$  with the following properties:

- (1)  $\|F - f\|_{\overline{X} \setminus U} < \epsilon$ ,
- (2)  $F(\overline{U \cap X}) \subset V$ , and
- (3)  $F(a_j) = \gamma_j(1)$ , and  $F(\overline{X})$  is  $\pi_1$ -exposed at  $F(a_j)$  for  $j = 1, \dots, m$ .

The following is essentially Lemma 1 in [23]. The difference is that Lemma 1 was stated for  $\pi_1$  instead of  $\pi_2$ , and for curves  $\lambda: [0, +\infty) \rightarrow \mathbb{C}^2$  instead of  $\lambda: (-\infty, +\infty) \rightarrow \mathbb{C}^2$  – neither make a difference for the proof.

**Lemma I.2.1.** Let  $K \subset \mathbb{C}^2$  be a polynomially convex compact set, and let

$$\Lambda = \{\lambda_j(t) : j = 1, \dots, m, \ t \in (-\infty, +\infty)\}$$

be a collection of pairwise disjoint smooth curves in  $\mathbb{C}^2 \setminus K$  without self intersection, enjoying the *immediate projection property* (with respect to  $\pi_2$ ):

- $\lim_{|t| \rightarrow \infty} |\pi_2(\lambda_j(t))| = \infty$  for all  $j$ , and
- there exists an  $M > 0$  such that  $\mathbb{C} \setminus (R\mathbb{B} \cup \pi_2(\Lambda))$  does not contain any relatively compact components for  $R \geq M$ .

Then for any  $r > 0$  and  $\epsilon > 0$  there exists  $\phi \in \text{Aut } \mathbb{C}^2$  such that the following are satisfied:

- (i)  $\|\phi - \text{Id}\|_K < \epsilon$ , and
- (ii)  $\phi(\Lambda) \subset \mathbb{C}^2 \setminus r\overline{\mathbb{B}^2}$ .

### I.3 The Induction Step

We will now describe an inductive procedure to construct a nested sequence of closed rectangles  $Q_n \subset Q$ , along with holomorphic embeddings  $f_n: \overline{\mathbb{P}^1 \setminus Q_n} \rightarrow \mathbb{C}^2$  that will be used to construct a proper holomorphic embedding

$$f: \mathbb{P}^1 \setminus \bigcap_n Q_n \hookrightarrow \mathbb{C}^2,$$

where  $C = \bigcap_n Q_n$  will be a Cantor set, where the construction will enable us to ensure that its Lebesgue measure  $\mu(C)$  is arbitrarily close to 4.

Set  $Q_1 := Q$  and set  $C_1 := \overline{\mathbb{P}^1 \setminus Q_1}$ . To construct  $Q_2$  from  $Q_1$  we let  $l_1$  be the vertical line segment dividing  $Q_1$  into two equal pieces. Then, for  $0 < \delta_2 < 1$ , we set

$$Q_2 := Q_1 \setminus l_1(\delta_2),$$

and we set  $C_2 := \overline{\mathbb{P}^1 \setminus Q_2}$ . Then  $C_2$  is the complement of the disjoint union of 2 open rectangles  $(Q_2^j)^\circ$ ,  $j = 1, 2$ , contained in  $Q_1$ .

Assume now that we have constructed a nested sequence  $\{Q_j\}_{j=1}^n$ ,  $n \geq 2$ , where

$$Q_n = \bigsqcup_{j=1}^{2^{n-1}} Q_n^j$$

is the disjoint union of  $2^{n-1}$  closed rectangles contained in  $Q_{n-1}$ , along with an increasing sequence of closed subsets  $C_n := \overline{\mathbb{P}^1 \setminus Q_n}$  in  $\mathbb{P}^1$ . We let  $l_n^j$  be the line segment – vertical for  $n$  odd, horizontal for  $n$  even – dividing  $Q_n^j$  into two equal pieces, we set  $l_n := \bigsqcup_{j=1}^{2^{n-1}} l_n^j$ , and for  $\delta_{n+1} > 0$  small enough we define

$$Q_{n+1} := Q_n \setminus l_n(\delta_{n+1}) =: \bigsqcup_{j=1}^{2^n} Q_{n+1}^j,$$

and

$$C_{n+1} := \overline{\mathbb{P}^1 \setminus Q_{n+1}}.$$

**Proposition I.3.1.** With the procedure above assume that we have constructed  $Q_n$  and  $C_n$  for  $n \geq 1$ . Let  $K_n \subset C_n^\circ$  be a compact set, let  $r_n > 0$ , and assume that  $f_n: C_n \hookrightarrow \mathbb{C}^2$  is a holomorphic embedding such that

$$f_n(\overline{C_n \setminus K_n}) \subset \mathbb{C}^2 \setminus r_n \overline{\mathbb{B}^2}. \quad (\text{I.3.1})$$

Then for any  $\epsilon_n > 0$  and any  $r_{n+1} > r_n$ , there exist  $\delta_{n+1} > 0$  arbitrarily close to zero and a holomorphic embedding  $f_{n+1}: C_{n+1} \hookrightarrow \mathbb{C}^2$  such that

## I. Proper Holomorphic Embeddings of complements of large Cantor sets in $\mathbb{C}^2$

- (a)  $\|f_{n+1} - f_n\|_{K_n} < \varepsilon_n$ ,
- (b)  $f_{n+1}(\overline{C_{n+1} \setminus K_n}) \subset \mathbb{C}^2 \setminus r_n \overline{\mathbb{B}^2}$ ,
- (c)  $f_{n+1}(\partial C_{n+1}) \subset \mathbb{C}^2 \setminus r_{n+1} \overline{\mathbb{B}^2}$ .

*Proof.* We extend  $f_n$  to a smooth embedding  $\tilde{f}_n: C_n \cup l_n \hookrightarrow \mathbb{C}^2$  with  $\tilde{f}_n(l_n)$  lying close enough to  $f_n(\partial C_n)$  so that by (I.3.1) we get

$$\tilde{f}_n(l_n) \subset \mathbb{C}^2 \setminus r_n \overline{\mathbb{B}^2}. \quad (\text{I.3.2})$$

Now Mergelyan's theorem (see e.g., [6]) ensures the existence of a holomorphic embedding  $\hat{f}_{n+1}: C_n \cup l_n \hookrightarrow \mathbb{C}^2$  such that

$$\|\hat{f}_{n+1} - \tilde{f}_n\|_{C_n \cup l_n} < \frac{\varepsilon_n}{4},$$

and

$$\hat{f}_{n+1}(\overline{(C_n \cup l_n) \setminus K_n}) \subset \mathbb{C}^2 \setminus r_n \overline{\mathbb{B}^2}. \quad (\text{I.3.3})$$

Then by choosing a preliminary  $\tilde{\delta}_{n+1} > 0$  sufficiently small (to be shrunk further later), and letting the set corresponding to  $C_{n+1}$  be denoted by  $\tilde{C}_{n+1}$  (and similarly for  $Q_{n+1}$ ), we have that  $f_{n+1} \in \mathcal{O}(\tilde{C}_{n+1})$ , and

$$\hat{f}_{n+1}(\overline{(\tilde{C}_{n+1} \setminus K_n)}) \subset \mathbb{C}^2 \setminus r_n \overline{\mathbb{B}^2}. \quad (\text{I.3.4})$$

Next, recall that  $\tilde{Q}_{n+1}$  is constructed from  $Q_n$  by splitting each  $Q_n^j$  into two smaller rectangles  $\tilde{Q}_n^{j,1}$  and  $\tilde{Q}_n^{j,2}$ , by removing the strip  $l_n^j(\tilde{\delta}_{n+1})$ . Choose smooth boundary points

$$\tilde{a}_{ji} \in \partial \tilde{Q}_n^{j,i} \cap \overline{l_n^j(\tilde{\delta}_{n+1})}, \quad (\text{I.3.5})$$

for  $i = 1, 2$ , and  $j = 1, \dots, 2^{n-1}$ , and relabel these to get  $2^n$  boundary points  $a_j$ , one in each  $\partial \tilde{Q}_{n+1}^j$ .

Now choose  $2^n$  pairwise disjoint smoothly embedded arcs  $\gamma_j: [0, 1] \hookrightarrow \mathbb{C}^2$  disjoint from  $r_n \overline{\mathbb{B}^2}$ , such that

$$\gamma_j([0, 1]) \cap \hat{f}_{n+1}(\tilde{C}_{n+1}) = \hat{f}_{n+1}(a_j) = \gamma_j(0),$$

and such that each point  $\gamma_j(1)$  is  $\pi_1$ -exposed for the surface

$$\hat{f}_{n+1}(\tilde{C}_{n+1}) \cup \bigcup_{j=1}^{2^n} \gamma_j([0, 1]).$$

Choose an open set  $V \subset \mathbb{C}^2$  containing the arcs  $\gamma_j([0, 1])$  with  $\bar{V} \cap r_n \overline{\mathbb{B}^2} = \emptyset$  and take  $U \subset \mathbb{P}^1$  to be the union of sufficiently small open balls centered at the points  $a_j$ , so that  $U \cap K_n = \emptyset$  and  $\hat{f}_{n+1}(U \cap \tilde{C}_{n+1}) \subset V$ . Then Theorem I.2.1 furnishes



a holomorphic embedding  $F_{n+1}: \tilde{C}_{n+1} \hookrightarrow \mathbb{C}^2$  such that  $p_j := F_{n+1}(a_j) = \gamma_j(1)$  is an exposed point for  $F_{n+1}(\tilde{C}_{n+1})$  for each  $j$ , and

$$\|F_{n+1} - \hat{f}_{n+1}\|_{K_n} < \frac{\varepsilon_n}{4}, \quad (\text{I.3.6})$$

and also

$$F_{n+1}(\overline{\tilde{C}_{n+1} \setminus K_n}) \subset \mathbb{C}^2 \setminus r_n \overline{\mathbb{B}^2}. \quad (\text{I.3.7})$$

Now choose  $\alpha_j \in \mathbb{C}$ ,  $j = 1, \dots, 2^n$ , such that setting

$$g_{n+1}(z, w) := \left( z, w + \sum_{j=1}^{2^n} \frac{\alpha_j}{\pi_1(p_j) - z} \right),$$

we have that

$$\|g_{n+1} \circ F_{n+1} - F_{n+1}\|_{K_n} < \frac{\varepsilon_n}{4},$$

and such that the conditions in Lemma I.2.1 are satisfied for the collection  $\Lambda$  of curves

$$\lambda_{ji} := g_{n+1} \circ F_{n+1}(\partial \tilde{Q}_n^{j,i}), \quad i = 1, 2, \quad j = 1, \dots, 2^{n-1},$$

that are the boundary of the unbounded complex curve

$$X_{n+1} := g_{n+1} \circ F_{n+1}(\tilde{C}_{n+1}).$$

Note that we still have

$$g_{n+1} \circ F_{n+1}(\overline{\tilde{C}_{n+1} \setminus K_n}) \subset \mathbb{C}^2 \setminus r_n \overline{\mathbb{B}^2}. \quad (\text{I.3.8})$$

Choose  $0 < \eta < 1$  such that  $(r_n + \eta) \overline{\mathbb{B}^2} \cap \Lambda = \emptyset$ . We may choose a compact polynomially convex set  $K' \subset X_{n+1}^\circ$  with  $g_{n+1} \circ F_{n+1}(K_n) \subset K'$  such that  $L = (r_n + \eta) \overline{\mathbb{B}^2} \cup K'$  is polynomially convex (see e.g., Theorem 4.14.6 in [8]). Then by Lemma I.2.1 there exists  $\phi_{n+1} \in \text{Aut } \mathbb{C}^2$  such that

$$\phi_{n+1}(\Lambda) \subset \mathbb{C}^2 \setminus r_{n+1} \overline{\mathbb{B}^2}, \quad (\text{I.3.9})$$

and

$$\|\phi_{n+1} - \text{Id}\|_L < \frac{\min\{\eta, \varepsilon_n\}}{4}. \quad (\text{I.3.10})$$

We consider the map  $f_{n+1}: \tilde{C}_{n+1} \rightarrow \mathbb{C}^2$  defined by

$$f_{n+1} := \phi_{n+1} \circ g_{n+1} \circ F_{n+1}.$$

We have that (a) and (b) (with  $\tilde{C}_{n+1}$  instead of  $C_{n+1}$ ) clearly hold, but now  $f_{n+1}$  has singularities on  $\partial \tilde{C}_{n+1}$ . However, we now consider  $0 < \delta_{n+1} < \tilde{\delta}_{n+1}$  to see what happens on  $C_{n+1}$ . As the points to expose are taken on the boundary components (see (I.3.5)), the singularities of  $f_{n+1}$  are not contained in  $C_{n+1}$  for any such  $\delta_{n+1}$ , and so  $f_n: C_{n+1} \rightarrow \mathbb{C}^2$  is holomorphic. Finally, since  $\partial C_{n+1}$  will converge to  $\partial \tilde{C}_{n+1}$  as  $\delta_{n+1} \rightarrow \tilde{\delta}_{n+1}$  we have (c) for  $\delta_{n+1}$  sufficiently close to  $\tilde{\delta}_{n+1}$ . ■

## I.4 Proof of Theorem I.1.1

We will prove Theorem I.1.1 via an inductive construction, where Proposition I.3.1 provides us with the inductive step. Without loss of generality, we may assume that  $\epsilon < 1$ .

### I.4.1 The Induction Scheme

To start the induction, with the notation as in Section I.3, we define  $f_1: C_1 \hookrightarrow \mathbb{C}^2$  by  $f_1(\zeta) := (2/\zeta, 0)$  for  $\zeta \in \mathbb{C}$ , and  $f_1(\infty) := (0, 0)$ . Setting  $r_1 = 1$  we note that  $f_1(\partial C_1) \subset \mathbb{C}^2 \setminus r_1 \overline{\mathbb{B}^2}$ , so if we choose  $0 < \delta'_1 < 1$  sufficiently close to zero, and set

$$K_1 := \mathbb{P}^1 \setminus Q_1(\delta'_1),$$

we have that  $K_1 \subset C_1^\circ$  and  $f_1(\overline{C_1 \setminus K_1}) \subset \mathbb{C}^2 \setminus r_1 \overline{\mathbb{B}^2}$ . Then the conditions in Proposition I.3.1 are satisfied with  $n = 1$ , and setting  $\delta_2 \leq \epsilon \cdot 2^{-4}$ ,  $r_2 = 2$ , we let  $f_2$  be the map furnished by the proposition, with  $\epsilon_1$  explained in the induction scheme below. We then choose  $\delta'_2 < \delta_2/2$  sufficiently close to zero such that if we set

$$K_2 := \mathbb{P}^1 \setminus Q_2(\delta'_2),$$

we have  $K_2 \subset C_2^\circ$  and  $f_2(\overline{C_2 \setminus K_2}) \subset \mathbb{C}^2 \setminus r_2 \overline{\mathbb{B}^2}$ .

Let us now state our induction hypothesis  $I_n$  for some  $n \geq 2$ . We assume that we have found and constructed the following.

- (i)<sub>n</sub> A decreasing sequence  $\delta_2 > \delta_3 > \dots > \delta_n$  of numbers with  $\delta_k \leq \epsilon \cdot 2^{-2k}$  such that  $\{Q_k\}_{k=1}^n$  is a nested sequence of rectangles.
- (ii)<sub>n</sub> A decreasing sequence  $\delta'_1 > \delta'_2 > \dots > \delta'_n$  of numbers with  $\delta'_1, \delta'_2$  as above, and  $\delta'_k < \delta_k/2$  for  $k = 1, \dots, n$ , and holomorphic embeddings  $f_k: C_k \hookrightarrow \mathbb{C}^2$  such that, setting  $K_k := \mathbb{P}^1 \setminus Q_k(\delta'_k)$ , we have that  $f_m(\overline{C_m \setminus K_k}) \subset \mathbb{C}^2 \setminus r_k \overline{\mathbb{B}^2}$  for  $1 \leq k \leq m \leq n$ , where  $r_k \geq k$ .
- (iii)<sub>n</sub> A sequence of positive numbers  $\{\eta_k\}_{k=2}^n$  such that if  $f: K_k \rightarrow \mathbb{C}^2$  is a holomorphic map with  $\|f - f_k\|_{K_k} < \eta_k$ , then  $f: K_{k-1} \hookrightarrow \mathbb{C}^2$  is an embedding.
- (iv)<sub>n</sub> A sequence of positive numbers  $\{\varepsilon_k\}_{k=1}^{n-1}$  such that  $\varepsilon_{k+j} < \eta_k \cdot 2^{-j-1}$ ,  $j \leq n - k - 1$ , with  $\|f_k - f_{k-1}\|_{K_{k-1}} < \varepsilon_{k-1}$  for  $k = 2, \dots, n$ .

Our constructions above gives (i)<sub>n</sub>, (ii)<sub>n</sub> and (iv)<sub>n</sub> in the case  $n = 2$  (possibly shrinking  $\epsilon_1$ ). Then, choosing  $\eta_2$  small enough, gives  $f$  and  $f'$  close to  $f_2$  and  $f'_2$  respectively (the latter by Cauchy estimates) on  $K_2$  such that  $f$  is injective and  $f'$  never vanishes on  $K_1$ . Being  $K_1$  compact, this is enough to achieve (iii)<sub>n</sub> when  $n = 2$ .

### I.4.2 Passing from $I_n$ to $I_{n+1}$

Let us assume that  $I_n$  is true and prove  $I_{n+1}$ . First of all we have that (i) $_{n+1}$ , (iii) $_{n+1}$  and the first part of (iv) $_{n+1}$  are just a matter of choosing respectively  $\delta_{n+1}$ ,  $\eta_{n+1}$  and  $\epsilon_n$  sufficiently small. By (ii) $_n$  with  $k = n$ , and with  $\epsilon_n$  above fixed, we may apply Proposition I.3.1 to get a holomorphic embedding  $f_{n+1}: C_{n+1} \hookrightarrow \mathbb{C}^2$  to obtain the second part of (iv) $_{n+1}$  and (ii) $_{n+1}$  with  $m = n + 1$  and  $k = n$ . Next, by choosing  $\delta'_{n+1}$  sufficiently small we get (ii) $_{n+1}$  for  $k = m = n + 1$ . It remains to explain how to achieve (ii) $_{n+1}$  for  $m = n + 1$  and  $k = 1, \dots, n - 1$ . Since

$$\overline{C_{n+1} \setminus K_k} = \overline{C_{n+1} \setminus K_n} \cup \overline{K_n \setminus K_k}$$

what is needed is  $f_{n+1}(\overline{K_n \setminus K_k}) \subset \mathbb{C}^2 \setminus r_k \overline{\mathbb{B}^2}$ . This follows from (ii) $_n$ , possibly after having decreased  $\epsilon_n$ .

### I.4.3 Proof of Theorem I.1.1

Consider the objects constructed in the inductive scheme above. Then by (iv) $_n$  we have that  $\lim_{j \rightarrow \infty} f_j = f$  exists on  $K_k$  for any  $k$ . We have that

$$\left( \bigcup_k C_k \right)^\circ = \mathbb{P}^1 \setminus \bigcap_k Q_k =: \mathbb{P}^1 \setminus C$$

and so  $\lim_{j \rightarrow \infty} f_j = f$  exists on  $\mathbb{P}^1 \setminus C$ . Now for any fixed  $k$  we get by (ii) $_n$  that  $f_n^{-1}(r_k \overline{\mathbb{B}^2}) \subset K_k$  for all  $n > k$  and therefore  $f^{-1}(r_k \overline{\mathbb{B}^2}) \subset K_k$ , so  $f$  is proper. By (iv) $_n$  we get that  $\|f - f_k\|_{K_k} < \eta_k$ , hence by (iii) $_n$  we have that  $f: K_{k-1} \hookrightarrow \mathbb{C}^n$  is an embedding for all  $k$ , so  $f$  is an embedding. Finally, note that when constructing a rectangle  $Q_{n+1}$  from  $Q_n$ , a crude estimate gives that one obtains  $Q_{n+1}$  by removing strips of total area bounded by  $2^n \cdot \delta_n$ . It follows that

$$\mu(C) = \mu\left(\bigcap_n Q_n\right) \geq 4 - \sum_{n=1}^{\infty} 2^n \cdot \delta_n \geq 4 - \epsilon \cdot \sum_{n=1}^{\infty} 2^{-n} > 4 - \epsilon.$$

■



# Bibliography

- [1] Alexander, H.; Explicit imbedding of the (punctured) disc into  $\mathbb{C}^2$ . *Comment. Math. Helv.*, **52**, 539–544, 1977.
- [2] Andrist, R., Forstnerič, F., Ritter, T. and Wold, E. F.; Proper holomorphic embeddings into Stein manifolds with the density property. *J. Anal. Math.*, **130** (2016), 135–150.
- [3] Bishop, E.; Mappings of partially analytic spaces. *Am. J. Math.*, **83**, 209–242, 1961.
- [4] Di Salvo, G. D.; Extended explanation of Orevkov’s paper on proper holomorphic embeddings of complements of Cantor spaces in  $\mathbb{C}^2$  and a discussion of their measure. Second article in author’s Ph.D. thesis, Universitetet i Oslo, 2022.
- [5] Eliashberg, Y. and Gromov, M.: Embeddings of Stein manifolds. *Ann. Math.*, (2), **136**, 123–135 (1992)
- [6] Fornæss, J. E., Forstnerič, F. and Wold, E. F.; Holomorphic approximation: the legacy of Weierstrass, Runge, Oka–Weil and Mergelyan. *Advancements in Complex Analysis – From Theory to Practice*. Breaz, Daniel & Rassias, Michael Th. (Ed.), Springer Nature. ISSN 978–3–030–40119–1. p. 133–192.
- [7] Forster, O; Plongements des variétés de Stein. *Comment. Math. Helv* **45** 170–184 (1970).
- [8] Forstnerič, F.; Stein manifolds and holomorphic mappings. The homotopy principle in complex analysis. Second edition. *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*, 56. Springer, Cham, 2017.
- [9] Forstnerič, F., Globevnik, J. and Stensønes, B.; Embedding holomorphic discs through discrete sets. *Math. Ann.*, **305** (1996), no. 3, 559–569.
- [10] Forstnerič, F. and Wold, E. F.; Bordered Riemann Surfaces in  $\mathbb{C}^2$ , *J. Math. Pures Appl.* **91** (2009) 100–114.
- [11] Forstnerič, F. and Wold, E. F.; Embeddings of infinitely connected planar domains into  $\mathbb{C}^2$ . *Anal. PDE* **6** (2013), no. 2, 499–514.
- [12] Globevnik, J. and Stensønes, B.; Holomorphic embeddings of planar domains into  $\mathbb{C}^2$  *Math. Ann.*, **303**, 579–597 (1995).

- [13] Kutzschebauch, F., Løv, E. and Wold, E. F.; Embedding some Riemann surfaces into  $\mathbb{C}^2$  with interpolation. *Math. Z.*, **262** (2009), no. 3, 603–611.
- [14] Laufer, H. B.; Imbedding annuli in  $\mathbb{C}^2$ . *J. Anal. Math.*, **26**, 187–215 (1973).
- [15] Majcen, I.; Proper holomorphic embeddings of finitely connected planar domains into  $\mathbb{C}^n$ . *Ark. Mat.* **51** (2013), no. 2, 329–343.
- [16] Majcen, I.; Embedding certain infinitely connected subsets of bordered Riemann surfaces properly into  $\mathbb{C}^2$ . *J. Geom. Anal.*, **19** (2009), no. 3, 695–707.
- [17] Narasimhan, R.; Holomorphic mappings of complex spaces. *Proc. Am. Math. Soc.*, **11**, 800–804 (1960).
- [18] Orevkov, S. Y.; Proper analytic embedding of  $\mathbb{CP}^1$  minus Cantor set into  $\mathbb{C}^2$ . *Uspekhi Mat. Nauk.* **63** (2008), no. 1(379), 155–156 (in Russian) English transl.: Russian Math. Surveys, **63** (2008), 168–169.
- [19] Remmert, R.; Sur les espaces analytiques holomorphiquement séparables et holomorphiquement convexes. *C. R. Acad. Sci. Paris* **243**, 211–221 (1956).
- [20] Ritter, T.; A strong Oka principle for embeddings of some planar domains into  $\mathbb{C} \times \mathbb{C}^*$ . *J. Geom. Anal.*, **23**(2), 571–597 (2013).
- [21] Schürmann, J.; Embeddings of Stein spaces into affine spaces of minimal dimension. *Math. Ann.*, **307**, 381–399, 1997.
- [22] Stehlé, J.-L.; Plongements du disque dans  $\mathbb{C}^2$ . In: Séminaire P. Lelong (Analyse). *Lect. Notes in Math.*, vol 275, pp. 119–130. Springer-Verlag, Berlin (1970).
- [23] Wold, E. F.; Proper holomorphic embeddings of finitely and some infinitely connected subsets of  $\mathbb{C}$  into  $\mathbb{C}^2$ . *Math. Z.* **252** (2006), no. 1, 1–9.

### Authors' addresses

**G. D. Di Salvo** University of Oslo, Postboks 1337 Blindern, 0316 Oslo, Norway,  
giovandd@math.uio.no

**E. F. Wold** University of Oslo, Postboks 1337 Blindern, 0316 Oslo, Norway,  
erlendfw@math.uio.no

Paper II

Extended explanation of Orevkov’s  
paper on proper holomorphic  
embeddings of complements of  
Cantor sets in  $\mathbb{C}^2$  and a discussion  
of their measure

G. D. Di Salvo

Available on arXiv: 2211.05046v2.

Abstract

We clarify the details of a cryptical paper by Orevkov in which a construction of a proper holomorphic embedding  $\gamma: \mathbb{P}^1 \setminus C \hookrightarrow \mathbb{C}^2$  is performed with techniques completely different from the ones exploited in Paper I; in particular, it is proved that such a construction can be done to get the Cantor set  $C$  to have zero Hausdorff dimension.

Contents

II.1	Introduction . . . . .	35
II.2	Definitions and preliminary results . . . . .	36
II.3	General Properties . . . . .	40
II.4	The Cantor set $C$ and the proper holomorphic embedding $\gamma$	45

II.1 Introduction

II.1.1 The main result and motivation

A very relevant open problem in complex geometry is to investigate whether every open Riemann surface  $\mathcal{R}$  admits a proper holomorphic embedding into  $\mathbb{C}^2$ . It has been suggested to search for a counterexample as the complement of a “thick” Cantor set  $C$  inside the Riemann Sphere, that is  $\mathcal{R} = \overline{\mathbb{C}} \setminus C$ . In paper

## II. Explanation of Orevkov's Paper on p.h.e. $\overline{\mathbb{C}} \setminus C \hookrightarrow \mathbb{C}^2$ and on the size of $C$ .

[5], the author proved the existence of a proper holomorphic embedding from the complement of a Cantor set  $C \subset \overline{\mathbb{C}}$  into  $\mathbb{C}^2$ . Nevertheless, no information dealing with the size of  $C$  is given; moreover, such a paper is quite cryptical. The motivations for the present paper are therefore two: to write in detail the construction of both the Cantor set  $C \subset \overline{\mathbb{C}}$  and the proper holomorphic embedding  $\gamma: \overline{\mathbb{C}} \setminus C \hookrightarrow \mathbb{C}^2$  originally presented in [5] and discuss the size of  $C$ , proving in particular that it can be constructed to have zero Hausdorff dimension. From the construction, it will be rather clear that it is really hard to prove that this  $C$  could have any kind of positive measure.

The main result of the present paper, originally presented by S. Orevkov in [5] without the estimate on the size of  $C$ , is the following

**Theorem II.1.1.** There exists a Cantor set  $C \subset \overline{\mathbb{C}}$  with  $\dim_H(C) = 0$  and a proper holomorphic embedding  $\gamma: \overline{\mathbb{C}} \setminus C \hookrightarrow \mathbb{C}^2$ .

It is worth mentioning that in [3] has been proved that the complement of a thick Cantor set inside  $\overline{\mathbb{C}}$  does not constitute a counterexample, as it is therein provided the construction of a proper holomorphic embedding  $\overline{\mathbb{C}} \setminus C \hookrightarrow \mathbb{C}^2$  where  $C \subset \overline{\mathbb{C}}$  is a Cantor set of arbitrarily large measure. Namely: such  $C$  is realized as a subset of a square whose side has length 2 and for every  $\epsilon > 0$  it can be built of Lebesgue measure greater than  $4 - \epsilon$ .

## II.2 Definitions and preliminary results

### II.2.1 Cantor sets

The Cantor set is defined as the intersection of the decreasing family of subsets of the unit interval of the real line  $E_0 = [0, 1] \subset \mathbb{R}$  defined inductively by removing at each step the open middle third from each interval of the previous set. Thus  $E_1 = [0, \frac{1}{3}] \cup [\frac{2}{3}, 1]$ ,  $E_2 = [0, \frac{1}{9}] \cup [\frac{2}{9}, \frac{1}{3}] \cup [\frac{2}{3}, \frac{7}{9}] \cup [\frac{8}{9}, 1]$  and so on. Finally one defines

$$E := \bigcap_{n \geq 0} E_n \subset [0, 1].$$

Given a topological space  $X$ , we define *Cantor set* any subset  $C \subset X$  homeomorphic to  $E$ . Hence a Cantor set is characterized by the topological properties of  $E$ ; thus, according to Brouwer's theorem in [2],  $C \subset X$  is a Cantor set if and only if it is

- (i)<sub>C</sub> not-empty ;
- (ii)<sub>C</sub> perfect (i.e. closed and with no isolated points) ;
- (iii)<sub>C</sub> compact ;
- (iv)<sub>C</sub> totally disconnected (i.e. every its connected component is a one-point set) ;
- (v)<sub>C</sub> metrizable .



## II.2.2 Spherical Distance

The one point compactification of  $\mathbb{R}^n$  is homeomorphic to  $\mathbb{S}^n = \{(x_1, \dots, x_{n+1}) \in \mathbb{R}^{n+1} : x_1^2 + \dots + x_{n+1}^2 = 1\}$  via stereographic projection:

$$\begin{aligned} \varphi_n : \overline{\mathbb{R}^n} &\rightarrow \mathbb{S}^n \\ \varphi_n(x) &:= \frac{2}{\|x\|^2 + 1} \left( x, \frac{\|x\|^2 - 1}{2} \right), \quad x \neq \infty; \\ \varphi_n(\infty) &:= (0, \dots, 0, 1). \end{aligned}$$

This allows us to define a notion of distance on any such  $\overline{\mathbb{R}^n}$  as

$$d_n(x, y) := \arccos(\varphi_n(x) \cdot \varphi_n(y)), \quad x, y \in \overline{\mathbb{R}^n},$$

(see e.g., [1], Theorem 5 pag. 444) where  $\cdot$  denotes the scalar product in  $\mathbb{R}^{n+1}$ . The spaces  $\overline{\mathbb{C}}$  and  $\overline{\mathbb{C}^2}$  are just the case  $n = 2$  and  $n = 4$  respectively, so we have a notion of distance on them, allowing us to talk about the diameter of their subsets. Namely, for a connected subset  $\Omega \subseteq \overline{\mathbb{R}^n}$  we denote its diameter as  $|\Omega| := \sup\{d_n(x, y) : x, y \in \Omega\}$  and if  $\Omega = \bigcup_j \omega_j$ , where  $\omega_j$  are pairwise disjoint connected subsets of  $\overline{\mathbb{R}^n}$  then we define  $|\Omega| := \sup_j |\omega_j|$ .

## II.2.3 Definitions and technical results

We will consider the Riemann Sphere  $\overline{\mathbb{C}} := \mathbb{C} \cup \{\infty\}$  and the one-point compactification of  $\mathbb{C}^2$  as  $\overline{\mathbb{C}^2} := \mathbb{C}^2 \cup \{\infty_2\}$ . Given a subset  $B \subset \mathbb{C}^2$  and a positive number  $\alpha > 0$ , we define the open subset

$$B(\alpha) := \{z \in \mathbb{C}^2 : \text{dist}(z, B) < \alpha\}.$$

Given a subset  $\Omega$  of  $\overline{\mathbb{C}}$  (resp.  $\overline{\mathbb{C}^2}$ ), we say that  $\Omega$  is *bounded* if  $\Omega \cap U = \emptyset$  for some  $U$  open neighborhood of  $\infty$  (resp. of  $\infty_2$ ), *unbounded* otherwise. We consider  $\{\epsilon_n\}_{n \geq 1}, \{R_n\}_{n \geq 0}$  strictly monotone sequences of positive real numbers, with  $\epsilon_n \rightarrow 0$  and  $R_n \rightarrow +\infty$ . Consider moreover  $\{a_n\}_{n \geq 1} \subset \mathbb{C}$  such that  $R_{n-1} < |a_n| < R_n$ . We will refer to these sequences as the *parameters* of the construction. Fixing these sequences, the sets

$$\begin{aligned} D_n &:= \{|z| < R_n\} \subset \mathbb{C} \\ C_n^v &:= D_n \times \mathbb{C} \\ C_n^h &:= \mathbb{C} \times D_n \\ C_n &:= C_n^v \cup C_n^h \\ B_n &:= \overline{C_n^v \cap C_n^h} = \overline{D_n} \times \overline{D_n} \end{aligned}$$

and the rational shears  $f_n : \mathbb{C}^2 \rightarrow \overline{\mathbb{C}^2}$ ,

$$f_n(x, y) := \begin{cases} (x, y + g_n(x)) & \text{for odd } n \\ (x + g_n(y), y) & \text{for even } n \end{cases},$$

## II. Explanation of Orevkov's Paper on p.h.e. $\overline{\mathbb{C}} \setminus C \hookrightarrow \mathbb{C}^2$ and on the size of $C$ .

where

$$g_n(z) := \frac{\epsilon_n}{z - a_n} ,$$

are defined. Finally define the sequence of continuous mappings  $\gamma_n : \overline{\mathbb{C}} \rightarrow \overline{\mathbb{C}^2}$  as

$$\begin{aligned} \gamma_n(z) &:= f_n \circ \dots \circ f_1(z, 0), \quad z \neq \infty \\ \gamma_n(\infty) &:= \infty_2 \end{aligned}$$

and consequently the sets

$$A_n := \gamma_n(\overline{\mathbb{C}}) = f_n \circ \dots \circ f_1(\mathbb{C} \times \{0\}) \cup \{\infty_2\} .$$

The construction of the proper holomorphic embedding  $\gamma$  and of the Cantor set  $C$  in Theorem II.1.1 relies on the sets and functions just defined: we will see that a suitable choice of the sequences  $\{\epsilon_n\}_n$ ,  $\{R_n\}_n$  and  $\{a_n\}_n$  allows to obtain  $C$  as the closure of the set  $\bigcup_n \gamma_n^{-1}(\infty_2)$  and the limit  $\lim_n \gamma_n$  to be a proper holomorphic embedding  $\gamma$  on  $\overline{\mathbb{C}} \setminus C$  into  $\mathbb{C}^2$ ; in particular the faster  $\{R_n\}_n$  diverges, the smaller the size of  $C$  so that it can be constructed to have zero Hausdorff dimension.

In the rest of this paper,  $\{k_n\}_{n \geq 0}$  will denote the Fibonacci sequence, that is the sequence of natural numbers defined by  $k_0 = k_1 = 1$  and

$$k_{n+2} = k_{n+1} + k_n, \quad \forall n \geq 0 .$$

The following remark gathers some known facts about the number of solutions of rational equations (see 4.23–4.25, [4]), which will be fundamental in the proof of the subsequent Lemma.

**Remark II.2.1.** Let  $r : \overline{\mathbb{C}} \rightarrow \overline{\mathbb{C}}$  be a rational function. There exists a natural number  $k$ , called *degree* of  $r$  and denoted as  $\deg r$ , such that for every  $a \in \overline{\mathbb{C}}$ , the equation  $r(z) = a$  has  $k$  solutions counted with their multiplicity, which are multiple only for a finite number of  $a \in \overline{\mathbb{C}}$ .

- (1) In particular, up to a slightly small perturbation of  $a$ , the zeros of  $r(z) - a$  are simple.
- (2) The zeros of  $r$  are as many as the poles, counted with their multiplicity, therefore the poles of  $r$  are all simple if and only if they are  $k$  distinct.
- (3) Let  $t$  be another rational function and  $w \in \overline{\mathbb{C}}$  fixed. Up to perturb  $a \in \mathbb{C}$  arbitrarily little, the equations  $t = w$  and  $r - a = 0$  have no common solutions, as it is easily seen observing that  $r^{-1}(a) \cap r^{-1}(a') = \emptyset$  for any  $a' \neq a$ .

**Lemma II.2.1.** Define a sequence of rational functions recursively:

$$\begin{aligned} r_0(z) &:= z \\ r_1(z) &:= g_1(z) \end{aligned}$$

$$r_{n+2}(z) := r_n(z) + g_{n+2}(r_{n+1}(z)) = r_n(z) + \frac{\epsilon_{n+2}}{r_{n+1}(z) - a_{n+2}}, \quad n \geq 0. \quad (\text{II.2.1})$$

Then, up to perturbing the elements of the sequence  $\{a_n\}_n \subset \mathbb{C}$  arbitrarily little, the following hold:

- (i) for every  $n \geq 0$ ,  $r_n$  and  $r_{n+1}$  do not share any pole and  $r_n^{-1}(\infty) \subsetneq r_{n+2}^{-1}(\infty)$ ; in particular  $\infty$  is a pole only for  $r_n$  with  $n$  even;
- (ii) the poles of  $r_n$  are  $k_n$  and all simple; therefore the equation

$$r_n(z) = a_{n+1}$$

has exactly  $k_n$  solutions, all distinct.

*Proof.* We see that  $r_n^{-1}(\infty) \cap r_{n+1}^{-1}(\infty) = \emptyset$  for all  $n \geq 0$  by induction on  $n$ . The base case follows immediately. Assume the claim true for  $n$ ; if by contradiction  $\exists w \in r_{n+1}^{-1}(\infty) \cap r_{n+2}^{-1}(\infty)$ , then by (II.2.1), it must be  $r_n(w) = \infty$ , so  $r_n^{-1}(\infty) \cap r_{n+1}^{-1}(\infty) \neq \emptyset$ , against inductive hypothesis.

The inclusion  $r_n^{-1}(\infty) \subsetneq r_{n+2}^{-1}(\infty)$  for all  $n \geq 0$  follows by looking at (II.2.1) as from (3) in Remark II.2.1, up to slightly perturbing  $a_{n+2}$ , one gets  $\{r_n = \infty\} \cap \{r_{n+1} - a_{n+2} = 0\} = \emptyset$ .

Clearly  $\infty$  is a pole for  $r_0$ , thus what said so far ensures that  $\infty$  is a pole for  $r_{2k}$  and not a pole for  $r_{2k+1}$ , for all  $k \geq 0$ . So (i) is proved.

Let us prove (ii) by induction on  $n$ . This is trivial for  $r_0$  and  $r_1$ . Assume the claim true up to  $n + 1$ . As observed in (3) in Remark II.2.1, up to a slightly small perturbation of  $a_{n+2}$ , the sets  $\{r_n = \infty\}$  and  $\{r_{n+1} - a_{n+2} = 0\}$  have empty intersection. Hence, by (II.2.1), the poles of  $r_{n+2}$  are precisely the poles of  $r_n$  (which are  $k_n$  and simple by inductive hypothesis) and the poles of  $\frac{\epsilon_{n+2}}{r_{n+1} - a_{n+2}}$  (which are  $k_{n+1}$  and simple as the zeros of the denominator are, up to a slightly small perturbation of  $a_{n+2}$  as pointed out in (1) in Remark II.2.1, in fact  $\deg r_{n+1} = k_{n+1}$  by inductive hypothesis). Therefore the poles of  $r_{n+2}$  are  $k_n + k_{n+1} = k_{n+2}$  and all simple, hence by (2) in Remark II.2.1 one has  $\deg r_{n+2} = k_{n+2}$ . So up to a slightly small perturbation of  $a_{n+3}$ , (1) guarantees that the equation  $r_{n+2}(z) = a_{n+3}$  has  $k_{n+2}$  solutions, all distinct. So (ii) is proved.  $\blacksquare$

We will assume from now on to choose  $R_n$  large enough, so that  $r_n(z) = a$  has  $k_n$  distinct solutions for every  $|a| \geq R_n$ . The following Lemma is used in Section II.3.3:

**Lemma II.2.2.** Let  $r$  be a rational function with a finite number of poles  $r^{-1}(\infty) = \{w_1, \dots, w_N\} \subset \overline{\mathbb{C}}$ , all simple. Assume that  $r(z) = a$  has  $k$  distinct solutions for every  $|a| \geq T_0$ , for some  $T_0$ . Then for any  $T \geq T_0$  large enough, the set

$$\Omega(r, T) := \{z \in \overline{\mathbb{C}} \setminus r^{-1}(\infty) : |r(z)| > T\}$$

is the disjoint union of  $k$  connected sets, each of which homeomorphic to the punctured open disk  $\{|z| < 1\} \setminus \{0\}$  and containing exactly one solution of

## II. Explanation of Orevkov's Paper on p.h.e. $\overline{\mathbb{C}} \setminus C \hookrightarrow \mathbb{C}^2$ and on the size of $C$ .

$r(z) = a$ , for every  $|a| > T$ . Moreover, such  $k$  sets can be taken to have arbitrarily small diameters, provided that  $T$  is large enough.

*Proof.* Assume the poles of  $r$  are all distinct; being them simple one has that  $N = k$ . We express  $r$  on a neighborhood  $V_j \subset \overline{\mathbb{C}}$  of a pole  $w_j$  as  $r(z) = h_j(z)(z - w_j)^{-1}$  or  $r(z) = h_j(z)z$ , accordingly with  $w_j$  to be different or equal to  $\infty$ , where  $h_j \in \mathcal{O}(V_j)$  is bounded and never vanishing. Let  $a \in \mathbb{C}$  with  $|a| > T$ . The larger  $T$ , the closer  $z$  has to be to the poles for  $r(z) = a$  to be solved. In particular, up to enlarging  $T$  we may assume that the  $V_j$  are pairwise disjoint, therefore  $X_j := \{z \in V_j \setminus \{w_j\} : |r(z)| > T\}$  are pairwise disjoint and connected. In particular for each  $\epsilon > 0$ , there exists  $T > 0$  sufficiently large such that the diameter of  $X_j$  is less than  $\epsilon$ . Moreover the union of the  $X_j$  is  $\Omega(r, T)$  and each  $X_j$  contains precisely one of the  $k$  solutions of the equation  $r(z) = a$  for any  $|a| > T$ , for  $T$  sufficiently large: if not, there must be one of them, say  $X_1$ , such that for any  $T$  sufficiently large it contains at least two solutions of  $r(z) = a$  for some  $|a| > T$ , leading  $w_1$  to be a multiple pole for  $r$ . ■

Finally, it is easily seen by induction that

$$\gamma_n = \begin{cases} (r_{n-1}, r_n) & \text{for odd } n \\ (r_n, r_{n-1}) & \text{for even } n \end{cases}, \quad (\text{II.2.2})$$

in fact the case  $n = 1$  is immediate and for  $n$  even, for example

$$\gamma_{n+1} = f_{n+1}(\gamma_n) = f_{n+1}(r_n, r_{n-1}) = (r_n, r_{n-1} + g_{n+1}(r_n)) = (r_n, r_{n+1}).$$

### II.3 General Properties

This section is devoted to the description of the objects introduced in Section II.2.3, which are the ingredients of the construction of the proper holomorphic embedding  $\gamma: \overline{\mathbb{C}} \setminus C \hookrightarrow \mathbb{C}^2$  presented in [5]. In particular we will focus on the choice of the three sequences  $\{\epsilon_n\}_{n \geq 1}$ ,  $\{R_n\}_{n \geq 0}$ ,  $\{a_n\}_{n \geq 1}$  upon which the whole construction is based.

#### II.3.1 On the set $\gamma_n^{-1}(\infty_2)$ and the sequence $\{R_n\}_{n \geq 0}$ . Definition and properties of vertical and horizontal components of $A_n$

It follows from (II.2.2) and (i) in Lemma II.2.1 that

$$\gamma_n^{-1}(\infty_2) \subsetneq \gamma_{n+1}^{-1}(\infty_2) \quad (\text{II.3.1})$$

holds true for all  $n \geq 0$ ; moreover  $\infty \in \gamma_n^{-1}(\infty_2)$  for every  $n \geq 0$ , so we can write

$$\gamma_n^{-1}(\infty_2) = \{\infty\} \cup \{t_j^{(n)}\}_{j=1}^{b_n-1} \quad (\text{II.3.2})$$

for some  $t_j^{(n)} \in \mathbb{C}$ , where  $b_n = k_{n+1}$ . Observe that  $A_n = \gamma_n(\overline{\mathbb{C}}) \subset \overline{\mathbb{C}^2}$  is compact with  $\infty_2 \in A_n$ ; moreover  $A_n \setminus \{\infty_2\} = \gamma_n(\mathbb{C})$  is a connected complex curve in  $\mathbb{C}^2$ .

**Proposition II.3.1.** Assume  $\epsilon_1, \dots, \epsilon_n$ ,  $a_1, \dots, a_n$  and  $R_1, \dots, R_{n-1}$  fixed,  $|a_{j+1}| > R_j > |a_j|$  for  $j = 1, \dots, n-1$ . Then there exist  $R_n > |a_n|$  such that what follows is true:

- (a)  $A_n \setminus (B_n \cup \{\infty_2\})$  has  $b_n$  disjoint components, and
- (b)  $A_n \subset C_n$ .

We refer to the components of  $A_n \setminus (B_n \cup \{\infty_2\})$  contained in  $C_n^h$  (resp. in  $C_n^v$ ) as the *horizontal* (resp. *vertical*) components of  $A_n$ . Correspondingly we split  $b_n$  as  $h_n + v_n$ . It turns out that

$$\begin{cases} v_n = k_{n-1} \\ h_n = k_n \end{cases} \quad (n \text{ even}), \quad \begin{cases} v_n = k_n \\ h_n = k_{n-1} \end{cases} \quad (n \text{ odd}). \quad (\text{II.3.3})$$

*Proof.* The components of  $A_n \setminus (B_n \cup \{\infty_2\})$  are the image, via  $\gamma_n$  of punctured neighborhoods in  $\overline{\mathbb{C}}$  of the points (II.3.2) (which are defined once  $a_n, \epsilon_n, a_j, \epsilon_j, R_j$ ,  $j \leq n-1$  are fixed). So, by Lemma II.2.2, the bigger  $R_n$ , the smaller these neighborhoods; hence these  $b_n$  neighborhoods are disjoint for  $R_n$  large enough, achieving (a), as  $\gamma_n$  is injective on  $\overline{\mathbb{C}} \setminus \gamma_n^{-1}(\infty_2)$  (see Section II.3.2). Looking at (II.2.2) it is clear that unbounded components of  $A_n = \gamma_n(\overline{\mathbb{C}})$  are due to the poles of  $r_n$  and  $r_{n-1}$ ; being  $r_n^{-1}(\infty) \cap r_{n-1}^{-1}(\infty) = \emptyset$  by (i) in Lemma II.2.1, it follows that when one component of  $A_n$  becomes unbounded, the other remains bounded. Hence, for  $R_n$  large enough, we get (b). Finally, (II.3.3) follows directly from Lemma II.2.1, once we express  $\gamma_n$  as in (II.2.2). ■

### II.3.2 Definition of certain sets $\Delta_n$ , $K_n$ and some properties of the mappings $\gamma_n$

Let us then define

$$\Delta_n := \gamma_n^{-1}(A_n \setminus B_n).$$

Given  $U \subset \overline{\mathbb{C}^2}$  open neighborhood of  $\infty_2$ ,  $\gamma_n^{-1}(U)$  is a neighborhood of  $\gamma_n^{-1}(\infty_2)$ ; this last set has  $b_n$  points, therefore Lemma II.2.2 guarantees that for  $U$  sufficiently small (meaning the points of  $U$  are sufficiently close to  $\infty_2$  with respect to the spherical measure introduced in Section II.2.2; equivalently  $U$  is contained in the complement of a closed ball in  $\mathbb{C}^2$  of sufficiently large radius),  $\gamma_n^{-1}(U)$  is the disjoint union of  $b_n$  open neighborhoods  $U_j^{(n)}$  of the points (II.3.2). In particular  $\forall \epsilon > 0 \exists U \subset \overline{\mathbb{C}^2}$  open neighborhood of  $\infty_2$  such that  $|U_j^{(n)}| < \epsilon$ . Since

$$\Delta_n = \gamma_n^{-1}(\overline{\mathbb{C}^2} \setminus B_n) \quad (\text{II.3.4})$$

## II. Explanation of Orevkov's Paper on p.h.e. $\overline{\mathbb{C}} \setminus C \hookrightarrow \mathbb{C}^2$ and on the size of $C$ .

(straightforward with double inclusion), we can always suppose to have  $R_n$  large enough such that

$$\Delta_n = \bigcup_{j=1}^{b_n} U_j^{(n)}$$

is a disjoint union. In particular for every  $n$ , there exists  $R_n$  so large that

$$|U_j^{(n)}| < \frac{1}{b_n^n}, \quad \forall j = 1, \dots, b_n. \quad (\text{II.3.5})$$

Let us now check that

$$\gamma_n: \overline{\mathbb{C}} \setminus \gamma_n^{-1}(\infty_2) \rightarrow \mathbb{C}^2$$

is a holomorphic embedding by induction on  $n$ . As  $\gamma_1: \overline{\mathbb{C}} \setminus \{a_1, \infty\} \rightarrow \mathbb{C}^2$ ,  $\gamma_1(z) = (z, \frac{\epsilon_1}{z-a_1})$ , the base case trivially follows. Assume the claim true for  $n$ . Then  $\gamma_{n+1} = f_{n+1} \circ \gamma_n$  is a holomorphic embedding in the intersection of the subset where  $\gamma_n$  is such (that is  $\overline{\mathbb{C}} \setminus \gamma_n^{-1}(\infty_2)$ ) and  $\{z \in \overline{\mathbb{C}} : \gamma_n(z) \notin f_{n+1}^{-1}(\infty_2)\} = \overline{\mathbb{C}} \setminus \gamma_{n+1}^{-1}(\infty_2)$ . Such an intersection is

$$\overline{\mathbb{C}} \setminus \gamma_n^{-1}(\infty_2) \cap \overline{\mathbb{C}} \setminus \gamma_{n+1}^{-1}(\infty_2) = \overline{\mathbb{C}} \setminus (\gamma_n^{-1}(\infty_2) \cup \gamma_{n+1}^{-1}(\infty_2)) = \overline{\mathbb{C}} \setminus \gamma_{n+1}^{-1}(\infty_2)$$

and we are done (the last equality follows from (II.3.1)).

Define then the following compacts in  $\overline{\mathbb{C}}$ :

$$\begin{aligned} K_n &:= \overline{\mathbb{C}} \setminus \Delta_n = \gamma_n^{-1}(A_n \cap B_n) = \gamma_n^{-1}(B_n), \quad n \geq 1 \\ K_0 &:= \emptyset. \end{aligned}$$

It is straightforward to see that  $K_n \subset \overline{\mathbb{C}} \setminus \gamma_n^{-1}(\infty_2)$ . It will follow from Proposition II.3.2 that  $K_n \subsetneq K_{n+1}^\circ$ ; therefore there exists  $\delta_n > 0$  such that if  $h: \overline{\mathbb{C}} \setminus \gamma_n^{-1}(\infty_2) \rightarrow \mathbb{C}^2$  is holomorphic with

$$\|\gamma_n - h\|_{K_n} < \delta_n,$$

then  $h$  is an embedding on  $K_{n-1}$ . Set  $\delta_0 := 1/4$  and assume without loss of generality  $\{\delta_n\}_{n \geq 0}$  to be strictly decreasing.

### II.3.3 From $\Delta_n$ to $\Delta_{n+1}$ : definition of pair of pants

We will prove that for a suitable choice of the parameters, the sequences  $\{\Delta_n\}_n$  and  $\{K_n\}_n$  are strictly decreasing and increasing respectively; in particular, we will focus on the topological behavior of the sequence  $\{\Delta_n\}_n$  when passing from step  $n$  to step  $n+1$ .

**Proposition II.3.2.** Assume  $\epsilon_1, \dots, \epsilon_{n+1}$ ,  $a_1, \dots, a_{n+1}$  and  $R_1, \dots, R_n$  fixed, with  $|a_{j+1}| > R_j > |a_j|$  for  $j = 1, \dots, n$  and rename the  $U_j^{(n)}$  to highlight which one comes from horizontal or vertical components of  $A_n \setminus B_n$  as

$$\Delta_n = \bigcup_{j=1}^{h_n} H_j^{(n)} \cup \bigcup_{j=1}^{v_n} V_j^{(n)} = H_n \cup V_n.$$

Then, for any  $R_{n+1} > |a_{n+1}|$  sufficiently large, we have that

$$\Delta_{n+1} \subsetneq \Delta_n, \quad (\text{II.3.6})$$

$$K_n \subsetneq K_{n+1}^\circ, \quad (\text{II.3.7})$$

and passing from  $\Delta_n$  to  $\Delta_{n+1}$ , for  $n$  even,  $V_j^{(n)}$  shrinks to  $V_j^{(n+1)}$ ,  $j = 1, \dots, v_n$ , while  $H_j^{(n)}$  splits itself into  $H_j^{(n+1)}$  and  $V_{v_n+j}^{(n+1)}$ ,  $j = 1, \dots, h_n$ , creating the so called *pair of pants*. For  $n$  odd, a similar conclusion holds, swapping vertical and horizontal components.

*Proof.* Let  $n$  be even. From (II.2.1) and (II.2.2) we have

$$\begin{aligned} \gamma_n &= (r_n, r_{n-1}), \text{ and} \\ \gamma_{n+1} &= (r_n, r_{n+1}) = (r_n, r_{n-1}) + \left(0, \frac{\epsilon_{n+1}}{r_n - a_{n+1}}\right), \end{aligned}$$

from which

$$\begin{aligned} H_n &= \{|r_n| > R_n\}, \\ V_n &= \{|r_{n-1}| > R_n\}, \text{ and} \\ H_{n+1} &= \{|r_n| > R_{n+1}\}. \end{aligned}$$

Moreover, up to slightly move  $a_{n+1}$ , (3) in Remark II.2.1 guarantees that  $r_{n-1} = \infty$  and  $r_n = a_{n+1}$  have disjoint solutions, so for  $R_{n+1}$  large enough we have

$$V_{n+1} = \{|r_{n-1}| > R_{n+1}\} \cup \{|r_n - a_{n+1}| < \epsilon_{n+1}/R_{n+1}\} =: V'_{n+1} \cup V''_{n+1},$$

with  $V'_{n+1} \cap V''_{n+1} = \emptyset$ . Since  $|a_{n+1}| > R_n$ , it follows that  $R_{n+1}$  sufficiently large implies  $V''_{n+1} \subseteq H_n$ ; the inclusions  $V'_{n+1} \subseteq V_n$  and  $H_{n+1} \subseteq H_n$  are trivial, so we get (II.3.6) and (II.3.7), where the strict inclusions follow from (II.3.4) and Lemma II.2.2, which guarantees the  $U_j^{(n+1)}$  shrunk arbitrarily little around the point they are neighborhood of, provided  $R_{n+1}$  to be large enough. Taking  $R_{n+1}$  large enough guarantees moreover that  $V''_{n+1}$  has  $k_n$  components (as the equation  $r_n = a_{n+1}$  has  $k_n$  simple solutions by (ii) in Lemma II.2.1) and  $H_{n+1} \cap V''_{n+1} = \emptyset$  (as  $R_{n+1} > |a_{n+1}|$ ). Now  $H_n$  has  $k_n$  components, each of which splits into a component of  $H_{n+1}$  and a component of  $V''_{n+1}$ . So the statement on the pair of pants follows.

For  $n$  odd the argument is the same, just switching the role of vertical and horizontal objects. ■

### II.3.4 The sequence $\{\epsilon_n\}_{n \geq 1}$

Describing constraints for  $R_n$  in Sections II.3.1, II.3.2 and II.3.3, we assumed to have already fixed  $a_1, \dots, a_n$ ,  $R_1, \dots, R_{n-1}$ , and in particular  $\epsilon_1, \dots, \epsilon_n$ . Let us now see how to choose the  $\epsilon_n$ .

## II. Explanation of Orevkov's Paper on p.h.e. $\overline{\mathbb{C}} \setminus C \hookrightarrow \mathbb{C}^2$ and on the size of $C$ .

**Proposition II.3.3.** There are sequences  $\{a_n\}_n$ ,  $\{\epsilon_n\}_n$  and  $\{R_n\}_n$  such that

$$\|f_n - \text{Id}\|_{\overline{B_j(\frac{1}{2})}} < \delta_j \cdot \frac{1}{2^{j+n}}, \quad 0 \leq j < n \quad (\text{II.3.8})$$

*Proof.* Let us construct the three sequences inductively on  $n$ . Case  $n = 1$ ; take any  $a_1 \in \mathbb{C}$  such that  $|a_1| > 1/2$ , define  $B_0$  as the origin  $(0, 0) \in \mathbb{C}^2$ . Then there exists  $\epsilon_1 > 0$  such that  $\|f_1 - \text{Id}\|_{\overline{B_0(\frac{1}{2})}} < \delta_0 \cdot \frac{1}{2^{1+0}}$  (note that  $B_0(\frac{1}{2})$  is the  $1/2$ -ball in  $\mathbb{C}^2$ ). Assume (II.3.8) true for  $n$ . In particular  $a_i, \epsilon_i, R_i$ ,  $i \leq n-1$  and  $a_n, \epsilon_n$  are defined, so we can take  $R_n$  to satisfy all the constraints discussed in the previous sections; in particular in Lemma II.2.1, Proposition II.3.1 and II.3.2, condition (II.3.5). So we can fix  $a_{n+1}$  such that  $|a_{n+1}| > \sqrt{2}R_n + 1/2$ . Define then  $\epsilon_{n+1} > 0$  such that  $\|f_{n+1} - \text{Id}\|_{\overline{B_n(\frac{1}{2})}} < \delta_n \cdot \frac{1}{2^{2n+1}}$ . Then (II.3.8) follows in the case  $n+1$ .  $\blacksquare$

With such a choice of parameters, we achieve

$$\|f_n \circ \cdots \circ f_j - \text{Id}\|_{B_{j-1}} < \frac{1}{2}, \quad 1 \leq j \leq n. \quad (\text{II.3.9})$$

The case  $n = 1$  follows from the base case of (II.3.8). Assume (II.3.9) true for  $n$  and let us see it holds for  $n+1$ :

$$\begin{aligned} \|f_{n+1} \circ \cdots \circ f_j - \text{Id}\|_{B_{j-1}} &\leq \|f_{n+1} - \text{Id}\|_{f_n \circ \cdots \circ f_j(B_{j-1})} \\ &\quad + \|f_n - \text{Id}\|_{f_{n-1} \circ \cdots \circ f_j(B_{j-1})} + \cdots + \|f_j - \text{Id}\|_{B_{j-1}} \\ &\leq \|f_{n+1} - \text{Id}\|_{\overline{B_{j-1}(\frac{1}{2})}} \\ &\quad + \|f_n - \text{Id}\|_{\overline{B_{j-1}(\frac{1}{2})}} + \cdots + \|f_j - \text{Id}\|_{\overline{B_{j-1}(\frac{1}{2})}} \end{aligned} \quad (\text{II.3.10})$$

$$\leq \frac{\delta_{j-1}}{2^{n+1+j-1}} + \cdots + \frac{\delta_{j-1}}{2^{j+j-1}} \quad (\text{II.3.11})$$

$$= \delta_{j-1} \cdot \sum_{i=2^{j-1}}^{n+j} \frac{1}{2^i} < \frac{1}{2}.$$

Hence (II.3.9) holds for any  $1 \leq j \leq n+1$  as promised. Just observe that (II.3.10) follows since

$$f_n \circ \cdots \circ f_j(B_{j-1}) \subset B_{j-1}(1/2) \subset \overline{B_{j-1}(1/2)}$$

holds true by inductive hypothesis for any  $1 \leq j \leq n$  and (II.3.11) follows from Proposition II.3.3.

**Remark II.3.1.** Since the three sequences underlying the whole construction depend one on the other, it is important to highlight the order with which their elements have to be taken. We have seen in Sections II.3.1, II.3.2 and II.3.3 that in order to define  $R_n$ , the parameters

$$a_j, \epsilon_j, R_j, \quad j \leq n-1, \quad \text{and} \quad a_n, \epsilon_n$$



need to be already fixed; then we immediately fix  $a_{n+1}$  such that  $|a_{n+1}| > \sqrt{2}R_n + 1/2$ . Similarly, in Section II.3.4 it turned out that in order to define  $\epsilon_n$ ,

$$a_j, \epsilon_j, R_j, \quad j \leq n-1, \quad \text{and} \quad a_n$$

need to be previously fixed; in particular, to choose  $\epsilon_1$ , some  $a_1 \in \mathbb{C}$ ,  $|a_1| > 1/2$ , has to be fixed. Namely, the order to follow for choosing the sequences of parameters to perform our construction is

$$a_1, \epsilon_1, R_1, a_2, \epsilon_2, R_2, \dots$$

## II.4 The Cantor set $C$ and the proper holomorphic embedding $\gamma$

### II.4.1 Definition of the Cantor set $C$

We are ready to define the object of our interest:

$$C := \bigcap_n \Delta_n \subset \overline{\mathbb{C}}. \quad (\text{II.4.1})$$

Let us see  $C$  is indeed a Cantor set by proving that it fulfills all the properties (i) $_C$ –(v) $_C$  stated in Section II.2. Being  $\overline{\mathbb{C}}$  metrizable, we get (v) $_C$ . Since for every  $n \in \mathbb{N}$  one has

$$\gamma_n^{-1}(\infty_2) \subset \gamma_{n+j}^{-1}(\infty_2) \subset \Delta_{n+j} \quad \forall j \in \mathbb{N},$$

it follows that

$$\bigcup_{n \geq 1} \gamma_n^{-1}(\infty_2) \subseteq C, \quad (\text{II.4.2})$$

from which  $C \neq \emptyset$  and thus we get (i) $_C$ . From the above construction, every component of  $\Delta_n$  eventually shrinks to a point (see e.g., (II.3.5)) and since the role of vertical and horizontal components switches at each step, it follows that  $\forall n \in \mathbb{N}$ ,  $\forall j \in \{1, \dots, b_n\}$  there will be infinitely many pair of pants inside each  $U_j^{(n)}$ , so there will be no isolated point. Moreover exploiting (II.3.7) one gets

$$(\overline{\mathbb{C}} \setminus C)^\circ = \left( \bigcup_n K_n \right)^\circ = \bigcup_n K_n^\circ = \bigcup_n K_n = \overline{\mathbb{C}} \setminus C \quad (\text{II.4.3})$$

from which  $C$  is closed, and thus (ii) $_C$  is verified. Being  $C$  closed and  $\overline{\mathbb{C}}$  compact, it follows that  $C$  is compact itself, so we have (iii) $_C$ . Finally, if any connected component of  $C$  had more than one point, then (II.3.5) would fail for  $n$  big enough. So  $C$  is totally disconnected, that is we have (iv) $_C$  and so we have proved  $C$  is indeed a Cantor set.

## II.4.2 Definition of the proper holomorphic embedding

$$\gamma: \overline{\mathbb{C}} \setminus C \hookrightarrow \mathbb{C}^2$$

From time to time we will use the following fact.

**Remark II.4.1.** If  $\{K_n\}_n$  is a normal exhaustion for a domain  $\Omega \subseteq \overline{\mathbb{C}}$  and  $\{z_n\}_n \subset \Omega$  converges to  $z_0 \in \partial\Omega$ , then we can suppose without loss of generality that  $z_n \in K_n \setminus K_{n-1}$ .

Consider  $\gamma_n$  as a mapping  $K_{n+1}^\circ \rightarrow \mathbb{C}^2$ ; thus the natural domain to define a limit mapping is  $\bigcup_{n \geq 1} K_n^\circ$  which equals  $\overline{\mathbb{C}} \setminus C$  from (II.4.3). For every  $n \geq k$  one has

$$\begin{aligned} \|\gamma_{n+1} - \gamma_n\|_{K_k} &= \|(f_{n+1} - \text{Id}) \circ \gamma_n\|_{\gamma_k^{-1}(B_k)} \\ &= \|(f_{n+1} - \text{Id}) \circ (f_n \circ \dots \circ f_{k+1} \circ \gamma_k)\|_{\gamma_k^{-1}(B_k)} \\ &\leq \|f_{n+1} - \text{Id}\|_{f_n \circ \dots \circ f_{k+1}(B_k)} \\ &\leq \|f_{n+1} - \text{Id}\|_{\overline{B_k(\frac{1}{2})}} \quad \text{by (II.3.9)} \\ &< \delta_k \cdot \frac{1}{2^{n+1+k}} \quad \text{by (II.3.8) ,} \end{aligned}$$

from which it follows that

$$\sum_{n \geq k} \|\gamma_{n+1} - \gamma_n\|_{K_k} < \sum_{n \geq k} \delta_k \cdot \frac{1}{2^{n+1+k}} < +\infty$$

for every fixed  $k \geq 1$ ; hence  $\{\gamma_n\}_n$  compactly converges to a holomorphic mapping

$$\gamma: \overline{\mathbb{C}} \setminus C \rightarrow \mathbb{C}^2,$$

since  $\{K_n\}_n$  is a normal exhaustion for the domain. It is now easily seen that  $\gamma$  is a local embedding (namely: an embedding on every  $K_{k-1}$ ) since for every  $k \geq 1$  one has

$$\|\gamma - \gamma_k\|_{K_k} \leq \sum_{n \geq k} \|\gamma_{n+1} - \gamma_n\|_{K_k} < \sum_{n \geq k} \delta_k \cdot \frac{1}{2^{n+1+k}} < \delta_k.$$

Finally, global embedding property of  $\gamma$  will follow automatically once properness is proved, achieving thus that  $\gamma$  is a proper holomorphic embedding from the complement of a Cantor set into  $\mathbb{C}^2$ , as wanted. Consider then  $\{z_n\}_n \subset \overline{\mathbb{C}} \setminus C = \bigcup_n K_n$  such that  $z_n \rightarrow \partial(\overline{\mathbb{C}} \setminus C) = \partial C$ . By Remark II.4.1 we may assume  $z_n \in K_n \setminus K_{n-1} \subseteq \Delta_{n-1}$ , from which  $|\gamma_{n-1}(z_n)| \rightarrow +\infty$  as  $n \rightarrow +\infty$ . Nevertheless this does not imply that  $|\gamma(z_n)| \rightarrow +\infty$ , as it just claims the divergence of  $\{\gamma_{n-1}(z_n)\}_n$  and no information on how the sequence  $\{\gamma_n\}_n$  acts on the elements of  $\{z_n\}_n$  is given. In particular, we need to achieve  $z_n$  to be set outside some  $R_n$ -ball (say  $B_{n-3}$ ) by a whole tail of  $\{\gamma_n\}_n$  (say  $\gamma_{n+j}$  for every  $j \in \mathbb{N}$ ). In this way the possibility for  $z_n$  to come back inside a ball is prevented

and we obtain properness of the limit function  $\gamma$ . Hence properness will be achieved if, for instance, the following condition is satisfied:

$$\gamma_{n+j}(K_n \setminus K_{n-1}) \subset \overline{\mathbb{C}}^2 \setminus B_{n-3} \quad \forall n \geq 3, j \in \mathbb{N}. \quad (\text{II.4.4})$$

As we assumed  $z_n \in K_n \setminus K_{n-1} \quad \forall n$ , one has

$$\gamma(z_n) = \lim_j \gamma_{j+n}(z_n) \stackrel{(\text{II.4.4})}{\in} \overline{\mathbb{C}}^2 \setminus B_{n-3},$$

which implies  $|\gamma(z_n)| \geq R_{n-3}$  and thus  $\lim_n |\gamma(z_n)| = +\infty$ , as wanted. Notice that

$$\gamma_n(\gamma_n^{-1}(B_n)) = A_n \cap B_n \quad (\text{II.4.5})$$

holds true for every  $n \in \mathbb{N}$  and this allows to prove that (II.4.4) holds, in fact

$$\begin{aligned} \gamma_{n+j}(K_n \setminus K_{n-1}) &= f_{n+j} \circ \cdots \circ f_{n+1} \circ \gamma_n(\gamma_n^{-1}(B_n) \setminus \gamma_n^{-1}(B_{n-1})) \quad \text{by (II.4.5)} \\ &= f_{n+j} \circ \cdots \circ f_{n+1}((A_n \cap B_n) \setminus (f_n(B_{n-1}) \cap A_n)) \\ &\subseteq f_{n+j} \circ \cdots \circ f_{n+1}(B_n \setminus f_n(B_{n-1})) \\ &\subseteq f_{n+j} \circ \cdots \circ f_{n+1}(B_n \setminus B_{n-2}) \quad \text{by (II.3.9)} \\ &\subseteq B_{n+1} \setminus B_{n-3} \quad \text{by (II.3.9)} \\ &\subseteq \overline{\mathbb{C}}^2 \setminus B_{n-3}, \end{aligned}$$

as promised.

### II.4.3 On the Hausdorff dimension of $C$ and proof of Theorem II.1.1

Fix  $\epsilon > 0$ ; the *Hausdorff*  $\epsilon$ -measure of  $C$  is, by definition

$$\mathcal{H}^\epsilon(C) := \liminf_{r \rightarrow 0} \left\{ \inf \left\{ \sum_i |T_i|^\epsilon : C \subseteq \bigcup_i T_i, |T_i| < r \right\} \right\}.$$

Since  $C \subset \Delta_n = \bigcup_{j=1}^{b_n} U_j^{(n)}$  holds true for any  $n$  and since for every  $r > 0$  there exists  $N_r$  such that  $1/b_n^n < r$  for all  $n \geq N_r$ , exploiting (II.3.5), it trivially follows that

$$\begin{aligned} \inf \left\{ \sum_i |T_i|^\epsilon : C \subseteq \bigcup_i T_i, |T_i| < r \right\} &\leq \inf \left\{ \sum_{j=1}^{b_n} |U_j^{(n)}|^\epsilon : n \geq N_r \right\} \\ &\leq \inf \left\{ \frac{1}{b_n^{\epsilon n - 1}} : n \geq N_r \right\} = 0, \end{aligned}$$

hence  $\mathcal{H}^\epsilon(C) = 0$  for any  $\epsilon > 0$  fixed. Therefore, recalling that the *Hausdorff dimension* of  $C$  is defined as

$$\dim_H(C) := \inf \{d \geq 0 : \mathcal{H}^d(C) = 0\},$$

## II. Explanation of Orevkov's Paper on p.h.e. $\overline{\mathbb{C}} \setminus C \hookrightarrow \mathbb{C}^2$ and on the size of $C$ .

we conclude that

$$\dim_H(C) = 0 .$$

This last observation, together with the construction of the proper holomorphic embedding  $\gamma: \overline{\mathbb{C}} \setminus C \hookrightarrow \mathbb{C}^2$  made in Section II.4.2, completes the proof of Theorem II.1.1.

### II.4.4 Two further characterizations of $C$

The original definition of the Cantor set  $C$  given in [5] is

$$C = \gamma^{-1}(\infty_2) . \quad (\text{II.4.6})$$

Actually,  $\gamma$  is not defined on the Cantor set  $C$ , therefore, for this last relation to make sense, we need to extend  $\gamma$  on  $C$ . Being  $C$  a Cantor set in  $\overline{\mathbb{C}}$ , it is the boundary of  $\overline{\mathbb{C}} \setminus C$ , domain on which  $\gamma$  is proper, therefore

$$\lim_{w \rightarrow C, w \in \overline{\mathbb{C}} \setminus C} |\gamma(w)| = \infty ,$$

in fact if  $z \in C$ , being  $C$  a Cantor set,  $z$  is an accumulation point of the complement of  $C$ , which is  $\bigcup_n K_n$ , thus  $z$  can be hit by a sequence  $\{z_n\}_n \subset \bigcup_n K_n$ ; by Remark II.4.1 let  $z_n \in K_n \setminus K_{n-1}$ . On one hand it is clear that  $\lim_j \gamma_{n+j}(z_n) = \gamma(z_n)$  for every  $n$ , on the other hand (II.4.4) implies  $\lim_n \lim_j \gamma_{n+j}(z_n) = \infty_2$ . So we can extend  $\gamma$  on  $C$  by continuity defining  $\gamma \equiv \infty_2$  on  $C$ . Since trivially  $\gamma \neq \infty_2$  on  $\overline{\mathbb{C}} \setminus C$ , we have that defining  $C$  as in (II.4.6) matches with the definition (II.4.1). We conclude by characterizing  $C$  in one last way:

$$C = \text{cl} \left( \bigcup_{n \geq 1} \gamma_n^{-1}(\infty_2) \right) . \quad (\text{II.4.7})$$

Being  $C$  closed, (II.4.2) implies that

$$\text{cl} \left( \bigcup_{n \geq 1} \gamma_n^{-1}(\infty_2) \right) \subseteq C .$$

Consider now the following inductive procedure: at step  $n$  we have  $b_n$  points, the elements of  $\gamma_n^{-1}(\infty_2)$ , and  $b_n$  neighborhoods  $U_j^{(n)}$  of them. At step  $n+1$  we have the old  $b_n$  points and  $k_n$  new points (that is, the elements of  $\gamma_{n+1}^{-1}(\infty_2)$ ) and  $b_{n+1} = b_n + k_n$  neighborhoods  $U_j^{(n+1)}$  of them,  $b_n$  of which are just the  $U_j^{(n)}$  shrunk around the points of  $\gamma_n^{-1}(\infty_2)$  and the  $k_n$  remaining ones are neighborhoods of the new points  $\gamma_{n+1}^{-1}(\infty_2) \setminus \gamma_n^{-1}(\infty_2)$ . Now  $\bigcup_{j=1}^{b_n} U_j^{(n)} = \Delta_n$  forms a decreasing sequence of open sets, whose intersection defines  $C$  and all the  $U_j^{(n)}$  eventually shrink to the point they are neighborhood of, hence  $C$  is the limit of the above inductive procedure. Therefore, given  $w \in C$ , either it appears at

some finite step of the procedure (hence  $w \in \gamma_n^{-1}(\infty_2)$  for some  $n$ ), or it appears in the limit: by construction, the points tend to accumulate (in fact there is no isolated point, see (ii) <sub>$C$</sub> ), that is  $w \in \text{Acc} \left( \bigcup_{n \geq 1} \gamma_n^{-1}(\infty_2) \right) \setminus \left( \bigcup_{n \geq 1} \gamma_n^{-1}(\infty_2) \right)$ .

So  $w \in \text{cl} \left( \bigcup_{n \geq 1} \gamma_n^{-1}(\infty_2) \right)$  and we are done.



# Bibliography

- [1] Brannan, D. A., Esplen, M. F. and Gray, J. J.; Geometry. Cambridge University Press, Second Edition (2012).
- [2] Brouwer, L. E. J.; On the structure of perfect sets of points. *KNAW, Proceedings* **12**, (1909–1910), Amsterdam, 1910, pp. 785–794.
- [3] Di Salvo, G. D., and Wold, E. F.; Proper holomorphic embeddings of complements of large Cantor sets in  $\mathbb{C}^2$ . First article in first author’s Ph.D. thesis, Universitetet i Oslo, 2022.
- [4] Forster, O.; Lectures on Riemann Surfaces. Springer–Verlag, New York Inc. 1981.
- [5] Orevkov, S. Y.; Proper analytic embedding of  $\mathbb{CP}^1$  minus Cantor set into  $\mathbb{C}^2$ . *Uspekhi Mat. Nauk.* **63** (2008), no. 1(379), 155–156 (in Russian) English transl.: Russian Math. Surveys, **63** (2008), 168–169.

## Author address

**G. D. Di Salvo** University of Oslo, Postboks 1337 Blindern, 0316 Oslo, Norway,  
giovandd@math.uio.no





# Families of Proper Holomorphic Embeddings and Carleman–type Theorems with parameters

G. D. Di Salvo, T. Ritter, and E. F. Wold

Published in *The Journal of Geometric Analysis*, January 2023, volume 33, issue 75.

Abstract

We solve the problem of simultaneously embedding properly holomorphically into  $\mathbb{C}^2$  a whole family of  $n$ –connected domains  $\Omega_r \subset \mathbb{P}^1$  such that none of the components of  $\mathbb{P}^1 \setminus \Omega_r$  reduces to a point, by constructing a continuous mapping  $\Xi: \bigcup_r \{r\} \times \Omega_r \rightarrow \mathbb{C}^2$  such that  $\Xi(r, \cdot): \Omega_r \hookrightarrow \mathbb{C}^2$  is a proper holomorphic embedding for every  $r$ . To this aim, a parametric version of both the Andersén–Lempert procedure and Carleman’s Theorem is formulated and proved.

Contents

III.1	Introduction . . . . .	53
III.2	The Setup . . . . .	54
III.3	Carleman approximation with parameters . . . . .	56
III.4	Andersén–Lempert Theory . . . . .	65
III.5	Proof of Theorem III.1.1 . . . . .	69

III.1 Introduction

Existence of proper holomorphic embeddings of Riemann surfaces  $\mathcal{R}$  into 2–dimensional complex manifolds  $X$ , e.g.,  $X = \mathbb{C}^2$ , with prescribed geometrical properties, e.g., being complete, has been an active area of research over the recent years. Various techniques have been developed, but in several cases, positive results have been obtained only at the cost of perturbing the complex structure of  $\mathcal{R}$  (see Černe–Forstnerič [4], Alarcón [1] and Alarcón–López [2]). It

### III. Families of Proper Holomorphic Embeddings and Carleman-type Theorems with parameters

---

can be hoped, however, that if you let  $r$  be a local parameter on the moduli space of Riemann surfaces of a given type, and you perform various constructions continuously with the parameter  $r$  near a given point  $r_0$ , then you will get a perturbation of the complex structure for each given  $r$ , but at least one perturbation will correspond to your initial  $r_0$ . Indeed this is the philosophy behind the embedding results of Globevnik–Stensønes [7]. The purpose of this article is to take a first step towards results of this type that may be generalized to larger classes of Riemann surfaces.

We will consider the following. It is known that any  $n$ -connected domain  $\Omega$  in the Riemann sphere may be mapped univalently onto a domain in the Riemann sphere whose complement consists of  $n$  parallel disjoint slits with a given inclination  $\Theta$  to the real axis. The univalent map achieving this is uniquely determined by  $\Theta$  and the choice of a certain normalization of the Laurent series expansion at a chosen point  $\zeta \in \Omega$  being sent to  $\infty$  (see Goluzin, [8], page 213). Considering a continuous family of  $n$ -connected domains, we obtain a continuously varying family of uniformizing slit-maps.

Let  $C_j \subset \mathbb{C}$  be compact disks and  $I_j \subset \mathbb{R}_{>0}$  be compact intervals,  $j = 1, \dots, n$ . Set  $B_j := C_j \times I_j$  and  $B := B_1 \times \dots \times B_n$ . Let  $r = ((a_1, b_1), \dots, (a_n, b_n))$  denote the coordinates on  $B$ , and letting  $l_{r,j}$  denote the closed straight line segment which is parallel to the real axis with right end-point  $a_j(r)$  and of length  $b_j(r)$ , we assume that  $L_r := \{l_{r,1}, \dots, l_{r,n}\}$  is a set of pairwise disjoint slits, and thus  $\mathbb{P}^1 \setminus L_r$  is an  $n$ -connected domain, none of whose boundary components are isolated points. After possibly having to apply the map  $z \mapsto (z - a_1(r))/b_1(r)$  we may assume that for all  $r$  we have that  $l_{r,1} = [-1, 0] \subset \mathbb{C}$ .

The goal is to prove the following.

**Theorem III.1.1.** In  $B \times \mathbb{P}^1$  set

$$\Omega = (B \times \mathbb{P}^1) \setminus \left( \bigcup_{r \in B} \{r\} \times L_r \right).$$

Then there exists a continuous map  $\Xi: \Omega \rightarrow \mathbb{C}^2$  such that for each  $r \in B$  we have that  $\Xi(r, \cdot): \Omega_r \rightarrow \mathbb{C}^2$  is a proper holomorphic embedding.

## III.2 The Setup

We will now introduce a setup to prove Theorem III.1.1. First, we need the notion of a certain directed family of curves.

Let  $C > 0$  and  $R > 1$ . Let  $\Gamma$  denote the half line  $\Gamma = \{x \in \mathbb{R} \subset \mathbb{C} : x \geq R-1\}$ , let  $B \subset \mathbb{R}^m$  be a compact set, and denote by  $(r, x)$  the coordinates on  $B \times \Gamma$ . Let  $h, h' = \frac{\partial h}{\partial x} \in \mathcal{C}(B \times \Gamma)$ , and assume that

$$|h(r, x)| < \frac{C}{2} \quad , \quad |h'(r, x)| < \frac{1}{2}.$$

**Definition III.2.1.** Let  $\theta \in [0, 2\pi)$ . Then the set of curves

$$e^{i\theta} \cdot \{x + ih(r, x) : r \in B, x \in \Gamma\}$$

is referred to as being  $\theta$ -directed, and subordinate to  $R, C$ . A family of curves is said to be  $\theta$ -directed if it is  $\theta$ -directed subordinate to  $R, C$  for sufficiently large  $R, C$ .

With the notation in the previous section, set  $\psi(z) := \frac{1}{z} + 1$ ,  $\lambda_{r,j} := \psi(l_{r,j})$ ,  $c_j(r) := \psi(a_j(r))$ . Then  $\Lambda_r := \{\lambda_{r,1}, \dots, \lambda_{r,n}\}$  is a set of disjoint slits in  $\mathbb{P}^1$ , where  $\lambda_{r,1}$  is the negative real axis and  $\lambda_{r,j}$  are circular slits (or possibly straight line segments along the real axis) for  $j = 2, \dots, n$ . We set  $e^{i\theta_{r,j}} := \psi'(a_j(r))/|\psi'(a_j(r))|$ , i.e., we have that  $e^{i\theta_{r,j}}$  is a unit tangent to the circle  $\Lambda_{r,j}$  on which  $\lambda_{r,j}$  lies at the point  $c_j(r)$ . Setting  $\alpha_{r,j}(z) := e^{-i\theta_{r,j}}(z - c_j(r))$  we have that  $\alpha_{r,j}(\Lambda_{r,j})$  is a circle which is tangent to the real axis at the origin, and we let  $\kappa_{r,j}$  denote the signed curvature of this circle; positive if the circle is in the upper half plane, negative if the circle is in the lower half plane, and zero if the circle is the real axis.

**Proposition III.2.1.** Fix  $j \in \{2, \dots, n\}$  and suppose that  $g_{r,j} \in \mathcal{O}(\Delta_\delta(c_j(r)))$  is a continuous family of functions, for  $r \in B$ . Let  $\theta \in [0, 2\pi)$ , and set

$$\varphi_j(r, z) := \frac{e^{i\theta}}{\alpha_{r,j}(z)} + g_{r,j}(z).$$

Then the family  $\Gamma_j$  of curves  $\varphi(r, \lambda_{r,j})$  is  $(\theta - \pi)$ -directed.

*Proof.* It suffices to prove this for  $\theta = 0$ . Then  $\alpha_{r,j}(\Lambda_{r,j})$  is parametrized near the origin by

$$\eta_{r,j}(x) = x + i \frac{\kappa_{r,j}}{2} x^2 + O(x^4).$$

Set  $\tilde{g}_{r,j}(z) = g_{r,j}(\alpha_{r,j}^{-1}(z))$ . We have that

$$\begin{aligned} \varphi_j(r, x) &= \frac{1}{x + i \frac{\kappa_{r,j}}{2} x^2 + O(x^4)} + \tilde{g}_{r,j}(\eta_{r,j}(x)) \\ &= \frac{x - i \frac{\kappa_{r,j}}{2} x^2 + O(x^4)}{x^2 + O(x^4)} + \tilde{g}_{r,j}(\eta_{r,j}(x)) \\ &= \left( \frac{1}{x} - i \frac{\kappa_{r,j}}{2} + O(x^2) \right) (1 + O(x^2)) + \tilde{g}_{r,j}(\eta_{r,j}(x)) \\ &= \frac{1}{x} - i \frac{\kappa_{r,j}}{2} + O(x) + \tilde{g}_{r,j}(\eta_{r,j}(x)). \end{aligned}$$

Since  $g_{r,j}(z)$  is close to a constant when  $z$  is close to  $c_j(r)$ , the uniform bound in the definition of  $(-\pi)$ -directed holds. Now

$$\varphi'_j(r, x) = \frac{-1}{x^2} + v_{r,j}(x),$$

### III. Families of Proper Holomorphic Embeddings and Carleman-type Theorems with parameters

where  $v_{r,j}(x)$  is bounded and scaling it to have almost unit length we see

$$x^2 \phi'_j(r, x) = -1 + x^2 v_{r,j}(x).$$

■

**Proposition III.2.2.** Fix  $\theta_2, \dots, \theta_n \in (0, 2\pi)$ . Define  $\phi_r: \mathbb{C} \setminus \{c_2(r), \dots, c_n(r)\} \rightarrow \mathbb{C}^2$  by

$$\phi_r(z) := \left( z, \sum_{j=2}^n \frac{e^{i\theta_j}}{\alpha_{r,j}(z)} \right).$$

Choose  $\delta > 0$  small, and let  $a, b \in \Delta_\delta(1/\sqrt{2})$ , and set  $A_{a,b}(z, w) := (az + bw, -bz + aw)$ . Write  $a = r_a e^{i\vartheta_a}, b = r_b e^{i\vartheta_b}$ . Then the family  $\Gamma_1$  defined by  $\Gamma_1 = \{\pi_1 \circ A_{a,b} \circ \phi_r(\lambda_{r,1}) : r \in B\}$  is  $(\vartheta_a - \pi)$ -directed, and each family  $\Gamma_j, j = 2, \dots, n$ , defined by  $\Gamma_j = \{\pi_1 \circ A_{a,b} \circ \phi_r(\lambda_{r,j}) : r \in B\}$  is  $(\vartheta_b + \theta_j - \pi)$ -directed.

*Proof.* For  $j = 2, \dots, n$  this is just Proposition III.2.1 since for any fixed  $j$  we have that  $\pi_1 \circ A_{a,b} \circ \phi_r(\lambda_{r,j})$  is parametrized by

$$\frac{r_b e^{i(\vartheta_b + \theta_j)}}{\alpha_{r,j}(z)} + \sum_{k \neq j} \left( \frac{r_b e^{i(\vartheta_b + \theta_k)}}{\alpha_{r,k}(z)} \right) + r_a e^{i\vartheta_a} z.$$

For  $j = 1$  this is because  $\pi_1 \circ A_{a,b} \circ \phi_r(\lambda_{r,j})$  is parametrized by  $r_a e^{i\vartheta_a} z + g_r(z)$  where  $g_r(z)$  is uniformly comparable to  $\frac{1}{z}$ . ■

### III.3 Carleman approximation with parameters

We will start by introducing some notation. Afterwards, we present Theorem III.3.1, a Carleman-type theorem (see e.g., [5]), which is the main result of the present section: families of smooth functions holomorphic on a disc can be approximated by entire functions on a smaller disc and on the union of several Lipschitz curves. The proof is obtained applying inductively Corollary III.3.1, which in turn easily follows from Proposition III.3.1, a tool that allows to approximate smooth functions on compact pieces of a Lipschitz curve; Corollary III.3.1 extends the result to several curves. Proposition III.3.1 relies on three technical lemmata that will be presented in Section III.3.3.

#### III.3.1 The setup

Recall that  $R > 1$ ,  $\Gamma$  is the half line  $\Gamma := \{x \in \mathbb{R} \subset \mathbb{C} : x \geq R-1\}$ ,  $B \subset \mathbb{R}^m$  is a compact and  $(r, x)$  are the coordinates on  $B \times \Gamma$ . For  $k = 1, \dots, n$  let  $h_k, h'_k = \frac{\partial h_k}{\partial x} \in \mathcal{C}(B \times \Gamma)$  be such that

$$|h_k(r, x)| < \frac{C}{2}, \quad |h'_k(r, x)| < \frac{1}{2} \quad (\text{III.3.1})$$

for some  $C > 0$ , for every  $(r, x) \in B \times \Gamma$ , and every  $k = 1, \dots, n$ . Then, setting  $l = 1/2$ , we have that

$$|h_k(r, x_1) - h_k(r, x_2)| \leq l|x_1 - x_2|, \quad \forall x_1, x_2 \in \Gamma, \quad r \in B, \quad (\text{III.3.2})$$

so  $h_k$  is  $l$ -Lipschitz and in this way we also call its graph. Let  $0 = \theta_1 < \theta_2 < \dots < \theta_n < 2\pi$  and define the Lipschitz curves

$$\Gamma_{k,r} := e^{i\theta_k} \cdot \{x + ih_k(r, x) : x \in \Gamma\}$$

and their union

$$\Gamma_r := \bigcup_{k=1}^n \Gamma_{k,r}.$$

If  $D \subseteq \Omega \subseteq \mathbb{C}$  are domains, a useful notation is given by setting

$$\mathcal{P}(B, \Omega, D) := \{f \in \mathcal{C}(B \times \Omega) : f(r, \cdot) \in \mathcal{O}(D) \quad \forall r \in B\}$$

and

$$\mathcal{P}(B, \Omega) := \mathcal{P}(B, \Omega, \Omega).$$

**Theorem III.3.1** (Carleman-type Theorem with parameters). Assume that  $f \in \mathcal{P}(B, \mathbb{C}, \overline{\Delta}_{\rho+3+\frac{3C}{2}})$  for some  $\rho > R$ . Then for any  $\epsilon \in \mathcal{C}(\mathbb{C})$ ,  $\epsilon > 0$ , there exists  $g \in \mathcal{P}(B, \mathbb{C})$  such that

$$|g(r, z) - f(r, z)| < \epsilon(z)$$

for all  $z \in \overline{\Delta}_\rho \cup \Gamma_r$ ,  $r \in B$ .

### III.3.2 Proof of Theorem III.3.1

Fix  $j \in \mathbb{N}$ ,  $j \geq R$  and let  $b$  be some real number such that

$$j + 3 + \frac{3C}{2} < b.$$

For  $\rho \geq C$  set

$$\psi(\rho) := \arcsin \frac{C}{\rho}$$

and define

$$S_\rho := \{se^{i\theta} : 0 < s < \infty, |\theta| < \psi(\rho)\} \quad \text{and} \quad A_{\rho,b} := \Delta_b \setminus \overline{S}_\rho.$$

Then  $S_\rho$  is the wedge in the right half-plane bounded by the straight lines passing through the origin and the intersection between  $\partial\Delta_\rho$  and the lines  $y = \pm C$ . Up

### III. Families of Proper Holomorphic Embeddings and Carleman-type Theorems with parameters

---

to consider a larger  $R$ , we assume  $e^{i\theta_j}S_\rho \cap e^{i\theta_k}S_\rho = \emptyset$  for all  $j \neq k$  for  $\rho \geq R$ . We define further the following sets

$$\begin{aligned}\omega_1 &:= \{z = x + iy : j + 1 < x, |z| < b, |y| < C\} \\ \omega_2 &:= \{z = x + iy : 0 < x < j + 2, |y| < C\} \cup A_{j,b} \\ \Omega &:= \omega_1 \cup \omega_2\end{aligned}$$

Given  $\delta > 0$ , we will denote the open  $\delta$ -neighborhood of  $D$  as

$$D(\delta) := \{z \in \mathbb{C} : d(z, D) < \delta\}.$$

The following proposition, or rather its corollary below, is the main technical ingredient in the proof of the Carleman Theorem III.3.1. The proposition follows from Lemma III.3.1, Lemma III.3.2, and finally Lemma III.3.3 below.

**Proposition III.3.1.** Assume that  $n = 1$ . Let  $\alpha: \bigcup_{r \in B} \{r\} \times \Gamma_r \rightarrow \mathbb{C}$  be continuous such that  $\alpha(r, \cdot) \in \mathcal{C}_c(\Gamma_r)$  for every  $r \in B$ , with

$$\text{supp } \alpha(r, \cdot) \subset \{z = x + iy \in \Gamma_r : j + 3 + \frac{3C}{2} < x, |z| < b, |y| < C/2\} \quad \forall r \in B.$$

Then for every  $\epsilon > 0$  there exists  $\{Q_t\}_{t>0} \subset \mathcal{P}(B, \mathbb{C})$  such that

$$\|\alpha(r, \cdot) - Q_t(r, \cdot)\|_{\Gamma_r \cap \overline{\Delta}_b} < \epsilon \quad (\text{III.3.3})$$

for every  $r \in B$ ,  $0 < t < t_0$ , and

$$Q_t \rightarrow 0 \text{ as } t \rightarrow 0 \quad (\text{III.3.4})$$

uniformly on  $B \times \omega_2(\delta)$ , for some  $\delta > 0$ .

**Corollary III.3.1.** Let  $\alpha: \bigcup_{r \in B} \{r\} \times \Gamma_r \rightarrow \mathbb{C}$  be continuous such that  $\alpha(r, \cdot) \in \mathcal{C}_c(\Gamma_r)$  for every  $r \in B$ , with

$$\text{supp } \alpha(r, \cdot) \subset \{z \in \Gamma_r : j + 3 + \frac{3C}{2} < |z| < b\}, \quad \forall r \in B.$$

Then for every  $\epsilon > 0$  there exists  $\{Q_t\}_{t>0} \subset \mathcal{P}(B, \mathbb{C})$  such that

$$\|\alpha(r, \cdot) - Q_t(r, \cdot)\|_{\Gamma_r \cap \overline{\Delta}_b} < \epsilon \quad (\text{III.3.5})$$

for every  $r \in B$ ,  $0 < t < t_0$ , and

$$Q_t \rightarrow 0 \text{ as } t \rightarrow 0 \quad (\text{III.3.6})$$

uniformly on  $B \times \overline{\Delta}_j(\delta)$ , for some  $\delta > 0$ .

*Proof.* On  $e^{-i\theta_k}\Gamma_{k,r}$  define  $\alpha_k(r, z) := \alpha(r, e^{i\theta_k}z)$ . Using the proposition we obtain approximations  $Q_{t,k}(r, z)$ . Then setting

$$Q_t(r, z) := \sum_{k=1}^n Q_{t,k}(r, e^{-i\theta_k}z)$$

will yield the result for sufficiently small  $t$ .  $\blacksquare$

*Proof of Theorem III.3.1:* The proof is by induction on  $k \geq 0$ , and the induction hypothesis is the following. For every  $j = 0, \dots, k$  there exist:

- (i)  $g_j \in \mathcal{P}(B, \mathbb{C}, \overline{\Delta}_{\rho+j+3+\frac{3C}{2}})$ ,
- (ii)  $|g_j(r, z) - f(r, z)| < \epsilon(z)/2$  for all  $z \in \overline{\Delta}_\rho \cup \Gamma_r$ ,  $r \in B$ , and
- (iii)  $\|g_j - g_{j-1}\|_{B \times \overline{\Delta}_{\rho+j-1}} < 2^{-j}$  for  $j \geq 1$ .

We start by setting  $g_0 := f$ ; then in the case  $k = 0$  we see that (i), (ii) hold, and (iii) is void. Assume now that the induction hypothesis holds for some  $k \geq 0$ . Fix  $\eta > 0$  such that

$$g_k(r, \cdot) \in \mathcal{O}(\overline{\Delta}_{\eta+\rho+k+3+\frac{3C}{2}}),$$

and choose a cutoff function  $\chi \in \mathcal{C}^\infty(\mathbb{C})$  such that  $0 \leq \chi \leq 1$ , such that  $\chi = 0$  near  $\overline{\Delta}_{\rho+k+3+\frac{3C}{2}}$ , and  $\chi = 1$  outside  $\overline{\Delta}_{\eta+\rho+k+3+\frac{3C}{2}}$ . Now  $g_k$  may be approximated on  $\overline{\Delta}_{\eta+\rho+k+3+\frac{3C}{2}}$  to arbitrary precision by  $h_k \in \mathcal{C}(B)[z]$  using Taylor series expansion, and so

$$h_k + \chi \cdot (g_k - h_k) =: h_k + \alpha_k$$

approximates  $g_k$  to arbitrary precision. Hence it suffices to approximate  $\alpha_k$  to arbitrary precision by a suitable function. Multiplying  $\alpha_k$  by a suitable cutoff function so that Corollary III.3.1 applies, we have that  $\alpha_k$  may be approximated to arbitrary precision on

$$\bigcup_r \Gamma_r \cap \overline{\Delta}_{\rho+k+2+3+\frac{3C}{2}}$$

by a function  $Q_k \in \mathcal{P}(B, \mathbb{C})$  which is arbitrarily small on  $\overline{\Delta}_{\rho+k}$ . Setting then  $g_{k+1} := h_k + Q_k + \tilde{\chi} \cdot (\alpha_k - Q_k)$  where  $\tilde{\chi}$  is a third cutoff function such that  $\tilde{\chi} = 0$  near  $\overline{\Delta}_{\rho+k+1+3+\frac{3C}{2}}$  and  $\tilde{\chi} = 1$  near  $\mathbb{C} \setminus \overline{\Delta}_{\rho+k+2+3+\frac{3C}{2}}$ , completes the induction step. We may finish the proof of Theorem III.3.1 by setting  $g := \lim_{j \rightarrow \infty} g_j$ , which exists by (iii), and the approximation holds by (ii).  $\blacksquare$

### III.3.3 Lemmata: Mergelyan–type and Runge’s Theorems with parameters

The three lemmata we present and prove in this section are fundamental ingredients to formulate a Mergelyan–type Theorem (see e.g., [5]). The first one of

### III. Families of Proper Holomorphic Embeddings and Carleman–type Theorems with parameters

---

them generalizes a theorem claimed and proved by P. Manne in his Ph.D. thesis [10] and is about the holomorphic (entire) approximation of a family of smooth functions, each of which is defined on a Lipschitz curve in the complex plane.

**Lemma III.3.1.** Assume that  $n = 1$ , and let  $\alpha$  be as in Proposition III.3.1. Then for every  $\epsilon > 0$  there exists  $\{H_t\}_{t>0} \subset \mathcal{P}(B, \mathbb{C})$  such that

$$\|\alpha(r, \cdot) - H_t(r, \cdot)\|_{\Gamma_r} < \epsilon \quad (\text{III.3.7})$$

for every  $r \in B$ ,  $0 < t < t_0$ , for some  $t_0 > 0$ , and

$$H_t \rightarrow 0 \text{ as } t \rightarrow 0 \quad (\text{III.3.8})$$

uniformly on  $B \times (\omega_1 \cap \omega_2)(\delta)$ , for some  $\delta > 0$ .

*Proof.* Extend  $h_1$  to a function  $h$  on the whole real line by setting  $h(r, x) := h_1(r, x)$  for  $x \geq R - 1$  and  $h(r, x) := h_1(r, 2R - 2 - x)$  for  $x < R - 1$ . Define  $S_r := \{s + ih(r, s) : s \in \mathbb{R}\}$ . Denote by  $z = x + ih(r, x)$  a point in  $\Gamma_r$  and let  $\zeta = \zeta(r, s) = s + ih(r, s)$  be a parametrization of  $S_r$ . Further,  $\zeta'(r, s) = \frac{\partial \zeta}{\partial s}(r, s)$ , extend  $\alpha(r, \cdot)$  to  $S_r \setminus \Gamma_r$  to be 0 for all  $r \in B$  and define

$$\begin{aligned} H_t(r, z) &:= \int_{S_r} \alpha(r, \zeta) K_t(\zeta, z) d\zeta \\ &= \int_{\mathbb{R}} \alpha(r, \zeta(r, s)) K_t(\zeta(r, s), z) \zeta'(r, s) ds \end{aligned}$$

for  $t > 0, r \in B, z \in \mathbb{C}$ , where

$$K_t(\zeta, z) := \frac{1}{t\sqrt{\pi}} e^{-\frac{(\zeta - z)^2}{t^2}}$$

is the Gaussian kernel.

We start by proving (III.3.8). Let  $z = x + iy \in (\omega_1 \cap \omega_2)(\delta)$ . We have that

$$\begin{aligned} |H_t(r, z)| &\leq \frac{1}{t\sqrt{\pi}} \int_{\mathbb{R}} |\alpha(r, \zeta(r, s))| e^{-\frac{(s-x)^2 - (h(r, s) - y)^2}{t^2}} |\zeta'(r, s)| ds \\ &= \frac{1}{t\sqrt{\pi}} \int_{j+3+\frac{3}{2}C < s < b} |\alpha(r, \zeta(r, s))| e^{-\frac{(s-x)^2 - (h(r, s) - y)^2}{t^2}} |\zeta'(r, s)| ds, \end{aligned}$$

and  $(s - x)^2 - (h(r, s) - y)^2 \geq (1 + \frac{3C}{2} - \delta)^2 - (\frac{3C}{2})^2$ , therefore (III.3.8) follows.

For any fixed  $\eta > 0$  we split  $S_r$  as

$$\begin{aligned} S_r^{(1)} &:= \{\zeta \in S_r : |\Re(\zeta - z)| \leq \eta\} = \{\zeta(r, s) : |s - x| \leq \eta\} \\ S_r^{(2)} &:= \{\zeta \in S_r : |\Re(\zeta - z)| > \eta\} = \{\zeta(r, s) : |s - x| > \eta\}. \end{aligned}$$



Since by (III.3.2) we have

$$|K_t(\zeta, z)| \leq \frac{1}{t\sqrt{\pi}} e^{-\frac{(s-x)^2(1-l^2)}{t^2}},$$

we immediately get the following upper bound:

$$\begin{aligned} \int_{S_r^{(1)}} |K_t(\zeta, z)| d|\zeta| &\leq \frac{1}{t\sqrt{\pi}} \int_{x-\eta}^{x+\eta} e^{-\frac{(s-x)^2(1-l^2)}{t^2}} ds \\ &= \frac{1}{\sqrt{\pi(1-l^2)}} \int_{|u| \leq \frac{\sqrt{1-l^2}}{t}\eta} e^{-u^2} du \\ &\leq \frac{1}{\sqrt{1-l^2}}, \end{aligned} \quad (\text{III.3.9})$$

which holds for every  $z \in \Gamma_r$ ,  $r \in B$  and  $t > 0$ . Similarly, one sees that for all  $\epsilon > 0, \eta > 0$  there exists  $t_0 > 0$  such that

$$\int_{S_r^{(2)}} |K_t(\zeta, z)| d|\zeta| \leq \frac{1}{\sqrt{\pi(1-l^2)}} \int_{|u| > \frac{\sqrt{1-l^2}}{t}\eta} e^{-u^2} du < \epsilon \quad (\text{III.3.10})$$

for every  $z \in \Gamma_r$ ,  $r \in B$ ,  $0 < t < t_0$ . We need one last property of the kernel, that is

$$\int_{S_r} K_t(\zeta, z) d\zeta = 1 \quad (\text{III.3.11})$$

for all  $z \in \Gamma_r$ ,  $r \in B$  and  $t > 0$ . Let us consider the function

$$F(z) := \int_{S_r} K_t(\zeta, z) d\zeta = \frac{1}{t\sqrt{\pi}} \int_{S_r} e^{-\frac{(\zeta-z)^2}{t^2}} d\zeta$$

which is holomorphic entire. Let  $z = x \in \mathbb{R}$  and define for  $T > 0$

$$\begin{aligned} A(T) &:= \{u + i0 : -T \leq u \leq T\}, \\ S_r(T) &:= \{\zeta \in S_r : -T \leq s \leq T\}, \end{aligned}$$

and let  $\rho_r^\pm(T)$  be the straight line segment between  $\pm T$  and  $\pm T + ih(r, \pm T)$ . Set

$$\gamma_r(T) := A(T) + \rho_r^+(T) - S_r(T) - \rho_r^-(T)$$

which is a piecewise  $\mathcal{C}^1$ -smooth closed curve which is nullhomotopic, hence we get

$$\frac{1}{t\sqrt{\pi}} \int_{\gamma_r(T)} e^{-\left(\frac{\zeta-x}{t}\right)^2} d\zeta = 0$$

for every  $t > 0$ ,  $r \in B$ ,  $x \in \mathbb{R}$  and  $T > 0$ . On the other hand

$$\begin{aligned} \frac{1}{t\sqrt{\pi}} \int_{\gamma_r(T)} e^{-\left(\frac{\zeta-x}{t}\right)^2} d\zeta &= \frac{1}{t\sqrt{\pi}} \left( \int_{A(T)} e^{-\left(\frac{u-x}{t}\right)^2} du + \int_{\rho_r^+(T)} e^{-\left(\frac{\zeta-x}{t}\right)^2} d\zeta \right. \\ &\quad \left. - \int_{S_r(T)} e^{-\left(\frac{\zeta-x}{t}\right)^2} d\zeta - \int_{\rho_r^-(T)} e^{-\left(\frac{\zeta-x}{t}\right)^2} d\zeta \right). \end{aligned}$$

### III. Families of Proper Holomorphic Embeddings and Carleman-type Theorems with parameters

---

Passing to the limit as  $T \rightarrow +\infty$ , the vertical contributions vanish (as  $h$  is bounded), while

$$\frac{1}{t\sqrt{\pi}} \int_{A(T)} e^{-(\frac{u-x}{t})^2} du \longrightarrow \frac{1}{t\sqrt{\pi}} \int_{\mathbb{R}} e^{-(\frac{u-x}{t})^2} du = 1$$

and

$$\frac{1}{t\sqrt{\pi}} \int_{S_r(T)} e^{-(\frac{\zeta-x}{t})^2} d\zeta \longrightarrow \frac{1}{t\sqrt{\pi}} \int_{S_r} e^{-(\frac{\zeta-x}{t})^2} d\zeta = F(x)$$

for every  $t > 0$ ,  $r \in B$  and  $x \in \mathbb{R}$ . This implies that the entire function  $F$  is identically 1 on the real line for every  $t > 0$  and  $r \in B$ , so by the identity principle it is constantly 1 on the whole  $\mathbb{C}$ ; in particular (III.3.11) holds true.

We gathered all the ingredients to prove (III.3.7). Let  $\epsilon > 0$ , let  $\eta > 0$  such that  $|\alpha(r, \zeta) - \alpha(r, z)| < \epsilon$  for all  $z \in \Gamma_r$ ,  $\zeta \in S_r^{(1)}$ , for all  $r \in B$ . Then

$$\begin{aligned} |H_t(r, z) - \alpha(r, z)| &= \left| \int_{S_r^{(1)}} \alpha(r, \zeta) K_t(\zeta, z) d\zeta + \int_{S_r^{(2)}} \alpha(r, \zeta) K_t(\zeta, z) d\zeta - \alpha(r, z) \right| \\ &= \left| \int_{S_r^{(1)}} \alpha(r, \zeta) K_t(\zeta, z) d\zeta + \right. \\ &\quad \left. \int_{S_r^{(2)}} \alpha(r, \zeta) K_t(\zeta, z) d\zeta - \alpha(r, z) \int_{S_r} K_t(\zeta, z) d\zeta \right| \\ &\leq \left| \int_{S_r^{(1)}} (\alpha(r, \zeta) - \alpha(r, z)) K_t(\zeta, z) d\zeta \right| + \\ &\quad \left| \int_{S_r^{(2)}} (\alpha(r, \zeta) - \alpha(r, z)) K_t(\zeta, z) d\zeta \right| \\ &\leq \epsilon \int_{S_r^{(1)}} |K_t(\zeta, z)| d|\zeta| + \\ &\quad \left( \|\alpha(r, \cdot)\|_{S_r^{(2)}} + |\alpha(r, z)| \right) \int_{S_r^{(2)}} |K_t(\zeta, z)| d|\zeta| \\ &\leq \frac{\epsilon}{\sqrt{1-l^2}} + 2\epsilon \|\alpha(r, \cdot)\|_{S_r}, \end{aligned}$$

where the second equality follows from (III.3.11) and the last inequality follows from (III.3.9) and (III.3.10). So we can conclude, since this last quantity can be taken arbitrarily small for  $\epsilon$  small, uniformly in  $z \in \Gamma_r$  and  $r \in B$ , for all  $0 < t < t_0$ , where  $t_0$  comes from (III.3.10). ■

The following Lemma shows how to modify the approximation constructed in Lemma III.3.1, so that, besides approximating the given smooth function, it becomes arbitrarily small on a suitable region. The price to pay is that the approximation obtained this way is no more entire; we will get “entireness” back with Lemma III.3.3, that is a parametric version of Runge’s Theorem (see e.g., [5]).

**Lemma III.3.2.** Assume  $n = 1$ , and let  $\alpha$  be as in Proposition III.3.1. Then for every  $\epsilon > 0$  there exists  $\{\xi_t\}_{t>0} \subset \mathcal{P}(B, \Omega(\delta))$  such that

$$\|\alpha(r, \cdot) - \xi_t(r, \cdot)\|_{\Gamma_r \cap \overline{\Delta_b}} < \epsilon \quad (\text{III.3.12})$$

for every  $r \in B$ ,  $0 < t < t_0$ , and

$$\xi_t \rightarrow 0 \text{ as } t \rightarrow 0 \quad (\text{III.3.13})$$

uniformly on  $B \times \omega_2(\delta)$ , for some  $\delta > 0$ .

*Proof.* Let  $H_t$  be the map defined in Lemma III.3.1 and  $\phi_i: \Omega(\delta) \rightarrow [0, 1]$  be smooth, such that

(i)  $\text{supp } \phi_i \subset \omega_i(\delta)$ , and

(ii)  $\phi_1 + \phi_2 \equiv 1$  on  $\Omega(\delta)$

for some  $\delta > 0$ . Define

$$g_{t,1}(r, z) := -H_t(r, z)\phi_2(z), \quad g_{t,2}(r, z) := H_t(r, z)\phi_1(z)$$

on  $B \times \Omega(\delta)$ ; then  $g_{t,i}(r, \cdot)$  is a smooth function on  $\Omega(\delta)$  and

$$g_{t,2} - g_{t,1} = H_t \quad (\text{III.3.14})$$

holds true on  $B \times \Omega(\delta)$ , therefore  $\frac{\partial g_{t,1}}{\partial \bar{z}}(r, z)$  and  $\frac{\partial g_{t,2}}{\partial \bar{z}}(r, z)$  are the same function; call it  $v_t$  and consider  $v_t(r, \cdot)$  smoothly extended on  $\overline{\Omega(\delta)}$ . Hence, defining

$$u_t(r, z) := \frac{1}{2\pi i} \iint_{\Omega(\delta)} \frac{v_t(r, \zeta)}{\zeta - z} d\zeta \wedge d\bar{\zeta},$$

we can assume without loss of generality to have smoothed the corners of  $\omega_i$  so that  $\Omega(\delta)$  is smoothly bounded, hence we are allowed to apply Theorem 2.2 in [3], which ensures that  $u_t(r, \cdot)$  is smooth on  $\overline{\Omega(\delta)}$  for every  $r \in B$  and solves  $\frac{\partial u_t}{\partial \bar{z}} = v_t$  on  $B \times \overline{\Omega(\delta)}$ , hence

$$\Theta_{t,i} := g_{t,i} - u_t \in \mathcal{P}(B, \Omega(\delta)), \quad i = 1, 2. \quad (\text{III.3.15})$$

Then (III.3.8) and (i) imply

- $g_{t,i} \rightarrow 0$  uniformly on  $P \times \omega_i(\delta)$  as  $t \rightarrow 0$ ,
- $\frac{\partial g_{t,1}}{\partial \bar{z}}(r, z) = -H_t(r, z)\frac{\phi_2}{\partial \bar{z}}(z) \rightarrow 0$  uniformly on  $B \times \omega_1(\delta)$  as  $t \rightarrow 0$ , and
- $\frac{\partial g_{t,2}}{\partial \bar{z}}(r, z) = H_t(r, z)\frac{\phi_1}{\partial \bar{z}}(z) \rightarrow 0$  uniformly on  $B \times \omega_2(\delta)$  as  $t \rightarrow 0$ .

### III. Families of Proper Holomorphic Embeddings and Carleman-type Theorems with parameters

---

The last two imply  $u_t \rightarrow 0$  uniformly on  $B \times \Omega(\delta)$ , hence

$$\Theta_{t,i} \rightarrow 0 \quad (\text{III.3.16})$$

uniformly on  $P \times \omega_i(\delta)$  as  $t \rightarrow 0$ . Since  $\Theta_{t,2} - \Theta_{t,1} = H_t$  on  $B \times \Omega(\delta)$ , it follows from (III.3.7), (III.3.15) and (III.3.16) that

$$\xi_t := \begin{cases} H_t + \Theta_{t,1} & B \times \omega_1(\delta) \\ \Theta_{t,2} & B \times \omega_2(\delta) \end{cases}$$

satisfies the stated properties. ■

**Lemma III.3.3** (Runge-type Theorem with parameters). With the notation of the previous lemma, for all  $\epsilon > 0$ , there is  $Q_t \in \mathcal{C}(B)[z]$  (polynomial with coefficients in  $\mathcal{C}(B)$ ); in particular  $\{Q_t\}_{t \geq 0} \subset \mathcal{P}(B, \mathbb{C})$  such that

$$\|\xi_t - Q_t\|_{B \times \overline{\Omega}} < \epsilon$$

for all  $t > 0, r \in B$ .

*Proof.* Observe that  $\overline{\Omega}$  is compact polynomially convex. Let  $\gamma = \partial\Omega(\frac{\delta}{2})$ . We have that

$$(r, \zeta, z) \mapsto \frac{\xi_t(r, \zeta)}{\zeta - z}$$

is uniformly continuous on  $B \times \gamma \times \overline{\Omega}$ , hence for every  $\epsilon > 0$  there exists  $\eta > 0$ , such that, dividing  $\gamma$  into  $N$  pieces  $\gamma_1, \dots, \gamma_N$  whose length  $L(\gamma_j)$  is less than  $\eta$  and fixing a point  $\zeta_j \in \gamma_j$  for every  $j$ ,

$$\left| \frac{\xi_t(r, \zeta)}{\zeta - z} - \frac{\xi_t(r, \zeta_j)}{\zeta_j - z} \right| < \frac{1}{N} \frac{2\pi}{L(\gamma_j)} \epsilon$$

holds  $\forall (r, \zeta, z) \in B \times \gamma_j \times \overline{\Omega}$ . Calling  $\gamma_j(1), \gamma_j(0)$  the final and initial points of  $\gamma_j$ , for every  $(r, z) \in B \times \overline{\Omega}$  one has

$$\begin{aligned} \xi_t(r, z) - \sum_{j=1}^N \overbrace{\frac{\gamma_j(1) - \gamma_j(0)}{2\pi i}}^{=: \beta_t(r, z)} \frac{\xi_t(r, \zeta_j)}{\zeta_j - z} &= \frac{1}{2\pi i} \int_{\gamma} \frac{\xi_t(r, \zeta)}{\zeta - z} d\zeta - \sum_{j=1}^N \frac{1}{2\pi i} \int_{\gamma_j} \frac{\xi_t(r, \zeta_j)}{\zeta_j - z} d\zeta \\ &= \frac{1}{2\pi i} \sum_{j=1}^N \int_{\gamma_j} \left( \frac{\xi_t(r, \zeta)}{\zeta - z} - \frac{\xi_t(r, \zeta_j)}{\zeta_j - z} \right) d\zeta \end{aligned}$$

thus

$$\|\xi_t - \beta_t\|_{B \times \overline{\Omega}} < \epsilon \quad (\text{III.3.17})$$

for every  $t > 0$ . The result now follows since each rational function  $z \mapsto \frac{1}{\zeta_j - z}$  may be approximated arbitrarily well on  $\overline{\Omega}$  by polynomials. ■

### III.4 Andersén–Lempert Theory

We will now apply Andersén–Lempert Theory in  $B \times \mathbb{C}^2$ . We have that  $B \subset \mathbb{R}^N$ , and when talking about analytic properties of sets and functions on  $B \times \mathbb{C}^2$  we will think of  $B \times \mathbb{C}^2 \subset \mathbb{C}^N \times \mathbb{C}^2$ ,  $\mathbb{C}^N = \mathbb{R}^N + i\mathbb{R}^N$ . For instance, by saying that  $K \subset B \times \mathbb{C}^2$  is polynomially convex compact we mean polynomially convex in  $\mathbb{C}^N \times \mathbb{C}^2$ ; this is, in fact, equivalent to  $K_r$  being polynomially convex in  $\{r\} \times \mathbb{C}^2$  for each  $r \in B$ .

With the setup introduced in Section III.2 we now set  $s_{r,j} := \phi_r(\lambda_{r,j})$  and we set

$$S_r := \bigcup_{j=1}^n s_{r,j}.$$

In the product space  $B \times \mathbb{C}^2$  we define  $S := \{(r, (z, w)) : (z, w) \in S_r, r \in B\}$ .

**Proposition III.4.1.** Let  $K \subset (B \times \mathbb{C}^2) \setminus S$  be a compact set such that  $K$  is polynomially convex. Let  $T > 0$ , and let  $\epsilon > 0$ . Then there exists a continuous map  $g: B \times \mathbb{C}^2 \rightarrow \mathbb{C}^2$  such that the following hold for all  $r \in B$ .

- (i)  $g(r, \cdot) \in \text{Aut } \mathbb{C}^2$ ,
- (ii)  $\|g(r, \cdot) - \text{Id}\|_{K_r} < \epsilon$ , and
- (iii)  $g(r, S_r) \subset \mathbb{C}^2 \setminus T\mathbb{B}^2$ .

For the following lemma we extend the map  $\psi$  defined in Section III.2 to a map  $\psi: \mathbb{P}^1 \times \mathbb{C} \rightarrow \mathbb{P}^1 \times \mathbb{C}$  by setting  $\psi(z, w) := (1/z + 1, w)$ , and we extend the map  $\phi_r(z)$  to a rational map on  $\mathbb{C}^2$  by setting

$$\phi_r(z, w) := \left( z, w + \sum_{j=2}^n \frac{e^{i\theta_j}}{\alpha_{r,j}(z)} \right). \quad (\text{III.4.1})$$

Moreover for  $T' < T''$  we set  $S_r(T', T'') := \{(z, w) \in S_r : T' \leq |(z, w)| \leq T''\}$ , and we let  $S(T', T'') := \bigcup_r \{r\} \times S_r(T', T'')$  which is the union over  $r$  in the product space  $B \times \mathbb{C}^2$ . Finally define  $S(T', T'')(\delta) := \bigcup_r \{r\} \times S_r(T', T'')(\delta)$  for  $\delta > 0$ .

**Lemma III.4.1.** There exist  $T'' > T' \gg T$  arbitrarily large,  $\delta > 0$ , such that for any  $\epsilon > 0$  there exists an open set  $U \subset B \times \mathbb{C}^2$  containing  $S \cup \overline{S(T', T'')(\delta)}$  and a smooth fiber preserving map  $\psi: [0, 1] \times U \rightarrow B \times \mathbb{C}^2$  such that, for each  $r \in B$  the following hold:

- (i)  $\psi_{r,t}(\cdot)$  is an isotopy of holomorphic embeddings, and  $\psi_{r,0}(\cdot) = \text{Id}$ ,
- (ii)  $\psi_{r,t}(S_r) \subset S_r$  for every  $t \in [0, 1]$ ,
- (iii)  $\|\psi_{r,t} - \text{Id}\|_{\mathcal{C}^2(S_r(T', T'')(\delta))} < \epsilon$  for every  $t \in [0, 1]$ , and
- (iv)  $\psi_{r,1}(S_r) \subset \mathbb{C}^2 \setminus T\mathbb{B}^2$ .

### III. Families of Proper Holomorphic Embeddings and Carleman-type Theorems with parameters

*Proof.* Set  $\gamma_{r,j}(z, w) := (b_j(r) \cdot z + a_j(r), w)$  such that  $\gamma_{r,j}[-1, 0]$  parametrizes  $l_{r,j}$ . Setting  $F_{r,j} := \phi_r \circ \psi \circ \gamma_{r,j}$  we have that  $F_{r,j}[-1, 0]$  parametrizes  $s_{r,j}$ . Fix  $T > 0$  and choose  $-1 < s < 0$  such that

$$\bigcup_{r \in B} \bigcup_{j=1}^m F_{r,j}^{-1}((T+1)\overline{\mathbb{B}^2} \cap s_{r,j}) \subset [-1, s].$$

Choose any pair  $T', T''$  such that  $F_{r,j}^{-1}(s_{r,j}(T', T'')) \subset (s, 0)$  for all  $r, j$ .

For  $N \in \mathbb{N}$  define

$$\eta_{N,t}(z, w) := \left( \frac{z - t(1+s)e^{-N(z-s)} + t(1+s)e^{-N(-s)}}{1 - t(1+s)e^{-N(-s)}}, w \right).$$

Then  $\eta_{N,t}$  is an isotopy of injective holomorphic maps near the real line in the  $z$ -plane, and leaves the real line invariant, fixing 0. We see that

$$\eta_{N,1}(x, 0) = \left( \frac{x - (1+s)e^{-N(x-s)} + (1+s)e^{-N(-s)}}{1 - (1+s)e^{-N(-s)}}, 0 \right)$$

from which  $\eta_{N,1}(s, 0) = (-1, 0)$  and  $\lim_{x \rightarrow +\infty} \eta_{N,1}(x, 0) = (+\infty, 0)$ , so the interval  $[s, \infty)$  is stretched to the interval  $[-1, \infty)$  when  $t = 1$ . Note that for any  $s' > s$  we have that  $\lim_{N \rightarrow \infty} \eta_{N,t} = \text{Id}$  uniformly on  $\{\Re(z) \geq s'\}$ .

Now let  $\sigma_{N,t}$  be the inverse isotopy to  $\eta_{N,t}$ , i.e.,  $\sigma_{N,t} = \eta_{N,1-t} \circ \eta_{N,1}^{-1}$ ; it is injective holomorphic near the real line in the  $z$ -plane, and by choosing  $N$  large, may be extended, arbitrarily close to the identity, to any set  $\{\Re(z) \geq s'\}$  for  $s' > s$ .

We may now define  $\psi_{r,t}(\cdot)$  on  $s_{r,j}$  by

$$\psi_{r,t} := F_{r,j} \circ \sigma_{N,t} \circ F_{r,j}^{-1}.$$

The claims of the lemma are satisfied by choosing  $N$  large, and  $\delta$  sufficiently small. ■

**Remark III.4.1.** If  $\delta$  is sufficiently small and  $\epsilon$  further sufficiently small we get that

$$K \cup \psi_t(S \cup \overline{S(T', T'')(\delta)})$$

is polynomially convex. Observe first that  $K_r \cup S_r(T', T'')$  is polynomially convex, since  $K_r$  is, and  $S_r(T', T'')$  is a collection of disjoint arcs. For a sufficiently small  $\delta'$  it is known that the tube  $\overline{S_r(T', T'')(\delta')}$  is polynomially convex, and for sufficiently small  $\delta'$  we have that  $K_r \cup \overline{S_r(T', T'')(\delta')}$  is polynomially convex. Then if  $\delta < \delta'$  and we consider  $\psi_{t,r}$  as in the lemma with  $\delta'$  instead of  $\delta$ , if  $\epsilon$  is small enough we get our claim, since the  $\delta, \delta'$  may be chosen independently of  $r$ .

*Proof of Proposition III.4.1:* Fix  $0 < \delta < 1$ . For each  $\eta \in \delta\mathbb{B}^2$  we set  $v_\eta = (0, 1) + \eta$ , and we let  $\pi_\eta$  denote the orthogonal projection onto the orthogonal complement of  $v_\eta$ . After applying the linear transformation

$$A(z, w) = ((1/\sqrt{2})z + (1/\sqrt{2})w, -(1/\sqrt{2})z + (1/\sqrt{2})w)$$

it follows from Proposition III.2.2 that the family  $\pi_\eta(s_{r,1})$  is  $(\vartheta_{1,\eta} - \pi)$ -directed and that the families  $\pi_\eta(s_{r,j})$  are  $(\vartheta_{2,\eta} + \theta_j - \pi)$ -directed, where the  $\vartheta_{j,\eta}$ 's vary continuously with  $\eta$ . From now on we will assume that have applied the transformation  $A$  without changing the notation for all sets considered above.

By increasing  $T > 0$  we may assume that  $K_r \subset T\mathbb{B}^2$  for all  $r$ , and we fix  $R$  as in Theorem III.3.1 such that  $\pi_\eta(T\mathbb{B}^2) \subset \Delta_R$ , and choose  $T' < T''$  such that  $\pi_\eta(S_r(T', T'')) \subset \mathbb{C} \setminus \Delta_{R+3+\frac{3}{2}C}$  for all  $r$  and all  $\eta$ .

Let  $\psi_t$  be the isotopy from Lemma III.4.1, extended to be the identity on some neighborhood of  $K$  which we regard as being included in  $U$ . On  $\psi_{t_0}(U)$  we define the vector field  $X_{t_0}(\zeta) = \frac{d}{dt} \big|_{t=t_0} \psi_t(\psi_{t_0}^{-1}(\zeta))$  (here  $\zeta = (r, x) = (r, z, w)$ ). The goal is to follow the standard Andersén–Lempert procedure parametrically for approximating the flow of the time dependent vector field  $X_t$  by compositions of flows of complete fields, but to modify these so that they do not move  $S \setminus S(0, T'')$ . The proof is the same as the corresponding proof in [9] where this was done without parameters, but we include here a sketch and some additional details. The reader is assumed to be familiar with the Andersén–Lempert–Forstnerič–Rosay construction.

*Step 1:* We will find flows  $\sigma_{r,j}(t, x)$ ,  $j = 1, \dots, m$ , such that the composition

$$\sigma_{r,m} \circ \dots \circ \sigma_{r,1}(t, x)$$

approximates  $\psi_t$ . The flows are of two forms:

$$\sigma_{r,j}(t, x) = x + ta_{r,j}(\pi_j(x))v_j \quad (\text{III.4.2})$$

or

$$\sigma_{r,j}(t, x) = x + (e^{ta_{r,j}(\pi_j(x))} - 1)\langle x, v_j \rangle v_j. \quad (\text{III.4.3})$$

We write  $\sigma_{r,j}(t, x) = x + b_{r,j}(t, x)v_j$ .

*Step 2:* The plan is then roughly to find a family of cutoff functions  $\chi_j \in \mathcal{C}^\infty(\mathbb{C}^2)$ ,  $0 \leq \chi_j \leq 1$ , such that  $\chi_j \equiv 1$  near  $T'\mathbb{B}^2$  and  $\chi_j \equiv 0$  near  $\mathbb{C}^2 \setminus T''\mathbb{B}^2$ , and define

$$\tilde{\sigma}_{r,j}(t, x) := x + \chi_j(x)b_{r,j}(t, x)v_j,$$

in such a way that all compositions

$$\tilde{\sigma}(j)_r := \tilde{\sigma}_{r,j} \circ \dots \circ \tilde{\sigma}_{r,1}$$

are as close to the identity as we like in  $\mathcal{C}^1$ -norm on  $S_r(T', T'')(\delta/2)$ . Note that  $\tilde{\sigma}_{r,j} = \sigma_{r,j}$  on  $T'\mathbb{B}^2$  and  $\tilde{\sigma}_{r,j} = \text{Id}$  outside  $T''\mathbb{B}^2$ . In particular the families  $\pi_j(\tilde{\sigma}(j)_r(S_r))$  are as close as we like to the original families  $\pi_j(S_r)$  and identical outside some compact set.

*Step 3:* For each  $j$  we may rewrite  $\tilde{\sigma}_{r,j}$  on  $\tilde{\sigma}(j-1)_r(S_r)$  as

$$\tilde{\sigma}_{r,j}(t, x) = x + c_{r,j}(t, \pi_j(x))v_j \text{ or } \tilde{\sigma}_{r,j}(t, x) = x + (e^{c_{r,j}(t, \pi_j(x))} - 1)\langle x, v_j \rangle v_j.$$

### III. Families of Proper Holomorphic Embeddings and Carleman–type Theorems with parameters

---

where the  $c_{r,j}$ 's extend to be holomorphic near  $\overline{\Delta}_{R+3+\frac{3}{2}C}$  and zero on  $\pi_j(\tilde{\sigma}(j-1)_r(S_r(T''), \infty))$ .

*Step 4:* Approximate the coefficients  $c_{r,j}$  in the sense of Carleman using Theorem III.3.1.

We now include some estimates explaining why the above scheme works (see also [9] where the construction is done without dependence of parameters).

Choose  $\chi \in \mathcal{C}^\infty(\mathbb{C}^2)$  nonnegative such that  $\chi \equiv 1$  near  $T'\overline{\mathbb{B}^2}$  and  $\chi \equiv 0$  near  $\mathbb{C}^2 \setminus T''\overline{\mathbb{B}^2}$ .

We may assume that the vector fields  $X_{r,t}$  satisfy  $\|X_{r,t}\| < \alpha$  on  $S(T', T'')(\delta)$  for any small  $\alpha > 0$ . Thus, freezing the vector field at time  $i/N$  to obtain a vector field  $X^i$  with a flow  $\gamma_{r,t}^i$ , we have that  $\|\gamma_{r,t}^i(t/N, x) - x\| \leq (t/N)\alpha$  on  $S_r(T', T'')(2\delta/3)$ . Thus, we may assume that the compositions

$$\gamma_{r,t/N}^i \circ \cdots \circ \gamma_{r,t/N}^1$$

exist and remain arbitrarily close to the identity on  $S_r(T', T'')(\delta/2)$  for  $i < N$ .

By Remark III.4.1 we may approximate each vector field  $X_r^i$  to arbitrary precision by a polynomial vector field, which we will still denote by  $X_r^i$ . By the parametric Andersén–Lempert observation, see Lemma 4.9.9 and the proof of Theorem 4.9.10 in [6], the family of vector fields  $X_r^i$  may be written as a sum of shear and over-shear vector fields  $X_r^i = \sum_{j=1}^m Y_{r,j}^i$  with  $Y_{r,j}^i(x) = g_{r,j}^i(x)v_j$  with  $v_j = v_{\eta_j}$ , with flows

$$\sigma_{r,j}^i(t, x) = x + b_{r,j}^i(t, x)v_j.$$

Write

$$\Theta_r^i(t, x) := \sigma_{r,m}^i(t, x) \circ \cdots \circ \sigma_{r,1}^i(t, x) =: x + f_r^i(t, x).$$

It is known that the composition

$$(\Theta_{r,t/nN}^N)^n \circ \cdots \circ (\Theta_{r,t/nN}^1)^n \tag{III.4.4}$$

converges to the flow of  $X_{r,t}$  on  $K_r \cup S_r$  as  $N$  and then  $n$  tends to infinity. Our first goal is to replace the maps  $\sigma_{r,j}^i(t, x)$  by maps  $x + \chi_j(x)b_{r,j}^i(t, x)v_j$  with  $\chi_j(x) = 1$  for  $|x| \leq T'$  and  $\chi_j(x) = 0$  for  $|x| \geq T''$  in the composition (III.4.4) or partial compositions of it, and show that we still get maps that are close to the identity on  $S_r(T', T'')(\delta/2)$ . The compositions thus obtained will remain the same on  $\{|x| < T'\}$  and be the identity map outside  $\{|x| \leq T''\}$ .

We will now modify the flows on  $S_r(T', T'')(\delta)$ . We have that  $\|f_r^i(t, x)\| \leq 2\alpha t$  for  $t$  sufficiently small. If we set

$$\tilde{\Theta}_r^i(t, x) = x + \chi(x)f_r^i(t, x)$$

we see that

$$\|(\tilde{\Theta}_{r,(t/n)}^i)^m - \text{Id}\| \leq 2\alpha(m/n)t$$

for  $n$  large.



We decompose in a natural way

$$\sigma(j)_r(t, x) := \sigma_{r,j} \circ \cdots \circ \sigma_{r,1}(t, x) = x + h_{r,1}(t, x)v_1 + \cdots + h_{r,j}(t, x)v_j;$$

then the  $h_{r,j}(t, \cdot)$  go to zero as  $t \rightarrow 0$ . Now for large  $n$  define

$$\tilde{\sigma}_{r,1}(t/n, x) = x + \chi_1(x)h_{r,1}(t/n, x)v_1,$$

and by induction

$$\tilde{\sigma}_{r,j+1}(t/n, x) = x + \chi_{j+1}(\tilde{\sigma}(j)_r^{-1}(t/n, x))h_{r,j+1}(\tilde{\sigma}(j)_r^{-1}(t/n, x))v_{j+1}.$$

Then

$$\tilde{\sigma}_{r,m,t/n}^i \circ \cdots \circ \tilde{\sigma}_{r,1,t/n}^i(x) = \tilde{\Theta}_r^i(t/n, x),$$

and we get that

$$\|(\tilde{\Theta}_{r,t/Nn}^N)^n \circ \cdots \circ (\tilde{\Theta}_{r,t/Nn}^1)^n - \text{Id}\| \leq 2\alpha t,$$

and corresponding estimates hold for partial compositions. Note that this shows that  $S_r(T', T'')(\delta/2)$  remains in  $S_r(T', T'')(2\delta/3)$  where we may assume that the  $\mathcal{C}^1$ -norms of the  $f_r^i$ 's are arbitrarily small, and so by arguments similar to those above we get that all partial compositions are close to the identity in  $\mathcal{C}^1$ -norm, which will allow us to use the implicit function theorem to rewrite as in Step 3 above.

Finally, Step 4 is carried out exactly as in [9]. ■

### III.5 Proof of Theorem III.1.1

*Proof of Theorem III.1.1.* Recall that  $\psi: \mathbb{P}^1 \rightarrow \mathbb{P}^1$  was originally defined by  $\psi(z) = \frac{1}{z} + 1$  and  $\psi(\mathbb{P}^1 \setminus L_r) = \mathbb{P}^1 \setminus \Lambda_r$ . Then, if  $0 < \theta_2 < \theta_3 < \cdots < \theta_n$ ,  $\phi_r: \mathbb{C} \setminus \{c_2(r), \dots, c_n(r)\} \rightarrow \mathbb{C}^2$  was defined by

$$\phi_r(z) = \left( z, \sum_{j=2}^n \frac{e^{i\theta_j}}{\alpha_{r,j}(z)} \right).$$

Let  $S_r$  and  $S$  be the sets defined in Section III.4 and set  $X_r := \phi_r \circ \psi(\mathbb{P}^1 \setminus L_r) = \phi_r(\mathbb{P}^1) \setminus S_r$  which is a 1-dimensional complex manifold with boundary  $\partial X_r = S_r$ . Define

$$C_j^r := \mathbb{P}^1 \setminus L_r(1/j),$$

such that  $\{C_j^r\}_{j=1}^\infty$  is a normal exhaustion of  $\mathbb{P}^1 \setminus L_r$  by  $\mathcal{O}(\mathbb{P}^1 \setminus L_r)$ -convex compact sets. It follows that  $K_j^r := \phi_r \circ \psi(C_j^r)$ ,  $j \geq 1$  is a normal exhaustion of  $X_r$  by  $\mathcal{O}(X_r)$ -convex compact sets. The proof of Proposition 1 in [11] ensures the following two crucial facts:

- (i)  $K_j^r$  are polynomially convex, and

### III. Families of Proper Holomorphic Embeddings and Carleman–type Theorems with parameters

---

- (ii) given any  $K \subset \mathbb{C}^2 \setminus S_r$  compact polynomially convex, the set  $K \cup K_j^r$  is polynomially convex for any  $j$  large enough .

We construct now inductively a sequence of continuous mappings, whose continuous limit

$$h: \bigcup_{r \in B} (\{r\} \times X_r) \rightarrow \mathbb{C}^2$$

will be a fiberwise proper holomorphic embedding that we will compose with suitable mappings to prove the statement.

Proposition III.4.1 provides a continuous  $g_1: B \times \mathbb{C}^2 \rightarrow \mathbb{C}^2$  such that, for every  $r \in B$

- $g_1(r, \cdot) \in \text{Aut } \mathbb{C}^2$ , and
- $g_1(r, S_r) \subset \mathbb{C}^2 \setminus 1\overline{\mathbb{B}^2}$ .

Assume that we have constructed  $H_j: B \times \mathbb{C}^2 \rightarrow B \times \mathbb{C}^2$ ,  $H_j(r, \cdot) = (r, h_j(r, \cdot))$ , continuous such that for every  $r \in B$  we have that

- $h_j(r, \cdot) \in \text{Aut } \mathbb{C}^2$ , and
- $h_j(r, S_r) \subset \mathbb{C}^2 \setminus j\overline{\mathbb{B}^2}$ .

It follows from (ii) that

$$L_j^r := h_j(r, K_{m_j}^r) \cup j\overline{\mathbb{B}^2} \subset \mathbb{C}^2 \setminus h_j(r, S_r) .$$

is polynomially convex for sufficiently large  $m_j$ . Then for every  $\epsilon_j > 0$ , Proposition III.4.1 gives us  $g_{j+1}: B \times \mathbb{C}^2 \rightarrow \mathbb{C}^2$  continuous, such that for every  $r \in B$  the following hold:

- $g_{j+1}(r, \cdot) \in \text{Aut } \mathbb{C}^2$ ,
- $\|g_{j+1}(r, \cdot) - \text{Id}\|_{L_j^r} < \epsilon_j$ , and
- $g_{j+1}(r, h_j(r, S_r)) \subset \mathbb{C}^2 \setminus (j+1)\overline{\mathbb{B}^2}$ .

Define  $G_{j+1}(r, \cdot) := (r, g_{j+1}(r, \cdot))$  and consequently  $H_{j+1} := G_{j+1} \circ H_j$ . Then letting  $\epsilon_j \rightarrow 0$  and  $m_j \rightarrow +\infty$  fast enough, the push–out method (see [6]) allows to conclude that, for every  $r \in B$ , the sequence  $\{h_j(r, \cdot)\}_j$  converges uniformly on compact subsets of  $D_r := \bigcup_{j \geq 1} h_j^{-1}(r, L_j^r)$  to a biholomorphism  $h_r: D_r \rightarrow \mathbb{C}^2$ . It is straightforward to check that  $X_r \subset D_r \subset \mathbb{C}^2 \setminus S_r$ , from which it follows that  $S_r = \partial X_r \subseteq \partial D_r$ , thus  $h_r: X_r \rightarrow \mathbb{C}^2$  is a proper holomorphic embedding for every  $r \in B$ . So setting  $H(r, \cdot) := h_r(\cdot)$ ,  $\Psi(r, \cdot) := (r, \psi(\cdot))$ , and  $\Phi(r, \cdot) := (r, \phi_r(\cdot))$ , the mapping  $\Xi := H \circ \Phi \circ \Psi: \Omega \rightarrow \mathbb{C}^2$  proves the claim of the Theorem. ■

# Bibliography

- [1] Alarcón, A.; *Complex curves in pseudoconvex Runge domains containing discrete subsets*. J. Anal. Math. **140** (2020), no. 1, 207–226.
- [2] Alarcón, A. and López, F. J.; *Proper holomorphic embeddings of Riemann surfaces with arbitrary topology into  $\mathbb{C}^2$* . J. Geom. Anal. **23** (2013), no. 4, 1794–1805.
- [3] Bell, S. R.; *The Cauchy Transform, Potential Theory and Conformal Mapping*, Second ed. CRC Press, Taylor & Francis Group, 2015
- [4] Černe, M. and Forstnerič, F.; *Embedding some bordered Riemann surfaces in the affine plane*. Math. Res. Lett., **9** (2002), 683–696
- [5] Fornæss, J. E., Forstnerič, F. and Wold, E. F.; *Holomorphic approximation: the legacy of Weierstrass, Runge, Oka–Weil and Mergelyan*. *Advancements in Complex Analysis – From Theory to Practice*. Breaz, Daniel & Rassias, Michael Th. (Ed.), Springer Nature. ISSN 978–3–030–40119–1. p. 133–192.
- [6] Forstnerič, F.; *Stein manifolds and holomorphic mappings. The homotopy principle in complex analysis. Second edition. Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*, 56. Springer, Cham, 2017.
- [7] Globevnik, J. and Stensønes, B.; *Holomorphic embeddings of planar domains into  $\mathbb{C}^2$* . Math. Ann., **303**, 579–597 (1995).
- [8] Goluzin, G. M.; *Geometric Theory of Functions of a Complex Variable*.
- [9] Kutzschebauch, F. and Wold, E. F.; *Carleman approximation by holomorphic automorphisms of  $\mathbb{C}^2$* . J. Reine Angew. Math. **738** (2018), 131–148.
- [10] Manne, P.; *Carleman approximation in several complex variables*, Ph.D. Thesis. University of Oslo, 1993.
- [11] Wold, E. F.; *Embedding Riemann surfaces properly into  $\mathbb{C}^2$* . Int. J. Math. **17**, pp. 963–974, 2007.

**Authors' addresses**

**G. D. Di Salvo** University of Oslo, Postboks 1337 Blindern, 0316 Oslo, Norway,  
giovandd@math.uio.no

**T. Ritter** University of Stavanger, Postboks 8600, 4036 Stavanger,  
tyson.ritter@uis.no

**E. F. Wold** University of Oslo, Postboks 1337 Blindern, 0316 Oslo, Norway,  
erlendfw@math.uio.no

# Approximation and accumulation results of holomorphic mappings with dense image

G. D. Di Salvo

Published in *Mathematica Scandinavica*, 2023, volume 129, issue 2, pp. 273–283

## Abstract

We display four approximation theorems for manifold-valued mappings. The first one approximates holomorphic embeddings on pseudoconvex domains in  $\mathbb{C}^n$  with holomorphic embeddings with dense images. The second theorem approximates holomorphic mappings on complex manifolds with bounded images with holomorphic mappings with dense images. The last two theorems work the other way around, constructing (in different settings) sequences of holomorphic mappings (embeddings in the first one) converging to a mapping with dense image defined on a given compact minus certain points (thus in general not holomorphic).

## Contents

IV.1	Introduction . . . . .	73
IV.2	Main Results . . . . .	74
IV.3	Technical Tools . . . . .	75
IV.4	Inductive procedure . . . . .	77
IV.5	Proofs . . . . .	82

## IV.1 Introduction

Calling  $Y$  a connected complex manifold and  $\Delta$  the unit open disc in  $\mathbb{C}$ , F. Forstnerič and J. Winkelmann proved in [5] that the set  $\mathcal{D}$  of all holomorphic maps  $f: \Delta \rightarrow Y$  with dense image, is dense in  $\mathcal{O}(\Delta, Y)$ .

We are going to present four results which generalize in different directions the above-mentioned one. The first one is the most straightforward generalization

---

DOI: 10.7146/math.scand.a-136450.

## IV. Approximation and accumulation results of holomorphic mappings with dense image

---

in higher dimension, which is stated for holomorphic embeddings. The second one still provides approximating holomorphic mappings with dense images but in a more general setting. Moreover, it is no more a “density result” as the approximating mappings do not belong to the set of mappings we approximate. The last two theorems work the other way around, describing mappings with dense images approximated by holomorphic mappings.

Another result accounting this topic can be found in [1]: given a closed complex submanifold  $X \subsetneq \mathbb{C}^n$ , for  $n > 1$ , there exists a *complete* (the image of every divergent path in  $X$  has infinite length in  $\mathbb{C}^n$ ) holomorphic embedding  $f: X \hookrightarrow \mathbb{C}^n$  with everywhere dense image; for  $n = 1$  the same result holds for complete holomorphic embeddings  $f: \mathbb{C} \hookrightarrow \mathbb{C}^2$ . If moreover  $X \cap \mathbb{B}^n \neq \emptyset$  and  $K \subset X \cap \mathbb{B}^n$  is compact, there exists a Runge domain  $\Omega \subset X$  containing  $K$  which admits a complete holomorphic embedding  $f: \Omega \hookrightarrow \mathbb{B}^n$  with dense image.

Finally in [3] it is proved the existence of a holomorphic injective mapping with dense image from the open unit polydisc in  $\mathbb{C}^m$  to an  $m$ -dimensional paracompact connected complex manifold  $M$ .

The paper is organized as follows: we start by presenting the main results, then a technical section follows. The subsequent section shows the inductive procedure on which the proofs of the main results rely, which in turn are provided in the last section.

I would like to warmly thank E. F. Wold for directing me to write this paper.

### IV.2 Main Results

All the manifolds in this paper are connected. We will denote with  $\mathcal{O}(X, Y)$  the set of all holomorphic mappings  $h: X \rightarrow Y$  and with  $\mathcal{O}_{\text{emb}}(X, Y)$  the set of all holomorphic embeddings.

**Theorem IV.2.1.** Let  $\Omega \subset \mathbb{C}^n$  open, bounded, star-shaped (with respect to 0) and pseudoconvex. Let  $Y$  be a complex manifold. Then

$$\{h \in \mathcal{O}_{\text{emb}}(\Omega, Y) : \overline{h(\Omega)} = Y\}$$

is dense inside  $\mathcal{O}_{\text{emb}}(\Omega, Y)$  (with respect the compact convergence topology).

Observe that the density is considered with respect to the topology of  $Y$ . The distance  $d$  on  $Y$  exploited throughout the paper to make computations (namely: it allows to define the distance between two mappings  $f, g: X \rightarrow Y$  on a given  $M \subset X$  by  $\|f - g\|_M := \sup\{d(f(x), g(x)) : x \in M\}$ ) is induced by a Riemannian metric. Such  $d$  induces the same topology  $Y$  is endowed with.

**Remark IV.2.1.** We will implicitly assume  $n \leq \dim Y$ , otherwise  $\mathcal{O}_{\text{emb}}(\Omega, Y) = \emptyset$ .

**Theorem IV.2.2.** Let  $X$  be a complex manifold,  $Y$  Stein manifold. Then

$$\mathcal{H}(X, Y) := \{h \in \mathcal{O}(X, Y) : \overline{h(X)} = Y\}$$

*compactly approximates*

$$\mathcal{G}(X, Y) := \{g \in \mathcal{O}(X, Y) : g \text{ non constant, } g(X) \subset\subset Y\}$$

meaning that, given  $g \in \mathcal{G}(X, Y)$ , for every  $M \subset X$  compact and for every  $\epsilon > 0$  there exists  $h \in \mathcal{H}(X, Y)$  such that  $\|g - h\|_M < \epsilon$ .

Observe that  $\mathcal{G}(X, Y) = \emptyset$  whenever  $X$  is either compact or euclidean: in the former case the holomorphic mappings are constant, in the latter Liouville holds, obtaining again  $g$  constant, as its image is bounded. An example of a domain  $X$  such that  $\mathcal{G}(X, Y) \neq \emptyset$  is given by any open relatively compact domain  $X \subset\subset Y$  in a Stein manifold; just consider any non-constant holomorphic bounded mapping  $g: X \rightarrow Y$ , e.g., the inclusion  $\iota: X \hookrightarrow Y$ .

To present the last two theorems, we need the following definition:

**Definition IV.2.1.** Let  $K$  be a compact in  $Y$  complex manifold. A point  $\zeta \in K$  is *locally exposable* if there exists a  $\mathcal{C}^2$ -smooth strictly plurisubharmonic function  $\rho$  on some open neighborhood  $U$  of  $\zeta$  such that

1.  $\rho(\zeta) = 0$  and  $d\rho(\zeta) \neq 0$ , and
2.  $\rho < 0$  on  $(K \cap U) \setminus \{\zeta\}$ .

Definition IV.2.1 was originally presented in [2]. It is a generalization of the more standard definition of *local peak point* (see e.g., [7] pag. 354).

**Theorem IV.2.3.** Let  $Y$  be a complex manifold,  $K \subset Y$  connected not finite Stein compact. Then the set of locally exposable points is non-empty and for any such point  $x_0 \in K$ , there exist  $U_k \subset Y$  open neighborhood of  $K$  and  $F_k \in \mathcal{O}_{\text{inj}}(U_k, Y)$  compactly convergent to  $F: K \setminus \{x_0\} \rightarrow Y$  such that  $\overline{F(K \setminus \{x_0\})} = Y$ .

**Theorem IV.2.4.** Let  $Y$  be a Stein manifold,  $K \subset Y$  connected not finite compact. Denote with  $\Gamma$  the closure in  $Y$  of the set of locally exposable points for  $K$ , which is non-empty. Then there exists  $U_k \subset Y$  open neighborhood of  $K$  and  $F_k \in \mathcal{O}(U_k, Y)$  compactly convergent to  $F: K \setminus \Gamma \rightarrow Y$  such that  $\overline{F(K \setminus \Gamma)} = Y$ .

In Theorems IV.2.3 and IV.2.4 the limit mapping  $F$  is not in general holomorphic, as the domain is not in general an open set. But being  $F$  obtained as uniform limit of holomorphic mappings, the realization of  $F$  could be a consistent definition for a holomorphic map on  $K \setminus \{x_0\}$  or  $K \setminus \Gamma$ .

### IV.3 Technical Tools

The proofs will extensively exploit Theorem 1.1 in [2] and a slightly different version of it (whose proof follows automatically from the original one) which is as follows:

**Theorem IV.3.1.** Let  $Y$  be a complex manifold and  $Y_0 \subset Y$  Stein compact. Let  $\zeta \in Y_0$  be locally exposable and  $\gamma: [0, 1] \rightarrow Y$  smoothly embedded curve such that  $\gamma(0) = \zeta$  and  $\gamma((0, \delta]) \subset Y \setminus Y_0$  for some  $\delta > 0$ . Then, for every  $V$  open neighborhood of  $\gamma$  and for every  $\epsilon > 0$ , there exist the following:

1.  $U \subset Y$  neighborhood of  $Y_0$ ,

#### IV. Approximation and accumulation results of holomorphic mappings with dense image

---

2. an arbitrarily small  $V' \subset V$  neighborhood of  $\zeta$  , and
3.  $f: U \rightarrow Y$  holomorphic mapping such that
  - $f(\zeta) = \gamma(1)$  ;
  - $f(V') \subset V$  ;
  - $\|f - \text{Id}\|_{Y_0 \setminus V'} < \epsilon$  .

If the whole curve  $\gamma((0, 1])$  lies in  $Y \setminus Y_0$  then  $f$  can be chosen to be injective on  $U$ .

In [2] the theorem is formulated in a slightly more general setting and considering only the case  $\gamma((0, 1]) \subset Y \setminus Y_0$ .

**Lemma IV.3.1.** Let  $X$  be a metrizable topological space,  $W \subset \subset X$  connected not finite. Let  $f_k: U_k \rightarrow X$  be a sequence of continuous mappings, where  $U_k \subset X$  is some open neighborhood of  $K_k$  and  $K_1 := \overline{W}$ ,  $K_{k+1} := f_k(K_k)$ . Define  $F_k := f_k \circ \dots \circ f_1: \overline{W} \rightarrow X$  and consider

- $\{\epsilon_k\}_k$  positive real numbers ,
- $\{C_k\}_k$  compact exhaustion of  $W$ , that is  $C_k \subset X$  compact,  $C_k \subsetneq C_{k+1}$  and  $\bigcup_k C_k = W$ , and
- $\{V'_k\}_k$  open sets in  $X$  ,

such that, setting  $r_k := \max\{d(x, F_k(C_k)) : x \in K_{k+1}\}$ , the following hold:

- (i)<sub>k</sub>  $V'_k \subset U_k$  and  $V'_k \cap K_k \neq \emptyset$  ,
- (ii)<sub>k</sub>  $\max\{r_k, \|f_k - \text{Id}\|_{K_k \setminus V'_k}\} \leq \epsilon_k$ , and
- (iii)<sub>k</sub>  $V'_{k+1} \cap F_k(C_k) = \emptyset$  .

If  $F_k$  converges uniformly on compacts of  $W$  to  $F: W \rightarrow X$ , then for every  $x \in K_{k+1}$

$$d(x, F(W)) \leq \sum_{n \geq k} \epsilon_n . \quad (\text{IV.3.1})$$

*Proof.* Let  $x \in K_{k+1}$ . Then by definition and from (ii)<sub>k</sub> one gets

$$d(x, F_k(C_k)) \leq r_k \leq \epsilon_k .$$

Then (iii)<sub>k</sub> implies  $F_k(C_k) \subset K_{k+1} \setminus V'_{k+1}$  hence (ii)<sub>k+1</sub> says that  $f_{k+1}$  moves  $F_k(C_k)$  less than  $\epsilon_{k+1}$ , therefore

$$d(x, f_{k+1}(F_k(C_k))) \leq \epsilon_k + \epsilon_{k+1} .$$



Since  $f_{k+1}(F_k(C_k)) = F_{k+1}(C_k) \subset F_{k+1}(C_{k+1}) \subset K_{k+2} \setminus V'_{k+2}$  we can repeat the argument getting

$$d(x, \underbrace{f_{k+2}f_{k+1}(F_k(C_k))}_{F_{k+2}(C_k)}) \leq \epsilon_k + \epsilon_{k+1} + \epsilon_{k+2}.$$

Inductively, for every  $m \geq 0$  we get

$$d(x, F_{k+m}(C_k)) \leq \sum_{j=0}^m \epsilon_{k+j}$$

and passing to  $\lim_{m \rightarrow \infty}$  (which is well defined in left hand side, since the distance is continuous and  $\{F_n\}_n$  uniformly converges to  $F$  on  $C_k$ ) we get  $d(x, F(C_k)) \leq \sum_{j \geq 0} \epsilon_{k+j}$ . Since  $F(C_k) \subset F(W)$ , we have  $d(x, F(W)) \leq d(x, F(C_k))$  and we conclude.  $\blacksquare$

Recall now a useful property of Stein manifolds ([7], §2, Proposition 2.21 and Theorem 3.24):

**Theorem IV.3.2.** Let  $Y$  be a Stein manifold. Then there exists  $\rho$ , a  $\mathcal{C}^2$ -smooth strictly plurisubharmonic exhausting function for  $Y$ , such that the set of critical points  $C := \{z \in Y : d\rho(z) = 0\}$  is discrete in  $Y$ . In particular, for every  $c \in \mathbb{R}$ ,  $\{\rho < c\} \subset\subset Y$  and  $Y_c := \{\rho \leq c\}$  is a Stein compact.

**Remark IV.3.1.** With the notation of Theorem IV.3.2, we see that every regular boundary point of a strictly pseudoconvex domain  $\{\rho < c\}$  is locally exposable: take  $\zeta \in \{\rho = c\} \setminus C$ ; we assume it is the origin in suitable local coordinates. Then considering  $\tilde{\rho}(z) := \rho(z) - c - \epsilon|z|^2$  for  $\epsilon$  small enough, we conclude.

## IV.4 Inductive procedure

Consider  $Q = \{q_n\}_n \subset Y$  such that  $\overline{Q} = Y$ . Fix  $\epsilon > 0$  and define  $\epsilon_k := \frac{\epsilon}{2^{k+1}}$ . For Theorems IV.2.1 and IV.2.2 we fix a compact subset  $M$  of the domain,  $\Omega$  and  $X$  respectively. In what follows, we exploit Theorem IV.3.1 to get a suitable sequence of holomorphic functions  $f_k$  allowing to reach all the points of  $Q$  which have not been already reached, so that the image of the composition tends to invade the whole codomain. Along with the  $f_k$ , we construct both a compact exhaustion  $\{C_k\}_k$  of the case domain  $\Omega$ ,  $X$ ,  $K \setminus \{x_0\}$  or  $K \setminus \Gamma$  so that it fulfills the hypothesis of Lemma IV.3.1, and all the other sequences needed. For the construction of the  $f_k$ , we will focus on Theorems IV.2.2 and IV.2.1; the procedure for the remaining two theorems is alike and it is afterwards explained. Similarly for the proofs: we worked out the details of the proof of Theorem IV.2.2, which is displayed by proving convergence of the composition of the  $f_k$ , approximation of the given function, and density of the image. The argument for the remaining three results is analogous and it is subsequently illustrated.

#### IV.4.1 Construction of $f_k$ for Theorem IV.2.2

##### IV.4.1.1 Existence of locally exposable points and base of the induction

Consider  $g: X \rightarrow Y$  holomorphic non-constant such that  $g(X) \subset\subset Y$ . Set  $K_1 := \overline{g(X)}$  and take  $c_1 \in \mathbb{R}$  such that  $K_1 \subseteq Y_{c_1}$  and  $K_1 \cap \partial Y_{c_1} \neq \emptyset$ . Consider  $\zeta_1 \in K_1 \cap \partial Y_{c_1}$ ; if  $\zeta_1 \notin C$  then it is locally exposable, otherwise we slightly move  $K_1$  by composing  $g$  with a suitable holomorphic small perturbation defined as follows. Assume  $\zeta_1$  is the only point in  $K_1 \cap \partial Y_{c_1}$ . Observe that in suitable local coordinates on  $\mathbb{C}^n$ , which we split as  $z = (z', z'') = x + iy = (x' + iy', x'' + iy'') \in \mathbb{C}^k \oplus \mathbb{C}^{n-k}$ , we can express  $\rho$  near the origin ([5], Lemma 3.10.1) as

$$\rho(z) = \rho(0) + E(x', x'', y', y'') + o(|z|^2)$$

where

$$E(x', x'', y', y'') := \sum_{j=1}^k (\lambda_j y_j^2 - x_j^2) + \sum_{j=k+1}^n (\lambda_j y_j^2 + x_j^2)$$

with  $\lambda_j > 1$  for  $j = 1, \dots, k$  and  $\lambda_j \geq 1$  for  $j = k+1, \dots, n$ , for some  $k \in \{0, 1, \dots, n\}$ . Assume  $\zeta_1$  to be the origin  $0 \in \mathbb{C}^n$  in these coordinates, so  $\rho(0) = \rho(\zeta_1) = c_1$ . The boundary of any  $Y_c$  is defined by  $\rho$ , so we want to move a small neighborhood of  $\zeta_1$  in a suitable direction allowing it to go across the level set  $\partial Y_{c_1}$ . Take then any nonzero vector  $v = (\xi' + i\eta', \xi'' + i\eta'')$  with  $\xi' = 0 \in \mathbb{R}^k$ . By standard results on Stein Manifolds ([6], Corollary 5.6.3), given  $w \in T_{\zeta_1} Y$  there exists a holomorphic vector field  $V: Y \rightarrow TY$  such that  $V(\zeta_1) = w$ . In our case we take  $w$  to correspond to  $v$ . The flow of  $V$  on some neighborhood  $W$  of  $K_1$  is a holomorphic mapping  $\phi_t: W \rightarrow Y$  defined for complex times sufficiently small in modulus, say  $|t| < T$ ,  $t \in \mathbb{C}$ . In local coordinates around  $\zeta_1$  it is  $\phi_t(0) = tv + o(|t|^2)$  and up to shrinking  $T$ , we have that

$$\rho(\phi_t(0)) = \rho(0) + t^2 E(0, \xi'', \eta', \eta'') + o(|t|^3) > \rho(0) = c_1$$

for any  $|t| < T$ , considering now  $t \in \mathbb{R}$ . By continuity of  $(z, t) \mapsto \phi_t(z)$  and of  $\rho$ , the flow will drag a whole small neighborhood of  $\zeta_1$  beyond  $\partial Y_{c_1}$  and for sufficiently small times it will be the only piece of  $K_1$  crossing the boundary (as we are assuming that  $K_1 \cap \partial Y_{c_1}$  is just one point), that is: there exist  $T' < T$  and  $U$  neighborhood of  $\zeta_1$  sufficiently small, so that

- $\rho(\phi_t(z)) > \rho(0)$ ,  $\forall z \in U \cap K_1$ ,  $T'/2 < t < T'$ ;
- $\rho(\phi_t(z)) \leq \rho(0)$ ,  $\forall z \in K_1 \setminus U$ ,  $0 \leq t < T'$ ;
- $\phi_t(U \cap K_1) \cap C = \emptyset$ ,  $\forall T'/2 < t < T'$ .

So, considering now  $\tilde{K}_1 := \phi_t(K_1)$  for any  $T'/2 < t < T'$ , there exists some  $c > c_1$  such that  $\tilde{K}_1 \subseteq Y_c$  and  $\tilde{K}_1 \cap \partial Y_c \setminus C \neq \emptyset$ .

If  $K_1 \cap \partial Y_{c_1}$  contains more than one point, either it is not discrete or it is a finite set. In the former case  $K_1 \cap \partial Y_{c_1} \setminus C \neq \emptyset$ , in the latter we may assume  $K_1 \cap \partial Y_{c_1} = \{\zeta_1, \zeta'_1\}$ ; then, applying the previous argument to one of

these points, the piece of  $K_1$  that is dragged across  $\partial Y_{c_1}$ , could be not only  $U \cap K_1$ , but also  $U' \cap K_1$ , for any small time (where  $U, U' \subset Y$  are suitably small neighborhoods of  $\zeta_1$  and  $\zeta'_1$  respectively), leading to a similar  $\tilde{K}_1$  and achieving the same conclusion. Therefore, up to consider  $\phi_t \circ g$  instead of  $g$ , we may assume that  $\exists \zeta_1 \in K_1 \cap \partial Y_{c_1} \setminus C$ , which is then locally exposable by Remark IV.3.1 and could be sent to any point of  $Y \setminus K_1$  by the holomorphic mapping  $f_1$  provided by Theorem IV.3.1 (see next section).

Finally define  $f_0 := \text{Id}_Y$ ,  $C_0 := \emptyset$ ,  $F_0 := f_0$  and  $n_0 := 0$ .

#### IV.4.1.2 Inductive step

Assume we have the following:  $K_k \subset Y$  compact,  $c_k \in \mathbb{R}$  such that  $K_k \subseteq Y_{c_k}$  and  $\exists \zeta_k \in K_k \cap \partial Y_{c_k} \setminus C$ ,  $F_{k-1}$  holomorphic on some neighborhood of  $K_1$ , with  $F_{k-1}(K_1) = K_k$ ,  $C_{k-1} \subset X$  compact and  $n_{k-1} \in \mathbb{N}$ . Consider then a smoothly embedded curve  $\gamma_k: [0, 1] \rightarrow Y$  such that

- (i)<sub>k</sub>  $\gamma_k(0) = \zeta_k$  ;
- (ii)<sub>k</sub>  $\gamma_k(1) = q_{n_k}$  where  $n_k := \min\{n > n_{k-1} : q_n \notin K_k\}$  ;
- (iii)<sub>k</sub>  $\gamma_k((0, \delta_k]) \subset Y \setminus Y_{c_k}$  for some  $\delta_k > 0$  .

Then for every  $V_k$  open neighborhood of  $\gamma_k$ , Theorem IV.3.1 guarantees there exist

- (1)<sub>k</sub>  $U_k \subset Y$  neighborhood of  $Y_{c_k}$  ;
- (2)<sub>k</sub>  $V'_k \subset (V_k \cap B(\zeta_k, \epsilon_k)) \setminus F_{k-1}(g(M \cup C_{k-1}))$  ;
- (3)<sub>k</sub>  $f_k: U_k \rightarrow Y$  holomorphic such that the following hold:
  - (a)<sub>k</sub>  $f_k(\zeta_k) = q_{n_k}$  ;
  - (b)<sub>k</sub>  $f_k(V'_k) \subset V_k$  ;
  - (c)<sub>k</sub>  $\|f_k - \text{Id}\|_{Y_{c_k} \setminus V'_k} < \epsilon_k$  .

Observe that to apply Theorem IV.3.1,  $V'_k$  needs to be a neighborhood of the locally exposable point  $\zeta_k$ ; moreover to have convergence (see next section) it has to avoid the image of (the fixed compact  $M$  and) the compact  $C_{k-1}$ . This cannot happen if  $g$  is constant, as  $F_{k-1}(g(M \cup C_{k-1})) = K_k = \{\zeta_k\}$  so  $V'_k$  would be a punctured neighborhood of  $\zeta_k$ ; moreover  $\zeta_k \in Y_{c_k} \setminus V'_k$  and (c)<sub>k</sub> would fail for  $\epsilon_k$  small enough. Up to perturbing  $f_k$  as we did with  $g$ , there exists  $c_{k+1} \in \mathbb{R}$  such that  $K_{k+1} \subseteq Y_{c_{k+1}}$  and  $\exists \zeta_{k+1} \in K_{k+1} \cap \partial Y_{c_{k+1}} \setminus C$ ; then  $K_{k+1} := f_k(K_k)$  is a compact in  $Y$ ,  $F_k := f_k \circ F_{k-1}$  is holomorphic on some neighborhood of  $K_1$  and  $F_k(K_1) = K_{k+1}$ ; we finally choose a compact  $C_k \subset X$  large so that  $\max\{d(x, F_k(g(C_k))) : x \in K_{k+1}\} \leq \epsilon_k$  and  $C_{k-1} \subsetneq C_k$ . The induction may proceed.

## IV.4.2 Construction of $f_k$ for Theorem IV.2.1

### IV.4.2.1 Existence of a locally exposable point

Let  $g \in \mathcal{O}_{\text{emb}}(\Omega, Y)$ . Exploiting sharshapedness and up to considering  $g_\delta(z) = g((1-\delta)z)$  for  $0 < \delta < 1$ , we can suppose without loss of generality  $g$  to be holomorphic and injective on  $U$  a Stein neighborhood of  $\bar{\Omega}$ . Call  $R = \max_{z \in \bar{\Omega}} |z|$ , let  $\zeta_0 \in \partial\Omega$  be such that  $|\zeta_0| = R$  and define  $\rho(z) := |z|^2 - R^2$  which is strictly plurisubharmonic, hence  $\zeta_0$  a locally exposable point by Remark IV.3.1. Define  $\zeta_1 := g(\zeta_0)$  and  $K_1 := g(\bar{\Omega})$ .

If  $\dim Y = n$ , then  $\zeta_1$  is locally exposable with respect to  $\rho_1 := \rho \circ g^{-1}$  and  $K_1$  is a Stein compact, in fact  $W_\alpha = g_\alpha(\Omega)$ ,  $0 < \alpha < \delta$  is a neighbourhood basis of Stein domains since each  $W_\alpha$  is biholomorphic to  $\Omega$  which is holomorphically convex. In this case,  $\zeta_1$  is locally exposable and  $K_1$  is a Stein compact asking  $g$  for mere injectivity.

Let us now prove that  $\zeta_1$  and  $K_1$  are still a locally exposable point and a Stein compact respectively even in the case  $\dim Y = m > n$ . Working with local charts we can assume to work on open subsets of  $\mathbb{C}^m$ . Since  $dg(\zeta)$  has full rank  $n$  at each point  $\zeta \in U$ , it is  $\text{Im } dg(\zeta) \simeq \mathbb{C}^n$  and up to a linear change of coordinates, we can assume  $\text{Im } dg(\zeta_0) = \mathbb{C}^n \times \{0\}^{m-n}$  and obviously  $\text{Im } dg(\zeta_0) \oplus \text{Span}_{\mathbb{C}}(e_{n+1}, \dots, e_m) = \mathbb{C}^m$ . Define then, for  $(z_1, \dots, z_m) \in U \times \mathbb{C}^{m-n}$

$$\tilde{g}(z_1, \dots, z_m) := g(z_1, \dots, z_n) + z_{n+1}e_{n+1} + \dots + z_me_m.$$

Clearly  $\tilde{g}(\zeta_0, 0) = \zeta_1$  and it is locally invertible near  $(\zeta_0, 0)$ . Call  $h: A \rightarrow B$  the inverse, where  $A, B \subset \mathbb{C}^m$  are open neighborhoods of  $\zeta_1$  and  $(\zeta_0, 0)$  respectively and since

$$\pi_j \circ h(z) = z_j \text{ for } j = n+1, \dots, m,$$

we have that

$$g(U) \cap A = \{z \in A : z_{n+1} = \dots = z_m = 0\};$$

we worked around  $\zeta_1$ , but the same argument holds for any other point (regardless of whether it is regular or singular) so  $g(U)$  is a complex subvariety; in particular, it is *locally closed* (every point in  $g(U)$  has an open neighborhood  $W$  such that  $g(U) \cap W$  is closed in  $W$ ), thus it admits a Stein neighborhood basis ([5], Theorem 3.1.1). The same holds for any  $g(U')$ , with  $\bar{\Omega} \subset U' \subset U$ , therefore  $K_1$  is a Stein compact. Define now

$$\rho_1(z) := \rho \circ \alpha(z) + \sum_{j=n+1}^m |z_j|^2,$$

where  $\alpha := \pi|_{\mathbb{C}^n} \circ h: A \rightarrow U$  and  $\pi|_{\mathbb{C}^n}: \mathbb{C}^m \rightarrow \mathbb{C}^n$  is the projection on the first  $n$  coordinates. The term  $\rho \circ \alpha$  is plurisubharmonic as  $\rho$  is such and  $\alpha$  is holomorphic; moreover

$$L_w(\rho \circ \alpha; t) = L_{\alpha(w)}(\rho; \alpha'(w)t) > 0$$

for every  $t \in \mathbb{C}^m$ ,  $\pi|_{\mathbb{C}^n}(t) \neq 0$ ,  $w \in A$ , in fact  $\ker \alpha'(w) = \{0\}^n \times \mathbb{C}^{m-n}$  for any  $w \in A$  and  $\rho$  is strictly plurisubharmonic. Therefore we add the plurisubharmonic term  $\beta(z) := \sum_{j=n+1}^m |z_j|^2$ . Clearly

$$L_w(\beta; t) = \beta(t) > 0$$

for every  $t \in \mathbb{C}^m$ ,  $\pi|_{\mathbb{C}^{m-n}}(t) \neq 0$ ,  $w \in A$ . Therefore  $\rho_1$  is strictly plurisubharmonic on  $\zeta_1$  (actually on the whole  $A$ ). It is clear that  $\rho_1(\zeta_1) = 0$  and  $\rho_1 < 0$  on  $A \cap K_1 \setminus \{\zeta_1\}$ ; then a computation shows that  $d\rho_1(\zeta_1) \neq 0$ , hence  $\zeta_1$  is locally exposable in  $K_1$  with respect to  $\rho_1$ .

#### IV.4.2.2 Inductive procedure

Let us observe that  $Y \setminus K_1$  is connected, in fact if  $n = 1$ ,  $\Omega$  is simply connected (being starshaped) thus its boundary is connected and so is its image.

Let  $n \geq 2$  and  $\dim Y = n$ ; then  $g(\Omega)$  is a Stein domain, which has connected boundary for  $\dim Y \geq 2$  (see [8], pag. 22).

The remaining case, by Remark IV.2.1, is  $n \geq 2$  and  $m = \dim Y > n$ . In this case just observe that  $g(\Omega)$  is a complex submanifold of complex codimension  $m - n \geq 1$ , so its complement is connected.

Assume we have  $K_k \subset Y$  Stein compact,  $\zeta_k \in K_k$  locally exposable with respect to some strictly plurisubharmonic  $\rho_k$ ,  $C_{k-1} \subset \Omega$  compact,  $F_{k-1}$  holomorphic injective on some neighborhood of  $K_1$  such that  $F_{k-1}(K_1) = K_k$ . The construction of  $f_k: U_k \rightarrow Y$  is as in Section IV.4.1.2, with  $Y_{c_k} = K_k$  (Stein compact) and  $\gamma_k((0, 1]) \subset Y \setminus Y_{c_k}$  (which can be achieved since  $Y \setminus Y_{c_k}$  is connected, as above), allowing  $f_k$  to be injective on  $U_k$  and setting  $\zeta_{k+1} := f_k(\zeta_k) = q_{n_k}$  (which is locally exposable with respect to  $\rho_{k+1} := \rho_k \circ f_k^{-1}$ ). Finally set  $K_{k+1} := f_k(K_k)$  Stein compact,  $F_k := f_k \circ F_{k-1}$  holomorphic injective on some open neighborhood of  $K_1$  and  $C_k \subset \Omega$  as in Section IV.4.1.2. In particular  $F_k(K_1) = K_{k+1}$  and  $\{C_k\}_k$  is a compact exhaustion of  $\Omega$ .

#### IV.4.3 Construction of $f_k$ for Theorem IV.2.3

Since  $K$  is a Stein compact, there exists  $U \subset Y$  Stein neighborhood of  $K$  and consequently a plurisubharmonic exhausting function  $\rho: U \rightarrow \mathbb{R}$  as recalled in Theorem IV.3.2. Then, the existence of at least one locally exposable point  $x_0 \in K$  is guaranteed by the argument of Section IV.4.1.1, with  $x_0, K, U$  playing the role of  $\zeta_1, K_1$  and  $Y$  respectively and being  $\{C_k\}_k$  compact exhaustion for  $K \setminus \{x_0\}$ . So, given any  $x_0 \in K$  locally exposable, the rest of the inductive procedure is as in Section IV.4.2.2, except for  $M$  and  $g$  which here just do not play any role, with  $x_0, K, \rho$  playing the role of  $\zeta_1, K_1$  and  $\rho_1$  respectively.

#### IV.4.4 Construction of $f_k$ for Theorem IV.2.4

The construction is as in Section IV.4.1, with no  $M$ , no  $g$ , with  $K$  playing the role of  $K_1$  and  $\{C_k\}_k$  compact exhaustion for  $K \setminus \Gamma$ . At each step we get a locally exposable point  $\zeta_k \in K_k \cap \partial Y_{c_k} \setminus C_k$ , sent to  $q_{n_k}$  by  $f_k$  and corresponding to some  $x_k \in K = K_1$ , which is locally exposable as well (thus  $\{x_k\}_k \subset \Gamma$ ).

## IV.5 Proofs

### IV.5.1 Proof of Theorem IV.2.2

*Proof.*  $F_k: K_1 = \overline{g(X)} \rightarrow Y$  is holomorphic. Then from  $(2)_{k+1}$  it follows that for every fixed  $j$

$$F_k(g(C_j)) \subset K_{k+1} \setminus V'_{k+1} \quad (\text{IV.5.1})$$

holds true for every  $k \geq j$ . Hence we get that, for every fixed  $j$

$$\|F_{k+1} - F_k\|_{g(C_j)} = \|f_{k+1} - \text{Id}\|_{F_k(g(C_j))} \leq \|f_{k+1} - \text{Id}\|_{K_{k+1} \setminus V'_{k+1}} < \epsilon_k \quad (\text{IV.5.2})$$

is true for every  $k \geq j$ , so  $\{F_k\}_k$  converges on compact subsets of  $g(X)$  to  $F: g(X) \rightarrow Y$  holomorphic. As above,  $(2)_{k+1}$  implies  $F_k(g(M)) \subset K_{k+1} \setminus V'_{k+1}$  for every  $k$ , hence inequality (IV.5.2) holds true for all  $k$ , thus

$$\|F - \text{Id}\|_{g(M)} \leq \sum_{k \geq 0} \|F_{k+1} - F_k\|_{g(M)} < \epsilon,$$

allowing us to conclude that  $\|h - g\|_M < \epsilon$ , where  $h = F \circ g: X \rightarrow Y$  is the approximating mapping. We now check it actually has dense image. If  $h(X)$  is not dense in  $Y$ , then there exists an open ball  $B \subset Y$  such that

$$\beta := d(B, h(X)) > 0.$$

The construction of the sets  $K_k$  and the sequence  $\{n_k\}_{k \geq 1}$  allows to consider a partition of  $Q$  as

$$q_{n_{k-1}}, \dots, q_{n_k-1} \in K_k, \quad k \geq 1.$$

In this way we can define the sequence

$$k(n) := j \quad \text{for } n = n_{j-1}, \dots, n_j - 1, \quad \text{for } j \geq 1$$

and we have  $q_n \in K_{k(n)}$  holds true for all  $n$ ; the sequence  $n \mapsto k(n)$  is increasing and such that  $k(n) \rightarrow +\infty$  as  $n \rightarrow +\infty$  (otherwise  $\exists \tilde{k}, N$  such that  $q_n \in K_{\tilde{k}}$  for all  $n \geq N$ , so  $Q$  would not be dense). Since  $g(X) \subset\subset Y$ , it follows by Lemma IV.3.1 that

$$d(q_n, h(X)) = d(q_n, F(g(X))) \leq \sum_{j \geq k(n)-1} \epsilon_j.$$

This last sum is less than  $\beta$  for any  $n \geq n_\beta$ , for a suitably large  $n_\beta$ . Therefore  $\{q_n\}_{n > n_\beta}$ , which is still dense in  $Y$ , does not meet an open ball, contradiction. ■

### IV.5.2 Proof of Theorem IV.2.1

*Proof.* It is the same as the previous proof. Just observe that now  $F_k$  is defined on  $K_1 = g(\overline{\Omega})$ , it is holomorphic injective and so is  $F$ . Since  $g$  is injective by assumption, then the approximating mapping  $h = F \circ g$  is holomorphic injective as well. ■

### IV.5.3 Proof of Theorem IV.2.3

*Proof.* The mappings  $F_k$  are defined, holomorphic and injective on some open neighborhood of  $K$  and converge to  $F: K \setminus \{x_0\} \rightarrow Y$  uniformly on compacts of  $K \setminus \{x_0\}$ . The construction of mappings  $f_k$  ensures, as for Theorem IV.2.2, to achieve  $\overline{F(K \setminus \{x_0\})} = Y$ . ■

### IV.5.4 Proof of Theorem IV.2.4

*Proof.* As for Theorem IV.2.3 (except for injectivity of  $F_k$ ), with  $K \setminus \Gamma$  instead of  $K \setminus \{x_0\}$ . ■





# Bibliography

- [1] Alarcón, A. and Forstnerič, F.; Complete densely embedded complex lines in  $\mathbb{C}^2$ . *Proc. Amer. Math. Soc.*, 146(3):1059–1067, 2018.
- [2] Deng, F., Fornæss, J. E. and Wold, E. F.; *Exposing Boundary Points of Strongly Pseudoconvex Subvarieties in Complex Spaces*. Proceedings Of The AMS, Volume 146, Number 6, June 2018, Pages 2473–2487, <http://dx.doi.org/10.1090/proc/13693>, Article electronically published on March 9, 2018.
- [3] Fornæss, J. E. and Stout, E. L.; *Polydiscs in complex manifolds.. Math. Ann.* 227 (1977), no. 2, 145–153.
- [4] Forstnerič, F.; Stein manifolds and holomorphic mappings. The homotopy principle in complex analysis. Second edition. *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*, 56. Springer, Cham, 2017.
- [5] Forstnerič, F. and Winkelmann, J.; *Holomorphic Discs with Dense Images*. *Mathematical Research Letters* 12, 265–268 (2005)
- [6] Hörmander, L.; *An introduction to Complex Analysis in Several Variables*. North-Holland, 3rd edition (1990)
- [7] Range, R. M.; *Holomorphic Functions and Integral Representations in Several Complex Variables*. Springer, Graduate Text in Mathematics 108, (1986)
- [8] Serre, J. P.; Un théorème de dualité. *Comment. Math. Helv.* **29** (1955), 9–26.

## Author address

**G. D. Di Salvo** University of Oslo, Postboks 1337 Blindern, 0316 Oslo, Norway,  
giovandd@math.uio.no