



# Study of a new kinematic weighting algorithm for the measurement of CP asymmetries in charm decays

LHCb Collaboration

Georgios Christou

Supervisors: Dr. Federico Betti, Prof. Angelo Carbone

August 2023

## Abstract

We investigate the asymmetries that occur in charm decays at the LHCb, specifically we study  $D^{*+} \rightarrow D^0 \pi^+$  and  $\bar{D}^{*-} \rightarrow D^0 \pi^-$  where  $D^0 \rightarrow K^- K^+$  or  $D^0 \rightarrow \pi^- \pi^+$ . We study the effect of  $CP$  and detection asymmetries on MC samples generated via RapidSim and implement a new kinematic weighting function which allows us to keep events that are otherwise discarded from LHCb data, since they are associated with large detection asymmetries.

# 1 Introduction

We investigate charm decays and specifically the  $D^*$  meson. By studying the differences between  $D^{*+}$  and  $D^{*-}$  decays shown in Fig. 1 we can estimate the  $CP$  asymmetry in charm decays.




Figure 1: Feynman diagram showing  $D^{*\pm} \rightarrow D^0 \pi^\pm$  decays.

The total asymmetry one observes at an experiment is a combination of multiple asymmetries. Namely, the total asymmetry consists of a  $CP$  asymmetry and a *detection asymmetry*. The former is associated with the difference between matter and anti-matter, while the latter is associated with the differences while detecting the soft pion ( $\pi_s$ ) that is produced during the decay.

The total asymmetry can be estimated using

$$A_{\text{total}} = \frac{A_{CP} + A_D}{1 + A_C A_D} \quad (1)$$

where  $A_{CP}$  and  $A_D$  are the  $CP$  and *integrated detection asymmetries*. The latter can be calculated through

$$A_D = \frac{\int d\vec{p} N(\vec{p}) A_D(\vec{p})}{\int d\vec{p} N(\vec{p})} \quad (2)$$

At the LHCb one observes large pion detection asymmetries that are associated with specific kinematic regions, which so far have been discarded, thus reducing the statistics. We can, however, introduce a *weighting function* that allows us to keep events associated with large asymmetries. Such a weighting function can be expressed as the following ratio

$$Q(\vec{p}_{D^*}, \vec{p}_{\pi_s}) \simeq \frac{\Gamma_{D^0}^{\pi\pi}(\vec{p}_{D^*} - \vec{p}_{\pi_s}) + \Gamma_{\bar{D}^0}^{\pi\pi}(\vec{p}_{D^*} - \vec{p}_{\pi_s})}{\Gamma_{D^0}^{KK}(\vec{p}_{D^*} - \vec{p}_{\pi_s}) + \Gamma_{\bar{D}^0}^{KK}(\vec{p}_{D^*} - \vec{p}_{\pi_s})} \quad (3)$$

where  $\Gamma_{D^0/\bar{D}^0}^{\pi\pi/KK}$  are the normalized distributions of  $D^0$  candidates. Unfortunately in Run-2 such candidates were discarded, thus, we do not have a large enough sample to accurately calculate the weighting function, thus we resort to Monte Carlo simulations.

## 2 Analysis

### 2.1 RapidSim

## References

- [1] Roel Aaij et al. “LHCb Detector Performance”. In: *Int. J. Mod. Phys. A* 30.07 (2015), p. 1530022. DOI: 10.1142/S0217751X15300227. arXiv: 1412.6352 [hep-ex].
- [2] G. A. Cowan, D. C. Craik, and M. D. Needham. “RapidSim: an application for the fast simulation of heavy-quark hadron decays”. In: *Comput. Phys. Commun.* 214 (2017), pp. 239–246. DOI: 10.1016/j.cpc.2017.01.029. arXiv: 1612.07489 [hep-ex].
- [3] Roel Aaij et al. “Observation of CP Violation in Charm Decays”. In: *Phys. Rev. Lett.* 122.21 (2019), p. 211803. DOI: 10.1103/PhysRevLett.122.211803. arXiv: 1903.08726 [hep-ex].
- [4] Federico Betti. “CP violation in  $D^0 \rightarrow K^+K^-$  and  $D^0 \rightarrow \pi^+\pi^-$  decays and lepton-flavour universality test with the decay  $B^0 \rightarrow D^{*-}\tau^+\nu_\tau$ ”. PhD thesis. Bologna U., 2019. DOI: 10.6092/unibo/amsdottorato/8769.