



Study of a new kinematic weighting algorithm for the measurement of CP asymmetries in charm decays

LHCb Collaboration

Georgios Christou

Supervisors: Dr Federico Betti, Prof. Angelo Carbone

August 2023

Abstract

We investigate the asymmetries that occur in charm decays at the LHCb, specifically we study $D^{*+} \rightarrow D^0 \pi^+$ and $\bar{D}^{*-} \rightarrow D^0 \pi^-$ where $D^0 \rightarrow K^- K^+$ or $D^0 \rightarrow \pi^- \pi^+$. We study the effect of CP and detection asymmetries on MC samples generated via RapidSim and implement a new kinematic weighting function which allows us to keep events that are otherwise discarded from LHCb data, since they are associated with large detection asymmetries.

1 Introduction




Figure 1: Feynman diagram showing $D^{*\pm} \rightarrow D^0 \pi^\pm$ decays.

2 Analysis

2.1 RapidSim

References

- [1] Roel Aaij et al. “LHCb Detector Performance”. In: *Int. J. Mod. Phys. A* 30.07 (2015), p. 1530022. DOI: 10.1142/S0217751X15300227. arXiv: 1412.6352 [hep-ex].
- [2] G. A. Cowan, D. C. Craik, and M. D. Needham. “RapidSim: an application for the fast simulation of heavy-quark hadron decays”. In: *Comput. Phys. Commun.* 214 (2017), pp. 239–246. DOI: 10.1016/j.cpc.2017.01.029. arXiv: 1612.07489 [hep-ex].
- [3] Roel Aaij et al. “Observation of CP Violation in Charm Decays”. In: *Phys. Rev. Lett.* 122.21 (2019), p. 211803. DOI: 10.1103/PhysRevLett.122.211803. arXiv: 1903.08726 [hep-ex].