

UNIVERSITY OF AMSTERDAM

MADTER THESIS

Simultaneous Evolution of Morphology and Locomotion of Soft Robots by Novelty Search

Author:

Georgios METHENITIS

Supervisor:

Arnoud VISSER, UvA

Dario IZZO, ESA

Daniel HENNES, ESA

July 2014

UNIVERSITY OF AMSTERDAM

Abstract

Faculty Name

Department or School Name

Master of Science

Simultaneous Evolution of Morphology and Locomotion of Soft Robots by Novelty Search

by Georgios METHENITIS

Soft robotics is a vivid research field on the science and engineering aspects of soft materials in mobile machines. Recent development in soft robotics and evolutionary optimization have shown the possibility to simultaneously evolve the morphology and locomotion of soft robots. Generative encoding coupled with neural evolution of augmented topologies (NEAT) shows promising results. It is of interest to study the development of different types of locomotion in low gravity environments while rewarding diversity at the behavioral level. Unlike most evolutionary algorithms, novelty search seeks novel behaviors and has no defined objective (e.g. distance traveled or speed). This research could result in a taxonomy of possible locomotion strategies at a given (low) gravity level that inspire new concepts for future robotic missions.

Contents

Abstract	ii
Contents	iii
List of Figures	v
List of Tables	vii
1 Introduction	1
2 Related Work	3
3 Results	5
3.1 Results	5
3.1.1 Fitness Based Search	6
3.1.2 Novelty Search	8
3.1.3 Novelty-Fitness Based Search Comparison	11
3.1.4 Local competition in fitness based evolution	19
3.1.5 Local competition in novelty search evolution	22
A Appendix Title Here	23

List of Figures

3.1	Caption	6
3.2	Best so far fitness in body lengths displacement of softbot's center of mass from 10 runs for fitness based search. The gravity acceleration for this experiment used was -27.468 m/s^2 , the size of the lattice $10 \times 10 \times 10$ and 4 available materials (2-actuated).	6
3.3	Best so far fitness in body lengths displacement of softbot's center of mass from 10 runs together with their mean (thick line) for fitness based search. The gravity acceleration for this experiment used was -27.468 m/s^2 , the size of the lattice $10 \times 10 \times 10$ and 4 available materials (2-actuated).	7
3.4	Best so far fitness in body lengths displacement of softbot's center of mass from 10 runs for novelty search. The gravity acceleration for this experiment used was -27.468 m/s^2 , the size of the lattice $10 \times 10 \times 10$ and 4 available materials (2-actuated).	8
3.5	Best so far fitness in body lengths displacement of softbot's center of mass from 10 runs together with their mean (thick line) for novelty search. The gravity acceleration for this experiment used was -27.468 m/s^2 , the size of the lattice $10 \times 10 \times 10$ and 4 available materials (2-actuated).	9
3.6	Novelty search - best so far fitness in body lengths displacement of softbot's center of mass from 10 runs together with their mean (thick line). The novelty is computed as the average distance from the K -nearest behaviors for $K \in \{1, 2, 5, 10, 20\}$. The gravity acceleration for this experiment used was -27.468 m/s^2 , the size of the lattice $5 \times 5 \times 5$ and 4 available materials (2-actuated).	10
3.7	Best so far fitness in body lengths displacement of softbot's center of mass averaged over 10 runs together with the standard deviation error. The gravity acceleration for this experiment used was -27.468 m/s^2 , the size of the lattice $5 \times 5 \times 5$ and 4 available materials (2-actuated).	11
3.8	Fitness of the generation champion (best individual) per generation in body lengths displacement of softbot's center of mass averaged over 10 runs together with the standard deviation error. The gravity acceleration for this experiment used was -27.468 m/s^2 , the size of the lattice $5 \times 5 \times 5$ and 4 available materials (2-actuated).	12
3.9	Best so far fitness in body lengths displacement of softbot's center of mass averaged over 10 runs together with the standard deviation error. The gravity acceleration for this experiment used was -27.468 m/s^2 , the size of the lattice $10 \times 10 \times 10$ and 4 available materials (2-actuated).	13

3.10	Distributions of average population fitness per generation over 10 runs. The gravity acceleration for this experiment used was -27.468 m/s^2 , the size of the lattice $5 \times 5 \times 5$ and 4 available materials (2-actuated). Blue (Novelty search), Green(Fitness based search).	14
3.11	Novel individual's behaviors upto generation averaged over 10 runs. The novelty is computed as the average distance from the 10-nearest behaviors. The gravity acceleration for this experiment used was -27.468 m/s^2 , the size of the lattice $5 \times 5 \times 5$ and 4 available materials (2-actuated). Blue (Novelty search), Green (Fitness based search).	15
3.12	Best so far fitness in body lengths displacement of softbot's center of mass averaged over 10 runs together with the standard deviation error. The gravity acceleration for this experiment used was -27.468 m/s^2 , the size of the lattice $5 \times 5 \times 5$ and 4 available materials (2-actuated).	16
3.13	Best so far fitness in body lengths displacement of softbot's center of mass averaged over 10 runs together with the standard deviation error. The gravity acceleration for this experiment used was -27.468 m/s^2 , the size of the lattice $5 \times 5 \times 5$ and 4 available materials (2-actuated).	17
3.15	Best so far fitness in body lengths displacement of softbot's center of mass averaged over 10 runs together with the standard deviation error. Local competition is held among the top 20% and in the complete population of each species. The gravity acceleration for this experiment used was -27.468 m/s^2 , the size of the lattice $5 \times 5 \times 5$ and 4 available materials (2-actuated).	19
3.16	Best so far fitness in body lengths displacement of softbot's center of mass averaged over 10 runs together with the standard deviation error. Local competition is held among the top 20% of each species population. The gravity acceleration for this experiment used was -27.468 m/s^2 , the size of the lattice $5 \times 5 \times 5$ and 4 available materials (2-actuated).	20
3.17	Best so far fitness in body lengths displacement of softbot's center of mass averaged over 10 runs together with the standard deviation error. Local competition is held among the complete population of each species. The gravity acceleration for this experiment used was -27.468 m/s^2 , the size of the lattice $5 \times 5 \times 5$ and 4 available materials (2-actuated).	21
3.18	Best so far fitness in body lengths displacement of softbot's center of mass averaged over 10 runs together with the standard deviation error. Local competition is held among the complete population of each species. The gravity acceleration for this experiment used was -27.468 m/s^2 , the size of the lattice $5 \times 5 \times 5$ and 4 available materials (2-actuated).	22

List of Tables

Chapter 1

Introduction

Chapter 2

Related Work

Chapter 3

Results

3.1 Results

3.1.1 Fitness Based Search




FIGURE 3.1: Caption

FIGURE 3.2: Best so far fitness in body lengths displacement of softbot's center of mass from 10 runs for fitness based search. The gravity acceleration for this experiment used was -27.468 m/s^2 , the size of the lattice $10 \times 10 \times 10$ and 4 available materials (2-actuated).




FIGURE 3.3: Best so far fitness in body lengths displacement of softbot's center of mass from 10 runs together with their mean (thick line) for fitness based search. The gravity acceleration for this experiment used was -27.468 m/s^2 , the size of the lattice $10 \times 10 \times 10$ and 4 available materials (2-actuated).

3.1.2 Novelty Search




FIGURE 3.4: Best so far fitness in body lengths displacement of softbot's center of mass from 10 runs for novelty search. The gravity acceleration for this experiment used was -27.468 m/s^2 , the size of the lattice $10 \times 10 \times 10$ and 4 available materials (2-actuated).




FIGURE 3.5: Best so far fitness in body lengths displacement of softbot’s center of mass from 10 runs together with their mean (thick line) for novelty search. The gravity acceleration for this experiment used was -27.468 m/s^2 , the size of the lattice $10 \times 10 \times 10$ and 4 available materials (2-actuated).




FIGURE 3.6: Novelty search - best so far fitness in body lengths displacement of soft-bot's center of mass from 10 runs together with their mean (thick line). The novelty is computed as the average distance from the K -nearest behaviors for $K \in \{1, 2, 5, 10, 20\}$. The gravity acceleration for this experiment used was -27.468 m/s^2 , the size of the lattice $5 \times 5 \times 5$ and 4 available materials (2-actuated).

3.1.3 Novelty-Fitness Based Search Comparison




FIGURE 3.7: Best so far fitness in body lengths displacement of softbot's center of mass averaged over 10 runs together with the standard deviation error. The gravity acceleration for this experiment used was -27.468 m/s^2 , the size of the lattice $5 \times 5 \times 5$ and 4 available materials (2-actuated).

¹Actuated materials penalize fitness:

$$f = (1 - (n_{actuated}/n_{total})^{1.5}) \times disp$$

, where $n_{actuated}$, is the number of actuated voxels, n_{total} total number of voxels and $disp$ the displacement of the softbot's center of mass.




FIGURE 3.8: Fitness of the generation champion (best individual) per generation in body lengths displacement of softbot's center of mass averaged over 10 runs together with the standard deviation error. The gravity acceleration for this experiment used was -27.468 m/s^2 , the size of the lattice $5 \times 5 \times 5$ and 4 available materials (2-actuated).




FIGURE 3.9: Best so far fitness in body lengths displacement of softbot's center of mass averaged over 10 runs together with the standard deviation error. The gravity acceleration for this experiment used was -27.468 m/s^2 , the size of the lattice $10 \times 10 \times 10$ and 4 available materials (2-actuated).




FIGURE 3.10: Distributions of average population fitness per generation over 10 runs. The gravity acceleration for this experiment used was -27.468 m/s^2 , the size of the lattice $5 \times 5 \times 5$ and 4 available materials (2-actuated). Blue (Novelty search), Green (Fitness based search).




FIGURE 3.11: Novel individual's behaviors upto generation averaged over 10 runs. The novelty is computed as the average distance from the 10-nearest behaviors. The gravity acceleration for this experiment used was -27.468 m/s^2 , the size of the lattice $5 \times 5 \times 5$ and 4 available materials (2-actuated). Blue (Novelty search), Green (Fitness based search).




FIGURE 3.12: Best so far fitness in body lengths displacement of softbot's center of mass averaged over 10 runs together with the standard deviation error. The gravity acceleration for this experiment used was -27.468 m/s^2 , the size of the lattice $5 \times 5 \times 5$ and 4 available materials (2-actuated).




FIGURE 3.13: Best so far fitness in body lengths displacement of softbot's center of mass averaged over 10 runs together with the standard deviation error. The gravity acceleration for this experiment used was -27.468 m/s^2 , the size of the lattice $5 \times 5 \times 5$ and 4 available materials (2-actuated).




FIGURE 3.14: Best so far fitness averaged over 10 runs together with the standard deviation error penalizing actuated materials¹. The gravity acceleration for this experiment used was -27.468 m/s^2 , the size of the lattice $5 \times 5 \times 5$ and 4 available materials (2-actuated).

3.1.4 Local competition in fitness based evolution




FIGURE 3.15: Best so far fitness in body lengths displacement of softbot's center of mass averaged over 10 runs together with the standard deviation error. Local competition is held among the top 20% and in the complete population of each species. The gravity acceleration for this experiment used was -27.468 m/s^2 , the size of the lattice $5 \times 5 \times 5$ and 4 available materials (2-actuated).




FIGURE 3.16: Best so far fitness in body lengths displacement of softbot's center of mass averaged over 10 runs together with the standard deviation error. Local competition is held among the top 20% of each species population. The gravity acceleration for this experiment used was -27.468 m/s^2 , the size of the lattice $5 \times 5 \times 5$ and 4 available materials (2-actuated).




FIGURE 3.17: Best so far fitness in body lengths displacement of softbot's center of mass averaged over 10 runs together with the standard deviation error. Local competition is held among the complete population of each species. The gravity acceleration for this experiment used was -27.468 m/s^2 , the size of the lattice $5 \times 5 \times 5$ and 4 available materials (2-actuated).

3.1.5 Local competition in novelty search evolution




FIGURE 3.18: Best so far fitness in body lengths displacement of softbot's center of mass averaged over 10 runs together with the standard deviation error. Local competition is held among the complete population of each species. The gravity acceleration for this experiment used was -27.468 m/s^2 , the size of the lattice $5 \times 5 \times 5$ and 4 available materials (2-actuated).

Appendix A

Appendix Title Here

Write your Appendix content here.

