

# 10907 Pattern Recognition

## Lecturers

Prof. Dr. Ivan Dokmanić (ivan.dokmanic@unibas.ch)

## Tutors

Alexandra Flora Spitzer (alexandra.spitzer@stud.unibas.ch)

Roman Fries (r.fries@unibas.ch)

Cheng Shi (cheng.shi@unibas.ch)

Vinith Kishore (vinith.kishore@unibas.ch)

Valentin Debarnot (valentin.debarnot@unibas.ch)

## Introductory Assignment

This assignment will not contribute to your final grade, but it is a requirement to earn at least 50 percent of the total points in order to be eligible for further. The purpose of this assignment is to get you refreshed on some basic math we will use in the course as well as on Python and Numpy. To submit your solutions, you will need to upload them to Gradescope, which we use throughout this course. Please follow the instructions in Section 3.

## 1 Math

### 1 Linear Algebra

#### Exercise 1 (1 point)

Consider the following system of linear equations,

$$\begin{aligned} 2x - 7y - 9z &= k_1 \\ 3x + 3y + 4z &= k_2 \\ 11x + 2y + 3z &= k_3 \end{aligned}$$

What are the values of  $k_1, k_2, k_3$ , for which the system above has solutions? If we set  $k_1 = k_2 = k_3 = 0$ , what are the possible values of  $x, y$  and  $z$ ?

#### Exercise 2 (Elementary Row and Column operations - 1 point)

Let  $M$  be a block matrix given as

$$M = \begin{bmatrix} I_n & \mathbf{0} \\ \mathbf{0} & AB \end{bmatrix},$$

where  $A \in \mathbb{R}^{s \times n}$ ,  $B \in \mathbb{R}^{n \times t}$  and  $I_n$  is an  $n \times n$  identity matrix. Use elementary row and column operations to transform this matrix into

$$N = \begin{bmatrix} B & I_n \\ \mathbf{0} & A \end{bmatrix}.$$

Provide an explanation of the row and column operations you used.

#### Exercise 3 (Rank of a Matrix - 2 points)

Let  $M$  and  $N$  be as in the previous exercise,  $M = \begin{bmatrix} I_n & \mathbf{0} \\ \mathbf{0} & AB \end{bmatrix}$  and  $N = \begin{bmatrix} B & I_n \\ \mathbf{0} & A \end{bmatrix}$ , with dimensions as before.



1. What conclusions can you draw regarding the rank of matrix  $\mathbf{N}$  from the ranks of matrices  $\mathbf{A}$  and  $\mathbf{B}$ ?
2. Using the results from part 1 of the exercise and the previous exercise, prove the inequality:  
 $\text{rank}(\mathbf{A}) + \text{rank}(\mathbf{B}) - \text{rank}(\mathbf{AB}) \leq n$ .
3. If  $\text{rank}(\mathbf{A}) = \text{rank}(\mathbf{AB}) = s$ , what is the rank of matrix  $\mathbf{B}$ ?

#### Exercise 4 (Eigenvalues and Positive Definiteness - 1 point)

Let  $\mathbf{A} \in \mathbb{R}^{s \times s}$  be an  $s \times s$  real matrix such that  $\mathbf{x}^T \mathbf{A} \mathbf{x} \geq 0$  for all  $\mathbf{x} \in \mathbb{R}^s$ . Show that  $\mathbf{A}$  does not have any negative eigenvalues.

## 2 Probability

#### Exercise 5 (1 point)

For the system of linear equations given below:

$$\begin{aligned} 2X - 7Y + 9Z &= K_1 \\ 3X + 3Y + 4Z &= K_2 \\ 5X + 2Y + 5Z &= K_3 \end{aligned}$$

The random variables  $X, Y, Z$  are independent and normally distributed such that  $X \sim \mathcal{N}(0, 1)$ ,  $Y \sim \mathcal{N}(1, 4)$  and  $Z \sim \mathcal{N}(0, 9)$ .

- What are the distributions of  $K_1, K_2$  and  $K_3$ ?
- What is the joint probability density function of  $K_1, K_2$  and  $K_3$ ,  $p(K_1 = k_1, K_2 = k_2, K_3 = k_3)$ ?

Recall that a normally distributed random variable  $X$  with mean  $\mu$  and variance  $\sigma^2$  (written  $X \sim \mathcal{N}(\mu, \sigma^2)$ ) has the probability density function (pdf)

$$p_X(x) = \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(-\frac{(x-\mu)^2}{2\sigma^2}\right)$$

so that for some  $a, b$  such that  $a \leq b$  we have

$$\mathbb{P}[a \leq X \leq b] = \int_a^b p_X(x) dx.$$

#### Exercise 6 (Sum of Bernoulli Distribution - 2 points)

Alice and Bob decide to play a game. They toss a biased coin  $n$  times. If it comes up heads at least  $n/2$  times, Bob wins; otherwise Alice wins. The total number of heads is

$$S = \sum_{i=1}^n X_i$$

where  $X_i$  is an independent Bernoulli random variable with mean  $p$ , that is,

$$\mathbb{P}(X_i = 1) = p, \quad \mathbb{P}(X_i = 0) = 1 - p.$$

Assume that  $p = 0.52$ .

1. What is the law of the random variable  $S$ ? Give its name, its probability mass function (p.m.f.) and its mean.

2. Bob only wants to play if he has a 90% chance of winning.

For a given  $n$ , what is the expected number of heads?

Compute the probability that Bob wins when  $n = 2$ .

Can you figure out for which values of  $n$  Bob will participate in the game?

*Hint:* One way to do this is by using Hoeffding's inequality: let  $Z_i$  be independent and identically distributed random variables such that  $0 \leq Z_i \leq 1$ , then

$$\mathbb{P}\left(\frac{1}{n} \sum_{i=1}^n Z_i - \mathbb{E}(Z_i) \geq t\right) \leq \exp(-nt^2).$$

3. Alice and Bob continue the game for a long, long time. What can you say about the value or distribution of  $S$  as  $n \rightarrow \infty$ ? What about the value or distribution of  $\frac{S}{n}$ ? And what about the value or distribution of  $\frac{S - np}{\sqrt{n}}$ ?

### Exercise 7 (Expectations - 1 point)

The expectation of a non-negative random variable  $X$  with probability density function  $f_X(x)$  is

$$\mathbb{E}(X) = \int_0^\infty x f_X(x) dx.$$

Show that an alternative way to compute it is by the following useful identity:

$$\mathbb{E}(X) = \int_0^\infty \mathbb{P}(X > t) dt.$$

## 2 Coding

To get your hands dirty with Python, you are asked to write some basic array operations related to linear algebra and probability. Specifically, you need to complete the blank functions in `sample.py`.

### Exercise 8 (Basic Computation — 2 points)

- Write a function `check_span` that takes as the inputs a vector  $\mathbf{y}$  and a matrix  $\mathbf{A} = [\mathbf{a}_1, \dots, \mathbf{a}_n]$ , where  $\mathbf{a}_i$ 's are the column vectors of the matrix. The function outputs a Boolean variable. The output is `True` if  $\mathbf{y}$  is in the span of column vectors in  $\mathbf{A}$ . Otherwise, the output is `False`.
- Write a function `estimate_mean_and_covariance` to compute the mean and the unbiased covariance matrix of a set of vectors  $\{\mathbf{x}_1, \dots, \mathbf{x}_n\}$ . The set of vectors are passed to the function as a matrix  $\mathbf{X} = [\mathbf{x}_1, \dots, \mathbf{x}_n]$ . The unbiased covariance matrix is defined as:

$$\mathbf{Q} = \frac{1}{n-1} \sum_{i=1}^n (\mathbf{x}_i - \bar{\mathbf{x}})(\mathbf{x}_i - \bar{\mathbf{x}})^T$$

where  $\bar{\mathbf{x}} = \frac{1}{n} \sum_{i=1}^n \mathbf{x}_i$ .

## 3 Submitting the Exercises

We use Gradescope to submit and evaluate the assignments. You should have received an invitation to the Gradescope course page. Go to the course page 10907 Pattern Recognition, you will see two assignment items, Sample Assignment: Coding and Sample Assignment: Math.

- For the Math part of this assignment, you upload the solution as a *single* **.pdf** file in the **Sample Assignment: Math**. For each exercise, you must to specify which pages containing the answer.
- For the Coding part of this assignment, the file **sample.py** contains the functions to be completed. You upload the completed file to the **Sample Assignment: Coding**.