



# **Geometry through Linear Algebra**



## G V V Sharma\*

1

4

9

### **CONTENTS**

1	Triangle	<b>Exercises</b>

#### 2 **Quadrilateral Exercises**

#### 3 Circle Exercises

#### 4 Miscellaneous Exercises

Abstract—This book provides a vector approach to geometry. The content and exercises are based on NCERT textbooks from Class 6-12.

### 1 Triangle Exercises

- 1.1. Each angle of an equilateral triangle is of 60°. **Solution:**
- 1.2. Triangles on the same base (or equal bases) and between the same parallels are equal in area.
- 1.3. Triangles on the same base (or equal bases) and having equal areas lie between the same parallels.
- 1.4. In  $\triangle ABC$ , D, E and F are respectively the midpoints of sides AB, BC and CA. Show that  $\triangle ABC$  is divided into four congruent triangles by joining D, E and F.

\*The author is with the Department of Electrical Engineering, Indian Institute of Technology, Hyderabad 502285 India e-mail: gadepall@iith.ac.in. All content in this manual is released under GNU 1.17. In an isosceles  $\triangle ABC$  with AB = AC, D and GPL. Free and open source.

- 1.5. The line-segment joining the mid-points of any two sides of a triangle is parallel to the third side and is half of it.
- 1.6. A line through the mid-point of a side of a triangle parallel to another side bisects the third side.
- 1.7. ABC is a triangle right angled at C. A line through the mid-point M of hypotenuse AB and parallel to BC intersects AC at D. Show that (i) D is the mid-point of AC (ii)  $MD \perp AC$ (iii)  $CM = MA = \frac{1}{2}AB$
- 1.8. Sides opposite to equal angles of a triangle are equal.
- 1.9. Each angle of an equilateral triangle is of 60°.
- 1.10. Using cosine formula in an equilateral  $\triangle$ , show that  $\cos 60^\circ = \frac{1}{2}$ .
- 1.11. Using (??), show that  $\sin 60^{\circ} = \frac{\sqrt{3}}{2}$ . 1.12. Find  $\sin 30^{\circ}$  and  $\sin 30^{\circ}$  using (??).
- 1.13. Triangles on the same base (or equal bases) and between the same parallels are equal in area.
- 1.14. Triangles on the same base (or equal bases) and having equal areas lie between the same parallels.
- 1.15. In  $\triangle ABC$ , the bisector AD of  $\angle A$  is perpendicular to side BC. Show that AB = AC and  $\triangle ABC$ is isosceles.
- 1.16. E and F are respectively the mid-points of equal sides AB and AC of  $\triangle ABC$ . Show that BF = CE.

E are points on BC such that BE = CD. Show that AD = AE.

- 1.18. AB is a line-segment. P and Q are points on 1.29. In an isosceles  $\triangle ABC$ , with AB = AC, the opposite sides of AB such that each of them is equidistant from the points A and B. Show that the line PQ is the perpendicular bisector
- 1.19. P is a point equidistant from two lines l and mintersecting at point A. Show that the line APbisects the angle between them.
- 1.20. D is a point on side BC of  $\triangle ABC$  such that AD = AC. Show that AB > AD
- 1.21. AB is a line segment and line l is its perpendicular bisector. If a point P lies on l, show that P is equidistant from A and B.
- 1.22. Line-segment AB is parallel to another linesegment CD. O is the mid-point of AD. Show that
  - a)  $\triangle AOB \cong \triangle DOC$
  - b) O is also the mid-point of BC.
- 1.23. In quadrilateral ACBD, AC = AD and ABbisects  $\angle A$ . Show that  $\triangle ABC \cong \triangle ABD$ . What can you say about BC and BD?
- 1.24. ABCD is a quadrilateral in which AD = BCand  $\angle DAB = \angle CBA$ . Prove that
  - a)  $\triangle ABD \cong \triangle BAC$
  - b) BD = AC
  - c)  $\angle ABD = \angle BAC$ .
- 1.25. l and m are two parallel lines intersected by another pair of parallel lines p and q to form the quadrilateral ABCD. Show that  $\triangle ABC \cong$  $\triangle CDA$ .
- 1.26. Line l is the bisector of  $\angle A$  and B is any point on *l. BP* and *BQ* are perpendiculars from *B* to the arms of  $\angle A$ . Show that:
  - a)  $\triangle APB \cong \triangle AQB$
  - b) BP = BQ or B is equidistant from the arms of  $\angle A$ .
- 1.27. ABCE is a quadrilateral and D is a point on BC such that, AC = AE, AB = AD and  $\angle BAD =$  $\angle EAC$ . Show that BC = DE.
- 1.28. In right triangle ABC, right angled at C, M is the mid-point of hypotenuse AB. C is joined to M and produced to a point D such that DM =CM. Point D is joined to point B. Show that:
  - a)  $\triangle AMC \cong \triangle BMD$
  - b)  $\angle DBC$  is a right angle.
  - c)  $\triangle DBC \cong \triangle ACB$

- d)  $CM = \frac{1}{2}AB$ See Fig. ??.
- bisectors of  $\angle B$  and  $\angle C$  intersect each other at O. Join A to O. Show that :
  - a) OB = OC
  - b) AO bisects  $\angle A$
- 1.30. In  $\triangle ABC$ , AD is the perpendicular bisector of BC. Show that  $\triangle ABC$  is an isosceles triangle in which AB = AC.
- 1.31. ABC is an isosceles triangle in which altitudes BE and CF are drawn to equal sides AC and AB respectively. Show that these altitudes are
- 1.32. ABC is a triangle in which altitudes BE and CF to sides AC and AB are equal. Show that
  - a)  $\triangle ABE \cong \triangle ACF$
  - b) AB = AC, i.e., ABC is an isosceles triangle.
- 1.33. ABC and DBC are two isosceles triangles on the same base BC. Show that  $\angle ABD = \angle ACD$ .
- 1.34.  $\triangle ABC$  and  $\triangle DBC$  are two isosceles triangles on the same base BC and vertices A and D are on the same side of BC. If AD is extended to intersect BC at P, show that
  - a)  $\triangle ABD \cong \triangle ACD$
  - b)  $\triangle ABP \cong \triangle ACP$
  - c) AP bisects  $\angle A$  as well as  $\angle D$ .
  - d) AP is the perpendicular bisector of BC.
- 1.35. AD is an altitude of an isosceles  $\triangle ABC$  in which AB = AC. Show that
  - a) AD bisects BC
  - b) AD bisects  $\angle A$ .
- 1.36. Two sides AB and BC and median AM of one triangle ABC are respectively equal to sides PQ and QR and median PN of  $\triangle PQR$ . Show that:
  - a)  $\triangle ABM \cong \triangle PQN$
  - b)  $\triangle ABC \cong \triangle PQR$
- 1.37. BE and CF are two equal altitudes of a triangle ABC. Using RHS congruence rule, prove that the triangle ABC is isosceles.
- 1.38. ABC is an isosceles triangle with AB = AC. Draw  $AP \perp BC$  to show that  $\angle B = \angle C$ .
- 1.39.  $\triangle ABC$  is an isosceles triangle in which AB =AC. Side BA is produced to D such that AD =AB. Show that  $\angle BCD$  is a right angle.
- 1.40. ABC is a right angled triangle in which  $\angle A =$ 90° and AB = AC. Find  $\angle B$  and  $\angle C$ .
- 1.41. Show that in a right angled triangle, the hy-

- potenuse is the longest side.
- 1.42. Sides AB and AC of  $\triangle ABC$  are extended to
- 1.43. Line segments AD and BC intersect at O and  $\angle D$ . Show that AD < BC.
- 1.44. AB and CD are respectively the smallest and that  $\angle A > \angle C$  and  $\angle B > \angle D$ .
- 1.45. In  $\triangle PQR$ , PR > PQ and PS bisects  $\angle QPR$ . Prove that  $\angle PSR > \angle PSQ$ .
- 1.46. Show that of all line segments drawn from a given point not on it, the perpendicular line segment is the shortest.
- 1.47. ABCD is a trapezium with  $AB \parallel DC$ . E and F are points on non-parallel sides AD and BC respectively such that EF is parallel to AB. Show that  $\frac{AE}{ED} = \frac{BF}{FC}$ .
- 1.48. ST is a line joining two points on PQ and PR in  $\triangle PQR$ . If  $\frac{PS}{SQ} = \frac{PT}{TR}$  and  $\angle PST = \angle PRQ$ , 1.63. ABC and AMP are two right triangles, right prove that PQR is an isosceles triangle.
- 1.49. If  $LM \parallel CB$  and  $LN \parallel CD$ , prove that  $\frac{AM}{AB} = \frac{AN}{AD}$ . 1.50. D is a point on AB and E, F are points on BC
- such that  $DE \parallel AC$  and  $DF \parallel AE$ . Prove that  $\frac{BF}{FE} = \frac{BE}{EC}$ .
- 1.51. O is a point in the interior of  $\triangle ABC$ . D is a point on OA. If  $DE \parallel OB$  and  $DF \parallel OC$ . Show that  $EF \parallel BC$ .
- 1.52. O is a point in the interior of  $\triangle PQR$ . A, BandC are points on OP, OQ and OR respectively such that  $AB \parallel PQ$  and  $AC \parallel PR$ . Show that  $BC \parallel$ QR.
- 1.53. ABCD is a trapezium in which  $AB \parallel DC$  and its diagonals intersect each other at the point O. Show that  $\frac{AO}{BO} = \frac{CO}{DO}$
- 1.54. The diagonals of a quadrilateral ABCD intersect each other at the point O such that  $\frac{AO}{BO} = \frac{CO}{DO}$ . Show that ABCD is a trapezium. 1.55.  $PQ \parallel RS$  and PS intersects QR at O. Show
- that  $\triangle OPQ \sim \triangle ORS$ .
- 1.56. CM and RN are respectively the medians of  $\triangle ABC$  and  $\triangle PQR$ . If  $\triangle ABC \sim \triangle PQR$ , prove that
  - a)  $\triangle AMC \sim \triangle PNR$

  - b)  $\frac{CM}{RN} = \frac{AB}{PQ}$ c)  $\triangle CMB \sim \triangle RNQ$
- 1.57. Diagonals AC and BD of a trapezium ABCD with  $AB \parallel DC$  intersect each other at the point

- O. Using a similarity criterion for two triangles,
- Sides AB and AC of  $\triangle ABC$  are extended to show that  $\frac{OA}{OC} = \frac{OB}{OD}$  points P and Q respectively. Also,  $\angle PBC < 1.58$ . In  $\triangle PQR$ , QP is extended to T and S is a point  $\angle QCB$ . Show that AC > AB. on QR such that  $\frac{QR}{QS} = \frac{QT}{PR}$ . If  $\angle PRQ = \angle PQS$ , show that that  $\triangle PQS \sim \triangle TQR$ .
- form  $\triangle OAB$  and  $\triangle ODC$ .  $\angle B < \angle A$  and  $\angle C < 1.59$ . S and T are points on sides PR and QR of  $\triangle PQR$  such that  $\angle P = \angle RTS$ . Show that  $\triangle RPQ \sim \triangle RTS$ .
- longest sides of a quadrilateral ABCD. Show 1.60. In  $\triangle ABC$ , D and E are points on the sides AB and AC respectively. If  $\triangle ABE \cong \triangle ACD$ , show that  $\triangle ADE \sim \triangle ABC$ .
  - 1.61. Altitudes AD and CE of  $\triangle ABC$  intersect each other at the point P. Show that:
    - a)  $\triangle AEP \sim \triangle CDP$
    - b)  $\triangle ABD \sim \triangle CBE$
    - c)  $\triangle AEP \sim \triangle ADB$
    - d)  $\triangle PDC \sim \triangle BEC$
  - 1.62. E is a point on the side AD produced of a parallelogram ABCD and BE intersects CD at *F*. Show that  $\triangle ABE \sim \triangle CFB$ .
  - angled at B and M respectively. M lies on AC and AB is extended to meet P. Prove that:
    - a)  $\triangle ABC \sim \triangle AMP$
    - b)  $\frac{CA}{PA} = \frac{BC}{MP}$
  - 1.64. CD and GH are respectively the bisectors of  $\angle ACB$  and  $\angle EGF$  such that D and H lie on sides AB and FE of  $\triangle ABC$  and  $\triangle EFG$ respectively. If  $\triangle ABC \sim \triangle FEG$ , show that:

  - 1.65.  $\frac{CD}{GH} = \frac{AC}{FG}$ 1.66.  $\triangle DCB \sim \triangle HGE$
  - 1.67.  $\triangle DCA \sim \triangle HGF$
  - 1.68. E is a point on side CB produced of an isosceles  $\triangle ABC$  with AB = AC. If  $AD \perp BC$ and  $EF \perp AC$ , prove that  $\triangle ABD \sim \triangle ECF$ .
  - 1.69. Sides AB and BC and median AD of a  $\triangle ABC$ are respectively proportional to sides PQ and QR and median PM of  $\triangle PQR$ . Show that  $\triangle ABC \sim \triangle POR$ .
  - 1.70. D is a point on the side BC of a  $\triangle ABC$  such that  $\angle ADC = \angle BAC$ . Show that  $CA^2 = CB.CD$ .
  - 1.71. Sides AB and AC and median AD of a  $\triangle ABC$ are respectively proportional to sides PQ and PR and median PM of another  $\triangle PQR$ . Show that  $\triangle ABC \sim \triangle PQR$ .
  - 1.72. If AD and PM are medians of  $\triangle sABC$  and PQR, respectively where  $\triangle ABC \sim \triangle PQR$ , prove that  $\frac{AB}{PQ} = \frac{\dot{A}D}{PM}$ 1.73. The line segment XY is parallel to side AC of

- $\triangle ABC$  and it divides the triangle into two parts of equal areas. Find the ratio  $\frac{AX}{AB}$
- 1.74. Diagonals of a trapezium ABCD with  $AB \parallel DC$ intersect each other at the point O. If AB =COD.
- 1.75. ABC and DBC are two triangles on the same base BC. If AD intersects BC at O, show that  $\frac{ar(ABC)}{ar(DBC)} = \frac{AO}{DO}.$ 1.76. If the areas of two similar triangles are equal,
- prove that they are congruent.
- 1.77. D, E and F are respectively the mid-points of sides AB, BC and CA of  $\triangle ABC$ . Find the ratio of the areas of  $\triangle DEF$  and  $\triangle ABC$ .
- 1.78. Prove that the ratio of the areas of two similar triangles is equal to the square of the ratio of their corresponding medians.
- 1.79. Prove that the area of an equilateral triangle 1.96. In an equilateral triangle, prove that three times described on one side of a square is equal to half the area of the equilateral triangle described on one of its diagonals.
- 1.80. ABC and BDE are two equilateral triangles such that D is the mid-point of BC. Find the ratio of the areas of triangles ABC and BDE.
- 1.81. The sides of two similar triangles are in the ratio 4: 9. Find the ratio the area of these triangles are in the ratio
- 1.82. In  $\triangle ABC$ ,  $\angle ACB = 90^{\circ}$  and  $CD \perp AB$ . Prove that  $\frac{BC^2}{AC^2} = \frac{BD}{AD}$ . 1.83. In  $\triangle ABC$ , if  $AD \perp BC$ , prove that  $AB^2 + CD^2 =$
- $BD^2 + AC^2$ .
- angled at A. Prove that  $4(BL^2 + CM^2) = 5BC^2$
- that  $OB^2 + OD^2 = OA^2 + OC^2$ .
- 1.86. PQR is a triangle right angled at P and M is a point on QR such that  $PM \perp QR$ . Show that  $PM^2 = QM.MR.$
- 1.87.  $\overrightarrow{ABD}$  is a triangle right angled at A and  $\overrightarrow{AC} \perp_{1.102}$ . Prove that the sum of the squares of the diag-BD. Show that
  - a)  $AB^2 = BC.BD$
  - b)  $AC^2 = BC.DC$
  - c)  $AD^2 = BD.CD$
- 1.88. ABC is an isosceles triangle right angled at C. Prove that  $AB^2 = 2AC^2$ .
- 1.89. ABC is an isosceles triangle with AC = BC. If  $AB^2 = 2AC^2$ , prove that ABC is a right triangle.
- 1.90. ABC is an equilateral triangle of side 2a. Find

- each of its altitudes.
- 1.91. Prove that the sum of the squares of the sides of a rhombus is equal to the sum of the squares of its diagonals.
- 2CD, find the ratio of the areas of  $\triangle sAOB$  and 1.92. O is a point in the interior of a  $\triangle ABC$ , OD  $\perp$  $BC, OE \perp AC and OF \perp AB$ . Show that
  - a)  $OA^2 + OB^2 + BD^2 OD2 OE2 OF2 = AF^2 +$  $BD^2 + CE^2$ .
  - b)  $AF^2 + BD^2 + CE^2 = AE^2 + CD^2 + BF^2$ .
  - 1.93. D and E are points on the sides CA and CB respectively of a  $\triangle ABC$  right angled at C. Prove that  $AE^2 + BD^2 = AB^2 + DE^2$ .
  - 1.94. The perpendicular from A on side BC of a  $\triangle ABC$  intersects BC at D such that DB = 3CD. Prove that  $2AB^2 = 2AC^2 + BC^2$ .
  - 1.95. In an equilateral  $\triangle ABC$ , D is a point on side BC such that  $BD = \frac{1}{2}BC$ . Prove that  $9AD^2 = 7AB^2$ .
  - the square of one side is equal to four times the square of one of its altitudes.
  - 1.97. PS is the bisector of  $\angle QPR$  of  $\triangle PQR$ . Prove that  $\frac{QS}{SR} = \frac{PQ}{PR}$ 1.98. *D* is a point on hypotenuse *AC* of  $\triangle ABC$ , such
  - that  $BD \perp AC, DM \perp BC$  and  $DN \perp AB$ . Prove that:
    - a)  $DM2 = DN \cdot MC$
    - b)  $DN2 = DM \cdot AN$
  - 1.99. ABC is a triangle in which  $\angle ABC > 90^{\circ}$  and  $AD \perp CB$  produced. Prove that  $AC^2 = AB^2 +$  $BC^2 + 2BC.BD.$
- 1.84. BL and CM are medians of a  $\triangle ABC$  right 1.100. ABC is a triangle in which  $\angle ABC < 90^{\circ}$ and  $AD \perp BC$ . Prove that  $AC^2 = AB^2 +$  $BC^2-2BC.BD.$
- 1.85. O is any point inside a rectangle ABCD. Prove 1.101. AD is a median of a  $\triangle ABC$  and  $AM \perp BC$ .
  - a)  $AC^2 = AD^2 + BC.DM + \left(\frac{BC}{2}\right)^2$ b)  $AB^2 = AD^2 BC.DM + \left(\frac{BC}{2}\right)^2$ c)  $AC^2 + AB^2 = 2AD^2 + \frac{1}{2}BC^2$

  - onals of parallelogram is equal to the sum of the squares of its sides.
  - 1.103. D is a point on side BC of  $\triangle ABC$  such that  $\frac{BD}{CD}\frac{AB}{AC}$ . Prove that AD is the bisector of  $\angle BAC$ .

### 2 Quadrilateral Exercises

2.1. Parallelograms on the same base (or equal bases) and between the same parallels are equal in area.

- 2.2. If a parallelogram and a triangle are on the same base and between the same parallels, then 2.14. In  $\triangle ABC$  and  $\triangle DEF, AB = DE, AB$ area of the triangle is half the area of the parallelogram.
- 2.3. The quadrilateral formed by joining the midpoints of the sides of a quadrilateral, in order, is a parallelogram.
- 2.4. Two parallel lines 1 and m are intersected by a transversal p. Show that the quadrilateral formed by the bisectors of interior angles is a rectangle.
- 2.5. Show that the bisectors of angles of a parallelogram form a rectangle.
- 2.6. ABCD is a parallelogram in which P and Q are mid-points of opposite sides AB and CD. If AQ intersects DP at S and BQ intersects CP at R, show that:
  - a) APCQ is a parallelogram.
  - b) DPBQ is a parallelogram.
  - c) PS QR is a parallelogram.
- 2.7. l, m and n are three parallel lines intersected by transversals p and q such that l, m and n cut off equal intercepts AB and BC on p. Show that l, m and n cut off equal intercepts DE and EFon q also.
- 2.8. Parallelograms on the same base (or equal bases) and between the same parallels are equal
- base and the corresponding altitude.
- 2.10. Parallelograms on the same base (or equal bases) and having equal areas lie between the same parallels.
- 2.11. If a parallelogram and a triangle are on the same base and between the same parallels, then area of the triangle is half the area of the parallelogram.
- 2.12. In parallelogram ABCD, two points P and Qare taken on diagonal BD such that DP = BQ. show that
  - a)  $\triangle APD \cong \triangle COB$
  - b) AP = CQ
  - c)  $\triangle AQB \cong \triangle CPD$
  - d) AQ = CP
  - e) APCQ is a parallelogram
- 2.13. ABCD is a parallelogram and AP and CQ are perpendiculars from vertices A and C on diagonal BD. Show that
  - a)  $\triangle APB \cong \triangle CQD$

- b) AP = CO
- DE, BC = EF and  $BC \parallel EF$ . Vertices A, Band C are joined to vertices D, E and F respectively. Show that
  - a) quadrilateral ABED is a parallelogram
  - b) quadrilateral *BEFC* is a parallelogram
  - c)  $AD \parallel CF$  and AD = CF
  - d) quadrilateral ACFD is a parallelogram
  - e) AC = DF
  - f)  $\triangle ABC \cong \triangle DEF$ .
- 2.15. ABCD is a trapezium in which  $AB \parallel CD$  and AD = BC. Show that
  - a)  $\angle A = \angle B$
  - b)  $\angle C = \angle D$
  - c)  $\triangle ABC \cong \triangle BAD$
  - d) diagonal AC = diagonal BD
- 2.16. ABCD is a quadrilateral in which P, Q, R and S are mid-points of the sides AB, BC, CD and DA AC is a diagonal. Show that
  - a)  $SR \parallel AC$  and  $SR = \frac{1}{2}AC$
  - b) PQ = SR
  - c) PQRS is a parallelogram.
- 2.17. ABCD is a rhombus and P, Q, R and S are the mid-points of the sides AB, BC, CD and DA respectively. Show that the quadrilateral PQRS is a rectangle.
- 2.9. Area of a parallelogram is the product of its 2.18. ABCD is a rectangle and P, Q, R and S are mid-points of the sides AB, BC, CD and DA respectively. Show that the quadrilateral PQRS is a rhombus.
  - 2.19. ABCD is a trapezium in which  $AB \parallel DC, BD$ is a diagonal and E is the mid-point of AD. A line is drawn through  $E \parallel AB$  intersecting BCat F. Show that F is the mid-point of BC.
  - 2.20. In a parallelogram ABCD, E and F are the mid-points of sides AB and CD respectively. Show that the line segments AF and EC trisect the diagonal BD.
  - 2.21. Show that the line segments joining the midpoints of the opposite sides of a quadrilateral bisect each other.
  - 2.22. ABCD is a parallelogram in which P and Q are mid-points of opposite sides AB and CD. If AQ intersects DP at S and BQ intersects CPat R, show that:
    - a) APCQ is a parallelogram.
    - b) *DPBQ* is a parallelogram.

- c) PS QR is a parallelogram.
- 2.23. *l, m* and *n* are three parallel lines intersected by transversals p and q such that l, m and n cut off 2.36. ABC and ABD are two triangles on the same equal intercepts AB and BC on p. Show that l, m and n cut off equal intercepts DE and EFon q also.
- 2.24. Diagonal AC of a parallelogram ABCD bisects  $\angle A$  . show that
  - a) it bisects  $\angle C$  also,
  - b) ABCD is a rhombus.
- 2.25. ABCD is a rhombus. Show that diagonal AC bisects  $\angle A$  as well as  $\angle C$  and diagonal BD bisects  $\angle B$  as well as  $\angle D$ .
- 2.26. ABCD is a rectangle in which diagonal AC bisects  $\angle A$  as well as  $\angle C$ . Show that
  - a) ABCD is a square
  - b) diagonal BD bisects  $\angle B$  as well as  $\angle D$ .
- 2.27. If E, F, G and H are respectively the mid-points of the sides of a parallelogram ABCD, show 2.40. XY is a line parallel to side BC of a  $\triangle ABC$ . If that

$$ar(EFGH) = \frac{1}{2}ar(ABCD). \qquad (2.27.1)$$

- 2.28. P and Q are any two points lying on the sides DC and AD respectively of a parallelogram ABCD. Show that ar(APB) = ar(BQC).
- 2.29. P is a point in the interior of a parallelogram 2.42. Diagonals AC and BD of a trapezium ABCD ABCD. Show that
  - a)  $ar(APB) + ar(PCD) = \frac{1}{2}ar(ABCD)$
  - b) ar(APD) + ar(PBC) = ar(APB) + ar(PCD)
- 2.30. PQRS and ABRS are parallelograms and X is any point on side BR. show that
  - a) ar(PQRS) = ar(ABRS)
  - b)  $ar(AXS) = \frac{1}{2}ar(PQRS)$
- 2.31. A farmer was having a field in the form of a 2.44. A villager Itwaari has a plot of land of the parallelogram *PQRS*. She took any point A on RS and joined it to points P and Q. In how many parts the fields is divided? What are the shapes of these parts? The farmer wants to sow wheat and pulses in equal portions of the field separately. How should she do it?
- 2.32. ABCD is a quadrilateral and  $BE \parallel AC$  and also BE meets DC produced at E. Show that area of  $\triangle ADE$  is equal to the area of the quadrilateral ABCD.
- 2.33. E is any point on median AD of a  $\triangle ABC$ . Show that ar(ABE) = ar(ACE).
- 2.34. In a  $\triangle ABC$ , E is the mid-point of median AD. 2.46. AP  $\parallel BQ \parallel CR$ . Prove that ar(AQC) =Show that  $ar(BED) = \frac{1}{4}ar(ABC)$ .

- 2.35. Show that the diagonals of a parallelogram divide it into four triangles of equal area.
- base AB. If line- segment CD is bisected by AB at O, show that ar(ABC) = ar(ABD).
- 2.37. D, E and F are respectively the mid-points of the sides BC, CA and AB of a  $\triangle ABC$ . show that
  - a) BDEF is a parallelogram.
  - b)  $ar(BDEF) = \frac{1}{2}ar(ABC)$
- 2.38. Diagonals AC and BD of quadrilateral ABCD intersect at O such that OB = OD. If AB = CD, then show that
  - a) ar(DOC) = ar(AOB)
  - b) ar(DCB) = ar(ACB)
  - c)  $ar(DEF) = \frac{1}{4}ar(ABC)$
- 2.39. D and E are points on sides AB and AC respectively of  $\triangle ABC$  such that ar(DBC) = ar(EBC). Prove that  $DE \parallel BC$ .
- $BE \parallel AC$  and  $CF \parallel AB$  meet XY at E and F respectively, show that ar(ABE) = ar(ACF).
- (2.27.1) 2.41. The side AB of a parallelogram ABCD is produced to any point P. A line through A and parallel to CP meets CB produced at Q and then parallelogram *PBQR* is completed. Show that ar(ABCD) = ar(PBQR).
  - with  $AB \parallel DC$  intersect each other at O. Prove that ar(AOD) = ar(BOC).
  - 2.43. ABCDE is a pentagon. A line through B parallel to AC meets DC produced at F. Show that
    - a) ar(ACB) = ar(ACF)
    - b) ar(AEDF) = ar(ABCDE).
  - shape of a quadrilateral. The Gram Panchayat of the village decided to take over some portion of his plot from one of the corners to construct a Health Centre. Itwaari agrees to the above proposal with the condition that he should be given equal amount of land in lieu of his land adjoining his plot so as to form a triangular plot. Explain how this proposal will be implemented.
  - 2.45. ABCD is a trapezium with  $AB \parallel DC$ . A line parallel to AC intersects AB at X and BC at Y. Prove that ar(ADX) = ar(ACY).
  - ar(PBR).

- 2.47. Diagonals AC and BD of a quadrilateral ABCD intersect at O in such a way that ar(AOD) = ar(BOC). Prove that ABCD is a trapezium.
- 2.48.  $AB \parallel DC \parallel RP$ . ar(DRC) = ar(DPC) and ar(BDP) = ar(ARC). Show that both the quadrilaterals ABCD and DCPR are trapeziums
- 2.49. Parallelogram *ABCD* and rectangle *ABEF* are on the same base *AB* and have equal areas. Show that the perimeter of the parallelogram is greater than that of the rectangle.
- 2.50. In  $\triangle ABC$ , D and E are two points on BC such that BD = DE = EC. Show that ar(ABD) = ar(ADE) = ar(AEC).
- 2.51. ABCD, DCFE and ABFE are parallelograms. Show that ar(ADE) = ar(BCF).
- 2.52. ABCD is a parallelogram and BC is produced to a point Q such that AD = CQ. If AQ intersect DC at P, show that ar(BPC) = ar(DPQ). ABC and BDE are two equilateral triangles such that D is the mid-point of BC. If AE intersects BC at P, show that
  - a)  $ar(BDE) = \frac{1}{4}ar(ABC)$
  - b)  $ar(BDE) = \frac{7}{2}ar(BAE)$
  - c)  $ar(ABC) = \tilde{2}ar(BEC)$
  - d) ar(BFE) = ar(AFD)
  - e) ar(BFE) = 2ar(FED)
  - f)  $ar(FED) = \frac{1}{8}ar(AFC)$
- 2.53. Diagonals AC and BD of a quadrilateral ABCD intersect each other at P. Show that  $ar(APB) \times ar(CPD) = ar(APD) \times ar(BPC)$ .
- 2.54. P and Q are respectively the mid-points of sides AB and BC of a  $\triangle ABC$  and R is the mid-point of AP, show that
  - a)  $ar(PRQ) = \frac{1}{2}ar(ARC)$
  - b) ar(PBQ) = ar(ARC)
  - c)  $ar(RQC) = \frac{3}{8}ar(ABC)$
- 2.55. ABC is a right triangle right angled at A. BCED, ACFG and ABMN are squares on the sides BC, CA and AB respectively. Line segment  $AX \perp DE$  meets BC at Y. Show that
  - a)  $\triangle MBC \cong \triangle ABD$
  - b) ar(BYXD) = ar(ABMN)
  - c) ar(CYXE) = 2ar(FCB)
  - d) ar(BYXD) = 2ar(MBC)
  - e)  $\triangle FCB \cong \triangle ACE$
  - f) ar(CYXE) = ar(ACFG)
  - g) ar(BCED) = ar(ABMN) + ar(ACFG)
- 2.56. L is a point on the diagonal AC of quadrilateral

- *ABCD*. If LM CB and LN CD, prove that  $\frac{AM}{AB} = \frac{AN}{AD}$  2.57. The angles of quadrilateral are in the ratio 3 : 5
- 2.57. The angles of quadrilateral are in the ratio 3:5: 9:13. Find all the angles of the quadrilateral.

### 3 Circle Exercises

- 3.1. Equal chords of a circle (or of congruent circles) subtend equal angles at the centre.
- 3.2. If the angles subtended by two chords of a circle (or of congruent circles) at the centre (corresponding centres) are equal, the chords are equal.
- 3.3. The perpendicular from the centre of a circle to a chord bisects the chord.
- 3.4. The line drawn through the centre of a circle to bisect a chord is perpendicular to the chord.
- 3.5. There is one and only one circle passing through three non-collinear points.
- 3.6. Equal chords of a circle (or of congruent circles) are equidistant from the centre (or corresponding centres).
- 3.7. Chords equidistant from the centre (or corresponding centres) of a circle (or of congruent circles) are equal.
- 3.8. If two arcs of a circle are congruent, then their corresponding chords are equal and conversely if two chords of a circle are equal, then their corresponding arcs (minor, major) are congruent.
- 3.9. Congruent arcs of a circle subtend equal angles at the centre.
- 3.10. The angle subtended by an arc at the centre is double the angle subtended by it at any point on the remaining part of the circle.
- 3.11. Angles in the same segment of a circle are equal.
- 3.12. Angle in a semicircle is a right angle.
- 3.13. If a line segment joining two points subtends equal angles at two other points lying on the same side of the line containing the line segment, the four points lie on a circle.
- 3.14. The sum of either pair of opposite angles of a cyclic quadrilateral is 180°.
- 3.15. If sum of a pair of opposite angles of a quadrilateral is 180°, the quadrilateral is cyclic.
- 3.16. AB is a diameter of the circle, CD is a chord equal to the radius of the circle. AC and BD when extended intersect at a point E. Prove that  $\angle AEB = 60^{\circ}$ .

- 3.17. ABCD is a cyclic quadrilateral in which AC and BD are its diagonals. If  $\angle DBC = 55^{\circ}$  and  $\angle BAC = 45^{\circ}$ , find  $\angle BCD$
- 3.18. Two circles intersect at two points AandB. AD and AC are diameters to the two circles. Prove that B lies on the line segment DC.
- 3.19. Prove that the quadrilateral formed (if possible) by the internal angle bisectors of any quadrilateral is cyclic.
- 3.20. Equal chords of a circle (or of congruent circles) subtend equal angles at the centre.
- 3.21. If the angles subtended by two chords of a (corresponding centres) are equal, the chords are equal.
- 3.22. The perpendicular from the centre of a circle to a chord bisects the chord.
- 3.23. The line drawn through the centre of a circle to bisect a chord is perpendicular to the chord.
- through three non-collinear points.
- 3.25. Equal chords of a circle (or of congruent circles) are equidistant from the centre (or corresponding centres).
- 3.26. Chords equidistant from the centre (or corresponding centres) of a circle (or of congruent circles) are equal.
- 3.27. If two arcs of a circle are congruent, then their if two chords of a circle are equal, then their corresponding arcs (minor, major) are congruent.
- at the centre.
- 3.29. The angle subtended by an arc at the centre is on the remaining part of the circle.
- 3.30. Angles in the same segment of a circle are equal.
- 3.31. Angle in a semicircle is a right angle.
- 3.32. If a line segment joining two points subtends equal angles at two other points lying on the same side of the line containing the line seg- 3.49. Let the vertex of an angle ABC be located ment, the four points lie on a circle.
- 3.33. The sum of either pair of opposite angles of a cyclic quadrilateral is 180°.
- 3.34. If sum of a pair of opposite angles of a quadrilateral is 180°, the quadrilateral is cyclic.
- equal to the radius of the circle. AC and BD

- when extended intersect at a point E. Prove that  $\angle AEB = 60^{\circ}$ .
- 3.36. Two circles intersect at two points A and B. AD and AC are diameters to the two circles. Prove that B lies on the line segment DC.
- 3.37. Prove that the quadrilateral formed (if possible) by the internal angle bisectors of any quadrilateral is cyclic.
- 3.38. If two equal chords of a circle intersect within the circle, prove that the segments of one chord are equal to corresponding segments of the other chord.
- circle (or of congruent circles) at the centre 3.39. If two equal chords of a circle intersect within the circle, prove that the line joining the point of intersection to the centre makes equal angles with the chords.
  - 3.40. If a line intersects two concentric circles (circles with the same centre) with centre O at A, B, C and D, prove that AB = CD.
- 3.24. There is one and only one circle passing 3.41. A chord of a circle is equal to the radius of the circle. Find the angle subtended by the chord at a point on the minor arc and also at a point on the major arc.
  - 3.42. If diagonals of a cyclic quadrilateral are diameters of the circle through the vertices of the quadrilateral, prove that it is a rectangle.
  - 3.43. If the non-parallel sides of a trapezium are equal, prove that it is cyclic.
  - corresponding chords are equal and conversely 3.44. Two circles intersect at two points B and C. Through B, two line segments ABD and PBQ are drawn to intersect the circles at A, D and P, Q respectively. Prove that  $\angle ACP = \angle QCD$ .
- 3.28. Congruent arcs of a circle subtend equal angles 3.45. If circles are drawn taking two sides of a triangle as diameters, prove that the point of intersection of these circles lie on the third side.
  - double the angle subtended by it at any point 3.46. ABC and ADC are two right triangles with common hypotenuse AC. Prove that  $\angle CAD =$  $\angle CBD$ .
    - 3.47. Prove that a cyclic parallelogram is a rectangle.
    - 3.48. Prove that the line of centres of two intersecting circles subtends equal angles at the two points of intersection.
    - outside a circle and let the sides of the angle intersect equal chords AD and CE with the circle. Prove that  $\angle ABC$  is equal to half the difference of the angles subtended by the chords AC and DE at the centre.
- 3.35. AB is a diameter of the circle, CD is a chord 3.50. Prove that the circle drawn with any side of a rhombus as diameter, passes through the point

- of intersection of its diagonals.
- 3.51. ABCD is a parallelogram. The circle through A, B and C intersect CD (produced if necessary) at E. Prove that AE = AD.
- 3.52. AC and BD are chords of a circle which bisect 3.68. Two chords AB and CD of a circle intersect each other. Prove that (i) AC and BD are diameters, (ii) ABCD is a rectangle.
- 3.53. Bisectors of angles A, B and C of a  $\triangle ABC$ intersect its circumcircle at D, E and F respectively. Prove that the angles of the  $\triangle DEF$  are  $90^{\circ} - \frac{A}{2}$ ,  $90^{\circ} - \frac{B}{2}$  and  $90^{\circ} - \frac{C}{2}$ .
- 3.54. Two congruent circles intersect each other at points A and B. Through A any line segment PAQ is drawn so that P, Q lie on the two circles. Prove that BP = BQ.
- 3.55. In any  $\triangle ABC$ , if the angle bisector of  $\angle A$  and perpendicular bisector of BC intersect, prove that they intersect on the circumcircle of the
- 3.56. The lengths of tangents drawn from an external point to a circle are equal.
- 3.57. Prove that in two concentric circles, the chord of the larger circle, which touches the smaller circle, is bisected at the point of contact.
- 3.58. Two tangents TP and TQ are drawn to a circle with centre O from an external point T. Prove that  $\angle PTQ = 2\angle OPQ$ .
- 3.59. Prove that the tangents drawn at the ends of a diameter of a circle are parallel.
- 3.60. Prove that the perpendicular at the point of contact to the tangent to a circle passes through the centre.
- 3.61. A quadrilateral *ABCD* is drawn to circumscribe a circle. Prove that AB + CD = AD + BC.
- 3.62. XY and X'Y' are two parallel tangents to a circle with centre O and another tangent ABwith point of contact C intersecting XY at Aand X'Y' at B. Prove that  $\angle AOB = 90^{\circ}$
- 3.63. Prove that the angle between the two tangents drawn from an external point to a circle is supplementary to the angle subtended by the line-segment joining the points of contact at the centre.
- 3.64. Prove that the parallelogram circumscribing a 4.10. A tangent PQ at a point P of a circle of radius circle is a rhombus.
- 3.65. Prove that opposite sides of a quadrilateral circumscribing a circle subtend supplementary angles at the centre of the circle.
- 3.66. Find the area of a sector of angle p (in degrees) of a circle with radius R.

- 3.67. Two chords AB and CD intersect each other at the point P. Prove that:
  - a)  $\triangle APC \sim \triangle DPB$
  - b) AP.PB = CP.DP
- each other at the point P (when produced) outside the circle. Prove that
  - a)  $\triangle PAC \sim \triangle PDB$
  - b) PA.PB = PC.PD

### 4 Miscellaneous Exercises

- 4.1. ABCD is a cyclic quadrilateral in which AC and BD are its diagonals. If  $\angle DBC = 55^{\circ}$  and  $\angle BAC = 45^{\circ}$ , find  $\angle BCD$
- 4.2. Two circles of radii 5 cm and 3 cm intersect at two points and the distance between their centres is 4 cm. Find the length of the common
- 4.3. A,B and C are three points on a circle with centre O such that  $\angle BOC = 30^{\circ}$  and  $\angle AOB =$ 60°. If D is a point on the circle other than the arc ABC, find  $\angle ADC$ .
- 4.4.  $\angle PQR = 100^{\circ}$ , where P, Q and R are points on a circle with centre O. Find  $\angle OPR$ .
- 4.5. A, B, C, D are points on a circle such that  $\angle ABC = 69^{\circ}, \angle ACB = 31^{\circ}, \text{ find } \angle BDC.$
- 4.6. A, B, C and D are four points on a circle. AC and BD intersect at a point E such that  $\angle BEC =$  $130^{\circ}$  and  $\angle ECD = 20^{\circ}$ . Find  $\angle BAC$ .
- 4.7. ABCD is a cyclic quadrilateral whose diagonals intersect at a point E. If  $\angle DBC =$  $70^{\circ}$ ,  $\angle BAC$  is  $30^{\circ}$ , find  $\angle BCD$ . Further, if AB =BC, find  $\angle ECD$ .
- 4.8. Two chords AB and CD of lengths 5 cm and 11 cm respectively of a circle are parallel to each other and are on opposite sides of its centre. If the distance between AB and CD is 6 cm, find the radius of the circle.
- 4.9. The lengths of two parallel chords of a circle are 6 cm and 8 cm. If the smaller chord is at distance 4 cm from the centre, what is the distance of the other chord from the centre?
- 5 cm meets a line through the centre O at a point Q so that OQ = 12 cm. Find the length of PO.
- 4.11. PQ is a chord of length 8 cm of a circle of radius 5 cm. The tangents at P and Q intersect at a point T. Find the length TP.

- circle is 24 cm and the distance of Q from the centre is 25 cm. Find the radius of the circle is
- 4.13. If TP and TQ are the two tangents to a circle  $\angle PTQ$
- 4.14. If tangents PA and PB from a point P to a circle with centre O are inclined to each other at angle of 80  $^{\circ}$  , then find  $\angle POA$
- 4.15. The length of a tangent from a point A at 4.28. In a circle of radius 21 cm, an arc subtends an distance 5 cm from the centre of the circle is 4 cm. Find the radius of the circle.
- 4.16. Two concentric circles are of radii 5 cm and 3 cm. Find the length of the chord of the larger circle which touches the smaller circle.
- 4.17. A  $\triangle ABC$  is drawn to circumscribe a circle of radius 4 cm such that the segments BD and DC into which BC is divided by the point of contact D are of lengths 8 cm and 6 cm respectively. Find the sides AB and AC.
- 4.18. The cost of fencing a circular field at the rate of ₹24 per metre is ₹5280. The field is to be ploughed at the rate of  $\ge 0.50$  per  $m^2$ . Find the cost of ploughing the field.
- 4.19. The radii of two circles are 19 cm and 9 cm respectively. Find the radius of the circle which has circumference equal to the sum of the circumferences of the two circles.
- 4.20. The radii of two circles are 8 cm and 6 cm respectively. Find the radius of the circle having area equal to the sum of the areas of the two 4.32. A brooch is made with silver wire in the form circles.
- 4.21. A circular archery target is marked with its five scoring regions from the centre outwards as Gold, Red, Blue, Black and White. The diameter of the region representing Gold score is 21 cm and each of the other bands is 10.5 cm 4.33. An umbrella has 8 ribs which are equally wide. Find the area of each of the five scoring regions.
- 4.22. The wheels of a car are of diameter 80 cm each. How many complete revolutions does 4.34. A car has two wipers which do not overlap. each wheel make in 10 minutes when the car is travelling at a speed of 66 km per hour?
- 4.23. Find the area of the sector of a circle with radius 4 cm and of angle 30 °. Also, find the 4.35. area of the corresponding major sector.
- 4.24. Find the area of the segment AYB, if radius of the circle is 21 cm and  $\angle AOB = 120^{\circ}$ .
- 4.25. Find the area of a sector of a circle with radius 6 cm if angle of the sector is 60  $^{\circ}$  .

- 4.12. From a point Q, the length of the tangent to a 4.26. Find the area of a quadrant of a circle whose circumference is 22 cm. 3. The length of the minute hand of a clock is 14 cm. Find the area swept by the minute hand in 5 minutes.
  - with centre O so that  $\angle POQ = 110^{\circ}$ , then find 4.27. A chord of a circle of radius 10 cm subtends a right angle at the centre. Find the area of the corresponding:
    - a) minor segment
    - b) major sector.
    - angle of 60  $^{\circ}$  at the centre. Find:
      - a) the length of the arc
      - b) area of the sector formed by the arc
      - c) area of the segment formed by the corresponding chord
    - 4.29. A chord of a circle of radius 15 cm subtends an angle of 60  $^{\circ}$  at the centre. Find the areas of the corresponding minor and major segments of the circle.
    - 4.30. A chord of a circle of radius 12 cm subtends an angle of 120  $^{\circ}$  at the centre. Find the area of the corresponding segment of the circle.
    - 4.31. A horse is tied to a peg at one corner of a square shaped grass field of side 15 m by means of a 5 m long rope. Find
      - a) the area of that part of the field in which the horse can graze.
      - b) the increase in the grazing area if the rope were 10 m long instead of 5 m.
    - of a circle with diameter 35 mm. The wire is also used in making 5 diameters which divide the circle into 10 equal sectors. Find:
      - a) the total length of the silver wire required.
      - b) the area of each sector of the brooch
    - spaced. Assuming umbrella to be a flat circle of radius 45 cm, find the area between the two consecutive ribs of the umbrella.
    - Each wiper has a blade of length 25 cm sweeping through an angle of 115  $^{\circ}$  . Find the total area cleaned at each sweep of the blades.
    - To warn ships for underwater rocks, a lighthouse spreads a red coloured light over a sector of angle 80 ° to a distance of 16.5 km. Find the area of the sea over which the ships are warned.

- 4.36. A round table cover has six equal designs. If the radius of the cover is 28 cm, find the cost of making the designs at the rate of ₹0.35 per  $cm^2$ .
- 4.37. Two circular flower beds are located on opposite sides of a square lawn ABCD of side 56 m. If the centre Of each circular flower bed is the point of intersection O of the diagonals of 4.49. The area of an equilateral  $\triangle ABC$  is 17320.5 the square lawn, find the sum of the areas of the lawn and the flower beds.
- 4.38. Four circles are inscribed inside a square ABCD of side 14 cm such that each one touches exernally two adjacent sides of the square and two other circles. Find the region 4.50. On a square handkerchief, nine circular designs between the circles and the square.
- 4.39. ABCD is a square of side 10 cm and semicircles are drawn with each side of the square as diameter. Find the area enclosed by the circular 4.51. OACB is a quadrant of a circle with centre O
- 4.40. P is a point on the semi-circle formed with diameter QR. Find the area between the semicircle and  $\triangle PQR$  if PQ = 24 cm, PR = 7 cm and O is the centre Of the circle.
- 4.41. AC and BD are two arcs on concentric circles with radii 14 cm and 7 cm respectively, such that  $\angle AOC = 40^{\circ}$ . Find the area of the region ABDC.
- 4.42. Find the area between a square ABCD of side 14cm and the semi circles APD and BPC.
- 4.43. Find the area of the region enclosed by a circular arc of radius 6 cm drawn with vertex O of an equilateral triangle OAB of side 12 cm as centre.
- 4.44. From each corner of a square of side 4 cm a quadrant of a circle of radius 1 cm is cut and also a circle of diameter 2 cm is cut. Find the area of the remaining portion of the square.
- 4.45. In a circular table cover of radius 32 cm, a design is formed leaving an equilateral  $\triangle ABC$ in the middle. Find the area of the design.
- 4.46. ABCD is a square of side 14 cm. With centres A, B, C and D, four circles are drawn such that each circle touches externally two of the remaining three circles. Find the area within the square that lies outside the circles.
- 4.47. The left and right ends of a racing track are semicircular. The distance between the two 4.58. Draw a triangle whose sides are 8cm and 11cm inner parallel line segments is 60 m and they are each 106 m long. If the track is 10 m wide, find:

- a) the distance around the track along its inner
- b) the area of the track.
- 4.48. AB and CD are two diameters of a circle (with centre O) perpendicular to each other and OD is the diameter of a smaller circle inside. If OA = 7 cm, find the area of the smaller circle.
- $cm^2$ . With each vertex of the triangle as centre, a circle is drawn with radius equal to half the length of the side of the triangle. Find the area of region within the triangle but outside the circles.
- are inscribed touching each other, each of radius 7 cm. Find the area of the remaining portion of the handkerchief.
- and radius 3.5 cm. D is a point on OA. If OD = 2 cm, find the area of the
  - a) quadrant OACB,
  - b) the region between the quadrant and  $\triangle OBD$ .
- 4.52. A square *OABC* is inscribed in a quadrant OPBQ. If OA = 20 cm, find the area between the square and the quadrant.
- 4.53. AB and CD are respectively arcs of two concentric circles of radii 21 cm and 7 cm and centre O. If  $\angle AOB = 30^{\circ}$ , find the area of the region ABCD.
- 4.54. ABC is a quadrant of a circle of radius 14 cm and a semicircle is drawn with BC as diameter. Find the area of the crescent formed.
- 4.55. Find the area common between the two quadrants of circles of radius 8 cm each if the centres of the circles lie on opposite sides of a square.
- 4.56. Find the area of the sector of a circle with radius 4 cm and of angle 30°. Also, find the area of the corresponding major sector.
- 4.57. A pole has to be erected at a point on the boundary of a circular park of diameter 13 metres in such a way that the differences of its distances from two diametrically opposite fixed gates A and B on the boundary is 7 metres. Is it possible to do so? If yes, at what distances from the two gates should the pole be erected?
- and the perimeter is 32 cm and find its area.
- 4.59. A triangular park ABC has sides 120m, 80m

- and 50m. A gardener Dhania has to put a fence all around it and also plant grass inside. Draw this park. How much area does she need to plant? Find the cost of fencing it with barbed wire at the rate of ₹20 per metre leaving a space 3m wide for a gate on one side.
- 4.60. The sides of a triangular plot are in the ratio of 3:5:7 and its perimeter is 300 m. Draw the plot and find its area.
- 4.61. A tower stands vertically on the ground. From a point on the ground, which is 15m away from 4.70. Find the area of a triangle two sides of which the foot of the tower, the angle of elevation of the height of the tower.
- 4.62. An electrician has to repair an electric fault 4.72. An isosceles triangle has perimeter 30 cm and pole of height 5m. She needs to reach a point 1.3m below the top of the pole to undertake the repair work. What should be the length of 4.73. A girl walks 4km west, then she walks 3km the ladder that she should use which, when inclined at an angle of 60° to the horizontal, would enable her to reach the required position? Also, how far from the foot of the pole 4.74. A circus artist is climbing a 20m long rope, should she place the foot of the ladder?
- 4.63. An observer 1.5m tall is 28.5m away from a chimney. The angle of elevation of the top of the chimney from her eyes is 45°. What is the height of the chimney?
- 4.64. From a point **P** on the ground the angle of elevation of the top of a 10m tall building is 30°. A flag is hoisted at the top of the building and the angle of elevation of the top of the flagstaff from P is 45°. Find the length of the flagstaff and the distance of the building from 4.76. A contractor plans to install two slides for the the point **P**.
- 4.65. The shadow of a tower standing on a level ground is found to be 40m longer when the Sun's altitude is 30° than when it is 60°. Find the height of the tower.
- 4.66. The angles of depression of the top and the bottom of an 8m tall building from the top of a multi-storeyed building are 30° and 45° respectively. Find the height of the multi- 4.77. storeyed building and the distance between the two buildings.
- 4.67. A traffic signal board, indicating 'SCHOOL 'a'. Find the area of the signal board, using Heron's formula. If its perimeter is 180 cm, what will be the area of the signal board?
- 4.68. The triangular side walls of a flyover have

- been used for advertisements. The sides of the walls are 122 m, 22 m and 120 m. The advertisements yield an earning of ₹5000 per  $m^2$  per year. A company hired one o its walls for 3 months. How much rent did it pay?
- 4.69. There is a slide in a park. One of its side walls has been painted in some colour with a message "KEEP THE PARK GREEN AND CLEAN". If the sides of the wall are 15 m, 11 m and 6 m, find the area painted in colour.
- are 18cm and 10cm and the perimeter is 42cm.
- the top of the tower is found to be 60°. Find 4.71. Sides of a triangle are in the ratio of 12:17: 25 and its perimeter is 540cm. Find its area.
  - each of the equal sides is 12 cm. Find the area of the triangle.
  - in a direction 30° east of north and stops. Determine the girl's displacement from her initial point of departure.
  - which is tightly stretched and tied from the top of a vertical pole to the ground. Find the height of the pole, if the angle made by the rope with the ground level is 30°.
  - 4.75. A tree breaks due to storm and the broken part bends so that the top of the tree touches the ground making an angle of 30° with it. The distance between the foot of the tree to the point where the top touches the ground is 8m. Find the height of the tree.
  - children to play in a park. For the children below the age of 5 years, she prefers to have a slide whose top is at a height of 1.5m, and is inclined at an angle of 30° to the ground, whereas for elder children she wants to have a steep slide at a height of 3m, and inclined at an angle of 60° to the ground. What should be the length of the slide in each case?
  - The angle of elevation of the top of a tower from a point on the ground, which is 30m away from the foot of the tower, is 30°. Find the height of the tower.
- AHEAD', is an equilateral triangle with side 4.78. A kite is flying at a height of 60m above the ground. The string attached to the kite is temporarily tied to a point on the ground. The inclination of the string with the ground is 60°. Find the length of the string, assuming that

- there is no slack in the string.
- 4.79. A 1.5m tall boy is standing at some distance from a 30m tall building. The angle of elevation from his eyes to the top of the building increases from 30° to 60° as he walks towards the building. Find the distance he walked towards the building.
- 4.80. From a point on the ground, the angles of elevation of the bottom and the top of a transmission tower fixed at the top of a 20 m high building are 45° and 60° respectively. Find the height of the tower.
- 4.81. A statue, 1.6 m tall, stands on the top of a pedestal. From a point on the ground, the angle of elevation of the top of the statue is 60° and 4.89. The angles of elevation of the top of a tower from the same point the angle of elevation of the top of the pedestal is 45°. Find the height of the pedestal.
- 4.82. The angle of elevation of the top of a building from the foot of the tower is  $30^{\circ}$  and the angle 4.90. E and F are points on the sides PQ and PR of elevation of the top of the tower from the foot of the building is 60°. If the tower is 50 m high, find the height of the building.
- 4.83. Two poles of equal heights are standing opposite each other on either side of the road, which is 80 m wide. From a point between them on the road, the angles of elevation of the top of the poles are 60° and 30°, respectively. Find the height of the poles and the distances of the 4.91. A girl of height 90 cm is walking away from point from the poles.
- 4.84. A TV tower stands vertically on a bank of a canal. From a point on the other bank directly opposite the tower, the angle of elevation of 4.92.  $\triangle ODC \sim \triangle OBA$ ,  $\angle BOC = 125^{\circ}$  and  $\angle CDO = 125^{\circ}$ the top of the tower is 60°. From another point 20 m away from this point on the line joing 4.93. Nazima is fly fishing in a stream. The tip of her this point to the foot of the tower, the angle of elevation of the top of the tower is 30°. Find the height of the tower and the width of the canal.
- 4.85. From the top of a 7 m high building, the angle of elevation of the top of a cable tower is 60° and the angle of depression of its foot is 45°. Determine the height of the tower.
- 4.86. As observed from the top of a 75 m high lighthouse from the sea-level, the angles of 4 94 depression of two ships are 30° and 45°. If one ship is exactly behind the other on the same side of the lighthouse, find the distance between the two ships.
- 4.87. A 1.2 m tall girl spots a balloon moving with

- the wind in a horizontal line at a height of 88.2 m from the ground. The angle of elevation of the balloon from the eyes of the girl at any instant is 60°. After some time, the angle of elevation reduces to 30°. Find the distance travelled by the balloon during the interval.
- 4.88. A straight highway leads to the foot of a tower. A man standing at the top of the tower observes a car at an angle of depression of 30°, which is approaching the foot of the tower with a uniform speed. Six seconds later, the angle of depression of the car is found to be 60°. Find the time taken by the car to reach the foot of the tower from this point.
- from two points at a distance of 4 m and 9 m from the base of the tower and in the same straight line with it are complementary. Prove that the height of the tower is 6 m.
- respectively of a  $\triangle PQR$ . For each of the following cases, state whether  $EF \parallel QR$ 
  - a) PE = 3.9cm, EQ = 3cm, PF = 3.6cm and FR = 2.4cm
  - b) PE = 4cm, QE = 4.5cm, PF = 8cm and RF = 9cm
  - c) PQ = 1.28cm, PR = 2.56cm, PE = 0.18cmand PF = 0.36cm
- the base of a lamp-post at a speed of 1.2 m/s. If the lamp is 3.6 m above the ground, find the length of her shadow after 4 seconds.
- 70°. Find  $\angle DOC$ ,  $\angle DCO$  and  $\angle OAB$ .
- fishing rod is 1.8 m above the surface of the water and the fly at the end of the string rests on the water 3.6 m away and 2.4 m from a point directly under the tip of the rod. Assuming that her string (from the tip of her rod to the fly) is taut, how much string does she have out? If she pulls in the string at the rate of 5 cm per second, what will be the horizontal distance of the fly from her after 12 seconds?
- A vertical pole of length 6 m casts a shadow 4 m long on the ground and at the same time a tower casts a shadow 28 m long. Find the height of the tower.
- 4.95. Let  $\triangle ABC \sim \triangle DEF$  and their areas be, respec-

- tively,  $64 \text{ cm}^2$  and  $121 \text{ cm}^2$ . If EF = 15.4 cm, 4.106. A field is in the shape of a trapezium whose find BC.
- 4.96. A ladder is placed against a wall such that its foot is at a distance of 2.5 m from the wall and its top reaches a window 6 m above the 4.107. ABCD is a parallelogram,  $AE \perp DC$  and  $CF \perp$ ground. Find the length of the ladder.
- 4.97. Sides of triangles are given below. Determine which of them are right triangles. In case of 4.108. Kamla has a triangular field with sides 240 a right triangle, write the length of its hypotenuse.
  - a) 7 cm, 24 cm, 25 cm
  - b) 3 cm, 8 cm, 6 cm
  - c) 50 cm, 80 cm, 100 cm
  - d) 13 cm, 12 cm, 5 cm
- 4.98. A ladder 10 m long reaches a window 8 m above the ground. Find the distance of the foot of the ladder from base of the wall.
- 4.99. A guy wire attached to a vertical pole of height 18 m is 24 m long and has a stake attached to the other end. How far from the base of the 4.109. Students of a school staged a rally for cleanlipole should the stake be driven so that the wire will be taut?
- 4.100. An aeroplane leaves an airport and flies due north at a speed of 1000 km per hour. At the same time, another aeroplane leaves the same airport and flies due west at a speed of 1200 km per hour. How far apart will be the two planes after  $1\frac{1}{2}$  hours?
- 4.101. Two poles of heights 6 m and 11 m stand on a plane ground. If the distance between the feet of the poles is 12 m, find the distance between 4.110. Sanya has a piece of land which is in the shape their tops.
- 4.102. In  $\triangle ABC$ ,  $AB = 6\sqrt{3}cm$ , AC = 12cm and BC = 12cm6*cm*. Find the angle *B*.
- 4.103. A park, in the shape of a quadrilateral ABCD, has  $\angle C = 90^{\circ}, AB = 9m, BC = 12m, CD = 5m$ and AD = 8m. How much area does it occupy? 2. Find the area of a quadrilateral ABCD in which AB = 3cm, BC = 4cm, CD = 4cm, DA = 4.111. 5cm and AC = 5cm.
- 4.104. A triangle and a parallelogram have the same base and the same area. If the sides of the triangle are 26 cm, 28 cm and 30 cm, and the parallelogram stands on the base 28 cm, find the height of the parallelogram.
- 4.105. A rhombus shaped field has green grass for 18 cows to graze. If each side of the rhombus is 304.112. A circular park of radius 20m is situated in m and its longer diagonal is 48 m, how much area of grass field will each cow be getting?

- parallel sides are 25 m and 10 m. The nonparallel sides are 14 m and 13 m. Find the area of the field.
- AD. If AB = 16cm, AE = 8 cm and CF = 10cm. find AD.
- m, 200 m, 360 m, where she grew wheat. In another triangular field with sides 240 m, 320 m, 400 m adjacent to the previous field, she wanted to grow potatoes and onions. She divided the field in two parts by joining the mid-point of the longest side to the opposite vertex and grew patatoes in one part and onions in the other part. Draw the figure for this problem. How much area (in hectares) has been used for wheat, potatoes and onions? (1 hectare  $= 10000 m^2$ ).
- ness campaign. They walked through the lanes in two groups. One group walked through the lanes AB, BC and CA; while the other through AC, CD and DA. Then they cleaned the area enclosed within their lanes. If AB = 9 m, BC = 40 m, CD = 15 m, DA = 28 m and  $\angle B = 90^{\circ}$ , which group cleaned more area and by how much? Draw the corresponding figure. Find the total area cleaned by the students (neglecting the width of the lanes).
  - of a rhombus. She wants her one daughter and one son to work on the land and produce different crops. She divided the land in two equal parts. If the perimeter of the land is 400 m and one of the diagonals is 160 m, how much area each of them will get for their crops? Draw the rhombus.
  - Three girls Reshma, Salma and Mandip are playing a game by standing on a circle of radius 5m drawn in a park. Reshma throws a ball to Salma, Salma to Mandip, Mandip to Reshma. If the distance between Reshma and Salma and between Salma and Mandip is 6m each, what is the distance between Reshma and Mandip?
- a colony. Three boys Ankur, Syed and David are sitting at equal distance on its boundary each having a toy telephone in his hands to

- talk each other. Find the length of the string of each phone.
- 4.113. The longest side of a triangle is 3 times the shortest side and the third side is 2 cm shorter than the longest side. If the perimeter of the 4.123. Two identical billiard balls strike a rigid wall triangle is at least 61 cm, find the minimum length of the shortest side.
- 4.114. A rectangular park is to be designed whose breadth is 3 m less than its length. Its area is to be 4 square metres more than the area of a park that has already been made in the shape of an isosceles triangle with its base as the breadth of the rectangular park and of altitude 12 m. Find its length and breadth.
- 4.115. The area of a rectangular plot is  $528 m^2$ . The length of the plot (in metres) is one more than twice its breadth. We need to find the length and breadth of the plot.
- 4.116. The altitude of a right triangle is 7 cm less than its base. If the hypotenuse is 13 cm, find the other two sides.
- 4.117. The diagonal of a rectangular field is 60 metres more than the shorter side. If the longer side is 30 metres more than the shorter side, find the 4.124. See Fig. 4.124. A mass of 6 kg is suspended by sides of the field.
- 4.118. Is it possible to design a rectangular mango grove whose length is twice its breadth, and the area is  $800 \text{ m}^2$ ? If so, find its length and breadth.
- 4.119. Is it possible to design a rectangular park of perimeter 80 m and area 400  $m^2$  If so, find its length and breadth.
- 4.120. On an open ground, a motorist follows a track that turns to his left by an angle of 600 after every 500 m. Starting from a given turn, specify the displacement of the motorist at the third, sixth and eighth turn. Compare the magnitude of the displacement with the total path length covered by the motorist in each case.
- 4.121. A passenger arriving in a new town wishes to go from the station to a hotel located 10 km away on a straight road from the station. A dishonest cabman takes him along a circuitous4.125. See Fig. 4.125. A mass of 4 kg rests on a path 23 km long and reaches the hotel in 28 min. What is
  - a) the average speed of the taxi,
  - b) the magnitude of average velocity? Are the two equal?

- above the ground. If the angle subtended at a ground observation point by the aircraft positions 10.0 s apart is 30°, what is the speed of the aircraft?
- with the same speed but at different angles, and get reflected without any change in speed, as shown in Fig. 4.123. What is
  - a) the direction of the force on the wall due to each ball?
- b) the ratio of the magnitudes of impulses imparted to the balls by the wall?





Fig. 4.123

a rope of length 2 m from the ceiling. A force of 50 N in the horizontal direction is applied at the midpoint P of the rope, as shown. What is the angle the rope makes with the vertical in equilibrium ? (Take  $g = 10 ms^{-2}$ ).



- horizontal plane. The plane is gradually inclined until at an angle  $\theta = 15^{\circ}$  with the horizontal, the mass just begins to slide. What is the coefficient of static friction between the block and the surface?
- 4.122. An aircraft is flying at a height of 3400 m<sup>4.126</sup>. A cyclist speeding at 18 km/h on a level road takes a sharp circular turn of radius 3 m




Fig. 4.125

without reducing the speed. The co-efficient of static friction between the tyres and the road is withstand a maximum tension of 200 N?

0.1. Will the cyclist slip while taking the turn?4.134. A woman pushes a trunk on a railway platform

- 4.127. A circular racetrack of radius 300 m is banked at an angle of 15°. If the coefficient of friction between the wheels of a race-car and the road is 0.2, what is the
  - a) optimum speed of the racecar to avoid wear and tear on its tyres, and
  - b) maximum permissible speed to avoid slipping ?
- 4.128. An aircraft executes a horizontal loop at a speed of 720 km/h with its wings banked at 4.135. A bob of mass m is suspended by a light string of length L. It is imparted a horizontal
- 4.129. A train runs along an unbanked circular track of radius 30 m at a speed of 54 km/h. The mass of the train is 10<sup>6</sup> kg. What provides the centripetal force required for this purpose The engine or the rails? What is the angle of banking required to prevent wearing out of the rail?
- 4.130. A disc revolves with a speed of  $33\frac{1}{3}$  rev/min and has a radius of 15 cm. Two coins are placed at 4 cm and 14 cm away from the centre of the record. If the co-efficient of friction between the coins and the record is 0.15, which of the 4.136. coins will revolve with the record?
- 4.131. A 70 kg man stands in contact against the inner wall of a hollow cylindrical drum of radius 3 m rotating about its vertical axis with 200 rev/min. The coefficient of friction between the wall and his clothing is 0.15. What is the 4.137. minimum rotational speed of the cylinder to enable the man to remain stuck to the wall (without falling) when the floor is suddenly removed?
- 4.132. A thin circular loop of radius R rotates about

its vertical diameter with an angular frequency  $\omega$ . Show that a small bead on the wire loop remains at its lowermost point for  $\omega \leq \sqrt{\frac{g}{R}}$ . What is the angle made by the radius vector joining the centre to the bead with the vertical downward direction for  $\omega = \sqrt{\frac{2g}{R}}$ ? Neglect friction.

4.133. A stone of mass 0.25 kg tied to the end of a string is whirled round in a circle of radius 1.5 m with a speed of 40 rev./min in a horizontal plane. What is the tension in the string? What is the maximum speed with which the stone can be whirled around if the string can withstand a maximum tension of 200 N?

A woman pushes a trunk on a railway platform which has a rough surface. She applies a force of 100 N over a distance of 10 m. Thereafter, she gets progressively tired and her applied force reduces linearly with distance to 50 N. The total distance through which the trunk has been moved is 20 m. Plot the force applied by the woman and the frictional force, which is 50 N versus displacement. Calculate the work done by the two forces over 20 m.

A bob of mass m is suspended by a light string of length L. It is imparted a horizontal velocity  $v_o$  such that it completes a semi-circular trajectory in the vertical plane with the string becoming slack only on reaching the topmost point, C. This is shown in Fig. 4.135. Obtain an expression for

- a) vo;
- b) the speeds at points B and C;
- c) the ratio of the kinetic energies  $(K_B/K_C)$  at B and C.

Comment on the nature of the trajectory of the bob after it reaches the point C.

The bob A of a pendulum released from 30°to the vertical hits another bob B of the same mass at rest on a table as shown in Fig. 4.136. How high does the bob A rise after the collision? Neglect the size of the bobs and assume the collision to be elastic.

The bob of a pendulum is released from a horizontal position. If the length of the pendulum is 1.5 m, what is the speed with which the bob arrives at the lowermost point, given that it dissipated 5% of its initial energy against air





Fig. 4.135




resistance?

4.138. The blades of a windmill sweep out a circle of area A.

- (a) If the wind flows at a velocity v perpendicular to the circle, what is the mass of the air passing through it in time t?
- (b) What is the kinetic energy of the air?
- (c) Assume that the windmill converts 25% of the wind's energy into electrical energy, and

that  $A = 30m^2$ , v = 36km/h and the density of air is  $1.2kgm^{-3}$ . What is the electrical power produced ?

- 4.139. A bullet of mass 0.012 kg and horizontal speed 70ms<sup>-1</sup>. strikes a block of wood of mass 0.4 kg and instantly comes to rest with respect to the block. The block is suspended from the ceiling by means of thin wires. Calculate the height to which the block rises. Also, estimate the amount of heat produced in the block.
- 4.140. Two inclined frictionless tracks, one gradual and the other steep meet at A from where two stones are allowed to slide down from rest, one on each track (Fig. 4.140). Will the stones reach the bottom at the same time? Will they reach there with the same speed? Explain. Given  $\theta_1 = 30^{\circ}\theta_2 = 60^{\circ}$  and h = 10m, what are the speeds and times taken by the two stones?



4.141. A 1 kg block situated on a rough incline is connected to a spring of spring constant 100 N m-1 as shown in Fig. 4.141. The block is released from rest with the spring in the unstretched position. The block moves 10 cm down the incline before coming to rest. Find the coefficient of friction between the block and the incline. Assume that the spring has a negligible mass and the pulley is frictionless.

4.142. Find the centre of mass of three particles at the vertices of an equilateral triangle. The masses of the particles are 100g, 150g, and 200g respectively. Each side of the equilateral triangle is 0.5m long.

4.143. Find the centre of mass of a triangular lamina.4.144. Find the centre of mass of a uniform L-shaped lamina (a thin flat plate) with dimensions as

shown in Fig. 4.144. The mass of the lamina




Fig. 4.141

is 3 kg.




Fig. 4.144

- 4.145. A metal bar 70 cm long and 4.00 kg in mass supported on two knife-edges placed 10 cm from each end. A 6.00 kg load is suspended at 30 cm from one end. Find the reactions at the knifeedges. (Assume the bar to be of uniform cross section and homogeneous.)
- 4.146. A 3m long ladder weighing 20 kg leans on a frictionless wall. Its feet rest on the floor 1 m from the wall as shown in Fig. 4.146. Find the reaction forces of the wall and the floor.
- 4.147. What is the moment of inertia of a rod of mass M, length 1 about an axis perpendicular to it through one end?
- 4.148. What is the moment of inertia of a ring about a tangent to the circle of the ring?
- 4.149. A cord of negligible mass is wound round the rim of a fly wheel of mass 20 kg and radius




Fig. 4.146

20 cm. A steady pull of 25 N is applied on the cord as shown in Fig. 4.149. The flywheel is mounted on a horizontal axle with frictionless bearings.

- (a) Compute the angular acceleration of the wheel.
- (b) Find the work done by the pull, when 2m of the cord is unwound.
- (c) Find also the kinetic energy of the wheel at this point. Assume that the wheel starts from rest
- (d) Compare answers to parts (b) and (c).
- 4.150. Three bodies, a ring, a solid cylinder and a solid sphere roll down the same inclined plane without slipping. They start from rest. The radii




Fig. 4.149

of the bodies are identical. Which of the bodies reaches the ground with maximum velocity?

- 4.151. In the HCl molecule, the separation between the nuclei of the two atoms is about 1.27  $\mathring{A}$  (1  $\mathring{A}$  = 10<sup>-10</sup> m). Find the approximate 4.156.  $\mathring{A}$  solid cylinder of mass 20 kg rotates about location of the CM of the molecule, given that a chlorine atom is about 35.5 times as massive as a hydrogen atom and nearly all the mass of an atom is concentrated in its nucleus.
- 4.152. A non-uniform bar of weight W is suspended at rest by two strings of negligible weight as 4.157 (a) A child stands at the centre of a turntable shown in Fig. 4.152. The angles made by the strings with the vertical are 36.9° and 53.1° respectively. The bar is 2 m long. Calculate the distance d of the centre of gravity of the bar from its left end.
- 4.153. A car weighs 1800 kg. The distance between its front and back axles is 1.8 m. Its centre of gravity is 1.05 m behind the front axle. Determine the force exerted by the level ground on each front wheel and each back wheel.
- 4.154.(a) Find the moment of inertia of a sphere about a tangent to the sphere, given the moment4.158. A rope of negligible mass is wound round a of inertia of the sphere about any of its diameters to be  $\frac{2MR^2}{5}$  where M is the mass of the sphere and R is the radius of the sphere.
  - (b) Given the moment of inertia of a disc of mass M and radius R about any of its diameters to be  $\frac{MR^2}{4}$ , find its moment of inertia4.159. To maintain a rotor at a uniform angular speed




Fig. 4.152

about an axis normal to the disc and passing through a point on its edge.

- 4.155. Torques of equal magnitude are applied to a hollow cylinder and a solid sphere, both having the same mass and radius. The cylinder is free to rotate about its standard axis of symmetry, and the sphere is free to rotate about an axis passing through its centre. Which of the two will acquire a greater angular speed after a given time.
  - its axis with angular speed 100 rads<sup>-1</sup>. The radius of the cylinder is 0.25 m. What is the kinetic energy associated with the rotation of the cylinder? What is the magnitude of angular momentum of the cylinder about its axis?
  - with his two arms outstretched. turntable is set rotating with an angular speed of 40 rev/min. How much is the angular speed of the child if he folds his hands back and thereby reduces his moment of inertia to 2/5 times the initial value? Assume that the turntable rotates without friction.
    - (b) Show that the child's new kinetic energy of rotation is more than the initial kinetic energy of rotation. How do you account for this increase in kinetic energy?
  - hollow cylinder of mass 3 kg and radius 40 cm. What is the angular acceleration of the cylinder if the rope is pulled with a force of 30 N? What is the linear acceleration of the rope? Assume that there is no slipping.

of 200 rads<sup>-1</sup>, an engine needs to transmit a torque of 180 N m. What is the power required by the engine? (Note: uniform angular velocity in the absence of friction implies zero torque. In practice, applied torque is needed to counter frictional torque). Assume that the engine is 100% efficient.

- 4.160. From a uniform disk of radius R, a circular hole of radius R/2 is cut out. The centre of the hole is at R/2 from the centre of the original disc. Locate the centre of gravity of the resulting flat body.
- 4.161. A metre stick is balanced on a knife edge at its centre. When two coins, each of mass 5 g are put one on top of the other at the 12.0 cm mark, the stick is found to be balanced at 45.0 cm. What is the mass of the metre stick?
- 4.162. A solid sphere rolls down two different inclined planes of the same heights but different angles of inclination.
  - (a) Will it reach the bottom with the same speed in each case?
  - (b) Will it take longer to roll down one plane
    4.167. A man stands on a rotating platform, with
  - (c) If so, which one and why?
- 4.163. A hoop of radius 2 m weighs 100 kg. It rolls along a horizontal floor so that its centre of mass has a speed of 20 cm/s. How much work has to be done to stop it?
- 4.164. The oxygen molecule has a mass of  $5.30 \times 10^{-26}$ kg and a moment of inertia of  $1.94 \times 10^{-46} kgm^2$ about an axis through its centre perpendicular to the lines joining the two atoms. Suppose the mean speed of such a molecule in a gas is 500 m/s and that its kinetic energy of rotation is two thirds of its kinetic energy of translation. Find the average angular velocity of the molecule.
- of angle of inclination 30°. At the bottom of the inclined plane the centre of mass of the cylinder has a speed of 5 m/s.
  - (a) How far will the cylinder go up the plane?
  - (b) How long will it take to return to the bottom?
- 4.166. As shown in Fig. 4.166, the two sides of a step ladder BA and CA are 1.6 m long and hinged at A. A rope DE, 0.5 m is tied half 4.169. Two discs of moments of inertia  $I_1$  and  $I_2$  about way up. A weight 40 kg is suspended from a point F, 1.2 m from B along the ladder BA. Assuming the floor to be frictionless and

neglecting the weight of the ladder, find the tension in the rope and forces exerted by the floor on the ladder. (Take  $g = 9.8m/s^2$ ) (Hint: Consider the equilibrium of each side of the ladder separately.)




Fig. 4.166

- his arms stretched horizontally holding a 5 kg weight in each hand. The angular speed of the platform is 30 revolutions per minute. The man then brings his arms close to his body with the distance of each weight from the axis changing from 90cm to 20cm. The moment of inertia of the man together with the platform may be taken to be constant and equal to  $7.6 \text{ kgm}^2$ .
  - (a) What is his new angular speed? (Neglect friction.)
  - (b) Is kinetic energy conserved in the process? If not, from where does the change come about?
- 4.165. A solid cylinder rolls up an inclined plane 4.168. A bullet of mass 10 g and speed 500 m/s is fired into a door and gets embedded exactly at the centre of the door. The door is 1.0 m wide and weighs 12 kg. It is hinged at one end and rotates about a vertical axis practically without friction. Find the angular speed of the door just after the bullet embeds into it. (Hint: The moment of inertia of the door about the vertical axis at one end is  $\frac{ML^2}{3}$ .)
  - their respective axes (normal to the disc and passing through the centre), and rotating with angular speeds  $\omega_1$  and  $\omega_2$  are brought into

contact face to face with their axes of rotation coincident.

- system?
- (b) Show that the kinetic energy of the combined system is less than the sum of the initial kinetic energies of the two discs. How do you account for this loss in energy? Take4.174. A cylinder of mass 10 kg and radius 15 cm is  $\omega_1 \neq \omega_2$ .
- 4.170. Prove the result that the velocity v of translation of a rolling body (like a ring, disc, cylinder or sphere) at the bottom of an inclined plane of a height h is given by

$$v^2 = \frac{2gh}{1 + \frac{k^2}{R^2}} \tag{4.170.1}$$

using dynamical consideration (i.e. by conthe radius of gyration of the body about its symmetry axis, and R is the radius of the body. The body starts from rest at the top of the plane.

4.171. A disc rotating about its axis with angular speed  $\omega_0$  is placed lightly (without any translational push) on a perfectly frictionless table. The radius of the disc is R. What are the linear velocities of the points A, B and C on the disc shown in Fig. 4.171? Will the disc roll in the direction indicated?




Fig. 4.171

- 4.172. Explain why friction is necessary to make the disc in Fig. 4.171 roll in the direction indicated.
  - (a) Give the direction of frictional force at B, and the sense of frictional torque, before4.177. Two uniform solid spheres of equal radii R, perfect rolling begins.

- (b) What is the force of friction after perfect rolling begins?
- (a) What is the angular speed of the two-disc4.173. A solid disc and a ring, both of radius 10 cm are placed on a horizontal table simultaneously, with initial angular speed equal to  $10 \pi rads^{-1}$ . Which of the two will start to roll earlier? The coefficient of kinetic friction is  $\mu_k = 0.2$ .
  - rolling perfectly on a plane of inclination 30°. The coefficient of static friction  $\mu_s = 0.25$ .
  - (a) How much is the force of friction acting on the cylinder?
  - (b) What is the work done against friction during rolling?
  - (c) If the inclination  $\theta$  of the plane is increased, at what value of  $\theta$  does the cylinder begin to skid, and not roll perfectly?
  - sideration of forces and torques). Note k is4.175. Three equal masses of m kg each are fixed at the vertices of an equilateral triangle ABC.
    - (a) What is the force acting on a mass 2m placed at the centroid G of the triangle?
    - (b) What is the force if the mass at the vertex A is doubled? Take AG = BG = CG = 1 m(see Fig. 4.175)




Fig. 4.175

- 4.176. Find the potential energy of a system of four particles placed at the vertices of a square of side l. Also obtain the potential at the centre of the square.
  - but mass M and 4 M have a centre to centre

separation 6 R, as shown in Fig. 4.177. The two spheres are held fixed. A projectile of mass m is projected from the surface of the sphere of mass M directly towards the centre of the second sphere. Obtain an expression for the minimum speed v of the projectile so that it reaches the surface of the second sphere.




Fig. 4.177