Date: 23/4/2024

Lab Assignment - 1

Signal Generation

Name: Girish S

Roll: AM.EN.U4AIE22044

Course Outcome:


a. Unit Impulse

CO1: Familiarise the basic concepts of communication systems

.1. Plot the following **elementary signals**.


```
b. Unit Step
   c. Ramp
   Continous Signal
t = -1:0.01:1;
impulse_continous = t==0;
step_continous = t >= 0;
ramp_continous = t .* (t >= 0);
subplot(3,1,1);
plot(t, impulse_continous);
title('Continous Time: Unit Impulse Signal');
xlabel('Time (seconds)');
ylabel('Amplitude');
subplot(3,1,2);
plot(t, step_continous);
title('Continous Time: Unit Step Signal');
xlabel('Time (seconds)');
ylabel('Amplitude');
subplot(3,1,3);
plot(t, ramp_continous);
title('Continous Time: Ramp Signal');
xlabel('Time (seconds)');
ylabel('Amplitude');
```

Continous Time: Unit Impulse Signal



Time (eeconde)


```
Discrete Signals
t = -1:0.2:1;
impulse_discrete = zeros(size(t));
impulse_discrete(t == 0) = 1;
step_discrete = zeros(size(t));
step_discrete(t >= 0) = 1;
ramp_discrete = t .* (t >= 0);
subplot(3,1,1);
stem(t, impulse_discrete);
title('Discrete Time: Unit Impulse Signal');
xlabel('Time (seconds)');
ylabel('Amplitude');
subplot(3,1,2);
stem(t, step_discrete);
title('Discrete Time: Unit Step Signal');
xlabel('Time (seconds)');
ylabel('Amplitude');
subplot(3,1,3);
stem(t, ramp_discrete);
title('Discrete Time: Ramp Signal');
xlabel('Time (seconds)');
ylabel('Amplitude');
```



- 2. Generate and plot the following continuous time signals (Define the time vector, t to range from 0 to 1 second, with a sampling rate of 1000 Hz) $\,$
- a. Sinusoidal Signal

Code:


- i. What happens if you increase the frequency? How does it affect the waveform?
- ii. How does changing the amplitude affect the waveform?
- iii. How does changing the phase affects the waveform? (Plot for phi values pi/2, pi/4, pi, -pi/2)



```
b. square signal
C. exponential signal
d. Sawtooth signal
e. Triangular signal
t = 0:0.001:1;
f =5;
A = 1;

A_square = 0.81;
T_square = 0.5;
y_square = A_square * square(2 * pi * (1 / T_square) * t);
subplot(2, 2, 1);
plot(t, y_square);
title('Square Signal');
xlabel('Time (seconds)');
ylabel('Amplitude');
```

```
y_{exp} = exp(-2 * t);
subplot(2, 2, 2);
plot(t, y_exp);
title('Exponential Signal');
xlabel('Time (seconds)');
ylabel('Amplitude');
y_sawtooth = sawtooth(2*pi*f*t);
subplot(2, 2, 3);
plot(t, y_sawtooth);
title('Sawtooth Signal');
xlabel('Time (seconds)');
ylabel('Amplitude');
y_triangular = sawtooth(2 * pi * f*t, 0.5);
subplot(2, 2, 4);
plot(t, y_triangular);
title('Triangular Signal');
xlabel('Time (seconds)');
ylabel('Amplitude');
```



```
3. Generate and plot the following signals. (Signal Operations)
      a. x1(t) = sin(2*pi*t/T).*exp(-2*t)
      b. x2(t) = 2*cos(2*pi*t/T2).*sin(2*pi*t/T3)
      c. x3(t) = sin(2*pi*t/T).*exp(-2*t)+sin(2*pi*t/T1).*exp(-4*t)
      d. x4(t) = aa*u(t), (take aa=5, u(t) is step signal, Plot both u(t) and
      x4(t))
figure;
t = 0:0.01:5;
T = 2;
T2 = 2;
T3 = 3;
T1 = 3;
aa = 5;
u_t = t >= 3;
x1 = \sin(2*pi*t/T) .* \exp(-2*t);
x2 = 2 * cos(2*pi*t/T2) .* sin(2*pi*t/T3);
x3 = \sin(2*pi*t/T) .* \exp(-2*t) + \sin(2*pi*t/T1) .* \exp(-4*t);
x4 = aa * u_t;
subplot(3, 2, 1);
plot(t, x1);
title('x1(t) = sin(2*pi*t/T).*exp(-2*t)');
xlabel('Time');
ylabel('Amplitude');
subplot(3, 2, 2);
plot(t, x2);
title('x2(t) = 2*cos(2*pi*t/T2).*sin(2*pi*t/T3)');
xlabel('Time');
ylabel('Amplitude');
subplot(3, 2, 3);
plot(t, x3);
title('x3(t) = sin(2*pi*t/T).*exp(-2*t)+sin(2*pi*t/T1).*exp(-4*t)');
xlabel('Time');
ylabel('Amplitude');
subplot(3, 2, 4);
plot(t, u_t, 'r', t, x4, 'b');
title('x4(t) = aa*u(t)');
xlabel('Time');
ylabel('Amplitude');
legend('u(t)', 'x4(t)');
```

