



Competitive Programming

From Problem 2 Solution in $O(1)$

Combinatorics

Pigeonhole Principle

Mostafa Saad Ibrahim

PhD Student @ Simon Fraser University



Pigeonhole Principle

- Assume we have $N=5$ boxes and $M=6$ apples
- Distribute apples in boxes in whatever way
- There must be 1 box with at least 2 apples
- If you want to avoid duplicate:
 - Put 1 apple per box. Remaining is 1 apple.
 - To put it, one box will have 2 apples
- What if we have 13 apples?
 - 1 box will have at least $\lceil 13/5 \rceil = 3$ apples
- Generally: $\lceil N/M \rceil$ per a box.

Many proofs based on it

- The set S has 10 integers, each between 0 and 100. Prove that there are two disjoint subsets of S that have the same sum.
- Given a set of N integers, there is a consecutive subset of them whose sum is divisible by N .
- Every sequence of $n^2 + 1$ distinct real numbers contains a subsequence of length $n + 1$ that is either strictly increasing or strictly decreasing.
- Show that among any $n + 1$ positive integers not exceeding $2n$ there must be an integer that divides one of the other integers.
- How many cards must be selected from a standard deck of 52 cards to guarantee that at least three cards of the same suit are chosen?
- Among any group of 367 people, there must be at least two with the same birthday
- Show that for every integer n there is a multiple of n that has only 0s and 1s in its decimal expansion.

Pigeonhole Principle: Example

- **Prove:** Among any N positive integers, there exists 2 whose difference is divisible by $N-1$.
- Recall: $|A-B| \% X = 0$ IFF $A \% X = B \% X$
- So, let's compute $\% N-1$?
 - Then we have $N-1$ values, each in range $[0-N-2]$
 - But, we have N numbers, then at least one mode will be duplicate
 - Given that 2 numbers at least has same $\% N-1$
 - Then, their difference must be divisible by $N-1$
- **Always** relate $\%$ with Pigeonhole

Pigeonhole Principle: Example

- Case for previous problem:
- Let A represents array of numbers
- $N = 5$ and $A = \{2, 3, 5, 7, 8\}$
- Compute $A \% 4 = \{2, 3, 1, 3, 0\}$
- 5 numbers, with values [0-3].
- Pick 2 with same mode - 3 is **repeated** mode
- Then, (3, 7) are the answer

Pigeonhole Principle: Example

- Prove: For any N positive integers, the **sum** of **subset** of them is divisible N
- Compute Accum array $\% N$
 - $\text{Accum}[i] = \{A[0] + A[1] \dots A[i]\} \% N$
- If any $\text{Accum}[i] = 0$, we are done
- Otherwise, we have N values in range $[1-N-1]$
- Then 2 positions will have **same mode**
- Then getting numbers between then is answer

Pigeonhole Principle: Example

- Let A represents array of numbers
- $N = 5$ and $A = \{2, 4, 8, 2, 7\}$
- Accumulate: $B = \{2, 6, 14, 16, 23\}$
- Mode 5: $C = \{2, 1, 4, 1, 3\}$
- Any zeros? No..remaining 4 values spread on 5 values...one of them must be repeated
 - if yes then $A[0]+A[1]...A[i]$ where $C[i] \% N = 0$
- 2nd and 4th have mode 1
- Then range from 3rd till 4th is answer: 8, 2

Pigeonhole and Competitions

- Most of time it helps in **proving**, rather than a technique to apply
 - Read many problem [examples](#) (web/books) + proofs
- In some problems, it can be the major trick
- Sometimes comes with Modular Arithmetic
- Some facts in graph:
 - A Path of M nodes ($M > N$) must have a node used more than once
 - A cycle of Length M ($M > N$ nodes), must be composed of cycles each of Length $\leq N$
 - Every graph contains two vertices of same degree

Let's try a competitive example..direct case

Powers tower % M

- Let's compute: $2^{3^4^5^6^7^8} \% 56$
 - We can solve it using Euler theorem
- let's simplify it, compute $2^x \% 56$
 - where x is very large, e.g. $x = 3^4^5^6^7^8$
- Imagine we compute $2^{i \% M}$ for $i [0 - \infty]$
 - We know we have M mod values: $[0 - M-1]$
 - Pigeonhole: values repeat in **maximum** $M + 1$ iterations
 - Then computing X should have **same value** as one of the first powers in range $[0-M-1]$
 - But which $2^{i \% M}$ correspond to $2^{x \% M}$?
- Let's simulate it

Powers tower % M

i	0	1	2	3	4	5	6	7
$2^i \% 56$	1	2	4	<u>8</u>	16	32	<u>8</u>	16

- 2^6 is same as 2^3 . Then 2^7 must = 2^4 ...etc
- $\{8,16,32\}$ is **cycle** and $\{1,2,4\}$ is **precycle**
 - Let length of the cycle be L, and length of precycle be P
- Given some X, we can compute its i position
 - $i = P + (L + \underline{X \% L} - P \% L) \% L$ [if $X > P$]
- Then?
 - Solve subproblem $X \% L$ in same manner
 - Then compute $2^i \% M$

تم بحمد الله

علمكم الله ما ينفعكم

ونفعكم بما تعلمتم

وزادكم علماً

Problems

- UVA (11237), SRM283-1-2(FactorialTower),
Hackerrank(A Perfect Set - GoDaddy
Hackathon),