

7 Factor Analysis: Theoretical Models

MA2003B Application of Multivariate Methods
in Data Science

Raul Gomez



Factor Analysis: Theoretical Models

Factor Analysis is a statistical technique that is widely used in applied fields to identify hidden patterns or latent structures behind large sets of observed variables.

Factor Analysis: Theoretical Models

Factor Analysis is a statistical technique that is widely used in applied fields to identify hidden patterns or latent structures behind large sets of observed variables.

The goal is to reduce complexity by grouping related variables into a smaller number of underlying **factors**, which are easier to interpret.

Some areas where factor analysis is commonly applied include:

- **Psychology and social sciences:** to uncover latent traits such as intelligence, anxiety, personality dimensions, or satisfaction levels, based on responses to questionnaires or surveys.

Some areas where factor analysis is commonly applied include:

- **Psychology and social sciences:** to uncover latent traits such as intelligence, anxiety, personality dimensions, or satisfaction levels, based on responses to questionnaires or surveys.
- **Marketing and consumer research:** to identify purchasing motives, brand perceptions, or market segments from customer behavior data.

Some areas where factor analysis is commonly applied include:

- **Psychology and social sciences:** to uncover latent traits such as intelligence, anxiety, personality dimensions, or satisfaction levels, based on responses to questionnaires or surveys.
- **Marketing and consumer research:** to identify purchasing motives, brand perceptions, or market segments from customer behavior data.
- **Finance and economics:** to reveal hidden drivers of stock returns, macroeconomic indicators, or risk factors.

Some areas where factor analysis is commonly applied include:

- **Psychology and social sciences:** to uncover latent traits such as intelligence, anxiety, personality dimensions, or satisfaction levels, based on responses to questionnaires or surveys.
- **Marketing and consumer research:** to identify purchasing motives, brand perceptions, or market segments from customer behavior data.
- **Finance and economics:** to reveal hidden drivers of stock returns, macroeconomic indicators, or risk factors.
- **Education:** to analyze test items and determine whether they measure common skills or abilities.

Some areas where factor analysis is commonly applied include:

- **Psychology and social sciences:** to uncover latent traits such as intelligence, anxiety, personality dimensions, or satisfaction levels, based on responses to questionnaires or surveys.
- **Marketing and consumer research:** to identify purchasing motives, brand perceptions, or market segments from customer behavior data.
- **Finance and economics:** to reveal hidden drivers of stock returns, macroeconomic indicators, or risk factors.
- **Education:** to analyze test items and determine whether they measure common skills or abilities.
- **Biological and health sciences:** to find latent structures in genetic data, symptom checklists, or patient-reported outcomes.

In practice, Factor Analysis is used when researchers suspect that a few unobserved dimensions are responsible for the correlations among many observed variables.

In practice, Factor Analysis is used when researchers suspect that a few unobserved dimensions are responsible for the correlations among many observed variables.

It is especially useful for building measurement scales, data reduction, and gaining insights into the structure of complex datasets.

Factor Analysis Setting

The theoretical context for factor analysis is a relation of the form

$$X = \mu + LF + \varepsilon$$

where:

- X is a random vector of p observed variables (indicators).

Factor Analysis Setting

The theoretical context for factor analysis is a relation of the form

$$X = \mu + LF + \varepsilon$$

where:

- X is a random vector of p observed variables (indicators).
- μ is a constant vector of dimension p .

Factor Analysis Setting

The theoretical context for factor analysis is a relation of the form

$$X = \mu + LF + \varepsilon$$

where:

- X is a random vector of p observed variables (indicators).
- μ is a constant vector of dimension p .
- L is a $p \times m$ constant matrix known as the matrix of **factor loadings**.

Factor Analysis Setting

The theoretical context for factor analysis is a relation of the form

$$X = \mu + LF + \varepsilon$$

where:

- X is a random vector of p observed variables (indicators).
- μ is a constant vector of dimension p .
- L is a $p \times m$ constant matrix known as the matrix of **factor loadings**.
- $F \sim \mathcal{N}(0, I_m)$ is a vector of m latent factors (unobserved variables).

Factor Analysis Setting

The theoretical context for factor analysis is a relation of the form

$$X = \mu + LF + \varepsilon$$

where:

- X is a random vector of p observed variables (indicators).
- μ is a constant vector of dimension p .
- L is a $p \times m$ constant matrix known as the matrix of **factor loadings**.
- $F \sim \mathcal{N}(0, I_m)$ is a vector of m latent factors (unobserved variables).
- $\varepsilon \sim \mathcal{N}(0, \Psi)$ is a vector of unique errors (specific variances), with Ψ diagonal.

Factor Analysis Setting

The theoretical context for factor analysis is a relation of the form

$$X = \mu + LF + \varepsilon$$

where:

- X is a random vector of p observed variables (indicators).
- μ is a constant vector of dimension p .
- L is a $p \times m$ constant matrix known as the matrix of **factor loadings**.
- $F \sim \mathcal{N}(0, I_m)$ is a vector of m latent factors (unobserved variables).
- $\varepsilon \sim \mathcal{N}(0, \Psi)$ is a vector of unique errors (specific variances), with Ψ diagonal.
- F and ε are assumed to be independent, in particular $\text{Cov}[F, \varepsilon] = 0$.

As a consequence of these assumptions, we have that

$$\mathbb{E}[X] = \mathbb{E}[\mu + LF + \varepsilon] = \mathbb{E}[\mu] + L\mathbb{E}[F] + \mathbb{E}[\varepsilon] = \mu$$

As a consequence of these assumptions, we have that

$$\mathbb{E}[X] = \mathbb{E}[\mu + LF + \varepsilon] = \mathbb{E}[\mu] + L\mathbb{E}[F] + \mathbb{E}[\varepsilon] = \mu$$

and

$$\begin{aligned}\text{Var}[X] &= \text{Var}[LF + \varepsilon] \\ &= \mathbb{E}[(LF + \varepsilon)(LF + \varepsilon)^T] \\ &= \mathbb{E}[LFF^T L^T] + \mathbb{E}[\varepsilon\varepsilon^T] + \mathbb{E}[LF\varepsilon^T] + \mathbb{E}[\varepsilon F^T L^T] \\ &= L\mathbb{E}[FF^T]L^T + \mathbb{E}[\varepsilon\varepsilon^T] + L\mathbb{E}[F\varepsilon^T] + \mathbb{E}[\varepsilon F^T]L^T \\ &= LI_m L^T + \Psi + L0 + 0L^T \\ &= LL^T + \Psi\end{aligned}$$

As a consequence of these assumptions, we have that

$$\mathbb{E}[X] = \mathbb{E}[\mu + LF + \varepsilon] = \mathbb{E}[\mu] + L\mathbb{E}[F] + \mathbb{E}[\varepsilon] = \mu$$

and

$$\begin{aligned}\text{Var}[X] &= \text{Var}[LF + \varepsilon] \\ &= \mathbb{E}[(LF + \varepsilon)(LF + \varepsilon)^T] \\ &= \mathbb{E}[LFF^T L^T] + \mathbb{E}[\varepsilon\varepsilon^T] + \mathbb{E}[LF\varepsilon^T] + \mathbb{E}[\varepsilon F^T L^T] \\ &= L\mathbb{E}[FF^T]L^T + \mathbb{E}[\varepsilon\varepsilon^T] + L\mathbb{E}[F\varepsilon^T] + \mathbb{E}[\varepsilon F^T]L^T \\ &= LL_m L^T + \Psi + L0 + 0L^T \\ &= LL^T + \Psi\end{aligned}$$

Furthermore, since X is a linear combination of normally distributed variables, it follows that

$$X \sim \mathcal{N}(\mu, LL^T + \Psi)$$

The variance decomposition $\text{Var}[X] = LL^T + \Psi$ shows that the total variance of the observed variables X can be decomposed into two components:

The variance decomposition $\text{Var}[X] = LL^T + \Psi$ shows that the total variance of the observed variables X can be decomposed into two components:

- **Common variance (LL^T):** This part of the variance is explained by the common factors F . It captures the shared variability among the observed variables that can be attributed to the underlying latent structure.

The variance decomposition $\text{Var}[X] = LL^T + \Psi$ shows that the total variance of the observed variables X can be decomposed into two components:

- **Common variance (LL^T):** This part of the variance is explained by the common factors F . It captures the shared variability among the observed variables that can be attributed to the underlying latent structure.
- **Unique variance (Ψ):** This part of the variance is specific to each observed variable and is not explained by the common factors. It includes measurement errors and any other variability that is unique to each variable.

The variance decomposition $\text{Var}[X] = LL^T + \Psi$ shows that the total variance of the observed variables X can be decomposed into two components:

- **Common variance (LL^T):** This part of the variance is explained by the common factors F . It captures the shared variability among the observed variables that can be attributed to the underlying latent structure.
- **Unique variance (Ψ):** This part of the variance is specific to each observed variable and is not explained by the common factors. It includes measurement errors and any other variability that is unique to each variable.

In particular, the variance of each individual observed variable X_i can be expressed as

$$\text{Var}[X_i] = \sum_{j=1}^m L_{ij}^2 + \sigma_i^2$$

where $h_i^2 = \sum_{j=1}^m L_{ij}^2$ is called the **communality** of the variable X_i , representing the portion of its variance explained by the common factors, and σ_i^2 is the unique variance specific to X_i .

Factor Score Estimation

Assume that we have a model

$$X = \mu + LF + \varepsilon$$

with the same conditions as before.

Factor Score Estimation

Assume that we have a model

$$X = \mu + LF + \varepsilon$$

with the same conditions as before.

Then the joint distribution of X and F is given by

$$\begin{bmatrix} X \\ F \end{bmatrix} \sim \mathcal{N} \left(\begin{bmatrix} \mu \\ 0 \end{bmatrix}, \begin{bmatrix} LL^T + \Psi & L \\ L^T & I_m \end{bmatrix} \right)$$

Factor Score Estimation

Assume that we have a model

$$X = \mu + LF + \varepsilon$$

with the same conditions as before.

Then the joint distribution of X and F is given by

$$\begin{bmatrix} X \\ F \end{bmatrix} \sim \mathcal{N} \left(\begin{bmatrix} \mu \\ 0 \end{bmatrix}, \begin{bmatrix} LL^T + \Psi & L \\ L^T & I_m \end{bmatrix} \right)$$

Notice that, in particular,

$$\text{Cov}[X, F] = \mathbb{E}[(X - \mu)(F - 0)^T] = \mathbb{E}[(LF + \varepsilon)F^T] = L\mathbb{E}[FF^T] + \mathbb{E}[\varepsilon F^T] = LI_m + 0 = L$$

Using the properties of the multivariate normal distribution, we can derive the conditional distribution of the factors F given the observed variables X .

Using the properties of the multivariate normal distribution, we can derive the conditional distribution of the factors F given the observed variables X .

This conditional distribution is also normal, with mean and covariance given by

$$[F|X = x] \sim \mathcal{N} \left(L^T (LL^T + \Psi)^{-1} (x - \mu), I_m - L^T (LL^T + \Psi)^{-1} L \right)$$

The mean of this conditional distribution,

$$\mathbb{E}[F|X = x] = L^T(LL^T + \Psi)^{-1}(x - \mu)$$

is known as the **factor score** for the observed data point $X = x$.

The mean of this conditional distribution,

$$\mathbb{E}[F|X = x] = L^T (LL^T + \Psi)^{-1}(x - \mu)$$

is known as the **factor score** for the observed data point $X = x$.

It provides an estimate of the latent factors based on the observed variables and the factor loadings.

The mean of this conditional distribution,

$$\mathbb{E}[F|X = x] = L^T (LL^T + \Psi)^{-1}(x - \mu)$$

is known as the **factor score** for the observed data point $X = x$.

It provides an estimate of the latent factors based on the observed variables and the factor loadings.

The covariance matrix

$$I_m - L^T (LL^T + \Psi)^{-1} L$$

represents the uncertainty associated with the factor score estimates.

Remark

Notice that, if

$$f = \mathbb{E}[F|X = x] = L^T(LL^T + \Psi)^{-1}(x - \mu)$$

then, in general, the **residual**

$$r(x) = x - \mu - Lf$$

is different from zero.

Remark

Notice that, if

$$f = \mathbb{E}[F|X = x] = L^T(LL^T + \Psi)^{-1}(x - \mu)$$

then, in general, the **residual**

$$r(x) = x - \mu - Lf$$

is different from zero.

In fact, as a function of the random variable X , the residual has probability distribution

$$r(X) \sim \mathcal{N}\left(0, L(I_m + L^T\Psi^{-1}L)^{-1}L^T + \Psi\right)$$

Rotation of Factors

An important aspect of factor analysis is that the factors are not uniquely defined.

Rotation of Factors

An important aspect of factor analysis is that the factors are not uniquely defined.

If F is a vector of factors, then any orthogonal transformation $F^* = QF$, where Q is an orthogonal matrix (i.e., $Q^T Q = I$), will yield a new set of factors that explain the same variance in the observed variables.

Rotation of Factors

An important aspect of factor analysis is that the factors are not uniquely defined.

If F is a vector of factors, then any orthogonal transformation $F^* = QF$, where Q is an orthogonal matrix (i.e., $Q^T Q = I$), will yield a new set of factors that explain the same variance in the observed variables.

In particular,

$$\text{Var}[F^*] = \text{Var}[QF] = Q \text{Var}[F] Q^T = Q I_m Q^T = I_m$$

Rotation of Factors

An important aspect of factor analysis is that the factors are not uniquely defined.

If F is a vector of factors, then any orthogonal transformation $F^* = QF$, where Q is an orthogonal matrix (i.e., $Q^T Q = I$), will yield a new set of factors that explain the same variance in the observed variables.

In particular,

$$\text{Var}[F^*] = \text{Var}[QF] = Q \text{Var}[F] Q^T = Q I_m Q^T = I_m$$

The corresponding factor loadings will be transformed as

$$L^* = LQ^T$$

In other words, if

$$X = \mu + LF + \varepsilon$$

is a valid factor model, then so is

$$X = \mu + L^*F^* + \varepsilon = \mu + LQ^T QF + \varepsilon = \mu + LF + \varepsilon$$

In other words, if

$$X = \mu + LF + \varepsilon$$

is a valid factor model, then so is

$$X = \mu + L^*F^* + \varepsilon = \mu + LQ^T QF + \varepsilon = \mu + LF + \varepsilon$$

This means that the solution to a factor analysis problem is not unique, and different rotations of the factors can lead to different interpretations.

In other words, if

$$X = \mu + LF + \varepsilon$$

is a valid factor model, then so is

$$X = \mu + L^*F^* + \varepsilon = \mu + LQ^T QF + \varepsilon = \mu + LF + \varepsilon$$

This means that the solution to a factor analysis problem is not unique, and different rotations of the factors can lead to different interpretations.

In the next presentation, we will explore various practical methods for factor analysis and rotation techniques geared towards achieving more interpretable factor solutions.