BENTHAM  
SCIENCE

## Next-generation Metabolomics in the Development of New Antidepressants: Using Albiflorin as an Example

Jian Han<sup>1#</sup>, Yonghong Xia<sup>2#</sup>, Lejun Lin<sup>3#</sup>, Zuoguang Zhang<sup>4</sup>, Hui Tian<sup>4</sup> and Kefeng Li<sup>5,\*</sup>

<sup>1</sup>Department of Orthopaedic Oncology, Yantaishan Hospital, Yantai, Shandong, 264000, China; <sup>2</sup>Intensive Care Unit, Yantai Yuhuangding Hospital, Yantai, Shandong, 264000, China; <sup>3</sup>Department of Nuclear Medicine, Yantai Yuhuangding Hospital, Yantai, Shandong, 264000, China; <sup>4</sup>Beijing Wonner Biotech. Co. Ltd., Beijing, 101111, China; <sup>5</sup>School of Medicine, University of California, San Diego, CA, 92103, USA

### ARTICLE HISTORY

Received: June 26, 2018  
Accepted: July 26, 2018

DOI:

10.2174/1381612824666180727114134

**Abstract:** Depression is a highly prevalent disorder that affects more than 300 million adults worldwide in 2015. Depression also frequently coexists with many other conditions such as osteoporosis and one-third of the Intensive Care Unit (ICU) survivors had depressive symptoms. Antidepressants have become the most commonly prescribed drugs in the United States. In addition to the regular process, drug discovery and development (R&D) for depression presents extra challenges because of the heterogeneity of the symptoms and various co-occurring disorders. Botanical medicine with multi-functional nature has been proposed to be more effective, providing rapid control of core and comorbid conditions of depression. With the technical advances in analytical instruments, metabolomics is entering into a “new generation”. Next-generation metabolomics (NGM) has the capability to comprehensively characterize drug-induced metabolic changes in the biological systems. NGM has demonstrated great potential in all the stages of pharmaceutical R&D in the last 10 years. Albiflorin isolated from Peony roots is a promising drug candidate with multi-target for depression and is currently under development by Beijing Wonner Biotech. In this work, we summarized the common analytical platforms for NGM and its main applications in drug R&D. We used albiflorin as an example to illustrate how NGM improves our understanding of drug candidate actions and facilitates drug safety evaluation. Future directions on how to expand the use of NGM for new antidepressant development in pharmaceutical industry were also discussed.


**Keywords:** Metabolomics, antidepressants, drug development and discovery, albiflorin, depression, next-generation metabolomics (NGM).

### 1. INTRODUCTION

Metabolomics is the new “omics” technology added to the system biology toolbox in the post-genomic era [1]. Metabolomics uses various advanced analytical chemistry approaches to characterize the metabolites and metabolism in biological systems. Metabolites are the direct “readout” of gene-environment interactions (G x E = metabolism). Unlike genomics, transcriptomics, and proteomics that can only tell us what might happen, metabolomics reflects what is currently happening. Metabolomics has broad applications in many diverse areas such as diseases pathogenesis and diagnosis, biomarker screening, drug discovery, precision medicine, plant biology, microbiology, nutrition and environmental monitoring [2-7].

Depression is one of the most common diseases and the leading cause of disability. It was reported that depression affects at least 4.4% of the world population in 2015 [8]. Depression is a multifactorial disorder with the abnormalities of cognitive functions, sleep and gut microbiome [9]. The development of drugs for major depressive disorder is particularly challenging due to the complexity of the symptoms, heterogeneity and comorbid diagnosis. The major classes of approved antidepressants include Selective Serotonin Reuptake Inhibitors (SSRIs), Tricyclic Antidepressants (TCAs) and monoamine oxidase inhibitors (MAOIs) [10]. However, current antidepressants are limited by their undesirable side effects and slow onset [11]. In addition, nearly 50% of patients do not respond to the first medication prescribed [12]. There is an urgent need for the development of new antidepressants.

Medicinal plants are rich reservoirs for new drugs and drug leads. Over the past decade, several medicinal plants and formulations have been used for the treatment of depression such as turmeric (*Curcuma longa*), St John's wort (*Hypericum perforatum*), white peony (*Paeonia lactiflora* Pall.) and lavender (*Lavandula* spp.) [13]. The dry root of white peony has been used as herbal medicine in China, Korea, and Japan for the treatment of major depressive disorder for hundreds of years [14]. Albiflorin is one of the main bioactive molecules isolated from the roots of white peony (Fig. 1). Recently, albiflorin was shown to be a promising new antidepressant [15]. Unlike the highly selective single target antidepressants, albiflorin is a novel serotonin (5-HT) and norepinephrine (NE) reuptake inhibitor in the hippocampus [16, 17]. The dopaminergic system also contributes to albiflorin-mediated antidepressant-like activity [17]. Given the heterogeneous nature of depression, multi-target agents like albiflorin have been proposed to be more effective to simultaneously control core and comorbidities of depression than the classical antidepressants with highly specific and selective single target on neurotransmitters [18].



Albiflorin

**Fig. (1).** Chemical structure of albiflorin isolated from the roots of *Paeonia lactiflora* Pall.

\*Address correspondence to this author at the School of Medicine, University of California, San Diego, CA, 92103, USA; E-mail: [kli@ucsd.edu](mailto:kli@ucsd.edu)

<sup>#</sup>These authors contributed equally.

The use of metabolomics in drug discovery and development is also called pharmacometabolomics. The traditional drug discovery and development process are costly and time-consuming. A standard process takes more than 10 years from the beginning to the market and costs over \$800 million with the success rate as low as 7% [19] (Fig. 2). Pharmacometabolomics dramatically facilitates this process and reduces the overall cost. It can be useful in all stages of drug discovery and development, providing holistic information for the determination of drug targets, models of actions, safety and toxicity evaluation (Fig. 2). Pharmacometabolomics is particularly essential for drugs with multi-targets which even take longer time than the development of a single target drug using traditional drug discovery approach.

In this work, we summarized the role of metabolomics in the discovery and development of antidepressants, particularly alflorin, in relation to novel antidepressants with multi-targets from medicinal plants. Since the use of metabolomics in target identification, chemical library screening and molecular mechanisms of drugs had been extensively discussed recently [1, 20, 21], here, we focused on its application in drug toxicity and safety evaluation.

## 2. NEXT-GENERATION METABOLOMICS (NGM)

The origins of metabolomics can be traced back to the newborn screening of inborn errors of metabolism in clinical chemistry since the 1960s, which was much further than genomics and proteomics [22]. The term “metabolome” was coined in 1998 by Steven Oliver in a review article [23]. The first metabolomics paper was published by the Nicholson group in 1999 [24]. In the first ten years of 21 century (2000-2010), the early applications of metabolomics were primarily conducted using  $^1\text{H}$  nuclear magnetic resonance spectroscopy ( $^1\text{H}$  NMR) and gas chromatography coupled to mass spectrometry (GC-MS) [25, 26]. However, metabolite measurements using NMR and GC-MS are restrained by the chemical diversity of metabolites and their broad dynamic range in cellular abundance [27]. NMR has relatively low sensitivity (limit of detection  $>5\ \mu\text{M}$ )

and cannot detect non-protonated metabolites [28]. GC-MS can only detect volatile compounds and require sophisticated sample derivatization steps for other non-volatile metabolites [4, 29]. In addition, lack of fragmentation information by GC-MS makes the identification of metabolites less accurate [4]. In the last 10 years (2008-now), with the advancements of separation and mass spectrometry detection techniques, NGM was developed on several different platforms [4, 30, 31]. These platforms provide a broad coverage of different classes of metabolites, superior detection sensitivity, high resolution and mass accuracy [31-35]. Single-cell metabolomics and imaging metabolomics were also developed in recent years [36, 37].

The brief workflow for NGM is shown in Fig. 3. A careful experimental design is essential for the success of a metabolomic study. Enough biological replicates should be prepared in order to draw the reliable conclusion. For human subjects, at least 15 subjects should be recruited for each group and gender. Age and gender should be matched between controls and diseases (treatment). The minimum sample size for animals and plants is 8 per group. The minimum sample size for *in vitro* cells is 6 per group. For cells, it usually requires at least one million cells depending on the sensitivity of the instruments. Metabolites are extracted from the samples using different extraction methods including organic solvents deproteinization and liquid-liquid extraction [38]. Metabolites are then separated and detected. The peaks are checked and statistical analysis is conducted.

Table 1 and Fig. 4 summarize the applications of different platforms for NGM. Due to the high chemical diversity of the metabolites and the limited scan speed for the detectors, separation is usually required for metabolomics. GC, Liquid Chromatography (LC) and Capillary Electrophoresis (CE) are the common separation techniques used in NGM [39]. Occasionally, two-dimensional separation by GC $\times$ GC or LC $\times$ LC was required [40]. In general, GC is a method of choice for nonpolar or volatile and thermal stable metabolites such as fatty acids. Polar metabolites can also be analyzed by GC after derivatization [41]. LC is the most versatile separation





Fig. (2). Metabolomics facilitates all the stages of drug discovery and development.

method, allowing the separation of compounds of a wide range of polarity [42]. HILIC phase is useful for the analysis of polar metabolites such as amino acids, organic acids, and sugars, while, nonpolar and weak polar metabolites can be separated by reverse phase columns (C18, biphenyl, *etc.*). Derivatization is generally not required for LC analysis. CE has high separation efficiency for polar ionic metabolites such as carbohydrates and amino acids. In addition, CE is especially suitable for volume-restricted biological samples such as spinal fluid and single cell extract [35, 43].



**Fig. (3).** The schematic representation of next-generation metabolomics workflow.

Mass Spectrometry (MS) is the most common detection approach for NGM due to its high sensitivity, selectivity and potential to identify the metabolites [44]. In the last few years, various MS analyzers were commercialized by different vendors including the time of flight MS (TOF/MS, LecoCorp), quadrupole TOF (Q-TOF,

Agilent), triple quadrupole (MS/MS, SCIEX, Shimadzu, and Waters) and orbitrap (Thermo) [45]. High-resolution TOF instruments coupled to GC, LC or CE are often used for untargeted metabolomics, while, triple quadrupole (QQQ) instruments are applied for targeted metabolomic analysis.

Metabolomics at the single-cell level is essential for the analysis of phenotypic heterogeneity between cells such as neurons in which individual differences are required for function [63]. The cells are sampled by fine-tip glass capillaries, Laser Capture Microdissection (LCM), optical trapping, Fluorescence-Activated Cell Sorting (FACS) and Magnetic-Activated Cell Sorting (MACS) depending on the cell types [64]. Due to the limited volume size for single cell extract, CE and nano-flow LC are the main front-end separation techniques for single cell metabolomics [65, 66]. MS is the major technique for detection as other metabolomics.

Imaging metabolomics is also a powerful tool to explore the model of actions of drugs, especially for drug candidates with multitarget [37]. Several well-developed mass spectrometry imaging (MSI) techniques are currently in use for the performance of imaging metabolomics such as high-resolution matrix-assisted laser desorption ionization (MALDI)-MSI [67, 68], TOF-secondary ionization mass spectrometry (TOF-SIMS) [69], desorption electrospray ionization (DESI)-MSI under ambient conditions [70, 71] and air-flow-assisted desorption electrospray ionization [72] (Fig. 4). Both molecular and spatial information can be obtained simultaneously by imaging metabolomics.

### 3. METABOLOMICS IN EXPLORING THE MECHANISM OF ACTION (MOA) OF ANTIDEPRESSANTS

To minimize the problems in downstream steps such as pre-clinical and clinical trials, it is desirable that MOA of any drug candidates is established during the drug discovery process [73]. Depression has a strong metabolic basis or a clear metabolic cause such as the metabolic perturbations in tryptophan, kynurenine and mitochondrial metabolism [74-76]. The discovery of the metabolic basis of depression often leads to the development of new therapeutic solutions. Metabolomics thus might provide a far more cost-effective and productive approach to elucidate MOA of antidepressants.


Some software and tools used for dissecting the MOA of drugs have been developed. Metaboanalyst ([www.metaboanalyst.ca](http://www.metaboanalyst.ca)) and Metlin (<https://metlin.scripps.edu>) are free online tools for comprehensive metabolomic data analysis [77, 78]. Ingenuity Pathway Analysis (IPA)/KEGG pathway analysis is a licensed software for the integration of multi-omic data including metabolomics, proteomics and transcriptomics.

Recently, metabolomics has been successfully applied to identify the molecular mechanisms of drug or drug candidates with

**Table 1.** Next-generation metabolomics platforms.

Separation Techniques		MS (Vendors)	Main Applications	References
GC or GC×GC	DB-5MS (Agilent); RTX-5MS (Restek)	TOF/MS (LecoCorp); Q-TOF (Agilent); Orbitrap (Thermo); oa-TOF(Waters)	Fatty acids, volatiles; nonpolar; polar compounds (requires derivatization)	[46-50]
LC	HILIC (amino, amide, diol, etc)	MS/MS (triple quadrupole, SCIEX, Shimadzu, Waters); TripleTOF (SCIEX); Q-TOF (Agilent); Orbitrap (Thermo)	Polar metabolites such as amino acids, organic acids, and sugars	[51, 52]
	RPLC (C18, C8, Biphenyl, etc)		Nonpolar and weak polar metabolites such as phospholipids and ceramides	[53-56]
CE	Fused-silica capillaries	TOFMS (Agilent); Orbitrap (Thermo); microTOF (Bruker); MS/MS (SCIEX)	Polar and charged metabolites such as carbohydrates and amino acids	[57-62]

Abbreviations: HILIC: Hydrophilic interaction liquid chromatography; RPLC: reverse phase; MS: mass spectrometry; TOF: Time of flight



**Fig. (4). The next-generation metabolomics platforms.** Gas chromatography: GC; Capillary electrophoresis: CE; Liquid chromatography: LC; Mass spectrometry: MS; Time of flight: TOF; Triple quadrupole: QQQ; matrix-assisted laser desorption/ionization: MALDI; TOF-secondary ionization mass spectrometry: TOF-SIMS; Desorption electrospray ionization: DESI; Laser ablation electrospray ionization: LAESI.

antidepressant activity. Ketamine is a promising fast-acting antidepressant. By GC-MS based metabolomics, Liang and the coauthors showed that ketamine exerts its antidepressant effects through an alpha-amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid (AMPA) inhibition-dependent mechanism [79, 80]. Metabolomics revealed that the antidepressant action of diterpene ginkgolides is mainly related to amino acid, energy and lipid metabolism [81, 82]. The mechanism of antidepressant activity of chlorogenic acid is found to be involved in regulating the abnormal pathways of nicotine and nicotinamide metabolism [83]. In addition, the antidepressive effects of Yangxinshi Tablet [84], Chaihu-Shu-Gan-San [85], venlafaxine [86] and scopolamine [87] were investigated by NGM.

### 3.1. Case Study 1-Metabolomic Profiling of Antidepressant Drug Action of Albiflorin


Here, we used albiflorin to demonstrate how metabolomics-based drug discovery may be applied to elucidate MOA of antidepressants. The rat model of depression was generated through 4-week exposure to Chronic Mild Stress (CMS) to male Sprague Dawley (SD) rats. Rats with CMS were then treated with albiflorin (7 mg/kg/d) or saline for 7 days. Plasma samples were collected before and after albiflorin treatment. Using the next-generation broad-spectrum targeted metabolomics [88], we found that plasma metabolites displayed dramatic differences between CMS + albiflorin and CMS + saline (Fig. 5A). We discovered a higher level of acetylcarnitine, 3-Hydroxydodecanoylcarnitine as well as myristoyl carnitine and a lower level of citric acid, succinic acid and creatine in the urine of rats exposed to CMS (Fig. 5B). Pathway analysis showed that these changes of metabolites were involved in fatty acid beta-oxidation and citric acid cycle in mitochondria. The dis-

turbed metabolic pathways were completely normalized by albiflorin treatment. Metabolomic analysis indicated that albiflorin might target on mitochondrial energy metabolism.

### 4. METABOLOMICS IN EVALUATING THE TOXICITY OF ANTIDEPRESSANTS

Although certain drugs can directly affect gene regulation, significant drug-induced effects are usually due to the changes of endogenous metabolites concentrations [89, 90]. Metabolite levels in the biological fluids are the readout of intra-tissue and intra-organ homeostasis [91]. The disturbances of metabolites by drugs lead to homeostasis imbalance. The consequences are recognized as drug toxicity. Metabolomics thus provides the information on the functional integrity of the whole organism over time after drug exposure. The main purpose of metabolomics is to quickly establish the toxicology and pharmacokinetics of compound candidates and reduce the drug failures that are due to toxicity in late-phase trials. The application of metabolomics for drug toxicity evaluation in the pharmaceutical industry started in the early 2000s [92, 93]. Studies before 2010 had been well summarized by several reviews written by the authors from both academics, US FDA and the leading pharmaceutical companies [92, 94, 95]. Perhaps, the greatest efforts were conducted by the Consortium for Metabonomic Toxicology (COMT), which was formed between five major pharmaceutical companies and academics from Imperial College London, UK [96]. Their work established a predictive expert system for predicting organ toxicity of drug candidates using metabolomics and nearly 150 studies were completed in 3 years [97]. The example of COMT clearly demonstrated that it is possible to use metabolomics to pro-





**Fig. (5).** Metabolomic analysis revealed that albiflorin might exert antidepressant-like activity through targeting on mitochondrial metabolism. **A:** OPLS-DA revealed the dramatic metabolites differences in urine between CMS rats with and without albiflorin treatment. **B:** Metabolites significantly altered after albiflorin treatment.

vide high throughput and inexpensive screening for large-scale drug toxicity assessment.

4.1. In vivo Toxicity Screening Models

In vivo drug toxicity evaluation using rodents (i.e., mice and rats) and non-rodents (i.e. dog, non-human primate) is used to assess organ toxicity, reproductive, developmental toxicity and genotoxicity and establish a safe starting dose for clinical studies [98]. Table 2 summarizes the published in vivo drug toxicity evaluation using metabolomics. Metabolomic profiling was showed to be more sensitive than the conventional techniques in the detection of potential drug hepatotoxicity in rats [99]. Griseofulvin is an antifungal medication used to treat a number of types of infections. Metabolomic analysis elucidated the biochemical basis of griseofulvin-induced liver injury, which was related to the significant accumulation of protoporphyrin IX (PPIX), N-methyl PPIX, bile acids, and glutathione (GSH) in the liver [100]. Using a targeted metabolomic approach, James, et al., found that glycodeoxycholic acid, taurodeoxycholic acid, and glycochenodeoxycholic acid were significantly elevated in the serum of children with acetaminophen overdose, indicating the injury of the liver [101]. Aranibar, et al., from Bristol-Myers Squibb laboratories identified ascorbic acid and gulonic acid in urine as the biomarkers of liver cytochrome P450 induction using NMR-based metabolomic profiling [102].

A metabolomics guided compound ranking system was developed to help the selection drug candidates with least toxic side effects on kidney in early drug development [103]. Urinary metabolomic profiling in combination with multivariate analysis led to the separation of post-dose time points and controls. Further analysis identified choline as a potential biomarker in urine for drug toxicity to the kidney. Doxorubicin is a chemotherapy medication widely used to treat different types of cancer such as breast cancer, bladder cancer and acute lymphocytic leukemia [104]. The systemic processes of doxorubicin-induced toxicity including onset and progression were elucidated using NGM. The increase of urinary uric acid, tryptophan, and phenylalanine and the decrease of hippuric acid and 2, 8-quinolinediol glucuronic acid were associated with the progression of doxorubicin-induced toxicity [105]. Tacrolimus is an immunosuppressant widely used to prevent rejection after the organ transplant. Its toxicity has to be carefully monitored during the treatment. Targeted metabolomics identified 3 potential human urinary biomarkers including glucose, sorbitol, and trimethylamine

oxide (TAO) which can be used for monitoring the toxicity onset of tacrolimus to the kidney [106]. Later, researchers reported the utility of urinary 1 and 3-methylhistidine as putative biomarkers of cerivastatin-induced skeletal muscle necrosis and hypertrophy in the rats [107]. An increase in urinary excretion of 1- and 3-methylhistidine was associated with the rat skeletal muscle injury caused by cerivastatin administration.

Metabolomics has also been used to evaluate the toxicity of medicinal plants used in traditional medicine. Caowu (*Radix Aconiti kusnezoffii*) is a medicinal plant that has been widely used in the treatment of rheumatic arthritis, painful joints, and tumors in China. Yan and the co-authors investigated the potential liver and heart toxicity after long-term administration of Caowu by LC-MS/MS based metabolomics [108]. The urinary metabolomics revealed the severe toxicity to heart and liver induced by Caowu. Total 13 phenotypic toxicity biomarkers were identified in urine such as palmitoyl glucuronide, D-glucuronic acid 1-phosphate, 3-indole carboxylic acid glucuronide and 3-Methyldioxyindole. Glucuronidation, catalyzed by UDP-glucuronosyltransferase (UGT) enzymes, is a major detoxification process that liver uses to remove unwanted substances from the body [109]. The decrease of glucuronidation products in rat urine after Caowu administration indicated the disturbance of normal hepatic function [108]. The toxicity of *Aconitum carmichaelii* Debx (Fu Zi in Chinese Traditional Medicine) has to be carefully monitored. Based on the metabolomic findings, a safe therapeutic range in the clinical use of Fu Zi was established using the serum betaine and phosphorylcholine as the biomarkers [110].

4.2. In vitro Toxicity Screening Models

In vitro drug toxicity screening using cells is high-throughput and inexpensive [111]. Traditionally, in vitro toxicity screenings heavily relied on the estimation of cell viability in response to the test drugs. Metabolomic profiles of cell metabolism can not only be used to discriminate between non-toxic and cytotoxic drugs, but also provide the potential molecular mechanisms of drug toxicity. A metabolomics cell-based approach was developed to evaluate drug-induced liver injury using human-derived hepatic cells like HepG2 [112]. The altered concentration of glutathione and gamma-glutamyl cycle were found to be associated with oxidative stress damage caused by hepatotoxic compounds. The changes of metabolites in fatty acids  $\beta$ -oxidation were related to drug-induced liver steatosis. To increase the throughput, a “metabolomics-on-a-chip”

**Table 2.** Representative studies on drug toxicity using metabolomics.

Drug Candidates	Target Organ Toxicity	Metabolomics Platforms	Biological Fluids	Biomarkers Increased (PubChem No.)	Biomarkers Decreased (PubChem No.)	References
Caowu	Liver and heart	UPLC-Q-TOF	Urine	151066; 439211; 68759; 440443	440650; 53481645; 161223; 5282997	[108]
Phenobarbital	Liver	NMR	Urine	54670067; 152304	N/A*	[102]
Cerivastatin	Skeletal muscle	NMR	Urine	64969; 92105	N/A	[107]
N/A	Kidney	NMR	Urine	305	N/A	[103]
Phenytoin	Liver	LC-MS/MS	Plasma	N/A	N/A	[99]
Doxorubicin	Whole body	UPLC-Q-TOF	Urine	1175; 6305; 221493	464; 76970196	[105]
Griseofulvin	Liver	UPLC-Q-TOF	Liver	4971; 124886	N/A	[100]
Acetaminophen	Liver	LC-MS/MS	Serum	3035026; 2733768	N/A	[101]
Tacrolimus	Kidney	LC-MS/MS	Urine	5793; 5780 ; 1145	N/A	[106]
Fu Zi	Heart, Liver, and Kidney	UPLC-Q-TOF	Serum	247	1014	[110]

\*N/A: Not applicable

approach was developed, where the cells were cultured on biochips and metabolic profiles in response to drug exposure were obtained by NMR [113].

Even though metabolomics has been used for drug toxicity evaluation for nearly 20 years, few reports were published on its application on antidepressants. This is partial because the whole society has not paid enough attention to depression and the pharmaceutical approaches for the treatment of depression are not really changed in the last 20 years. However, it is difficult to assess how many pharmaceutical companies are using metabolomics for the toxicity evaluation of potential antidepressant drug candidates or lead compounds since many cases might not be published due to the intellectual property concerns. Here, we provided a case study using albiflorin to demonstrate the feasibility of NGM in combination with traditional approaches in the toxicity assessments of antidepressant drug candidates during the early stages of drug development.

#### 4.3. Case Study 2: Urinary Metabolomic Analysis in Combination with Conventional Approaches Revealed no Remarkable Toxicity of Albiflorin


No severe side effects were reported for Peony in thousands of years' TCM clinical practice. Albiflorin is the bioactive molecule in the roots of white peony. Studies conducted by Beijing Wonner Biotech demonstrated the antidepressant-like activities of albiflorin, which might be developed as a new class of antidepressant [15]. The potential toxicity of albiflorin was first evaluated using the conventional approaches including in vivo acute oral toxicity (Rats and dogs), in vitro mammalian cells chromosome aberration test and bone marrow cell micronucleus test (Fig. 6A). The albiflorin dose for the conventional toxicity evaluation was 5000 mg/kg, which was 3000-fold higher than the dose for human clinical use. The results showed that albiflorin administration did not cause the significant change of Beagle dog food intake (Fig. 6B). The blood markers such as blood urea, creatinine, lipids panel,  $\text{Na}^+$ ,  $\text{K}^+$ , and  $\text{Cl}^-$  were within normal range in the albiflorin treated group. No toxic effects were observed in dog heart and kidney. Similarly, there was no significant difference in the body weight of SD rats between albiflorin treated group and controls (Fig. 6C). The chromosome

aberration test showed that albiflorin overdose did not induce any structural chromosomal abnormalities such as breaks and exchanges. The micronucleus test is the widely used in vivo genotoxicity test [114]. Albiflorin did not show any genotoxicity as there was no significant increase of micronucleated reticulocytes and micronucleated polychromatic erythrocytes.

Even though the conventional techniques did not identify any obvious toxicity caused by albiflorin overdose, the subtle detrimental toxicity can not be excluded. More sensitive NGM was then used to explore the potential toxicity of albiflorin. SD rats were administrated with 7 mg/kg of albiflorin and the urine samples were collected 12 h pre-administration and 12 h post-administration. Urine metabolites were extracted using extraction buffer containing methanol-acetonitrile-water (40:40:20, v/v/v). Briefly, 100  $\mu\text{l}$  of urine was mixed with 400  $\mu\text{l}$  of extraction buffer. The mixture was then incubated on ice for 10 min and then centrifuged for 10 min at 16000 g. The supernatant was then taken for further metabolomic analysis. LC-MS/MS-based metabolomics analysis was conducted on the urine samples as described before [88]. It was shown that urine metabolomic profile in post albiflorin treatment group was significantly distinct from that of the pre-administration (Fig. 7A). Enrichment analysis was conducted using the top 30 most differential metabolites between the two groups and the search was against disease-associated metabolites database in metaboanalyst ([www.metaboanalyst.ca](http://www.metaboanalyst.ca)). A total of 384 different diseases covering all the main organs were included in the search. No statistical significant hits were obtained, which suggested that albiflorin did not produce any toxic effects on rat metabolism (Fig. 7B).

#### 5. FUTURE DIRECTIONS

Clearly, the literature summary and cases provided in this paper have demonstrated the tremendous potential of metabolomics for drug discovery and development applications. However, as the newest addition to the "omics", metabolomics has not been widely used for new antidepressants development by the pharmaceutical industry. Indeed, only one new drug (Brexipiprazole) was approved for the treatment of depression by US FDA in the last 10 years (2008-2018) (<https://www.centerwatch.com/drug-information/fda-approved-drugs/year/2018>). More efforts are urgently needed to



**Fig. (6). The conventional *in vivo* albiflorin toxicity evaluation.** **A:** The schematic diagram of the convention toxicity tests and next-generation metabolomics based test. **B:** Acute oral toxicity test showed that the food intake of Beagle dogs was not significantly changed after albiflorin administration (5000 mg/kg). **C:** Acute oral toxicity indicated that there was no significant difference on the body weight of Sprague Dawley rats between albiflorin treated group (5000 mg/kg) and controls. Student's t test was conducted. N = 5 rats per group.


enhance the productivity and creativity of antidepressants discovery and development (R&D) by metabolomics.

### 5.1. Looking Beyond the Monoamine Hypothesis of Depression

The monoamine hypothesis has dominated antidepressants R&D in the pharmaceutical companies for over 20 years, leading to the development of current first-line drugs for the treatment of depression such as SSRIs, TCAs and MOAs [115]. Recent studies revealed that the pathogenesis of depression is far beyond the monoamine deficiency. Depression is a multi-factorial complex disorder with multiple symptoms and disturbances in the brain and other parts of the body including gut microbiome [116]. In addition, depression is often accompanied by other diseases such as osteoporosis, diabetes, cancers and cardiovascular diseases [117]. Compared to the highly selective neurotransmitter-based antidepressant drugs, multi-target strategies have been proposed to be more effective control of the core and concurrent symptoms of depression [118, 119]. Botanical medicine like albiflorin has anti-inflammatory [120], anti-obesitic [121], anti-PTSD [122] and analgesic effects [123]. The multitarget nature of albiflorin makes it a strong candidate for the treatment of patients with co-existing diseases or resistant to classical antidepressant drugs.

### 5.2. Standardization of Metabolomic Procedures

Standardization is essential for broad application of metabolomics in drug R&D, so that the results generated by one lab and on one instrument can be validated by different labs and on different instruments. As we described in Table 2, NGM may be conducted on different platforms using the instruments from different vendors. Moreover, the workstations and data formats are also different between vendors. Therefore, harmonization of metabolomic procedures is extremely challenging. The Wishart group from the University of Alberta provided some recommendations on NMR-based metabolomic studies using urine samples including urine collection, sample preparation and data acquisition [124]. Recently, a white paper published by "Precision Medicine and Pharmacometabolomics Task Group" on Metabolomics journal offered recommendations on the selection of metabolomic platforms, samples collection, preservation and standardization of measurements for precision medicine [125]. Standardization is still needed for each metabolomic platform, each biological fluid, data analysis, and deposition.



**Fig. (7).** The evaluation of the potential toxicity of albiflorin using next-generation metabolomics. **A:** PLS-DA showed the separation of urine metabolic profile between pre and post albiflorin treatment group. **B:** Enrichment analysis did not yield any significant hits indicating no potential albiflorin induced toxicity. The enrichment analysis was performed using Metaboanalyst 3.0 ([www.metaboanalyst.ca](http://www.metaboanalyst.ca)).

### 5.3. Understanding the Biology Behind Metabolomic Big Data

Like other omic techniques, metabolomics usually generates a large set of raw data. How to obtain meaningful biological information behind the data is the key. Nowadays, for a typical metabolomic study, more than half of the total project time is devoted to the data processing and bioinformatics analysis. A comprehensive list of free metabolomic analysis tools and their usage have been summarized in several recent reviews [126, 127]. However, metabolomics is a young discipline. The tools for the functional interpretation of metabolomic experiments are still at least 10 years behind genomics, transcriptomics, and proteomics. This bottleneck limits the use of metabolomics in the pharmaceutical industry [128]. Moreover, many current metabolomic tools are run under R, Python or Java and require the users to have computer programming skills. More user-friendly tools are needed, which could provide the ability to walk through the whole data analysis process from data normalization, statistical analysis to biological pathways. In addition, in the context of omics era, tools for multi-scale and multi-omics integration will be very useful to link the phenotypes with genome, transcriptome, proteome, metabolome and gut microbiome.

### CONCLUSION

Globally, investment in new drug research and discovery (R&D) falls sharply, especially for new antidepressants due to the dramatical increase of R&D expenditure and the decline of the world's economy [20, 129]. Metabolomics is entering into a "new generation" towards more sensitive, broader coverage, cheaper and more quantitative methods. NGM might be the "deal breaker" for pharmaceutical R&D.

In this review, we discussed the next-generation metabolomics from its basic concept, set up its application in antidepressant discovery and development. We also provided 2 cases to demonstrate its utility. Perhaps, with the efforts from academics, healthcare providers, pharmaceutical industries, and governments, NGM will eventually become an indispensable tool in the discovery and development of new antidepressants and really facilitate this time-consuming and pricy cycle.

### CONSENT FOR PUBLICATION

Not applicable.

### CONFLICT OF INTEREST

Z.Z. is holding US, Europe and China patents for the use of albiflorin in the treatment of depression (China patent no. ZL2009102654220, ZL201180051467.3, ZL201080069710.X; US patent no. US9023817B2, US9453041B2, US20130231469A1, US20130316966 A1, US9555055B2 and Europe patent no. EP2491934B1). Other authors declare no competing financial interests.

### ACKNOWLEDGEMENTS

This work was partially supported by grants from the National Natural Science Foundation of China (No. 81628008), the Natural Science Foundation of Tianjin City (No. 16JCQNJC14400), Foundation of Science and Technology for overseas returnees of Tianjin and Tianjin "131" innovative talents project. The authors thank Academy of Military Medical Sciences, China and Tianjin University of Traditional Chinese Medicine for conducting some of the animal studies described in the paper. Shapes for some of the figures were from SlidesCarnival (<https://www.slidescarnival.com/>).

### REFERENCES

- [1] Wishart DS. Emerging applications of metabolomics in drug discovery and precision medicine. *Nat Rev Drug Discov* 2016; 15(7): 473-84.
- [2] Lin L, Cao B, Xu Z, *et al.* *In vivo* HMRS and lipidomic profiling reveals comprehensive changes of hippocampal metabolism during aging in mice. *Biochem Biophys Res Commun* 2016; 470(1): 9-14.
- [3] Li K, Wang X, Pidatala VR, Chang CP, Cao X. Novel quantitative metabolomic approach for the study of stress responses of plant root metabolism. *J Proteome Res* 2014; 13(12): 5879-87.
- [4] Yu L, Li K, Zhang X. Next-generation metabolomics in lung cancer diagnosis, treatment and precision medicine: mini review. *Oncotarget* 2017; 8(70): 115774-86.
- [5] Cui S, Li K, Ang L, *et al.* Plasma phospholipids and sphingolipids identify stent restenosis after percutaneous coronary intervention. *JACC Cardiovasc Interv* 2017; 10(13): 1307-16.



- [6] Li K, Pidatala RR, Ramakrishna W. Mutational, proteomic and metabolomic analysis of a plant growth promoting copper-resistant *Pseudomonas* spp. *FEMS Microbiol Lett* 2012; 335(2): 140-8.
- [7] Li K, Pidatala VR, Shaik R, Datta R, Ramakrishna W. Integrated metabolomic and proteomic approaches dissect the effect of metal-resistant bacteria on maize biomass and copper uptake. *Environ Sci Technol* 2014; 48(2): 1184-93.
- [8] Friedrich MJ. Depression is the leading cause of disability around the world. *JAMA* 2017; 317(15): 1517.
- [9] Otte C, Gold SM, Penninx BW, et al. Major depressive disorder. *Nat Rev Dis Primers* 2016; 2: 16065.
- [10] Pereira VS, Hiroaki-Sato VA. A brief history of antidepressant drug development: from tricyclics to beyond ketamine. *Acta Neuropsychiatr* 2018; 1-16.
- [11] Harmer CJ, Duman RS, Cowen PJ. How do antidepressants work? New perspectives for refining future treatment approaches. *Lancet Psychiatry* 2017; 4(5): 409-18.
- [12] Duman RS, Aghajanian GK, Sanacora G, Krystal JH. Synaptic plasticity and depression: new insights from stress and rapid-acting antidepressants. *Nat Med* 2016; 22(3): 238-49.
- [13] Sarris J. Herbal medicines in the treatment of psychiatric disorders: 10-year updated review. *Phytother Res* 2018; 32(7): 1147-62.
- [14] Mao QQ, Ip SP, Xian YF, Hu Z, Che CT. Anti-depressant-like effect of peony: A mini-review. *Pharm Biol* 2012; 50(1): 72-7.
- [15] Wang YL, Wang JX, Hu XX, et al. Antidepressant-like effects of albiflorin extracted from *Radix paeoniae* Alba. *J Ethnopharmacol* 2016; 179: 9-15.
- [16] Jin ZL, Gao N, Xu W, et al. Receptor and transporter binding and activity profiles of albiflorin extracted from *Radix paeoniae* Alba. *Sci Rep* 2016; 6: 33793.
- [17] Song J, Hou X, Hu X, et al. Not only serotonergic system, but also dopaminergic system involved in albiflorin against chronic unpredictable mild stress-induced depression-like behavior in rats. *Chem Biol Interact* 2015; 242: 211-7.
- [18] Millan MJ. Multi-target strategies for the improved treatment of depressive states: Conceptual foundations and neuronal substrates, drug discovery and therapeutic application. *Pharmacol Ther* 2006; 110(2): 135-370.
- [19] Rautio J, Meanwell NA, Di L, Hageman MJ. The expanding role of prodrugs in contemporary drug design and development. *Nat Rev Drug Discov* 2018; 19(8): 559-87.
- [20] Cuperlovic-Culp M, Culp AS. Applied metabolomics in drug discovery. *Expert Opin Drug Discov* 2016; 11(8): 759-70.
- [21] Nassar AF, Wu T, Nassar SF, Wisniewski AV. UPLC-MS for metabolomics: A giant step forward in support of pharmaceutical research. *Drug Discov Today* 2017; 22(2): 463-70.
- [22] Martínez-Morillo E, Prieto García B, Alvarez Menéndez FV. Challenges for worldwide harmonization of newborn screening programs. *Clin Chem* 2016; 62(5): 689-98.
- [23] Oliver SG, Winson MK, Kell DB, Baganz F. Systematic functional analysis of the yeast genome. *Trends Biotechnol* 1998; 16(9): 373-8.
- [24] Nicholson JK, Lindon JC, Holmes E. 'Metabonomics': understanding the metabolic responses of living systems to pathophysiological stimuli via multivariate statistical analysis of biological NMR spectroscopic data. *Xenobiotica* 1999; 29(11): 1181-9.
- [25] Bingol K. Recent advances in targeted and untargeted metabolomics by NMR and MS/NMR methods. *High Throughput* 2018; 7(2): E9.
- [26] Koek MM, Jellema RH, van der Greef J, Tas AC, Hankemeier T. Quantitative metabolomics based on gas chromatography mass spectrometry: status and perspectives. *Metabolomics* 2011; 7(3): 307-28.
- [27] Fernie AR, Aharoni A, Willmitzer L, et al. Recommendations for reporting metabolite data. *Plant Cell* 2011; 23(7): 2477-82.
- [28] Wishart DS. Quantitative metabolomics using NMR. *Trends Anal Chem* 2008; 27: 228-37.
- [29] Zhang A, Sun H, Wang P, Han Y, Wang X. Modern analytical techniques in metabolomics analysis. *Analyst (Lond)* 2012; 137(2): 293-300.
- [30] Coene KLM, Kluijtmans LAJ, van der Heeft E, et al. Next-generation metabolic screening: targeted and untargeted metabolomics for the diagnosis of inborn errors of metabolism in individual patients. *J Inher Metab Dis* 2018; 41(3): 337-53.
- [31] Zhang W, Hankemeier T, Ramautar R. Next-generation capillary electrophoresis-mass spectrometry approaches in metabolomics. *Curr Opin Biotechnol* 2017; 43: 1-7.
- [32] Warth B, Raffener P, Granados A, et al. Metabolomics reveals that dietary xenoestrogens alter cellular metabolism induced by palbociclib/letrozole combination cancer therapy. *Cell Chem Biol* 2018; 25(3): 291-300.e3.
- [33] Johnson CH, Santidrian AF, LeBoeuf SE, et al. Metabolomics guided pathway analysis reveals link between cancer metastasis, cholesterol sulfate, and phospholipids. *Cancer Metab* 2017; 5: 9.
- [34] Zhang B, Zhang H, Du C, et al. Metabolic responses of the growing *Daphnia similis* to chronic AgNPs exposure as revealed by GC-Q-TOF/MS and LC-Q-TOF/MS. *Water Res* 2017; 114: 135-43.
- [35] Wang X, Li K, Adams E, Van Schepdael A. Capillary electrophoresis-mass spectrometry in metabolomics: the potential for driving drug discovery and development. *Curr Drug Metab* 2013; 14(7): 807-13.
- [36] Zenobi R. Single-cell metabolomics: Analytical and biological perspectives. *Science* 2013; 342(6163): 1243259.
- [37] He J, Luo Z, Huang L, et al. Ambient mass spectrometry imaging metabolomics method provides novel insights into the action mechanism of drug candidates. *Anal Chem* 2015; 87(10): 5372-9.
- [38] Masson P, Alves AC, Ebbels TM, Nicholson JK, Want EJ. Optimization and evaluation of metabolite extraction protocols for untargeted metabolic profiling of liver samples by UPLC-MS. *Anal Chem* 2010; 82(18): 7779-86.
- [39] Kuehnbaum NL, Britz-McKibbin P. New advances in separation science for metabolomics: resolving chemical diversity in a post-genomic era. *Chem Rev* 2013; 113(4): 2437-68.
- [40] Ralston-Hooper K, Jannasch A, Adamec J, Sepúlveda M. The use of two-dimensional gas chromatography-time-of-flight mass spectrometry (GC×GC-TOF-MS) for metabolomic analysis of polar metabolites. *Methods Mol Biol* 2011; 708: 205-11.
- [41] Wood PL. Mass spectrometry strategies for clinical metabolomics and lipidomics in psychiatry, neurology, and neuro-oncology. *Neuropsychopharmacology* 2014; 39(1): 24-33.
- [42] Xiao JF, Zhou B, Ransom HW. Metabolite identification and quantitation in LC-MS/MS-based metabolomics. *Trends Anal Chem* 2012; 32: 1-14.
- [43] Zhang W, Hankemeier T, Ramautar R. Next-generation capillary electrophoresis-mass spectrometry approaches in metabolomics. *Curr Opin Biotechnol* 2017; 43: 1-7.
- [44] Dettmer K, Aronov PA, Hammock BD. Mass spectrometry-based metabolomics. *Mass Spectrom Rev* 2007; 26(1): 51-78.
- [45] Snyder DT, Pulliam CJ, Ouyang Z, Cooks RG. Miniature and fieldable mass spectrometers: recent advances. *Anal Chem* 2016; 88(1): 2-29.
- [46] Ji J, Sun J, Pi F, et al. GC-TOF/MS-based metabolomics approach to study the cellular immunotoxicity of deoxynivalenol on murine macrophage ANA-1 cells. *Chem Biol Interact* 2016; 256: 94-101.
- [47] Chang KL, Ho PC. Gas chromatography time-of-flight mass spectrometry (GC-TOF-MS)-based metabolomics for comparison of caffeinated and decaffeinated coffee and its implications for Alzheimer's disease. *PLoS One* 2014; 9(8): e104621.
- [48] Fiehn O. Metabolomics by gas chromatography-mass spectrometry: combined targeted and untargeted profiling. *Curr Protoc Mol Biol* 2016; 114: 30.4.1-32.
- [49] Weidt S, Haggarty J, Kean R, et al. A novel targeted/untargeted GC-Orbitrap metabolomics methodology applied to *Candida albicans* and *Staphylococcus aureus* biofilms. *Metabolomics* 2016; 12(12): 189.
- [50] Woo HM, Noack S, Seibold GM, Willbold S, Eikmanns BJ, Bott M. Link between phosphate starvation and glycogen metabolism in *Corynebacterium glutamicum*, revealed by metabolomics. *Appl Environ Microbiol* 2010; 76(20): 6910-9.
- [51] Virgiliou C, Gika HG, Theodoridis GA. HILIC-MS/MS multi-targeted method for metabolomics applications. *Methods Mol Biol* 2018; 1738: 65-81.
- [52] Tang DQ, Zou L, Yin XX, Ong CN. HILIC-MS for metabolomics: An attractive and complementary approach to RPLC-MS. *Mass Spectrom Rev* 2016; 35(5): 574-600.
- [53] Fu J, Shen M, Shen Y, et al. LC-MS/MS-based metabolome analysis of biochemical pathways altered by food limitation in larvae of ivory shell, *babylonia areolata*. *Mar Biotechnol (NY)* 2018; 20(4): 451-66.

- [54] Ciborowski M, Teul J, Martin-Ventura JL, Egido J, Barbas C. Metabolomics with LC-QTOF-MS permits the prediction of disease stage in aortic abdominal aneurysm based on plasma metabolic fingerprint. *PLoS One* 2012; 7(2): e31982.
- [55] Zou W, Wen X, Xie C, *et al.* LC-Q-TOF-MS based plasma metabolomic profile of subclinical pelvic inflammatory disease: A pilot study. *Clin Chim Acta* 2018; 483: 164-9.
- [56] Saigusa D, Okamura Y, Motoike IN, *et al.* Establishment of protocols for global metabolomics by lc-ms for biomarker discovery. *PLoS One* 2016; 11(8): e0160555.
- [57] Liu JX, Aerts JT, Rubakhin SS, Zhang XX, Sweedler JV. Analysis of endogenous nucleotides by single cell capillary electrophoresis-mass spectrometry. *Analyst (Lond)* 2014; 139(22): 5835-42.
- [58] Ramautar R. Sheathless capillary electrophoresis-mass spectrometry for the profiling of charged metabolites in biological samples. *Methods Mol Biol* 2018; 1738: 183-92.
- [59] Tokunaga M, Kami K, Ozawa S, *et al.* Metabolome analysis of esophageal cancer tissues using capillary electrophoresis-time-of-flight mass spectrometry. *Int J Oncol* 2018; 52(6): 1947-58.
- [60] Dai C, Arceo J, Arnold J, *et al.* Metabolomics of oncogene-specific metabolic reprogramming during breast cancer. *Cancer Metab* 2018; 6: 5.
- [61] Boizard F, Brunchault V, Moulos P, *et al.* A capillary electrophoresis coupled to mass spectrometry pipeline for long term comparable assessment of the urinary metabolome. *Sci Rep* 2016; 6: 34453.
- [62] MacLennan MS, Kok MGM, Soliman L, So A, Hurtado-Coll A, Chen DDY. Capillary electrophoresis-mass spectrometry for targeted and untargeted analysis of the sub-5 kDa urine metabolome of patients with prostate or bladder cancer: A feasibility study. *J Chromatogr B Analyt Technol Biomed Life Sci* 2018; 1074-1075: 79-85.
- [63] Fessenden M. Metabolomics: Small molecules, single cells. *Nature* 2016; 540(7631): 153-5.
- [64] Qi M, Philip MC, Yang N, Sweedler JV. Single Cell Neurometabolomics. *ACS Chem Neurosci* 2018; 9(1): 40-50.
- [65] Li Q, Chen P, Fan Y, *et al.* Multicolor fluorescence detection-based microfluidic device for single-cell metabolomics: simultaneous quantitation of multiple small molecules in primary liver cells. *Anal Chem* 2016; 88(17): 8610-6.
- [66] Emara S, Amer S, Ali A, Abouleila Y, Oga A, Masujima T. Single-Cell Metabolomics. *Adv Exp Med Biol* 2017; 965: 323-43.
- [67] Sarabia LD, Boughton BA, Rupasinghe T, *et al.* High-mass-resolution MALDI mass spectrometry imaging reveals detailed spatial distribution of metabolites and lipids in roots of barley seedlings in response to salinity stress. *Metabolomics* 2018; 14(5): 63.
- [68] Smith A, Galli M, L'Imperio V, Pagni F, Magni F. MALDI-MS imaging in the study of glomerulonephritis. *Methods Mol Biol* 2017; 1618: 85-94.
- [69] Tian H, Fletcher JS, Thuret R, *et al.* Spatiotemporal lipid profiling during early embryonic development of *Xenopus laevis* using dynamic ToF-SIMS imaging. *J Lipid Res* 2014; 55(9): 1970-80.
- [70] Clendinen CS, Monge ME, Fernández FM. Ambient mass spectrometry in metabolomics. *Analyst (Lond)* 2017; 142(17): 3101-17.
- [71] Jarmusch AK, Pirro V, Baird Z, Hattab EM, Cohen-Gadol AA, Cooks RG. Lipid and metabolite profiles of human brain tumors by desorption electrospray ionization-MS. *Proc Natl Acad Sci USA* 2016; 113(6): 1486-91.
- [72] Song X, Luo Z, Li X, *et al.* In situ hydrogel conditioning of tissue samples to enhance the drug's sensitivity in ambient mass spectrometry imaging. *Anal Chem* 2017; 89(12): 6318-23.
- [73] Kelloff GJ, Sigman CC. Cancer biomarkers: selecting the right drug for the right patient. *Nat Rev Drug Discov* 2012; 11(3): 201-14.
- [74] Zhou X, Liu L, Lan X, *et al.* Polyunsaturated fatty acids metabolism, purine metabolism and inosine as potential independent diagnostic biomarkers for major depressive disorder in children and adolescents. *Mol Psychiatry* 2018; 1-11.
- [75] Mocking RJT, Assies J, Ruhé HG, Schene AH. Focus on fatty acids in the neurometabolic pathophysiology of psychiatric disorders. *J Inherit Metab Dis* 2018; 41(4): 597-611.
- [76] Liu L, Zhou X, Zhang Y, *et al.* Hippocampal metabolic differences implicate distinctions between physical and psychological stress in four rat models of depression. *Transl Psychiatry* 2018; 8(1): 4.
- [77] Chong J, Soufan O, Li C, *et al.* MetaboAnalyst 4.0: towards more transparent and integrative metabolomics analysis. *Nucleic Acids Res* 2018; 46(W1): W486-94.
- [78] Guijas C, Montenegro-Burke JR, Domingo-Almenara X, *et al.* METLIN: A technology platform for identifying knowns and unknowns. *Anal Chem* 2018; 90(5): 3156-64.
- [79] Lian B, Xia J, Yang X, *et al.* Mechanisms of ketamine on mice hippocampi shown by gas chromatography-mass spectrometry-based metabolomic analysis. *Neuroreport* 2018; 29(9): 704-11.
- [80] Zanos P, Gould TD. Mechanisms of ketamine action as an antidepressant. *Mol Psychiatry* 2018; 23(4): 801-11.
- [81] Hu Q, Shen P, Bai S, *et al.* Metabolite-related antidepressant action of diterpene ginkgolides in the prefrontal cortex. *Neuropsychiatr Dis Treat* 2018; 14: 999-1011.
- [82] Bai S, Zhang X, Chen Z, *et al.* Insight into the metabolic mechanism of Diterpene Ginkgolides on antidepressant effects for attenuating behavioural deficits compared with venlafaxine. *Sci Rep* 2017; 7(1): 9591.
- [83] Ma WN, Zhou MM, Gou XJ, *et al.* Urinary metabolomic study of chlorogenic acid in a rat model of chronic sleep deprivation using gas chromatography-mass spectrometry. *Int J Genomics* 2018; 2018: 1361402.
- [84] Du H, Zhang H, Zhao Y, *et al.* Metabolic analysis of the antidepressive effects of Yangxinshi Tablet in a vascular depression model in mice. *Biomed Chromatogr* 2018; 32(3): 32.
- [85] Jia HM, Yu M, Ma LY, Zhang HW, Zou ZM. Chaihu-Shu-Gan-San regulates phospholipids and bile acid metabolism against hepatic injury induced by chronic unpredictable stress in rat. *J Chromatogr B Analyt Technol Biomed Life Sci* 2017; 1064: 14-21.
- [86] Shen P, Hu Q, Dong M, *et al.* Venlafaxine exerts antidepressant effects possibly by activating MAPK-ERK1/2 and PI3K-AKT pathways in the hippocampus. *Behav Brain Res* 2017; 335: 63-70.
- [87] Martin AE, Schober DA, Nikolayev A, *et al.* Further evaluation of mechanisms associated with the antidepressantlike signature of scopolamine in mice. *CNS Neurol Disord Drug Targets* 2017; 16(4): 492-500.
- [88] Yuan M, Breitkopf SB, Yang X, Asara JM. A positive/negative ion-switching, targeted mass spectrometry-based metabolomics platform for bodily fluids, cells, and fresh and fixed tissue. *Nat Protoc* 2012; 7(5): 872-81.
- [89] Arbuckle JH, Gardina PJ, Gordon DN, *et al.* Inhibitors of the histone methyltransferases EZH2/1 induce a potent antiviral state and suppress infection by diverse viral pathogens. *MBio* 2017; 8(4): e01141-17.
- [90] Nicholson JK, Connelly J, Lindon JC, Holmes E. Metabonomics: A platform for studying drug toxicity and gene function. *Nat Rev Drug Discov* 2002; 1(2): 153-61.
- [91] Ivanisevic J, Elias D, Deguchi H, *et al.* Arteriovenous Blood Metabolomics: A readout of intra-tissue metabostasis. *Sci Rep* 2015; 5: 12757.
- [92] Griffin JL. The potential of metabonomics in drug safety and toxicology. *Drug Discov Today Technol* 2004; 1(3): 285-93.
- [93] Nicholson JK, Wilson ID. Opinion: understanding 'global' systems biology: metabonomics and the continuum of metabolism. *Nat Rev Drug Discov* 2003; 2(8): 668-76.
- [94] Waters NJ. The role of metabonomics at the interface between drug metabolism and safety assessment. *Curr Drug Metab* 2010; 11(8): 686-92.
- [95] Bai JP, Abernethy DR. Systems pharmacology to predict drug toxicity: integration across levels of biological organization. *Annu Rev Pharmacol Toxicol* 2013; 53: 451-73.
- [96] Lindon JC, Keun HC, Ebbels TM, Pearce JM, Holmes E, Nicholson JK. The Consortium for Metabonomic Toxicology (COMET): Aims, activities and achievements. *Pharmacogenomics* 2005; 6(7): 691-9.
- [97] Lindon JC, Nicholson JK, Holmes E, *et al.* Contemporary issues in toxicology the role of metabonomics in toxicology and its evaluation by the COMET project. *Toxicol Appl Pharmacol* 2003; 187(3): 137-46.
- [98] Anadón A, Martínez MA, Castellano V, Martínez-Larrañaga MR. The role of in vitro methods as alternatives to animals in toxicity testing. *Expert Opin Drug Metab Toxicol* 2014; 10(1): 67-79.
- [99] Mattes W, Davis K, Fabian E, *et al.* Detection of hepatotoxicity potential with metabolite profiling (metabolomics) of rat plasma. *Toxicol Lett* 2014; 230(3): 467-78.
- [100] Liu K, Yan J, Sachar M, *et al.* A metabolomic perspective of griseofulvin-induced liver injury in mice. *Biochem Pharmacol* 2015; 98(3): 493-501.

- [101] James L, Yan K, Pence L, *et al.* Comparison of bile acids and acetaminophen protein adducts in children and adolescents with acetaminophen toxicity. *PLoS One* 2015; 10(7): e0131010.
- [102] Aranibar N, Bhaskaran V, Ott KH, *et al.* Modulation of ascorbic acid metabolism by cytochrome P450 induction revealed by metabolomics and transcriptional profiling. *Magn Reson Chem* 2009; 47(Suppl. 1): S12-9.
- [103] Dieterle F, Schlotterbeck G, Ross A, Niederhauser U, Senn H. Application of metabolomics in a compound ranking study in early drug development revealing drug-induced excretion of choline into urine. *Chem Res Toxicol* 2006; 19(9): 1175-81.
- [104] Gorini S, De Angelis A, Berrino L, Malara N, Rosano G, Ferraro E. Chemotherapeutic drugs and mitochondrial dysfunction: focus on doxorubicin, trastuzumab, and sunitinib. *Oxid Med Cell Longev* 2018; 2018: 7582730.
- [105] Wang J, Reijmers T, Chen L, *et al.* Systems toxicology study of doxorubicin on rats using ultra performance liquid chromatography coupled with mass spectrometry based metabolomics. *Metabolomics* 2009; 5(4): 407-18.
- [106] Klepacki J, Klawitter J, Klawitter J, Thurman JM, Christians U. A high-performance liquid chromatography-tandem mass spectrometry-based targeted metabolomics kidney dysfunction marker panel in human urine. *Clin Chim Acta* 2015; 446: 43-53.
- [107] Aranibar N, Vassallo JD, Rathmacher J, *et al.* Identification of 1- and 3-methylhistidine as biomarkers of skeletal muscle toxicity by nuclear magnetic resonance-based metabolic profiling. *Anal Biochem* 2011; 410(1): 84-91.
- [108] Yan Y, Zhang A, Dong H, *et al.* Toxicity and detoxification effects of herbal caowu via ultra performance liquid chromatography/mass spectrometry metabolomics analyzed using pattern recognition method. *Pharmacogn Mag* 2017; 13(52): 683-92.
- [109] Yang G, Ge S, Singh R, *et al.* Glucuronidation: driving factors and their impact on glucuronide disposition. *Drug Metab Rev* 2017; 49(2): 105-38.
- [110] Tan Y, Ko J, Liu X, *et al.* Serum metabolomics reveals betaine and phosphatidylcholine as potential biomarkers for the toxic responses of processed *Aconitum carmichaelii* Debx. *Mol Biosyst* 2014; 10(9): 2305-16.
- [111] Shukla SJ, Huang R, Austin CP, Xia M. The future of toxicity testing: A focus on *in vitro* methods using a quantitative high-throughput screening platform. *Drug Discov Today* 2010; 15(23-24): 997-1007.
- [112] García-Cañaveras JC, Castell JV, Donato MT, Lahoz A. A metabolomics cell-based approach for anticipating and investigating drug-induced liver injury. *Sci Rep* 2016; 6: 27239.
- [113] Choucha Snouber L, Bunesco A, Naudot M, *et al.* Metabolomics-on-a-chip of hepatotoxicity induced by anticancer drug flutamide and its active metabolite hydroxyflutamide using HepG2/C3a microfluidic biochips. *Toxicol Sci* 2013; 132(1): 8-20.
- [114] Hayashi M. The micronucleus test-most widely used *in vivo* genotoxicity test. *Genes Environ* 2016; 38: 18.
- [115] Boku S, Nakagawa S, Toda H, Hishimoto A. Neural basis of major depressive disorder: Beyond monoamine hypothesis. *Psychiatry Clin Neurosci* 2018; 72(1): 3-12.
- [116] Vuong HE, Yano JM, Fung TC, Hsiao EY. The microbiome and host behavior. *Annu Rev Neurosci* 2017; 40: 21-49.
- [117] Cizza G, Primma S, Csako G. Depression as a risk factor for osteoporosis. *Trends Endocrinol Metab* 2009; 20(8): 367-73.
- [118] Millan MJ. On 'polypharmacy' and multi-target agents, complementary strategies for improving the treatment of depression: A comparative appraisal. *Int J Neuropsychopharmacol* 2014; 17(7): 1009-37.
- [119] Tang SW, Tang WH, Leonard BE. Multitarget botanical pharmacotherapy in major depression: A toxic brain hypothesis. *Int Clin Psychopharmacol* 2017; 32(6): 299-308.
- [120] Bi X, Han L, Qu T, *et al.* Anti-Inflammatory effects, SAR, and action mechanism of monoterpenoids from *Radix paeoniae* Alba on LPS-stimulated RAW 264.7 cells. *Molecules* 2017; 22(5): E715.
- [121] Jeong MY, Park J, Youn DH, *et al.* Albiflorin ameliorates obesity by inducing thermogenic genes via AMPK and PI3K/AKT *in vivo* and *in vitro*. *Metabolism* 2017; 73: 85-99.
- [122] Qiu ZK, He JL, Liu X, Zeng J, Chen JS, Nie H. Anti-PTSD-like effects of albiflorin extracted from *Radix paeoniae* Alba. *J Ethnopharmacol* 2017; 198: 324-30.
- [123] Zhou J, Wang L, Wang J, *et al.* Paeoniflorin and albiflorin attenuate neuropathic pain via MAPK pathway in chronic constriction injury rats. *Evid Based Complement Alternat Med* 2016; 2016: 8082753.
- [124] Emwas AH, Luchinat C, Turano P, *et al.* Standardizing the experimental conditions for using urine in NMR-based metabolomic studies with a particular focus on diagnostic studies: A review. *Metabolomics* 2015; 11(4): 872-94.
- [125] Beger RD, Dunn W, Schmidt MA, *et al.* Metabolomics enables precision medicine: "A White Paper, Community Perspective". *Metabolomics* 2016; 12(10): 149.
- [126] Spicer R, Salek RM, Moreno P, Cañueto D, Steinbeck C. Navigating freely-available software tools for metabolomics analysis. *Metabolomics* 2017; 13(9): 106.
- [127] Misra BB, Fahrman JF, Grapov D. Review of emerging metabolomic tools and resources: 2015-2016. *Electrophoresis* 2017; 38(18): 2257-74.
- [128] Robertson DG, Watkins PB, Reilly MD. Metabolomics in toxicology: preclinical and clinical applications. *Toxicol Sci* 2011; 120(Suppl. 1): S146-70.
- [129] Schuhmacher A, Gassmann O, Hinder M, Chang R. Changing R&D models in research-based pharmaceutical companies. *J Transl Med* 2016; 14(1): 105.