

Computation of the Ground State Energy and Wave Function of the Hydrogen Molecule using the Variational Monte Carlo Method

JAAP WESDORP[†], BAS DIRKSE[†]

[†]Delft University of Technology

j.j.wesdorp@student.tudelft.nl

b.dirkse@student.tudelft.nl

March 28, 2016

Abstract

Some abstract

1. INTRODUCTION

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Ut purus elit, vestibulum ut, placerat ac, adipiscing vitae, felis. Curabitur dictum gravida mauris. Nam arcu libero, nonummy eget, consectetur id, vulputate a, magna. Donec vehicula augue eu neque. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Mauris ut leo. Cras viverra metus rhoncus sem. Nulla et lectus vestibulum urna fringilla ultrices. Phasellus eu tellus sit amet tortor gravida placerat. Integer sapien est, iaculis in, pretium quis, viverra ac, nunc. Praesent eget sem vel leo ultrices bibendum. Aenean faucibus. Morbi dolor nulla, malesuada eu, pulvinar at, mollis ac, nulla. Curabitur auctor semper nulla. Donec varius orci eget risus. Duis nibh mi, congue eu, accumsan eleifend, sagittis quis, diam. Duis eget orci sit amet orci dignissim rutrum.

2. METHODS

In the variational method for finding the ground state of a system, we start with a trial wave function of a specific form, which depends on (several) variational parameters. We then compute the expectation value of the energy, given the Hamiltonian of the system and optimize the parameters to find the ground state within our variational space. The computation of the expectation value of the energy is done using the Metropolis Monte Carlo method, thereby avoiding the need to normalize the trial wave function. The optimization of the variational parameters is done by a damped steepest descent method, where the derivative of the energy with respect to each variational parameter is computed using some statistical averaging algorithm, thereby avoiding the use of finite differences of stochastic variables.

Natural Units of Computation

In our computation we naturally use atomic units. For the distance, we use the unit of Bohr radius

$$a_0 = \frac{4\pi\epsilon_0\hbar^2}{me^2} \approx 0.529 \times 10^{-10} \text{ m}, \quad (1)$$

and for energy we use twice the ionization energy of a Hydrogen atom

$$E_0 = \frac{m}{\hbar^2} \left(\frac{e^2}{4\pi\epsilon_0} \right)^2 \approx 27.2 \text{ eV} \quad (2)$$

The Hamiltonian of the Hydrogen Molecule

In this paper we approximate the Hydrogen molecule system as two stationary protons separated by a distance s , supplemented with two electrons. We choose our coordinate system such that both protons are symmetrically located around the origin and positioned on the x-axis, i.e. the protons are at $[\pm s/2, 0, 0]^T$. The Hamiltonian of this system includes the kinetic energy of the electrons, the potential energy of each electron due to the two protons and an electron-electron interaction term. Let the positions of the electrons be given by \mathbf{r}_1 and \mathbf{r}_2 respectively. In our natural units, the Hamiltonian is then given by

$$H = -\frac{1}{2}(\nabla_1^2 + \nabla_2^2) - \left[\frac{1}{r_{1L}} + \frac{1}{r_{1R}} + \frac{1}{r_{2L}} + \frac{1}{r_{2R}} \right] + \frac{1}{r_{12}}, \quad (3)$$

where

$$\begin{aligned} \mathbf{r}_{iL} &= \mathbf{r}_i + \frac{s}{2} \hat{\mathbf{x}}; & \mathbf{r}_{iR} &= \mathbf{r}_i - \frac{s}{2} \hat{\mathbf{x}}; & i &= 1, 2; \text{ and} \\ \mathbf{r}_{12} &= \mathbf{r}_1 - \mathbf{r}_2. \end{aligned} \quad (4)$$

The Variational Wave Function

The variational form of the wave functions is chosen to be of the following form

$$\Psi(\mathbf{r}_1, \mathbf{r}_2) = \phi_1(\mathbf{r}_1) \phi_2(\mathbf{r}_2) \psi(\mathbf{r}_1, \mathbf{r}_2), \quad (5)$$

where

$$\phi_i(\mathbf{r}_i) = e^{-r_{iL}/a} + e^{-r_{iR}/a} = \phi_{iL}(\mathbf{r}_i) + \phi_{iR}(\mathbf{r}_i), \quad i = 1, 2. \quad (6)$$

Note that these parts are just the sum of the wave functions of each electron being in the ground state of the left or right hydrogen atom. The interaction part of the wave function is chosen to be the Jastrow function

$$\psi(\mathbf{r}_1, \mathbf{r}_2) = \exp \left[\frac{r_{12}}{\alpha(1 + \beta r_{12})} \right]. \quad (7)$$

Note that we now have four variational parameters a , s , α and β . Later we will put constraints on two of these parameters to ensure that the expectation value of the energy remains bounded.

The Local Energy

The expectation value of the energy is computed using the local energy and a weight function. The weight function is then incorporated using the Metropolis Monte Carlo method, so that the expectation value of the energy is the integral of the local energy using points sampled from the weight function distribution. We thus compute the energy as

$$\langle E \rangle = \frac{\langle \Psi | H | \Psi \rangle}{\langle \Psi | \Psi \rangle} = \iint \frac{\Psi^* H \Psi}{\langle \Psi | \Psi \rangle} d^3 \mathbf{r}_1 d^3 \mathbf{r}_2 = \iint \frac{\Psi \Psi^*}{\langle \Psi | \Psi \rangle} \frac{H \Psi}{\Psi} d^3 \mathbf{r}_1 d^3 \mathbf{r}_2 = \iint \omega \varepsilon d^3 \mathbf{r}_1 d^3 \mathbf{r}_2, \quad (8)$$

where

$$\omega(\mathbf{r}_1, \mathbf{r}_2) = \frac{|\Psi|^2}{\langle \Psi | \Psi \rangle}, \quad (9)$$

is the weight function and

$$\varepsilon(\mathbf{r}_1, \mathbf{r}_2) = \frac{H \Psi}{\Psi} \quad (10)$$

is the local energy. Note that both expressions depend on the positions \mathbf{r}_1 and \mathbf{r}_2 , and also on the variational parameters. Furthermore, note that ω is a normalized probability distribution, which is very useful for the integration of (8).

Plugging in the variational wave function and the Hydrogen molecule Hamiltonian into (10) we find an expression for the local energy in our problem

$$\begin{aligned} \epsilon(\mathbf{r}_1, \mathbf{r}_2) = & -\frac{1}{a^2} + \frac{1}{a\phi_1} \left(\frac{\phi_{1L}}{r_{1L}} + \frac{\phi_{1R}}{r_{1R}} \right) + \frac{1}{a\phi_2} \left(\frac{\phi_{2L}}{r_{2L}} + \frac{\phi_{2R}}{r_{2R}} \right) - \left(\frac{1}{r_{1L}} + \frac{1}{r_{1R}} + \frac{1}{r_{2L}} + \frac{1}{r_{2R}} \right) \\ & + \frac{1}{r_{12}} + \left(\frac{\phi_{1L}\hat{\mathbf{r}}_{1L} + \phi_{1R}\hat{\mathbf{r}}_{1R}}{\phi_1} - \frac{\phi_{2L}\hat{\mathbf{r}}_{2L} + \phi_{2R}\hat{\mathbf{r}}_{2R}}{\phi_2} \right) \cdot \frac{\hat{\mathbf{r}}_{12}}{2a(1+\beta r_{12})^2} - \frac{(4\beta+1)r_{12}+4}{4(1+\beta r_{12})^4 r_{12}}. \end{aligned} \quad (11)$$

At this point we can eliminate two variational parameters by imposing the so-called Coulomb cusp conditions. These conditions are required to ensure that the energy does not blow up when either electron approaches either proton, or when the two electrons approach each other. The four cases of an electron approaching a proton leads to the same condition $a(1+e^{-s/a}) = 1$. The case of the two electrons approaching each other leads to the condition $\alpha = 2$.

Metropolis Monte Carlo Integration

To evaluate the integral (8) we use the Metropolis Monte Carlo method. In this method we choose random positions \mathbf{r}_1^i and \mathbf{r}_2^i from the distribution ω and approximate the integral as

$$\iint \omega \epsilon d^3\mathbf{r}_1 d^3\mathbf{r}_2 \approx \frac{1}{N} \sum_{i=1}^N \epsilon(\mathbf{r}_1^i, \mathbf{r}_2^i), \quad (12)$$

where $(\mathbf{r}_1^i, \mathbf{r}_2^i) \sim \omega$ are N randomly chosen points from the weight function. The key is to generate a lot of points from the distribution ω , which depends on Ψ , without having to normalize it. We can do this by generating a large set of points using a Markov Chain.

3. RESULTS AND DISCUSSION

Fusce mauris. Vestibulum luctus nibh at lectus. Sed bibendum, nulla a faucibus semper, leo velit ultricies tellus, ac venenatis arcu wisi vel nisl. Vestibulum diam. Aliquam pellentesque, augue quis sagittis posuere, turpis lacus congue quam, in hendrerit risus eros eget felis. Maecenas eget erat in sapien mattis porttitor. Vestibulum porttitor. Nulla facilisi. Sed a turpis eu lacus commodo facilisis. Morbi fringilla, wisi in dignissim interdum, justo lectus sagittis dui, et vehicula libero dui cursus dui. Mauris tempor ligula sed lacus. Duis cursus enim ut augue. Cras ac magna. Cras nulla. Nulla egestas. Curabitur a leo. Quisque egestas wisi eget nunc. Nam feugiat lacus vel est. Curabitur consectetur.

Suspendisse vel felis. Ut lorem lorem, interdum eu, tincidunt sit amet, laoreet vitae, arcu. Aenean faucibus pede eu ante. Praesent enim elit, rutrum at, molestie non, nonummy vel, nisl. Ut lectus eros, malesuada sit amet, fermentum eu, sodales cursus, magna. Donec eu purus. Quisque vehicula, urna sed ultricies auctor, pede lorem egestas dui, et convallis elit erat sed nulla. Donec luctus. Curabitur et nunc. Aliquam dolor odio, commodo pretium, ultricies non, pharetra in, velit. Integer arcu est, nonummy in, fermentum faucibus, egestas vel, odio.

4. CONCLUSIONS

Sed commodo posuere pede. Mauris ut est. Ut quis purus. Sed ac odio. Sed vehicula hendrerit sem. Duis non odio. Morbi ut dui. Sed accumsan risus eget odio. In hac habitasse platea dictumst.

Pellentesque non elit. Fusce sed justo eu urna porta tincidunt. Mauris felis odio, sollicitudin sed, volutpat a, ornare ac, erat. Morbi quis dolor. Donec pellentesque, erat ac sagittis semper, nunc dui lobortis purus, quis congue purus metus ultricies tellus. Proin et quam. Class aptent taciti sociosqu ad litora torquent per conubia nostra, per inceptos hymenaeos. Praesent sapien turpis, fermentum vel, eleifend faucibus, vehicula eu, lacus.

REFERENCES

- [1] L. Verlet (1967). Computer experiments on classical fluids, Thermodynamical properties of Lennard-Jones molecules, *Phys. Rev.* 159 (1967), 89 - 103.
- [2] J. L. Lebowitz, J. K. Percus, and L. Verlet, (1967). Ensemble dependence of fluctuations with application, to machine calculations, *Phys. Rev.*, 253 (1967), 250 - 254.
- [3] J. M. Thijssen (2007). Computational Physics, Ch. 7-8 *Cambridge University Press*