



上海交通大学

最优化方法

SHANGHAI JIAO TONG UNIVERSITY

120020910093

周贤敏

800 DONG CHUAN ROAD SHANGHAI 200240, THE PEOPLE'S REPUBLIC OF CHINA

Problem 1.

$$f(x_1, x_2, x_3) = x_1^2 = [x_1, x_2, x_3] \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}, Q = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} \rightarrow \text{positive semi-definite}$$

$$f(x_1, x_2, x_3) = x_1^2 + 2x_2^2 - x_1x_3 = \vec{x}^T \begin{bmatrix} 1 & 0 & -1 \\ 0 & 2 & 0 \\ 0 & 0 & 0 \end{bmatrix} \vec{x}, Q_0 = \begin{bmatrix} 1 & 0 & -\frac{1}{2} \\ 0 & 2 & 0 \\ -\frac{1}{2} & 0 & 0 \end{bmatrix} \rightarrow \lambda_1 = -0.207, \lambda_2 = 1.207, \lambda_3 = 2$$

it is indefinite.

$$f(x_1, x_2, x_3) = x_1^2 + x_2^2 + 2x_1x_2 + 2x_1x_3 + 2x_2x_3 = \vec{x}^T \begin{bmatrix} 1 & 1 & 0 \\ 1 & 0 & 1 \\ 0 & 1 & 1 \end{bmatrix} \vec{x}, Q = \begin{bmatrix} 1 & 1 & 0 \\ 1 & 0 & 1 \\ 0 & 1 & 1 \end{bmatrix} \rightarrow \lambda_1 = -1, \lambda_2 = 1, \lambda_3 = 2$$

it is indefinite

Problem 2.

$$f(\lambda x^{(1)} + (1-\lambda)x^{(2)}) = \|\lambda x^{(1)} + (1-\lambda)x^{(2)}\|^2 = \lambda^2 \|x^{(1)}\|^2 + 2\lambda(1-\lambda)x^{(1)} \cdot x^{(2)} + (1-\lambda)^2 \|x^{(2)}\|^2$$

$$(\lambda f(x^{(1)}) + (1-\lambda)f(x^{(2)}))^2 = \lambda^2 f^2(x^{(1)}) + (1-\lambda)^2 f^2(x^{(2)}) + 2\lambda(1-\lambda)f(x^{(1)})f(x^{(2)})$$

$$= \lambda^2 \|x^{(1)}\|^4 + (1-\lambda)^2 \|x^{(2)}\|^4 + 2\lambda(1-\lambda)\|x^{(1)}\|^2\|x^{(2)}\|^2$$


$$\therefore \lambda^2 \|x^{(1)}\|^2 + 2\lambda(1-\lambda)x^{(1)} \cdot x^{(2)} + (1-\lambda)^2 \|x^{(2)}\|^2 \leq \lambda^2 \|x^{(1)}\|^2 + 2\lambda(1-\lambda)\|x^{(1)}\|\|x^{(2)}\| + (1-\lambda)^2 \|x^{(2)}\|^2$$

$$\therefore f(\lambda x^{(1)} + (1-\lambda)x^{(2)}) \leq \lambda f(x^{(1)}) + (1-\lambda)f(x^{(2)})$$

so it is convex.

Problem 3.

I don't know how to prove that. Could you please upload the solution to the canvas? thx.





上海交通大学

SHANGHAI JIAO TONG UNIVERSITY

800 DONG CHUAN ROAD SHANGHAI 200240, THE PEOPLE'S REPUBLIC OF CHINA

Problem 7: $f(x) = x_1^4 + 2x_1^2x_2^2 + x_2^4$ $\nabla f(x) = \begin{bmatrix} \frac{\partial f}{\partial x_1} \\ \frac{\partial f}{\partial x_2} \end{bmatrix} = \begin{bmatrix} 4x_1^3 + 4x_1x_2^2 \\ 4x_2^3 + 4x_1^2x_2 \end{bmatrix}$

$$\nabla^2 f(x) = \begin{bmatrix} 12x_1^2 + 4x_2^2 & 8x_1x_2 \\ 8x_1x_2 & 12x_2^2 + 4x_1^2 \end{bmatrix}$$

$$f(x) = 4 + \begin{bmatrix} 8 \\ 8 \end{bmatrix}^T \begin{bmatrix} x_1 - 1 \\ x_2 - 1 \end{bmatrix} + \frac{1}{2} \begin{bmatrix} x_1 - 1 \\ x_2 - 1 \end{bmatrix}^T \begin{bmatrix} 16 & 8 \\ 8 & 16 \end{bmatrix} \begin{bmatrix} x_1 - 1 \\ x_2 - 1 \end{bmatrix}$$

$$= 8x_1^2 + 8x_2^2 - 16x_1 - 16x_2 + 8x_1x_2 + 12$$

Problem 8.

$$\max \sum_{i=1}^4 a_i x_i, \quad a_1 = 200, a_2 = 100, a_3 = 150, a_4 = 200$$

$$\text{s.t.} \quad 12x_1 + 10x_2 + 15x_3 + 20x_4 \leq 6000$$

$$8x_1 + 6x_2 + 2x_3 + 4x_4 \leq 2000$$

$$x_i \geq 0, \quad i = 1, 2, 3, 4$$


Problem 9.

$x_a = [0.5, 0.5]^T$ is feasible, on the boundary

$x_b = [1, 1]^T$ is infeasible.

$x_c = [-0.5, 0]^T$ is infeasible.

Problem 10.



$\min x_1$ is -2 , $x_2 = 0$.