## Assignment 2

## Gitanshu Arora

## Problem 11.16.5.8(exemplar):-

A die is loaded in such a way that each odd number is twice as likely to occur as each even number. Find P(G), where G is the event that a number greater than 3 occurs on a single roll of the die.

## Solution:-

Let *X* be a random variable denoting the number obtained on the die.

Let m be any natural number such that  $m \in \{1, 2, 3\}$ .

$$\Pr(X = k) = \begin{cases} 2p, & \text{if } k = 2m - 1 \\ p, & \text{if } k = 2m \end{cases} \tag{1}$$

Let  $F_X(k)$  be the cumulative distribution function such that,

$$F_X(k) = \Pr\left(X \le k\right) \tag{2}$$

If k = 2m - 1,

$$F_X(k) = \underbrace{\Pr(X=1) + \Pr(X=3) + ... \Pr(X=2m-1)}_{m \text{ terms}} + \underbrace{\Pr(X=2) + \Pr(X=4) + ... \Pr(X=2m-2)}_{m-1 \text{ terms}}$$
(3)

$$= (m)(2p) + (m-1)(p)$$
 (4)

$$= p(3m-1) \tag{5}$$

$$=\frac{p(3k+1)}{2}\tag{6}$$

If k = 2m,

$$F_X(k) = \underbrace{\Pr(X=1) + \Pr(X=3) + ... \Pr(X=2m-1)}_{m \text{ terms}} + \underbrace{\Pr(X=2) + \Pr(X=4) + ... \Pr(X=2m)}_{m \text{ terms}}$$
(7)

$$= (m)(2p) + (m)(p)$$
 (8)

$$= p(3m) \tag{9}$$

$$=\frac{3pk}{2}\tag{10}$$

So,

$$F_X(k) = \begin{cases} \frac{p(3k+1)}{2}, & \text{if } k = 2m - 1\\ \frac{3pk}{2}, & \text{if } k = 2m \end{cases}$$
 (11)

Since  $1 \le X \le 6$ ,

$$F_X(6) = 1 \tag{12}$$

$$\implies \frac{3p(6)}{2} = 1 \tag{13}$$

$$\implies p = \frac{1}{9} \tag{14}$$

$$Pr(G) = Pr(X > 3) \tag{15}$$

$$= F_X(6) - F_X(3) \tag{16}$$

$$=\frac{3p(6)}{2} - \frac{p\{3(3)+1\}}{2} \tag{17}$$

$$= 9p - 5p = 4p \tag{18}$$

$$=\frac{4}{9}\tag{19}$$

1