

The bootstrap

Here is a series of exercises on the non-parametric bootstrap. We will first empirically derive the probability that a given observation is part of a bootstrap sample.

```
# sampling with replacement
p.not.in.sample <- function(n){
  return((1-(1/n))**n)
}
```

Example: what is the probability that a given observation is not in the sample, if it has size 100?


```
p.not.in.sample(100)
```

```
## [1] 0.3660323
```

```
# now let's simulate and plot
```

```
plot(100:10000, p.not.in.sample(100:10000), main="The probability of not sampling a given observation as n grows")
```

The probability of not sampling a given observation as n grows



Looking at the above result we can answer the following question: what proportion of the original observations do you expect to be in a bootstrap sample of size n?

On average, weighting each observation as $1/n$, we would have that the number of observations left out would be approximately equal to $1/3$ (take the expectation of the indicator variable to see it). Hence, on average, we

would expect to have $2/3$ of the original dataset in a bootstrap sample.

Empirical coverage of Bootstrap confidence intervals

We want to estimate the trimmed mean of the Gamma distribution where the 10% largest and 10% smallest observations are trimmed.

```
set.seed(0)
# approximate true parameter value with a huge sample
true.tm <- mean(rgamma(100000000, shape = 2, rate = 1), trim = 0.1)
true.tm
```

```
## [1] 1.820736
```

So true.tm is our ground truth. We're now going to draw a small sample from the true distribution to use it to do inference.

```
n<-40
small.sample <- rgamma(n,shape=2,rate=1)
# our sample estimate
hat.tm <- mean(small.sample, trim = 0.1)
hat.tm
```


```
## [1] 2.171702
```

We'll now use Bootstrap to build 95% confidence intervals around our estimate.

```
# First of all, we need to create B bootstrap sets.
B <- 50
boot.sets <- matrix(nrow=n,ncol=B)
for(b in 1:B){
  b.set <- sample(small.sample,size=n, replace = T)
  boot.sets[,b] <- b.set
}
```


Let's visualize the first five datasets

```
par(mfrow=c(2,3))
hist(small.sample, main="original sample")
hist(boot.sets[,1], main="Boot sample 1")
hist(boot.sets[,2], main="Boot sample 2")
hist(boot.sets[,3], main="Boot sample 3")
hist(boot.sets[,4], main="Boot sample 4")
hist(boot.sets[,5], main="Boot sample 5")
```



Okay now we're ready to create the confidence intervals. We're going to create four different CIs.

```
hat.star <- matrix(nrow = B, ncol=1)
for(b in 1:B){
  hat.star[b]<-mean(boot.sets[,b],trim=0.1)
}
hist(hat.star, probability = T, breaks=20)
abline(v=hat.tm, col="red")
lines(density(hat.star), col="red")
```




The reversed quantile CI

```
# find the empirical quantiles of the bootstrap distribution
p.star <- hat.star - hat.tm
lower.q <- quantile(p.star, probs = 0.025)
upper.q <- quantile(p.star, probs = 0.975)
lower <- hat.tm - upper.q
upper <- hat.tm - lower.q
c(lower, upper)
```

```
##      97.5%      2.5%
## 1.643181 2.536170
```

```
hist(hat.star, xlim = c(1,3))
abline(v=lower, col="blue")
abline(v=upper, col="blue")
abline(v=hat.tm, col="red")
abline(v=true.tm, col="pink")
```



Let's check our results against R results.

```
require("boot")
```

```
## Loading required package: boot
```

```
# statistic functions
tm.fun <- function(x, ind){return(mean(x[ind], trim=0.1))}
tm.var <- function(x, ind){
  tm.value <- tm.fun(x[ind])
  # second level bootstrap to obtain variance of our estimate
  tm.variance <- var(boot(data=x[ind],R=50,statistic=tm.fun)$t)
  return(c(tm.value,tm.variance))
}
```

```
boot.res <- boot(data=small.sample,statistic=tm.fun, R=50, sim="ordinary")
boot.ci(boot.res, conf=0.95, type="perc")
```

```
## BOOTSTRAP CONFIDENCE INTERVAL CALCULATIONS
## Based on 50 bootstrap replicates
##
## CALL :
## boot.ci(boot.out = boot.res, conf = 0.95, type = "perc")
##
## Intervals :
## Level      Percentile
## 95%      ( 1.818,  2.714 )
## Calculations and Intervals on Original Scale
```

```
## Some percentile intervals may be unstable
```


The normal approximation CI

this one is base on the assumption that the estimate distribution tends to a Gaussian as n grows

```
sd.hat.tm <- sqrt(var(hat.star))
lower.q <- qnorm(0.025)*sd.hat.tm
lower <- hat.tm + lower.q
upper <- hat.tm - lower.q
c(lower,upper)
```

```
## [1] 1.683751 2.659654
```

```
hist(hat.star, xlim = c(1,3))
abline(v=lower, col="blue")
abline(v=upper, col="blue")
abline(v=hat.tm, col="red")
abline(v=true.tm, col="pink")
```



Now let's check with the R results:

```
boot.ci(boot.res, conf=0.95, type="norm")
```

```
## BOOTSTRAP CONFIDENCE INTERVAL CALCULATIONS
```

```
## Based on 50 bootstrap replicates
```

```
##
```

```
## CALL :
```

```
## boot.ci(boot.out = boot.res, conf = 0.95, type = "norm")
```

```
##
## Intervals :
## Level      Normal
## 95%   ( 1.742,  2.534 )
## Calculations and Intervals on Original Scale
```


The naive CI

The naive CI directly uses the `hat.star` quantiles. Note: no theoretical justification unless it's distribution is symmetric.

```
lower.q <- quantile(hat.star, probs = 0.025)
upper.q <- quantile(hat.star, probs = 0.975)
lower <- lower.q
upper <- upper.q
c(lower,upper)
```

```
##      2.5%      97.5%
## 1.807235 2.700224
```

```
hist(hat.star, xlim = c(1,3))
abline(v=lower, col="blue")
abline(v=upper, col="blue")
abline(v=hat.tm, col="red")
abline(v=true.tm, col="pink")
```



Again let's check with the R results:

```
boot.ci(boot.res, conf=0.95, type="basic")


## BOOTSTRAP CONFIDENCE INTERVAL CALCULATIONS
## Based on 50 bootstrap replicates
##
## CALL :
## boot.ci(boot.out = boot.res, conf = 0.95, type = "basic")
##
## Intervals :
## Level      Basic
## 95%      ( 1.629,  2.525 )
## Calculations and Intervals on Original Scale
## Some basic intervals may be unstable
```

The bootstrap T CI

Now, to obtain the bootstrap T CI we need a second level bootstrap.

```
C <- 50
hat.star.T <- matrix(nrow=B, ncol=1)
for(b in 1:B){
  hat.star.b<-matrix(nrow=C,ncol=1)
  for(c in 1:C){
    c.set <- sample(boot.sets[,b],size=n, replace = T)
    hat.star.b[c]<-mean(c.set,trim=0.1)
  }
  sd.b<-sqrt(var(hat.star.b))
  hat.star.T[b]<-(hat.star[b]-hat.tm)/sd.b
}
hist(hat.star.T, probability = T, breaks=20)
lines(density(hat.star.T), col="red")
```


Histogram of hat.star.T




We'll now use the empirical quantiles of this simil-T distribution to estimate the CIs.

```
lower.q <- quantile(hat.star.T, probs = 0.025)*sd.hat.tm
upper.q <- quantile(hat.star.T, probs = 0.975)*sd.hat.tm
lower <- hat.tm + lower.q
upper <- hat.tm - lower.q
c(lower,upper)
```

```
## [1] 1.747416 2.595989
```

```
hist(hat.star, xlim = c(1,3))
abline(v=lower, col="blue")
abline(v=upper, col="blue")
abline(v=hat.tm, col="red")
abline(v=true.tm, col="pink")
```



And finally, let's check with the R results:

```
boot.res <- boot(data=small.sample, statistic=tm.var, R=50, sim="ordinary")
boot.ci(boot.res, conf=0.95, type="stud", index = c(1,2))
```

```
## BOOTSTRAP CONFIDENCE INTERVAL CALCULATIONS
## Based on 50 bootstrap replicates
##
## CALL :
## boot.ci(boot.out = boot.res, conf = 0.95, type = "stud", index = c(1,
## 2))
##
## Intervals :
## Level      Studentized
## 95%      ( 1.807,  2.628 )
## Calculations and Intervals on Original Scale
## Some studentized intervals may be unstable
```

Okay, as you have probably noticed, the bootstrap CI are variable. We'll now study their coverage exploiting this variability with a simulation.

```
nsim<-1000
```