

## PRACTICE PROBLEMS CHAPTER 6

1.

- (i) (a)  $\det(A - \lambda I) = \lambda^2 - 1 \Rightarrow \lambda_1 = -1, \lambda_2 = 1$ .  
 $A - \lambda_1 I = A + I = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix} \rightarrow \text{RREF} \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix} \Rightarrow$  e-vector associated to  $\lambda_1 = -1$  is  $\mathbf{x}_1 = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$ ,  
 $A - \lambda_2 I = A - I = \begin{bmatrix} -1 & 1 \\ 1 & -1 \end{bmatrix} \rightarrow \text{RREF} \begin{bmatrix} 1 & -1 \\ 0 & 0 \end{bmatrix} \Rightarrow$  e-vector associated to  $\lambda_2 = 1$  is  $\mathbf{x}_2 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$ .
- (b)  $\det(A - \lambda I) = \lambda^2 + 1 \Rightarrow \lambda_1 = i, \lambda_2 = -i$ .  
 $A - \lambda_1 I = A - iI = \begin{bmatrix} -i & 1 \\ -1 & -i \end{bmatrix} \rightarrow \text{RREF} \begin{bmatrix} 1 & i \\ 0 & 0 \end{bmatrix} \Rightarrow$  e-vector associated to  $\lambda_1 = i$  is  $\mathbf{x}_1 = \begin{bmatrix} -i \\ 1 \end{bmatrix}$ ,  
 The e-vector associate to  $\lambda_2 = -i$  is the complex conjugate of  $\mathbf{x}_1$ :  $\mathbf{x}_2 = \begin{bmatrix} i \\ 1 \end{bmatrix}$ .
- (c)  $\det(A - \lambda I) = (1 - \lambda)^2 \Rightarrow \lambda_1 = \lambda_2 = 1, \lambda_1 = 1$  (AM = 2)  
 $A - \lambda_1 I = A - I = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} \Rightarrow$  two free variables and possible e-vectors are  $\mathbf{x}_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \mathbf{x}_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$ . (GM = 2)
- (d)  $\det(A - \lambda I) = (1 - \lambda)^2 \Rightarrow \lambda_1 = \lambda_2 = 1, \lambda_1 = 1$  (AM = 2)  
 $A - \lambda_1 I = A - I = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} \Rightarrow$  e-vector associated to  $\lambda_1 = 1$  is  $\mathbf{x} = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ . (GM = 1)
- (ii) (a) The matrix is diagonalizable (distinct e-vectors):  $X = \begin{bmatrix} 1 & 1 \\ -1 & 1 \end{bmatrix}$  and  $D = \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix}$   
 (b) The matrix is diagonalizable (distinct e-vectors):  $X = \begin{bmatrix} -i & i \\ 1 & 1 \end{bmatrix}$  and  $D = \begin{bmatrix} i & 0 \\ 0 & -i \end{bmatrix}$   
 (c) The matrix is diagonalizable (AM = GM = 2):  $X = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$  and  $D = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$   
 (d) Since the matrix has one linearly independent eigenvector (GM = 1) associated to the repeated eigenvalue  $\lambda = 1$  (AM = 2), we have GM < AM and the matrix is defective (non diagonalizable).

2. (a)  $\det(A - \lambda I) = (1 - \lambda)(-\lambda)(3 - \lambda) \Rightarrow$  the eigenvalues of  $A$  are  $\lambda_1 = 1, \lambda_2 = 0, \lambda_3 = 3$ .

The eigenvalues of  $A^2$  are 1, 0, 9 and the eigenvalues of  $A^n$  are  $1, 0, 3^n$ .

- (b)  $A - \lambda_1 I = A - I = \begin{bmatrix} 0 & 2 & 1 \\ 0 & -1 & 6 \\ 0 & 0 & 2 \end{bmatrix} \rightarrow \text{RREF} \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix} \Rightarrow$  e-vector associated to  $\lambda_1 = 1$  is  $\mathbf{x}_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$ ,  
 $A - \lambda_2 I = A - 0I = A = \begin{bmatrix} 1 & 2 & 1 \\ 0 & 0 & 6 \\ 0 & 0 & 3 \end{bmatrix} \rightarrow \text{RREF} \begin{bmatrix} 1 & 2 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix} \Rightarrow$  e-vector associated to  $\lambda_2 = 0$  is  $\mathbf{x}_2 = \begin{bmatrix} -2 \\ 1 \\ 0 \end{bmatrix}$ ,

$$A - \lambda_3 I = A - 3I = \begin{bmatrix} -2 & 2 & 1 \\ 0 & -3 & 6 \\ 0 & 0 & 0 \end{bmatrix} \rightarrow \text{RREF} \begin{bmatrix} 1 & 0 & -\frac{5}{2} \\ 0 & 1 & -2 \\ 0 & 0 & 0 \end{bmatrix} \Rightarrow \text{e-vector associated to } \lambda_3 = 3 \text{ is } \mathbf{x}_3 = \begin{bmatrix} 5 \\ 4 \\ 2 \end{bmatrix}$$

The eigenspaces for the matrix  $A$  have basis  $\mathbf{x}_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \mathbf{x}_2 = \begin{bmatrix} -2 \\ 1 \\ 0 \end{bmatrix}, \mathbf{x}_3 = \begin{bmatrix} 5 \\ 4 \\ 2 \end{bmatrix}$

These are also the basis for the eigenspaces for the matrix  $A^2$  and  $A^n$ .

(c) Since  $A = XDX^{-1} = \begin{bmatrix} 1 & -2 & 5 \\ 0 & 1 & 4 \\ 0 & 0 & 2 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 3 \end{bmatrix} \begin{bmatrix} 1 & 2 & -\frac{13}{2} \\ 0 & 1 & -2 \\ 0 & 0 & \frac{1}{2} \end{bmatrix}$

$$\text{we have } A^n = X D^n X^{-1} = \begin{bmatrix} 1 & -2 & 5 \\ 0 & 1 & 4 \\ 0 & 0 & 2 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 3^n \end{bmatrix} \begin{bmatrix} 1 & 2 & -\frac{13}{2} \\ 0 & 1 & -2 \\ 0 & 0 & \frac{1}{2} \end{bmatrix} = \begin{bmatrix} 1 & 2 & -\frac{13}{2} + \frac{5(3^n)}{2} \\ 0 & 0 & 2(3^n) \\ 0 & 0 & 3^n \end{bmatrix}$$

(d) Substituting  $n = 7$  in the formula from part (c) gives

$$A^7 = \begin{bmatrix} 1 & 2 & -\frac{13}{2} + \frac{5(3^7)}{2} \\ 0 & 0 & 2(3^7) \\ 0 & 0 & 3^7 \end{bmatrix} = \begin{bmatrix} 1 & 2 & 5461 \\ 0 & 0 & 4374 \\ 0 & 0 & 2187 \end{bmatrix}$$

3.

(a) The eigenvalues of  $A$  are  $\lambda_1 = 2$  and  $\lambda_2 = 3$ .

(b) The eigenvalues of  $A^2 + \alpha A + \beta I$  are  $2^2 + 2\alpha + \beta$  and  $3^2 + 3\alpha + \beta$

4. The characteristic equation of the matrix is given by  $\lambda^2 - 2\lambda + 9k - 35 = 0$ . From the quadratic formula we have that this equation has two distinct solutions if and only if  $b^2 - 4ac = 4 - 4(9k - 35) > 0$ . Solving the inequality gives  $k < 4$ .

5.  $\text{tr}(A) = \lambda_1 + \lambda_2 = 5$ ,  $\det(A) = \lambda_1 \lambda_2 = -14 \Rightarrow \lambda_1 = -2, \lambda_2 = 7$

6. (a) Putting the vectors into a matrix:  $\begin{bmatrix} 1 & 3 & 19 \\ -1 & 2 & 6 \\ -2 & -1 & -13 \end{bmatrix}$  RREF  $\begin{bmatrix} 1 & 0 & 4 \\ 0 & 1 & 5 \\ 0 & 0 & 0 \end{bmatrix}$ . Thus  $\mathbf{v} = 4\mathbf{x}_1 + 5\mathbf{x}_2$

$$(b) A\mathbf{v} = A(4\mathbf{x}_1 + 5\mathbf{x}_2) = 4(A\mathbf{x}_1) + 5(A\mathbf{x}_2) = 4(\lambda_1 \mathbf{x}_1) + 5(\lambda_2 \mathbf{x}_2) = (4)(-2) \begin{bmatrix} 1 \\ -1 \\ -2 \end{bmatrix} + (5)(3) \begin{bmatrix} 3 \\ 2 \\ -1 \end{bmatrix} = \begin{bmatrix} 37 \\ 38 \\ 1 \end{bmatrix}$$

7. (a)  $\det(A - \lambda I) = (3 - \lambda)^3$ , thus the matrix has a repeated eigenvalue  $\lambda = 3$  of Algebraic Multiplicity three

$$A - \lambda_1 I = A - 3I = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix} \quad \text{The matrix is already in RREF and a basis of eigenspace is } \mathbf{x} = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix},$$

Since  $\text{GM} = 1 < \text{AM} = 3$ , the matrix is defective and not diagonalizable.

(b)  $\det(A - \lambda I) = (4 - \lambda)(1 - \lambda)^2$ . The matrix has the eigenvalue  $\lambda_1 = 1$  with  $\text{AM} = 2$  and  $\lambda_2 = 4$  with  $\text{AM} = 1$ . In order to determine whether the matrix is diagonalizable we need to determine the Geometric Multiplicity of the repeated eigenvalue  $\lambda_1 = 1$ .

$$A - \lambda_1 I = A - I = \begin{bmatrix} 3 & -3 & 0 \\ 0 & 0 & 0 \\ -4 & 4 & 0 \end{bmatrix} \Rightarrow \text{RREF} \begin{bmatrix} 1 & -1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} \quad \text{Basis of eigenspace associated to } \lambda_1 = 1 \text{ is}$$

$$\mathbf{x}_1 = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \quad \mathbf{x}_2 = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}. \quad \text{Thus } \text{GM} = \text{AM} = 2 \text{ and the matrix is diagonalizable.}$$

The eigenvalue  $\lambda = 4$  has associated eigenvector  $[-3, 0, 4]^T$  and a possible diagonalization is:

$$X = \begin{bmatrix} 1 & 0 & -3 \\ 1 & 0 & 0 \\ 0 & 1 & 4 \end{bmatrix} \quad \text{and} \quad D = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 4 \end{bmatrix}$$

8. In Problem 7 part (b) we found that the matrix  $A$  is diagonalizable as

$$A = XDX^{-1} = \begin{bmatrix} 1 & 0 & -3 \\ 1 & 0 & 0 \\ 0 & 1 & 4 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 4 \end{bmatrix} \begin{bmatrix} 0 & 1 & 0 \\ 4/3 & -4/3 & 1 \\ -1/3 & 1/3 & 0 \end{bmatrix}$$

The matrix  $B$  is given by  $B = X \sqrt{D} X^{-1} = \begin{bmatrix} 1 & 0 & -3 \\ 1 & 0 & 0 \\ 0 & 1 & 4 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 2 \end{bmatrix} \begin{bmatrix} 0 & 1 & 0 \\ 4/3 & -4/3 & 1 \\ -1/3 & 1/3 & 0 \end{bmatrix} = \begin{bmatrix} 2 & -1 & 0 \\ 0 & 1 & 0 \\ -4/3 & 4/3 & 1 \end{bmatrix}$

9. If  $\lambda_1 = 3 + 2i$  is an eigenvalue, then the complex conjugate  $\lambda_2 = 3 - 2i$  is also an eigenvalue.

If the matrix is singular, then  $\lambda_3 = 0$  is an eigenvalue. The sum of the eigenvalues equals the trace of the matrix which is given by  $a_{11} + a_{22} + a_{33} + a_{44} = 4$ , thus  $(3 + 2i) + (3 - 2i) + 0 + \lambda_4 = 4$  which gives  $\lambda_4 = -2$ .

10.

(a) If  $\lambda = 0$  then  $\det(A - \lambda I) = \det(A) = 0$  and the matrix is singular which contradicts the assumption, thus  $\lambda \neq 0$ .

(b) If  $A\mathbf{x} = \lambda\mathbf{x}$  then  $A^{-1}A\mathbf{x} = \lambda A^{-1}\mathbf{x}$  which gives  $\mathbf{x} = \lambda A^{-1}\mathbf{x}$  or  $A^{-1}\mathbf{x} = \frac{1}{\lambda}\mathbf{x}$  thus  $\frac{1}{\lambda}$  is an eigenvalue of  $A^{-1}$  with  $\mathbf{x}$  the corresponding eigenvector.

11. The matrix has the eigenvalue  $\lambda = a$  with algebraic multiplicity 3. The basis of the corresponding eigenspace consists of two linearly independent eigenvectors:  $[1, 0, 0]^T$  and  $[0, 1, 0]^T$ . Since there are only two linearly independent eigenvectors, we have  $\text{GM} = 2 < \text{AM} = 3$  and the matrix is defective.

12. If  $A$  is diagonalizable, then  $A = XDX^{-1}$  where  $D$  is a diagonal matrix. If  $B$  is similar to  $A$ , then there exists a nonsingular matrix  $S$  such that  $B = S^{-1}AS$ . It follows that

$$\begin{aligned} B &= S^{-1}(XDX^{-1})S \\ &= (S^{-1}X)D(S^{-1}X)^{-1} \end{aligned}$$

Therefore  $B$  is diagonalizable with diagonalizing matrix  $S^{-1}X$ .

13.

(a) The rank of  $A$  is given by the number of nonzero singular values, thus the rank is 3.

(b) An orthonormal basis for  $R(A^T)$  is given by  $\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3$ , where  $\mathbf{v}_i$  is the  $i$ th column of  $V$ .

(c) An orthonormal basis is given by  $\mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3$  where  $\mathbf{u}_i$  is the  $i$ th column of  $U$ .

(d) The rank-1 matrix  $B$  that is the closest matrix of rank-1 to  $A$  is given by

$$B = \sigma_1 \mathbf{u}_1 \mathbf{v}_1^T = 100 \begin{bmatrix} 2/5 \\ 2/5 \\ 2/5 \\ 2/5 \\ 3/5 \end{bmatrix} \begin{bmatrix} 1/2 & 1/2 & 1/2 & 1/2 \end{bmatrix} = \begin{bmatrix} 20 & 20 & 20 & 20 \\ 20 & 20 & 20 & 20 \\ 20 & 20 & 20 & 20 \\ 20 & 20 & 20 & 20 \\ 30 & 30 & 30 & 30 \end{bmatrix}$$

(e) From Theorem 6.5.3 we have that  $\|B - A\|_F = \sqrt{\sigma_2^2 + \sigma_3^2} = 10\sqrt{2}$ .

$$(f) \quad C = \sigma_1 \mathbf{u}_1 \mathbf{v}_1^T + \sigma_2 \mathbf{u}_2 \mathbf{v}_2^T = B + 10 \begin{bmatrix} -2/5 \\ -2/5 \\ -2/5 \\ 3/5 \\ 2/5 \end{bmatrix} \begin{bmatrix} 1/2 & -1/2 & -1/2 & 1/2 \end{bmatrix} = \begin{bmatrix} 18 & 22 & 22 & 18 \\ 18 & 22 & 22 & 18 \\ 18 & 22 & 22 & 18 \\ 23 & 17 & 17 & 23 \\ 32 & 28 & 28 & 32 \end{bmatrix}$$

(g)  $\|C - A\|_F = \sigma_3 = 10$ .