

Sam Grayson's Notebook (with L^AT_EX)

January 14, 2015

- 1.1 $ma = b$
 $na = c$
 $na + ma = b + c$
 $(n + m)a = b + c$
 $a|(b + c)$
 Definition of 'divides'
 Definition of 'divides'
 Algebra
 Algebra
 Definition of 'divides' ■
- 1.2 Let $d = -c$
 $a|(b + d)$
 $a|(b - c)$
 Theorem 1.1
 substitution ■
- 1.3 $ma = b$
 $na = c$
 $mana = bc$
 $a|bc$
 Definition of 'divides'
 Definition of 'divides'
 Algebra
 Definition of 'divides' ■
- 1.4 $mana = bc$
 $a^2|bc$
 see last proof
 Definition of 'divides' ■
- 1.5 If $a|b$ then $a|b^n$
 $b = ka$
 $b^n = (ka)^n = k^nk^{(n-1)}a^n$
 $k|b^n$
 Definition of 'divides'
 Algebra
 Definition of 'divides' ■
- 1.6 $ka = b$
 $ack = bc$
 $a|bc$
 Definition of 'divides'
 Algebra
 Definition of 'divides' ■
- 1.7 1. $45 - 9 = 36 = 9 \cdot 4$. True
 2. $37 - 2 = 35 = 7 \cdot 5$. True
 3. $37 - 3 = 34$. False
 4. $37 - (-3) = 40 = 8 \cdot 5$. True
- 1.8 let k be all the numbers
 where $k \equiv b \pmod{3}$
 $3|(k - b)$
 $3n = k - b$
 $3n + k = n$
 Definition of 'mod'
 Definition of 'divides'
 Algebra ■
- 1.9 $a - a = 0 = 0n$
 $n|(a - a)$
 $a \equiv 0 \pmod{n}$
 Arithmetic
 Definition of 'divides'
 Definition of 'mod' ■

1.10	$n (a-b)$ $kn = a-b$ $-kn = b-a$ $n (b-a)$ $b \equiv a \pmod{n}$ ■	Definition of 'mod' Definition of 'divides' Algebra Definition of 'divides'
1.11	$n (a-b)$ $n (b-c)$ $n (a-b+b-c)$ $n (a-c)$ $a \equiv c \pmod{n}$	Definition of 'mod' Definition of 'mod' Theorem 1.1 Algebra Definition of 'mod' ■
1.12	$n (a-b)$ $n (c-d)$ $n (a+c-b-d)$ $n ((a+c)-(b+d))$ $a+c \equiv b+d \pmod{n}$	Definition of 'mod' Definition of 'mod' Theorem 1.1 Algebra definition 'mod' ■
1.13	let $e = -c$ and $f = -d$ $a+e \equiv b+f$ $a-c \equiv b-d$	Theorem 1.12 substitution ■
1.14	$n (a-b)$ $n (c-d)$ $n (a-b)(c-d)$	Definition of 'mod' Definition of 'mod' Theorem 1.3 ■
1.15	$a \equiv b \pmod{n}$ $a^2 \equiv b^2 \pmod{n}$	Premise Theorem 1.14 ■
1.16	$a \equiv b \pmod{n}$ $a^2 \equiv b^2 \pmod{n}$ $a^2a \equiv b^2b \pmod{n}$ $a^3 \equiv b^3 \pmod{n}$	Premise Theorem 1.15 Theorem 1.14 Algebra ■
1.17	$a \equiv b \pmod{n}$ $a^{k-1} \equiv b^{k-1} \pmod{n}$ $a^{k-1}a \equiv b^{k-1}b \pmod{n}$ $a^k \equiv b^k \pmod{n}$	Premise Premise Theorem 1.14 Algebra ■
1.18	Base case: $a \equiv b \pmod{n}$ Inductive Hypothesis: $a^{k-1} \equiv b^{k-1} \pmod{n}$ Inductive step: $a^{k-1}a \equiv b^{k-1}b \pmod{n}$ $a^k \equiv b^k \pmod{n}$ Conclusion: $a^k \equiv b^k \pmod{n}$	Premise (assumption) Theorem 1.14 Algebra inductively ■
1.19	12. $6 \equiv 2 \pmod{4}$ $5 \equiv 1 \pmod{4}$	

$$6 + 5 \equiv 2 + 1 \pmod{4}$$

$$13. \quad 6 - 5 \equiv 2 - 1 \pmod{4}$$

$$14. \quad 6 \cdot 5 \equiv 2 \cdot 1$$

$$15. \quad 6^2 \equiv 2^2 \pmod{4}$$

$$16. \quad 6^3 \equiv 2^3 \pmod{4}$$

$$17. \quad 6^4 \equiv 2^4 \pmod{4}$$

$$18. \quad 6^k \equiv 2^k \pmod{4}$$

1.20 No

Consider the case where $n = 4$, $c = 0$, $a = 1$, and $b = 2$.

$$ac \equiv bc \pmod{n}$$

$$a \neq b$$

1.21 See 1.22 and 1.23

$$1.22 \quad 3|a$$

$$3|b$$

$$3|9$$

$$3|(9b_k 10^{k-1})$$

$$3|(b - 9b_k 10^{k-1})$$

$$3|(b_{k-1} + b_k)b_{k-2} \dots b_0$$

$$3|(a_k + a_{k-1} + a_{k-2} + \dots a_1 + a_0)$$

Here is the algebra I used in the step labeled ‘Algebra*’:

$$\begin{array}{rcccccc}
 & & & & & b - b_k 9 10^{k-1} & = \\
 & & & & & b - b_k (10 - 1) 10^{k-1} & = \\
 & & & & & b + (-b_k 10 \cdot 10^{k-1} + b_k 1 10^{k-1}) & = \\
 & & & & & b + (-b_k 10^k + b_k 10^{k-1}) & = \\
 + & \begin{array}{cccccc} b_k & b_{k-1} & b_{k-2} & \dots & b_0 \\ (-b_k) & b_k & 0 & \dots & 0 \end{array} & = \\
 \hline
 & (b_k + b_{k-1}) & b_{k-2} & \dots & b_0 &
 \end{array}$$

$$1.23 \quad 3|a$$

$$3|(b_k + b_{k-1} + \dots + b_0)$$

$$3|9$$

$$3|(b_k 9c) \text{ where } c \text{ is } k \text{ ones in a row}$$

$$3|(b_k + b_{k-1} + \dots + b_0 + b_k 9c)$$

$$3|(b_k 10^k + b_{k-1} + \dots + b_0)$$

$$3|(a_k 10^k + a_{k-1} 10^{k-1} + \dots + a_0 10^0)$$

$$3|(a_k a_{k-1} \dots a_0)$$

Premise (Base Case)

Let b be an integer where... (Inductive Hypothesis)

Arithmetic

Theorem 1.3

Theorem 1.2

Algebra* (Inductive Step)

Inductive axiom ■

Premise (Base Case)

Assumption (Inductive Hypothesis)

Arithmetic

Theorem 1.3

Theorem 1.2

Algebra*

Inductive Axiom

Definition of digits ■

Here is the algebra I used in the step labeled ‘Algebra*’:

$$\begin{aligned}
 b_k + b_{k-1} + \dots + b_0 + b_k 9c &= \\
 b_k + b_{k-1} + \dots + b_0 + b_k d &= \text{ where } d \text{ is a number with } k \text{ nines} \\
 b_k + b_{k-1} + \dots + b_0 + b_k(10^k - 1) &= \\
 b_k + b_{k-1} + \dots + b_0 + b_k 10^k - b_k &= \\
 b_{k-1} + \dots + b_0 + b_k 10^k &
 \end{aligned}$$

$$1.24 \quad 4|a \text{ if and only if } 4|(a_1 + a_3 + \dots)(a_0 + a_2 + a_4 + \dots)$$

$$1.25 \quad 1. \quad m = nq + r \text{ where } m = 25, n = 7, q = 3, \text{ and } r = 4$$

$$2. \quad m = 277, n = 4, q = 66, \text{ and } r = 1$$

$$3. \quad m = 33, n = 11, q = 3, r = 0$$

$$4. \quad m = 33, n = 45, q = 0, r = 33$$

1.26 Setup:

(Make a list of multiples of n that are greater than m and choose the smallest one to define $n(q + 1)$.)

$$A := \{k | k \in \mathbb{N} \wedge kn > m\}$$

$$\exists a \ni (a \in A \wedge an > m \wedge \forall k \in A (a \leq k))$$

Well-ordering Principle

$$q := a - 1$$

$$r := m - nq$$

Proving r satisfies upper bound

(If it didn't, then a wouldn't be an element of A , but we know that a is in A .) $r > n - 1$ Assume for contradiction

$$r \geq n$$

Inequality over integers

$$\exists j \ni (r - n = j \wedge j \geq 0)$$

Property of inequalities

$$nq + r = m$$

Algebra (from definition of r)

$$nq + (n + j) = m$$

Algebra

$$n(q + 1) + j = m$$

Algebra

$$n(q + 1) \leq m$$

Property of inequalities

$$n(q + 1) > m$$

Algebra (from definition of a)

$$\therefore r \leq n - 1$$

Contradiction

Proving r satisfies lower bound

(If it didn't, then there would be another element smaller than a in A , but a is the least element in A .) $r < 0$ Assume for contradiction

$$nq + r = m$$

Algebra (from definition of r)

$$nq > m$$

Property of inequalities

$$q \in A$$

$q \in \mathbb{N} \wedge nq > m$ is the c

$$\forall k (k \in A \rightarrow q + 1 \leq k)$$

Definition of a (it's the smallest

$$q \leq q + 1$$

Universal instantiation

$$\therefore r \geq 0$$

Contradiction

■