

$$\frac{2.267061\text{grams}}{1.39 \times 10^{24}} \approx \frac{\frac{c}{L_F} T_{SI} M_F}{1.39 \times 10^{24}} = \sqrt{\frac{k_e^2 q_e^4 T}{c G M L^2}} = \sqrt{\frac{c \alpha^2 \hbar^2 T}{G M L^2}} = M_F$$

©Roy Pfund* (CC BY-NC-SA 4.0) 2022

<https://creativecommons.org/licenses/by-nc-sa/4.0/>

April 9, 2022

2 Speed of Sound = c_0

$$\frac{h \alpha}{c q_e^2} \frac{2c^2}{4\pi} = \frac{\mu_0 c^2}{4\pi} = \frac{1}{4\pi\epsilon_0} = k_e; \quad \frac{\mu_0}{2} \frac{c q_e^2}{2\pi \hbar} = \alpha$$

= 0.00729735257 “It has been a mystery ever since it was discovered more than fifty years ago, and all good theoretical physicists put this number up on their wall and worry about it.” [Feynman, 1985, p. 129]. Some CFD software uses plank units to reduce computation by reducing the billions of multiplications of k_b ; we should care about people as much as machines. Given how the constants above are involved with α , it is possible to reduce the number of constants people need to know drastically $M_F^W L_F^X T_F^Y Q_F^Z =$

$$\left(\sqrt{\frac{c \alpha^2 \hbar^2 T}{G M L^2}} \right)^W \left(\frac{G}{c^2} M_F \frac{M L^2}{\alpha^2 \hbar T} \right)^X \left(\frac{G}{c^3} M_F \frac{M L^2}{\alpha^2 \hbar T} \right)^Y (q_e)^Z$$

$$\begin{aligned} k_e &= \alpha M_F^1 L_F^3 T_F^{-2} Q_F^{-2} & \hbar &= 1 M_F^1 L_F^2 T_F^{-1} \\ \mu_0 &= \alpha 4 \pi M_F^1 L_F^1 Q_F^{-2} & c &= 1 L_F^1 T_F^{-1} \\ \epsilon_0 &= \frac{1}{\alpha 4 \pi} M_F^{-1} L_F^{-3} T_F^2 Q_F^2 & N_A &= 1^{**} \\ R_\infty &= \frac{m_e}{M_F} \frac{\alpha^2}{4 \pi} L_F^{-1} & M_F c^2 &= E_F = 1 M_F^1 L_F^2 T_F^{-2} \\ & & \frac{2 \pi \hbar}{c q_e} &= k_b = 2 \pi M_F^1 L_F^1 Q_F^{-1} \end{aligned}$$

Given $c_0 = \sqrt{\frac{\gamma_0 N_A k_B K}{MolarMass}}$ and $39.947g \text{ mol}^{-1}$ = molar mass of the argon gas from the experiment measuring c_0 in a purified isotope of argon gas at the Triple-Point of water = 273.16K [de-Podesta et al., 2013] Where $U_F = \frac{M L^2}{k_b T^2}$, $K = \frac{Kelvin}{U_F} U_F$, and $\gamma_0 = 5/3$ for monotonic gases. Let's see how this matches up with the $c_0^2 = 94756.245 m^2 s^{-2}$ from the experiment in 2013. e.g. on how to correct N_A being based on the Dalton = $\frac{1}{12}$ the mass of Carbon isotope C^{12} over to M_F

$$**\text{adjusted argon gas molar mass} = 40.671 M_F = \frac{39.947 kg}{M_F N_A}$$

$$c_0 = \sqrt{\frac{\frac{5}{3} N_A k_B \frac{273.16 Kelvin}{U_F} U_F}{40.671 M_F}} = 307.701 m s^{-1} \approx \sqrt{c_0^2}$$

3 Time is On Our Side(& Distance)

With a Sympathetic Constant = D_C = light-second / L_F to save our wallets, watches, measuring wheels, and road signage we can still use existing definitions of distance and time. Perhaps one day for the sake of simplicity, Bureau international des poids et mesures might redefine the second and meter such that D_C is exactly 1.39×10^{24} rather than approx. 5 almost 6 nines.

$$\frac{1.3899982 \times 10^{24} \approx \frac{c T_{SI}}{L_F} = D_C}{1.39 \times 10^{24}} = 0.999998675$$

1 Dimensionality

Stoney-Mass $M_S = \sqrt{\frac{\hbar \alpha c}{G}}$ units are correct, but with $M_F = M_S \sqrt{\hbar \alpha}$ alone gives correct arithmetic; for all the equations above to not only work arithmetic wise, $\sqrt{\frac{T}{M^1 L^2}}$ is needed to cancel any wrong units put into M_F from \hbar , M , L , and T with no subscripts are meant to mean from whichever unit system you are converting from or currently using, in this case, SI units 1kg, 1m, and 1s respectively. $\sqrt{\frac{T}{M^1 L^2}}$ reappears in the solution for G . However G is wrong to begin with, because with Relativity it doesn't really exist, as it is the curvature of space time.


$$\alpha^2 \hbar = \alpha^2 \frac{M_F L_F^2}{T_F} = \frac{G}{M_F^3 L_F^{-1} T_F^{-2}}$$

*e-mail: r0ypfund@gm411.c0m

4 There and back again

Calibrating scales to $D_C M_F \approx 2.267061\text{grams}$ and ammeter to $D_C q_e \approx 222702.26$ Coulombs will result in $M_F^W L_F^X T_F^Y Q_F^Z = \left(\frac{p = D_C M_F}{1.39 \times 10^{24}} \right)^W \left(\frac{299792458 \text{Meters}}{1.39 \times 10^{24}} \right)^X \left(\frac{1 \text{Second}}{1.39 \times 10^{24}} \right)^Y \left(\frac{q = D_C q_e}{1.39 \times 10^{24}} \right)^Z$

$$\begin{aligned} \mu_0 &= \alpha 4 \pi \left(c \frac{s}{m} \right) p^1 m^1 q^{-2} & k_b &= 2 \pi D_C^{-1} \left(c \frac{s}{m} \right) p^1 m^1 q^{-1} \\ \epsilon_0 &= \frac{1}{\alpha 4 \pi} \left(c \frac{s}{m} \right)^{-3} p^{-1} m^{-3} s^2 q^2 & k_e &= \alpha \left(c \frac{s}{m} \right)^3 p^1 m^3 s^{-2} q^{-2} \\ R_\infty &= \frac{\alpha^2}{4 \pi} D_C^2 \left(c \frac{s}{m} \right)^{-1} \frac{m_e}{p} m^{-1} & \hbar &= D_C^{-2} \left(c \frac{s}{m} \right)^2 p^1 m^2 s^{-1} \\ G &= \frac{\hbar T}{M L^2} \alpha^2 \left(c \frac{s}{m} \right)^3 p^3 m^{-1} s^{-2} & c &= \left(c \frac{s}{m} \right) m^1 s^{-1} \\ & & N_A &= D_C^{**} \end{aligned}$$



5 Constants

SI Avogadro constant $N_A = 6.02214076 \times 10^{23} \approx \frac{0.012 \text{ kg}}{\text{mole } ^{12}\text{C}}$

Planck constant $h = 6.62607015 \times 10^{-34} \text{ kg m}^2 \text{ s}^{-1}$

lightspeed constant $c = 299792458 \text{ m s}^{-1}$

electron charge $q_e = 1.602176634 \times 10^{-19} \text{ Coulombs}$

gravity constant $G = 6.67430 \times 10^{-11} \text{ kg}^{-1} \text{ m}^3 \text{ s}^{-2}$

s = “duration of 9192631770 periods of the radiation corresponding to the transition between the two hyperfine levels of the ground state of the cesium 133 atom”[SI, 1968] to finish the definition of c , An international agreement in Paris on Oct. 20 1983 defines the meter as $\frac{1}{299792458}$ the distance light travels in a vacuum in 1 second[Times, 1983], 2018 Codata values of q_e , and G , come from [Tiesinga et al., 2021]. On May 20, 2019 the values of $\hbar = \frac{h}{2\pi}$, and h , were fixed to their current value along with N_A to the Dalton = $\frac{1}{12}$ the mass ^{12}C isotope of Carbon[Bettin].

5.1 How the Avogadro constant was measured for the last time

This section is paraphrased almost verbatim from [Bettin]. N_A and h were measured using a incredibly round & pure ball of Si^{28} and a Kibble balance. Where $\alpha^2 m_e c / 2 h = R_\infty$ is the Rydberg constant, $\sum_{i=28}^{30} x_i A_r(^i\text{Si}) = A_r(\text{Si})$ average molar mass of a silicon atom in the crystal is calculated using the proportions x_i of the various isotopes ^iSi , V is Volume of Silicon Sphere, a Lattice parameter of the silicon crystal, 8 is the number of atoms in an elementary cell of the lattice(cube with edge length a). M Molar mass of silicon contained in sphere. m mass of sphere.

$$N = \frac{8V}{a^3} = \text{Number of atoms in silicon sphere}$$

$$N_A = \frac{M}{m} \frac{8V}{a^3} = \text{Avogadro constant}$$

$$m(\text{Si}) = \frac{m}{N} = \frac{m a^3}{8V} = m(e) \frac{A_r(\text{Si})}{A_r(e)}$$

$$m(e) = \frac{2 h R_\infty}{c \alpha^2} = \frac{2 (2\pi \hbar) R_\infty}{c \alpha^2}$$

$$h = \frac{c \alpha^2 m a^3}{2 R_\infty 8 V} \frac{A_r(e)}{\sum_{i=28}^{30} x_i A_r(^i\text{Si})}$$

References

Horst Bettin. How the avogadro constant was measured for the last time. URL <https://q-more.chemeurope.com/q-more-articles/287/how-the-avogadro-constant-was-measured-for-the-last-time.html>.

Michael dePodesta, Robin Underwood, Gavin Sutton, Paul Morantz, Peter Harris, Darren F Mark, Finlay M Stuart, Gergely Vargha, and Graham Machin. A low-uncertainty measurement of the boltzmann constant. *Metrologia*, 50(4): 354–376, jul 2013. doi: 10.1088/0026-1394/50/4/354. URL <https://doi.org/10.1088/0026-1394/50/4/354>.

Richard Feynman. *QED : the strange theory of light and matter*. Princeton University Press, Princeton, N.J, 1985. ISBN 978-0-691-08388-9.

SI. *CGPM13 CGPM : Comptes rendus de la 13e reunion (1968)*. Bureau international des poids et mesures, F-92312 Sèvres Cedex, France, 1968. URL <https://www.bipm.org/documents/20126/17314988/CGPM13.pdf/ff522dd4-7c97-9b8d-127b-4fe77f3fe2bc?version=1.2&t=1587104721230&download=true#page=103>.

George Johnstone Stoney. On the physical units of nature. *The Scientific Proceedings of the Royal Dublin Society*, 3:51–60, 1883. URL <https://books.google.com/books?id=R79WAAAAIAAJ&pg=PA51>.

Eite Tiesinga, Peter J. Mohr, David B. Newell, and Barry N. Taylor. Codata recommended values of the fundamental physical constants: 2018. *Rev. Mod. Phys.*, 93:025010, Jun 2021. doi: 10.1103/RevModPhys.93.025010. URL <https://link.aps.org/doi/10.1103/RevModPhys.93.025010>.

NY Times. Science watch. *NY Time*, page 6, Nov 1983. URL <https://www.nytimes.com/1983/11/01/science/science-watch-011004.html>.

Supplementary Materials

Only reason I found this was for some reason or another I had just looked at $\sqrt{\hbar \alpha}$, and I had Avogadro constant for Stoney-Mass $M_S = \sqrt{\frac{\hbar \alpha c}{G}}$ mass popping up in a script I had running with some notes to see how the Planck-Units, and the Stoney-Units[Stoney, 1883], baked out the need for certain constants. But when I had left the speed of sound calc to see how the planck units canceled out the Boltzman constant, and the Avogadro constant for M_S also popped up, not exact match but so close I had to do more tests.