

SI Unit Posters

Version 1.1 (June 12, 2020)

Created by Glenwing in June of 2020

<https://github.com/Glenwing/SI-Unit-Posters>



This work is licensed under a Creative Commons Attribution-ShareAlike 4.0 International license. It may be shared and distributed freely, and modified or used for derivative works, including for commercial purposes, as long as due credit is given and any derivative works are licensed under these same terms.

<http://creativecommons.org/licenses/by-sa/4.0/>

This work was compiled from L^AT_EX source code. The source code itself is licensed under the GNU General Public License (GPL). Any modifications to the source code must also be made open-source and licensed under GPL.

Some of the text in this work (particularly the description for the weber and lumen) is based on the text of *Wikipedia* articles with very little modification. The text of *Wikipedia* is licensed under the CC BY-SA license. Other portions (such as the candela description) are based on the text of the *SI Brochure 9th Edition* with very little modification. The *SI Brochure* is licensed under the CC BY license.

This work uses the following typefaces:

- *Computer Modern* (Serif) by Donald E. Knuth, licensed under the Knuth License
- *IBM Plex* (Thin) by IBM, licensed under the SIL Open Font License
- *Timeless* by Manfred Klein, a freely-distributed font
- *Aquiline Two* by Manfred Klein, a freely-distributed font, modified by Glenwing
- *Times New Roman* (Bold) by Monotype, whose proprietary license allows free usage of the font in non-commercial documents and static images

LE
SYSTÈME INTERNATIONAL
D'UNITÉS

A tous les temps, à tous les peuples

For all times, for all peoples

S

Second

The span of time that passes during 9 192 631 770
unperturbed ground state hyperfine transitions
of a caesium-133 atom

m

Metre

The distance traveled by light in vacuum
over a time interval of exactly $1/299\,792\,458$ s

kg

Kilogram

The amount of mass such that the Planck constant h
is exactly equal to $6.626\,070\,15 \times 10^{-34} \text{ J}\cdot\text{s}$ ($\text{kg}\cdot\text{m}^2\cdot\text{s}^{-1}$)

K

Kelvin

The change in temperature which results in a change
in thermal energy of exactly $1.380\,649 \times 10^{-23}$ J
and 0 K is the temperature of absolute zero

A

Ampere

The movement of electric charge at a rate of exactly
 $1/1.602\,176\,634 \times 10^{19}$ times the elementary charge e per second

cd

Candela

The luminous intensity, in a given direction, of a source
emitting monochromatic light of frequency 540×10^{12} Hz
with a radiant intensity of $1/683$ W per steradian in that direction

mol

Mole

A collection of atoms, molecules, or particles
in the amount of exactly $6.022\,140\,76 \times 10^{23}$

N

Newton

The force required to accelerate a 1 kg object
to a velocity of 1 m/s at a uniform rate in 1 second

Pa

Pascal

The pressure applied by 1 N of force acting on a 1 m² area

J

Joule

The energy required to accelerate a 1 kg object
at 1 m/s^2 through a distance of 1 m

W

Watt

The transfer of energy at a rate of 1 J/s

C

Coulomb

The electric charge of exactly $1/1.602\,176\,634 \times 10^{19}$ protons

V

Volt

The difference in electric potential between two points
in an electric field which imparts 1 J of energy
to 1 C of charge moving between the two points

Ω

Ohm

The electrical resistance between two points such that
a 1 V potential difference produces a 1 A electric current

F

Farad

The capacitance between two points such that
an electric potential difference of 1 V
results in a buildup of 1 C of stored charge

H

Henry

The inductance between two points such that
an electric current changing at a rate of 1 A/s
produces an electric potential difference of 1 V

Wb

Weber

The magnetic flux that, linking a circuit of one turn,
produces an electric potential difference of 1 V
when it is reduced to 0 at a uniform rate in 1 s

T

Tesla

The flux density of a magnetic field that applies 1 N of force
to a 1 C charge moving through the field at 1 m/s

lm

Lumen

The concentration of visible light
passing through a solid angle of 1 steradian
emitted from a source with a luminous intensity of 1 cd

lx

Lux

The concentration of visible light on a surface equal to
a luminous flux of 1 lumen spread across an area of 1 m²