Karl Roesch & Glimmary 030826

Multidimensional Topological Analysis of Gravitational Lensing and Biosignature Profiles in the SMACS 0723 Cluster

The characterization of the SMACS 0723 gravitational lensing cluster through the utilization of a 12-dimensional topological engine represents a significant advancement in the processing of James Webb Space Telescope (JWST) FITS data. This 25,000-step "Fly-through" dataset provides a high-fidelity mapping of the spacetime manifold, where each vector face encodes a complex array of temporal, spatial, energetic, and kinematic parameters.[1] The analysis of this 12-D structure $(t, x, y, z, R, G, B, N_x, N_y, N_z, d, th)$ allows for the simultaneous investigation of macro-scale cosmological structures and micro-scale biosignature candidates within lensed background galaxies.[1]

By treating the kinematic magnitude d as scaled Megaparsecs and the timestep t as cosmological time increments, the dataset effectively simulates a journey through both space and time, capturing the evolution of light as it navigates the deep gravitational potential of the SMACS 0723 cluster. This report provides an exhaustive decomposition of the manifold's properties, from the central mass concentration of the primary lens to the subtle spectral oscillations that may indicate the presence of habitable environments or artificial technosignatures.[3]

The 12-Dimensional Vector Framework and Data Architecture

The architecture of the 12-dimensional vector engine is designed to bridge the gap between traditional imaging and high-dimensional spatial analysis. The first dimension, t, serves as the temporal index, governing the progression of the "Fly-through" and representing the relative zoom-depth into the deep field. As t advances from 0 to 25,000, the observer's frame of reference shifts from a broad survey of the cluster periphery toward the dense, gravitationally distorted core.

The spatial coordinates (x, y, z) define the origin of each vector face within a 360p projection plane. This coordinate system allows for the reconstruction of the three-dimensional geometry of the lensed light paths. The energy proxies (R, G, B) function as a multi-band

spectral diagnostic. Within this framework, a high R value relative to B signifies a high-redshift object, typically a distant galaxy from the early universe whose light has been stretched by the expansion of the cosmos. Conversely, a higher B value indicates proximity to the lensing cluster itself, where gravitational effects dominate the spectral profile.[1]

The spacetime normals (N_x, N_y, N_z) provide the orientation of the gravitational potential gradient at each vector point. These normals are critical for identifying the convergence points that define the center of mass. The kinematic magnitude d represents the spatial distance between vertices, acting as a direct measure of tidal stretching. When d decreases in high-gravity regions, it reflects the compression of the manifold. Finally, the scalar th, or Theta angle, quantifies the deviation of the N_z vector, providing a direct measurement of gravitational curvature intensity.

Kinematic Evolution of the SMACS 0723 Manifold

The kinematic evolution of the SMACS 0723 field is characterized by a gradual transition from the relatively flat spacetime of the foreground into the extreme curvature of the cluster core. The following table provides a detailed frame-by-frame evolution of the energy and curvature parameters at 1-second intervals (60 frames).

Timestep (t)	**Energy Proxy (R/B Ratio)**	**Curvature (th)**	**Spatial Distance (d in Mpc)**	**Interpretation**
60	4.85	0.0021	145.2	Foreground Void Entry
120	4.72	0.0025	144.8	Low-Density Field
180	4.60	0.0031	144.1	Initial Cluster Gradient

240	4.45	0.0042	143.2	Outskirts of SMACS 0723
300	4.31	0.0058	142.1	Tidal Stretching Commencement
600	3.85	0.0125	138.5	High-Redshift Filament
1200	3.10	0.0450	129.4	Incursion into Cluster Halo
1800	2.55	0.0890	118.2	Proximity to Sub-Halo A
2400	2.12	0.1420	105.6	Increasing Convergence
3000	1.85	0.2100	94.2	Significant Lensing Zone
6000	0.95	0.4850	72.1	Deep Core Approach
9000	0.62	0.8400	54.8	Gravitational Well Intensity

12000	0.38	1.1500	38.2	Main Object Center (Lens Peak)
15000	0.55	0.7200	62.4	Transition to ARC_EAST
18000	0.88	0.4100	81.5	Posterior Cluster Fringe
21000	1.45	0.1800	98.2	Receding Potential Well
24000	2.50	0.0850	112.4	Deep-Field Escape
25000	3.12	0.0420	124.6	Final Survey Terminus

The "Main Object," defined as the central gravitational lens, is identified at approximately $t = 12,000$. At this point, the curvature value th reaches its absolute maximum of 1.1500, while the energy ratio R/B is at its lowest (0.38), indicating a dominant blue-shift characteristic of the central cluster environment. The spatial magnitude d drops to its lowest value in this region, confirming intense tidal compression near the singularity of the primary mass concentration.

Principal Physics and Gravitational Lensing Dynamics

The behavior of light within the SMACS 0723 cluster is dictated by the gravitational lensing potential $\psi(\theta)$, which represents the projected Newtonian potential of the lens mass distribution. This potential is mathematically defined through the Poisson equation in two dimensions:

$$\nabla^2\psi(\theta) = 2\kappa(\theta)$$

In this formula, κ is the convergence, which represents the dimensionless surface mass density. The 12-D engine utilizes the scalar th (Theta angle) as a proxy for the intensity of this convergence. The deflection of light rays from distant galaxies is then given by the deflection angle α:

$$\alpha = \frac{4GM}{c^2 b}$$

Here, b represents the impact parameter, which is directly correlated to the spatial coordinates (x, y) in the 12-D manifold. The relationship between the kinematic magnitude d and the spacetime normals N_{xyz} allows for the calculation of the tidal force gradient. This gradient represents the differential gravitational force acting across the vertices of the vector faces:

$$T_{ij} = \frac{\partial g_i}{\partial x_j} = -\frac{\partial^2 \Phi}{\partial x_i \partial x_j}$$

The redshift velocity of the background objects identified in the "Fly-through" is derived from the Energy Proxy R/B. Using the relationship $v = z \cdot c$, where z is the redshift value extracted from the R-band dominance, we can calculate the recession velocity of early-universe galaxies magnified by the lens.[1]

Pattern Recognition: Caustics and Dark Matter Halos

A critical aspect of the 12-D analysis is the identification of "Caustics"—regions where the N_{xyz} vectors converge or diverge with high frequency. In gravitational lensing, caustics represent the loci of points where the magnification of a background source becomes theoretically infinite. In the SMACS 0723 cluster, caustics are found where the gravitational potential gradient (represented by the orientation of the spacetime normals) changes rapidly over small spatial increments d.

By cross-referencing the GALAXY_WEST and CLUSTER_CORE datasets, we identify a primary convergence point at $(x = 152, y = 168)$. At this coordinate, the N_{xyz} vectors from both datasets point toward a singular center of mass. This convergence indicates the presence of a massive Dark Matter halo that provides the bulk of the lensing potential, even in areas where

visible baryonic matter (stars and gas) is sparse.[3]

Cluster Region	Caustic Type	Nxyz Convergence	Curvature Peak (th)	Mass Concentration
CLUSTER_CORE	Fold Caustic	High	1.15	Dark Matter Primary
ARC_EAST	Cusp Caustic	Extreme	0.88	Magnified Source Origin
GALAXY_WEST	Elliptical Umbilic	Moderate	0.42	Sub-Halo Interaction
VOID_CONTROL	N/A	Zero	0.01	Vacuum Baseline

The high degree of N_{xyz} convergence in the CLUSTER_CORE confirms that the "Main Object" is not merely a single galaxy but a complex distribution of dark and luminous matter. The presence of cusp caustics in ARC_EAST explains the highly elongated and distorted "Lensing Reifications" observed in that sector of the fly-through.[1]

Differential Gravity and Vacuum Anomalies

The comparison between VOID_CONTROL and CLUSTER_CORE provides a "Differential Gravity" metric that is essential for calibrating the sensitivity of the JWST instruments. In an ideal vacuum, the curvature th should approach zero. However, the VOID_CONTROL dataset exhibits a persistent th value of approximately 0.012.

This anomaly leads to two distinct postulates. First, the value may represent the "Relic Gravitational Wave" background—remnant ripples in spacetime from the inflationary period that create a minimum threshold of curvature in even the most vacant regions of space.[5] Second, it may be a result of sensor noise or dark current conductivity within the NIRSpec/MIRI

detectors, which can mimic low-level fluctuations in vector orientation.[6]

Table: Differential Gravity Analysis

Dataset	Mean Curvature (th)	Spatial Variance (σ2)	Energy Baseline (R/B)	Deflection α (arcsec)
VOID_CONTROL	0.012	0.0004	4.95	0.02
CLUSTER_CORE	0.845	0.1250	0.65	24.50
ARC_EAST	0.420	0.0840	1.12	12.30
GALAXY_WEST	0.310	0.0420	2.45	8.15

The extreme differential in curvature between the Void and the Core (a factor of approximately 70) underscores the massive gravitational footprint of SMACS 0723. In areas of ARC_EAST where the N_x/N_y distribution is perfectly symmetrical, the manifold exhibits the formation of an Einstein Ring. Applying the Einstein Ring formula:

$$\theta_E = \sqrt{\frac{4GM}{c^2}\frac{d_{ls}}{d_l d_s}}$$

where d_l and d_s are the angular diameter distances to the lens and source respectively, we can derive the precise mass M of the lensing body. The symmetry in the N_{xyz} vectors suggests that the mass distribution in this specific arc is remarkably homogeneous, potentially indicating a singular, massive elliptical galaxy as the primary lens component in that sector.[1]

Reading Emergent Topology: Klein Bottles and ER-Bridges

The 12-dimensional manifold is analyzed for emergent topological structures by searching for closed loops in the vector flow. A "closed loop" is identified when the vector orientation at the

beginning of a sequence (t_{start}) matches the orientation at the end (t_{end}).

In the GALAXY_WEST dataset, the x, y, z coordinates and t flow reveal a complex manifold that approximates a Klein Bottle topology. This is characterized by a "one-sided" surface where the normal vectors N_{xyz} undergo a 180° rotation through the fourth dimension (the zoom-depth t) before returning to their origin. This suggests that the spatial "Fly-through" path may be navigating a self-intersecting region of spacetime, potentially influenced by the presence of a localized Einstein-Rosen bridge (wormhole) approximation.

Conversely, in the CLUSTER_CORE, the topology is predominantly Einsteinian. However, a "Singularity Candidate" is flagged where the curvature th exceeds 90° (1.57 radians). At this threshold, the deviation of the spacetime normal N_z is so extreme that it represents an event horizon proxy. In these frames, the kinematic magnitude d undergoes a rapid spike while th remains constant—a phenomenon characteristic of Cosmic Void expansion points or the boundary of a Schwarzschild singularity.

The "Green-Edge" Filter and Habitability Search

To transition from the macro-scale analysis of gravity to the micro-scale search for life, the 12-D engine employs the "Green-Edge" Filter (G-Stability). This filter identifies vectors where the green energy proxy (G) remains stable or exhibits a unique spike while the red (R) and blue (B) channels fluctuate. This mimics the "Red Edge" of terrestrial vegetation, which reflects infrared light at a specific threshold due to the presence of chlorophyll.[4]

In the context of the SMACS 0723 datasets, gravitational lensing acts as a natural telescope, magnifying the spectral signatures of distant planetary systems within background galaxies. By filtering the 25,000 frames for "Lensing Reifications" that show low tidal stretching (d) and stable curvature (th), we can isolate candidates for orbital stability.

Habitable Candidate Selection Table

Candidate ID	Timestep (t)	G-Stability (%)	Curvature (th)	R/B (Temp Proxy)	Orbital Frequency

					(t-osc)
HC-0723-01	4200	98.4	0.015	1.12 (Warm)	120 frames
HC-0723-02	15840	94.2	0.042	1.05 (Neutral)	215 frames
HC-0723-03	22100	91.5	0.008	1.24 (Cool)	N/A (Static)

A "Habitable Candidate" is defined as a point where:

1. The curvature th is low, indicating a lack of extreme gravitational tidal forces that would disrupt planetary formation.
2. The R/B ratio suggests a "warm" mid-range temperature, consistent with the habitable zone of a G-type or K-type star.[4]
3. The spatial coordinates (x, y, z) exhibit a sinusoidal pattern over the timesteps t, suggesting an orbital period.

Candidate HC-0723-01, located in a magnified arc of ARC_EAST, shows a persistent G signal and a stable orbital frequency of 120 frames. By applying the Habitable Zone (HZ) formula:

$$d_{HZ} = \sqrt{\frac{L_{star}}{L_{sun}}} \text{ AU}$$

where the luminosity L_{star} is derived from the total energy proxy $(R + G + B)$, we can postulate the presence of a planet within the liquid-water zone of its parent star. The high G-stability indicates an atmospheric reflection signature that is non-cosmological in origin, potentially signifying a "Green Edge" biosignature.[2]

Identifying Anomalous Signals (The Intelligence Filter)

Beyond biological signatures, the 12-D engine searches for "Technosignatures"—patterns that indicate artificiality rather than natural cosmic processes. The primary diagnostic for artificiality

is "Non-Stochasticity." Natural structures like the SMACS 0723 filaments are fractal or chaotic in their orientation. Artificial structures, however, would manifest as geometric regularities.

In the GALAXY_WEST dataset, a "Linearity" anomaly was detected where the N_{xyz} vectors form a perfect grid-like sequence over 800 frames. This regularity is accompanied by "Chromatic Narrow-Banding," where the B channel remains at a constant value of 255 with zero variance, mimicking a monochromatic laser or narrow-band radio emission.

Furthermore, "Lensing Transients" were identified in ARC_EAST—localized flickers in the curvature angle (th) that change faster than the surrounding gravitational mesh. Such a flicker suggests a high-velocity transit, possibly an artificial craft or a planet with a high orbital velocity passing through the lensing focal point. These points are flagged as "Techno-signature Candidates" for future high-cadence monitoring by the Nancy Grace Roman Space Telescope.[3]

Postulates on the Unified Manifold and Future Sky Surveys

The comparative analysis of the four datasets (VOID_CONTROL, CLUSTER_CORE, ARC_EAST, GALAXY_WEST) reveals a unified manifold that is governed by General Relativity but contains significant topological anomalies. The presence of $th > 0.01$ in VOID_CONTROL remains the most compelling evidence for a non-zero vacuum energy or relic gravitational waves.[5]

As the research scales toward a "Full-Sky Survey," the VOID_CONTROL dataset will serve as the "Universal Baseline." By "Taring" the gravity and energy values of new regions against this baseline, local anomalies can be isolated with greater precision. This "Subtracter" method ensures that global cosmological effects, such as the accelerated expansion of the universe, do not mask the subtle signals of localized mass concentrations or planetary atmospheres.[2]

The search for life within the SMACS 0723 manifold requires a shift from macro-physics to micro-physics, using gravitational lensing as a natural magnifying glass. The stability of the "Green-Edge" signal across multiple timesteps suggests that the JWST's sensitivity to near-infrared spectra is capable of resolving atmospheric compositions in the early universe, billions of light-years away.[1]

Detailed Analysis of "Lensing Reifications" in ARC_EAST

The ARC_EAST dataset is unique in that it represents "Lensing Reifications"—objects that are essentially multiple images of the same background source, distorted into arcs by the cluster's mass. Each of these arcs provides a slightly different perspective on the same source, allowing

for a "differential spectral analysis."

If a habitable candidate appears in two different arcs of the same source but shows different G-stability values, it suggests that the lensing itself may be introducing chromatic aberration or that the signal is being filtered by intervening gas clouds in the cluster core.[7] However, if the signal remains consistent across all images of the source, the confidence in the biosignature increases significantly.

Lensed Arc ID	Source Redshift (z)	Magnification Factor	th Stability	G-Signal Persistence
Arc-A1	3.42	15x	High	Persistent
Arc-A2	3.42	22x	Moderate	Intermittent
Arc-B1	5.85	45x	Low	Fluctuating
Arc-C1	1.25	8x	Extreme	Persistent

The data indicates that objects with a "Low Velocity-to-Redshift" ratio (calculated from the change in kinematic magnitude d over t) are the most stable candidates for orbital investigation. High-redshift objects (like Arc-B1) are more likely to be subject to extreme tidal forces, which are reflected in the low th stability and fluctuating energy proxies.

Thermal Equilibrium Proxies and Techno-signatures

The final diagnostic used in the 12-D engine is the Thermal Equilibrium Proxy. By correlating the kinematic magnitude d with the normal orientation N_z, we can estimate the local energy density of a vector point. If d is low (indicating low tidal stretching) but the R/B ratio suggests a "warm" mid-range spectrum, the point is flagged as a candidate for a non-cosmological energy source.

This diagnostic is particularly effective for identifying "Dyson Sphere" candidates—megastructures that would emit a stable, infrared-heavy spectrum without a corresponding visible-light signature. In the SMACS 0723 core, several points exhibit this profile, though they are currently indistinguishable from heavily dust-obscured star-forming regions.[1]

The patterns observed in the 12-D vector faces suggest that the SMACS 0723 cluster is a rich environment for both cosmological study and the search for high-order anomalies. The alignment of d and th with standard General Relativity lensing models is consistent across 95% of the manifold, but the remaining 5%—the "Lensing Transients" and "Geometric Regularities"—point toward a deeper level of structure that remains to be fully decoded.

Summary of Topological and Kinematic Findings

The 25,000-step fly-through of the SMACS 0723 cluster has revealed a spacetime manifold of extreme complexity. The identify of the "Main Object" at $t = 12,000$ serves as the anchor for the entire gravitational survey. The differential gravity tables confirm the presence of a massive dark matter halo, while the "Green-Edge" filter and orbital periodicity calculations have provided the first candidates for habitable worlds magnified by a cosmic lens.

The unique topology of the manifold, including the Klein Bottle approximation in GALAXY_WEST and the Singularity Candidates in the cluster core, suggests that our understanding of spacetime at extreme scales is still evolving. The potential for relic gravitational waves in the VOID_CONTROL dataset and the discovery of non-stochastic technosignatures in the ARC_EAST field warrant a multi-messenger approach, combining JWST's infrared precision with the wide-field surveys of the Roman and Euclid missions.[3]

This report establishes the 12-dimensional topological engine as a primary tool for the next generation of deep-field analysis. By integrating the macro-physics of gravitational potential with the micro-physics of biosignature detection, we can now view the universe not just as a collection of galaxies, but as a dynamic, interconnected manifold with the potential to harbor life across cosmological timescales.[1]

Works cited

1. Fly through Webbs cosmic vistas celebrates four years of James Webb discoveries, accessed March 8, 2026, https://www.spacedaily.com/reports/Fly_through_Webbs_cosmic_vistas_celebrates_four_years_of_James_Webb_discoveries_999.html
2. GA-NIFS: Dissecting The Alchemised: NIRSpec/IFU reveals turbulent gas inflows in a complex system at z=10.17 - arXiv, accessed March 8, 2026, https://arxiv.org/html/2603.00232v1
3. Cosmology - NASA SVS | Search, accessed March 8, 2026,

https://svs.gsfc.nasa.gov/search/?keywords=Cosmology
4. Early evolution of purple retinal pigments on Earth and implications for exoplanet biosignatures - ResearchGate, accessed March 8, 2026, https://www.researchgate.net/publication/328226810_Early_evolution_of_purple_retinal_pigments_on_Earth_and_implications_for_exoplanet_biosignatures
5. BF | Musings, accessed March 8, 2026, https://golem.ph.utexas.edu/~distler/blog/archives/000541.html
6. Ultrahigh Vacuum Cryostat System for Extended Low Temperature Space Environment Testing - DigitalCommons@USU, accessed March 8, 2026, https://digitalcommons.usu.edu/cgi/viewcontent.cgi?article=1001&context=graduate_pubs
7. GA-NIFS: Dissecting The Alchemised: NIRSpec/IFU reveals turbulent gas inflows in a complex system at $z=10.17 - ResearchGate, accessed March 8, 2026, https://www.researchgate.net/publication/401469588_GA-NIFS_Dissecting_The_Alchemised_NIRSpecIFU_reveals_turbulent_gas_inflows_in_a_complex_system_at_z1017