

Highlights

Short Paper

A Short Subtitle

Alice Anonymous, Bob Security, Cat Memes, Derek Zoolander

- Highlight 1
- Highlight 2
- Highlight 3

Short Paper

A Short Subtitle

Alice Anonymous^{a,1,*}, Bob Security^{b,2}, Cat Memes^{b,3}, Derek Zoolander

^a*Some Institute of Technology, Department Name, Street Address, City, Postal Code*

^b*Another University, Department Name, Street Address, City, Postal Code*

Abstract

This is the abstract. Lorem ipsum dolor sit amet, consectetur adipiscing elit. Vestibulum augue turpis, dictum non malesuada a, volutpat eget velit. Nam placerat turpis purus, eu tristique ex tincidunt et. Mauris sed augue eget turpis ultrices tincidunt. Sed et mi in leo porta egestas. Aliquam non laoreet velit. Nunc quis ex vitae eros aliquet auctor nec ac libero. Duis laoreet sapien eu mi luctus, in bibendum leo molestie. Sed hendrerit diam diam, ac dapibus nisl volutpat vitae. Aliquam bibendum varius libero, eu efficitur justo rutrum at. Sed at tempus elit.

Keywords: keyword1, keyword2

1. Introduction

Built in 1964, Malawi's year of independence, Queen Elizabeth Hospital (QECH), or Queen's, as it's known locally, is the country's largest public hospital. Designed in the Tropical Modernism style of architecture popular within Africa in the late colonial period, Queen's open-plan form sprawls across a broad swath of central Blantyre, with dozens of wards, specialised facilities, and administrative buildings, linked together by a bewildering maze of covered walkways. A hive of around-the-clock activity, Queen's bustles from dawn to dusk with a constant stream of patients, drawn from across the country, and maintains a city-like feel even after dark as staff and stay-in family members cook and care for their in-patient dependants. Yet, for the first-time visitor, it's not the architecture or the crowds that are the most striking when entering the grounds, but the smell. Immediately noticeable, even from the road outside the hospital, Queen's is

*Corresponding author

Email addresses: `alice@example.com` (Alice Anonymous), `bob@example.com` (Bob Security), `cat@example.com` (Cat Memes), `derek@example.com` (Derek Zoolander)

¹This is the first author footnote.

²Another author footnote, this is a very long footnote and it should be a really long footnote. But this footnote is not yet sufficiently long enough to make two lines of footnote text.

³Yet another author footnote.


smothered by an acrid-smelling, white smoke, which hangs over the grounds day and night: the product of several constantly burning fires spread across the campus. Breathing this air, when a significant amount of burning is occurring, primarily at night when the smoke is less visible, can be incredibly uncomfortable. Moreover, the tropical architecture, designed to let air and light circulate, means that even indoors, within patient wards, surgery, and treatment areas, the air quality can also be unbearable. Poor air quality associated with burning, through released particulate matter (PM), has been linked to multiple negative health outcomes like asthma, heart disease, heart failure, stroke, and cancer among others ([Anderson et al., 2012](#); [Bell and HEI Health Review Committee, 2012](#)). In a space of healing and recovery, why is such burning occurring, and how can it continue to persist?

Queen's, simply, produces an immense amount of waste that it is unable to efficiently manage. The dozens of wards, offices, and kitchens, serving the hundreds of patients and staff, generates considerable infectious medical waste, mixed domestic and office waste, and kitchen waste, which needs to be removed and disposed of daily. Some individual wards implement waste separation, binning medical wastes separately from general waste, however separation ends at the point of collection. Queen's is not served by regular municipal waste collection. What municipal collection that is done is ad-hoc, sporadic, and based on the hospital's available financial resources. Rather, most hospital waste (infectious and non-infectious) is gathered by grounds staff and burnt at the hospital's incinerator, located at a central point within the hospital campus ([Kalina and Tilley, 2020](#)). However, the vast majority of the deposited waste is not incinerated, as a new incinerator provided by the Ministry of Health in late 2019, is only able to handle a small percentage of the hospital's waste. Furthermore, the incinerator is frequently inoperable, under repair, or without power,. Rather, the waste is openly burned, in a constantly smouldering pile to the side of the incinerator building, sending up the aforementioned clouds of smoke, which choke the grounds (Figure 1). Innumerable other, smaller, fires, scattered across the QECH campus (Figure 2) contribute their part, as grounds and maintenance staff habitually burn leaves and other garden refuse. Furthermore, caregivers who reside separately on hospital grounds, cook for themselves throughout the day and night using biomass (charcoal or wood), and burn their own domestic waste. As a result, air quality at the hospital is a constant source of discussion, with patients, staff, and caregivers struggling to cope, when just breathing is, as the quote used in the title of this article, 'unbearable.'

Challenges with solid waste management (SWM) are not unique to Queen's, and remain persistent globally, particularly within African nations, and the Global South more broadly ([Bundhoo, 2018](#); [Kubanza et al., 2022](#); [Muheirwe et al., 2022](#); [Srivastava et al., 2015](#)). Explanations for these disparities have centred on the on waste generation outstripping waste management infrastructure ([Muheirwe et al., 2022](#)), and the high costs of waste management systems, which often prove unaffordable for many low-and-middle-income countries (LMICs) ([Guerrero et al., 2013](#); [Kubanza et al., 2022](#)). Waste collection is an important step

within SWM systems, and is a common barrier within LMIC contexts (Bundhoo, 2018). Waste inequalities between high-income countries (HICs) and low-income countries (LICs) is often most pernicious and visible at the point of collection, with some HICs achieving 100% collection rates, while some African nations have rates well below 50% (Shi et al., 2021). Inequalities persist beyond collection, however. Even once waste is collected, municipalities in LMICs may burn or dump it in non-sanitary landfills due to the absence of further treatment or disposal options (Cook and Velis, 2021). According to Kaza et al. (2018), 90% of waste in LIC is disposed of in unregulated dumps or openly burned, while the quantity of waste generation in these countries is expected to triple by 2050. As a result, open waste burning, in nations like Malawi, can be expected to become more prevalent.

```
# Figure 1: Open burning at QECH, with the old incinerator building in the background (Authors).  
knitr::include_graphics(here::here("manuscript/img/image-replace-tree.png"))
```



Although hospital waste is among the common types of solid waste openly burned in LICs (in addition to municipal solid waste, sewage sludge, market or commercial waste, agricultural residues), the open burning of hospital waste has not garnered as much discussion as the burning of other waste fractions (Cook and Velis, 2021). The World Health Organisation (WHO) defines hospital, or health-care waste, as waste generated by health-care activities, of which approximately 85% is general, non-hazardous waste, but the remaining 15% can be considered hazardous, and may be toxic, infectious, or radioactive. These infectious fractions are commonly incinerated in both high and middle-income countries, however in the absence of capacity, or costly incineration or autoclave infrastructure, open burning is often a last resort option for disposal in LIC contexts (World Health Organization, 2018). The open burning of biomedical waste may reduce infection risk from potentially harmful pathogens, however, such waste may contain sharps, radioactive waste, mercury-containing instruments, and plastics, the open burning of which may result in the emission of dioxins, furans, and toxic particulate matter (PM) (World Health Organization, 2018; Cook and Velis, 2021).

Poor air quality in African cities can be attributed to multiple factors, including vehicular emissions, dust, or the use of low-quality fuels for cooking, lighting, and heating, however, the open burning of waste has been identified as one of the key contributors to air pollution within urban areas (Amegah, 2018; Bulto, 2020). Although open trashing burning is a problem in LICs broadly, for instance, Wiedinmyer et al. (2014) estimate that open trash burning contributes to 29% of the total global PM_{2.5} anthropogenic emissions, it is of particular concern in Sub-Saharan African (SSA) cities, which are home to 19 of the world's 50 biggest dumpsites (United Nations Development Programme, 2021). Particulate matter, or PM, is not specific to trash burning, but rather the product of any type of incomplete combustion and is one of the most commonly measured indicators for air quality. PM is measured in terms of its particle size and generally classified into PM with particle sizes less than 10 μ m (PM₁₀) and those with sizes less than 2.5 μ m (PM_{2.5}), while one way of measuring the severity of air pollution is by expressing the mass of PM per volume of air (μ g PM_x/m³ air). PM has been linked to multiple negative health outcomes. Amongst particulate air pollution, PM₁₀ and PM_{2.5} are of particular concern as they can penetrate deep into the lungs, and their exposure has been associated with asthma, heart disease, heart failure, stroke, and cancer (Mohapatra and Biswal, 2014; Xing et al., 2016). As a result, the health impacts of ambient air particle pollution may be significant, especially for the most vulnerable. For instance, the IHME (2019) estimates PM related air pollution resulted in 4.14 million deaths worldwide (in 2019), of which, 394,000 occurred in Africa. Yet, despite its importance, the air quality and health picture in Africa is worryingly incomplete. For instance, the WHO only collects data from 10 African countries (covering 39 cities) (World Health Organization, 2016). Likewise, robust, epidemiological data from and for the African context is also lacking, though emerging data suggests strong correlations between PM and poor health outcomes (Coker and Kizito, 2018; Heft-Neal et al., 2018). Moving

forward, it is not just the air at Queen’s which lacks clarity.

Utilising a mixed-methods approach, including a network of custom-designed air quality monitors and qualitative fieldwork with hospital staff, patients, and caregivers, the purpose of this investigation was to gain a multi-dimensional understanding of the impacts of the open burning of the various solid and organic waste fractions at QECH. Specifically, this work aimed to intensively and longitudinally measure the air quality, in particulate matter (PM), at multiple locations at and surrounding the central burning point. Furthermore, we also aimed to qualitatively understand how affected individuals perceive air quality at QECH and understand potential health impacts.

This research responds to a several specific gaps within the body of academic air quality literature: Although there has been ample discussion of air quality challenges in African cities ([Mbow-Diokhane, 2019](#); [Mir Alvarez et al., 2020](#); [Petkova et al., 2013](#); [Singh et al., 2021](#)), there is a paucity of data on air quality impacts linked to trash burning. In particular, within African cities, there has been a total dearth of scholarship on the burning of medical waste, or on air quality within hospital contexts. Furthermore, while the voices of the citizens of Northern cities ([Deguen et al., 2012](#); [Howel et al., 2002](#); [Reames and Bravo, 2019](#)) have been well documented on localised air quality issues, the voices of African urban dwellers have not been given the same consideration, and there remains a pressing need to understand how these populations experience the impacts of the open burning of trash within their communities and understand potential risks.

Findings suggest that particulate matter concentrations are routinely above the WHO guidelines and are especially worrisome at several locations. Over the course of the two-month study, the hazardous limits for both PM10 and PM2.5 were exceeded at all locations except for the Lions Sight hospital. The limits were exceeded fewer than 50 times at five of the locations, but the monitors at the Lighthouse Clinic and the Guardian Shelter recorded more than 50 instances above the hazardous limits. The extremely hazardous air quality at the Guardian shelter is mostly from cooking over open fires, while the PM at the Lighthouse Clinic is a function of being located directly adjacent to the smouldering waste pile.

Interviews show that patients, staff, and caregivers alike, are keenly aware that the air quality at Queen’s is poor, with most respondents reporting frequent respiratory-related illness. Many also linked the smoke to potential long-term health complications and expressed a belief that the pollution could be contributing to other, more potentially life-threatening diseases, such as asthma, cancer, and tuberculosis. Moreover, the tropical design of hospital buildings have rendered most coping mechanisms ineffective, with most staff only finding relief at home: relief that is not available to the hundreds of patients and caregivers who sleep at QECH or are unable to leave.

In many contexts, waste burning is a necessary, but sporadic event with limited exposure. However, in the context of a large hospital, serving the sickest and poorest, this study shows that the consistent, and highly toxic smoke produced is an unrelenting and unnecessary burden that must be addressed, lest the patients leave in worse condition than in which they arrived.

2. Material and methods

This study utilised a mixed-methods approach to both quantitatively measure the air quality at eight locations around QECH, and to qualitatively investigate the perceived impacts amongst staff and caregivers. Although the work was conducted over one sustained period in late 2019, it must be contextualised within five years' experience of research and activism within QECH by the authors. All relevant permissions were obtained from QECH beforehand, through consultations with administration and staff, and the research was approved by the National Committee On Research in The Social Sciences And Humanities in Malawi, Protocol NO. P.03/19/356.

2.1. Study site

The sensors were located within the QECH campus in consultation with hospital management. The goal was to locate sensors across the campus with a range of distances and directions from the incinerator, though the final decision was based on accessibility, the permission of the unit's head, and convenience. Each of the locations is briefly described below and shown in Figure 2:

- **Lions Sight First Eye Hospital**, often shortened to **Lions Sight**, is the largest eye hospital in Malawi and serves as the main teaching eye hospital for the Kamuzu University of Health Sciences (formerly College of Medicine). It is staffed by 5 consultants and a team of clinical officers and nurses. It provides a mix of public (no cost) and private (at cost) services.
- **The Blantyre Malaria Project (BMP)**, established by Professors Terrie Taylor and Malcolm Molyneux, has carried out clinical research and patient care in the area of paediatric malaria, specifically cerebral malaria, for more than 30 years. Its research infrastructure includes an administrative team, an inpatient research ward, an MRI centre, and an outpatient research clinic in Ndirande township and in other districts outside of Blantyre.
- **Ward 6B** is a male ward for trauma and orthopaedic patients. It was established as part of the original hospital design. It is a 60-bed ward with a dedicated nursing station and treatment rooms.
- **The Blantyre Lighthouse Trust Clinic**, also known as **Umodzi Family Centre** or simply, the **Lighthouse Clinic**, is one of four operating across Malawi. The clinics work in collaboration with

the Ministry of Health (MOH) to provide integrated HIV testing, treatment and care for people living with HIV.

- The **Moyo Nutritional Rehabilitation and Research Unit**, known on campus as **uMoyo**, is a 57-bed nutritional rehabilitation unit (NRU) for treating infants and children with severe malnutrition and acute illnesses. It is one of 104 operational NRUs in Malawi. It also has an outpatient therapeutic feeding program (OTP) for children with malnutrition who can be treated outside of the hospital.
- **Mercy James Institute for Pediatric Surgery and Intensive Care (MJC)**, shortened on campus to **Mercy James**, was opened in 2017 by the NGO Raising Malawi, which was founded by Madonna. With 3 operating rooms and 50 beds, it is the first and only first dedicated paediatric surgery hospital in Malawi. Prior to its opening, QECH had fewer than 10 intensive care beds.
- The **Administration Building** is the operational hub of the campus, containing mostly offices and meeting rooms.
- When a patient arrives at QECH, she or he will usually arrive with a guardian: someone to cook for them, buy medicine, do their laundry, and help them bathe. The **Guardian Shelter** is a gender-separated concrete floor shelter for sleeping along with a cooking pavilion and toilet/shower block are maintained by a local NGO. Though simple, the facility is secure and offers one of the only free/accessible toilet facilities for visitors on the QECH campus.

```
# map
```

```
# Figure 2: Sensor locations within QECH
```

2.2. Waste management

As the largest public hospital in the country, Queen's produces an immense amount of waste. Hospital staff, such as doctors and nurses, primarily generate and handle medical waste (dry, wet, sharps), which is disposed of in separate waste bins labelled accordingly within the ward. Hospital visitors (caretakers and patients) generate food waste and are advised by staff to discard such waste in waste bins outside, on hospital grounds. Cleaning staff are responsible for emptying bins and maintaining the cleanliness of indoor and outdoor spaces. Collected bags are temporarily stored in the sluice room, and cleaners during the night shift are responsible for transporting the waste on wheelchairs or trolleys to the incinerator for burning by waste disposers.

Yet, according to respondents and our own observations, waste management procedures within the hospital are fraught with challenges which impact the implementation of best practices. These include the lack of

a coherent hospital-wide waste management policy, insufficient waste management training for staff, and a persistent lack of material and financial resources. For instance, although hospital administration was able to articulate a set of standard operating principles (SoPs) for waste management, below the administration, doctors and nurses were generally limited in their understanding about its contents. Furthermore, cleaners, who are responsible for day-to-day cleansing and waste disposal, yet rate at the bottom of the institutional hierarchy at Queens, were unable to access or analyse hospital policy, and generally tended to rely on “experience” and “human judgement” when handling potentially hazardous waste. Moreover, although nearly all doctors and nurses had prior education on waste management best practices, cleaners began their waste handling duties without any prior waste management experience, and although there is an orientation programme for new hires, there is no regular refresher training or capacity building regarding waste for existing staff, at any level.

Despite its national prominence, and international reputation for academic medical research, Queen’s is plagued by constant shortages of human and material resources. Yet, these shortages are not shared equally across the space, with a sharp division in cleanliness, resources, and quality of care between public wards, supported by the Malawi Ministry of Health, and the private wads with international funding, such as the aforementioned Mercy James Institute. For staff, these shortages have manifested in poor salaries, missed pay cheques, failing infrastructure and limited supplies ([Kalina and Tilley, 2020](#)). Regarding waste management, these shortages have manifested in a chronic shortage of black bags, waste containers and cleaning staff. As a result, due to a lack of bags and storage containers, there is minimal segregation of waste, and highly infectious items (bandages, gloves, syringes, and other sharps) are disposed of along with non-hazardous waste (food, newspapers, packaging). As a result, rather than facilitating different disposal pathways for hazardous and non-hazardous waste, all waste is mixed, necessitating more burning than would be necessary if non-hazardous waste was diverted, and contributing to workplace risk, with reports of accidental exposure to potentially infected waste, including accidental pokes from improperly disposed of sharps, common amongst hospital staff. Regardless, separation ends at the point of collection. The hospital is not serviced by regular municipal waste collection, and as a result, nearly all of the waste produced by the hospital, hazardous and non-hazardous, has to be disposed of on-site, either within the limited capacity of the hospital incinerator, or more commonly, through open burning. As a result, the constantly smouldering pile of waste puts forth a continuous cloud of grey smoke, which mingling with the dozens of other fires on the grounds, from cooking fires and burning garden waste, blankets Queen’s in a permanent cloud of foul smelling haze.

2.3. Measuring Air Quality

2.3.1. Hardware and Software

The PM monitoring device centres around a Raspberry Pi 3 Model B single-board computer complete with an operating system and storage space. The added advantage of storage space is that the data are not lost if the internet connection breaks down. Because the Pi can only read digital signals, we needed to include an analog-to-digital (ADC) converter. The amount of dust is measured by a Nova PM SDS011 High Precision Laser sensor. This sensor measures particles at 2.5 and 10 micrometres in diameter in mg/m³. A list of the specific hardware and software components can be found in Appendix TODO.

2.3.2. Installation

The choice of where the sensors were installed was done in collaboration with the QECH administration, but the position of the sensors at the individual buildings were decided by the research team. Four air quality sensors were installed on the outside of buildings (Guardian Shelter, Mercy James, Malaria Project, Lighthouse Clinic) while the remaining four sensors were installed inside (Ward 6B, Administration, Lions Sight, uMoyo). All of the sensors were mounted to a wall at a height of approximately two metres from the ground with the help of QECH maintenance department personnel. This height was chosen to prevent the public from tampering with the equipment when they visit the hospital. Additionally, we assured that each pipe that inhaled ambient air was freely protruding in the building (room) in order to capture the air quality.

Because connecting each unit to a power supply was not possible, each unit was equipped with an external battery (5000mAh) which powered the sensor unit for three days continuously. To avoid any down time, we changed the batteries every two days. Data were also collected from the sensor at the time of battery replacement. The sensors were capable of being connected to WIFI, a feature which enabled us to download the data into our computers wirelessly.

2.3.3. Qualitative Methods

Qualitative data collection consisted of 26 interviews with caregivers and hospital staff (including janitorial and maintenance staff, nurses, doctors, and administrators) conducted in October and November 2019, as well as an additional 31 interviews conducted with staff, patients, and caregivers in January and February 2021. Interviews were semi-structured, and included questions on hospital waste management practices, perceived best practices, and workplace health, safety, and risk. However, the semi-structured nature of the interviews allowed for the participants to raise topics of interest, and for unexpected themes to emerge. These interviews are supplemented by participatory observation recorded within detailed field notes, and supported by nearly a decade of sustained involvement and activism by the authors in the waste management practices of the hospital.

Interview respondents were chosen using a purposive or judgement sampling regimen, i.e. a subjective sampling method in which respondents are selected based on their ability to effectively contribute to the study's research objectives (Kitchin and Tate, 2013). All respondents were purposively selected based on availability, willingness to provide written, informed consent, and their individual insight into the questions posed within the study. Interviews were conducted in the local language (Chichewa), audio recorded, and transcribed into English. Participation was voluntary, and responses were recorded anonymously. Written, informed consent was obtained from each respondent prior to the interview. The study was approved by the National Committee on Research in the Social Sciences and Humanities (NCRSH) of Malawi; Protocol NO. P.03/19/356. Collected data were analysed thematically and was coded within the software programme Nvivo, which organises materials and assists with the coding process.

2.3.4. Limitations

Though extensive and useful in understanding the conditions at QECH, the data set would benefit from a comparison with background values. As trash-burning is extensive throughout the city, there were no obvious locations that could be confidently used as representative background values. And although there were several possible areas upwind and outside of town, the logistics and safety involved in accessing them to change the batteries meant that the measured values at Queens could not be compared to a stable background concentration. Knowing the wind direction and velocity would have helped in better identifying the potential sources of the measured PM. However, the necessary equipment was not easily available in Blantyre at the time, and as an exploratory study, the immediate hazards for the staff and patients at QECH are relevant regardless of the burning source.

Knowing the exact burning locations and burning times, especially at the incinerator and the Guardian Shelter, would have helped in better identifying the sources and movements of the plumes. However, given the size of the campus and its 24-hour schedule, a much larger research team would have been required to quantify all the burning. Finally, The humidity, and the potential for interference was not accounted for; the measurements were not adjusted for humidity which could affect the values, though not enough to significantly alter the key findings and need for immediate action.

3. Theory/calculation (not applicable)

4. Results and Discussion

The Malawi Bureau of Standards is the national agency responsible for setting and publishing all standards in the country. At the time of writing, the official website (www.mbsmw.org) was not available. However, published work that references Malawian standards (Kutlar Joss et al., 2017; Mapoma et al., 2014) indicate that the maximum 24-h PM₁₀ value is 25 g/m³ and the maximum annual PM_{2.5} value is 8 g/m³. There

is no daily maximum value for PM2.5. It should be noted that both of these values are at, or below, the level-4 interim WHO targets.

Therefore, for international comparisons and due to the lack of comprehensive Malawian Standards, we use the WHO recommendations as a basis of comparison for the measured values ([World Health Organization, 2021](#)). The relevant particulate matter targets are presented in Table 1.

```
# Table 1. Particulate matter targets from @worldhealthorganization2021who
knitr::include_graphics(here::here("manuscript/img/image-replace-table.png"))
```

Pollutant	Averaging time	Interim target				AQG level
		1	2	3	4	
PM _{2.5} , µg/m ³	Annual	35	25	15	10	5
	24-hour	75	50	37.5	25	15
PM ₁₀ , µg/m ³	Annual	70	50	30	20	15
	24-hour	150	100	75	50	45

4.1. Particulate Measurements

4.1.1. Peaks

Due to localised variation (air movement in the immediate area) and the density of the observations, the plotted 5-minute data obscure persistent trends. The full set of 5-minute data are plotted and presented in Appendix XA1 (TODO). However, the number of measured values that exceeded the hazardous limit for both parameters are summarised in Table 2.

```
# Table 2. Peaks for both PM2.5 and PM10 are the number of data points above the WHO interim target
```

Over the course of the two-month study, the hazardous limits for both parameters were exceeded at all locations except for the Lions Sight hospital. The limits were exceeded fewer than 50 times at five of the locations, and only the monitors at the Lighthouse Clinic and the Guardian Shelter recorded more than 50 instances above the hazardous limits. At the Lighthouse Clinic, the PM10 limit was exceeded almost twice as often as the PM2.5 limit, while at the Guardian Shelter, the reverse is true. Because the Lighthouse Clinic is within 50 m of the incinerator, the number and predominance of PM10 peaks is characteristic of incomplete combustion and dust that is typical for the area and the open burning that occurs there. The residents at

the Guardian Shelter however are mostly cooking within contained clay stoves that have been designed to improve combustion of the coal and wood that is burned within them. The fewer number of PM10 peaks is a testament to this intervention, though the frequency of PM2.5 peaks is still beyond acceptable.

4.1.2. 24-hour averages

The 24-hour averaged values (logged every 5 minutes) are presented for each location, for both PM10 and PM2.5 in Figure 3. As well as dampening local variation and peaks, the WHO air quality guidelines are also based on 24-hour averages which are the standard against which the health risks can be judged.

Overall, PM2.5 values remained below 100 g/m³ at 6 of 8 locations (Administration, Lions Sight, Malaria Project, uMoyo, Ward 6B, and Mercy James); PM10 values were consistently below 100 g/m³ at the same locations, but with several average values extending slightly above, and then infrequently.

The daily averages at both the Lighthouse Clinic and the Guardian Shelter are both consistently higher for both parameters and the two averages closely followed the same general trends. Though the peaks at the Lighthouse Clinic were higher than the Guardian Shelter, the low values were consistently lower, indicating more times of little or no burning, unlike the Guardian Shelter emissions which were relatively constant. However, further analysis of 12 hour averages (8:00-15:59 (working hours) and 16:00-7:59 (evening)) did not indicate clear differences between the time periods; stated differently, the data did not clearly point to more or less burning in the day or at night.

Figure 3. Average 24-hour PM10 and PM2.5 at 8 monitoring stations over 2 months

4.1.3. PM ratios

Given that the type of fuel and the type of burning (contained vs. open) produce very different particulate “fingerprints”, the ratio of PM10 to PM2.5 values were examined to determine if a clear difference between locations, and therefore source, could be identified. The results are presented in Figure 4.

Figure 4. PM10:PM2.5 values for each location. Each line shows the minimum and maximum ratio for

The results presented in Figure 5 illustrate both the presence of PM10 relative to PM2.5 as well as the range of values over which that ratio is observed.

Unlike the peak values presented in Table 2, the values presented in Figure 5 indicate the relative presence of larger (PM10) particles to smaller ones (PM2.5) regardless of their concentration: the ratio of two low concentrations can be the same as the ratio of two high concentrations and is therefore more indicative of source than of proximity.

In general, the calculated ratios are concentrated between 1.25 with few values exceeding 7.5. Three from each the Guardian Shelter and the Ward 6B exceeded 7.5 though the most values exceeding 5 were at the Guardian Shelter. It is interesting to note that despite the large daily variations at the Lighthouse Clinic, the parameter ratios there were fairly consistent with few spikes, and only 2 days with ratios above 5.

An examination of the range of ratios is used to better understand the variability within a single day. For example, several of the calculated values at the Guardian Shelter have a range of more than 5, and one day had a range of close to 10 i.e. the minimum ratio recorded for that day was close to 1 and the maximum value was over 10. The composition of emissions recorded at that location ranged significantly, and likely reflects a range of burning styles and/or fuel type.

4.2. Exposure Data

The relative amount of time exposed to a given category of air quality is shown in Figure 5. Each square represents 1% of the measured time.

Figure 5. Percentage of measured values categorised according to WHO 2021 targets

Unsurprisingly, a majority of the air quality at the Lighthouse Clinic and the Guardian Shelter falls at, or below, interim target 1. What is less expected is that even though there are fewer days of hazardous or very bad air quality at the other locations, most locations do not have target 4 values for even half of the time. Mercy James, the Malaria Project and Lions Sight have the largest percentages of days with air quality meeting the interim target 4. Interim target 2 rated air occupies an unexpectedly large proportion of the days at all locations, indicating that although there are a few very toxic, and a few quite good days, the majority of the air is actually neither, but still cause for concern.

4.3. “The Air We Breathe is Not Good”: Perspectives from QECH

As the previous section has described, the open incineration of waste at QECH has created hazardous conditions for those occupying the space. These risks were not lost on staff and caregivers, as interviews demonstrated broad and near universal awareness of the poor air quality within the hospital grounds. Overwhelmingly, within interviews, both caregivers and staff were quick to decry the poor quality of the air, generally without prompting. The few exceptions were those staff posted on the peripheries of the hospital grounds, at a distance from the spaces of incineration. However, even those who did not experience the impacts of the burning, were aware of it, and considered themselves fortunate to be posted in a section of the hospital where it was less of a problem. Furthermore, according to caregivers and staff who work night shifts, air quality can be particularly bad at night and in the early morning, despite the smoke being less visible, due to the habit of janitorial staff concentrating their burning during the late hours. The poor

air quality on hospital grounds is also a frequent cause for complaint by patients and visitors, with nearly every staff member interviewed being able to recall having received a complaint, and in turn, complaining to administration. One of the staff members responsible for the burning said that he personally, had received hundreds of complaints, but was powerless to affect meaningful change, aside from burning at different hours, until the incinerator could be repaired.

Nonetheless, despite this consensus that air quality caused by the burning of waste within the hospital was a problem, significant differences emerged between respondents over their understandings of potential impacts, the effectiveness of various coping mechanisms, and their problematisations linked to the burning of specific waste materials. In addition, interviews revealed that these understandings were informed by a significant amount of misinformation, even amongst trained medical staff, which may contribute to them being less able to mitigate potential risks for themselves and those who rely on their care.

4.3.1. Impacts, Problematisations, and Misinformation

The poor air quality within QECH was responsible, according to respondents, for a wide array of health impacts. The most common ones cited included: coughing and sneezing, sore throat, stinging eyes, breathing difficulties, and persistent cold and flu. Nausea was also mentioned, but was not a commonly described impact. Only one respondent, of the 26 total interviewed, did not describe lingering health impacts which they could ascribe to the smoke, however, they also described having chronic eye irritation, but did not believe the smoke was a contributing factor.

In addition to these impacts, which respondents bear on a daily basis, many also believed that the smoke could contribute to a number of more serious, long-term health complications. For instance, nearly a quarter of respondents raised concerns of the potential impact that the smoke could have on patients or staff with asthma. Others flagged poor air quality as a potential risk factor for certain cancers, lung disease, or heart problems. For a few, the smoke posed an unknown danger, they were not sure what types of impacts it could have, but they were sure it was harmful in some way.

Also, understandably, given the large tuberculosis ward present within the hospital grounds, and the high prevalence of the disease within Malawi ([World Health Organization, 2019](#)) there was significant concern (more than half of respondents) about the impact the poor air quality could have on the infected. However, there also persisted a belief among several respondents, including several nurses, that the smoke could be a cause of the disease itself. As one staff member (20/11/2019) stated, “I believe breathing this air for a long time can cause Tuberculosis.” This, however, was only one of the few instances of misinformation which staff members held regarding air quality and health. Another example, voiced by several respondents, included a belief that some staff members were immune to ill-health impacts of the smoke, because they had

received vaccinations from the hospital. Nonetheless, they were concerned about the impacts of the smoke on patients and visitors, as one staff member (18/11/2019) expressed:

Personally, I have never experienced [eye discomfort] because I get vaccinated and I am protected including other staff. However, we realize that the air can badly affect other people and patients who come to this place.

Another interesting misconception that emerged, which may be tied partly to translation and transcription, was a different cultural understanding of smoke versus smells. More than half of the respondents appeared to conflate the two, with some expressing a belief that it was the odour of what was being burnt that was harmful, rather than the smoke being given off. This has led many to specifically problematise the burning of certain wastes, such as plastics, medicines, and other medical wastes, which give off distinctive or less pleasant odours, as opposed to the burning of other items, such as garden refuse, which may produce significant smoke, and contribute to higher recorded values of particulate matter, but produce a less pungent, or more normalised, odour.

4.3.2. Coping Mechanisms

Finally, in order to manage the impacts of QECH's persistently poor air quality, staff and visitors reported having developed a number of coping mechanisms, designed to help them get through their daily routines. These included staying indoors, blocking doors and windows, and taking breaks away from hospital grounds in order to catch some breaths of fresher air. However, for janitorial staff, inside was not necessarily better, as several reported their indoor workspaces as being also insufferable for long periods of time from the smell of cleaning agents and other chemicals. Furthermore, amongst respondents there was a general disagreement over the effectiveness of personal protective equipment (PPE), such as face masks, towards mitigating the impacts of the smoke. A few staff described pleading to hospital administration for such equipment, to no avail. However, other staff members, who do have access to PPE, noted that even face masks do little to mitigate the impacts of the smoke, describing them as ineffective.

Most staff, however, have been unable to find anyway to mitigate the impacts of the smoke, and only found relief once they reached home at the end of their shift, as one of the janitorial staff described, "we only feel safe when we are home". Of course, this relief is not available for the hundreds of patients and caregivers who sleep at QECH or are unable to leave. Ultimately, most place their hope in the construction of the new incinerator (which had not yet been activated at the time of the interviews), and biding their time as construction drags on; coping as best they can. As one staff member (25/11/2019) described, "we are just hoping we will start breathing good air soon, when the new incinerator is opened."

5. Conclusions

6. Appendix

If there is more than one appendix, they should be identified as A, B, etc. Formulae and equations in appendices should be given separate numbering: Eq. (A.1), Eq. (A.2), etc.; in a subsequent appendix, Eq. (B.1) and so on. Similarly for tables and figures: Table A.1; Fig. A.1, etc.

7. Figures and tables

Figure 1 is generated using an R chunk.




Figure 1: A meaningless scatterplot

8. Tables coming from R

Tables can also be generated using R chunks, as shown in Table 1 example.

```
knitr::kable(head(mtcars)[,1:4])
```

Table 1: Caption centered above table

	mpg	cyl	disp	hp
Mazda RX4	21.0	6	160	110

	mpg	cyl	disp	hp
Mazda RX4 Wag	21.0	6	160	110
Datsun 710	22.8	4	108	93
Hornet 4 Drive	21.4	6	258	110
Hornet Sportabout	18.7	8	360	175
Valiant	18.1	6	225	105

References

- Amegah, A.K., 2018. Proliferation of low-cost sensors. What prospects for air pollution epidemiologic research in Sub-Saharan Africa? *Environmental Pollution* 241, 1132–1137. doi:[10.1016/j.envpol.2018.06.044](https://doi.org/10.1016/j.envpol.2018.06.044).
- Anderson, J.O., Thundiyil, J.G., Stolbach, A., 2012. Clearing the Air: A Review of the Effects of Particulate Matter Air Pollution on Human Health. *Journal of Medical Toxicology* 8, 166–175. doi:[10.1007/s13181-011-0203-1](https://doi.org/10.1007/s13181-011-0203-1).
- Bell, M.L., HEI Health Review Committee, 2012. Assessment of the health impacts of particulate matter characteristics. *Res Rep Health Eff Inst* , 5–38.
- Bulto, T.W., 2020. Impact of Open Burning Refuse on Air Quality: In the Case of “Hidar Sitaten” at Addis Ababa, Ethiopia. *Environmental Health Insights* 14, 117863022094320. doi:[10.1177/1178630220943204](https://doi.org/10.1177/1178630220943204).
- Bundhoo, Z.M.A., 2018. Solid waste management in least developed countries: Current status and challenges faced. *Journal of Material Cycles and Waste Management* 20, 1867–1877. doi:[10.1007/s10163-018-0728-3](https://doi.org/10.1007/s10163-018-0728-3).
- Coker, E., Kizito, S., 2018. A Narrative Review on the Human Health Effects of Ambient Air Pollution in Sub-Saharan Africa: An Urgent Need for Health Effects Studies. *International Journal of Environmental Research and Public Health* 15, 427. doi:[10.3390/ijerph15030427](https://doi.org/10.3390/ijerph15030427).
- Cook, E., Velis, C., 2021. Global Review on Safer End of Engineered Life. Technical Report. Royal Academy of Engineering. doi:[10.5518/100/58](https://doi.org/10.5518/100/58).
- Deguen, S., Ségala, C., Pédrone, G., Mesbah, M., 2012. A New Air Quality Perception Scale for Global Assessment of Air Pollution Health Effects. *Risk Analysis* 32, 2043–2054. doi:[10.1111/j.1539-6924.2012.01862.x](https://doi.org/10.1111/j.1539-6924.2012.01862.x).
- Guerrero, L.A., Maas, G., Hogland, W., 2013. Solid waste management challenges for cities in developing countries. *Waste Management* 33, 220–232. doi:[10.1016/j.wasman.2012.09.008](https://doi.org/10.1016/j.wasman.2012.09.008).
- Heft-Neal, S., Burney, J., Bendavid, E., Burke, M., 2018. Robust relationship between air quality and infant mortality in Africa. *Nature* 559, 254–258. doi:[10.1038/s41586-018-0263-3](https://doi.org/10.1038/s41586-018-0263-3).
- Howel, D., Moffatt, S., Prince, H., Bush, J., Dunn, C.E., 2002. Urban Air Quality in North-East England: Exploring the Influences on Local Views and Perceptions. *Risk Analysis* 22, 121–130. doi:[10.1111/0272-4332.t01-1-00010](https://doi.org/10.1111/0272-4332.t01-1-00010).
- IHME, 2019. Ambient Particulate Matter Pollution—Level 4 Risk. Technical Report. Institute for Health Metrics and Evaluation. Seattle, WA.
- Kalina, M., Tilley, E., 2020. “This is our next problem”: Cleaning up from the COVID-19 response. *Waste Management* 108, 202–205. doi:[10.1016/j.wasman.2020.05.006](https://doi.org/10.1016/j.wasman.2020.05.006).
- Kaza, S., Yao, L., Bhada-Tata, P., Van Woerden, F., 2018. What a Waste 2.0: A Global Snapshot of Solid Waste Management to 2050. Technical Report. World Bank Group. Washington D.C.
- Kitchin, R., Tate, N., 2013. *Conducting Research in Human Geography: Theory, Methodology and Practice*. Routledge.
- Kubanza, N.S., Matsika, R., Magha, C.G., 2022. Exploring the role of local authorities and community participation in solid

- waste management in sub-saharan Africa: A study of Alexandra, Johannesburg, South Africa. *Local Environment* 27, 197–214. doi:[10.1080/13549839.2021.2010186](https://doi.org/10.1080/13549839.2021.2010186).
- Kutlar Joss, M., Eeftens, M., Gintowt, E., Kappeler, R., Künzli, N., 2017. Time to harmonize national ambient air quality standards. *International Journal of Public Health* 62, 453–462. doi:[10.1007/s00038-017-0952-y](https://doi.org/10.1007/s00038-017-0952-y).
- Mapoma, H.W.T., Tenthani, C., Tsakama, M., Kosamu, I.B.M., 2014. Air quality assessment of carbon monoxide, nitrogen dioxide and sulfur dioxide levels in Blantyre, Malawi: A statistical approach to a stationary environmental monitoring station. *African Journal of Environmental Science and Technology* 8, 330–343. doi:[10.4314/ajest.v8i6](https://doi.org/10.4314/ajest.v8i6).
- Mbow-Diokhane, A., 2019. Air Quality in African Cities, in: Mboup, G., Oyelaran-Oyeyinka, B. (Eds.), *Smart Economy in Smart African Cities: Sustainable, Inclusive, Resilient and Prosperous*. Springer, Singapore. *Advances in 21st Century Human Settlements*, pp. 297–311. doi:[10.1007/978-981-13-3471-9_9](https://doi.org/10.1007/978-981-13-3471-9_9).
- Mir Alvarez, C., Hourcade, R., Lefebvre, B., Pilot, E., 2020. A Scoping Review on Air Quality Monitoring, Policy and Health in West African Cities. *International Journal of Environmental Research and Public Health* 17, 9151. doi:[10.3390/ijerph17239151](https://doi.org/10.3390/ijerph17239151).
- Mohapatra, D.K., Biswal, S.K., 2014. Effect of Particulate Matter (PM) on plants, climate, ecosystems, and human health , 12.
- Muheirwe, F., Kombe, W., Kihila, J.M., 2022. The paradox of solid waste management: A regulatory discourse from Sub-Saharan Africa. *Habitat International* 119, 102491. doi:[10.1016/j.habitatint.2021.102491](https://doi.org/10.1016/j.habitatint.2021.102491).
- Petkova, E.P., Jack, D.W., Volavka-Close, N.H., Kinney, P.L., 2013. Particulate matter pollution in African cities. *Air Quality, Atmosphere & Health* 6, 603–614. doi:[10.1007/s11869-013-0199-6](https://doi.org/10.1007/s11869-013-0199-6).
- Reames, T.G., Bravo, M.A., 2019. People, place and pollution: Investigating relationships between air quality perceptions, health concerns, exposure, and individual- and area-level characteristics. *Environment International* 122, 244–255. doi:[10.1016/j.envint.2018.11.013](https://doi.org/10.1016/j.envint.2018.11.013).
- Shi, Y., Wang, Y., Yue, Y., Zhao, J., Maraseni, T., Qian, G., 2021. Unbalanced status and multidimensional influences of municipal solid waste management in Africa. *Chemosphere* 281, 130884. doi:[10.1016/j.chemosphere.2021.130884](https://doi.org/10.1016/j.chemosphere.2021.130884).
- Singh, A., Ng'ang'a, D., Gatari, M.J., Kidane, A.W., Alemu, Z.A., Derrick, N., Webster, M.J., Bartington, S.E., Thomas, G.N., Avis, W., Pope, F.D., 2021. Air quality assessment in three East African cities using calibrated low-cost sensors with a focus on road-based hotspots. *Environmental Research Communications* 3, 075007. doi:[10.1088/2515-7620/ac0e0a](https://doi.org/10.1088/2515-7620/ac0e0a).
- Srivastava, V., Ismail, S.A., Singh, P., Singh, R.P., 2015. Urban solid waste management in the developing world with emphasis on India: Challenges and opportunities. *Reviews in Environmental Science and Bio/Technology* 14, 317–337. doi:[10.1007/s11157-014-9352-4](https://doi.org/10.1007/s11157-014-9352-4).
- Wiedinmyer, C., Yokelson, R.J., Gullett, B.K., 2014. Global Emissions of Trace Gases, Particulate Matter, and Hazardous Air Pollutants from Open Burning of Domestic Waste. *Environmental Science & Technology* 48, 9523–9530. doi:[10.1021/es502250z](https://doi.org/10.1021/es502250z).
- World Health Organization, 2016. *Ambient Air Pollution: A Global Assessment of Exposure and Burden of Disease*. World Health Organization.
- World Health Organization, 2018. *Health-Care Waste. Technical Report 1*. World Health Organization. Geneva, Switzerland.
- World Health Organization, 2019. *Global Tuberculosis Report 2019. Technical Report WHO/CDS/TB/2019.15*.
- World Health Organization, 2021. *WHO Global Air Quality Guidelines: Particulate Matter (PM_{2.5} and PM₁₀), Ozone, Nitrogen Dioxide, Sulfur Dioxide and Carbon Monoxide*. World Health Organization.
- Xing, Y.F., Xu, Y.H., Shi, M.H., Lian, Y.X., 2016. The impact of PM_{2.5} on the human respiratory system. *Journal of Thoracic Disease* 8, 6.