

PhD

Autor: Lukas Johannsen
Erstgutachter: Prof. Dr. Gleb Arutyunov
Zweitgutachter: ?
Ort und Datum: Hamburg im ? 2027

Abstract

Acknowledgments

⁰This thesis has benefited from large language models.

Contents

1	Notes	3
1.1	Basics of elliptic structures	3
1.1.1	Elliptic functions and theta functions	3
1.1.2	Belavin's elliptic R -matrix	4
1.1.3	RTT representations	4
1.1.4	Elliptic Drinfeld functor	5
1.1.5	Generalized Schur-Weyl duality	6
1.2	Loop Yangian	6
1.2.1	As quantization of rational spin RS Poisson algebra	6
1.2.2	Fock space representation	7
	Notation	8
	Bibliography	9

Chapter 1

Notes

1.1 Basics of elliptic structures

1.1.1 Elliptic functions and theta functions

Definition 1.1.1. An elliptic curve E (over \mathbb{C}) is a smooth projective curve or Riemann surface of genus 1. These are of the form $E \cong \mathbb{C}/\Lambda$ for $\Lambda = \mathbb{Z} \oplus \tau\mathbb{Z}$ for $\tau \in \mathbb{H} = \{z \in \mathbb{C} \mid \Im z > 0\}$. This does not faithfully parametrize elliptic curves, but $M_{1,1} := \mathbb{H}/SL_2(\mathbb{Z})$ does, where $SL_2(\mathbb{Z})$ acts by Möbius transformations. Algebraically, every elliptic curve may be brought into the form

$$Y^2Z = 4X(X - Z)(X - \lambda Z),$$

where λ is the λ -invariant of E , which is also not faithful up to an action of $SL_2(\mathbb{Z})/\Gamma(2) \cong S_3$. The invariant combination

$$j = 2^8 \frac{(\lambda^2 - \lambda + 1)^3}{\lambda^2(\lambda - 1)^2}$$

is the j -invariant, yielding a bijection $j : M_{1,1} \rightarrow \mathbb{C}$. Stacky points are at $j = 0$ and $j = 12^3 = 1728$, where Λ becomes the lattice of the Eisenstein and Gauss integers, respectively, with automorphism groups \mathbb{Z}_2 and \mathbb{Z}_3 (plus the usual involution $Y \mapsto -Y$, giving \mathbb{Z}_4 and \mathbb{Z}_6). There is an isomorphism

$$\mathbb{C}/\Lambda \rightarrow E(\mathbb{C}), \quad z + \Lambda \mapsto \begin{cases} [\wp(z|\tau) : \wp'(z|\tau) : 1], & z \notin \Lambda \\ [0 : 1 : 0], & z \in \Lambda. \end{cases}$$

An elliptic function is a meromorphic function $f : \mathbb{C} \rightarrow \mathbb{C}$ that is Λ -periodic such that it descends to a meromorphic function on E . Theta functions are sections of certain line bundles over E , which may be represented as entire functions $\vartheta : \mathbb{C} \rightarrow \mathbb{C}$ satisfying

$$\vartheta(z + 1|\tau) = \vartheta(z|\tau), \quad \vartheta(z + \tau|\tau) = \exp(-\pi i \tau - 2\pi i z) \vartheta(z|\tau)$$

This line bundle has in fact only one section up to a prefactor, and this is the Jacobi theta function.

1.1.2 Belavin's elliptic R -matrix

[ES98]

Definition 1.1.2. Let $\xi = \exp(2\pi i/\ell)$. Define a projectively flat rank ℓ vector bundle on $E = \mathbb{C}/\Lambda$ by the two monodromies

$$A = \begin{pmatrix} 1 & 0 & \cdots & 0 \\ 0 & \xi & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \xi^{\ell-1} \end{pmatrix}, \quad B = \exp(-\pi i \frac{\ell-1}{\ell}) \begin{pmatrix} 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 1 \\ 1 & 0 & \cdots & 0 \end{pmatrix},$$

satisfying $A^\ell, B^\ell = 1$ and $BA = \xi AB$ giving a flat/holomorphic PGL_ℓ -bundle $P \rightarrow E$. Belavin's elliptic R -matrix is the unique R -matrix satisfying the QYBE and unitarity as well as

- (i) $R^B(z)$ has simple poles only at $\eta + \Lambda$,
- (ii) $R^B(0) = P$,
- (iii) $R^B(z+1) = A_1 R^B(z) A_1^{-1} = A_2^{-1} R^B(z) A_2$, $R^B(z+\tau) = B_1 R^B(z) B_1^{-1} = B_2^{-1} R^B(z) B_2$.

In particular, the R -matrix lives on the ℓ -fold cover $\bar{E} = \mathbb{C}/\ell\Lambda$, where P is trivialized. We may view the R -matrix as an element of

$$\text{End } \mathbb{C}^\ell \otimes \Gamma(E, P \ltimes \text{End } \mathbb{C}^\ell),$$

where $\Gamma(E, P \ltimes \text{End } \mathbb{C}^\ell)$ are meromorphic sections of the associated bundle $P \ltimes \text{End } \mathbb{C}^\ell$.

1.1.3 RTT representations

[ES98]

Definition 1.1.3. Define the category \mathcal{C}_B whose objects are vector spaces V equipped with an invertible element

$$L(z) \in \text{Mat}_\ell(\text{End } V \otimes \mathcal{M}(\bar{E})),$$

thus having values in $\text{End } \mathbb{C}^\ell \otimes \text{End } V$, satisfying

$$R_{12}^B(z-w) L_1(z) L_2(w) = L_2(w) L_1(z) R_{12}^B(z-w).$$

Morphisms are linear maps $f : V \rightarrow V'$ satisfying $\varphi L(z) = L'(z) \varphi$. There is a tensor structure via

$$(V, L(z)) \otimes (V', L'(z)) := (V \otimes V', L_{12}(z) L'_{13}(z))$$

and finite-dimensional objects $(V, L(z))$ have duals $(V^*, L^*(z))$ with $L^*(z) = (L(z)^{-1})^{t_2}$. Belavin's R -matrix ensures the existence of a vector representation $(\mathbb{C}^\ell, R^B(z))$.

1.1.4 Elliptic Drinfeld functor

In order to define an elliptic Drinfeld functor, we first need an analog of the Yangian representation on $\mathbb{C}^\ell[y]$. For fixed z , we can view the coefficients of $R^B(z-y)$ as meromorphic $\text{End } \mathbb{C}^\ell$ -valued functions $f_{ij}(y)$ with at most a simple pole at $z - \eta + \Lambda$. They satisfy $f_{ij}(y+1) = \text{Ad}_A^{-1} f_{ij}(y)$ and $f_{ij}(y+\tau) = \text{Ad}_B^{-1} f_{ij}(y)$ and are thus meromorphic sections of the adjoint bundle $\text{Ad } P$. These naturally act on sections of the bundle associated to \mathbb{C}^ℓ . Let us abbreviate the space of such sections as

$$\Theta := \{f : \mathbb{C} \rightarrow \mathbb{C}^\ell \text{ meromorphic} \mid f(y+1) = A^{-1}f(y), f(y+\tau)B^{-1}f(y)\}.$$

We obtain an RTT representation on Θ . More generally, we may define

$$L_N(z) := R_{01}^B(z-y_1) \cdots R_{0N}^B(z-y_N)$$

whose coefficients act on the space

$$\Theta_N := \{f : \mathbb{C}^N \rightarrow (\mathbb{C}^\ell)^{\otimes N} \text{ meromorphic} \mid f(y_i+1) = A_i^{-1}f(y_i), f(y_i+\tau)B_i^{-1}f(y_i)\}.$$

These are the sections of a vector bundle $V_N \rightarrow E^N$, which can be pulled back to the (η -deformed) configuration space of points of E or even on $M_{1,1+N}$ such that the R -matrix allows us to put an RTT representation on its sections.

There is a commuting action of S_N from the right on Θ_N via R -matrices: $(i\ j) \mapsto R_{ij}^B(y_i - y_j)$, as well as sections of the structure sheaf of the configuration space of points on E . Together, these form a generalization of the degenerate affine Hecke algebra. If these sections of the structure sheaf are replaced by the theta functions of [Has95], we may let the elliptic difference operators act. This space is

$$Th^l = \{f : \mathbb{C}^N \rightarrow \mathbb{C} \text{ holomorphic} \mid f(y+e_i) = f(y), f(y+\tau e_i) = \exp(-\pi i l \tau - 2\pi i l y_i) f(y)\}.$$

Then the S_N -invariant subspace is spanned by the $\binom{N+l}{l}$ $\hat{\mathfrak{gl}}_N$ -characters of level l . This defines a line bundle L on E^N .

All in all, we may twist the vector bundle V_N by L , obtaining $V_N \otimes L \rightarrow E^N$ and we have an action of S_N giving the descent data for a vector bundle W on the quotient stack $E^N // S_N$. Its space of meromorphic sections gives an object in \mathbb{C}_B via $L_N(z)$. This defines a functor from quasi-coherent modules on $E^N // S_N$ to \mathbb{C}_B .

Define the space

$$\Theta_{\ell,N}^l = \{f : \mathbb{C}^N \xrightarrow{\text{mer.}} (\mathbb{C}^\ell)^{\otimes N} \mid f(y_i+1) = A_i^{-1}f(y_i), f(y_i+\tau) = \exp(-\pi i l \tau - 2\pi i l y_i) B_i^{-1}f(y_i)\}.$$

These are sections of a vector bundle on E^N and we have actions

$$E(\mathfrak{gl}_\ell) \curvearrowright \Theta_{\ell,N}^\bullet \curvearrowright S_N \ltimes Th^\bullet$$

via $L_N(z)$, permutations acting via R -matrices, and theta functions acting by scalar multiplication. This is graded by the level l . We now want to compute $\Theta_{\ell,N}^l \otimes_{S_N \ltimes Th^l} Th^l$. This is done by projecting out the action of S_N on $\Theta_{\ell,N}^\bullet$. This is the Hilbert space of the elliptic spin RS model, you might call it the space of non-abelian characters of $A_{N-1}^{(1)}$ graded by level. Then we let Ruijsenaars difference operators act after Hasegawa. The reason this also acts on the sections of the vector bundle is the existence of a connection.

1.1.5 Generalized Schur-Weyl duality

In general, we would like to construct a Schur-Weyl duality for any bundle of conformal blocks for any genus. Braid/Hecke generators are obtained as monodromies along the configuration space, which become R -matrices, the coordinates become one set of generators and tangent vectors give a second set of generators. For genus zero, this gives the Schur-Weyl duality between the loop Yangian and the degenerate double affine Hecke algebra, while for genus one, this gives a Schur-Weyl duality between a degenerate elliptic double affine Hecke algebra and a loop elliptic quantum group $LE(\mathfrak{gl}_\ell)$.

Instead of coordinates plus tangent vectors, the generators can also come from the product of two surfaces, going into 4d CS theory.

1.2 Loop Yangian

1.2.1 As quantization of rational spin RS Poisson algebra

[AF98]

The Hamiltonian reduction in the classical case is done as follows: Start with $(A, g, S) \in \mathfrak{gl}_N^* \times GL_N \times \mathfrak{gl}_N^*$ and factorize $S_{ij} = \sum_\alpha a_i^\alpha b_j^\alpha$, defining $S_{ij}^{\alpha\beta} := a_i^\alpha b_j^\beta$, where α, β can range in $1, \dots, \ell$. Then

$$T^{\alpha\beta}(z) = \delta^{\alpha\beta} + \text{tr} \frac{S^{\alpha\beta}}{z - A} = \delta^{\alpha\beta} + \sum_{n \geq 0} T_n^{\alpha\beta} z^{-n-1}, \quad T_n^{\alpha\beta} = \text{tr} A^n S^{\alpha\beta}$$

generates the classical Yangian. Letting $J_n^{\alpha\beta} := \text{tr} g^n S^{\alpha\beta}$, we define

$$J^{\alpha\beta}(z) = \sum_{n=-\infty}^{\infty} J_n^{\alpha\beta} z^{-n-1},$$

which generates the classical loop algebra. On the reduced phase space, they are also given by

$$J_n^{\alpha\beta} = \sum_{ij} (\mathbf{L}^{n-1})_{ij} \mathbf{a}_j^\alpha \mathbf{c}_i^\beta,$$

where $\mathbf{L}, \mathbf{a}, \mathbf{c}$ are the invariant versions of $L = TgT^{-1}$, a , and c (T begin the diagonalizer for A). This makes it clear how the Lax matrix corresponds to the monodromy around a loop. Thus, it

is known that

$$J_1^{\alpha\beta} = \sum_i S_i^{\beta\alpha}, \quad S_i^{\alpha\beta} = \mathbf{c}_i^\alpha \mathbf{a}_i^\beta, \quad \mathbf{c}_i^\alpha = \sum_\beta S_i^{\alpha\beta}, \quad \mathbf{a}_i^\alpha = \frac{S_i^{\beta\alpha}}{\sum_\gamma S_i^{\beta\gamma}}$$

Recall that the loop Yangian $LY(\mathfrak{gl}_\ell)$ is Schur-Weyl dual to the degenerate double affine Hecke algebra \ddot{H}_N and that the center of the Yangian $Y(\mathfrak{gl}_\ell)$ generated by the quantum determinant gives Hamiltonians for the quantum trigonometric spin CM model, while the center of the loop algebra $L(\mathfrak{gl}_\ell)$ gives Hamiltonians for the quantum rational spin RS model. This gives natural quantizations to $T_n^{\alpha\beta}$ and $J_n^{\alpha\beta}$. The formula for $J_1^{\alpha\beta}$ suggests the quantization rule

$$S_i^{\alpha\beta} \rightarrow e_i^{\alpha\beta} \otimes X_i,$$

where $e_i^{\alpha\beta}$ is a matrix unit acting on the i th tensorand and X_i is the i th Laurent generator of \ddot{H}_N . The following Poisson rules remain to be checked:

$$\{S_i^{\alpha\beta}, S_j^{\mu\nu}\} = \frac{1}{y_i - y_j} (S_i^{\mu\beta} S_j^{\alpha\nu} + S_i^{\alpha\nu} S_j^{\mu\beta}) - \frac{\delta^{\beta\mu}}{y_i - y_j + \eta} (S_i S_j)^{\alpha\nu} + \frac{\delta^{\alpha\nu}}{y_j - y_i + \eta} (S_j S_i)^{\mu\beta}$$

and $\{y_i, S_j^{\alpha\beta}\} = S_j^{\alpha\beta} \delta_{ij}$. This second rule follows directly, the first is very non-trivial and should be checked using Mathematica. The first term seems unusual, but the last two terms look like the usual commutation relations in \mathfrak{gl}_ℓ .

1.2.2 Fock space representation

[Kod16]

We construct the level k Fock space. Let $U = \mathbb{C}[X^{\pm 1}] \otimes \mathbb{C}^\ell \otimes \mathbb{C}^k$. Given an integer K and $N < M$, there is a natural map $\bigwedge^N U \rightarrow \bigwedge^M U$. Define the Fock space

$$F_K = \varprojlim_N \bigwedge^N U.$$

Note that we have an isomorphism

$$\bigwedge^N U \cong (\mathbb{C}^\ell)^{\otimes N} \otimes_{S_N} (\mathbb{C}[X_1^{\pm 1}, \dots, X_N^{\pm 1}] \otimes (\mathbb{C}^k)^{\otimes N})$$

This clearly has an action of the affine Yangian by Schur-Weyl duality, and this extends to the limit F_K . For $k = 1$, we recover the trigonometric Calogero-Moser system.

Notation

Bibliography

- [AF98] G. Arutyunov and S. Frolov. On the Hamiltonian structure of the spin Ruijsenaars-Schneider model. *Journal of Physics A: Mathematical and General*, 31(18):4203–4216, May 1998. doi:10.1088/0305-4470/31/18/010.
- [ES98] P. Etingof and O. Schiffmann. A link between two elliptic quantum groups, 1998. arXiv:math/9801108.
- [Has95] K. Hasegawa. Ruijsenaars’ commuting difference operators as commuting transfer matrices, 1995. arXiv:q-alg/9512029.
- [Kod16] R. Kodera. Higher level Fock spaces and affine Yangian, 2016. arXiv:1607.03237.

Erklärung