

***Pseudarthria panii* (Fabaceae: Desmodieae), a new species from Asia, 120 years after its first collection**

RONG ZHANG^{1,2}, TING-SHUANG YI¹ & BO PAN^{3*}

¹ Germplasm Bank of Wild Species, Kunming Institute of Botany, Chinese Academy of Sciences, Kunming, 650201, Yunnan, China

² University of Chinese Academy of Sciences, Beijing, 100049, China

³ Centre for Integrative Conservation, Xishuangbanna Tropical Botanical Garden, Chinese Academy of Sciences, Mengla, 666303, Yunnan, China

*Author for Correspondence: pb@xtbg.org.cn

Abstract

A new trifoliate shrub, *Pseudarthria panii* (Fabaceae: Desmodieae), is described and illustrated. It resembles *Pseudarthria viscida*, but differs by its taller height, upright habit, late flowering, leaflets with an acuminate apex, longer light purple flowers, longer glabrous fruits, and more seeds. It occurs in the montane forest of China (Yunnan), Laos, Myanmar, Thailand, and Vietnam. Molecular evidence from the concatenated chloroplast fragments of *rbcL* and *matK* confirms its placement in *Pseudarthria*, which is a newly recorded genus from China, Laos, Thailand and Vietnam. The first collection of the new species can be dated back to ca. 120 years ago, i.e. Augustine Henry's collection in Yunnan during 1896–1899. A key to *Pseudarthria* is also presented.

Keywords: China, history of collection, legume, Leguminosae, taxonomy

Introduction

Pseudarthria Wight & Arnott (1834: 209) is a small genus of 4–6 species distributed in the Old World (Verdcourt 2000). Most of these species occur in seasonally dry tropical areas of Africa (Verdcourt 1971; Lock 1989), while only one species, *P. viscida* (L.) Wight & Arnott (1834: 209), is found in South and Southeast Asia (Baker 1879; Baker *et al.* 1963), where it is used in the treatment of fever, bronchial asthma, hemorrhoids, edema, diabetes mellitus, diarrhea and tuberculosis (Deepa *et al.* 2004). Traditionally, the genus was placed in the tribe Pseudarthrieae (Hutchinson 1964), subtribe Pseudarthriinae (Pramanik & Thothathri 1989) or subtribe Desmodiinae (Ohashi *et al.* 1981) of the tribe Desmodieae (Hutchinson 1964). However, molecular evidence revealed that it should be ascribed to the *Desmodium* group, one of three groups in the tribe Desmodieae (Nemoto & Ohashi 2003, Ohashi 2005, Jabbour *et al.* 2018).

The species of *Pseudarthria* are all pinnately 3-foliate herbs, shrubs or subshrubs and closely resemble *Desmodium* Desvaux (1813: 122), but differ in their unsegmented fruits (Baker 1871). During a field survey in Yunnan Province, China, we found a distinct ‘*Desmodium*-like’ species that morphologically did not match any of the known species from China, due to its non-articulated fruits with nearly straight sutures. Further morphological observations revealed that it represents an undescribed *Pseudarthria* species as described and illustrated below. Molecular phylogenetic analysis of the concatenated chloroplast fragments of *rbcL* and *matK* confirmed its position within *Pseudarthria*.

Materials and methods

Morphological studies

All morphological characters were examined in detail and compared with possible relatives based on the specimens from the Global Plants of JSTOR (<http://plants.jstor.org/>), HITBC, KUN, A, US and Kew herbaria.

Molecular methods and phylogenetic analyses

For the phylogenetic analyses we used the two concatenated chloroplast fragments of *rbcL* and *matK* from 40 species from 17 genera of the tribe Desmodieae. All sequences except that of the new species were acquired from GenBank (Table S1 in Appendix). The voucher specimens (PZ02) of the new species were deposited in the herbarium of the Kunming Institute of Botany, Chinese Academy of Sciences (KUN). Total genomic DNA was extracted from 100 mg of silica gel-dried leaflets with a modified CTAB protocol (Doyle 1987) and sent to Novogene, Nanjing, China, for library construction and next generation sequencing. A DNA 250 bp insert size library was constructed. More than 3 Gb paired-end reads of 2×150 bp for this species were generated on the Illumina HiSeq XTen platform. The reads were assembled using Linux-OS SPAdes v3.10.1 (Bankevich *et al.* 2012) with k-mer = 127 into contigs. Contigs were filtered by python script *slim_fastg_by_blast.py* (Jin *et al.* 2018) and later connected into plastome using Bandage Ubuntu dynamic v.8.0 (Wick *et al.* 2015). The connected plastome sequence and reference *Lotus japonicus* downloaded from GenBank (NC002694) were aligned with MAFFT (Katoh *et al.* 2013) using default parameters to copy annotations, coupled with manual adjustment in Geneious v.9.1.4 (Kearse *et al.* 2012). We then extracted *rbcL* and *matK* sequences from the assembled plastome for phylogenetic analyses. Sequences of *rbcL* and *matK* were deposited in GenBank under the accessions MG674205 and MG674206.

Phylogenetic analyses

Sequences were aligned with MAFFT (Katoh *et al.* 2013) using default parameters (L-INS-I algorithm) in Geneious v.9.1.4. Alignments were subsequently analyzed visually and adjusted manually. Phylogenetic analyses were performed with standard Bayesian inference (BI) and maximum-likelihood (ML) methods. The best-fit model was determined by MODELTEST v.3.7 (Posada & Crandall 1998) with the Akaike Information Criterion (AIC) (Posada & Buckley 2004). The BI analysis was performed with MrBayes v.3.2 (Ronquist *et al.* 2012) with four Markov Chain Monte Carlo (MCMC) runs using a random starting tree, an invgamma rate model with six discrete categories and 10 million generations, with a sampling frequency of one every 1000 generations and 25% of the trees were discarded as burn-in. Stationarity was considered to be reached when the average standard deviation of split frequencies was below 0.01. The ML analysis was performed with command line RAxML v.7.2.8 (Stamatakis 2006) in Linux OS, including tree robustness assessment using 1000 replicates of rapid bootstrap (the “-f a” option) with the GTRGAMMA substitution model to assess branch support (Stamatakis 2006).

Results

Phylogenetic analyses

The ML and Bayesian analyses of the combined data set of two chloroplast fragments of *matK* and *rbcL* generated congruent topologies. The ML tree with ML bootstrap (BS) and Bayesian posterior probability (PP) values is shown in Fig. 1. The new species *Pseudarthria panii* and *P. hookeri* Wight & Arnott (1834: 209) formed a clade strongly supported in the ML analysis (BS = 98%, PP = 0.75; Fig.1). The phylogeny showed that *Pseudarthria* was retrieved as sister to a clade containing *Alysicarpus*, *Christia*, *Codoriocalyx*, *Desmodium*, *Hegnera*, *Mecopus*, *Melliniella* and *Uraria*, and that *Desmodium* was not monophyletic.

Taxonomy

Pseudarthria panii Rong Zhang, Ting-Shuang Yi & Bo Pan (潘勃), *sp. nov.* 百年假节蚂蝗 (Fig. 2–3)

Type:—CHINA. Yunnan: Xishuangbanna, Menghai County, Xiding, 21°55'36"N, 100°7'39"E, 1800 m, 13 December 2014, *Bo Pan XD001* (holotype: Barcode 1262264, KUN! isotype: HITBC!)

Diagnosis:—This species is similar to *Pseudarthria viscida* in the rhomboid-ovate leaflets and unjointed pods, but differs by its taller and upright growth form (vs. 0.6–1.2 m in height, diffuse or prostrate), late flowering in September (vs. May), leaflets with an acuminate (vs. acute) apex, light purple (vs. pink) and longer (14 vs. 5 mm) flowers, longer (30–60 vs. 9–20 mm) and glabrous (vs. sticky pubescent) pods, and more (5–12 vs. 2–6) seeds. This new species also resembles *Desmodium megaphylla*, a species widely distributed in Asia including China, especially under the flowering condition, but differs from it by the linear and unsegmented fruits.




FIGURE 1. Phylogenetic relationships among 40 species from 17 genera of Desmodieae based on the combined data set of chloroplast *matK* and *rbcL*. The tree was constructed using the Maximum likelihood method. Numbers near the nodes are posterior probabilities and bootstrap percentages (PP, BP) from Bayesian analysis and Maximum likelihood, respectively. A dash (--) indicates the bootstrap <50% for ML tree. The new species in this study is shown in bold type.

Description:—Shrub with several clustered stems rising from one rootstock, 1.5–3.0 m tall, slightly zigzagged, and few branched. Old stem erect, terete, and glabrescent, while the young branchlets bending, more or less angled, and finely pubescent with small hooked hairs. Leaves two ranked, obviously getting smaller along the branch. Stipules triangulate, ca. 7 × 2 mm, caduceus late. Petioles 20–60 mm, covered with very short hooked hairs. Stipels linear, 2–4 mm. Petiolules shorter than the stipels, ca. 3 mm, pubescent with both long hairs and mixed with short hooked hairs. Leaflets 3, ovate, 30–120 × 15–60 mm, acuminate at apex, round at base, adaxial surface thin hairy with hooked hairs or glabrescent, but densely short pubescent abaxially; margins entire, revolved when dry. Terminal leaflets slightly bigger than or four times as big as the lateral ones. Secondary veins 4–6 pairs. Veins all sunken on the adaxial side. Main veins reach the margin. Inflorescences 50–140 mm long, much longer than the subtending leaf, 1–3 pseudoracemes clustered at the axil, ascending or pendulous. Each branchlet ends with a terminal inflorescence. Rachis of the inflorescence slender and short, 10–30 mm, pubescent with small hooked hairs. Bracts caducous. 2–3 flowers clustered at each node. Pedicels long and gracile, 10–15 mm, hooked hairy and soon glabrous. Bracteoles 2 below the calyx, minute. Calyx campanulate, 4 lobed, 4–5 mm, upper two teeth almost connate, lower lobe the longest. Corolla papilionaceous, light or pale purple. Standard orbicular to obovate, ca. 10 mm, with no auricles; wings lanceolate and incurved, ca. 14 mm, much larger than the keel; keel oblanceolate, ca. 10 mm; both wings and keel auriculate and clawed. Stamens 10, ca. 12 mm, vexillary stamen connate with the other 9 at base. Pistil linear, ca. 10 mm, with a dilated up-curved part and a




FIGURE 2. *Pseudarthria panii*. A. Flowering branch. B. Fruiting branch. C. Stipule on angulate stem. D. Fruits. E. *Conapium schenklingi*, weevil from the pod. A–D. Photos by Bo Pan. E. Courtesy of Dr. Zhi-Liang Wang.




FIGURE 3. *Pseudarthria panii*. A. abaxial side of leaflet (revolved margin), B. fruiting branch, C. petiole clothed with small hooked hairs, D. flower, E. stamens, F. pistil, G. calyx, H. standard, I. wing, K. keel, L. habit. Illustrations by Bo Pan, from the holotype—CHINA. Yunnan: Xishuangbanna, Menghai County, Xiding, 13 December 2014, Barcode 1262264, *Bo Pan XD001* (KUN).

nectary disc at base; stigma capitate; pods linear and glabrous, $30\text{--}60 \times 3\text{--}4$ mm, with 5–12 seeds; both sutures almost straight, valves with conspicuous reticulate veins when young; indehiscent when ripe, persistent on the branches for a long time. Valves break into squares when torn but are still connected along the sutures. Seeds elliptic and compressed, ca. 2×3 mm.

Phenology and distribution:—The flowering period is very short, from late September to early October and fruiting occurs in October to December, with the fruits persistent on the branches until the following May. This species is distributed in the broad-leaved montane forests in China (Yunnan Province), Laos, Myanmar, Thailand and Vietnam (Fig. 4), at an elevation of 900–2000 m. Though we found a very limited number of specimens, we presume that it has more populations and a wider distribution in the wild because there have been few field surveys in this region and a very low rate of identification. *Pseudarthria panii* is highly scattered, and each known location has only 3–5 individuals. Its young leaves and fruits are always infested by pests, e.g. the pods of the Menghai population are seriously damaged by a weevil *Conapium schenklingi* Wagner (1914) (Fig. 3E), identified by Professor Marek Wanat (University of Wrocław, Poland).



FIGURE 4. Geographical distribution of *Pseudarthria panii*. Prepared by Dr. Rui-Wu Zhou.

Etymology:—the epithet '*panii*' honours Lao-Cang Pan (1894–1962), who was the grandfather of the corresponding author Bo Pan. He was a carriage driver and horse veterinarian and has a large family and nurtured the love of nature among his children. His year of birth was during the period of Augustine Henry, an Irish Botanist, collecting in China. Henry collected plants in Taiwan in 1893–1894, and then Yunnan in 1896–1899 (Nelson 1983). Henry collected this *Pseudarthria* species (No. 12483B) in Simao, Yunnan Province, during 1898–1899, around 120 years ago, and considered it as '*Pueraria peduncularis*'. Here we propose a Chinese name '百年假节蚂蝗' for it; '百年' which means 'one hundred years', referring to the fact that it took more than 100 years to get a proper name. '假节蚂蝗' (<http://duocet.ibiodiversity.net/index.php?title=豆科>) derives from the Latin generic name, '*Pseud*—false' and '*arthria*—article', which means 'pods without articles'.

Additional specimens examined:—**CHINA. Yunnan:** Damenglong, Mengsheng, under thick woods, 1850 m, September 1936, *C.W. Wang* 78285 (PE). Lüchun, Huanglianshan, Qimaba, 900–1200 m, 1 November 1995, *Su-Gong Wu et al.* 1068 (KUN). No precise location record, hilly slopes, 1670 m, 21 November 1986, 86 Team (KUN). Simao, 5000 ft, *A. Henry* 12483B (A). Ximeng, broad-leaved forest, 2000 m, 22 October 1989, *Guo-Da Tao & Xi-Wen Li* 39759 (HITBC). Xishuangbanna, Mengla, Shangyong, 900 m, 22 October 1974, *Zeng-Hong Yang* 12055 (HITBC). Yingjiang County, Xima, Dazhai, 1700 m, 2 November 1974, *Guo-Da Tao* 13076, 12786 (HITBC). **LAOS.** 20 km south to the China-Laos border, 1700 m, 3 May 1993, *Guo-Da Tao* 930222 (HITBC). **MYANMAR. Chin State:** Par Kung Village, Kanpetlet. 4000 ft, 18 December 2011, *Ling Shein Man* 88022 (QBG). **THAILAND.** Tak, Tha Song Yang, Near Mon Kieo Lom, Mae Moei National Park, 17°31'25"N, 98°04'06"E, 820 m, edge of montane forest, 6 November 2010, *Pooma P. et al.* 7590 (BKF). **VIETNAM.** Annam, 1500 m, 13 September 1940, *M. Poilane* 30391 (P).

Key to *Pseudarthria* species:

1. Stems angulate; terminal leaflet ovate or rhomboid-ovate; plant from Asia2
1. Stems sulcate; terminal leaflet oblong, oblong-lanceolate or obovate; plant from Africa3
2. Shrub 1.5–3.0 m, upright; leaflet acuminate at apex; flowering in September to October, flower light purple and 14 mm long; pod glabrous and 30–60 mm long, seeds 5–12*P. panii*
2. Shrub 0.6–1.2 m, diffuse to prostrate; leaflet acute at apex; flowering in May, flower pink and 5 mm long; pod sticky pubescent and 9–20 mm long, seeds 2–6*P. viscida*
3. Lateral and terminal leaflets oblong-lanceolate*P. crenata*
3. Terminal leaflet obovate, lateral leaflets ovate to elliptic4
4. Racemes lax; pod 1–3-seeded*P. fagifolia*
4. Racemes dense; pod many-seeded5
5. Leaves glabrescent beneath; flowers in compact racemes*P. confertiflora*
5. Leaves densely persistently pubescent beneath; flowers dense but not in compact racemes5
6. Pod 4–5 mm wide, 5–6-seeded*P. macrophylla*
6. Pod 3 mm wide, 6–10-seeded*P. hookeri*

Acknowledgement

We are grateful to Ms Belinda (Wei Cao), Ms Shuying Zhong, Ms Lu Liu, Dr. Chengxi Ye, Dr. Li Wang, and Dr. Yu Song for their assistance with fieldwork, lab work and accommodation. We would like to thank Dr. Ashley Egan for the invitation to the Smithsonian Institution and Dr. Ruth Clark for the check of Kew herbarium specimens. We also thank the HITBC herbarium and the Central Laboratory of XTBG for access to specimens and equipment, the Plant Germplasm and Genomics Center, Kunming Institute of Botany, Chinese Academy of Sciences (KIB, CAS) for DNA extraction. Thanks to Prof. Richard Corlett for improving the manuscript, Dr. Rui-Wu Zhou for his preparation of the distribution map, Prof. Marek Wanat for the identification of the weevil, and Dr. Zhi-Liang Wang for the photo. We also gratefully acknowledge the anonymous reviewers and editor for their valuable comments and suggestions. This work is funded by the Recruitment Program of Foreign Experts (WQ20110491035).

References

- Baker, C.A. & Bakhuizen Van Den Brink, R.C. (1963) *Pseudarthria*. In: *Flora of Java*. NVP Noordhoff. Groningen, pp. 609.
- Baker, J.G. (1871) *Pseudarthria*. In: Oliver, D. (Eds.) *Flora of tropical Africa* 2. L. Reeve. London, pp. 167.
- Baker, J.G. (1879) *Pseudarthria*. In: Hooker, J.D. (Eds.) *The Flora of British India* 2. L. Reeve. London, pp. 153–154.
- Bankevich, A., Nurk, S., Antipov, D., Gurevich, A.A., Dvorkin, M., Kulikov, A.S., Lesin, V.M., Nikolenko, S.I., Pham, S., Prjibelski, A.D., Pyshkin, A.V., Sirotkin, A.V., Vyahhi, N., Tesler, G., Alekseyev, M.A. & Pevzner, P.A. (2012) SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *Journal of Computational Biology* 19 (5): 455–477.
<https://doi.org/10.1089/cmb.2012.0021>
- Deepa, M.A., Bai, V.N. & Basker, S. (2004) Antifungal properties of *Pseudarthria viscida*. *Fitoterapia* 75 (6): 581–584.
<https://doi.org/10.1016/j.fitote.2004.04.008>
- Desvaux, N.A. (1813) *Journal de Botanique, Appliquée à l'Agriculture, à la Pharmacie, à la Médecine et aux Arts* 1 (3): 122, pl. 5, f. 15.
- Doyle, J.J. & Doyle, J.L. (1987) A rapid DNA isolation procedure for small quantities of fresh leaf tissue. *Phytochemical Bulletin* 19:11–15.
- Hutchinson J. (1964) Fabaceae. In: *The genera of flowing plants*. Vol. 1. The Clarendon Press, Oxford.
- Jabbour, F., Gaudeul, M., Lambourdière, J., Ramstein, G., Hassanin, A., Labat, J.N. & Sarthou, C. (2018) Phylogeny, biogeography and character evolution in the tribe Desmodieae (Fabaceae: Papilionoideae), with special emphasis on the New Caledonian endemic genera. *Molecular Phylogenetics and Evolution* 118 (2018): 108–121.
<https://doi.org/10.1016/j.ympev.2017.09.017>
- Jin, J.J., Yu, W.B., Yang, J.B., Song, Y., Yi, T.S. & Li, D.Z. (2018) GetOrganelle: a simple and fast pipeline for de novo assembly of a complete circular chloroplast genome using genome skimming data. *bioRxiv* 256479.
<https://doi.org/10.1101/256479>
- Katoh, K. & Standley, D.M. (2013) MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Molecular Biology and Evolution* 30 (4):772–780.
<https://doi.org/10.1093/molbev/mst010>
- Kearse, M., Moir, R., Wilson, A., Stones-Havas, S., Cheung, M., Sturrock, S., Buxton, S., Cooper, A., Markowitz, S., Duran, C., Thierer, T., Ashton, B., Meintjes, P. & Dremmond, A. (2012) Geneious basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics* 28 (12): 1647–1649.
<https://doi.org/10.1093/bioinformatics/bts199>
- Lock, J.M. (1989) *Legumes of Africa: a check-list*. Royal Botanic Garden, Kew, pp. 210–251.
- Nelson, E.C. (1983) Augustine Henry and the exploration of the Chinese Flora. *Arnoldia* 43 (1): 21–38.
- Nemoto, T. & Ohashi, H. (2003) Diversity and Evolution of Anatomical Structure of Loment in Tribe Desmodieae (Papilionoideae). In: Klitgaard, B.B. & Bruneau, A. (Eds.) *Advances in Legume Systematics*, part 10. Royal Botanic Gardens, Kew, pp. 395–412.
- Ohashi, H. (2005) Desmodieae. In: Lewis, G., Schrire, B., Mackinder, B. & Lock, M. (Eds.) *Legumes of the World*. Royal Botanic Gardens, Kew, pp. 433–445.
- Ohashi, H., Polhill, R.H. & Schubert, B.G. (1981) Desmodieae. In: Polhill, R.M. & Raven, P.H. (Eds.) *Advances in Legume Systematics*, Part 1. Royal Botanic Gardens, Kew, pp. 292–300.
- Posada, D. & Buckley, T.R. (2004) Model selection and model averaging in phylogenetics: advantages of Akaike information criterion and Bayesian approaches over likelihood ratio tests. *Systematic Biology* 53 (5): 793–808.
<https://doi.org/10.1080/10635150490522304>
- Posada, D. & Crandall K.A. (1998) MODELTEST: testing the model of DNA substitution. *Bioinformatics* 14 (9): 817–818.
<https://doi.org/10.1093/bioinformatics/14.9.817>
- Pramanik, A. & Thothathri, K. (1989) Pseudarthriinae, a new status for tribe Pseudarthrieae (Leguminosae). *Taxon* 38 (1): 109–110.
<https://doi.org/10.2307/1220907>
- Ronquist, F., Teslenko, M., Van der Mark, P., Ayres, D.L., Darling, A., Höhna, S., Larget, B., Liu, L., Suchard, M.A. & Huelsenbeck, J.P. (2012) MrBayes 3.2: Efficient Bayesian phylogenetic inference and model choice across a large model space. *Systematic Biology* 61 (3): 539–542.
<https://doi.org/10.1093/sysbio/sys029>
- Schidler, A.K. (1914) Das genus *Pseudarthria* Wight et Arn. *Repertorium specierum novarum regni vegetabilis, Beihefte* 2: 3–17.
- Stamatakis, A. (2006) RAXML-VI-HPC: maximum likelihood-based phylogenetic analysis with thousands of taxa and mixed models. *Bioinformatics* 22 (21): 2688–2690
<https://doi.org/10.1093/bioinformatics/btl446>

- Verdcourt, B. (1971) *Pseudarthria*. In: Milne-Redhead, E. & Polhill, R.M. (Eds.) *Flora of Tropical East Africa. Leguminosae Part3, subfamily Papilionoideae 1*. Crown Agents for Overseas Governments & Administrations. London, pp. 483–487.
- Verdcourt, B. (2000) Leguminosae-Papilionoideae. In: *Flora Zambesiaca* 3 (6). Royal Botanic Gardens. Kew, pp. 25–27.
- Wick, R.R., Schultz, M.B., Zobel, J. & Holt, K.E. (2015) Bandage: interactive visualization of de novo genome assemblies. *Bioinformatics* 31 (20): 3350–3352.
<https://doi.org/10.1093/bioinformatics/btv383>
- Wight, R. & Arnott, G.A.W. (1834) *Pseudarthria*. In: *Prodromus Florae Peninsulae Indiae Orientalis*. Parbury, Allen, & CO. London, pp. 209.

TABLE S1. Accession number of *matK* and *rbcL* sequences of sampled species obtained from GenBank.

species name	<i>matK</i>	<i>rbcL</i>
<i>Akschindlium godefroyanum</i> (Kuntze) H. Ohashi	KX713079	KY702639
<i>Alysicarpus heyneanus</i> Wight & Arn.	KT222213	KT222229
<i>Alysicarpus pubescens</i> J.S.Law	KT222218	KT222236
<i>Alysicarpus vaginalis</i> (L.) DC.	JQ587509	KJ773259
<i>Arthroclianthus deplanchei</i> Hochr.	KX213147	KY702603
<i>Campylotropis macrocarpa</i> (Bunge) Rehder	AY386870	EU717277
<i>Christia vespertilionis</i> (L.f.) Bakh.f.	KF621098	KF621119
<i>Codoriocalyx motorius</i> (Houtt.) H. Ohashi	KF621099	KX527051
<i>Codoriocalyx motorius</i> (Houtt.) H. Ohashi	HM049526	KX344706
<i>Dendrolobium lanceolatum</i> (Dunn) Schindl.	AB924975	KY702645
<i>Desmodium barbatum</i> (L.) Benth.	EU717420	JQ591689
<i>Desmodium canadense</i> (L.) DC.	HQ593266	HQ590061
<i>Desmodium ciliare</i> (Willd.) DC.	KJ772716	KJ773442
<i>Desmodium distortum</i> (Aubl.) J.F.Macbr.	JQ587590	JQ591694
<i>Desmodium floridanum</i> Chapman	EF549994	KJ773443
<i>Desmodium gangeticum</i> (L.) DC.	KP093702	KX344694
<i>Desmodium heterocarpon</i> (L.) DC.	AB925045	KY702613
<i>Desmodium illinoense</i> A.Gray	KT456906	KX385961
<i>Desmodium incanum</i> DC.	JF501110	KJ773444
<i>Desmodium laxiflorum</i> DC.	KP094091	KP095045
<i>Desmodium paniculatum</i> (L.) DC.	KJ772717	KX385962
<i>Desmodium psilocarpum</i> A.Gray	AY386896	LT576814
<i>Desmodium purpusii</i> Brandegees	JQ587603	JQ591711
<i>Desmodium styracifolium</i> (Osbeck) Merr.	JN407126	JN407286
<i>Desmodium triflorum</i> (L.) DC.	KJ772719	KJ773446
<i>Hyloidesmum pauciflorum</i> (Nutt.) H. Ohashi & R.R. Mill	EU717421	EU717280
<i>Hegnera obcordata</i> (Miq.) Schindl.	AB924977	KY702617
<i>Hyloidesmum glutinosum</i> (Muhl. ex Willd.) H. Ohashi & R.R. Mill	KJ592990	KJ593475
<i>Hyloidesmum laxum</i> (DC.) H.Ohashi & R.R.Mill	KF621102	KF621124
<i>Hyloidesmum nudiflorum</i> (L.) H. Ohashi & R.R. Mill	KP642947	KP402657
<i>Lespedeza capitata</i> Michx.	KT456912	KJ773621
<i>Lespedeza cuneate</i> (Dum.Cours.) G.Don	EU717416	U74215
<i>Mecopus nidulans</i> Benn.	KX713102	KY702622
<i>Melliniella micrantha</i> Harms	KX713103	KY702623
<i>Nephrodesmus ferrugineus</i> Däniker	KX213176	KY702651
<i>Ohwia caudata</i> (Thunb.) H.Ohashi	KX713109	KX527365
<i>Phyllodium pulchellum</i> (L.) Desv.	HM049524	KY702658
<i>Pseudarthria hookeri</i> Wight & Arn.	JF270902	JF265559
<i>Pseudarthria panii</i> Rong Zhang, Ting-Shuang Yi & Bo Pan	MG674205	MG674206
<i>Tadehagi triquetrum</i> (L.) H.Ohashi	JN407127	KX527094
<i>Uraria lagopodoides</i> Roxb.	KF621107	KF621128

Note: Accession numbers HM049526 and KX344706 were recorded as *Desmodium gyrans* (L. f.) DC. in GenBank, we use the accepted name *Codoriocalyx motorius* (Thunb.) H. Ohashi in this study (<http://www.tropicos.org/Name/100356174>); similarly, EU717421 and EU717280 as *Desmodium pauciflorum* (Nutt.) DC. in Genbank, we use the accepted name *Hedysarum pauciflorum* (Nutt.) H. Ohashi & R.R. Mill (<http://www.tropicos.org/Name/50327477>).