

Insects

1227 species

[Home](#)

Insects

[Use Filters](#)

Sort By:

Popularity

A to Z

Group By:

None

Family

Population Status

Abisara echerius

[Abisara echerius](#)

[Abisara echerius](#), the plum Judy, is a small but striking butterfly found in Asia belonging to the [Punches and Judies family \(Riodinidae\)](#). It is difficult to distinguish it from [Abisara bifasciata](#). This [active butterfly](#) is usually seen at the tops...

Acalymma vittatum

[Acalymma vittatum](#)

[Acalymma vittatum](#), the striped cucumber beetle, is a beetle of the family [Chrysomelidae](#) and a serious [pest of cucurbit crops](#) in both larval and adult stages. It is distributed from eastern North America to the [Rocky Mountains](#). It is replaced in...

Acanthops falcata

Acanthops falcata

Acanthops falcata, common name South American dead leaf mantis or boxer mantis, is a species of praying mantis in the subfamily Acanthopinae of the family Acanthopidae and is one of many praying mantises from various genera that resembles a dead...



Acherontia atropos

Acherontia atropos

Acherontia atropos, the African death's-head hawkmoth, is the most widely recognized of three species within the genus Acherontia (the other two being Acherontia lachesis and Acherontia styx). It is most commonly identified by the vaguely...



Acherontia lachesis

Acherontia lachesis

Acherontia lachesis, the greater death's head hawkmoth or bee robber, is a large (up to 13 cm wingspan) sphingid moth found in India, Sri Lanka and much of the Oriental region. It is one of the three species of death's-head hawkmoth genus,...



Acilius sulcatus

Acilius sulcatus

Acilius sulcatus is a species of water beetle in the family Dytiscidae. It is fairly large (14.4 – 18.2 mm), with color variation shown throughout its range. Typically it is yellow and black.



Acraea horta

Acraea horta

Acraea horta or the garden acraea is a butterfly of the family Nymphalidae. It is found in South Africa and Zimbabwe.



Acraea terpsicore

Acraea terpsicore

Acraea terpsicore, the tawny coster, is a small, 53–64 millimetres (2.1–2.5 in), leathery-winged butterfly common in grassland and scrub habitats. It belongs to the Nymphalidae or brush-footed butterfly family. It has a weak fluttery flight. It...



Acronicta afflicta

Acronicta afflicta

Acronicta afflicta, the afflicted dagger moth, is a moth of the family Noctuidae. It is found in Canada (Nova Scotia, Quebec and Ontario), the United States (including Alabama, North Carolina, Oklahoma, Georgia, Maryland, New York and Ohio) as...



Acronicta alni

Acronicta alni

Acronicta alni, the alder moth, is a moth of the family Noctuidae. It is found in Europe (from southern Fennoscandia to Spain, Italy and the Balkans), Turkey, the European part of Russia and the neighbouring countries, the Caucasus, the Ural,...



Acronicta americana

Acronicta americana

Acronicta americana, the American dagger moth, is a moth of the family Noctuidae. It was originally described by Thaddeus William Harris in 1841 and is native to North America.



Acronicta insita

Acronicta insita

Acronicta insita, the large gray dagger or fingered dagger, is a moth of the family Noctuidae. The species was first described by Augustus Radcliffe Grote in 1874. It is found from Newfoundland west to the Pacific coast and Vancouver Island and...



Acronicta lobeliae

Acronicta lobeliae

Acronicta lobeliae, the lobelia dagger moth or greater oak dagger moth, is a moth of the family Noctuidae. The species was first described by Achille Guenée in 1852. It is found in North America. The larvae feed on Quercus species.



Acronicta rumicis

Acronicta rumicis

Acronicta rumicis, the knot grass moth, is a species of moth which is part of the genus Acronicta and family Noctuidae. It was first described by Carl Linnaeus in his 1758 10th edition of Systema Naturae. It is found in the Palearctic region. A....



Acrophylla titan

Acrophylla titan

Acrophylla titan, the titan stick insect, is the second-longest stick insect found in Australia. First described by William Sharp Macleay in 1826, it was considered to be the longest stick insect in the world until the discovery of Ctenomorpha...



Actaeon beetle

Megasoma actaeon

Actaeon beetle (Megasoma actaeon) is a rhinoceros beetle of the family Scarabaeidae.



Actias artemis

Actias artemis

Actias artemis is a moth native to Japan, Korea, China, India and Malaysia. The species was first described by Otto Vasilievich Bremer and William Grey in 1853. Actias artemis is a close relative and look-alike of Actias luna, the American Luna...



Actias isis

Actias isis

Actias isis is a moth of the family Saturniidae first described by Léon Sonthonnax in 1899.



Actias ningpoana

Actias ningpoana

Actias ningpoana, the Chinese moon moth, is a moth of the family Saturniidae. The species was first described by father-and-son entomologists Cajetan and Rudolf Felder in 1862. It is quite large, and has long, curved, hindwing tails. There are...



Actias selene

Actias selene

Actias selene, the Indian moon moth or Indian luna moth, is a species of saturniid moth from Asia. It was first described by Jacob Hübner in 1807. This species is popular among amateur entomologists and is often reared from eggs or cocoons that...



Actias sinensis

Actias sinensis

Actias sinensis, the Golden moon moth, is a moth of the Family Saturniidae. It is found in China, Taiwan, Vietnam, Myanmar, India and Thailand. The species was first described by Francis Walker in 1855.



Adalia bipunctata

Adalia bipunctata

Adalia bipunctata, the two-spot ladybird, two-spotted ladybug or two-spotted lady beetle, is a carnivorous beetle of the family Coccinellidae that is found throughout the holarctic region. It is very common in western and central Europe. It is...



Adelpha californica

Adelpha californica

Adelpha californica, the California sister, is a species of butterfly in the family Nymphalidae. They are common in California, but can also be found in western Nevada and Oregon, as well as in northern Baja California. The upper surfaces of...



Adetomyrma venatrix

Adetomyrma venatrix

Adetomyrma venatrix, more commonly known as the Dracula ant, so named because of its grisly feeding habits of drinking the blood of its young, is an endangered species of ants endemic to Madagascar. Workers of this species are blind. The species...



Adhemarius gannascus

Adhemarius gannascus

Adhemarius gannascus is a moth of the family Sphingidae first described by Caspar Stoll in 1790.



Adonis blue

Lysandra bellargus

The Adonis blue (Lysandra bellargus, also known as Polyommatus bellargus) is a butterfly in the family Lycaenidae. It inhabits the Palearctic realm (Western Europe, Central Europe, Southern Europe, Southern Russia, Iraq, Iran, Caucasus,....



Adoxophyes orana

Adoxophyes orana

Adoxophyes orana, the summer fruit tortrix, is a moth of the family Tortricidae. It is found in the Palearctic realm and Taiwan. The wingspan is 17–22 mm. The moth flies in two generations from May to November. The larvae overwinter in loosely...



Aedes aegypti

Aedes aegypti

Aedes aegypti (/ˈiːdiːz/ from Greek ἀηδής: "hateful" and /aɪˈdʒɛpti/ from Latin, meaning "of Egypt"). the yellow fever mosquito, is a mosquito that can spread dengue fever, chikungunya, Zika fever, Mayaro and yellow fever viruses, and other...



Aedes albopictus

Aedes albopictus

Aedes albopictus (Stegomyia albopicta), from the mosquito (Culicidae) family, also known as the (Asian) tiger mosquito or forest mosquito, is a mosquito native to the tropical and subtropical areas of Southeast Asia. In the past few centuries,...



Aedes cinereus

Aedes cinereus

Aedes cinereus is a mosquito species in the genus Aedes.



Aedes vexans

Aedes vexans

Aedes vexans, the inland floodwater mosquito or tomguito, is a cosmopolitan and common pest mosquito. This species has been collected in southern California.



Aegiale hesperiaris

Aegiale hesperiaris

Aegiale hesperiaris, commonly known as the tequila giant skipper, is a species of butterfly in the family Hesperinae. It is the only species in the monotypic genus Aegiale. Its caterpillar is one of two varieties of edible "maguey worms" that...



Aeshna affinis

Aeshna affinis

Aeshna affinis, the southern migrant hawker or blue-eyed hawker, is a dragonfly found in southern Europe and Asia. It is in the family Aeshnidae and is very similar in appearance to A. mixta.



Aeshna canadensis

Aeshna canadensis

Aeshna canadensis, the Canada darner, is a species of dragonfly in the family Aeshnidae. It is common throughout southern Canada and the northern United States.



African armyworm

Spodoptera exempta

The African armyworm (Spodoptera exempta), also called okalombo, kommandowurm, or nutgrass armyworm, is a species of moth of the family Noctuidae. The larvae often exhibit marching behavior when traveling to feeding sites, leading to the common...



Agathidium vaderi

Agathidium vaderi

Agathidium vaderi is a species of round fungus beetle in the family Leiodidae. It is found in North America. The beetle was named after the fictional character Darth Vader by Cornell University entomologists due to its shiny helmetlike head that...



Agelastica alni

Agelastica alni

Agelastica alni, the alder leaf beetle, is a species of leaf beetle (Chrysomelidae) in the genus Agelastica. Agelastica alni is distributed in Europe, the Caucasus, Siberia, north-eastern Kazakhstan, and in the 19th century was introduced to the...



Aglae

Aglae caerulea

Aglae is a genus of euglossine bees, with the only described species Aglae caerulea. Like all orchid bees, it is restricted to the Neotropics. They are metallic blue. This species, like the genus Exaerete, is a nest parasite on free-living...



Aglais io

Aglais io

Aglais io, the European peacock, or the peacock butterfly, is a colourful butterfly, found in Europe and temperate Asia as far east as Japan. It was formerly classified as the only member of the genus Inachis (the name is derived from Greek...



Aglais milberti

Aglais milberti

Aglais milberti, the fire-rim tortoiseshell or Milbert's tortoiseshell, is considered the only species of the proposed Aglais genus that occurs in North America.



Aglossa cuprina

Aglossa cuprina

Aglossa cuprina, the grease moth, is a snout moth, family Pyralidae, described by Philipp Christoph Zeller in 1872. The grease moth is closely related to the genus Pyralis, and as a result, is usually associated with the meal moth, Pyralis...



Agreeable tiger moth

Spilosoma congrua

The agreeable tiger moth (*Spilosoma congrua*) is one of three species of white tiger moth which are common in the United States. It has pronounced black eyes, white abdomen, and orange "bib" which set it apart from its cousin the Virginia tiger...



Agriades optilete

Agriades optilete

Agriades optilete, the cranberry blue, is a butterfly of the family Lycaenidae. It is found in north eastern Europe, the Alps, North Asia, Japan, Korea and north western North America. The length of the forewings is about 14 mm. The butterfly...



Agrias amydon

Agrias amydon

Agrias amydon, the Amydon agrias or white-spotted agrias, is a butterfly of the family Nymphalidae. It is found in Mexico, Central America, and South America. The larvae feed on Erythroxylum species including E. havanense.



Agrias narcissus

Agrias narcissus

Agrias narcissus is a butterfly of the family Nymphalidae. It is found in South America.



Agrilus anxius

Agrilus anxius

Agrilus anxius, the bronze birch borer, is a wood-boring buprestid beetle native to North America, more numerous in warmer parts of the continent and rare in the north. It is a serious pest on birch trees (Betula), frequently killing them. The...



Agrius cingulata

Agrius cingulata

Agrius cingulata, the pink-spotted hawkmoth or sweetpotato hornworm, is a moth in the family Sphingidae. The species was first described by Johan Christian Fabricius in 1775.



Agrius convolvuli

Agrius convolvuli

Agrius convolvuli, the convolvulus hawk-moth, is a large hawk-moth. It is common throughout Europe, Asia, Africa, Australia and New Zealand, partly as a migrant. In New Zealand, it is also known as the kumara moth, and in the Māori language as...



Agrochola circellaris

Agrochola circellaris

Agrochola circellaris, or The Brick, is a species of moth of the family Noctuidae. The species was first described by Johann Siegfried Hufnagel in 1766. It is distributed throughout most of Europe, Asia Minor and Armenia. This is a variable...



Ailanthus webworm

Atteva aurea

The ailanthus webworm (Atteva aurea) is an ermine moth now found commonly in the United States. It was formerly known under the scientific name Atteva punctella (see Taxonomy section). This small, very colorful moth resembles a true bug or beetle...



Alabama argillacea

Alabama argillacea

Alabama argillacea, the cotton leafworm or cotton worm, is a moth of the family Erebidae. It is native to the New World, but has been extirpated from the United States and Canada, having not been recorded since 1998. In the Neotropics, it can be...



Alkali bee

Nomia melanderi

The alkali bee, Nomia melanderi, is a ground-nesting bee native to deserts and semi-arid desert basins of the western United States. It was described by Theodore Dru Alison Cockerell in 1906. While solitary, these bees nest near each other and...



Allegheny mound ant

Formica exsectoides

The Allegheny mound ant (Formica exsectoides) is a species of ant native to the Atlantic area of North America. Its range extends from Nova Scotia to parts of Georgia. Like other field ants, the Allegheny mound ant builds large mounds, however...



Allograpta obliqua

Allograpta obliqua

Allograpta obliqua is a common North American species of hoverfly, commonly known as oblique streaktail. The larvae are important predators of aphids while adults are pollinators.



Allomerus decemarticulatus

Allomerus decemarticulatus

Allomerus decemarticulatus is an Amazonian ant species found in the tropics of South America. This species is most notable for the workers' complex and extreme predatory behavior, which involves a symbiosis with both a plant and fungal...



Almond moth

Cadra cautella

The almond moth or tropical warehouse moth (*Cadra cautella*) is a small, stored-product pest. Almond moths infest flour, bran, oats, and other grains, as well as dried fruits. It belongs to the family of snout moths (*Pyralidae*), and more...



Alsophila pometaria

Alsophila pometaria

Alsophila pometaria, the fall cankerworm, is a moth of the family Geometridae. It is found in North America from Nova Scotia west to Alberta, south to Colorado and California and zones of Spain



Alucita hexadactyla

Alucita hexadactyla

Alucita hexadactyla (twenty-plume moth) is a "micromoth" of the many-plumed moth family (Alucitidae). It is found in Eurasia. It was previously thought to also occur in North America, but a 2004 study showed that the North American species are...



Alypia octomaculata

Alypia octomaculata

Alypia octomaculata, the eight-spotted forester, is a moth of the family Noctuidae first described by Johan Christian Fabricius in 1775. It is native to Canada, but can be found today throughout Northern America, ranging between Nova Scotia to...



Amara aenea

Amara aenea

Amara aenea is a ground beetle common in almost the whole of Europe and Northern Asia. Its range covers also parts of Northern Africa. It is known as the common sun beetle. A. aenea adults are predators that eat other insects, such as the apple...



Amara aulica

Amara aulica

Amara aulica is a species of beetle of the genus Amara in the Harpalinae subfamily. It is native to Europe.



Amata huebneri

Amata huebneri

Amata huebneri, commonly known as the wasp moth, is a species of moth in the family Erebidæ (subfamily Arctiinae - "woolly bears" or "tiger moths"). The species was first described by Jean Baptiste Boisduval in 1829. It is found from the Indo...



Amblyptilia acanthadactyla

Amblyptilia acanthadactyla

Amblyptilia acanthadactyla, also known as the beautiful plume, is a moth of the family Pterophoroidea found in across the Palearctic including Europe. The species was first described by the German entomologist, Jacob Hübner in 1813.



Amegilla bombiformis

Amegilla bombiformis

Amegilla bombiformis, commonly known as the teddy bear bee or golden haired mortar bee, is an Australian native bee in the family Apidae. It was originally described by F. Smith in 1854 as Saropoda bombiformis from a collection near the Richmond...



American cockroach

Periplaneta americana

The American cockroach (Periplaneta americana) is the largest species of common cockroach, and often considered a pest. In certain regions of the U.S. it is colloquially known as the waterbug, though it is not a true waterbug since it is not...



American lady

Vanessa virginiensis

The American painted lady or American lady (Vanessa virginiensis) is a butterfly found throughout North America. The larvae feed on various Asteraceae, such as the cudweeds (genus Gnaphalium), the pussytoes (Antennaria), and the everlastings...



American snout

Libytheana carinenta

The American snout or common snout butterfly (Libytheana carinenta) is a member of the subfamily Libytheinae in the brush-footed butterfly family, Nymphalidae. This species is found in both North and South America. The larval host plants are...



Ammopelmatus cahuilaensis

Ammopelmatus cahuilaensis

Ammopelmatus cahuilaensis (commonly known as the Coachella Valley Jerusalem cricket) is a species of insect in the family Stenopelmatidae. The species is found in the Coachella Valley and was described by Ernest R. Tinkham in 1968, in The Great...



Ammopelmatus kelsoensis

Ammopelmatus kelsoensis

Ammopelmatus kelsoensis, commonly known as the Kelso Jerusalem cricket, is a species of insect in the family Stenopelmatidae. It is endemic to the Kelso Dunes in the United States.



Amplipterus panopus

Amplipterus panopus

Amplipterus panopus, the mango hawkmoth, is a moth of the family Sphingidae. The species was first described by Pieter Cramer in 1779. It is found in Sri Lanka, southern and northern India (including the Andaman Islands and Nicobar Islands),...



Anagrapha falcifera

Anagrapha falcifera

Anagrapha falcifera, the celery looper, is a moth of the family Noctuidae. The species was first described by William Kirby in 1837. It is found in North America from Newfoundland, Labrador and southern Canada to Georgia, Mississippi, Texas,...



Anartia amathea

Anartia amathea

Anartia amathea, the brown peacock (scarlet peacock, red peacock), is a species of nymphalid butterfly, found primarily in South America. This butterfly is very similar to the banded peacock or Anartia fatima, which primarily exists north of the...



Anartia fatima

Anartia fatima

Anartia fatima, the banded peacock, is a butterfly in the family Nymphalidae. It is commonly found in south Texas, Mexico, and Central America but most studied in Costa Rica. This butterfly prefers subtropical climates and moist areas, such as...



Anasa tristis

Anasa tristis

Anasa tristis is a species of bug in the family Coreidae. It is a major pest of squash and pumpkins, found throughout North America, and is a vector of the cucurbit yellow vine disease bacterium. These bugs can emit an unpleasant odor when...



Anax guttatus

Anax guttatus

Anax guttatus, the pale-spotted emperor or lesser green emperor, is a dragonfly of the family Aeshnidae.



Anax parthenope

Anax parthenope

Anax parthenope, the lesser emperor, is a dragonfly of the family Aeshnidae. It is found in Southern Europe, North Africa, Middle East and Asia.



Anchiale austrotessulata

Anchiale austrotessulata

Anchiale austrotessulata (synonyms include Ctenomorphodes tessulatus), the tessellated stick insect, tessellated phasmid or tessulata stick insect, is a medium-sized, stick insect found in the Brisbane area of Australia. Fully grown males in...



Ancistrocerus gazella

Ancistrocerus gazella

The European potter wasp or European tube wasp (Ancistrocerus gazella) is a species of potter wasp. As an imago (adult), the female collects as many as 20 caterpillars for each nest, which consists of a single cell. Her larval offspring then feed...



Ancistroides folus

Ancistroides folus

Ancistroides folus, the grass demon, is a small but prominent butterfly found in India & Nepal that belongs to the skipper family, Hesperiidae. It is regarded as an occasional pest of ginger and turmeric plants.



Andricus grossulariae

Andricus grossulariae

Andricus grossulariae is a gall wasp species inducing agamic acorn cup galls on oak tree acorn cups and sexual phase galls on catkins. Synonyms include Andricus fructuum (Trotter, 1899), Andricus gemellus (Belizin & Maisuradze, 1961), Andricus...



Andricus quercuscalicis

Andricus quercuscalicis

Andricus quercuscalicis is a gall wasp species inducing knopper galls. Knopper galls develop as a chemically induced distortion of growing acorns on pedunculate oak (Quercus robur L.) trees, caused by gall wasps, which lay eggs in buds with their...



Angle shades

Phlogophora meticulosa

The angle shades (Phlogophora meticulosa) is a moth of the family Noctuidae. The species was first described by Carl Linnaeus in his 1758 10th edition of Systema Naturae. It is distributed throughout Europe as far east as the Urals and also in...



Angoumois grain moth

Sitotroga cerealella

The Angoumois grain moth (*Sitotroga cerealella*) is a species of the Gelechiidae moth family, commonly referred to as the "rice grain moth". It is most abundant in the temperate or tropical climates of India, China, South Africa, Indonesia,...



Anicla infecta

Anicla infecta

Anicla infecta is a moth of the family Noctuidae, described by Ochsenheimer in 1816. It is known as the green cutworm when a caterpillar and the green cutworm moth when mature. It is found from south-eastern Canada (Nova Scotia and Quebec)...



Anisolabis maritima

Anisolabis maritima

Anisolabis maritima, commonly known as the maritime earwig or the seaside earwig, is a species of earwig in the family Anisolabididae. Similar to the seashore earwig, this species can be found near the shore line, and is cosmopolitan. It can be...



Anisota senatoria

Anisota senatoria

Anisota senatoria, the orangestriped oakworm, also known as the orange-tipped oakworm, is a Nearctic moth of the family Saturniidae and subfamily Ceratocampinae. It is one of the more common Saturniids, reaching pest status occasionally in the...



Anisota stigma

Anisota stigma

Anisota stigma, the spiny oakworm moth, is a moth of the family Saturniidae. The species was first described by Johan Christian Fabricius in 1775. It is found in North America from Massachusetts and southern Ontario to Florida, west to Minnesota,...



Anisota virginiensis

Anisota virginiensis

Anisota virginiensis, the pink-striped oakworm moth, is a species of silk moth of the family Saturniidae.



Anomis flava

Anomis flava

Anomis flava, the cotton looper, tropical anomis or white-pupiled scallop moth, is a moth of the family Erebiidae. The species was first described by Johan Christian Fabricius in 1775. It is found in large parts of the world, including China,...



Anomis sabulifera

Anomis sabulifera

Anomis sabulifera, the angled gem or jute semi-looper, is a moth of the family Erebiidae. The species was first described by Achille Guenée in 1852. It has a Palearctic distribution and ranges from Africa eastwards to India, Sri Lanka and...



Anophthalmus hitleri

Anophthalmus hitleri

Anophthalmus hitleri (Slovene: Hitlerjev brezokec) is a species of blind cave beetle found only in about fifteen humid caves in Slovenia. The blind cave beetle shares its genus with 41 other species and 95 different subspecies. Members of its...



Anoplognathus pallidicollis

Anoplognathus pallidicollis

Anoplognathus pallidicollis is a relatively large beetle of the family Scarabaeidae, native to Australia. The beetle has a life span of 24 months and it grows up to 20 mm in length. Adults were once common in the summer months, particularly...



Anotogaster sieboldii

Anotogaster sieboldii

Anotogaster sieboldii, as known as golden-ringed dragonfly, jumbo dragonfly, Siebold's dragonfly or oniyanma (オニヤンマ、鬼蜻蜓、馬大頭) in Japanese, 無霸勾蜓 in Chinese and 장수잠자리('jang-su-jamjari', means 'General Dragonfly') in Korean is the largest species of...



Anteos clorinde

Anteos clorinde

Anteos clorinde, the white angled-sulphur, is a butterfly of the family Pieridae. The species was originally described by Jean-Baptiste Godart in 1824.



Anthene amarah

Anthene amarah

Anthene amarah, the black-striped hairtail, leaden hairtail or leaden ciliate blue, is a butterfly of the family Lycaenidae. It is found in tropical Africa and Arabia (up to Aqaba). The habitat consists of savanna and occasionally open areas in...



Antheraea polyphemus

Antheraea polyphemus

Antheraea polyphemus, the Polyphemus moth, is a North American member of the family Saturniidae, the giant silk moths. It is a tan-colored moth, with an average wingspan of 15 cm (6 in). The most notable feature of the moth is its large, purplish...



Anthidium florentinum

Anthidium florentinum

Anthidium florentinum, one of several European wool carder bees, is a territorial species of bee in the family Megachilidae, the leaf-cutter, carder, or mason bees.



Anthocharis cardamines

Anthocharis cardamines

Anthocharis cardamines, the orange tip, is a butterfly in the family Pieridae, which contains about 1,100 species. A. cardamines is mainly found throughout Europe and temperate Asia (Palearctic) The males feature wings with a signature orange...



Anthonomus pomorum

Anthonomus pomorum

Anthonomus pomorum or the apple blossom weevil is a univoltine herbivore of apple trees, Malus domestica.



Anthonomus rubi

Anthonomus rubi

Anthonomus rubi, the strawberry-blossom weevil or strawberry blossom weevil, is a species of weevil found in Europe and western Asia. It feeds on plants of the family Rosaceae and is an important pest of strawberry (Fragaria × ananassa Duchesne)...



Apantesis arge

Apantesis arge

Apantesis arge, the arge moth or arge tiger moth, is a moth of the family Erebidæ. The species was first described by Dru Drury in 1773. It is found in North America from Quebec and Maine to Florida, West to New Mexico, North to North Dakota and...



Apantesis phalerata

Apantesis phalerata

Apantesis phalerata, also known as the harnessed tiger moth, is a moth species of the family Erebidæ. It was first described by Thaddeus William Harris in 1841.



Apatura ilia

Apatura ilia

Apatura ilia, the lesser purple emperor, is a species of butterfly native to most of Europe and east across the Palearctic. It is named for its similarity to the purple emperor butterfly.



Apatura iris

Apatura iris

Apatura iris, the purple emperor, is a Palearctic butterfly of the family Nymphalidæ.



Aphomia sociella

Aphomia sociella

Aphomia sociella, also known as the bee moth and the bumble bee wax moth, is a small moth of the family Pyralidæ (snout moths) and subfamily Galleriinae. Its body and forewings are typically reddish brown, tan, or dark green in color and females...



Aphrissa statira

Aphrissa statira

Aphrissa statira, the statira sulphur, is a species of Lepidoptera in the family Pieridae. The species is a medium-sized yellow butterfly, with females more pale than males. They are found from southern regions of Florida and Texas through...



Aphytis melinus

Aphytis melinus

Aphytis melinus is an internal parasite of the California red scale, Aonidiella aurantii, which is a pest of citrus in California and elsewhere. This wasp drums its antennae against the scale insect to find out if it is healthy, if it is already...



Apis andreniformis

Apis andreniformis

Apis andreniformis, or the black dwarf honey bee, is a relatively rare species of honey bee whose native habitat is the tropical and subtropical regions of Southeast Asia. A. andreniformis was the fifth honey bee species to be described of the...



Apis cerana

Apis cerana

Apis cerana, the eastern honey bee, Asiatic honey bee or Asian honey bee, is a species of honey bee native to South, Southeast and East Asia. This species is the sister species of Apis koschevnikovi and both are in the same subgenus as the...



Apis dorsata

Apis dorsata

Apis dorsata, the rock bee or giant honey bee, is a honey bee of South and Southeast Asia. They are typically around 17–20 mm (0.7–0.8 in) long and nests are mainly built in exposed places far off the ground, like on tree limbs, under cliff...



Apis florea

Apis florea

The dwarf honey bee (or red dwarf honey bee), *Apis florea*, is one of two species of small, wild honey bees of southern and southeastern Asia. It has a much wider distribution than its sister species, *Apis andreniformis*. First identified in the...



Apis mellifera cecropia

Apis mellifera cecropia

Apis mellifera cecropia, the Greek bee, is the subspecies of honey bee that is native to southern Greece. It is found also in southern Albania, and it is very similar to Apis mellifera ligustica, the Italian bee. it is favored for its extreme...



Apis mellifera iberiensis

Apis mellifera iberiensis

Apis mellifera iberiensis, or the Spanish bee, is a western honey bee subspecies native to the Iberian Peninsula. It is also found on the Balearic Islands.



Apis mellifera lamarckii

Apis mellifera lamarckii

Lamarck's honey bee or the Egyptian honey bee, *Apis mellifera lamarckii*, is a subspecies of honey bee occurring in a narrow range along the Egyptian Nile Valley of Egypt and Sudan, named after Jean-Baptiste Lamarck and is considered the first...



Apis nigrocincta

Apis nigrocincta

Apis nigrocincta is a species of honey bee that inhabits the Philippine island of Mindanao as well as the Indonesian islands of Sangihe and Sulawesi. The species is known to have queens with the highest mating frequencies of any species of the...



Apodemia mormo

Apodemia mormo

Apodemia mormo (Mormon metalmark) is a principally Nearctic butterfly in the family Riodinidae. It is a particularly fascinating species for ecological and evolutionary research, as evidenced by its shifting taxonomic classifications. Conflicting...



Apodemia mormo langei

Apodemia mormo langei

Apodemia mormo langei, the Lange's metalmark butterfly, is an endangered North American butterfly. It is a subspecies of the Mormon metalmark and belongs to the family Riodinidae. The butterfly is endemic to California, where it is known from one...



Apollo

Parnassius apollo

The Apollo or mountain Apollo (Parnassius apollo), is a butterfly of the family Papilionidae.



Aporia crataegi

Aporia crataegi

Aporia crataegi, the black-veined white, is a large butterfly of the family Pieridae. A. crataegi is widespread and common. Its range extends from northwest Africa in the west to Transcaucasia and across the Palearctic to Siberia and Japan in the...



Appias albina

Appias albina

Appias albina, the common albatross, is a small butterfly of the family Pieridae. It is found in south and southeast Asia to Australia.



Appias libythea

Appias libythea

Appias libythea, the striped albatross, is a small butterfly of the family Pieridae, that is, the yellows and whites, which is found in south and southeast Asia.



Appias lyncida

Appias lyncida

Appias lyncida, the chocolate albatross, is a butterfly of the family Pieridae, that is, the yellows and whites, which is found in south and southeast Asia.



Appias nero

Appias nero

Appias nero, the orange albatross, is a butterfly of the family Pieridae, that is, the yellows and whites. The species occurs from northern India to the Sunda Islands, the Philippines, Sulawesi and eastwards. Subspecies Appias nero galba is found...



Apple maggot

Rhagoletis pomonella

The apple maggot (Rhagoletis pomonella), also known as the railroad worm (but distinct from the Phrixothrix beetle larva, also called railroad worm), is a species of fruit fly, and a pest of several types of fruits, especially apples. This...



Aptera fusca

Aptera fusca

Aptera fusca, the Cape Mountain cockroach, giant cockroach or Table Mountain cockroach, is a large cockroach which is widespread on low vegetation in open areas in the fynbos biome of the Western Cape region of South Africa. Adult females can be...



Arachnocampa luminosa

Arachnocampa luminosa

Arachnocampa luminosa (Skuse, 1891), commonly known as New Zealand glowworm or simply glowworm, is a species of fungus gnat endemic to New Zealand. The larval stage and the imago produce a blue-green bioluminescence. The species is known to dwell...



Archaeoprepona demophon

Archaeoprepona demophon

Archaeoprepona demophon, the one-spotted prepona, banded king shoemaker, or demophon shoemaker is a butterfly belonging to the family Nymphalidae.



Archimantis latistyla

Archimantis latistyla

Archimantis latistyla, commonly known as the large brown mantis (also known as the bunny mantis or Australian ghost hopper mantis) is a species of mantid native to Australia. The large brown mantis has two subspecies, a widespread subspecies and...



Archimantis monstrosa

Archimantis monstrosa

Archimantis monstrosa is a species of mantid in the family Mantidae. A. monstrosa, or monster mantis, commonly reaches a length of 90mm or more. It is less common than the often-seen large brown mantis. A. monstrosa that live near the coast can...



Arctia plantaginis

Arctia plantaginis

Arctia plantaginis, the wood tiger, is a moth of the family Erebidae. Several subspecies are found in the Holarctic ecozone south to Anatolia, Transcaucasus, northern Iran, Kazakhstan, Mongolia, China, Korea and Japan. One subspecies is endemic...



Arctia villica

Arctia villica

Arctia villica, the cream-spot tiger, is a moth of the family Erebidae. The species was first described by Carl Linnaeus in his 1758 10th edition of Systema Naturae. It is distributed from the Iberian Peninsula, Anatolia, western and northern...



Arenivaga investigata

Arenivaga investigata

Arenivaga investigata, also known as the desert cockroach, is a member of the cockroach family Corydiidae.



Arge pagana

Arge pagana

Arge pagana is a sawfly in the family Argidae. It is known by the name "large rose sawfly" although the related species Arge ochropus is also known by this name.



Argema mimosae

Argema mimosae

Argema mimosae, the African moon moth, is a giant silk moth of the family Saturniidae. Similar in appearance to the giant Madagascan moon moth (Argema mittrei), but smaller, this moth can be found widely in Eastern Africa and more locally in...



Argentine ant

Linepithema humile

The Argentine ant (Linepithema humile, formerly Iridomyrmex humilis) is an ant native to northern Argentina, Uruguay, Paraguay, Bolivia and southern Brazil. This invasive species was inadvertently introduced by humans on a global scale and has...



Argina astrea

Argina astrea

Argina astrea, the crotalaria podborer, is a moth of the family Erebidæ. The species was first described by Dru Drury in 1773. It is found in eastern Africa, southern Asia of India, Sri Lanka, and Indo-Australia, including the Pacific Islands...



Argynnis hyperbius

Argynnis hyperbius

The Indian fritillary (Argynnis hyperbius) is a species of butterfly of the nymphalid or brush-footed family. It is usually found from south and southeast Asia to Australia.



Argynnis laodice

Argynnis laodice

Argynnis laodice, Pallas' fritillary, is a butterfly of the family Nymphalidae. It occurs in damp forested places in southern Scandinavia, Eastern Europe, northwestern Kazakhstan, and across the Palearctic to Siberia, Amur, Korea and Japan.



Argynnis pandora

Argynnis pandora

Argynnis pandora, the cardinal, is a butterfly of the family Nymphalidae. It is common throughout southern Europe and is also found in northern Africa and the Middle east and then east across the Palearctic to the Tian-Shan and northwestern India.



Ariadne merione

Ariadne merione

Ariadne merione, the common castor, is an orange butterfly with brown lines whose larvae feed almost exclusively on castor (Ricinus communis). It is similar in appearance to Ariadne ariadne, the angled castor. This species is found in South and...



Aricia agestis

Aricia agestis

Aricia agestis, the brown argus, is a butterfly in the family Lycaenidae. It is found throughout the Palearctic realm, north to northern Jutland (Denmark) and east to Siberia and the Tian Shan.



Arna pseudoconspersa

Arna pseudoconspersa

Arna pseudoconspersa, the tea tussock moth or Japanese browntail moth, is a moth of the family Erebiidae. The species was first described by Embrik Strand in 1914. It is found in Japan, Taiwan, Korea, China, Vietnam and India.Both the larvae and...



Arocatus roeselii

Arocatus roeselii

Arocatus roeselii is a species of lygaeid bug.



Ascalapha odorata

Ascalapha odorata

The erebid moth Ascalapha odorata, commonly known as the black witch, is a large bat-shaped, dark-colored nocturnal moth, normally ranging from the southern United States to Brazil. Ascalapha odorata is also migratory into Canada and most states...



Ash borer

Podosesia syringae

The ash borer (*Podosesia syringae*), or lilac borer, is a clearwing moth in the family Sesiidae. It is found throughout North America and can be a pest of ash and lilac.



Asian giant hornet

Vespa mandarinia

The Asian giant hornet (*Vespa mandarinia*) or northern giant hornet, including the color form referred to as the Japanese giant hornet, is the world's largest hornet. It is native to temperate and tropical East Asia, South Asia, Mainland Southeast...



Asian hornet

Vespa velutina

The Asian hornet (*Vespa velutina*), also known as the yellow-legged hornet or Asian predatory wasp, is a species of hornet indigenous to Southeast Asia. It is of concern as an invasive species in some other countries.



Asian long-horned beetle

Anoplophora glabripennis

The Asian long-horned beetle (*Anoplophora glabripennis*), also known as the starry sky, sky beetle, or ALB, is native to the Korean Peninsula, northern and southern China, and disputably in northern Japan. This species has now been accidentally...



Asota caricae

Asota caricae

Asota caricae, the tropical tiger moth, is a species of noctuid moth in the family Erebidae. It is found from the Indo-Australian tropics of India and Sri Lanka to Queensland and Vanuatu.



Asota heliconia

Asota heliconia

Asota heliconia is a moth in the family Erebidæ. It is found from the Indo-Australian tropics east to Queensland and the Solomons. The wingspan is 52 mm. Larvae have been recorded on Averrhoa species.



Asota plagiata

Asota plagiata

Asota plagiata, the two-spots tiger moth, is a moth of the family Erebidae first described by Francis Walker in 1854. It is found in the northern half of Australia. The wingspan is 49–58 mm. The larvae feed on Ficus macrophylla.



Atalopedes campestris

Atalopedes campestris

Atalopedes campestris (called sachem in the United States and Canada) is a small grass skipper butterfly. It has a wingspan...



Athyma nefte

Athyma nefte

Athyma nefte, the colour sergeant, is a species of brush-footed butterfly found in tropical South and Southeast Asia.



Atta mexicana

Atta mexicana

Atta mexicana is a species of leaf-cutter ant, a New World ant of the subfamily Myrmicinae of the genus Atta. This species is from one of the two genera of advanced attines (fungus-growing ants) within the tribe Attini.



Atta sexdens

Atta sexdens

Atta sexdens is a species of leafcutter ant belonging to the tribe Attini, native to America, from the southern United States (Texas) to northern Argentina. They are absent from Chile. They cut leaves to provide a substrate for the fungus farms...



Attacus atlas

Attacus atlas

Attacus atlas, the Atlas moth, is a large saturniid moth endemic to the forests of Asia. The species was described by Carl Linnaeus in his 1758 10th edition of Systema Naturae. The Atlas moth is one of the largest lepidopterans, with a wingspan...



Australian cockroach

Periplaneta australasiae

The Australian cockroach (Periplaneta australasiae) is a common species of tropical cockroach, with a length of 23–35 mm (0.91–1.38 in). It is brown overall, with the tegmina having a conspicuous lateral pale stripe or margin, and the pronotum...



Australian hornet

Abispa ephippium

The Australian hornet (Abispa ephippium) is not a true hornet, it is a type of potter wasp or "mason wasp", is a vespid native to the Australian states and territories of the Australian Capital Territory, New South Wales, Northern Territory...



Austrophlebia costalis

Austrophlebia costalis

Austrophlebia costalis, the southern giant darner, is a species of dragonfly in the family Telephlebiidae endemic to eastern Australia. Austrophlebia costalis is an enormous dark dragonfly with strong yellow markings on its body and a brown band...



Autographa precationis

Autographa precationis

Autographa precationis (common looper moth) is a moth of the family Noctuidae. It is found in eastern and central North America. The wingspan is 30–38 mm. Adults are on wing from April to October depending on the location. There are three or more...



Automeris io

Automeris io

Automeris io, the Io moth (EYE-oh) or peacock moth, is a colorful North American moth in the family Saturniidae. The Io moth is also a member of the subfamily Hemileucinae. The name Io comes from Greek mythology in which Io was a mortal lover of...



Automeris metzli

Automeris metzli

Automeris metzli is a moth of the family Saturniidae. It is found from Mexico to Venezuela, Colombia and Ecuador and can also be found in Trinidad. The larvae feed on Quercus species.



Autumnal moth

Epirrita autumnata

The autumnal moth (Epirrita autumnata) is a moth of the family Geometridae. The species was first described by Moritz Balthasar Borkhausen in 1794. It is found throughout the Palearctic region and the Near East and has a much wider distribution...



Axylia putris

Axylia putris

The flame (Axylia putris) is a moth of the family Noctuidae. It is found throughout Europe then east across the Palearctic to Armenia, western Siberia and Amur, Korea and Japan. The range extends into northern India. This species has creamy-buff...



Azure damselfly

Coenagrion puella

The azure damselfly (Coenagrion puella) is a species of damselfly found in most of Europe. It is notable for its distinctive black and blue colouring. They are commonly found around ponds and lakesides during the summer.



Azure hawker

Aeshna caerulea

The azure hawker (Aeshna caerulea) is one of the smaller species of hawker dragonflies, (family Aeshnidae). The flight period is from late May to August. It is about 62 mm long. Both sexes have azure blue spots on each abdominal segment and the...



Bacillus rossius

Bacillus rossius

The European stick insect (Bacillus rossius) also called the European stick bug, European walking stick, or Mediterranean stick insect, is a species of stick insect, common in southern Europe. The species is endemic to the northwestern...



Bactrocera correcta

Bactrocera correcta

Bactrocera correcta is a species of tephritid fruit flies that is widely distributed in Southeast Asia. It is a serious pest species with a broad host range and has caused major infestations in Vietnam and Thailand.



Bactrocera tryoni

Bactrocera tryoni

The Queensland fruit fly (Bactrocera tryoni) is a species of fly in the family Tephritidae in the insect order Diptera. B. tryoni is native to subtropical coastal Queensland and northern New South Wales. They are active during the day, but mate...



Badamia exclamationis

Badamia exclamationis

Badamia exclamationis, commonly known as the brown awl or narrow-winged awl, is a butterfly belonging to the family Hesperidae. It is found in south and southeast Asia, Australia, and Oceania.



Banded demoiselle

Calopteryx splendens

The banded demoiselle (Calopteryx splendens) is a species of damselfly belonging to the family Calopterygidae. It is often found along slow-flowing streams and rivers. It is a Eurasian species occurring from the Atlantic coast eastwards to Lake...



Banded sugar ant

Camponotus consobrinus

The banded sugar ant (Camponotus consobrinus), also known as the sugar ant, is a species of ant native to Australia. A member of the genus Camponotus in the subfamily Formicinae, it was described by German entomologist Wilhelm Ferdinand Erichson...



Baronia

Baronia brevicornis

Baronia brevicornis, commonly known as the short-horned baronia, is a species of butterfly in the monotypic genus Baronia and is placed in a subfamily of its own, the Baroniinae, a sister group of the remainder of the swallowtail butterflies. It...



Battus philenor

Battus philenor

Battus philenor, the pipevine swallowtail or blue swallowtail, is a swallowtail butterfly found in North America and Central America. This butterfly is black with iridescent-blue hindwings. They are found in many different habitats, but are most...



Beautiful demoiselle

Calopteryx virgo

The beautiful demoiselle (Calopteryx virgo) is a species of damselfly belonging to the family Calopterygidae. It is found in Europe, North Africa, and Western Asia. It is often found along fast-flowing waters.



Beech blight aphid

Grylloprociphilus imbricator

The beech blight aphid (Grylloprociphilus imbricator) is a small insect in the order Hemiptera that feed primarily on the sap of American beech trees. The aphids form dense colonies on small branches and the undersides of leaves. A secondary...



Beet armyworm

Spodoptera exigua

The beet armyworm or small mottled willow moth (*Spodoptera exigua*) is one of the best-known agricultural pest insects. It is also known as the asparagus fern caterpillar. It is native to Asia, but has been introduced worldwide and is now found...



Belenois aurota

Belenois aurota

Belenois aurota, the pioneer or pioneer white or caper white, is a small to medium-sized butterfly of the family Pieridae, that is, the yellows and whites, which is found in South Asia and Africa. In Africa, it is also known as the brown-veined...



Belenois java

Belenois java

Belenois java, the caper white or common white, is a small butterfly of the family Pieridae found in Australia, Indonesia, and Melanesia. It is highly migratory and is often confused with the cabbage white (*Pieris rapae*).



Belgica antarctica

Belgica antarctica

Belgica antarctica, the Antarctic midge, is a species of flightless midge, endemic to the continent of Antarctica. At 2–6 mm (0.08–0.2 in) long, it is the largest purely terrestrial animal native to the continent. It also has the smallest known...



Bhutanitis lidderdalii

Bhutanitis lidderdalii

Bhutanitis lidderdalii, the Bhutan glory, is a species of swallowtail butterfly (family Papilionidae), which is found in Bhutan, parts of northeastern India and of Southeast Asia. A spectacular insect much sought after by collectors, the species...



Bibasis gomata

Bibasis gomata

Bibasis gomata, commonly known as the pale green awlet, is a butterfly belonging to the family Hesperiidae. It is found in Northeast India, the Western Ghats and parts of Southeast Asia. The butterfly was reassigned to genus Burara by Vane-Wright...



Bill Gates' flower fly

Eristalis gatesi

Bill Gates' flower fly (Eristalis gatesi) is a flower fly endemic to Costa Rica named after Bill Gates. Another fly found in similar habitats was named after Gates' associate Paul Allen, called Paul Allen's flower fly (Eristalis alleni);...



Blaberus discoidalis

Blaberus discoidalis

Blaberus discoidalis, commonly known as the discoid cockroach, tropical cockroach, West Indian leaf cockroach, false death's head cockroach, Haitian cockroach, and drummer, is a cockroach native to Central America of the “giant cockroach” family,...



Blaberus giganteus

Blaberus giganteus

Blaberus giganteus, the Central American giant cave cockroach or Brazilian cockroach, is a cockroach belonging to the family Blaberidae. One of the world's largest cockroaches, it is native to the warm parts of the Neotropical realm.



Black arches

Lymantria monacha

The black arches or nun moth (Lymantria monacha) is a small Palearctic moth. It is considered a forest pest.



Black carpet beetle

Attagenus unicolor

The black carpet beetle (*Attagenus unicolor*) is a 3–5-millimetre-long (0.12–0.20 in) beetle that can be a serious household pest. The larvae grow to 7 mm (0.28 in) in length, are reddish brown in colour and covered with bristles. The larval form...



Black garden ant

Lasius niger

The black garden ant (*Lasius niger*), also known as the common black ant, is a formicine ant, the type species of the subgenus *Lasius*, which is found across Europe and in some parts of North America, South America, Asia and Australasia. The...



Black hairstreak

Satyrrium pruni

The black hairstreak (*Satyrrium pruni*) is a butterfly in the family Lycaenidae.



Black imported fire ant

Solenopsis richteri

The black imported fire ant (*Solenopsis richteri*), or simply BIFA, is a species of ant in the genus *Solenopsis* (fire ants). It was long thought to either be a subspecies or a color variation of *Solenopsis invicta* (the red imported fire ant, or...



Black saddlebags

Tamea lacerata

The black saddlebags (*Tamea lacerata*) is a species of skimmer dragonfly found throughout North America. It has distinctive wings with characteristic black blotches at their proximal ends, which make the dragonfly look as though it is wearing...



Black-tailed skimmer

Orthetrum cancellatum

The black-tailed skimmer (Orthetrum cancellatum) is a dragonfly belonging to the family Libellulidae.



Blattella asahinai

Blattella asahinai

The Asian cockroach (Blattella asahinai) is a species of cockroach that was first described in 1981 from insects collected on Okinawa Island, Japan. It is a small species of cockroach,...



Blera fallax

Blera fallax

Blera fallax, the pine hoverfly or roodkapje, is a rare species of hoverfly normally associated with mature pine trees in Northern and Central Europe.



Blissus leucopterus

Blissus leucopterus

Blissus leucopterus, also known as the true chinch bug, is a small North American insect in the order Hemiptera and family Blissidae. It is the most commonly encountered species of the genus Blissus, which are all known as chinch bugs. A closely...



Blood-vein

Timandra comae

The blood-vein (Timandra comae) is a moth of the family Geometridae. The species was first described by Anton Schmidt in 1931.



Bloody-nosed beetle

Timarcha tenebricosa

The bloody-nosed beetle (Timarcha tenebricosa), also called blood spewer or blood-spewing beetle, is a leaf beetle native to Europe.



Blue ant

Diamma bicolor

The blue ant (Diamma bicolor), also known as the blue-ant or bluebottle, is a species of flower wasp in the family Thynnidae. It is the sole member of the genus Diamma and of the subfamily Diamminae. Despite its common name and wingless body, it...



Blue riverdamsel

Pseudagrion microcephalum

The blue riverdamsel, Pseudagrion microcephalum is a common species of damselfly in the family Coenagrionidae. It is also known as the blue sprite and blue grass dart.



Blue-eyed darner

Rhionaeschna multicolor

The blue-eyed darner (Rhionaeschna multicolor, syn. Aeshna multicolor) is a common dragonfly of the family Aeshnidae; native to the western United States, it is commonly sighted in the sagebrush steppe of the Snake River Plain, occurring east to...



Blue-tailed damselfly

Ischnura elegans

The blue-tailed damselfly or common bluetail (Ischnura elegans) is a damselfly, belonging to the family Coenagrionidae.



Body louse

Pediculus humanus humanus

The body louse (Pediculus humanus humanus, also known as Pediculus humanus corporis) is a hematophagic ectoparasite louse that infests humans. It is one of three lice which infest humans, the other two being the head louse, and the crab louse or...



Bogong moth

Agrotis infusa

The bogong moth (Agrotis infusa) is a temperate species of night-flying moth, notable for its biannual long-distance seasonal migrations towards and from the Australian Alps, similar to the diurnal monarch butterfly. During the autumn and winter...



Boll weevil

Anthonomus grandis

The boll weevil (Anthonomus grandis) is a species of beetle in the family Curculionidae. The boll weevil feeds on cotton buds and flowers. Thought to be native to Central Mexico, it migrated into the United States from Mexico in the late 19th...



Boloria titania

Boloria titania

Boloria titania, the Titania's fritillary or purple bog fritillary, is a butterfly of the subfamily Heliconiinae of the family Nymphalidae.



Bombus auricomus

Bombus auricomus

Bombus auricomus is a species of bumblebee known by the common name black and gold bumblebee. It is native to eastern North America, including Ontario in Canada and much of the eastern United States, as far west as the Great Plains.This species...



Bombus balteatus

Bombus balteatus

Bombus balteatus, the golden-belted bumble bee, is a species of bumblebee with a boreal and high altitude distribution in northern Eurasia and North America.



Bombus bohemicus

Bombus bohemicus

Bombus bohemicus, also known as the gypsy's cuckoo bumblebee, is a species of socially parasitic cuckoo bumblebee found in most of Europe with the exception of the southern Iberian Peninsula and Iceland. B. bohemicus practices inquilinism, or...



Bombus campestris

Bombus campestris

Bombus campestris is a very common cuckoo bumblebee found in most of Europe.



Bombus citrinus

Bombus citrinus

Bombus citrinus is a species of bumblebee known commonly as the lemon cuckoo bumblebee due to its lemon-yellow color. It is native to eastern North America. This is a cuckoo bumblebee, one that invades the colonies of other bumblebees, kills the...



Bombus cryptarum

Bombus cryptarum

Bombus cryptarum is a species of bumblebee. It is native to the northern hemisphere, where it is "one of the most widespread bumblebees in the world." It occurs throughout Europe, Asia, and western North America. It is known commonly as the...



Bombus distinguendus

Bombus distinguendus

Bombus distinguendus, the great yellow bumblebee, is a species of bumblebee found in Austria, Belgium, the Czech Republic, Denmark, Finland, France, Germany, Great Britain, Hungary, Ireland, Lithuania, Poland, Romania, Slovakia, Switzerland,...



Bombus fraternus

Bombus fraternus

Bombus fraternus is an endangered species of bumblebee known commonly as the Southern Plains bumblebee. It is native to the United States east of the Rocky Mountains. It is most often encountered in the southern Great Plains and along the Gulf...



Bombus hortorum

Bombus hortorum

Bombus hortorum, the garden bumblebee or small garden bumblebee, is a species of bumblebee found in most of Europe north to 70°N, as well as parts of Asia and New Zealand. It is distinguished from most other bumblebees by its long tongue used for...



Bombus hypnorum

Bombus hypnorum

The tree bumblebee or new garden bumblebee (Bombus hypnorum) is a species of bumblebee common in the European continent and parts of Asia. Since the start of the twenty-first century, it has spread to Great Britain. These bumblebees prefer...



Bombus lapidarius

Bombus lapidarius

Bombus lapidarius is a species of bumblebee in the subgenus Melanobombus. Commonly known as the red-tailed bumblebee, B. lapidarius can be found throughout much of Central Europe. Known for its distinctive black and red body, this social bee is...



Bombus monticola

Bombus monticola

Bombus monticola, the bilberry bumblebee, blaeberry bumblebee or mountain bumblebee, is a species of bumblebee found in Europe.



Bombus nevadensis

Bombus nevadensis

Bombus nevadensis, the Nevada bumblebee, is a species of bumblebee. It is native to North America, where it occurs from Alaska to California in the west, and east to Wisconsin, and in Arizona, New Mexico, and Mexico.



Bombus occidentalis

Bombus occidentalis

Bombus occidentalis, the western bumblebee, is one of around 30 bumblebee species present in the western United States and western Canada. A recent review of all of its close relatives worldwide appears to have confirmed its status as a separate...



Bombus pensylvanicus

Bombus pensylvanicus

Bombus pensylvanicus, the American bumblebee, is a threatened species of bumblebee native to North America. It occurs in eastern Canada, throughout much of the Eastern United States, and much of Mexico. Once the most prevalent bumblebee in the...



Bombus polaris

Bombus polaris

Bombus polaris is a common Arctic bumblebee species. B. polaris is one of two bumblebees that live above the Arctic Circle. The other is its social parasite Bombus hyperboreus. B. polaris is a social bee that can survive at near freezing...



Bombus rupestris

Bombus rupestris

Bombus rupestris is a species of cuckoo bumblebee present in most of Europe except Iceland. In the Balkans it is found in montane and alpine habitats northwards from Central Greece. It is also found in Turkey.



Bombus suckleyi

Bombus suckleyi

Bombus suckleyi is a species of bumblebee known commonly as Suckley's cuckoo bumblebee, named after biologist George Suckley. Suckley's bumble bee is a generalist pollinator and represents a rare group of obligate, parasitic bumble bees (cuckoo...



Bombus sylvarum

Bombus sylvarum

Bombus sylvarum, the shrill carder bee or knapweed carder-bee, is a species of bumblebee with a wide distribution across Europe, east to the Ural Mountains, and north to Great Britain, Ireland, and southern Scandinavia.



Bombus ternarius

Bombus ternarius

Bombus ternarius, commonly known as the orange-belted bumblebee or tricolored bumblebee, is a yellow, orange and black bumblebee. It is a ground-nesting social insect whose colony cycle lasts only one season, common throughout the northeastern...



Bombus terrestris

Bombus terrestris

Bombus terrestris, the buff-tailed bumblebee or large earth bumblebee, is one of the most numerous bumblebee species in Europe. It is one of the main species used in greenhouse pollination, and so can be found in many countries and areas where it...



Bombus vestalis

Bombus vestalis

Bombus vestalis, the vestal cuckoo bumblebee, is a species of cuckoo bumblebee that lives in most of Europe, as well as North Africa and western Asia. It is a brood parasite that takes over the nests of other bee species. Its primary host is...



Bombylius major

Bombylius major

Bombylius major (commonly named the large bee-fly, the dark-edged bee-fly or the greater bee fly) is a parasitic bee mimic fly. B. major is the most common type of fly within the Bombylius genus. The fly derives its name from its close...



Bombyx mandarina

Bombyx mandarina

Bombyx mandarina, the wild silk moth, is a species of moth in the family Bombycidae. It is the closest relative of Bombyx mori, the domesticated silk moth. The silkworm is the larva or caterpillar of a silk moth. Unlike the domesticated relative...



Bombyx mori

Bombyx mori

The domestic silk moth (Bombyx mori) is an insect from the moth family Bombycidae. It is the closest relative of Bombyx mandarina, the wild silk moth. The silkworm is the larva (or caterpillar) of a silk moth. The silkworm is of particular...



Borbo bevani

Borbo bevani

Borbo bevani, the Beavan's swift, is a butterfly belonging to the family Hesperidae. It is named after Captain Robert Cecil Beavan. It is found throughout India.



Borbo cinnara

Borbo cinnara

Borbo cinnara, commonly known as the rice swift, Formosan swift or rice leaf folder, is a butterfly belonging to the family Hesperidae. It is found in Sri Lanka, India, Myanmar, Vietnam, Cambodia, Taiwan, and Australia.



Boxelder bug

Boisea trivittata

The boxelder bug (Boisea trivittata), also called box bug, maple bug or, inaccurately, box beetle, is a species of true bug native to eastern North America. The western boxelder bug Boisea rubrolineata is a relative of this species and is native...



Brachygastra mellifica

Brachygastra mellifica

Brachygastra mellifica, commonly known as the Mexican honey wasp, is a neotropical social wasp. It can be found in North America. B. mellifica is one of few wasp species that produces honey. It is also considered a delicacy in some cultures in...



Brahmaea certhia

Brahmaea certhia

Brahmaea certhia, the Sino-Korean owl moth, is a moth from the family Brahmaeidae, the Brahmin moths. It is found in the Korean Peninsula and China. The wingspan is 100 mm (3.9 in) to 120 mm (4.7 in). The larvae feed on privet, Fraxinus mandshurica...



Brenthis ino

Brenthis ino

Brenthis ino, the lesser marbled fritillary, is a butterfly of the family Nymphalidae.



Bright-line brown-eye

Lacanobia oleracea

The bright-line brown-eye (Lacanobia oleracea) is a moth of the family Noctuidae. The species was first described by Carl Linnaeus in his 1758 10th edition of Systema Naturae. It is a common species throughout Europe, but is also found in North...



Brimstone moth

Opisthograptis luteolata

The brimstone moth (Opisthograptis luteolata) is a moth of the family Geometridae. The species was first described by Carl Linnaeus in his 1758 10th edition of Systema Naturae. It should not be confused with the brimstone butterfly Gonepteryx...



Brown cockroach

Periplaneta brunnea

The brown cockroach (Periplaneta brunnea) is a species of cockroach in the family Blattidae. It is probably originally native to Africa, but today it has a circumtropical distribution, having been widely introduced. In cooler climates it can only...



Brown hairstreak

Thecla betulae

The brown hairstreak (Thecla betulae) is a butterfly in the family Lycaenidae. The range includes most of the Palearctic.



Brown hawker

Aeshna grandis

The brown hawker (Aeshna grandis) is a large dragonfly about 73 millimetres (2.9 in) long. It is a distinctive species and is easily recognised, even in flight, by its brown body and bronze wings. At rest, blue spots on the second and third...



Brown marmorated stink bug

Halyomorpha halys

The brown marmorated stink bug (Halyomorpha halys) is an insect in the family Pentatomidae, native to China, Japan, Korea, and other Asian regions. In September 1998, it was collected in Allentown, Pennsylvania, where it is believed to have been...



Brown planthopper

Nilaparvata lugens

The brown planthopper (BPH), Nilaparvata lugens (Stål) (Hemiptera: Delphacidae) is a planthopper species that feeds on rice plants (Oryza sativa L.). These insects are among the most important pests of rice, which is the major staple crop for...



Brown-banded cockroach

Supella longipalpa

The brown-banded cockroach (Supella longipalpa) is a species of small cockroach, measuring about 10 to 14 mm long and the most well-known in the genus Supella. It is tan to light brown and has two light-colored bands across the wings and abdomen,...



Brown-tail moth

Euproctis chrysorrhoea

The brown-tail moth (Euproctis chrysorrhoea) is a moth of the family Erebidæ. It is native to Europe, neighboring countries in Asia, and the north coast of Africa. Descriptions of outbreaks, i.e., large population increases of several years...



Buck moth

Hemileuca maia

The buck moth (Hemileuca maia) is a common insect found in oak forests, stretching in the United States from peninsular Florida to New England, and as far west as Texas and Kansas. It was first described by Dru Drury in 1773. The larvae typically...



Buff ermine

Spilarctia luteum

The buff ermine (Spilarctia luteum) is a moth of the family Erebidæ. It is sometimes placed in the genus Spilosoma. The species was first described by Johann Siegfried Hufnagel in 1766. It is found throughout the temperate belt of the Palearctic...



Buff-tip

Phalera bucephala

The buff-tip (Phalera bucephala) is a moth of the family Notodontidæ. It is found throughout Europe and in Asia to eastern Siberia. The species was first described by Carl Linnaeus in his 1758 10th edition of Systema Naturæ.



Bunaea alcinoe

Bunaea alcinoe

Bunaea alcinoe, the cabbage tree emperor moth, is an African moth species belonging to the family Saturniidae. It was first described by Caspar Stoll in 1780.



Burnsius communis

Burnsius communis

Burnsius communis, the common checkered-skipper, formerly known as Pyrgus communis, is a species of butterfly in the family Hesperidae. It is known as the frequently seen Pyrginae species in the northern United States by collectors and watchers...



Byasa polyeuctes

Byasa polyeuctes

Byasa polyeuctes, the common windmill, is the most common member in India of the windmills genus (Byasa), comprising tailed black swallowtail butterflies with white spots and red submarginal crescents.



Cabbage looper

Trichoplusia ni

The cabbage looper (Trichoplusia ni) is a medium-sized moth in the family Noctuidae, a family commonly referred to as owlet moths. Its common name comes from its preferred host plants and distinctive crawling behavior. Cruciferous vegetables,...



Cabbage moth

Mamestra brassicae

The cabbage moth (Mamestra brassicae) is primarily known as a pest that is responsible for severe crop damage of a wide variety of plant species. The common name, cabbage moth, is a misnomer as the species feeds on many fruits, vegetables, and...



Cactoblastis cactorum

Cactoblastis cactorum

Cactoblastis cactorum, the cactus moth, South American cactus moth or nopal moth, is native to Argentina, Paraguay, Uruguay and southern Brazil. It is one of five species in the genus Cactoblastis that inhabit South America, where many...



California tortoiseshell

Nymphalis californica

The California tortoiseshell (Nymphalis californica) is a butterfly of the family Nymphalidae. The wings of the California tortoiseshell have ragged edges. The upper sides of the wings are orange with black spots and a wide black margin. The...



Caligo telamonius memnon

Caligo telamonius memnon

Caligo telamonius memnon, commonly known as the giant owl or pale owl, is a subspecies of butterfly of the family Nymphalidae. This subspecies can be found in rainforests and secondary forests from Mexico to the Amazon rainforest in South...



Calinaga buddha

Calinaga buddha

Calinaga buddha, the freak, is a species of butterfly in the Calinaginae subfamily that is found in parts of Southeast Asia, China and India.



Calliphora livida

Calliphora livida

Calliphora livida is a member of the family Calliphoridae, the blow flies. This large family includes the genus Calliphora, the "blue bottle flies". This genus is important in the field of forensic entomology because of its value in post-mortem...



Calliphora vicina

Calliphora vicina

Calliphora vicina is a member of the family Calliphoridae, which includes blow flies and bottle flies. These flies are important in the field of forensic entomology, being used to estimate the time of a person's death when a corpse is found and...



Callistege mi

Callistege mi

Callistege mi, the Mother Shipton moth, is a moth of the family Erebidæ. It was classified by Carl Alexander Clerck in 1759, and is also known under the name of Euclidia mi. In Finnish it is known as piirtoyökkönen and in German as Scheck-Tageule.



Callosobruchus maculatus

Callosobruchus maculatus

Callosobruchus maculatus is a species of beetles known commonly as the cowpea weevil or cowpea seed beetle. It is a member of the leaf beetle family, Chrysomelidae, and not a true weevil. This common pest of stored legumes has a cosmopolitan...



Calophasia lunula

Calophasia lunula

Calophasia lunula is a Palearctic species of noctuid moth known by the common names toadflax moth and toadflax brocade moth.



Calosoma sycophanta

Calosoma sycophanta

Calosoma sycophanta, the forest caterpillar hunter, is a species of ground beetle belonging to the family Carabidae.



Calyptra canadensis

Calyptra canadensis

Calyptra canadensis, the Canadian owlet or meadow rue owlet moth, is a moth of the family Erebidæ. The species was first described by Charles J. S. Bethune in 1865. It is found in North America from Nova Scotia to North Carolina in mountains,...



Calyptra thalictri

Calyptra thalictri

Calyptra thalictri is a moth of the family Erebidæ. It is native to the area ranging from Japan and Korea to China and Malaysia, west through the Urals to Southern Europe, but it has recently expanded its range to northern Europe. In 2000, it...



Campaea margaritata

Campaea margaritata

Campaea margaritata, commonly known in the UK as the light emerald, is a moth of the family Geometridæ. The species was first described by Carl Linnaeus in his 1767 12th edition of Systema Naturæ. It is widely distributed throughout Europe, the...



Campaea perlata

Campaea perlata

Campaea perlata, the pale beauty, is a species of moth in the family Geometridæ. The average wingspan is about 28–51 mm, and the female tends to be much larger than the male.



Camponotus compressus

Camponotus compressus

Camponotus compressus is a type of ground-nesting species of ant found in India and Southeast Asia. It is a frequent visitor to toilets as it consumes urea. It is one of the many species which tends plant-sap-sucking insects like aphids and tree...



Camponotus sericeus

Camponotus sericeus

Camponotus sericeus is a species of carpenter ant (genus Camponotus) widely distributed in the Afrotropical and oriental regions.



Cantharis fusca

Cantharis fusca

Cantharis fusca is a species of soldier beetle.C. fusca reaches a length of 10–15 millimetres (0.39–0.59 in). Except for parts of the head and thorax, which are red or orange, this species is completely black. The body is flat and long, with a...



Cape honey bee

Apis mellifera capensis

The Cape honey bee or Cape bee (Apis mellifera capensis) is a southern South African subspecies of the western honey bee. They play a major role in South African agriculture and the economy of the Western Cape by pollinating crops and producing...



Carabus auratus

Carabus auratus

Carabus auratus, the golden ground beetle, is a species of ground beetle in the genus Carabus. This species is native to central and western Europe and has been introduced into North America.



Carabus nemoralis

Carabus nemoralis

Carabus nemoralis (commonly called the "bronze carabid") is a ground beetle common in central and northern Europe, as well as Iceland and Canada. While native to Europe, it has been introduced to and is expanding its range throughout North America.



Cardinal beetle

Pyrochroa coccinea

The name cardinal beetle typically refers to three different species of beetles. The red-headed or 'common' cardinal beetle (Pyrochroa serraticornis) is a red to orange beetle with, as the name suggests, a red head. It is about 20 millimetres...



Carniolan honey bee

Apis mellifera carnica

The Carniolan honey bee (Apis mellifera carnica, Pollmann) is a subspecies of the western honey bee. The Carniolan honey bee is native to Slovenia, southern Austria, and parts of Albania, Croatia, Bosnia and Herzegovina, Montenegro, Serbia,...



Carolina mantis

Stagmomantis carolina

The Carolina mantis (Stagmomantis carolina) is a species of praying mantis of the subfamily Stagmomantinae. Sexual cannibalism occurs in roughly one quarter of all intersexual encounters of this species, though specimens of this species will...



Carrot fly

Chamaepsila rosae

The carrot fly (*Chamaepsila rosae*) is a pest of gardens and farms, and mainly affects the crop of carrots, but can also attack parsnips, parsley and celery. It is a member of the family Psilidae (order Diptera).



Castalius rosimon

Castalius rosimon

Castalius rosimon, the common Pierrot, is a small butterfly found in India that belongs to the lycaenids, or blues family.



Cat flea

Ctenocephalides felis

The cat flea (scientific name *Ctenocephalides felis*) is an extremely common parasitic insect whose principal host is the domestic cat, although a high proportion of the fleas found on dogs also belong to this species. This is despite the...



Catochrysops panormus

Catochrysops panormus

Catochrysops panormus, the silver forget-me-not, is a small butterfly found from India to the Philippines and south to Australia that belongs to the lycaenids or blues family. The species was first described by Cajetan Felder in 1860.



Catopsilia pomona

Catopsilia pomona

Catopsilia pomona, the common emigrant or lemon emigrant, is a medium-sized pierid butterfly found in Asia, Cambodia and parts of Australia. The species gets its name from its habit of migration. Some early authors considered them as two distinct...



Caucasian honey bee

Apis mellifera caucasia

The Caucasian honey bee (Apis mellifera caucasia) is a subspecies of the western honey bee.



Celastrina ladon

Celastrina ladon

Celastrina ladon, the spring azure or echo blue, is a butterfly of the family Lycaenidae. It is found in North America from Alaska and Canada south of the tundra, through most of the United States except the Texas coast, southern plain and...



Cepora nerissa

Cepora nerissa

Cepora nerissa, the common gull, is a small to medium-sized butterfly of the family Pieridae, that is, the yellows and whites, which is native to Sri Lanka, India, China, southeast Asia, and Indonesia.



Ceratitis capitata

Ceratitis capitata

Ceratitis capitata, commonly known as the Mediterranean fruit fly or medfly, is a yellow-and-brown fly native to sub-Saharan Africa. It has no near relatives in the Western Hemisphere and is considered to be one of the most destructive fruit...



Ceratomia amyntor

Ceratomia amyntor

Ceratomia amyntor, the elm sphinx or four-horned sphinx, is a North American moth in the family Sphingidae. The species was first described by Carl Geyer in 1835. It has a wingspan...



Ceratomia catalpae

Ceratomia catalpae

Ceratomia catalpae, the catalpa sphinx, is a hawk moth of the family Sphingidae. The species was first described by Jean Baptiste Boisduval in 1875. Other common names are the Catawba worm, or Catalpa sphinx.



Cereal leaf beetle

Oulema melanopus

The cereal leaf beetle (Oulema melanopus) is a significant crop pest, described by Carl Linnaeus in 1758.



Ceriagrion cerinorubellum

Ceriagrion cerinorubellum

Ceriagrion cerinorubellum, commonly known as the orange-tailed marsh dart or bi-coloured damsel, is a medium-sized damselfly in the family Coenagrionidae. It is a very common species of damselflies in Asia.



Cethosia cyane

Cethosia cyane

Cethosia cyane, the leopard lacewing, is a species of heliconiine butterfly found from India to southern China (southern Yunnan), and Indochina. Its range has expanded in the last few decades, and its arrival in the southern part of the Malay...



Cethosia nietneri

Cethosia nietneri

Cethosia nietneri, the Tamil lacewing, is a species of nymphalid butterfly found in Sri Lanka and south India. The species name is after John Nietner who obtained specimens of the butterfly from Ceylon from which it was described.



Cetonia aurata

Cetonia aurata

Cetonia aurata, called the rose chafer or the green rose chafer, is a...



Chalkhill blue

Lysandra coridon

The chalkhill blue (Lysandra coridon) is a butterfly in the family Lycaenidae. It is a small butterfly that can be found throughout the Palearctic realm, where it occurs primarily in grasslands rich in chalk. Males have a pale blue colour, while...



Charaxes brutus

Charaxes brutus

Charaxes brutus, the white-barred emperor or white-barred Charaxes, is a butterfly of the family Nymphalidae. It is found in Africa (see subspecies section for detailed information).Its flight period is year-round. Notes on the biology of brutus...



Charaxes candiope

Charaxes candiope

Charaxes candiope, the green-veined emperor or green-veined charaxes, is a butterfly of the family Nymphalidae. It is common in sub-Saharan Africa.



Charaxes varanes

Charaxes varanes

Charaxes varanes, the pearl emperor, Karkloof emperor, or pearl charaxes, is a butterfly of the family Nymphalidae, found in Africa from Saudi Arabia to South Africa. The wingspan is 65–70 millimetres (2.6–2.8 in) in males and 70–90 millimetres...



Chilades lajus

Chilades lajus

Chilades lajus, the lime blue, is a small butterfly found in India, Sri Lanka, Myanmar, Taiwan, Hong Kong, Hainan, Mangulam Island, Sulawesi and the Philippines that belongs to the lycaenids or blues family.



Chilades parrhasius

Chilades parrhasius

Chilades parrhasius, the small Cupid, is a small butterfly that belongs to the lycaenids or blues family. It is found in Nepal, southern Turan, southern Ghissar, Iran, Afghanistan, Pakistan, Sri Lanka, United Arab Emirates, Oman and southern,...



Chinese mantis

Tenodera sinensis

The Chinese mantis (Tenodera sinensis) is a species of mantis native to Asia and the nearby islands. In 1896, this species was accidentally introduced by a nursery tender at Mt. Airy near Philadelphia, United States. Tenodera sinensis often is...



Chlidonoptera lestoni

Chlidonoptera lestoni

Chlidonoptera lestoni is a species of praying mantis in the family Hymenopodidae.



Chloridea virescens

Chloridea virescens

Chloridea virescens, commonly known as the tobacco budworm, is a moth of the family Noctuidae found throughout the eastern and southwestern United States along with parts of Central America and South America.It is a major pest of field crops...



Choaspes benjaminii

Choaspes benjaminii

Choaspes benjaminii,also known as the Indian awlking or common awlking, is a species of butterfly in the family Hesperiiidae. The species is named after Benjamin Delessert and was described on the basis of a specimen collected by Adolphe Delessert...



Choeradodis rhomboidea

Choeradodis rhomboidea

Choeradodis rhomboidea, common names tropical shield mantis, hood mantis (or hooded mantis), and leaf mantis, is a South American species of praying mantis, restricted to the Amazon and Guiana Shield.



Choeradodis stalii

Choeradodis stalii

Choeradodis stalii is a species of praying mantis with common names that include tropical shield mantis, hooded mantis, and leaf mantis. It is found in Brazil, Ecuador, French Guiana, Panama, and Peru. As described by one insect-breeding hobbyist,...



Choristoneura fumiferana

Choristoneura fumiferana

Choristoneura fumiferana, the eastern spruce budworm, is a species of moth of the family Tortricidae native to the eastern United States and Canada. The caterpillars feed on the needles of spruce and fir trees. Eastern spruce budworm populations...



Chorthippus biguttulus

Chorthippus biguttulus

Chorthippus biguttulus, the bow-winged grasshopper, is one of the most common species of grasshopper found in the dry grassland of northern and central Europe. It is part of a group of species (biguttulus-group) that are very difficult to...



Chrysina resplendens

Chrysina resplendens

Chrysina resplendens is a scarab beetle found in mid-elevation forests in Costa Rica and western Panama. This beetle is typically shiny golden and has a length of 25–29 mm (0.98–1.14 in).



Chrysolina americana

Chrysolina americana

Chrysolina americana, common name rosemary beetle, is a species of beetle belonging to the family Chrysomelidae.



Chrysolina hyperici

Chrysolina hyperici

Chrysolina hyperici, the Saint John's wort beetle is a species of beetles of the family Chrysomelidae.



Chrysomya megacephala

Chrysomya megacephala

Chrysomya megacephala, more commonly known as the oriental latrine fly or oriental blue fly, is a member of the family Calliphoridae (blowflies). It is a warm-weather fly with a greenish-blue metallic box-like body. The fly infests corpses soon...



Chrysomya putoria

Chrysomya putoria

Chrysomya putoria, also known as the tropical African latrine blowfly, is a fly species belonging to the blowfly family, Calliphoridae. C. putoria is native to Africa and has recently spread to the Americas. These flies pose significant health...



Cicindela sexguttata

Cicindela sexguttata

The six-spotted tiger beetle, also known as the six-spotted green tiger beetle (Cicindela sexguttata), is a common North American species of Tiger beetles in the Cicindelinae subfamily. It is common in many areas of the states, and is well known....



Cimex lectularius

Cimex lectularius

Cimex lectularius is a species of Cimicidae. Its primary hosts are humans, and it is one of the world's major "nuisance pests".Although bed bugs can be infected with at least 28 human pathogens, no studies have found that the insects are capable...



Cinnabar moth

Tyria jacobaeae

The cinnabar moth (Tyria jacobaeae) is a brightly coloured arctiid moth found as a native species in Europe and western and central Asia then east across the Palearctic to Siberia to China. It has been introduced into New Zealand, Australia and...



Cirrochroa thais

Cirrochroa thais

Cirrochroa thais, also known as the Tamil yeoman, is a species of nymphalid butterfly found in forested areas of tropical Sri Lanka and India. It is the state insect of Tamil Nadu, an Indian state.



Cissusa spadix

Cissusa spadix

Cissusa spadix, the black-dotted brown moth, is a species of moth in the family Erebidae. The species is found from Ontario and Quebec, south through most of the United States, to Arizona and Georgia. The wingspan is about 35 mm.



Citheronia regalis

Citheronia regalis

Citheronia regalis, the regal moth or royal walnut moth, is a North American moth in the family Saturniidae. The caterpillars are called hickory horned devils. The adult (imago) has a wingspan of 3.75–6.1 in (9.5–15.5 cm). The species was first...



Citrine forktail

Ischnura hastata

The citrine forktail (Ischnura hastata) is a damselfly of the family Coenagrionidae.



Citrus long-horned beetle

Anoplophora chinensis

The citrus long-horned beetle (Anoplophora chinensis, also appearing in many sources as Anoplophora malasiaca) is a long-horned beetle native to Japan, China, Korea, Taiwan and Southeast Asia where it is considered a serious pest. This beetle has...



Clouded Apollo

Parnassius mnemosyne

The clouded Apollo (Parnassius mnemosyne) is a butterfly species of the family of swallowtail butterflies (Papilionidae) found in the Palearctic realm. Clouded Apollos inhabit meadows and woodland clearings with plenty of flowering plants, both in...



Coccinella novemnotata

Coccinella novemnotata

Coccinella novemnotata, the nine-spotted ladybug or nine-spotted lady beetle or C9, is a species of ladybug in the family Coccinellidae native to North America. This beetle was once ubiquitous across the continent but experienced a sharp and...



Coccinella septempunctata

Coccinella septempunctata

Coccinella septempunctata, the seven-spot ladybird (or, in North America, seven-spotted ladybug or "C-7"), is a carnivorous beetle native to the Old World and is the most common ladybird in Europe. Its elytra are of a red colour, but each...



Coccinella transversalis

Coccinella transversalis

Coccinella transversalis, commonly known as the transverse ladybird or transverse lady beetle, is a species of ladybird beetle found from India across southern and southeastern Asia to Malesia and Australia. It is not to be confused with...



Coccinella undecimpunctata

Coccinella undecimpunctata

Coccinella undecimpunctata, the eleven-spot ladybird or eleven-spotted lady beetle, it is native to central Asia, though commonly found in Europe, and formerly North America as its populations are decreasing. It is of the family Coccinellidae,...



Cochliomyia hominivorax

Cochliomyia hominivorax

Cochliomyia hominivorax, the New World screw-worm fly, or simply screw-worm, is a species of parasitic fly that is well known for the way in which its larvae (maggots) eat the living tissue of warm-blooded animals. It is present in the New World...



Codling moth

Cydia pomonella

The codling moth (Cydia pomonella) is a member of the Lepidopteran family Tortricidae. They are major pests to agricultural crops, mainly fruits such as apples and pears, and a codling moth larva is often called an "apple worm". Because the...



Coenonympha tullia

Coenonympha tullia

Coenonympha tullia, the large heath or common ringlet, is a butterfly in the family Nymphalidae. It flies in a variety of grassy habitats, including roadsides, woodland edges and clearings, prairies, bogs, and arctic and alpine taiga and tundra....



Colias alfacariensis

Colias alfacariensis

Colias alfacariensis, Berger's clouded yellow, is a butterfly of the family Pieridae. It was separated from the pale clouded yellow, C. hyale, in 1905. Berger's clouded yellow is a Palearctic species (South and Central Europe, South Russia,...



Colias croceus

Colias croceus

Colias croceus, clouded yellow, is a small butterfly of the family Pieridae, the yellows and whites.



Colias erate

Colias erate

Colias erate, commonly known as the eastern pale clouded yellow, is a species of butterfly in the family Pieridae. It is found from south-eastern Europe, through Turkey over central Asia up to Japan and Taiwan. To the south, its range stretches...



Colias hyale

Colias hyale

Colias hyale, the pale clouded yellow, is a butterfly of the family Pieridae, (also known as the yellows and whites) which is found in most of Europe and large parts of the Palearctic. It is a rare migrant to the British Isles and Scandinavia....



Colias interior

Colias interior

Colias interior, the pink-edged sulphur, is a species of North American butterfly in the family Pieridae. It is the State Butterfly of Maine.



Colletes inaequalis

Colletes inaequalis

Colletes inaequalis is a common species of plasterer bee (family Colletidae), native to North America. Like other species in the genus, it builds cells in underground nests that are lined with a polyester secretion, earning the genus the nickname...



Colobopsis saundersi

Colobopsis saundersi

Colobopsis saundersi (synonym Camponotus saundersi), also called the Malaysian exploding ant, is a species of ant found in Malaysia and Brunei, belonging to the genus Colobopsis. A worker can explode suicidally and aggressively as an ultimate act...



Colorado potato beetle

Leptinotarsa decemlineata

The Colorado potato beetle (Leptinotarsa decemlineata), also known as the Colorado beetle, the ten-striped spearman, the ten-lined potato beetle, or the potato bug, is a major pest of potato crops. It is...



Colotis danae

Colotis danae

Colotis danae, the crimson tip or scarlet tip, is a small butterfly of the family Pieridae, that is, the yellows and whites. It is found in Asia and Africa.



Colotis phisadia

Colotis phisadia

Colotis phisadia, the blue-spotted Arab, is a small butterfly of the family Pieridae, that is, the yellows and whites, which is found in Mauritania, Senegal, Mali, Nigeria, Niger, Chad, Sudan, Ethiopia, Kenya, Kuwait, Uganda, Arabia, the Near...



Comet moth

Argema mittrei

The comet moth or Madagascan moon moth (Argema mittrei). is a moth native to the rain forests of Madagascar. The species was first described by Félix Édouard Guérin-Méneville in 1847. The adult moth cannot feed and only lives for 4 to 5 days....



Common asparagus beetle

Crioceris asparagi

The common asparagus beetle (Crioceris asparagi) is an important pest of asparagus crops both in Europe and in North America. Asparagus is its only food plant. The beetle is 6.0 mm to 9.5 mm long and slightly elongated. It is metallic blue-black...



Common blue

Polyommatus icarus

The common blue butterfly or European common blue (Polyommatus icarus) is a butterfly in the family Lycaenidae and subfamily Polyommatinae. The butterfly is found throughout the Palearctic and has been introduced to North America. Butterflies in...



Common darter

Sympetrum striolatum

The common darter (Sympetrum striolatum) is a dragonfly of the family Libellulidae native to Eurasia. It is one of the most common dragonflies in Europe, occurring in a wide variety of water bodies, though with a preference for breeding in still...



Common emerald

Hemithea aestivaria

The common emerald (*Hemithea aestivaria*) is a moth of the family Geometridae. The species is found throughout the Nearctic and Palearctic regions and the Near East. It is mostly commonly found in the southern half of the Ireland and Britain. It...



Common footman

Manulea lurideola

The common footman (*Manulea lurideola*) is a moth of the family Erebidae. The species was first described by Johann Leopold Theodor Friedrich Zincken in 1817. It is distributed throughout Europe and east through the Palearctic to Lake Baikal. This...



Common furniture beetle

Anobium punctatum

The common furniture beetle or common house borer (*Anobium punctatum*) is a woodboring beetle originally from Europe but now distributed worldwide. In the larval stage it bores in wood and feeds upon it. Adult *Anobium punctatum* measure 2.7–4.5...



Common green bottle fly

Lucilia sericata

The common green bottle fly (*Lucilia sericata*) is a blowfly found in most areas of the world and is the most well-known of the numerous green bottle fly species. Its body is 10–14 mm (0.39–0.55 in) in length – slightly larger than a house fly –...



Common hawker

Aeshna juncea

The common hawker, moorland hawker or sedge darner (*Aeshna juncea*) is one of the larger species of hawker dragonflies. It is native to Palearctic (from Ireland to Japan) and northern North America. The flight period is from June to early...



Common Quaker

Orthosia cerasi

The common Quaker (Orthosia cerasi) is a moth of the family Noctuidae first described by Johan Christian Fabricius in 1775. Some authors prefer the synonym Orthosia stabilis (Denis & Schiffermüller, 1775). It is distributed throughout Europe and...



Common swift moth

Korscheltellus lupulina

The common swift (Korscheltellus lupulina) is a moth of the family Hepialidae. It was previously placed in the genus Hepialus. It is a common, often abundant European species. The species was described by Carl Linnaeus in his 1758 10th edition of...



Cooloola monster

Cooloola propator

The Cooloola monster (Cooloola propator) is a large burrowing orthopteran of the family Cooloolidae, a family erected to accommodate it because it is so dissimilar to other ensiferans. It was discovered in 1980 in the Great Sandy National Park in...



Copper underwing

Amphipyra pyramidea

The copper underwing, humped green fruitworm or pyramidal green fruitworm (Amphipyra pyramidea) is a moth of the family Noctuidae. The species was first described by Carl Linnaeus in his 1758 10th edition of Systema Naturae.



Corydalus cornutus

Corydalus cornutus

The eastern dobsonfly, *Corydalus cornutus*, is a large insect in the Corydalidae family. It is found in eastern North America in regions with fast-flowing streams where its aquatic larvae develop. These are known as hellgrammites and are among the...



Corythucha ciliata

Corythucha ciliata

Corythucha ciliata, the sycamore lace bug, is a species of lace bug in the family Tingidae that is associated with sycamore trees. It originates in the New World, but has been introduced and has spread in Europe. It is a small, whitish insect and...



Coscinocera hercules

Coscinocera hercules

Coscinocera hercules, the Hercules moth, is a moth of the family Saturniidae, endemic to New Guinea and northern Australia. The species was first described by William Henry Miskin in 1876.



Costarica

Costarica costaricensis

Costarica is a genus of grasshoppers in the subfamily Romaleinae; described by Koçak & Kemal in 2008. Its only species is Costarica costaricensis.



Cotesia congregata

Cotesia congregata

Cotesia congregata is a parasitoid wasp of the genus Cotesia. The genus is particularly noted for its use of polydnaviruses. Parasitoids are distinct from true parasites in that a parasitoid will ultimately kill its host or otherwise sterilize it.



Cotinis nitida

Cotinis nitida

Cotinis nitida, commonly known as the green June beetle, June bug or June beetle, is a beetle of the family Scarabaeidae. It is found in the eastern United States and Canada, where it is most abundant in the South. It is sometimes confused with...



Crab louse

Pthirus pubis

The crab louse or pubic louse (Pthirus pubis) is an insect that is an obligate ectoparasite of humans, feeding exclusively on blood. The crab louse usually is found in the person's pubic hair. Although the louse cannot jump, it can also live in...



Crataerina pallida

Crataerina pallida

Crataerina pallida, the swift lousefly, is a species of biting fly in the family of louse flies Hippoboscidae. These flies are commonly encountered in the nests of the common swift (*Apus apus*) in Europe and Asia. The lousefly spends its entire...



Cream wave

Scopula floslactata

The cream wave (*Scopula floslactata*) is a moth of the family Geometridae. The species was first described by Adrian Hardy Haworth in 1809. It is found in forest and woodland regions, feeding on grasses and small plants such as dandelion.



Creobroter gemmatus

Creobroter gemmatus

Creobroter gemmatus, common name jeweled flower mantis, is a species of praying mantis native to Asia.



Creobroter pictipennis

Creobroter pictipennis

Creobroter pictipennis, with the common name Indian flower mantis, is a species of praying mantis native to Asia. Males grow to about 1.5 in long (3.8 cm) and females are slightly larger.



Ctenolepisma lineatum

Ctenolepisma lineatum

Ctenolepisma lineatum is a species of insect of the order Zygentoma. It is similar to the closely related silverfish but can be distinguished by being rather stouter and less shiny with all appendages (antennae and 3 "tails") noticeably longer....



Ctenucha virginica

Ctenucha virginica

Ctenucha virginica, the Virginia ctenucha, is a moth of the family Erebidæ. The species was first described by Eugenius Johann Christoph Esper in 1794.



Culex tritaeniorhynchus

Culex tritaeniorhynchus

Culex (Culex) tritaeniorhynchus is a species of mosquito and is the main vector of the disease Japanese encephalitis. This mosquito is a native of northern Asia, and parts of Africa (northeast and sub-Saharan). Females target large animals for...



Culicoides imicola

Culicoides imicola

Culicoides imicola (Culicoides imicola Kieffer, former name C. pallidipennis Carter) is a species of Ceratopogonidae that transmits the bluetongue virus (BTV) and the African horse sickness virus. This particular species has been recorded in...



Curetis thetis

Curetis thetis

Curetis thetis, the Indian sunbeam, is a species of lycaenid or red butterfly found in Indomalayan realm.



Cyaniris semiargus

Cyaniris semiargus

Cyaniris semiargus, the Mazarine blue, is a Palearctic butterfly in the family Lycaenidae.



Cybister chinensis

Cybister chinensis

Cybister chinensis is a species of diving beetle native to East Asia. It is predatory, feeding on tadpoles, small fish and aquatic insects, and adults are about 3.3–4.2 cm (1.3–1.7 in) long.



Cybister lateralimarginalis

Cybister lateralimarginalis

Cybister lateralimarginalis is a species of beetle native to the Palearctic, including Europe, the Middle East and North Africa.



Cyclommatus metallifer

Cyclommatus metallifer

Cyclommatus metallifer is a species of stag beetle in the family Lucanidae. There is sexual dimorphism within the species. Males are generally larger in size and have enlarged mandibles. It is named for its metallic coloration, which ranges in...



Cycnia tenera

Cycnia tenera

Cycnia tenera, the dogbane tiger moth or delicate cycnia, is a moth in the family Erebidæ. It occurs throughout North America, from southern British Columbia to Nova Scotia southwards to Arizona and Florida. The species is distasteful and there...



Cydia latiferreana

Cydia latiferreana

Cydia latiferreana, the filbertworm moth, is a moth of the family Tortricidae. It was formerly (and sometimes is still) separated in a monotypic genus Melissopus. It is found in most of North America. The wingspan is about 18 mm. Adults are on...



Cydia saltitans

Cydia saltitans

Cydia saltitans or jumping bean moth is a moth from Mexico that is most widely known as its larva, where it inhabits the carpels of seeds from several related shrubby trees, mainly Sebastiania pavoniana or Sapium biloculare (syn. Pleradenophora...



Cydia splendana

Cydia splendana

Cydia splendana, the chestnut tortrix, is a moth of the family Tortricidae. It is found in Europe. It is also known as the acorn moth, but this can also refer to Blastobasis glandulella from North America, which belongs to the more primitive...



Cymbalophora pudica

Cymbalophora pudica

Cymbalophora pudica, the discrete chaperon, is a moth of the family Erebidae. The species was first described by Eugenius Johann Christoph Esper in 1784.



Cynomya mortuorum

Cynomya mortuorum

Cynomya mortuorum belongs to the order Diptera, sometimes referred to as "true flies". In English, the only common name occasionally used is "fly of the dead" (Totenfliege in German). It has a bluish-green appearance, similar to other...



Danaus affinis

Danaus affinis

Danaus affinis, the Malay tiger, mangrove tiger or swamp tiger, is a butterfly found in tropical Asia. It belongs to the "crows and tigers", the danaine group of the brush-footed butterflies family. This is a highly variable species. In its range....



Danaus chrysippus

Danaus chrysippus

Danaus chrysippus, also known as the plain tiger, African queen, or African monarch, is a medium-sized butterfly widespread in Asia, Australia and Africa. It belongs to the Danainae subfamily of the brush-footed butterfly family Nymphalidae....



Danaus eresimus

Danaus eresimus

Danaus eresimus, the soldier or tropical queen, is a North American, Caribbean, and South American butterfly in the family Nymphalidae. Their flight is slow and they are reasonably easy to approach, but will fly for some distance if approached too...



Danaus genutia

Danaus genutia

Danaus genutia, the common tiger, is one of the common butterflies of India. It belongs to the "crows and tigers", that is, the Danainae group of the brush-footed butterflies family. The butterfly is also called striped tiger in India to...



Daphnis hypothous

Daphnis hypothous

Daphnis hypothous, the jade hawkmoth, is a moth of the family Sphingidae described by Pieter Cramer in 1780. It is known from Sri Lanka, southern and northern India, Nepal, Myanmar, southern China, Taiwan, Thailand, Malaysia, and Indonesia. It is...



Daphnis nerii

Daphnis nerii

Daphnis nerii, the oleander hawk-moth or army green moth, is a moth of the family Sphingidae. It was described by Carl Linnaeus in his 1758 10th edition of Systema Naturae.



Dark dagger

Acronicta tridens

The dark dagger (Acronicta tridens) is a moth of the family Noctuidae. The species was first described by Michael Denis and Ignaz Schiffermüller in 1775. It is distributed throughout Europe (from southern Fennoscandia to the Balkans and Italy),...



Dark green fritillary

Speyeria aglaja

The dark green fritillary (Speyeria aglaja) is a species of butterfly in the family Nymphalidae. The insect has a wide range in the Palearctic realm - Europe, Morocco, Iran, Siberia, Central Asia, China, Korea, and Japan.



Deathwatch beetle

Xestobium rufovillosum

The deathwatch beetle (Xestobium rufovillosum) is a species of woodboring beetle that sometimes infests the structural timbers of old buildings. The adult beetle is brown and measures on average 7 mm (0.3 in) long. Eggs are laid in dark crevices...



Deilephila elpenor

Deilephila elpenor

Deilephila elpenor, the elephant hawk moth or large elephant hawk moth, is a moth in the family Sphingidae. Its common name is derived from the caterpillar's resemblance to an elephant's trunk. It is most common in central Europe and is...



Deilephila porcellus

Deilephila porcellus

Deilephila porcellus, the small elephant hawk-moth, is a moth of the family Sphingidae. The species was first described by Carl Linnaeus in his 1758 10th edition of Systema Naturae.



Deinacrida fallai

Deinacrida fallai

Deinacrida fallai or the Poor Knights giant wētā is a species of insect in the family Anostomatidae. It is endemic to the Poor Knights Islands off northern New Zealand. D. fallai are commonly called giant wētā due to their large size. They are...



Deinacrida heteracantha

Deinacrida heteracantha

Deinacrida heteracantha, also known as the Little Barrier giant wētā or wētāpunga (Māori: wētāpunga), is a wētā in the order Orthoptera and family Anostomatidae. It is endemic to New Zealand, where it survived only on Little Barrier Island,...



Deinacrida parva

Deinacrida parva

Deinacrida parva is a species of insect in the family Anostomatidae, the king crickets and weta. It is known commonly as the Kaikoura wētā or Kaikoura giant wētā. It was first described in 1894 from a male individual then rediscovered in 1966...



Deinacrida rugosa

Deinacrida rugosa

Deinacrida rugosa, commonly called the Cook Strait giant weta or Stephens Island weta, is a species of insect in the family Anostomatidae. The scientific name *Deinacrida* means "terrible grasshopper" and *rugosa* means "wrinkled". It is endemic...



Delhi Sands flower-loving fly

Rhaphiomidas terminatus abdominalis

The Delhi Sands flower-loving fly (*Rhaphiomidas terminatus abdominalis*) is a subspecies of flower-loving fly, a member of the genus *Rhaphiomidas* in the Mydidae family. It was the first fly added to the Endangered Species List in the United...



Delias eucharis

Delias eucharis

Delias eucharis, the common Jezebel, is a medium-sized pierid butterfly found in many areas of south and southeast Asia, especially in the non-arid regions of India, Bangladesh, Sri Lanka, Indonesia, Myanmar and Thailand. The common Jezebel is...



Delias harpalyce

Delias harpalyce

Delias harpalyce, the imperial white, is a butterfly in the family Pieridae. It is endemic to Australia.



Delias pasithoe

Delias pasithoe

Delias pasithoe, the redbase Jezebel is a medium-sized butterfly of the family Pieridae, that is, the yellows and whites. The species is found in parts of South Asia and Southeast Asia. There has been some dispute for which species the specific...



Dendroctonus rufipennis

Dendroctonus rufipennis

Dendroctonus rufipennis, the spruce beetle, is a species of bark beetle native to British Columbia, Newfoundland and Labrador, Nova Scotia, Ontario, Quebec, Northern Manitoba, the Yukon, Alaska, Colorado, Wyoming, Montana, and Maine. They are...



Dendrolimus pini

Dendrolimus pini

Dendrolimus pini, the pine-tree lappet, is a moth of the family Lasiocampidae. The species was first described by Carl Linnaeus in his 1758 10th edition of Systema Naturae. It is found in most of Europe ranging to eastern Asia. The wingspan is...



Dendroplectron

Dendroplectron aucklandensis

Dendroplectron aucklandensis the Auckland Island wētā, is a cave wētā in the family Rhaphidophoridae, the only member of the genus Dendroplectron. It is endemic to the subantarctic Auckland Islands of New Zealand.



Dermatobia hominis

Dermatobia hominis

The human botfly, Dermatobia hominis (Greek δέρμα, skin + βίος, life, and Latin hominis, of a human), is a species of botfly whose larvae parasitise humans (in addition to a wide range of other animals, including other primates). It is also known...



Deroplatys desiccata

Deroplatys desiccata

Deroplatys desiccata, known by the common name giant dead leaf mantis, is a praying mantis from Southeast Asia. This is the type species of genus Deroplatys.



Desert locust

Schistocerca gregaria

The desert locust (Schistocerca gregaria) is a species of locust, a periodically swarming, short-horned grasshopper in the family Acrididae. They are found primarily in the deserts and dry areas of northern and eastern Africa, Arabia, and...



Deudorix epijarbas

Deudorix epijarbas

Deudorix epijarbas, the cornelian or hairy line blue, is a species of lycaenid or blue butterfly found in south and southeast Asia from India to Fiji, including the Philippines, and also the tropical coast of Queensland in Australia. The species...



Devil's coach horse beetle

Ocypus olens

The devil's coach-horse beetle (Ocypus olens) is a species of beetle belonging to the large family of the rove beetles (Staphylinidae). It was originally included in the genus Staphylinus in 1764, and some authors and biologists still use this...



Diabrotica undecimpunctata

Diabrotica undecimpunctata

Diabrotica undecimpunctata, the spotted cucumber beetle or southern corn rootworm, is a species of cucumber beetle that is native to North America. The species can be a major agricultural pest insect in North America. Spotted cucumber beetles...



Diaethria phlogea

Diaethria phlogea

Diaethria phlogea, the 89/98 butterfly, is a species of butterfly of the family Nymphalidae. It is found in Colombia, South America. It has been given the nickname "89/98" because of the markings on its wings resembling an 89 and 98. Some authors...



Diamondback moth

Plutella xylostella

The diamondback moth (Plutella xylostella), sometimes called the cabbage moth, is a moth species of the family Plutellidae and genus Plutella. The small, grayish-brown moth sometimes has a cream-colored band that forms a diamond along its back....



Diaphania nitidalis

Diaphania nitidalis

Diaphania nitidalis, the pickleworm, is a serious agricultural pest insect in the family Crambidae. It damages squash primarily, but it is also a common pest of other cucurbits such as cucumbers and melons. It is a tropical species which can be...



Dichomeris ligulella

Dichomeris ligulella

The palmerworm (Dichomeris ligulella) is a moth of the family Gelechiidae. It is found in eastern North America. The wingspan is 15–18 mm. Adults are on wing from April to October. There is one generation per year. The larvae feed on apple....



Dicopomorpha echmepterygis

Dicopomorpha echmepterygis

Dicopomorpha echmepterygis is the smallest known insect and a species of parasitoid wasp in the family Mymaridae, which exhibits strong sexual dimorphism. The males are blind, apterous, and their body length is only 40% that of females. With a...



Didymuria violescens

Didymuria violescens

Didymuria violescens, the spur legged phasmid, also known as the violet-winged stick insect, or violet-winged phasma, is a common phasmid native to Australia. Males have distinct spurs and thicker hind legs (the hind have the spurs) which is where...



Differential grasshopper

Melanoplus differentialis

The differential grasshopper (Melanoplus differentialis) is a species of grasshopper belonging to the genus Melanoplus. It is found throughout northern Mexico, the central United States and southern Ontario, Canada. It is considered a pest over...



Dingy skipper

Erynnis tages

The dingy skipper (Erynnis tages) is a species of butterfly in the family Hesperiiidae.



Dioryctria sylvestrella

Dioryctria sylvestrella

Dioryctria sylvestrella, the new pine knot-horn or maritime pine borer, is a moth of the family Pyalidae. It is found in Europe, parts of Asia and North Africa. The adult is a small mottled brown and white insect with a wingspan of 28 to 35 mm...



Diphthera festiva

Diphthera festiva

Diphthera festiva, the hieroglyphic moth, is a species of moth in the family Nolidae and is the only moth in its subfamily Diphtherinae. It is found in the tropical and subtropical areas of South America (as far south as Bolivia and Brazil),...



Diplacodes haematodes

Diplacodes haematodes

Diplacodes haematodes, the scarlet percher, is a species of dragonfly in the family Libellulidae. It occurs throughout Australia (except Tasmania), Timor, New Guinea, Vanuatu, and New Caledonia. It is locally common in habitats with hot sunny...



Diplolepis rosae

Diplolepis rosae

Diplolepis rosae is a gall wasp which causes a gall known as the rose bedeguar gall, bedeguar gall wasp, Robin's pincushion, mossy rose gall, or simply moss gall. The gall develops as a chemically induced distortion of an unopened leaf axillary...



Dolba

Dolba hyloeus

Dolba is a monotypic moth genus in the family Sphingidae erected by Francis Walker in 1856. Its only species, Dolba hyloeus, the pawpaw sphinx, was first described by Dru Drury in 1773.



Doleschallia bisaltide

Doleschallia bisaltide

Doleschallia bisaltide, the autumn leaf, is a nymphalid butterfly found in South Asia, Southeast Asia, and Australasia. In Australia it is also known as the leafwing.



Dolichovespula maculata

Dolichovespula maculata

Dolichovespula maculata is a species of wasp in the genus Dolichovespula and a member of the eusocial, cosmopolitan family Vespidae. It is known by many colloquial names, primarily bald-faced hornet, but also including bald-faced aerial...



Dorcus curvidens

Dorcus curvidens

Dorcus curvidens is a species of stag beetle.



Dorcus parallelipipedus

Dorcus parallelipipedus

Dorcus parallelipipedus, the lesser stag beetle, is a species of stag beetle found in Europe.



Doru aculeatum

Doru aculeatum

Doru aculeatum, the spine-tailed earwig, is a species of earwig in the family Forficulidae.:144 It is found in the woods and grassy areas of eastern North America and occurs at outdoor lights at night.:55 The adults have a brown body with pale...



Dot moth

Melanchra persicariae

The dot moth (Melanchra persicariae) is a moth of the family Noctuidae. The species was first described by Carl Linnaeus in 1761. It is a very distinctive species with very dark brown, almost black, forewings marked with a large white stigma from...



Double-striped pug

Gymnoscelis rufifasciata

The double-striped pug (Gymnoscelis rufifasciata) is a moth of the family Geometridae. It is a widespread and common species, being found throughout the Palearctic region, including the Near East and North Africa. This is a variable species but...



Downy emerald

Cordulia aenea

The downy emerald (Cordulia aenea) is a species of dragonfly. It is metallic green and bronze in color, and its thorax is coated with fine hairs, hence its name. Like most other emeralds, the downy emerald has bright shiny green eyes. Adults are...



Drugstore beetle

Stegobium paniceum

The drugstore beetle (*Stegobium paniceum*), also known as the bread beetle, biscuit beetle, and misnamed as the biscuit weevil (despite not being a weevil), is a tiny, brown beetle that can be found infesting a wide variety of dried plant...



Dryadula phaetusa

Dryadula phaetusa

Dryadula is a monotypic genus of the butterfly family Nymphalidae. Its single species, Dryadula phaetusa, known as the banded orange heliconian, banded orange, or orange tiger, is native from Brazil to central Mexico, and in summer can be found...



Dryas iulia

Dryas iulia

Dryas iulia (often incorrectly spelled julia), commonly called the Julia butterfly, Julia heliconian, the flame, or flambeau, is a species of brush-footed (or nymphalid) butterfly. The sole representative of its genus Dryas, it is native from...



Dryocampa rubicunda

Dryocampa rubicunda

Dryocampa rubicunda, the rosy maple moth, is a small North American moth in the family Saturniidae, also known as the great silk moths. It was first described by Johan Christian Fabricius in 1793. The species is known for its wooly body and pink...



Dun-bar

Cosmia trapezina

The dun-bar (*Cosmia trapezina*) is a moth of the family Noctuidae. It is a common Palearctic species.



Dusky large blue

Phengaris nausithous

The dusky large blue (Phengaris nausithous) is a species of butterfly in the family Lycaenidae. It is found in Armenia, Austria, Azerbaijan, Belarus, Bulgaria, Czech Republic, France, Georgia, Germany, Hungary, Kazakhstan, Moldova, Montenegro,....



Dynastes satanas

Dynastes satanas

Dynastes satanas, the Satanas beetle, is a species of beetle belonging to the family Scarabaeidae. The name is sometimes misspelled as "satanus".Because of collecting and habitat loss this rare species is protected and included in the list of...



Dynastes tityus

Dynastes tityus

Dynastes tityus, the eastern Hercules beetle, is a species of rhinoceros beetle native to the Eastern United States. The adult's elytra are green, gray or tan, with black markings, and the whole animal, including the male's horns, may reach 60 mm...



Dysphania militaris

Dysphania militaris

Dysphania militaris is a species of moth of the family Geometridae that is found from in the tropical regions of South and Southeast Asian countries such as China, India, Myanmar, Andaman Islands, Sumatra, Philippines and Java. It was first...



Dytiscus latissimus

Dytiscus latissimus

Dytiscus latissimus is a large species of aquatic beetle in family Dytiscidae. It is native to Europe and considered threatened.



Eacles imperialis

Eacles imperialis

Eacles imperialis, the imperial moth, is a member of the family Saturniidae and subfamily Ceratocampinae. It is found mainly in the East of South America and North America, from the center of Argentina to south Canada. The species was first...



Early bumblebee

Bombus pratorum

The early bumblebee or early-nesting bumblebee (Bombus pratorum) is a small bumblebee with a wide distribution in most of Europe and parts of Asia. It is very commonly found in the UK and emerges to begin its colony cycle as soon as February...



Eastern carpenter bee

Xylocopa virginica

Xylocopa virginica, sometimes referred to as the eastern carpenter bee, extends through the eastern United States and into Canada. They are sympatric with Xylocopa micans in much of southeastern United States. They nest in various types of wood...



Eastern tent caterpillar

Malacosoma americanum

The eastern tent caterpillar (Malacosoma americanum) is a species of moth in the family Lasiocampidae, the tent caterpillars or lappet moths. It is univoltine, producing one generation per year. It is a tent caterpillar, a social species that...



Eciton hamatum

Eciton hamatum

Eciton hamatum is a species of army ant in the subfamily Dorylinae; it is found from Mexico to central Brazil and Bolivia. The species differs from Eciton burchellii, in that it does not fan out into the underbrush when foraging. Rather, it...



El Segundo blue

Euphilotes battoides allyni

The El Segundo blue (Euphilotes allyni) is an endangered species of butterfly. It is endemic to a small dune ecosystem in Southern California that used to be a community called Palisades del Rey, close to the Los Angeles International Airport (LAX).



Elaphrus viridis

Elaphrus viridis

Elaphrus viridis, the Delta green ground beetle, is a species of ground beetle restricted to a small region within Solano County, California. Its color is a metallic-green, usually with bronze spots on its elytra, though some lack these spots....



Electric ant

Wasmannia auropunctata

The little fire ant (Wasmannia auropunctata), also known as the electric ant, is a small...



Elephant beetle

Megasoma elephas

The elephant beetle (Megasoma elephas) is a member of the family Scarabaeidae and the subfamily Dynastinae. Elephant beetles are Neotropical rhinoceros beetles.



Elm leaf beetle

Xanthogaleruca luteola

Xanthogaleruca luteola, commonly known as the elm-leaf beetle, is a beetle species in the family Chrysomelidae that is native to Europe but invasive in other parts of the world.



Eloria noyesi

Eloria noyesi

Eloria noyesi, the coca tussock moth, is a moth of the subfamily Lymantriinae first described by Schaus in 1927. It is beige and its larvae feed on coca plants. It is found mostly in Peru and Colombia. The government of Colombia has proposed a...



Elymnias hypermnestra

Elymnias hypermnestra

Elymnias hypermnestra, the common palmfly, is a species of satyrine butterfly found in South and Southeast Asia.



Emerald ash borer

Agrilus planipennis

The emerald ash borer (Agrilus planipennis), also known by the acronym EAB, is a green buprestid or jewel beetle native to north-eastern Asia that feeds on ash species (Fraxinus spp.). Females lay eggs in bark crevices on ash trees, and larvae...



Emerald cockroach wasp

Ampulex compressa

The emerald cockroach wasp or jewel wasp (Ampulex compressa) is a solitary wasp of the family Ampulicidae. It is known for its unusual reproductive behavior, which involves stinging a cockroach and using it as a host for its larvae. It thus...



Emperor

Anax imperator

The emperor dragonfly or blue emperor (*Anax imperator*) is a large species of hawker dragonfly of the family Aeshnidae. It is the bulkiest dragonfly in most of Europe, including the United Kingdom, although exceeded by the magnificent emperor (*A....*



Empusa pennata

Empusa pennata

Empusa pennata, or the conehead mantis, is a species of praying mantis in genus Empusa native to the Mediterranean Region. It can be found in Portugal, Spain, southern France, Italy and on the mediterranean coasts of Morocco, Algeria, Tunisia,...



Enallagma cyathigerum

Enallagma cyathigerum

Enallagma cyathigerum (common blue damselfly, common bluet, or northern bluet) is a species found mainly between latitudes 40°N and 72°N; It is widely distributed in the Palearctic, and the Nearctic species Enallagma annexum was at one time...



Engrailed

Ectropis crepuscularia

The engrailed and small engrailed (*Ectropis crepuscularia*) are moths of the family Geometridae found from the British Isles through central and eastern Europe to the Russian Far East and Kazakhstan. The western Mediterranean and Asia Minor and...



Epiophlebia laidlawi

Epiophlebia laidlawi

Epiophlebia laidlawi, the Himalayan relict dragonfly, is one of four species of Eiprocta in the family Epiophlebiidae. They have at one time been classified as a suborder Anisozygoptera, considered as intermediate between the dragonflies and the...



Erinnyis alope

Erinnyis alope

Erinnyis alope, the Alope sphinx, is a moth of the family Sphingidae. It lives from the northern part of South America, through Central America, up to Northern Mexico and the very south of the United States, although strays have been recorded as...



Erinnyis ello

Erinnyis ello

Erinnyis ello, the ello sphinx, is a moth of the family Sphingidae. The species was first described by Carl Linnaeus in his 1758 10th edition of Systema Naturae. It is distributed from Argentina through Central America to the United States as far...



Erionota thrax

Erionota thrax

Erionota thrax, the palm redeye or the banana skipper, is a species of butterfly belonging to the family Hesperidae. It is found from India, through south-eastern Asia to Papua New Guinea. In the north it is found up to southern China. It is an...



Ernobius mollis

Ernobius mollis

Ernobius mollis, also known as the pine bark anobiid, pine knot borer, bark borer, or waney edge borer, is a species of beetle in the family Ptinidae.



Erratic ant

Tapinoma erraticum

The erratic ant (Tapinoma erraticum) is a species of dolichoderine ant first described in 1789 by Latreille. This species ranges throughout Central Europe from the mountains of south Italy to north Germany. It is present in coastal areas of...



Essex skipper

Thymelicus lineola

Thymelicus lineola, known in Europe as the Essex skipper and in North America as the European skipper, is a species of butterfly in the family HesperIIDae. With a wingspan of 2.5 to 2.9 cm, it is very similar in appearance to the small skipper,...



Euchaetes egle

Euchaetes egle

Euchaetes egle, the milkweed tiger moth or milkweed tussock moth, is a moth in the family ErebidAe and the tribe Arctiini, the tiger moths. The species was first described by Dru Drury in 1773. It is a common mid- through late summer feeder on...



Euchloe ausonides

Euchloe ausonides

Euchloe ausonides, the large marble or creamy marbledwing, is a species of butterfly that occurs in western North America. It lays eggs on the terminal flower buds of a variety of plants in the mustard family, including introduced Eurasian...



Euchloe ausonides insulana

Euchloe ausonides insulanus

Euchloe ausonides insulanus, the island marble butterfly, is a subspecies of the Euchloe ausonides (large marble butterfly) and is found in the San Juan Islands in Washington in the northwestern United States. The butterfly was thought to be...



Euchloe olympia

Euchloe olympia

Euchloe olympia, the Olympia marble, is a butterfly in the family Pieridae. Its range is southern Canada and the Midwest, down into the southwestern United States. E. olympia is related to E. guaymasensis, but has a distinctive phenotype and...



Eudocima apta

Eudocima apta

Eudocima apta is a moth of the family Erebidae. It is found in large parts of Brazil. At times it migrates north into the United States. The wingspan is about 45 mm.



Eudocima homaena

Eudocima homaena

Eudocima homaena is a moth of the family Erebidae first described by Jacob Hübner in 1816. It is found in the Indian subregion, Sri Lanka, Myanmar, Taiwan, the Nicobars, Peninsular Malaysia, Borneo, the Philippines and on Christmas Island. It is...



Eudocima phalonia

Eudocima phalonia

Eudocima phalonia, the common fruit-piercing moth, is a fruit piercing moth of the family Erebidae. The species was first described by Carl Linnaeus in his 1763 Centuria Insectorum. It is found in large parts of the tropics, mainly in Asia,...



Eudocima salamina

Eudocima salamina

Eudocima salamina, the green fruit-piercing moth, is a moth of the family Erebidae. The species was first described by Pieter Cramer in 1777. It is found from India, and across south-east Asia to the Pacific Islands. In Australia it occurs in...



Eudryas grata

Eudryas grata

Eudryas grata is a moth known as the beautiful wood nymph. They are known for their mimicry of bird droppings. Found in abundance, predominantly across the entire eastern United States. Hosts for the caterpillar include Ampelopsis, buttonbush,...



Eumaeus atala

Eumaeus atala

Eumaeus atala, also known as the Atala butterfly or coontie hairstreak, is a small colorful butterfly in the family Lycaenidae. It is found in southeastern Florida (including the Florida Keys) in the United States, and the Bahamas, Cuba, and the...



Eumeta crameri

Eumeta crameri

Eumeta crameri is a bagworm moth of the family Psychidae. It was described by John O. Westwood in 1854 and has worldwide distribution in tropical and subtropical habitats, including India, Bangladesh, Sri Lanka, New Zealand and Puerto Rico.



Eumorpha achemon

Eumorpha achemon

Eumorpha achemon, the Achemon sphinx, is a moth of the family Sphingidae. The species was first described by Dru Drury in 1773.



Eumorpha labruscae

Eumorpha labruscae

Eumorpha labruscae, the gaudy sphinx, is a moth in the family Sphingidae.



Eumorpha vitis

Eumorpha vitis

Eumorpha vitis, known as the vine sphinx, is a moth of the family Sphingidae.



Eupeodes americanus

Eupeodes americanus

Eupeodes americanus, the American hoverfly, is found throughout North America and inhabits meadows, and fields with flowers and foliage. Adults feed on nectar, whereas their larvae feed on aphids. The adult fly is black to metallic green, and has...



Eupithecia orichloris

Eupithecia orichloris

Eupithecia orichloris is a species of moth that is native to Kauai, Oahu, Maui, Lanai and Hawaii. It is known for having its larval form as a caterpillar which is insectivorous. The caterpillar has two abdominal appendages which serve as triggers...



Euplagia quadripunctaria

Euplagia quadripunctaria

Euplagia quadripunctaria, the Jersey tiger, or Spanish flag, is a diurnal moth of the family Erebidæ. The species was first described by Nikolaus Poda von Neuhaus in 1761. The adult wingspan is 52–65 millimetres (2.0–2.6 in), and they fly from...



Euploea core

Euploea core

Euploea core, the common crow, is a common butterfly found in South Asia to Australia. In India it is also sometimes referred to as the common Indian crow, and in Australia as the Australian crow. It belongs to the crows and tigers subfamily...



Euploea eunice

Euploea eunice

Euploea eunice, commonly called blue-banded king crow, is a butterfly found in the Indomalayan realm that belongs to the crows and tigers, that is, the Danaid group of the brush-footed butterflies family. The larva feeds on Ficus, Flacourtia...



Euploea klugii

Euploea klugii

Euploea klugii, the brown king crow or king crow, is a butterfly from the family Nymphalidae found in India and Southeast Asia. The species was first described by the entomologist Frederic Moore in 1858.



Euploea mulciber

Euploea mulciber

Euploea mulciber, the striped blue crow, is a butterfly found in India and Southeast Asia that belongs to the crows and tigers, that is, the danaid group of the brush-footed butterflies family.



Euploea radamanthus

Euploea radamanthus

Euploea radamanthus, the magpie crow, is a butterfly found in India and Southeast Asia that belongs to the danaid group of the brush-footed butterflies family. Euploea radamanthus is found in the Eastern Himalayas and into the Malay region.



Euploea sylvester

Euploea sylvester

Euploea sylvester, the double-branded crow, also known as the two-brand crow in Australia, is a butterfly found in South Asia, Southeast Asia and parts of Australia that belongs to the crows and tigers, that is, the danaid group of the...



Euptoieta hegesia

Euptoieta hegesia

Euptoieta hegesia, the Mexican fritillary, is a North and South American butterfly in the family Nymphalidae.



Eurema blanda

Eurema blanda

Eurema blanda, the three-spot grass yellow, is a small butterfly of the family Pieridae which is found in Sri Lanka, India and southeast Asia.



Eurema hecabe

Eurema hecabe

Eurema hecabe, the common grass yellow, is a small pierid butterfly species found in Asia, Africa and Australia. They are found flying close to the ground and are found in open grass and scrub habitats. It is simply known as "the grass yellow" in...



Eurema mexicana

Eurema mexicana

Eurema mexicana, the Mexican yellow, sometimes called the wolf-face sulphur, is a North and South American butterfly in the family Pieridae. It occurs mainly in Mexico but occasionally is found in central and southwestern United States and rarely...



Euroleon nostras

Euroleon nostras

Euroleon nostras is a species of antlion found over most of Europe. The scientific name can be translated as "our European lion". Adults resemble dragonflies or damselflies and may reach up to 30 mm (1.2 in) long, with a wingspan of 70 mm (2.8...



European beewolf

Philanthus triangulum

The European beewolf (Philanthus triangulum), also known as the bee-killer wasp or the bee-eating philanthus (from the now obsolete synonym Philanthus apivorus), is a solitary wasp that lives in the Western Palearctic and Afrotropics. Although...



European corn borer

Ostrinia nubilalis

The European corn borer (Ostrinia nubilalis), also known as the European corn worm or European high-flyer, is a moth of the family Crambidae. It is a pest of grain, particularly maize (Zea mays). The insect is native to Europe, originally...



European dark bee

Apis mellifera mellifera

The European dark bee (Apis mellifera mellifera) is a subspecies of the western honey bee, evolving in central Asia, with a proposed origin of the Tien Shan Mountains and later migrating into eastern and then northern Europe after the last ice...



European hornet

Vespa crabro

The European hornet (Vespa crabro) is the largest eusocial wasp native to Europe. It is also the only true hornet (genus Vespa) found in North America, having been introduced to the United States and Canada from Europe as early as 1840. Vespines,...



European mantis

Mantis religiosa

The European mantis (*Mantis religiosa*) is a large hemimetabolic insect in the family of the Mantidae ('mantids'), which is the largest family of the order Mantodea (mantises). Their common name praying mantis is derived from the distinctive...



European paper wasp

Polistes dominula

The European paper wasp (*Polistes dominula*) is one of the most common and well-known species of social wasps in the genus *Polistes*. Its diet is more diverse than those of most *Polistes* species—many genera of insects versus mainly caterpillars in...



Eurytides marcellus

Eurytides marcellus

Eurytides marcellus, the zebra swallowtail (formerly listed under genera *Protographium*, *Iphiclides*, *Graphium* and *Papilio* by some authorities), is a swallowtail butterfly native to the eastern United States and south-eastern Canada. It is the...



Euschistus ictericus

Euschistus ictericus

Euschistus ictericus is a North American species of shield bug. It grows up to 12 mm (0.5 in) long, and lives in damp areas.



Euthalia aconthea

Euthalia aconthea

Euthalia aconthea, the common baron, often called simply baron, is a medium-sized nymphalid butterfly native to Sri Lanka, India and southeast Asia. It flies with stiff wing beats and often glides. The wing is not flapped very far below the...



Euthalia lubentina

Euthalia lubentina

Euthalia lubentina, the gaudy baron, is a species of nymphalid butterfly found in South, Cambodia, and Southeast Asia. It was first described by Pieter Cramer in 1777.



Evergreen bagworm

Thyridopteryx ephemeraeformis

The evergreen bagworm (Thyridopteryx ephemeraeformis), commonly known as bagworm, eastern bagworm, common bagworm, common basket worm, or North American bagworm, is a moth that spins its cocoon in its larval life, decorating it with bits of plant...



Extatosoma tiaratum

Extatosoma tiaratum

Extatosoma tiaratum, commonly known as the spiny leaf insect, the giant prickly stick insect, Macleay's spectre, or the Australian walking stick, is a large species of Australian stick insect. The species has the Phasmid Study Group number PSG9.



Fall webworm

Hyphantria cunea

The fall webworm (Hyphantria cunea) is a moth in the family Erebidae known principally for its larval stage, which creates the characteristic webbed nests on the tree limbs of a wide variety of hardwoods in the late summer and fall. It is...



False garden mantis

Pseudomantis albofimbriata

The false garden mantis (Pseudomantis albofimbriata) is a species of praying mantis in the family Mantidae, and was first described in 1860 by Carl Stål as Mantis albofimbriata. Females reach 70 mm while males reach 50 mm.



False potato beetle

Leptinotarsa juncta

The false potato beetle (Leptinotarsa juncta) is a beetle found primarily in the Mid-Atlantic and Southeastern United States. Its distribution extends to Maine. Adult beetles emerge from the soil in the late spring or early summer and begin...



Fannia scalaris

Fannia scalaris

Fannia scalaris, also known as the latrine fly, is a fly species in the Fanniidae family. This species is smaller and more slender than the house fly, Musca domestica, and is similar in appearance to the lesser house fly, Fannia canicularis. The...



Feathered thorn

Colotois pennaria

The feathered thorn (Colotois pennaria) is a moth of the family Geometridae. It was first described by Carl Linnaeus in 1761.



Fender's blue butterfly

Icaricia icarioides fenderi

Fender's blue butterfly (Icaricia icarioides fenderi) is an endangered subspecies of Boisduval's blue (Icaricia icarioides) endemic to the Willamette Valley of northwestern Oregon, United States. The potential range of the butterfly extends from...



Feniseca

Feniseca tarquinius

Feniseca tarquinius, the harvester, is a butterfly of the family Lycaenidae, and the only member of the monotypic genus Feniseca. It is found in eastern North America. This butterfly is the only carnivorous (i.e., insectivorous) butterfly in North...



Fiery skipper

Hylephila phyleus

The fiery skipper (*Hylephila phyleus*) is a species of butterfly in the family Hesperiidae. Reaching approximately 1 inch (2.5 cm) in length, male fiery skippers are orange or yellow while the females are dark brown. Small brown spots may be...



Firebrat

Thermobia domestica

The firebrat (*Thermobia domestica*) is a small insect (typically 1–1.5 cm), in the order Zygentoma.



Flame skimmer

Libellula saturata

The flame skimmer or firecracker skimmer (*Libellula saturata*) is a common dragonfly of the family Libellulidae, native to western North America.



Florida leaf-footed bug

Acanthocephala femorata

The Florida leaf-footed bug (*Acanthocephala femorata*) is a species of insect. The genus name *Acanthocephala* means "spiny head" and comes from the pointed tylus at the tip of the head. *Acanthocephala femorata* is found in the continental United...



Florida woods cockroach

Eurycotis floridana

The Florida woods cockroach is a large cockroach species which typically grows to a length of 30–40 mm (1.2–1.6 in). When alarmed, adults can eject an extremely foul-smelling directional spray up to 1 m, which inspired several of its other common...



Foreign grain beetle

Ahasverus advena

The foreign grain beetle (Ahasverus advena) is a species of beetle in the family Silvanidae. It is related to the sawtoothed grain beetle.



Forest bug

Pentatoma rufipes

The forest bug or red-legged shieldbug (Pentatoma rufipes) is a species of shield bug in the family Pentatomidae, commonly found in most of Europe. It inhabits forests, woodlands, orchards, and gardens.



Forest ringlet

Dodonidia helmsii

The forest ringlet (Dodonidia helmsii), also known as Helms' butterfly, or te pēpepe pōuri or pepe pouri in the Māori language, is a rare butterfly of the family Nymphalidae endemic to New Zealand. It is the only species in the genus Dodonidia.



Forest tent caterpillar moth

Malacosoma disstria

The forest tent caterpillar moth (Malacosoma disstria) is a moth found throughout North America, especially in the eastern regions. Unlike related tent caterpillar species, the larvae of forest tent caterpillars do not make tents, but rather,...



Forficula auricularia

Forficula auricularia

Forficula auricularia is a species complex comprising the common earwig or European earwig, an omnivorous insect in the family Forficulidae. The European earwig survives in a variety of environments and is a common household insect in North...



Forficula smyrnensis

Forficula smyrnensis

Forficula smyrnensis is a species of earwig in the family Forficulidae. It is mostly found in the Palearctic realm.



Formica aquilonia

Formica aquilonia

Formica aquilonia, also called the Scottish wood ant, is a species of wood ant of the genus *Formica* which are widely distributed in Europe and Asia, occurring from Scandinavia in the north to Bulgaria and Italy in the south, and from the UK...



Formica exsecta

Formica exsecta

Formica exsecta (the narrow-headed ant or excised wood ant) is a species of ant found from Western Europe to Asia. A rare formicine ant with a deeply excised head, *F. exsecta* forms small mounds up to around a foot in height consisting of much...



Formica fusca

Formica fusca

Formica fusca is a black-colored ant commonly found throughout Europe as well as parts of southern Asia and Africa. It has the common names silky ant or dusky ant. The range within the palaeartic region extends from Portugal in the west to Japan...



Formica polyctena

Formica polyctena

Formica polyctena is a species of European red wood ant in the genus Formica and large family Formicidae. The species was first described by Arnold Förster in 1850. The latin species name polyctena is from Greek and literally means 'many cattle'...



Formica rufibarbis

Formica rufibarbis

Formica rufibarbis is a European formicine ant of the Formica fusca group. In the classification by Auguste Forel, it is treated in the subgenus Serviformica. F. rufibarbis is subject to a Species Action Plan (SAP) in England, where it is known...



Formosan subterranean termite

Coptotermes formosanus

The Formosan termite (Coptotermes formosanus) is a species of termite local to southern China and introduced to Taiwan (formerly known as Formosa, where it gets its name), Japan, South Africa, Sri Lanka, Hawaii, and the continental United...



Four-spotted chaser

Libellula quadrimaculata

Libellula quadrimaculata, known in Europe as the four-spotted chaser and in North America as the four-spotted skimmer, is a dragonfly of the family Libellulidae found widely throughout Europe, Asia, and North America. The adult stage is found...



Freyeria trochylus

Freyeria trochylus

Freyeria trochylus, the grass jewel, is a small butterfly found in Africa, Arabia (United Arab Emirates, Oman, Saudi Arabia), southern Europe (Bulgaria and Greece), India and southern Asia that belongs to the lycaenids or blues family.



Fulgora laternaria

Fulgora laternaria

Fulgora laternaria (often misspelled "lanternaria") is a species of Neotropical fulgorid planthopper. It is known by a large variety of common names, among them lantern fly, peanut bug, peanut-headed lanternfly, alligator bug, jequitiranaboia,...



Galeruca tanaceti

Galeruca tanaceti

Galeruca tanaceti is a species of leaf beetle found in the Palearctic realm, and is the type species of the genus Galeruca. It feeds on various plants in both its adult and larval stages. The species was first described by Carl Linnaeus in his...



Galleria mellonella

Galleria mellonella

Galleria mellonella, the greater wax moth or honeycomb moth, is a moth of the family Pyralidae. G. mellonella is found throughout the world. It is one of two species of wax moths, with the other being the lesser wax moth. G. mellonella eggs are...



Gangara thyrsis

Gangara thyrsis

Gangara thyrsis, commonly known as the giant redeye, is a species of butterfly belonging to the family Hesperidae. It breeds on a number of palm species.



Garden carpet

Xanthorhoe fluctuata

The garden carpet (Xanthorhoe fluctuata) is a moth of the family Geometridae. It is an abundant and familiar species across its huge range which covers the whole Palearctic region from Ireland to Japan and including the Near East and North...



Gatekeeper

Pyronia tithonus

The gatekeeper or hedge brown (Pyronia tithonus) is a European species of butterfly. Given its preference for warmer weather, the restriction of range expansion can be assumed to be due to climate. Colonies vary in size depending on the available...



Geotrupes stercorarius

Geotrupes stercorarius

Geotrupes stercorarius is a species of earth-boring dung beetle. Its common name is the dor beetle, or, the dumbledore, and is common throughout Europe. The beetle is up to 2.5 cm (1 in) long. The whole beetle is weakly lustrous and darkly...



Geranium bronze

Cacyreus marshalli

The geranium bronze or brun des pélargoniums in French (Cacyreus marshalli), is a butterfly in the family Lycaenidae. The geranium bronze butterfly is native to South Africa. The butterfly was first introduced to Europe in the late 20th century....



German cockroach

Blattella germanica

The German cockroach (Blattella germanica), colloquially known as the croton bug, is a species of small cockroach, typically about 1.1 to 1.6 cm (0.43 to 0.63 in) long. In color it varies from tan to almost black, and it has two dark, roughly...



Ghost moth

Hepialus humuli

The ghost moth or ghost swift (Hepialus humuli) is a moth of the family Hepialidae. It is common throughout Europe, except for the far south-east. Female ghost moths are larger than males, and exhibit sexual dimorphism with their differences in...



Giant burrowing cockroach

Macropanesthia rhinoceros

The giant burrowing cockroach (Macropanesthia rhinoceros) is also known as the rhinoceros cockroach, and Queensland giant cockroach. These cockroaches are native to Australia and mostly found in tropical and subtropical parts of Queensland. They...



Giant Fijian long-horned beetle

Xixuthrus heros

The giant Fijian long-horned beetle (Xixuthrus heros) is native to the island of Viti Levu in Fiji, and is one of the largest living insect species, with specimens around 15 cm long, excluding legs, antennae, or jaws. It is closely related to the...



Giant Hawaiian darner

Anax strenuus

The giant Hawaiian darner (Anax strenuus), also known as the giant Hawaiian dragonfly or pinao, is a species of dragonfly in the family Aeshnidae. It is one of two species of dragonfly that is endemic to the Hawaiian Islands (the other is...



Giant leopard moth

Hypercompe scribonia

The giant leopard moth (Hypercompe scribonia) is a moth of the family Erebidae. They are distributed through North America from southern Ontario, and southern and eastern United States through New England, Mexico, and south to Colombia. The...



Giraffe weevil

Trachelophorus giraffa

The giraffe weevil (Trachelophorus giraffa) is a weevil endemic to Madagascar, however there are other unrelated weevil species vernacularly named 'giraffe weevil' in Asia, such as Cynotrachelus flavotuberosus in Vietnam.



Glanville fritillary

Melitaea cinxia

The Glanville fritillary (Melitaea cinxia) is a butterfly of the family Nymphalidae. It is named for the naturalist who discovered it and the checkerboard pattern on its wings. These butterflies live in almost all of Europe, especially Finland,....



Glassy-winged sharpshooter

Homalodisca vitripennis

The glassy-winged sharpshooter (Homalodisca vitripennis, formerly known as H. coagulata) is a large leafhopper (family Cicadellidae), similar to other species of sharpshooter.



Glaucopsyche lygdamus

Glaucopsyche lygdamus

Glaucopsyche lygdamus, the silvery blue, is a small butterfly native to North America.



Golden spider beetle

Niptus hololeucus

The golden spider beetle, Niptus hololeucus, is a species of spider beetle in the family Ptinidae.



Goliathus cacicus

Goliathus cacicus

Goliathus cacicus, the chief goliath, is a species of beetles of the family Scarabaeidae.



Goliathus orientalis

Goliathus orientalis

Goliathus orientalis is a species of beetles belonging to the family Scarabaeidae.



Gonatista grisea

Gonatista grisea

Gonatista grisea, common name grizzled mantis, Florida bark mantid or lichen mimic mantid, is a species of praying mantis native to the southern United States, primarily Florida. They commonly perch on trees facing down to wait for prey, and will...



Gonepteryx cleopatra

Gonepteryx cleopatra

Gonepteryx cleopatra, the Cleopatra or Cleopatra butterfly, is a medium-sized butterfly of the family Pieridae.



Gonepteryx rhamni

Gonepteryx rhamni

Gonepteryx rhamni, commonly named the common brimstone, is a butterfly of the family Pieridae. It lives throughout the Palearctic zone and is commonly found across Europe, Asia, and North Africa. Across much of its range, it is the only species...



Gongylus gongylodes

Gongylus gongylodes

Gongylus gongylodes, also known as the wandering violin mantis, ornate mantis, or Indian rose mantis, is a species of praying mantis in the family Empusidae. Characterized by extremely slender limbs with large appendages, it is not a particularly...



Gonimbrasia belina

Gonimbrasia belina

Gonimbrasia belina is a species of emperor moth which is native to the warmer parts of southern Africa. Its large edible caterpillar, known as the mopane worm, madora, amacimbi “pigeon moth” or masontja, feeds primarily but not exclusively on...



Gothic

Naenia typica

The Gothic (Naenia typica) is a moth of the family Noctuidae. The species was first described by Carl Linnaeus in his 1758 10th edition of Systema Naturae. It is distributed in temperate Eurasia, in the Palearctic realm, including Europe, Turkey,...



Graphium agamemnon

Graphium agamemnon

Graphium agamemnon, the tailed jay, is a predominantly green and black tropical butterfly that belongs to the swallowtail family. The butterfly is also called the green-spotted triangle, tailed green jay, or green triangle. It is a common,...



Graphium antiphates

Graphium antiphates

Graphium antiphates, the five-bar swordtail, is a species of papilionid butterfly found in south and southeast Asia. The species was first described by Pieter Cramer in 1775.



Graphium cloanthus

Graphium cloanthus

Graphium cloanthus, the glassy bluebottle, is a common, non-threatened tropical butterfly of the family Papilionidae.



Graphium doson

Graphium doson

Graphium doson, the common jay, is a black, tropical papilionid (swallowtail) butterfly with pale blue semi-transparent central wing bands that are formed by large spots. There is a marginal series of smaller spots. The underside of wings is...



Graphium nomius

Graphium nomius

Graphium nomius, the spot swordtail, is a butterfly found in South and Southeast Asia that belongs to the swallowtail family. The species was first described by Eugenius Johann Christoph Esper in 1793. One of the grandest sights is a host of spot...



Graphium sarpedon

Graphium sarpedon

Graphium sarpedon, the common bluebottle or blue triangle in Australia, is a species of swallowtail butterfly that is found in South and Southeast Asia, as well as eastern Australia. There are approximately sixteen subspecies with differing...



Graphocephala coccinea

Graphocephala coccinea

Graphocephala coccinea is a meadow and woodland-dwelling species of brightly colored leafhopper native to North and Central America, from Canada south to Panama. Common names include candy-striped leafhopper, red-banded leafhopper,...



Graphocephala fennahi

Graphocephala fennahi

Graphocephala fennahi (rhododendron leafhopper) is a species of leafhopper native to the United States. Its common name derives from it feeding on the sap of rhododendrons. The species was introduced to Great Britain in the 1930s and continental...



Graphoderus bilineatus

Graphoderus bilineatus

Graphoderus bilineatus is a species of beetle in family Dytiscidae. The IUCN Red List reports it from Austria, Belgium, Bosnia and Herzegovina, Croatia, the Czech Republic, Denmark, Finland, France, Germany, Hungary, Italy, Latvia, Lithuania,...



Gray hairstreak

Strymon melinus

The gray hairstreak (Strymon melinus) is also called the bean lycaenid or cotton square borer. It is a member of the Lycaenidae family, known as the gossamer-winged butterflies and the second-largest family of butterflies. It is one of the most...



Grayling

Hipparchia semele

The grayling or rock grayling (Hipparchia semele) is a species in the brush-footed butterfly family Nymphalidae. Although found all over Europe, the grayling mostly inhabits coastal areas, with inland populations declining significantly in recent...



Great diving beetle

Dytiscus marginalis

The great diving beetle (Dytiscus marginalis) is an aquatic diving beetle native to Europe and northern Asia, and in the UK is common in Wales, much of England and southern Scotland but less common on chalk and in the far north. The great diving...



Great purple hairstreak

Atlides halesus

The great purple hairstreak (Atlides halesus), also called the great blue hairstreak, is a common gossamer-winged butterfly species in parts of the United States. It is actually a Neotropical species; its North American range only includes the...



Great spangled fritillary

Speyeria cybele

The great spangled fritillary (Speyeria cybele) is a North American butterfly of the family Nymphalidae.



Green darner

Anax junius

The green darner or common green darner (Anax junius), after its resemblance to a darning needle, is a species of dragonfly in the family Aeshnidae. One of the most common and abundant species throughout North America, it also ranges south to...



Green hairstreak

Callophrys rubi

The green hairstreak (Callophrys rubi) is a small butterfly in the family Lycaenidae.



Green shield bug

Palomena prasina

The green shield bug (*Palomena prasina*) is a European shield bug species in the family Pentatomidae. The name might equally apply to several other species in the tribe Nezarini, or if referred-to as a "green stink bug", it might more...



Green stink bug

Chinavia hilaris

The green stink bug or green soldier bug (*Chinavia hilaris*) is a stink bug of the family Pentatomidae.



Green-veined white

Pieris napi

The green-veined white (*Pieris napi*) is a butterfly of the family Pieridae.



Greenhouse whitefly

Trialeurodes vaporariorum

Trialeurodes vaporariorum, commonly known as the glasshouse whitefly or greenhouse whitefly, is an insect that inhabits the world's temperate regions. Like various other whiteflies, it is a primary insect pest of many fruit, vegetable and...



Greta oto

Greta oto

Greta oto is a species of brush-footed butterfly and member of the subfamily Danainae, tribe Ithomiini, and subtribe Godyridina. It is known by the common name glasswing butterfly for its transparent wings, which allow it to camouflage without...



Grey dagger

Acronicta psi

The grey dagger (Acronicta psi) is a moth of the family Noctuidae.



Grey pug

Eupithecia subfuscata

The grey pug (Eupithecia subfuscata) is a moth of the family Geometridae. It is found throughout the Palearctic region. It is also found in North America. Since it does not place any special demands on climatic conditions, special caterpillar...



Gryllotalpa gryllotalpa

Gryllotalpa gryllotalpa

Gryllotalpa gryllotalpa, commonly known as the European mole cricket, is widespread in Europe and has been introduced to the eastern United States. The scientific name is 'mole cricket', derived from the Latin 'gryllus' (cricket); and 'talpa'...



Gryllus texensis

Gryllus texensis

Gryllus texensis is a species of cricket described by William H. Cade and Otte in (2000). Cade and Otte clarify that field crickets collected in the Southeastern United States from Florida to Texas identified as Gryllus integer, were in fact...



Gryllus veletis

Gryllus veletis

Gryllus veletis, commonly known as the spring field cricket, is abundant throughout eastern North America. G. veletis is a solitary, aggressive, omnivorous, burrow-inhabiting species of cricket. This species is commonly confused with Gryllus...



Gulf fritillary

Dione vanillae

The Gulf fritillary or passion butterfly (Dione vanillae) is a bright orange butterfly in the subfamily Heliconiinae of the family Nymphalidae. That subfamily was formerly set apart as a separate family, the Heliconiidae. The Heliconiinae are...



Gynaephora groenlandica

Gynaephora groenlandica

Gynaephora groenlandica, the Arctic woolly bear moth, is an erebid moth native to the High Arctic in the Canadian archipelago, Greenland and Wrangel Island in Russia. It is known for its slow rate of development, as its full caterpillar life...



Habropoda laboriosa

Habropoda laboriosa

Habropoda laboriosa, the southeastern blueberry bee, is a bee in the family Apidae. It is native to the eastern United States. It is regarded as the most efficient pollinator of southern rabbiteye blueberries, because the flowers require buzz...



Hairy dragonfly

Brachytron pratense

Brachytron is a monotypic genus of European dragonfly of the family Aeshnidae containing the hairy dragonfly (Brachytron pratense), also known as the hairy hawker or spring hawker.



Halictus rubicundus

Halictus rubicundus

Halictus rubicundus, the orange-legged furrow bee, is a species of sweat bee found throughout the Northern Hemisphere. H. rubicundus entered North America from the Old World during one of two main invasions of Halictus subgenera. These invasions...



Halloween pennant

Celithemis eponina

The Halloween pennant (Celithemis eponina) is a species of dragonfly in the family Libellulidae. It is native to eastern North America, as well as the West Indies.



Halysidota tessellaris

Halysidota tessellaris

Halysidota tessellaris, also called the pale tiger moth, banded tussock moth, and tessellated halisidota, is in the family Erebiidae and the tribe Arctiini, the tiger moths. The species was first described by James Edward Smith in 1797. Like many...



Hamearis lucina

Hamearis lucina

Hamearis lucina, the Duke of Burgundy, the only member of the genus Hamearis, is a European butterfly in the family Riodinidae. For many years, it was known as the "Duke of Burgundy fritillary", because the adult's chequered pattern is strongly...



Harmonia axyridis

Harmonia axyridis

Harmonia axyridis is a large lady beetle or ladybug species that is most commonly known as the harlequin, Asian, or multicoloured asian lady beetle. This is one of the most variable species in the world, with an exceptionally wide range of colour...



Harpegnathos saltator

Harpegnathos saltator

Harpegnathos saltator, sometimes called the Indian jumping ant or Jerdon's jumping ant, is a species of ant found in India. They have long mandibles and have the ability to leap a few inches. They are large-eyed and active predators that hunt...



Hasora chromus

Hasora chromus

Hasora chromus, the common banded awl, is a butterfly belonging to the family HesperIIDae which is found in the Indian subcontinent, Southeast Asia and Australia.



Head louse

Pediculus humanus capitis

The head louse (Pediculus humanus capitis) is an obligate ectoparasite of humans. Head lice are wingless insects that spend their entire lives on the human scalp and feed exclusively on human blood. Humans are the only known hosts of this...



Heart and dart

Agrotis exclamationis

The heart and dart (*Agrotis exclamationis*) is a moth of the family Noctuidae. The species was first described by Carl Linnaeus in his 1758 10th edition of Systema Naturae. A familiar moth to many, it is considered one of the most common of the...



Heath fritillary

Melitaea athalia

The heath fritillary (*Melitaea athalia*) is a species of butterfly in the family Nymphalidae. It is found throughout the Palaearctic from western Europe to Japan, in heathland, grassland, and in coppiced woodland. Its association with coppiced...



Hebomoia glaucippe

Hebomoia glaucippe

Hebomoia glaucippe, the great orange-tip, is a butterfly belonging to the family Pieridae, that is the yellows and whites. It is found in the Indomalayan realm and Wallacea.



Hebrew character

Orthosia gothica

The Hebrew character (*Orthosia gothica*) is a moth in the family Noctuidae. The species was first described by Carl Linnaeus in his 1758 10th edition of Systema Naturae. It is found throughout Europe. Both the common and binomial names of this moth...



Heliconius charithonia

Heliconius charithonia

Heliconius charithonia, the zebra longwing or zebra heliconian, is a species of butterfly belonging to the subfamily Heliconiinae of the family Nymphalidae. It was first described by Carl Linnaeus in his 1767 12th edition of Systema Naturae. The...



Heliconius doris

Heliconius doris

Heliconius doris, the Doris longwing or Doris is a species of butterfly in the family Nymphalidae. It is known for being a polymorphic species which participates in various Müllerian mimicry rings throughout Central America and the Amazon...



Heliconius erato

Heliconius erato

Heliconius erato, or the red postman, is one of about 40 neotropical species of butterfly belonging to the genus Heliconius. It is also commonly known as the small postman, the red passion flower butterfly, or the crimson-patched longwing. It was...



Heliconius hecale

Heliconius hecale

Heliconius hecale, the tiger longwing, Hecale longwing, golden longwing or golden heliconian, is a heliconiid butterfly that occurs from Mexico to the Peruvian Amazon. Hecale, was an old woman who gave shelter to Theseus on his way to capture the...



Helicoverpa armigera

Helicoverpa armigera

Helicoverpa armigera is a species of Lepidoptera in the family Noctuidae. It is known as the cotton bollworm, corn earworm, Old World (African) bollworm, or scarce bordered straw (the lattermost in the UK, where it is a migrant). The larvae feed...



Helicoverpa zea

Helicoverpa zea

Helicoverpa zea, commonly known as the corn earworm, is a species (formerly in the genus Heliothis) in the family Noctuidae. The larva of the moth Helicoverpa zea is a major agricultural pest. Since it is polyphagous (feeds on many different...



Heliophorus epicles

Heliophorus epicles

Heliophorus epicles, commonly known as the purple sapphire, is a species of lycaenid or blue butterfly found in Asia. The species was first described by Jean Baptiste Godart in 1823.



Heliothis virescens

Heliothis virescens

Heliothis virescens, the marbled clover, is a moth of the family Noctuidae. It is found in Europe and across the Palearctic to Central Asia then to Japan, Korea and Sakhalin. In the south, it penetrates to Kashmir and Myanmar. As a migratory...



Helophilus pendulus

Helophilus pendulus

Helophilus pendulus is a European hoverfly. Its scientific name means "dangling marsh-lover" (from Greek helo-, "marsh", -phil, "love", Latin pend-, "hang"). It is a very common species in Britain, where it is the commonest Helophilus species. It...



Hemaris diffinis

Hemaris diffinis

Hemaris diffinis, the snowberry clearwing, is a moth of the family Sphingidae. This moth is sometimes called "hummingbird moth" or "flying lobster". This moth should not be confused with the hummingbird hawk-moth of Europe.



Hemiceratoides hieroglyphica

Hemiceratoides hieroglyphica

Hemiceratoides hieroglyphica is a moth from Madagascar. It was discovered in 2006 that it frequents sleeping birds at night, and drinks their tears, using a specialized, harpoon-like proboscis. Tear-feeding moths outside of Madagascar (for...



Hemiphlebia mirabilis

Hemiphlebia mirabilis

Hemiphlebia mirabilis, commonly known as the ancient greenling, is a species of damselfly, the only living species of the genus Hemiphlebia and the family Hemiphlebiidae. It is very small with a long, metallic-green body and clear wings. It is...



Hercules beetle

Dynastes hercules

The Hercules beetle (Dynastes hercules) is a species of rhinoceros beetle native to the rainforests of southern Mexico, Central America, South America, and the Lesser Antilles. It is the longest extant species of beetle in the world, and is also...



Hermetia illucens

Hermetia illucens

Hermetia illucens, the black soldier fly, is a common and widespread fly of the family Stratiomyidae. Since the late 20th century, H. illucens has increasingly been gaining attention because of its usefulness for recycling organic waste and...



Hesperia comma

Hesperia comma

Hesperia comma, the silver-spotted skipper or common branded skipper, is a butterfly of the family Hesperidae. It is known as silver-spotted skipper in Europe and common branded skipper in North America where the butterfly Epargyreus clarus, a...



Hesperia dacotae

Hesperia dacotae

Hesperia dacotae, the Dakota skipper, is a small to medium-sized North American butterfly. It has a wingspan of approximately one inch and the antennae form a hook. The male's wings are a tawny-orange to brown on the forewings with a prominent...



Hessian fly

Mayetiola destructor

The Hessian fly or barley midge, Mayetiola destructor, is a species of fly that is a significant pest of cereal crops including wheat, barley and rye. Though a native of Asia, upon its discovery it was believed to have been transported into North...



Heteropteryx

Heteropteryx dilatata

Heteropteryx is a monotypic genus of stick insects containing Heteropteryx dilatata as the only described species, and gives its name to the family of the Heteropterygidae. Their only species may be known as jungle nymph, Malaysian stick insect,...



Hierodula majuscula

Hierodula majuscula

Hierodula majuscula is a species of praying mantis in the genus Hierodula. It is also known as the giant rainforest mantis and the Australian giant mantis. It is found in coastal northern Australia, usually in rainforest and adjacent habitats....



Hierodula papua

Hierodula anglisz

Hierodula papua is a species of praying mantis in the family Mantidae.



Hierodula tenuidentata

Hierodula tenuidentata

Hierodula tenuidentata is a species of praying mantis, sharing its common name giant Asian mantis with other large members of genus Hierodula in the family Mantidae. It is native to India, and has been introduced to Greece.



Hierodula venosa

Hierodula venosa

Hierodula venosa is a species of praying mantis in the family Mantidae.



Hieroglyphus daganensis

Hieroglyphus daganensis

The African rice grasshopper, Hieroglyphus daganensis is a medium-sized grasshopper species found in the Sahel region. Although not called a locust in English, this species shows gregarious behaviour and some morphological change (phase...



High brown fritillary

Fabriciana adippe

Fabriciana adippe, the high brown fritillary, is a large and brightly colored butterfly of the family Nymphalidae, native to Europe and across the Palearctic to Japan. It is known for being Great Britain's most threatened butterfly and is listed...



Highland midge

Culicoides impunctatus

The highland midge (scientific name: Culicoides impunctatus; Scots: Midgie; Scottish Gaelic: Meanbh-chuileag) is a species of biting midge found across the Palearctic (throughout the British Isles, Scandinavia, other regions of Northern Europe....



Hine's emerald

Somatochlora hineana

The Hine's emerald (*Somatochlora hineana*) is an endangered dragonfly species found in the United States and Canada. Populations exist in Illinois, Michigan, Missouri, Ontario, and Wisconsin. Larvae are found in shallow, flowing water in fens and...



Hippodamia convergens

Hippodamia convergens

Hippodamia convergens, commonly known as the convergent lady beetle, is one of the most common lady beetles in North America and is found throughout the continent. Aphids form their main diet and they are used for the biological control of these...



Hippodamia tredecimpunctata

Hippodamia tredecimpunctata

Hippodamia tredecimpunctata, commonly known as the thirteen-spot ladybeetle, is a species of lady beetle.



Holly blue

Celastrina argiolus

The holly blue (*Celastrina argiolus*) is a butterfly that belongs to the lycaenids or blues family and is native to the Palearctic.



Horse guard wasp

Stictia carolina

The horse guard wasp (*Stictia carolina*) is a type of sand wasp (*Bembicini*) from the eastern United States which preys primarily upon horse-flies (*Tabanidae*). It is a large, colorful, fast-flying wasp, one of 28 species in the genus *Stictia* (which...



Horse-chestnut leaf miner

Cameraria ohridella

The horse-chestnut leaf miner (Cameraria ohridella) is a leaf-mining moth of the family Gracillariidae. The horse-chestnut leaf miner was first observed in North Macedonia in 1984, and was described as a new species in 1986. Its larvae are leaf...



House cricket

Acheta domesticus

Acheta domesticus, commonly called the house cricket, is a cricket most likely native to Southwestern Asia, but between 1950 and 2000 it became the standard feeder insect for the pet and research industries and spread worldwide. They can be kept...



Huhu beetle

Prionoplus reticularis

The huhu beetle (Prionoplus reticularis) is a longhorn beetle endemic to New Zealand. It is the heaviest beetle found in New Zealand.



Hummingbird hawk-moth

Macroglossum stellatarum

The hummingbird hawk-moth (Macroglossum stellatarum) is a species of hawk moth found across temperate regions of Eurasia. The species is named for its similarity to hummingbirds, as they feed on the nectar of tube-shaped flowers using their long...



Hyalophora cecropia

Hyalophora cecropia

Hyalophora cecropia, the cecropia moth, is North America's largest native moth. It is a member of the family Saturniidae, or giant silk moths. Females have been documented with a wingspan of five to seven inches (13 to 18 cm) or more. These moths...



Hyalophora columbia

Hyalophora columbia

Hyalophora columbia, the Columbia silkmoth or larch silkmoth, is a moth of the family Saturniidae. In the east it is found from Quebec and Ontario to Michigan, northern Wisconsin, and south-eastern Manitoba. In the west it is found from Alberta...



Hyles euphorbiae

Hyles euphorbiae

Hyles euphorbiae, the spurge hawk-moth, is a European moth of the family Sphingidae. It has been found in Pontresina, Switzerland (altitude of 1,805m). This hawk moth is used as an agent of biological pest control against the noxious weed leafy...



Hyles gallii

Hyles gallii

Hyles gallii, the bedstraw hawk-moth or galium sphinx, is a moth of the family Sphingidae. The species was first described by S. A. von Rottemburg in 1775.



Hyles livornica

Hyles livornica

Hyles livornica, the striped hawk-moth, is a moth of the family Sphingidae.



Hyles livornicoides

Hyles livornicoides

Hyles livornicoides, the Australian striped hawk moth, is a moth of the family Sphingidae first described by Lucas in 1892. The larvae are known by the common name Yeperenye caterpillar, in the Arrernte language the caterpillars are referred to...



Hyles tithymali

Hyles tithymali

Hyles tithymali, the Barbary spurge hawk-moth, is a species of moth in the family Sphingidae first described by Jean Baptiste Boisduval in 1834. Genomic analysis places the entire species as a subspecies of Hyles euphorbiae. It is found in North...



Hylobius abietis

Hylobius abietis

Hylobius abietis or the large pine weevil is a beetle belonging to family Curculionidae. This species is widely regarded as the most important pest of most commercially important coniferous trees in European plantations. Seedlings planted or...



Hylotrupes

Hylotrupes bajulus

Hylotrupes is a monotypic genus of woodboring beetles in the family Cerambycidae, the longhorn beetles. The sole species, Hylotrupes bajulus, is known by several common names, including house longhorn beetle, old house borer, and European house...



Hymenoepimecis argyraphaga

Hymenoepimecis argyraphaga

Hymenoepimecis argyraphaga is a Costa Rican parasitoid wasp whose host is the spider Plesiomete argyra. The wasp is unusual in modifying the spider's web building behavior to make a web made of very strong lines designed to support the wasp's...



Hypolimnas bolina

Hypolimnas bolina

Hypolimnas bolina, the great eggfly, common eggfly, varied eggfly, or in New Zealand the blue moon butterfly, is a species of nymphalid butterfly found from Madagascar to Asia and Australia.



Hyposmocoma molluscivora

Hyposmocoma molluscivora

Hyposmocoma molluscivora is a Hawaiian moth whose larvae are predators, capturing snails in their silk, much like a hunting spider's web, and then crawling inside the snail's shell to eat it alive. It has been called the snail-eating caterpillar...



Iambrix salsala

Iambrix salsala

Iambrix salsala, the chestnut bob, is a butterfly belonging to the family Hesperiiidae, that is found in parts of South Asia and Southeast Asia.



Icaricia icarioides

Icaricia icarioides

Icaricia icarioides, or Boisduval's blue, is a butterfly of the family Lycaenidae found in North America. This butterfly has 25 recognized subspecies. This species has been classified in at least four different genera since it was named by Jean...



Idea leuconoe

Idea leuconoe

Idea leuconoe, also known as the paper kite butterfly, rice paper butterfly, large tree nymph, is a butterfly known especially for its presence in butterfly houses and live butterfly expositions. It has a wingspan of 12 to 14 cm. The paper kite...



Idea malabarica

Idea malabarica

Idea malabarica, the Malabar tree nymph, is a large butterfly found in peninsular India. that belongs to the danaid group of the family Nymphalidae. It is found in forest clearings and above the forest canopy.



Ideopsis vulgaris

Ideopsis vulgaris

Ideopsis vulgaris, the blue glassy tiger, is a butterfly that belongs to the crows and tigers, that is, the danaid group of the brush-footed butterflies family.



Idia concisa

Idia concisa

Idia concisa, the pale-winged idia (also known as Idia sp. nr. aemula since it has not been formally described) is a litter moth of the family Erebidae. The species was first described by William Trowbridge Merrifield Forbes in 1954. It is...



Ignelater luminosus

Ignelater luminosus

Ignelater luminosus is a bioluminescent species of click beetle native to the island of Puerto Rico, one of several Caribbean species in the genus Ignelater that are known as cucubanos. Cucubanos are often confused with fireflies, which are in a...



Indianmeal moth

Plodia interpunctella

The Indianmeal moth (Plodia interpunctella), also spelled Indian meal moth and Indian-meal moth, is a pyraloid moth of the family Pyralidae. Alternative common names are hanger-downers, weevil moth, pantry moth, flour moth or grain moth. The...



Iphiclides feisthamelii

Iphiclides feisthamelii

Iphiclides feisthamelii, the southern scarce swallowtail, southern swallowtail or Iberian scarce swallowtail, is a butterfly found in Italy, Slovenia, southern France, Spain, Portugal, Morocco, Algeria and Tunisia. It is sometimes considered a...



Ischnura aurora

Ischnura aurora

Ischnura aurora, the gossamer damselfly or golden dartlet and also known as the aurora bluetail, is a species of damselfly in the family Coenagrionidae.



Ischnura heterosticta

Ischnura heterosticta

Ischnura heterosticta, one of at least two species with the common name common bluetail, is an Australian damselfly of the family Coenagrionidae. It is the largest of the three Ischnura species in Australia. They are generally found near...



Ischnura senegalensis

Ischnura senegalensis

Ischnura senegalensis, also known variously as common bluetail, marsh bluetail, ubiquitous bluetail, African bluetail, and Senegal golden dartlet, is a widespread damselfly of the family Coenagrionidae. It is native from Africa, through the...



Italian bee

Apis mellifera ligustica

Apis mellifera ligustica is the Italian bee which is a subspecies of the western honey bee (Apis mellifera).



Ixodiphagus hookeri

Ixodiphagus hookeri

Ixodiphagus hookeri, the tick wasp, is an encyrtid wasp which lays its eggs into ticks. It seems to use a symbiotic bacteria, Wolbachia pipientis, to weaken the tick's immune system.



Jack jumper ant

Myrmecia pilosula

The jack jumper ant (Myrmecia pilosula), also known as the jack jumper, jumping jack, hopper ant, or jumper ant, is a species of venomous ant native to Australia. Most frequently found in Tasmania and southeast mainland Australia, it is a member...



Japanese beetle

Popillia japonica

The Japanese beetle (Popillia japonica) is a species of scarab beetle. The adult measures 15 mm (0.6 in) in length and 10 mm (0.4 in) in width, has iridescent copper-colored elytra, and a green thorax and head. Due to natural predators, the...



Japanese rhinoceros beetle

Allomyrina dichotoma

Allomyrina dichotoma, also known as Japanese rhinoceros beetle, Japanese horned beetle, or kabutomushi (カブトムシ, lit. 'Kabuto insect'), is a species of rhinoceros beetle.



Juniper carpet

Thera juniperata

The juniper carpet (Thera juniperata) is a moth of the family Geometridae. The species was first described by Carl Linnaeus in his 1758 10th edition of Systema Naturae. It is found throughout Europe and the Near East, but is rather uncommon and...



Junonia atlites

Junonia atlites

Junonia atlites, the grey pansy, is a species of nymphalid butterfly found in South Asia.



Junonia coenia

Junonia coenia

Junonia coenia, known as the common buckeye or buckeye, is a butterfly in the family Nymphalidae. Its range covers much of North America and some of Central America, including most of the eastern half of the US, the lower to middle Midwest, the...



Junonia genoveva

Junonia genoveva

Junonia genoveva, the mangrove buckeye, is a butterfly of the family Nymphalidae. The species was first described by Pieter Cramer in 1780. It is found in South America, and possibly into Central America. The wingspan is 45–57 mm. The butterfly is...



Junonia lemonias

Junonia lemonias

Junonia lemonias, the lemon pansy, is a common nymphalid butterfly found in Cambodia and South Asia. It is found in gardens, fallow land, and open wooded areas.



Kallima inachus

Kallima inachus

Kallima inachus, the orange oakleaf, Indian oakleaf or dead leaf, is a nymphalid butterfly found in Tropical Asia from India to Japan. With wings closed, it closely resembles a dry leaf with dark veins and is a commonly cited example of camouflage.



Kamehameha butterfly

Vanessa tameamea

The Kamehameha butterfly (Vanessa tameamea) is one of the two species of butterfly endemic to Hawaii, the other is Udara blackburni. The Hawaiian name is pulelehua. This is today a catch-all native term for all butterflies; its origin seems to be...



Kaniska canace

Kaniska canace

Kaniska canace, the blue admiral, is a nymphalid butterfly, the only species of the genus Kaniska. It is found in south and southeast Asia.



Karner blue

Plebejus samuelis

The Karner blue (Plebejus samuelis) is an endangered species of small blue butterfly found in some Great Lakes states, small areas of New Jersey, the Capital District region of New York, and southern New Hampshire (where it is the official state...



Khapra beetle

Trogoderma granarium

The khapra beetle (Trogoderma granarium), also called cabinet beetle, which originated in South Asia, is one of the world's most destructive pests of grain products and seeds. It is considered one of the 100 worst invasive species in the world....



Kricogonia lyside

Kricogonia lyside

Kricogonia lyside, the lyside sulphur or guayacan sulphur, is a North American, Caribbean, and South American butterfly in the family Pieridae. In seasons with heavy monsoons, this butterfly is seen in massive migrations which are frequent in...



Labidura riparia

Labidura riparia

Labidura riparia is a species of earwig in the family Labiduridae characterized by their modified cerci as forceps, and light tan color. They are commonly known as the shore earwig, tawny earwig, riparian earwig, or the striped earwig due to two...



Lampides boeticus

Lampides boeticus

Lampides boeticus, the pea blue, or long-tailed blue, is a small butterfly that belongs to the lycaenids or gossamer-winged family.



Lamproptera meges

Lamproptera meges

Lamproptera meges, the green dragontail, is a species of swallowtail butterfly (family Papilionidae) found in parts of South Asia and Southeast Asia. There are ten subspecies. A specimen from Java is the type species of the genus Lamproptera.



Laothoe populi

Laothoe populi

Laothoe populi, the poplar hawk-moth, is a moth of the family Sphingidae. The species was first described by Carl Linnaeus in his 1758 10th edition of Systema Naturae. It is found throughout the Palearctic region and the Near East and is one of...



Large blue

Phengaris arion

The large blue (Phengaris arion) is a species of butterfly in the family Lycaenidae. The species was first defined in 1758 and first recorded in Britain in 1795. In 1979 the species became mostly extinct in Britain but has been successfully...



Large chequered skipper

Heteropterus morpheus

The large chequered skipper (Heteropterus morpheus) is a species of butterfly in the family Hesperiidae. It is the single member of the monotypic genus Heteropterus. The species can be found in isolated populations in Europe and east across the...



Large copper

Lycaena dispar

The large copper (Lycaena dispar) is a butterfly of the family Lycaenidae. L. dispar has been commonly arranged into three subspecies: L. dispar dispar, (single-brooded) which was commonly found in England, but is now extinct, L. d. batavus,...



Large emerald

Geometra papilionaria

The large emerald (Geometra papilionaria) is a moth which is the type species for the family Geometridae. It is found throughout the Palearctic region and the Near East in and around deciduous forests, heathlands, marshland and in settlements...



Large milkweed bug

Oncopeltus fasciatus

Oncopeltus fasciatus, known as the large milkweed bug, is a medium-sized hemipteran (true bug) of the family Lygaeidae. It is distributed throughout North America, from Central America through Mexico and the Caribbean to southern areas in Canada....



Large red damselfly

Pyrrosoma nymphula

The large red damselfly (Pyrrosoma nymphula) is a species of damselflies belonging to the family Coenagrionidae. It is native to the western Palearctic.



Large skipper

Ochlodes sylvanus

The large skipper (Ochlodes sylvanus) is a butterfly of the family Hesperidae.



Large tortoiseshell

Nymphalis polychloros

The large tortoiseshell or blackleg tortoiseshell (Nymphalis polychloros) is a butterfly of the family Nymphalidae.



Large yellow underwing

Noctua pronuba

The large yellow underwing (Noctua pronuba) is a moth, the type species for the family Noctuidae. It is an abundant species throughout the Palearctic realm, one of the most common and most familiar moths of the region. In some years the species...



Lasiommata maera

Lasiommata maera

Lasiommata maera, the large wall brown, is a butterfly in the family Nymphalidae.



Lasiommata megera

Lasiommata megera

Lasiommata megera, the wall or wall brown, is a butterfly in the family Nymphalidae (subfamily Satyrinae). It is widespread in the Palearctic realm with a large variety of habitats and number of generations a year.



Laysan noctuid moth

Agrotis laysanensis

The Laysan noctuid moth (Agrotis laysanensis) is a species of moth in the family Noctuidae. It is endemic to Laysan, in the Northwestern Hawaiian Islands. This moth was one, if not the main species, eaten by the extinct Laysan millerbird. Agrotis...



Leek moth

Acrolepiopsis assectella

The leek moth or onion leaf miner (Acrolepiopsis assectella) is a species of moth of family Acrolepiidae (formerly Glyphipterigidae) and the genus Acrolepiopsis. The species is native to Europe and Siberia, but is also found in North America,...



Lemonia dumi

Lemonia dumi

Lemonia dumi is a species of moth of the family Brahmaeidae (older classifications placed it in Lemoniidae). It is found in scattered populations in Central Europe. The wingspan is 45–65 mm. The moth flies from October to November depending on the...



Lepidosaphes ulmi

Lepidosaphes ulmi

Lepidosaphes ulmi also known as apple mussel scale or oystershell scale is a widely invasive scale insect that is a pest of trees and woody plants. The small insects attach themselves to bark and cause injury by sucking the tree's sap; this...



Leptidea sinapis

Leptidea sinapis

Leptidea sinapis, or the wood white butterfly of the family Pieridae, is a small white butterfly that is mainly found in England, Ireland, and Northern Europe. The butterfly has white wings with grey or yellow markings near the center or tip of...



Leptoconops torrens

Leptoconops torrens

Leptoconops torrens (commonly known as the biting midge fly) is a species of small biting flies in the no-see-um family Ceratopogonidae. They were first mentioned in writing by Charles Henry Tyler Townsend in 1893. The name Leptoconops carteri is...



Leptocybe invasa

Leptocybe invasa

Leptocybe invasa, the blue gum chalcid wasp or eucalyptus gall wasp, is a chalcid wasp which is the only species in the monotypic genus Leptocybe in the subfamily Tetrastichinae, of the family Eulophidae. It is a gall wasp which causes the...



Leptoglossus phyllopus

Leptoglossus phyllopus

Leptoglossus phyllopus or eastern leaf-footed bug is a species of leaf-footed bugs in the same genus as the western conifer seed bug (L. occidentalis. The Eastern leaf-footed bug is found throughout the southern United States, from Florida to...



Leptosia nina

Leptosia nina

Leptosia nina, the psyche, is a small butterfly of the family Pieridae (the sulphurs, yellows and whites) and is found in Indian subcontinent, southeast Asia and Australia. The upper forewing has a black spot on a mainly white background. The...



Leptotes plinius

Leptotes plinius

Leptotes plinius, the zebra blue or plumbago blue, is a species of blue butterfly (Lycaenidae) found in Sri Lanka, India to Australia. The species was first described by Johan Christian Fabricius in 1793.



Lesser water boatman

Corixa punctata

The lesser waterboatman or lesser water boatman (Corixa punctata) is a water-dwelling insect of the order Hemiptera. Adults normally range in size from 5 to 15 mm long, and are found in ponds, lakes and sometimes even swimming pools. The boatman...



Lesser yellow underwing

Noctua comes

The lesser yellow underwing (Noctua comes) is a moth of the family Noctuidae.



Lexias dirtea

Lexias dirtea

Lexias dirtea, the archduke, is a species of butterfly of the family Nymphalidae.



Libellula depressa

Libellula depressa

Libellula depressa, the broad-bodied chaser or broad-bodied darter, is one of the most common dragonflies in Europe and central Asia. It is very distinctive with a very broad flattened abdomen, four wing patches and, in the male, the abdomen...



Light brown apple moth

Epiphyas postvittana

The light brown apple moth (Epiphyas postvittana) (often abbreviated to LBAM) is a leafroller moth belonging to the lepidopteran family Tortricidae.



Lime-speck pug

Eupithecia centaureata

The lime-speck pug (Eupithecia centaureata) is a moth of the family Geometridae. It is a common species throughout the Palearctic region (where it is found in Europe, Central Asia, Mongolia, southern Siberia, eastern China (Guangdong) and...



Limenitis arthemis

Limenitis arthemis

Limenitis arthemis, the red-spotted purple or white admiral, is a North American butterfly species in the cosmopolitan genus Limenitis. It has been studied for its evolution of mimicry, and for the several stable hybrid wing patterns within this...



Limenitis camilla

Limenitis camilla

Limenitis camilla, the (Eurasian) white admiral, is a butterfly of the family Nymphalidae. It is found in woodland throughout southern Britain and much of Europe and the Palearctic, extending as far east as Japan. Adult white admirals have dark...



Limenitis reducta

Limenitis reducta

Limenitis reducta, the southern white admiral, is a butterfly of the family Nymphalidae.



Liphyra brassolis

Liphyra brassolis

Liphyra brassolis, the moth butterfly, is a butterfly found in South Asia, Southeast Asia and Australia that belongs to the lycaenid family. The larvae are predatory and feed on ant larvae. This is one of the largest species of lycaenid...



Lipoptena cervi

Lipoptena cervi

Lipoptena cervi, the deer ked or deer fly, is a species of biting fly in the family of louse flies, Hippoboscidae. These flies are commonly encountered in temperate areas of Europe, Siberia, and northern China. They have been introduced to North...



Lipoptena depressa

Lipoptena depressa

Lipoptena depressa, or the Western American deer ked, is species of fly in the family Hippoboscidae. The flies are blood-feeding ectoparasites of mule deer, Odocoileus hemionus and white-tailed deer, Odocoileus virginianus. They are found in the...



Loepa katinka

Loepa katinka

Loepa katinka, the golden emperor moth, is a species of moth belonging to the genus Loepa of the family Saturniidae. It is found in Southeast Asia and South Asia. Its wingspan measures 95 mm and its body is 30 mm long.



Long-tailed skipper

Urbanus proteus

The long-tailed skipper (Urbanus proteus) is a spread-winged skipper butterfly found throughout tropical and subtropical South America, south to Argentina and north into the eastern United States and southern Ontario. It cannot live in areas with...



Lophocampa caryae

Lophocampa caryae

Lophocampa caryae, the hickory tiger moth, hickory tussock moth, or hickory halisidota, is a moth in the family Erebididae and the tribe Arctiini, the tiger moths. The species is widely distributed in the eastern half of North America. In other...



Lorquin's admiral

Limenitis lorquini

Lorquin's admiral (Limenitis lorquini) is a butterfly in the Nymphalinae subfamily. The butterfly is named after Pierre Joseph Michel Lorquin, a French naturalist who came to California from France during the Gold Rush, and made important...



Losaria coon

Losaria coon

Losaria coon, the common clubtail, is a butterfly belonging to the swallowtail family, Papilionidae. The butterfly belongs to the clubtails, genus Losaria. It includes several subspecies and is found from the Nicobar Islands and Assam in India,...



Lovebug

Plecia nearctica

The lovebug (Plecia nearctica) is a species of march fly found in parts of Central America and the southeastern United States, especially along the Gulf Coast. It is also known as the honeymoon fly or double-headed bug. During and after mating,...



Lucanus capreolus

Lucanus capreolus

Lucanus capreolus, the reddish-brown stag beetle or pinching beetle, is a beetle of the family Lucanidae. The specific name capreolus is derived from Latin, meaning "roe deer" (capreolus literally means "little goat"). The name refers to the...



Lucanus cervus

Lucanus cervus

Lucanus cervus, known as the European stag beetle, or the greater stag beetle, is one of the best-known species of stag beetle (family Lucanidae) in Western Europe, and is the eponymous example of the genus. L. cervus is listed as Near Threatened...



Lucanus maculifemoratus

Lucanus maculifemoratus

Lucanus maculifemoratus is a beetle of the family Lucanidae. This species is known as Miyama stag beetle (Deep Mountain stag beetle) in Japan. These beetles have a light covering of golden hair on their back, mostly around the rear edges.Lucanus...



Lucilia bufonivora

Lucilia bufonivora

Lucilia bufonivora is a member of the fly family Calliphoridae which are commonly known as blow flies. L. bufonivora is commonly referred to as a toadfly. The adult flies will typically feed on pollen and nectar of flowers, while the larvae are...



Lucilia coeruleiviridis

Lucilia coeruleiviridis

Lucilia coeruleiviridis, formerly Phaenecia coeruleiviridis, is commonly known as a green bottle fly, because of its metallic blue-green thorax and abdomen. L. coeruleiviridis was first discovered by French entomologist Pierre-Justin-Marie...



Lucilia illustris

Lucilia illustris

Lucilia illustris is a member of the fly family Calliphoridae, commonly known as a blow fly. Along with several other species, L. illustris is commonly referred to as a green bottle fly. Lucilia illustris is typically 6–9 mm in length and has a...



Lulworth skipper

Thymelicus acteon

The Lulworth skipper (Thymelicus acteon) is a butterfly of the family Hesperiidae. Its name is derived from Lulworth Cove in the county of Dorset, England, where the first specimens in Great Britain were collected in 1832 by English naturalist...



Luna moth

Actias luna

The luna moth (Actias luna), also called the American moon moth, is a Nearctic moth in the family Saturniidae, subfamily Saturniinae, a group commonly named the giant silk moths. The moth has lime-green wings and a white body. Its caterpillars are...



Luthrodes pandava

Luthrodes pandava

Luthrodes pandava (formerly Chilades pandava), the plains Cupid or cycad blue, is a species of lycaenid butterfly found in Pakistan, India, Sri Lanka, Myanmar, United Arab Emirates, Indochina, Peninsular Malaysia, Singapore, Taiwan, Java, Sumatra...



Lycaena epixanthe

Lycaena epixanthe

Lycaena epixanthe, also known as the bog copper or cranberry-bog copper, is a North American species of butterfly in the family Lycaenidae. Adults like to sip drops of dew clinging to leaves and almost exclusively nectar on their host plant...



Lycaena hyllus

Lycaena hyllus

Lycaena hyllus, the bronze copper, is a butterfly of the lycaenids family found in North America.



Lycaena phlaeas

Lycaena phlaeas

Lycaena phlaeas, the small copper, American copper, or common copper, is a butterfly of the Lycaenids or gossamer-winged butterfly family. According to Guppy and Shepard (2001), its specific name phlaeas is said to be derived either from the...



Lycaena tityrus

Lycaena tityrus

Lycaena tityrus, the sooty copper, is a butterfly of the family Lycaenidae. It is found in Europe.



Lychnis

Hadena bicruris

The lychnis (Hadena bicruris) is a moth of the family Noctuidae. It is found in northern and western Europe and Turkey. It has an Atlantic-Mediterranean distribution. In the East Palearctic it is replaced by Hadena capsincola.



Lymantria dispar

Lymantria dispar

Lymantria dispar, also known as the gypsy moth or the spongy moth, is an Eurasian species of moth in the family Erebidae. Lymantria dispar is subdivided into several subspecies, with subspecies such as L. d. dispar and L. d. japonica being...



Maconellicoccus hirsutus

Maconellicoccus hirsutus

Maconellicoccus hirsutus, (also known as the pink, grape or hibiscus mealybug) is a pest of many plants, trees, and shrubs. It infests hibiscus, citrus, coffee, sugar cane, annonas, plums, guava, mango (although, mango mealybug is more...



Macrodonia cervicornis

Macrodonia cervicornis

Macrodonia cervicornis (Linnaeus, 1758), also known as the sabertooth longhorn beetle, is one of the largest beetles, if one allows for the enormous mandibles of the males, from which it derives both of the names in its binomen: Macrodonia...



Madagascar hissing cockroach

Gromphadorhina portentosa

The Madagascar hissing cockroach (Gromphadorhina portentosa), also known as the hissing cockroach or simply hisser, is one of the largest species of cockroach, reaching 5 to 7.5 centimetres (2 to 3 inches) at maturity. They are native to the...



Madeiran large white

Pieris brassicae wollastoni

The Madeiran large white (Pieris brassicae wollastoni) is a subspecies of the large white butterfly, endemic to Madeira. It was described by the English entomologist, Arthur Gardiner Butler in 1886.



Maladera insanabilis

Maladera insanabilis

Maladera insanabilis, sometimes colloquially called "Khomeini beetle", is a species of a beetle in the family Scarabaeidae. It is prevalent in Iran and other countries in the Middle-East. Adults are active in the summer and in the spring. Males...



Manduca quinquemaculata

Manduca quinquemaculata

Manduca quinquemaculata, the five-spotted hawkmoth, is a brown and gray hawk moth of the family Sphingidae. The caterpillar, often referred to as the tomato hornworm, can be a major pest in gardens; they get their name from a dark projection on...



Manduca rustica

Manduca rustica

Manduca rustica, the rustic sphinx, is a moth of the family Sphingidae. The species was first described by Johan Christian Fabricius in 1775.



Manduca sexta

Manduca sexta

Manduca sexta is a moth of the family Sphingidae present through much of the Americas. The species was first described by Carl Linnaeus in his 1763 Centuria Insectorum. Commonly known as the Carolina sphinx moth and the tobacco hawk moth (as...



Manticora scabra

Manticora scabra

Manticora scabra is a species of tiger beetle native to Mozambique, Transvaal and Zimbabwe.



Map

Araschnia levana

The map (Araschnia levana) is a butterfly of the family Nymphalidae.



Marchalina hellenica

Marchalina hellenica

Marchalina hellenica is a scale insect that lives in the eastern Mediterranean region, mainly in Greece and Turkey. It is an invasive species in Melbourne, Australia. It lives by sucking the sap of pine trees, mainly the Turkish Pine (Pinus...



Marsh fritillary

Euphydryas aurinia

The marsh fritillary (Euphydryas aurinia) is a butterfly of the family Nymphalidae. Commonly distributed in the Palearctic region, the marsh fritillary's common name derives from one of its several habitats, marshland. The prolonged larval stage...



Maruca vitrata

Maruca vitrata

Maruca vitrata is a pantropical insect pest of leguminous crops like pigeon pea, cowpea, mung bean and soybean. Its common names include the maruca pod borer, bean pod borer, soybean pod borer, mung moth, and the legume pod borer. The species was...



Mastotermes darwiniensis

Mastotermes darwiniensis

Mastotermes darwiniensis, common names giant northern termite and Darwin termite, is a termite species found only in northern Australia. It is the most primitive extant termite species. Contrary to common belief, this species does not form mounds...



Meadow argus

Junonia villida

The meadow argus (*Junonia villida*) is a butterfly in the family Nymphalidae, commonly found in Australia and Nelsons Island. It is also known as Albin's Hampstead eye in the United Kingdom, where it has occurred only as an accidental import.



Meadow brown

Maniola jurtina

The meadow brown (*Maniola jurtina*) is a butterfly found in the Palearctic realm. Its range includes Europe south of 62°N, Russia eastwards to the Urals, Asia Minor, Iraq, Iran, North Africa and the Canary Islands. The larvae feed on grasses.



Mealworm

Tenebrio molitor

Mealworms are the larval form of the yellow mealworm beetle, *Tenebrio molitor*, a species of darkling beetle. Like all holometabolic insects, they go through four life stages: egg, larva, pupa, and adult. Larvae typically measure about 2.5...



Meandrusa payeni

Meandrusa payeni

Meandrusa payeni, the yellow gorgon, is a species of swallowtail found in parts of South Asia and Southeast Asia. It belongs to the hooked swallowtails genus, *Meandrusa*, of the family Papilionidae. It is also called the outlet sword or the sickle.



Median wasp

Dolichovespula media

The median wasp (*Dolichovespula media*) is a species of social wasp of the family Vespidae found throughout Europe and Asia. It builds aerial paper nests often in shrubs or trees, and occasionally under the eaves of buildings. It is most common to...



Mediterranean flour moth

Ephestia kuehniella

The Mediterranean flour moth or mill moth (Ephestia kuehniella) is a moth of the family Pyralidae. It is a common pest of cereal grains, especially flour. This moth is found throughout the world, especially in countries with temperate climates....



Megachile pluto

Megachile pluto

Megachile pluto, also known as Wallace's giant bee or raja ofu (lit. 'king of the bees'), is a large resin bee found in Indonesia. With a wingspan of 63.5 mm (2.5 in) it is the largest known living bee species. It was believed to be extinct until...



Megachile rotundata

Megachile rotundata

Megachile rotundata, the alfalfa leafcutting bee, is a European bee that has been introduced to various regions around the world. As a solitary bee species, it does not build colonies or store honey, but is a very efficient pollinator of alfalfa,...



Megalopyge crispata

Megalopyge crispata

Megalopyge crispata, the black-waved flannel moth, crinkled flannel moth or white flannel moth, is a moth of the Megalopygidae family. It is found along the east coast of the United States, and as far inland as Oklahoma. This wingspan is 25–40 mm....



Megalopyge opercularis

Megalopyge opercularis

Megalopyge opercularis is a moth of the family Megalopygidae. It has numerous common names, including southern flannel moth for its adult form, and puss caterpillar, asp, Italian asp, fire caterpillar, woolly slug, opossum bug, puss moth, tree...



Megaselia scalaris

Megaselia scalaris

The fly Megaselia scalaris is a member of the order Diptera and the family Phoridae, and it is widely distributed in warm regions of the world. The family members are commonly known as the "humpbacked fly", the "coffin fly", and the "scuttle..."



Melanargia galathea

Melanargia galathea

Melanargia galathea, the marbled white, is a medium-sized butterfly in the family Nymphalidae. Despite its common name and appearance, this butterfly is one of the "browns", of the subfamily Satyrinae. This species can be found across most of...



Melangyna viridiceps

Melangyna viridiceps

Melangyna viridiceps is an Australian hoverfly, known as the common hover fly.



Melanitis leda

Melanitis leda

Melanitis leda, the common evening brown, is a common species of butterfly found flying at dusk. The flight of this species is erratic. They are found in Africa, South Asia and South-east Asia extending to parts of Australia.



Melanoplus bivittatus

Melanoplus bivittatus

Melanoplus bivittatus, the two-striped grasshopper, is a poikilothermic species of grasshopper belonging to the genus Melanoplus. It is commonly found in North America, with high quantities inhabiting Canadian prairies and farmland.



Mellinus arvensis

Mellinus arvensis

Mellinus arvensis, the field digger wasp, is a species of solitary wasp. The wasp can commonly be found from July to late September or October in sandy places. In hard soil however the female will often try to steal a nest from another member of...



Meloimorpha japonica

Meloimorpha japonica

Meloimorpha japonica, also known as suzumushi (鈴虫, lit. 'bell cricket'), the bell cricket, and the bell-ring cricket, is a species of cricket widespread in Asia (from India, through Indochina to Japan). It is known particularly for its chiming...



Metallyticus splendidus

Metallyticus splendidus

Metallyticus splendidus (informally known as the iridescent bark mantis) is a rare species of praying mantis found in Southeast Asia. It has an iridescent appearance.



Mexican bean beetle

Epilachna varivestis

The Mexican bean beetle (Epilachna varivestis) is a species of lady beetle that can be an agricultural pest. It is one of the few North American lady beetles that feed on plants rather than other insects. It is found throughout Mexico and the...



Mezium americanum

Mezium americanum

Mezium americanum, the American spider beetle or black spider beetle, is a species of beetle in the subfamily Ptininae, the spider beetles. These are sometimes mistaken for spiders or mites because of their rounded abdomens and long legs. It has...



Migrant hawker

Aeshna mixta

The migrant hawker (Aeshna mixta) is one of the smaller species of hawker dragonflies. It can be found away from water but for breeding it prefers still or slow-flowing water and can tolerate brackish sites. The flight period is from July to the...



Migratory locust

Locusta migratoria

The migratory locust (Locusta migratoria) is the most widespread locust species, and the only species in the genus Locusta. It occurs throughout Africa, Asia, Australia and New Zealand. It used to be common in Europe but has now become rare...



Miller

Acronicta leporina

The miller (Acronicta leporina) is a moth of the family Noctuidae. It is found throughout Europe apart from the far south-east. The range extends from the South of Spain, Central Italy and Bulgaria to Scotland and Central Scandinavia, crossing...



Mimas tiliae

Mimas tiliae

Mimas tiliae, the lime hawk-moth, is a moth of the family Sphingidae. It is found throughout the Palearctic region and the Near East, and has also been identified in Canada's east and western provinces and in northern Spain (Europe). The species...



Mint moth

Pyrausta aurata

The mint moth (Pyrausta aurata) is a small moth from the family Crambidae, also known by the common name Small Purple and Gold.



Miomantis caffra

Miomantis caffra

Miomantis caffra (common name: springbok mantis) is a species of praying mantis native to southern Africa. It appeared in New Zealand in 1978, and was found more recently in Portugal and Los Angeles, USA, likely spread through the exotic pet...



Miomantis paykullii

Miomantis paykullii

Miomantis paykullii is a species of praying mantis in the family Miomantidae. It is one of several species sometimes known as the Egyptian praying mantis, along with Miomantis abyssinica. The species is found in a number of African countries....



Mission blue butterfly

Icaricia icarioides missionensis

The Mission blue (Icaricia icarioides missionensis) is a blue or lycaenid butterfly subspecies native to the San Francisco Bay Area of the United States. The butterfly has been declared as endangered by the US federal government. It is a...



Moduza procris

Moduza procris

Moduza procris, the commander, sometimes included in the genus Limenitis, is a medium-sized, strikingly coloured brush-footed butterfly found in South Asia and Southeast Asia. It is notable for the mode of concealment employed by its caterpillar...



Monarch butterfly

Danaus plexippus

The monarch butterfly or simply monarch (Danaus plexippus) is a milkweed butterfly (subfamily Danainae) in the family Nymphalidae. Other common names, depending on region, include milkweed, common tiger, wanderer, and black-veined brown. It is...



Monomorium santschii

Monomorium santschii

Monomorium santschii is a species of ant that is native to Tunisia. The most famous species in the genus Monomorium is the highly invasive pharaoh ant, Monomorium pharaonis. It is a parasitic ant that has no worker caste. The queen enters the...



Moorhen flea

Dasypsyllus gallinulae

The moorhen flea (Dasypsyllus gallinulae) is a flea originating from South America. It is now globally widespread. It is a large flea, easily identified because the male has two heavy horn-like spines on one of the genital flaps, and the female...



Mormon cricket

Anabrus simplex

The Mormon cricket (Anabrus simplex) is a large insect native to western North America in rangelands dominated by sagebrush and forbs. Anabrus is a genus in the shield-backed katydid subfamily in the Tettigoniidae family, commonly called...



Mormotomyiidae

Mormotomyia hirsuta

The family Mormotomyiidae (Diptera: Ephydroidea) contains only one known species, Mormotomyia hirsuta, commonly known as the frightful hairy fly or terrible hairy fly, which is found in Kenya. The fly was first described by English entomologist...



Morpho cypris

Morpho cypris

Morpho cypris, the Cypris morpho, is a Neotropical butterfly. It is found in Panama, Costa Rica, Nicaragua, Colombia, Venezuela, Trinidad and Tobago and Ecuador. Several subspecies and many forms have been described.



Morpho menelaus

Morpho menelaus

The Menelaus blue morpho (Morpho menelaus) is one of thirty species of butterfly in the subfamily Morphinae. Its wingspan is approximately 12 cm (4.7"), and its dorsal forewings and hindwings are a bright, iridescent blue edged with black, while...



Morpho peleides

Morpho peleides

Morpho peleides, the Peleides blue morpho, common morpho or the emperor is an iridescent tropical butterfly found in Mexico, Central America, northern South America, Paraguay and Trinidad. Most authorities believe that peleides is a subspecies of...



Morpho polyphemus

Morpho polyphemus

Morpho polyphemus, the white morpho or Polyphemus white morpho, is a white butterfly of Mexico and Central America, ranging as far south as Costa Rica. As suggested by its name, this is one of the relatively few Morphos that is white rather than...



Mottled umber

Erannis defoliaria

The mottled umber (Erannis defoliaria) is a moth of the family Geometridae. It is common throughout much of the Palearctic region. The species was first described by Carl Alexander Clerck in 1759.



Mountain pine beetle

Dendroctonus ponderosae

The mountain pine beetle (Dendroctonus ponderosae) is a species of bark beetle native to the forests of western North America from Mexico to central British Columbia. It has a hard black exoskeleton, and measures...



Mouse moth

Amphipyra tragopoginis

The mouse moth (Amphipyra tragopoginis) is a moth of the family Noctuidae. It is a widespread species with a Holarctic distribution.



Musk beetle

Aromia moschata

The musk beetle (Aromia moschata) is a Eurasian species of longhorn beetle belonging to the subfamily Cerambycinae, tribe Callichromatini. Its name comes from the delicate musky smell it emits when menaced. The beetle has a shell with an...



Myathropa florea

Myathropa florea

Myathropa florea, sometimes referred to as the Batman hoverfly, is a very common European and North African species of hoverfly. Adults may be seen on flowers from May to September. It is of a similar size to the common drone fly (Eristalis...



Mycalesis patnia

Mycalesis patnia

Mycalesis patnia, the glad-eye bushbrown, is a satyrid butterfly found in southern India and Sri Lanka.



Mycalesis perseus

Mycalesis perseus

Mycalesis perseus, the dingy bushbrown or common bushbrown, is a species of satyrine butterfly found in south Asia and southeast Asia.



Mycalesis visala

Mycalesis visala

Mycalesis visala, the long-brand bushbrown, is a species of satyrine butterfly found in south Asia. It might include the Tamil bushbrown as a subspecies.



Myrmecia gulosa

Myrmecia gulosa

Myrmecia gulosa, the red bull ant, also known as the giant bull ant or "hoppy joe", is a species of bulldog ant from the genus Myrmecia. It is abundant throughout Eastern Australia.



Myrmecia nigriceps

Myrmecia nigriceps

Myrmecia nigriceps, also known as the black-headed bull ant, is a species of ant endemic to Australia. A member of the genus Myrmecia in the subfamily Myrmeciinae, it was first described by Austrian entomologist Gustav Mayr in 1862. These ants...



Myrmecophilus acervorum

Myrmecophilus acervorum

Myrmecophilus acervorum is an orthopteran insect belonging to the family Myrmecophilidae (the ant-loving crickets). This continental European species is probably the most widespread and most frequently encountered member of this rather obscure...



Myrmelachista schumanni

Myrmelachista schumanni

Myrmelachista schumanni, also known as the lemon ant, is a species of ant from South America. It is notable for the creation of devil's gardens. Using its own herbicide, they kill off all the plants in an area except for the myrmecophytes, or...



Myrmica rubra

Myrmica rubra

Myrmica rubra, also known as the common red ant or erroneously the European fire ant, is a species of ant of the genus Myrmica, found all over Europe and is now invasive in some parts of North America and Asia. It is mainly red in colour, with...



Myrmicaria brunnea

Myrmicaria brunnea

Myrmicaria brunnea is a species of ant that is native to southern Asia. They have a distinctive down-curved abdomen and spines on the thorax.



Myzus persicae

Myzus persicae

Myzus persicae, known as the green peach aphid, greenfly, or the peach-potato aphid, is a small green aphid belonging to the order Hemiptera. It is the most significant aphid pest of peach trees, causing decreased growth, shrivelling of the...



Nasutitermes corniger

Nasutitermes corniger

Nasutitermes corniger is a species of arboreal termite that is endemic to the neotropics. It is very closely related to Nasutitermes ephratae. The species has been studied relatively intensively, particularly on Barro Colorado Island, Panama....



Nematus ribesii

Nematus ribesii

Nematus ribesii is a species of sawfly in the family Tenthredinidae. English names include common gooseberry sawfly and imported currantworm. This insect is best known as a pest of gooseberries. The larvae feed on the foliage of the plant, often...



Neonympha mitchellii

Neonympha mitchellii

Neonympha mitchellii is an endangered species of nymphalid butterfly of the eastern United States. There are two known subspecies: N. m. mitchellii, the nominate subspecies, commonly called Mitchell's satyr or Mitchell's marsh satyr, is found in...



Nepa cinerea

Nepa cinerea

Nepa cinerea is a species of water scorpion (Nepidae), found in most of Europe, including the British Isles, as well as North Africa and southern and northern Asia.



Nephrotoma appendiculata

Nephrotoma appendiculata

Nephrotoma appendiculata, the spotted crane fly, is a species of crane fly.



Nephrotoma crocata

Nephrotoma crocata

Nephrotoma crocata is a species of crane fly found in most of Europe and northern Russia. The subspecies N. c. luteata is found in southwest France, Portugal, Spain, Morocco, and Algeria.



Neptis columella

Neptis columella

Neptis columella, the short banded sailer, is a species of nymphalid butterfly found in South and Southeast Asia.



Neptis hylas

Neptis hylas

Neptis hylas, the common sailor, is a species of nymphalid butterfly found in the Indian subcontinent and southeast Asia. It has a characteristic stiff gliding flight achieved by short and shallow wingbeats just above the horizontal.



Neuroterus albipes

Neuroterus albipes

Neuroterus albipes is a gall wasp that forms chemically induced leaf galls on oak trees which has both bisexual and agamic generations and therefore forms two distinct galls, the smooth spangle gall and Schenck's gall. Neuroterus laeviusculus and...



Neuroterus quercusbaccarum

Neuroterus quercusbaccarum

The common spangle gall on the underside of leaves and the currant gall on the male catkins or occasionally the leaves, develop as chemically induced distortions on pedunculate oak (*Quercus robur*), or sessile oak (*Quercus petraea*) trees, caused...



New Zealand bat fly

Mystacinobia zelandica

The New Zealand bat fly (*Mystacinobia zelandica*) is a small, wingless insect which lives in a commensal relationship with the New Zealand lesser short-tailed bat. It is a true fly, in the order Diptera, placed in its own genus, *Mystacinobia*, and...



New Zealand red admiral

Vanessa gonerilla

The New Zealand red admiral (*Vanessa gonerilla*) is a butterfly endemic to New Zealand. Its Māori name is kahukura, which means "red cloak". The red admiral is a member of the family Nymphalidae, the subfamily Nymphalinae and the tribe Nymphalini....



Nicrophorus americanus

Nicrophorus americanus

Nicrophorus americanus, also known as the American burying beetle or giant carrion beetle, is a critically endangered species of beetle endemic to North America. It belongs to the order Coleoptera and the family Silphidae. The carrion beetle in...



Nicrophorus defodiens

Nicrophorus defodiens

Nicrophorus defodiens is a burying beetle described by Mannerheim in 1846. In 2012, *N. defodiens* was found to be one of at least two burying beetles which can breed in the forest canopy.



Nicrophorus interruptus

Nicrophorus interruptus

Nicrophorus interruptus is a species of burying beetle or sexton beetle belonging to the family Silphidae subfamily Nicrophorinae.



Nicrophorus marginatus

Nicrophorus marginatus

Nicrophorus marginatus is a burying beetle described by Johan Christian Fabricius in 1801.



Nicrophorus orbicollis

Nicrophorus orbicollis

Nicrophorus orbicollis is a nearctic burying beetle first described by Thomas Say in 1825. It is a member of the genus Nicrophorus or sexton beetles, comprising the most common beetles in the family Silphidae. This species is a decomposer feeding...



Nicrophorus vespilloides

Nicrophorus vespilloides

Nicrophorus vespilloides is a burying beetle described by Johann Friedrich Wilhelm Herbst in 1783. The beetles are 10 – 18 mm long. They have two conspicuous orange-yellow bands on the elytra. The color of the antennae are an important...



Nihoa conehead katydid

Banza nihoa

The Nihoa conehead katydid (Banza nihoa) is a species of katydid which is endemic to the Hawaiian island of Nihoa (Northwestern Hawaiian Islands). It is one of the ten species in the genus Banza, all of them native to Hawaii, although it is the...



Niobe fritillary

Fabriciana niobe

The Niobe fritillary (Fabriciana niobe) is a species of butterfly in the family Nymphalidae.



Noble chafer

Gnorimus nobilis

The noble chafer (Gnorimus nobilis) is a species of beetles belonging to the family Scarabaeidae, subfamily Cetoniinae.



Norfolk damselfly

Coenagrion armatum

The Norfolk damselfly or dark bluet (Coenagrion armatum) is a species of blue damselfly of the family Coenagrionidae native to Eurasia.



Nosopsyllus fasciatus

Nosopsyllus fasciatus

Nosopsyllus fasciatus, the northern rat flea, is a species of flea found on domestic rats and house mice. Northern rat fleas are external parasites, living by hematophagy off the blood of rodents. N. fasciatus can bite humans, but they are more...



Notocrypta curvifascia

Notocrypta curvifascia

Notocrypta curvifascia, the restricted demon, is a butterfly belonging to the family Hesperiiidae. N. curvifascia is commonly found in many regions of temperate and tropical East Asia, Indonesia, and the Indian subcontinent. Among butterflies, it...



November moth

Epirrita dilutata

The November moth (Epirrita dilutata) is a moth of the family Geometridae. The species was first described by Michael Denis and Ignaz Schiffermüller in 1775. It can be found in the Palearctic realm in western Europe from central Scandinavia to...



Novius cardinalis

Novius cardinalis

Novius cardinalis (common names vedalia beetle or cardinal ladybird) is a species of ladybird beetle native to Australia. It was formerly placed in the genus Rodolia, but that genus was synonymized under the genus Novius in 2020.



Nutmeg

Hadula trifolii

The nutmeg (Hadula trifolii or Anarta trifolii), also known as the clover cutworm, is a moth of the family Noctuidae.



Nyctemera adversata

Nyctemera adversata

Nyctemera adversata, the marbled white moth, is a moth of the family Erebidae first described by Johann Gottlieb Schaller in 1788. It is found in Sri Lanka, Bangladesh, India, Nepal, Myanmar, China territories like Tibet, Sichuan, Yunnan,...



Nyctemera annulata

Nyctemera annulata

Nyctemera annulata, the magpie moth, is a moth of the family Erebidae. The species was first described by Jean Baptiste Boisduval in 1832. It is endemic to New Zealand and found in all parts of the country.



Nymphalis antiopa

Nymphalis antiopa

Nymphalis antiopa, known as the mourning cloak in North America and the Camberwell beauty in Britain, is a large butterfly native to Eurasia and North America. The immature form of this species is sometimes known as the spiny elm caterpillar....



Nymphalis vaualbum

Nymphalis vaualbum

Nymphalis vaualbum or N. l-album, the Compton tortoiseshell, or false comma, is a species of butterfly in the family Nymphalidae. An assertion that the name Nymphalis l-album is in fact the correct name over the widely used Nymphalis vaualbum...



Nymphalis xanthomelas

Nymphalis xanthomelas

Nymphalis xanthomelas, the scarce tortoiseshell, is a species of nymphalid butterfly found in eastern Europe and Asia. This butterfly is also referred as yellow-legged tortoiseshell or large tortoiseshell (however, in Europe, "large...



Nymphes myrmeleonides

Nymphes myrmeleonides

Nymphes myrmeleonides is an Australian insect in the order Neuroptera, known as the blue eyes lacewing. It is found in areas of New South Wales and Queensland. The species have a body length of up to 4 centimetres (1.6 in) and a wingspan of up to...



Nysius raphanus

Nysius raphanus

Nysius raphanus (also known as the false chinch bug) is a small (no more...



Nysius wekiuicola

Nysius wekiuicola

Nysius wekiuicola is a species of wēkiu bug (a type of seed bug in the genus Nysius) endemic to the area surrounding the summit of Mauna Kea, on the island of Hawai‘i. It is closely related to Nysius aa, which is endemic to the summit of the...



Oak processionary

Thaumetopoea processionea

The oak processionary (OPM) (Thaumetopoea processionea) is a moth whose caterpillars can be found in oak forests, where they feed on oak leaves, causing significant damage. They travel in nose-to-tail processions (hence their name), often...



Odontomachus bauri

Odontomachus bauri

Odontomachus bauri is a species of ponerinae ant known as trap jaw ants. The trap jaw consists of mandibles which contain a spring-loaded catch mechanism. This mechanism permits the ants to accumulate energy before striking or releasing the...



Odontotaenius disjunctus

Odontotaenius disjunctus

Odontotaenius disjunctus, the patent-leather beetle or horned passalus, is a saproxylic beetle in the family Passalidae (bess beetles) which can grow to just over an inch-and-a-half long, weigh 1-2 grams and are capable of pulling 50 times their...



Oenochroma vinaria

Oenochroma vinaria

Oenochroma vinaria, the pink-bellied moth, is a moth of the family Geometridae. It is found in most of Australia. The wingspan is 50–70 mm. The larvae feed on Grevillea, Banksia and Hakea species.



Olive fruit fly

Bactrocera oleae

The olive fruit fly (Bactrocera oleae) is a species of fruit fly which belongs to the subfamily Dacinae. It is a phytophagous species whose larvae feed on the fruit of olive trees, hence the common name. It is considered a serious pest in the...



Onchestus rentzi

Onchestus rentzi

Onchestus rentzi is an Australian species of stick insect, commonly named the crowned stick insect, described in 2006. It lives in rainforests along the coast of Queensland and the Northern Territory. It is named after the Australian orthopterist...



Oncideres cingulata

Oncideres cingulata

Oncideres cingulata, the twig girdler, is a brownish-gray beetle,...



Operophtera fagata

Operophtera fagata

Operophtera fagata, the northern winter moth, is a moth of the family Geometridae. It is widespread through much of Europe.



Ophiusa tirhaca

Ophiusa tirhaca

Ophiusa tirhaca, the green drab, is a moth of the family Erebidæ. The species was first described by Pieter Cramer in 1777. It is found in southern Europe, Africa, Australia and the southern parts of Asia. The wingspan is about 50 mm. The larvae...



Opodiphthera eucalypti

Opodiphthera eucalypti

Opodiphthera eucalypti, the emperor gum moth, is a species of moth in the family Saturniidae native to Australia. This species was formerly placed in the genus Antheraea.



Orachrysops niobe

Orachrysops niobe

Orachrysops niobe, the Brenton blue, is a species of butterfly in the family Lycaenidae and is endemic to South Africa. The wingspan is 24–38 mm for males and 22–42 mm for females. Adults are on wing from October to November and from February to...



Orange-spotted emerald

Oxygastra curtisii

The orange-spotted emerald (Oxygastra curtisii) is a dragonfly in the family Corduliidae. It is the only species in its genus. The orange-spotted emerald is about 53 millimetres (2.1 in) long. It has bright green eyes and a bronzy-green body with...



Organ pipe mud dauber

Trypoxylon politum

The organ pipe mud dauber (Trypoxylon politum) is a predatory wasp in the family Crabronidae. It is fairly large, ranging from 3.9–5.1 cm, and has been recorded to fly from May to September. Females and males are similar in colour, a shiny black,...



Orgyia antiqua

Orgyia antiqua

Orgyia antiqua, the rusty tussock moth or vapourer, is a moth in the family Erebidæ.



Orgyia leucostigma

Orgyia leucostigma

Orgyia leucostigma, the white-marked tussock moth, is a moth in the family Erebidae. The species was first described by James Edward Smith in 1797. The caterpillar is very common especially in late summer in eastern North America, extending as...



Oriental cockroach

Blatta orientalis

The oriental cockroach (Blatta orientalis), also known as the waterbug (as they live in damp areas) or black cockroaches (as their bodies are mostly dark), is a large species of cockroach, adult males being 18–29 mm (0.71–1.14 in) and adult...



Oriental hornet

Vespa orientalis

The Oriental hornet (Vespa orientalis) is a social insect species of the family Vespidae. It can be found in Southwest Asia, Northeast Africa, the island of Madagascar (but no reports have been made of its presence on the island for many years),...



Ormia ochracea

Ormia ochracea

Ormia ochracea is a small yellow nocturnal fly in the family Tachinidae. It is notable for its parasitism of crickets and its exceptionally acute directional hearing. The female is attracted to the song of the male cricket and deposits larvae on...



Ornithoptera aesacus

Ornithoptera aesacus

Ornithoptera aesacus, the Obi Island birdwing, is a rare species of birdwing butterfly, endemic to the Island of Obira (formerly Obi), Indonesia. The specific epithet of Ornithoptera aesacus, is named after Æsacus, the eldest son of Priam.



Ornithoptera croesus

Ornithoptera croesus

Ornithoptera croesus, the Wallace's golden birdwing, is a species of birdwing butterfly found in northern Maluku in Indonesia. It is a member of the Ornithoptera priamus species group which, including croesus, is only found east of the Wallace...



Ornithoptera euphorion

Ornithoptera euphorion

Ornithoptera euphorion, the Cairns birdwing, is a species of birdwing butterfly endemic to Queensland, and is Australia's largest endemic butterfly species. Other common names include Cooktown birdwing and northern birdwing. The names Cairns and...



Ornithoptera goliath

Ornithoptera goliath

Ornithoptera goliath, the Goliath birdwing, is a birdwing butterfly found in New Guinea. It is the second largest butterfly in the world, after the Queen Alexandra's birdwing.



Ornithoptera meridionalis

Ornithoptera meridionalis

Ornithoptera meridionalis, the southern tailed birdwing, is the smallest species of the genus Ornithoptera. It is known from a handful of localities in southeast Papua, New Guinea (O. meridionalis meridionalis) and several localities along the...



Ornithoptera victoriae

Ornithoptera victoriae

Ornithoptera victoriae, the Queen Victoria's birdwing, is a birdwing butterfly of the family Papilionidae, found in the Solomon Islands and Papua New Guinea (Bougainville Island only).What was originally described as Ornithoptera allotei, is a...



Orthetrum trinacria

Orthetrum trinacria

Orthetrum trinacria, the Long Skimmer, is a species of dragonfly in the family Libellulidae. It is found in Algeria, Angola, Botswana, Burkina Faso, Cameroon, Ivory Coast, Egypt, Ethiopia, France (Corsica), Gambia, Ghana, Italy (Sicily and...



Orthodera novaezealandiae

Orthodera novaezealandiae

Orthodera novaezealandiae, (Māori: whe) known as the New Zealand mantis or the New Zealand praying mantis, is a species of praying mantis which is, as both the scientific name and common names suggest, indigenous and endemic to New Zealand.



Osmia bicornis

Osmia bicornis

Osmia bicornis (synonym Osmia rufa) is a species of mason bee, and is known as the red mason bee due to its covering of dense gingery hair. It is a solitary bee that nests in holes or stems and is polylectic, meaning it forages pollen from...



Osmia californica

Osmia californica

Osmia californica is a megachilid bee, or mason bee. Native to North America, the mason bees are important pollinators. O. californica generally emerges a little later in the spring than the better known orchard mason bee (O. lignaria). Like the...



Osmia cornifrons

Osmia cornifrons

Osmia cornifrons, also known as the horned-face bee, is a species of solitary bee indigenous to Northern Asia. Physically, this species of bee is recognized for its horn-like extensions originating from its lower face. Populations of O....



Osmia cornuta

Osmia cornuta

Osmia cornuta, the European orchard bee, is a species of bee in the genus Osmia.



Osmia lignaria

Osmia lignaria

Osmia lignaria, commonly known as the orchard mason bee or blue orchard bee, is a megachilid bee that makes nests in natural holes and reeds, creating individual cells for its brood that are separated by mud dividers. Unlike carpenter bees, it...



Osmia ribifloris

Osmia ribifloris

Osmia ribifloris, one of several species referred to as a blueberry bee, is a megachilid bee native to western North America, including Oregon, California, Nevada, Utah, Arizona, New Mexico, Texas and northern Mexico. This solitary bee normally...



Otiorhynchus ovatus

Otiorhynchus ovatus

Otiorhynchus ovatus, the strawberry root weevil, is one of the many species in the weevil family (Curculionidae), occurring across Canada and the northern United States. The species was first described by Carl Linnaeus in his landmark 1758 10th...



Pachliopta aristolochiae

Pachliopta aristolochiae

Pachliopta aristolochiae, the common rose, is a swallowtail butterfly belonging to the genus Pachliopta, the roses, or red-bodied swallowtails. It is a common butterfly which is extensively distributed across south and southeast Asia.



Pachliopta hector

Pachliopta hector

Pachliopta hector, the crimson rose, is a large swallowtail butterfly belonging to the genus Pachliopta (roses) of the red-bodied swallowtails. It is recorded as a species of "Least Concern (LC)" by IUCN.



Pachliopta jophon

Pachliopta jophon

Pachliopta jophon, the Ceylon rose or Sri Lankan rose, is a butterfly found in Sri Lanka that belongs to the swallowtail family. It was earlier classified as a subspecies of Pachliopta hector, the crimson rose.



Pachliopta kotzebuea

Pachliopta kotzebuea

Pachliopta kotzebuea, the pink rose, is a butterfly of the family Papilionidae (the swallowtails). It is found in the Philippines. Food plants include birthwort (Aristolochia) species.



Pallid-winged grasshopper

Trimerotropis pallidipennis

The pallid-winged grasshopper (Trimerotropis pallidipennis) is a common grasshopper of the family Acrididae, native to the deserts of western North America along with South America, ranging from British Columbia to Argentina. They are more active...



Palos Verdes blue

Glaucopsyche lygdamus palosverdesensis

The Palos Verdes blue (Glaucopsyche lygdamus palosverdesensis) is a small endangered butterfly native to the Palos Verdes Peninsula in southwest Los Angeles County, California, United States. As its distribution has been proven to be limited to...



Pameridea roridulae

Pameridea roridulae

Pameridea roridulae is a species of jumping tree bug in the family Miridae. It has a symbiotic relationship with Roridula, where it feeds on trapped insects and its excretions are absorbed through the leaves of the Roridula, giving the plant much...



Panchlora nivea

Panchlora nivea

Panchlora nivea, the Cuban cockroach or green banana cockroach, is a small species of cockroach in the subfamily Panchlorinae. It is found in the subtropical or tropical climates of Cuba, the Caribbean and southern US: along the Gulf Coast from...



Pandora moth

Coloradia pandora

The Pandora moth or Pandora pinemoth (Coloradia pandora) is an insect belonging to the moth genus Coloradia. The species was first described by C. A. Blake in 1863. It is native to the western United States. The larvae of the Pandora moth feed on...



Panstrongylus geniculatus

Panstrongylus geniculatus

Panstrogylus geniculatus is a blood-sucking sylvatic insect noted as a putative vector of minor importance in the transmission of Trypanosoma cruzi to humans; this is a parasite, which causes Chagas disease. The insect is described as sylvatic:...



Pantala flavescens

Pantala flavescens

Pantala flavescens, the globe skimmer, globe wanderer or wandering glider, is a wide-ranging dragonfly of the family Libellulidae. This species and Pantala hymenaea, the "spot-winged glider", are the only members of the genus Pantala. It was...



Pantoporia hordonia

Pantoporia hordonia

Pantoporia hordonia, the common lascar, is a species of nymphalid butterfly found in Cambodia, tropical and subtropical Asia.



Papilio aegeus

Papilio aegeus

Papilio aegeus, the orchard swallowtail butterfly or large citrus butterfly is a species of butterfly from the family Papilionidae, that is found in eastern Australia and Papua New Guinea. The larvae of this species are sometimes considered a...



Papilio agestor

Papilio agestor

Papilio (Chilasa) agestor, the tawny mime, is a swallowtail butterfly, native to Indian subcontinent and widely found across Asia. The butterfly belongs to the mime subgenus, Chilasa, of the genus Papilio or the black-bodied swallowtails.



Papilio alexanor

Papilio alexanor

Papilio alexanor, the Alexanor or southern swallowtail, is a butterfly species in the Papilionidae or swallowtails. The insect has a wingspan of 62–70 mm. It flies from April to July in a single generation, in mountainous regions (1700 m) of...



Papilio breviceauda

Papilio breviceauda

Papilio breviceauda, the short-tailed swallowtail, is a North American butterfly in the family Papilionidae.



Papilio buddha

Papilio buddha

Papilio buddha, the Malabar banded peacock, is a species of swallowtail butterfly found in the Western Ghats of India. The Government of Kerala declared it as the official Kerala state butterfly.



Papilio castor

Papilio castor

Papilio castor, the common raven, is a species of swallowtail butterfly found in Cambodia and South Asia.



Papilio cresphontes

Papilio cresphontes

The giant swallowtail (Papilio ctesiphontes) is the largest butterfly in North America. It is abundant through many parts of eastern North America; populations from western North America and down into Panama are now (as of 2014) considered to...



Papilio crino

Papilio crino

Papilio crino, the common banded peacock, is a species of swallowtail (Papilionidae) butterfly found in parts of the Indian subcontinent, including India, Nepal, Bhutan and Sri Lanka.



Papilio demodocus

Papilio demodocus

Papilio demodocus, the citrus swallowtail or Christmas butterfly, is a swallowtail butterfly which commonly occurs over the entirety of sub-Saharan Africa, including Madagascar, besides the southern Arabian Peninsula. The caterpillars feed on...



Papilio demoleus

Papilio demoleus

Papilio demoleus is a common and widespread swallowtail butterfly. The butterfly is also known as the lime butterfly, lemon butterfly, lime swallowtail, and chequered swallowtail. These common names refer to their host plants, which are usually...



Papilio dravidarum

Papilio dravidarum

Papilio dravidarum, the Malabar raven, is an endemic species of swallowtail butterfly found in the Western Ghats of India.



Papilio elephenor

Papilio elephenor

Papilio elephenor, the yellow-crested spangle, is a species of swallowtail butterfly found in Northeast India. Following decades without confirmed sightings, it was rediscovered in 2009 in Assam.



Papilio glaucus

Papilio glaucus

Papilio glaucus, the eastern tiger swallowtail, is a species of butterfly native to eastern North America. It is one of the most familiar butterflies in the eastern United States, ranging north to southern Ontario, Canada, and is common in many...



Papilio helenus

Papilio helenus

Papilio helenus, the red Helen, is a large swallowtail butterfly found in forests of southern India and parts of southeast Asia.



Papilio krishna

Papilio krishna

Papilio krishna, the Krishna peacock, is a large swallowtail butterfly found in forests in China, Nepal, north east India, Myanmar and Vietnam.



Papilio machaon

Papilio machaon

Papilio machaon, the Old World swallowtail, is a butterfly of the family Papilionidae. The butterfly is also known as the common yellow swallowtail or simply the swallowtail (a common name applied to all members of the family, but this species...



Papilio machaon oregonius

Papilio machaon oregonius

The Oregon swallowtail (Papilio machaon oregonius, sometimes classified as Papilio oregonius or Papilio bairdii oregonius) is a subspecies of swallowtail butterfly native to the United States of America, Oregon, Washington, and Idaho and...



Papilio mayo

Papilio mayo

Papilio mayo, the Andaman Mormon, is a species of swallowtail butterfly that is endemic to the Andamans in the Bay of Bengal. It is a species that is protected by Indian Law. The scientific name honours Richard Bourke, 6th Earl of Mayo, who was...



Papilio memnon

Papilio memnon

Papilio memnon, the great Mormon, is a large butterfly native to southern Asia that belongs to the swallowtail family. It is widely distributed and has thirteen subspecies. The female is polymorphic and with mimetic forms.



Papilio multicaudata

Papilio multicaudata

Papilio multicaudata, the two-tailed swallowtail, is a species of the family Papilionidae found in western North America from British Columbia to Central America.



Papilio nephelus

Papilio nephelus

Papilio nephelus is a species of swallowtail butterfly belonging to the family Papilionidae. Subspecies include P. n. chaon, the yellow Helen, and P. n. sunatus, the black and white Helen.



Papilio ophidicephalus

Papilio ophidicephalus

Papilio ophidicephalus, the emperor swallowtail, is a butterfly of the family Papilionidae. It is found in Sub-Saharan Africa. The wingspan is 90–110 mm in males and 100–120 mm in females. It has two broods, one from August to December and the...



Papilio palinurus

Papilio palinurus

Papilio palinurus, the emerald swallowtail, emerald peacock, or green-banded peacock, is a butterfly of the genus Papilio of the family Papilionidae. It is native to Southeast Asia, but is regularly kept in butterfly houses around the world.



Papilio paris

Papilio paris

Papilio paris, the Paris peacock (Myanmar name: ဒေါင်းစိမ်းလိပ်ပြာ), is a species of swallowtail butterfly found in the Indian subcontinent and southeast Asia.



Papilio polyctor

Papilio polyctor

Papilio polyctor, the common peacock, is a swallowtail butterfly found in the Indian subcontinent. It is found in the Himalayas and parts of India from the foothills to 7,000 feet (2,100 m) between March and October. It has distinct dry- and...



Papilio polytes

Papilio polytes

Papilio polytes, the common Mormon, is a common species of swallowtail butterfly widely distributed across Asia. This butterfly is known for the mimicry displayed by the numerous forms of its females which mimic inedible red-bodied swallowtails,...



Papilio polyxenes

Papilio polyxenes

Papilio polyxenes, the (eastern) black swallowtail, American swallowtail or parsnip swallowtail, is a butterfly found throughout much of North America. An extremely similar-appearing species, Papilio joanae, occurs in the Ozark Mountains region,...



Papilio protenor

Papilio protenor

Papilio protenor, the spangle, is a butterfly found in India belonging to the swallowtail family.



Papilio rutulus

Papilio rutulus

Papilio rutulus, the western tiger swallowtail, is a swallowtail butterfly belonging to the Papilionidae family. The species was first described by Hippolyte Lucas in 1852. Like the other tiger swallowtails, the western tiger swallowtail was...



Papilio troilus

Papilio troilus

Papilio troilus, the spicebush swallowtail or green-clouded butterfly, is a common black swallowtail butterfly found in North America. It has two subspecies, Papilio troilus troilus and Papilio troilus ilioneus, the latter found mainly in the...



Papilio zalmoxis

Papilio zalmoxis

Papilio zalmoxis, the giant blue swallowtail, is an African butterfly belonging to the family Papilionidae. The name of the species is given in honor of Zalmoxis – a divinity of the Getae (a people of the lower Danube), mentioned by Herodotus in...



Papilio zelicaon

Papilio zelicaon

Papilio zelicaon, the anise swallowtail, is a common swallowtail butterfly of western North America. Both the upper and lower sides of its wings are black, but the upper wing has a broad yellow stripe across it, giving the butterfly an overall...



Parablepharis

Parablepharis kuhlii

Parablepharis is a genus of praying mantises in the family Hymenopodidae. It is monotypic, being represented by the single species, Parablepharis kuhlii and is found in India, Myanmar, Vietnam, Borneo, Java and Yunnan, China.



Parantica aglea

Parantica aglea

Parantica aglea, the glassy tiger, is a butterfly found in Indomalayan realm that belongs to the crows and tigers, that is, the danaid group of the brush-footed butterflies family.



Parantica melaneus

Parantica melaneus

Parantica melaneus (known colloquially as the chocolate tiger), is a butterfly found in Asia that belongs to the crows and tigers, that is, the danaid group of the brush-footed butterflies family.



Parantica sita

Parantica sita

Parantica sita, the chestnut tiger, is a butterfly found in Asia that belongs to the crows and tigers, that is, the danaid group of the brush-footed butterflies family.



Paraponera clavata

Paraponera clavata

Paraponera clavata, commonly known as the bullet ant, is a species of ant named for its extremely painful sting. It inhabits humid lowland rainforests in Central and South America.



Parcoblatta fulvescens

Parcoblatta fulvescens

Parcoblatta fulvescens, the fulvous wood cockroach, is a species of cockroach endemic to the United States and possibly Canada that measures around 13 mm (0.5 in) long.



Parcoblatta virginica

Parcoblatta virginica

Parcoblatta virginica, the Virginia wood cockroach, is a small cockroach species of the genus Parcoblatta, measuring about a centimeter long as an adult.



Parent bug

Elasmucha grisea

Elasmucha grisea, common name parent bug, is a species of shield bugs or stink bugs belonging to the family Acanthosomatidae. The term parent bugs includes also the other species of the genus Elasmucha and some species of the family...



Pareronia valeria

Pareronia valeria

Pareronia valeria, the common wanderer or Malayan wanderer, is a medium-sized butterfly of the family Pieridae, that is, the yellows and whites, and is found in India and Southeast Asia. The butterfly found in India is sometimes considered as a...



Parktown prawn

Libanasidus vittatus

The Parktown prawn, African king cricket or tusked king cricket (Libanasidus vittatus) is a species of king cricket endemic to Southern Africa. It is unrelated to prawns, Libanasidus being insects in the order Orthoptera – crickets, locusts and...



Parnara guttata

Parnara guttata

Parnara guttata, the common straight swift, is a butterfly of the family HesperIIDae. It is found in the Indomalayan realm, Amur (Russian Far East), eastern China, and Japan. Its wingspan is about 40 mm. Common straight swift larvae feed on...



Parnassius actius

Parnassius actius

Parnassius actius, also known as the scarce red apollo, is a high-altitude butterfly found in Central Asia. It is a member of the snow Apollo genus (Parnassius) of the swallowtail family (Papilionidae).



Parnassius clodius

Parnassius clodius

Parnassius clodius is a white butterfly which is found in the United States and Canada. It is a member of the snow Apollo genus (Parnassius) of the swallowtail family (Papilionidae). Its elevation range is 0–7,000 ft (0–2,134 m).



Parnassius eversmanni

Parnassius eversmanni

Parnassius eversmanni, or Eversmann's parnassian, is a high-altitude butterfly which is found in eastern Russia, Mongolia, Japan, Alaska, and the Yukon. It is a member of the snow Apollo genus (Parnassius) of the swallowtail family, Papilionidae....



Parnassius phoebus

Parnassius phoebus

Parnassius phoebus, known as the Phoebus Apollo or small Apollo, is a butterfly species of the swallowtail butterfly family, Papilionidae, found in the Palearctic and North America.P. phoebus is found in the Alps, Urals, Siberia, Kazakhstan,...



Paysandisia archon

Paysandisia archon

Paysandisia archon is a moth of the family Castniidae. It is native to Uruguay and central Argentina and has been accidentally introduced to Europe, where it is spreading rapidly. It is considered the only member of the genus Paysandisia.German...



Peach blossom

Thyatira batis

The peach blossom (Thyatira batis) is a moth of the family Drepanidae. It was first described by Carl Linnaeus in his 1758 10th edition of Systema Naturae.It is found throughout Europe and east through the Palearctic to Japan and Mongolia. It is...



Pearl-bordered fritillary

Boloria euphrosyne

The pearl-bordered fritillary (Boloria euphrosyne) is a butterfly of the family Nymphalidae found in Europe and through Russia across the Palearctic to the north of Kazakhstan.



Pediculus humanus

Pediculus humanus

Pediculus humanus is a species of louse that infects humans. It comprises two subspecies: Pediculus humanus humanus Linnaeus, 1758 – body louse Pediculus humanus capitis De Geer, 1767 – head louse



Pelecinus polyturator

Pelecinus polyturator

Pelecinus polyturator is a species of wasp in the family Pelecinidae. The large (up to 7 cm) glossy black insects are the most common and well-known members of the family. The adults drink nectar, and they live in crop fields, woods, and suburban...



Pelopidas mathias

Pelopidas mathias

Pelopidas mathias, the dark small-branded swift, small branded swift, lesser millet skipper or black branded swift, is a butterfly belonging to the family Hesperiiidae. It is found throughout much of south, southeast and East Asia, and as far as...



Pennsylvania wood cockroach

Parcoblatta pensylvanica

The Pennsylvania wood cockroach (Parcoblatta pensylvanica) or Pennsylvanian cockroach is a common species of cockroach in eastern and central North America.



Peppered moth

Biston betularia

The peppered moth (Biston betularia) is a temperate species of night-flying moth. It is mostly found in the northern hemisphere in places like Asia, Europe and North America. Peppered moth evolution is an example of population genetics and...



Peruphasma schultei

Peruphasma schultei

Peruphasma schultei, known as the black beauty stick insect and the golden-eyed stick insect, is a species of phasmid found in the Cordillera del Condor region of northern Peru. In the wild the insect feeds on Schinus plants, but will feed on...



Petalura gigantea

Petalura gigantea

Petalura gigantea, the giant dragonfly or south-eastern petaltail, is a species of dragonfly in the family Petaluridae from southeastern Australia. Males have body length of 6–7.5 cm (2.4–3.0 in) and a wingspan up to 11 cm (4.3 in), while females...



Petalura ingentissima

Petalura ingentissima

Petalura ingentissima, the giant petaltail, has been described as the world's largest dragonfly, with a wingspan of 160 mm. It is found in Queensland, Australia. Dr R.J. Tillyard described the giant petaltail in 1908. Its species name is derived...



Petrobius maritimus

Petrobius maritimus

Petrobius maritimus, the shore bristletail or sea bristletail, is a species of Archaeognatha found on rocky shores from the Mediterranean Sea to the North Sea. Individuals may grow up to 15 mm and are grey in colour, with long bristly antennae and...



Phalaenoides glycinae

Phalaenoides glycinae

Phalaenoides glycinae, the Australian grapevine moth, is a moth of the family Noctuidae that is native to southeastern Australia. The species was first described by John Lewin in 1805.



Phalanta phalantha

Phalanta phalantha

Phalanta phalantha, the common leopard or spotted rustic, is a sun-loving butterfly of the nymphalid or brush-footed butterfly family.



Phanaeus vindex

Phanaeus vindex

Phanaeus vindex, the rainbow scarab (a name also used for the entire genus), is a North American species of true dung beetle in the family Scarabaeidae. It is found in eastern and central United States (Florida and New England to Arizona and...



Pharaoh ant

Monomorium pharaonis

The pharaoh ant (Monomorium pharaonis) is a small (2 mm) yellow or light brown, almost transparent ant notorious for being a major indoor nuisance pest, especially in hospitals. A cryptogenic species, it has now been introduced to virtually every...



Phengaris alcon

Phengaris alcon

Phengaris alcon, the Alcon blue or Alcon large blue, is a butterfly of the family Lycaenidae and is found in Europe and across the Palearctic to Siberia and Mongolia.



Philaethria dido

Philaethria dido

Philaethria dido, the scarce bamboo page or dido longwing, is a butterfly in the family Nymphalidae. It is found in Central America and tropical South America, both east and west of the Andes, from Brazil and Ecuador northwards to Mexico. Strays...



Phobaeticus kirbyi

Phobaeticus kirbyi

Phobaeticus kirbyi is species of stick insect native to Borneo. It is one of the world's longest insects. The holotype deposited at the Natural History Museum in London measures 32.8 cm (12.9 in) in body length and 54.6 cm (21.5 in) total length,...



Phobetron pithecium

Phobetron pithecium

Phobetron pithecium, the hag moth, is a moth of the family Limacodidae. Its larva is known as the monkey slug.



Phoebis sennae

Phoebis sennae

Phoebis sennae, the cloudless sulphur, is a mid-sized butterfly in the family Pieridae found in the Americas. There are several similar species such as the yellow angled-sulphur (Anteos maerula), which has angled wings, statira sulphur (Aphrissa...



Photuris pensylvanica

Photuris pensylvanica

Photuris pensylvanica, known by the common names Pennsylvania firefly, lightning bug, dot-dash firefly and (in its larval state) glowworm, is a species of firefly from the United States and Canada. It is also widely known under the Latin name...



Phthorimaea operculella

Phthorimaea operculella

Phthorimaea operculella, also known as the potato tuber moth or tobacco splitworm, is a moth of the family Gelechiidae. It is an oligophagous insect that feeds on the plant family Solanaceae and is especially known for being a major pest of...



Phyllocrania paradoxa

Phyllocrania paradoxa

Phyllocrania paradoxa, common name ghost mantis, is a small species of mantis from Africa remarkable for its leaf-like body. It is one of the three species in the genus Phyllocrania. It is known for its distinct and exclusive camouflaged...



Phyllodes imperialis

Phyllodes imperialis

Phyllodes imperialis, the imperial fruit-sucking moth or pink underwing moth, is a noctuid moth in the family Erebidae, subfamily Calpinae. It was first described by Herbert Druce in 1888. The species can be found in north-eastern Queensland to...



Phylloxera

Daktulosphaira vitifoliae

Grape phylloxera is an insect pest of grapevines worldwide, originally native to eastern North America. Grape phylloxera (Daktulosphaira vitifoliae (Fitch 1855) belong to the family Phylloxeridae, within the order Hemiptera, bugs); originally...



Pieris brassicae

Pieris brassicae

Pieris brassicae, the large white, also called cabbage butterfly, cabbage white, cabbage moth (erroneously), or in India the large cabbage white, is a butterfly in the family Pieridae. It is a close relative of the small white, Pieris rapae.The...



Pieris canidia

Pieris canidia

Pieris canidia, the Indian cabbage white, is a butterfly in the family Pieridae found in India, Nepal and Indochina. Pieris rapae is one of the most closely related species in the Pieridae.



Pieris oleracea

Pieris oleracea

Pieris oleracea, or more commonly known as the mustard white, is a butterfly in the family Pieridae native to a large part of Canada and the northeastern United States. The nearly all-white butterfly is often found in wooded areas or open plains....



Pieris rapae

Pieris rapae

Pieris rapae is a small- to medium-sized butterfly species of the whites-and-yellows family Pieridae. It is known in Europe as the small white, in North America as the cabbage white or cabbage butterfly, on several continents as the small cabbage...



Pieris virginiensis

Pieris virginiensis

Pieris virginiensis, the West Virginia white, is a butterfly found in North America in the Great Lakes states, along the Appalachians from New England to Alabama, and in southern Ontario. They are typically found in moist deciduous forests....



Pine beauty

Panolis flammea

The pine beauty (Panolis flammea) is a moth of the family Noctuidae. It is a common species of pine woods in Europe. The distribution area extends from Portugal to western Siberia, the Caucasus and Asia Minor. In the north it extends to the...



Pine processionary

Thaumetopoea pityocampa

The pine processionary (Thaumetopoea pityocampa) is a moth of the subfamily Thaumetopoeinae in the family Notodontidae, known for the irritating hairs of its caterpillars, their processions, and the economic damage they cause in coniferous...



Pink bollworm

Pectinophora gossypiella

The pink bollworm (*Pectinophora gossypiella*; Spanish: lagarta rosada) is an insect known for being a pest in cotton farming. The adult is a small, thin, gray moth with fringed wings. The larva is a dull white caterpillar with eight pairs of legs...



Plum curculio

Conotrachelus nenuphar

The plum curculio (*Conotrachelus nenuphar*) is a true weevil native to the regions east of the Rocky Mountains in the United States and Canada. It is notorious for destroying fruits if left uncontrolled.



Poko noctuid moth

Agrotis crinigera

The poko noctuid moth (Agrotis crinigera) is a moth in the family Noctuidae. The species was first described by Arthur Gardiner Butler in 1881. It is endemic to the Hawaiian islands of Maui, Hawai‘i, and O‘ahu. It was said to be sometimes very...



Polish cochineal

Porphyrophora polonica

Polish cochineal (Porphyrophora polonica, Margarodes polonicus), also known as Polish carmine scales (Polish: czerwiec polski), is a scale insect formerly used to produce a crimson dye of the same name, colloquially known as "Saint John's blood"....



Polistes chinensis

Polistes chinensis

Polistes chinensis is a polistine vespid wasp in the cosmopolitan genus Polistes, and is commonly known as the Asian, Chinese or Japanese paper wasp. It is found in East Asia, in particular China and Japan. The subspecies P. chinensis antennalis...



Polyergus lucidus

Polyergus lucidus

Polyergus lucidus is a species of slave-making ant in the subfamily Formicinae endemic to the eastern United States. It is an obligatory social parasite, unable to feed itself or look after its brood and reliant on ants of another species of the...



Polyergus samurai

Polyergus samurai

Polyergus samurai is a species of slave-making ant in the subfamily Formicinae.



Polygonia c-album

Polygonia c-album

Polygonia c-album, the comma, is a food generalist (polyphagous) butterfly species belonging to the family Nymphalidae. The angular notches on the edges of the forewings are characteristic of the genus Polygonia, which is why species in the genus...



Polygonia faunus

Polygonia faunus

Polygonia faunus, the green comma, Faunus comma, or Faunus anglewing is a butterfly of the family Nymphalidae.



Polygonia interrogationis

Polygonia interrogationis

Polygonia interrogationis, commonly called the question mark butterfly, is a North American nymphalid butterfly. It lives in wooded areas, city parks, generally in areas with a combination of trees and open space. The color and textured...



Polyrhachis sokolova

Polyrhachis sokolova

Polyrhachis sokolova is a species of ant from Australia and New Guinea that recently was discovered to be capable of surviving tidal inundations. Researchers at the James Cook University gained attention in 2006 after finding and reporting this...



Polyura athamas

Polyura athamas[1]

Polyura athamas, the common nawab, is a species of fast-flying canopy butterfly found in tropical Asia. It belongs to the Charaxinae (rajahs and nawabs) in the brush-footed butterfly family (Nymphalidae).It occurs in the Himalayas from Kashmir to...



Polyura schreiber

Polyura schreiber

Polyura schreiber, the blue nawab, is a butterfly species found in tropical Asia. It belongs to the Charaxinae (rajahs and nawabs) in the brush-footed butterfly family (Nymphalidae). It occurs from south India and Assam through Myanmar,...



Pontia daplidice

Pontia daplidice

Pontia daplidice, the Bath white, is a small butterfly of the family Pieridae, the yellows and whites, which occurs in the Palearctic region. It is common in central and southern Europe, migrating northwards every summer, often reaching southern...



Popa spurca

Popa spurca

Popa spurca, also known as the African twig mantis, is a species of mantis native to Africa. It takes its common name from its resemblance to twig from a woody plant and grows up to 8 centimetres (3.1 in) long if female or 7 cm (2.8 in) long if...



Poplar admiral

Limenitis populi

The poplar admiral (*Limenitis populi*) is a butterfly in the Limenitidine clade of the family Nymphalidae.



Prioneris thestylis

Prioneris thestylis

Prioneris thestylis, the spotted sawtooth, is a small butterfly of the family Pieridae, that is, the yellows and whites, which is found in India and Southeast Asia.



Prionus coriarius

Prionus coriarius

Prionus coriarius (sometimes referred to as "the tanner" or "the sawyer") is a species of longhorn beetle.



Proceratium google

Proceratium google

Proceratium google, also known as the Google ant, was discovered in Madagascar by Brian L. Fisher, Associate Curator of Entomology at the California Academy of Sciences. The ant has an oddly shaped abdomen, adapted for hunting its exclusive meal...



Prosopocoilus giraffa

Prosopocoilus giraffa

Prosopocoilus giraffa, the giraffe stag beetle, is the world's largest stag beetle and is a member of the family Lucanidae within the order Coleoptera. They have very long, toothed and notched mandibles that protrude about half the size of their...



Protambulyx strigilis

Protambulyx strigilis

Protambulyx strigilis, the streaked sphinx, is a moth of the family Sphingidae prevalent in the Americas from Florida to Central and South America. The species was first described by Carl Linnaeus in 1771.



Pseudococcus viburni

Pseudococcus viburni

Pseudococcus viburni (formerly Pseudococcus affinis (Maskell), and commonly known as the obscure mealybug and tuber mealybug) is a close relative of the grape mealybug (P. maritimus) and a pest of the vineyards around the world.



Pseudocreobotra wahlbergi

Pseudocreobotra wahlbergi

Pseudocreobotra wahlbergi, or the spiny flower mantis, is a small flower mantis...



Pseudophilotes sinaicus

Pseudophilotes sinaicus

Pseudophilotes sinaicus, the Sinai baton blue, one of the world's smallest butterflies, lives only on mountainside patches of Sinai thyme in an arid corner of the Sinai Peninsula in Egypt called Saint Katherine Protectorate.



Pseudovates arizonae

Pseudovates arizonae

Pseudovates arizonae, common name Arizona unicorn mantis, is a species of praying mantis native to North America and is only found in the state of Arizona. At least three other related Pseudovates are found in Mexico, and a similar-looking...



Pseudovates chlorophaea

Pseudovates chlorophaea

Pseudovates chlorophaea, with the common name Texas unicorn mantis, is a species of praying mantis in the family Mantidae. It is native to the southern United States, Mexico, Central America, and northern South America.



Psilogramma increta

Psilogramma increta

Psilogramma increta, the plain grey hawkmoth, is a moth of the family Sphingidae.



Psilogramma menephron

Psilogramma menephron

Psilogramma menephron, the privet hawk moth or large brown hawkmoth, is a member of the family Sphingidae. It was described by Pieter Cramer in 1780. It is usually found in Sri Lanka, India (including the Andaman Islands), Nepal, central and...



Psylla pyri

Psylla pyri

Psylla pyri, commonly known as the pear psylla or pear psyllid, is an insect in the family Psyllidae. Originating in Europe and Asia, it has spread to North America. It is a pest of pear trees, sucking the sap, damaging the foliage, flowers and...



Psyllobora vigintiduopunctata

Psyllobora vigintiduopunctata

Psyllobora vigintiduopunctata (often abbreviated to Psyllobora 22-punctata), the 22-spot ladybird, (earlier known as Thea vigintiduopunctata) is a common, 3–5 mm long ladybird native to Europe. The elytra are yellow in colour with 22 black spots....



Pteronarcys californica

Pteronarcys californica

Pteronarcys californica is a species of insect in the family Pteronarcyidae, the giant stoneflies and salmonflies. It is known commonly as a salmonfly. Salmonflies are an important aquatic insect for fly anglers and many nymph and adult fly...



Pterophorus pentadactyla

Pterophorus pentadactyla

Pterophorus pentadactyla, commonly known as the white plume moth, is a moth in the family Pterophoridae. It is found in the West Palearctic including North Africa and Europe. The wingspan is 26–34 mm (1.0–1.3 in). It is uniformly white, with the...



Purple hairstreak

Favonius quercus

The purple hairstreak (*Favonius quercus*) is a butterfly in the family Lycaenidae distributed throughout much of Europe, North Africa, Anatolia, Caucasus, and Transcaucasia. The larva feeds on *Quercus robur*, *Quercus petraea*, *Quercus cerris* and...



Pyrgus malvae

Pyrgus malvae

Pyrgus malvae, the grizzled skipper, is a butterfly species from the family Hesperiidae. It is a small skipper (butterfly) with a chequered pattern on its wings that appears to be black and white. This butterfly can be found throughout Europe and...



Pyrgus ruralis

Pyrgus ruralis

Pyrgus ruralis, the two-banded checkered skipper, is a species of skipper butterfly (family Hesperiidae). It is found from southern British Columbia and the Rocky Mountains of SW Alberta, with populations south to central California, Nevada,...



Pyrops candelaria

Pyrops candelaria

Pyrops candelaria (*Latarnaria candelaria* and *Fulgora candelaria* in older literature) is a species of planthopper often placed in the tribe Latarnariini. This species has been recorded from: Guangdong, Guangxi, Cambodia, Vietnam, Hong Kong, Laos,...



Pyrrharctia isabella

Pyrrharctia isabella

Pyrrharctia isabella, the Isabella tiger moth, whose larval form is called the banded woolly bear, woolly bear, or woolly worm, occurs in the United States and southern Canada. It was first formally named by James Edward Smith in 1797.



Pyrrhocoris apterus

Pyrrhocoris apterus

The firebug, Pyrrhocoris apterus, is a common insect of the family Pyrrhocoridae. Easily recognizable due to its striking red and black coloration, it may be confused with the similarly coloured though unrelated Corizus hyoscyami (cinnamon bug or...



Queen

Danaus gilippus

The queen butterfly (Danaus gilippus) is a North and South American butterfly in the family Nymphalidae with a wingspan...



Queen Alexandra's birdwing

Ornithoptera alexandrae

Ornithoptera alexandrae, the Queen Alexandra's birdwing, is the largest species of butterfly in the world, with females reaching wingspans slightly in excess of 25 cm to 28 cm (9.8 inches to 11 inches). This birdwing is restricted to the forests...



Queen of Spain fritillary

Issoria lathonia

The Queen of Spain fritillary (Issoria lathonia) is a butterfly of the family Nymphalidae.



Quino checkerspot

Euphydryas editha quino

The Quino checkerspot (*Euphydryas editha quino*) is a butterfly native to southern California and northwestern Baja California. It is a subspecies of the common Edith's checkerspot (*Euphydryas editha*) and the second such subspecies to be listed...



Rabdophaga rosaria

Rabdophaga rosaria

Rabdophaga rosaria is a gall midge which forms Camellia galls or terminal rosette gall on willow species. It was first described by Hermann Loew in 1850.



Raspberry crazy ant

Nylanderia fulva

The tawny crazy ant or Raspberry crazy ant, *Nylanderia fulva*, is an ant originating in South America. Like the longhorn crazy ant (*Paratrechina longicornis*), this species is called "crazy ant" because of its quick, unpredictable movements (the...



Raspberry beetle

Byturus tomentosus

The raspberry beetle (*Byturus tomentosus*) is a species of beetle in the fruitworm family Byturidae. It is a major pest that is widespread in north-central Europe, affecting raspberry, blackberry and loganberry plants. This species is related to...



Red flour beetle

Tribolium castaneum

The red flour beetle (*Tribolium castaneum*) is a species of beetle in the family Tenebrionidae, the darkling beetles. It is a worldwide pest of stored products, particularly food grains, and a model organism for ethological and food safety research.



Red harvester ant

Pogonomyrmex barbatus

Pogonomyrmex barbatus is a species of harvester ant from the genus Pogonomyrmex. Its common names include red ant and red harvester ant. These large (5– to 7-mm) ants prefer arid chaparral habitats and are native to the Southwestern United...



Red imported fire ant

Solenopsis invicta

Solenopsis invicta, the fire ant, or red imported fire ant (RIFA), is a species of ant native to South America. A member of the genus Solenopsis in the subfamily Myrmicinae, it was described by Swiss entomologist Felix Santschi as a variant of S....



Red locust

Nomadacris septemfasciata

The red locust (Nomadacris septemfasciata) is a large grasshopper species found in sub-Saharan Africa. Its name refers to the colour of its hind wings. It is sometimes called the criquet nomade in French, due to its nomadic movements in the dry...



Red underwing

Catocala nupta

The red underwing (Catocala nupta) is a moth of the family Erebidæ. The species was first described by Carl Linnaeus in his 1767 12th edition of Systema Naturae. This is a large (80 mm wingspan) nocturnal Palearctic (including Europe) species...



Red-veined darter

Sympetrum fonscolombii

The red-veined darter or nomad (Sympetrum fonscolombii) is a dragonfly of the genus Sympetrum.



Reticulitermes flavipes

Reticulitermes flavipes

Reticulitermes flavipes, the eastern subterranean termite, is the most common termite found in North America. These termites are the most economically important wood destroying insects in the United States and are classified as pests. They feed...



Rhagoletis cerasi

Rhagoletis cerasi

Rhagoletis cerasi (the cherry fruit fly or European cherry fruit fly) is a species of fruit fly in the family Tephritidae.



Rhagoletis juglandis

Rhagoletis juglandis

Rhagoletis juglandis, also known as the walnut husk fly, is a species of tephritid or fruit fly in the family Tephritidae. It is closely related to the walnut husk maggot Rhagoletis suavis (Loew, 1862). This species of fly belongs to the R....



Rhingia campestris

Rhingia campestris

Rhingia campestris is a species of hoverfly, 7–11 millimetres (0.3–0.4 in) long, with a wingspan of 12–18 mm (0.5–0.7 in). It is common across the Palearctic from March until November. It has a broad orange abdomen with a black line along the...



Rhinocyllus conicus

Rhinocyllus conicus

Rhinocyllus conicus is a species of true weevil. It is best known as a controversial agent of biological pest control which has been used against noxious thistles in the genera Carduus, Cirsium, Onopordum, and Silybum. The adult weevil is black...



Rhombodera extensicollis

Rhombodera extensicollis

Rhombodera extensicollis is a species of praying mantises in the family Mantidae, found in Indomalaya.



Rhombodera megaera

Rhombodera megaera

Rhombodera megaera is a species of praying mantises in the family Mantidae, found in China and Thailand.



Rhyothemis variegata

Rhyothemis variegata

Rhyothemis variegata, known as the common picture wing or variegated flutterer, is a species of dragonfly of the family Libellulidae, found in South Asia.



Ringlet

Aphantopus hyperantus

The ringlet (Aphantopus hyperantus) is a butterfly in the family Nymphalidae. It is only one of the numerous "ringlet" butterflies in the tribe Satyrini.



Roesel's bush-cricket

Roeseliana roeselii

Roesel's bush-cricket, Roeseliana roeselii (synonym Metrioptera roeselii) is a European bush-cricket, named after August Johann Rösel von Rosenhof, a German entomologist.



Romalea

Romalea microptera

Romalea is a genus of grasshoppers native to the Southeastern and South-central United States. As traditionally defined, it contains a single species, Romalea microptera, known commonly as the Georgia Thumper, eastern lubber grasshopper, Florida...



Ropalidia marginata

Ropalidia marginata

Ropalidia marginata is an Old World species of paper wasp. It is primitively eusocial, not showing the same bias in brood care seen in other social insects with greater asymmetry in relatedness. The species employs a variety of colony founding...



Rosalia longicorn

Rosalia alpina

The Rosalia longicorn (Rosalia alpina) or Alpine longhorn beetle, is a large longicorn (family Cerambycidae) that is distinguished by its distinctive markings.



Roseate skimmer

Orthemis ferruginea

The roseate skimmer (Orthemis ferruginea) is a species of dragonfly in the family Libellulidae. It is native to the Americas, where its distribution extends from the United States to Brazil. It is common and widespread. It is an introduced...



Rothschild's birdwing

Ornithoptera rothschildi

Rothschild's birdwing (Ornithoptera rothschildi) is a large birdwing butterfly, endemic to the Arfak Mountains in Western New Guinea. What was originally described as Ornithoptera akakeae, is a supposed natural hybrid between Ornithoptera...



Rough prominent

Nadata gibbosa

The rough prominent (Nadata gibbosa) is a moth of the family Notodontidae, subfamily Phalerinae. It is also known as the white-dotted prominent and the tawny prominent. The species was first described by James Edward Smith in 1797. This common...



Ruddy darter

Sympetrum sanguineum

The ruddy darter (Sympetrum sanguineum) is a species of dragonfly of the family Libellulidae.



Russian wheat aphid

Diuraphis noxia

The Russian wheat aphid (Diuraphis noxia) is an aphid that can cause significant losses in cereal crops. The species was introduced to the United States in 1986 and is considered an invasive species there. This aphid is pale green and up to 2 mm...



Saddleback caterpillar

Acharia stimulea

The saddleback caterpillar (Acharia stimulea, formerly Sibine stimulea) is the larva of a species of moth native to eastern North America. It is also found in Mexico. The species belongs to the family of slug caterpillars, Limacodidae. The larva...



Saga pedo

Saga pedo

Saga pedo is a species of wingless bush cricket from the southern half of Europe and western and central Asia. This brown or green bush cricket typically has a total length, from the head to the tip of the ovipositor, of up to 10.5 cm (4.1 in),...



Saharan silver ant

Cataglyphis bombycina

The Saharan silver ant (Cataglyphis bombycina) is a species of insect that lives in the Sahara Desert. It is the fastest of the world's 12,000 known ant species, clocking a velocity of 855 millimetres per second (over 1.9 miles per hour or 3.1...



Saint Francis's satyr

Neonympha mitchellii francisci

The Saint Francis's satyr (Neonympha mitchellii francisci) is an endangered butterfly subspecies found only in the US state of North Carolina. First discovered in 1983, it was first described by David K. Parshall and Thomas W. Kral in 1989 and...



Salt Creek tiger beetle

Cicindela nevadica lincolniana

The Salt Creek tiger beetle (Cicindela nevadica lincolniana) is a critically endangered subspecies of tiger beetle endemic to the saline wetlands of northern Lancaster County, Nebraska, adjacent to and immediately to the north of the city of...



Samia cynthia

Samia cynthia

Samia cynthia, the ailanthus silkworm, is a saturniid moth, used to produce silk fabric but not as domesticated as the silkworm, Bombyx mori. The moth has very large wings of 113–125 mm (4.4–4.9 in), with a quarter-moon shaped spot on both the...



Sara longwing

Heliconius sara

The Sara longwing (Heliconius sara) is a species of neotropical heliconiid butterfly found from Mexico to the Amazon Basin and southern Brazil. It is a colourful species: the dorsal wing surface is black with a large medial patch of metallic blue...



Sarcophaga africa

Sarcophaga africa

Sarcophaga (Bercaea) africa is a species of fly belonging to the family Sarcophagidae, the flesh-flies. It is the best known species in its genus. S. africa feeds on living and dead tissue, including snails, and other decomposing matter, and...



Sarcophaga pernix

Sarcophaga pernix

Sarcophaga pernix, also known as the red-tailed flesh fly, is a fly in the Sarcophagidae family. This fly often breeds in carrion and feces, making it a possible vector for disease. The larvae of this species can cause myiasis, as well as...



Saturnia pavonia

Saturnia pavonia

Saturnia pavonia, the small emperor moth, is a moth of the family Saturniidae. It was first described by Carl Linnaeus in his 1758 10th edition of Systema Naturae. Sometimes, the incorrect genus name Pavonia is still used for this species. This...



Saturnia pyri

Saturnia pyri

Saturnia pyri, the giant peacock moth, great peacock moth, giant emperor moth or Viennese emperor, is a Saturniid moth which is native to Europe. The species was first described by Michael Denis and Ignaz Schiffermüller in 1775. It is the largest...



Scaeva pyrastris

Scaeva pyrastris

Scaeva pyrastris, common name the pied hoverfly, is a species of hoverfly.



Scaeva selenitica

Scaeva selenitica

Scaeva selenitica is a species of hoverfly.



Scarce large blue

Phengaris teleius

The scarce large blue (Phengaris teleius) is a species of butterfly in the family Lycaenidae. It is found in Austria, Slovenia, Croatia, the Czech Republic, France, Georgia, Germany, Hungary, Italy, Japan, Kazakhstan, Mongolia, the Netherlands,...



Scarce swallowtail

Iphiclides podalirius

The scarce swallowtail (Iphiclides podalirius) is a butterfly belonging to the family Papilionidae. It is also called the sail swallowtail or pear-tree swallowtail.



Scarlet dragonfly

Crocothemis erythraea

The scarlet dragonfly (Crocothemis erythraea) is a species of dragonfly in the family Libellulidae. Its common names include broad scarlet, common scarlet-darter, and scarlet darter.



Scarlet dwarf

Nannophya pygmaea

Nannophya pygmaea, known variously as the scarlet dwarf, northern pygmyfly, or tiny dragonfly, is a dragonfly of the family Libellulidae, native from Southeast Asia to China and Japan, occasionally found south to Australia.



Scarlet lily beetle

Lilioceris lili

The scarlet lily beetle, red lily beetle, or lily leaf beetle (Lilioceris lili), is a leaf beetle that eats the leaves, stem, buds, and flowers, of lilies, fritillaries and other members of the family Liliaceae. It lays its eggs most often on...



Scarlet skimmer

Crocothemis servilia

The scarlet skimmer or ruddy marsh skimmer, Crocothemis servilia, is a species of dragonfly of the family Libellulidae, native to east and southeast Asia and introduced to Jamaica, Florida, and Hawaii.



Scarlet tiger moth

Callimorpha dominula

The scarlet tiger moth (Callimorpha dominula, formerly Panaxia dominula) is a colorful moth belonging to the tiger moth subfamily, Arctiinae. The species was first described by Carl Linnaeus in his 1758 10th edition of Systema Naturae.



Scathophaga stercoraria

Scathophaga stercoraria

Scathophaga stercoraria, commonly known as the yellow dung fly or the golden dung fly, is one of the most familiar and abundant flies in many parts of the Northern Hemisphere. As its common name suggests, it is often found on the feces of large...



Sceliphron caementarium

Sceliphron caementarium

Sceliphron caementarium, also known as the yellow-legged mud-dauber wasp, black-and-yellow mud dauber (within the US), or black-waisted mud-dauber (outside of the US), is a species of sphecid wasp. There are some 30 other species of Sceliphron...



Sceliphron spirifex

Sceliphron spirifex

Sceliphron spirifex is a species of sphecid wasp. It has a medium-sized body (17–27 millimetres or 0.7–1.1 inches), which is dull black with a long, yellow petiole (waist). The legs are black with yellow bands, the antennae are black and the...



Schizodactylus monstrosus

Schizodactylus monstrosus

Schizodactylus monstrosus or the maize cricket, is a species of large, robust cricket found in Asia, belonging to the family Schizodactylidae. It is found mainly in sandy habitats along rivers, and has large flattened tarsal extensions and wings...



Scirpophaga innotata

Scirpophaga innotata

Scirpophaga innotata, the rice white stemborer, is a species of moth of the family Crambidae. The species was described by Francis Walker in 1863. It is found in Indonesia, Pakistan, the Philippines and the tropical north of Australia. The larvae...



Scolitantides orion

Scolitantides orion

Scolitantides orion, the chequered blue, is a species of butterfly in the family Lycaenidae (gossamer-winged butterflies). It is found in Europe, Russia and east across the Palearctic to Japan.



Scotch argus

Erebia aethiops

The Scotch argus (Erebia aethiops) is a butterfly of the family Nymphalidae. In spite of its English name argus, it is not a close relation of the brown argus nor the northern brown argus.



Senegalese grasshopper

Oedaleus senegalensis

The Senegalese grasshopper (Oedaleus senegalensis) is a medium-sized grasshopper species found in the Sahel region of Africa, the Canary Islands, Cape Verde Islands, and West Asia. Although not called a locust in English, this species shows...



Setaceous Hebrew character

Xestia c-nigrum

The setaceous Hebrew character (*Xestia c-nigrum*) is a moth of the family Noctuidae. The species was first described by Carl Linnaeus in his 1758 10th edition of Systema Naturae. It is found in the Palearctic realm. It is a common species...



Shark

Cucullia umbratica

The shark (*Cucullia umbratica*) is a moth of the family Noctuidae. The species was first described by Carl Linnaeus in his landmark 1758 10th edition of Systema Naturae.



Sheep moth

Hemileuca eglanderina

The sheep moth, or common sheep moth (*Hemileuca eglanderina*), is a member of the family Saturniidae of silk moths and is native to western North America. In California, its range is west of the Sierran crest and the mountains of Southern...



Short-haired bumblebee

Bombus subterraneus

The short-haired bumblebee (*Bombus subterraneus*), or short-haired humble-bee, is a species of bumblebee found in Eurasia, as well as in New Zealand, where it is an introduced species. It lived in the British Isles among other parts of Europe. This...



Short-tailed blue

Cupido argiades

The short-tailed blue or tailed Cupid (*Cupido argiades*) is a butterfly that forms part of the family Lycaenidae. It is found from Europe to Japan and in India.



Silver Y

Autographa gamma

The silver Y (Autographa gamma) is a migratory moth of the family Noctuidae which is named for the silvery Y-shaped mark on each of its forewings.



Silver-studded blue

Plebejus argus

The silver-studded blue (Plebejus argus) is a butterfly in the family Lycaenidae. It has bright blue wings rimmed in black with white edges and silver spots on its hindwings, lending it the name of the silver-studded blue. P. argus can be found...



Silver-washed fritillary

Argynnis paphia

The silver-washed fritillary (Argynnis paphia) is a common and variable butterfly found over much of the Palearctic realm – Algeria, Europe and across the Palearctic to Japan.



Silverleaf whitefly

Bemisia tabaci

The silverleaf whitefly (Bemisia tabaci, also informally referred to as the sweet potato whitefly) is one of several species of whitefly that are currently important agricultural pests. A review in 2011 concluded that the silverleaf whitefly is...



Simosyrphus grandicornis

Simosyrphus grandicornis

Simosyrphus grandicornis is an Australasian species of hoverfly, and is one of the two most common hoverflies in Australia, alongside Melangyna viridiceps. It has been introduced to a number of Polynesian Islands and Hawaii.



Sinea diadema

Sinea diadema

Sinea diadema is a species of assassin bug family (Reduviidae), in the subfamily Harpactorinae. Sinea diadema is bivoltine, preys on small bugs and beetles, and overwinters in the egg stage.



Siproeta epaphus

Siproeta epaphus

Siproeta epaphus, the rusty-tipped page or brown siproeta, is a New World butterfly that lives all year in tropical habitats. It has large wings, averaging 7.0–7.5 cm (2.8–3.0 in), that are black in the inner portion of the top surface and brown...



Siproeta stelenes

Siproeta stelenes

Siproeta stelenes (malachite) is a Neotropical brush-footed butterfly (family Nymphalidae). The malachite has large wings that are black and brilliant green or yellow-green on the upperside and light brown and olive green on the underside. It is...



Six-spot burnet

Zygaena filipendulae

The six-spot burnet (Zygaena filipendulae) is a day-flying moth of the family Zygaenidae.



Small angle shades

Euplexia lucipara

The small angle shades (Euplexia lucipara) is a moth of the family Noctuidae. It is distributed throughout the Palearctic. The species was first described by Carl Linnaeus in his 1758 10th edition of Systema Naturae. As the common name suggests,...



Small blue

Cupido minimus

The small blue (Cupido minimus) is a Palearctic butterfly in the family Lycaenidae. Despite its common name, it is not particularly blue. The male has some bluish suffusion at the base of its upper wings but is mostly dark brown like the female....



Small dusty wave

Idaea seriata

The small dusty wave (Idaea seriata) is a moth of the family Geometridae first described by Franz von Paula Schrank in 1802. It is found throughout Western, Central and Northern Europe. In the north, its range extends as far as Denmark and...



Small heath

Coenonympha pamphilus

The small heath (Coenonympha pamphilus) is a butterfly species belonging to the family Nymphalidae, classified within the subfamily Satyrinae (commonly known as "the browns"). It is the smallest butterfly in this subfamily. The small heath is...



Small hive beetle

Aethina tumida

The small hive beetle (*Aethina tumida*) is a beekeeping pest. It is native to sub-Saharan Africa, but has spread to many other locations, including North America, Australia, and the Philippines. The small hive beetle can be a destructive pest of...



Small pearl-bordered fritillary

Boloria selene

Boloria selene, known in Europe as the small pearl-bordered fritillary and in North America as the silver-bordered fritillary, is a species of butterfly of the family Nymphalidae. It is found across Europe, Asia and North America, and feeds...



Small red damselfly

Ceriagrion tenellum

The small red damselfly (*Ceriagrion tenellum*) is a small damselfly flying in heathland bogs and streams. It is in the family Coenagrionidae.



Small skipper

Thymelicus sylvestris

The small skipper (*Thymelicus sylvestris*) is a butterfly of the family Hesperiidae.



Small tortoiseshell

Aglais urticae

The small tortoiseshell (*Aglais urticae*) is a colourful Eurasian butterfly in the family Nymphalidae. Adults feed on nectar and may hibernate over winter; in warmer climates they may have two broods in a season. While the dorsal surface of the...



Smerinthus cerisyi

Smerinthus cerisyi

Smerinthus cerisyi, the one-eyed sphinx or Cerisy's sphinx, is a moth of the family Sphingidae. The species was first described by William Kirby who named the species in honor of Alexandre Louis Lefèbvre de Cérisy in 1837.



Smerinthus jamaicensis

Smerinthus jamaicensis

Smerinthus jamaicensis, the twin-spotted sphinx, is a moth of the family Sphingidae. The species was first described by Dru Drury in 1773.



Smerinthus ocellatus

Smerinthus ocellatus

Smerinthus ocellatus, the eyed hawk-moth, is a European moth of the family Sphingidae. The species was first described by Carl Linnaeus in his 1758 10th edition of Systema Naturae. The eyespots are not visible in resting position, where the...



Smokybrown cockroach

Periplaneta fuliginosa

The smokybrown cockroach (Periplaneta fuliginosa) is a large species of cockroach, winged, and growing to a length of 32–35 millimetres (1.3–1.4 in).



Solenopsis daguerrei

Solenopsis daguerrei

Solenopsis daguerrei is a species of parasitic ants native to Argentina and Uruguay. The young queens of the species invade the nests of other species, such as the red imported fire ant (RIFA) (Solenopsis invicta). This is made possible by the...



Solenopsis fugax

Solenopsis fugax

Solenopsis fugax is a myrmicine ant of the genus Solenopsis. It is the only member of its genus to be native to the British Isles, and although rare, it has been taken by Horace Donisthorpe in a number of localities on England's southern coast,...



Solenopsis geminata

Solenopsis geminata

Solenopsis geminata or tropical fire ant is a species of fire ants, described by Fabricius in 1804, in the tribe Solenopsidini; it was originally placed in the Atta genus. This species has a pan-tropical distribution.



Southern fire ant

Solenopsis xyloni

The southern fire ant (Solenopsis xyloni), also known as the Californian fire ant or cotton ant,:89 is a stinging fire ant native to southern parts of the United States. Its behaviour is similar to the red imported fire ant (S. invicta),...



Southern hawker

Aeshna cyanea

The southern hawker or blue hawker (Aeshna cyanea) is a species of hawker dragonfly.



Spanish fly

Lytta vesicatoria

The Spanish fly (Lytta vesicatoria) is an aposematic emerald-green beetle in the blister beetle family (Meloidae). It is distributed across Eurasia. The species and others in its family were used in traditional apothecary preparations as...



Speckled bush-cricket

Leptophyes punctatissima

The speckled bush-cricket (Leptophyes punctatissima) is a flightless species of bush-cricket belonging to the family Tettigoniidae. The species was originally described as Locusta punctatissima in 1792.



Speckled wood

Pararge aegeria

The speckled wood (Pararge aegeria) is a butterfly found in and on the borders of woodland areas throughout much of the Palearctic realm. The species is subdivided into multiple subspecies, including Pararge aegeria aegeria, Pararge aegeria...



Speiredonia cthulhui

Speiredonia cthulhui

Speiredonia cthulhui is a species of moth of the family Erebidæ first described by Alberto Zilli and Jeremy D. Holloway in 2005. It is found on New Guinea. The length of the forewings is 26.5 mm for males and 25.5 mm for females. It is named after...



Speiredonia spectans

Speiredonia spectans

Speiredonia spectans, the granny's cloak moth, is a moth of the family Erebidæ. The species was first described by Achille Guenée in 1852. It is found in north-eastern Australia and Tasmania. Strays have been recorded on Norfolk Island and in...



Speyeria atlantis

Speyeria atlantis

Speyeria atlantis, the Atlantis fritillary, is a butterfly of the family Nymphalidae of North America. It is from the Avalon Peninsula of Newfoundland and Labrador to northern British Columbia, across the northern United States south as far as...



Sphaerophoria scripta

Sphaerophoria scripta

Sphaerophoria scripta, the long hoverfly, is a species of hoverfly belonging to the family Syrphidae.



Sphecius speciosus

Sphecius speciosus

Sphex speciosus, often simply referred to as the cicada killer or the cicada hawk, is a large, solitary digger wasp species in the family Crabronidae. The name may be applied to any species of crabronid that preys on cicadas, though in North...



Sphenarches anisodactylus

Sphenarches anisodactylus

Sphenarches anisodactylus, commonly known as the geranium plume moth, is a species of moth in the family Pterophoridae. It is found in western Africa, Madagascar, India, Sri Lanka, Thailand, Japan, the New Hebrides and Central and South America,...



Sphex pensylvanicus

Sphex pensylvanicus

Sphex pensylvanicus, the great black wasp, is a species of digger wasp. It lives across most of North America and grows to a size of 20–35 mm (0.8–1.4 in). The larvae feed on living insects that the females paralyze and carry to the underground...



Sphinx chersis

Sphinx chersis

Sphinx chersis, the great ash sphinx or northern ash sphinx, is a moth that belongs to the family Sphingidae.



Sphinx ligustri

Sphinx ligustri

Sphinx ligustri, the privet hawk moth, is a moth found in most of the Palearctic realm. The species was first described by Carl Linnaeus in his 1758 10th edition of Systema Naturae.



Sphinx pinastri

Sphinx pinastri

Sphinx pinastri, the pine hawk-moth, is a moth of the family Sphingidae. It is found in Palearctic realm and sometimes the Nearctic realm. This species has been found in Scotland but is usually found in England. The species was first described by...



Sphodromantis gastrica

Sphodromantis gastrica

Sphodromantis gastrica, with the common names African mantis or common green mantis, is a species of praying mantis from Africa. It averages 65–75 millimetres (2.6–3.0 in) in body length, and has a very diverse diet, hunting any prey of reasonable...



Spined soldier bug

Podisus maculiventris

The spined soldier bug (Podisus maculiventris) is a species of stink bug common in North America. They are predators of gypsy moth caterpillars and the larvae of beetles such as the Colorado potato beetle and the Mexican bean beetle. Since the...



Spitfire sawfly

Perga affinis

The spitfire sawfly (Perga affinis, family Pergidae) is a hymenopteran insect found in Australia. It is up to 22 mm long, has two pairs of wings, with a wingspan up to 40 mm, and its wings are honey colored. Its larvae are up to 80 mm long and...



Spodoptera eridania

Spodoptera eridania

Spodoptera eridania (southern armyworm) is a moth that is known to be a pest. They are one of the most important defoliators in the tropical and subtropical regions of the western hemisphere that feed heavily on plants while they are young, often...



Spodoptera ornithogalli

Spodoptera ornithogalli

Spodoptera ornithogalli (yellow-striped armyworm, cotton cutworm) is a moth of the family Noctuidae. When first discovered this particular species was thought to be the American representative of S. littoralis as the two species have very similar...



Square-spot rustic

Xestia xanthographa

The square-spot rustic (Xestia xanthographa) is a moth of the family Noctuidae. It is found in Europe, North Africa and east across the Palearctic (excluding China) and in North America. The species is quite variable in appearance, the forewings...



Squash vine borer

Melittia cucurbitae

The squash vine borer (Melittia cucurbitae) is a diurnal species of sesiid moth. The moth is often mistaken for a bee or wasp because of its movements, and the bright orange hind leg scales. The females typically lay their eggs at the base of...



Stable fly

Stomoxys calcitrans

Stomoxys calcitrans is commonly called the stable fly, barn fly, biting house fly, dog fly, or power mower fly. Unlike most members of the family Muscidae, Stomoxys calcitrans ('sharp mouth' + 'kicking') and others of its genus suck blood from...



Stagmomantis limbata

Stagmomantis limbata

Stagmomantis limbata, common name bordered mantis, bosque mantis, Arizona mantis, or New Mexico praying mantis, is a species of praying mantis native to North America, most prevalent in the south-western United States. This beneficial insect is...



Stagmomantis montana

Stagmomantis montana

Stagmomantis montana (Rehn, 1935), common name Mountain mantis, is a species of praying mantis in the family Mantidae. They are native to Mexico and Central America. S. montana sinaloae has been identified as a subspecies.



Stalk borer

Papaipema nebris

The stalk borer (Papaipema nebris) is a moth of the family Noctuidae. It is found from southern Canada, through the Eastern United States (East of the Rocky Mountains) to the Gulf of Mexico, although it is absent from Florida. This wingspan is...



Statilia maculata

Statilia maculata

Statilia maculata, common name Asian jumping mantis or "小螳螂", ko-kamakiri (Japanese meaning "small mantis") or "좀사마귀", joem-sa-ma-gui (Korean meaning "small mantis"), is a species of mantis native to Asia that can be found in Romania, China and...



Stenocara gracilipes

Stenocara gracilipes

Stenocara gracilipes is a species of beetle that is native to the Namib Desert in southern Africa. This is one of the most arid areas of the world, receiving only 1.4 centimetres (0.55 in) of rain per year. The beetle is able to survive by...



Stibochiona nicea

Stibochiona nicea

Stibochiona nicea, the popinjay, is a species of nymphalid butterfly found in Asia.



Strategus aloeus

Strategus aloeus

Strategus aloeus, the ox beetle, is a species of rhinoceros beetle native to the Americas.



Strigiphilus garylarsoni

Strigiphilus garylarsoni

Strigiphilus garylarsoni is a species of chewing louse found only on owls. The species has no common name. The species was first described by biologist Dale H. Clayton in 1990, who named it after cartoonist Gary Larson.



Suastus gremius

Suastus gremius

Suastus gremius, the Indian palm bob or palm bob, is a butterfly belonging to the family Hesperiiidae. It is found in the Indomalayan realm.



Sunset morpho

Morpho hecuba

Morpho hecuba, the sunset morpho, is a Neotropical butterfly and the largest species in the genus Morpho. Its wingspan can reach 20 centimetres (7.9 in), but is usually from 13–15 centimetres (5.1–5.9 in). "M. hecuba is the largest known Morpho...



Surinam cockroach

Pycnoscelus surinamensis

The Surinam cockroach or greenhouse cockroach (Pycnoscelus surinamensis) is a species of burrowing cockroach. It is a common plant pest endemic to the Indomalayan realm that has spread to tropical and into subtropical regions around the world,...



Svensson's copper underwing

Amphipyra berbera

Svensson's copper underwing (*Amphipyra berbera*) is a moth of the family Noctuidae. The species was first described by Charles E. Rungs in 1949. It is distributed throughout Europe including Russia east to the Urals. This species has a wingspan of...



Swallow-tailed moth

Ourapteryx sambucaria

The swallow-tailed moth (*Ourapteryx sambucaria*) is a moth of the family Geometridae. The species was first described by Carl Linnaeus in his 1758 10th edition of *Systema Naturae*. It is a common species across Europe and the Near East.



Sycamore

Acronicta aceris

The sycamore (*Acronicta aceris*) is a moth of the family Noctuidae. The species was first described by Carl Linnaeus in his 1758 10th edition of *Systema Naturae*. It is distributed through most of Europe, from central England south to Morocco. To...



Sympetrum frequens

Sympetrum frequens

Sympetrum frequens, the Autumn darter (アキアカネ, akiakane), is a species of dragonfly endemic to Japan. It lives near grasslands and can grow up to 40 mm in length. After emerging, these dragonflies migrate to high mountains where they feed until...



Symphaedra

Euthalia nais

Symphaedra nais, also known as the baronet, is a species of Nymphalid butterfly found in South Asia. It was formerly included in the genus *Euthalia* but it is a sister to members of the genus *Bassarona*.



Synanthedon exitiosa

Synanthedon exitiosa

Synanthedon exitiosa, the peachtree borer, is a species of moth in the family Sesiidae that is native to North America. The adult female of the species has a wingspan of about 3.5 centimeters. It is dark metallic blue in color with an orange band...



Synanthedon tipuliformis

Synanthedon tipuliformis

Synanthedon tipuliformis, known as the currant clearwing, is a moth of the family Sesiidae. It is endemic to the Palearctic realm, but is an invasive species in the Nearctic realm and the Australasian realm.



Synemon plana

Synemon plana

Synemon plana, commonly known as the golden sun moth, is a diurnal moth native to Australia and throughout its range, it is currently classified as vulnerable under the Environment Protection and Biodiversity Conservation Act 1999.



Syritta pipiens

Syritta pipiens

Syritta pipiens, sometimes called the thick-legged hoverfly, is one of the most common species in the insect family Syrphidae. This fly originates from Europe and is currently distributed across Eurasia and North America. They are fast and nimble...



Syrphus ribesii

Syrphus ribesii

Syrphus ribesii is a very common Holarctic species of hoverfly. Its larvae feed on aphids. In common with many other species of hoverfly, males have the eyes meeting on the top of the head, whilst females have their eyes widely separated.



Tabanus lineola

Tabanus lineola

Tabanus lineola, also known as the striped horse fly, is a species of biting horse-fly. It is known from the eastern coast of North America and the Gulf coast of Mexico.



Talicauda nyseus

Talicauda nyseus

Talicauda nyseus, the red Pierrot, is a small but striking butterfly found in the Indian subcontinent and South-East Asia belonging to the lycaenids, or blues family. The red Pierrots, often found perching on its larva host plant, Kalanchoe, are...



Tanaecia lepidea

Tanaecia lepidea

Tanaecia lepidea, the grey count, is a species of nymphalid butterfly found in South and Southeast Asia.



Tansy beetle

Chrysolina graminis

The tansy beetle (Chrysolina graminis) is a species of leaf beetle. The common name derives from its main foodplant, Tansy (Tanacetum vulgare), but it can also use other wetland plants such as Gypsywort (Lycopus europaeus) and Water Mint (Mentha...



Tapinoma melanocephalum

Tapinoma melanocephalum

Tapinoma melanocephalum is a species of ant that goes by the common name ghost ant. They are recognised by their dark head and pale or translucent legs and gaster (abdomen). This colouring makes this tiny ant seem even smaller.



Tapinoma sessile

Tapinoma sessile

Tapinoma sessile is a species of small ant that goes by the common names odorous house ant, sugar ant, stink ant, and coconut ant. Their colonies are polydomous (consist of multiple nests) and polygynous (contain multiple reproducing queens)....



Tarnished plant bug

Lygus lineolaris

The tarnished plant bug (TPB), Lygus lineolaris, is a species of plant-feeding insect in the family Miridae. It has piercing-sucking mouthparts and has become a serious pest on small fruits and vegetables in North America. It is considered a...



Tarucus balkanicus

Tarucus balkanicus

Tarucus balkanicus, the Balkan Pierrot or little tiger blue, is a small butterfly that belongs to the lycaenids or blues family. It is found in Mauritania, Niger (the Aïr region), Sudan (Khartoum), Uganda, Saudi Arabia, the United Arab Emirates,....



Teinopalpus imperialis

Teinopalpus imperialis

Teinopalpus imperialis, the Kaiser-i-Hind, is a rare species of swallowtail butterfly found from Nepal and north India east to north Vietnam. The common name literally means "emperor of India". The Kaiser-i-Hind is much sought after by butterfly...



Telegonus fuligator

Telegonus fuligator

Telegonus fuligator, the two-barred flasher, is a species of spread-wing skipper butterfly in the family Hesperiidae which may constitute a possible cryptic species complex. It ranges all over the Americas, from the southern United States to...



Telephone-pole beetle

Micromalthus debilis

The telephone-pole beetle (Micromalthus debilis) is a beetle native to the eastern United States, and the only living representative of the otherwise extinct family Micromalthidae. They have an unusual life cycle involving asexually reproducing...



Temnostoma vespiforme

Temnostoma vespiforme

Temnostoma vespiforme is a species of hoverfly. Larva of this species feed in decaying wood of deciduous trees.



Ten-lined June beetle

Polyphylla decemlineata

The ten-lined June beetle (Polyphylla decemlineata), also known as the watermelon beetle, is a scarab beetle found in the western United States and Canada. The adults are attracted to light and feed on foliage. They can make a hissing sound when...



Tenebrio obscurus

Tenebrio obscurus

Tenebrio obscurus, or the dark mealworm beetle, is a species of darkling beetle whose larvae are known as mini mealworms. These insects should not be confused with younger mealworms (Tenebrio molitor) or with the confused flour beetle (Tribolium...



Tenodera angustipennis

Tenodera angustipennis

Tenodera angustipennis is a species of mantis native to Asia and nearby areas of Oceania. The species was introduced and became established in the eastern United States. Tenodera angustipennis was noticed as early as 1921 in Aberdeen, Maryland,...



Tenodera australasiae

Tenodera australasiae

Tenodera australasiae, the purple-winged mantis, is species of praying mantis. Found throughout Australia, it is common in the eastern regions. Both males and females are capable of flight. The species has not been shown to be parthenogenetic.



Tetragonula carbonaria

Tetragonula carbonaria

Tetragonula carbonaria (previously known as Trigona carbonaria) is a stingless bee, endemic to the north-east coast of Australia. Its common name is sugarbag bee. They are also occasionally referred to as bush bees. The bee is known to pollinate...



Tetramorium caespitum

Tetramorium caespitum

Tetramorium caespitum, also known as the pavement ant, is a species of ant in the family Formicidae.



Tetramorium immigrans

Tetramorium immigrans

Tetramorium immigrans—also known as the immigrant pavement ant, pavement ant, and the sugar ant in parts of North America—is an ant native to Europe, which also occurs as an introduced pest in North America. Its common name comes from the fact...



Tettigonia viridissima

Tettigonia viridissima

Tettigonia viridissima, the great green bush-cricket, is a large species of bush-cricket belonging to the subfamily Tettigoniinae.



Texas leafcutter ant

Atta texana

The Texas leafcutter ant (Atta texana) is a species of fungus-farming ant in the subfamily Myrmicinae. It is found in Texas, Louisiana, and north-eastern Mexico. Other common names include town ant, parasol ant, fungus ant, cut ant, and night...



Theretra oldenlandiae

Theretra oldenlandiae

Theretra oldenlandiae, the impatiens hawkmoth, taro hornworm or white-banded hunter hawkmoth, is a member of the family Sphingidae.



Theretra silhetensis

Theretra silhetensis

Theretra silhetensis, the brown-banded hunter hawkmoth, is a moth of the family Sphingidae described by Francis Walker in 1856. It lives in Indo-Australia, India, Sri Lanka, Papua New Guinea, East Australia, Solomon Islands, Fiji Islands, Vanuatu...



Tholymis tillarga

Tholymis tillarga

Tholymis tillarga, the coral-tailed cloudwing, is a species of dragonfly in the family Libellulidae. It is found from tropical West Africa to Asia, Australia and the Pacific Islands. Common names include old world twister, evening skimmer....



Thyreus nitidulus

Thyreus nitidulus

Thyreus nitidulus, commonly known as the neon cuckoo bee, is a parasitic bee of the genus Thyreus, called cuckoo bees. It is a stocky bee, notable for its brilliant metallic blue- and black-banded colors.Originally described by Danish...



Thysania agrippina

Thysania agrippina

Thysania agrippina is a species of moth in the family Erebidæ. It was described by Maria Sibylla Merian in her 1705 publication Metamorphosis insectorum Surinamensium, and Pieter Cramer provided the formal description of the species in 1776. The...



Timberman beetle

Acanthocinus aedilis

The timberman beetle (Acanthocinus aedilis) is a species of woodboring beetle belonging to the longhorn beetle family.It is found in woodlands, with a large distribution through Europe, Russia and Central Asia. It is also known as the Siberian...



Tinea pellionella

Tinea pellionella

Tinea pellionella, the case-bearing clothes moth, is a species of tineoid moth in the family Tineidae, the fungus moths. This species has a cosmopolitan distribution, occurring nearly worldwide.



Tineola bisselliella

Tineola bisselliella

Tineola bisselliella, known as the common clothes moth, webbing clothes moth, or simply clothing moth, is a species of fungus moth (family Tineidae, subfamily Tineinae). It is the type species of its genus Tineola and was first described by the...



Tirumala limniace

Tirumala limniace

Tirumala limniace, the blue tiger, is a butterfly found in South Asia, and Southeast Asia that belongs to the brush-footed butterfly family, Nymphalidae. More specifically it is part of the crows and tigers or danaid group. This butterfly shows...



Tirumala septentrionis

Tirumala septentrionis

Tirumala septentrionis, the dark blue tiger, is a danaid butterfly found in the Indian subcontinent and Southeast Asia.



Titan beetle

Titanus giganteus

The titan beetle (Titanus giganteus) is a Neotropical longhorn beetle, the sole species in the genus Titanus, and one of the largest known beetles.



Toxorhynchites rajah

Toxorhynchites rajah

Toxorhynchites rajah is a species of mosquito in the genus Toxorhynchites. It is endemic to Sabah, Malaysian Borneo. In its larval stage, T. rajah is found exclusively in the pitchers of Nepenthes rajah (hence the name), a species of pitcher...



Tramea basilaris

Tramea basilaris

Tramea basilaris, also known as the keyhole glider, red marsh trotter, or wheeling glider, is a species of dragonfly in the family Libellulidae. It is found throughout most of Africa, Arabia and in Asia. Capable of very long migration and nomadic...



Triatoma infestans

Triatoma infestans

Triatoma infestans, commonly called winchuka or vinchuca in Argentina, Bolivia, Uruguay and Chile, barbeiro in Brazil, chipo in Venezuela and also known as "kissing bug" or "barber bug" in English, is a blood-sucking bug (like virtually all the...



Triatoma protracta

Triatoma protracta

Triatoma protracta is a species of any mid sized insect in the family Reduviidae. It is known commonly as the western bloodsucking conenose. It is distributed in the western United States and Mexico. This species and other "kissing bugs" are...



Trichius fasciatus

Trichius fasciatus

Trichius fasciatus, the Eurasian bee beetle, is a beetle species belonging to the family Scarabaeidae, subfamily Cetoniinae.



Troides aeacus

Troides aeacus

Troides aeacus, the golden birdwing, is a large tropical butterfly belonging to the swallowtail family, Papilionidae.



Troides minos

Troides minos

Troides minos, the southern birdwing, also called Sahyadri birdwing, is a large and striking swallowtail butterfly endemic to South India. With a wingspan of 140–190 mm, it is the second largest butterfly of India. It is listed as Least Concern...



Tropidoderus childrenii

Tropidoderus childrenii

Tropidoderus childrenii, the Children's stick insect, was first described in 1833 by Gray.



Tunga penetrans

Tunga penetrans

Tunga penetrans is a species of flea also known as the jigger, jigger flea, chigoe, chigo, chigoe flea, chigo flea, nigua, niknik, sand flea, or burrowing flea. It is a parasitic insect found in most tropical and sub-tropical climates. In its...



Turnip moth

Agrotis segetum

Agrotis segetum, sometimes known as the turnip moth, is a moth of the family Noctuidae. The species was first described by Michael Denis and Ignaz Schiffermüller in 1775. It is a common European species and it is found in Africa and across...



Two-spotted bumble bee

Bombus bimaculatus

The two-spotted bumble bee (Bombus bimaculatus) is a species of social bumble bee found in the eastern half of the United States and the adjacent south-eastern part of Canada. In older literature this bee is often referred to as Bremus...



Tyta

Tyta luctuosa

Tyta luctuosa is a noctuid ("owlet") moth. Its common names include four-spotted moth and field bindweed moth. It is the only member of the genus Tyta, which belongs to the Metoponiinae subfamily. The species was first described by Michael Denis...



Udea ferrugalis

Udea ferrugalis

Udea ferrugalis, the rusty dot pearl, is a moth of the family Crambidae. The species was first described by Jacob Hübner in 1796.



Urania leilus

Urania leilus

Urania leilus, the green-banded urania, is a day-flying moth of the family Uraniidae. The species was first described by Carl Linnaeus in his 1758 10th edition of Systema Naturae. It is found in tropical South America east of the Andes,...



Utetheisa pulchelloides

Utetheisa pulchelloides

Utetheisa pulchelloides, the heliotrope moth, is a moth of the family Erebidae. It is found in the Indo-Australian region including Borneo, Hong Kong, New Zealand, Papua, Seychelles, most of Australia, Tenerife and La Línea de la Concepción. The...



V-pug

Chloroclystis v-ata

The v-pug (Chloroclystis v-ata) is a moth of the family Geometridae. It is found throughout the Palearctic region, the Near East and North Africa. It is well distributed in the British Isles except for the north of Scotland. The species was first...



Vagrant darter

Sympetrum vulgatum

The vagrant darter (Sympetrum vulgatum) is a European dragonfly. The species takes its English name from its habit of occasionally appearing as a rare vagrant north of its normal range. However, it is likely to be under-recorded because it is...



Valley elderberry longhorn beetle

Desmocerus californicus dimorphus

The valley elderberry longhorn beetle (Desmocerus californicus dimorphus) is a subspecies of longhorn beetle native to the riparian forests of the Central Valley of California from Redding to Bakersfield. It is listed as a federally threatened...



Vanessa atalanta

Vanessa atalanta

Vanessa atalanta, the red admiral or, previously, the red admirable, is a well-characterized, medium-sized butterfly with black wings, red bands, and white spots. It has a wingspan of about 2 inches (5 cm). It was first described by Carl Linnaeus...



Vanessa cardui

Vanessa cardui

Vanessa cardui is the most widespread of all butterfly species. It is commonly called the painted lady, or formerly in North America the cosmopolitan.



Vanessa indica

Vanessa indica

Vanessa indica, called the Indian red admiral or the Asian admiral in the United States, is a butterfly found in the higher altitude regions of India, primarily the Himalayas and the Nilgiri Hills. It is also found in Sri Lanka and Myanmar,...



Vanessa vulcania

Vanessa vulcania

Vanessa vulcania, the Canary red admiral, is a butterfly of the family Nymphalidae. It is found on the Canary Islands (except Lanzarote) and Madeira. Previously, it was considered a subspecies of Vanessa indica, but has been raised to species...



Varied carpet beetle

Anthrenus verbasci

The varied carpet beetle (Anthrenus verbasci) is a 3 mm-long beetle belonging to the family Dermestidae. They are a common species, often considered a pest of domestic houses and, particularly, natural history museums, where the larvae may damage...



Variegated meadowhawk

Sympetrum corruptum

The variegated meadowhawk (Sympetrum corruptum) is a dragonfly of the family Libellulidae, native to North America.



Velleius dilatatus

Velleius dilatatus

Velleius dilatatus, the hornet rove beetle, is a species of rove beetle belonging to the family Staphylinidae. This beetle is commensal with the European hornet (Vespa crabro), living in its nests.



Vespa simillima

Vespa simillima

Vespa simillima, the yellow hornet, including the color form known as the Japanese hornet or Japanese yellow hornet (キイロスズメバチ), is a common hornet species in the Eastern Hemisphere. The typical mainland color form (Japanese:ケブカスズメバチ, Korean:...



Vespula flaviceps

Vespula flaviceps

Vespula flaviceps is a species of social wasp in the genus Vespula. It found in Eastern Asia and Japan. Studies have suggested that the queens of this species may mate with more males and use sperm more evenly. The reason for this is not yet well...



Vespula germanica

Vespula germanica

Vespula germanica, the European wasp, German wasp, or German yellowjacket, is a species of wasp found in much of the Northern Hemisphere, native to Europe, Northern Africa, and temperate Asia. It has spread and become well-established in many...



Vespula vulgaris

Vespula vulgaris

Vespula vulgaris, known as the Common wasp, is a species found in regions that include the United Kingdom, Ireland, Germany, India, China, New Zealand and Australia. It is sometimes known in English as the European wasp, but the same name is used...



Viburnum leaf beetle

Pyrrhalta viburni

Pyrrhalta viburni is a species of leaf beetle native to Europe and Asia, commonly known as the viburnum leaf beetle. It was first detected in North America in 1947 in Ontario, Canada. However, specimens had been collected in Annapolis Royal, Nova...



Viceroy

Limenitis archippus

The viceroy (Limenitis archippus) is a North American butterfly. It was long thought to be a Batesian mimic of the monarch butterfly, but since the viceroy is also distasteful to predators, it is now considered a Müllerian mimic instead. The...



Vindula erota

Vindula erota

Vindula erota, the common cruiser, is a species of nymphalid butterfly found in forested areas of tropical South Asia and Southeast Asia.



Vine weevil

Otiorhynchus sulcatus

The black vine weevil (Otiorhynchus sulcatus) is an insect native to Europe but common in North America as well. It is a pest of many garden plants.



Violet click beetle

Gambrinus violaceus

The violet click beetle (Gambrinus violaceus, formerly Limoniscus violaceus) is a black beetle, 12 mm (0.5 in) long, with a faint blue/violet reflection. It gets its name from the family habit of springing upwards with an audible click if it...



Volucella pellucens

Volucella pellucens

Volucella pellucens, the pellucid fly, is a hoverfly.



West Coast lady

Vanessa annabella

The West Coast lady (Vanessa annabella) is one of three North American species of brush-footed butterflies known colloquially as the "painted ladies". V. annabella occurs throughout much of the western US and southwestern Canada. The other two...



Western conifer seed bug

Leptoglossus occidentalis

The western conifer seed bug (Leptoglossus occidentalis), often mistakenly referred to as stink bug, sometimes abbreviated as WCSB, is a species of true bug (Hemiptera) in the family Coreidae. It is native to North America west of the Rocky...



Western corn rootworm

Diabrotica virgifera virgifera

The Western corn rootworm, Diabrotica virgifera virgifera, is one of the most devastating corn rootworm species in North America, especially in the midwestern corn-growing areas such as Iowa. A related species, the Northern corn rootworm, D....



Western pygmy blue

Brephidium exilis

The western pygmy blue (Brephidium exilis or Brephidium exile) is one of the smallest butterflies in the world and the smallest in North America. It has reached Hawaii, as well as the Persian Gulf, including eastern Saudi Arabia, Bahrain and the...



Wheat weevil

Sitophilus granarius

The wheat weevil (Sitophilus granarius), also known as the grain weevil or granary weevil, is an insect that feeds on cereal grains, and is a common pest in many places. It can cause significant damage to harvested stored grains and may...



White-faced darter

Leucorrhinia dubia

The white-faced darter or small whiteface (Leucorrhinia dubia) is a dragonfly belonging to the genus Leucorrhinia in the family Libellulidae, characterised by red and black markings and a distinctive white patch on the head. It is found in...



White-letter hairstreak

Satyrium w-album

The white-letter hairstreak (*Satyrion w-album*) is a butterfly in the family *Lycaenidae*.



Widow skimmer

Libellula luctuosa

The widow skimmer (*Libellula luctuosa*) is one of the group of dragonflies known as king skimmers. The nymphs live in the water, molting and growing until they are ready to emerge from the water and then molting a final time to reveal their wings.



Willow beauty

Peribatodes rhomboidaria

The willow beauty (*Peribatodes rhomboidaria*) is a moth of the family *Geometridae*. It is a common species of Europe and adjacent regions (Near East and the Maghreb). While it is found widely throughout Scandinavian countries, which have a maritime...



Winter moth

Operophtera brumata

The winter moth (*Operophtera brumata*) is a moth of the family *Geometridae*. It is an abundant species in Europe and the Near East and a famous study organism for evaluating insect population dynamics. It is one of very few lepidopterans of...



Witch-hazel cone gall aphid

Hormaphis hamamelidis

The witch-hazel cone gall aphid (*Hormaphis hamamelidis*) is a minuscule insect, a member of the aphid superfamily, whose presence on a witch-hazel (*Hamamelis virginiana*) plant is easily recognizable by a conical gall structure. The gall is green...



Xylocopa aerata

Xylocopa aerata

Xylocopa aerata, the golden-green carpenter bee or green carpenter bee, is one of two species of carpenter bee found only in the conservation areas around Sydney, and in the Great Dividing Range in New South Wales in Australia. Its only other...



Xylocopa bombylans

Xylocopa bombylans

Xylocopa bombylans, the peacock carpenter bee, is a species of carpenter bee found in Australia. It gets its common name by its habit of burrowing into wood. It was originally described by Danish naturalist Johan Christian Fabricius in 1775. Its...



Xylophanes tersa

Xylophanes tersa

Xylophanes tersa, the tersa sphinx, is a moth of the family Sphingidae. The species was first described by Carl Linnaeus in 1771. It is found from the United States (Massachusetts south to southern Florida, west to Nebraska, New Mexico and...



Xylota segnis

Xylota segnis

Xylota segnis, The Brown-toed Forest Fly, is a common species of hoverfly.



Yellow admiral

Vanessa itea

The yellow admiral or Australian admiral (*Vanessa itea*) is a butterfly native to Australia, New Zealand, Lord Howe Island, and Norfolk Islands. The Māori name is kahukōwhai, which means "yellow cloak". The yellow admiral is a member of the family...



Yellow crazy ant

Anoplolepis gracilipes

The yellow crazy ant (*Anoplolepis gracilipes*), also known as the long-legged ant or Maldive ant, is a species of ant, thought to be native to West Africa or Asia. They have been accidentally introduced to numerous places in the world's...



Yellow meadow ant

Lasius flavus

The yellow meadow ant (*Lasius flavus*), also known as the yellow hill ant, is a species of ant occurring in Europe (where it is one of the most common ants), Asia, and North Africa. Populations in North America are now considered a different...



Yellow-tail

Euproctis similis

The yellow-tail, goldtail moth or swan moth (*Sphrageidus similis*) is a moth of the family Erebidæ. The species was first described by Johann Kaspar Füssli in 1775, and has commonly been placed within the related genus *Euproctis*. It is...



Yellow-winged darter

Sympetrum flaveolum

The yellow-winged darter (*Sympetrum flaveolum*) is a dragonfly found in Europe and mid and northern China. Breeding is confined to stagnant water, usually in peat bogs. Although not resident in the United Kingdom it occasionally migrates there in...



Yponomeuta malinellus

Yponomeuta malinellus

Yponomeuta malinellus, the apple ermine, is a moth of the family Yponomeutidae. It is native to Europe and Asia, and it has spread to North America. The wingspan...



Zabulon skipper

Lon zabulon

The Zabulon skipper (Lon zabulon) (sometimes called the southern dimorphic skipper) is a North American butterfly first described by the French naturalists Jean Baptiste Boisduval and John Eatton Le Conte from the state of Georgia, United States.



Zebra caterpillar

Melanchra picta

The zebra caterpillar is the larva of an American noctuid moth (Melanchra picta) that feeds on cabbages, beets and other cultivated plants. The head, thorax, and forewings of adults are chestnut- or reddish-brown, usually with purplish brown...



Zelotypia

Zelotypia stacyi

Zelotypia is a monotypic moth genus of the family Hepialidae. The only described species is Z. stacyi, the bentwing ghost moth, which is only found in Queensland and New South Wales, Australia. This is a very large species with a wingspan of up...



Zerene cesonia

Zerene cesonia

Zerene cesonia, the southern dogface, is a North and South American butterfly in the family Pieridae, subfamily Coliadinae (until recently the species was sometimes placed in the related genus Colias instead of Zerene).



Zerene eurydice

Zerene eurydice

Zerene eurydice, the California dogface butterfly, belongs to the family Pieridae and is a sister genus to Colias. The Zerene eurydice and the Colias both share the "characteristic of having yellow-orange and black wing coordination."...



Zerynthia polyxena

Zerynthia polyxena

Zerynthia polyxena, the southern festoon, is a butterfly belonging to the butterfly family Papilionidae.



Zizeeria karsandra

Zizeeria karsandra

Zizeeria karsandra, the dark grass blue, is a small butterfly first described by Frederic Moore in 1865. It is found from the southern Mediterranean, in a broad band to India, Sri Lanka, the Andaman and Nicobar islands, Myanmar, Thailand, Malaysia,...



Zizeeria knysna

Zizeeria knysna

Zizeeria knysna, the dark grass blue or African grass blue, is a species of blue butterfly (Lycaenidae) found in Africa, on Cyprus and the Iberian Peninsula. Sometimes mistakenly called Zizeeria lysimon (Hübner), that species name is invalid as it is...



Zizina labradus

Zizina labradus

Zizina labradus, the common grass blue, grass blue, or clover blue, is a small Australian butterfly of the family Lycaenidae.



Zizina otis

Zizina otis

Zizina otis, the lesser grass blue, is a species of blue (Lycaenidae) butterfly found in south and southeast Asia. The species was first described by Johan Christian Fabricius in 1787. The lesser grass blue is often misidentified as Zizina labradus,...



Zizula hylax

Zizula hylax

Zizula hylax, 'the Tiny grass blue' is a species of blue butterfly.



Zophobas morio

Zophobas morio

Zophobas morio is a species of darkling beetle, whose larvae are known by the common name superworm, kingworm, morio worm or simply Zophobas. Superworms are common in the reptile pet industry as food, along with giant mealworms, which are Tenebrio...



Zygaena carniolica

Zygaena carniolica

Zygaena carniolica, sometimes described as the crepuscular burnet or eastern burnet, is a member of the family Zygaenidae.



Zygaena lonicerae

Zygaena lonicerae

Zygaena lonicerae, the narrow-bordered five-spot burnet, is a moth of the family Zygaenidae. The species was first described by Theodor Gottlieb von Scheven in 1777.