

? Nycticebus linglom
? Nycticebus linglom
Nycticebus linglom

? Nycticebus linglom is a fossil strepsirrhine primate from the Miocene of Thailand. Known only from a single tooth, an upper third molar, it is thought to be related to the living slow lorises, but the material is not sufficient to assign the...

Aardvark
Aardvark
Orycteropus afer

The aardvark (*Orycteropus afer*) is a medium-sized, burrowing, nocturnal mammal native to Africa. Unlike most other insectivores, it has a long pig-like snout, which is used to sniff out food. It roams over most of the southern two-thirds of the...

Aardwolf
Aardwolf
Proteles cristata

The aardwolf (*Proteles cristata*) is an insectivorous mammal, native to Africa. It belongs to the same family as the hyena; however, unlike many of its carnivorous relatives, the aardwolf does not hunt large animals. Its name means "earth-wolf" in...

Aba roundleaf bat
Aba roundleaf bat
Hipposideros abae

The Aba roundleaf bat (*Hipposideros abae*), also known as the Aba leaf-nosed bat is a species of bat in the family Hipposideridae. It is found in west Africa along the southern coast from Nigeria to Senegal. Populations have also been noted in...

Abbott's duiker
Abbott's duiker
Cephalophus spadix

The Abbott's duiker (*Cephalophus spadix*), also known as minde in Swahili, is a large, forest-dwelling duiker (small antelope) found only in a few scattered enclaves in Tanzania. It may be a subspecies of the yellow-backed duiker. It is very...

Aberdare mole shrew
Aberdare mole shrew
Surdisorex norae

The Aberdare mole shrew (*Surdisorex norae*) is a species of mammal in the family Soricidae endemic to the Aberdare Mountains in Kenya. Its natural habitat is tropical high-altitude bamboo and grassland.

Aberdare Mountains African mole-rat
Aberdare Mountains African mole-rat
Tachyoryctes audax

The Aberdare Mountains African mole-rat (*Tachyoryctes audax*) is a species of rodent in the family Spalacidae endemic to Kenya. Its natural habitats are subtropical or tropical moist montane forests and subtropical or tropical high-altitude...

Abert's Squirrel
Abert's Squirrel
Sciurus aberti

Abert's squirrel (*Sciurus aberti*) is a tree squirrel native to North America. It lives in pine forests and favors the seeds and cones of the Mexican pinyon and the ponderosa pine. It is named in honor of the American naturalist John James Abert.

Abo bat

Abo bat

Glauconycteris poensis

The Abo bat (*Glauconycteris poensis*) is a species of vesper bat in the family Vespertilionidae. It is found in several countries in West Africa and Central Africa. It is found in these habitats: subtropical or tropical dry forest and subtropical...

Abrothrix andinus

Abrothrix andinus

Abrothrix andina

Abrothrix andina, also known as the Andean Altiplano mouse or Andean akodont, is a species of rodent in the genus *Abrothrix* of family Cricetidae. It is found in the Altiplano habitat of the Andes from central Peru through Bolivia, south to...

Abrothrix herskovitzi

Abrothrix herskovitzi

Abrothrix herskovitzi

Abrothrix herskovitzi, also known as Herskovitz's grass mouse or Herskovitz's akodont, is a species of rodent in the genus *Abrothrix* of family Cricetidae. It is found only on some remote islands in southern Chile. It is named after American...

Abrothrix illuteus

Abrothrix illuteus

Abrothrix illuteus

Abrothrix illuteus, also known as the gray akodont, gray grass mouse, or gray soft-haired mouse, is a species of small rodent in the genus *Abrothrix* of family Cricetidae. It is found only in northwestern Argentina.

Abrothrix jelskii

Abrothrix jelskii

Abrothrix jelskii

Abrothrix jelskii, also known as Jelski's Altiplano mouse, Jelski's grass mouse, or the ornate akodont, is a species of rodent in the genus *Abrothrix* of family Cricetidae. It is found in the altiplano habitat of the Andes from central Peru...

Abrothrix lanosus

Abrothrix lanosus

Abrothrix lanosus

Abrothrix lanosus, also known as the woolly grass mouse or woolly akodont, is a species of rodent in the family Cricetidae. It is found in southern Argentina and Chile. It was previously classified in the genus *Akodon* rather than *Abrothrix*.

Abrothrix longipilis

Abrothrix longipilis

Abrothrix longipilis

Abrothrix longipilis, also known as the long-haired grass mouse or long-haired akodont, is a species of rodent in the family Cricetidae. It is found in central and southern Argentina and Chile. The southern Chilean *Abrothrix sanborni* may not be...

Abrothrix olivaceus

Abrothrix olivaceus

Abrothrix olivacea

Abrothrix olivacea, also known as the olive grass mouse or olive akodont, is a

species of rodent in the genus *Abrothrix* of family Cricetidae. It is found from northern Chile into southern Chile and Argentina, including the islands of Tierra del...

Abrothrix olivaceus markhami
Abrothrix olivaceus markhami
Abrothrix olivaceus markhami

Abrothrix olivaceus markhami, also known as the Wellington Akodont or Markham's Grass Mouse, is a subspecies of the South American rodent *Abrothrix olivaceus*. It occurs on Wellington Island and the nearby Southern Patagonian Ice Field in southern...

Abrothrix sanborni
Abrothrix sanborni
Abrothrix sanborni

Abrothrix sanborni, also known as Sanborn's grass mouse or Sanborn's akodont, is a species of rodent in the genus *Abrothrix* of family Cricetidae. It is found in southern Argentina and Chile, but may not be distinct from *A. longipilis*.

Abyssinian Genet
Abyssinian Genet
Genetta abyssinica

The Abyssinian genet (*Genetta abyssinica*) is an African carnivoran listed as Data Deficient on the IUCN Red List. It is one of the least-known genet species.

Abyssinian grass rat
Abyssinian grass rat
Arvicanthis abyssinicus

The Abyssinian grass rat (*Arvicanthis abyssinicus*) is a species of rodent in the family Muridae. It is found in Ethiopia. Its natural habitats are subtropical or tropical high-altitude grassland, arable land, and pastureland.

Abyssinian hare
Abyssinian hare
Lepus habessinicus

The Abyssinian hare (*Lepus habessinicus*) is a species of mammal in the family Leporidae. It is almost entirely restricted to the nations of the Horn of Africa, though it extends marginally into eastern Sudan and may also occur in far northern...

Acacia rat
Acacia rat
Thallomys paedulcus

The acacia rat (*Thallomys paedulcus*) is a species of rodent in the family Muridae. It is found in Botswana, Democratic Republic of the Congo, Ethiopia, Kenya, Malawi, Mozambique, Namibia, Somalia, South Africa, Eswatini, Tanzania, Zambia, and...

Aceh rat
Aceh rat
Rattus blangorum

The Aceh rat (*Rattus blangorum*) is a species of rodent from the family Muridae. The Aceh rat is listed as Data Deficient on the IUCN Red List because only two specimens have ever been recorded, and little is known about the species. The species...

Aceramarca gracile opossum
Aceramarca gracile opossum
Gracilinanus aceramarcae

The *Aceramarca gracile* opossum or Bolivian gracile opossum (*Gracilinanus aceramarcae*) is a species of opossum. It is native to Bolivia and Peru, where it occurs in tropical elfin forest habitat. This opossum is mostly arboreal, but it may forage...

Acrobatic cavy
Acrobatic cavy
Kerodon acrobata

The acrobatic cavy (*Kerodon acrobata*) also known as the Acrobatic Moco and Climbing Cavy is a cavy species native to Brazil in the Amazon rainforest. It is found from Goiás state to Tocantins state, west of the Espigão Mestre, Serra Geral de...

Acuminate horseshoe bat
Acuminate horseshoe bat
Rhinolophus acuminatus

The acuminate horseshoe bat (*Rhinolophus acuminatus*) is a species of bat in the family Rhinolophidae. It is found in Brunei, Cambodia, Indonesia, Laos, Malaysia, the Philippines, and Thailand. It lives in forests and urban areas.

Adam's horseshoe bat
Adam's horseshoe bat
Rhinolophus adami

Adam's horseshoe bat (*Rhinolophus adami*) is a species of bat in the family Rhinolophidae. It is endemic to Republic of the Congo. Its natural habitats include caves.

Addax
Addax
Addax nasomaculatus

The addax (*Addax nasomaculatus*) is an antelope native to the Sahara Desert. The only member of the genus Addax, it was first described scientifically by Henri de Blainville in 1816. Addax are well-adapted to exist in their desert habitat, as they...

Aders's duiker
Aders's duiker
Cephalophus adersi

The Aders's duiker (*Cephalophus adersi*), also known as nunga in Swahili, kunga marara in Kipokomo and harake in Giriama, is a small, forest-dwelling duiker found only in Zanzibar and Kenya. It may be a subspecies of the red, Harvey's, or...

Admiralty flying fox
Admiralty flying fox
Pteropus admiralitatum

The Admiralty flying fox (*Pteropus admiralitatum*) is a species of fruit bat in the family Pteropodidae, the megabats. It is found in Papua New Guinea and the Solomon Islands.

Admiralty Island cuscus
Admiralty Island cuscus
Spilocuscus kraemeri

The Admiralty Island cuscus or Manus Island spotted cuscus (*Spilocuscus kraemeri*) is a species of marsupial in the family Phalangeridae. It is endemic to the Admiralty Islands of Papua New Guinea. It is the smallest member of the genus...

AEEL's sportive lemur
AEEL's sportive lemur
Lepilemur aeeclis

The Antafia sportive lemur, or red-shouldered sportive lemur (*Lepilemur aeeclis*) is a sportive lemur endemic to Madagascar. It has a total length of about 52 to 59 cm (20 to 23 in), of which 24–26 cm (9.4–10.2 in) are tail. The AEECL's sportive...
Aegialomys galapagoensis
Aegialomys galapagoensis
Aegialomys galapagoensis

The Galápagos rice rat (*Aegialomys galapagoensis*), also known as the Galápagos oryzomys, is a species of rodent that is endemic to the Galápagos Islands.
Aegialomys xantheolus
Aegialomys xantheolus
Aegialomys xantheolus

Aegialomys xantheolus, also known as the yellowish oryzomys or yellowish rice rat, is a species of rodent in the family Cricetidae. It belongs to the genus *Aegialomys* in tribe Oryzomyini, which was not recognized as distinct from *Oryzomys* until...
Aellen's pipistrelle
Aellen's pipistrelle
Pipistrellus inexpectatus

Aellen's pipistrelle (*Pipistrellus inexpectatus*) is a species of vesper bat. It can be found in possibly Benin, Cameroon, Ghana, Nigeria, and Sierra Leone. It is found in dry and moist savanna.
Aellen's roundleaf bat
Aellen's roundleaf bat
Hipposideros marisae

Aellen's roundleaf bat (*Hipposideros marisae*) is a species of bat in the family Hipposideridae. It is found in Ivory Coast, Guinea, and Liberia. Its natural habitats are subtropical or tropical dry forest, subtropical or tropical moist lowland...
Afghan flying squirrel
Afghan flying squirrel
Eoglaucomys fimbriatus baberi

The Afghan flying squirrel (*Eoglaucomys fimbriatus baberi*) is a subspecies of rodent in the family Sciuridae. It is endemic to Afghanistan.
Afghan hedgehog
Afghan hedgehog
Hemiechinus auritus megalotis

The Afghan hedgehog (*Hemiechinus auritus megalotis*) is a subspecies of the long-eared hedgehog native to Afghanistan and western Pakistan. It is larger and reddish in colour, but otherwise similar in appearance to the long-eared hedgehog.
Afghan pika
Afghan pika
Ochotona rufescens

The Afghan pika (*Ochotona rufescens*) is a species of small mammal in the pika family, Ochotonidae. It is found in Afghanistan, Iran, Pakistan and Turkmenistan and the IUCN lists it as being of "least concern".
Afghan vole
Afghan vole
Blanfordimys afghanus

The Afghan vole (*Microtus afghanus*) is a species of rodent in the family Cricetidae. It is found in south-central Asia.
African black shrew
African black shrew

Crocidura nigrofusca

The African black shrew (*Crocidura nigrofusca*) is a species of shrew. It is native to Africa, where it is widespread. Other common names include tenebrous shrew. This shrew can be found in several types of lower-elevation moist tropical forest...

African brush-tailed porcupine

African brush-tailed porcupine

Atherurus africanus

The African brush-tailed porcupine (*Atherurus africanus*) is a species of rat-like Old World porcupine, indigenous to a broad belt of Africa ranging from Guinea on the west coast to Kenya on the east. This is a common species with a very wide...

African Buffalo

African Buffalo

Syncerus caffer

African buffalo (*Syncerus caffer*) are large, heavy cow-like animals native to Africa. They are widely regarded as among the most dangerous animals on the African continent, and according to some estimates they gore, trample, and kill over 200...

African Bush Elephant

African Bush Elephant

Loxodonta africana

The African bush elephant (*Loxodonta africana*) is one of two living African elephant species. It is the largest living terrestrial animal and is distributed across 37 African countries. Since 2021, it has been listed as Endangered on the IUCN Red...

African Civet

African Civet

Civettictis civetta

The African civet (*Civettictis civetta*) is a large type of civet that lives throughout sub-Saharan Africa. It is the last surviving member of its genetic group and is regarded as the largest civet-like species in the African continent.

African...

African Clawless Otter

African Clawless Otter

Aonyx capensis

The African clawless otter (*Aonyx capensis*), also known as the Cape clawless otter or groot otter, is the second-largest freshwater otter species. It inhabits permanent water bodies in savannah and lowland forest areas through most of...

African dusky shrew

African dusky shrew

Crocidura caliginea

The African dusky shrew or African foggy shrew (*Crocidura caliginea*) is a species of shrew. It is native to the Democratic Republic of the Congo, where it lives in forests.

African forest buffalo

African forest buffalo

Syncerus caffer nanus

The African forest buffalo (*Syncerus caffer nanus*), also known as the dwarf buffalo or the Congo buffalo, is the smallest subspecies of the African buffalo. It is related to the Cape buffalo (*Syncerus caffer caffer*), the Sudan buffalo (*Syncerus*...

African Forest Elephant

African Forest Elephant

Loxodonta cyclotis

The African forest elephant (*Loxodonta cyclotis*) is one of the two living African elephant species. It is the smallest of the three living elephant species. The first scientific description of the species was published in 1900. During the 20th...

African giant free-tailed bat

African giant free-tailed bat

Tadarida ventralis

The African giant free-tailed bat (*Tadarida ventralis*), or African free-tailed bat is a species of bat in the family Molossidae. It is found in Democratic Republic of the Congo, Eritrea, Ethiopia, Kenya, Malawi, Mozambique, South Sudan,...

African Giant Shrew

African Giant Shrew

Crocidura olivieri

The African giant shrew (*Crocidura olivieri*) is a species of large white-toothed shrew. It is native to Africa, where it has a widespread distribution and occurs in many types of habitat. In the Nile Valley it is found near human habitation,...

African Golden Cat

African Golden Cat

Caracal aurata

The African golden cat (*Caracal aurata*) remains little known by science, though within Africa there is much tribal superstition about it. It is also called the 'leopard's brother' because, according to local people, it follows the leopard. It is...

African Golden Wolf

African Golden Wolf

Canis anthus

The African golden wolf (*Canis lupaster*) is a canine that plays a prominent role in some African cultures. It was previously classified as an African variant of the Golden jackal. In 2015, a series of analyses on the species' mitochondrial DNA...

African grass rat

African grass rat

Arvicanthis niloticus

The African grass rat (*Arvicanthis niloticus*) is a species of rodent in the family Murinae.

African groove-toothed rat

African groove-toothed rat

Mylomys dybowski

The African groove-toothed rat or mill rat (*Mylomys dybowski*) is a species of rodent in the family Muridae found in Angola, Cameroon, the Central African Republic, the Republic of the Congo, the Democratic Republic of the Congo, Ivory Coast,...

African Leopard

African Leopard

Panthera pardus pardus

The African leopard (*Panthera pardus pardus*) is the nominate subspecies of the leopard, native to many countries in Africa. It is widely distributed in most of sub-Saharan Africa, but the historical range has been fragmented in the course of...

African long-fingered bat

African long-fingered bat

Miniopterus africanus

The African long-fingered bat (*Miniopterus africanus*) is a species of vesper bat in the family Vespertilionidae. It is found only in Kenya. It is found in subtropical or tropical moist montane forests. This species is often considered a synonym...

African Manatee

African Manatee

Trichechus senegalensis

The African manatee (*Trichechus senegalensis*), also known as the West African manatee, is a species of manatee that inhabits much of the western region of Africa – from Senegal to Angola. It is the only manatee species to be found in the Old...

African marsh rat

African marsh rat

Dasymys incomtus

The African marsh rat or common dasymys (*Dasymys incomtus*) is a species of rodent in the family Muridae. It is found in Angola, Democratic Republic of the Congo, Ethiopia, Kenya, Malawi, South Africa, South Sudan, Tanzania, Uganda, Zambia, and...

African Palm Civet

African Palm Civet

Nandinia binotata

The African palm civet (*Nandinia binotata*), also known as the two-spotted palm civet, is a small feliform mammal widely distributed in sub-Saharan Africa. It is listed as Least Concern on the IUCN Red List.

African Pygmy Mouse

African Pygmy Mouse

Mus minutoides

The African pygmy mouse (*Mus minutoides*) is one of the smallest rodents. It is a member of the enormous superfamily Muroidea, which includes over 1000 different species and is kept as a pet in many parts of the world.

African pygmy squirrel

African pygmy squirrel

Myosciurus pumilio

The African pygmy squirrel (*Myosciurus pumilio*) is a species of rodent in the family Sciuridae. It is monotypic within the genus *Myosciurus*. It is found in tropical rainforests in Cameroon, the Republic of the Congo, Equatorial Guinea, and...

African Savanna Hare

African Savanna Hare

Lepus victoriae

The African savanna hare (*Lepus victoriae*) is a medium-sized mammal found only in Africa. It looks very similar to the Cape hare in appearance but can be told apart by its distinctively grooved incisors.

African sheath-tailed bat

African sheath-tailed bat

Coleura afra

The African sheath-tailed bat (*Coleura afra*) is a species of sac-winged bat in the family Emballonuridae found in Angola, Benin, the Central African Republic, the Republic of the Congo, the Democratic Republic of the Congo, Ivory Coast,...

African smoky mouse

African smoky mouse

Heimyscus fumosus

The African smoky mouse or smokey heimyscus (*Heimyscus fumosus*) is a species of rodent in the family Muridae. It is the only species in the genus *Heimyscus*. It is native to Central Africa, where it occurs in Cameroon, Central African Republic, ...
African Striped Weasel
African Striped Weasel
Poecilogale albinucha

The African striped weasel (*Poecilogale albinucha*) is one of the smallest carnivorous mammals in Africa. It is a nocturnal hunter and finds its prey primarily by scent.
African trident bat
African trident bat
Triaenops afer

The African trident bat (*Triaenops afer*) is a species of bat found in Africa.
African wading rat
African wading rat
Colomys goslingi

The African wading rat or African water rat (*Colomys goslingi*) is a species of rodent in the family Muridae. It is native to Africa, where it occurs in Angola, Burundi, Cameroon, Republic of the Congo, Democratic Republic of the Congo, ...
African Wild Ass
African Wild Ass
Equus africanus

The African wild ass (*Equus africanus*) is a wild member of the horse family, Equidae. This species is thought to be the ancestor of the domestic donkey, which is sometimes placed within the same species.
African Wild Dog
African Wild Dog
Lycaon pictus

The African wild dog (*Lycaon pictus*) is the largest wild canine in Africa, and the only extant member of the genus *Lycaon*; it is distinguished from *Canis* by dentition highly specialized for a hypercarnivorous diet, and by a lack of dewclaws....
African Wildcat
African Wildcat
Felis lybica

The African wildcat (*Felis lybica*) is a small wildcat species. In Cyprus, an African wildcat was found in a burial site next to a human skeleton in the Pre-Pottery Neolithic B settlement Shillourokambos. The graves are estimated to have been...
African yellow bat
African yellow bat
Scotophilus dinganii

The African yellow bat (*Scotophilus dinganii*) is a species of bat in the family Vespertilionidae, the vesper bats. Other common names include African yellow house bat, yellow-bellied house bat, and Dingan's Bat. It is one of fifteen species in...
Afroalpine vlei rat
Afroalpine vlei rat
Otomys orestes

The afroalpine vlei rat (*Otomys orestes*) is a species of rodent in the family Muridae. It is found in the high moorlands of Kenya.

Agag gerbil
Agag gerbil
Gerbillus agag

The Agag gerbil (*Gerbillus agag*) is distributed mainly in southern Mauritania to northern Nigeria and Sudan. IUCN lists the junior synonyms *Gerbillus cosensi* and *G. dalloni* as critically endangered.

Agile antechinus
Agile antechinus
Antechinus agilis

The agile antechinus (*Antechinus agilis*) is a species of small carnivorous marsupial of the family Dasyuridae. It is found in Australia.

Agile Gibbon
Agile Gibbon
Hylobates agilis

The Agile gibbon (*Hylobates agilis*), a is an endangered primate found in Southeast Asia. Like all gibbons, they move by swinging from branch to branch for distances up to 15 m (50 ft) and can also make leaps up to 8 m (26 ft), and walk bipedally...

Agile gracile opossum
Agile gracile opossum
Gracilinanus agilis

The agile gracile opossum (*Gracilinanus agilis*), is an opossum species from South America. It is found in Argentina, Bolivia, Brazil, Colombia, Peru, Paraguay, and Uruguay. It is nocturnal, arboreal and frequents the forest understory, where they...

Agile kangaroo rat
Agile kangaroo rat
Dipodomys agilis

The agile kangaroo rat (*Dipodomys agilis*) is a species of rodent in the family Heteromyidae. It is endemic to southern California in the United States. Relatively little information has been published on the natural history, life history,...

Agile mangabey
Agile mangabey
Cercocebus agilis

The agile mangabey (*Cercocebus agilis*) is an Old World monkey of the white-eyelid mangabey group found in swampy forests of Central Africa in Equatorial Guinea, Cameroon, Gabon, Central African Republic, Republic of Congo, and DR Congo. Until...

Agile Wallaby
Agile Wallaby
Macropus agilis

The Agile wallaby (*Notamacropus agilis*) is a species of wallaby found in northern Australia and New Guinea. It is the most common wallaby in north Australia.

Agricola's gracile opossum
Agricola's gracile opossum
Cryptonanus agricolai

Agricola's gracile opossum (*Cryptonanus agricolai*) is a species of opossum in the family Didelphidae endemic to eastern Brazil. Its habitat is the caatinga and cerrado. While its conservation status has not been determined, expansion of...

Ahmanson's sportive lemur
Ahmanson's sportive lemur
Lepilemur ahmansonorum

Ahmanson's sportive lemur (*Lepilemur ahmansonorum*), or the Tsiombikibo sportive lemur, is a sportive lemur endemic to Madagascar. It is a relatively small sportive lemur with a total length of about 47 to 54 cm (19 to 21 in), of which 23–25 cm...

Ailuropoda baconi

Ailuropoda baconi

Ailuropoda baconi

Ailuropoda baconi is an extinct panda from the Late Pleistocene, 750 thousand years ago, and was preceded by *A. wulingshanensis* and *A. microta* as an ancestor of the giant panda (*A. melanoleuca*). Very little is known about this animal; however,...

Ailuropoda microta

Ailuropoda microta

Ailuropoda microta

Ailuropoda microta is the earliest known ancestor of the giant panda. It measured 1 m (3 ft) in length; the modern giant panda grows to a size in excess of 1.5 m (5 ft). Wear patterns on its teeth suggest it lived on a diet of bamboo, the primary...

Akishima whale

Akishima whale

Eschrichtius akishimaensis

Eschrichtius akishimaensis, described in 2017, is one of two species, with the modern day gray whale, of the genus *Eschrichtius*, and is the first and only fossil species of the genus, dating to around 1.77–1.95 million years ago (mya) in the...

Akodon aerosus

Akodon aerosus

Akodon aerosus

Akodon aerosus, also known as the highland grass mouse or Yungas akodont, is a species of rodent in the family Cricetidae. It is found in the eastern Andes from eastern Ecuador through Peru into central Bolivia.

Akodon affinis

Akodon affinis

Akodon affinis

Akodon affinis, also known as the Colombian grass mouse or Cordillera Occidental akodont, is a species of rodent in the family Cricetidae. It is found only in the Cordillera Occidental of Colombia.

Akodon albiventer

Akodon albiventer

Akodon albiventer

Akodon albiventer, also known as the white-bellied grass mouse or white-bellied akodont, is a species of rodent in the family Cricetidae. It is found in the Andean highlands from southeastern Peru to southwestern Bolivia, northwestern

Argentina,...

Akodon azarae

Akodon azarae

Akodon azarae

Akodon azarae, also known as Azara's akodont or Azara's grass mouse, is a rodent species from South America. It is found from southernmost Brazil through Paraguay and Uruguay into eastern Argentina. It is named after Spanish naturalist Félix de...

Akodon boliviensis

Akodon boliviensis

Akodon boliviensis

Akodon boliviensis, also known as the Bolivian grass mouse or Bolivian akodont, is a species of rodent in the family Cricetidae. It is found in the Andes from southeastern Peru through Bolivia into northwestern Argentina.

Akodon budini

Akodon budini

Akodon budini

Akodon budini, also known as Budin's akodont or Budin's grass mouse, is a species of rodent in the family Cricetidae. It is found in the Andes of northwestern Argentina and adjacent Bolivia. The species is named after Emilio Budin, an Argentine...

Akodon caenosus

Akodon caenosus

Akodon caenosus

Akodon caenosus is a rodent in the genus Akodon found in northwestern Argentina and south-central Bolivia. Since its description in 1918, it has been alternatively classified as a separate species or a subspecies of Akodon lutescens (formerly...

Akodon spegazzinii

Akodon spegazzinii

Akodon spegazzinii

Akodon spegazzinii, also known as Spegazzini's akodont or Spegazzini's grass mouse, is a rodent in the genus Akodon found in northwestern Argentina. It occurs in grassland and forest at 400 to 3,500 m (1,300 to 11,500 ft) above sea level.

After...

Akodon sylvanus

Akodon sylvanus

Akodon sylvanus

Akodon sylvanus, also known as the forest grass mouse or woodland akodont, is a species of rodent in the family Cricetidae. It is found only in a small part of northwestern Argentina.

Alabama beach mouse

Alabama beach mouse

Peromyscus polionotus ammobates

The Alabama beach mouse (Peromyscus polionotus ammobates) is a federally endangered species that lives along the Alabama coast.

Alai mole vole

Alai mole vole

Ellobius alaicus

The Alai mole vole (Ellobius alaicus) is a species of rodent in the family Cricetidae. It is known only from Kyrgyzstan, where it has been found in temperate grassland in the Alai Mountains. Little else is known about the vole. It is threatened...

Alashan ground squirrel

Alashan ground squirrel

Spermophilus alashanicus

The Alashan ground squirrel (Spermophilus alashanicus) is a species of squirrel. It is native to China and Mongolia. This squirrel lives at elevations up to 3200 meters. It lives on the steppe and in foothills and alpine meadows. In China it can...

Alashan wapiti

Alashan wapiti

Cervus canadensis alashanicus

The Alashan wapiti is a subspecies of *Cervus canadensis* (named "elk" or "wapiti" in North America), found in Northern China and Mongolia. It is the smallest subspecies of elk, has the lightest color, and is the least studied, other than the...

Alaska Marmot
Alaska Marmot
Marmota broweri

The Alaska marmot (*Marmota broweri*) is a rodent that likes to live on rocky, mountainous terrain, often near lakes. It was once considered to be the same species as the Hoary marmot but is now known to be unique.

Alaska moose
Alaska moose
Alces alces gigas

The Alaska moose (*Alces alces gigas*), or Alaskan moose in Alaska, or giant moose and Yukon moose in Canada, is a subspecies of moose that ranges from Alaska to western Yukon. The Alaska moose is the largest subspecies of moose. Alaska moose...

Alaska Peninsula brown bear
Alaska Peninsula brown bear
Ursus arctos middendorffi / *horribilis*

The Alaska Peninsula brown bear or "peninsular grizzly" is a colloquial nomenclature for a brown bear that lives in the coastal regions of southern Alaska, although according to other sources, it is a population of the mainland grizzly bear...

Alaska tiny shrew
Alaska tiny shrew
Sorex yukonicus

The Alaska tiny shrew (*Sorex yukonicus*) is a species of shrew. It is endemic to Alaska. This species was formerly included under the description of the Eurasian least shrew (*S. minutissimus*). It was described as a separate species in 1997. It...

Alaskan Hare
Alaskan Hare
Lepus othus

The Alaskan hare (*Lepus othus*) is one of the largest species of hares. It is one of two species of hares native to the state of Alaska; the other is the more widespread Snowshoe hare (*Lepus americanus*).

Alaskan Tundra Wolf
Alaskan Tundra Wolf
Canis lupus tundrarum

The Alaskan tundra wolf (*Canis lupus tundrarum*) is a North American subspecies of Grey wolf. It was named in 1912 by Gerrit Smith Miller Jr., who noted that it closely approaches the Great Plains wolf in skull and tooth morphology.

Alcathoe bat
Alcathoe bat
Myotis alcathoe

The Alcathoe bat (*Myotis alcathoe*) is a European bat in the genus *Myotis*. Known only from Greece and Hungary when it was first described in 2001, its known distribution has since expanded to Spain, England, Sweden, and Azerbaijan, among other...

Alcorn's pocket gopher
Alcorn's pocket gopher
Pappogeomys alcorni

Alcorn's pocket gopher (*Pappogeomys bulleri alcorni*) is a subspecies of rodent in the family Geomyidae. It is endemic to Mexico. Its natural habitat is montane pine-oak forest (910-3,010 m) in the Sierra del Tigre of south Jalisco. It is reputed...
Aldabra flying fox
Aldabra flying fox
Pteropus aldabrensis

The Aldabra flying fox (*Pteropus aldabrensis*) is a species of megabat in the genus *Pteropus*. It is endemic to the Aldabra Atoll in the Seychelles, like *Chaerephon pusilla*, though the latter may be the same species as the little free-tailed bat.
Alexander Archipelago Wolf
Alexander Archipelago Wolf
Canis lupus ligoni

The Alexander Archipelago wolf (*Canis lupus ligoni*) is a subspecies of the Gray wolf. It first arrived in Alaska sometime between 7,000 and 8,000 years ago, after the end of the Wisconsin glaciation period. It is thought that the wolves were...
Alexander's bush squirrel
Alexander's bush squirrel
Paraxerus alexandri

Alexander's bush squirrel (*Paraxerus alexandri*) is a species of squirrel native to the Democratic Republic of the Congo and Uganda. It is arboreal and lives in tropical moist forests, especially undisturbed mature forests. It is a common species...
Alexander's kusimanse
Alexander's kusimanse
Crossarchus alexandri

Alexander's kusimanse (*Crossarchus alexandri*) is a mongoose species native to Central African rainforests up to an elevation of 2,900 m (9,500 ft). It has a body length of 30 to 45 cm (12 to 18 in) and weighs between 0.45 and 1.4 kg (0.99 and...
Alexandria false antechinus
Alexandria false antechinus
Pseudantechinus mimulus

The Alexandria false antechinus (*Pseudantechinus mimulus*), also known as the Carpentarian false antechinus or Carpentarian pseudantechinus, is a small carnivorous marsupial, found only in a number of small, isolated localities in northern...
Algerian gerbil
Algerian gerbil
Gerbillus garamantis

Algerian gerbil (*Gerbillus garamantis*) is a species of rodent distributed mainly in Algeria. It is sometimes considered a subspecies of the dwarf gerbil.
Algerian mouse
Algerian mouse
Mus spretus

The Algerian mouse (*Mus spretus*), also known as the western Mediterranean mouse, is a wild species of mouse closely related to the house mouse, native to open habitats around the western Mediterranean.
Allegheny Woodrat
Allegheny Woodrat
Neotoma magister

The Allegheny woodrat (*Neotoma magister*) is a species of "pack rat" from North

America. It is the second-largest member of the native North American rats and was once believed to be a subspecies of the Eastern woodrat (*Neotoma floridana*).

Allen's big-eared bat

Allen's big-eared bat

Idionycteris phyllotis

Allen's big-eared bat (*Idionycteris phyllotis*) is a species of vesper bat in the monotypic genus *Idionycteris*. It occurs in Mexico and in Arizona, California, Nevada, New Mexico, Utah, and Colorado in the United States.

Allen's chipmunk

Allen's chipmunk

Neotamias senex

Allen's chipmunk (*Neotamias senex*) is a species of chipmunk. It is also known as the shadow chipmunk. It is native to the western United States, where it occurs in California, Nevada, and Oregon. It is a common species of the Sierra Nevada....

Allen's cotton rat

Allen's cotton rat

Sigmodon alleni

Allen's cotton rat (*Sigmodon alleni*) is a species of rodent in the family Cricetidae. It is endemic to western Mexico, where its distribution extends from Sinaloa to Oaxaca. The formerly recognized *S. planifrons* and *S. vulcani* are now considered...

Allen's spotted bat

Allen's spotted bat

Glauconycteris humeralis

Allen's spotted bat (*Glauconycteris humeralis*) is a species of vesper bat in the family Vespertilionidae found in the following countries: the Central African Republic, the Democratic Republic of the Congo, Kenya, and Uganda. It is found in...

Allen's squirrel

Allen's squirrel

Sciurus alleni

Allen's squirrel (*Sciurus alleni*) is a tree squirrel in the genus *Sciurus* endemic to northern Mexico. It has no subspecies.

Allen's striped bat

Allen's striped bat

Glauconycteris alboguttata

Allen's striped bat (*Glauconycteris alboguttata*) is a species of bat in the family Vespertilionidae, the vesper bats. It is native to Africa, where it occurs in Cameroon and the Democratic Republic of the Congo. This species can be found in...

Allen's Swamp Monkey

Allen's Swamp Monkey

Allenopithecus nigroviridis

The Allen's swamp monkey (*Allenopithecus nigroviridis*) is a species of Old World monkey and the only member of the genus *Allenopithecus*. Phylogenetically, it is a sister clade to the guenons, but differs in dentition and habits. Allen's swamp...

Allen's wood mouse

Allen's wood mouse

Hylomyscus alleni

Allen's hylomyscus or Allen's wood mouse (*Hylomyscus alleni*) is a species of rodent in the family Muridae. It is native to West and Central Africa, where it is widely distributed. It occurs in deciduous forest habitat.

Allen's woodrat
Allen's woodrat
Hodomys alleni

Allen's woodrat (*Hodomys alleni*) is a species of rodent in the family Cricetidae. It is the only species in the genus *Hodomys*.

Allen's yellow bat
Allen's yellow bat
Rhogeessa alleni

Allen's yellow bat (*Rhogeessa alleni*) is a species of vesper bat. There is some taxonomic debate surrounding this species, with some authors considering *Baeodon* a genus rather than a subgenus. It is endemic to Mexico.

Allied rock-wallaby
Allied rock-wallaby
Petrogale assimilis

The allied rock-wallaby or Weasel rock-wallaby (*Petrogale assimilis*) is a species of rock-wallaby found in northeastern Queensland, Australia. It forms part of the *P. lateralis/penicillata* species complex and is very similar to six other species...

Alpaca
Alpaca
Vicugna pacos

The alpaca (*Lama pacos*) is a species of South American camelid mammal. It is similar to, and often confused with, the llama. However, alpacas are often noticeably smaller than llamas. The two animals are closely related and can successfully...

Alpine Chipmunk
Alpine Chipmunk
Neotamias alpinus

The alpine chipmunk (*Neotamias alpinus*) is a species of chipmunk native to the high elevations of the Sierra Nevada of California.

Alpine field mouse
Alpine field mouse
Apodemus alpicola

The alpine field mouse (*Apodemus alpicola*) is a species of rodent in the family Muridae. It is found in Austria, France, Italy, Germany, and Switzerland.

Alpine Ibex
Alpine Ibex
Capra ibex

The Alpine ibex (*Capra ibex*) is a species of wild goat that lives in the mountains of the European Alps. After being extirpated from most areas by the 19th century, the Alpine ibex was successfully reintroduced to parts of its historical range....

Alpine long-eared bat
Alpine long-eared bat
Plecotus macrobullaris

The Alpine long-eared bat or mountain long-eared bat (*Plecotus macrobullaris*) is a species of long-eared bat. It was originally described from Switzerland and Austria as a species intermediate between the brown long-eared bat and the grey...

Alpine Marmot
Alpine Marmot
Marmota marmota

The Alpine marmot (*Marmota marmota*) is a large ground-dwelling squirrel that lives in mountainous areas of central and southern Europe, in the Alps, Carpathians, Tatras and Northern Apennines. In 1948 they were reintroduced with success in the...

Alpine Musk Deer

Alpine Musk Deer

Moschus chrysogaster

The Alpine musk deer (*Moschus chrysogaster*) is an endangered musk deer species found only in Asia. It is not true deer, but rather its family is closely related to Bovidae, the group that contains antelopes, bovines, sheep, and goats.

Alpine pika

Alpine pika

Ochotona alpina

The alpine pika (*Ochotona alpina*) is a species of small mammal in the pika family, Ochotonidae. The summer pelage of different subspecies varies drastically but, in general, it is dark or cinnamon brown, turning to grey with a yellowish tinge...

Alpine pine vole

Alpine pine vole

Microtus multiplex

The alpine pine vole (*Microtus multiplex*) is a species of rodent in the family Cricetidae. It is found in Austria, France, Italy, Serbia, Montenegro, Bosnia and Herzegovina (Una National Park) and Switzerland.

Alpine shrew

Alpine shrew

Sorex alpinus

The alpine shrew (*Sorex alpinus*) is a species of mammal in the family Soricidae. It is found in the alpine meadows and coniferous forests of central and southern European mountain ranges.

Alpine woolly rat

Alpine woolly rat

Mallomys gunung

The alpine woolly rat (*Mallomys gunung*) is a species of rodent in the family Muridae. It is found only in West Papua, Indonesia. It is found only at high elevations, has an extent of occurrence less than 5,000 km², and is known from only two...

Alston's brown mouse

Alston's brown mouse

Scotinomys teguina

Alston's brown mouse, also called Alston's singing mouse, short-tailed singing mouse, or singing mouse (*Scotinomys teguina*), is a species of rodent in the family Cricetidae. It is found in Central America, from Chiapas, Mexico, to western...

Alston's cotton rat

Alston's cotton rat

Sigmodon alstoni

Alston's cotton rat (*Sigmodon alstoni*) is a rodent species from South America. It is found in Brazil, Colombia, Guyana, Suriname, and Venezuela, where it inhabits lowland savannas.

Alston's mouse opossum

Alston's mouse opossum

Marmosa alstoni

Alston's mouse opossum (*Marmosa alstoni*), also known as Alston's opossum, is a

medium-sized pouchless marsupial of the family Didelphidae. It is arboreal and nocturnal, inhabiting forests from Belize to northern Colombia. The main components of...

Alta Floresta titi

Alta Floresta titi

Plecturocebus grovesi

The Alta Floresta titi (*Plecturocebus grovesi*) is a species of titi, a type of New World monkey, endemic to Brazil. It was described from the municipality of Alta Floresta in the state of Mato Grosso. It is a member of the Eastern Amazonian clade...

Altai argali

Altai argali

Ovis ammon ammon

Altai argali (*Ovis ammon ammon*) is a traditional subspecies of argali, a wild sheep that roams the highlands of the Altai Mountains in Central Asia. Altai argali are the largest sheep in the world and also have the heaviest horns. The horns of...

Altai birch mouse

Altai birch mouse

Sicista napaea

The Altai birch mouse (*Sicista napaea*) is a species of rodent in the family Sminthidae. It is native to Russia and Kazakhstan. A baby Altai birch mouse is called a 'pinkie, kitten or pup'. The females are called 'doe' and males 'buck'. A Altai...

Altai mole

Altai mole

Talpa altaica

The Altai mole or Siberian mole (*Talpa altaica*) is a species of mole in the family Talpidae. It is found throughout the taiga zone of south-central Siberia in Russia, as far south as northern Mongolia and Kazakhstan. This mole lives in forested...

Altai Wapiti

Altai Wapiti

Cervus canadensis sibiricus

The Altai wapiti (*Cervus canadensis sibiricus*) is a subspecies of Elk. It is sometimes called the Altai elk by North Americans.

Altiplano chinchilla mouse

Altiplano chinchilla mouse

Chinchillula sahamae

The Altiplano chinchilla mouse or achallo (*Chinchillula sahamae*) is a species of rodent in the family Cricetidae. It is the only species in the genus Chinchillula. It is found in Argentina, Bolivia, Chile, and Peru.

Altiplano grass mouse

Altiplano grass mouse

Akodon lutescens

The Altiplano grass mouse (*Akodon lutescens*) is a species of rodent in the family Cricetidae. It is found in Bolivia, and Peru.

Amami rabbit

Amami rabbit

Pentalagus furnessi

The Amami rabbit (*Pentalagus furnessi* ; Amami:), or Amami no-kuro-usagi (アマミノクロウサギ 奄美野黒兔, lit. "Amami wild black rabbit"), also known as the Ryukyu rabbit

is a dark-furred rabbit which is only found in Amami Ōshima and Toku-no-Shima, two small...

Amargosa vole

Amargosa vole

Microtus californicus scirpensis

The Amargosa vole (*Microtus californicus scirpensis*) is one of 17 subspecies of the California vole (*M. californicus*). The most closely related subspecies is *M. californicus vallicola*.

Amazon bamboo rat

Amazon bamboo rat

Dactylomys dactylinus

For Asian bamboo rats (family: Spalacidae), see Bamboo rat. The Amazon bamboo rat (*Dactylomys dactylinus*) is a species of spiny rat from the Amazon Basin of South America. It is also referred to as coro-coro, Toró, Rato-do-Bambú, or Rata del...

Amazon black howler

Amazon black howler

Alouatta nigerrima

The Amazon black howler (*Alouatta nigerrima*) is a species of howler monkey, a type of New World monkey, endemic to the south-central Amazon in Brazil. Until 2001, most authorities included it as a subspecies (or simply a taxonomically...

Amazon dwarf squirrel

Amazon dwarf squirrel

Microsciurus flaviventer

The Amazon dwarf squirrel (*Microsciurus flaviventer*) is a chipmunk-sized tree squirrel native to South America.

Amazon River Dolphin

Amazon River Dolphin

Inia geoffrensis

The Amazon river dolphin (*Inia geoffrensis*) is a species of toothed whale. It is the largest species of river dolphin, and it has one of the widest-ranging diets among toothed whales. The Amazon river dolphin has been treated with suspicion by...

Amazon Weasel

Amazon Weasel

Mustela africana

The Amazon weasel (*Neogale africana*) is a South American weasel described by Anselme Gaëtan Desmarest in 1818. It is the largest of the three species of South American weasel.

Amazonian brown brocket

Amazonian brown brocket

Mazama nemorivaga

The Amazonian brown brocket (*Mazama nemorivaga*), also known as the small brown brocket, is a small species of deer that is almost entirely restricted to South America.

Amazonian hocicudo

Amazonian hocicudo

Oxymycterus amazonicus

The Amazonian hocicudo (*Oxymycterus amazonicus*) is a species of rodent in the family Cricetidae from South America. It is found in the Amazon basin in Brazil where it lives in moist lowland forest. It is a common species with a large range and...

Amazonian Manatee
Amazonian Manatee
Trichechus inunguis

The Amazonian manatee (*Trichechus inunguis*) is a large aquatic mammal with a bizarre appearance, first described as "a curious combination of a seal and a hippopotamus". It is the smallest of the three extant species of manatee. Manatees are...

Amazonian marsh rat
Amazonian marsh rat
Holochilus sciureus

The Amazonian marsh rat (*Holochilus sciureus*), also known as the common marsh rat, or simply the marsh rat, is a rodent species from South America.

Amazonian red-sided opossum
Amazonian red-sided opossum
Monodelphis glirina

The Amazonian red-sided opossum (*Monodelphis glirina*) is a South American opossum species of the family Didelphidae, formerly viewed as part of *M. brevicaudata*. It is found in Bolivia, Brazil and Peru, where it inhabits the Amazon rainforest. It...

Amazonian sac-winged bat
Amazonian sac-winged bat
Saccopteryx gymnura

The Amazonian sac-winged bat (*Saccopteryx gymnura*) is a bat species of the family Emballonuridae from South America. It is found in northern Brazil, Guyana, French Guiana, Suriname and possibly Venezuela.

Amber Mountain fork-marked lemur
Amber Mountain fork-marked lemur
Phaner electromontis

The Amber Mountain fork-marked lemur (*Phaner electromontis*), also known as the Montagne d'Ambre fork-crowned lemur or Tanta, is a small primate, and like the other lemurs, can only be found on the island of Madagascar. The species is named after...

American Beaver
American Beaver
Castor canadensis

The North American beaver (*Castor canadensis*) is a semiaquatic animal native to North America. It is one of the official national wildlife of Canada symbols and is the official state mammal of Oregon and New York.

American Badger
American Badger
Taxidea taxus

The American badger (*Taxidea taxus*) is a member of the Mustelidae, a diverse family of carnivorous mammals that also includes weasels, otters, ferrets, and the wolverine. It is similar in appearance to the European badger, although not closely...

American Bison
American Bison
Bison bison

The American bison (*Bison bison*) is a species of bison native to North America. Sometimes colloquially referred to as buffalo (a distinct species of bovine), it is one of two extant species of bison, alongside the European bison. Once roaming

in...

American Black Bear
American Black Bear
Ursus americanus

The American black bear (*Ursus americanus*) is a medium-sized bear endemic to North America. It is the continent's smallest and most widely distributed bear species. American black bears are omnivores, with their diets varying greatly depending on...

American Hog-Nosed Skunk
American Hog-Nosed Skunk
Conepatus leuconotus

The American hog-nosed skunk (*Conepatus leuconotus*) is a species of hog-nosed skunk from Central and North America, and is one of the largest skunks in the world, growing to lengths of up to 2.7 feet (82 cm). Recent work has concluded the...

American Marten
American Marten
Martes americana

The American marten (*Martes americana*) is a small North American mammal, a member of the family Mustelidae. It is sometimes called simply the Pine marten; the name "pine marten" is derived from the common name of the distinct Eurasian species,...

American Mink
American Mink
Neovison vison

The American mink is a semiaquatic species of mustelid native to North America. It is considered by some as the cutest member of the weasel family. Its name is derived from the Swedish word 'meank'. Being part of the weasel family, American minks...

American Pika
American Pika
Ochotona princeps

American pikas (*Ochotona princeps*) are mountain-dwelling mammals found in the mountains of western North America. They are smaller relatives of rabbits and hares. They have two different ways of foraging; they either directly consume food or they...

American Pygmy Shrew
American Pygmy Shrew
Sorex hoyi

The American pygmy shrew (*Sorex hoyi*) is a small mole-like mammal that was first discovered in 1831 by naturalist William Cane in Georgian Bay, Parry Sound. It is believed to be the second-smallest mammal in the world. American pygmy shrew has an...

American red fox
American red fox
Vulpes vulpes fulvus

The North American red fox (*Vulpes vulpes fulva*) is a North American subspecies of the red fox. It is the largest of the true foxes and one of the most widely distributed members of the order Carnivora, occurring in North America. It is listed...

American Red Squirrel
American Red Squirrel
Tamiasciurus hudsonicus

The American red squirrel (*Tamiasciurus hudsonicus*) is one of three species of tree squirrels currently classified in the genus *Tamiasciurus*, known as the pine squirrels. The American red squirrel is variously known as the Pine squirrel, North...

American Shrew Mole
American Shrew Mole
Neurotrichus gibbsii

The American shrew mole (*Neurotrichus gibbsii*) is the smallest species of mole. It is not closely related to the Asian shrew mole (*Uropsilus* in *Urotrichini*). Its common name "shrew mole" comes from shrew-like fur and typical mole head.

American Water Shrew
American Water Shrew
Sorex palustris

The American water shrew (*Sorex palustris*) is a mole-like mammal found throughout North America. This tiny animal lives in semi-aquatic habitats and is known for being the smallest mammalian diver.

Ammodile
Ammodile
Ammodillus imbellis

The ammodile, walo or Somali gerbil (*Ammodillus imbellis*) is a species of rodent in the family Muridae. It is the only species in the genus *Ammodillus*. It is found in Ethiopia and Somaliland. Its natural habitat is subtropical or tropical dry...

Amphinectomys savamis
Amphinectomys savamis
Amphinectomys savamis

Amphinectomys savamis, also known as the Ucayali water rat or amphibious rat, is a rodent from the Peruvian Amazon. It is placed as the only member of genus *Amphinectomys* in the tribe *Oryzomyini* of family *Cricetidae*. It is similar to *Nectomys*,...

Amur Hedgehog
Amur Hedgehog
Erinaceus amurensis

The Amur hedgehog (*Erinaceus amurensis*) is a species of hedgehog similar to the European hedgehog in appearance and behavior, although it is more lightly colored. The name "hedgehog" refers to the pig-like grunts it makes as it forages.

Amur lemming
Amur lemming
Lemmus amurensis

The Amur lemming (*Lemmus amurensis*) is a species of lemmings found near the Amur River in Siberia.

Amur Leopard
Amur Leopard
Panthera pardus orientalis

The Amur leopard (*Panthera pardus orientalis*) is a leopard subspecies native to Russia and China. It is listed as Critically Endangered on the IUCN Red List and is considered one of the rarest cats on Earth.

Anastasia Island beach mouse
Anastasia Island beach mouse
Peromyscus polionotus phasma

The Anastasia Island beach mouse (*Peromyscus polionotus phasma*) is a subspecies of

the oldfield mouse of the southeastern United States. It occurs in the sand dunes of Florida and Alabama beaches. It has been classified by the United States Fish...
Anatolian boar
Anatolian boar
Sus scrofa libycus

The Anatolian boar (*Sus scrofa libycus*) is a subspecies of wild boar endemic to Turkey, Levant, Israel and Transcaucasia. It is likely to be one of the ancestors of today's domestic pigs.
Anatolian vole
Anatolian vole
Microtus anatolicus

The Anatolian vole (*Microtus anatolicus*) is a species of vole found in Turkey, specifically Konya Province. The population is unknown but it lives in small colonies. The species exists in an area that is a remnant zone of Tethys Ocean, and this...
Anchieta's pipistrelle
Anchieta's pipistrelle
Pipistrellus anchietae

Anchieta's serotine (*Neoromicia anchieta*), formerly known as Anchieta's pipistrelle, is a species of vesper bat. It is found in Angola, Democratic Republic of the Congo, South Africa, Zambia, Zimbabwe and Madagascar. The species inhabits dry...
Andaman horseshoe bat
Andaman horseshoe bat
Rhinolophus cognatus

The Andaman horseshoe bat (*Rhinolophus cognatus*) is a species of bat in the family Rhinolophidae. It is endemic to the Andaman Islands. During the day, it roosts in caves, but may also choose tree hollows.
Andaman rat
Andaman rat
Rattus stoicus

The Andaman rat (*Rattus stoicus*) is a species of rodent in the family Muridae. It is endemic to the Andaman Islands, where it has been recorded on Henry Lawrence Island, South Andaman, and Middle Andaman. It is a nocturnal species that lives in...
Andaman shrew
Andaman shrew
Crocidura andamanensis

The Andaman shrew or Andaman white-toothed shrew (*Crocidura andamanensis*) is a critically endangered species of mammal in the family Soricidae. It is endemic to the South Andaman Island of India. They are usually active by twilight or in the...
Andaman spiny shrew
Andaman spiny shrew
Crocidura hispida

The Andaman spiny shrew or Andaman shrew (*Crocidura hispida*) is a species of mammal in the family Soricidae. It is endemic to India. Its natural habitat is subtropical or tropical dry forests. It is threatened by habitat loss.
Andean big-eared mouse
Andean big-eared mouse
Auliscomys sublimis

The Andean big-eared mouse (*Auliscomys sublimis*) is a species of rodent in the

family Cricetidae. It is found in Argentina, Bolivia, Chile, and Peru.

Andean caenolestid

Andean caenolestid

Caenolestes condorensis

The Andean caenolestid (*Caenolestes condorensis*), also known as the Andean shrew opossum or Condor caenolestid, is a shrew opossum known only from Cordillera del Cóndor (Ecuador), its type locality. It was first described by zoologists Bruce D....

Andean gerbil mouse

Andean gerbil mouse

Eligmodontia puerulus

The Andean gerbil mouse or Altiplano laucha (*Eligmodontia puerulus*) is a species of rodent in the family Cricetidae. It is found in Argentina, Bolivia, Chile, and Peru.

Andean hairy armadillo

Andean hairy armadillo

Chaetophractus nationi

The Andean hairy armadillo (*Chaetophractus nationi*) is an armadillo found mainly in Bolivia, in the region of the Puna; the departments of Oruro, La Paz, and Cochabamba (Gardner, 1993). Nowark (1991) describes it as distributed in Bolivia and...

Andean leaf-eared mouse

Andean leaf-eared mouse

Phyllotis andium

The Andean leaf-eared mouse (*Phyllotis andium*) is a species of rodent in the family Cricetidae. It is found in Ecuador and Peru.

Andean Mountain Cat

Andean Mountain Cat

Leopardus jacobita

The Andean mountain cat (*Leopardus jacobita*) is a small wild cat native to the high Andes. It is regarded as one of the world's most endangered wild cats and perhaps South America's rarest felid, yet, as far as cat species go, it is amongst the...

Andean mountain cavy

Andean mountain cavy

Microcavia niata

The Andean mountain cavy (*Microcavia niata*) is a species of rodent in the family Caviidae. It is found in Bolivia, Chile and Peru.

Andean mouse

Andean mouse

Andinomys edax

The Andean mouse (*Andinomys edax*) is a species of rodent in the family Cricetidae. It is the only species in the genus *Andinomys*. It is found in Argentina, Bolivia, Chile, and Peru.

Andean porcupine

Andean porcupine

Coendou quichua

The Andean porcupine (*Coendou quichua*) or Quichua porcupine is a species of rodent in the family Erethizontidae. It is found in the Andes of northern Ecuador and Colombia as well as in Panama. This porcupine is little known, but is probably...

Andean rat

Andean rat

Lenoxus apicalis

The Andean rat (*Lenoxus apicalis*), or white-tailed akodont, is the only species in the genus *Lenoxus*. It is a rodent in the tribe Akodontini found on the eastern slopes of the Andes of eastern Peru and western Bolivia. Examination of its genome...

Andean squirrel

Andean squirrel

Sciurus pucheranii

The Andean squirrel (*Sciurus pucheranii*) is a tree squirrel endemic to Colombia where it inhabits montane rain forest and cloud forests of the Cordillera Occidental and Cordillera Central ranges of the Colombian Andes, at elevations between...

Andean swamp rat

Andean swamp rat

Neotomys ebriosus

The Andean swamp rat (*Neotomys ebriosus*) is a species of rodent in the family Cricetidae. It is the only species in the genus *Neotomys*. It is found in Argentina, Bolivia, Chile, and Peru.

Andean vesper mouse

Andean vesper mouse

Calomys lepidus

The Andean vesper mouse (*Calomys lepidus*) is a species of rodent in the family Cricetidae. It is found in Argentina, Bolivia, Chile, and Peru. Andean vesper mice are typically found at high elevations in the altiplano region. Due to their...

Andean white-eared opossum

Andean white-eared opossum

Didelphis pernigra

The Andean white-eared opossum (*Didelphis pernigra*) is an opossum species from South America. It is found in the Andes Mountains, ranging from Venezuela to Bolivia. Together with the Guianan white-eared opossum (*D. imperfecta*), this species was...

Andersen's flying fox

Andersen's flying fox

Pteropus intermedius

Andersen's flying fox (*Pteropus intermedius*) is a species of flying fox in the family Pteropodidae found in south Burma and west Thailand. Although it is hunted for both food and as a pest, it is not known if this has a significant impact on the...

Andersen's fruit-eating bat

Andersen's fruit-eating bat

Dermanura anderseni

Andersen's fruit-eating bat (*Dermanura anderseni*) is a bat species from South America. It is found in Bolivia, Brazil, Colombia, Ecuador, French Guiana, and Peru.

Andersen's naked-backed fruit bat

Andersen's naked-backed fruit bat

Dobsonia anderseni

Andersen's naked-backed fruit bat or Andersen's bare-backed fruit bat (*Dobsonia anderseni*) is a large cave-dwelling species of megabat in the family Pteropodidae. It is endemic to the Bismarck Archipelago including the Admiralty Islands in Papua...

Andersen's slit-faced bat
Andersen's slit-faced bat
Nycteris aurita

Andersen's slit-faced bat (*Nycteris aurita*) is a slit-faced bat species found in East Africa. It has been recorded in Somaliland, through South Sudan and Ethiopia, into Kenya and Tanzania. No information is available on the population size of...

Anderson's four-eyed opossum
Anderson's four-eyed opossum
Philander andersoni

Anderson's four-eyed opossum (*Philander andersoni*) is an opossum species from South America. It is found in Brazil, Colombia, Ecuador, Peru and Venezuela. Its dorsal fur is dark, with a black stripe, about 3-4 cm wide, going vertically down the...

Anderson's gerbil
Anderson's gerbil
Gerbillus andersoni

Anderson's gerbil (*Gerbillus andersoni*) is a species of rodent distributed from Tunisia to Israel. Their habitats and diets are similar to other gerbils. The gestation period is 20-22 days and the average litter size is four or five. The IUCN...

Anderson's Oldfield mouse
Anderson's Oldfield mouse
Thomasomys andersoni

Anderson's Oldfield mouse (*Thomasomys andersoni*) is a species of rodent in the family Cricetidae.

Anderson's shrew
Anderson's shrew
Suncus stoliczkanus

Anderson's shrew (*Suncus stoliczkanus*) is a medium-sized species of shrew. It is light gray in color with yellow fur around the throat and pectoral region, comparatively large ears and a tail that measures about 50 - 70% of body length. This...

Anderson's shrew mole
Anderson's shrew mole
Uropsilus andersoni

The Anderson's shrew mole (*Uropsilus andersoni*) is a species of mammal in the family Talpidae. It is endemic to China. Its species name "andersoni " was chosen to honor American scientific collector Malcolm Playfair Anderson.

Anderson's squirrel
Anderson's squirrel
Callosciurus quinquestriatus

Anderson's squirrel (*Callosciurus quinquestriatus*) is a species of rodent in the family Sciuridae. It is found in forests in China (Yunnan only) and Myanmar. It is threatened by habitat loss.

Anderson's white-bellied rat
Anderson's white-bellied rat
Niviventer andersoni

Anderson's white-bellied rat (*Niviventer andersoni*) is a species of rodent in the family Muridae. It is endemic to China and known from Yunnan, Sichuan, and Shaanxi provinces. Its range might extend to northern Guizhou. It inhabits montane...

Andrew Rebori's house bat

Andrew Rebori's house bat
Scotophilus andrewreborii

Andrew Rebori's house bat (*Scotophilus andrewreborii*) is a species of bat found in Africa.

Andrew's hill rat
Andrew's hill rat
Bunomys andrewsi

Andrew's hill rat (*Bunomys andrewsi*) is a species of rodent in the family Muridae. It is found only in Sulawesi and Buton Island, Indonesia.

Andrews' beaked whale
Andrews' beaked whale
Mesoplodon bowdoini

Andrews' beaked whale (*Mesoplodon bowdoini*), sometimes known as the deep-crest beaked whale or splay-toothed whale, is one of the most poorly known members of a poorly known genus. The species has never been observed in the wild, and is known...

Andrews's three-toed jerboa
Andrews's three-toed jerboa
Stylodipus andrewsi

Andrews's three-toed jerboa, or the Mongolian jerboa, (*Stylodipus andrewsi*) is a species of rodent in the family Dipodidae. It is found in China (Inner Mongolia, Gansu, and Ningxia provinces) and Mongolia.

Angel Island mouse
Angel Island mouse
Peromyscus guardia

The Angel Island mouse (*Peromyscus guardia*), or La Guarda deermouse, is a species of rodent in the family Cricetidae.

Angola Colobus
Angola Colobus
Colobus angolensis

The Angola colobus (*Colobus angolensis*), Angolan black-and-white colobus or Angolan colobus, is a primate species of Old World monkey belonging to the genus *Colobus*.

Angolan African dormouse
Angolan African dormouse
Graphiurus angolensis

The Angolan African dormouse (*Graphiurus angolensis*) is a species of rodent in the family Gliridae. Found in central and north Angola and western Zambia, it has been recorded from seven localities over an altitudinal range from 1,000 to 2,000 m...

Angolan brush-furred rat
Angolan brush-furred rat
Lophuromys angolensis

The Angolan brush-furred rat (*Lophuromys angolensis*) is a species of brush-furred mouse found in Angola and the southwest of the Democratic Republic of Congo.

Angolan epauletted fruit bat
Angolan epauletted fruit bat
Epomophorus angolensis

The Angolan epauletted fruit bat (*Epomophorus angolensis*) is a species of megabat in the family Pteropodidae. It is found in Angola and Namibia. Its natural habitats are dry savanna and moist savanna. It is threatened by habitat loss.

Angolan free-tailed bat
Angolan free-tailed bat
Mops condylurus

The Angolan free-tailed bat (*Mops condylurus*) is a species of bat in the family Molossidae. It is found in Angola, Benin, Botswana, Burkina Faso, Burundi, Cameroon, the Republic of the Congo, the Democratic Republic of the Congo, Ivory Coast,...

Angolan genet
Angolan genet
Genetta angolensis

The Angolan genet or miombo genet (*Genetta angolensis*) is a genet species endemic to Southern Africa. It is considered common in this region and therefore listed as Least Concern in the IUCN Red List. Little is known about its ecology.

Angolan Giraffe
Angolan Giraffe
Giraffa camelopardalis angolensis

The Angolan giraffe (*Giraffa camelopardalis angolensis* or *Giraffa giraffa angolensis*) is a subspecies of giraffe. It is also known as the Namibian giraffe or Smokey giraffe.

Angolan hairy bat
Angolan hairy bat
Cistugo seabrae

The Angolan hairy bat (*Cistugo seabrae*) also known as Angolan wing-gland bat is a species of bat in the Cistugidae family. It can be found in the following countries: Angola, Namibia, and South Africa. It is found in these habitats: hot deserts.

Angolan kusimanse
Angolan kusimanse
Crossarchus ansorgei

The Angolan kusimanse (*Crossarchus ansorgei*), also known as Ansorge's kusimanse, is a species of small mongoose. There are two recognized subspecies: *C. a. ansorgei*, found in Angola; and *C. a. nigricolor*, found in DR Congo, which do not have...

Angolan long-eared bat
Angolan long-eared bat
Laephotis angolensis

The Angolan long-eared bat (*Laephotis angolensis*) is a species of vesper bat in the Vespertilionidae family. It can be found in moist savanna in Angola and Democratic Republic of the Congo.

Angolan marsh rat
Angolan marsh rat
Dasymys nudipes

The Angolan marsh rat (*Dasymys nudipes*) is a species of rodent in the family Muridae. It is found in Angola, Botswana, Namibia, and Zambia. Its natural habitats are subtropical or tropical seasonally wet or flooded lowland grassland and swamps. It...

Angolan multimammate mouse
Angolan multimammate mouse
Myomyscus angolensis

The Angolan multimammate mouse (*Mastomys angolensis*) is a species of rodent in the family Muridae. It is found in Angola and Democratic Republic of the Congo. It was formerly classified in the genus *Myomyscus* but has been reclassified into the...

Angolan rousette
Angolan rousette
Myonycteris angolensis

The Angolan fruit bat, Angolan rousette or Silky bat (*Myonycteris angolensis*) is a species of megabat in the family Pteropodidae. It is found in Angola, Burundi, Cameroon, Central African Republic, Republic of Congo, Democratic Republic of...

Angolan slender mongoose
Angolan slender mongoose
Galerella flavescens

The Angolan slender mongoose (*Herpestes flavescens*) is a mongoose native to southwestern Africa, specifically southwestern Angola and northwestern Namibia. It has been listed as "Least Concern" on the IUCN Red List, as it is not threatened and...

Angolan talapoin
Angolan talapoin
Miopithecus talapoin

The Angolan talapoin (*Miopithecus talapoin*), also known as the southern talapoin, is a species of primate in the family Cercopithecidae. Talapoin is the smallest species of Old-World monkeys.

Angolan vlei rat
Angolan vlei rat
Otomys anchietae

The Angolan vlei rat (*Otomys anchietae*) is a species of rodent in the family Muridae. It is found only in Angola. Its natural habitats are dry savanna, moist savanna, and subtropical or tropical seasonally wet or flooded lowland grassland. It is...

Angolan wood mouse
Angolan wood mouse
Hylomyscus carillus

The Angolan hylomyscus or Angolan wood mouse (*Hylomyscus carillus*) is a species of rodent in the family Muridae. It is found only in Angola. Its natural habitat is subtropical or tropical dry forest.

Angoni vlei rat
Angoni vlei rat
Otomys angoniensis

The Angoni vlei rat (*Otomys angoniensis*) is a species of rodent in the family Muridae. It is found in Botswana, Kenya, Malawi, Mozambique, South Africa, Eswatini, Tanzania, Zambia, and Zimbabwe. Its natural habitats are moist savanna, temperate...

Angular hocicudo
Angular hocicudo
Oxymycterus angularis

The angular hocicudo (*Oxymycterus angularis*) is a rodent species from South America. It is found in Brazil.

Angulate pipistrelle
Angulate pipistrelle
Pipistrellus angulatus

The angulate pipistrelle (*Pipistrellus angulatus*), also known as the New Guinea pipistrelle, is a species of vesper bat found in Papua New Guinea and the Solomon Islands.

Anhui musk deer
Anhui musk deer

Moschus anhuiensis

The Anhui musk deer (*Moschus anhuiensis*) is an endangered species of musk deer that is endemic to the Dabie Mountains of western Anhui province, China. It was formerly described as a subspecies of *Moschus berezovskii* and *Moschus moschiferus*, but...

Anita's leaf-eared mouse

Anita's leaf-eared mouse

Phyllotis anitae

Anita's leaf-eared mouse (*Phyllotis anitae*) is a species of rodent in the family Cricetidae. It was discovered in Tucumán Province in northwestern Argentina in alder forest of the upper part of the Southern Andean Yungas ecoregion, on the...

Anjouan myotis

Anjouan myotis

Myotis anjouanensis

The Anjouan myotis (*Myotis anjouanensis*) is a species of vesper bat. It is found only in Comoros.

Ankarana dwarf lemur

Ankarana dwarf lemur

Cheirogaleus shethi

The Ankarana dwarf lemur or Sheth's dwarf lemur (*Cheirogaleus shethi*) is a species of dwarf lemur, part of the *C. medius* group, known only from northern Madagascar. It has been found in two reserves (Ankarana Special Reserve and Analamerana...

Ankarana Special Reserve tufted-tailed rat

Ankarana Special Reserve tufted-tailed rat

Eliurus carletoni

The Ankarana Special Reserve tufted-tailed rat (*Eliurus carletoni*) (synonymous with *Eliurus* sp. SAJ-2009a) is a species of rodent in the family Nesomyidae. It was first described in 2009. It is endemic to Madagascar, in the Ankarana Special...

Ankarana Sportive Lemur

Ankarana Sportive Lemur

Lepilemur ankaranensis

The Ankarana sportive lemur (*Lepilemur ankaranensis*), or Ankarana weasel lemur, is a sportive lemur endemic to Madagascar. It is one of the smaller sportive lemurs with a total length of about 53 cm (21 in), including 25 cm (9.8 in) of tail....

Ankole African mole-rat

Ankole African mole-rat

Tachyoryctes ankoliae

The Ankole mole-rat or Ankole African mole-rat (*Tachyoryctes ankoliae*) is a species of rodent in the family Spalacidae found in Angola and Uganda. Its natural habitats are moist savanna and arable land. Some taxonomic authorities consider it to...

Annamit myotis

Annamit myotis

Myotis annamiticus

The Annamit myotis (*Myotis annamiticus*) is a species of mouse-eared bat in the family Vespertilionidae, described in 2001, and indigenous to the Minh Hóa District on the northern coast of Vietnam. Following its description, investigators...

Annamite Striped Rabbit

Annamite Striped Rabbit

Nesolagus timminsi

The Annamite striped rabbit (*Nesolagus timminsi*) is found in Southeast Asia and only recently became known to Western scientists. It is known as the tho van in Vietnamese and ka tai lai sua in Lao.

Annandale's rat

Annandale's rat

Sundamys annandalei

Annandale's rat (*Sundamys annandalei*) is a species of rodent in the family Muridae. It is found in Indonesia (Sumatra), Peninsular Malaysia, and Singapore. It was classified as *Rattus annandalei* until 2017, but mitochondrial and nuclear DNA show...

Anosy mouse lemur

Anosy mouse lemur

Microcebus tanosi

The Anosy mouse lemur (*Microcebus tanosi*) is a species of mouse lemur known only from the Manantantely Forest and Ivorona Forest in southeastern Madagascar, near Tôlanaro. Specimens were first collected in April 2007, and its discovery was...

Anoura aequatoris

Anoura aequatoris

Anoura aequatoris

Anoura aequatoris (also known as the equatorial tailless bat) is a species of microbat that lives in South America in the countries of Bolivia, Colombia, Ecuador, and Peru.

Anoura peruana

Anoura peruana

Anoura peruana

Anoura peruana is a species of bat from Colombia and Peru. It was elevated to a species in 2010, after previously being considered a subspecies of Geoffroy's tailless bat (*A. geoffroyi*). The females are larger than the males.

Ansell's epauletted fruit bat

Ansell's epauletted fruit bat

Epomophorus anelli

Ansell's epauletted fruit bat (*Epomophorus anelli*) is a species of megabat in the family Pteropodidae.

Ansell's mole-rat

Ansell's mole-rat

Fukomys anelli

Ansell's mole-rat (*Fukomys anelli*) is a species of rodent in the family Bathyergidae. It is endemic to Zambia. Its natural habitats are moist savanna and miombo forests. It is noted for its very long tunnels, up to 2.8 km (1.7 mi) for a single...

Ansell's shrew

Ansell's shrew

Crocidura anellorum

Ansell's shrew (*Crocidura anellorum*) is a species of mammal in the family Soricidae. It is endemic to Zambia. Its natural habitat is subtropical or tropical moist lowland forests. It is threatened by habitat loss.

Ansell's wood mouse

Ansell's wood mouse

Hylomyscus anelli

Ansell's wood mouse (*Hylomyscus anelli*) is a species of rodent in the genus

Hylomyscus. It was described in 1979.

Ansorge's brush-furred rat

Ansorge's brush-furred rat

Lophuromys ansorgei

Ansorge's brush-furred rat (*Lophuromys ansorgei*) is a species of rodent in the family Muridae. It was identified in 1896 by de Winton as *L. ansorgei*. However, it was widely regarded as *L. sicapusi* until 2000, when Walter Verheyen, Theo

Dierckx,...

Ansorge's free-tailed bat

Ansorge's free-tailed bat

Chaerephon ansorgei

Ansorge's free-tailed bat (*Chaerephon ansorgei*) is a species of bat in the family Molossidae native to sub-Saharan Africa. It is named for W.J. Ansorge, who collected the first formally described specimen.

Antarctic Fur Seal

Antarctic Fur Seal

Arctocephalus gazella

The Antarctic fur seal (*Arctocephalus gazella*), is one of eight seals in the genus *Arctocephalus*, and one of nine fur seals in the subfamily Arctocephalinae. Despite what its name suggests, the Antarctic fur seal is mostly distributed in...

Antarctic Minke Whale

Antarctic Minke Whale

Balaenoptera bonaerensis

The Antarctic minke whale or southern minke whale (*Balaenoptera bonaerensis*) is a species of minke whale within the suborder of baleen whales. It is the second smallest rorqual after the common minke whale and the third smallest baleen whale....

Antechinus arktos

Antechinus arktos

Antechinus arktos

Antechinus arktos, the black-tailed antechinus, is a species of small carnivorous marsupial native to Australia.

Antechinus mysticus

Antechinus mysticus

Antechinus mysticus

Antechinus mysticus (common name buff-footed antechinus) is a species of marsupial.

Antelope Jackrabbit

Antelope Jackrabbit

Lepus alleni

The antelope jackrabbit (*Lepus alleni*) is a species of North American hare found in southern Arizona and northwestern Mexico that occupies dry desert areas.

Antillean fruit-eating bat

Antillean fruit-eating bat

Brachyphylla cavernarum

The Antillean fruit-eating bat (*Brachyphylla cavernarum*) is one of two leaf-nosed bat species belonging to the genus *Brachyphylla*. The species occurs in the Caribbean from Puerto Rico to St. Vincent and Barbados. Fossil specimens have also been...

Antillean ghost-faced bat

Antillean ghost-faced bat

Mormoops blainvillei

The Antillean ghost-faced bat (*Mormoops blainvillei*) is a species of bat in the family Mormoopidae. It is found in Cuba, the Dominican Republic, Haiti, Jamaica, and Puerto Rico.

Antilopine Kangaroo

Antilopine Kangaroo

Macropus antilopinus

The antilopine kangaroo (*Osphranter antilopinus*), also known as the antilopine wallaroo or the antilopine wallaby, is a species of macropod found in northern Australia: in Cape York Peninsula in Queensland, the Top End of the Northern Territory,...

Antioquian sac-winged bat

Antioquian sac-winged bat

Saccopteryx antioquensis

The Antioquian sac-winged bat (*Saccopteryx antioquensis*) is a species of bat in the family Emballonuridae found in Colombia.

Aonyx capensis capensis

Aonyx capensis capensis

Aonyx capensis capensis

The Cape clawless otter (*Aonyx capensis capensis*) is a subspecies of African clawless otter found in sub-Saharan Africa near permanent bodies of freshwater and along the seacoast. It is the largest of the Old World otters and the third largest...

Apeco Oldfield mouse

Apeco Oldfield mouse

Thomasomys apeco

The Apeco Oldfield mouse (*Thomasomys apeco*) is a species of rodent in the family Cricetidae. It is known only from a single locality in north central Peru, which includes Rio Abiseo National Park, where it was found in cloud forest at an...

Apennine shrew

Apennine shrew

Sorex samniticus

The Apennine shrew (*Sorex samniticus*) is a species of shrew in the family Soricidae. The mammal is endemic to Italy.

Apodemus avicennicus

Apodemus avicennicus

Apodemus avicennicus

Apodemus avicennicus is a species of rodent found in Iran.

Appalachian Cottontail

Appalachian Cottontail

Sylvilagus obscurus

The Appalachian cottontail (*Sylvilagus obscurus*) is a species of cottontail rabbit in the family Leporidae. It is a rare species found in the upland areas of the eastern United States. The species was only recognized as separate from the New...

Aquatic Genet

Aquatic Genet

Genetta piscivora

The aquatic genet (*Genetta piscivora*) is a genet that has only been recorded in the northeast of the Democratic Republic of the Congo. Since it is only known from about 30 specimens in zoological collections, it had been listed as Data Deficient...

Aquatic rat
Aquatic rat
Anotomys leander

The aquatic rat, Ecuador fish-eating rat, fish-eating rat or Ecuadoran ichthyomyine (*Anotomys leander*) is a South American species of semiaquatic rodent in the family Cricetidae. It is the only species in the genus *Anotomys*. This species is...

Arabian Gazelle
Arabian Gazelle
Gazella arabica

The Arabian gazelle (*Gazella arabica*) is an antelope species in the genus *Gazella*. Gazelle is derived from Arabic: غزال *ġazāl*, Maghrebi pronunciation *ġazēl*. The Arab people traditionally hunted the gazelle, however, later appreciated for their...

Arabian jird
Arabian jird
Meriones arimalius

The Arabian jird (*Meriones arimalius*) is a species of rodent in the family Muridae. It is found in Oman, Saudi Arabia and United Arab Emirates.

Arabian Leopard
Arabian Leopard
Panthera pardus nimr

The Arabian leopard (*Panthera pardus nimr*) is a leopard subspecies native to the Arabian Peninsula. It has been listed as Critically Endangered on the IUCN Red List since 1996 as fewer than 200 wild individuals were estimated to be alive in 2006....

Arabian Oryx
Arabian Oryx
Oryx leucoryx

The Arabian oryx (*Oryx leucoryx*) is amongst the largest desert mammals of the Arab region and throughout history has been an important aspect of its heritage. Once extinct in the wild, it once again inhabits the dry Arabian Peninsula. This...

Arabian pipistrelle
Arabian pipistrelle
Hypsugo arabicus

The Arabian pipistrelle (*Hypsugo arabicus*) is a species of vesper bat in the family Vespertilionidae. It is found only in Iran and Oman.

Arabian Red Fox
Arabian Red Fox
Vulpes vulpes arabica

The Arabian red fox (*Vulpes vulpes arabica*) is a subspecies of the Red fox native to the Arabian Peninsula.

Arabian Sand Gazelle
Arabian Sand Gazelle
Gazella marica

The Arabian sand gazelle (*Gazella marica*) is a species of gazelle found only in the West Asia. These elegant animals are very swift and can run at can run at very high speeds. Gazelles tend to live in herds, and eat fine, easily digestible...

Arabian shrew
Arabian shrew
Crocidura arabica

The Arabian shrew (*Crocidura arabica*) is a species of mammal in the family

Soricidae. It is found in Oman and Yemen. They are solitary carnivores.

Arabian Tahr

Arabian Tahr

Arabitragus jayakari

The Arabian tahr (*Arabitragus jayakari*) is a species of tahr native to eastern Arabia. Until recently, it was placed in the genus *Hemitragus*, but genetic evidence supports its removal to a separate monotypic genus, *Arabitragus*.The Arabian tahr...

Arabian trident bat

Arabian trident bat

Asellia arabica

The Arabian trident bat (*Asellia arabica*) is a species of Old World leaf-nosed bat found in the Middle East.

Arabian Wildcat

Arabian Wildcat

Felis lybica lybica

The Arabian wildcat (*Felis lybica lybica*) is an African wildcat subspecies that inhabits the Arabian Peninsula.

Arabian wolf

Arabian wolf

Canis lupus arabs

The Arabian wolf (*Canis lupus arabs*) is a subspecies of gray wolf native to the Arabian Peninsula, the Negev Desert, the Sinai Peninsula, and Jordan. It is the smallest wolf subspecies, and a desert-adapted subspecies that normally lives in...

Aracá uakari

Aracá uakari

Cacajao ayresi

The Aracá uakari (*Cacajao ayresi*), also known as the Ayres black uakari, is a newly described species of monkey from the northwest Brazilian Amazon. It was found by Jean-Phillipe Boubli of the University of Auckland after following native...

Araguaian River Dolphin

Araguaian River Dolphin

Inia araguaiaensis

The Araguaian river dolphin (*Inia araguaiaensis*) is a recently discovered species of river dolphin from South America. It is closely related to the Amazon river dolphin and was originally distinguished from the latter in January 2014.

Aratathomas's yellow-shouldered bat

Aratathomas's yellow-shouldered bat

Sturnira aratathomasi

Aratathomas's yellow-shouldered bat (*Sturnira aratathomasi*) is a species of bat in the family Phyllostomidae. It is found in Colombia, Ecuador, Peru, and Venezuela.

Archaeophocaena

Archaeophocaena

Archaeophocaena teshioensis

Archaeophocaena teshioensis is a species of extinct porpoise from the Late Miocene Koetoi Formation of Japan living around 6.4–5.5 million years ago (mya). The holotype specimen comprises a partial skull. The animal, along with *Miophocaena* and...

Arctic Fox

Arctic Fox

Vulpes lagopus

The Arctic fox (*Vulpes lagopus*) is a small fox native to the Arctic regions of the Northern Hemisphere and common throughout the Arctic tundra biome. It is well adapted to living in cold environments and is best known for its thick, warm fur that...

Arctic Ground Squirrel
Arctic Ground Squirrel
Spermophilus parryii

The Arctic ground squirrel (*Urocitellus parryii*) (Inuktitut: ᐱᓄᓂᓂᓂ, siksik) is a species of ground squirrel native to the Arctic and Subarctic of North America and Asia. People in Alaska, particularly around the Aleutians, refer to them as...

Arctic Hare
Arctic Hare
Lepus arcticus

The Arctic hare (*Lepus arcticus*) is a highly adapted hare to living in the Arctic tundra and other icy biomes. They can travel together with many other hares and can run up to 60 kilometers per hour (40 mph).

Arctic Lemming
Arctic Lemming
Dicrostonyx torquatus

The Arctic lemming (*Dicrostonyx torquatus*) is a species of rodent native to Russian Federation. This species was once found in England, but it is now extirpated.

Arctic reindeer
Arctic reindeer
Rangifer tarandus eogroenlandicus

The Arctic reindeer (*Rangifer tarandus eogroenlandicus*), also known as the East Greenland caribou or Greenland caribou, was a subspecies of the reindeer (or the caribou in North America) that once lived in eastern Greenland. It has been extinct...

Arctic Shrew
Arctic Shrew
Sorex arcticus

The Arctic shrew (*Sorex arcticus*), also known as the blackback shrew or saddlebacked shrew, is a medium-sized shrew found in Canada and the northern United States. Separate species status has been proposed for the maritime shrew (*Sorex...*

Arctic Wolf
Arctic Wolf
Canis lupus arctos

The Arctic wolf (*Canis lupus arctos*) is a sub-species of grey wolves. These wolves spend their lives in the Arctic tundra, higher than the northern tree line. They are the only wolves in the world with their coloring, unique due to the...

Arcuate horseshoe bat
Arcuate horseshoe bat
Rhinolophus arcuatus

The arcuate horseshoe bat (*Rhinolophus arcuatus*) is a species of bat in the family Rhinolophidae. It is found in Indonesia, Malaysia, Papua New Guinea, and the Philippines.

Arends's golden mole
Arends's golden mole
Carpitalpa arendsi

Arends' golden mole (*Carpitalpa arendsi*) is a species of mammal in the family Chrysochloridae. It is found in Mozambique and Zimbabwe. Its natural habitats are

temperate forests, subtropical or tropical dry forest, subtropical or tropical moist...

Arfak pygmy bandicoot

Arfak pygmy bandicoot

Microperoryctes aplini

The Arfak pygmy bandicoot (*Microperoryctes aplini*) is a species of marsupial in the family Peramelidae. It is endemic to the Arfak mountains in the Vogelkop Peninsula of West Papua, in Indonesia. Its natural habitat is subtropical or tropical...

Argali

Argali

Ovis ammon

The argali (*Ovis ammon*) is a wild sheep that roams the highlands of western East Asia, the Himalayas, Tibet, and the Altai Mountains. It is the largest species of wild sheep.

Argentine bolo mouse

Argentine bolo mouse

Necromys benefactus

The Argentine bolo mouse (*Necromys benefactus*) is a species of rodent in the family Cricetidae. It is endemic to central Argentina, where it is found in the pampas and the drier espinal (lowland thorn brush).

Argentine brown bat

Argentine brown bat

Eptesicus furinalis

The Argentine brown bat (*Eptesicus furinalis*), is a bat species from South and Central America.

Argentine fat-tailed mouse opossum

Argentine fat-tailed mouse opossum

Thylamys sponsorius

The Argentine fat-tailed mouse opossum (*Thylamys sponsorius*) was formerly considered a species opossum in the family Didelphidae. It is found in the eastern foothills of the Andes in northern Argentina and southern Bolivia. Its dorsal fur is...

Argentine hocicudo

Argentine hocicudo

Oxymycterus akodontius

The Argentine hocicudo (*Oxymycterus akodontius*) is a species of rodent in the family Cricetidae. It is found only in northern Argentina in yungas (eastern Andean mountain forest) habitat. The exact population is unknown as only three specimens...

Argentine swamp rat

Argentine swamp rat

Scapteromys aquaticus

The Argentine swamp rat (*Scapteromys aquaticus*) is a semiaquatic rodent species from South America. It is found in northeastern Argentina and Paraguay, where it lives in freshwater marshes and along the southern coast of the Río de la Plata...

Argentine tuco-tuco

Argentine tuco-tuco

Ctenomys argentinus

The Argentine tuco-tuco (*Ctenomys argentinus*) is a species of rodent in the family Ctenomyidae. It is endemic to Argentina.

Arianus's rat
Arianus's rat
Rattus omichlodes

Arianus's rat (*Rattus omichlodes*), also known as Arianus' New Guinea mountain rat, is a species of rat native to the mountains of Papua Province, Indonesia.

Arid thicket rat
Arid thicket rat
Grammomys aridulus

The arid thicket rat (*Grammomys aridulus*) is a species of rodent in the family Muridae. It is found only in Sudan. Its natural habitat is subtropical or tropical dry shrubland. This species is currently listed on the "Nearly Threatened" list in...

Arizona cotton rat
Arizona cotton rat
Sigmodon arizonae

The Arizona cotton rat or Colorado River cotton rat (*Sigmodon arizonae*) is a species of rodent in the family Cricetidae. It is found in Mexico and the United States.

Arizona gray squirrel
Arizona gray squirrel
Sciurus arizonensis

The Arizona gray squirrel (*Sciurus arizonensis*) is a tree squirrel, in the genus *Sciurus*, endemic to the canyons and valleys surrounded by deciduous and mixed forests in eastern Arizona and northern Mexico. It is threatened by habitat loss. The...

Arizona myotis
Arizona myotis
Myotis occultus

The Arizona myotis (*Myotis occultus*) is a vesper bat species inhabiting much of the southwestern United States and central Mexico as far south as the Distrito Federal.

Arizona pocket mouse
Arizona pocket mouse
Perognathus amplus

The Arizona pocket mouse (*Perognathus amplus*) is a rodent native to the Sonoran desert. It is a small mouse with a thinly furred tail that is smooth from base to tip (i.e. it has no tuft). In color it ranges from tan to orange. It is a...

Arizona shrew
Arizona shrew
Sorex arizonae

The Arizona shrew (*Sorex arizonae*) is a species of shrew native to North America.

Armbruster's wolf
Armbruster's wolf
Canis armbrusteri

Armbruster's wolf (*Canis armbrusteri*) is an extinct species that was endemic to North America and lived during the Irvingtonian stage of the Pleistocene epoch, spanning from 1.9 Mya–250,000 years BP. It is notable because it is proposed as the...

Armenian birch mouse
Armenian birch mouse
Sicista armenica

The Armenian birch mouse (*Sicista armenica*) is a species of rodent in the family Sminthidae.

Armenian Mouflon

Armenian Mouflon

Ovis gmelini gmelini

The Armenian mouflon (*Ovis gmelini gmelini*) is an endangered subspecies of mouflon. It was first described in 1840 by Edward Blyth, who equated it with the "Orientalische Schaaf" (Oriental sheep) described by Samuel Gottlieb Gmelin in 1774.

Armenian shrew

Armenian shrew

Crocidura armenica

The Armenian shrew (*Crocidura armenica*) is a species of mammal in the family Soricidae. It is endemic to Armenia.

Armenian whiskered bat

Armenian whiskered bat

Myotis hajastanicus

The Armenian whiskered bat (*Myotis hajastanicus*), also known as the Hajastan myotis or the Armenian myotis, is a species of bat from the family Vespertilionidae. The Armenian whiskered bat was formerly included as a part of the whiskered bat,...

Armored rat

Armored rat

Hoplomys gymnurus

The armored rat (*Hoplomys gymnurus*) is a species of rodent in the family Echimyidae. It is monotypic within the genus *Hoplomys*. It is found in Latin America, from northern Honduras to northwest Ecuador. It possesses a range of spines on its back...

Arnhem Land rock rat

Arnhem Land rock rat

Zyzomys maini

The Arnhem Land rock rat (*Zyzomys maini*) also known as the Arnhem rock-rat and by the Indigenous Australian name of kodjperre is a species of rodent in the family Muridae. It is found only in the Top End Region of the Northern Territory in Australia.

Arnhem leaf-nosed bat

Arnhem leaf-nosed bat

Hipposideros inornatus

The Arnhem leaf-nosed bat (*Hipposideros inornatus*) is a species of bat in the family Hipposideridae. It lives in the sandstone areas of Kakadu National Park (Northern Territory, Australia).

Arnhem sheath-tailed bat

Arnhem sheath-tailed bat

Taphozous kapalgensis

The Arnhem sheath-tailed bat, species *Taphozous kapalgensis*, is an emballonurid bat found at the Top End of Australia. The species is also referred to as the white-striped sheath-tail for the distinguishing marks at the flank, a feature observable...

Arnhold's mouse lemur

Arnhold's mouse lemur

Microcebus arnholdi

Arnhold's mouse lemur or the Montagne d'Ambre mouse lemur (*Microcebus arnholdi*) is a species of mouse lemur endemic to Madagascar. Its holotype was first collected on 27 November 2005, and was first described in 2008. According to genetic tests,...

Arnoux's Beaked Whale

Arnoux's Beaked Whale

Berardius arnuxii

Arnoux's beaked whale (*Berardius arnuxii*) is one of the species of beaked whales found in great tracts of the Southern Ocean. Little is known about their behavior due to their secretive nature and infrequent encounters with live individuals.

Arrow flying squirrel

Arrow flying squirrel

Hylopetes sagitta

The arrow flying squirrel (*Hylopetes sagitta*) is a species of flying squirrel. It is endemic to Java and Bangka, Indonesia. The population is unknown as it has only been collected from a few localities. It is nocturnal and arboreal and may be...

Arroyo of Paradise brucie

Arroyo of Paradise brucie

Bucepattersonius paradisus

The Arroyo of Paradise brucie (*Bucepattersonius paradisus*), also known as the Arroyo of Paradise akodont is a South American rodent species of the family Cricetidae. It is known only from northeastern Argentina. No population data is available...

Artibeus schwartzi

Artibeus schwartzi

Artibeus schwartzi

Artibeus schwartzi is a species of bat found in the Lesser Antilles. It was previously considered a subspecies of the Jamaican fruit bat, (*A. jamaicensis*). It has been hypothesized that it arose from hybridization of three *Artibeus* species:

A...

Aru flying fox

Aru flying fox

Pteropus aruensis

The Aru flying fox (*Pteropus aruensis*) is a Critically Endangered species of megabat found in the Aru Islands in Indonesia. It was described by Wilhelm Peters in 1867. It was formerly considered a subspecies of the black-bearded flying fox. The...

Arunachal macaque

Arunachal macaque

Macaca munzala

The Arunachal macaque (*Macaca munzala*) is a macaque native to Arunachal Pradesh in North-east India. It is listed as Endangered on the IUCN Red List. It was scientifically described in 2005. It is a relatively large brown primate with a...

Ash-colored Oldfield mouse

Ash-colored Oldfield mouse

Thomasomys cinereus

The ash-colored Oldfield mouse (*Thomasomys cinereus*) is a species of rodent in the family Cricetidae. It is found in Ecuador and Peru.

Ash-grey mouse

Ash-grey mouse

Pseudomys albocinereus

The ash-grey mouse (*Pseudomys albocinereus*) is a rodent in the family Muridae. Larger and more robust than *Mus musculus*, the common house mouse, it is found only in Southwest Australia.
Asháninka arboreal chinchilla rat
Asháninka arboreal chinchilla rat
Cuscomys ashaninka

The Asháninka arboreal chinchilla rat (*Cuscomys ashaninka*) is a large species of chinchilla rats from the Andes of far northern Cusco in Peru. It was first scientifically described in 1999. The first specimen of this species was discovered by...
Ashaninka Oldfield mouse
Ashaninka Oldfield mouse
Thomasomys onkiro

The Ashaninka Oldfield mouse (*Thomasomys onkiro*) is a species of rodent in the family Cricetidae. It is known only from a single locality (which includes Otishi National Park) in the Cordillera Oriental of the southern Peruvian Andes, in montane...
Ashy black titi
Ashy black titi
Plecturocebus cinerascens

The ashy black titi, *Plecturocebus cinerascens*, is a species of titi, a type of New World monkey, endemic to Brazil. It was originally described as *Callicebus cinerascens* in 1823.
Ashy chinchilla rat
Ashy chinchilla rat
Abrocoma cinerea

The ashy chinchilla rat (*Abrocoma cinerea*) is a species of chinchilla rats in the family Abrocomidae found in Argentina, Bolivia, Chile, and Peru.
Ashy roundleaf bat
Ashy roundleaf bat
Hipposideros cineraceus

The ashy roundleaf bat (*Hipposideros cineraceus*) is a species of bat in the family Hipposideridae found in Bhutan, Cambodia, India, Indonesia, Laos, Malaysia, Myanmar, Nepal, Pakistan, Thailand, and Vietnam.
Ashy-bellied Oldfield mouse
Ashy-bellied Oldfield mouse
Thomasomys cinereiventris

The ashy-bellied Oldfield mouse (*Thomasomys cinereiventris*) is a species of rodent in the family Cricetidae. It is found in Colombia and Ecuador.
Ashy-gray tube-nosed bat
Ashy-gray tube-nosed bat
Murina cineracea

The ashy-gray tube-nosed bat (*Murina cineracea*) is a species in the vesper bat family Vespertilionidae, found in southern Asia, including Pakistan, northern India, Burma, Thailand, Laos, and Vietnam. They have tube-shaped nostrils (hence the...
Ashy-headed flying fox
Ashy-headed flying fox
Pteropus caniceps

The ashy-headed flying fox or North Moluccan flying fox (*Pteropus caniceps*) is a species of bat in the family Pteropodidae. It is endemic to Indonesia.

Asia Minor ground squirrel
Asia Minor ground squirrel
Spermophilus xanthoprymnus

The Asia Minor ground squirrel (*Spermophilus xanthoprymnus*), also known as Anatolian souslik, Anatolian ground squirrel, is a species of rodent in the family Sciuridae. It is found in Armenia, Iran, and Turkey. The scientific name roughly...
Asia Minor spiny mouse
Asia Minor spiny mouse
Acomys cilicicus

The Asia Minor spiny mouse (*Acomys cilicicus*) is a species of rodent in the family Muridae.
Asian Badger
Asian Badger
Meles leucurus

The Asian badger (*Meles leucurus*), also known as the sand badger, is a species of badger native to Mongolia, China, Kazakhstan, Kyrgyzstan, the Korean Peninsula and Russia.
Asian barbastelle
Asian barbastelle
Barbastella leucomelas

The eastern barbastelle or Asian barbastelle (*Barbastella darjelingensis*) is a species of vesper bat found throughout much of Asia, from Afghanistan to Taiwan.
Asian Elephant
Asian Elephant
Elephas maximus

The Asian elephant (*Elephas maximus*) is one of only three living species of elephants or elephantids anywhere in the world, the others being the African bush elephant and African forest elephant. The Asian elephant is the largest living land...
Asian garden dormouse
Asian garden dormouse
Eliomys melanurus

The Asian garden dormouse or large-eared garden dormouse, (*Eliomys melanurus*) is a species of rodent in the family Gliridae. It is found in Egypt, Iraq, Israel, Jordan, Lebanon, Libya, Saudi Arabia, Syria and Turkey. Its natural habitats are...
Asian Golden Cat
Asian Golden Cat
Catopuma temminckii

The Asian golden cat (*Catopuma temminckii*) is a medium-sized wild cat native to Asia. In China, the Asian golden cat is thought to be a kind of leopard and is known as "rock cat" or "yellow leopard". Different color phases have different names;...
Asian gray shrew
Asian gray shrew
Crocidura attenuata

The Asian gray shrew (*Crocidura attenuata*) is a species of mammal in the family Soricidae. It is found in Bhutan, Cambodia, China, India, Laos, Malaysia, Myanmar, Nepal, Pakistan, the Philippines, Thailand, and Vietnam. It is one of the...
Asian highland shrew
Asian highland shrew
Suncus montanus

The Asian highland shrew (*Suncus montanus*) is a species of mammal in the family Soricidae. It is found in India and Sri Lanka. Its habitat is subtropical or tropical dry forests. It is threatened by habitat loss. It is known as ග්‍රී ලංකා කුඳු...

Asian House Shrew
Asian House Shrew
Suncus murinus

The Asian house shrew (*Suncus murinus*) is a shrew species native to South and Southeast Asia that has been listed as Least Concern on the IUCN Red List since 2008 because of its large population and wide distribution. It has been introduced in...

Asian lesser white-toothed shrew
Asian lesser white-toothed shrew
Crocidura shantungensis

The Asian lesser white-toothed shrew (*Crocidura shantungensis*) is a species of mammal in the family Soricidae. It is found in Siberia, China, Korea, and Japan.

Asian Lion
Asian Lion
Panthera leo persica

The Asian lion is a population of *Panthera leo leo* that today survives in the wild only in India. The first scientific description of the Asian lion was published in 1826 by the Austrian zoologist Johann N. Meyer, who named it *Felis leo persicus*....

Asian Palm Civet
Asian Palm Civet
Paradoxurus hermaphroditus

The Asian palm civet (*Paradoxurus hermaphroditus*) is a viverrid native to South and Southeast Asia. In Indonesia, this animal is threatened by poaching and illegal wildlife trade and buyers use it for the increasing production of kopi luwak, also...

Asian particolored bat
Asian particolored bat
Vespertilio sinensis

The Asian parti-colored bat (*Vespertilio sinensis*) is a species of parti-coloured bat. An adult Asian parti-colored bat has a body length of 6–7 cm, a tail of 4.3–4.5 cm, and a wing length of 5 cm. Asian parti-colored bats are distributed across...

Asian red-cheeked squirrel
Asian red-cheeked squirrel
Dremomys rufigenis

The Asian red-cheeked squirrel (*Dremomys rufigenis*) is a species of rodent in the family Sciuridae. It is found in south-eastern Asia.

Asian Wild Water Buffalo
Asian Wild Water Buffalo
Bubalus arnee

The Asian wild water buffalo (*Bubalus arnee*) is among the heaviest living wild bovid species. It is the most likely ancestor of the domestic water buffalo.

Asiatic Black Bear
Asiatic Black Bear
Ursus thibetanus

The Asian black bear (*Ursus thibetanus*) is a medium-sized bear species native to

Asia. It is unique among bears in that it is largely adapted to an arboreal lifestyle. It is listed as vulnerable on the IUCN Red List, and is threatened by...
Asiatic brush-tailed porcupine
Asiatic brush-tailed porcupine
Atherurus macrourus

The Asiatic brush-tailed porcupine (*Atherurus macrourus*) is a species of rodent in the family Hystricidae. It is found in China, Bhutan, India, Laos, Malaysia, Myanmar, Thailand, and Vietnam.
Asiatic Cheetah
Asiatic Cheetah
Acinonyx jubatus venaticus

The Asiatic cheetah (*Acinonyx jubatus venaticus*) is a critically endangered cheetah subspecies currently only surviving in Iran. It once occurred from the Arabian Peninsula and the Near East to the Caspian region, Transcaucasus, Kyzylkum Desert,...
Asiatic long-tailed climbing mouse
Asiatic long-tailed climbing mouse
Vandeleuria oleracea

The Asiatic long-tailed climbing mouse (*Vandeleuria oleracea*) is a species of rodent found in South and Southeast Asia. It is known as ගස් මීයා by Sinhalese people. It is known to spread the Ixodes tick-borne viral Kyasanur Forest Disease (KFD).
Asiatic short-tailed shrew
Asiatic short-tailed shrew
Blarinella quadraticauda

The Asiatic short-tailed shrew (*Blarinella quadraticauda*) is one of three species of shrew in the genus *Blarinella*. It is in the family Soricidae and is endemic to China.
Asiatic Wildcat
Asiatic Wildcat
Felis lybica ornata

The Asiatic wildcat (*Felis lybica ornata*) is an African wildcat subspecies. It is also known as the Asian steppe wildcat and Indian desert cat. There is no information on the current status or population numbers for the Asiatic wildcat's entire...
Assam Macaque
Assam Macaque
Macaca assamensis

The Assam macaque (*Macaca assamensis*) is a macaque of the Old World monkey family native to South and Southeast Asia. Since 2008, it has been listed as Near Threatened on the IUCN Red List as it is experiencing significant declines.
Assam mole shrew
Assam mole shrew
Anourosorex assamensis

The Assam mole shrew (*Anourosorex assamensis*) is a species of red-toothed shrew endemic to northeast India.
Astor markhor
Astor markhor
Capra falconeri falconeri

The Astor markhor or flare-horned markhor (*Capra falconeri falconeri*) is a subspecies of the markhor, native to Kashmir and northern Pakistan. To the west it

reaches the easternmost parts of Afghanistan. The range of the Astor markhor is very...

Atacama myotis

Atacama myotis

Myotis atacamensis

The Atacama myotis (*Myotis atacamensis*) is a species of vesper bat in the family Vespertilionidae. It is found in Chile and Peru, an example ecoregion of occurrence being the Chilean matorral.

Atherton antechinus

Atherton antechinus

Antechinus godmani

The Atherton antechinus (*Antechinus godmani*), also known as Godman's antechinus, is a species of small carnivorous marsupial native to Australia. It is one of the rarest members of its genus, and differs from other antechinuses in its more...

Atlantic bamboo rat

Atlantic bamboo rat

Kannabateomys amblyonyx

The Atlantic bamboo rat (*Kannabateomys amblyonyx*), or southern bamboo rat, is a spiny rat species from South America, found in humid tropical forests in Argentina, Brazil and Paraguay. It is the only member of the genus *Kannabateomys*.

Atlantic Forest climbing mouse

Atlantic Forest climbing mouse

Rhipidomys mastacalis

The Atlantic Forest climbing mouse (*Rhipidomys mastacalis*) is an arboreal rodent species in the family Cricetidae from South America. It is found in the Atlantic Forest of southeastern Brazil at elevations from sea level to 1500 m. Its karyotype...

Atlantic Forest hocicudo

Atlantic Forest hocicudo

Oxymycterus dasytrichus

The Atlantic Forest hocicudo (*Oxymycterus dasytrichus*) is a species of rodent in the family Cricetidae. It is found only in southeastern Brazil, where it lives in the Atlantic Forest and associated wetlands at altitudes from sea level to 800 m...

Atlantic humpback dolphin

Atlantic humpback dolphin

Sousa teuszii

The Atlantic humpback dolphin (*Sousa teuszii*) is a species of humpback dolphin that is found in coastal areas of West Africa. It is regarded as critically endangered by the IUCN.

Atlantic Spotted Dolphin

Atlantic Spotted Dolphin

Stenella frontalis

The Atlantic spotted dolphin (*Stenella frontalis*) is a dolphin found in warm temperate and tropical waters of the Atlantic Ocean. Older members of the species have a very distinctive spotted coloration all over their bodies.

Atlantic titi

Atlantic titi

Callicebus personatus

The Atlantic titi or masked titi (*Callicebus personatus*) is a species of titi, a type of New World monkey, endemic to Brazil.

Atlantic White-Sided Dolphin

Atlantic White-Sided Dolphin
Lagenorhynchus acutus

The Atlantic white-sided dolphin (*Lagenorhynchus acutus*) is a distinctively coloured dolphin found in the cool to temperate waters of the North Atlantic Ocean.
Atlas wild ass
Atlas wild ass
Equus africanus atlanticus

The Atlas wild ass (*Equus africanus atlanticus*), also known as Algerian wild ass, is a purported extinct subspecies of the African wild ass that was once found across North Africa and parts of the Sahara. It was last represented in a villa mural...
Attwater's pocket gopher
Attwater's pocket gopher
Geomys attwateri

Attwater's pocket gopher (*Geomys attwateri*) is a species of rodent in the family Geomyidae. It is endemic to the Texas Coastal Bend in the southcentral United States. This species was named in honor of English naturalist Henry Philemon Attwater.
Australasian bent-wing bat
Australasian bent-wing bat
Miniopterus orianae

The Australasian bent-wing bat (*Miniopterus orianae*) is a species of vesper bat in the family Miniopteridae. It is found in Australia and in Southeast Asia.
Australian Feral Camel
Australian Feral Camel
Camelus dromedarius

Australian feral camels are feral populations of Dromedary camels (*Camelus dromedarius*) in Australia. They were imported from British India and Afghanistan during the 19th century for transport and construction during the colonization of the...
Australian Fur Seal
Australian Fur Seal
Arctocephalus pusillus

The brown fur seal (*Arctocephalus pusillus*), also known as the Cape fur seal, South African fur seal and Australian fur seal, is a species of fur seal.
Australian humpback dolphin
Australian humpback dolphin
Sousa sahalensis

The Australian humpback dolphin (*Sousa sahalensis*) is a species of humpback dolphin and the fourth recognized humpback dolphin species chronologically. The specific name sahalensis is derived from the Sahul Shelf, located between northern...
Australian myotis
Australian myotis
Myotis australis

The Australian myotis (*Myotis australis*) is a species of vesper bat. It is found only in Australia. This taxa may not represent a valid species. Only one specimen has ever been documented, supposedly from New South Wales. This specimen may have...
Australian Sea Lion
Australian Sea Lion
Neophoca cinerea

The Australian sea lion (*Neophoca cinerea*), also known as the Australian sea-lion or Australian sealion, is a species of sea lion that is the only endemic pinniped in Australia. It is currently monotypic in the genus *Neophoca*, with the extinct...
Australian Snubfin Dolphin
Australian Snubfin Dolphin
Orcaella heinsohni

The Australian snubfin dolphin (*Orcaella heinsohni*) is a dolphin found off the northern coasts of Australia. It closely resembles the Irrawaddy dolphin (of the same genus, *Orcaella*) and was not described as a separate species until 2005.
The...
Australian Swamp Rat
Australian Swamp Rat
Rattus lutreolus

The Australian swamp rat (*Rattus lutreolus*) is a rodent that lives on the coasts of southern and eastern Australia. These rats are known for creating tunnels through the vegetation through which they can constantly move.
Austro-Hungarian wolf
Austro-Hungarian wolf
Canis lupus minor

The Austro-Hungarian wolf (*Canis lupus minor*), also called the reed wolf, is a proposed subspecies of the gray wolf that is thought to have once inhabited Hungary and eastern Austria. Its subspecies status is uncertain, as there are few records...
Awash multimammate mouse
Awash multimammate mouse
Mastomys awashensis

The Awash multimammate mouse or Awash mastomys (*Mastomys awashensis*) is a species of rodent in the family Muridae found only in Ethiopia. Phylogenetically the Awash multimammate mouse is the sister taxon of the Natal multimammate mouse (M...
Axis Deer
Axis Deer
Axis axis

The Axis deer (*Axis axis*) is a deer species native to the Indian subcontinent. It was first described and given a binomial name by German naturalist Johann Christian Polycarp Erxleben in 1777.
Aye-Aye
Aye-Aye
Daubentonia madagascariensis

The Aye-aye (*Daubentonia madagascariensis*) is a long-fingered lemur that inhabits the rainforests in Madagascar. It is the world's largest nocturnal primate and has a unique appearance. When first discovered, it was thought to be a type of large...
Azagnyi fruit bat
Azagnyi fruit bat
Megaloglossus azagnyi

The Azagnyi fruit bat (*Megaloglossus azagnyi*), also called the western Woermann's fruit bat, is a species of megabat found in Africa.
Azara's Agouti
Azara's Agouti
Dasyprocta azarae

Azara's agouti (*Dasyprocta azarae*) is a South American agouti species from the

family Dasyproctidae. Found in Brazil, Paraguay, and Argentina, it is named after Spanish naturalist Félix de Azara. The population is unknown and may have gone...
Azara's night monkey
Azara's night monkey
Aotus azarae

Azara's night monkey (*Aotus azarae*), also known as the southern night monkey, is a night monkey species from South America. It is found in Argentina, Bolivia, Brazil, Peru and Paraguay. The species is monogamous, with the males providing a large...
Azara's tuco-tuco
Azara's tuco-tuco
Ctenomys azarae

The Mendoza tuco-tuco (*Ctenomys mendocinus*) is a species of rodent in the family Ctenomyidae.
Azaras's capuchin
Azaras's capuchin
Sapajus cay

Azaras's capuchin or hooded capuchin (*Sapajus cay*) is a species of robust capuchin. It occurs in eastern Paraguay, southeastern Bolivia, northern Argentina, and Brazil, at Mato Grosso do Sul and Mato Grosso states, in Pantanal. Its habitat...
Azores noctule
Azores noctule
Nyctalus azoreum

The Azores noctule (*Nyctalus azoreum*) is a species of bat found in the dry forests of the Azores. It is the only species of mammal endemic to the Azores. It has been recorded on most of the islands of the Azores, and remains common on some but...
Aztec fruit-eating bat
Aztec fruit-eating bat
Dermanura azteca

The Aztec fruit-eating bat (*Dermanura azteca*) is a species of bat in the family Phyllostomidae. It is found in Costa Rica, El Salvador, Guatemala, Honduras, Mexico, and Panama.
Aztec mastiff bat
Aztec mastiff bat
Molossus aztecus

The Aztec mastiff bat (*Molossus aztecus*) is a species of bat in the family Molossidae. It is found from Jalisco and Cozumel Island in southern Mexico to Nicaragua (it has been reported from Guatemala but not from El Salvador or Honduras) in a...
Aztec mouse
Aztec mouse
Peromyscus aztecus

The Aztec mouse (*Peromyscus aztecus*) is a species of rodent in the family Cricetidae, native to southern Mexico and parts of Central America.
Azuero howler
Azuero howler
Alouatta coibensis trabeata

The Azuero howler (*Alouatta coibensis trabeata*) a type of monkey that is a subspecies of the Coiba Island howler *A. coibensis*. This subspecies is endemic to the Azuero Peninsula in Panama. The Azuero howler is distinguished primarily by its...

Azuero spider monkey
Azuero spider monkey
Ateles geoffroyi azuerensis

The Azuero spider monkey (*Ateles geoffroyi azuerensis*) is a possible subspecies of spider monkey that is in critical danger of extinction according to the IUCN Red List of Threatened Species. Common names of this subspecies include mono charro,...

Azumi shrew
Azumi shrew
Sorex hosonoi

The Azumi shrew (*Sorex hosonoi*) is a species of mammal in the family Soricidae. It is endemic to Japan, where it is found in the mountainous regions of central Honshu Island. It is a close relative of the Eurasian least shrew (*Sorex...*

Babault's mouse shrew
Babault's mouse shrew
Myosorex babaulti

Babault's mouse shrew (*Myosorex babaulti*) is a species of mammal in the family Soricidae found in Burundi, the Democratic Republic of the Congo, and Uganda. Its natural habitat is subtropical or tropical moist montane forests. It is threatened...

Babyrousa bolabatuensis
Babyrousa bolabatuensis
Babyrousa bolabatuensis

Babyrousa bolabatuensis, the Bola Batu babirusa, is a species of babirusa from the Indonesian island of Sulawesi. It was first described in 1950 as a subspecies of *Babyrousa babyrussa*, then the only recognized species of babirusa, and raised to...

Back-Striped Weasel
Back-Striped Weasel
Mustela strigidorsa

The Back-striped weasel (*Mustela strigidorsa*) is a small mammal widely distributed in Southeastern Asia. It is listed as Least Concern on the IUCN Red List because of its presumed large population, occurrence in many protected areas, apparent...

Bactrian Camel
Bactrian Camel
Camelus bactrianus

The Bactrian camel (*Camelus bactrianus*) is a large even-toed ungulate native to the steppes of Central Asia. It exists mainly in the domesticated form. Their name comes from the ancient historical region of Bactria. Domesticated Bactrian camels...

Bactrian Deer
Bactrian Deer
Cervus elaphus bactrianus

The Bactrian deer (*Cervus hanglu bactrianus*), also called the Bukhara deer, Bokhara deer, or Bactrian wapiti, is a lowland subspecies of Central Asian red deer native to Central Asia. It is similar in ecology to the related Yarkand deer (*C. h....*

Badlands bighorn
Badlands bighorn
Ovis canadensis auduboni

The Badlands bighorn, *Ovis canadensis auduboni*, also commonly known as Audubon's bighorn sheep, is an extinct subspecies of bighorn sheep of the northern Great Plains in North America. Its existence as a separate subspecies is disputed.

Baer's wood mouse
Baer's wood mouse
Hylomyscus baeri

Baer's hylomyscus or Baer's wood mouse (*Hylomyscus baeri*) is a species of rodent in the family Muridae. It is found in Ivory Coast, Ghana, and Sierra Leone. Its natural habitat is subtropical or tropical moist lowland forests. It is threatened by...

Baffin Island wolf
Baffin Island wolf
Canis lupus manningi

The Baffin Island wolf (*Canis lupus manningi*), also known as the Baffin Island tundra wolf, is a subspecies of grey wolf which resides exclusively on Baffin Island and several nearby islands. It was not formally recognized as a subspecies until...

Bagobo rat
Bagobo rat
Bullimus bagobus

The Bagobo rat (*Bullimus bagobus*) is one of three species of rodent in the genus *Bullimus*, and in the diverse family Muridae. It is found in the Visayas and Mindanao, Philippines.

Bahaman funnel-eared bat
Bahaman funnel-eared bat
Chilonatalus tumidifrons

The Bahaman funnel-eared bat (*Chilonatalus tumidifrons*) is a species of bat in the family Natalidae. This bat is endemic to the Bahamas, in the northeastern Caribbean. It was first discovered by Miller in 1903. Their category list and criteria is...

Bahamian hutia
Bahamian hutia
Geocapromys ingrahami

The Bahamian hutia or Ingraham's hutia (*Geocapromys ingrahami*) is a species of hutia in the subfamily Capromyinae native to the Bahamas. Its natural habitats are subtropical or tropical moist lowland forest, subtropical or tropical dry...

Bahamian raccoon
Bahamian raccoon
Procyon lotor maynardi

The Bahamian raccoon (*Procyon lotor maynardi*), also called Bahama raccoon or Bahamas raccoon, is a subspecies of the common raccoon endemic on the New Providence Island in the Bahamas. The binomial name, *maynardi*, comes from Charles Johnson...

Bahia porcupine
Bahia porcupine
Coendou insidiosus

The Bahia porcupine, *Coendou insidiosus*, is a New World porcupine species in the family Erethizontidae endemic to the Atlantic Forest of southeastern Brazil. It was formerly sometimes assigned to *Sphiggurus*, a genus no longer recognized since...

Baiji
Baiji
Lipotes vexillifer

The baiji (*Lipotes vexillifer*) is a possibly extinct species of freshwater dolphin. It is thought to be the first dolphin species driven to extinction due to the impact of humans. This dolphin is listed as "critically endangered: possibly..."

Baikal Seal
Baikal Seal
Pusa sibirica

Baikal seals (*Pusa sibirica*) are tiny seals that are members of the family Phocidae. They are native to Siberia and live only within Lake Baikal, from where their common name comes. They are the only type of freshwater seal type and it is still...

Bailey's pocket mouse
Bailey's pocket mouse
Chaetodipus baileyi

Bailey's pocket mouse (*Chaetodipus baileyi*) is a species of rodent of the subfamily Perognathinae, family Heteromyidae. It is found in Baja California, Sinaloa and Sonora in Mexico and in California, Arizona and New Mexico in the United States.

Bailey's shrew
Bailey's shrew
Crocidura baileyi

Bailey's shrew (*Crocidura baileyi*) is a species of mammal in the family Soricidae. The name honours American naturalist and museum director Alfred Marshall Bailey.

Baird's Beaked Whale
Baird's Beaked Whale
Berardius bairdii

Four-toothed whales or giant beaked whales are beaked whales in the genus *Berardius*. They include Arnoux's beaked whale (*Berardius arnuxii*) in cold Southern Hemisphere waters, and Baird's beaked whale (*Berardius bairdii*) in the cold temperate...

Baird's pocket gopher
Baird's pocket gopher
Geomys breviceps

Baird's pocket gopher or the Louisiana pocket gopher (*Geomys breviceps*) is a species of pocket gopher that is native to the southern United States. In total, there are three almost identical species of eastern pocket gopher; *Geomys attwateri*, *G....*

Baird's shrew
Baird's shrew
Sorex bairdi

Baird's shrew (*Sorex bairdi*) is a species of mammal in the family Soricidae. It is endemic to northwest Oregon. Baird's shrew inhabits moist conifer forests. Its fur is darker brown in winter than in summer, when it is brownish-chestnut or olive...

Baird's Tapir
Baird's Tapir
Tapirus bairdii

The Baird's tapir (*Tapirus bairdii*), also known as the Central American tapir, is a species of tapir native to Mexico, Central America, and northwestern South America. It is the largest of the three species of tapir native to the Americas, as...

Baja California pronghorn
Baja California pronghorn
Antilocapra americana peninsularis

The Baja California pronghorn or peninsular pronghorn (*Antilocapra americana peninsularis*) is a subspecies of pronghorn, endemic to Baja California in Mexico.

The wild population is estimated at 200.
Baja California rock squirrel
Baja California rock squirrel
Otospermophilus atricapillus

The Baja California rock squirrel (*Otospermophilus atricapillus*) is a species of rodent in the family Sciuridae. It is endemic to Baja California, Mexico.

Baja pocket mouse
Baja pocket mouse
Chaetodipus rudinoris

The Baja pocket mouse (*Chaetodipus rudinoris*) is a species of rodent in the family Heteromyidae. The species occurs in southern California, Baja California and on islands in the Gulf of California.

Baker's small-toothed harvest mouse
Baker's small-toothed harvest mouse
Reithrodontomys bakeri

Baker's small-toothed harvest mouse (*Reithrodontomys bakeri*) is a species of rodent in the family Cricetidae. It is one of about 20 species of the genus *Reithrodontomys*. This species was discovered in 2004 and is found in Central Mexico.

Bala tube-nosed bat
Bala tube-nosed bat
Murina balaensis

The Bala tube-nosed bat (*Murina balaensis*) is a critically endangered species of bat found in Thailand.

Balaenoptera bertae
Balaenoptera bertae
Balaenoptera bertae

Balaenoptera bertae is an extinct species of baleen whale that lived from 3.35 to 2.5 Mya during the Pliocene in the region of today's San Francisco Bay Area. It held, also during the early to middle Neogene, a diverse assembly of cetaceans....

Bald Uakari
Bald Uakari
Cacajao calvus

The bald uakari (*Cacajao calvus*) or bald-headed uakari is a small New World monkey characterized by a very short tail; bright, crimson face; a bald head; and long coat. The bald uakari is restricted to várzea forests and other wooded habitats...

Bale Mountains vervet
Bale Mountains vervet
Chlorocebus djamdjamensis

The Bale Mountains vervet (*Chlorocebus djamdjamensis*) is a terrestrial Old World monkey endemic to Ethiopia, found in the bamboo forests of the Bale Mountains. All species in *Chlorocebus* were formerly in the genus *Cercopithecus*. The Bale...

Bale shrew
Bale shrew
Crocidura bottegoides

The Bale shrew (*Crocidura bottegoides*) is a species of mammal in the family Soricidae. It is endemic to Ethiopia, in the Bale Mountains of the Ethiopian Highlands. Its natural habitats are subtropical or tropical moist montane forests and...

Bali cattle
Bali cattle
Bos javanicus domesticus

The Bali cattle (*Bos javanicus domesticus*), also known as Balinese cattle, domestic banteng or Bali banteng, are a domesticated species of bovine which originated from the banteng (*Bos javanicus*). Bali cattle are an important source of meat and...

Balikun jerboa

Balikun jerboa

Allactaga balikunica

The Balikun jerboa (*Allactaga balikunica*) is a species of rodent in the family Dipodidae. It is found in arid areas of northwestern China and Mongolia. It eats green plants, plant roots, seeds, grasshoppers and beetles.

Balkan Lynx

Balkan Lynx

Lynx lynx balcanicus

The Balkan lynx (*Lynx lynx balcanicus*) is a subspecies of the Eurasian lynx. It is considered a national symbol in North Macedonia and appears on the Macedonian five-denar coin. This cat is considered to be the largest cat in the Balkans. It has...

Balkan mole

Balkan mole

Talpa stankovici

The Balkan mole (*Talpa stankovici*) is a species of mammal in the family Talpidae. It is found in Montenegro, Albania, North Macedonia and Greece, including the island of Corfu.

Balkan mole-rat

Balkan mole-rat

Spalax graecus

The Bukovina blind mole-rat or (erroneously) the Balkan blind mole-rat (*Spalax graecus*) is a species of rodent in the family Spalacidae found in Romania, Moldova, and Ukraine.

Balkan snow vole

Balkan snow vole

Dinaromys bogdanovi

The Balkan snow vole (*Dinaromys bogdanovi*), also known as Martino's snow vole, is the only member of the genus *Dinaromys*. Eight subspecies of this vole have been recognized from southern parts of Europe. The genus name means "Dinaric mouse",...

Balochistan black bear

Balochistan black bear

Ursus thibetanus gedrosianus

The Balochistan (or Baluchistan) black bear (*Ursus thibetanus gedrosianus*) is a subspecies of the Asian black bear occurring in the Balochistan Mountains of southern Pakistan and Iran. It has an unusually thin coat for an Asian black bear, but...

Balochistan forest dormouse

Balochistan forest dormouse

Dryomys niethammeri

The Balochistan forest dormouse (*Dryomys niethammeri*) is a species of rodent in the family Gliridae. It is native to Pakistan. It is an endangered animal

Balochistan gerbil

Balochistan gerbil

Gerbillus nanus

The Balochistan gerbil or dwarf gerbil (*Gerbillus nanus*), is distributed mainly

from Morocco across north Africa, the Arabian Peninsula, the Middle East, and western Asia. This is a common species with a wide distribution which faces no obvious...

Baluchi mouse-like hamster
Baluchi mouse-like hamster
Calomyscus baluchi

The Baluchi mouse-like hamster (*Calomyscus baluchi*) is a species of rodent in the family Calomyscidae. It is found in Afghanistan and Pakistan.

Baluchistan Pygmy Jerboa
Baluchistan Pygmy Jerboa
Salpingotulus michaelis

The Baluchistan pygmy jerboa (*Salpingotulus michaelis*) is a very small rodent in the family Dipodidae. In the 1999 Guinness Book of Records, it was listed as tied for the smallest rodent in the world with the African pygmy mouse.

Banahao shrew-rat
Banahao shrew-rat
Rhynchomys banahao

The Banahao shrew-rat (*Rhynchomys banahao*) is a species of rodent in the genus *Rhynchomys*. It was described in 2007.

Banahaw tree mouse
Banahaw tree mouse
Musseromys gulantang

Banahaw tree mouse (*Musseromys gulantang*) is a species of rodent in the family of Muridae. It is named after Mount Banahaw in Luzon, Philippines.

Banana bat
Banana bat
Musonycteris harrisoni

The banana bat (*Musonycteris harrisoni*) is an endangered species of bat in the family Phyllostomidae. It is also commonly known as the trumpet-nosed bat or the Colima long-nosed bat.

Banana climbing mouse
Banana climbing mouse
Dendromus messorius

The banana climbing mouse (*Dendromus messorius*) is a species of rodent in the family Nesomyidae. It is found in Benin, Cameroon, Democratic Republic of the Congo, Nigeria, and Togo. Its natural habitat is subtropical or tropical dry lowland...

Banana pipistrelle
Banana pipistrelle
Neoromicia nana

The banana serotine (*Afronycteris nanus*), formerly known as the banana pipistrelle, is a species of vesper bat found throughout much of Africa. It was previously known as *Pipistrellus nanus*, but genetic analysis later reclassified it in...

Banded Hare-Wallaby
Banded Hare-Wallaby
Lagostrophus fasciatus

The banded hare-wallaby, mernine, or munning (*Lagostrophus fasciatus*) is a marsupial currently found on the Islands of Bernier and Dorre off western Australia. Reintroduced populations have recently been established on islands and fenced...

Banded Linsang
Banded Linsang
Prionodon linsang

The Banded linsang (*Prionodon linsang*) is a carnivorous mammal native to the Sundaic region of Southeast Asia. The name of these animals originated in the Javanese language as linsang or wlinsang, and previously, was translated incorrectly in...

Banded Mongoose
Banded Mongoose
Mungos mungo

The Banded mongoose (*Mungos mungo*) is a mongoose species native from the Sahel to Southern Africa. While most mongoose species live solitary lives, the Banded mongoose lives in colonies with a complex social structure and forages in groups.

Banded Palm Civet
Banded Palm Civet
Hemigalus derbyanus

The Banded palm civet (*Hemigalus derbyanus*) is a viverrid found in Southeast Asia. It is listed as Near Threatened on the IUCN Red List because of its large geographic and elevation range and tolerance to some habitat disturbance.

Banded Pig
Banded Pig
Sus scrofa vittatus

The Banded pig (*Sus scrofa vittatus*) is a subspecies of Wild boar found in Southeast Asia. It is known as the Wild boar in Singapore.

Banded surili
Banded surili
Presbytis femoralis

The Raffles' banded langur (*Presbytis femoralis*), also known as the banded leaf monkey or banded surili, is a species of primate in the family Cercopithecidae. It is endemic to Singapore and southern Peninsular Malaysia. The species underwent...

Banggai cuscus
Banggai cuscus
Strigocuscus pelengensis

The Banggai cuscus (*Strigocuscus pelengensis*) is a species of marsupial in the family Phalangeridae. It is endemic to Indonesia. It is a species of cuscus, a type of possum. It is found in the Peleng and Sula Islands to the east of Sulawesi in...

Bangs's mountain squirrel
Bangs's mountain squirrel
Syntheosciurus brochus

Bangs's mountain squirrel (*Syntheosciurus brochus*) is a poorly known species of tree squirrel, that only lives in Costa Rica and Panama. It can be found in mountain rain forests at an altitude between 1,900 and 2,600 metres (6,200 and 8,500 ft),...

Bangweulu tsessebe
Bangweulu tsessebe
Damaliscus lunatus superstes

The Bangweulu tsessebe (*Damaliscus lunatus superstes*) is a population and possible taxon of *Damaliscus lunatus*] (topi or sassaby), which are large African antelopes of the grasslands. This population is presently restricted to northern Zambia in...

Bank Vole
Bank Vole

Myodes glareolus

The bank vole (*Myodes glareolus*) is a small vole with red-brown fur and some grey patches, with a tail about half as long as its body. A rodent, it lives in woodland areas and is around 100 millimetres (3.9 in) in length. The bank vole is found...

Banka shrew

Banka shrew

Crocidura vosmaeri

The Banka shrew (*Crocidura vosmaeri*) is a species of mammal in the family Soricidae. It is only known from the Bangka Island in Indonesia and possibly Sumatra. It lives in primary and secondary lowland forest and it is not clear if can adapt to...

Banks flying fox

Banks flying fox

Pteropus fundatus

The Banks flying fox (*Pteropus fundatus*) is a species of megabat in the family Pteropodidae. It is endemic to Vanuatu. Its natural habitats are subtropical or tropical dry forests and subtropical or tropical swamps. These small fruit bats are...

Banner-tailed kangaroo rat

Banner-tailed kangaroo rat

Dipodomys spectabilis

The banner-tailed kangaroo rat (*Dipodomys spectabilis*) is a species of rodent in the family Heteromyidae. It is found in arid environments in the southwestern United States and Mexico where it lives in a burrow by day and forages for seeds and...

Bannister's rat

Bannister's rat

Melomys bannisteri

Bannister's rat, or great Key Island mosaic-tailed rat, or great Key Island melomys or great Kai Island melomys (*Melomys bannisteri*) is a species of rodent of the genus *Melomys*. It is endemic to the Kai Besar, one of the islands which are part...

Banteng

Banteng

Bos javanicus

The banteng (*Bos javanicus*) is a species of cattle found in Southeast Asia. Banteng occur in a variety of habitats throughout their range and can survive without water for long periods during droughts. The wild banteng is classified as Endangered...

Baoule's mouse

Baoule's mouse

Mus baoulei

Baoule's mouse (*Mus baoulei*) is a species of rodent in the family Muridae. It is found in Benin, Ivory Coast, Guinea, possibly Ghana, and possibly Sierra Leone. Its natural habitat is dry savanna.

Baptista Lake titi

Baptista Lake titi

Plecturocebus baptista

The Baptista Lake titi, *Plecturocebus baptista*, is a species of titi, a type of New World monkey, endemic to Brazil. It was originally described as *Callicebus baptista* in 1939.

Barasingha

Barasingha

Rucervus duvaucelii

Barasingha is a deer species with conspicuously large antlers. Overall, this mammal has as much as 12 antlers. In fact, the name of this species has Hindi origin and means '12-antlered deer'. Unfortunately, Barasingha is nowadays among the most...

Barbados raccoon

Barbados raccoon

Procyon lotor gloveralleni

The Barbados raccoon (*Procyon lotor gloveralleni*) is an extinct subspecies of the common raccoon (*Procyon lotor*), that was endemic on Barbados in the Lesser Antilles until 1964.

Barbara Brown's titi

Barbara Brown's titi

Callicebus barbarabrownae

Barbara Brown's titi (*Callicebus barbarabrownae*), also popularly known as the blond titi monkey or northern Bahian blond titi, is a species of titi, a type of New World monkey. This critically endangered species is endemic to the Caatinga in...

Barbary Ground Squirrel

Barbary Ground Squirrel

Atlantoxerus getulus

The Barbary ground squirrel (*Atlantoxerus getulus*) is a small rodent native to North Africa and has been introduced into the Canary Islands. They live in open areas in burrows and form colonies similar to North American prairie dogs, and have...

Barbary Macaque

Barbary Macaque

Macaca sylvanus

The unique Barbary macaques (*Macaca sylvanus*) are the only primates, occurring north of the Sahara in Africa. In addition, these animals are the only macaques, found outside of Asia. As highly adaptable creatures, they are capable of surviving in...

Barbary Sheep

Barbary Sheep

Ammotragus lervia

The Barbary sheep (*Ammotragus lervia*) is a species of caprine native to rocky mountains in North Africa. Although it is rare in its native North Africa, it has been introduced to North America, southern Europe, and elsewhere. Aoudad is another...

Barbary stag

Barbary stag

Cervus elaphus barbarus

The Barbary stag (*Cervus elaphus barbarus*), also known as the Atlas deer or African elk, is a subspecies of the red deer that is native to North Africa. It is the only deer known to be native to Africa, aside from *Megaceroides algericus*, which...

Barbary striped grass mouse

Barbary striped grass mouse

Lemniscomys barbarus

The Barbary striped grass mouse (*Lemniscomys barbarus*) is a small rodent of the suborder Myomorpha. This monotypic species is native to coastal Morocco, Algeria and Tunisia in northwest Africa. In the past it was believed to also occur...

Barbour's rock mouse
Barbour's rock mouse
Petromyscus barbouri

Barbour's rock mouse (*Petromyscus barbouri*) is a species of rodent in the family Nesomyidae. It is found only in South Africa. Its natural habitats are subtropical or tropical dry shrubland and rocky areas. It is threatened by habitat loss.

Barbour's vlei rat
Barbour's vlei rat
Otomys barbouri

Barbour's vlei rat (*Otomys barbouri*) is a species of rodent in the family Muridae. It is found in Kenya and Uganda. Its natural habitat is subtropical or tropical high-altitude shrubland. It is threatened by habitat loss.

Bare-backed fruit bat
Bare-backed fruit bat
Dobsonia moluccensis

The bare-backed fruit bat or Moluccan naked-backed fruit bat (*Dobsonia moluccensis*) is a fruit bat in the family Pteropodidae.

Bare-backed rousette
Bare-backed rousette
Rousettus spinalatus

The bare-backed rousette (*Rousettus spinalatus*) is a species of megabat.

Bare-bellied hedgehog
Bare-bellied hedgehog
Paraechinus nudiventris

The bare-bellied hedgehog (*Paraechinus nudiventris*), also known as the Madras hedgehog, is a species of hedgehog that is endemic to dry arid regions and scrubby jungles in southeastern India. As it was believed to be rare, it was formerly listed...

Bare-eared squirrel monkey
Bare-eared squirrel monkey
Saimiri ustus

The bare-eared squirrel monkey (*Saimiri ustus*) is a squirrel monkey endemic to Brazil and possibly eastern Bolivia.

Bare-tailed armored tree-rat
Bare-tailed armored tree-rat
Pattonomys occasius

The bare-tailed armored tree-rat (*Pattonomys occasius*) is a species of arboreal rodent in the family Echimyidae. It is found in lowland tropical rainforest east of the Andes in Ecuador and Peru.

Bare-tailed woolly mouse opossum
Bare-tailed woolly mouse opossum
Marmosa regina

The bare-tailed woolly mouse opossum (*Marmosa regina*) or short-furred woolly mouse opossum is a South American marsupial of the family Didelphidae. Its range includes Brazil, Colombia, Ecuador, Peru, and Bolivia. It is found in tropical...

Bare-Tailed Woolly Opossum
Bare-Tailed Woolly Opossum
Caluromys philander

The Bare-tailed woolly opossum (*Caluromys philander*) is a marsupial from South America. It was first described by Swedish zoologist Carl Linnaeus in 1758. This

species is also called mucura-xixica in Portuguese, zarigüeya lanuda parda in Spanish,...

Barren ground shrew
Barren ground shrew
Sorex ugyunak

The barren ground shrew (*Sorex ugyunak*) is a small shrew found in northern Canada west of Hudson Bay and in Alaska. At one time, this species was considered to be a subspecies of the similar cinereus shrew (*S. cinereus*). It is similar in...

Barren-ground caribou
Barren-ground caribou
Rangifer tarandus groenlandicus

The barren-ground caribou (*Rangifer tarandus groenlandicus*) is a subspecies of the reindeer (or the caribou in North America) that is found mainly in the Canadian territories of Nunavut and the Northwest Territories, as well as in Kitaa,...

Bartel's flying squirrel
Bartel's flying squirrel
Hylopetes bartelsi

The Bartel's flying squirrel (*Hylopetes bartelsi*) is a rodent species in the family Sciuridae endemic to West Java, Indonesia. It inhabits subtropical and tropical forest.

Bartels's rat
Bartels's rat
Sundamys maxi

Bartels's rat (*Sundamys maxi*) is a species of rodent in the family Muridae. It is found only in West Java, Indonesia, including in Gunung Gede Pangrango National Park.

Bartels's spiny rat
Bartels's spiny rat
Maxomys bartelsii

Bartels's spiny rat (*Maxomys bartelsii*) is a species of rodent in the family Muridae. It is found only in Indonesia. This species is indigenous to the volcano forests of west and central Java, the main island of Indonesia. The species is able to...

Basilan flying squirrel
Basilan flying squirrel
Petinomys crinitus

The Basilan flying squirrel (Philippines flying squirrel or Mindanao flying squirrel) (*Petinomys crinitus*) is a species of rodent in the family Sciuridae. It is endemic to the Philippines. It is often confused with mindanao flying squirrel...

Bastard big-footed mouse
Bastard big-footed mouse
Macrotarsomys bastardi

The bastard big-footed mouse (*Macrotarsomys bastardi*) is a species of rodent in the family Nesomyidae. It is found only in Madagascar. *Macrotarsomys bastardi* is the smallest-bodied of the three species recognized within the genus *Macrotarsomys*....

Bat-Eared Fox
Bat-Eared Fox
Otocyon megalotis

The Bat-eared fox (*Otocyon megalotis*) is a species of fox that lives in the African

savanna. Fossil records indicate this canid first appeared during the middle Pleistocene. It is named for its large ears, which have a role in thermoregulation....

Batak shrew

Batak shrew

Crocidura batakorum

The Batak shrew (*Crocidura batakorum*) is a species of shrew from the Philippines.

Bates's pygmy antelope

Bates's pygmy antelope

Nesotragus batesi

Bates's pygmy antelope (*Nesotragus batesi*), also known as the dwarf antelope, pygmy antelope or Bates' dwarf antelope, is a very small antelope living in the moist forest and brush of Central and West Africa. It is in the same genus as the suni.

Bates's shrew

Bates's shrew

Crocidura batesi

Bates's shrew (*Crocidura batesi*) is a species of mammal in the family Soricidae. It is found in Cameroon, Republic of the Congo, Equatorial Guinea, and Gabon. Its natural habitat is subtropical or tropical moist lowland forests. This large black...

Bates's slit-faced bat

Bates's slit-faced bat

Nycteris arge

Bate's slit-faced bat (*Nycteris arge*) is a species of slit-faced bat frequently confused with *Nycteris major*. It is broadly distributed and common, living throughout many parts of Africa in forests and savannas.

Batomys uragon

Batomys uragon

Batomys uragon

Batomys uragon (proposed common name Mount Isarog hairy-tailed rat) is a species of cloud rat endemic to the Philippines. It was first discovered in 1988, and the full description was published in 2015, in the Proceedings of the Biological...

Bavarian pine vole

Bavarian pine vole

Microtus bavaricus

The Bavarian pine vole (*Microtus bavaricus*) is a vole from the Austrian, Italian, and Bavarian Alps of Europe. It lives in moist meadows at elevations of 600 to 1,000 meters. There are 23 museum specimens of this species. This rodent was...

Bawean Deer

Bawean Deer

Hyelaphus kuhlii

The Bawean deer (*Axis kuhlii*) is a highly threatened species of deer native to Indonesia. Due to ongoing habitat loss, small population size, and limited range, the Bawean deer is evaluated as critically endangered on the IUCN Red List of...

Bay Cat

Bay Cat

Catopuma badia

The Bay cat (*Catopuma badia*) is one of the world's rarest and least-studied cats. These cats are so elusive that it was more than a century before researchers were able to study a live cat in detail. They have striking, rust-red colored fur,

with...

Bay Duiker

Bay Duiker

Cephalophus dorsalis

The Bay duiker (*Cephalophus dorsalis*) is a forest-dwelling antelope native to western and southern Africa. It was first described by British zoologist John Edward Gray in 1846. A notable feature of this duiker is the well-pronounced solid stripe...

Beach vole

Beach vole

Microtus breweri

The beach vole or Muskeget vole (*Microtus breweri*) is a rodent in the family Cricetidae endemic to Muskeget Island, Massachusetts. It is a close relative of the eastern meadow vole.

Beaded wood mouse

Beaded wood mouse

Hylomyscus aeta

The beaded hylomyscus or beaded wood mouse (*Hylomyscus aeta*) is a species of rodent in the family Muridae. It is found in Burundi, Cameroon, Central African Republic, Republic of the Congo, Democratic Republic of the Congo, Equatorial Guinea,...

Beady-eyed mouse

Beady-eyed mouse

Thomasomys baeops

The beady-eyed mouse (*Thomasomys baeops*) is a species of rodent in the family Cricetidae. It is found only in Ecuador.

Bearded emperor tamarin

Bearded emperor tamarin

Saguinus imperator subgriseus

The bearded emperor tamarin (*Saguinus imperator subgriseus*) is one of the two subspecies of the emperor tamarin. It is similar to the black-chinned emperor tamarin, with the beard. It usually lives in groups of 3-8, but can be found alone.

Bearded Seal

Bearded Seal

Erignathus barbatus

Bearded seals (*Erignathus barbatus*) are known as 'ice seals'. They are members of a group of four Arctic seal species that live in Alaskan waters and use sea ice for feeding, resting, and pupping. Bearded seals get their generic name from two...

Beatrice's bat

Beatrice's bat

Glauconycteris beatrix

Beatrice's bat (*Glauconycteris beatrix*) is a species of vesper bat in the family Vespertilionidae. It can be found in the following countries: Angola, Central African Republic, Democratic Republic of the Congo, Ivory Coast, Equatorial Guinea,...

Beaufort's naked-backed fruit bat

Beaufort's naked-backed fruit bat

Dobsonia beauforti

Beaufort's naked-backed fruit bat (*Dobsonia beauforti*) is a species of megabat in the family Pteropodidae. Its natural habitat is in caves.

Beccari's margareta rat

Beccari's margareta rat
Margaretamys beccarii

Beccari's margareta rat (*Margaretamys beccarii*) is a species of rodent in the family Muridae. It is endemic to the island of Sulawesi in Indonesia.
Beccari's sheath-tailed bat
Beccari's sheath-tailed bat
Emballonura beccarii

Beccari's sheath-tailed bat (*Emballonura beccarii*) is a species of sac-winged bat in the family Emballonuridae. It is found in New Guinea and in some nearby islands in both Indonesia and Papua New Guinea.
Beccari's shrew
Beccari's shrew
Crocidura beccarii

Beccari's shrew (*Crocidura beccarii*) is a species of mammal in the family Soricidae. It is found in the Barisan Mountains of Sumatra. It is threatened by habitat loss.
Bechstein's bat
Bechstein's bat
Myotis bechsteinii

Bechstein's bat (*Myotis bechsteinii*) is a species of vesper bat found in Europe and western Asia, living in extensive areas of woodland.
Beech Marten
Beech Marten
Martes foina

The Beech marten (*Martes foina*) is a species of marten native to much of Europe and Central Asia, though it has established a feral population in North America. It is superficially similar to the European pine marten but differs from it by its...
Beecroft's flying squirrel
Beecroft's flying squirrel
Anomalurus beecrofti

Beecroft's flying squirrel (*Anomalurus beecrofti*) or Beecroft's scaly-tailed squirrel, is a species of rodent in the family Anomaluridae. Some authorities consider it to be monotypic within the genus *Anomalurops*. Its natural habitats are...
Beelzebub's tube-nosed bat
Beelzebub's tube-nosed bat
Murina beelzebub

Beelzebub's tube-nosed bat (*Murina beelzebub*), also Beelzebub bat or demon bat, is a species in the vesper bat family Vespertilionidae, found in the Greater Mekong region of Southeast Asia, specifically the Quảng Trị and Gia Lai provinces of...
Behn's bat
Behn's bat
Glyphoncycteris behnii

Behn's bat, Behn's big-eared bat, or Behn's graybeard bat (*Glyphoncycteris behnii*) is a species of bat in the family Phyllostomidae found in Brazil and Peru. It is known only from six specimens so it is considered rare. It feeds on insects and...
Beijing barbastelle
Beijing barbastelle
Barbastella beijingensis

The Beijing barbastelle (*Barbastella beijingensis*), also known as the Beijing

wide-eared bat, is a species of vesper bat endemic to Beijing Municipality, China. The species was discovered by Chinese zoology students in 2001 in caves of Fangshan...

Beijing mouse-eared bat
Beijing mouse-eared bat
Myotis pequinus

The Beijing mouse-eared bat, or Peking myotis (*Myotis pequinus*) is a species of vesper bat. It is found only in China.

Beira
Beira
Dorcatragus megalotis

The beira (*Dorcatragus megalotis*) is a small arid adapted antelope that inhabits arid regions of the Horn of Africa. The term 'beira' is derived from its Somali name. The only captive breeding group of beira was at Al Wabra Wildlife Preservation,...

Belding's Ground Squirrel
Belding's Ground Squirrel
Urocitellus beldingi

Belding's ground squirrel (*Urocitellus beldingi*) is a squirrel that lives on mountains in the western United States. This species is not of conservation concern, and its range includes some protected areas.

Bell groove-toothed swamp rat
Bell groove-toothed swamp rat
Pelomys campanae

The bell groove-toothed swamp rat (*Pelomys campanae*) is a species of rodent in the family Muridae found in Angola and the Democratic Republic of the Congo. Its natural habitats are moist savanna, subtropical or tropical dry lowland grassland,...

Bellier's striped grass mouse
Bellier's striped grass mouse
Lemniscomys bellieri

Bellier's lemniscomys or Bellier's striped grass mouse (*Lemniscomys bellieri*) is a species of rodent in the family Muridae. It is found in Burkina Faso, Ivory Coast, Ghana, Guinea, and possibly Sierra Leone. Its natural habitat is typically a dry...

Beluga Whale
Beluga Whale
Delphinapterus leucas

The Beluga whale (*Delphinapterus leucas*) is a cetacean adapted to life in the Arctic, with anatomical and physiological characteristics that differentiate it from other cetaceans. Amongst these are its all-white color and the absence of a dorsal...

Bemaraha Woolly Lemur
Bemaraha Woolly Lemur
Avahi cleesei

The Bemaraha woolly lemur (*Avahi cleesei*) is a rare species of woolly lemur from Madagascar. It was first recorded in 1990 by a team of scientists from Zurich University but was formally described as a species only on 11 November 2005. The local...

Bengal Fox
Bengal Fox
Vulpes bengalensis

The Bengal fox (*Vulpes bengalensis*), also known as the Indian fox, is a fox

endemic to the Indian subcontinent from the Himalayan foothills and Terai of Nepal through southern India, and from southern and eastern Pakistan to eastern India and...

Bengal mongoose

Bengal mongoose

Herpestes javanicus palustris

The small Indian mongoose (*Urva auropunctata*) is a mongoose species native to Iraq and northern South Asia; it has also been introduced to many regions of the world, such as several Caribbean and Pacific islands.

Bengal Slow Loris

Bengal Slow Loris

Nycticebus bengalensis

The Bengal slow loris (*Nycticebus bengalensis*) is a strepsirrhine primate native to the Indian subcontinent and Indochina. Its geographic range is larger than that of any other slow loris species. Bengal slow lorises favor rainforests with dense...

Bengal Tiger

Bengal Tiger

Panthera tigris tigris

The Bengal tiger (*Panthera tigris tigris*) is the nominate Tiger subspecies. It ranks among the biggest wild cats alive today. It is considered to belong to the world's charismatic megafauna. The Bengal tiger is estimated to have been present in...

Benito roundleaf bat

Benito roundleaf bat

Hipposideros beatus

The Benito roundleaf bat (*Hipposideros beatus*) is a species of bat in the family Hipposideridae found in Cameroon, Central African Republic, Republic of the Congo, Democratic Republic of the Congo, Ivory Coast, Equatorial Guinea, Gabon, Ghana,...

Benkeith's short-tailed bat

Benkeith's short-tailed bat

Carollia benkeithi

Benkeith's short-tailed bat (*Carollia benkeithi*) is a leaf-nosed bat species found in Peru, Bolivia, and Brazil. It very closely resembles the chestnut short-tailed bat, and the two species are likely often confused.

Bennett's chinchilla rat

Bennett's chinchilla rat

Abrocoma bennettii

Bennett's chinchilla rat (*Abrocoma bennettii*) is a species of chinchilla rat in the family Abrocomidae. It is found only in Chile where its habitat is Mediterranean-type scrub on the western side of the Andes. The IUCN has assessed its...

Bennett's Tree-Kangaroo

Bennett's Tree-Kangaroo

Dendrolagus bennettianus

Bennett's tree-kangaroo (*Dendrolagus bennettianus*) is a large tree-kangaroo. Males can weigh from 11.5 kg up to almost 14 kg (25 to 31 lbs), while the females range between about 8 to 10.6 kg (17.6 to 23 lbs). They are very agile and are able to...

Berbera gerbil

Berbera gerbil

Gerbillus acticola

The Berbera gerbil (*Gerbillus acticola*) is distributed mainly in Somalia, but its

range may extend into Ethiopia and Djibouti. It is endemic to the Ethiopian xeric grasslands and shrublands ecoregion.

Berdmore's ground squirrel

Berdmore's ground squirrel

Menetes berdmorei

The Berdmore's ground squirrel (*Menetes berdmorei*) is a ground squirrel found in Southeast Asia, from the east of Myanmar to Vietnam. It is however absent on the Malay peninsula, as well as the islands. The squirrel has a grey-brown back and a...

Berg's tuco-tuco

Berg's tuco-tuco

Ctenomys bergi

Berg's tuco-tuco (*Ctenomys bergi*) is a species of rodent in the family Ctenomyidae, named after the Latvian-Argentine biologist Frederico Guillermo Carlos Berg. It is endemic to northwestern Córdoba Province in central Argentina. Its habitat is...

Bernard's wolf

Bernard's wolf

Canis lupus bernardi

Bernard's wolf (*Canis lupus bernardi*), also known as the Banks Island wolf or the Banks Island tundra wolf, is an extinct subspecies of the gray wolf that was limited to Banks and Victoria Island of the Arctic Archipelago.

Betsileo short-tailed rat

Betsileo short-tailed rat

Brachyuromys betsileoensis

The Betsileo short-tailed rat (*Brachyuromys betsileoensis*) is a species of rodent in the family Nesomyidae. It is found only in Madagascar. Its natural habitats are subtropical or tropical dry forests and subtropical or tropical dry shrubland.

Betsileo sportive lemur

Betsileo sportive lemur

Lepilemur betsileo

The betsileo sportive lemur (*Lepilemur betsileo*) is a sportive lemur endemic to Madagascar. It is a relatively large sportive lemur with a total length of about 58 to 27 cm (23 to 11 in), of which 32 to 33 cm (12.6 to 13 in) are tail. The...

Betsileo woolly lemur

Betsileo woolly lemur

Avahi betsileo

The Betsileo woolly lemur or Betsileo avahi (*Avahi betsileo*) is a species of woolly lemur native to southeastern Madagascar, in the District of Fandriana. It weighs about 1 kg. The pelage differs significantly from other southeastern woolly...

Bettongia anhydra

Bettongia anhydra

Bettongia anhydra

Bettongia anhydra, also known as desert bettong, is a recently extinct species of potoroine marsupial.

Bettongia moyesi

Bettongia moyesi

Bettongia moyesi

Bettongia moyesi is a fossil species of potoroid marsupial.

Bezoar ibex

Bezoar ibex

Capra aegagrus aegagrus

The bezoar ibex (*Capra aegagrus aegagrus*) is a wild goat subspecies that is native to the montane forested areas in the Caucasus and the Zagros Mountains.

Bharal

Bharal

Pseudois nayaur

The bharal (*Pseudois nayaur*) is a medium-sized bovid native to the high Himalayas. The Helan Mountains of Ningxia have the highest concentration of these animals in the world.

Bhutan giant flying squirrel

Bhutan giant flying squirrel

Petaurista nobilis

The Bhutan giant flying squirrel (*Petaurista nobilis*), also known as the gray's giant flying squirrel or noble giant flying squirrel, is a species of rodent in the family Sciuridae. This species lives in Himalayan forests and it is one of the...

Bhutan takin

Bhutan takin

Budorcas taxicolor whitei

The Bhutan Takin (*Budorcas taxicolor whitei*) is a vulnerable subspecies of Takin native to Bhutan, North Eastern India, Western part of China, and Tibet. The main threats to the Bhutan Takin are hunting and habitat loss.

Biak giant rat

Biak giant rat

Uromys boeadii

The Biak Island uromys or Biak giant rat (*Uromys boeadii*) is a species of rodent in the family Muridae. It is known only from a single specimen collected on the Indonesian island of Biak.

Biak glider

Biak glider

Petaurus biacensis

The Biak glider (*Petaurus biacensis*) is a species of marsupial in the family Petauridae. It is endemic to the Schouten Islands in the western region of Papua Province, Indonesia. It was formerly considered to be a subspecies of *Petaurus*...

Biak naked-backed fruit bat

Biak naked-backed fruit bat

Dobsonia emersa

The Biak naked-backed fruit bat (*Dobsonia emersa*) is a species of megabat in the family Pteropodidae. It is endemic to Indonesia.

Biak roundleaf bat

Biak roundleaf bat

Hipposideros papua

The Biak roundleaf bat or Geelvinck Bay leaf-nosed bat (*Hipposideros papua*) is a species of bat in the family Hipposideridae. It is endemic to the Biak Islands (Schouten Islands) and Cenderawasih Bay (Geelvinck Bay) area of Papua Province,...

Bibimys chacoensis

Bibimys chacoensis

Bibimys chacoensis

Bibimys chacoensis is a species of small rodent of the family Cricetidae living in the north-central part of Southern Cone of South America. The common name of this species is Chaco crimson-nosed rat (in Spanish: *ratón de hocico rosado*)

norteño)....
Bibimys labiosus
Bibimys labiosus
Bibimys labiosus

Bibimys labiosus), also known as the large-lipped crimson-nosed rat is a species of rodent in the family Cricetidae. It is found in Brazil and Argentina.

Bibimys torresi
Bibimys torresi
Bibimys torresi

Bibimys torresi), also known as Torres's crimson-nosed rat is a species of rodent in the family Cricetidae. It is found in Argentina.

Bibundi bat
Bibundi bat
Glauconycteris egeria

The Bibundi bat (*Glauconycteris egeria*) is a species of vesper bat in the family Vespertilionidae. It can be found in the following countries: Cameroon, Republic of the Congo, and Uganda. It is also found in Dzanga-Sangha Special Reserve.

Bickham's little yellow bat
Bickham's little yellow bat
Rhogeessa bickhami

Bickham's little yellow bat (*Rhogeessa bickhami*) is a species of vesper bat found in Central America.

Bicolored musk shrew
Bicolored musk shrew
Crocidura fuscomurina

The bicolored musk shrew (*Crocidura fuscomurina*) is a species of mammal in the family Soricidae. It is found in Angola, Benin, Botswana, Burkina Faso, Burundi, Cameroon, Central African Republic, Chad, Democratic Republic of the Congo, Ivory...

Bicolored roundleaf bat
Bicolored roundleaf bat
Hipposideros bicolor

The bicolored roundleaf bat (*Hipposideros bicolor*) is a species of bat in the family Hipposideridae found in Indonesia, Malaysia, the Philippines, Thailand, and Timor-Leste. This bat inhabits caves, rock crevices and tunnels among lowland...

Bicolored shrew
Bicolored shrew
Crocidura leucodon

The bicolored shrew or bicoloured white-toothed shrew (*Crocidura leucodon*) is a species of mammal in the family Soricidae. It is found in eastern, central and southern Europe and in western Asia. It is a nocturnal species and feeds on insects...

Bicolored tube-nosed bat
Bicolored tube-nosed bat
Murina bicolor

The bicolored tube-nosed bat (*Murina bicolor*) is a species of vesper bat endemic to Taiwan.

Bicolored-spined porcupine
Bicolored-spined porcupine
Coendou bicolor

The bicolored-spined porcupine (*Coendou bicolor*) is a species of nocturnal and arboreal rodent in the family Erethizontidae. It is found in Bolivia, Colombia, Ecuador, and Peru. The head and body of *Coendou bicolor* measure about 543 mm, and...

Bidentate yellow-eared bat
Bidentate yellow-eared bat
Vampyriscus bidens

The bidentate yellow-eared bat (*Vampyriscus bidens*) is a species of bat in the family Phyllostomidae, native to South America. Formerly classified in the *Vampyressa* genus, phylogenetic analyses support its inclusion in *Vampyriscus*.

Bidentate yellow-shouldered bat
Bidentate yellow-shouldered bat
Sturnira bidens

The bidentate yellow-shouldered bat (*Sturnira bidens*) is a species of bat in the family Phyllostomidae. It is found in Colombia, Ecuador, Peru, and Venezuela.

Big bonneted bat
Big bonneted bat
Eumops dabbenei

The big bonneted bat, or Dabbene's mastiff bat (*Eumops dabbenei*) is a species of bat in the family Molossidae, native to South America. It is named for a former conservator at the Buenos Aires National Museum.

Big Brown Bat
Big Brown Bat
Eptesicus fuscus

The Big brown bat (*Eptesicus fuscus*) is a species of vesper bat found throughout the Americas. It was first described as a species in 1796. Compared to other microbats, Big brown bats are relatively large and consume a diverse array of...

Big crested mastiff bat
Big crested mastiff bat
Promops centralis

The big crested mastiff bat (*Promops centralis*), is a bat species from South and Central America.

Big deer mouse
Big deer mouse
Peromyscus grandis

The big deer mouse (*Peromyscus grandis*) is a species of rodent in the family Cricetidae. It is found only in Guatemala.

Big free-tailed bat
Big free-tailed bat
Nyctinomops macrotis

The big free-tailed bat (*Nyctinomops macrotis*) is a bat species found in South, North and Central America, as well as the Caribbean.

Big Hairy Armadillo
Big Hairy Armadillo
Chaetophractus villosus

The Big hairy armadillo (*Chaetophractus villosus*) is one of the largest and most numerous armadillos in South America. It is an accomplished digger and spends most of its time below ground in burrows. In Spanish this animal is colloquially known...

Big Mexican small-eared shrew
Big Mexican small-eared shrew
Cryptotis magna

The big Mexican small-eared shrew (*Cryptotis magna*) is a species of mammal in the family Soricidae. It is endemic to Mexico.

Big naked-backed bat

Big naked-backed bat

Pteronotus gymnonotus

The big naked-backed bat (*Pteronotus gymnonotus*), is a bat species from South and Central America.

Big pocket gopher

Big pocket gopher

Orthogeomys lanius

The big pocket gopher (*Heterogeomys lanius*) is a species of rodent in the family Geomyidae. It is endemic to Veracruz state in eastern Mexico. It has only been found on the southeastern slopes of Pico de Orizaba, at elevations of 1,300 metres...

Big red bat

Big red bat

Lasiurus egregius

The big red bat (*Aeorestes egregius*) is a species of vesper bat from South and Central America.

Big-eared brown bat

Big-eared brown bat

Histiotus macrotus

The big-eared brown bat (*Histiotus macrotus*) is a species of vesper bat found in Argentina, Paraguay, and Chile.

Big-eared climbing rat

Big-eared climbing rat

Otodylomys phyllotis

The big-eared climbing rat (*Otodylomys phyllotis*) is a species of rodent in the family Cricetidae. It is found in Central America from Mexico to Costa Rica.

Big-eared flying fox

Big-eared flying fox

Pteropus macrotis

The big-eared flying fox (*Pteropus macrotis*) is a species of bat in the family Pteropodidae, larger bats who subsist largely on fruits. The species is distributed across a range in Indonesia, Papua New Guinea and islands nearing the Cape York...

Big-eared horseshoe bat

Big-eared horseshoe bat

Rhinolophus macrotis

The big-eared horseshoe bat (*Rhinolophus macrotis*) is a bat species within the Rhinolophidae native to China, India, Indonesia, Laos, Malaysia, Nepal, Pakistan, the Philippines, Thailand, and Vietnam.

Big-eared kangaroo rat

Big-eared kangaroo rat

Dipodomys elephantinus

The big-eared kangaroo rat (*Dipodomys elephantinus*) is a kangaroo rat that lives in chaparral. It is named for and distinguished by its comparatively large ears.

Big-eared mastiff bat

Big-eared mastiff bat

Otomops papuensis

The big-eared mastiff bat (*Otomops papuensis*) is a species of bat in the family Molossidae endemic to Papua New Guinea, known from Gulf Province and Oro Province. It is only known from 11 specimens and it is not easy to survey. It forages in...

Big-eared opossum
Big-eared opossum
Didelphis aurita

The big-eared opossum (*Didelphis aurita*) also known as a saruê is an opossum species from South America. It is found in Argentina, Brazil and Paraguay. This species, which was considered a population of the common opossum (*D. marsupialis*) for...

Big-eared pipistrelle
Big-eared pipistrelle
Hypsugo macrotis

The big-eared pipistrelle (*Hypsugo macrotis*) is a species of vesper bat in the family Vespertilionidae. It can be found in Indonesia and Malaysia. It forages over mud flats over Peninsula Malaysia but its roosting activities are unknown. Its...

Big-eared roundleaf bat
Big-eared roundleaf bat
Hipposideros macrobullatus

The big-eared roundleaf bat (*Hipposideros macrobullatus*) is a species of bat in the family Hipposideridae. It is endemic to Indonesia, known from Kangean Islands, southwestern Sulawesi and Seram Island. It roosts in caves and tree hollows and...

Big-eared swamp rat
Big-eared swamp rat
Malacomys longipes

The big-eared swamp rat (*Malacomys longipes*) is a species of rodent in the family Muridae. It is found in sub-Saharan Africa from Nigeria to Kenya and south to Zambia and Angola. Its natural habitat is subtropical or tropical moist lowland forests.

Big-eared woolly bat
Big-eared woolly bat
Chrotopterus auritus

The big-eared woolly bat or (Peters's) woolly false vampire bat (*Chrotopterus auritus*) is a species of bat, belonging to the Order Chiroptera and Family Phyllostomidae. The name *Chrotopterus* is derived from Greek roots *ch* ariots (skin, color),...

Big-Headed Mole-Rat
Big-Headed Mole-Rat
Tachyoryctes macrocephalus

The big-headed African mole rat, (*Tachyoryctes macrocephalus*), also known as the giant root-rat, Ethiopian African mole rat, or giant mole rat, is a rodent species in the family Spalacidae. It is endemic to Ethiopia's Bale Mountains. Its natural...

Bighorn Sheep
Bighorn Sheep
Ovis canadensis

The Bighorn sheep (*Ovis canadensis*) is a species of sheep native to North America. Sheep originally crossed to North America over the Bering Land Bridge from Siberia; the population in North America peaked in the millions, and the bighorn sheep...

Bini free-tailed bat
Bini free-tailed bat
Myotis whitleyi

The Bini free-tailed bat (*Myopterus whitleyi*) is a species of bat in the family Molossidae. It is found in Cameroon, Central African Republic, Democratic Republic of the Congo, Gabon, Ghana, Nigeria, and Uganda. Its natural habitat is...

Binturong

Binturong

Arctictis binturong

The binturong (*Arctictis binturong*) is a carnivore of a medium size found in dense forests in South-East Asia. This animal is thought to be a close relative of the palm civet. It is the biggest member of this family.

Bioko Allen's bushbaby

Bioko Allen's bushbaby

Sciurocheirus alleni

Bioko Allen's bushbaby (*Sciurocheirus alleni*), also known as the Bioko squirrel galago, is a species of primate in the galago family found in Bioko, Equatorial Guinea. Its natural habitat is subtropical or tropical dry forests. The bushbaby is...

Bioko drill

Bioko drill

Mandrillus leucophaeus poensis

The Bioko drill (*Mandrillus leucophaeus poensis*) is a subspecies of the drill, an Old World monkey. It is endemic to Bioko Island, Equatorial Guinea, located off the west coast of Africa. The drill is one of the largest monkey species, and is...

Bioko forest shrew

Bioko forest shrew

Sylvisorex isabellae

The Bioko forest shrew or Isabella's shrew (*Sylvisorex isabellae*) is a species of mammal in the family Soricidae found in Cameroon and Equatorial Guinea. Its natural habitats are subtropical or tropical moist montane forest and subtropical or...

Birdlike noctule

Birdlike noctule

Nyctalus aviator

The birdlike noctule (*Nyctalus aviator*) is a species of bat. An adult birdlike noctule has a body length of 7.1-9.5 cm, a tail of 5.5-6.4 cm, and a wing length of 5.8-5.95 cm. It nests in the holes in old trees and buildings, and sometimes in...

Bishop's slender opossum

Bishop's slender opossum

Marmosops bishopi

Bishop's slender opossum (*Marmosops bishopi*) is a small, arboreal marsupial opossum native to Brazil, Peru, and Bolivia. It somewhat resembles a placental rat or shrew.

Bismarck giant rat

Bismarck giant rat

Uromys neobritannicus

The Bismarck giant rat (*Uromys neobritannicus*) is a species of rodent in the family Muridae. It is endemic to the island of New Britain, Papua New Guinea.

Bismarck masked flying fox

Bismarck masked flying fox

Pteropus capistratus

The Bismarck masked flying fox (*Pteropus capistratus*) is a species of flying fox in the family Pteropodidae found in Papua New Guinea and named after the Bismarck Archipelago. It was once considered a subspecies of *Pteropus temminckii* before...

Bismarck trumpet-eared bat
Bismarck trumpet-eared bat
Kerivoula myrella

The Bismarck trumpet-eared bat (*Kerivoula myrella*), also known as the Manus Island woolly bat, is a species of vesper bat in the family Vespertilionidae. It is found in subtropical or tropical dry forests.

Biting chinchilla mouse
Biting chinchilla mouse
Euneomys mordax

The biting chinchilla mouse (*Euneomys mordax*) is a species of rodent in the family Cricetidae. It is found in west central Argentina and nearby areas of Chile, where it inhabits highland meadows, probably at elevations of 1740 to 3000 m.

Black Agouti
Black Agouti
Dasyprocta fuliginosa

The black agouti (*Dasyprocta fuliginosa*) is a South American species of agouti from the family Dasyproctidae.

Black and red bush squirrel
Black and red bush squirrel
Paraxerus lucifer

The black and red bush squirrel (*Paraxerus lucifer*) is a species of rodent in the family Sciuridae occurring in Malawi, Tanzania, and Zambia. In Malawi, it is found in Misuku Hills and Nyika Plateau and in Tanzania Poroto Mountains and Mount...

Black and Rufous Elephant Shrew
Black and Rufous Elephant Shrew
Rhynchocyon petersi

The black and rufous elephant shrew, (*Rhynchocyon petersi*) the black and rufous sengi, or the Zanj elephant shrew is one of the 17 species of elephant shrew found only in Africa. It is native to the lowland montane and dense forests of Kenya and...

Black bearded saki
Black bearded saki
Chiropotes satanas

The black bearded saki (*Chiropotes satanas*) is a species of bearded saki, native to the Amazon rainforest in South America, specifically to an area of north-eastern Brazil. Black Bearded saki are of the Platyrrhine infraorder, also known as New...

Black bonneted bat
Black bonneted bat
Eumops auripendulus

The black bonneted bat, also known as Shaw's mastiff bat, (*Eumops auripendulus*), is a species of bat from the Americas.

Black Capuchin
Black Capuchin
Sapajus nigritus

The black capuchin, (*Sapajus nigritus*), also known as the black-horned capuchin, is a capuchin monkey from the Atlantic Forest in south-eastern Brazil and far north-eastern Argentina. Historically, it was included as a subspecies of the tufted...

Black colobus
Black colobus
Colobus satanas

The black colobus (*Colobus satanas*), or satanic black colobus, is a species of Old World monkey belonging to the genus *Colobus*. The species is found in a small area of western central Africa. Black colobuses are large, completely covered with...

Black Crested Gibbon

Black Crested Gibbon

Nomascus concolor

The Black crested gibbon (*Nomascus concolor*) is a Critically Endangered primate found in Asia. They live in forests and are well known for their habit of swinging between the branches of trees on their long arms.

Black crested mangabey

Black crested mangabey

Lophocebus aterrimus

The black crested mangabey (*Lophocebus aterrimus*) is a species of primate in the family *Cercopithecidae*. It is only found in Democratic Republic of the Congo with a small habitat extending to Angola. Its natural habitat is subtropical or...

Black dorcopsis

Black dorcopsis

Dorcopsis atrata

The black dorcopsis or black forest wallaby (*Dorcopsis atrata*) is a species of marsupial in the family *Macropodidae*. It is endemic to an island at the eastern end of New Guinea where its natural habitat is subtropical or tropical dry forests. It...

Black Duiker

Black Duiker

Cephalophus niger

The black duiker (*Cephalophus niger*), also known as tuba in Dyula, is a forest-dwelling duiker found in the southern parts of Sierra Leone, Liberia, Côte d'Ivoire, Ghana, Benin, and Nigeria. Black duikers stand around 500 mm (20 in) tall at the...

Black dwarf porcupine

Black dwarf porcupine

Coendou nycthemera

The black dwarf porcupine, also known as Koopman's porcupine, *Coendou nycthemera*, is a porcupine species from the New World porcupine family endemic to northern Brazil. It occurs in the Amazon rainforest east of the Madeira River and south of the...

Black Flying Fox

Black Flying Fox

Pteropus alecto

The Black flying fox (*Pteropus alecto*) is among the largest bats in the world but is considerably smaller than the largest species in its genus, *Pteropus*. The black flying fox is found only in Australia, Papua New Guinea, and Indonesia. It is not...

Black flying squirrel

Black flying squirrel

Aeromys tephromelas

The black flying squirrel or large black flying squirrel (*Aeromys tephromelas*) is a species of rodent in the family *Sciuridae*. It is found in Brunei, Indonesia, and Malaysia; its habitat is primary and secondary forests and gardens where it uses...

Black Giant Squirrel

Black Giant Squirrel

Ratufa bicolor

The Black giant squirrel (*Ratufa bicolor*) is a large tree squirrel native to the Indomalayan zoogeographic region. It is one of the largest species of squirrel in the world.

Black Howler Monkey

Black Howler Monkey

Alouatta caraya

The black howler (*Alouatta caraya*) or black-and-gold howler, is among the largest New World monkeys and a member of the *Alouatta* genus. The black howler is distributed in areas of South America such as Paraguay, southern Brazil, eastern Bolivia,...

Black Jackrabbit

Black Jackrabbit

Lepus insularis

The black jackrabbit (*Lepus insularis*) is a species of mammal in the family Leporidae. Endemic to Mexico, its only known location is Espiritu Santo Island in the Gulf of California. The IUCN has listed this species as a "vulnerable species"...

Black Lemur

Black Lemur

Eulemur macaco

The black lemur (*Eulemur macaco*) is a species of lemur from the family Lemnidae. Like all lemurs, it is endemic to Madagascar. Originally, the species was thought to have two subspecies, *Eulemur macaco macaco* and *Eulemur macaco flavifrons*, both...

Black Lion Tamarin

Black Lion Tamarin

Leontopithecus chrysopygus

The black lion tamarin (*Leontopithecus chrysopygus*), also known as the golden-rumped lion tamarin, is a lion tamarin endemic to the Brazilian state of São Paulo, almost exclusively at the Morro do Diabo State Park. Its limited geographical range...

Black mastiff bat

Black mastiff bat

Molossus rufus

The black mastiff bat (*Molossus rufus*) is a bat species. It ranges from the northern region of South America (excluding Chile), most of Central America (excluding Belize) and parts of southern Mexico.

Black Mongoose

Black Mongoose

Galerella nigrata

The Black mongoose (*Herpestes flavescens nigratus*) is a subspecies of the Angolan slender mongoose. Although originally described as a separate species by Oldfield Thomas in 1928, it has often been considered a subspecies of the Slender mongoose...

Black musk deer

Black musk deer

Moschus fuscus

The black musk deer or dusky musk deer (*Moschus fuscus*) is a species of even-toed ungulate in the family Moschidae. It is found in Bhutan, China, India, Myanmar, and Nepal.

Black myotis

Black myotis
Myotis nigricans

The black myotis (*Myotis nigricans*) is a vesper bat species from South and Central America. Its body is dark brown/black. The head to body length, not including the tail, is about 5 cm (2 in.). The black myotis is a tiny bat with a small pointed...

Black pika
Black pika
Ochotona nigritia

The black pika or silver pika (*Ochotona nigritia*) is a species of mammal in the family Ochotonidae. It was thought to be common to the Yunnan Province of China where it was first discovered in 2000, it is only known from four specimens. The...

Black Rat
Black Rat
Rattus rattus

The Black rat (*Rattus rattus*) has an extremely long tail that is longer than its head and body. It likely originated in the Indian subcontinent but is now found worldwide. Romans were the ones who introduced this species to Britain. Black rats...

Black Rhinoceros
Black Rhinoceros
Diceros bicornis

Rhinos are very ancient animals. In fact, these mammals do resemble some prehistoric species. They appeared millions of years ago, during the Miocene era. One of the most conspicuous characteristics of these animals is the two horns of their...

Black Sea field mouse
Black Sea field mouse
Apodemus ponticus

The Black Sea field mouse (*Apodemus ponticus*) is a species of rodent in the family Muridae. It is found in Armenia, Azerbaijan, Georgia, possibly Iran, Iraq, Russian Federation, and Turkey.

Black shrew
Black shrew
Suncus ater

The black shrew (*Suncus ater*) is a white-toothed shrew only known from Mount Kinabalu in the Malaysian state of Sabah on the island of Borneo. It is listed as a critically endangered species due to habitat loss and a restricted range. It is the...

Black Snub-Nosed Monkey
Black Snub-Nosed Monkey
Rhinopithecus bieti

The black-and-white snub-nosed monkey (*Rhinopithecus bieti*), also known as the Yunnan snub-nosed monkey, is a large black and white primate that lives only in the southern Chinese province of Yunnan, where it is known to the locals as the Yunnan...

Black squirrel monkey
Black squirrel monkey
Saimiri vanzolinii

The black squirrel monkey (*Saimiri vanzolinii*), also known as the blackish squirrel monkey or black-headed squirrel monkey, is a small New World primate, endemic to the central Amazon in Brazil. It largely resembles the female of the far

more...

Black tamarin

Black tamarin

Saguinus niger

The black tamarin (*Saguinus niger*) or western black-handed tamarin, is a species of tamarin endemic to Brazil.

Black titi

Black titi

Cheracebus lugens

The black titi (*Cheracebus lugens*), is a species of titi, a type of New World monkey, from South America. It is found in Brazil, Colombia, and Venezuela. It was described in 1811 as *Calicebus lugens*.

Black Wallaroo

Black Wallaroo

Osphranter bernardus

The Black wallaroo (*Osphranter bernardus*) is a species of macropod that lives in a small mountainous area in Arnhem Land, Northern Territory, Australia. It is classified as near threatened, mostly due to its limited distribution. A large...

Black Wildebeest

Black Wildebeest

Connochaetes gnou

The Black wildebeest (*Connochaetes gnou*) is one of the two closely related wildebeest species. It was first described in 1780 by Eberhard August Wilhelm von Zimmermann. Black wildebeest live in three distinct social groups and communicate using a...

Black-and-White Ruffed Lemur

Black-and-White Ruffed Lemur

Varecia variegata

The black-and-white ruffed lemur (*Varecia variegata*) is an endangered species of ruffed lemur, one of two which are endemic to the island of Madagascar. Despite having a larger range than the red ruffed lemur, it has a much smaller population...

Black-Backed Jackal

Black-Backed Jackal

Canis mesomelas

The Black-backed jackal (*Lupulella mesomelas*) is a medium-sized canine native to eastern and southern Africa. These regions are separated by roughly 900 kilometers. Compared to other members of the genus *Canis*, the Black-backed jackal is a very...

Black-bearded flying fox

Black-bearded flying fox

Pteropus melanopogon

The black-bearded flying fox (*Pteropus melanopogon*) is an endangered species of megabat in the genus *Pteropus*. It is endemic to Indonesia, found on the islands of Ambon, Buru, Seram, Banda, and Yamdena. Currently considered monotypic, it...

Black-bearded tomb bat

Black-bearded tomb bat

Taphozous melanopogon

The black-bearded tomb bat (*Taphozous melanopogon*) is a species of sac-winged bat. It is found in South and South East Asia.

Black-bellied fruit bat

Black-bellied fruit bat

Melonycteris melanops

The black-bellied fruit bat (*Melonycteris melanops*) is a species of order Chiroptera in the family Pteropodidae.

Black-Capped Capuchin

Black-Capped Capuchin

Sapajus apella

The tufted capuchin (*Sapajus apella*), also known as brown capuchin, black-capped capuchin, or pin monkey is a New World primate from South America and the Caribbean islands of Trinidad and Margarita. As traditionally defined, it is one of the...

Black-capped fruit bat

Black-capped fruit bat

Chironax melanocephalus

The black-capped fruit bat (*Chironax melanocephalus*) is a species of megabat in the monotypic genus *Chironax*.

Black-capped marmot

Black-capped marmot

Marmota camtschatica

The black-capped marmot (*Marmota camtschatica*) is a species of rodent in the family Sciuridae. It is endemic to the Russian Far East, but its range is discontinuous and divided into three main parts, each with its own subspecies. The...

Black-Capped Squirrel Monkey

Black-Capped Squirrel Monkey

Saimiri boliviensis

The black-capped squirrel monkey (*Saimiri boliviensis*) is a species of New-World monkey native to the upper Amazon basin in Bolivia, western Brazil and eastern Peru. They weigh between 365 and 1135 grams and measure, from the head to the base of...

Black-chinned emperor tamarin

Black-chinned emperor tamarin

Saguinus imperator imperator

The black-chinned emperor tamarin (*Saguinus imperator imperator*) is one of the two subspecies of the emperor tamarin. Unlike the bearded emperor tamarin, it has no beard. It is distributed throughout the rainforests of Brazil, Peru, and Bolivia....

Black-clawed brush-furred rat

Black-clawed brush-furred rat

Lophuromys melanonyx

The black-clawed brush-furred rat (*Lophuromys melanonyx*) is a species of rodent in the family Muridae. It is found only in Ethiopia. Its natural habitat is subtropical or tropical high-altitude grassland. It is threatened by habitat loss.

Black-crowned Central American squirrel monkey

Black-crowned Central American squirrel monkey

Saimiri oerstedii oerstedii

The black-crowned Central American squirrel monkey (*Saimiri oerstedii oerstedii*) is a subspecies of the Central American squirrel monkey. Its range is restricted to the Pacific coast of western Panama to the western portion of the Chiriquí...

Black-eared flying fox

Black-eared flying fox

Pteropus melanotus

The black-eared flying fox, species *Pteropus melanotus*, is a bat of the family Pteropodidae (megabats). Also known as Blyth's flying fox, it is found on the Andaman Islands and Nicobar Islands (India), and in Sumatra (Indonesia). A population on...

Black-eared mouse

Black-eared mouse

Peromyscus melanotis

The black-eared mouse (*Peromyscus melanotis*), or black-eared deer mouse, is a species of rodent in the family Cricetidae, native to North America.

Black-eared squirrel

Black-eared squirrel

Nannosciurus melanotis

The black-eared squirrel (*Nannosciurus melanotis*) is a species of rodent in the family Sciuridae. It is monotypic within the genus *Nannosciurus*. This tiny squirrel is found in forests in Borneo, Sumatra and Java. Except for its striking whitish...

Black-Flanked Rock-Wallaby

Black-Flanked Rock-Wallaby

Petrogale lateralis

The Black-flanked rock-wallaby (*Petrogale lateralis*) is a small wallaby, one of several rock-wallabies in the genus *Petrogale*. It was first described by John Gould in 1842 and is an important symbol in Anangu mythology.

Black-Footed Cat

Black-Footed Cat

Felis nigripes

The Black-footed cat (*Felis nigripes*) is the smallest wild cat in Africa. The first black-footed cat known to science was discovered in the northern Karoo of South Africa and described in 1824. This rare and secretive cat usually rests in burrows...

Black-Footed Ferret

Black-Footed Ferret

Mustela nigripes

The endangered Black-footed ferret (*Mustela nigripes*) belongs to the weasel family and is the only ferret that is native to North America. Domestic ferrets are a different species and of European origin. The Black-footed ferret was once found in...

Black-footed gray langur

Black-footed gray langur

Semnopithecus hypoleucos

The black-footed gray langur (*Semnopithecus hypoleucos*) is an Old World monkey, one of the species of langurs. Found in southern India, this, like other gray langurs, is a leaf-eating monkey.

Black-footed mongoose

Black-footed mongoose

Bdeogale nigripes

The black-footed mongoose (*Bdeogale nigripes*) is a mongoose species native to Central Africa, where it inhabits deep deciduous forests from eastern Nigeria to the southern Democratic Republic of the Congo. It has been listed as Least Concern on...

Black-footed shrew

Black-footed shrew

Crocidura nigripes

The black-footed shrew (*Crocidura nigripes*) is a species of mammal in the family Soricidae. It is endemic to northern and central Sulawesi, Indonesia where it lives on the floor of the tropical forests. The International Union for Conservation...
Black-footed tree-rat
Black-footed tree-rat
Mesembriomys gouldii

The black-footed tree-rat also known as Djintamoonga (*Mesembriomys gouldii*) is one of two endemic arboreal rat species from the genus *Mesembriomys* found in the northern regions of Australia. The species is one of the largest murids found in...
Black-fronted duiker
Black-fronted duiker
Cephalophus nigrifrons

The black-fronted duiker (*Cephalophus nigrifrons*) is a small antelope found in central and west-central Africa.
Black-fronted titi
Black-fronted titi
Callicebus nigrifrons

The black-fronted titi (*Callicebus nigrifrons*) is a species of titi, a type of New World monkey. The black-fronted titi is a small diurnal primate. The body of this primate is covered in grey to brown fur with black fur concentrated around the...
Black-gilded pipistrelle
Black-gilded pipistrelle
Arielulus circumdatus

The bronze sprite (*Arielulus circumdatus*), also known as the black-gilded pipistrelle, is a species of vesper bat found in China, India, Myanmar, and Nepal.
Black-headed marmoset
Black-headed marmoset
Mico nigriceps

The black-headed marmoset (*Mico nigriceps*) is a marmoset species endemic to Brazil. It inhabits humid tropical rainforest, mostly second growth and edge; the distribution is not exactly known but is thought to be Rio dos Marmelos in the north...
Black-headed night monkey
Black-headed night monkey
Aotus nigriceps

The black-headed night monkey (*Aotus nigriceps*) is a night monkey species from South America. It is found in Brazil, Bolivia and Peru. The *A. nigriceps* in Peru were notably inhabiting areas that were degraded, and often these areas were...
Black-Headed Spider Monkey
Black-Headed Spider Monkey
Ateles fusciceps

The black-headed spider monkey (*Ateles fusciceps*) is a type of New World monkey, from Central and South America. It is found in Colombia, Ecuador, and Panama. Although primatologists such as Colin Groves (1989) follow Kellogg and Goldman (1944)...
Black-mantled tamarin
Black-mantled tamarin
Leontocebus nigricollis

The black-mantled tamarin, *Leontocebus nigricollis*, is a species of saddle-back tamarin from the northwestern Amazon in far western Brazil, southeastern Colombia, northeastern Peru and eastern Ecuador. There are 3 subspecies: Spix's black mantle...

Black-rumped agouti
Black-rumped agouti
Dasyprocta prymnolopha

The black-rumped agouti (*Dasyprocta prymnolopha*) is an agouti species from the family Dasyproctidae. It is endemic to Brazil, and its range roughly equals the Northeast Region. It is named after its black rump which contrast clearly with the...

Black-shanked douc
Black-shanked douc
Pygathrix nigripes

The black-shanked douc (*Pygathrix nigripes*) is an endangered species of douc found mostly in the forests of eastern Cambodia, with some smaller populations in Vietnam. This species is unique among the doucs in having a largely greyish-blue face....

Black-shouldered opossum
Black-shouldered opossum
Caluromysiops irrupta

The black-shouldered opossum (*Caluromysiops irrupta*), also known as the white-eared opossum, is an opossum known from western Brazil and southeastern Peru. It was first described by Colin Campbell Sanborn, curator of Field Museum of Natural...

Black-spined Atlantic tree-rat
Black-spined Atlantic tree-rat
Phyllomys nigrispinus

The black-spined Atlantic tree-rat, (*Phyllomys nigrispinus*), is a South American spiny rat species in the family Echimyidae. It is found in southeastern Brazil, where it inhabits moist broadleaf forest and semideciduous forest in the Atlantic...

Black-spotted cuscus
Black-spotted cuscus
Spilocuscus rufoniger

The black-spotted cuscus (*Spilocuscus rufoniger*) is a species of marsupial in the family Phalangeridae. It is among the largest members of the family, only being surpassed by the bear cuscus. It is a relatively colourful species found in forests...

Black-striped capuchin
Black-striped capuchin
Sapajus libidinosus

The black-striped capuchin (*Sapajus libidinosus*), also known as the bearded capuchin, is a New World monkey in the family Cebidae. They can be found in northern and central Brazil. These capuchins mostly live in dry forests, and savannah...

Black-striped squirrel
Black-striped squirrel
Callosciurus nigrovittatus

The black-striped squirrel (*Callosciurus nigrovittatus*) is a species of rodent in the family Sciuridae. It is found throughout Java, Sumatra, southern Thailand, the Malay Peninsula, and numerous small islands. This taxon consists of four...

Black-striped wallaby
Black-striped wallaby
Macropus dorsalis

The black-striped wallaby (*Notamacropus dorsalis*), also known as the scrub wallaby or eastern brush wallaby, is a medium-sized wallaby found in Australia, from Townsville in Queensland to Narrabri in New South Wales. In New South Wales, it is...

Black-tailed dasyure

Black-tailed dasyure

Murexia melanurus

The black-tailed dasyure (*Murexia melanurus*) is a species of marsupial in the family Dasyuridae.

Black-tailed deer

Black-tailed deer

Odocoileus hemionus columbianus

Two forms of black-tailed deer or blacktail deer that occupy coastal woodlands in the Pacific Northwest of North America are subspecies of the mule deer (*Odocoileus hemionus*). They have sometimes been treated as a species, but virtually all...

Black-tailed gerbil

Black-tailed gerbil

Gerbilliscus nigricaudus

The black-tailed gerbil (*Gerbilliscus nigricaudus*) is a species of rodent found in Ethiopia, Kenya, Somalia, and Tanzania. Its natural habitats are dry savanna, subtropical or tropical dry shrubland, and arable land.

Black-tailed hairy dwarf porcupine

Black-tailed hairy dwarf porcupine

Coendou melanurus

The black-tailed hairy dwarf porcupine, *Coendou melanurus*, is a South American porcupine species from the family Erethizontidae. It is found in Brazil, Colombia, Ecuador, French Guiana, Guyana, Suriname and Venezuela. This species was formerly...

Black-tailed hutia

Black-tailed hutia

Mysateles melanurus

The black-tailed hutia (*Mesocapromys melanurus*), also known as the bushy-tailed hutia, is a species of rodent in the subfamily Capromyinae that is endemic to lowland moist forests in Cuba. It is threatened by habitat loss. Although it was...

Black-Tailed Jackrabbit

Black-Tailed Jackrabbit

Lepus californicus

The Black-tailed jackrabbit (*Lepus californicus*) is a common hare of the western United States and Mexico. It is one of the largest North American hares. The Black-tailed jackrabbit does not migrate or hibernate during winter and uses the same...

Black-tailed marmoset

Black-tailed marmoset

Mico melanurus

The black-tailed marmoset (*Mico melanurus*) is a species of New World monkey from central South America, where it ranges from the south-central Amazon in Brazil, south through the Pantanal and eastern Bolivia, to the Chaco in far northern...

Black-tailed mosaic-tailed rat

Black-tailed mosaic-tailed rat

Melomys rufescens

The black-tailed mosaic-tailed rat (*Melomys rufescens*) is a species of rodent in the family Muridae. It is endemic to the island of New Guinea.

Black-tailed mouse

Black-tailed mouse
Peromyscus melanurus

The black-tailed mouse (*Peromyscus melanurus*) is a species of rodent in the family Cricetidae endemic to Mexico.

Black-Tailed Prairie Dog
Black-Tailed Prairie Dog
Cynomys ludovicianus

The black-tailed prairie dog (*Cynomys ludovicianus*) is a rodent of the family Sciuridae found in the Great Plains of North America from about the United States-Canada border to the United States-Mexico border. Unlike some other prairie dogs,...

Black-tailed tree rat
Black-tailed tree rat
Thallomys nigricauda

The black-tailed tree rat, also called black-tailed acacia rat or black-tailed thallomys, (*Thallomys nigricauda*), is a species of rodent in the family Muridae. It is found in Angola, Botswana, Namibia and South Africa, where its natural habitat...

Black-tufted gerbil
Black-tufted gerbil
Gerbillus famulus

The black-tufted gerbil (*Gerbillus famulus*) is distributed mainly across the southwestern Arabian Peninsula.

Black-Tufted Marmoset
Black-Tufted Marmoset
Callithrix penicillata

The black-tufted marmoset (*Callithrix penicillata*), also known as Mico-estrela in Portuguese, is a species of New World monkey that lives primarily in the Neotropical gallery forests of the Brazilian Central Plateau. It ranges from Bahia to...

Black-winged little yellow bat
Black-winged little yellow bat
Rhogeessa tumida

The black-winged little yellow bat (*Rhogeessa tumida*) is a species of vesper bat native to Central America.

Blackbuck
Blackbuck
Antilope cervicapra

The blackbuck (*Antilope cervicapra*) is an antelope that inhabits grassy plains and lightly forested areas with perennial water sources. It is the sole living member of the genus *Antilope* and was scientifically described by Carl Linnaeus in 1758....

Blackish deer mouse
Blackish deer mouse
Peromyscus furvus

The blackish deer mouse (*Peromyscus furvus*) is a species of rodent in the family Cricetidae found only in Mexico, and is relatively poorly studied.

Blackish grass mouse
Blackish grass mouse
Thaptomys nigrita

The blackish grass mouse (*Thaptomys nigrita*) also formerly called the ebony

akodont, is a rodent species from South America. It is found in Argentina, Brazil, Paraguay and Uruguay. It is the only species in the genus Thaptomys.
Blackish small-eared shrew
Blackish small-eared shrew
Cryptotis nigrescens

The blackish small-eared shrew (*Cryptotis nigrescens*) is a species of shrew in the family Soricidae. It is found in parts of Costa Rica, El Salvador, Guatemala, Honduras, Mexico, and Panama. An example specific habitat is the Petenes mangroves...
Blackish white-toothed shrew
Blackish white-toothed shrew
Crocidura nigricans

The blackish white-toothed shrew (*Crocidura nigricans*) is a species of mammal in the family Soricidae. It is endemic to Angola.
Blainville's Beaked Whale
Blainville's Beaked Whale
Mesoplodon densirostris

Blainville's beaked whale (*Mesoplodon densirostris*), or the dense-beaked whale, is believed to be the widest ranging mesoplodont whale. The French zoologist Henri de Blainville first described the species in 1817 from a small piece of jaw – the...
Blanford's bat
Blanford's bat
Hesperoptenus blanfordi

Blanford's bat (*Hesperoptenus blanfordi*), also known as the least false-serotine bat, is a species of vesper bat. It can be found in Brunei, Cambodia, Indonesia, Laos, Malaysia, Myanmar and Thailand, where it lives in various different forested...
Blanford's Fox
Blanford's Fox
Vulpes cana

Blanford's fox (*Vulpes cana*) is a small fox native to the Middle East and Central Asia. It is listed as Least Concern on the IUCN Red List.
Blanford's fruit bat
Blanford's fruit bat
Sphaerias blanfordi

Blanford's fruit bat (*Sphaerias blanfordi*) is a mountain species of megabat. It is found in several countries in South and Southeast Asia.
Blanford's jerboa
Blanford's jerboa
Jaculus blanfordi

Blanford's jerboa (*Jaculus blanfordi*) is a species of rodent in the family Dipodidae. It is native to Central Asia and is found in Turkmenistan, Uzbekistan, Afghanistan, Iran and Pakistan.
Blanford's rat
Blanford's rat
Madromys blanfordi

Blanford's rat (*Madromys blanfordi*) is a species of rodent in the family Muridae. It is the only species in the genus Madromys. Known as වළිග සුදු වන මීයා (meaning white-tailed wood rat) in Sinhala language. It is found throughout India, Sri...
Blasius's horseshoe bat
Blasius's horseshoe bat

Rhinolophus blasii

Blasius's horseshoe bat (*Rhinolophus blasii*) is a species of insectivorous bat in the family Rhinolophidae found throughout large parts of the Mediterranean, Middle East and Northern Africa.

Blazed Luzon shrew-rat

Blazed Luzon shrew-rat

Chrotomys silaceus

The blazed Luzon shrew rat (*Chrotomys silaceus*) is a species of rodent in the family Muridae, endemic to the Philippines.

Blesbok

Blesbok

Damaliscus pygargus phillipsi

The blesbok or blesbuck (*Damaliscus pygargus phillipsi*) is an subspecies of the bontebok antelope endemic to South Africa, Eswatini and Namibia. It has a distinctive white face and forehead which inspired the name, because bles is the Afrikaans...

Blick's grass rat

Blick's grass rat

Arvicanthis blicki

Blick's grass rat (*Arvicanthis blicki*) is a species of rodent in the family Muridae. It is found only in Ethiopia. Its natural habitat is subtropical or tropical high-altitude grassland. It is threatened by habitat loss.

Blind mole

Blind mole

Talpa caeca

The blind mole (*Talpa caeca*), also known as the Mediterranean mole, is a mole found in the Mediterranean region. It is similar to the European mole, differing most prominently in having eyes covered with skin. It is found in Albania, France,...

Blond Capuchin

Blond Capuchin

Sapajus flavius

The Blond capuchin (*Sapajus flavius*) is a species of the capuchin monkeys group to northeastern Brazil. They are considered to be a critically endangered species.

Blue Duiker

Blue Duiker

Philantomba monticola

The blue duiker (*Philantomba monticola*) is a small antelope found in central, southern and eastern Africa. It is the smallest duiker. The species was first described by Swedish naturalist Carl Peter Thunberg in 1789. 12 subspecies are...

Blue Monkey

Blue Monkey

Cercopithecus mitis

The blue monkey or diademed monkey (*Cercopithecus mitis*) is a species of Old World monkey native to Central and East Africa, ranging from the upper Congo River basin east to the East African Rift and south to northern Angola and Zambia. It...

Blue Nile patas monkey

Blue Nile patas monkey

Erythrocebus poliophaeus

The Blue Nile patas monkey (*Erythrocebus poliophaeus*) or Heuglin's patas monkey is

a species of Old World monkey found in Africa along the Blue Nile river valley in Ethiopia, Sudan, and potentially South Sudan. While first described in 1862, it...
Blue Whale
Blue Whale
Balaenoptera musculus

The Blue whale (*Balaenoptera musculus*) is a marine mammal and a baleen whale. It is the largest animal known to have ever existed. The Blue whale was once abundant in nearly all the Earth's oceans until the end of the 19th century. It was hunted...
Blue Wildebeest
Blue Wildebeest
Connochaetes taurinus

The Blue wildebeest (*Connochaetes taurinus*) is a large antelope and one of the two species of wildebeest. Blue wildebeest are found in short-grass plains, thriving in areas that are neither too wet nor too arid. Three African populations of this...
Blue-Eyed Black Lemur
Blue-Eyed Black Lemur
Eulemur flavifrons

The blue-eyed black lemur (*Eulemur flavifrons*), also known as the Sclater's lemur, is a species of true lemur. It can attain a body length of 39–45 cm, a tail length of 51–65 cm- a total length of 90–100 cm, and a weight of 1.8–1.9 kg. Being a...
Blue-eyed cuscus
Blue-eyed cuscus
Phalanger matabiru

The blue-eyed cuscus (*Phalanger matabiru*) is a species of marsupial in the family Phalangeridae. It is endemic to the two small islands of Ternate and Tidore, west of the island of Halmahera in North Maluku province, Indonesia.
Blue-eyed spotted cuscus
Blue-eyed spotted cuscus
Spilocuscus wilsoni

The blue-eyed spotted cuscus or Biak spotted cuscus is a species of marsupial in the family Phalangeridae. It is endemic to the islands of Biak and Supiori in the Cenderawasih Bay in West Papua of Indonesia. Only scientifically described in 2004,...
Blunt-eared bat
Blunt-eared bat
Tomopeas ravus

The blunt-eared bat or Peruvian crevice-dwelling bat (*Tomopeas ravus*) is a species of bat in the family Molossidae. It is monotypic within the genus *Tomopeas* and subfamily Tomopeatinae. It is endemic to Peru, where it is considered critically...
Blyth's horseshoe bat
Blyth's horseshoe bat
Rhinolophus lepidus

Blyth's horseshoe bat (*Rhinolophus lepidus*) is a species of bat in the family Rhinolophidae. It is found across southern Asia from Afghanistan to Vietnam. The species can be identified from its pointed, bifid sella.
Blyth's vole
Blyth's vole
Phaiomys leucurus

Blyth's vole (*Phaiomys leucurus*) is a species of rodent in the family Cricetidae. It is the only species in the genus *Phaiomys*. It is found in mountainous regions in northern India, Nepal and China. It is a burrowing rodent and lives in small...

Bobak Marmot
Bobak Marmot
Marmota bobak

The Bobak marmot (*Marmota bobak*) is a social animal that hibernates for more than half the year. It is a symbol of Luhansk Oblast, Ukraine, as is found on its coat of arms and the coats of arms of some of its districts. It is also depicted on the...

Bobcat
Bobcat
Lynx rufus

The bobcat (*Lynx rufus*) is a medium-sized cat native to North America. It is featured in some stories of the indigenous peoples of North and Central America and in the folklore of European-descended inhabitants of the Americas.

Bobrinski's jerboa
Bobrinski's jerboa
Allactodipus bobrinskii

Bobrinski's jerboa (*Allactodipus bobrinskii*) is a species of rodent in the family Dipodidae. It is monotypic within the genus *Allactodipus*. It is found in Turkmenistan and Uzbekistan.

Bobrinski's serotine
Bobrinski's serotine
Eptesicus bobrinskoi

Bobrinski's serotine (*Eptesicus bobrinskoi*) is a species of vesper bat. It can be found in Iran and Kazakhstan.

Bocage's African fat mouse
Bocage's African fat mouse
Steatomys bocagei

Bocage's African fat mouse (*Steatomys bocagei*) is an extant species of rodent indigenous to Angola, the Republic of the Congo, and the Democratic Republic of the Congo (DRC). Given the distribution of the species across the Angolan Plateau and...

Bocage's mole-rat
Bocage's mole-rat
Fukomys bocagei

Bocage's mole-rat (*Fukomys bocagei*) is a species of rodent in the family Bathyergidae. It is found in Angola, Namibia, possibly Democratic Republic of the Congo, and possibly Zambia. Its natural habitats are subtropical or tropical dry forests,...

Bocage's rock rat
Bocage's rock rat
Aethomys bocagei

Bocage's rock rat (*Aethomys bocagei*) is a species of rodent in the family Muridae. It is found in Angola and Democratic Republic of the Congo. Its natural habitats are subtropical or tropical dry forests, dry savanna, and moist savanna.

Bocharic myotis
Bocharic myotis
Myotis bucharensis

The Bocharic myotis or Bokhara whiskered bat (*Myotis bucharensis*) is a species of mouse-eared bat in the family Vespertilionidae, described in 1950, and indigenous to Tajikistan, and Uzbekistan.
Boeadi's roundleaf bat

Boeadi's roundleaf bat
Hipposideros boeadii

Boeadi's roundleaf bat (*Hipposideros boeadii*) is a species of roundleaf bat found in Indonesia.

Boehm's bush squirrel
Boehm's bush squirrel
Paraxerus boehmi

Boehm's bush squirrel (*Paraxerus boehmi*) is a species of rodent in the family Sciuridae found in Burundi, the Democratic Republic of the Congo, Kenya, Rwanda, South Sudan, Tanzania, Uganda, and Zambia. Its natural habitats are subtropical or...

Boehm's gerbil
Boehm's gerbil
Gerbilliscus boehmi

Boehm's gerbil (*Gerbilliscus boehmi*) is a species of rodent found in Angola, Burundi, Democratic Republic of the Congo, Kenya, Malawi, Mozambique, Rwanda, Tanzania, Uganda, and Zambia. Its natural habitats are dry savanna, moist savanna, and...

Bogotá grass mouse
Bogotá grass mouse
Neomicroxus bogotensis

The Bogotá grass mouse or Bogotá akodont, (*Neomicroxus bogotensis*) is a species of rodent in the family Cricetidae. It is found in the Andes eastern and central Colombia and northwestern Venezuela. Alavarado-Serrano and D'Elía (2013) have...

Bogotá yellow-shouldered bat
Bogotá yellow-shouldered bat
Sturnira bogotensis

The Bogotá yellow-shouldered bat (*Sturnira bogotensis*) is a species of bat in the family Phyllostomidae. It is found in Colombia, Ecuador, Peru, and Venezuela at altitudes from 300 m to above 2000 m, particularly in cloud forest. The species is...

Bohor Reedbuck
Bohor Reedbuck
Redunca redunca

The bohor reedbuck (*Redunca redunca*) is an antelope native to central Africa. The animal is placed under the genus *Redunca* and in the family Bovidae. It was first described by German zoologist and botanist Peter Simon Pallas in 1767. The bohor...

Bokermann's nectar bat
Bokermann's nectar bat
Lonchophylla bokermanni

Bokermann's nectar bat (*Lonchophylla bokermanni*) is a bat species from South America. It is endemic to Brazil. It feeds on nectar, and is listed as an endangered species.

Bokhara horseshoe bat
Bokhara horseshoe bat
Rhinolophus bocharicus

The Bokhara horseshoe bat (*Rhinolophus bocharicus*) is a species of bat in the family Rhinolophidae. It is found in Afghanistan, Turkmenistan, Uzbekistan, and possibly in Iran and Pakistan.

Bolam's mouse
Bolam's mouse

Pseudomys bolami

Bolam's mouse (*Pseudomys bolami*) is a species of nocturnal, burrowing rodent in the family Muridae that inhabits the semi arid and southern arid regions of Australia. It has a number of physiological and behavioural adaptations developed to cope...

Bolivian bamboo rat

Bolivian bamboo rat

Dactylomys boliviensis

The Bolivian bamboo rat (*Dactylomys boliviensis*), is a species of spiny rat from South America.

Bolivian big-eared mouse

Bolivian big-eared mouse

Auliscomys boliviensis

The Bolivian big-eared mouse (*Auliscomys boliviensis*) is a species of rodent in the family Muridae. It is found in Bolivia, Chile, and Peru.

Bolivian chinchilla rat

Bolivian chinchilla rat

Abrocoma boliviensis

The Bolivian chinchilla rat (*Abrocoma boliviensis*) is a species of chinchilla rat in the family Abrocomidae. It is found only in Manuel María Caballero Province, Bolivia. Its natural habitat is the rocky areas of cloud forests in Bolivia's...

Bolivian red howler

Bolivian red howler

Alouatta sara

The Bolivian red howler (*Alouatta sara*) is a species of howler monkey, a type of New World monkey, endemic to Bolivia. It can be found in rain forests, including riverine and seasonally flooded forests.

Bolivian River Dolphin

Bolivian River Dolphin

Inia boliviensis

The Bolivian river dolphin (*Inia boliviensis*) is one of four freshwater river dolphin species in South America. Locally known as bufeo, the Bolivian river dolphin was briefly thought to be a subspecies (as *I. geoffrensis boliviensis*) of the...

Bolivian squirrel

Bolivian squirrel

Sciurus ignitus

The Bolivian squirrel (*Sciurus ignitus*) is a tree squirrel that is endemic to South America. Little is known of the species, which may represent a species complex.

Bolivian tuco-tuco

Bolivian tuco-tuco

Ctenomys boliviensis

The Bolivian tuco-tuco (*Ctenomys boliviensis*) is a species of rodent in the family Ctenomyidae. It is found in Argentina, Bolivia, and Paraguay.

Bolivian vesper mouse

Bolivian vesper mouse

Calomys boliviae

The Bolivian vesper mouse (*Calomys boliviae*) is a species of rodent in the family Cricetidae. It is found in Argentina, Bolivia and perhaps Peru.

Bonda mastiff bat
Bonda mastiff bat
Molossus currentium

The Bonda mastiff bat or Thomas's mastiff bat (*Molossus currentium*), is a species of mastiff bat from South and Central America. It is named for the town of Bonda, near Santa Marta in Colombia.

Bonetto's tuco-tuco
Bonetto's tuco-tuco
Ctenomys bonettoi

Bonetto's tuco-tuco (*Ctenomys bonettoi*) is a species of rodent in the family Ctenomyidae. It is endemic to Argentina.

Bongo Antelope
Bongo Antelope
Tragelaphus eurycerus

The bongo (*Tragelaphus eurycerus*) is a herbivorous, mostly nocturnal forest ungulate. Bongos are characterised by a striking reddish-brown coat, black and white markings, white-yellow stripes and long slightly spiralled horns. They are the only...

Bongolava mouse lemur
Bongolava mouse lemur
Microcebus bongolavensis

The Bongolava mouse lemur (*Microcebus bongolavensis*) is a species of mouse lemur endemic to Madagascar. It lives in western deciduous forest within a limited range, including Bongolava Forest and Ambodimahabibo Forest between the Sofia River and...

Bonin flying fox
Bonin flying fox
Pteropus pselaphon

The Bonin flying fox, Bonin fruit bat (*Pteropus pselaphon*), or in Japanese Ogasawara giant bat (オガサワラオコウモリ, Ogasawara ōkōmori) is a species of flying fox in the family Pteropodidae. It is endemic to four islands (Chichijima, Hahajima, North Iwo...

Bonnet Macaque
Bonnet Macaque
Macaca radiata

The bonnet macaque (*Macaca radiata*), also known as zati, is a species of macaque endemic to southern India. Its distribution is limited by the Indian Ocean on three sides and the Godavari and Tapti Rivers, along with its related competitor the...

Bonobo
Bonobo
Pan paniscus

The bonobo (*Pan paniscus*) is an endangered species of great ape found in a 500,000 km² (190,000 sq mi) area of the Congo Basin in the Democratic Republic of the Congo, Central Africa. Bonobos are one of the closest living relatives to humans,...

Bontebok
Bontebok
Damaliscus pygargus

The bontebok (*Damaliscus pygargus*) is an antelope found in South Africa, Lesotho and Namibia. *D. pygargus* has two subspecies; the nominate subspecies (*D. p. pygargus*), occurring naturally in the Fynbos and Renosterveld areas of the Western...

Bonthain rat
Bonthain rat
Rattus bontanus

The Bonthain rat (*Rattus bontanus*) is a species of rodent in the family Muridae. It is found only in southwestern Sulawesi, Indonesia.

Boodie
Boodie
Bettongia lesueur

The boodie (*Bettongia lesueur*) is a small, furry, rat-like mammal native to Australia. It is a member of the rat-kangaroo family (Potoroidae). The boodie lives in burrows and, like most marsupials, carries its young in a pouch.

Booted macaque
Booted macaque
Macaca ochreata

The booted macaque (*Macaca ochreata*) is a macaque of the Sulawesi Island, Indonesia. This Old World monkey is diurnal and spends most of the day in the trees. It is 50-59 cm long plus a tail of 35-40 cm. The booted macaque feeds on figs, buds,...

Boraha mouse lemur
Boraha mouse lemur
Microcebus boraha

The Nosy Boraha mouse lemur (*Microcebus boraha*) is a species of mouse lemur described in 2016 from Madagascar. It was discovered by a team of researchers at the German Primate Center. It was initially discovered among closely related species...

Boreal woodland caribou
Boreal woodland caribou
Rangifer tarandus caribou

The boreal woodland caribou (*Rangifer tarandus caribou*), also known as woodland caribou, boreal forest caribou and forest-dwelling caribou, is a North American subspecies of reindeer (or caribou in North America) found only in Canada. Unlike the...

Bornean Bearded Pig
Bornean Bearded Pig
Sus barbatus

The Bornean bearded pig (*Sus barbatus*), also known ambiguously as the Sunda bearded pig or simply bearded pig, is a species in the pig genus, *Sus*. It can be recognized by its prominent beard. It also sometimes has tassels on its tail. It is found...

Bornean Clouded Leopard
Bornean Clouded Leopard
Neofelis diardi borneensis

The Bornean clouded leopard (*Neofelis diardi borneensis*) is a subspecies of the Sunda clouded leopard. It differs from the Batu-Sumatran clouded leopard in the shape and frequency of spots, as well as in cranio-mandibular and dental characters.

Bornean Ferret Badger
Bornean Ferret Badger
Melogale everetti

The Bornean ferret badger (*Melogale everetti*) is a small, nocturnal mammal that is endemic to the island of Borneo. It is a member of the family Mustelidae and is one

of five species of the genus *Melogale*.

Bornean horseshoe bat

Bornean horseshoe bat

Rhinolophus borneensis

The Bornean horseshoe bat (*Rhinolophus borneensis*) is a species of bat in the family *Rhinolophidae*. It is found in Brunei, Cambodia, Indonesia, Laos, Malaysia, and Vietnam.

Bornean mountain ground squirrel

Bornean mountain ground squirrel

Dremomys everetti

The Bornean mountain ground squirrel (*Sundasciurus everetti*) is a species of rodent in the family *Sciuridae*. The scientific name commemorates British colonial administrator and zoological collector Alfred Hart Everett.

Bornean Orangutan

Bornean Orangutan

Pongo pygmaeus

The Bornean orangutan (*Pongo pygmaeus*) is the world's largest mammal that lives in a tree and is also one of the world's greatest apes. Together with the Sumatran orangutan and Tapanuli orangutan, it belongs to the only genus of great apes native...

Bornean pithecheirops

Bornean pithecheirops

Pithecheirops otion

The Bornean pithecheirops (*Pithecheirops otion*) is a species of rodent found in Borneo. It is monotypic in the genus *Pithecheirops*. *Pithecheirops* is closely related to the genus *Pithecheir*, and the name is derived from *Pithecheir* and the Greek...

Bornean pygmy shrew

Bornean pygmy shrew

Suncus hosei

The Bornean pygmy shrew (*Suncus hosei*) is a species of shrew in the family *Soricidae*. It was named for zoologist Charles Hose.

Bornean rhinoceros

Bornean rhinoceros

Dicerorhinus sumatrensis harrissoni

The Bornean rhinoceros (*Dicerorhinus sumatrensis harrissoni*), also known as the eastern Sumatran rhinoceros or eastern hairy rhinoceros, is one of three subspecies of Sumatran rhinoceros. The subspecies may be functionally extinct, with only one...

Bornean shrew

Bornean shrew

Crocidura foetida

The Bornean shrew (*Crocidura foetida*) is a species of mammal in the family *Soricidae*. It is found only on Borneo, throughout most of the island; it may or may not be present in Brunei. Multilocus phylogenies reveal paraphyly in *C. foetida*....

Bornean smooth-tailed treeshrew

Bornean smooth-tailed treeshrew

Dendrogale melanura

The Bornean smooth-tailed treeshrew (*Dendrogale melanura*) is a species of treeshrew in the family *Tupaiaidae*. It is endemic to Borneo. Its natural habitat is subtropical or tropical dry forests. It is threatened by habitat loss.

Bornean water shrew
Bornean water shrew
Chimarrogale phaeura

The Bornean water shrew (*Chimarrogale phaeura*) is a species of mammal in the family Soricidae. It is endemic to Malaysia. Its natural habitat is rivers. It is threatened by habitat loss.

Bornean White-Bearded Gibbon
Bornean White-Bearded Gibbon
Hylobates albibarbis

The Bornean white-bearded gibbon, (*Hylobates albibarbis*), also known as the Bornean agile gibbon or southern gibbon, is a species of gibbon endemic to southern Borneo. It is an endangered species, due to the undergoing logging of the tropical...

Bornean yellow muntjac
Bornean yellow muntjac
Muntiacus atherodes

The Bornean yellow muntjac (*Muntiacus atherodes*) is a muntjac species endemic to the moist forests of Borneo.

Borneo black-banded squirrel
Borneo black-banded squirrel
Callosciurus orestes

The Borneo black-banded squirrel (*Callosciurus orestes*) is a species of rodent in the family Sciuridae. It is endemic to northern Borneo.

Borneo Elephant
Borneo Elephant
Elephas maximus borneensis

The smallest elephant in Asia, the Borneo elephant (*Elephas maximus borneensis*), or otherwise known as the Pygmy elephant is a rather charming and adorable creature. Compared to other Asian elephants, Borneo elephants are more gentle-natured...

Borneo fruit bat
Borneo fruit bat
Aethalops aequalis

The Borneo fruit bat (*Aethalops aequalis*) is a species of megabat found in the mountains of Borneo, specifically East Malaysia and Brunei. It is considered a subspecies of *Aethalops alecto* by some authors. The Borneo fruit bat typically roosts in...

Borneo roundleaf bat
Borneo roundleaf bat
Hipposideros doriae

The Borneo roundleaf bat or Bornean leaf-nosed bat (*Hipposideros doriae*) is a species of bat in the family Hipposideridae. It is found in Borneo, Sumatra and Peninsular Malaysia. *Hipposideros sabanus* is a synonym of this species.

Bosavi woolly rat
Bosavi woolly rat
Mallomys sp.

The Bosavi woolly rat is a species of rodent that was discovered deep in the jungle of Papua New Guinea in 2009. It is believed to belong to the genus *Mallomys*, within the family Muridae, according to initial investigation, although this has yet...

Botswanan long-eared bat
Botswanan long-eared bat
Laephotis botswanae

The Botswanan long-eared bat (*Laephotis botswanae*) is a species of vesper bat in the family Vespertilionidae. It can be found in the following countries: Botswana, Democratic Republic of the Congo, Malawi, Namibia, South Africa, Tanzania, Zambia,...

Botta's gerbil
Botta's gerbil
Dipodillus bottai

Botta's gerbil (*Dipodillus bottai*) is a species of gerbil endemic to Sudan and possibly northern Kenya. According to the International Union for Conservation of Nature, it was listed as Least Concern in 1996 until it was listed as Data Deficient...

Botta's Pocket Gopher
Botta's Pocket Gopher
Thomomys bottae

Botta's pocket gopher (*Thomomys bottae*) is a burrowing rodent native to western North America. It is also known in some sources as Valley pocket gopher, particularly in California. Both the specific and common names of this species honor...

Botta's serotine
Botta's serotine
Eptesicus bottae

Botta's serotine (*Eptesicus bottae*) is a species of vesper bat, one of 25 in the genus *Eptesicus*. It is found in rocky areas and temperate desert.

Bottego's shrew
Bottego's shrew
Crocidura bottegi

Bottego's shrew (*Crocidura bottegi*) is a species of mammal in the family Soricidae. It is found 1,750 metres (5,740 ft) above sea level in Ethiopia and possibly Kenya particularly northeast of Lake Turkana and one in Kenya from Marsabit.

Bougainville monkey-faced bat
Bougainville monkey-faced bat
Pteralopex anceps

The Bougainville monkey-faced bat or Bougainville flying monkey (*Pteralopex anceps*) is a megabat endemic to Bougainville Island of Papua New Guinea and Choiseul Island of the Solomon Islands in Melanesia. It inhabits mature forests in upland...

Bougainville mosaic-tailed rat
Bougainville mosaic-tailed rat
Melomys bougainville

The Bougainville mosaic-tailed rat (*Melomys bougainville*) is a species of rodent in the family Muridae. It is found only in Papua New Guinea.

Bougainville naked-tailed rat
Bougainville naked-tailed rat
Solomys salebrosus

The Bougainville naked-tailed rat (*Solomys salebrosus*) or Bougainville giant rat is a species of rodent in the family Muridae. It is found in Papua New Guinea and the Solomon Islands.

Bourlon's genet
Bourlon's genet
Genetta bourloni

Bourlon's genet (*Genetta bourloni*) is a genet species native to the Upper Guinean forests. It is known from only 29 zoological specimens in natural history museum and has been described as a new *Genetta* species in 2003. It is listed as Vulnerable...

Bourret's horseshoe bat

Bourret's horseshoe bat

Rhinolophus paradoxolophus

Bourret's horseshoe bat (*Rhinolophus paradoxolophus*) is a species of horseshoe bat native to Southeast Asia. The name "paradoxolophus " is derived from the Greek words *paradoxos*, meaning "contrary to expectation", and *lophos*, meaning "crest"....

Bouvier's Red Colobus

Bouvier's Red Colobus

Piliocolobus bouvieri

Bouvier's red colobus (*Piliocolobus bouvieri*) is a species of colobus monkey rediscovered in the Republic of the Congo in 2015, after four decades without a confirmed sighting.

Bower's white-toothed rat

Bower's white-toothed rat

Berylmys bowersi

Bower's white-toothed rat (*Berylmys bowersi*) is a species of rodent in the family Muridae native to southeast Asia.

Bowhead Whale

Bowhead Whale

Balaena mysticetus

The Bowhead whale (*Balaena mysticetus*) is a species of baleen whale and the only living representative of the genus *Balaena*. They are the only baleen whale endemic to the Arctic and subarctic waters and are named after their characteristic...

Boyacá spiny rat

Boyacá spiny rat

Proechimys chrysaeolus

The Boyacá spiny rat (*Proechimys chrysaeolus*) is a species of rodent in the family Echimyidae. It is endemic to Colombia; it is found primary forest in Carare River valley, Boyacá Department, 100 to 500 meters above sea level. Nocturnal,...

Brachyprotoma

Brachyprotoma

Brachyprotoma obtusata

Brachyprotoma obtusata (also known as the short-faced skunk) is an extinct genus of skunk of the Pleistocene epoch what is now North America.

Brachytarsomys mahajambaensis

Brachytarsomys mahajambaensis

Brachytarsomys mahajambaensis

Brachytarsomys mahajambaensis is an extinct rodent from northwestern Madagascar. It is known from nine isolated molars found in several sites during fieldwork that started in 2001. First described in 2010, it is placed in the genus

Brachytarsomys...

Brahma white-bellied rat

Brahma white-bellied rat

Niviventer brahma

The Brahma white-bellied rat (*Niviventer brahma*) is a species of rodent in the family Muridae. It is found in northeastern India, northern Myanmar, and

southwestern China (Yunnan). It lives in various forest habitats at elevations of...

Brandt's Bat

Brandt's Bat

Myotis brandtii

Brandt's bat (*Myotis brandtii*) is a species of vesper bat native to Europe and parts of western Asia. It is named for the German zoologist Johann Friedrich von Brandt.

Brandt's guirara

Brandt's guirara

Euryzomatomys guirara

Brandt's guirara, *Euryzomatomys guirara*, is a Brazilian spiny rat species. It was formerly considered conspecific with *E. spinosus*.

Brandt's hedgehog

Brandt's hedgehog

Paraechinus hypomelas

Brandt's hedgehog (*Paraechinus hypomelas*) is a species of desert hedgehog native to parts of the Middle East and Central Asia. Its common name derives from its having first been described by Johann Friedrich von Brandt, a director of the...

Brandt's vole

Brandt's vole

Lasiopodomys brandtii

Brandt's vole, (*Lasiopodomys brandtii*), also known as the steppe vole, is a species of rodent in the family Cricetidae. It is native to shrublands and grasslands in Russia, Mongolia and northern China.

Brants's climbing mouse

Brants's climbing mouse

Dendromus mesomelas

Brants's climbing mouse (*Dendromus mesomelas*) is a species of rodent in the family Nesomyidae. It is found in Botswana, Democratic Republic of the Congo, Kenya, Malawi, Mozambique, Namibia, South Africa, Tanzania, and Zambia. Its natural habitats...

Brants's Whistling Rat

Brants's Whistling Rat

Parotomys brantsii

Brants's whistling rat (*Parotomys brantsii*) is a rodent from Africa. It was first described in 1834 by the Scottish zoologist Andrew Smith who named it in honour of the Dutch zoologist and author Anton Brants.

Brazilian arboreal mouse

Brazilian arboreal mouse

Rhagomys rufescens

The Brazilian arboreal mouse (*Rhagomys rufescens*) is a South American rodent species of the family Cricetidae. It is found in the Atlantic Forest of southeast Brazil, often close to bamboo thickets. It can be distinguished from *Rhagomys*...

Brazilian big-eyed bat

Brazilian big-eyed bat

Chiroderma doriae

The Brazilian big-eyed bat (*Chiroderma doriae*) is a species of phyllostomid bat from South America. The scientific name honours Italian naturalist Giacomo Doria.

Brazilian brown bat

Brazilian brown bat

Eptesicus brasiliensis

The Brazilian brown bat (*Eptesicus brasiliensis*), is a bat species from South and Central America.

Brazilian funnel-eared bat

Brazilian funnel-eared bat

Natalus macrourus

The Brazilian funnel-eared bat (*Natalus macrourus*) is a South American bat species found in eastern Brazil and in Paraguay. It roosts in caves, which makes it vulnerable to disturbance of these scarce sites, and in particular, to extermination...

Brazilian gracile opossum

Brazilian gracile opossum

Gracilinanus microtarsus

The Brazilian gracile opossum (*Gracilinanus microtarsus*) is a species of small opossum from Brazil.

Brazilian Guinea Pig

Brazilian Guinea Pig

Cavia aperea

The Brazilian guinea pig (*Cavia aperea*) (preá in Portuguese) is a guinea pig species found in Argentina, Brazil, Bolivia, Colombia, Ecuador, Guyana, Paraguay, Uruguay and Venezuela. *Cavia aperea* has been successfully mated to the domestic guinea...

Brazilian shrew mouse

Brazilian shrew mouse

Blarinomys breviceps

The Brazilian shrew mouse (*Blarinomys breviceps*), also known as the blarinine akodont, is a rodent in the tribe Akodontini from the Atlantic Forest of eastern and southeastern Brazil. It is the only species in the genus *Blarinomys*.

Phylogenetic...

Brazilian slender opossum

Brazilian slender opossum

Marmosops paulensis

The Brazilian slender opossum (*Marmosops paulensis*) is an opossum species from South America. It is found in moist montane forest in the Atlantic Forest region of southeastern Brazil, including the states of Minas Gerais, Rio de Janeiro, São...

Brazilian spiny tree-rat

Brazilian spiny tree-rat

Makalata didelphoides

The Brazilian spiny tree rat (*Makalata didelphoides*) is a species of rodent in the family Echimyidae. It is found in Bolivia, Brazil, French Guiana, Guyana, Suriname, Venezuela, and Trinidad and Tobago where it lives in lowland tropical...

Brazilian squirrel

Brazilian squirrel

Sciurus aestuans

The Brazilian squirrel (or Guianan squirrel) (*Sciurus aestuans*) is a tree squirrel in the genus *Sciurus* endemic to South America. It is found in South-eastern Colombia, Brazil, Guyana, French Guiana, Suriname and Venezuela. It is a dark brown...

Brazilian Tapir

Brazilian Tapir

Tapirus terrestris

Brazilian tapir (*Tapirus terrestris*) is a species of rhinoceros family with a characteristic fleshy trunk, which is prehensile and able to grasp objects such as leaves. It is the largest surviving native terrestrial mammal in the Amazon....

Brazilian Three-Banded Armadillo

Brazilian Three-Banded Armadillo

Tolypeutes tricinctus

The Brazilian three-banded armadillo (*Tolypeutes tricinctus*) is an armadillo species endemic to eastern Brazil, where it is known as tatu-bola (Portuguese pronunciation: , lit. ball armadillo). It is one of only two species of armadillo that can...

Brazilian tuco-tuco

Brazilian tuco-tuco

Ctenomys brasiliensis

The Brazilian tuco-tuco (*Ctenomys brasiliensis*) is a tuco-tuco species from South America. It is found mainly in the state of Minas Gerais in southeastern Brazil, though Charles Darwin mentions it during his trip through present-day Uruguay.

Brazilian yellow-toothed cavy

Brazilian yellow-toothed cavy

Galea flavidens

The Brazilian yellow-toothed cavy (*Galea flavidens*) is a cavy species from South America. It is found in Brazil. *Galea flavidens* is a yellow-toothed-cavy. *G. flavidens* lives in Brazil. This species is listed by the IUCN as Least Concern, in spite...

Bridges's degu

Bridges's degu

Octodon bridgesi

Bridges's degu (*Octodon bridgesi*) is a species of rodent in the family Octodontidae. It is found in southern Chile. The species was named after Thomas Bridges.

Bridled Nail-Tail Wallaby

Bridled Nail-Tail Wallaby

Onychogalea fraenata

The bridled nail-tail wallaby (*Onychogalea fraenata*), also known as the bridled nail-tailed wallaby, bridled nailtail wallaby, bridled wallaby, merrin, and flashjack, is a vulnerable species of macropod. It is a small wallaby found in three...

Bristle-spined rat

Bristle-spined rat

Chaetomys subspinosus

The bristle-spined rat (*Chaetomys subspinosus*) is an arboreal rodent from Atlantic forest in eastern Brazil. Also known as the bristle-spined porcupine or thin-spined porcupine, it is the only member of the genus *Chaetomys* and the subfamily...

British Columbia wolf

British Columbia wolf

Canis lupus columbianus

The British Columbia wolf (*Canis lupus columbianus*) is a subspecies of gray wolf which lives in a narrow region that includes those parts of the mainland coast and near-shore islands that are covered with temperate rainforest, which extends from...

Broad-eared bat

Broad-eared bat

Nyctinomops laticaudatus

The broad-eared bat or broad-tailed bat (*Nyctinomops laticaudatus*) is a species of free-tailed bat from the Americas.

Broad-eared horseshoe bat

Broad-eared horseshoe bat

Rhinolophus euryotis

The broad-eared horseshoe bat (*Rhinolophus euryotis*) is a species of bat in the family Rhinolophidae. It is found in Indonesia and Papua New Guinea.

Broad-footed climbing mouse

Broad-footed climbing mouse

Rhipidomys latimanus

The broad-footed climbing mouse (*Rhipidomys latimanus*) is a species of rodent in the family Cricetidae. It is found in Colombia, Ecuador, Panama and Peru.

Broad-Footed Mole

Broad-Footed Mole

Scapanus latimanus

The broad-footed mole (*Scapanus latimanus*) was a former mole species that has since been split into three distinct species in the genus *Scapanus* : Northern broad-footed mole, *Scapanus latimanus*, ranging from southern Oregon to central...

Broad-headed pipistrelle

Broad-headed pipistrelle

Parahypsugo crassulus

The broad-headed serotine (*Nycticeinops crassulus*), formerly known as the broad-headed pipistrelle, is a species of vesper bat in the family Vespertilionidae. It is found in Angola, Cameroon, Republic of the Congo, Democratic Republic of the...

Broad-headed spiny rat

Broad-headed spiny rat

Clyomys laticeps

The broad-headed spiny rat (*Clyomys laticeps*) is a spiny rat species from South America. The etymology of the species name is the Latin word *laticeps* meaning "wide-headed".

Broad-striped dasyure

Broad-striped dasyure

Paramurexia rothschildi

The broad-striped dasyure (*Murexia rothschildi*) is a species of marsupial in the family Dasyuridae. It is endemic to Papua New Guinea. Its natural habitat is subtropical or tropical dry forests. Some authorities place it in the monotypic genus...

Broad-striped Malagasy mongoose

Broad-striped Malagasy mongoose

Galidictis fasciata

The broad-striped Malagasy mongoose or broad-striped vontsira (*Galidictis fasciata*) is a species of Galidiinae, a subfamily of mongoose-like euplerids native to Madagascar. The species contains two known subspecies: *Galidictis fasciata fasciata*...

Broad-striped tube-nosed fruit bat

Broad-striped tube-nosed fruit bat

Nyctimene aello

The broad-striped tube-nosed fruit bat (*Nyctimene aello*), also known as the greater tube-nosed bat is a species of megabat in the genus *Nyctimene*. It is found

in West Papua, Indonesia, Papua New Guinea, and central Philippines. In 1912, Andersen...
Broad-toothed mouse
Broad-toothed mouse
Mastacomys fuscus

The broad-toothed rat (*Mastacomys fuscus*) is a species of rodent in the family Muridae.
Broad-toothed tailless bat
Broad-toothed tailless bat
Anoura latidens

The broad-toothed tailless bat (*Anoura latidens*) is a species of bats in the family Phyllostomidae. It is found in Colombia, Peru, and Venezuela.
Brock's yellow-eared bat
Brock's yellow-eared bat
Vampyriscus brocki

Brock's yellow-eared bat (*Vampyriscus brocki*) is a species of bat in the family Phyllostomidae, the leaf-nosed bats. It is native to South America, where it occurs in Brazil, Colombia, Guyana, Suriname, French Guiana, and Peru. This bat lives in...
Brockman's gerbil
Brockman's gerbil
Gerbillus brockmani

Brockman's gerbil (*Gerbillus brockmani*) is distributed mainly in Somaliland and is only known from the type locality, Burao in central Somaliland.
Brockman's rock mouse
Brockman's rock mouse
Myomyscus brockmani

Brockman's rock mouse, or Brockman's myomyscus (*Ochromyscus brockmani*) is a species of rodent in the family Muridae. It is found in Central African Republic, Ethiopia, Kenya, Somalia, Sudan, Tanzania, Uganda, and possibly Republic of the...
Bronze quoll
Bronze quoll
Dasyurus spartacus

The bronze quoll (*Dasyurus spartacus*) is a species of quoll found only in the Trans-Fly savanna and grasslands of New Guinea and West Papua.
Bronze tube-nosed bat
Bronze tube-nosed bat
Murina aenea

The bronze tube-nosed bat (*Murina aenea*) is a species of vesper bat in the family Vespertilionidae. It is found only in Malaysia.
Brooke's duiker
Brooke's duiker
Cephalophus brookei

The Brooke's duiker (*Cephalophus brookei*) is a species of antelope. It is distributed throughout Liberia, Sierra Leone and Côte d'Ivoire. It was elevated to species status.
Brooke's squirrel
Brooke's squirrel
Sundasciurus brookei

Brooke's squirrel (*Sundasciurus brookei*) is a species of rodent in the family Sciuridae. It is found in Indonesia and Malaysia. Its natural habitat is

subtropical or tropical dry forests. It is threatened by habitat loss. It was named for...

Brooks's dyak fruit bat

Brooks's dyak fruit bat

Dyacopterus brooksi

Brooks's dyak fruit bat (*Dyacopterus brooksi*) is a species of megabat in the family Pteropodidae endemic to Sumatra. It was formerly included as a subspecies of the dayak fruit bat. It is named for Cecil Joslin Brooks, who collected the type...

Broom hare

Broom hare

Lepus castroviejoi

The broom hare (*Lepus castroviejoi*) is a species of hare endemic to northern Spain.

Brosset's big-eared bat

Brosset's big-eared bat

Micronycteris brosseti

Brosset's big-eared bat (*Micronycteris brosseti*) is a bat species from South America. It is found in Brazil, French Guiana, Guyana and Peru. It feeds on insects and sometimes fruit and the exact population is unknown. The only listed threat is...

Brown Antechinus

Brown Antechinus

Antechinus stuartii

The brown antechinus (*Antechinus stuartii*), also known as Stuart's antechinus and Macleay's marsupial mouse, is a species of small carnivorous marsupial of the family Dasyuridae. The males die after their first breeding season, and the species...

Brown Bear

Brown Bear

Ursus arctos

The Brown bear (*Ursus arctos*) is a large bear species found across Eurasia and North America. In North America, the populations of Brown bears are called Grizzly bears, while the subspecies that inhabits the Kodiak Islands of Alaska is known as...

Brown deer mouse

Brown deer mouse

Peromyscus megalops

The brown deer mouse (*Peromyscus megalops*) is a species of rodent in the family Cricetidae found only in Mexico.

Brown dorcopsis

Brown dorcopsis

Dorcopsis muelleri

The brown dorcopsis (*Dorcopsis muelleri*), also known as the brown forest wallaby, is a species of marsupial in the family Macropodidae. It is endemic to the lowlands of West New Guinea and the nearby Indonesian islands in West Papua of Misool,...

Brown flower bat

Brown flower bat

Erophylla bombifrons

The brown flower bat (*Erophylla bombifrons*) is a species of bat from the family Phyllostomidae. It is native to the island of Hispaniola, present in both Haiti and the Dominican Republic, and Puerto Rico. In most cases, the brown flower bat is...

Brown four-eyed opossum
Brown four-eyed opossum
Metachirus nudicaudatus

The brown four-eyed opossum (*Metachirus nudicaudatus*) is a pouchless marsupial of the family Didelphidae. It is found in different forested habitats of Central and South America, from Nicaragua to Brazil and northern Argentina, including...

Brown fruit-eating bat
Brown fruit-eating bat
Artibeus concolor

The brown fruit-eating bat (*Artibeus concolor*) is a bat species from South America. It is found in Brazil, Colombia, Ecuador, French Guiana, Guyana, Peru, Suriname and Venezuela.

Brown Fur Seal
Brown Fur Seal
Arctocephalus pusillus

The brown fur seal (*Arctocephalus pusillus*), also known as the Cape fur seal, South African fur seal and Australian fur seal, is a species of fur seal.

Brown Greater Galago
Brown Greater Galago
Otolemur crassicaudatus

The Brown greater galago (*Otolemur crassicaudatus*) is a nocturnal primate, the largest in the family of Galagos. As opposed to smaller galago species it would climb, walk, or run rather than leap.

Brown hairy dwarf porcupine
Brown hairy dwarf porcupine
Coendou vestitus

The brown hairy dwarf porcupine (*Coendou vestitus*) is a species of rodent in the family Erethizontidae. Found in the Andes in Colombia and Venezuela, its natural habitat is subtropical or tropical moist lowland forests. It is not easy to study...

Brown Howler
Brown Howler
Alouatta guariba

The brown howler (*Alouatta guariba*), also known as brown howler monkey, is a species of howler monkey, a type of New World monkey that lives in forests in southeastern Brazil and far northeastern Argentina (Misiones). It lives in groups of two...

Brown Hyena
Brown Hyena
Hyaena brunnea

The Brown hyena (*Parahyaena brunnea*) is a species of hyena found in Africa. It is currently the rarest species of hyena. The largest remaining Brown hyena population is located in the southern Kalahari Desert and coastal areas in Southwest Africa.

Brown Long-Eared Bat
Brown Long-Eared Bat
Plecotus auritus

The brown long-eared bat or common long-eared bat (*Plecotus auritus*) is a small Eurasian insectivorous bat. It has distinctive ears, long and with a distinctive fold. It is extremely similar to the much rarer grey long-eared bat which was only...

Brown mastiff bat

Brown mastiff bat
Promops nasutus

The brown mastiff bat (*Promops nasutus*), is a South American bat species, found in Venezuela, Trinidad, Guyana, Surinam, Brazil, Ecuador, Peru, Bolivia, Paraguay, and northern Argentina.

Brown Mouse Lemur
Brown Mouse Lemur
Microcebus rufus

The brown mouse lemur (*Microcebus rufus*) is a small primate, and like the other mouse lemurs can only be found on the island of Madagascar. They are known also as the rufous mouse lemur, eastern rufous mouse lemur, red mouse lemur, or russet...

Brown Palm Civet
Brown Palm Civet
Paradoxurus jerdoni

The Brown palm civet (*Paradoxurus jerdoni*) is a viverrid from India. It is about as large as the Common palm civet, but with a long and sleek tail.

Brown pipistrelle
Brown pipistrelle
Hypsugo imbricatus

The brown pipistrelle (*Hypsugo imbricatus*) is a species of vesper bat in the family Vespertilionidae. It is found in Indonesia and Malaysia.

Brown Rat
Brown Rat
Rattus norvegicus

The Brown rat (*Rattus norvegicus*) is a widespread species of common rat. One of the largest muroids, it weighs between 140 and 500 g (5 and 17+3/4 oz). Thought to have originated in northern China and neighboring areas, this rodent has now spread...

Brown Spider Monkey
Brown Spider Monkey
Ateles hybridus

The brown spider monkey or variegated spider monkey (*Ateles hybridus*) is a critically endangered species of spider monkey, a type of New World monkey, from forests in northern Colombia and northwestern Venezuela. Like all spider monkeys, it has...

Brown tent-making bat
Brown tent-making bat
Uroderma magnirostrum

The brown tent-making bat (*Uroderma magnirostrum*) is a bat species from South and Central America.

Brown titi
Brown titi
Plecturocebus brunneus

The brown titi (*Plecturocebus brunneus*) is a species of titi, a type of New World monkey, from South America. It is endemic to Brazil. It was originally described as *Callicebus brunneus* in 1842 and transferred to the newly erected genus...

Brown tube-nosed bat
Brown tube-nosed bat
Murina suilla

The brown tube-nosed bat (*Murina suilla*) is a species of vesper bat in the family Vespertilionidae. It can be found in the following countries: Brunei Darussalam,

Indonesia, Malaysia, and the Philippines.
Brown weeper capuchin
Brown weeper capuchin
Cebus brunneus

The brown weeper capuchin (*Cebus brunneus*) or Venezuelan brown capuchin is a species of gracile capuchin monkey endemic to Venezuela, although some sources also consider it to occur on Trinidad.

Brown Woolly Monkey
Brown Woolly Monkey
Lagothrix lagothricha

The common woolly monkey, brown woolly monkey, or Humboldt's woolly monkey (*Lagothrix lagothricha*) is a woolly monkey from Colombia, Ecuador, Peru, Bolivia, Brazil, and Venezuela. It lives in groups of two to 70 individuals, usually splitting...

Brown-backed bearded saki
Brown-backed bearded saki
Chiropotes israelita

The brown-backed bearded saki (*Chiropotes israelita*) is one of five species of bearded saki, a type of New World monkey. It is endemic to the Amazon in north-western Brazil (north of the Rio Negro and west of the Branco River) and southern...

Brown-bellied broad-nosed bat
Brown-bellied broad-nosed bat
Platyrrhinus fusciventris

The brown-bellied broad-nosed bat (*Platyrrhinus fusciventris*) is a species of bat in the family Phyllostomidae. As a phyllostomid bat, it is characterized by a narrow uropatagium which is fringed with hair; a white dorsal stripe; large inner...

Brown-eared woolly opossum
Brown-eared woolly opossum
Caluromys lanatus

The brown-eared woolly opossum (*Caluromys lanatus*), also known as the western woolly opossum, is an opossum from South America. It was first described by German naturalist Ignaz von Olfers in 1818. The opossum is characterized by a brown to...

Brown-headed spider monkey
Brown-headed spider monkey
Ateles fusciceps fusciceps

The brown-headed spider monkey (*Ateles fusciceps fusciceps*), is a critically endangered subspecies of the black-headed spider monkey, a type of New World monkey, found in northwestern Ecuador. Its type locality is at 1500 m in the Hacienda...

Brown-mantled tamarin
Brown-mantled tamarin
Leontocebus fuscicollis

The brown-mantled tamarin (*Leontocebus fuscicollis*), also known as Spix's saddle-back tamarin, is a species of saddle-back tamarin from South America. It is found in Bolivia, Brazil and Peru. The brown-mantled tamarin is sympatric with the pygmy...

Brown-tailed mongoose
Brown-tailed mongoose
Salanoia concolor

The brown-tailed mongoose, brown-tailed vuntsira, Malagasy brown-tailed mongoose, or salano (*Salanoia concolor*) is a species of mammal in the family Eupleridae. It is endemic to Madagascar. Its natural habitat is subtropical or tropical dry...

Brown-Throated Three-Toed Sloth

Brown-Throated Three-Toed Sloth

Bradypus variegatus

The Brown-throated sloth (*Bradypus variegatus*) is the most common of the four species of three-toed sloth. It is native to rainforest canopies of Central and South America. The slowest mammal in the world, the Brown-throated three-toed sloth is...

Brown's pademelon

Brown's pademelon

Thylogale browni

Brown's pademelon (*Thylogale browni*) is a species of marsupial in the family Macropodidae. It is found in New Guinea. Its natural habitats are subtropical or tropical dry forests, dry savanna, subtropical or tropical dry shrubland, and...

Bucepattersonius griserufescens

Bucepattersonius griserufescens

Bucepattersonius griserufescens

Bucepattersonius griserufescens, also known as the gray-bellied akodont or grey-bellied brucie, is a rodent in the cricetid genus *Bucepattersonius* from southeastern Brazil, specifically the states of Minas Gerais, Espírito Santo and Rio de...

Bucepattersonius guarani

Bucepattersonius guarani

Bucepattersonius guarani

Bucepattersonius guarani, also known as the Guaraní akodont or Guaraní brucie, is a South American rodent in the genus *Bucepattersonius*. It is known from a single specimen from Misiones Province in northeastern Argentina, at the southern end of...

Bucepattersonius igniventris

Bucepattersonius igniventris

Bucepattersonius igniventris

Bucepattersonius igniventris, also known as the red-bellied akodont or red-bellied brucie, is a South American rodent in the genus *Bucepattersonius*. It is known only from a few specimens from the type locality in Iporanga, São Paulo, Brazil....

Bucepattersonius iheringi

Bucepattersonius iheringi

Bucepattersonius iheringi

Bucepattersonius iheringi, also known as Ihering's akodont, Ihering's hocicudo, or Ihering's brucie, is a South American rodent in the genus *Bucepattersonius*. It occurs in Santa Catarina and Rio Grande do Sul, southern Brazil, and nearby...

Bucepattersonius misionensis

Bucepattersonius misionensis

Bucepattersonius misionensis

Bucepattersonius misionensis, also known as the Misiones akodont or Misiones brucie, is a rodent species from South America in the genus *Bucepattersonius*. It is known from a single individual collected in Misiones Province in northeastern...

Brukkaros pygmy rock mouse

Brukkaros pygmy rock mouse

Petromyscus monticularis

The Brukkaros pygmy rock mouse (*Petromyscus monticularis*) is a species of rodent in the family Nesomyidae. It is found in Namibia and South Africa. Its natural habitat is subtropical or tropical dry shrubland.

Brumback's night monkey

Brumback's night monkey

Aotus brumbacki

Brumback's night monkey (*Aotus brumbacki*) is a species of night monkey found in Colombia. It has traditionally been considered a subspecies of gray-bellied night monkey, *Aotus lemurinus*. but it has recently been argued that it should be...

Brush mouse

Brush mouse

Peromyscus boylii

The brush mouse (*Peromyscus boylii*) is a species of rodent in the family Cricetidae. It is found in mountainous areas of Mexico and the western United States at altitudes over 2,000 m (6,600 ft).

Brush Rabbit

Brush Rabbit

Sylvilagus bachmani

The brush rabbit (*Sylvilagus bachmani*), or western brush rabbit, or Californian brush rabbit, is a species of cottontail rabbit found in western coastal regions of North America, from the Columbia River in Oregon to the southern tip of the Baja...

Brush-tailed mulgara

Brush-tailed mulgara

Dasymercus blythi

The brush-tailed mulgara (*Dasymercus blythi*), previously the mulgara *Dasymercus cristicauda* is a medium sized carnivorous Australian marsupial species weighing approximately 100 g. The brush-tailed mulgara is sexually dimorphic with males being...

Brush-Tailed Phascogale

Brush-Tailed Phascogale

Phascogale tapoatafa

The brush-tailed phascogale (*Phascogale tapoatafa*), also known by its Australian native name tuan, the common wambenger, the black-tailed mousesack or the black-tailed phascogale, is a rat-sized arboreal carnivorous marsupial of the family...

Brush-tailed rabbit rat

Brush-tailed rabbit rat

Conilurus penicillatus

The brush-tailed rabbit rat (*Conilurus penicillatus*) is a species of rodent in the family Muridae. It is found in Australia and Papua New Guinea.

Brush-Tailed Rock-Wallaby

Brush-Tailed Rock-Wallaby

Petrogale penicillata

The brush-tailed rock-wallaby or small-eared rock-wallaby (*Petrogale penicillata*) is a kind of wallaby, one of several rock-wallabies in the genus *Petrogale*. It inhabits rock piles and cliff lines along the Great Dividing Range from about 100 km...

Bryde's Whale

Bryde's Whale

Balaenoptera brydei

Bryde's whale (BRUU-dəz Brooder's), or the Bryde's whale complex, putatively comprises three species of rorqual and maybe four. The "complex" means the number

and classification remains unclear because of a lack of definitive information and...

Bubal hartebeest

Bubal hartebeest

Alcelaphus buselaphus buselaphus

The bubal hartebeest, also known as northern hartebeest or bubal antelope or simply bubal (*Alcelaphus buselaphus buselaphus*) is the extinct nominal (i.e., first described) subspecies of hartebeest. It was formerly found north of the Saharan...

Buchara shrew

Buchara shrew

Sorex buchariensis

The Buchara shrew or Pamir shrew (*Sorex buchariensis*) is a species of mammal in the family Soricidae. It is found in the Pamir Mountains in Tajikistan, Kyrgyzstan, and Uzbekistan. It lives in mountain birch and poplar forests on slopes and...

Bucharian vole

Bucharian vole

Blanfordimys bucharicus

The Bucharian vole (*Microtus bucharicus*) is a species of rodent in the family Cricetidae. It is found in Afghanistan, Tajikistan, Uzbekistan, and Turkmenistan. Its natural habitat is temperate desert.

Budin's chinchilla rat

Budin's chinchilla rat

Abrocoma budini

Budin's chinchilla rat (*Abrocoma budini*) is a species of chinchilla rat in the family Abrocomidae. Found only in Argentina, the categorization of this species was based on analysis of four specimens which were caught among the rocks in the...

Budin's tuco-tuco

Budin's tuco-tuco

Ctenomys budini

Budin's tuco-tuco (*Ctenomys budini*) was formerly considered a species of rodent in the family Ctenomyidae. It is endemic to southeast Jujuy Province in northwest Argentina. Given the extensive human presence in its limited range, it has been...

Buenos Aires leaf-eared mouse

Buenos Aires leaf-eared mouse

Phyllotis bonariensis

The Buenos Aires leaf-eared mouse (*Phyllotis bonariensis*) is a species of rodent in the family Cricetidae. It is found only in Argentina.

Buergers' tree-kangaroo

Buergers' tree-kangaroo

Dendrolagus goodfellowi buergersi

The Buergers' tree-kangaroo (*Dendrolagus goodfellowi buergersi*) is a subspecies of the Goodfellow's tree-kangaroo from Papua New Guinea, where they dwell mainly in tropical rainforests. Their diet consists of mostly leaves and fruit, which they...

Buettikofer's Epauletted Fruit Bat

Buettikofer's Epauletted Fruit Bat

Epomops buettikoferi

Buettikofer's epauletted fruit bat (*Epomops buettikoferi*) is a species of megabat in the family Pteropodidae. It is found in Ivory Coast, Ghana, Guinea, Guinea-Bissau, Liberia, Nigeria, Senegal, and Sierra Leone. Its natural habitats are...

Buettikofer's shrew

Buettikofer's shrew

Crocidura buettikoferi

Buettikofer's shrew (*Crocidura buettikoferi*) is a species of mammal in the family Soricidae. It is found in southern Nigeria and scantily present in Ivory Coast, Ghana, Guinea, Liberia and Sierra Leone. Its natural habitat is subtropical or...

Buff-bellied climbing mouse

Buff-bellied climbing mouse

Rhipidomys fulviventer

The buff-bellied climbing mouse (*Rhipidomys fulviventer*) is a species of rodent in the family Cricetidae. It is found in Colombia and Venezuela.

Buff-bellied fat-tailed mouse opossum

Buff-bellied fat-tailed mouse opossum

Thylamys venustus

The buff-bellied fat-tailed mouse opossum (*Thylamys venustus*) is a species of opossum in the family Didelphidae. It is found in the transitional and humid forests of northern Argentina and southern Bolivia. Its dorsal fur is cinnamon brown. Most...

Buffoon striped grass mouse

Buffoon striped grass mouse

Lemniscomys macculus

The Buffoon lemniscomys or Buffoon striped grass mouse (*Lemniscomys macculus*) is a species of rodent in the family Muridae. It is found in the Democratic Republic of the Congo, Ethiopia, Kenya, South Sudan, Uganda, and possibly Rwanda. Its natural...

Buffy broad-nosed bat

Buffy broad-nosed bat

Platyrrhinus infuscus

The buffy broad-nosed bat (*Platyrrhinus infuscus*) is a bat species from South America. It is found in Bolivia, western Brazil, Colombia, Ecuador, and Peru.

Buffy flower bat

Buffy flower bat

Erophylla sezekorni

The buffy flower bat (*Erophylla sezekorni*) is a species of bat in the leaf-nosed bat family, Phyllostomidae. It is found in the Bahamas, the Cayman Islands, Cuba, and Jamaica.

Buffy-Headed Marmoset

Buffy-Headed Marmoset

Callithrix flaviceps

The buffy-headed marmoset (*Callithrix flaviceps*) is a rare species of marmoset endemic to the rainforests of south-eastern Brazil. It occurs in southern Espírito Santo and possibly northern Rio de Janeiro and its distribution extends into Minas...

Buffy-tufted marmoset

Buffy-tufted marmoset

Callithrix aurita

The buffy-tufted marmoset (*Callithrix aurita*), also known as the buffy tufted-ear marmoset or white-eared marmoset, is a New World monkey that lives in the forests on the Atlantic coast of southeast Brazil. Of all the marmosets, it has the...

Buka Island mosaic-tailed rat

Buka Island mosaic-tailed rat

Melomys spechti

The Buka Island mosaic-tailed rat or Buka Island melomys (*Melomys spechti*) is a

species of rat in Oceania. It is endemic to Buka Island, in the Autonomous Region of Bougainville in northeastern Papua New Guinea.

Buka Island solomys

Buka Island solomys

Solomys spriggsarum

The Buka Island solomys (*Solomys spriggsarum*), also known as the Buka naked-tailed rat or Buka Island naked-tailed rat, is an extinct species of rodent known only from subfossil remains. This species occurred on Buka Island, the second largest...

Bukharan markhor

Bukharan markhor

Capra falconeri heptneri

The Bukharan markhor, or Tadjik markhor (*Capra falconeri heptneri*) is an endangered goat-antelope, native to Tajikistan, Turkmenistan and Uzbekistan, possibly also Afghanistan. Today it is found in few scattered populations, for example in...

Buller's chipmunk

Buller's chipmunk

Neotamias bulleri

Buller's chipmunk (*Neotamias bulleri*) is a species of rodent in the family Sciuridae. It is endemic to Mexico.

Buller's pocket gopher

Buller's pocket gopher

Pappogeomys bulleri

Buller's pocket gopher (*Pappogeomys bulleri*) is a species of gopher that is endemic to Mexico. It is monotypic within the genus *Pappogeomys*. It includes the Alcorn's pocket gopher (*P. b. alcorni*) as a subspecies. Previously, this subspecies was...

Bulmer's fruit bat

Bulmer's fruit bat

Aproteles bulmerae

Bulmer's fruit bat (*Aproteles bulmerae*) is a megabat endemic to New Guinea. It is listed as a critically endangered species due to habitat loss and hunting. It is the only member of the genus *Aproteles*. Due to its imperiled status, it is...

Bunchgrass leaf-eared mouse

Bunchgrass leaf-eared mouse

Phyllotis osilae

The bunchgrass leaf-eared mouse (*Phyllotis osilae*) is a species of rodent in the family Cricetidae. It is found in Argentina, Bolivia, and Peru.

Bunn's short-tailed bandicoot rat

Bunn's short-tailed bandicoot rat

Nesokia bunnii

Bunn's short-tailed bandicoot rat (*Nesokia bunnii*) is a species of rodent in the family Muridae. It is found only in the marshes of southeastern Iraq and is named for the Iraqi zoologist Dr. Munir K. Bunni.

Bunny rat

Bunny rat

Reithrodon auritus

The bunny rat, or hairy-soled conyrat (*Reithrodon auritus*) is a species of rodent in the family Cricetidae, native to southern South America.

Bunting's thicket rat

Bunting's thicket rat

Grammomys buntingi

Bunting's thicket rat (*Grammomys buntingi*) is a species of rodent in the family Muridae. It is found in Ivory Coast, Guinea, Liberia, Senegal, and Sierra Leone. Its natural habitats are subtropical or tropical moist lowland forests and subtropical...

Bunyoro rabbit

Bunyoro rabbit

Poelagus marjorita

The Bunyoro rabbit or Central African rabbit (*Poelagus marjorita*) is a species of mammal in the family Leporidae. It is monotypic within the genus *Poelagus*. It is found in central Africa and its typical habitat is damp savannah, often with rocky...

Burchell's zebra

Burchell's zebra

Equus quagga burchellii

Burchell's zebra (*Equus quagga burchellii*) is a southern subspecies of the plains zebra. It is named after the British explorer and naturalist William John Burchell. Common names include the bontequagga, Damaraland zebra, and Zululand zebra...

Burma pipistrelle

Burma pipistrelle

Hypsugo lophurus

The Burma pipistrelle (*Hypsugo lophurus*) is a species of vesper bat in the family Vespertilionidae found in Myanmar and Thailand. It is known from Maliwun in Tanintharyi Region (Tenasserim).

Burmeister's porpoise

Burmeister's porpoise

Phocoena spinipinnis

Burmeister's porpoise (*Phocoena spinipinnis*) is a species of porpoise endemic to the coast of South America. It was first described by Hermann Burmeister, for whom the species is named, in 1865. Locally it is known as *marsopa espinosa* (thorny...

Burmese Ferret-Badger

Burmese Ferret-Badger

Melogale personata

Burmese ferret-badgers (*Melogale personata*) are carnivorous mammals that live in Southeast Asia. They are nocturnal animals, that live on the ground and sleep most of the day in their burrows.

Burmese hare

Burmese hare

Lepus peguensis

The Burmese hare (*Lepus peguensis*) is a species of mammal in the family Leporidae. It is found in Cambodia, Laos, Myanmar, Thailand, and Vietnam.

Burmese short-tailed shrew

Burmese short-tailed shrew

Blarinella wardi

The Burmese short-tailed shrew (*Blarinella wardi*) is one of three species of shrew in the genus *Blarinella*. It is in the family Soricidae and is found in China and Myanmar. Its natural habitat is subtropical or tropical dry forests.

Burmese whiskered bat

Burmese whiskered bat

Myotis montivagus

The Burmese whiskered myotis or Burmese whiskered bat (*Myotis montivagus*) is a species of vesper bat. It is found in China, India, Myanmar, Laos, and Vietnam.
Burrowing chinchilla mouse
Burrowing chinchilla mouse
Euneomys fessor

The burrowing chinchilla mouse (*Euneomys fessor*) is a species of rodent in the family Cricetidae. It is found in Salta Province, Argentina.
Burrunan Dolphin
Burrunan Dolphin
Tursiops australis

The Burrunan dolphin (*Tursiops australis*) is a proposed species of bottlenose dolphin found in parts of Victoria, Australia first described in 2011. Its exact taxonomy is debated: numerous studies support it as being a separate species within...
Burt's deer mouse
Burt's deer mouse
Peromyscus caniceps

Burt's deer mouse (*Peromyscus caniceps*) is a species of rodent in the family Cricetidae. It is endemic to Mexico, where it is found only on Montserrat Island off the east coast of Baja California Sur. The species is threatened by predation by...
Burton's gerbil
Burton's gerbil
Gerbillus burtoni

Burton's gerbil (*Gerbillus burtoni*) is distributed mainly in Darfur, Sudan. Less than 250 individuals of this species of rodent are thought to persist in the wild. It may have been named after Edward Burton, who had the gerbil in his menagerie,...
Burton's vlei rat
Burton's vlei rat
Otomys burtoni

Burton's vlei rat (*Otomys burtoni*) is a species of rodent in the family Muridae. It is found only in Cameroon. Its natural habitats are subtropical or tropical high-altitude shrubland, subtropical or tropical high-altitude grassland, and swamps. It...
Burton's yellow-shouldered bat
Burton's yellow-shouldered bat
Sturnira burtonlimi

Burton's yellow-shouldered bat (*Sturnira burtonlimi*) is a species of leaf-nosed bat found in Panama and Costa Rica.
Buru Babirusa
Buru Babirusa
Babirusa babyrussa

Buru babirusas (*Babirusa babyrussa*) are wild pig-like animals native to Southeast Asia. They have a remarkable "prehistoric" appearance due to the prominent upwards incurving canine tusks of the males, which actually pierce the flesh in the snout.
Bush Dog
Bush Dog
Speothos venaticus

The Bush dog (*Speothos venaticus*) is a canine found in Central and South America. It is very rare in most areas except in Suriname, Guyana, and Peru; it was first identified by Peter Wilhelm Lund from fossils in Brazilian caves and was

believed...

Bush Rat

Bush Rat

Rattus fuscipes

The bush rat (*Rattus fuscipes*) is a small Australian nocturnal animal. It is one of the most common indigenous species of rat on the continent, found in many heathland areas of Victoria and New South Wales.

Bush vlei rat

Bush vlei rat

Myotomys unisulcatus

The bush vlei rat or Karoo bush rat (*Myotomys unisulcatus*, formerly *Otomys unisulcatus*) is a species of rodent in the family Muridae. It is found in Namibia and South Africa. Its natural habitat is temperate shrubland. The Karoo rat uses...

Bushbuck

Bushbuck

Tragelaphus sylvaticus

The bushbuck (*Tragelaphus scriptus*) is a common and a widespread species of antelope in Sub-Saharan Africa. Bushbucks are found in a wide range of habitats, such as rain forests, montane forests, forest-savanna mosaic, savanna, bushveld and...

Bushpig

Bushpig

Potamochoerus larvatus

The bushpig (*Potamochoerus larvatus*) is a member of the pig family that lives in East and Southern Africa. Bushpigs are very aggressive and extremely powerful. They are also fast and can swim well.

Bushveld elephant shrew

Bushveld elephant shrew

Elephantulus intufi

The bushveld elephant shrew or bushveld sengi (*Elephantulus intufi*) is a species of elephant shrew in the family Macroscelididae. It is found in Angola, Botswana, Namibia, and South Africa. Its natural habitats are subtropical or tropical dry...

Bushveld gerbil

Bushveld gerbil

Gerbilliscus leucogaster

The bushveld gerbil (*Gerbilliscus leucogaster*) is a species of rodent found in Angola, Botswana, Democratic Republic of the Congo, Kenya, Malawi, Mozambique, Namibia, South Africa, Eswatini, Tanzania, Uganda, Zambia, and Zimbabwe. Its natural...

Bushveld horseshoe bat

Bushveld horseshoe bat

Rhinolophus simulator

The Bushveld horseshoe bat (*Rhinolophus simulator*) is a species of bat in the family Rhinolophidae. It is found in Botswana, Cameroon, Ivory Coast, Eswatini, Ethiopia, Guinea, Kenya, Liberia, Malawi, Mozambique, Nigeria, South Africa, South...

Bushy-tailed hairy-footed gerbil

Bushy-tailed hairy-footed gerbil

Gerbillurus vallinus

The bushy-tailed hairy-footed gerbil (*Gerbillurus vallinus*) is a species of rodent found in Namibia and South Africa. Its natural habitats are dry savanna, temperate

grassland, and hot deserts. It is threatened by habitat loss.

Bushy-tailed jird

Bushy-tailed jird

Sekeetamys calurus

The bushy-tailed jird or bushy-tailed dipodil (*Sekeetamys calurus*) is a species of rodent in the family Muridae. It is the only species in the genus *Sekeetamys*. It is found in Egypt, Israel, Jordan, Saudi Arabia, and Sudan. Its natural habitat...

Bushy-tailed mongoose

Bushy-tailed mongoose

Bdeogale crassicauda

The bushy-tailed mongoose (*Bdeogale crassicauda*) is a mammal in the family Herpestidae found in central Africa, from southern Kenya to central Mozambique.

Bushy-Tailed Opossum

Bushy-Tailed Opossum

Glironia venusta

The bushy-tailed opossum (*Glironia venusta*) is an opossum from South America. It was first described by English zoologist Oldfield Thomas in 1912. It is a medium-sized opossum characterized by a large, oval, dark ears, fawn to cinnamon coat with...

Bushy-Tailed Woodrat

Bushy-Tailed Woodrat

Neotoma cinerea

The bushy-tailed woodrat, or packrat (*Neotoma cinerea*) is a species of rodent in the family Cricetidae found in Canada and the United States. Its natural habitats are boreal forests, temperate forests, dry savanna, temperate shrubland, and...

Busuanga squirrel

Busuanga squirrel

Sundasciurus hoogstraali

The Busuanga squirrel (*Sundasciurus hoogstraali*) is a species of rodent in the family Sciuridae. It is endemic to the Philippines.

Butiaba naked-tailed shrew

Butiaba naked-tailed shrew

Crocidura littoralis

The Butiaba naked-tailed shrew (*Crocidura littoralis*) is a species of mammal in the family Soricidae. It is found in Cameroon, Central African Republic, Republic of the Congo, Democratic Republic of the Congo, Kenya, and Uganda. Its natural...

Buxton's jird

Buxton's jird

Meriones sacramenti

Buxton's jird (*Meriones sacramenti*) is a species of rodent in the family Muridae. It is found in the Negev desert and the northern coastal parts of the Sinai peninsula.

Caatinga vesper mouse

Caatinga vesper mouse

Calomys expulsus

The caatinga vesper mouse (*Calomys expulsus*) (also known as the caatinga laucha or rejected vesper mouse) is a rodent species in the family Cricetidae from South America. It is endemic to eastern Brazil, where it is found in open savanna...

Cabrera's hutia

Cabrera's hutia

Mesocapromys angelcabrerai

Cabrera's hutia (*Mesocapromys angelcabrerai*) is a species of rodent in the subfamily Capromyinae. It is endemic to Cuba. Its natural habitats are subtropical or tropical mangrove forests and swamps. It is threatened by habitat loss. Its species...

Cabrera's vole
Cabrera's vole
Microtus cabrerai

Cabrera's vole (*Microtus cabrerai*) is a species of vole native to Spain and Portugal. It is named for Ángel Cabrera, a mammalogist then working at the Museo Nacional de Ciencias Naturales in Madrid. It is the only living member of the subgenus...

Cacomistle
Cacomistle
Bassariscus sumichrasti

The cacomistle (*Bassariscus sumichrasti*) is an arboreal relative of the raccoon. It is often confused with its cousin the Ring-tailed cat because of the similarity of their appearance, but unlike the Ring-tail cat, the cacomistle does not have...

Cactus Mouse
Cactus Mouse
Peromyscus eremicus

The Cactus mouse (*Peromyscus eremicus*) is a species of rodent native to western North America. It is a closely related group of New World mice often called "deermice".

Cadena's tailless bat
Cadena's tailless bat
Anoura cadenai

Cadena's tailless bat (*Anoura cadenai*) is a species of bat native to Colombia. In 2006 it was described as a separate species from the tailed tailless bat species complex.

Cadorna's pipistrelle
Cadorna's pipistrelle
Hypsugo cadornae

Cadorna's pipistrelle (*Hypsugo cadornae*) is a species of vesper bat in the family Vespertilionidae. It is found in India, Laos, Myanmar, Thailand, and Vietnam.

Cairo Spiny Mouse
Cairo Spiny Mouse
Acomys cahirinus

The Cairo spiny mouse (*Acomys cahirinus*), also known as the common spiny mouse, Egyptian spiny mouse, or Arabian spiny mouse, is a nocturnal species of rodent in the family Muridae. It is found in Africa north of the Sahara, where its natural...

Cajamarca Oldfield mouse
Cajamarca Oldfield mouse
Thomasomys praetor

The Cajamarca Oldfield mouse (*Thomasomys praetor*) is a species of rodent in the family Cricetidae. It is present in the Andes of northwestern Peru, where its habitats include shrubby páramo, montane forest, and secondary forest. The rodent is...

Calabar Angwantibo
Calabar Angwantibo
Arctocebus calabarensis

The Calabar angwantibo (*Arctocebus calabarensis*) is a rare species of primate found only in Africa. It takes its name from the Nigerian city of Calabar. It is closely related to the potto (*Perodicticus potto*) and the various lorises.

Calabria pine vole

Calabria pine vole

Microtus brachycercus

The Calabria pine vole (*Microtus brachycercus*) is a vole found in southern and central Italy initially described by Lehmann as a subspecies of *M. savii*. Genetic tests in the Calabrian region found, although similar, the X chromosome is larger...

Calabrian black squirrel

Calabrian black squirrel

Sciurus meridionalis

The Calabrian black squirrel (*Sciurus meridionalis*) is a species of tree squirrel in the genus *Sciurus*, endemic to the forests of the regions of Calabria and Basilicata, in the south of the Italian Peninsula.

Calaby's pademelon

Calaby's pademelon

Thylogale calabyi

Calaby's pademelon (*Thylogale calabyi*), also known as the alpine wallaby, is a species of marsupial in the family Macropodidae. It is endemic to Papua New Guinea. It is threatened by habitat loss and hunting. Pademelons share many similarities in...

Calamian Deer

Calamian Deer

Hyelaphus calamianensis

The Calamian deer (*Axis calamianensis*) is an endangered species of deer found only in the Philippines. It is known as the "hog deer" due to its manner of fleeing from danger.

Calamian treeshrew

Calamian treeshrew

Tupaia moellendorffi

The Calamian treeshrew (*Tupaia palawanensis*) is a treeshrew species found in the islands of Busuanga and Culion, which are part of the Calamian Islands group in the Philippines. The German zoologist Paul Matschie first described a Calamian...

California chipmunk

California chipmunk

Neotamias obscurus

The California chipmunk or chaparral chipmunk (*Neotamias obscurus*) is a species of rodent in the squirrel family Sciuridae. It is found in Baja California, Mexico and in southern California in the United States.

California Deermouse

California Deermouse

Peromyscus californicus

The California deermouse (*Peromyscus californicus*) is the largest species of deermouse in the United States. They are New World mice only distantly related to the Common house and laboratory mouse (*Mus musculus*). They are accomplished jumpers and...

California grizzly bear

California grizzly bear

Ursus arctos californicus

The California grizzly bear (*Ursus arctos californicus*) is an extinct population or subspecies of the brown bear, generally known (together with other North American brown bear populations) as the grizzly bear. "Grizzly" could have meant...
California Ground Squirrel
California Ground Squirrel
Otospermophilus beecheyi

The California ground squirrel (*Otospermophilus beecheyi*) is a common ground squirrel of the western United States and the Baja California Peninsula. It was named after Frederick William Beechey, an early 19th-century British explorer and naval...
California kangaroo rat
California kangaroo rat
Dipodomys californicus

The California kangaroo rat (*Dipodomys californicus*) is a species of rodent in the family Heteromyidae. However, populations are declining, having not fully recovered after the drought in California from 2013 to 2015 destroyed their habitat (the...
California leaf-nosed bat
California leaf-nosed bat
Macrotus californicus

The California leaf-nosed bat (*Macrotus californicus*) is a species of bat in the family Phyllostomidae. It is found in Mexico and the United States. Its natural habitat is hot deserts. It is threatened by habitat loss.
California Mule Deer
California Mule Deer
Odocoileus hemionus californicus

The California mule deer (*Odocoileus hemionus californicus*) is a subspecies of Mule deer whose range covers much of the state of California.
California myotis
California myotis
Myotis californicus

The California myotis (*Myotis californicus*) is a species of vesper bat. It is found in British Columbia in Canada, Guatemala, Mexico, and in the western United States, including California.
California pocket mouse
California pocket mouse
Chaetodipus californicus

The California pocket mouse (*Chaetodipus californicus*) is a species of rodent in the family Heteromyidae.
California red tree mouse
California red tree mouse
Arborimus pomo

The Sonoma tree vole or California red tree mouse (*Arborimus pomo*) is a species of rodent in the family Cricetidae. The species is found in northwest California. The preferred habitat for this primarily arboreal vole is old-growth Douglas-fir...
California Sea Lion
California Sea Lion
Zalophus californianus

The California sea lion (*Zalophus californianus*) is one of six species of sea lions. They mainly haul-out on sandy or rocky beaches, but they also frequent manmade environments such as marinas and wharves. California sea lions are particularly...

California Vole
California Vole
Microtus californicus

The California vole (*Microtus californicus*) is a small rodent that has numerous natural predators. It is also known as the "California meadow mouse", a misnomer as this species is a vole, not a mouse.

Callewaert's mouse
Callewaert's mouse
Mus callewaerti

Callewaert's mouse (*Mus callewaerti*) is a species of rodent in the family Muridae. It is found in Angola and Democratic Republic of the Congo. Its natural habitat is dry savanna.

Calomys cerqueirai
Calomys cerqueirai
Calomys cerqueirai

Calomys cerqueirai is a species of rodent in the genus *Calomys* from southeastern Brazil. Distinct from other *Calomys* in its karyotype and characters of the fur, it is known only from two places in Minas Gerais. The karyotype was first described...

Calomyscus elburzensis
Calomyscus elburzensis
Calomyscus elburzensis

Calomyscus elburzensis, also known as Goodwin's *calomyscus* or Goodwin's brush-tailed mouse is a species of mouse-like hamster. It is found in northeastern and northern Iran, western Afghanistan, and southern Turkmenistan, where it is found in...

Camas Pocket Gopher
Camas Pocket Gopher
Thomomys bulbivorus

The Camas pocket gopher (*Thomomys bulbivorus*) is a large burrowing rodent, first described in 1829. This is a herbivorous animal that collects plant matter in large cheek pouches and then hoards it in an extensive system of tunnels.

Cambodian striped squirrel
Cambodian striped squirrel
Tamiops rodolphii

The Cambodian striped squirrel (*Tamiops rodolphii*) is a species of rodent in the family Sciuridae. It is found in eastern Thailand, Cambodia, southern Laos, and southern Vietnam.

Cameron Highlands white-bellied rat
Cameron Highlands white-bellied rat
Niviventer cameroni

The Cameron Highlands white-bellied rat (*Niviventer cameroni*), also known as the Cameron Highlands *niviventer*, is a species of rodent in the family Muridae. It has only been found in the mountain forests of the Cameron Highlands on the Malay...

Cameroon clawless otter
Cameroon clawless otter
Aonyx congicus

The Congo clawless otter (*Aonyx congicus*), also known as the Cameroon clawless otter, is a species in the family Mustelidae. It was formerly recognised as a subspecies (*Aonyx capensis congicus*) of the African clawless otter. This clawless otter...

Cameroon climbing mouse

Cameroon climbing mouse
Dendromus oreas

The Cameroon climbing mouse (*Dendromus oreas*) is a species of rodent in the family Nesomyidae which is endemic to the montane grasslands on three mountains in Cameroon.

Cameroon scaly-tail
Cameroon scaly-tail
Zenkerella insignis

The Cameroon scaly-tail (*Zenkerella insignis*), also referred to as the Cameroon anomalure, flightless anomalure or flightless scaly-tail, is a rodent species endemic to West Central Africa. The scientific literature has never (or possibly only...

Cameroon soft-furred mouse
Cameroon soft-furred mouse
Praomys morio

The Cameroon soft-furred mouse or Cameroon praomys (*Praomys morio*) is a species of rodent in the family Muridae. It is found in Cameroon and Equatorial Guinea. Its natural habitat is subtropical or tropical moist montane forests. It is threatened...

Cameroonian forest shrew
Cameroonian forest shrew
Sylvisorex camerunensis

The Cameroonian forest shrew (*Sylvisorex camerunensis*) is a species of mammal in the family Soricidae. It is found in the Montane forests of Western Cameroon around Mount Oku and Lake Manengouba, and in Southeastern Nigeria in the Gote...

Cameroonian shrew
Cameroonian shrew
Crocidura picea

The Cameroonian shrew (*Crocidura picea*) is a species of mammal in the family Soricidae. It is endemic to Cameroon. Its natural habitat is subtropical or tropical moist montane forests. It is threatened by habitat loss.

Camiguin forest mouse
Camiguin forest mouse
Apomys camiguinensis

The Camiguin forest mouse (*Apomys camiguinensis*) is a forest mouse endemic to the island of Camiguin in the southern Philippines. It has large ears and eyes, a long tail and rusty-brown fur, and it feeds mostly on insects and seeds. This...

Camiguin forest rat
Camiguin forest rat
Bullimus gamay

Camiguin forest rat, or Camiguin bullimus (*Bullimus gamay*) is one of three species of rodents in the genus *Bullimus*. It is endemic to the island of Camiguin, the Philippines.

Campbell's Dwarf Hamster
Campbell's Dwarf Hamster
Phodopus campbelli

The Campbell's dwarf hamster (*Phodopus campbelli*) is native to Eastern and Central Asia. The species is named after W.C. Campbell, who discovered this animal in Mongolia in 1902. Endemic to the Djungaria region of Mongolia, these animals are...

Campbell's mona monkey
Campbell's mona monkey

Cercopithecus campbelli

Campbell's mona monkey, also known as Campbell's guenon and Campbell's monkey (*Cercopithecus campbelli*) is a species of primate in the family Cercopithecidae found in the Ivory Coast, Gambia, Ghana, Guinea, Guinea-Bissau, Liberia, Senegal, and...

Campo-Ma'an fruit bat

Campo-Ma'an fruit bat

Casinonycteris campomaanensis

The Campo-Ma'an fruit bat is a species of bat found in Cameroon. The first individual encountered was in 2007. It was described as a new species in 2014. Only three individuals have ever been encountered.

Canadian Lynx

Canadian Lynx

Lynx canadensis

The Canadian lynx (*Lynx canadensis*) is a medium-sized North American lynx. It was first described by Robert Kerr in 1792. The Canadian lynx occurs predominantly in dense boreal forests, and its range strongly coincides with that of the Snowshoe...

Canarian shrew

Canarian shrew

Crocidura canariensis

The Canarian shrew (*Crocidura canariensis*) is a species of mammal in the family Soricidae. It is endemic to the Canary Islands, specifically the eastern islands of Lanzarote, Fuerteventura, Lobos, and Mount Clara. It used to be found on...

Canary big-eared bat

Canary big-eared bat

Plecotus teneriffae

The Canary long-eared bat (*Plecotus teneriffae*), also known as the Canary big-eared bat or Tenerife long-eared bat, is a species of vesper bat endemic to the Canary Islands. According to the IUCN, it is vulnerable to the risk of extinction.The...

Candango mouse

Candango mouse

Juscelinomys candango

The candango mouse or candango akodont (*Juscelinomys candango*) is an extinct rodent species from South America. It was found around Brasilia in 1960, but its habitat has been overtaken by urban sprawl, and is now presumed extinct.

Canis arnensis

Canis arnensis

Canis arnensis

Canis arnensis (or Arno River dog) is an extinct species of canine that was endemic to Mediterranean Europe during the Early Pleistocene. The Arno River dog has been described as a small jackal-like dog. Its anatomy and morphology relate it more...

Canis cedazoensis

Canis cedazoensis

Canis cedazoensis

Canis cedazoensis is an extinct species of smaller canid which was endemic to North America during the Pleistocene epoch, 1.8 Ma–300,000 years ago.The morphology and dentition of *C. cedazoensis* suggests a jackal-like animal that was more...

Canis ferax

Canis ferax

Canis ferax

Canis ferox (Latin: *canis* : dog, *ferox* : fierce; hence fierce dog) is a species of canid which was endemic to North America and lived during the Late Miocene and Early Pliocene. Existing nearly 6 million years ago, this animal is thought to be an...

Canis lupus dingo
Canis lupus dingo
Canis lupus dingo

In the taxonomic treatment presented in the third (2005) edition of *Mammal Species of the World*, *Canis lupus dingo* is a taxonomic rank that includes both the dingo that is native to Australia and the New Guinea singing dog that is native to the...

Cansdale's swamp rat
Cansdale's swamp rat
Malacomys cansdalei

Cansdale's swamp rat (*Malacomys cansdalei*) is a species of rodent in the family Muridae. It is found in Ivory Coast, Ghana, and possibly Liberia. Its natural habitat is subtropical or tropical moist lowland forests.

Cantabrian chamois
Cantabrian chamois
Rupicapra pyrenaica parva

The Cantabrian chamois (*Rupicapra pyrenaica parva*) is a slim mountain goat antelope, and is one of the 10 subspecies of the genus *Rupicapra*. It ranges the Cantabrian Mountains in northern Spain, with a population of 17,000 animals in 2007-2008.

Cantor's roundleaf bat
Cantor's roundleaf bat
Hipposideros galeritus

Cantor's roundleaf bat (*Hipposideros galeritus*) is a species of bat in the family Hipposideridae. It is found in Bangladesh, Brunei, Cambodia, India, Indonesia, Laos, Malaysia, Sri Lanka, Thailand and Vietnam. It is named after Theodore Edward...

Canut's horseshoe bat
Canut's horseshoe bat
Rhinolophus canuti

Canut's horseshoe bat (*Rhinolophus canuti*) is a species of bat in the family Rhinolophidae. It is endemic to Indonesia.

Canyon bat
Canyon bat
Parastrellus hesperus

The canyon bat (*Parastrellus hesperus*), also known as the western pipistrelle, is a species of vesper bat. It is found in Mexico and in the western United States. The species has historically been placed in the genus *Pipistrellus*, but molecular...

Canyon mouse
Canyon mouse
Peromyscus crinitus

The canyon mouse (*Peromyscus crinitus*) is a gray-brown mouse found in many states of the western United States and northern Mexico. Its preferred habitat is arid, rocky desert. It is the only species in the *Peromyscus crinitus* species...

Caparaó grass mouse
Caparaó grass mouse
Akodon mystax

The Caparaó grass mouse (*Akodon mystax*) is a rodent species from South America. It is found in Brazil.

Caparaó hocicudo

Caparaó hocicudo

Oxymycterus caparaoe

The Caparaó hocicudo (*Oxymycterus caparaoe*) is a rodent species from South America. It is endemic to Brazil where it is found in the Pico da Bandeira mountain range.

Cape dune mole-rat

Cape dune mole-rat

Bathyergus suillus

The Cape dune mole-rat (*Bathyergus suillus*) is a species of solitary burrowing rodent in the family Bathyergidae. It is endemic to South Africa and named for the Cape of Good Hope.

Cape elephant shrew

Cape elephant shrew

Elephantulus edwardii

The Cape elephant shrew (*Elephantulus edwardii*), also known as the Cape rock elephant-shrew, or Cape rock sengi, is a species of elephant shrew in the family Macroscelididae. It is endemic to South Africa, although it is a relatively common...

Cape Fox

Cape Fox

Vulpes chama

The Cape fox (*Vulpes chama*), also called the asse, cama fox or the silver-backed fox, is a small species of fox, native to southern Africa. It is also called a South African version of a fennec fox due to its similarly big ears. It is the only...

Cape Genet

Cape Genet

Genetta tigrina

The Cape genet (*Genetta tigrina*), also known as the South African large-spotted genet, is a genet species endemic to South Africa. As it is common and not threatened, it is listed as Least Concern on the IUCN Red List. Like other genets, it is...

Cape gerbil

Cape gerbil

Gerbilliscus afra

The Cape gerbil, *Gerbilliscus afra*, is a species of rodent found only in South Africa. Its natural habitats are subtropical or tropical dry shrubland and temperate desert.

Cape golden mole

Cape golden mole

Chrysochloris asiatica

The Cape golden mole (*Chrysochloris asiatica*) is a small, insectivorous mammal of the family Chrysochloridae, the golden moles. It is native to certain south western regions of South Africa. The species is a solitary subterranean insectivore,...

Cape Gray Mongoose

Cape Gray Mongoose

Galerella pulverulenta

The Cape grey mongoose (*Herpestes pulverulentus*) is a small mammal found only in South Africa. These energetic terrestrial mammals have a varied diet and are often spotted by humans when they cross roads.

Cape Ground Squirrel
Cape Ground Squirrel
Xerus inauris

The Cape ground squirrel or South African ground squirrel (*Geosciurus inauris*) is found in most of the drier parts of southern Africa from South Africa, through to Botswana, and into Namibia, including Etosha National Park. The name Cape ground...

Cape Grysbok
Cape Grysbok
Raphicerus melanotis

The Cape grysbok (*Raphicerus melanotis*) is a small antelope that is native to South Africa. It has an interesting habit of fluffing out the fur at its rear end to make itself look bigger.

Cape hairy bat
Cape hairy bat
Myotis tricolor

The Cape hairy bat, also known as little brown bat, Temminck's mouse-eared bat, Cape myotis, tricoloured mouse-eared bat, Cape hairy myotis, Temminck's hairy bat and three-coloured bat (*Myotis tricolor*) is a species of vesper bat that is found...

Cape Hare
Cape Hare
Lepus capensis

The Cape hare (*Lepus capensis*), also called the desert hare, is a hare native to Africa and Arabia extending into India.

Cape horseshoe bat
Cape horseshoe bat
Rhinolophus capensis

The Cape horseshoe bat (*Rhinolophus capensis*) is a species of bat in the family Rhinolophidae. It is endemic to South Africa, and is potentially threatened by habitat loss and disturbance of its roosting sites, although it is present in large...

Cape mole-rat
Cape mole-rat
Georychus capensis

The Cape mole-rat (*Georychus capensis*) is a species of mole-rat endemic to South Africa. It is the only extant species currently described as belonging in the genus *Georychus*.

Cape Mountain Zebra
Cape Mountain Zebra
Equus zebra zebra

The Cape mountain zebra (*Equus zebra zebra*) is a subspecies of Mountain zebra that is found in certain mountainous regions of South Africa. It is the smallest of all existing zebra species and also the most geographically restricted.

Cape Porcupine
Cape Porcupine
Hystrix africaeaustralis

The Cape porcupine (*Hystrix africaeaustralis*) or South African porcupine, is a species of Old World porcupine native to central and southern Africa.

Cape serotine
Cape serotine
Neoromicia capensis

The Cape serotine (*Laephotis capensis*) is a species of vesper bat occurring in Sub-Saharan Africa. 'Serotine' is from Latin 'serotinus' meaning 'of the evening'. It is found in Angola, Benin, Botswana, Burundi, Cameroon, Central African Republic,...

Cape short-eared gerbil
Cape short-eared gerbil
Desmodillus auricularis

The Cape short-eared gerbil (*Desmodillus auricularis*) is a species of rodent in the family Muridae. It is the only species in the genus *Desmodillus*. It is found in Angola, Botswana, Namibia, and South Africa. Its natural habitats are hot deserts...

Cape spiny mouse
Cape spiny mouse
Acomys subspinosus

The Cape spiny mouse (*Acomys subspinosus*) is a murid rodent found in the Western Cape Province of South Africa. They have a dorsal covering of spiny hairs with dark grey-brown coloration, and a white underbelly. The Cape Spiny Mouse has large...

Cape wild dog
Cape wild dog
Lycaon pictus pictus

The African wild dog (*Lycaon pictus*), also called the African painted dog and the African hunting dog, is a wild canine which is a native species to sub-Saharan Africa. It is the largest wild canine in Africa, and the only extant member of the...

Cape York melomys
Cape York melomys
Melomys capensis

The Cape York mosaic-tailed rat, or Cape York melomys (*Melomys capensis*) is a species of rodent in the family Muridae. It is found only in Australia, on the Cape York Peninsula.

Cape York rat
Cape York rat
Rattus leucopus

The Cape York rat (*Rattus leucopus*) is a species of rodent in the family Muridae. It is found in southern New Guinea, in both Indonesia and Papua New Guinea, and in Cape York Peninsula in Australia.

Cape York rock-wallaby
Cape York rock-wallaby
Petrogale coenensis

The Cape York rock-wallaby (*Petrogale coenensis*) is a species of rock-wallaby restricted to Cape York Peninsula in northeastern Queensland, Australia. It is a member of a group of seven very closely related rock-wallabies, all found in...

Capped Langur
Capped Langur
Trachypithecus pileatus

The capped langur (*Trachypithecus pileatus*) is a species of primate in the family Cercopithecidae. It is native to Bangladesh, Bhutan, India, and Myanmar. Its natural habitat is subtropical or tropical dry forests. It is threatened by habitat...

Capricorn leaf-eared mouse
Capricorn leaf-eared mouse
Phyllotis caprinus

The capricorn leaf-eared mouse (*Phyllotis caprinus*) is a species of rodent in the family Cricetidae. It is found in Argentina and Bolivia.

Capromys gundlachianus
Capromys gundlachianus
Capromys gundlachianus

The Archipiélago de Sabana hutia (*Capromys pilorides gundlachianus*) is a subspecies of the Desmarest's hutia endemic to the Sabana-Camagüey Archipelago of Cuba.

Capybara
Capybara
Hydrochoerus hydrochaeris

The capybara (*Hydrochoerus hydrochaeris*) is a giant cavy rodent native to South America. It is the largest living rodent and a member of the genus *Hydrochoerus*. Its close relatives include guinea pigs and Rock cavy, and it is more distantly...

Caquetá titi
Caquetá titi
Plecturocebus caquetensis

The Caquetá titi, *Plecturocebus caquetensis*, also known as the red-bearded titi or the bushy-bearded titi, is a species of titi monkey endemic to Colombia found in the Department of Caquetá region. Taxonomically, it is a member of the "*Callicebus*...

Caracal
Caracal
Caracal caracal

The caracal (*Caracal caracal*) is a medium-sized wild cat. It was first scientifically described by German naturalist Johann Christian Daniel von Schreber in 1776. Typically nocturnal, the caracal is highly secretive and difficult to observe. It...

Cariri climbing mouse
Cariri climbing mouse
Rhipidomys cariri

The Cariri climbing mouse (*Rhipidomys cariri*) is a partly arboreal rodent species from South America. It is known from two mesic localities in Ceara, northeastern Brazil, within the semiarid caatinga ecoregion. It has been found in areas under...

Carmen Mountain shrew
Carmen Mountain shrew
Sorex milleri

The Carmen Mountain shrew (*Sorex milleri*) is a species of mammal in the family Soricidae. It is endemic to Mexico.

Caroline flying fox
Caroline flying fox
Pteropus molossinus

The Caroline flying fox or Pohnpei flying fox (*Pteropus molossinus*) is a species of megabat in the genus *Pteropus*, endemic to Micronesia. Its natural habitat is subtropical or tropical dry forests. Prior to a ban on commercial exploitation, it...

Carpathian Lynx

Carpathian Lynx
Lynx lynx carpathicus

The Carpathian lynx (*Lynx lynx carpathicus*) is a subspecies of the Eurasian lynx found in the Carpathian Basin. It was once common throughout Europe but is now extinct in some areas.

Carpathian wisent
Carpathian wisent
Bison bonasus hungarorum

The Carpathian wisent (*Bison bonasus hungarorum*) was a subspecies of the European bison that inhabited the Carpathian Mountains, Moldavia and Transylvania. It may also have lived in what are today Ukraine and Hungary. It began to die out about a...

Carpentarian dunnart
Carpentarian dunnart
Sminthopsis butleri

The Carpentarian dunnart (*Sminthopsis butleri*) also known as the Butler's dunnart with a puffy brown or mouse grey colour above and the underside of white, similar to its close relative the Kakadu dunnart. Head to anus length is 75-88mm with a...

Carpentarian rock rat
Carpentarian rock rat
Zyzomys palatalis

The Carpentarian rock rat (*Zyzomys palatalis*) is a species of rodent in the family Muridae. It is found only in Australia.

Carpenter's lar gibbon
Carpenter's lar gibbon
Hylobates lar carpenteri

Carpenter's lar gibbon, *Hylobates lar carpenteri*, is an endangered subspecies of white-handed gibbon, also known as the lar gibbon. It is listed as an endangered species because it is believed to have undergone a decline of more than 50% in the...

Carriker's round-eared bat
Carriker's round-eared bat
Lophostoma carrikeri

Carriker's round-eared bat (*Lophostoma carrikeri*) is a bat species from South America. It is found in Bolivia, Brazil, Colombia, Guyana, Peru, Suriname and Venezuela. It was discovered by and named for Melbourne A. Carriker, Jr.

Carruther's mountain squirrel
Carruther's mountain squirrel
Funisciurus carruthersi

The Carruther's mountain squirrel (*Funisciurus carruthersi*) is a species of rodent in the family Sciuridae. It is found in Burundi, Democratic Republic of the Congo, Rwanda, and Uganda. Its natural habitat is subtropical or tropical moist...

Cascade golden-mantled ground squirrel
Cascade golden-mantled ground squirrel
Callospermophilus saturatus

The Cascade golden-mantled ground squirrel (*Callospermophilus saturatus*) is a species of rodent in the family Sciuridae, in the order Rodentia. It is the largest species of the three within the genus *Callospermophilus*. It is found in the Cascade...

Cascade mountain wolf
Cascade mountain wolf

Canis lupus fuscus

The Cascade mountain wolf (*Canis lupus fuscus*) is an extinct subspecies of the gray wolf that was once found in the Pacific Northwest (British Columbia, Oregon, and Washington), but became extinct in 1940.

Cascade red fox

Cascade red fox

Vulpes vulpes cascadiensis

The Cascade red fox (*Vulpes vulpes cascadiensis*) is a subspecies of red fox native to Washington in the United States.

Caspian Red Deer

Caspian Red Deer

Cervus elaphus maral

The Caspian red deer (*Cervus elaphus maral*) is one of the easternmost subspecies of Red deer. It is sometimes called maral, Noble deer, or Eastern red deer.

Caspian Seal

Caspian Seal

Pusa caspica

The Caspian seal (*Pusa caspica* or *Phoca caspica*) is one of the smallest members of the earless seal family and unique in that it is found exclusively in the brackish Caspian Sea. They are found not only along the shorelines, but also on the many...

Caspian shrew

Caspian shrew

Crocidura caspica

The Caspian shrew (*Crocidura caspica*) is a species of mammal in the family Soricidae. It is found along the southern coast of the Caspian Sea in Iran and Azerbaijan.

Catamarca tuco-tuco

Catamarca tuco-tuco

Ctenomys knighti

The Catamarca tuco-tuco (*Ctenomys knighti*) is a species of rodent in the family Ctenomyidae. It is endemic to Argentina.

Cauca climbing mouse

Cauca climbing mouse

Rhipidomys caucensis

The Cauca climbing mouse (*Rhipidomys caucensis*) is a species of arboreal rodent in the family Cricetidae. It is endemic to Colombia, where it is found in montane forest at elevations from 2200 to 3500 m.

Caucasian birch mouse

Caucasian birch mouse

Sicista caucasica

The Caucasian birch mouse (*Sicista caucasica*) is a species of rodent in the family Sminthidae, that is endemic to Russia. Reports that it occurs in Turkey probably refer to Armenian birch mouse (*Sicista armenica*), from which it can only be...

Caucasian lynx

Caucasian lynx

Lynx lynx dinniki

The Caucasian lynx (*Lynx lynx dinniki*), also known as the Caucasus lynx or the eastern lynx, is a subspecies of Eurasian lynx native in the Caucasus, Iran, Turkey, and European Russia. It is proposed for the Caucasian lynx to be listed as...

Caucasian mole
Caucasian mole
Talpa caucasica

The Caucasian mole (*Talpa caucasica*) is a mammal in the family Talpidae that is endemic to the Caucasus Mountains of Russia and Georgia. Ognev's mole (*T. ognevi*) was formerly classified as a subspecies, but is now thought to be a distinct species.

Caucasian moose
Caucasian moose
Alces alces caucasicus

The Caucasian moose, also known as the Caucasian elk (*Alces alces caucasicus*) is an extinct subspecies of moose found in the Caucasus Mountains of Eastern Europe and Asia Minor, in modern-day European Russia, Armenia, Azerbaijan, Georgia, and...

Caucasian Mountain ground squirrel
Caucasian Mountain ground squirrel
Spermophilus musicus

The Caucasian Mountain ground squirrel (*Spermophilus musicus*) is a species of rodent in the family Sciuridae. It is endemic to the North Caucasus.

Caucasian pygmy shrew
Caucasian pygmy shrew
Sorex volnuchini

The Caucasian pygmy shrew or Ukrainian shrew (*Sorex volnuchini*) is a species of mammal in the family Soricidae. It is found in Armenia, Azerbaijan, Georgia, Russia, Ukraine, East Azerbaijan Province in Iran, and Turkey.

Caucasian shrew
Caucasian shrew
Sorex satunini

The Caucasian shrew (*Sorex satunini*) is a species of mammal in the family Soricidae. It is found in Armenia, Russia and Turkey.

Caucasian snow vole
Caucasian snow vole
Chionomys gud

The Caucasian snow vole (*Chionomys gud*) is a species of rodent in the family Cricetidae. It is found in Georgia, Russian Federation, and Turkey. Its natural habitats are temperate forests and temperate grassland.

Caucasian Squirrel
Caucasian Squirrel
Sciurus anomalus

The Caucasian squirrel (*Sciurus anomalus*) is a tree squirrel found in temperate forests in south-western Asia. The species is traditionally said to have first been described in 1778 by Johann Friedrich Gmelin, however, the true first description...

Caucasian Wildcat
Caucasian Wildcat
Felis silvestris caucasica

The Caucasian wildcat (*Felis silvestris caucasica*) is a European wildcat subspecies. It was described by Konstantin Satunin in 1905 on the basis of a skin of a female cat collected near Borjomi in Georgia.

Caucasian wisent
Caucasian wisent
Bison bonasus caucasicus

The Caucasian wisent (*Bison bonasus caucasicus*) or dombay (домбай) was a subspecies of European bison that inhabited the Caucasus Mountains of Eastern Europe.

Caucasus field mouse

Caucasus field mouse

Apodemus hyrcanicus

The Caucasus field mouse (*Apodemus hyrcanicus*) is a species of rodent in the family Muridae. It is found in Talysh region in SE Azerbaijan, northern Iran and possibly SW Turkmenistan.

Cave hyena

Cave hyena

Crocota crocuta spelaea

The cave hyena (*Crocota crocuta spelaea*), also known as the Ice Age spotted hyena, was a paleosubspecies of spotted hyena which ranged from the Iberian Peninsula to eastern Siberia. It is one of the best known mammals of the Ice Age and is well...

Cave lynx

Cave lynx

Lynx pardinus spelaeus

The cave lynx or Mediterranean cave lynx (*Lynx spelaeus* or *Lynx pardinus spelaeus*) is an extinct felid species that lived during the Pleistocene. It is controversially discussed to be a subspecies of the modern Iberian lynx (*Lynx pardinus*) and...

Cave myotis

Cave myotis

Myotis velifer

The cave myotis (*Myotis velifer*) is a species of vesper bat (*Vespertilionidae*) in the genus *Myotis*.

Cave nectar bat

Cave nectar bat

Eonycteris spelaea

The cave nectar bat, dawn bat, common dawn bat, common nectar bat or lesser dawn bat (*Eonycteris spelaea*) is a species of megabat within the genus *Eonycteris*. The scientific name of the species was first published by Dobson in 1871.

Cave wolf

Cave wolf

Canis lupus spelaeus

The cave wolf (*Canis lupus spelaeus*) is an extinct subspecies of wolf that lived during the Late Pleistocene Ice Age. It inhabited what is now modern-day western Europe. The Don wolf (*C. l. brevis*) from eastern Europe is regarded as a taxonomic...

Cavia anolaimae

Cavia anolaimae

Cavia anolaimae

Cavia anolaimae is a guinea pig species from South America. It is found in Colombia near Bogotá. It is believed to be a feral offshoot of the domestic guinea pig, *Cavia porcellus*, and is often treated as a synonym of *C. porcellus*, but Zúñiga et...

Cavia guianae

Cavia guianae

Cavia guianae

Cavia guianae is a guinea pig species from South America. It is found in southern Venezuela, Guyana, and portions of northern Brazil. Some biologists believe it to be a feral offshoot of the domestic guinea pig, *Cavia porcellus* ; others subsume...

Cavia patzelti

Cavia patzelti

Cavia patzelti

Cavia patzelti, commonly known as the Sacha guinea pig, is a species of *Cavia* native to Ecuador. Little is known about its population.

Cebu warty pig

Cebu warty pig

Sus cebifrons cebifrons

The Cebu warty pig (*Sus cebifrons cebifrons*) previously lived in Cebu, Philippines before becoming extinct in modern times, primarily due to habitat destruction. This pig was assessed to be extinct in 2000. The main species, *Sus cebifrons*, the...

Cedros Island mule deer

Cedros Island mule deer

Odocoileus hemionus cerrosensis

The Cedros Island mule deer (*Odocoileus hemionus cerrosensis*) is a subspecies of mule deer found only on Cedros Island off the coast of Baja California. Only about 50 individuals remain, with no captive population. Its behavior is similar to...

Celebes Crested Macaque

Celebes Crested Macaque

Macaca nigra

The Celebes crested macaque (*Macaca nigra*), also known as the crested black macaque, Sulawesi crested macaque, or the black ape, is an Old World monkey that lives in the Tangkoko reserve in the northeastern tip of the Indonesian island of...

Celebes dwarf squirrel

Celebes dwarf squirrel

Prosciurillus murinus

The Celebes dwarf squirrel (*Prosciurillus murinus*) is a species of rodent in the family Sciuridae. It is endemic to northeast and central Sulawesi, Indonesia, and is also found on nearby islands including Sangir Island.

Celebes rat

Celebes rat

Taeromys celebensis

The Celebes rat (*Taeromys celebensis*) is a species of rodent in the family Muridae. It is found only in Sulawesi, Indonesia.

Celebes shrew-rat

Celebes shrew-rat

Crunomys celebensis

The Celebes shrew-rat (*Crunomys celebensis*) is a species of rodent in the family Muridae. It is found only in Sulawesi, Indonesia. Its natural habitat is subtropical or tropical dry forests. It is threatened by habitat loss.

Celebes Warty Pig

Celebes Warty Pig

Sus celebensis

The Celebes warty pig (*Sus celebensis*) is a species of wild pig that lives on Sulawesi in Indonesia. It is the only pig species that has been domesticated apart from the Wild boar.

Central African oyan

Central African oyan
Poiana richardsonii

The Central African oyan (*Poiana richardsonii*), also called Central African linsang, is a linsang species native to Central Africa.

Central African Potto
Central African Potto
Perodicticus edwardsi

The Central African potto (*Perodicticus edwardsi*) is a nocturnal primate found in Central Africa. It is also known as Milne-Edwards's potto, after Alphonse Milne-Edwards.

Central American Agouti
Central American Agouti
Dasyprocta punctata

The Central American agouti (*Dasyprocta punctata*) is a species of agouti from the family Dasyproctidae. The main portion of its range is from Chiapas and the Yucatan Peninsula (southern Mexico), through Central America, to northwestern Ecuador,...

Central American dwarf squirrel
Central American dwarf squirrel
Microsciurus alfari

The Central American dwarf squirrel, also known as Alfaro's pygmy squirrel, (*Microsciurus alfari*) is a small tree squirrel in the genus *Microsciurus* and tribe Sciurini found in Colombia, Costa Rica, Nicaragua, and Panama. No species of squirrel...

Central American least shrew
Central American least shrew
Cryptotis orophila

The Central American least shrew (*Cryptotis orophila*) is a species of mammal in the family Soricidae. It is found in the highlands and mid-elevations from the western part of the northern coast of Honduras to central Costa Rica. It is found in...

Central American red brocket
Central American red brocket
Mazama temama

The Central American red brocket (*Mazama temama*) is a species of brocket deer ranging from southern Mexico, through Central America, to northwestern Colombia. In 1792 Robert Kerr originally described it as a unique separate species as opposed to...

Central American Squirrel Monkey
Central American Squirrel Monkey
Saimiri oerstedii

The Central American squirrel monkey (*Saimiri oerstedii*), also known as the red-backed squirrel monkey, is a squirrel monkey species from the Pacific coast of Costa Rica and Panama. It is restricted to the northwestern tip of Panama near the...

Central Asian boar
Central Asian boar
Sus scrofa davidi

The Central Asian boar (*Sus scrofa davidi*) is a small long maned subspecies of wild boar indigenous to Southeastern Iran, Pakistan and Northwest India.

Central Chimpanzee
Central Chimpanzee

Pan troglodytes troglodytes

The Central chimpanzee (*Pan troglodytes troglodytes*) is a subspecies of the chimpanzee closely related to the other great apes such as gorillas, orangutans, and humans. Central chimpanzees are considered highly intelligent apes. They are also...

Central European Boar

Central European Boar

Sus scrofa scrofa

The Central European boar (*Sus scrofa scrofa*) is a subspecies of Wild boar.

Central European red deer

Central European red deer

Cervus elaphus hippelaphus

The Central European red deer or common red deer (*Cervus elaphus hippelaphus*) is a subspecies of red deer native to central Europe. The deer's habitat ranges from France, Switzerland, Austria, Germany, and Denmark to the western Carpathians. It...

Central Kashmir vole

Central Kashmir vole

Alticola montosa

The central Kashmir vole (*Alticola montosa*) is a species of rodent in the family Cricetidae. It is found only in India and Pakistan.

Central leaf-eared mouse

Central leaf-eared mouse

Graomys chacoensis

The central leaf-eared mouse (*Graomys chacoensis*) is a species of rodent in the family Cricetidae. It is known only from central Argentina. Prior to 1994, it was considered a subspecies of *G. griseoflavus*. Then referred to as *G. centralis*, as of...

Central Mexican broad-clawed shrew

Central Mexican broad-clawed shrew

Cryptotis alticola

The Central Mexican broad-clawed shrew (*Cryptotis alticola*) is a species of mammal in the family Soricidae. It is found in the highlands above 2000 m in the Mexican states of Colima, Hidalgo, Jalisco, Michoacán, Mexico, and Puebla, Morelos, and...

Central pebble-mound mouse

Central pebble-mound mouse

Pseudomys johnsoni

The central pebble-mound mouse (*Pseudomys johnsoni*) is a species of rodent in the family Muridae, native to Australia. The Kimberley mouse (*Pseudomys laborifex*) was, until recently, considered distinct from *P. johnsoni*, but they are now known...

Central rock rat

Central rock rat

Zyzomys pedunculatus

The central rock rat (*Zyzomys pedunculatus*), also known as the central thick-tailed rock-rat, Macdonnell Range rock-rat, Australian native mouse, rat à grosse queue (French), or rata coligorda (Spanish), is a species of rodent in the family...

Central Sulawesi echiothrix

Central Sulawesi echiothrix

Echiothrix centrosa

The central Sulawesi echiothrix (*Echiothrix centrosa*) is a species of rodent in the family Muridae. It is found only in Sulawesi, Indonesia.

Central Texas pocket gopher

Central Texas pocket gopher

Geomys texensis

The central Texas pocket gopher or Llano pocket gopher (*Geomys texensis*) is a species of rodent in the family Geomyidae. It is endemic to central Texas in the United States.

Ceram fruit bat

Ceram fruit bat

Pteropus ocularis

The Ceram fruit bat or Seram flying fox (*Pteropus ocularis*) is a species of megabat in the family Pteropodidae. It is endemic to the mountainous forests of two Maluku islands, Buru and Seram, including the Manusela National Park on Seram.

They...

Ceram rat

Ceram rat

Nesoromys ceramicus

The Ceram rat (*Nesoromys ceramicus*) is a species of rodent in the family Muridae. It is found only in Seram Island, Indonesia, where it has been recorded on Mount Mansuela.

Cerrado climbing mouse

Cerrado climbing mouse

Rhipidomys macrurus

The cerrado climbing mouse or long-tailed rhipidomys (*Rhipidomys macrurus*) is an arboreal rodent species in the family Cricetidae from South America. It is found in primary or secondary forests of the cerrado and caatinga in central and eastern...

Cerrado mouse

Cerrado mouse

Thalpomys cerradensis

The cerrado mouse (*Thalpomys cerradensis*) is a rodent species from South America. It is found in the cerrado of Brazil.

Cerrado red-nosed mouse

Cerrado red-nosed mouse

Wiedomys cerradensis

The cerrado red-nosed mouse (*Wiedomys cerradensis*) is a rodent species from South America. It is known from one locality in Bahia, eastern Brazil. The species was found in semi-deciduous forest in the cerrado ecoregion.

Cerradomys goytaca

Cerradomys goytaca

Cerradomys goytaca

Cerradomys goytaca is a species of rodent from South America in the genus *Cerradomys*. It occurs only in sandy plains in southeastern Brazil.

Cerradomys langguthi

Cerradomys langguthi

Cerradomys langguthi

Cerradomys langguthi is a species of rodent from South America in the genus *Cerradomys*. It occurs only in northeastern Brazil and was formerly included in *C. subflavus*.

Cerradomys maracajuensis

Cerradomys maracajuensis
Cerradomys maracajuensis

Cerradomys maracajuensis, also known as the Maracaju oryzomys, is a rodent species from South America. It is terrestrial and is found in gallery forests in Bolivia, Paraguay and nearby Brazil and Peru. It was first discovered near the Brazilian...

Cerradomys marinhos
Cerradomys marinhos
Cerradomys marinhos

Cerradomys marinhos, also known as Marinho's rice rat, is a rodent species from South America. It is found in Minas Gerais, Brazil. It was formerly known as *Oryzomys marinhos*, but was transferred to the new genus *Cerradomys* in 2006.

Cerradomys scotti
Cerradomys scotti
Cerradomys scotti

Cerradomys scotti, also known as Lindbergh's oryzomys, is a rodent species from South America in the genus *Cerradomys*. It is terrestrial and is found in the cerrado (savanna) ecozone of south central Brazil, Bolivia and Paraguay. The species is...

Cerradomys subflavus
Cerradomys subflavus
Cerradomys subflavus

Cerradomys subflavus, also known as the terraced rice rat or flavescent oryzomys, is a rodent species from South America in the genus *Cerradomys*. It is found in the states of Goiás, São Paulo, and Minas Gerais, Brazil. Populations in Bolivia,...

Cerradomys vivoi
Cerradomys vivoi
Cerradomys vivoi

Cerradomys vivoi is a species of rodent from South America in the genus *Cerradomys*. It occurs only in eastern Brazil, in the states of Bahia, Goiás, and Sergipe, and was formerly included in *C. subflavus*.

Cervus elaphus acoronatus
Cervus elaphus acoronatus
Cervus elaphus acoronatus

Cervus elaphus acoronatus is an extinct subspecies of the red deer belonging to the family Cervidae. Some authors consider it a distinct species, *Cervus acoronatus*.

Ceylon spiny mouse
Ceylon spiny mouse
Mus fernandoni

The Ceylon spiny mouse (*Mus fernandoni*) is a species of rodent in the family Muridae. It is found only in Sri Lanka, where it is known as ශ්‍රී ලංකා කටු හීන් මීයා in Sinhala language.

Chacma Baboon
Chacma Baboon
Papio ursinus

The Chacma baboon (*Papio ursinus*) is a member of the Old World monkey family. It is one of the largest of all monkeys. Located primarily in southern Africa, the chacma baboon has a wide variety of social behaviors, including a dominance...

Chaco grass mouse
Chaco grass mouse
Akodon toba

The Chaco grass mouse (*Akodon toba*) is a species of rodent in the family Cricetidae. It is found in Argentina, Bolivia, and Paraguay.
Chacoan gracile opossum
Chacoan gracile opossum
Cryptonanus chacoensis

The chacoan gracile opossum (*Cryptonanus chacoensis*) is a species of opossum in the family Didelphidae. It is native to Argentina, Brazil and Paraguay. Its habitat is seasonally flooded grasslands and forests in and near the Gran Chaco.
Chacoan mara
Chacoan mara
Dolichotis salinicola

The Chacoan mara (*Dolichotis salinicola*) is a relatively large South American rodent of the cavy family. They are a close relative of the better known Patagonian mara.
Chacoan naked-tailed armadillo
Chacoan naked-tailed armadillo
Cabassous chacoensis

The Chacoan naked-tailed armadillo (*Cabassous chacoensis*) is a species of South American armadillo. It is the smallest of the naked-tailed armadillos, having an average head-body length of 30 cm (12 in), while the other species range from 32 to...
Chacoan Peccary
Chacoan Peccary
Catagonus wagneri

The Chacoan peccary or tagua (*Catagonus wagneri* or *Parachoerus wagneri*) is the last extant species of the genus *Catagonus* ; it is a peccary found in the Gran Chaco of Paraguay, Bolivia, and Argentina. Approximately 3,000 remain in the world. It...
Chacoan pygmy opossum
Chacoan pygmy opossum
Chacodelphys formosa

The Chacoan pygmy opossum (*Chacodelphys formosa*) is a recently described genus and species of didelphimorph marsupial. The only species in *Chacodelphys*, *C. formosa*, was known until 2004 from only one specimen collected in 1920 in the Chaco of...
Chacoan tuco-tuco
Chacoan tuco-tuco
Ctenomys dorsalis

The Chacoan tuco-tuco (*Ctenomys dorsalis*) is a species of rodent in the family Ctenomyidae. It lives in Bolivia and Paraguay.
Chadian wild dog
Chadian wild dog
Lycaon pictus sharicus

The African wild dog (*Lycaon pictus*), also called the African painted dog and the African hunting dog, is a wild canine which is a native species to sub-Saharan Africa. It is the largest wild canine in Africa, and the only extant member of the...
Chadwick Beach cotton mouse
Chadwick Beach cotton mouse
Peromyscus gossypinus restrictus

The Chadwick Beach cotton mouse (*Peromyscus gossypinus restrictus*) is a presumed extinct subspecies of the cotton mouse (*Peromyscus gossypinus*). It was confined to

a small area on the Manasota Key peninsula in Florida.

Chaerephon atsinanana
Chaerephon atsinanana
Chaerephon atsinanana

Chaerephon atsinanana is a free-tailed bat found on Madagascar. It was considered a subspecies of the little free-tailed bat until 2010. During the day, they are known to roost in man-made structures such as the roofs or attics of buildings.

Chaerephon jobimena
Chaerephon jobimena
Chaerephon jobimena

Chaerephon jobimena, commonly known as the black and red free-tailed bat, is a species of bat in the family Molossidae. It is endemic to western Madagascar. With a forearm length of 45 to 48 mm (1.8 to 1.9 in), *C. jobimena* is somewhat larger than...

Chalchalero viscacha rat
Chalchalero viscacha rat
Tympanoctomys loschalchalerosorum

The Chalchalero viscacha rat or Chalchalero vizcacha rat (*Tympanoctomys loschalchalerosorum*) is a species of caviomorph rodent in the family Octodontidae. It was formerly considered to be monotypic within the genus *Salinoctomys*, but has been...

Chamois
Chamois
Rupicapra rupicapra

The chamois (*Rupicapra rupicapra*) is a species of goat-antelope native to mountains in Europe. It has also been introduced to the South Island of New Zealand. Some subspecies of chamois are strictly protected in the EU under the European Habitats...

Champion's tree mouse
Champion's tree mouse
Pogonomys championi

Champion's tree mouse (*Pogonomys championi*) is a species of rodent in the family Muridae. It is found only in Papua New Guinea.

Channel Islands spotted skunk
Channel Islands spotted skunk
Spilogale gracilis amphialus

The island spotted skunk (*Spilogale gracilis amphialus*) is an insular endemic carnivore and a subspecies of the western spotted skunk (*Spilogale gracilis*). Little is known about their exact variations from the mainland spotted skunk and...

Chaotung vole
Chaotung vole
Eothenomys olitor

The Chaotung vole (*Eothenomys olitor*) is a species of rodent in the family Cricetidae. It is found only in Yunnan, Sichuan, and Guizhou, China.

Chapa pygmy dormouse
Chapa pygmy dormouse
Typhlomys cinereus chapensis

The Chapa pygmy dormouse or Vietnamese pygmy dormouse (*Typhlomys cinereus chapensis*) is a rodent endemic to Vietnam. It is listed as a critically endangered species. It is often considered a separate species, but is now recognized as a...

Chapin's free-tailed bat

Chapin's free-tailed bat
Chaerephon chapini

Chapin's free-tailed bat (*Chaerephon chapini*) is a species of bat in the family Molossidae. It is found in central and southern Africa.

Chapman's prehensile-tailed hutia
Chapman's prehensile-tailed hutia
Mysateles prehensilis gundlachi

The prehensile-tailed hutia (*Mysateles prehensilis*) is a species of hutia in the subfamily Capromyinae endemic to Cuba. It is the only member of the genus *Mysateles*. It is an arboreal foliovore, found in both primary and secondary forest.

Chapman's Zebra
Chapman's Zebra
Equus quagga chapmani

Chapman's zebra (*Equus quagga chapmani*) is a subspecies of the Plains zebra, named after its discoverer James Chapman. Like the other subspecies of plains zebra, it is a herbivore and undertakes a migration during the wet season to find fresh...

Charming climbing mouse
Charming climbing mouse
Rhipidomys venustus

The charming climbing mouse (*Rhipidomys venustus*) is a species of nocturnal rodent in the family Cricetidae. It is endemic to Venezuela.

Charming thicket rat
Charming thicket rat
Thamnomys venustus

The charming thicket rat (*Thamnomys venustus*) is a species of rodent in the family Muridae. It is described as data deficient as *Thamnomys schoutedeni*. It is found in Democratic Republic of the Congo, Rwanda, and Uganda. Its natural habitat is...

Checkered elephant shrew
Checkered elephant shrew
Rhynchocyon cirnei

The checkered elephant shrew or checkered sengi (*Rhynchocyon cirnei*) is a species of elephant shrew in the family Macroscelididae.

Cheesman's gerbil
Cheesman's gerbil
Gerbillus cheesmani

Cheesman's gerbil (*Gerbillus cheesmani*) is a small rodent in the subfamily Gerbillinae of the family Muridae. It is distributed mainly in Arabian Peninsula to southwestern Iran. It has orange-brown fur, white underparts, large eyes and a very...

Cheetah
Cheetah
Acinonyx jubatus

The cheetah (*Acinonyx jubatus*) is a large cat native to Africa and central Iran. It is the fastest land animal, estimated to be capable of running at 80 to 128 km/h (50 to 80 mph) with the fastest reliably recorded speeds being 93 and 98 km/h (58...

Chelemys delfini
Chelemys delfini
Chelemys delfini

Chelemys delfini, also known as the Magellanic long-clawed akodont, is a species of

rodent in the genus *Chelemys* of family Cricetidae. It is endemic to the areas of Punta Arenas and Torres del Paine National Park in southern Chile, where it lives...

Chelemys macronyx
Chelemys macronyx
Chelemys macronyx

Chelemys macronyx, also known as the Andean long-clawed mouse or Andean long-clawed akodont, is a species of rodent in the genus *Chelemys* of family Cricetidae. It is native to Argentina and Chile, where it is found in Patagonian Nothofagus forest...

Chelemys megalonyx
Chelemys megalonyx
Chelemys megalonyx

Chelemys megalonyx, also known as the large long-clawed mouse or large long-clawed akodont is a species of rodent in the genus *Chelemys* of family Cricetidae. It is endemic to central Chile.

Cheng's jird
Cheng's jird
Meriones chengi

Cheng's jird (*Meriones chengi*) is a species of rodent in the family Muridae. It was named in honour of the Chinese zoologist Professor Tso-hsin Cheng. It is found only in the Turpan Depression of eastern Xinjiang, China.

Cherrie's pocket gopher
Cherrie's pocket gopher
Orthogeomys cherriei

Cherrie's pocket gopher (*Heterogeomys cherriei*) is a species of rodent in the family Geomyidae. It is found in Honduras, Nicaragua, and Costa Rica. It is threatened by habitat loss. Some authors classify it in the genus *Orthogeomys*, but recent...

Chestnut climbing mouse
Chestnut climbing mouse
Dendromus mystacalis

The chestnut climbing mouse (*Dendromus mystacalis*) is a species of rodent in the family Nesomyidae. It is found in Angola, Democratic Republic of the Congo, Ethiopia, Kenya, Lesotho, Malawi, Mozambique, Rwanda, South Africa, Eswatini, Tanzania,...

Chestnut dunnart
Chestnut dunnart
Sminthopsis archeri

The chestnut dunnart (*Sminthopsis archeri*) is a dunnart that was described by Van Dyck in 1986 and is named because of its chestnut colour in the upperparts of its body. The length from snout to tail is 167–210 mm, of which head to anus is...

Chestnut long-tongued bat
Chestnut long-tongued bat
Lionycteris spurrelli

The chestnut long-tongued bat (*Lionycteris spurrelli*) is a bat species from South and Central America. It is monotypic within its genus.

Chestnut sac-winged bat
Chestnut sac-winged bat
Cormura brevirostris

The chestnut sac-winged bat, or Wagner's sac-winged bat (*Cormura brevirostris*) is a species of sac-winged bat native to South and Central America. It is the only

species within its genus.
Chestnut short-tailed bat
Chestnut short-tailed bat
Carollia castanea

The chestnut short-tailed bat (*Carollia castanea*) is a bat species from South and Central America. The species is often confused with the Benkeith's short-tailed bat. It is a mainly frugivorous species that has been known to consume insects (its...
Chestnut tree mouse
Chestnut tree mouse
Pogonomys macrourus

The chestnut tree mouse (*Pogonomys macrourus*) is a species of rodent in the family Muridae. It is found in Indonesia and Papua New Guinea.
Chestnut white-bellied rat
Chestnut white-bellied rat
Niviventer fulvescens

The chestnut white-bellied rat (*Niviventer fulvescens*) is a species of rodent in the family Muridae. It is a small rodent with a distinct bright chestnut upper-coat and a white under-coat. The colour of the upper-coat is variable from very bright...
Chestnut-bellied shrew
Chestnut-bellied shrew
Sorex ventralis

The chestnut-bellied shrew (*Sorex ventralis*) is a species of mammal in the family Soricidae. It is endemic to Mexico.
Chestnut-bellied spiny rat
Chestnut-bellied spiny rat
Maxomys ochraceiventer

The chestnut-bellied spiny rat (*Maxomys ochraceiventer*) is a species of rodent in the family Muridae. It is found in Indonesia and Malaysia.
Chestnut-bellied titi
Chestnut-bellied titi
Plecturocebus caligatus

The chestnut-bellied titi (*Plecturocebus caligatus*) is a species of titi, a type of New World monkey, endemic to Brazil. It was originally described as *Callicebus caligatus* in 1842.
Chestnut-striped opossum
Chestnut-striped opossum
Monodelphis rubida

The chestnut-striped opossum (*Monodelphis rubida*) is an opossum species from South America. It is found in Brazil. More specifically, it is found in East and Central Brazil, including areas such as Goiás, Minas Gerais, and São Paulo. It is...
Chevrier's field mouse
Chevrier's field mouse
Apodemus chevrieri

Chevrier's field mouse (*Apodemus chevrieri*) is a species of rodent in the family Muridae. It is found only in China.
Chiapan climbing rat
Chiapan climbing rat
Tylomys bullaris

The Chiapan climbing rat (*Tylomys bullaris*) is a species of rodent in the family Cricetidae. It is found only in Mexico. The species is known from only one location in Tuxtla Gutiérrez, Chiapas. The habitat in the region is being converted to...

Chiapan deer mouse

Chiapan deer mouse

Peromyscus zarhynchus

The Chiapan deer mouse (*Peromyscus zarhynchus*) is a species of rodent in the family Cricetidae. It is found only in Mexico.

Chibchan water mouse

Chibchan water mouse

Chibchanomys trichotis

The Chibchan water mouse (*Chibchanomys trichotis*) is a species of rodent in the family Cricetidae. It is found in the Andean highlands of Colombia, Peru, and Venezuela. Its natural habitats are tropical cloud forests and streams at elevations from...

Chihuahuan mouse

Chihuahuan mouse

Peromyscus polius

The Chihuahuan mouse (*Peromyscus polius*) is a species of rodent in the family Cricetidae. It is found only in Mexico.

Chihuahuan pocket mouse

Chihuahuan pocket mouse

Chaetodipus eremicus

The Chihuahuan pocket mouse (*Chaetodipus eremicus*) is a species of heteromyid rodent found in the southwestern United States and Mexico. It was formerly considered a subspecies of the desert pocket mouse (*C. pencillatus*), but was determined to...

Chilean Dolphin

Chilean Dolphin

Cephalorhynchus eutropia

The Chilean dolphin (*Cephalorhynchus eutropia*) is a small dolphin found only off the coast of Chile, where it is commonly referred to as tonina. It is perhaps one of the least studied of all cetaceans.

Chilean myotis

Chilean myotis

Myotis chiloensis

The Chilean myotis (*Myotis chiloensis*) is a species of vesper bat found in southern South America.

Chilean rock rat

Chilean rock rat

Aconaemys fuscus

The Chilean rock rat (*Aconaemys fuscus*) is a species of rodent in the family Octodontidae. It is found in the high Andes of Argentina and Chile.

Chilonatalus macer

Chilonatalus macer

Chilonatalus macer

Chilonatalus macer is a species of bat endemic to Cuba.

Chiltan ibex

Chiltan ibex

Capra aegagrus chialtanensis

The Chiltan ibex or Chiltan goat (*Capra aegagrus chialtanensis*) is a wild goat endemic to Chiltan, Balochistan, Pakistan.

Chinanteco deer mouse
Chinanteco deer mouse
Habromys chinanteco

The Chinanteco deer mouse (*Habromys chinanteco*) is a species of rodent in the family Cricetidae. It is endemic to—found only in Mexico.

Chinese Bamboo Rat
Chinese Bamboo Rat
Rhizomys sinensis

The Chinese bamboo rat (*Rhizomys sinensis*) is a species of rodent found in Asia. It has a very wide range and is common in some localities. It is considered a plantation pest in parts of China and is presumed to have a large total population.

Chinese dormouse
Chinese dormouse
Chaetocauda sichuanensis

The Chinese dormouse or Sichuan dormouse (*Chaetocauda sichuanensis*) is a species of dormouse found in subalpine mixed forests in northern Sichuan, China, where it is known from Jiuzhaigou and Wanglang Nature Reserves. It is known only from two...

Chinese Ferret-Badger
Chinese Ferret-Badger
Melogale moschata

The Chinese ferret-badger (*Melogale moschata*), also known as the small-toothed ferret-badger is a member of the Mustelidae, and widely distributed in Southeast Asia. It is listed as Least Concern on the IUCN Red List and considered tolerant of...

Chinese giant flying squirrel
Chinese giant flying squirrel
Petaurista xanthotis

The Chinese giant flying squirrel (*Petaurista xanthotis*) is a species of rodent in the family Sciuridae. It is endemic to China. This squirrel inhabits high elevation spruce forests in China. It feeds nocturnally on young shoots, leaves, and pine...

Chinese Goral
Chinese Goral
Naemorhedus griseus

The Chinese goral (*Naemorhedus griseus*), also known as the grey long-tailed goral or central Chinese goral, is a species of goral, a small goat-like ungulate, native to mountainous regions of Myanmar, China, India, Thailand, Vietnam, and...

Chinese Hamster
Chinese Hamster
Cricetulus griseus

The Chinese hamster (*Cricetulus griseus* or *Cricetulus barabensis griseus*) is a rodent in the genus *Cricetulus* of the subfamily Cricetidae that originated in the deserts of northern China and Mongolia. They are distinguished by an uncommonly long...

Chinese hare
Chinese hare
Lepus sinensis

The Chinese hare (*Lepus sinensis*) is a species of mammal in the family Leporidae. It is found in China, Taiwan and Vietnam.

Chinese highland shrew

Chinese highland shrew
Sorex excelsus

The Chinese highland shrew (*Sorex excelsus*) is a species of shrew in the family Soricidae. It is found in China.

Chinese jumping mouse
Chinese jumping mouse
Eozapus setchuanus

The Chinese jumping mouse (*Eozapus setchuanus*) is a species of rodent in the family Dipodidae. It is monotypic within the genus *Eozapus*. It is endemic to China where its natural habitat is temperate forests, steppes and meadows in mountainous...

Chinese mole shrew
Chinese mole shrew
Anourosorex squamipes

The Chinese mole shrew (*Anourosorex squamipes*) is one of four species of Asian mole shrew in the genus *Anourosorex*. Despite the name, it is found not only in China but also in northeast India, Myanmar, Thailand, and Vietnam. Like all other...

Chinese Mountain Cat
Chinese Mountain Cat
Felis bieti

The Chinese mountain cat (*Felis bieti*), also known as Chinese desert cat and Chinese steppe cat, is a small wild cat endemic to western China that has been listed as Vulnerable on the IUCN Red List since 2002, as the effective population size...

Chinese noctule
Chinese noctule
Nyctalus plancyi

The Chinese noctule (*Nyctalus plancyi*) is a common and widespread species of bat belonging to the family Vespertilionidae.

Chinese Pangolin
Chinese Pangolin
Manis pentadactyla

The Chinese pangolin (*Manis pentadactyla*) is a pangolin native to the northern Indian subcontinent, northern parts of Southeast Asia and southern China. It has been listed as Critically Endangered on the IUCN Red List since 2014, as the wild...

Chinese pipistrelle
Chinese pipistrelle
Hypsugo pulveratus

The Chinese pipistrelle (*Hypsugo pulveratus*) is a species of vesper bat in the family Vespertilionidae. It is found in China, Laos, Thailand, and Vietnam.

Chinese pygmy dormouse
Chinese pygmy dormouse
Typhlomys cinereus

The Chinese pygmy dormouse (*Typhlomys cinereus*) is a species of rodent of the family Platacanthomyidae found in China and Vietnam.

Chinese red pika
Chinese red pika
Ochotona erythrotis

The Chinese red pika (*Ochotona erythrotis*) is a species of mammal in the family

Ochotonidae. Typical of a pika it has short limbs, a small tail and round ears. Specific to the Chinese red pika has distinctive red color in its pelt. The Chinese...

Chinese rufous horseshoe bat
Chinese rufous horseshoe bat
Rhinolophus sinicus

The Chinese rufous horseshoe bat (*Rhinolophus sinicus*) is a species of bat in the family Rhinolophidae. It is found in Bhutan, China, India, Nepal, and Vietnam. The species is most easily confused with *R. affinis*, from which it is best...

Chinese scrub vole
Chinese scrub vole
Neodon irene

The Chinese scrub vole, or Irene's mountain vole, (*Neodon irene*) is a species of rodent in the family Cricetidae. It is endemic to mountainous parts of southern China and is very similar to the Sikkim mountain vole in appearance. The...

Chinese shrew
Chinese shrew
Sorex sinalis

The Chinese shrew or dusky shrew (*Sorex sinalis*) is a species of mammal in the family Soricidae. It is endemic to China, where it occurs in Gansu, Sichuan (including Jiuzhaigou), and Shaanxi. Its natural habitat is subtropical or tropical dry...

Chinese shrew mole
Chinese shrew mole
Uropsilus soricipes

The Chinese shrew mole (*Uropsilus soricipes*) is a species of mammal in the family Talpidae. It is endemic to China, Sichuan Province. Its natural habitat is temperate forests.

Chinese striped hamster
Chinese striped hamster
Cricetulus barabensis

The Chinese striped hamster (*Cricetulus barabensis*), also known as the striped dwarf hamster, is a species of hamster. It is distributed across Northern Asia, from southern Siberia through Mongolia and northeastern China to northern North Korea....

Chinese water myotis
Chinese water myotis
Myotis laniger

The Chinese water myotis (*Myotis laniger*) is a species of vesper bat. It is native to eastern India, Vietnam and China where it is found in forest habitats. It is believed to roost in caves and hollow trees.

Chinese water shrew
Chinese water shrew
Chimarrogale styani

The Chinese water shrew (*Chimarrogale styani*) is a species of mammal in the family Soricidae. It is found in China and Myanmar.

Chinese white-bellied rat
Chinese white-bellied rat
Niviventer confucianus

The Chinese white-bellied rat (*Niviventer confucianus*) is a species of rodent in the family Muridae. It is widely spread in China and also occurs in northern

Myanmar, northwest Thailand, and northwest Vietnam. It might also occur in northern Laos.

Chinese white-toothed shrew
Chinese white-toothed shrew
Crocidura rapax

The Chinese white-toothed shrew (*Crocidura rapax*) is a species of mammal in the family Soricidae.

Chinese zokor
Chinese zokor
Eospalax fontanierii

The Chinese zokor (*Eospalax fontanierii*) is a species of rodent in the family Spalacidae. It is endemic to China, ranging from Qinghai Province eastwards to Beijing in steppe and alpine grasslands. Henri Milne-Edwards first described it in 1867....

Chinkara
Chinkara
Gazella bennettii

The chinkara (*Gazella bennettii*) is a gazelle species native to Asia. Gazelles are known as swift animals. Appreciated for their grace, gazelles are a symbol most commonly associated in Arabic literature with female beauty.

Chiriqui brown mouse
Chiriqui brown mouse
Scotinomys xerampelinus

The Chiriqui brown mouse (*Scotinomys xerampelinus*), also known as the long-tailed singing mouse, is a species of rodent in the family Cricetidae. It is found in cloud forest and paramo at elevations of 2100 to 3400 m in Costa Rica and...

Chiriqui harvest mouse
Chiriqui harvest mouse
Reithrodontomys creper

The Chiriqui harvest mouse (*Reithrodontomys creper*) is a species of rodent in the family Cricetidae. It is found in Costa Rica and Panama.

Chiriqui pocket gopher
Chiriqui pocket gopher
Orthogeomys cavator

The Chiriqui pocket gopher (*Heterogeomys cavator*) is a species of rodent in the family Geomyidae. It is found in Costa Rica and Panama. Some authors classify it in the genus *Orthogeomys*, but recent research has allowed this and its related...

Chiriquinan serotine
Chiriquinan serotine
Eptesicus chiriquinus

The Chiriquinan serotine (*Eptesicus chiriquinus*) is a species of house bat. The Chiriquinan serotine is listed as Least Concern on the IUCN Red List due to its wide distribution and the unlikelihood of its speedy decline. However, the species is...

Chiroderma vizottoi
Chiroderma vizottoi
Chiroderma vizottoi

Chiroderma vizottoi is a species of frugivorous bat found in the northeast of Brazil.

Chiruromys lamia
Chiruromys lamia

Chiruromys lamia

Chiruromys lamia, also known as the lamia or the broad-headed tree mouse, is a species of rodent found chiefly in southeastern New Guinea. It is arboreal, living in hollow tree nests, and is found at elevations of 1,200–2,300 metres...

Chisel-toothed kangaroo rat

Chisel-toothed kangaroo rat

Dipodomys microps

The chisel-toothed kangaroo rat (*Dipodomys microps*) is a species of rodent in the family Heteromyidae. There are 13 sub-species. Saltbush leaves are a major dietary component, requiring specialized physiology to eliminate the salt while retaining...

Choco broad-nosed bat

Choco broad-nosed bat

Platyrrhinus chocoensis

The Choco broad-nosed bat (*Platyrrhinus chocoensis*) is a species of bat in the family Phyllostomidae. It is native to Colombia, Panama, and Ecuador, where it is found in the Choco region lowlands. It is threatened by habitat loss. In 2013, Bat...

Chocoan long-tongued bat

Chocoan long-tongued bat

Lonchophylla chocoana

The Chocoan long-tongued bat (*Lonchophylla chocoana*) is a species of bat found in South America. It was described as a new species in 2004.

Chocolate pipistrelle

Chocolate pipistrelle

Falsistrellus affinis

The chocolate pipistrelle (*Hypsugo affinis*) is a species of vesper bat in the family Vespertilionidae. It is found in China, India, Myanmar, Nepal, and Sri Lanka.

Chocolate wattled bat

Chocolate wattled bat

Chalinolobus morio

The chocolate wattled bat, species *Chalinolobus morio*, is a bat allied to the family Vespertilionidae. It is found only in Australia, including the island Tasmania, and widespread in southern regions. It is known to reside from sea level to at...

Christie's big-eared bat

Christie's big-eared bat

Plecotus christii

Christie's long-eared bat (*Plecotus christii*), also known as the Egyptian long-eared bat, Christie's big-eared bat, or gray long-eared bat, is a species of vesper bat in the family Vespertilionidae. It is known from North Africa and the Middle...

Christine's margareta rat

Christine's margareta rat

Margaretamys christinae

Christine's margareta rat (*Margaretamys christinae*) is a species of rodent in the family Muridae. It is found only in Indonesia on the island of Sulawesi, in a mountainous part of the southeastern peninsula. It was first discovered by an...

Christmas Island flying fox

Christmas Island flying fox

Pteropus melanotus natalis

The Christmas Island flying fox or Christmas Island fruit bat, as the name suggests, is a flying fox endemic to Christmas Island. It is unclear if it should be considered a distinct species (*Pteropus natalis*), or a subspecies of the black-eared...

Christmas Island Shrew
Christmas Island Shrew
Crocidura trichura

The Christmas Island shrew (*Crocidura trichura*) is an extremely rare or possibly extinct shrew. There is conclusive evidence that this species has declined dramatically since 1900, yet the reason is unproven.

Christy's dormouse
Christy's dormouse
Graphiurus christyi

Christy's dormouse (*Graphiurus christyi*) is a species of rodent in the family Gliridae. It is found in Cameroon and Democratic Republic of the Congo. Its natural habitat is subtropical or tropical moist lowland forests.

Chudeau's spiny mouse
Chudeau's spiny mouse
Acomys chudeaui

Chudeau's spiny mouse (*Acomys chudeaui*) is a species of rodent in the family Muridae found in Mauritania and Morocco. Its natural habitats are rocky areas and hot deserts.

Cinderella fat-tailed mouse opossum
Cinderella fat-tailed mouse opossum
Thylamys cinderella

The Cinderella fat-tailed mouse opossum (*Thylamys cinderella*) is a species of opossum in the family Didelphidae. It is found in northern Argentina and southern Bolivia, in the eastern foothills of the Andes. Its dorsal fur is gray brown to dark...

Cinderella shrew
Cinderella shrew
Crocidura cinderella

The cinderella shrew (*Crocidura cinderella*) is a species of mammal in the family Soricidae. It is found in Burkina Faso, Gambia, Guinea-Bissau, Mali, Mauritania, Niger, and Senegal. Its natural habitat is dry savanna.

Cinereus ringtail possum
Cinereus ringtail possum
Pseudochirulus cinereus

The Daintree River ringtail possum (*Pseudochirulus cinereus*), also known as the cinereus ringtail possum, is a species of possum found in northeastern Queensland, Australia. It was long believed to be the same as the Herbert River ringtail...

Cinereus Shrew
Cinereus Shrew
Sorex cinereus

The Cinereus shrew (*Sorex cinereus*) is a small mole-like mammal found in the northern parts of North America. This is the most widely distributed shrew in North America, where it is also known as the Common shrew.

Cinnamon antechinus
Cinnamon antechinus
Antechinus leo

The cinnamon antechinus (*Antechinus leo*), also known as the Iron Ranges antechinus and the Cape York antechinus, is a species of small carnivorous marsupial of the family Dasyuridae. It is the only mammal endemic to Cape York Peninsula, being...

Cinnamon bear

Cinnamon bear

Ursus americanus cinnamomum

The cinnamon bear (*Ursus americanus cinnamomum*) is both a highly variable color morph and a subspecies of the American black bear, native to the central, eastern, and western areas of the United States and Canada. Established populations are...

Cinnamon dog-faced bat

Cinnamon dog-faced bat

Cynomops abrasus

The cinnamon dog-faced bat (*Cynomops abrasus*), is a South American bat species of the family Molossidae. It is found in northern and central South America.

Cinnamon myotis

Cinnamon myotis

Myotis fortidens

The cinnamon myotis (*Myotis fortidens*) is a species of vesper bat. It is found in Guatemala and Mexico.

Cinnamon red bat

Cinnamon red bat

Lasiurus varius

The cinnamon red bat (*Lasiurus varius*) is a species of bat in the family Vespertilionidae. It was first described from a specimen that had been collected in Chile. For more than one hundred years after its initial description, it was largely...

Cinnamon-colored Oldfield mouse

Cinnamon-colored Oldfield mouse

Thomasomys cinnameus

The cinnamon-colored Oldfield mouse (*Thomasomys cinnameus*), also called the "cinnamon-colored *Thomasomys*", is a species of rodent in the family Cricetidae. It is present in the Cordillera Oriental of the Andes from north central Ecuador to...

Ciscaucasian hamster

Ciscaucasian hamster

Mesocricetus raddei

The Ciscaucasian hamster (*Mesocricetus raddei*) is a species of rodent in the family Cricetidae. It is also known as the Georgian hamster and is found only in Georgia and Russia.

Claire's mouse lemur

Claire's mouse lemur

Microcebus mambiratra

Claire's mouse lemur (*Microcebus mambiratra*), or the Nosy Be mouse lemur, is a newly described species of lemur from the genus of the mouse lemurs (*Microcebus*). It lives on the island Nosy Bé in the Antsiranana province of Madagascar, and on the...

Clara's echymipera

Clara's echymipera

Echymipera clara

Clara's echymipera (*Echymipera clara*), or Clara's spiny bandicoot or white-lipped bandicoot, is a species of marsupial in the family Peramelidae. It is found in West Papua, Indonesia and Papua New Guinea. Its natural habitat is subtropical or...

Clarke's vole
Clarke's vole
Microtus clarkei

Clarke's vole (*Microtus clarkei*) is a species of rodent in the family Cricetidae. It is found only in China.

Clear-winged woolly bat
Clear-winged woolly bat
Kerivoula pellucida

The clear-winged woolly bat (*Kerivoula pellucida*) is a species of vesper bat in the family Vespertilionidae. It is found in Brunei, Indonesia, Malaysia, and the Philippines. Members of this species are relatively small, typically weighing about...

Cliff Chipmunk
Cliff Chipmunk
Neotamias dorsalis

The cliff chipmunk (*Neotamias dorsalis*) is a small, bushy-tailed squirrel that typically lives along cliff walls or boulder fields bordering Pinyon-juniper woodlands in the Western United States and Mexico (commonly spotted in northern Arizona...

Climbing shrew
Climbing shrew
Suncus megalura

The climbing shrew (*Suncus megalurus*) is a species of mammal in the family Soricidae, which is found in subtropical Africa. It is found in Angola, Benin, Burundi, Cameroon, Central African Republic, Republic of the Congo, Democratic Republic of...

Cloud forest grass mouse
Cloud forest grass mouse
Akodon torques

The cloud forest grass mouse (*Akodon torques*) is a species of rodent in the family Cricetidae. It is found only in Peru.

Clouded Leopard
Clouded Leopard
Neofelis nebulosa

The Clouded leopard (*Neofelis nebulosa*) is a wild cat inhabiting dense forests of the Himalayas through mainland Southeast Asia into South China. In the early 19th century, a Clouded leopard was brought to London from China and described in 1821....

Clymene Dolphin
Clymene Dolphin
Stenella clymene

The Clymene dolphin (*Stenella clymene*) is a dolphin native to the Atlantic Ocean. It is the only confirmed case of hybrid speciation in marine mammals, descending from the Spinner dolphin and the Striped dolphin.

Coast mole
Coast mole
Scapanus orarius

The coast mole or Pacific mole (*Scapanus orarius*) is a medium-sized North American mole found in forested and open areas with moist soils along the Pacific coast from southwestern British Columbia to northwestern California.
Coastal black-handed titi

Coastal black-handed titi
Callicebus melanochir

The coastal black-handed titi (*Callicebus melanochir*) is a species of titi, a type of New World monkey, endemic to Brazil.

Coastal sheath-tailed bat
Coastal sheath-tailed bat
Taphozous australis

The coastal sheath-tailed bat (*Taphozous australis*), or coastal tomb bat, is a species of sheath-tailed bat in the family Emballonuridae. It is found in Australia and Papua New Guinea.

Coastal topi
Coastal topi
Damaliscus lunatus topi

The coastal topi (*Damaliscus lunatus topi*) is a highly social antelope of the genus *Damaliscus*. It is a subspecies of the common tsessebe.

Coccymys kirrhos
Coccymys kirrhos
Coccymys kirrhos

Coccymys kirrhos is a rodent in the family Muridae that is native to New Guinea. The species was described in 2009.

Coccymys shawmayeri
Coccymys shawmayeri
Coccymys shawmayeri

Coccymys shawmayeri is a rodent in the family Muridae native to New Guinea. The species was described in 2009.

Cochabamba grass mouse
Cochabamba grass mouse
Akodon siberiae

The Cochabamba grass mouse (*Akodon siberiae*) is a species of rodent in the family Cricetidae. It is found only in Bolivia. Its natural habitat is subtropical or tropical moist lowland forests.

Cohen's horseshoe bat
Cohen's horseshoe bat
Rhinolophus cohenae

Cohen's horseshoe bat (*Rhinolophus cohenae*) is a species of bat belonging to the family Rhinolophidae, endemic to South Africa. It was first described in 2012. The species was named after Lientjie Cohen who collected the type specimen in 2004....

Coiba Island howler
Coiba Island howler
Alouatta coibensis

The Coiba Island howler (*Alouatta coibensis*) is a type of howler monkey, a type of New World monkey, endemic to Panama. Although the Coiba Island howler has been recognized as a separate species by a number of authorities since a 1987 study of...

Coiban agouti
Coiban agouti
Dasyprocta coibae

The Coiban agouti (*Dasyprocta coibae*) is a species of rodent in the family Dasyproctidae. It is endemic to the island of Coiba (Panama) and resembles the more widespread Central American agouti. It is threatened by habitat loss.

Coiban mastiff bat
Coiban mastiff bat
Molossus coibensis

The Coiban mastiff bat (*Molossus coibensis*) is a species of bat in the family Molossidae. Its range extends from Chiapas in southern Mexico to Mato Grosso in Brazil, including Peru, Ecuador, Venezuela, Panama, Costa Rica and El Salvador. The...

Coimbra Filho's titi
Coimbra Filho's titi
Callicebus coimbrai

Coimbra Filho's titi (*Callicebus coimbrai*) or just Coimbra's titi is a species of titi, a type of New World monkey, endemic to forests in the Brazilian states of Bahia and Sergipe. It was first discovered by Shuji Kobayashi. It is considered one...

Coke's hartebeest
Coke's hartebeest
Alcelaphus buselaphus cokii

Coke's hartebeest (*Alcelaphus buselaphus cokii*) or kongoni is an antelope native to Kenya and Tanzania. It can breed with Lelwel hartebeest to produce a hybrid known as the Kenya Highland hartebeest (*Alcelaphus buselaphus lelwel* x *cokii*).
Colburn's tuco-tuco
Colburn's tuco-tuco
Ctenomys colburni

Colburn's tuco-tuco (*Ctenomys colburni*) is a species of rodent in the family Ctenomyidae. It is known only from Argentina.
Collared Brown Lemur
Collared Brown Lemur
Eulemur collaris

The collared brown lemur (*Eulemur collaris*), also known as the red-collared brown lemur or red-collared lemur, is a medium-sized strepsirrhine primate and one of twelve species of brown lemur in the family Lemnidae. It is only found in...
Collared Mangabey
Collared Mangabey
Cercopithecus torquatus

The collared mangabey (*Cercopithecus torquatus*), also called red-capped mangabey and white-collared mangabey is a species of primate in the family Cercopithecidae of Old World monkeys. It formerly included the sooty mangabey as a subspecies. As...
Collared mongoose
Collared mongoose
Herpestes semitorquatus

The collared mongoose (*Urva semitorquata*) is a mongoose species native to Borneo and Sumatra; its presence in the Philippines is uncertain. It is listed as Near Threatened on the IUCN Red List.
Collared Peccary
Collared Peccary
Pecari tajacu

The Collared peccary (*Dicotyles tajacu*) is a species of even-toed mammal found in North, Central, and South America. They are commonly referred to as javelina, saíno, or báquiro, although these terms are also used to describe other species in the...
Collared Pika

Collared Pika
Ochotona collaris

The collared pika (*Ochotona collaris*) is a species of mammal in the pika family, Ochotonidae, and part of the order Lagomorpha, which comprises rabbits, hares, and pikas. It is a small (about 160 g) alpine lagomorph that lives in boulder fields...

Collared pipistrelle
Collared pipistrelle
Arielulus aureocollaris

The collared sprite or collared pipistrelle (*Thainycteris aureocollaris*) is a species of vesper bat found in Laos and Thailand.

Collared titi
Collared titi
Cheracebus torquatus

The collared titi (*Cheracebus torquatus*) is a species of titi, a type of New World monkey. It is endemic to northern Brazil.

Collared tuco-tuco
Collared tuco-tuco
Ctenomys torquatus

The collared tuco-tuco (*Ctenomys torquatus*) is a tuco-tuco species from South America. It is found in southern Brazil, Uruguay and northern Argentina where it lives underground in a burrow it digs in savannah habitats. It is a relatively common...

Collie's squirrel
Collie's squirrel
Sciurus colliaei

Collie's squirrel (*Sciurus colliaei*) is a tree squirrel in the genus *Sciurus* endemic to Mexico.

Collins' squirrel monkey
Collins' squirrel monkey
Saimiri collinsi

Collins' squirrel monkey (*Saimiri collinsii*) is a species of squirrel monkey from Brazil. It had been considered a subspecies of the common squirrel monkey (*S. sciureus*) until a genetic study by Jessica Lynch Alfaro et al. elevated it to...

Colombian black-handed titi
Colombian black-handed titi
Cheracebus medemi

The Colombian black-handed titi (*Cheracebus medemi*) is a species of titi, a type of New World monkey, endemic to Colombia. It was originally described in the genus *Callicebus* in 1963.

Colombian bonneted bat
Colombian bonneted bat
Eumops trumbulli

The Colombian bonneted bat (*Eumops trumbulli*), also known as Trumbull's bonneted bat, is a bat species from South America. It is found in Bolivia, Brazil, Colombia, French Guiana, Guyana, Peru, Suriname and Venezuela.

Colombian forest mouse
Colombian forest mouse
Chilomys instans

The Colombian forest mouse (*Chilomys instans*) is a species of rodent in the family Cricetidae. Some authorities consider it to be the only species in the genus

Chilomys, while others accept *Chilomys fumeus* as being a valid species, and it may...

Colombian small-eared shrew
Colombian small-eared shrew
Cryptotis colombiana

The Colombian small-eared shrew (*Cryptotis colombiana*) is a species of mammal in the family Soricidae. It is endemic to Colombia, where it is known from the Cordillera Central in Antioquia Department at elevations from 1750 to 2800 m. It is...

Colombian soft-furred spiny rat
Colombian soft-furred spiny rat
Diplomys caniceps

The Colombian soft-furred spiny-rat (*Diplomys caniceps*), or arboreal soft-furred spiny-rat, is a species of rodent in the family Echimyidae. It is found in Colombia and Ecuador. Its natural habitat is subtropical or tropical moist lowland forests.

Colombian spider monkey
Colombian spider monkey
Ateles fusciceps rufiventris

The Colombian spider monkey (*Ateles fusciceps rufiventris*), is a subspecies of the Black-headed spider monkey, a type of New World monkey, found in Colombia and Panama. Some authorities, such as Froelich (1991), Collins and Dubach (2001) and...

Colombian spiny rat
Colombian spiny rat
Proechimys canicollis

The Colombian spiny-rat (*Proechimys canicollis*) is a species of rodent in the family Echimyidae. It is found in Colombia and Venezuela.

Colombian Weasel
Colombian Weasel
Mustela felipei

The Colombian weasel (*Neogale felipei*) is a very rare small mammal. It is the second-smallest living carnivoran and probably the rarest carnivoran in South America. The Colombian weasel is also one of the least studied carnivore species in the...

Colombian woolly monkey
Colombian woolly monkey
Lagothrix lugens

The Colombian woolly monkey (*Lagothrix lagothricha lugens*) is a critically endangered subspecies of the common woolly monkey from Colombia. It was initially thought to be a subspecies of the common woolly monkey (*L. lagothricha*), but was later...

Colorado Chipmunk
Colorado Chipmunk
Neotamias quadrivittatus

The Colorado chipmunk (*Neotamias quadrivittatus*) is a species of chipmunk in the squirrel family Sciuridae. It is endemic to Colorado, Utah, Arizona and New Mexico in the United States.

Colombian Ground Squirrel
Colombian Ground Squirrel
Urocitellus columbianus

The Columbian ground squirrel (*Urocitellus columbianus*) is a species of rodent common in certain regions of Canada and the northwestern United States. It is the

second largest member of the genus *Urocitellus*, which is part of the tribe...
Columbian White-Tailed Deer
Columbian White-Tailed Deer
Odocoileus virginianus leucurus

The Columbian white-tailed deer (*Odocoileus virginianus leucurus*) is one of the several subspecies of White-tailed deer in North America. It is named after the Columbia River in Oregon and Washington where it congregates.

Comb-toed jerboa
Comb-toed jerboa
Paradipus ctenodactylus

The comb-toed jerboa (*Paradipus ctenodactylus*) is a species of rodent in the family *Dipodidae*. It is monotypic within the genus *Paradipus*. It is found in Kazakhstan, Turkmenistan, and Uzbekistan.

Commerson's Dolphin
Commerson's Dolphin
Cephalorhynchus commersonii

Commerson's dolphin (*Cephalorhynchus commersonii*), also referred to by the common names jacobita, skunk dolphin, piebald dolphin, panda dolphin, or tonina overa (in South America), is a small oceanic dolphin of the genus *Cephalorhynchus*....

Commerson's roundleaf bat
Commerson's roundleaf bat
Macronycteris commersoni

Commerson's roundleaf bat (*Macronycteris commersoni*), also known as Commerson's leaf-nosed bat, is a species of bat endemic to Madagascar. It is named after French naturalist Philibert Commerson (1727-1773). Bat populations of Africa or São Tomé...

Commissaris's long-tongued bat
Commissaris's long-tongued bat
Glossophaga commissarisi

Commissaris's long-tongued bat (*Glossophaga commissarisi*) is a bat species from South and Central America.

Common beisa oryx
Common beisa oryx
Oryx beisa beisa

The common beisa oryx (*Oryx beisa beisa*), also known as the beisa oryx, is the nominate subspecies of the East African oryx native to the Horn of Africa and Kenya. It is closely related to the fringe-eared oryx. There are four species of oryx,...

Common bent-wing bat
Common bent-wing bat
Miniopterus schreibersii

The common bent-wing bat (*Miniopterus schreibersii*), also known as the Schreibers's long-fingered bat or Schreibers's bat, is a species of insectivorous bat. They appear to have dispersed from a subtropical origin and distributed throughout the...

Common big-eared bat
Common big-eared bat
Micronycteris microtis

The common big-eared bat (*Micronycteris microtis*) is a bat species from South and Central America. It is a neotropical leaf-nosed bat (a member of the family *Phyllostomidae*). Recent research has demonstrated this bat's ability to find

motionless...

Common blossom bat

Common blossom bat

Syconycteris australis

The common blossom bat (*Syconycteris australis*) also known as the southern blossom bat or Queensland blossom bat, is a megabat in the family Pteropodidae. The common blossom bat feeds mostly on nectar and pollen rather than fruit. It is one of...

Common Bottlenose Dolphin

Common Bottlenose Dolphin

Tursiops truncatus

The Common bottlenose dolphin (*Tursiops truncatus*) is a wide-ranging marine mammal of the family Delphinidae. The common bottlenose dolphin is a very familiar dolphin species due to the wide exposure it receives in captivity in marine parks and...

Common Brown Lemur

Common Brown Lemur

Eulemur fulvus

The common brown lemur (*Eulemur fulvus*) is a species of lemur in the family Lemnidae. It is found in Madagascar and has been introduced to Mayotte.

Common Brushtail Possum

Common Brushtail Possum

Trichosurus vulpecula

The Common brushtail possum (*Trichosurus vulpecula*) is a nocturnal, semiarboreal marsupial native to Australia. It is the second-largest of the possums. It is mainly a folivore and in most Australian habitats, eucalyptus leaves are a significant...

Common Chimpanzee

Common Chimpanzee

Pan troglodytes

The chimpanzee (*Pan troglodytes*) is a species of great ape native to the forest and savannah of tropical Africa. When its close relative the bonobo was more commonly known as the Pygmy chimpanzee, this species was often called the Common...

Common Duiker

Common Duiker

Sylvicapra grimmia

The common duiker (*Sylvicapra grimmia*), also known as the grey or bush duiker, is a small antelope found everywhere in Africa south of the Sahara, excluding the Horn of Africa and the rainforests of the central and western parts of the...

Common Dwarf Mongoose

Common Dwarf Mongoose

Helogale parvula

The common dwarf mongoose (*Helogale parvula*) is a mongoose species native to Angola, northern Namibia, KwaZulu-Natal in South Africa, Zambia and East Africa. It is part of the genus *Helogale* and as such is related to *H. hirtula*.

Common echymipera

Common echymipera

Echymipera kalubu

The common echymipera (*Echymipera kalubu*), or common spiny bandicoot, is a bandicoot. It is long-snouted even by bandicoot standards. The upper parts are a coarse reddish brown, flecked with spiny buff and black hairs. The tail is short and...

Common Eland

Common Eland
Taurotragus oryx

The Common eland (*Taurotragus oryx*) is an antelope species from East and Southern Africa. It was scientifically described by Peter Simon Pallas in 1766. The Common eland is used by humans for leather, meat, and milk, and has been domesticated in...

Common fat-tailed mouse opossum
Common fat-tailed mouse opossum
Thylamys pusillus

The common fat-tailed mouse opossum (*Thylamys pusillus*) is a species of opossum in the family Didelphidae. It occurs in Argentina, Bolivia, Brazil, and Paraguay in chaco and Andean foothill habitats. Its head-and-body length is about 75 to 120...

Common Genet
Common Genet
Genetta genetta

The Common genet (*Genetta genetta*) is a small feliform mammal. It is native to Africa and was introduced to southwestern Europe.

Common gundi
Common gundi
Ctenodactylus gundi

The common gundi (*Ctenodactylus gundi*) is a species of rodent in the family Ctenodactylidae. It is found in Algeria, Libya, Morocco, and Tunisia. The parasitic organism *Toxoplasma gondii* was first described in 1908 in Tunis by Charles Nicolle...

Common Hippo
Common Hippo
Hippopotamus amphibius

The hippopotamus (*Hippopotamus amphibius*) is a large semiaquatic mammal native to sub-Saharan Africa. It is one of only two extant species in the family Hippopotamidae, the other being the Pygmy hippopotamus. Its name comes from the ancient Greek...

Common Kusimanse
Common Kusimanse
Crossarchus obscurus

The common kusimanse (*Crossarchus obscurus*), also known as the long-nosed kusimanse or simply kusimanse, is a small, diurnal member of the Mungotinae.

Common Marmoset
Common Marmoset
Callithrix jacchus

The Common marmoset (*Callithrix jacchus*) is a unique New World Monkey with a rather unusual appearance and a number of special adaptations. Marmosets have an arboreal locomotion similar to squirrels. They can hang onto trees vertically leap...

Common Minke Whale
Common Minke Whale
Balaenoptera acutorostrata

The Common minke whale (*Balaenoptera acutorostrata*) is a small member of the suborder of baleen whales. It is the smallest species of the rorquals and the second smallest species of baleen whale. American whalers in the 19th century simply...

Common mole-rat
Common mole-rat

Cryptomys hottentotus

The common mole-rat, African mole-rat, or Hottentot mole-rat, (*Cryptomys hottentotus*) is a burrowing rodent found in Southern Africa, in particular in the Western Cape province of South Africa. It also occurs in Eswatini, Lesotho, Malawi,...

Common Noctule

Common Noctule

Nyctalus noctula

The Common noctule (*Nyctalus noctula*) is an insectivorous bat that is widespread throughout Europe, Asia, and North Africa. These bats hibernate in large groups and females migrate during early pregnancy to return to their natal maternity...

Common Opossum

Common Opossum

Didelphis marsupialis

The Common opossum (*Didelphis marsupialis*) is a marsupial species living in the Americas. It is a cat-sized omnivorous animal, which can be recognized by its prehensile tail and its tendency to feign death when cornered.

Common Pipistrelle

Common Pipistrelle

Pipistrellus pipistrellus

The common pipistrelle (*Pipistrellus pipistrellus*) is a small pipistrelle microbat whose very large range extends across most of Europe, North Africa, South Asia, and may extend into Korea. It is one of the most common bat species in the British...

Common planigale

Common planigale

Planigale maculata

The common planigale (*Planigale maculata*), also known as the pygmy planigale or the coastal planigale, is one of many small marsupial carnivores known as "marsupial mice" found in Australia. There they fill a similar niche to the insectivores of...

Common punaré

Common punaré

Thrichomys apereoides

The common punaré, *Thrichomys apereoides*, is a species of spiny rat endemic to Brazil.

Common Ringtail Possum

Common Ringtail Possum

Pseudocheirus peregrinus

The Common ringtail possum (*Pseudocheirus peregrinus*) is an Australian marsupial. It lives in a variety of habitats and eats a variety of leaves, flowers, and fruits. This possum also consumes a special type of faeces that is produced during the...

Common rock rat

Common rock rat

Zyzomys argurus

The common rock rat (*Zyzomys argurus*) is a species of rodent in the family Muridae. It is found only in Australia, where it lives in the rocky areas of woodlands, grasslands and low open forests, particularly on the talus, or scree, of cliff...

Common rufous-nosed rat

Common rufous-nosed rat

Oenomys hypoxanthus

The common rufous-nosed rat (*Oenomys hypoxanthus*) is a species of rodent in the family Muridae. It is found in Angola, Burundi, Cameroon, Central African Republic, Republic of the Congo, Democratic Republic of the Congo, Equatorial Guinea,...

Common sheath-tailed bat

Common sheath-tailed bat

Taphozous georgianus

The common sheath-tail bat (*Taphozous georgianus*), is a bat in the family Emballonuridae, occurring in northern Australia.

Common Shrew

Common Shrew

Sorex araneus

The Common shrew (*Sorex araneus*) is among the most frequently found mammals in Northern Europe and Great Britain but doesn't occur in Ireland. It is one of the rare venomous mammals and lives for approximately 14 months.

Common Spotted Cuscus

Common Spotted Cuscus

Spilocuscus maculatus

The common spotted cuscus (*Spilocuscus maculatus*), also known as the white cuscus, is a cuscus, a marsupial that lives in the Cape York region of Australia, New Guinea, and nearby smaller islands.

Common thick-thumbed bat

Common thick-thumbed bat

Glischropus tylopus

The common thick-thumbed bat (*Glischropus tylopus*) is a species of vesper bat found in Brunei, Indonesia, Malaysia, Myanmar, Philippines, and Thailand. It has two subspecies: *G. t. tylopus* *G. t. batjanus*

Common Treeshrew

Common Treeshrew

Tupaia glis

The Common treeshrew (*Tupaia glis*) is a small mammal native to Southeast Asia. It is one of the largest among treeshrews and was first described in 1820 by the French explorers Pierre-Médard Diard and Alfred Duvaucel.

Common tube-nosed fruit bat

Common tube-nosed fruit bat

Nyctimene albiventer

The common tube-nosed fruit bat (*Nyctimene albiventer*) is a species of megabat in the family Pteropodidae. It is found at islands north of Australia, and in Indonesia, Papua New Guinea, the Philippines and the Solomon Islands.

Common Vampire Bat

Common Vampire Bat

Desmodus rotundus

The Common vampire bat (*Desmodus rotundus*) is a small, leaf-nosed bat native to Latin America. It is one of three extant species of vampire bat. The common vampire bat mainly feeds on the blood of livestock and usually approaches its prey at...

Common Vole

Common Vole

Microtus arvalis

The common vole (*Microtus arvalis*) is a European rodent.

Common Wallaroo

Common Wallaroo
Macropus robustus

The common wallaroo (*Osphranter robustus*), also known as the euro, hill wallaroo, or simply wallaroo, is a species of macropod. The word euro is particularly applied to one subspecies (*O. r. erubescens*).The eastern wallaroo is mostly nocturnal...

Common Warthog
Common Warthog
Phacochoerus africanus

The Common warthog (*Phacochoerus africanus*) is a wild member of the pig family (Suidae) found in sub-Saharan Africa. It is the only pig species that has adapted to grazing and savanna habitats.

Common Wombat
Common Wombat
Vombatus ursinus

The Common wombat (*Vombatus ursinus*) is a native Australian species. The early settlers called it a 'badger' due to its excellent burrowing skills, though the closest relative of the wombat is the koala. As a matter of fact, the Common wombat is...

Common yellow-toothed cavy
Common yellow-toothed cavy
Galea musteloides

The common yellow-toothed cavy (*Galea musteloides*) is a species of rodent in the family Caviidae, closely related to the domesticated guinea pig. It is found in Argentina, Bolivia, Chile, and Peru. Its karyotype has $2n = 68$ and $FN = 136$. G....

Comoro rousette
Comoro rousette
Rousettus obliviosus

The Comoro rousette (*Rousettus obliviosus*) is a species of megabat in the family Pteropodidae endemic to the Comoros Islands. Its natural habitats are subtropical or tropical moist lowland forests, caves, plantations, and urban areas. It is...

Complex-toothed flying squirrel
Complex-toothed flying squirrel
Trogopterus xanthipes

The complex-toothed flying squirrel (*Trogopterus xanthipes*) occurs in the southern Chinese provinces Hubei, Hunan, Guizhou, Sichuan, and Yunnan. The common name refers to the teeth, which differ from those of other species of flying squirrels.

Congo gerbil
Congo gerbil
Taterillus congicus

The Congo gerbil or Congo tateril (*Taterillus congicus*) is a species of rodent found in Cameroon, Central African Republic, Chad, Democratic Republic of the Congo, Sudan, and possibly Uganda. Its natural habitat is dry savanna.

Congo golden mole
Congo golden mole
Huetia leucorhina

The Congo golden mole (*Huetia leucorhina*) is a species of mammal in the family Chrysochloridae. It is found in Angola, Cameroon, Central African Republic, Republic of the Congo, and Democratic Republic of the Congo. Its natural habitats are...

Congo rope squirrel
Congo rope squirrel

Funisciurus congicus

The Congo rope squirrel (*Funisciurus congicus*) is a species of rodent in the family *Sciuridae*. It is found in Angola, Democratic Republic of the Congo, and Namibia. Its natural habitats are moist savanna, subtropical or tropical dry shrubland, and...

Congo white-toothed shrew

Congo white-toothed shrew

Crocidura congobelgica

The Congo white-toothed shrew (*Crocidura congobelgica*) is a species of mammal in the family *Soricidae*. It is found in Democratic Republic of the Congo and Uganda. Its natural habitat is subtropical or tropical moist lowland forests. It is...

Conilurus capricornensis

Conilurus capricornensis

Conilurus capricornensis

Conilurus capricornensis commonly known as Capricorn rabbit rat is an extinct species of rabbit rat (*Conilurus*). It was described as a new species in 2010 and is only known by Pleistocene and Holocene dental remains. The species' epithet refers...

Conover's tuco-tuco

Conover's tuco-tuco

Ctenomys conoveri

Conover's tuco-tuco (*Ctenomys conoveri*) is a species of rodent in the family *Ctenomyidae*. It is found in Argentina, Bolivia, and Paraguay.

Convex horseshoe bat

Convex horseshoe bat

Rhinolophus convexus

The convex horseshoe bat (*Rhinolophus convexus*) is a species of bat in the family *Rhinolophidae*. It is found in Malaysia and Laos.

Cook's hocicudo

Cook's hocicudo

Oxymycterus josei

Cook's hocicudo (*Oxymycterus josei*) is a species of rodent in the family *Cricetidae*. It is found only in southern Uruguay, where it lives in wetlands, moist grasslands and scrub. The species is named after American zoologist Joseph "José" A....

Cook's mouse

Cook's mouse

Mus cookii

Cook's mouse (*Mus cookii*) is a species of rodent in the family *Muridae*. It is found in southern and south-eastern Asia from India to Vietnam.

Cooper's mountain squirrel

Cooper's mountain squirrel

Paraxerus cooperi

Cooper's mountain squirrel (*Paraxerus cooperi*) is a species of rodent in the family *Sciuridae* found in Cameroon and Nigeria. Its natural habitat is subtropical or tropical moist montane forests.

Copper woolly bat

Copper woolly bat

Kerivoula cuprosa

The copper woolly bat (*Kerivoula cuprosa*) is a species of vesper bat in the family

Vespertilionidae. It is found in Cameroon, Democratic Republic of the Congo, Ivory Coast, and Guinea. Its natural habitats are subtropical or tropical dry forests,...

Coppery brushtail possum
Coppery brushtail possum
Trichosurus johnstonii

The coppery brushtail possum (*Trichosurus johnstonii*) is a species of marsupial possum in the family Phalangeridae. Coppery brushtails are found within the Atherton Tablelands area of Queensland, in northeastern Australia. These mammals inhabit...

Coppery pipistrelle
Coppery pipistrelle
Arielulus cuprosus

The coppery sprite (*Arielulus cuprosus*), also known as the coppery pipistrelle, is a species of vesper bat found only in Malaysia.

Coppery ringtail possum
Coppery ringtail possum
Pseudochirops cupreus

The coppery ringtail possum (*Pseudochirops cupreus*) is a species of marsupial in the family Pseudocheiridae. It is found in Indonesia and Papua New Guinea.

Coppery titi
Coppery titi
Plecturocebus cupreus

The coppery titi (*Plecturocebus cupreus*) is a species of titi, a type of New World monkey, from South America. It is found in the Amazon of Brazil and Peru, and perhaps northern Bolivia. It was described as *Callicebus cupreus* in 1823.

Coquerel's giant mouse lemur
Coquerel's giant mouse lemur
Mirza coquereli

Coquerel's giant mouse lemur (*Mirza coquereli*), also known as Coquerel's dwarf lemur or the southern giant mouse lemur, is a small nocturnal lemur endemic to Madagascar. This species can be found in parts of the Madagascar dry deciduous forests.

Coquerel's Sifaka
Coquerel's Sifaka
Propithecus coquereli

Coquerel's sifaka (*Propithecus coquereli*) is a diurnal, medium-sized lemur of the sifaka genus *Propithecus*. It is native to northwest Madagascar. Coquerel's sifaka was once considered to be a subspecies of Verreaux's sifaka, but was eventually...

Córdoba vesper mouse
Córdoba vesper mouse
Calomys venustus

The Córdoba vesper mouse or Córdoba laucha (*Calomys venustus*) is a South American rodent species of the family Cricetidae. It is endemic to the area of Córdoba Province, central Argentina, where it is found in the espinal (dry lowland thorn...

Corsac Fox
Corsac Fox
Vulpes corsac

The Corsac foxes (*Vulpes corsac*) are medium-sized foxes found in Asia. They live in open grassy steppes and semideserts and try to stay far away from human habitations. Fossils of Corsac foxes date back to the mid-Pleistocene and show they once...

Corsican hare
Corsican hare
Lepus corsicanus

The Corsican hare (*Lepus corsicanus*), also known as the Apennine hare or Italian hare, is a species of hare found in southern and central Italy and Corsica.

Corsican Red Deer
Corsican Red Deer
Cervus elaphus corsicanus

The Corsican red deer (*Cervus elaphus corsicanus*) is a subspecies of Red deer (*Cervus elaphus*) found in Italy and France. The subspecies gets its name from the island of Corsica from where it was, however, extirpated in the early 1970s.

Corsican Wildcat
Corsican Wildcat
Felis catus

The Corsican wildcat is an isolated feral cat (*Felis catus*) population that used to be considered a subspecies of the African wildcat (*Felis lybica*). Now it is thought to have been introduced to Corsica around the beginning of the first...

Coruro
Coruro
Spalacopus cyanus

The coruro (*Spalacopus cyanus*) is a species of rodent in the family Octodontidae. It is monotypic within the genus *Spalacopus*. The species is endemic to central Chile, where it has been found in a wide variety of habitats, from coastal to...

Cotton Mouse
Cotton Mouse
Peromyscus gossypinus

The cotton mouse (*Peromyscus gossypinus*) is a species of rodent in the family Cricetidae found in the woodlands of the US South.

Cotton-Top Tamarin
Cotton-Top Tamarin
Saguinus oedipus

The Cotton-top tamarin (*Saguinus oedipus*) is a small New World monkey found in South America. Its name comes from the white hair that spans its head and flows down past the neck. When moving through their forest range, these small tamarins run...

Coues's climbing mouse
Coues's climbing mouse
Rhipidomys couesi

Coues's climbing mouse (*Rhipidomys couesi*) is a species of rodent in the family Cricetidae. It is found in Colombia, Ecuador, Peru, Trinidad and Tobago, and Venezuela. It is named in honour of the American zoologist Elliott Coues who studied...

Cougar
Cougar
Puma concolor

The cougar (*Puma concolor*) is a large cat native to the Americas. It is the most widespread of any large wild terrestrial mammal in the Western Hemisphere. Due to its wide range, the cougar has many names, including puma, Mountain lion,...

Country mouse
Country mouse
Pseudomys patrius

The country mouse (*Pseudomys patrius*) also known as the pebble-mound mouse or eastern pebble mound mouse is a species of rodent in the family Muridae. It lives only in Australia, where it is considered rare. It was described by Thomas and...
Cowan's shrew tenrec
Cowan's shrew tenrec
Microgale cowani

Cowan's shrew tenrec (*Microgale cowani*) is a species of mammal in the family Tenrecidae.
Cox's roundleaf bat
Cox's roundleaf bat
Hipposideros coxi

Cox's roundleaf bat or Cox's leaf-nosed bat (*Hipposideros coxi*) is a species of bat in the family Hipposideridae. It is endemic to Borneo. All confirmed records are from Sarawak (Malaysia), but it might also occur in Kalimantan (Indonesia).
Coxing's white-bellied rat
Coxing's white-bellied rat
Niviventer coninga

Coxing's white-bellied rat or spiny Taiwan niviventer (*Niviventer coninga*) is a rodent in the family Muridae. The species was first described by Robert Swinhoe in 1864. It is endemic to Taiwan and occurs in broad-leaf forests and their edges and...
Coyhaique tuco-tuco
Coyhaique tuco-tuco
Ctenomys coyhaiquensis

The Coyhaique tuco-tuco (*Ctenomys coyhaiquensis*) is a species of rodent in the family Ctenomyidae. It is endemic to southern Chile. The name comes from the Chilean province and municipality of Coyhaique.
Coyote
Coyote
Canis latrans

The coyote (*Canis latrans*) is a species of canine native to North America. It fills much of the same ecological niche as the Golden jackal does in Eurasia. The coyote is larger and more predatory and was once referred to as the American jackal by...
Coypu
Coypu
Myocastor coypus

The coypu (*Myocastor coypus*), otherwise known as nutria or River rat, is a large, semi-aquatic rodent. In spite of being a separate species, the animal is sometimes mistaken for a beaver or otter. These rodents are 'a boon and a bane', in the...
Cozumel harvest mouse
Cozumel harvest mouse
Reithrodontomys spectabilis

The Cozumel harvest mouse (*Reithrodontomys spectabilis*) is a species of rodent in the family Cricetidae. It is endemic to the Mexican island of Cozumel off the Yucatán Peninsula. It is nocturnal and semiareboreal, and lives in dense secondary...
Cozumel Island Coati
Cozumel Island Coati
Nasua narica nelsoni

The Cozumel Island coati (*Nasua narica nelsoni*) has been treated as a separate

species, but the vast majority of recent authorities treat it as a subspecies of the White-nosed coati. It is also known as the Cozumel Island coati.

Cozumel Raccoon

Cozumel Raccoon

Procyon pygmaeus

The Cozumel raccoon (*Procyon pygmaeus*), is a critically endangered species of island raccoon endemic on Cozumel Island off the coast of the Yucatan Peninsula, Mexico. It is also called the pygmy raccoon, dwarf raccoon, Cozumel Island raccoon,...

Cozumelan golden bat

Cozumelan golden bat

Mimon cozumelae

The Cozumelan golden bat (*Mimon cozumelae*) is a bat species from Central and northern South America. It is found in Mexico, Guatemala, Belize, Honduras, Nicaragua, Costa Rica, Panama and Colombia. At one time this species was considered to be a...

Crab-Eating Fox

Crab-Eating Fox

Cerdocyon thous

The Crab-eating fox (*Cerdocyon thous*) is a species of medium-sized canid native to the central part of South America. It is not closely related to true foxes. The part of its scientific name comes from the Greek words *kerdo* (meaning fox) and *kyon*...

Crab-Eating Mongoose

Crab-Eating Mongoose

Herpestes urva

The Crab-eating mongoose (*Urva urva*) is a species of mongoose found in Asia. These animals are very skillful hunters and good swimmers. They are not considered threatened and are listed as Least Concern on the IUCN Red List.

Crab-Eating Raccoon

Crab-Eating Raccoon

Procyon cancrivorus

The crab-eating raccoon or South American raccoon (*Procyon cancrivorus*) is a species of raccoon native to marshy and jungle areas of Central and South America (including Trinidad and Tobago). It is found from Costa Rica south through most areas...

Crab-eating rat

Crab-eating rat

Ichthyomys hydrobates

The crab-eating rat (*Ichthyomys hydrobates*) is a species of semiaquatic rodent in the family Cricetidae. It is found in Colombia, Ecuador, and Venezuela. Its natural habitats are rivers and swamps.

Crabeater Seal

Crabeater Seal

Lobodon carcinophaga

Crabeater seals (*Lobodon carcinophaga*) are considered to be the most abundant seals in the world. It is believed by some scientists that there are more of them than all the other seal species combined. Crabeater seals are "true" or "earless"...

Crafty vesper mouse

Crafty vesper mouse

Calomys callidus

The crafty vesper mouse (*Calomys callidus*) is a species of rodent in the family Cricetidae. It is found in Argentina and Paraguay.
Crawford-Cabral's shaggy rat
Crawford-Cabral's shaggy rat
Dasymys cabrali

Crawford-Cabral's shaggy rat or Crawford-Cabral's shaggy marsh rat (*Dasymys cabrali*) is a species of shaggy marsh rat endemic to north-eastern Namibia, near the Okavango River.
Crawford's gray shrew
Crawford's gray shrew
Notiosorex crawfordi

Crawford's gray shrew (*Notiosorex crawfordi*) also known as the desert shrew, is a small shrew found in the southwestern United States and northern Mexico. It is a member of the family Soricidae of the order Eulipotyphla. It was the only known...
Crawshay's Zebra
Crawshay's Zebra
Equus quagga crawshayi

Crawshay's zebra (*Equus quagga crawshayi*) is a subspecies of the Plains zebra.
Creagh's horseshoe bat
Creagh's horseshoe bat
Rhinolophus creaghi

Creagh's horseshoe bat (*Rhinolophus creaghi*) is a species of bat in the family Rhinolophidae. It is found in Indonesia and Malaysia.
Cream-Colored Giant Squirrel
Cream-Colored Giant Squirrel
Ratufa affinis

The Cream-coloured giant squirrel (*Ratufa affinis*) is a large tree squirrel that lives in forests of Southeast Asia. The species is near threatened and vulnerable to habitat degradation, and it has probably been extirpated in Singapore where the...
Creek groove-toothed swamp rat
Creek groove-toothed swamp rat
Pelomys fallax

The creek groove-toothed swamp rat (*Pelomys fallax*) is a species of rodent in the family Muridae. It is found in Angola, Burundi, Democratic Republic of the Congo, Kenya, Malawi, Mozambique, Namibia, Rwanda, Tanzania, Uganda, Zambia, and...
Creeping vole
Creeping vole
Microtus oregoni

The creeping vole (*Microtus oregoni*), sometimes known as the Oregon meadow mouse, is a small rodent in the family Cricetidae. Ranging across the Pacific Northwest of North America, it is found in forests, grasslands, woodlands, and chaparral...
Creighton's slender opossum
Creighton's slender opossum
Marmosops creightoni

Creighton's slender opossum (*Marmosops creightoni*), also known commonly as Voss' slender opossum is a species of South American opossum of the family Didelphidae. It is known only from the valley of the Rio Zongo in La Paz Department, Bolivia,...
Crest-tailed mulgara
Crest-tailed mulgara
Dasycercus cristicauda

The crest-tailed mulgara (*Dasyrctus cristicauda*), is a small to medium-sized Australian carnivorous marsupial and a member of the family Dasyuridae (meaning "hairy tail") which includes quolls, dunnarts, the numbat, Tasmanian devil and extinct...

Crested agouti
Crested agouti
Dasyprocta cristata

The crested agouti (*Dasyprocta cristata*) is a species of rodent in the family Dasyproctidae. It is endemic to Guyana and Suriname. Its taxonomic status is uncertain and it may be synonymous with *Dasyprocta leporina*, leading the IUCN to rate it...

Crested capuchin
Crested capuchin
Sapajus robustus

The crested capuchin or robust tufted capuchin (*Sapajus robustus*) is a species of robust capuchin monkey. It is endemic to Brazil. It was formerly considered a subspecies of the black capuchin but is now considered by some to be a separate species.

Crested mona monkey
Crested mona monkey
Cercopithecus pogonias

The crested mona monkey, also known as the crowned guenon, crowned monkey, golden-bellied guenon, or golden-bellied monkey, (*Cercopithecus pogonias*), is a species of African primate in the family Cercopithecidae found in west central Africa.

Crested Porcupine
Crested Porcupine
Hystrix cristata

The Crested porcupine (*Hystrix cristata*) is a species of rodent native to Italy, North Africa and sub-Saharan Africa. It is one of the largest rodents in the world. Crested porcupines live up to 28 years, the second-longest of any rodent after...

Crested roundleaf bat
Crested roundleaf bat
Hipposideros inexpectatus

The crested roundleaf bat (*Hipposideros inexpectatus*) is a species of bat in the family Hipposideridae. It is endemic to Indonesia.

Crested servaline genet
Crested servaline genet
Genetta cristata

The crested servaline genet (*Genetta cristata*), also known as the crested genet, is a genet species endemic to Nigeria and Cameroon. As the population has declined due to loss of habitat, it is listed as Vulnerable on the IUCN Red List. It was...

Crested-tailed deer mouse
Crested-tailed deer mouse
Habromys lophurus

The crested-tailed deer mouse (*Habromys lophurus*) is a species of rodent in the family Cricetidae found in El Salvador, Guatemala, and Mexico.

Cretan shrew
Cretan shrew
Crocidura zimmermanni

The Cretan shrew (*Crocidura zimmermanni*) is a species of mammal in the family

Soricidae. It is endemic and exclusive to the island of Crete, Greece. Its natural habitat is temperate shrubland, and the animal is threatened by habitat loss. It is...

Cretan wildcat

Cretan wildcat

Felis silvestris cretensis

The Cretan wildcat is a member of the genus *Felis*. Its taxonomic status is unclear at present, as some biologists consider it probably introduced, or a European wildcat (*Felis silvestris silvestris*), or a hybrid between European wildcat and...

Crete spiny mouse

Crete spiny mouse

Acomys minous

The Crete spiny mouse (*Acomys minous*) is a species of mouse endemic to Crete. It is characterized by the coarse, stiff hairs on its back and tail and a notably grayer coloration and more pointed face than other species of spiny mice. Its fur...

Crocidura annamitensis

Crocidura annamitensis

Crocidura annamitensis

Crocidura annamitensis is a small species of shrew from Vietnam, where it is found in the Annamite Range. It is a small shrew- no more than 60 mm in head and body length, and is gray in color. It has a relatively short tail.

Crocidura cranbrookii

Crocidura cranbrookii

Crocidura cranbrookii

Crocidura cranbrookii is a species of shrew from Northern Myanmar.

Crocidura grandiceps

Crocidura grandiceps

Crocidura grandiceps

Crocidura grandiceps is a species of mammal in the family Soricidae. It is found in Benin, Ivory Coast, Ghana, Guinea, Liberia, Nigeria, and Togo. Its natural habitat is subtropical or tropical moist lowland forests. It is threatened by habitat...

Crocidura guy

Crocidura guy

Crocidura guy

Crocidura guy is of species of shrew from Northeastern Vietnam, Viet Bac karst formation.

Crocidura hikmiya

Crocidura hikmiya

Crocidura hikmiya

Crocidura hikmiya (Sinhharaja shrew or Sri Lankan rain forest shrew) is a species of shrew described from the rainforests of Sri Lanka, based on both morphological and molecular data. Its closest sister species is the Sri Lankan long-tailed shrew,...

Crocidura kegoensis

Crocidura kegoensis

Crocidura kegoensis

Crocidura kegoensis, also known as the Ke Go shrew or Ke Go white-toothed shrew, is a species of shrew in the genus *Crocidura* described in 2004. It is smaller than other *Crocidura* species known from Vietnam, brownish-grey in colour with black...

Crocidura phanluongi

Crocidura phanluongi

Crocidura phanluongi

Crocidura phanluongi is a species of shrew in the genus *Crocidura* from southern Vietnam and nearby Cambodia. It is a somewhat small, gray shrew with an ecologically diverse distribution.

Crocidura sapaensis
Crocidura sapaensis
Crocidura sapaensis

Crocidura sapaensis is a species of white-toothed shrew native to northern Vietnam. It was first described in 2013 and its specific name derives from the Sa Pa where it was discovered.

Croslet horseshoe bat
Croslet horseshoe bat
Rhinolophus coelophyllus

The croslet horseshoe bat (*Rhinolophus coelophyllus*) is a species of bat in the family Rhinolophidae. It is found in Laos, Malaysia, Myanmar, and Thailand.

Cross River bushbaby
Cross River bushbaby
Sciurocheirus cameronensis

The Cross River bushbaby, also known as the Cross River squirrel galago, (*Sciurocheirus cameronensis*) is a species of prosimian primate in the family Galagidae which is endemic to a restricted area of West Africa. It is one of four species of...

Cross River gorilla
Cross River gorilla
Gorilla gorilla diehli

The Cross River gorilla (*Gorilla gorilla diehli*) is a Critically Endangered subspecies of the western gorilla (*Gorilla gorilla*). It was named a new species in 1904 by Paul Matschie, a mammalian taxonomist working at the Humboldt University...

Crosse's shrew
Crosse's shrew
Crocidura crossei

Crosse's shrew (*Crocidura crossei*) is a species of mammal in the family Soricidae. It is found in Benin, Cameroon, Ivory Coast, Ghana, Guinea, Liberia, Nigeria, Sierra Leone, and Togo. Its natural habitat is subtropical or tropical moist lowland...

Crowned Lemur
Crowned Lemur
Eulemur coronatus

The crowned lemur (*Eulemur coronatus*) is a lemur that is 31–36 cm (12–14 in) long and weighs 2 kg (4.4 lb). Its tail is about 42–51 cm (17–20 in) long.

Crowned shrew
Crowned shrew
Sorex coronatus

The crowned shrew or Millet's shrew (*Sorex coronatus*) is a species of mammal in the family Soricidae. It is found in Austria, Belgium, France, Germany, Liechtenstein, the Netherlands, Spain, Switzerland, and the British island of Jersey. It is...

Crowned Sifaka
Crowned Sifaka
Propithecus coronatus

The crowned sifaka (*Propithecus coronatus*) is a sifaka endemic to western Madagascar. It is of comparable size to the Golden-crowned sifaka and up to a meter in length, of which 47-57 centimeters are tail. The species is an arboreal vertical...

Crump's mouse

Crump's mouse

Diomys crumpi

Crump's mouse (*Diomys crumpi*) is a species of rodent in the family Muridae. It is the only species in the genus *Diomys*, and is found in northern India (recorded in Bihar and Manipur), southern Nepal (recorded in the Central Terai), and northern...

Cryptic myotis

Cryptic myotis

Myotis crypticus

The cryptic myotis (*Myotis crypticus*) is a European vespertilionid bat. It is a member of the Natterer's bat (*M. nattereri*) species complex, and is also the closest living relative of *M. nattereri*. It is mostly distributed across European...

Csorba's mouse-eared bat

Csorba's mouse-eared bat

Myotis csorbai

The Csorba's mouse-eared bat (*Myotis csorbai*) is a species of vesper bat. It is found only in Nepal.

Ctenomys andersoni

Ctenomys andersoni

Ctenomys andersoni

Ctenomys andersoni, also called Anderson's cujuchi, is a species of tuco-tuco native to Bolivia. Found only in Cerro Itahuaticua, Department of Santa Cruz, at an elevation of around 810 metres (2,700 ft), the species measures 271 millimetres...

Ctenomys yatesi

Ctenomys yatesi

Ctenomys yatesi

Ctenomys yatesi, also called Yates' tuco-tuco, is a species of tuco-tuco native to Bolivia. Only found near Roboré, Department of Santa Cruz, at an elevation of around 550 metres (1,800 ft), the species measures about 220 millimetres (8+3/4 in)...

Cuanza vlel rat

Cuanza vlel rat

Otomys cuanzensis

The Cuanza vlel rat (*Otomys cuanzensis*) is a species of rodent in the family Muridae. It is found only in Angola. Its natural habitats are subtropical or tropical high-altitude grassland and swampland.

Cuban coney

Cuban coney

Geocapromys columbianus

The Cuban coney (*Geocapromys columbianus*) is an extinct species of rodent in the subfamily Capromyinae. It was endemic to Cuba. Its natural habitats were lowlands moist forests, xeric shrublands and rocky areas. Some scientists indicate that this...

Cuban evening bat

Cuban evening bat

Nycticeius cubanus

The Cuban evening bat (*Nycticeius cubanus*) is a species of bat in the vesper bat family, *Vespertilionidae*, that is endemic to western Cuba. It is a small bat, even smaller than congener *Nycticeius humeralis*. It is insectivorous, but otherwise...

Cuban fig-eating bat

Cuban fig-eating bat

Phyllops falcatus

The Cuban fig-eating bat, or white-shouldered bat, (*Phyllops falcatus*) is a species of bat in the family *Phyllostomidae*, found only in the Caribbean. It is the sole extant species in the genus *Phyllops*, although two other species, *P. vetus* and...

Cuban flower bat

Cuban flower bat

Phyllonycteris poeyi

The Cuban flower bat (*Phyllonycteris poeyi*), also called Poey's flower bat, is a species of bat in the family *Phyllostomidae*. It is found on the Caribbean islands of Cuba and Hispaniola (in both Haiti and the Dominican Republic).

Cuban fruit-eating bat

Cuban fruit-eating bat

Brachyphylla nana

The Cuban fruit-eating bat (*Brachyphylla nana*) is a species of bat in the family *Phyllostomidae* found in the Cayman Islands, Cuba, and Hispaniola (both the Dominican Republic and Haiti). It has been extirpated from the Bahamas and Jamaica.

Cuban funnel-eared bat

Cuban funnel-eared bat

Chilonatalus micropus

The Cuban funnel-eared bat (*Chilonatalus micropus*) is a species of bat in the family *Natalidae*. It is one of two species within the genus *Chilonatalus*. It is found in Colombia's Archipelago of San Andrés, Providencia and Santa Catalina,...

Cuban greater funnel-eared bat

Cuban greater funnel-eared bat

Natalus primus

The Cuban greater funnel-eared bat (*Natalus primus*) is a species of funnel-eared bat. It is endemic to a cave in westernmost Cuba. It is a member of the family *Natalidae*.

Cuban Solenodon

Cuban Solenodon

Solenodon cubanus

The Cuban solenodon, or almiquí (*Atopogale cubana*), is a species of eulipotyphlan endemic to Cuba. It is the only species in the genus *Atopogale*. It belongs to the family *Solenodontidae* along with a similar species, the Hispaniolan solenodon...

Cuban yellow bat

Cuban yellow bat

Lasiurus insularis

The Cuban yellow bat (*Dasypterus insularis*) is a species of bat from the family *Vespertilioninae*. It was previously included as a subspecies of the northern yellow bat, a species that has a similar ecology and biology. The species is endemic to...

Culion tree squirrel

Culion tree squirrel

Sundasciurus moellendorffi

The Culion tree squirrel (*Sundasciurus moellendorffi*) is a species of rodent in the family *Sciuridae*. It is endemic to the Philippines. Its natural habitat is

subtropical or tropical dry forests. It is threatened by habitat loss.

Culpeo

Culpeo

Lycalopex culpaeus

The culpeo (*Lycalopex culpaeus*) is a species of South American fox. Regardless of the name, it is not a true fox, but more closely related to wolves and jackals. It is the second-largest native canid on the continent after the Maned wolf.

Cuniculus hernandezi

Cuniculus hernandezi

Cuniculus hernandezi

Cuniculus hernandezi is a name coined for a proposed species of paca described as an endemic to the Central Andes in Colombia, the only place it has been identified as such to date. Individuals may weigh over 14 pounds (6.4 kg), making C....

Curacao myotis

Curacao myotis

Myotis nesopolus

The *Curacao myotis* (*Myotis nesopolus*) is a species of vesper bat (family *Vespertilionidae*). It is found in Bonaire (Netherlands), Colombia, Curaçao and Venezuela.

Curry's bat

Curry's bat

Glauconycteris curryae

Curry's bat (*Glauconycteris curryae*) is a species of vesper bat in the family *Vespertilionidae*. It is found in Cameroon and Democratic Republic of the Congo. Its natural habitats are subtropical or tropical moist lowland forests and subtropical or...

Cursor grass mouse

Cursor grass mouse

Akodon cursor

The cursor grass mouse or cursorial akodont (*Akodon cursor*), is a sigmodontine rodent from South America.

Cuscomys oblativus

Cuscomys oblativus

Cuscomys oblativus

The Machu Picchu arboreal chinchilla rat (*Cuscomys oblativus*) is a large species of South American chinchilla rats, known from skeletal remains found by members of the Peruvian Expedition of 1912. The animals were buried alongside people in...

Cushioned gerbil

Cushioned gerbil

Gerbillus pulvinatus

The cushioned gerbil (*Gerbillus pulvinatus*) is distributed mainly in Somalia, Ethiopia, Djibouti and Kenya.

Cutch rat

Cutch rat

Cremnomys cutchicus

The Cutch rat (*Cremnomys cutchicus*) is a species of rodent in the family *Muridae*. It is found in India, where it is widely distributed in the states of Andhra Pradesh, Bihar, Gujarat, Jharkhand, Karnataka, and Rajasthan.

Cuvier's Beaked Whale

Cuvier's Beaked Whale

Ziphius cavirostris

The Cuvier's beaked whale (*Ziphius cavirostris*) is the most widely distributed of all beaked whales in the family Ziphiidae. It is smaller than most baleen whales yet large among beaked whales. Cuvier's beaked whale is pelagic and occurs in...

Cuvier's Gazelle

Cuvier's Gazelle

Gazella cuvieri

Cuvier's gazelle (*Gazella cuvieri*) is a species of gazelle native to Algeria, Morocco, Western Sahara, and Tunisia. It is also known as the edmi. It is one of the darkest gazelle species, possibly an adaptation to its partial woodland habitat....

Cuvier's spiny rat

Cuvier's spiny rat

Proechimys cuvieri

Cuvier's spiny-rat, *Proechimys cuvieri*, is a spiny rat species from South America. It is found in Brazil, French Guiana, Guyana, Peru and Suriname.

Cyclops roundleaf bat

Cyclops roundleaf bat

Doryrhina cyclops

The cyclops roundleaf bat or cyclops leaf-nosed bat (*Doryrhina cyclops*) is a species of bat in the family Hipposideridae found in the forests of equatorial Africa.

Cynomops milleri

Cynomops milleri

Cynomops milleri

Cynomops milleri is a species of bat that is native to South America. It was previously considered a subspecies of the Para dog-faced bat. It is considered a small- to medium-sized member of its genus. It is classified as least concern by the...

Cypriot mouse

Cypriot mouse

Mus cypriacus

The Cypriot mouse (*Mus cypriacus*) is a species of mouse endemic to Cyprus. Its primary habitat seems to be the vineyards and fields of the Troödos Mountains region. The mouse was recognized as a new species in 2004 by Thomas Cucchi, a research...

Cyprus spiny mouse

Cyprus spiny mouse

Acomys nesiotus

The Cyprus spiny mouse (*Acomys nesiotus*) is a little-known rodent endemic to Cyprus. These nocturnal animals are generally found in arid areas. After the last reliable record in 1980 no considerable effort has been made until 2007 when four...

Cyrenaica shrew

Cyrenaica shrew

Crocidura aleksandrisi

The Cyrenaica shrew or Alexander's shrew (*Crocidura aleksandrisi*) is a species of white-toothed shrew in the family Soricidae which is endemic to Libya. The Cyrenaica shrew is a small shrew which is very similar to the widespread lesser...

D'Albertis' ringtail possum

D'Albertis' ringtail possum

Pseudochirops albertisii

D'Albertis' ringtail possum (*Pseudocheirops albertisii*) is a species of marsupial in the family Pseudocheiridae. It is found in Indonesia and Papua New Guinea. Its natural habitat is subtropical or tropical dry forests.

D'Anchieta's fruit bat

D'Anchieta's fruit bat

Plerotes anchietae

D'Anchieta's fruit bat or D'Anchieta's epauletted bat (*Plerotes anchietae*) is a species of megabat in the family Pteropodidae. It is the only species in the genus *Plerotes*. It is found in Angola, Democratic Republic of the Congo, Malawi, and...

D'Entrecasteaux Archipelago pogonomys

D'Entrecasteaux Archipelago pogonomys

Pogonomys fergussoniensis

The D'Entrecasteaux Archipelago pogonomys (*Pogonomys fergussoniensis*), also known as the D'Entrecasteaux Archipelago tree mouse, is a species of prehensile-tailed rat from the family Muridae that is endemic to Papua New Guinea. It was once...

D'Orbigny's tuco-tuco

D'Orbigny's tuco-tuco

Ctenomys dorbignyi

D'Orbigny's tuco-tuco (*Ctenomys dorbignyi*) is a species of rodent in the family Ctenomyidae, named after French naturalist Alcide d'Orbigny. It is found in northeast Argentina. Its karyotype has $2n = 70$, $FN = 84-88$, which is cytogenetically...

Daghestan pine vole

Daghestan pine vole

Microtus daghestanicus

The Daghestan pine vole (*Microtus daghestanicus*) is a species of rodent in the family Cricetidae. It is found in Russia, Georgia, Armenia, and Azerbaijan.

Dahl's jird

Dahl's jird

Meriones dahli

Dahl's jird (*Meriones dahli*) is a species of rodent in the family Muridae. It is found only in Armenia and the neighboring Agri province of Turkey.

Dainty fat mouse

Dainty fat mouse

Steatomys cuppedius

The dainty fat mouse (*Steatomys cuppedius*) is a species of rodent in the family Nesomyidae. It is found in Benin, Burkina Faso, Mali, Niger, Nigeria, and Senegal. Its natural habitat is subtropical or tropical dry shrubland.

Dall's Porpoise

Dall's Porpoise

Phocoenoides dalli

Dall's porpoise (*Phocoenoides dalli*) is a species of porpoise endemic to the North Pacific. It is the largest of porpoises and the only member of the genus *Phocoenoides*. The species is named after American naturalist W. H. Dall.

Dalquest's pocket mouse

Dalquest's pocket mouse

Chaetodipus dalquesti

Dalquest's pocket mouse (*Chaetodipus dalquesti*) is a species of rodent in the family Heteromyidae, sometimes viewed as a subspecies of *Chaetodipus ammophilus*. It is endemic to Mexico. The pocket mouse is named after Walter W. Dalquest...

Dalton's mouse
Dalton's mouse
Praomys daltoni

Dalton's mouse or Dalton's praomys (*Praomys daltoni*) is a species of rodent in the family Muridae. It is found in Benin, Burkina Faso, Cameroon, Central African Republic, Chad, Ivory Coast, Gambia, Ghana, Guinea, Guinea-Bissau, Liberia, Mali,...

Dama Gazelle
Dama Gazelle
Nanger dama

The dama gazelle (*Nanger dama*) also known as the addra gazelle or mhorra gazelle, is a species of gazelle. It lives in Africa, in the Sahara desert and the Sahel. A critically endangered species, it has disappeared from most of its former range...

Damara horseshoe bat
Damara horseshoe bat
Rhinolophus damarensis

The *Damara horseshoe bat*, *Rhinolophus damarensis*, is a species of bat found in Africa.

Damara woolly bat
Damara woolly bat
Kerivoula argentata

The *Damara woolly bat* (*Kerivoula argentata*) is a nocturnal insectivorous species of vesper bat in the family Vespertilionidae found in Africa. This species typically has reddish brown fur on its back and white fur on its abdomen. Its natural...

Damaraland Mole-Rat
Damaraland Mole-Rat
Fukomys damarensis

The *Damaraland mole-rat* (*Fukomys damarensis*), *Damara mole rat* or *Damaraland blesmole*, is a burrowing rodent found in southern Africa. Along with the smaller, less hairy, naked mole rat, it is a species of eusocial mammal.

Danfoss's mouse lemur
Danfoss's mouse lemur
Microcebus danfossi

Danfoss's mouse lemur (*Microcebus danfossi*), or the *Ambarijeby mouse lemur*, is a species of mouse lemur endemic to Madagascar. It lives in western deciduous forest within a limited range between the Sofia River and the Maevarano River. It is a...

Daniel's tufted-tailed rat
Daniel's tufted-tailed rat
Eliurus danieli

Daniel's tufted-tailed rat (*Eliurus danieli*) is a species of rodent in the family Nesomyidae. It was discovered in 2003 in the Parc National de l'Isalo in south-central Madagascar. It is named for Daniel Rakotonindravony, professor of animal...

Daovantien
Daovantien
Tonkinomys daovantieni

Daovantien
Daphne's Oldfield mouse
Daphne's Oldfield mouse
Thomasomys daphne

Daphne's Oldfield mouse (*Thomasomys daphne*) is a species of rodent in the family Cricetidae. It is found in Bolivia and Peru.
Dar es Salaam pipistrelle
Dar es Salaam pipistrelle
Pipistrellus permixtus

The Dar es Salaam pipistrelle (*Pipistrellus permixtus*) is a species of vesper bat. It is found only in Tanzania. Its natural habitat is subtropical or tropical dry forests.
Daraina sportive lemur
Daraina sportive lemur
Lepilemur milanoii

The *Daraina sportive lemur* (*Lepilemur milanoii*) is a sportive lemur endemic to Madagascar. It is a relatively small sportive lemur with a total length of about 49 to 56 cm (19 to 22 in), of which 24–27 cm (9.4–10.6 in) are tail. *L. milanoii* is...
Darfur gerbil
Darfur gerbil
Gerbillus muriculus

The Darfur gerbil (*Gerbillus muriculus*) is distributed mainly in western Sudan.
Darien harvest mouse
Darien harvest mouse
Reithrodontomys darienensis

The Darien harvest mouse (*Reithrodontomys darienensis*) is a species of rodent in the family Cricetidae. It is found only in Panama.
Darien pocket gopher
Darien pocket gopher
Orthogeomys dariensis

The Darien pocket gopher (*Heterogeomys dariensis*) is a species of rodent in the family Geomyidae. It occurs in Panama and Colombia. Some authors classify it in the genus *Orthogeomys*, but recent research has allowed this and its related...
Darién small-eared shrew
Darién small-eared shrew
Cryptotis merus

The Darién small-eared shrew (*Cryptotis merus*) is a species of mammal in the family Soricidae. It is known only from montane regions along the border between Colombia and Darién Province, Panama, where it has been found in rainforest at...
Dark bolo mouse
Dark bolo mouse
Necromys obscurus

The dark bolo mouse or dark-furred akodont (*Necromys obscurus*) is a species of rodent in the family Cricetidae. There are two subspecies; one (ssp. *scagliarum*) is found in eastern and central parts of Buenos Aires Province, Argentina, and the...
Dark fruit-eating bat
Dark fruit-eating bat
Artibeus obscurus

The dark fruit-eating bat (*Artibeus obscurus*), is a bat species from South America.
Dark kangaroo mouse
Dark kangaroo mouse
Microdipodops megacephalus

The dark kangaroo mouse (*Microdipodops megacephalus*) is a species of rodent in the family Heteromyidae. It is found in California, Idaho, Nevada, Oregon and Utah in the United States.

Dark long-tongued bat

Dark long-tongued bat

Lichonycteris obscura

The dark long-tongued bat (*Lichonycteris obscura*) is a species of bat from South and Central America. It was formerly considered monotypic within the genus *Lichonycteris*, but is now recognized as one of two species in that genus, along with the...

Dark sheath-tailed bat

Dark sheath-tailed bat

Mosia nigrescens

The dark sheath-tailed bat (*Mosia nigrescens*) is a species of sac-winged bat in the family Emballonuridae. It is the only species in the genus *Mosia*. It is found in Indonesia, Papua New Guinea, and Solomon Islands.

Dark spiny tree-rat

Dark spiny tree-rat

Echimys saturnus

The dark spiny tree-rat (*Echimys saturnus*) is a species of rodent in the family Echimyidae. It is a rarely encountered and nocturnal species, found in eastern Ecuador and central Peru. As compared to *Echimys chrysurus*, the main diagnostic...

Dark-brown serotine

Dark-brown serotine

Neoromicia brunnea

The dark-brown serotine (*Pseudoromicia brunnea*) is a species of vesper bat found in Central and West Africa.

Dark-caped Atlantic spiny rat

Dark-caped Atlantic spiny rat

Trinomys mirapitanga

The dark-caped Atlantic spiny rat or Mirapitanga spiny rat (*Trinomys mirapitanga*), is a spiny rat species from South America. It is found in Brazil. Also known as the Pau Brazil Spiny Rat, it is known only from specimens found at two locations on...

Dark-footed mouse shrew

Dark-footed mouse shrew

Myosorex cafer

The dark-footed mouse shrew (*Myosorex cafer*) is a species of mammal in the family Soricidae found in Mozambique, South Africa, Eswatini, and Zimbabwe. Its natural habitat is subtropical or tropical moist montane forests. It is threatened by...

Dark-nosed small-footed myotis

Dark-nosed small-footed myotis

Myotis melanorhinus

The dark-nosed small-footed myotis (*Myotis melanorhinus*) is a species of mouse-eared bat in the family Vespertilionidae, described in 1890, and indigenous to Canada, Mexico, and the United States. While the International Union for Conservation...

Dark-tailed tree rat

Dark-tailed tree rat

Niviventer cremoriventer

The dark-tailed tree rat or Sundaic arboreal niviventer (*Niviventer cremoriventer*) is a species of rodent in the family Muridae. It is found on the Malay Peninsula

(Thailand, Malaysia), including some offshore islands (Myanmar: Mergui...
Dark-winged lesser house bat
Dark-winged lesser house bat
Scotoecus hirundo

The dark-winged lesser house bat (*Scotoecus hirundo*) is a species of vesper bat. It can be found in Angola, Benin, Cameroon, Central African Republic, Chad, Democratic Republic of the Congo, Republic of Côte d'Ivoire, Ethiopia, Gambia, Ghana, ...
Darling's horseshoe bat
Darling's horseshoe bat
Rhinolophus darlingi

Darling's horseshoe bat (*Rhinolophus darlingi*) is a species of bat in the family Rhinolophidae found in Africa. Its natural habitats are dry savanna, caves, and subterranean habitats (other than caves).
Darwin's Fox
Darwin's Fox
Lycalopex fulvipes

Darwin's fox or Darwin's zorro (*Lycalopex fulvipes*) is an endangered canid from the genus *Lycalopex*. It is also known as the zorro chilote or zorro de Darwin in Spanish and lives on Nahuelbuta National Park, the Cordillera de Oncol, Cordillera...
Darwin's leaf-eared mouse
Darwin's leaf-eared mouse
Phyllotis darwini

Darwin's leaf-eared mouse (*Phyllotis darwini*) is a species of rodent in the family Cricetidae. It has terrestrial habits and is endemic to coastal central and northern Chile. It is also found in the Atacama Desert. Members of the species have...
Dassie rat
Dassie rat
Petromus typicus

The dassie rat (*Petromus typicus*) is an African rodent found among rocky outcroppings. It is the only living member of its genus, *Petromus*, and family, *Petromuridae*. The name "dassie" means "hyrax" in Afrikaans, and the two animals are found in...
Daubenton's Bat
Daubenton's Bat
Myotis daubentonii

Daubenton's bat (*Myotis daubentonii*) is a Eurasian bat that is considered to be increasing its numbers in many areas. It was first described in 1817 by Heinrich Kuhl, who named it in honor of French naturalist Louis-Jean-Marie Daubenton.
Daubenton's free-tailed bat
Daubenton's free-tailed bat
Myotis daubentonii

The Daubenton's free-tailed bat or Daubenton's winged-mouse bat (*Myotis daubentonii*) is a species of bat in the family Molossidae. It is found in Central African Republic, Democratic Republic of the Congo, Ivory Coast, and Senegal. Its...
Daurian ground squirrel
Daurian ground squirrel
Spermophilus dauricus

The Daurian ground squirrel (*Spermophilus dauricus*) is a species of rodent in the

family Sciuridae. It is found in China, Mongolia, and Russia.

Daurian hedgehog

Daurian hedgehog

Mesechinus dauuricus

The Daurian hedgehog (*Mesechinus dauuricus*) is a solitary small hedgehog. It is listed in the Red book of Russian Federation as a protected species with an unclear status, generally considered to be endangered, although the IUCN lists it as...

Daurian pika

Daurian pika

Ochotona dauurica

The Daurian pika (*Ochotona dauurica*) is a small relative of rabbits and hares in the order Lagomorpha. It is well known for its "barking" alarm call, and for its peculiar habit of making hay to help survive the winter. There are 4 recognized...

Davao squirrel

Davao squirrel

Sundasciurus davensis

The Davao squirrel (*Sundasciurus davensis*) is a species of rodent in the family Sciuridae. It is endemic to the Philippines.

David's echymipera

David's echymipera

Echymipera davidi

David's echymipera (*Echymipera davidi*), or David's spiny bandicoot, is a species of marsupial in the family Peramelidae. It is found on the island of Kiriwina, in the Trobriand Islands of Papua New Guinea, and may be present on other nearby...

David's myotis

David's myotis

Myotis davidii

David's myotis (*Myotis davidii*) is a species of vesper bat. It is endemic to China where it is found in the provinces of Hebei, Hubei, Inner Mongolia, Shaanxi, Shanxi and Hong Kong. It was previously considered to be a subspecies of the...

Davies's big-eared bat

Davies's big-eared bat

Glyphonycteris daviesi

Davies's big-eared bat or the graybeard bat (*Glyphonycteris daviesi*) is a species of bat in the family Phyllostomidae. It is named after James (Jim) Noel Davies (B. 1936) who discovered it whilst on an expedition in British Guiana, South America...

Davis's round-eared bat

Davis's round-eared bat

Lophostoma evotis

Davis's round-eared bat (*Lophostoma evotis*) is a species of Central American bat in the family Phyllostomidae. It is found in Belize, Guatemala, Honduras, and southeastern Mexico.

Davy's naked-backed bat

Davy's naked-backed bat

Pteronotus davyi

Davy's (lesser) naked-backed bat (species *Pteronotus davyi*) is a small, insect-eating, cave-dwelling bat of the Order Chiroptera and Family Mormoopidae. It is found throughout South and Central America, including Trinidad, but not Tobago,...

Day's grass mouse

Day's grass mouse

Akodon dayi

Day's grass mouse (*Akodon dayi*) is a species of rodent in the family Cricetidae. It is found only in Bolivia.

Day's shrew

Day's shrew

Suncus dayi

Day's shrew (*Suncus dayi*) is a species of mammal in the family Soricidae. It is endemic to India. Its natural habitat is subtropical or tropical dry forests. It is threatened by habitat loss.

Dayak Fruit Bat

Dayak Fruit Bat

Dyacopterus spadiceus

The Dayak fruit bat (*Dyacopterus spadiceus*) is a relatively rare frugivorous megabat species found only in southeast Asia. Few specimens of any of the three species exist, due not only to their rarity but also because they rarely enter the...

Dayak roundleaf bat

Dayak roundleaf bat

Hipposideros dyacorum

The Dayak roundleaf bat (*Hipposideros dyacorum*), also known as the least roundleaf bat, is a species of bat in the family Hipposideridae. It is endemic to Indonesia and Malaysia.

De Brazza's Monkey

De Brazza's Monkey

Cercopithecus neglectus

De Brazza's monkey (*Cercopithecus neglectus*) is an Old World monkey endemic to the riverine and swamp forests of central Africa. The largest species in the guenon family, it is one of the most widespread arboreal African primates. Aside from...

De Graaff's soft-furred mouse

De Graaff's soft-furred mouse

Praomys degraaffi

De Graaff's soft-furred mouse (*Praomys degraaffi*) is a species of rodent in the family Muridae. It is found in Burundi, Rwanda, and Uganda. Its natural habitat is subtropical or tropical moist montane forests. It is threatened by habitat loss.

De Vis's woolly rat

De Vis's woolly rat

Mallomys aroaensis

De Vis's woolly rat (*Mallomys aroaensis*) is a species of rodent in the family Muridae. It is found only in Papua New Guinea.

De Vivo's disk-winged bat

De Vivo's disk-winged bat

Thyroptera devivoi

De Vivo's disk-winged bat (*Thyroptera devivoi*) is a species of disc-winged bat found in South America. It was described as a new species in 2006.

De Winton's golden mole

De Winton's golden mole

Cryptochloris wintoni

De Winton's golden mole (*Cryptochloris wintoni*) is a species of mammal in the family Chrysochloridae. It is endemic to South Africa. Its natural habitats are subtropical dry shrubland, Mediterranean-type shrubby vegetation, and sandy shores. It...

De Winton's long-eared bat
De Winton's long-eared bat
Laephotis wintoni

De Winton's long-eared bat (*Laephotis wintoni*) is a species of vesper bat in the family Vespertilionidae. It can be found in the following countries: Ethiopia, Kenya, Lesotho, South Africa, and Tanzania. It is found in these habitats: dry savanna, ...

De Winton's shrew
De Winton's shrew
Chodsigoa hypsibius

The De Winton's shrew (*Chodsigoa hypsibius*) is a species of mammal in the family Soricidae. It is endemic to China.

Decken's horseshoe bat
Decken's horseshoe bat
Rhinolophus deckenii

Decken's horseshoe bat (*Rhinolophus deckenii*) is a species of horseshoe bat. It is found in Kenya and Tanzania. Its natural habitats are subtropical or tropical moist lowland forest, subtropical or tropical moist montane forest, moist savanna, ...

Definitive leaf-eared mouse
Definitive leaf-eared mouse
Phyllotis definitus

The definitive leaf-eared mouse (*Phyllotis definitus*) is a species of rodent in the family Cricetidae. It is found only in west central Peru, in rocky and shrub-covered areas at elevations between 2600 and 3000 m.

Defua rat
Defua rat
Dephomys defua

The Defua rat (*Dephomys defua*) is a species of rodent in the family Muridae. It is found in Ivory Coast, Ghana, Guinea, Liberia, and Sierra Leone. Its natural habitats are subtropical or tropical moist lowland forests, subtropical or tropical ...

Degu
Degu
Octodon degus

The degu (*Octodon degus*) is a small rodent found only in Chile. The closest relatives of these highly social and curious animals are chinchilla and guinea pig.

Dekeyser's nectar bat
Dekeyser's nectar bat
Lonchophylla dekeyseri

Dekeyser's nectar bat (*Lonchophylla dekeyseri*) is a bat species from South America. It is found in Brazil and Bolivia.

Delacour's Langur
Delacour's Langur
Trachypithecus delacouri

Delacour's langur (*Trachypithecus delacouri*) is a critically endangered species of Old World monkeys. It is named for French-American ornithologist Jean Théodore Delacour.

Delacour's marmoset rat
Delacour's marmoset rat
Hapalomys delacouri

Delacour's marmoset rat (*Hapalomys delacouri*), also known as the lesser marmoset

rat, is an arboreal species of rodent in the family Muridae. It is found in China, Laos, and Vietnam. Its natural habitat is montane subtropical or tropical dry...

Delany's mouse

Delany's mouse

Delanymys brooksi

Delany's mouse or Delany's swamp mouse (*Delanymys brooksi*) is a species of rodent in the family Nesomyidae. It is the only species in the genus *Delanymys* and the only extant member of subfamily Delanymyinae, which also contains the fossil genus...

Delectable soft-furred mouse

Delectable soft-furred mouse

Praomys delectorum

The delectable soft-furred mouse or East African praomys (*Montemys delectorum*) is a species of rodent in the family Muridae. It is the only member of the genus *Montemys*; it was formerly classified in the genus *Praomys*. It is found in Kenya,...

Delicate deer mouse

Delicate deer mouse

Habromys delicatulus

The delicate deer mouse (*Habromys delicatulus*) is a species of rodent in the family Cricetidae. It was first described by Michael Carlton, Oscar Sanchez and Guillermina Urbano-Vidales after being discovered in a patch of cloud forest on the...

Delicate mouse

Delicate mouse

Mus tenellus

The delicate mouse (*Mus tenellus*) is a species of rodent in the family Muridae. It is found in Ethiopia, Kenya, Somalia, Sudan, Tanzania, and possibly Eritrea. Its natural habitat is dry savanna.

Delicate salt flat mouse

Delicate salt flat mouse

Salinomys delicatus

The delicate salt flat mouse, *Salinomys delicatus*, (in Spanish, *el ratón delicado de los salares*) is a sigmodontine rodent species in the family Cricetidae from South America. It is the only species in the genus *Salinomys*. Its habitat is...

Delicate slender opossum

Delicate slender opossum

Marmosops parvidens

The delicate slender opossum (*Marmosops parvidens*) is a small pouchless marsupial of the family Didelphidae that occurs in French Guiana, Guyana, Suriname, and adjacent Venezuela and Brazil. *Marmosops pinheiroi*, *Marmosops bishopi* and

Marmosops...

Delicate vesper mouse

Delicate vesper mouse

Calomys tener

The delicate vesper mouse (*Calomys tener*) is a South American rodent species of the family Cricetidae. It is found in Argentina, Bolivia and Brazil. Its karyotype has $2n = 66$ and $FN = 66$.

Delmarva Fox Squirrel

Delmarva Fox Squirrel

Sciurus niger cinereus

The Delmarva fox squirrel (*Sciurus niger cinereus*) is a subspecies of the Eastern

fox squirrel from the eastern United States. The Delmarva fox squirrel was formerly endangered but was removed from the Endangered Species List effective December...
Deltaic four-eyed opossum
Deltaic four-eyed opossum
Philander deltae

The deltaic four-eyed opossum or delta opossum (*Philander deltae*) is a species of opossum found in the Orinoco River delta region of Venezuela, South America, first described in 2006. It inhabits perennially flooded swamp forest or seasonally...
Demonic tube-nosed fruit bat
Demonic tube-nosed fruit bat
Nyctimene masalai

The demonic tube-nosed fruit bat (*Nyctimene masalai*) is a species of bat in the family Pteropodidae. It is endemic to Papua New Guinea. The holotype specimen was collected in 1979 on New Ireland, in the Bismarck Archipelago. It was described as...
Denebola brachycephala
Denebola brachycephala
Denebola brachycephala

Denebola is an extinct genus of whale belonging to the family Monodontidae. It contains a single known species, *Denebola brachycephala*. It is the earliest known ancestor of the beluga, *Delphinapterus leucas*, and dates from the Upper Miocene...
Dent's horseshoe bat
Dent's horseshoe bat
Rhinolophus denti

Dent's horseshoe bat (*Rhinolophus denti*) is a species of bat in the family Rhinolophidae. It is found in Angola, Botswana, Ghana, Guinea, Guinea-Bissau, Namibia, South Africa, and Zimbabwe. The bat's natural habitats are dry savannah country and...
Dent's mona monkey
Dent's mona monkey
Cercopithecus denti

Dent's mona monkey (*Cercopithecus denti*) is an Old World monkey in the family Cercopithecidae found in the Democratic Republic of the Congo, the Congo, Rwanda, western Uganda, and the Central African Republic. It was previously classified as a...
Dent's shrew
Dent's shrew
Crocidura denti

Dent's shrew (*Crocidura denti*) is a species of mammal in the family Soricidae. It is found in Cameroon, Central African Republic, Republic of the Congo, Democratic Republic of the Congo, Equatorial Guinea, Gabon, Guinea, Liberia, Nigeria, Sierra...
Dent's vlei rat
Dent's vlei rat
Otomys denti

Dent's vlei rat (*Otomys denti*) is a species of rodent in the family Muridae. It is found in Burundi, Democratic Republic of the Congo, Malawi, Tanzania, Uganda, and Zambia. Its natural habitats are subtropical or tropical moist montane forests,...
Deppe's squirrel
Deppe's squirrel
Sciurus deppei

Deppe's squirrel (*Sciurus deppei*) is a species of tree squirrel in the genus *Sciurus* native to Belize, Costa Rica, El Salvador, Guatemala, Honduras, Mexico, and Nicaragua.

Deraniyagala's beaked whale

Deraniyagala's beaked whale

Mesoplodon hotaula

Deraniyagala's beaked whale (*Mesoplodon hotaula*) is a species of mesoplodont whale.

Derby's woolly opossum

Derby's woolly opossum

Caluromys derbianus

Derby's woolly opossum (*Caluromys derbianus*), or the Central American woolly opossum, is an opossum found in deciduous and moist evergreen forests of Central America, from southern Mexico to western Ecuador and Colombia. It was first described...

Dermanura rava

Dermanura rava

Dermanura rava

Dermanura rava is a species of leaf-nosed bat found in Central and South America.

Deroo's mouse

Deroo's mouse

Praomys derooi

Deroo's mouse or Deroo's praomys (*Praomys derooi*) is a species of rodent in the family Muridae. It is found in Benin, Ghana, Nigeria, and Togo. Its natural habitats are dry savanna and urban areas.

Desert bandicoot

Desert bandicoot

Perameles eremiana

The desert bandicoot (*Perameles eremiana*) is an extinct bandicoot of the arid country in the centre of Australia.

Desert bighorn sheep

Desert bighorn sheep

Ovis canadensis nelsoni

The desert bighorn sheep (*Ovis canadensis nelsoni*) is a subspecies of bighorn sheep (*Ovis canadensis*) that is native to the deserts of the United States' intermountain west and southwestern regions, as well as northwestern Mexico. The Bureau of...

Desert Cottontail

Desert Cottontail

Sylvilagus audubonii

The desert cottontail (*Sylvilagus audubonii*), also known as Audubon's cottontail, is a New World cottontail rabbit, and a member of the family Leporidae. Unlike the European rabbit, they do not form social burrow systems, but compared with some...

Desert dormouse

Desert dormouse

Selevinia betpakdalaensis

The desert dormouse (*Selevinia betpakdalaensis*) is a species of rodent in the dormouse family, Gliridae. This species was formerly placed in its own family, Seleviniidae, but it is now considered to be a dormouse, monotypic within the genus...

Desert hare

Desert hare
Lepus tibetanus

The desert hare (*Lepus tibetanus*) is a species of hare found in Central Asia, Northwest China, and the western Indian subcontinent. Little is known about this species except that it inhabits grassland and scrub areas of desert and semi-desert....

Desert Hedgehog
Desert Hedgehog
Paraechinus aethiopicus

The desert hedgehog (*Paraechinus aethiopicus*) is a species of mammal in the family Erinaceidae.

Desert Kangaroo Rat
Desert Kangaroo Rat
Dipodomys deserti

The Desert kangaroo rat (*Dipodomys deserti*) is an adorable rodent with noticeably large hind legs, helping it escape from predators. The Desert kangaroo rat resembles a tiny kangaroo and is the size of a mouse. The species is so called due to its...

Desert long-eared bat
Desert long-eared bat
Otonycteris hemprichii

The desert long-eared bat (*Otonycteris hemprichii*) is a species of vesper bat found in North Africa and the Middle East.

Desert mouse
Desert mouse
Pseudomys desertor

The desert mouse (*Pseudomys desertor*), also known as the brown desert mouse, is a species of rodent in the family Muridae. It is endemic to Australia. The first desert mouse specimen was collected by Australian zoologist Gerard Krefft on the...

Desert musk shrew
Desert musk shrew
Crocidura smithii

The desert musk shrew (*Crocidura smithii*) is a species of mammal in the family Soricidae. It is found in Ethiopia, Senegal, and possibly Somalia. Its natural habitat is dry savanna.

Desert pipistrelle
Desert pipistrelle
Hypsugo ariel

The desert pipistrelle (*Hypsugo ariel*) is a species of vesper bat in the genus *Hypsugo*. It is found in Egypt, Israel, Jordan, Oman, Saudi Arabia, Sudan, and Yemen. Some of the Arabian populations were previously regarded as a separate species,...

Desert pocket gopher
Desert pocket gopher
Geomys arenarius

The desert pocket gopher (*Geomys arenarius*) is a species of rodent in the family Geomyidae. It is found in the state of Chihuahua in Mexico and in Texas and New Mexico in the United States.

Desert Pocket Mouse
Desert Pocket Mouse
Chaetodipus penicillatus

The Desert pocket mouse (*Chaetodipus penicillatus*) is a North American species of heteromyid rodent. True to its common name, the Desert pocket mouse prefers sandy, sparsely vegetated desert environments.

Desert pygmy mouse

Desert pygmy mouse

Mus indutus

The desert pygmy mouse (*Mus indutus*) is a species of rodent in the family Muridae. It is found in Angola, Botswana, Namibia, South Africa, Zambia, and Zimbabwe. Its natural habitats are dry savanna and subtropical or tropical high-altitude grassland.

Desert red bat

Desert red bat

Lasiurus blossevillii

The western red bat or desert red bat (*Lasiurus frantzii*) is a species of microbat in the family Vespertilionidae. It is found in western North America and Central America.

Desert Warthog

Desert Warthog

Phacochoerus aethiopicus

The desert warthog (*Phacochoerus aethiopicus*) is a species of even-toed ungulate in the pig family (Suidae), found in northern Kenya and Somalia, and possibly Djibouti, Eritrea, and Ethiopia. This is the range of the extant subspecies, commonly...

Desert Woodrat

Desert Woodrat

Neotoma lepida

The desert woodrat (*Neotoma lepida*) is a species of pack rat native to desert regions of western North America.

Desert yellow bat

Desert yellow bat

Scotoecus pallidus

The desert yellow bat (*Scotoecus pallidus*) is a species of vesper bat. It is found in India, Pakistan, and Bangladesh. Its natural habitats are subtropical or tropical dry forests, subtropical or tropical dry shrubland, rural gardens, and urban...

Desmarest's Hutia

Desmarest's Hutia

Capromys pilorides

The Desmarest's hutia (*Capromys pilorides*) is a species of rodent endemic to Cuba. It is the largest of the extant species of hutia (although the extinct Giant hutias were far larger).

Desmarest's spiny pocket mouse

Desmarest's spiny pocket mouse

Heteromys desmarestianus

Desmarest's spiny pocket mouse (*Heteromys desmarestianus*) is a species of rodent in the family Heteromyidae. It is found in Colombia, Costa Rica, El Salvador, Guatemala, Honduras, Mexico, Nicaragua, and Panama.

Desperate shrew

Desperate shrew

Crocidura desperata

The desperate shrew (*Crocidura desperata*) is a species of mammal in the family Soricidae. It is endemic to Tanzania. Its natural habitat is subtropical or tropical moist montane forests. It is threatened by habitat loss.

Dhofar shrew

Dhofar shrew

Crocidura dhofarensis

The Dhofar shrew (*Crocidura dhofarensis*) is a white-toothed shrew found only in the Dhofar region of Oman.

Dhole

Dhole

Cuon alpinus

The dhole (*Cuon alpinus*) is a canid native Asia. During the Pleistocene, the dhole ranged throughout Asia, Europe, and North America but became restricted to its historical range 12,000-18,000 years ago. The dhole is listed as Endangered on the...

Diadem leaf-nosed bat

Diadem leaf-nosed bat

Hipposideros diadema

The diadem leaf-nosed bat or diadem roundleaf bat (*Hipposideros diadema*) is one of the most widespread species of bat in the family Hipposideridae. It is probably most closely related to *Hipposideros demissus* from Makira and to *Hipposideros*...

Diademed Sifaka

Diademed Sifaka

Propithecus diadema

The diademed sifaka (*Propithecus diadema*), or diademed simpona, is an endangered species of sifaka, one of the lemurs endemic to certain rainforests in eastern Madagascar. Along with the indri, this species is one of the two largest living...

Dian's tarsier

Dian's tarsier

Tarsius dentatus

Dian's tarsier (*Tarsius dentatus*), also known as the Diana tarsier, is a nocturnal primate endemic to central Sulawesi, Indonesia. Its head-body length is 11.5-12 centimetres (4.5-4.7 in) and it has a tail of 22 centimetres (8.7 in). Dian's...

Diana Monkey

Diana Monkey

Cercopithecus diana

The Diana monkey (*Cercopithecus diana*) is an Old World monkey found in the high canopy forests in Sierra Leone, Liberia, and western Côte d'Ivoire. Named for its white brow which is said to resemble the bow of the Roman goddess Diana, this...

Dibatag

Dibatag

Ammodorcas clarkei

The dibatag (*Ammodorcas clarkei*) is a slender antelope native to Africa. It is often confused with the gerenuk due to their striking resemblance. Dibatag are very secretive due to their brown coat that provides excellent camouflage. They are...

Dibbler

Dibbler

Parantechinus apicalis

Dibbler is the common name for *Parantechinus apicalis*, an endangered species of marsupial. It is an inhabitant of the southwest mainland of Western Australia and some offshore islands. It is a member of the order Dasyuromorphia, and the only...

Dice's cottontail
Dice's cottontail
Sylvilagus dicei

Dice's cottontail (*Sylvilagus dicei*) is a species of cottontail rabbit in the family Leporidae. It is found in Costa Rica and Panama, in páramo and cloud forest habitats.

Dickey's deer mouse
Dickey's deer mouse
Peromyscus dickeyi

Dickey's deer mouse (*Peromyscus dickeyi*) is a species of rodent in the family Cricetidae. It is endemic to Mexico, being found only on a small island in the Gulf of California. The species is named for Donald Dickey, who sponsored the expedition...

Dieterlen's brush-furred mouse
Dieterlen's brush-furred mouse
Lophuromys dieterleni

Dieterlen's brush-furred mouse, Mt Oku brush-furred mouse, or Mount Oku brush-furred rat (*Lophuromys dieterleni*) is a species of rodent in the family Muridae. It is endemic to Mount Oku, Cameroon. Its natural habitat is montane forest at...

Diminutive serotine
Diminutive serotine
Eptesicus diminutus

The diminutive serotine (*Eptesicus diminutus*) is a bat species from South America. It is found in Argentina, Colombia, Brazil, Paraguay, Uruguay, and Venezuela.

Diminutive woodrat
Diminutive woodrat
Nelsonia neotomodon

The diminutive woodrat (*Nelsonia neotomodon*) is a species of rodent in the family Cricetidae. It is endemic to Mexico.

Dinagat bushy-tailed cloud rat
Dinagat bushy-tailed cloud rat
Crateromys australis

The Dinagat bushy-tailed cloud rat, Dinagat crateromys or Dinagat cloud rat (*Crateromys australis*) is a species of cloud rat in the family Muridae. It is one of the eight cloud rats found only in the Philippines, and is specifically endemic to...

Dinagat gymnure
Dinagat gymnure
Podogymnura aureospinula

The Dinagat gymnure (*Podogymnura aureospinula*) is a species of mammal in the family Erinaceidae. It is endemic to the Philippines. Its natural habitat is subtropical or tropical dry forests. It is threatened by habitat loss. The mammal is an...

Dinagat hairy-tailed rat
Dinagat hairy-tailed rat
Batomys russatus

Dinagat hairy-tailed rat or Russet Batomys (*Batomys russatus*) is one of five species of rodent in the genus *Batomys*. It is in the diverse family Muridae. This species is endemic to the Philippines.

Dingiso
Dingiso

Dendrolagus mbaiso

The dingiso (*Dendrolagus mbaiso*) is an endangered tree-kangaroo found only on the island of New Guinea (in Indonesia). Though sacred to the local Moni people, it is still threatened by hunting and habitat loss.

Dingo

Dingo

Canis dingo

The dingo (*Canis familiaris*, *Canis familiaris dingo*, *Canis dingo*, or *Canis lupus dingo*) is an ancient (basal) lineage of dog found in Australia. It is variously considered a form of domestic dog not warranting recognition as a subspecies, a...

Disk-footed bat

Disk-footed bat

Eudiscopus denticulus

The disk-footed bat (*Eudiscopus denticulus*) is a species of vesper bat in the Vespertilionidae family found in Laos and Myanmar. The disc-footed bat was recorded in the northeastern Indian state's (Meghalaya), Lailad area near the Nongkhylllem...

Distinguished Oldfield mouse

Distinguished Oldfield mouse

Thomasomys notatus

The distinguished Oldfield mouse (*Thomasomys notatus*) is a species of rodent in the family Cricetidae. It is found only in Peru.

Djungarian Hamster

Djungarian Hamster

Phodopus sungorus

The Djungarian hamster (*Phodopus sungorus*) is one of three species of hamster in the genus *Phodopus*. It is ball-shaped and typically half the size of the Syrian hamster, so is called a dwarf hamster along with all *Phodopus* species. As winter...

Dneper common shrew

Dneper common shrew

Sorex averini

The Dneper common shrew (*Sorex averini*) is a species of mammal in the family Soricidae.

Dobson's epauletted fruit bat

Dobson's epauletted fruit bat

Epomophorus dobsonii

Dobson's epauletted fruit bat, or Dobson's fruit bat (*Epomophorus dobsonii*) is a species of megabat in the family Pteropodidae. It is found in Angola, Democratic Republic of the Congo, Malawi, Mozambique, Rwanda, Tanzania, and Zambia. Its...

Dobson's horseshoe bat

Dobson's horseshoe bat

Rhinolophus yunanensis

Dobson's horseshoe bat (*Rhinolophus yunanensis*) is a species of bat in the family Rhinolophidae. It is found in China, India, Myanmar, and Thailand.

Dobson's shrew tenrec

Dobson's shrew tenrec

Nesogale dobsoni

Dobson's shrew tenrec (*Nesogale dobsoni*) is a species of mammal in the family Tenrecidae. It is endemic to Madagascar. Its natural habitats are subtropical or tropical moist lowland forest, subtropical or tropical moist montane forest,...

Doğramacı's vole

Doğramaci's vole
Microtus dogramacii

Doğramaci's vole (*Microtus dogramacii*) is a species of rodent in the family Cricetidae. It is found only in central Turkey, and is similar to the social vole, although different in terms of cranial proportions and karyotype.

Dollman's melomys
Dollman's melomys
Melomys dollmani

Dollman's melomys (*Melomys dollmani*) is a species of rodent from the family Muridae. It lives in the eastern highlands of Papua New Guinea at an elevation of at least 1,200 metres (3,900 ft) and on the slopes of Mount Hagen and Mount Sisa....

Dollman's spiny rat
Dollman's spiny rat
Maxomys dollmani

Dollman's spiny rat (*Maxomys dollmani*) is a species of rodent in the family Muridae. It is found only in Indonesia.

Dollman's tree mouse
Dollman's tree mouse
Prionomys batesi

Dollman's tree mouse (*Prionomys batesi*) is a poorly understood climbing mouse from central Africa. It is unique enough that it has been placed in a genus of its own, *Prionomys*, since its discovery in 1910.

Dollman's vlei rat
Dollman's vlei rat
Otomys dollmani

Dollman's vlei rat (*Otomys dollmani*) is a species of rodent in the family Muridae. It is found only in Kenya. Its natural habitats are subtropical or tropical high-altitude grassland and swamps. It is threatened by habitat loss. Some authorities,...

Dolorous grass mouse
Dolorous grass mouse
Akodon dolores

The dolorous grass mouse or Córdoba akodont (*Akodon dolores*) is a species of rodent in the family Cricetidae. It is found only in Argentina. It is an invertivore and can grow as large as 50.5g.

Domestic Guinea Pig
Domestic Guinea Pig
Cavia porcellus

The Guinea pig or Domestic guinea pig (*Cavia porcellus*) is a species of rodent belonging to the genus *Cavia* in the family Caviidae. Breeders tend to use the word cavy to describe the animal, while in scientific and laboratory contexts, it is far...

Domestic Pig
Domestic Pig
Sus scrofa domesticus

Domestic pigs (*Sus domesticus*) are commonly known as approachable, loyal, and highly intelligent creatures. These animals are rather large representatives of even-toed ungulates. Additionally, they are noticeably larger and slower than the wild...

Domestic Sheep

Domestic Sheep
Ovis aries

Sheep or Domestic sheep (*Ovis aries*) are domesticated, ruminant mammals typically kept as livestock. Like all ruminants, sheep are even-toed ungulates. Numbering a little over one billion, Domestic sheep are also the most numerous species of...

Domestic Yak
Domestic Yak
Bos grunniens

The Domestic yak (*Bos grunniens*) is a species of long-haired domesticated cattle that is descended from the wild yak. The yak may have diverged from cattle at any point between one and five million years ago, and there is some suggestion that it...

Domesticated Ferret
Domesticated Ferret
Mustela putorius furo

The ferret (*Mustela furo*) is a small, domesticated species belonging to the family Mustelidae. A ferret is most likely a domesticated form of the wild European ferret or polecat (*Mustela putorius*). Ferrets were bred for sport, hunting rabbits in...

Dominican myotis
Dominican myotis
Myotis dominicensis

The Dominican myotis (*Myotis dominicensis*) is a species of vesper bat. It is found in Dominica and Guadeloupe.

Dong Bac's trident bat
Dong Bac's trident bat
Aselliscus dongbacana

Dong Bac's trident bat (*Aselliscus dongbacana*) is a species of bat in the family Hipposideridae. It is found in northeastern Vietnam. Its type locality is Na Phong cave, Ba Be National Park, Bac Kan Province, Vietnam. *Aselliscus dongbacana* is...

Dongola gerbil
Dongola gerbil
Gerbillus dongolanus

The Dongola gerbil (*Gerbillus dongolanus*) is a rodent distributed mainly in Dongola, Sudan. It is sometimes considered conspecific with the greater Egyptian gerbil.

Donkey
Donkey
Equus africanus asinus

The Domestic donkey is a hoofed mammal in the family Equidae, the same family as the horse. It derives from the African wild ass, *Equus africanus*, and may be classified either as a subspecies thereof, *Equus africanus asinus*, or as a separate...

Dorcas Gazelle
Dorcas Gazelle
Gazella dorcas

The dorcas gazelle (*Gazella dorcas*), also known as the ariel gazelle, is a small and common gazelle. The dorcas gazelle stands about 55-65 cm (1.8-2.1 feet) at the shoulder, with a head and body length of 90-110 cm (3-3.5 feet) and a weight of...

Doria's tree-kangaroo
Doria's tree-kangaroo
Dendrolagus dorianus

Doria's tree-kangaroo (*Dendrolagus dorianus*) is a tree-kangaroo endemic to Papua New Guinea.

Dormer's bat

Dormer's bat

Scotozous dormeri

Dormer's bat or Dormer's pipistrelle (*Scotozous dormeri*) is a species of vesper bat. It is monotypical for its genus. It is found in Bangladesh, Bhutan, India, and Pakistan. Its natural habitats are subtropical or tropical dry forests, arable...

Dormouse tufted-tailed rat

Dormouse tufted-tailed rat

Eliurus myoxinus

The Dormouse tufted-tailed rat (*Eliurus myoxinus*) is a species of rodent in the family Nesomyidae. It is found only in Madagascar.

Dorothy's slender opossum

Dorothy's slender opossum

Marmosops dorothea

Dorothy's slender opossum (*Marmosops dorothea*) is a species of opossum in the family Didelphidae. It is found in Brazil and Bolivia. It is threatened by habitat loss. Recent research suggests that *M. dorothea* is a synonym of *M. noctivagus*, the...

Doucet's musk shrew

Doucet's musk shrew

Crocidura douceti

Doucet's musk shrew (*Crocidura douceti*) is a species of mammal in the family Soricidae. It is found in Ivory Coast, Liberia, and possibly Nigeria. Its natural habitat is subtropical or tropical dry forests.

Douglas Squirrel

Douglas Squirrel

Tamiasciurus douglasii

The Douglas squirrel (*Tamiasciurus douglasii*) is a pine squirrel found in the Pacific Northwest of North America. It is sometimes known as the chickaree or Pine squirrel, although these names are also used for the American red squirrel. The...

Drab Atlantic tree-rat

Drab Atlantic tree-rat

Phyllomys dasythrix

The Drab Atlantic tree-rat, *Phyllomys dasythrix*, is a spiny rat species from South America. It is found in Brazil.

Dragon tube-nosed fruit bat

Dragon tube-nosed fruit bat

Nyctimene draconilla

The dragon tube-nosed fruit bat (*Nyctimene draconilla*) is a species of bat in the family Pteropodidae. It is found on both sides of New Guinea: West Papua, Indonesia and Papua New Guinea. It is slightly smaller and very similar in appearance to...

Dressy Oldfield mouse

Dressy Oldfield mouse

Thomasomys vestitus

The dressy Oldfield mouse (*Thomasomys vestitus*) is a species of rodent in the family Cricetidae. It is found only in Venezuela.

Drill

Drill

Mandrillus leucophaeus

The drill (*Mandrillus leucophaeus*) is one of Africa's most endangered mammals. It is related to baboons and even more closely to mandrills.

Dromedary Camel

Dromedary Camel

Camelus dromedarius

The dromedary camel (*Camelus dromedaries*) is a large even-toed ungulate, of the genus *Camelus*, with one hump on its back. It is the tallest of the three species of camel. It has not occurred naturally in the wild for nearly 2,000 years. It was...

Drouhard's shrew tenrec

Drouhard's shrew tenrec

Microgale drouhardi

Drouhard's shrew tenrec (*Microgale drouhardi*), also known as the striped shrew tenrec, is a species of mammal in the family Tenrecidae. It is endemic to Madagascar. Its natural habitats are subtropical or tropical moist lowland forest and...

Dryad shrew tenrec

Dryad shrew tenrec

Microgale dryas

The dryad shrew tenrec (*Microgale dryas*), also known as the tree shrew tenrec, is a species of mammal in the family Tenrecidae. It is endemic to Madagascar. Its natural habitat is subtropical or tropical moist lowland forests. It is vulnerable...

Dryadonycteris capixaba

Dryadonycteris capixaba

Dryadonycteris capixaba

Dryadonycteris capixaba is a species of leaf-nosed bat found in South America. It is monotypic within the genus *Dryadonycteris*

Dryas monkey

Dryas monkey

Cercopithecus dryas

The Dryas monkey (*Chlorocebus dryas*), also known as Salonga monkey, ekele, or inoko, is a little-known species of Old World monkey found only in the Congo Basin, restricted to the left bank of the Congo River. It is now established that the...

Drylands vesper mouse

Drylands vesper mouse

Calomys musculus

The drylands vesper mouse (*Calomys musculus*) is a species of rodent in the family Cricetidae. It is found in Argentina and Paraguay.

Drymoreomys

Drymoreomys

Drymoreomys albimaculatus

Drymoreomys is a rodent genus in the tribe Oryzomyini that lives in the Atlantic Forest of Brazil. The single species, *D. albimaculatus*, is known only from the states of São Paulo and Santa Catarina and was not named until 2011. It lives in the...

Dsinezumi shrew

Dsinezumi shrew

Crocidura dsinezumi

The *Dsinezumi* shrew (*Crocidura dsinezumi*), also known as the Japanese white-

toothed shrew, is a species of musk shrew found in Japan and on Korea's Jeju Island. It is widespread, and considered to be of "least concern" by the IUCN. There has been...

Dubious trumpet-eared bat
Dubious trumpet-eared bat
Phoniscus aerosa

The dubious trumpet-eared bat (*Phoniscus aerosa*) is a species of vesper bat in the family Vespertilionidae. It is found only in possibly South Africa.

Dudu's brush-furred rat
Dudu's brush-furred rat
Lophuromys dudui

Dudu's brush-furred rat (*Lophuromys dudui*) is a rodent belonging to the genus *Lophuromys*. It is native to the northeast of the Democratic Republic of Congo, from Kisangani to the eastern mountains of Garamba, Blukwa and Djugu to Irangi. It has a...

Dugong
Dugong
Dugong dugon

The dugong (*Dugong dugon*) is one of four living species of the order Sirenia, which also includes three species of manatees. It is the only living representative of the once-diverse family Dugongidae; its closest modern relative, Steller's sea...

Duke of Abruzzi's free-tailed bat
Duke of Abruzzi's free-tailed bat
Chaerephon aloysiisabaudiae

Duke of Abruzzi's free-tailed bat (*Chaerephon aloysiisabaudiae*) is a species of bat in the family Molossidae. It is found in Cameroon, Central African Republic, Democratic Republic of the Congo, Ivory Coast, Gabon, Ghana, South Sudan, and...

Duke of Bedford's vole
Duke of Bedford's vole
Proedromys bedfordi

The Duke of Bedford's vole (*Proedromys bedfordi*) is a species of rodent in the family Cricetidae. It is found only in mountainous parts of central China. It is a rare species and the International Union for Conservation of Nature has assessed...

Dulzura kangaroo rat
Dulzura kangaroo rat
Dipodomys simulans

The Dulzura kangaroo rat, or San Diego kangaroo rat (*Dipodomys simulans*) is a species of rodent in the family Heteromyidae. It is found in Baja California, Mexico, and in the Colorado Desert and elsewhere in California in the United States. It...

Dune hairy-footed gerbil
Dune hairy-footed gerbil
Gerbillurus tytonis

The dune hairy-footed gerbil, or the Namib dune gerbil (*Gerbillurus tytonis*) is a species of rodent found only in Namibia. Its natural habitat is temperate desert where it lives in loose sand among sand dunes, feeding opportunistically on...

Durango chipmunk
Durango chipmunk
Neotamias durangae

The Durango chipmunk (*Neotamias durangae*) is a species of rodent in the family Sciuridae. It is endemic to Mexico.

Durrell's Vontsira
Durrell's Vontsira
Salanoia durrelli

Durrell's vontsira (*Salanoia durrelli*) is a small mammal discovered in 2004. It belongs to a group of meat-eating, cat- or fox-like mammals found only in Madagascar. The species is closely related to the Brown-tailed mongoose (*Salanoia concolor*).

Dusky Antechinus
Dusky Antechinus
Antechinus swainsonii

The dusky antechinus (*Antechinus swainsonii*), also known as Swainson's antechinus or the dusky marsupial mouse, is a species of small marsupial carnivore, a member of the family Dasyuridae. It is found in Australia.

Dusky bushbaby
Dusky bushbaby
Galago matschiei

The dusky bushbaby (*Galago matschiei*) is a species of primate in the family Galagidae. It is also known as Matschie's galago, in honour of the German zoologist Paul Matschie, curator of mammals at the Museum für Naturkunde in Berlin. Native to...

Dusky caenolestid
Dusky caenolestid
Caenolestes fuliginosus

The dusky caenolestid (*Caenolestes fuliginosus*), also known as Tate's shrew opossum, is a shrew opossum from South America. The dusky caenolestid is characterized by a dark brown coat with a lighter underbelly, soft and thick fur, and a loosely...

Dusky Dolphin
Dusky Dolphin
Lagenorhynchus obscurus

The dusky dolphin (*Lagenorhynchus obscurus*) is a dolphin found in coastal waters in the Southern Hemisphere. Its specific epithet is Latin for "dark" or "dim". It is very closely genetically related to the Pacific white-sided dolphin, but...

Dusky elephant shrew
Dusky elephant shrew
Elephantulus fuscus

The dusky elephant shrew or dusky sengi (*Elephantulus fuscus*) is a species of elephant shrew in the family Macroscelididae. It is found in Malawi, Mozambique, and Zambia. Its natural habitat is dry savanna.

Dusky field rat
Dusky field rat
Rattus sordidus

The dusky field rat (*Rattus sordidus*), also known as the canefield rat, is a species of rodent in the family Muridae. It is found in Australia, Indonesia, and Papua New Guinea. In Australia it is found in northern Queensland and along the east...

Dusky fruit bat
Dusky fruit bat
Penthetor lucasi

The dusky fruit bat, (*Penthetor lucasi*) is a species of bat found in Indonesia and Malaysia.

Dusky hopping mouse
Dusky hopping mouse
Notomys fuscus

The dusky hopping mouse (*Notomys fuscus*), is a small rodent endemic to Australia, inhabiting desert regions characterised by sand dunes. Populations have experienced significant declines since the arrival of Europeans, and continue to be subject...

Dusky Leaf Monkey
Dusky Leaf Monkey
Trachypithecus obscurus

The dusky leaf monkey, spectacled langur, or spectacled leaf monkey (*Trachypithecus obscurus*) is a species of primate in the family Cercopithecidae. It is found in Peninsular Malaysia, Myanmar and Thailand, and can occasionally be found in...

Dusky leaf-nosed bat
Dusky leaf-nosed bat
Hipposideros ater

The dusky leaf-nosed bat (*Hipposideros ater*) is a bat from the genus *Hipposideros* whose habitat extends from India and Sri Lanka to the Philippines, New Guinea and Northern Australia. This species is counted in the *H. bicolor* species group and...

Dusky mosaic-tailed rat
Dusky mosaic-tailed rat
Melomys aereus

The dusky mosaic-tailed rat (*Melomys aereus*) is a species of rodent in the family Muridae.

Dusky Pademelon
Dusky Pademelon
Thylogale brunii

The Dusky pademelon (*Thylogale brunii*) is a species of marsupial found in New Guinea. The scientific name of this pademelon honors Cornelis de Bruijn, the Dutch painter who first described it in the second volume of his Travels, originally...

Dusky pipistrelle
Dusky pipistrelle
Pipistrellus hesperus

The dusky pipistrelle (*Pipistrellus hesperus*) is a small pipistrelle bat found in Africa.

Dusky rat
Dusky rat
Rattus colletti

The dusky rat (*Rattus colletti*) is an indigenous species of rodent in the family Muridae found in Australia.

Dusky slender opossum
Dusky slender opossum
Marmosops fuscatus

The dusky slender opossum (*Marmosops fuscatus*) is a species of opossum in the family Didelphidae. It is found in Colombia, Trinidad and Tobago, and Venezuela. It is threatened by habitat loss.

Dusky spiny tree-rat
Dusky spiny tree-rat
Makalata obscura

The dusky spiny tree rat (*Makalata obscura*) is a species of rodent in the family Echimyidae. It is endemic to Brazil. The etymology of the species name corresponds

to the Latin word *obscurus* meaning dark.

Dusky tube-nosed bat

Dusky tube-nosed bat

Murina fusca

The dusky tube-nosed bat (*Murina fusca*) is a species of vesper bat in the family *Vespertilionidae*. It is found only in China.

Dusky-footed elephant shrew

Dusky-footed elephant shrew

Elephantulus fuscipes

The dusky-footed elephant shrew or dusky-footed sengi (*Elephantulus fuscipes*) is a species of elephant shrew in the family *Macroscelididae*. It is found in Democratic Republic of the Congo, South Sudan, and Uganda. Its natural habitat is...

Dusky-Footed Woodrat

Dusky-Footed Woodrat

Neotoma fuscipes

The Dusky-footed woodrat (*Neotoma fuscipes*) is a species of nocturnal rodent from North America. Woodrats are nicknamed "packrats" or "trade rats" because they tend to hoard things, build large domed dens, and "trade" by dropping then picking up...

Duthie's golden mole

Duthie's golden mole

Chlorotalpa duthieae

Duthie's golden mole (*Chlorotalpa duthieae*) is a species of mammal in the family *Chrysochloridae*. It is endemic to South Africa. Its natural habitats are subtropical or tropical moist lowland forests, moist savanna, temperate grassland, arable...

Dwarf blue sheep

Dwarf blue sheep

Pseudois schaeferi

The bharal (*Pseudois nayaur*), also called the blue sheep, is a caprine native to the high Himalayas. It is the only member of the genus *Pseudois*. It occurs in India, Bhutan, China (in Gansu, Ningxia, Sichuan, Tibet, and Inner Mongolia), Myanmar,...

Dwarf bonneted bat

Dwarf bonneted bat

Eumops bonariensis

The dwarf bonneted bat, or Peters' mastiff bat, (*Eumops bonariensis*), is a bat species from South and Central America.

Dwarf brocket

Dwarf brocket

Mazama chunyi

The dwarf brocket (*Mazama chunyi*), or chunyi, is a small species of deer native to the Andean highlands in western Bolivia and southeastern Peru, where it is found in forest and páramo. Its pelage is reddish-brown with dark grey foreparts and...

Dwarf dog-faced bat

Dwarf dog-faced bat

Molossops temminckii

The dwarf dog-faced bat (*Molossops temminckii*) is a species of free-tailed bat from South America. It is found in Argentina, Bolivia, Brazil, Colombia, Ecuador, Guyana, Peru, Paraguay and Uruguay, typically at lower elevations. It is one of two...

Dwarf fat-tailed jerboa

Dwarf fat-tailed jerboa
Pygeretmus pumilio

The dwarf fat-tailed jerboa (*Pygeretmus pumilio*) is a species of rodent in the family Dipodidae. It is found in China, Iran, Kazakhstan, Mongolia, and Russia. They mostly reside in desert environments which can explain their size. They go long...

Dwarf fat-tailed mouse opossum
Dwarf fat-tailed mouse opossum
Thylamys velutinus

The dwarf fat-tailed mouse opossum (*Thylamys velutinus*), also known as the velvety fat-tailed opossum is an opossum species from South America. It is endemic to Brazil, where it is found in cerrado and caatinga habitats. Its head-and-body length...

Dwarf flying fox
Dwarf flying fox
Pteropus woodfordi

The dwarf flying fox, least flying fox, or least fruit bat (*Pteropus woodfordi*) is a species of flying fox in the family Pteropodidae. It is endemic to the Solomon Islands. It is threatened by deforestation, which is believed to damage roosting...

Dwarf free-tailed bat
Dwarf free-tailed bat
Mops nanulus

The dwarf free-tailed bat (*Mops nanulus*) is a species of bat in the family Molossidae. It is found in Cameroon, Democratic Republic of the Congo, Ivory Coast, Ethiopia, Ghana, Guinea, Kenya, Nigeria, Sierra Leone, South Sudan, and Uganda. Its...

Dwarf gymnure
Dwarf gymnure
Hylomys parvus

The dwarf gymnure (*Hylomys parvus*) is a gymnure found only at Mount Kerinci, Sumatra, Indonesia. It is listed by the International Union for Conservation of Nature as a critically endangered species due to a restricted range. The dwarf gymnure...

Dwarf hutia
Dwarf hutia
Mesocapromys nanus

The dwarf hutia (*Mesocapromys nana*) is a possibly extinct species of hutia known only from Cuba. It was last seen in 1937.

Dwarf little fruit bat
Dwarf little fruit bat
Rhinophylla pumilio

The dwarf little fruit bat (*Rhinophylla pumilio*) is a species of leaf-nosed bat from South America.

Dwarf multimammate mouse
Dwarf multimammate mouse
Mastomys pernanus

The dwarf multimammate mouse (*Serengetimys pernanus*) is a species of rodent in the family Muridae. It is the only member of the genus *Serengetimys* ; it was formerly classified in the genus *Mastomys*. It is found in Kenya, Rwanda, and Tanzania. Its...

Dwarf musk deer
Dwarf musk deer

Moschus berezovskii

The dwarf musk deer or Chinese forest musk deer (*Moschus berezovskii*, Chinese: 林麝; pinyin: Lín shè) is an artiodactyl native to southern and central China and northernmost Vietnam. The species name is after the collector Mikhail Mikhailovich...

Dwarf scaly-tailed squirrel

Dwarf scaly-tailed squirrel

Anomalurus pusillus

The dwarf scaly-tailed squirrel (*Anomalurus pusillus*) is a species of rodent in the family Anomaluridae. It is found in Cameroon, Central African Republic, Republic of the Congo, Democratic Republic of the Congo, Equatorial Guinea, Gabon and...

Dwarf shrew

Dwarf shrew

Sorex nanus

The dwarf shrew (*Sorex nanus*) is a species of mammal in the family Soricidae endemic to Arizona, Colorado, Montana, Nebraska, New Mexico, South Dakota, Utah, and Wyoming in the United States. The type locality is Estes Park (Larimer County),...

Dwarf slit-faced bat

Dwarf slit-faced bat

Nycteris nana

The dwarf slit-faced bat (*Nycteris nana*) is a species of slit-faced bat living in forest and savanna regions of Central Africa. Two subspecies have been noted: *N. n. nana* and *N. n. tristis*.

Dwarf Sperm Whale

Dwarf Sperm Whale

Kogia sima

A small whale (smaller than some dolphins), the Dwarf sperm whale (*Kogia sima*) is a rare and insufficiently explored species. Most of the available information about this animal is taken from occasional strandings. The Dwarf sperm whale is quite...

Ear-spot squirrel

Ear-spot squirrel

Callosciurus adamsi

The ear-spot squirrel (*Callosciurus adamsi*) is a species of rodent in the family Sciuridae. It is endemic to Borneo (Indonesia and Malaysia). Diurnal and active mainly in small trees. Closely resembles the plantain squirrel (*Callosciurus notatus*)...

Eared hutia

Eared hutia

Mesocapromys auritus

The eared hutia or large-eared hutia (*Mesocapromys auritus*) is a species of rodent in the subfamily Capromyinae.

Earless water rat

Earless water rat

Crossomys moncktoni

The earless water rat (*Crossomys moncktoni*) is a New Guinea rodent, part of the *Hydromys* group of the subfamily of Old World rats and mice (Murinae). It is the only species of the genus *Crossomys*. This species is probably most closely related to...

Earth-colored mouse

Earth-colored mouse
Mus terricolor

The earth-colored mouse (*Mus terricolor*) is a species of rodent in the family Muridae. It is found in India, possibly Indonesia, Nepal, and Pakistan. The earth-colored mouse lives in cultivated fields in raised moist mounds of Earth, where they...

East African epauletted fruit bat
East African epauletted fruit bat
Epomophorus minimus

The East African epauletted fruit bat (*Epomophorus minimus*) is a species of megabat in the family Pteropodidae. It is found in Ethiopia, Kenya, Somalia, Tanzania, and Uganda. Its natural habitats are dry savanna and rocky areas. It is threatened...

East African highland shrew
East African highland shrew
Crocidura allex

The East African highland shrew (*Crocidura allex*) is a species of mammal in the family Soricidae. It is found in Kenya and Tanzania. Its natural habitats are subtropical or tropical moist montane forests, subtropical or tropical high-altitude...

East African little collared fruit bat
East African little collared fruit bat
Myonycteris relicta

The East African little collared fruit bat (*Myonycteris relicta*) is a species of megabat in the family Pteropodidae. It is found in Kenya, Tanzania, and Zimbabwe. Its natural habitats are subtropical or tropical dry forests and moist savanna. It...

East African Oryx
East African Oryx
Oryx beisa

The East African oryx (*Oryx beisa*), also known as the beisa is a species of antelope from East Africa. It has two subspecies: the common beisa oryx (*Oryx beisa beisa*) found in steppe and semidesert throughout the Horn of Africa and north of the...

East African Potto
East African Potto
Perodicticus ibeanus

The East African potto (*Perodicticus ibeanus*) is a nocturnal primate found in Central and East Africa. It is thought to be the sister species to the Central African potto, from which it diverged during the late Miocene, about 5.5 million years ago.

East African springhare
East African springhare
Pedetes surdaster

The East African springhare (*Pedetes surdaster*), is not closely related to the hare, which is a lagomorph, but is a member of the Pedetidae, a rodent family.

East African wild dog
East African wild dog
Lycaon pictus lupinus

The African wild dog (*Lycaon pictus*), also called the African painted dog and the African hunting dog, is a wild canine which is a native species to sub-Saharan

Africa. It is the largest wild canine in Africa, and the only extant member of the...

East Asian Finless Porpoise
East Asian Finless Porpoise
Neophocaena sunameri

The East Asian finless porpoise (*Neophocaena sunameri*) is a species of porpoise found in East Asia. The Yangtze finless porpoise was formerly considered a subspecies, but is now thought to be a distinct species.

East Asian free-tailed bat
East Asian free-tailed bat
Tadarida insignis

The East Asian free-tailed bat (*Tadarida insignis*) was formerly considered a synonym of the European free-tailed bat. Its range includes China, Japan, and the Korean Peninsula.

East Asian tailless leaf-nosed bat
East Asian tailless leaf-nosed bat
Coelops frithii

The East Asian tailless leaf-nosed bat or tail-less leaf-nosed bat (*Coelops frithii*) is a species of bat in the family Hipposideridae. It is found in Bangladesh, China, India, Indonesia, Laos, Malaysia, Myanmar, Taiwan, and Vietnam. The species...

East Caucasian Tur
East Caucasian Tur
Capra cylindricornis

The East Caucasian tur (*Capra cylindricornis*) is a mountain-dwelling caprine living in the eastern half of the Greater Caucasus mountains. The ancestors of the modern goats are thought to have moved into mountainous regions where they adapted to...

East Siberian brown bear
East Siberian brown bear
Ursus arctos collaris

The East Siberian brown bear (*Ursus arctos collaris*) is a population or subspecies of brown bear which ranges from eastern Siberia, beginning at the Yenisei river, north to the Arctic Circle, as far as Trans-Baikaliya, the Stanovoy Range, the...

Eastern Amazon climbing mouse
Eastern Amazon climbing mouse
Rhipidomys emiliae

The eastern Amazon climbing mouse (*Rhipidomys emiliae*) is a rodent species from South America. It is endemic to central Brazil, where it is found in the eastern fringe of the Amazon rainforest, as well as in gallery forest and tropical dry...

Eastern Barred Bandicoot
Eastern Barred Bandicoot
Perameles gunnii

The Eastern barred bandicoot (*Perameles gunnii*) is a nocturnal, rabbit-sized marsupial endemic to southeastern Australia. Once the Eastern barred bandicoot was widespread and abundant throughout Victoria's volcanic plains and grasslands but now...

Eastern bent-wing bat
Eastern bent-wing bat
Miniopterus fuliginosus

The eastern bent-wing bat, (*Miniopterus fuliginosus*) is a species of vesper bat in the family Vespertilionidae. It is found in South Asia, Far-east Asia, the east

Caucasus Mountains and also in Southeast Asian regions.

Eastern Bettong

Eastern Bettong

Bettongia gaimardi

The eastern bettong (*Bettongia gaimardi*), also known as the Balbo (by the Ngunnawal People who used to keep them as pets), southern bettong and Tasmanian bettong, is a bettong whose natural range includes southeastern Australia and eastern...

Eastern black crested gibbon

Eastern black crested gibbon

Nomascus nasutus

The eastern black-crested gibbon (*Nomascus nasutus*), also known as the Cao-vit black crested gibbon or the Cao-vit crested gibbon, is a species of gibbon from southeast China and northern Vietnam. The term "Cao-vit" originated from the sounds of...

Eastern black rhinoceros

Eastern black rhinoceros

Diceros bicornis michaeli

The eastern black rhinoceros (*Diceros bicornis michaeli*) is also known as the East African black rhinoceros or eastern hook-lipped rhinoceros. It is a subspecies of the black rhinoceros. Its numbers are very low due to poaching for its horn and...

Eastern broad-nosed bat

Eastern broad-nosed bat

Scotorepens orion

The eastern broad-nosed bat or Orion broad-nosed bat (*Scotorepens orion*) is a species of vespertilionid bat. It is found only in Australia, east of the Great Dividing Range, from about Rockhampton to Melbourne, with a small isolated population...

Eastern broad-toothed field mouse

Eastern broad-toothed field mouse

Apodemus mystacinus

The eastern broad-toothed field mouse (*Apodemus mystacinus*) is a species of rodent in the family Muridae.

Eastern caenolestid

Eastern caenolestid

Caenolestes sangay

The eastern caenolestid (*Caenolestes sangay*) is a shrew opossum found on the eastern slopes of the Andes in southern Ecuador.

Eastern cave bat

Eastern cave bat

Vespadelus troungtoni

The eastern cave bat or Troungton's forest bat (*Vespadelus troungtoni*) is a species of vesper bat in the family Vespertilionidae. It is found only in Australia. This bat lives in caves along the east coast of Australia and in adjacent inland...

Eastern chestnut mouse

Eastern chestnut mouse

Pseudomys gracilicaudatus

The eastern chestnut mouse (*Pseudomys gracilicaudatus*) is a species of rodent in the family Muridae. It is found only in Australia, along the eastern coast from northern Queensland and into New South Wales as far as Jervis Bay.

Eastern Chimpanzee
Eastern Chimpanzee
Pan troglodytes schweinfurthii

The Eastern chimpanzee (*Pan troglodytes schweinfurthii*) is a subspecies of the Common chimpanzee.

Eastern Chipmunk
Eastern Chipmunk

Tamias striatus

The Eastern chipmunk (*Tamias striatus*) is a small striped rodent found in North America. The name "chipmunk" comes from the Ojibwe word *ajidamoo* (or possibly *ajidamoonh*, the same word in the Ottawa dialect of Ojibwe), which translates literally...

Eastern common cuscus
Eastern common cuscus
Phalanger intercastellanus

The eastern common cuscus (*Phalanger intercastellanus*) is a species of marsupial in the family *Phalangeridae* found in eastern Papua New Guinea. Until recently, it was considered conspecific with *P. mimicus*, and before that also with *P....*

Eastern Cordillera small-footed shrew
Eastern Cordillera small-footed shrew
Cryptotis brachyonyx

The Eastern Cordillera small-footed shrew (*Cryptotis brachyonyx*) is a species of mammal in the family *Soricidae*. It is endemic to Colombia, where it is known from the western slopes of the central Cordillera Oriental at elevations from 1300 to...

Eastern Cottontail
Eastern Cottontail
Sylvilagus floridanus

The Eastern cottontail (*Sylvilagus floridanus*) is a New World cottontail rabbit, a member of the family *Leporidae*. It is the most common rabbit species in North America.

Eastern deer Mouse
Eastern deer Mouse
Peromyscus maniculatus

Endemic to North America, the Eastern deer mouse (*Peromyscus maniculatus*) is one of the most common organisms of the biotic community in nearly all woodlands throughout the continent. However, even the most attentive observers often cannot spot...

Eastern elk
Eastern elk
Cervus canadensis canadensis

The eastern elk (*Cervus canadensis canadensis*) is an extinct subspecies or distinct population of elk that inhabited the northern and eastern United States, and southern Canada. The last eastern elk was shot in Pennsylvania on September 1, 1877....

Eastern falanouc
Eastern falanouc
Eupleres goudotii

The Eastern falanouc (*Eupleres goudotii*) is a rare mongoose-like mammal in the carnivorous family *Eupleridae* endemic to Madagascar. It is classified alongside the Western falanouc (*Eupleres major*), recognized only in 2010, in the genus *Eupleres*....

Eastern false pipistrelle
Eastern false pipistrelle
Falsistrellus tasmaniensis

The eastern false pipistrelle (*Falsistrellus tasmaniensis*) is a vespertilionid bat that occurs in eastern and south-eastern Australia, including the island of Tasmania.

Eastern forest bat
Eastern forest bat
Vespadelus pumilus

The eastern forest bat (*Vespadelus pumilus*) is a species of vesper bat in the family Vespertilionidae. It is found only in Australia, where it has been recorded from Queensland to New South Wales. The population is in decline, with the number of...

Eastern Fox Squirrel
Eastern Fox Squirrel
Sciurus niger

The Eastern fox squirrel (*Sciurus niger*) is the largest species of tree squirrel native to North America. Despite the differences in size and coloration, it is sometimes mistaken for American red squirrels or Eastern gray squirrels in areas where...

Eastern Gorilla
Eastern Gorilla
Gorilla beringei

The Eastern gorilla (*Gorilla beringei*) is a critically endangered species of Gorilla and the largest living primate. Illegal hunting threatens the species.

Eastern Gray Squirrel
Eastern Gray Squirrel
Sciurus carolinensis

The Eastern gray squirrel (*Sciurus carolinensis*) is a tree squirrel native to eastern North America; there it is the most prodigious and ecologically essential natural forest regenerator. Widely introduced to certain places around the world, the...

Eastern Grey Kangaroo
Eastern Grey Kangaroo
Macropus giganteus

The Eastern grey kangaroo (*Macropus giganteus*) is a marsupial found only in Australia. It is the second largest and heaviest living marsupial and native land mammal in Australia. It is also known as the Great grey kangaroo and the Forester kangaroo.

Eastern harvest mouse
Eastern harvest mouse
Reithrodontomys humulis

The eastern harvest mouse (*Reithrodontomys humulis*) is a species of rodent in the family Cricetidae. It is endemic to the Southeastern United States. Its natural habitats are subtropical or tropical seasonally wet or flooded lowland grassland,...

Eastern heather vole
Eastern heather vole
Phenacomys ungava

The eastern heather vole, (*Phenacomys ungava*), is a small North American vole. Until recently, this species was considered to belong to the same species as the

western heather vole, *Phenacomys intermedius*. It is also called the Ungava...
Eastern hoolock gibbon
Eastern hoolock gibbon
Hoolock leuconedys

The eastern hoolock gibbon (*Hoolock leuconedys*) is a primate from the gibbon family, Hylobatidae. It is one of three species of hoolock gibbon. This species is found in east of the Chindwin River, such as the Mahamyaing Wildlife Sanctuary, and...
Eastern lesser bamboo lemur
Eastern lesser bamboo lemur
Hapalemur griseus

The eastern lesser bamboo lemur (*Hapalemur griseus*), also known as the gray bamboo lemur, the gray gentle lemur, and the Mahajanga lemur is a small lemur endemic to Madagascar, with three known subspecies. As its name suggests, the eastern...
Eastern Long-Beaked Echidna
Eastern Long-Beaked Echidna
Zaglossus bartoni

The eastern long-beaked echidna (*Zaglossus bartoni*), also known as Barton's long-beaked echidna, is one of three species from the genus *Zaglossus* to occur in New Guinea. It is found mainly in the eastern half at elevations between 2,000 and...
Eastern long-eared bat
Eastern long-eared bat
Nyctophilus bifax

The eastern long-eared bat, species *Nyctophilus bifax*, is a small flying mammal, a vespertilionid bat. It is found in eastern Australia and Papua New Guinea.
Eastern long-fingered bat
Eastern long-fingered bat
Myotis macrodactylus

The eastern long-fingered bat, or big-footed myotis (*Myotis macrodactylus*) is a species of vesper bat. An adult big-footed myotis has a body length of 4.1-4.8 cm, a tail of 3.1-4.9 cm, and a wing length of 3.7-4.2 cm. It nests in groups, and...
Eastern lowland gorilla
Eastern lowland gorilla
Gorilla beringei graueri

The eastern lowland gorilla (*Gorilla beringei graueri*) or Grauer's gorilla is a Critically Endangered subspecies of eastern gorilla endemic to the mountainous forests of eastern Democratic Republic of the Congo. Important populations of this...
Eastern lowland olingo
Eastern lowland olingo
Bassaricyon alleni

The eastern lowland olingo (*Bassaricyon alleni*) is a species of olingo from South America, where it is known from the lowlands east of the Andes in Bolivia, Brazil, Colombia, Ecuador, Guyana, Peru and Venezuela. It is the only olingo species...
Eastern Meadow Vole
Eastern Meadow Vole
Microtus pennsylvanicus

The Eastern meadow vole (*Microtus pennsylvanicus*) is a North American rodent. It is active year-round, usually at night. It also digs burrows, where it stores food for the winter and females give birth to their young.
Eastern Mole

Eastern Mole
Scalopus aquaticus

The eastern mole or common mole (*Scalopus aquaticus*) is a medium-sized, overall grey North American mole and the only member of the genus *Scalopus*. Its large, hairless, spade-shaped forefeet are adapted for digging. The species is native to...

Eastern moose
Eastern moose
Alces alces americana

The eastern moose (*Alces alces americana*) is a subspecies of moose that currently ranges throughout Eastern Canada, New England and northern New York State. It inhabits boreal forests and mixed deciduous forests. It is the third largest North...

Eastern puna mouse
Eastern puna mouse
Punomys kofordi

The eastern puna mouse (*Punomys kofordi*) is a species of rodent in the family Cricetidae. It is endemic to southern Peru, where it is found in moist areas within puna grassland at elevations of 4500 to 4800 m in the Kallawaya mountain range, a...

Eastern Pygmy Possum
Eastern Pygmy Possum
Cercartetus nanus

The eastern pygmy possum (*Cercartetus nanus*) is a diprotodont marsupial of south-eastern Australia. Occurring from southern Queensland to eastern South Australia and also Tasmania, it is found in a range of habitats, including rainforest,...

Eastern Quoll
Eastern Quoll
Dasyurus viverrinus

The eastern quoll (*Dasyurus viverrinus*), found only in Australia, and formerly known as the eastern native cat, is a medium-sized carnivorous dasyurid marsupial. They are widespread and even locally common in Tasmania. They have been considered...

Eastern rainforest grammomys
Eastern rainforest grammomys
Grammomys kuru

The eastern rainforest grammomys (*Grammomys kuru*), also known as the eastern rainforest thicket rat, is a species of rodent from the family Muridae. The eastern rainforest grammomys is distributed in the Congo Basin in the Central African...

Eastern rat
Eastern rat
Rattus mordax

The eastern rat (*Rattus mordax*) is a species of rodent in the family Muridae. It is found only in Papua New Guinea.

Eastern Red Bat
Eastern Red Bat
Lasiurus borealis

The eastern red bat (*Lasiurus borealis*) is a species of microbat in the family Vespertilionidae. Eastern red bats are widespread across eastern North America, with additional records in Bermuda.

Eastern rock elephant shrew

Eastern rock elephant shrew
Elephantulus myurus

The eastern rock elephant shrew or eastern rock sengi (*Elephantulus myurus*) is a species of elephant shrew in the family Macroscelididae. It is found in Botswana, Mozambique, South Africa, and Zimbabwe. Its natural habitats are subtropical or...
Eastern short-eared rock-wallaby
Eastern short-eared rock-wallaby
Petrogale wilkinsi

The eastern short-eared rock-wallaby or Wilkins' rock-wallaby (*Petrogale wilkinsi*) is a species of rock-wallaby found in the northernmost parts of the Northern Territory of Australia, and is common in the Kakadu and Litchfield National Parks. It...
Eastern shrew mouse
Eastern shrew mouse
Pseudohydromys murinus

The eastern shrew mouse (*Pseudohydromys murinus*) is a species of rodent in the family Muridae. It is found only in Papua New Guinea.
Eastern small-footed myotis
Eastern small-footed myotis
Myotis leibii

The eastern small-footed bat (*Myotis leibii*) is a species of vesper bat. It can be found in southern Ontario and Quebec in Canada and in mountainous portions of the eastern United States from New England to northern Georgia, and westward to...
Eastern small-toothed rat
Eastern small-toothed rat
Macruromys major

The eastern small-toothed rat (*Macruromys major*) is a species of rodent in the family Muridae. It is found in West Papua, Indonesia and Papua New Guinea.
Eastern spiny mouse
Eastern spiny mouse
Acomys dimidiatus

The eastern spiny mouse or Arabian spiny mouse (*Acomys dimidiatus*) is a species of rodent in the family Muridae. They have a wide range, having been found in Middle Eastern deserts, as well as being prevalent in riverine forests in Africa. This...
Eastern Spotted Skunk
Eastern Spotted Skunk
Spilogale putorius

The eastern spotted skunk (*Spilogale putorius*) is a small, relatively slender skunk found in North America, throughout the eastern United States and in small areas of Canada and Mexico. This small skunk is more weasel-like in body shape than the...
Eastern striped bandicoot
Eastern striped bandicoot
Microperoryctes ornata

The eastern striped bandicoot (*Microperoryctes ornata*) is a species of marsupial in the family Peramelidae. It is found in eastern Papua New Guinea in the Enga Province. It is endemic to mountain forested habitats ranging from 1,000 – 3,600 m in...
Eastern tube-nosed bat
Eastern tube-nosed bat
Nyctimene robinsoni

The eastern or Queensland tube-nosed bat (*Nyctimene robinsoni*) is a megabat in the family Pteropodidae that lives in north-eastern Australia. *N. robinsoni* is one of the few species in Pteropodidae that roosts solitarily. They get their common...

Eastern voalavo

Eastern voalavo

Voalavo antsahabensis

The eastern voalavo (*Voalavo antsahabensis*) is a rodent in the family Nesomyidae which occurs in the Anjozorobe forest of eastern Madagascar. Although surveys before 2002 failed to record the species, it is common in some places. However, it is...

Eastern water bat

Eastern water bat

Myotis petax

The eastern water bat or Sakhalin bat (*Myotis petax*) is a species of mouse-eared bat. It was for a long time considered to be a subspecies of *Myotis daubentonii*.

Eastern white-eared giant rat

Eastern white-eared giant rat

Hyomys goliath

The eastern white-eared giant rat (*Hyomys goliath*) is a species of rodent in the family Muridae. It is found only in Papua New Guinea. The species has been known to eat karuka nuts (*Pandanus julianettii*), and growers will put platforms or other...

Eastern Wolf

Eastern Wolf

Canis lycaon

The Eastern wolf (*Canis lupus lycaon* or *Canis lycaon* or *Canis rufus lycaon*) is considered to be either a unique subspecies of Grey wolf or Red wolf or a separate species from both. Regardless of its status, it is regarded as unique and therefore...

Eastern Woolly Lemur

Eastern Woolly Lemur

Avahi laniger

The eastern woolly lemur (*Avahi laniger*), also known as the eastern avahi or Gmelin's woolly lemur, is a species of woolly lemur native to eastern Madagascar, where it lives in the wet tropical rainforest at low elevations along the eastern...

Echigo mole

Echigo mole

Mogera etigo

The Echigo mole (*Mogera etigo*) is an endangered species of mammal in the family Talpidae. It is endemic to Japan, being found only on the Echigo plain, Niigata prefecture.

Echimys vieirai

Echimys vieirai

Echimys vieirai

Echimys vieirai is a spiny rat species from South America, described in 2005. It is found in Brazil. The species name *vieirai* is a patronym for the Brazilian mammalogist Carlos Octaviano da Cunha Vieira, curator of the Mammal Collection at the...

Ecuadoran spiny pocket mouse

Ecuadoran spiny pocket mouse

Heteromys teleus

The Ecuadoran spiny pocket mouse (*Heteromys teleus*) is a species of rodent in the family Heteromyidae. It is endemic to central western Ecuador, where it is found at elevations from sea level to 2000 m on the coastal plain and western slopes of...

Ecuadorian capuchin

Ecuadorian capuchin

Cebus aequatorialis

The Ecuadorian capuchin (*Cebus aequatorialis*), or Ecuadorian white-fronted capuchin is a species of gracile capuchin monkey of the family Cebidae. It was formerly classified as a subspecies of the white-fronted capuchin. Mittermeier and Rylands...

Ecuadorian grass mouse

Ecuadorian grass mouse

Neomicroxus latebricola

The Ecuadorian grass mouse (*Neomicroxus latebricola*) is a species of rodent in the family Cricetidae. It is found only in Ecuador.

Ecuadorian mantled howler

Ecuadorian mantled howler

Alouatta palliata aequatorialis

The Ecuadorian mantled howler (*Alouatta palliata aequatorialis*) is a subspecies of the mantled howler, *A. palliata*. It ranges from Panama (or possibly the eastern tip of Costa Rica) through Colombia and Ecuador into northern Peru. The range...

Ecuadorian sac-winged bat

Ecuadorian sac-winged bat

Balantiopteryx infusca

The Ecuadorian sac-winged bat (*Balantiopteryx infusca*) is a species of sac-winged bat in the family Emballonuridae. It is found in Colombia and Ecuador. According to the IUCN Red List of Threatened Species, the population trend is decreasing for...

Ecuadorian small-eared shrew

Ecuadorian small-eared shrew

Cryptotis equatoris

The Ecuadorian small-eared shrew (*Cryptotis equatoris*) is a species of shrew in the family Soricidae. It is found on the western and eastern slopes of the Andes in central Ecuador.

Ecuadorian squirrel monkey

Ecuadorian squirrel monkey

Saimiri cassiquiarensis macrodon

The Ecuadorian squirrel monkey (*Saimiri cassiquiarensis macrodon*) is a type of squirrel monkey. It had been considered a subspecies of the Guianan squirrel monkey, *Saimiri sciureus*, but was elevated to a full species, *S. macrodon*, based on a...

Edible Dormouse

Edible Dormouse

Glis glis

The Edible dormouse (*Glis glis*) is a large dormouse and one of only two living species in the genus *Glis*, found in most of Europe and parts of western Asia. Its name comes from the Romans, who ate them as a delicacy.

Edith's leaf-eared mouse

Edith's leaf-eared mouse

Graomys edithae

Edith's leaf-eared mouse (*Graomys edithae*) is a species of rodent in the family

Cricetidae. It is known only from Argentina, where it was found in a montane grassland on Otro Cerro at an elevation of 3000 m.
Edward's swamp rat
Edward's swamp rat
Malacomys edwardsi

Edward's swamp rat (*Malacomys edwardsi*) is a species of rodent in the family Muridae. It is found in Ivory Coast, Ghana, Guinea, Liberia, Nigeria, and Sierra Leone. Its natural habitats are subtropical or tropical moist lowland forests and...
Edwards's long-tailed giant rat
Edwards's long-tailed giant rat
Leopoldamys edwardsi

Edwards's long-tailed giant rat (*Leopoldamys edwardsi*) is a species of rodent in the family Muridae. It is found in China, India, Indonesia, Laos, Malaysia, Burma, Thailand, and Vietnam.
Ega long-tongued bat
Ega long-tongued bat
Scleronycteris ega

The Ega long-tongued bat (*Scleronycteris ega*) is a bat species from South America. It is found in northwestern Brazil and southern Venezuela. It is monotypic within its genus.
Eger's long-fingered bat
Eger's long-fingered bat
Miniopterus egeri

Eger's long-fingered bat (*Miniopterus egeri*) is a species of long-fingered bat found in Madagascar.
Egyptian free-tailed bat
Egyptian free-tailed bat
Tadarida aegyptiaca

The Egyptian free-tailed bat, also known as Egyptian guano bat or Egyptian nyctinome, (*Tadarida aegyptiaca*) is a species of bat in the family Molossidae.
Egyptian Fruit Bat
Egyptian Fruit Bat
Rousettus aegyptiacus

The Egyptian fruit bat or Egyptian rousette (*Rousettus aegyptiacus*) is a species of megabat that is found in Africa, the Middle East, the Mediterranean, and the Indian subcontinent. It is one of three *Rousettus* species with an African-Malagasy...
Egyptian Mongoose
Egyptian Mongoose
Herpestes ichneumon

The Egyptian mongoose (*Herpestes ichneumon*), also known as ichneumon, is a mongoose species native to the coastal regions along the Mediterranean Sea between North Africa and Turkey, tropical and subtropical grasslands, savannas, and shrublands...
Egyptian mouse-tailed bat
Egyptian mouse-tailed bat
Rhinopoma cystops

The Egyptian mouse-tailed bat (*Rhinopoma cystops*) is a species of mouse-tailed bat found in North Africa and the Middle East.
Egyptian pygmy shrew
Egyptian pygmy shrew

Crocidura religiosa

The Egyptian pygmy shrew or sacred shrew (*Crocidura religiosa*) is a species of mammal in the family Soricidae. It is endemic to Egypt. Its natural habitat is arable land. It is threatened by habitat loss, partially caused by the construction of...

Egyptian slit-faced bat

Egyptian slit-faced bat

Nycteris thebaica

The Egyptian slit-faced bat (*Nycteris thebaica*) is a species of slit-faced bat broadly distributed throughout Africa and the Middle East. It is a species of microbat in the family Nycteridae. Six subspecies are known.

Egyptian tomb bat

Egyptian tomb bat

Taphozous perforatus

The Egyptian tomb bat (*Taphozous perforatus*) is a species of sac-winged bat in the family Emballonuridae. It is a medium- to large-sized microbat with a mass of approximately 30 g (1.1 oz). It is an aerial insectivore, foraging in open space....

Egyptian Weasel

Egyptian Weasel

Mustela subpalmata

The Egyptian weasel (*Mustela nivalis subpalmata*) is a unique population of the Least weasel found only in northern Egypt. It was formerly considered a distinct species, as *Mustela subpalmata*.

Egyptian Wolf

Egyptian Wolf

Canis lupaster lupaster

The Egyptian wolf (*Canis lupaster lupaster*) is a subspecies of African golden wolf.

Eisentraut's mouse shrew

Eisentraut's mouse shrew

Myosorex eisentrauti

Eisentraut's mouse shrew (*Myosorex eisentrauti*) is a Myosoricinae shrew found only on the island of Bioko, Equatorial Guinea. It is listed as a critically endangered species due to habitat loss and a restricted range.

Eisentraut's pipistrelle

Eisentraut's pipistrelle

Parahypsugo eisentrauti

Eisentraut's serotine (*Nycticeinops eisentrauti*), formerly known as Eisentraut's pipistrelle, is a species of vesper bat in the family Vespertilionidae. It is found in Cameroon, Democratic Republic of the Congo, Kenya, Somalia, and Uganda. Its...

Eisentraut's shrew

Eisentraut's shrew

Crocidura eisentrauti

Eisentraut's shrew (*Crocidura eisentrauti*) is a species of mammal in the family Soricidae. It is endemic to Cameroon. Its natural habitat is subtropical or tropical high-altitude grassland.

Eisentraut's striped mouse

Eisentraut's striped mouse

Hybomys badius

Eisentraut's striped mouse or Eisentraut's hybomys (*Hybomys badius*) is a species

of rodent in the family Muridae. It is found only in Cameroon. Its natural habitat is subtropical or tropical moist montane forests. It is threatened by habitat loss.

Ejeta's house bat

Ejeta's house bat

Scotophilus ejetai

Ejeta's yellow bat or Ejeta's house bat (*Scotophilus ejetai*) is a species of vesper bat endemic to Ethiopia. It was described as a new species of bat in 2014.

El Carrizo deer mouse

El Carrizo deer mouse

Peromyscus ochraventer

The El Carrizo deer mouse (*Peromyscus ochraventer*) is a species of rodent in the family Cricetidae. It is found only in Mexico.

El Dorado grass mouse

El Dorado grass mouse

Akodon orophilus

The El Dorado grass mouse or Utcubamba akodont (*Akodon orophilus*) is a species of rodent in the family Cricetidae. It is found in Ecuador and Peru.

Eld's Deer

Eld's Deer

Panolia eldii

Eld's deer (*Rucervus eldii* (or *Panolia eldii*)), also known as the thamin or brow-antlered deer, is an endangered species of deer endemic to South Asia.

Eldorado broad-nosed bat

Eldorado broad-nosed bat

Platyrrhinus aurarius

The Eldorado broad-nosed bat (*Platyrrhinus aurarius*) is a species of bat in the family Phyllostomidae. It is found in Guyana, Suriname, northern Brazil, and southern Venezuela.

Elegant fat-tailed mouse opossum

Elegant fat-tailed mouse opossum

Thylamys elegans

The elegant fat-tailed mouse opossum (*Thylamys elegans*), also known as the Chilean mouse opossum, is an opossum from central Chile. The type species of *Thylamys*, it was first described by English naturalist George Robert Waterhouse in 1839. This...

Elegant margareta rat

Elegant margareta rat

Margaretamys elegans

The elegant margareta rat (*Margaretamys elegans*) is a species of rodent in the family Muridae. It is endemic to the island of Sulawesi in Indonesia.

Elegant myotis

Elegant myotis

Myotis elegans

The elegant myotis (*Myotis elegans*) is a species of vesper bat found in Belize, Costa Rica, El Salvador, Guatemala, Honduras, Mexico, and Nicaragua.

Elegant water shrew

Elegant water shrew

Nectogale elegans

The elegant water shrew (*Nectogale elegans*) is a species of mammal in the subfamily Soricinae of the family Soricidae. It is monotypic within the genus

Nectogale. It lives in Sikkim and China.
Elephant Seal
Elephant Seal
Genus Mirounga

Elephant seals are large marine mammals and are the largest extant carnivorans that can weigh up to 5,000 kilograms (11,000 lb). Elephant seals are classified under the order Pinnipedia, which, in Latin, means feather- or fin-footed. Both...

Elephas beyeri
Elephas beyeri
Elephas beyeri

Elephas beyeri is an extinct species of dwarf elephant belonging to the Elephantidae family of the Middle Pleistocene. It was named after the anthropologist H. Otley Beyer. The type specimen was discovered on Cabarruyan Island in The Philippines...

Elephas celebensis
Elephas celebensis
Elephas celebensis

Elephas celebensis or the Sulawesi dwarf elephant is an extinct species of elephant.

Elephas ekorensis
Elephas ekorensis
Elephas ekorensis

Elephas ekorensis is an extinct species of large herbivorous mammals belonging to the family Elephantidae. Fossils have been found in East Africa dating as far back as the Early Pliocene age, between 5.3 and 3.6 million years ago. It is the...

Elephas hysudricus
Elephas hysudricus
Elephas hysudricus

Elephas hysudricus is an extinct elephant species and was described from fossil remains found in the Siwalik hills. It lived during the Pliocene and Pleistocene epochs.

Elephas hysudrindicus
Elephas hysudrindicus
Elephas hysudrindicus

Elephas hysudrindicus, commonly known as Gajah Blora (Blora Elephant) in Indonesia is a species of extinct elephant of the Pleistocene of Java and is anatomically distinct from the extant Asian elephant (*E. maximus*). The head of the Bandung...

Elephas iolensis
Elephas iolensis
Elephas iolensis

Elephas iolensis is an extinct species of large herbivorous mammal belonging to the Elephantidae family. The type specimen is located in the National Museum of Natural History in Paris. The species is thought to have lived in the African Savannah...

Elery's tube-nosed bat
Elery's tube-nosed bat
Murina eleryi

Elery's tube-nosed bat (*Murina eleryi*), also known as the Mekong bat, is a species of common bats first discovered in a forest of northern Vietnam.

Elgon shrew
Elgon shrew
Crocidura elgonius

The Elgon shrew (*Crocidura elgonius*) is a species of mammal in the family Soricidae. It is found in Kenya and Tanzania. Its natural habitat is subtropical or tropical moist montane forests.

Elias's Atlantic spiny rat

Elias's Atlantic spiny rat

Trinomys eliasi

Elias's Atlantic spiny-rat or the Rio de Janeiro spiny rat (*Trinomys eliasi*), is a spiny rat species from South America. It is found in Brazil.

Eligmodontia typus

Eligmodontia typus

Eligmodontia typus

Eligmodontia typus is a species of rodent in the family Cricetidae. It is found in Argentina and possibly also Chile. The northernmost population might represent a distinct species, *E. bolsonensis*, to which the common name highland gerbil mouse...

Elk

Elk

Cervus canadensis

The elk (*Cervus canadensis*) is one of the largest species within the deer family, Cervidae, and one of the largest terrestrial mammals in its native range. The common name elk, used in North America, creates confusion because the larger *Alces*...

Ellerman's tufted-tailed rat

Ellerman's tufted-tailed rat

Eliurus ellermani

Ellerman's tufted-tailed rat (*Eliurus ellermani*) is a species of rodent in the family Nesomyidae. It is endemic to eastern Madagascar. As only two specimens of this species have been observed, questions have been raised regarding whether this...

Elliot's short-tailed shrew

Elliot's short-tailed shrew

Blarina hylophaga

Elliot's short-tailed shrew (*Blarina hylophaga*) is a small, slate grey, short-tailed species of shrew. Its common name comes from Daniel Giraud Elliot, who first described the species in 1899.

Elongated shrew

Elongated shrew

Crocidura elongata

The elongated shrew (*Crocidura elongata*) is a species of mammal in the family Soricidae. It is endemic to the island of Sulawesi in Indonesia. It lives in the forests of central, northern, and eastern Sulawesi from 200 to 2000 meters elevation.

Eloquent horseshoe bat

Eloquent horseshoe bat

Rhinolophus eloquens

The eloquent horseshoe bat (*Rhinolophus eloquens*) is a species of bat in the family Rhinolophidae. It is found in Ethiopia, Kenya, Rwanda, Somalia, South Sudan, Tanzania, and Uganda. Its natural habitats are subtropical or tropical moist lowland...

Elvira rat

Elvira rat

Cremnomys elvira

The Elvira rat (*Cremnomys elvira*) is a critically endangered species of rodent in the family Muridae. The species was first described by Sir John Ellerman in 1946. It is found only in the Eastern Ghats of Tamil Nadu, India.

Embi African mole-rat

Embi African mole-rat

Tachyoryctes spalacinus

The Embi mole-rat (*Tachyoryctes spalacinus*) is a species of rodent in the family Spalacidae endemic to Kenya. Its natural habitats are subtropical or tropical moist montane forests, moist savanna, arable land, pastureland, plantations, and...

Emilia's gracile opossum

Emilia's gracile opossum

Gracilinanus emiliae

Emilia's gracile opossum (*Gracilinanus emiliae*) is an opossum species from South America. It is found in Brazil, Colombia, French Guiana, Surinam.

Emilia's marmoset

Emilia's marmoset

Mico emiliae

The Emilia's marmoset (*Mico emiliae*), also known as Snethlage's marmoset, is a marmoset endemic to Brazil. It is found only in the Brazilian states of Pará and Mato Grosso. It was named to honour German-born Brazilian ornithologist Emilie...

Emilia's short-tailed opossum

Emilia's short-tailed opossum

Monodelphis emiliae

Emilia's short-tailed opossum (*Monodelphis emiliae*) is an opossum species from South America. It is found in Bolivia, Brazil and Peru.

Emily's tuco-tuco

Emily's tuco-tuco

Ctenomys emilianus

Emily's tuco-tuco (*Ctenomys emilianus*) is a species of rodent in the family Ctenomyidae. It is endemic to Argentina.

Emin's gerbil

Emin's gerbil

Taterillus emini

Emin's gerbil or Emin's tateril (*Taterillus emini*) is a species of rodent found in the Democratic Republic of the Congo, Ethiopia, Kenya, Sudan, and Uganda. Its natural habitats are dry savanna, subtropical or tropical dry lowland grassland, and...

Emin's pouched rat

Emin's pouched rat

Cricetomys emini

Emin's pouched rat (*Cricetomys emini*), also known as the African pouched rat, is a large rat of the muroid superfamily. It is related to *Cricetomys gambianus*, the Gambian pouched rat. Both species belong to *Cricetomys*, the genus of the giant...

Emma's giant rat

Emma's giant rat

Uromys emmae

Emma's uromys or Emma's giant rat (*Uromys emmae*) is a species of rodent in the family Muridae. It is only known from Owi Island, Indonesia.

Emperor rat

Emperor rat

Uromys imperator

The emperor rat (*Uromys imperator*) is a large species of rodent in the family Muridae. It is endemic to the island of Guadalcanal in the Solomon Islands. It is classified as critically endangered by the IUCN but may already be extinct.

Emperor Tamarin

Emperor Tamarin

Saguinus imperator

The small Emperor tamarin (*Saguinus imperator*) is probably the most adorable species of tamarin. This New World monkey is euphoniously called due to its long and white-colored mustache, which is very similar to that of Wilhelm II, Emperor of...

Enders's small-eared shrew

Enders's small-eared shrew

Cryptotis endersi

Enders's small-eared shrew (*Cryptotis endersi*) is a species of mammal in the family Soricidae. It is endemic to Panama.

Endo's pipistrelle

Endo's pipistrelle

Pipistrellus endoi

Endo's pipistrelle (*Pipistrellus endoi*) is a species of vesper bat that is endemic to Japan. It is found in temperate forests.

Enggano rat

Enggano rat

Rattus enganus

The Enggano rat (*Rattus enganus*) is a species of rodent in the family Muridae. It is found only on Enggano Island, Indonesia. It is critically endangered.

Epixerus

Epixerus

Epixerus ebii

Epixerus ebii, also known as Ebian's palm squirrel, Temminck's giant squirrel, or the western palm squirrel, is a species of rodent in the family Sciuridae. It is the only species in the genus *Epixerus*, although eastern populations (subspecies...

Epixerus ebii wilsoni

Epixerus ebii wilsoni

Epixerus ebii wilsoni

Epixerus ebii wilsoni is a subspecies of the squirrel *Epixerus ebii* from Cameroon, Republic of the Congo, Equatorial Guinea, and Gabon. It has also been treated as a separate species. Its natural habitat is subtropical or tropical moist lowland...

Eptesicus anatolicus

Eptesicus anatolicus

Eptesicus anatolicus

The Anatolian serotine bat (*Eptesicus anatolicus*) is a species of bat found in the Middle-East, Cyprus and Rhodes Island, Greece.

Eptesicus lobatus

Eptesicus lobatus

Eptesicus lobatus

Eptesicus lobatus is a species of bat of genus *Eptesicus* and family Vespertilionidae.

Eptesicus taddeii

Eptesicus taddeii

Eptesicus taddeii

Taddei's serotine (*Eptesicus taddeii*) is a species of medium-sized bat belonging to the family Vespertilionidae. It is restricted to the Atlantic Forest of southern Brazil. This newly discovered species, which has a total length of up to 120 mm, ...
Equatorial dog-faced bat
Equatorial dog-faced bat
Cabreramops aequatorianus

The equatorial dog-faced bat (*Cabreramops aequatorianus*) is a species of bat in the family Molossidae. It is endemic to Ecuador. They are found in dry, tropical forests. The species is now endangered. The equatorial dog-faced bat feeds on insects.
Equatorial saki
Equatorial saki
Pithecia aequatorialis

The equatorial saki (*Pithecia aequatorialis*), also called the red-bearded saki, is a species of saki monkey, a type of New World monkey. It is found in northeastern Peru and Ecuador. Not much is known about the equatorial saki, its range being ...
Equivalent teeth shrew mole
Equivalent teeth shrew mole
Uropsilus aequodonenia

The equivalent-teeth shrew mole (*Uropsilus aequodonenia*) is a species of mammal in the family Talpidae. It is endemic to Sichuan, China. It is characterized by having nine teeth in the row above and nine teeth in the lower row. The data indicate ...
Equus capensis
Equus capensis
Equus capensis

Equus capensis (*E. capensis*) is an extinct species of zebra that lived during the Pleistocene of South Africa. *E. capensis* was first described from the Cape Town region of South Africa in 1909. *E. capensis* can be estimated to have grown to about ...
Equus fraternus
Equus fraternus
Equus fraternus

Equus fraternus is an extinct species of *Equus*, which was native to North America. Specimens of *E. fraternus* have been found in Florida, Louisiana and Virginia.
Equus namadicus
Equus namadicus
Equus namadicus

Equus namadicus is a prehistoric equid, known from remains dating to the Middle and Late Pleistocene from across the Indian subcontinent, with its last dated records being approximately 29-14,000 years ago. It appears to be closely related to the ...
Equus sivalensis
Equus sivalensis
Equus sivalensis

Equus sivalensis is an extinct equid, discovered in the Siwalik hills. Remains date to 2.6 million years ago, and it is assumed that it was extinct during the last ice age, between 75,000 and 10,000 years ago, as part of the late Pleistocene ...
Equus stenonis
Equus stenonis
Equus stenonis

Equus stenonis or the Stenon zebra, is an extinct species of equine closely related to modern zebras and asses that inhabited Eurasia in the Pleistocene. The African species *Equus koobiforensis* and *E. oldowayensis* are closely related to...

Eremoryzomys

Eremoryzomys

Eremoryzomys polius

Eremoryzomys polius, also known as the gray rice rat or the Marañon oryzomys, is a rodent species in the tribe Oryzomyini of the family Cricetidae. Discovered in 1912 and first described in 1913 by Wilfred Osgood, it was originally placed in...

Erika's tuco-tuco

Erika's tuco-tuco

Ctenomys erikacuellarae

Erika's tuco-tuco (*Ctenomys erikacuellarae*), is a species of tuco-tuco native to Bolivia. Found only in the Cordillera Oriental mountain ranges in the Santa Cruz and Chuquisaca Departments, at elevations of around 810 to 1,800 metres (2,660 to...

Ermine

Ermine

Mustela erminea

The ermine (*Mustela erminea*) is a mustelid native to Eurasia and the northern portions of North America. Introduced in the late 19th century into New Zealand to control rabbits, the ermine has had a devastating effect on native bird populations....

Ernst Mayr's water rat

Ernst Mayr's water rat

Leptomys ernstmayri

Ernst Mayr's water rat (*Leptomys ernstmayri*) is a species of rodent in the family Muridae, named for evolutionary biologist Ernst Mayr. It is found in the Foja Mountains of Papua Province, Indonesia, and in the mountains of northeastern Papua New...

Escalera's bat

Escalera's bat

Myotis escaleraei

Escalera's bat (*Myotis escaleraei*) is a European bat in the genus *Myotis*, found in Spain (including the Balearic Islands), Portugal, and far southern France. Although the species was first named in 1904, it was included in Natterer's bat (*Myotis*...

Espíritu Santo antelope squirrel

Espíritu Santo antelope squirrel

Ammospermophilus insularis

The Espiritu Santo antelope squirrel (*Ammospermophilus insularis*) is a species of antelope squirrel in the family Sciuridae. It is endemic to Mexico, where it is known only from the island of Espiritu Santo in the Gulf of California. The species...

Ethiopian amphibious rat

Ethiopian amphibious rat

Nilopegamys plumbeus

The Ethiopian amphibious rat also known as the Ethiopian water mouse (*Nilopegamys plumbeus*) is an insectivorous and semiaquatic species of rodent in the monotypic genus *Nilopegamys* of the family Muridae. There has only been one known specimen....

Ethiopian big-eared bat

Ethiopian big-eared bat

Plecotus balensis

The Ethiopian long-eared bat or Ethiopian big-eared bat (*Plecotus balensis*) is a recently described species of long-eared bat in the family Vespertilionidae.

Ethiopian dwarf mongoose

Ethiopian dwarf mongoose

Helogale hirtula

The Ethiopian dwarf mongoose (*Helogale hirtula*), also known as the desert dwarf mongoose or Somali dwarf mongoose, is a mongoose native to East Africa, particularly Ethiopia, Kenya, and Somalia. The Ethiopian dwarf mongoose will send out warning...

Ethiopian epauletted fruit bat

Ethiopian epauletted fruit bat

Epomophorus labiatus

The Ethiopian epauletted fruit bat (*Epomophorus labiatus*) is a species of megabat in the family Pteropodidae. It is found in Burundi, Chad, Republic of the Congo, Democratic Republic of the Congo, Eritrea, Ethiopia, Kenya, Malawi, Mozambique,...

Ethiopian forest brush-furred rat

Ethiopian forest brush-furred rat

Lophuromys chrysopus

The Ethiopian forest brush-furred rat or golden-footed brush-furred rat (*Lophuromys chrysopus*), is a species of rodent in the family Muridae. It is endemic to Ethiopia where its natural habitat is subtropical or tropical moist montane forests....

Ethiopian hare

Ethiopian hare

Lepus fagani

The Ethiopian hare (*Lepus fagani*) is a species of mammal in the family Leporidae. It was first described in 1903, by the British mammalogist Oldfield Thomas. The dorsal pelage is brownish buff, and is finely grizzled with black. The ventral...

Ethiopian highland hare

Ethiopian highland hare

Lepus starcki

The Ethiopian highland hare (*Lepus starcki*) or Starck's hare is a medium-sized species of mammal in the rabbit and hare family, Leporidae. Its dorsal pelage is grizzled, buff white and spotted and streaked with black, while its belly fur is pure...

Ethiopian large-eared roundleaf bat

Ethiopian large-eared roundleaf bat

Hipposideros megalotis

The Ethiopian large-eared roundleaf bat (*Hipposideros megalotis*) is a species of bat in the family Hipposideridae. It is found in Djibouti, Eritrea, Ethiopia, Kenya, Saudi Arabia, and Somalia. Its natural habitats are dry savanna, subtropical or...

Ethiopian mylomys

Ethiopian mylomys

Mylomys rex

The Ethiopian mylomys (*Mylomys rex*) is a species of rodent in the family Muridae. It is only found in Ethiopia.

Ethiopian narrow-headed rat

Ethiopian narrow-headed rat

Stenocephalemys albicaudata

The Ethiopian narrow-headed rat (*Stenocephalemys albicaudata*) is a species of rodent in the family Muridae. It is found only in Ethiopia. Its natural habitats are subtropical or tropical high-altitude shrubland and subtropical or tropical...
Ethiopian striped mouse
Ethiopian striped mouse
Muriculus imberbis

The Ethiopian striped mouse or striped-back mouse (*Mus imberbis*) is a species of rodent in the family Muridae. It is only found in Ethiopia. It was formerly classified in the monotypic genus *Muriculus*, but phylogenetic evidence supports *Muriculus*...
Ethiopian thicket rat
Ethiopian thicket rat
Grammomys minnae

The Ethiopian thicket rat (*Grammomys minnae*) is a species of rodent in the family Muridae. It is found only in Ethiopia. Its natural habitat is subtropical or tropical dry shrubland. It is threatened by habitat loss.
Ethiopian white-footed mouse
Ethiopian white-footed mouse
Stenocephalemys albipes

The Ethiopian white-footed mouse or white-footed stenocephalemys (*Stenocephalemys albipes*) is a species of rodent in the family Muridae. It lives in Ethiopia and Eritrea. Its natural habitats are tropical moist montane forest and tropical...
Ethiopian Wolf
Ethiopian Wolf
Canis simensis

The Ethiopian wolf (*Canis simensis*) is a canine native to the Ethiopian Highlands. In southeastern Ethiopia, it is also known as the Horse jackal. Unlike most large canids, which are widespread, generalist feeders, the Ethiopian wolf is a highly...
Ethiopian woolly bat
Ethiopian woolly bat
Kerivoula eriophora

The Ethiopian woolly bat (*Kerivoula eriophora*) is a species of vesper bat in the family Vespertilionidae. It is found only in Ethiopia.
Etruscan Shrew
Etruscan Shrew
Suncus etruscus

The Etruscan shrew (*Suncus etruscus*) is the smallest known extant mammal by mass, weighing only about 1.8 g (0.063 oz) on average. It is characterized by very rapid movements and a fast metabolism, eating about 1.5-2 times its own body weight per...
Eumops delticus
Eumops delticus
Eumops delticus

Eumops delticus is a species of free-tailed bat found in South America.
Eumops ferox
Eumops ferox
Eumops ferox

Eumops ferox, the fierce bonneted bat or the chestnut mastiff bat, is a species of free-tailed bat found in the Caribbean and Mexico. Until recently, it was synonymous with Wagner's bonneted bat.
Eumops nanus

Eumops nanus
Eumops nanus

Eumops nanus is a species of bat found in Central and South America.

Eumops wilsoni
Eumops wilsoni
Eumops wilsoni

Eumops wilsoni is a species of bat native to Ecuador and Peru. The bat has a distinct karyotype, sequence of the mitochondrial cytochrome-b gene, and other distinct genetic markers that distinguish it from closely related bats such as Eumops...

Euphrates jerboa
Euphrates jerboa
Allactaga euphratica

The Euphrates jerboa (*Allactaga euphratica*) is a rodent of the family Dipodidae and genus *Allactaga*. They are characteristically known as hopping rodents. They have been found in Pakistan, Afghanistan, Iran, Iraq, Jordan, Kuwait, Saudi Arabia,...

Eupleres major
Eupleres major
Eupleres major

The western falanouc (*Eupleres major*) is a rare mongoose-like mammal endemic to Madagascar. Until recently, the eastern falanouc (*Eupleres goudotii*) was the only recognized species in the genus *Eupleres*. In 2010, Goodman and Helgen provided...

Eurasian Beaver
Eurasian Beaver
Castor fiber

The Eurasian beaver (*Castor fiber*) is a beaver species that was once widespread in Eurasia but was hunted to near-extinction for both its fur and castoreum. At the turn of the 20th century, only about 1,200 beavers survived in eight relict...

Eurasian brown bear
Eurasian brown bear
Ursus arctos arctos

The Eurasian brown bear (*Ursus arctos arctos*) is one of the most common subspecies of the brown bear, and is found in much of Eurasia. It is also known as the European brown bear, common brown bear, common bear, and colloquially by many other...

Eurasian Harvest Mouse
Eurasian Harvest Mouse
Micromys minutus

The Eurasian harvest mouse (*Micromys minutus*) is a small rodent native to Europe and Asia. It is typically found in fields of cereal crops, such as wheat and oats, in reed beds, and in other tall ground vegetation, such as long grass and...

Eurasian least shrew
Eurasian least shrew
Sorex minutissimus

The Eurasian least shrew (*Sorex minutissimus*), also called the lesser pygmy shrew, is the second-smallest mammal by mass after the Etruscan shrew.

Eurasian Lynx
Eurasian Lynx
Lynx lynx

The Eurasian lynx (*Lynx lynx*) is a medium-sized wild cat. It is Europe's third-largest predator after Brown bears and wolves and is the biggest of the species of lynx. The Eurasian lynx lives in forests and despite its wide distribution, it is...

Eurasian Pygmy Shrew

Eurasian Pygmy Shrew

Sorex minutus

The Eurasian pygmy shrew (*Sorex minutus*), often known simply as the pygmy shrew, is a widespread shrew of the northern Palearctic.

Eurasian Water Shrew

Eurasian Water Shrew

Neomys fodiens

The Eurasian water shrew (*Neomys fodiens*), known in the United Kingdom as the water shrew, is a relatively large shrew, up to 10 cm (4 in) long, with a tail up to three-quarters as long again. It has short, dark fur, often with a few white...

Eurasian wolf

Eurasian wolf

Canis lupus lupus

The Eurasian wolf (*Canis lupus lupus*), also known as the common wolf, is a subspecies of grey wolf native to Europe and Asia. It was once widespread throughout Eurasia prior to the Middle Ages. Aside from an extensive paleontological record,...

European ass

European ass

Equus hemionus hydruntinus

The European wild ass (*Equus hemionus hydruntinus*) or hydruntine is an extinct equine from the Middle Pleistocene to Late Holocene of Western Eurasia. It appeared first in the fossil record around 350–300,000 years before present. In the late...

European Badger

European Badger

Meles meles

The European badger (*Meles meles*) is a mustelid native to almost all of Europe. It has a wide range and a large stable population size and is thought to be increasing in some regions. In Europe, where no other badger species commonly occurs, it...

European free-tailed bat

European free-tailed bat

Tadarida teniotis

The European free-tailed bat (*Tadarida teniotis*, sometimes given as *Tadarida insignis*) is a species of free-tailed bat found in the Old World. Other common names include the bulldog bat and the mastiff bat because of the presence of wrinkling on...

European Ground Squirrel

European Ground Squirrel

Spermophilus citellus

The European ground squirrel (*Spermophilus citellus*), also known as the European souslik, is a species from the squirrel family, *Sciuridae*. It is among the few European species in the genus *Spermophilus*. Like all squirrels, it is a member of the...

European Hamster

European Hamster

Cricetus cricetus

The European hamster (*Cricetus cricetus*), also known as the Eurasian hamster, black-bellied hamster or common hamster, is the only species of hamster in the genus *Cricetus*. It is native to grassland and similar habitats in a large part of...

European Hare
European Hare
Lepus europaeus

The European hare (*Lepus europaeus*) is a species of hare native to Europe and parts of Asia. It is among the largest hare species and is adapted to temperate, open country. Generally nocturnal and shy in nature, hares change their behavior in the...

European Hedgehog
European Hedgehog
Erinaceus europaeus

The European hedgehog (*Erinaceus europaeus*) is a hedgehog species native to Europe. It is a generally common and widely distributed species that can survive across a wide range of habitat types. It is a well-known species, and a favorite in...

European Jackal
European Jackal
Canis aureus moreoticus

The European jackal (*Canis aureus moreoticus*) is a subspecies of the Golden jackal. It was first described by French naturalist Isidore Geoffroy Saint-Hilaire during the Morea expedition.

European Mink
European Mink
Mustela lutreola

The European minks (*Mustela lutreola*) are small carnivores once widely distributed over almost the whole of the European continent and currently surviving in a few enclaves as fragmented populations. The huge change in its numbers and...

European Mole
European Mole
Talpa europaea

The European mole (*Talpa europaea*) is a small fossorial mammal that lives across most of Europe. It is also known as the Common mole and the Northern mole. This mole lives in a tunnel system, which it constantly extends. It uses these tunnels to...

European mouflon
European mouflon
Ovis aries musimon

The European mouflon (*Ovis aries musimon*) is a feral subspecies of the primitive domestic sheep. It was originally found only on the Mediterranean islands of Corsica and Sardinia, but has since been introduced into many other regions of Europe....

European Otter
European Otter
Lutra lutra

The Eurasian otter (*Lutra lutra*) is a semiaquatic mammal native to Eurasia. It is the most widely distributed member of the otter subfamily (*Lutrinae*) of the weasel family (*Mustelidae*). It is found in the waterways and coasts of Europe, many...

European pine vole
European pine vole
Microtus subterraneus

The European pine vole (*Microtus subterraneus*), also known as the common pine vole, is a species of rodent in the family Cricetidae. It is native to much of Europe and parts of Asia.

European Polecat

European Polecat

Mustela putorius

The European polecat (*Mustela putorius*) is a species of mustelid native to western Eurasia and North Africa. It originated in Western Europe during the Middle Pleistocene, with its closest living relatives being the Steppe polecat, the...

European Rabbit

European Rabbit

Oryctolagus cuniculus

The European rabbit (*Oryctolagus cuniculus*) is a small mammal native to southwestern Europe and northwest Africa. It is known as an invasive species because it has been introduced to countries on all continents with the exception of Antarctica,...

European snow vole

European snow vole

Chionomys nivalis

The European snow vole or snow vole (*Chionomys nivalis*, previously *Arvicola nivalis* and *Microtus nivalis*) is a species of rodent in the family Cricetidae. It has dense, pale grey fur and a pale-coloured tail and can reach about 14 cm (5.5 in)...

European Water Vole

European Water Vole

Arvicola amphibius

European water voles > (*Arvicola amphibius*) are semiaquatic rodents. They are often informally called water rats, though they only superficially resemble a true rat. Their dark color allows them to blend in well in the densely vegetated areas and...

European Wildcat

European Wildcat

Felis silvestris silvestris

The European wildcat (*Felis silvestris silvestris*) is a small wildcat subspecies found in Europe. Wildcats and the other members of the cat family had a common ancestor about 10-15 million years ago. The European wildcat evolved during the...

Euroscaptor subanura

Euroscaptor subanura

Euroscaptor subanura

The Vietnamese mole (*Euroscaptor subanura*) is a species of talpine mole found in Vietnam. The species was first identified in December 2008 in the foothills of the Tam Dao mountains in northern Vietnam. At the time of the formal description of...

Euryoryzomys emmonsae

Euryoryzomys emmonsae

Euryoryzomys emmonsae

Euryoryzomys emmonsae, also known as Emmons' rice rat or Emmons' oryzomys, is a rodent from the Amazon rainforest of Brazil in the genus *Euryoryzomys* of the family Cricetidae. Initially misidentified as *E. macconnelli* or *E. nitidus*, it was...

Euryoryzomys lamia

Euryoryzomys lamia

Euryoryzomys lamia

Euryoryzomys lamia, also known as the buffy-sided oryzomys or monster rice rat, is

a species of rodent in the family Cricetidae. It is found only in central Brazil, where it lives in forest enclaves within the cerrado. The species' known...

Euryoryzomys legatus
Euryoryzomys legatus
Euryoryzomys legatus

Euryoryzomys legatus, also known as the Tarija oryzomys or big-headed rice rat, is a species of rodent in the family Cricetidae. It now belongs to the genus Euryoryzomys, having previously been placed in Oryzomys. It is found in the eastern Andes...

Euryoryzomys macconnelli
Euryoryzomys macconnelli
Euryoryzomys macconnelli

Euryoryzomys macconnelli, also known as MacConnell's rice rat or MacConnell's oryzomys, is a rodent species from South America. It is found in Brazil, Colombia, Ecuador, French Guiana, Guyana, Peru, Suriname and Venezuela, where it lives in...

Euryoryzomys nitidus
Euryoryzomys nitidus
Euryoryzomys nitidus

Euryoryzomys nitidus, also known as the elegant oryzomys or elegant rice rat, is a rodent species in the family Cricetidae. Previously it was known as Oryzomys nitidus, but it is not closely related to Oryzomys as that genus is now constructed....

Euryoryzomys russatus
Euryoryzomys russatus
Euryoryzomys russatus

Euryoryzomys russatus, also known as the russet oryzomys, russet rice rat, or big-headed rice rat, is a species of rodent in the family Cricetidae. It is a member of the genus Euryoryzomys, which was split off from Oryzomys in 2006. It was first...

Eva's desert mouse
Eva's desert mouse
Peromyscus eva

Eva's desert mouse (*Peromyscus eva*) is a species of rodent in the genus *Peromyscus* of the family Cricetidae found only in the Baja California peninsula of Mexico.

Evening Bat
Evening Bat
Nycticeius humeralis

The Evening bat (*Nycticeius humeralis*) is a small member of a widely distributed bat family. The species owes its name to the genus *Vespertilio*, which takes its name from a word for bat, *vespertilio*, derived from the Latin term *vesper* meaning...

Everglades short-tailed shrew
Everglades short-tailed shrew
Blarina peninsulae

The Everglades short-tailed shrew, (*Blarina peninsulae*) is a species of shrew in the genus *Blarina*. It is endemic to Florida.

Eversmann's hamster
Eversmann's hamster
Allocricetulus eversmanni

Eversmann's hamster (*Allocricetulus eversmanni*) is a species of hamster in the family Cricetidae, and is one of two members of the genus *Allocricetulus*. It has been named after the Russian naturalist, zoologist, and explorer Eduard

Friedrich...
Evorsk vole
Evorsk vole
Microtus evoronensis

The Evorsk vole (*Microtus evoronensis*) is a species of rodent in the family Cricetidae. It is found only in Russia. They are stout rodents with short ears, legs and tails. These animals live in grassy areas where they eat green vegetation such as...

Ezo Red Fox
Ezo Red Fox
Vulpes vulpes schrencki

The Ezo red fox (*Vulpes vulpes schrencki*) is a subspecies of Red fox. Ezo is a Japanese word meaning "foreigner" and referred to the historical lands of the Ainu people to the north of Honshu, which the Japanese named Ezo-chi. In the Ainu...

Fair brocket
Fair brocket
Mazama tienhoveni

The fair brocket (*Mazama tienhoveni*), also known as the white brocket deer or veado branco to locals, is a possible species of brocket deer from the south-central Amazon near the Aripuanã River in Brazil. The coloration of the fair brocket is...

Fallow Deer
Fallow Deer
Dama dama

Fallow deer (*Dama dama*) are amongst the prettiest deer to be seen in Europe. They are historically native to Turkey and possibly the Italian Peninsula, the Balkan Peninsula, and the island of Rhodes in Europe. Prehistorically native to and...

False canyon mouse
False canyon mouse
Peromyscus pseudocrinitus

The false canyon mouse or Coronados deer mouse, (*Peromyscus pseudocrinitus*) is a species of rodent in the family Cricetidae. It is known only from Coronados Island, a small island in the Gulf of California, part of Baja California Sur, Mexico....

False Killer Whale
False Killer Whale
Pseudorca crassidens

The False killer whale (*Pseudorca crassidens*) is a species of oceanic dolphin that is the only extant representative of the genus *Pseudorca*. It is found in oceans worldwide but mainly frequents tropical regions. The name "false killer whale"...

False potto
False potto
Pseudopotto martini

The false potto (*Pseudopotto martini*) is a lorisoid primate of uncertain taxonomic status found in Africa. Anthropologist Jeffrey H. Schwartz named it in 1996 as the only species of the genus *Pseudopotto* on the basis of two specimens (consisting...

False serotine bat
False serotine bat
Hesperoptenus doriae

The false serotine bat (*Hesperoptenus doriae*) is a species of vesper bat. It is found only in Malaysia. It is threatened by habitat loss.
False water rat

False water rat
Xeromys myoides

Xeromys myoides, also known as the water mouse, marine mouse, mangrove mouse, false water rat, manngay and yirrkoo, is a species of rodent native to waterways of Australia and Papua New Guinea.

False zokor
False zokor
Myospalax aspalax

The false zokor (*Myospalax aspalax*) is a species of rodent in the family Spalacidae. It is found in Mongolia, China and Russia.

Famatina chinchilla rat
Famatina chinchilla rat
Abrocoma famatina

The Famatina chinchilla rat (*Abrocoma famatina*) is a species of chinchilla rats in the family Abrocomidae found only in Argentina.

Famatina tuco-tuco
Famatina tuco-tuco
Ctenomys famosus

The Famatina tuco-tuco (*Ctenomys famosus*) is a species of rodent in the family Ctenomyidae. It is endemic to northern Argentina. The common name of the species comes from the municipality, department and mountain range of the same name at the...

Fancy mouse
Fancy mouse
Mus musculus domestica

A fancy mouse (fancy means 'hobby' in this context) is a domesticated form of the house mouse (*Mus musculus*), one of many mouse species, usually kept as a type of pocket pet. Fancy mice have also been specially bred for exhibiting, with shows...

Fancy rat
Fancy rat
Rattus norvegicus domestica

The fancy rat (*Rattus norvegicus domestica*) is the domesticated form of *Rattus norvegicus*, the brown rat, and the most common species of rat kept as a pet. The name fancy rat derives from the idea of animal fancy (the promotion of domesticated...

Far Eastern myotis
Far Eastern myotis
Myotis bombinus

The Far Eastern myotis or bombinus bat (*Myotis bombinus*) is a species of mouse-eared bat found in East Asia. It is widespread but uncommon across the Korean Peninsula, and spends the winter hibernating in caves.

Fardoulis's blossom bat
Fardoulis's blossom bat
Melonycteris fardoulisi

Fardoulis's blossom bat (*Melonycteris fardoulisi*) is a species of bat in the Pteropodidae family. It is endemic to the Solomon Islands. Its natural habitat is subtropical or tropical moist lowland forests. It is threatened by habitat loss.

Fat mouse
Fat mouse
Steatomys pratensis

The fat mouse (*Steatomys pratensis*) is a species of rodent in the family Nesomyidae. It is found in Angola, Botswana, Cameroon, Democratic Republic of the Congo, Malawi, Mozambique, Namibia, South Africa, Tanzania, Zambia, and Zimbabwe. Its...

Fat Sand Rat

Fat Sand Rat

Psammomys obesus

The Fat sand rat is a gerbil species, endemic to desert areas, where it maintains a usual rodent diet, primarily consisting of leaves. This species was accidentally found in Egypt by the US Naval Medical Research Unit. Despite the common name,...

Fat-nosed spiny rat

Fat-nosed spiny rat

Maxomys inflatus

The fat-nosed spiny rat (*Maxomys inflatus*) is a species of rodent in the family Muridae. It is found only in Indonesia.

Fat-Tailed Dunnart

Fat-Tailed Dunnart

Sminthopsis crassicaudata

The fat-tailed dunnart (*Sminthopsis crassicaudata*) is a species of mouse-like marsupial of the Dasyuridae, the family that includes the little red kaluta, quolls, and the Tasmanian devil. It has an average body length of 60–90 millimeters...

Fat-Tailed Dwarf Lemur

Fat-Tailed Dwarf Lemur

Cheirogaleus medius

The fat-tailed dwarf lemur (*Cheirogaleus medius*), also known as the lesser dwarf lemur, western fat-tailed dwarf lemur, or spiny forest dwarf lemur, is endemic to Madagascar.

Fat-tailed false antechinus

Fat-tailed false antechinus

Pseudantechinus macdonnellensis

The fat-tailed false antechinus (*Pseudantechinus macdonnellensis*), also called the fat-tailed pseudantechinus and red-eared antechinus, is a member of the order Dasyuromorphia. It is an inhabitant of western and central Australia.

Fat-Tailed Gerbil

Fat-Tailed Gerbil

Pachyuromys duprasi

The Fat-tailed gerbil (*Pachyuromys duprasi*) is a medium-sized rodent belonging to subfamily Gerbillinae. It is the most docile species of the gerbil subfamily. Fat-tailed gerbils are frequently kept as pets.

Father Basilio's striped mouse

Father Basilio's striped mouse

Hybomys basillii

Father Basilio's striped mouse or Bioko hybomys (*Hybomys basillii*) is a species of rodent in the family Muridae. It is found only in Equatorial Guinea. Its natural habitat is subtropical or tropical moist montane forest. It is threatened by habitat...

Fawn antechinus

Fawn antechinus

Antechinus bellus

The fawn antechinus (*Antechinus bellus*) is a species of small carnivorous

marsupial found in northern Australia. It is the only Antechinus to be found in the Northern Territory and has a patchy, restricted range.

Fawn hopping mouse
Fawn hopping mouse
Notomys cervinus

The fawn hopping mouse (*Notomys cervinus*) is a rodent native to the central Australian desert. Like all hopping mice it has strong front teeth, a long tail, dark eyes, big ears, well-developed haunches and very long, narrow hind feet. It weighs...

Fawn leaf-nosed bat
Fawn leaf-nosed bat
Hipposideros cervinus

The fawn leaf-nosed bat - *Hipposideros cervinus* - is a species of bat in the family Hipposideridae found in Australia, Indonesia, Malaysia, the Philippines and Vanuatu.

Fawn-colored mouse
Fawn-colored mouse
Mus cervicolor

The fawn-colored mouse (*Mus cervicolor*) is a species of rodent in the family Muridae. It is found in Cambodia, India, and possibly Indonesia, Laos, Myanmar, Nepal, Thailand, and Vietnam.

Fawn-footed mosaic-tailed rat
Fawn-footed mosaic-tailed rat
Melomys cervinipes

The fawn-footed mosaic-tailed rat, or fawn-footed melomys (*Melomys cervinipes*) is a species of rodent in the family Muridae. It is found along eastern coastal regions of Australia. The upper parts of this species are a sandy-brown colour, the...

Fea's muntjac
Fea's muntjac
Muntiacus feae

The Fea's muntjac or Tenasserim muntjac (*Muntiacus feae*) is a rare species of muntjac native to southern Myanmar and Thailand. It is a similar size to the common muntjac (adult weight is 18 - 21 kg (40 - 46 lb)). It is diurnal and solitary,...

Fea's tree rat
Fea's tree rat
Chiromyscus chiropus

Fea's tree rat or Indochinese chiromyscus (*Chiromyscus chiropus*) is a species of rodent in the family Muridae. It is found in Yunnan (China), eastern Myanmar, northern Thailand, Laos, and Vietnam.

Feather-tailed possum
Feather-tailed possum
Distoechurus pennatus

The feather-tailed possum (*Distoechurus pennatus*) is a species of marsupial in the family Acrobatidae. It is found in West Papua, Indonesia and Papua New Guinea. It is not to be confused with the feathertail glider, the only other species in the...

Feathertail Glider
Feathertail Glider
Acrobates pygmaeus

The feathertail glider (*Acrobates pygmaeus*), also known as the pygmy gliding possum, pygmy glider, pygmy phalanger, flying phalanger and flying mouse, is a

species of marsupial native to eastern Australia. It is the world's smallest gliding...

Fecund vesper mouse
Fecund vesper mouse
Calomys fecundus

The fecund vesper mouse (*Calomys fecundus*) is a species of rodent in the family Cricetidae. It is known only from Bolivia, where it is found in the yungas and Tucumano boliviano forest ecoregions at elevations from 600 to 2700 m. It is regarded...

Felis attica
Felis attica
Felis attica

Felis attica is an extinct cat, of which the first fossil skull was excavated near Pikermi in Attica, Greece. Fossils were also excavated near the Moldovan city of Taraclia. It was also discovered in Maragheh, northwestern Iran. *F. attica* was...

Felis chaus affinis
Felis chaus affinis
Felis chaus affinis

Felis chaus affinis is a jungle cat subspecies. It was described by British zoologist John Edward Gray in 1830 based on an illustration by Thomas Hardwicke.

Felis chaus chaus
Felis chaus chaus
Felis chaus chaus

Felis chaus chaus is the nominate subspecies of the jungle cat. The Baltic-German naturalist Johann Anton Gldenstdt was the first scientist who observed a jungle cat in the southern frontier of the Russian empire during his travels in 1768-1775...

Felis chaus fulvidina
Felis chaus fulvidina
Felis chaus fulvidina

Felis chaus fulvidina is a jungle cat subspecies. The mammal collector of the Natural History Museum Oldfield Thomas described the first jungle cat from Indochina in 1928. It occurs mainly in deciduous forests rich in dipterocarp trees. Since the...

Felis margarita margarita
Felis margarita margarita
Felis margarita margarita

Felis margarita margarita, sometimes called the Saharan sand cat, is a subspecies of the sand cat native to the Sahara.

Felis margarita thinobia
Felis margarita thinobia
Felis margarita thinobia

Felis margarita thinobia, known as the Turkestan sand cat, Arabian sand cat, and Pakistan sand cat, is a sand cat subspecies native to deserts in the Arabian Peninsula, Iran, Pakistan and Central Asia.

Felou gundi
Felou gundi
Felovia vae

The felou gundi (*Felovia vae*) is a species of rodent in the family Ctenodactylidae. It is monotypic within the genus *Felovia*. It is found in Mali, Mauritania, and Senegal. Its natural habitats are dry savanna, subtropical or

tropical dry...
Felten's myotis
Felten's myotis
Myotis punicus

The Felten's myotis (*Myotis punicus*) is a species of vesper bat.

Felten's vole
Felten's vole
Microtus felteni

Felten's vole (*Microtus felteni*) is a species of rodent in the family Cricetidae. It is found in Albania, Greece, North Macedonia, and Serbia.

Fennec Fox
Fennec Fox
Vulpes zerda

The Fennec fox (*Vulpes zerda*) is a small crepuscular fox native to the deserts of North Africa. It is the smallest species of fox. Its coat, ears, and kidney functions have adapted to the desert environment with high temperatures and little...

Fernandez's sword-nosed bat
Fernandez's sword-nosed bat
Lonchorhina fernandezi

Fernandez's sword-nosed bat (*Lonchorhina fernandezi*) is a species of bat in the family Phyllostomidae. It is the smallest species of the *Lonchorhina* genus. It is endemic to Venezuela. In 2013, Bat Conservation International listed this species...

Ferreira's fish-eating rat
Ferreira's fish-eating rat
Neusticomys ferreirai

Ferreira's fish-eating rat (*Neusticomys ferreirai*) is a species of rodent in the family Cricetidae. It is known only from primary lowland tropical rainforest. It is named after 18th-century Brazilian naturalist Alexandre Rodrigues Ferreira.

Ferreira's spiny tree-rat
Ferreira's spiny tree-rat
Mesomys hispidus

Ferreira's spiny tree-rat, *Mesomys hispidus*, is a spiny rat species from South America. It is found in Bolivia, Brazil, Colombia, Ecuador, French Guiana, Guyana, Peru, Suriname and Venezuela.

Field Vole
Field Vole
Microtus agrestis

The short-tailed field vole, short-tailed vole, or simply field vole (*Microtus agrestis*) is a grey-brown vole, around 10 cm in length, with a short tail. It is one of the most common mammals in Europe, with a range extending from the Atlantic...

Fierce roundleaf bat
Fierce roundleaf bat
Hipposideros dinops

The fierce roundleaf bat (*Hipposideros dinops*) is a species of bat in the family Hipposideridae. It is found in Indonesia, Papua New Guinea, and the Solomon Islands.

Fiery spiny mouse
Fiery spiny mouse

Acomys ignitus

The fiery spiny mouse (*Acomys ignitus*) is a species of rodent in the family Muridae found in Kenya and Tanzania. Its natural habitats are dry savanna and rocky areas. It may be found as a commensal in human habitations.

Fiery squirrel

Fiery squirrel

Sciurus flammifer

The fiery squirrel (*Sciurus flammifer*) is a rodent in the family Sciuridae. The taxon is endemic to the area south of the Orinoco River in the state of Bolívar, Venezuela.

Fijian mastiff bat

Fijian mastiff bat

Chaerephon bregullae

The Fijian mastiff bat (*Chaerephon bregullae*), also known as the Fijian free-tailed bat, is a species of bat in the family Molossidae. It is found in Fiji and Vanuatu. In 2013, Bat Conservation International listed this species as one of the 35...

Fijian monkey-faced bat

Fijian monkey-faced bat

Mirimiri acrodonta

The Fijian monkey-faced bat (*Mirimiri acrodonta*) Also known as Fijian flying fox or Fijian flying monkey, is a megabat endemic to Fiji. It was discovered in old-growth cloud forest on Des Vœux Peak, the second highest mountain peak (1,195 m) on...

Fin Whale

Fin Whale

Balaenoptera physalus

The Fin whale (*Balaenoptera physalus*) is a cetacean belonging to the parvorder of baleen whales. It is the second-longest species of cetacean on Earth after the Blue whale. American naturalist Roy Chapman Andrews called the fin whale "the...

Findley's myotis

Findley's myotis

Myotis findleyi

Findley's myotis (*Myotis findleyi*) is a species of vesper bat. It is found only on the Tres Mariás Islands off the west coast of Mexico.

Finlayson's cave bat

Finlayson's cave bat

Vespadelus finlaysoni

Finlayson's cave bat (*Vespadelus finlaysoni*) is a species of vespertilionid bat found only in Australia.

Finlayson's squirrel

Finlayson's squirrel

Callosciurus finlaysonii

Finlayson's squirrel or the variable squirrel (*Callosciurus finlaysonii*, sometimes misspelled *C. finlaysoni*) is a species of rodent in the family Sciuridae. It is found in Cambodia, Laos, Myanmar, Thailand, and Vietnam. The species occurs in a...

Finnish Forest Reindeer

Finnish Forest Reindeer

Rangifer tarandus fennicus

The Finnish forest reindeer (*Rangifer tarandus fennicus*) is a rare subspecies of

the reindeer native to Finland and northwestern Russia. It is also known as Eurasian or European forest reindeer.
Fire-bellied brush-furred rat
Fire-bellied brush-furred rat
Lophuromys nudicaudus

The fire-bellied brush-furred rat (*Lophuromys nudicaudus*) is a species of rodent in the family Muridae. It is found in Cameroon, Central African Republic, the Republic of the Congo, the Democratic Republic of the Congo, Equatorial Guinea, and...
Fire-footed rope squirrel
Fire-footed rope squirrel
Funisciurus pyrropus

The fire-footed rope squirrel (*Funisciurus pyrropus*) is a species of rodent in the family Sciuridae. It is found in Angola, Burundi, Cameroon, Republic of the Congo, Democratic Republic of the Congo, Ivory Coast, Equatorial Guinea, Gabon, Gambia,...
Fischer's guara
Fischer's guara
Euryzygomatomys spinosus

Fischer's guara (*Euryzygomatomys spinosus*), is a South American spiny rat species found in Argentina, Brazil and Paraguay. It is one of only two species in the genus *Euryzygomatomys*. Its karyotype has $2n=46$ and $FN=88$.
Fischer's little fruit bat
Fischer's little fruit bat
Rhinophylla fischerae

Fischer's little fruit bat (*Rhinophylla fischerae*) is a bat species from South America. It is found in Brazil, Colombia, Ecuador, Peru and Venezuela.
Fischer's pygmy fruit bat
Fischer's pygmy fruit bat
Haplonycteris fischeri

The Fischer's pygmy fruit bat or Philippine pygmy fruit bat (*Haplonycteris fischeri*) is a species of megabat in the family Pteropodidae. It is monotypic within the genus *Haplonycteris*. It is endemic to the Philippines. Its natural habitat is...
Fischer's shrew
Fischer's shrew
Crocidura fischeri

Fischer's shrew (*Crocidura fischeri*) is a species of mammal in the family Soricidae. It is found in Kenya and Tanzania. Its natural habitat is dry savanna. It is threatened by habitat loss.
Fisher
Fisher
Pekania pennanti

The fisher (*Pekania pennanti*) is a small, carnivorous mammal native to North America. It is a member of the mustelid family (commonly referred to as the weasel family), and is sometimes misleadingly referred to as a fisher cat, although it is not...
Fishing Cat
Fishing Cat
Prionailurus viverrinus

The Fishing cat (*Prionailurus viverrinus*) is a medium-sized wild cat of South and Southeast Asia. It is a threatened species whose population has declined severely over the last decade. Its scientific name comes from its viverrine or civet-like...

Five-toed pygmy jerboa
Five-toed pygmy jerboa
Cardiocranius paradoxus

The five-toed pygmy jerboa (*Cardiocranius paradoxus*) is a species of rodent in the family Dipodidae. It is monotypic within the genus *Cardiocranius*. It is found in China, Kazakhstan, and Mongolia. Its natural habitat is temperate desert. This...

Flamarion's tuco-tuco
Flamarion's tuco-tuco
Ctenomys flamarioni

Flamarion's tuco-tuco or the tuco-tuco of the dunes (*Ctenomys flamarioni*) is a South American rodent species of the family Ctenomyidae. Its karyotype has $2n = 48$ and $FN = 50-78$. It is found in coastal dunes of Rio Grande do Sul state, southern...

Flat-faced fruit-eating bat
Flat-faced fruit-eating bat
Artibeus planirostris

The flat-faced fruit-eating bat (*Artibeus planirostris*) is a South American species of bat in the family Phyllostomidae. It is sometimes considered a subspecies of the Jamaican fruit bat, but can be distinguished by its larger size, the presence...

Flat-haired mouse
Flat-haired mouse
Mus platythrix

The flat-haired mouse (*Mus platythrix*) is a species of rodent in the family Muridae. It is found only in India, where it is widely distributed throughout South India and central India.

Flat-Headed Cat
Flat-Headed Cat
Prionailurus planiceps

The Flat-headed cat (*Prionailurus planiceps*) is a small wild cat native to Southeast Asia. It was first described in 1827 based on a zoological specimen collected in Sumatra. This rare cat inhabits wetlands and preys mainly on fish, frogs, and...

Flat-headed kusimanse
Flat-headed kusimanse
Crossarchus platycephalus

The flat-headed kusimanse (*Crossarchus platycephalus*) is a mongoose species endemic to West African rainforests. It is listed as Least Concern on the IUCN Red List since 2008.

Flat-headed myotis
Flat-headed myotis
Myotis planiceps

The flat-headed myotis (*Myotis planiceps*) is a species of vesper bat. It is endemic to Mexico where it is found in certain montane forests in the Sierra Madre Oriental in the northeast of the country. Once thought to be extinct, this bat was...

Flat-headed shrew
Flat-headed shrew
Crocidura planiceps

The flat-headed shrew (*Crocidura planiceps*) is a species of mammal in the family Soricidae. It is found in Democratic Republic of the Congo, Ethiopia, Nigeria, and

Uganda. The flat-headed shrew is listed as Data Deficient (DD), inadequate...
Flat-headed vole
Flat-headed vole
Alticola strelzowi

The flat-headed vole (*Alticola strelzowi*) is a species of rodent in the family Cricetidae. It is found in China, Kazakhstan, Mongolia, and Russian Federation.
Flat-skulled shrew
Flat-skulled shrew
Sorex roboratus

The flat-skulled shrew (*Sorex roboratus*) is a species of mammal in the family Soricidae. It is found in Mongolia, China and Russia.
Fleurete's sportive lemur
Fleurete's sportive lemur
Lepilemur fleuretae

Fleurete's sportive lemur (*Lepilemur fleuretae*), or the Andohahela sportive lemur, is a sportive lemur endemic to Madagascar. It is a medium-sized sportive lemur with a total length of about 58 to 67 cm (23 to 26 in), of which about 30 cm...
Flores Giant Rat
Flores Giant Rat
Papagomys armandvillei

The Flores giant rat (*Papagomys armandvillei*) is a rodent of the family Muridae that occurs on the island of Flores in Indonesia. It has been recorded in Rutong Protection Forest. The species is found in primary, secondary and disturbed forest...
Flores long-nosed rat
Flores long-nosed rat
Paulamys naso

The Flores long-nosed rat, also known as Paula's long-nosed rat, (*Paulamys naso*) is a species of rodent endemic to Flores Island, Indonesia. This species was first described from subfossil fragments collected in the 1950s by Theodor Verhoeven,...
Flores shrew
Flores shrew
Suncus mertensi

The Flores shrew (*Suncus mertensi*) is a white-toothed shrew found only on Flores Island, Indonesia. It is listed as a critically endangered species due to habitat loss and a restricted range.
Flores warty pig
Flores warty pig
Sus heurni

The Flores warty pig (*Sus celebensis floresianus*), is a subspecies in the pig genus (*Sus*) found in southern Asia.
Flores woolly bat
Flores woolly bat
Kerivoula flora

The Flores woolly bat (*Kerivoula flora*) is a species of vesper bat in the family Vespertilionidae.
Florida black bear
Florida black bear
Ursus americanus floridanus

The Florida black bear (*Ursus americanus floridanus*) is a subspecies of the

American black bear that has historically ranged throughout most of Florida and the southern portions of Georgia, Alabama, and Mississippi. The large black-furred bears...

Florida black wolf

Florida black wolf

Canis lupus floridanus

The Florida black wolf (*Canis rufus floridanus*), also known as the Florida wolf and the black wolf, is an extinct subspecies of red wolf. that was endemic to Florida. This subspecies became extinct in 1934 due to crowding out of its habitat and...

Florida bonneted bat

Florida bonneted bat

Eumops floridanus

The Florida bonneted bat or Florida mastiff bat (*Eumops floridanus*) is a species of bat in the genus *Eumops*, the bonneted bats or mastiff bats. Until recently, it was classified as a subspecies of Wagner's bonneted bat (*Eumops glaucinus*). It is...

Florida Mouse

Florida Mouse

Podomys floridanus

The Florida mouse (*Podomys floridanus*) is a species of rodent in the family Cricetidae. It is the only species in the genus *Podomys*, which is the only mammal genus endemic to Florida. The Florida mouse (also known as the big-eared deer mouse, the...

Florida naked-tailed rat

Florida naked-tailed rat

Solomys salomonis

The Florida naked-tailed rat (*Solomys salomonis*) is a poorly known and possible extinct species of rodent in the family Muridae. It was confined to the Nggela Islands (previously known as Florida Islands) in the Solomon Islands. The originally...

Florida Panther

Florida Panther

Puma concolor cougar

The Florida panther is known under a number of common names including Costa Rican puma, Florida cougar, and Florida puma. It is the only confirmed cougar population in the eastern United States, and currently occupies 5% of its historic range. In...

Flower-faced bat

Flower-faced bat

Anthops ornatus

The flower-faced bat (*Anthops ornatus*) is a species of bat in the family Hipposideridae. It is in the monotypic genus *Anthops*. It is found in the Autonomous Region of Bougainville of Papua New Guinea and in the Solomon Islands. This rare and...

Flower's gerbil

Flower's gerbil

Gerbillus floweri

Flower's gerbil (*Gerbillus floweri*) is a large gerbil distributed mainly in Egypt, from the eastern fringes of the Nile delta to the Sinai Peninsula, south of El Arish. Less than 250 individuals of this species are thought to persist in the... Flower's shrew

Flower's shrew
Crocidura floweri

Flower's shrew (*Crocidura floweri*) is a species of mammal in the family Soricidae. It is endemic to Egypt. Its natural habitat is arable land. It is threatened by habitat loss.

Flute-nosed bat
Flute-nosed bat
Murina florium

The flute-nosed bat (*Murina florium*) is a vespertilionid bat with an unusually shaped nose, the tubular nostrils facing outward from the end of the muzzle. They occur in the north of the Australian state of Queensland, in Indonesia, and on Papua...

Fly River roundleaf bat
Fly River roundleaf bat
Hipposideros muscinus

The Fly River roundleaf bat (*Hipposideros muscinus*) is a species of bat in the family Hipposideridae. It is found in West Papua, Indonesia and Papua New Guinea.

Fly River trumpet-eared bat
Fly River trumpet-eared bat
Kerivoula muscina

The Fly River trumpet-eared bat (*Kerivoula muscina*) is a species of vesper bat in the family Vespertilionidae. It is found only in Papua New Guinea. Its natural habitat is subtropical or tropical dry forests.

Fly River water rat
Fly River water rat
Leptomys signatus

The Fly River water rat (*Leptomys signatus*) is a species of rodent in the family Muridae. It is found only in Papua New Guinea. Its natural habitat is subtropical or tropical dry forests. It is threatened by habitat loss.

Foa's red colobus
Foa's red colobus
Piliocolobus foai

Foa's red colobus (*Piliocolobus foai*) or the Central African red colobus, is a species of red colobus monkey found in the Democratic Republic of the Congo.

Foch's tuco-tuco
Foch's tuco-tuco
Ctenomys fochi

Foch's tuco-tuco (*Ctenomys fochi*) is a species of rodent in the family Ctenomyidae. It is endemic to northwestern Argentina, where it is known from southwestern Catamarca Province. The species is named after World War I general Ferdinand Foch.

Fog shrew
Fog shrew
Sorex sonomae

The fog shrew (*Sorex sonomae*) is a species of mammal in the family Soricidae. It is endemic to northern California and Oregon in the United States.

Forest Dormouse
Forest Dormouse
Dryomys nitedula

The Forest dormouse (*Dryomys nitedula*) is a small rodent from eastern Europe, the

Balkans and parts of western Central Asia. The species competes in artificial and natural nests with Hazel dormice, Fat dormice, and birds. Their biggest...
Forest giant squirrel
Forest giant squirrel
Protoxerus stangeri

The forest giant squirrel (*Protoxerus stangeri*) or Stanger's squirrel is a species of rodent in the family Sciuridae found in Angola, Benin, Burundi, Cameroon, the Central African Republic, the Republic of the Congo, the Democratic Republic of...
Forest horseshoe bat
Forest horseshoe bat
Rhinolophus silvestris

The forest horseshoe bat (*Rhinolophus silvestris*) is a species of bat in the family Rhinolophidae. It is found in Republic of the Congo and Gabon. Its natural habitats are subtropical or tropical moist lowland forest and caves. It is threatened...
Forest Oldfield mouse
Forest Oldfield mouse
Thomasomys silvestris

The forest Oldfield mouse (*Thomasomys silvestris*) is a species of rodent in the family Cricetidae. It is found only in Ecuador.
Forest pipistrelle
Forest pipistrelle
Pipistrellus adamsi

The forest pipistrelle (*Pipistrellus adamsi*) is a species of vesper bat found in Australia, in the northernmost parts of Queensland and the Northern Territory.
Forest soft-furred mouse
Forest soft-furred mouse
Praomys rostratus

The forest soft-furred mouse or West African praomys (*Praomys rostratus*) is a species of rodent in the family Muridae. It is found in Ivory Coast, Ghana, Guinea, and Liberia. Its natural habitats are subtropical or tropical moist lowland forest...
Forest thicket rat
Forest thicket rat
Grammomys dryas

The forest thicket rat (*Grammomys dryas*) is a species of rodent in the family Muridae. It is found in Burundi, Democratic Republic of the Congo, and Uganda. Its natural habitat is subtropical or tropical moist montane forests.
Forest tuco-tuco
Forest tuco-tuco
Ctenomys sylvanus

The forest tuco-tuco (*Ctenomys sylvanus*) was formerly considered a species of rodent in the family Ctenomyidae. It is endemic to Salta and southeast Jujuy Provinces in northwest Argentina. The IUCN currently recognizes it as a subspecies of C....
Forest-steppe marmot
Forest-steppe marmot
Marmota kastschenkoi

The forest-steppe marmot (*Marmota kastschenkoi*) is a rodent species of the marmot genus found in south-central Russia. It lives in wooded forest steppe at an altitude of 180–450 m (590–1,480 ft) in a relatively small region located directly

east...

Formosan black bear

Formosan black bear

Ursus thibetanus formosanus

The Formosan black bear (臺灣黑熊, *Ursus thibetanus formosanus*), also known as the Taiwanese black bear or white-throated bear, is a subspecies of the Asiatic black bear. It was first described by Robert Swinhoe in 1864. Formosan black bears are...

Formosan golden tube-nosed bat

Formosan golden tube-nosed bat

Harpiola isodon

The Formosan golden tube-nosed bat (*Harpiola isodon*) is native to the high-altitude regions of Taiwan.

Formosan Rock Macaque

Formosan Rock Macaque

Macaca cyclopis

The Formosan rock macaque (*Macaca cyclopis*), also known as the Formosan rock monkey or Taiwanese macaque, is a macaque endemic to the island of Taiwan, which has also been introduced to Japan. Besides humans, Formosan rock macaques are the only...

Formosan sambar deer

Formosan sambar deer

Rusa unicolor swinhoei

The Formosan sambar (*Rusa unicolor swinhoei*) is a subspecies of the sambar found in Taiwan. It is the largest native herbivore there. Its fur color changes with season to provide camouflage. It is yellowish-brown in the summer and dark brown in...

Formosan Sika Deer

Formosan Sika Deer

Cervus nippon taiouanus

The Formosan sika deer (*Cervus nippon taiouanus*) is a subspecies of Sika deer found only on the island of Taiwan.

Formosan woolly horseshoe bat

Formosan woolly horseshoe bat

Rhinolophus formosae

The Formosan woolly horseshoe bat (*Rhinolophus formosae*) is a species of bat from the family Rhinolophidae. It is endemic to Taiwan and occurs in primary forests in low to middle altitudes of central areas of Taiwan. Its roosting locations...

Forrest's mountain vole

Forrest's mountain vole

Neodon forresti

Forrest's mountain vole (*Neodon forresti*) is a species of rodent in the family Cricetidae found within China in northwest Yunnan. The initial study by Hinton in 1923 identified it as morphologically close to the Chinese scrub vole (*N. irene*)...

Forrest's mouse

Forrest's mouse

Leggadina forresti

The Forrest's mouse (*Leggadina forresti*), or desert short-tailed mouse, is a small species of rodent in the family Muridae. It is a widespread but sparsely distributed species found across arid and semi-arid inland Australia, commonly found in...

Forrest's pika
Forrest's pika
Ochotona forresti

The Forrest's pika (*Ochotona forresti*) is a species of mammal in the pika family, Ochotonidae. It is found in Bhutan, China, India, and Myanmar. The summer dorsal pelage and ventral pelage are dark rufous or blackish brown, and the winter dorsal...

Forrest's rock squirrel
Forrest's rock squirrel
Sciurotamias forresti

Forrest's rock squirrel (*Sciurotamias forresti*) is a species of rodent in the family Sciuridae. It is endemic to Sichuan and Yunnan in China, where it inhabits cliffs covered in shrubby vegetation at an altitude around 3,000 m (9,800 ft)....

Fossa
Fossa
Cryptoprocta ferox

The unusual-looking fossa (*Cryptoprocta ferox*) from Madagascar looks like a puma and has both canine and feline features. It is the largest mammalian carnivore on the island, and preys mainly on lemurs, pursuing them through the trees with...

Fossorial giant rat
Fossorial giant rat
Gyldenstolpia fronto

The fossorial giant rat (*Gyldenstolpia fronto*) is a species of rodent in the family Cricetidae. It is found in Argentina and Brazil but was determined extinct following a recent assessment of the conservation status of Sigmodontine rodents. Its...

Four-Horned Antelope
Four-Horned Antelope
Tetracerus quadricornis

The four-horned antelope (*Tetracerus quadricornis*), or chousingha, is a small antelope found in India and Nepal. Its four horns distinguish it from most other bovids, which have two horns (with a few exceptions, such as the Jacob sheep). The...

Four-striped grass mouse
Four-striped grass mouse
Rhabdomys pumilio

The four-striped grass mouse (*Rhabdomys pumilio*) or four-striped grass rat, is a species of rodent in the family Muridae. It is found throughout the southern half of Africa up to 2,300 metres (7,500 ft) above sea level, extending as far north as...

Four-striped ground squirrel
Four-striped ground squirrel
Lariscus hosei

The four-striped ground squirrel (*Lariscus hosei*) is a species of rodent in the family Sciuridae. It is endemic to Borneo. Its natural habitat is subtropical or tropical dry forests. It is threatened by habitat loss.

Four-toed elephant shrew
Four-toed elephant shrew
Petrodromus tetradactylus

The four-toed elephant shrew or four-toed sengi is the only living species in the genus *Petrodromus*, which together with five other extant genera *Rhynchocyon*, *Macroscelides*, *Petrosaltator*, *Galegeeska* and *Elephantulus* constitutes the order...

Four-Toed Hedgehog
Four-Toed Hedgehog
Atelerix albiventris

The Four-toed hedgehog (*Atelerix albiventris*) is a spiny mammal found in Africa. Its common name is derived from the number of toes found on its hind feet. The Four-toes hedgehog is a popular small pet in various countries, and has been...
Four-toed jerboa
Four-toed jerboa
Allactaga tetradactyla

The four-toed jerboa (*Allactaga tetradactyla*) is a rodent of the family Dipodidae and genus *Allactaga* that has four digits. It is the sole species in the subgenus *Scarturus*. Four-toed jerboas are native to Egypt and Libya. They live in coastal...
Four-toed rice tenrec
Four-toed rice tenrec
Oryzorictes tetradactylus

The four-toed rice tenrec (*Oryzorictes tetradactylus*) is a species of mammal in the family Tenrecidae. It is endemic to Madagascar. Its natural habitats are subtropical or tropical moist lowland forest, subtropical or tropical moist montane...
Fox's shaggy rat
Fox's shaggy rat
Dasymys foxi

Fox's shaggy rat (*Dasymys foxi*) is a species of rodent in the family Muridae. It is found only in Nigeria. Its natural habitats are moist savanna, subtropical or tropical seasonally wet or flooded lowland grassland, swampland, and plantations. It...
Fox's shrew
Fox's shrew
Crocidura foxi

Fox's shrew (*Crocidura foxi*) is a species of mammal in the family Soricidae. It is found in Benin, Burkina Faso, Cameroon, Central African Republic, Chad, Ivory Coast, Gambia, Ghana, Guinea, Guinea-Bissau, Mali, Nigeria, Senegal, South Sudan,...
François' Langur
François' Langur
Trachypithecus francoisi

The François' langur (*Trachypithecus francoisi*), also known as the Francois' leaf monkey, Tonkin leaf monkey, or white side-burned black langur is a species of lutung and the type species of its species group. It is one of the least studied of...
Franklin's Ground Squirrel
Franklin's Ground Squirrel
Poliocitellus franklinii

Franklin's ground squirrel (*Poliocitellus franklinii*) is a species of squirrel native to North America, and the only member of the genus *Poliocitellus*. Due to the destruction of prairie, the populations of Franklin's ground squirrel have...
Franquet's epauletted fruit bat
Franquet's epauletted fruit bat
Epomops franqueti

Franquet's epauletted fruit bat (*Epomops franqueti*) is a species of megabat in the family Pteropodidae, and is one of three different species of epauletted bats.

Franquet's epauletted fruit bat has a range of habitats, varying from sub-saharan...

Fraser's Dolphin

Fraser's Dolphin

Lagenodelphis hosei

Fraser's dolphin (*Lagenodelphis hosei*) is a cetacean that lives in deep oceans. In 1895, Charles E. Hose found a skull on a beach in Sarawak, Borneo but the species was discovered only in 1971. At that time washed-up specimens were found on Cocos...

Fraser's musk shrew

Fraser's musk shrew

Crocidura poensis

Fraser's musk shrew (*Crocidura poensis*) is a species of mammal in the family Soricidae. It is found in Benin, Cameroon, Ivory Coast, Equatorial Guinea, Ghana, Guinea, Liberia, Nigeria, São Tomé and Príncipe, Sierra Leone, and Togo. Its natural...

Fraternal fruit-eating bat

Fraternal fruit-eating bat

Artibeus fraterculus

The fraternal fruit-eating bat (*Artibeus fraterculus*) is a species of bat in the family Phyllostomidae that is found in drier habitats in Ecuador and Peru. It was formerly considered to be a subspecies of the Jamaican fruit bat, but was raised...

Fraternal hill rat

Fraternal hill rat

Bunomys fratorum

The fraternal hill rat (*Bunomys fratorum*) is a species of rodent in the family Muridae. It is found only in northeastern Sulawesi, Indonesia. Its natural habitat is tropical dry forest. It is threatened by habitat loss.

Fraternal myotis

Fraternal myotis

Myotis frater

The fraternal myotis (*Myotis frater*) is a species of vesper bat native to East Asia.

Fraternal squirrel

Fraternal squirrel

Sundasciurus fraterculus

The fraternal squirrel (*Sundasciurus fraterculus*) is a species of rodent in the family Sciuridae. It is endemic to Indonesia, found only in the Mentawai Archipelago (Siberut, Sipora, North and South Pagai islands), located off the west coast of...

Free State pygmy mouse

Free State pygmy mouse

Mus orangiae

The Free State pygmy mouse or Orange mouse (*Mus orangiae*) is a species of rodent in the family Muridae. It is found in Lesotho and South Africa. Its natural habitats are subtropical or tropical high-altitude grassland, arable land, and pastureland.

Fresno kangaroo rat

Fresno kangaroo rat

Dipodomys nitratoides

The Fresno kangaroo rat or San Joaquin kangaroo rat (*Dipodomys nitratoides*) is a species of rodent in the family Heteromyidae. It is endemic to areas within and near the San Joaquin Valley of California in the United States. Habitat

destruction...

Friendly leaf-eared mouse
Friendly leaf-eared mouse
Phyllotis amicus

The friendly leaf-eared mouse (*Phyllotis amicus*) is a species of rodent in the family Cricetidae. It is found only in Peru.

Fringe-Eared Oryx
Fringe-Eared Oryx
Oryx beisa callotis

The Fringe-eared oryx (*Oryx callotis* or *Oryx beisa callotis*) is a large species or subspecies of oryx antelope native to East Africa. It was originally described as a distinct species by Oldfield Thomas in 1892, but was subsequently re-evaluated...

Fringe-lipped bat
Fringe-lipped bat
Trachops cirrhosus

The fringe-lipped bat (*Trachops cirrhosus*) is a leaf-nosed bat from southern Mexico to Bolivia and southern Brazil. It has three subspecies and no known fossils. It is the only species within its genus.

Fringe-tailed gerbil
Fringe-tailed gerbil
Gerbilliscus robustus

The fringe-tailed gerbil (*Gerbilliscus robustus*) is a species of rodent found in Central African Republic, Chad, Eritrea, Ethiopia, Kenya, Niger, Somalia, Sudan, Tanzania, Uganda, possibly Cameroon, and possibly Nigeria. Its natural habitats are...

Fringed fruit-eating bat
Fringed fruit-eating bat
Artibeus fimbriatus

The fringed fruit-eating bat (*Artibeus fimbriatus*), is a species of bat native to South America.

Fringed myotis
Fringed myotis
Myotis thysanodes

The fringed myotis (*Myotis thysanodes*) is a species of vesper bat that is found in British Columbia, Mexico, and the western United States.

Frosted hairy dwarf porcupine
Frosted hairy dwarf porcupine
Coendou pruinosus

The frosted hairy dwarf porcupine, *Coendou pruinosus*, is a porcupine species in the family Erethizontidae from Colombia and northern and eastern Venezuela. It was formerly sometimes assigned to *Sphiggurus*, a genus no longer recognized since...

Frosted myotis
Frosted myotis
Myotis pruinosus

The frosted myotis (クロホオヒゲコウモリ in Japanese) (*Myotis pruinosus*) is a species of vesper bat. It is found only in Japan.

Frosted sac-winged bat
Frosted sac-winged bat
Saccopteryx canescens

The frosted sac-winged bat (*Saccopteryx canescens*) is a bat species of the family

Emballonuridae from South America. It is found in northern Brazil, Colombia, Ecuador, French Guiana, Guyana, Peru, Suriname, Venezuela and possibly Bolivia.
Fukomys ilariae
Fukomys ilariae
Fukomys iliariae

Fukomys ilariae, commonly known as the Somali striped mole rat, is a species of small mole rat that is endemic to Africa.

Fukomys vandewoestijneae
Fukomys vandewoestijneae
Fukomys vandewoestijneae

Fukomys vandewoestijneae or Caroline's mole rat was first noticed in 2002 by a research team from the University of Ghent. This new species, distinguished by a distinctive skull shape, was recently described by Paul Van Daele and his team in...

Fulvous harvest mouse
Fulvous harvest mouse
Reithrodontomys fulvescens

The fulvous harvest mouse (*Reithrodontomys fulvescens*) is a species of rodent in the family Muridae. It is found in El Salvador, Guatemala, Honduras, Mexico, Nicaragua, and the United States.
Fulvous-bellied climbing rat
Fulvous-bellied climbing rat
Tylomys fulviventer

The fulvous-bellied climbing rat (*Tylomys fulviventer*) is a species of rodent in the family Cricetidae. It is found only in Panama.
Fulvus roundleaf bat
Fulvus roundleaf bat
Hipposideros fulvus

The fulvus roundleaf bat or fulvus leaf-nosed bat (*Hipposideros fulvus*) is a species of bat in the family Hipposideridae. It is found in Afghanistan, Bangladesh, Cambodia, China, India, Laos, Myanmar, Pakistan, Sri Lanka and Thailand and...
Funambulus obscurus
Funambulus obscurus
Funambulus obscurus

The dusky striped squirrel (*Funambulus obscurus*) is a species of rodent, a small squirrel (Sciuridae) from Sri Lanka where largely confined to rainforests in the southwestern "wet zone" with higher rainfall than the rest of the island. It was...
Furry-eared dwarf lemur
Furry-eared dwarf lemur
Cheirogaleus crossleyi

The furry-eared dwarf lemur (*Cheirogaleus crossleyi*), or Crossley's dwarf lemur, only found on the island of Madagascar, as with all other lemurs. It has a pelage coloration that is red-brown dorsally and gray ventrally. The eye-rings of this...
Furtive tuco-tuco
Furtive tuco-tuco
Ctenomys occultus

The furtive tuco-tuco (*Ctenomys occultus*) is a species of rodent in the family Ctenomyidae. It is endemic to Argentina.
Fynbos golden mole
Fynbos golden mole

Amblysomus corriae

The fynbos golden mole (*Amblysomus corriae*) is a species of mammal in the golden mole family, Chrysochloridae. It is endemic to South Africa. Its natural habitats are fynbos Mediterranean shrubland vegetation, temperate forests, subtropical or...

Gabon bushbaby

Gabon bushbaby

Sciurocheirus gabonensis

The Gabon bushbaby (*Sciurocheirus gabonensis*) is a species of primate in the family Galagidae found in Cameroon, Gabon, and the Republic of the Congo. Its head and body length is 8.5 in with a 10-in tail, and it weighs about 10 oz. It lives in...

Gabon Talapoin

Gabon Talapoin

Miopithecus ogouensis

The Gabon talapoin (*Miopithecus ogouensis*), also known as the northern talapoin, is a small species of African monkey native to riparian habitats in Cameroon, Equatorial Guinea, Gabon, the western Republic of the Congo and the far western...

Gairdner's shrewmouse

Gairdner's shrewmouse

Mus pahari

Gairdner's shrewmouse (*Mus pahari*) is a species of rodent in the family Muridae found in China, India, Laos, Myanmar, Thailand, and Vietnam.

Galagoides kumbirensis

Galagoides kumbirensis

Galagoides kumbirensis

The Angolan dwarf galago (*Galagoides kumbirensis*) is a species of dwarf galago native to Angola, and was named after western Angolan Kumbira Forest. Though 36 individuals of the Angolan dwarf galago were identified in September 2013, it was...

Galápagos Fur Seal

Galápagos Fur Seal

Arctocephalus galapagoensis

The Galápagos fur seal (*Arctocephalus galapagoensis*) is one of eight seals in the genus *Arctocephalus* and one of nine seals in the subfamily Arctocephalinae. It is the smallest of all of the eared seals. They are endemic to the Galápagos Islands...

Galápagos Sea Lion

Galápagos Sea Lion

Zalophus wollebaeki

Galápagos sea lions (*Zalophus wollebaeki*) are the smallest sea lion species and one of the most numerous species in the Galápagos archipelago. They are fairly social and can often be spotted sunning on sandy shores or rocks or gliding through the...

Gallagher's free-tailed bat

Gallagher's free-tailed bat

Chaerephon gallagheri

Gallagher's free-tailed bat (*Chaerephon gallagheri*) is a free-tailed bat endemic to the Democratic Republic of the Congo. Only one individual has ever been documented.

Gambian epauletted fruit bat

Gambian epauletted fruit bat

Epomophorus gambianus

The Gambian epauletted fruit bat (*Epomophorus gambianus*) is a species of megabat in the family Pteropodidae. The species is known to travel in packs and are also known to be tropical. It is found in Benin, Burkina Faso, Cameroon, Central African...

Gambian mongoose
Gambian mongoose
Mungos gambianus

The Gambian mongoose (*Mungos gambianus*) is a mongoose species native to the Guinean forest-savanna mosaic from Gambia to Nigeria. It is listed as Least Concern on the IUCN Red List since 2008.

Gambian Pouched Rat
Gambian Pouched Rat
Cricetomys gambianus

The Gambian pouched rat (*Cricetomys gambianus*) is a species of pouched rat of the giant pouched rat genus *Cricetomys*. It is among the largest muroids in the world, growing to about 0.9 m (3 ft) long, including the tail, which makes up half of its...

Gambian slit-faced bat
Gambian slit-faced bat
Nycteris gambiensis

The Gambian slit-faced bat (*Nycteris gambiensis*) is a species of bat in the family Nycteridae found in Benin, Burkina Faso, Cameroon, Ivory Coast, Gambia, Ghana, Guinea, Guinea-Bissau, Mali, Mauritania, Niger, Nigeria, Senegal, Sierra Leone, and...

Gambian sun squirrel
Gambian sun squirrel
Heliosciurus gambianus

The Gambian sun squirrel (*Heliosciurus gambianus*) is a species of rodent in the family Sciuridae. It is found in Angola, Benin, Burkina Faso, Central African Republic, Chad, Democratic Republic of the Congo, Ivory Coast, Eritrea, Ethiopia,...

Ganges River Dolphin
Ganges River Dolphin
Platanista gangetica

The Ganges River dolphin (*Platanista gangetica*) is a species of toothed whale found in South Asia. It is also known by the name susu (popular name) or "Sisu" (Assamese language) and shushuk (Bengali). The Ganges river dolphin has been recognized...

Gansu hamster
Gansu hamster
Cansumys canus

The Gansu hamster (*Cansumys canus*) is a species of rodent in the family Cricetidae, endemic to China. It is the only species in the genus *Cansumys*.

Gansu mole
Gansu mole
Scapanulus oweni

The Gansu mole (*Scapanulus oweni*) is a species of mammal in the family Talpidae endemic to central China, where it occurs in Shaanxi, Gansu, Sichuan, and Qinghai. It is the only species in the genus *Scapanulus*. Along with the Medog mole...

Gansu pika
Gansu pika
Ochotona cansus

The Gansu pika (*Ochotona cansus*) is a species of mammal in the pika family, Ochotonidae. It is endemic to China.

Gansu shrew

Gansu shrew

Sorex cansulus

The Gansu shrew (*Sorex cansulus*) is a red-toothed shrew found only in a small part of Gansu province and other adjacent areas of China. With its very limited range, it is sensitive to habitat loss and is listed as a "vulnerable species" in the...

Ganzhorn's mouse lemur

Ganzhorn's mouse lemur

Microcebus ganzhorni

Ganzhorn's mouse lemur (*Microcebus ganzhorni*) is a species of lemur described in 2016. The discovery was made by researchers at the German Primate Center. It was discovered in Madagascar among closely related species such as Madame Berthe's...

Ganzu vole

Ganzu vole

Caryomys eva

The Ganzu vole, Eva's red-backed vole, Eva's vole, Gansu vole, or Taozhou vole (*Caryomys eva*) is a species of rodent in the family Cricetidae. It is found in mountain forests in China. The IUCN has assessed it as being of "least concern".

Gaoligong forest hedgehog

Gaoligong forest hedgehog

Mesechinus wangi

The Gaoligong forest hedgehog (*Mesechinus wangi*) or Wang's forest hedgehog is a species of hedgehog in the family Erinaceidae found only in China. It is endemic to the slopes of Mt. Gaoligong in the Yunnan Province, where it lives in subtropical...

Gaoligong pika

Gaoligong pika

Ochotona gaoligongensis

The Gaoligong pika (*Ochotona gaoligongensis*) is a species of mammal in the family Ochotonidae. It is endemic to China. Many of the general physical characteristics of the pika species, are shared by Gaoligong pikas. However, the Gaoligong pika...

Garden Dormouse

Garden Dormouse

Eliomys quercinus

Garden dormice (*Eliomys quercinus*) are rodents in the dormouse family. They are named for their long, dormant hibernation period.

Gardner's climbing mouse

Gardner's climbing mouse

Rhipidomys gardneri

Gardner's climbing mouse (*Rhipidomys gardneri*) is a rodent species from South America. It is found in western Brazil, southeastern Peru and northeastern Bolivia. It is an uncommon, arboreal species, but faces no particular threats so the...

Gardner's spiny rat

Gardner's spiny rat

Proechimys gardneri

Gardner's spiny-rat, *Proechimys gardneri*, is a spiny rat species from South America. It is found in Bolivia and Brazil.

Garlepp's mouse

Garlepp's mouse
Galenomys garleppi

Garlepp's mouse (*Galenomys garleppi*) is a species of rodent in the family Cricetidae. It is the only species in the genus *Galenomys*. It is found in western Bolivia, southern Peru and possibly Chile at elevations over 3,000 m in the Altiplano.

Garrido's hutia
Garrido's hutia
Mysateles garridoi

Garrido's hutia (*Capromys garridoi*) is a critically endangered or possibly extinct species of hutia that is found in the Cuban moist forests Global 200 ecoregion.

Gaskell's false serotine
Gaskell's false serotine
Hesperoptenus gaskelli

Gaskell's false serotine (*Hesperoptenus gaskelli*) is a species of vesper bat. It is found only in Indonesia. It is threatened by habitat loss.

Gaumer's spiny pocket mouse
Gaumer's spiny pocket mouse
Heteromys gaumeri

Gaumer's spiny pocket mouse (*Heteromys gaumeri*) is a species of rodent in the family Heteromyidae. It ranges over northern Belize and Guatemala and the Yucatán Peninsula of Mexico, where it lives in lowland semideciduous forest and thorn scrub....

Gaur
Gaur
Bos gaurus

The gaur (*Bos gaurus*) is a bovine native to South and Southeast Asia. It is the largest species among the wild cattle and the Bovidae. In Malaysia, this adorable giant is called seladang, and pyaung in Myanmar.

Gayal
Gayal
Bos frontalis

The gayal (*Bos frontalis*) is a large domestic cattle found in Southeast Asia.

Gazella erlangeri
Gazella erlangeri
Gazella erlangeri

Erlanger's gazelle or Neumann's gazelle, (*Gazella erlangeri*), is a small dark gazelle with a stout body and short legs. It is described from Saudi Arabia and Yemen. It is considered a threatened species, although no recent information is...

Geata mouse shrew
Geata mouse shrew
Myosorex geata

The Geata mouse shrew (*Myosorex geata*) is a species of mammal in the family Soricidae endemic to Tanzania. Its natural habitat is subtropical or tropical moist montane forests.

Gebe cuscus
Gebe cuscus
Phalanger alexandrae

The Gebe cuscus (*Phalanger alexandrae*) is a species of marsupial in the family Phalangeridae. It is endemic to the island of Gebe, North Maluku province,

Indonesia, where it lives at elevations from sea level to 300 m. It is named after the...

Gee's Golden Langur
Gee's Golden Langur
Trachypithecus geei

Gee's golden langur (*Trachypithecus geei*), also known as simply the golden langur, is an Old World monkey found in a small region of Western Assam, India and in the neighboring foothills of the Black Mountains of Bhutan. Long considered sacred...

Geelvink Bay flying fox
Geelvink Bay flying fox
Pteropus pohlei

The Geelvink Bay flying fox or Geelvink Bay fruit bat (*Pteropus pohlei*) is a species of flying fox in the family Pteropodidae. It is endemic to the islands of Yapen, Numfor, and Rani, which lie north of New Guinea in Indonesia's Papua Province....

Gelada
Gelada
Theropithecus gelada

The gelada (*Theropithecus gelada*) is a species of Old World monkey found only in the Ethiopian Highlands. Like its close relatives the baboons (genus *Papio*), it is largely terrestrial, spending much of its time foraging in grasslands. They are...

Gemsbok
Gemsbok
Oryx gazella

The gemsbok (*Oryx gazella*) is a large antelope native to the arid regions of Southern Africa. This rare animal is depicted on the coat of arms of Namibia, where the current population of the species is estimated at 373,000 individuals. In the...

Genoways's yellow bat
Genoways's yellow bat
Rhogeessa genowaysi

Genoways's yellow bat (*Rhogeessa genowaysi*) is a species of vesper bat. It is found only in Mexico. It is threatened by habitat loss. Due to its imperiled status, it is identified by the Alliance for Zero Extinction as a species in danger of...

Geoffroy's bat
Geoffroy's bat
Myotis emarginatus

Geoffroy's bat (*Myotis emarginatus*) is a species of vesper bat.

Geoffroy's Cat
Geoffroy's Cat
Leopardus geoffroyi

Geoffroy's cat (*Leopardus geoffroyi*) is a small wild cat native to South America. It is about the size of a domestic cat and is widespread and abundant over most of its range.

Geoffroy's horseshoe bat
Geoffroy's horseshoe bat
Rhinolophus clivosus

Geoffroy's horseshoe bat (*Rhinolophus clivosus*) is a species of bat in the family Rhinolophidae found in Africa. Its natural habitats are subtropical or tropical moist lowland forest, Mediterranean-type shrubby vegetation, caves, subterranean...

Geoffroy's rousette
Geoffroy's rousette
Rousettus amplexicaudatus

Geoffroy's rousette (*Rousettus amplexicaudatus*) is a species of megabat or Old World fruit bats. It is one of 10 species in the genus *Rousettus*.

Geoffroy's Spider Monkey
Geoffroy's Spider Monkey
Ateles geoffroyi

Geoffroy's spider monkey (*Ateles geoffroyi*), also known as the black-handed spider monkey or the Central American spider monkey is a species of spider monkey, a type of New World monkey, from Central America, parts of Mexico and possibly a small...

Geoffroy's tailless bat
Geoffroy's tailless bat
Anoura geoffroyi

Geoffroy's tailless bat (*Anoura geoffroyi*) is a species of phyllostomid bat from the American tropics.

Geoffroy's Tamarin
Geoffroy's Tamarin
Saguinus geoffroyi

Geoffroy's tamarin (*Saguinus geoffroyi*), also known as the Panamanian, Red-crested or Rufous-naped tamarin, is a tamarin, a type of small monkey, found in Panama and Colombia. The species name *geoffroyi* is in honor of French naturalist Étienne...

Geoxus annectens
Geoxus annectens
Geoxus annectens

Geoxus annectens, also known as Pearson's long-clawed akodont or Pearson's long-clawed mouse, is a species of rodent in the tribe Abrotrichini of family Cricetidae. Molecular data suggests that its closest relative is *Geoxus valdivianus*. Formerly...

Geoxus valdivianus
Geoxus valdivianus
Geoxus valdivianus

Geoxus valdivianus, also known as the long-clawed mole mouse or Valdivian long-clawed akodont, is a species of rodent in the tribe Abrotrichini of family Cricetidae found in the Valdivian temperate rain forests and Magellanic subpolar forests of...

Gerbe's vole
Gerbe's vole
Microtus gerbii

Gerbe's vole (*Microtus gerbii*) is a species of rodent in the family Cricetidae found in France and Spain.

Gerbil leaf-eared mouse
Gerbil leaf-eared mouse
Phyllotis gerbillus

The gerbil leaf-eared mouse (*Phyllotis gerbillus*) is a species of rodent in the family Cricetidae. It is found only in Peru. It has been discovered that *P. amicus* and *P. gerbillus* are closely related as sister species. However, *P. gerbillus*...

Gerbil mouse
Gerbil mouse
Malacothrix typica

The gerbil mouse or long-eared mouse (*Malacothrix typica*) is a species of rodent in the family Nesomyidae. It is found in Angola, Botswana, Namibia, and South Africa. Its natural habitats are dry savanna, subtropical or tropical dry shrubland,...

Gerenuk

Gerenuk

Litocranius walleri

The gerenuk (*Litocranius walleri*) is an unusually looking long-necked antelope found in the Horn of Africa and the drier parts of East Africa. The gerenuk was first described by the naturalist Victor Brooke in 1879. The common name derives from...

German's one-toothed moss mouse

German's one-toothed moss mouse

Pseudohydromys germani

German's one-toothed moss mouse (*Pseudohydromys germani*) is a species of rodent in the family Muridae which occurs in the mountains of southeastern New Guinea.

Gerp's mouse lemur

Gerp's mouse lemur

Microcebus gerpi

Gerp's mouse lemur (*Microcebus gerpi*) is a species of mouse lemur known only from the Sahafina Forest in eastern Madagascar, near Mantadia National Park. Its discovery was announced in 2012 by a German and Malagasy research team. The Sahafina...

Gervais's Beaked Whale

Gervais's Beaked Whale

Mesoplodon europaeus

Gervais's beaked whale (*Mesoplodon europaeus*), sometimes known as the Antillean beaked whale, Gulf Stream beaked whale, or European beaked whale (from which its scientific name is derived) is the most frequently stranding type of mesoplodont...

Gervais's fruit-eating bat

Gervais's fruit-eating bat

Dermanura cinerea

Gervais's fruit-eating bat (*Dermanura cinerea*) is a bat species from South America. It is found in Brazil, French Guiana, Guyana, eastern Peru, Suriname and eastern Venezuela.

Gervais's funnel-eared bat

Gervais's funnel-eared bat

Nyctiellus lepidus

Gervais's funnel-eared bat (*Nyctiellus lepidus*) is a species of bat in the family Natalidae. It is monotypic within the genus *Nyctiellus*. It is found in Bahamas and Cuba.

Ghana mole-rat

Ghana mole-rat

Fukomys zechi

The Ghana mole-rat or Togo mole-rat (*Fukomys zechi*) is a species of rodent in the family Bathyergidae. It is endemic to Ghana. Its natural habitats are moist savanna, subtropical or tropical dry shrubland, subtropical or tropical dry lowland...

Ghana rufous-nosed rat

Ghana rufous-nosed rat

Oenomys ornatus

The Ghana rufous-nosed rat (*Oenomys ornatus*), also known as the West African

Oenomys, is a species of rodent in the family Muridae. It is found in Ivory Coast, Ghana, Guinea, Liberia, and Sierra Leone. Its natural habitats are subtropical or...

Ghost Bat

Ghost Bat

Macroderma gigas

The ghost bat (*Macroderma gigas*) is a flying mammal found in northern Australia. The species is the only Australian bat that preys on large vertebrates - birds, reptiles and other mammals - which they detect using acute sight and hearing,...

Ghost-faced bat

Ghost-faced bat

Mormoops megalophylla

The ghost-faced bat (*Mormoops megalophylla*) is a bat in the genus *Mormoops*. It occurs in Belize, Colombia, Ecuador, El Salvador, Guatemala, Honduras, Mexico, Peru, Trinidad and Tobago, Venezuela, and Texas in the United States. *Mormoops*...

Giant Anteater

Giant Anteater

Myrmecophaga tridactyla

The Giant anteater (*Myrmecophaga tridactyla*) is an insectivorous mammal native to Central and South America. It is one of four living species of anteaters, of which is the largest member. The Giant anteater is mostly terrestrial, in contrast to...

Giant Armadillo

Giant Armadillo

Priodontes maximus

The Giant armadillo (*Priodontes maximus*) is the largest living species of armadillo that lives in South America. It prefers termites and some ants as prey and often consumes the entire population of a termite mound. This species is considered...

Giant Atlantic tree-rat

Giant Atlantic tree-rat

Phyllomys thomasi

The giant Atlantic tree-rat (*Phyllomys thomasi*) is a spiny rat species from Brazil. The species is endemic to São Sebastião Island off the coast of São Paulo State. The island is largely protected by the Ilhabela State Park.

Giant bandicoot

Giant bandicoot

Peroryctes broadbenti

The giant bandicoot (*Peroryctes broadbenti*) is a species of marsupial in the family Peramelidae endemic to Papua New Guinea. Its natural habitat is subtropical or tropical dry forests. The type locality is Papua New Guinea, Central Prov., banks...

Giant bushy-tailed cloud rat

Giant bushy-tailed cloud rat

Crateromys schadenbergi

The giant bushy-tailed cloud rat or Luzon bushy-tailed cloud rat (*Crateromys schadenbergi*) is a large, endangered species of rodent in the family Muridae. It is only found in pine and mossy forest at altitudes of 2,000-2,740 metres...

Giant Eland

Giant Eland

Taurotragus derbianus

The Giant eland (*Taurotragus derbianus*) is the largest species of antelope. It is also called "Lord Derby's eland" in honor of Edward Smith-Stanley, 13th Earl of Derby. It was due to his efforts that the Giant eland was first introduced to...

Giant forest genet
Giant forest genet
Genetta victoriae

The giant forest genet (*Genetta victoriae*), also known as the giant genet, is a genet species endemic to the Congo Basin. As it is considered as widely distributed and common, it is listed as Least Concern on the IUCN Red List.

Giant Forest Hog
Giant Forest Hog
Hylochoerus meinertzhageni

The Giant forest hog (*Hylochoerus meinertzhageni*) is considered the largest wild member of the pig family. It is native to wooded habitats in Africa. Despite its large size and relatively wide distribution, it was first described only in 1904.

Giant ghost-faced bat
Giant ghost-faced bat
Mormoops magna

The giant ghost-faced bat (*Mormoops magna*) is an extinct species of bat endemic to the Caribbean. It is only known from fragmental humeri remains, which physically resemble those of *Mormoops megalophylla* but are larger in size.

Giant golden mole
Giant golden mole
Chrysospalax trevelyani

The giant golden mole (*Chrysospalax trevelyani*) is a small mammal found in Africa. At 23 centimetres (9.1 in) in length, it is the largest of the golden mole species. The mole has dark, glossy brown fur; the name golden comes from the Greek word...

Giant Golden-Crowned Flying Fox
Giant Golden-Crowned Flying Fox
Acerodon jubatus

The Giant golden-crowned flying fox (*Acerodon jubatus*) is native and endemic exclusively to the Philippines. Otherwise called the Golden-capped Fruit Bat, this animal is the largest and one of the rarest bats around the globe, currently...

Giant Kangaroo Rat
Giant Kangaroo Rat
Dipodomys ingens

The giant kangaroo rat (*Dipodomys ingens*) is an endangered species of heteromyid rodent endemic to California.

Giant mole shrew
Giant mole shrew
Anourosorex schmidi

The giant mole shrew (*Anourosorex schmidi*) is a species of red-toothed shrew native to the southeastern slopes of the Himalaya of Bhutan and India.

Giant mole-rat
Giant mole-rat
Spalax giganteus

The giant blind mole-rat (*Spalax giganteus*) is a species of rodent in the family Spalacidae endemic to the North Caucasus region of Russia. It feeds on roots and tubers and lives underground in a burrow that it digs with its teeth.

Giant Muntjac
Giant Muntjac
Muntiacus vuquangensis

The Giant muntjac (*Muntiacus vuquangensis*) is a species of muntjac deer. It is the

largest muntjac species and is most closely related to the Indian muntjac.
Giant naked-tailed rat
Giant naked-tailed rat
Uromys anak

The giant naked-tailed rat (*Uromys anak*) is a species of rodent in the family Muridae. It is found in West Papua, Indonesia and Papua New Guinea. It lives in tropical forests, wetlands, and in degraded forests.

Giant Otter
Giant Otter
Pteronura brasiliensis

The Giant otter (*Pteronura brasiliensis*) is a South American carnivorous mammal. It is the longest member of the weasel family, Mustelidae, a globally successful group of predators. The Giant otter has no serious natural predators other than...

Giant Otter Shrew
Giant Otter Shrew
Potamogale velox

The giant otter shrew (*Potamogale velox*) is a semiaquatic, carnivorous afrotherian mammal. It is found in the main rainforest block of central Africa from Nigeria to Zambia, with a few isolated populations in Kenya and Uganda. It lives in...

Giant Panda
Giant Panda
Ailuropoda melanoleuca

The Giant panda (*Ailuropoda melanoleuca*) is a bear species endemic to China. The name "giant panda" is sometimes used to distinguish it from the Red panda, a neighboring musteloid. The Giant panda has often served as China's national symbol,...

Giant Pangolin
Giant Pangolin
Smutsia gigantea

The Giant pangolin (*Smutsia gigantea*) is the biggest and the rarest of the African species of pangolin. It looks like an armored animal and behaves in the secretive manner of a spy. Its footprints look the same as a small elephant's. This unusual...

Giant peccary
Giant peccary
Pecari maximus

The collared peccary (*Dicotyles tajacu*) is a species of artiodactyl (even-toed) mammal in the family Tayassuidae found in North, Central, and South America. It is the only member of the genus *Dicotyles*. They are commonly referred to as javelina,...

Giant pika
Giant pika
Ochotona whartoni

The giant pika or Wharton's pika (*Ochotona whartoni*) is an extinct mammal species in the family Ochotonidae. It lived during the Pleistocene and early Holocene in northern parts of North America (Alaska, US and Canada). Very similar forms have...

Giant pocket gopher
Giant pocket gopher
Orthogeomys grandis

The giant pocket gopher (*Orthogeomys grandis*) is a species of rodent in the family Geomyidae. It is found in Mexico, Guatemala, El Salvador and Honduras. It is the

type species of the genus *Orthogeomys* ; some zoologists also include in this...
Giant roundleaf bat
Giant roundleaf bat
Macronycteris gigas

The giant roundleaf bat (*Macronycteris gigas*) is a species of bat in the family Hipposideridae found in western tropical Africa. Its natural habitats are subtropical or tropical moist lowland forests and caves. The species was formerly...
Giant Sable Antelope
Giant Sable Antelope
Hippotragus niger variani

The Giant sable antelope (*Hippotragus niger variani*) is a large, rare subspecies of the Sable antelope native and endemic to Angola. It is the national symbol of Angola and is held in great regard by its people. This was perhaps one of the...
Giant solenodon
Giant solenodon
Solenodon arredondoi

The giant solenodon (*Solenodon arredondoi*) is an extinct species of soricomorph that occurred in western Cuba. A larger animal than the still-extant Cuban solenodon (*Atopogale cubana*), the species is believed to have been eradicated through...
Giant tapir
Giant tapir
Tapirus augustus

The giant tapir (*Tapirus augustus*) is an extinct species of tapir that lived in southern China, Vietnam and Laos, with reports suggesting it also lived in Taiwan, Java, and potentially on Borneo. The species has been recorded from Middle and...
Giant thicket rat
Giant thicket rat
Grammomys gigas

The giant thicket rat (*Grammomys gigas*) is a species of rodent in the family Muridae which is restricted to a small area of Kenya.
Giant tree-rat
Giant tree-rat
Toromys grandis

The giant tree-rat (*Toromys grandis*) is a species in the family Echimyidae, the spiny rats. It is the only species in the monotypic genus *Toromys*. It is endemic to Brazil, where it occurs in the flooded forest along the banks of the Amazon River...
Giant white-tailed rat
Giant white-tailed rat
Uromys caudimaculatus

The giant white-tailed rat (*Uromys caudimaculatus*) is an Australian rodent native to tropical rainforest of north Queensland, with subspecies occurring in New Guinea and the Aru Islands. It is one of the largest rodents in Australia, reaching up...
Gilbert's dunnart
Gilbert's dunnart
Sminthopsis gilberti

Gilbert's dunnart (*Sminthopsis gilberti*) is a recently discovered dunnart, described in 1984. The length from snout to tail being 155–180 mm of which head to anus is 80–90 mm and tail 75–90 mm long. Hind foot size is 18 mm, ear length of 21

mm...

Gilbert's Potoroo
Gilbert's Potoroo
Potorous gilbertii

Gilbert's potoroo (*Potorous gilbertii*; formerly *Hypsiprymnus gilbertii*) is Australia's most endangered marsupial. It is the rarest marsupial in the world and one of the world's rarest critically endangered mammals. Gilbert's potoroo was thought...

Gilded tube-nosed bat
Gilded tube-nosed bat
Murina rozendaali

The gilded tube-nosed bat (*Murina rozendaali*) is a species of vesper bat in the family *Vespertilionidae*.

Gilliard's flying fox
Gilliard's flying fox
Pteropus gilliardorum

Gilliard's flying fox (*Pteropus gilliardorum*) is a species of flying fox in the family *Pteropodidae*. In Spanish, the common name is *zorro volador de Gilliard*. It is endemic to Papua New Guinea. It is known from only three specimens.

Giluwe rat
Giluwe rat
Rattus giluwensis

The Giluwe rat (*Rattus giluwensis*) is a species of rodent in the family *Muridae*. It is found only in Papua New Guinea, on Mount Giluwe and the subalpine grasslands of the Kaijende Highlands.

Ginkgo-toothed beaked whale
Ginkgo-toothed beaked whale
Mesoplodon ginkgodens

The ginkgo-toothed beaked whale (*Mesoplodon ginkgodens*) is a poorly known species of whale even for a beaked whale, and was named for the unusual shape of its dual teeth. It is a fairly typical-looking species, but is notable for the males not...

Giraffa priscilla
Giraffa priscilla
Giraffa priscilla

Giraffa priscilla is an extinct species of Miocene giraffe that lived on the Indian subcontinent. In 2014, new fossil remains were described in Pakistan.

Giraffa pygmaea
Giraffa pygmaea
Giraffa pygmaea

Giraffa pygmaea is an extinct species of giraffe from Africa during the Pliocene, and died out during the Pleistocene about 0.781 million years ago.

Giraffa sivalensis
Giraffa sivalensis
Giraffa sivalensis

Giraffa sivalensis is an extinct species of giraffe occurring in Asia during the Pliocene and Pleistocene epochs. Almost perfectly preserved cervical vertebrae have been found, as well as humeri, radii, metacarpals and teeth. The size of G....

Giraffa stillei
Giraffa stillei
Giraffa stillei

Giraffa stillei is an extinct species of giraffe endemic to Africa during the Pliocene to Pleistocene periods. It had a range from Malawi to Central Africa.
Giraffe
Giraffe
Giraffa camelopardalis

The Northern giraffe (Giraffa camelopardalis) is the world's tallest living animal native to North Africa. The giraffe has intrigued various ancient and modern cultures for its peculiar appearance and has often been featured in paintings, books, ...
Glacier Bay water shrew
Glacier Bay water shrew
Sorex alaskanus

The Glacier Bay water shrew (Sorex alaskanus) is a species of mammal in the family Soricidae. It is endemic to Alaska in the United States. It can swim underwater, and when it stops swimming, air trapped in its fur lets it float back up to the...
Glacier bear
Glacier bear
Ursus americanus emmonsii

The glacier bear (Ursus americanus emmonsii), sometimes referred to as the "blue bear", is a subspecies of American black bear with silver-blue or gray hair endemic from Southeast Alaska, to the extreme northwestern tip of British Columbia, and...
Glacier rat
Glacier rat
Rattus richardsoni

The glacier rat (Rattus richardsoni) is a species of rodent in the family Muridae. It is endemic to the New Guinea Highlands near Puncak Trikora (=Mt. Wilhelmina) and Puncak Jaya (=Carstensz Pyramid), West Papua, Indonesia. Its altitudinal range...
Gland-tailed free-tailed bat
Gland-tailed free-tailed bat
Chaerephon bemmeleni

The gland-tailed free-tailed bat (Chaerephon bemmeleni) is a species of bat in the family Molossidae. Its natural habitats are subtropical or tropical moist montane forests, dry savanna, and caves. It is threatened by habitat loss.
Glass's shrew
Glass's shrew
Crocidura glassi

The Glass's shrew (Crocidura glassi) is a species of mammal in the family Soricidae. It is endemic to Ethiopia. The mammal's natural habitats are subtropical or tropical high-altitude shrubland, subtropical or tropical high-altitude grassland, ...
Gleaning mouse
Gleaning mouse
Peromyscus spicilegus

The gleaning mouse (Peromyscus spicilegus) is a species of rodent in the family Cricetidae. It is found only in Mexico.
Glen's long-fingered bat
Glen's long-fingered bat
Miniopterus gleni

Glen's long-fingered bat (Miniopterus gleni) is a species of vesper bat in the family Vespertilionidae and it belongs to the class Mammalia. It is found only in Madagascar.

Glen's wattled bat
Glen's wattled bat
Glauconycteris gleni

Glen's wattled bat (*Glauconycteris gleni*) is a species of vesper bat in the family Vespertilionidae. It is found in Cameroon and Uganda. Its natural habitat is subtropical or tropical moist lowland forests. It is threatened by habitat loss.

Globicephala etruriae
Globicephala etruriae
Globicephala etruriae

Globicephala etruriae is an extinct species of pilot whale from the Pliocene of Italy. The type specimen was found in the Piacenzian coastal claystone at Volterra. It was named in 1987 by G. Pilleri.

Gloomy tube-nosed bat
Gloomy tube-nosed bat
Murina tenebrosa

The gloomy tube-nosed bat (*Murina tenebrosa*) is a species of vesper bat in the family Vespertilionidae. It is only known by the holotype, an old female, collected on Tsushima Island in 1962. It might be possibly extinct as Tsushima is badly...

Glover Allen's dasymys
Glover Allen's dasymys
Dasymys alleni

Glover Allen's dasymys (Glover Allen's shaggy rat or *Dasymys alleni*) is a species of shaggy marsh rat described in 1953, and indigenous to Mount Rungwe in south-western Tanzania. Contemporary studies of the taxonomy diverge, with the...

Glover's pika
Glover's pika
Ochotona gloveri

Glover's pika (*Ochotona gloveri*) is a species of mammal in the family Ochotonidae. It was first described in 1922, by Michael Rogers Oldfield Thomas. The summer dorsal pelage is grayish rufous, grayish brown, or tea brown in colour. The winter...

Gmelin's white-toothed shrew
Gmelin's white-toothed shrew
Crocidura gmelini

Gmelin's white-toothed shrew (*Crocidura gmelini*) is a species of mammal in the family Soricidae. It is found in Afghanistan, China, Iran, and Pakistan.

Gnome fruit-eating bat
Gnome fruit-eating bat
Dermanura gnoma

The gnome fruit-eating bat (*Dermanura gnoma*) is a bat species from South America. It is found in Bolivia, Brazil, Colombia, Ecuador, French Guiana, Guyana, Peru, Suriname and Venezuela. This species was originally discovered to be different from...

Goa
Goa
Procapra picticaudata

The goa (*Procapra picticaudata*) is a relatively small but graceful antelope that lives in the Tibetan plateau. It is also known as the Tibetan gazelle.

Goat
Goat
Capra aegagrus hircus

The goat or domestic goat (*Capra hircus*) is a domesticated species of goat-antelope typically kept as livestock. It was domesticated from the wild goat (*C. aegagrus*) of Southwest Asia and Eastern Europe. The goat is a member of the animal...

Gobi Altai mountain vole
Gobi Altai mountain vole
Alticola barakshin

The Gobi Altai mountain vole (*Alticola barakshin*) is a species of rodent in the family Cricetidae. It can be found in China, Mongolia, and the Russian Federation.
Gobi bear
Gobi bear
Ursus arctos gobiensis

The Gobi bear (*Ursus arctos gobiensis*), known in Mongolian as the Mazaalai (Мазаалай), is a subspecies of the brown bear (*Ursus arctos*) that is found in the Gobi Desert of Mongolia. It is listed as critically endangered by the Mongolian Redbook...
Gobi big brown bat
Gobi big brown bat
Eptesicus gobiensis

The Gobi big brown bat (*Eptesicus gobiensis*) is a species of vesper bat. It is found in Afghanistan, China, India, Mongolia, Pakistan, and Russia. Russian zoologist Professor Count Nikolay Alekseyevich Bobrinski first described it in 1926, the...
Gobi jerboa
Gobi jerboa
Allactaga bullata

The Gobi jerboa (*Allactaga bullata*) is a species of rodent in the family Dipodidae. It is found in China and Mongolia. Its natural habitats are temperate grassland and temperate desert. The Gobi jerboa was discovered in 1925 by Glover Morrill...
Godman's long-tailed bat
Godman's long-tailed bat
Choeroniscus godmani

Godman's long-tailed bat (*Choeroniscus godmani*) is a species of bat in the family Phyllostomidae. It is found in Colombia, Costa Rica, El Salvador, Guatemala, Guyana, Honduras, Mexico, Nicaragua, Suriname, and Venezuela.
Godman's rock-wallaby
Godman's rock-wallaby
Petrogale godmani

Godman's rock-wallaby (*Petrogale godmani*) is a diprotodont marsupial, and a typical rock-wallaby. It is found in northern and north-eastern Queensland, Australia. This rock-wallaby is found in low open forest, open scrub, or montane regions, ...
Goeldi's Marmoset
Goeldi's Marmoset
Callimico goeldii

The Goeldi's marmoset or Goeldi's monkey (*Callimico goeldii*) is a small, South American New World monkey that lives in the upper Amazon basin region of Bolivia, Brazil, Colombia, Ecuador, and Peru. It is the only species classified in the genus...
Goeldi's spiny rat

Goeldi's spiny rat
Proechimys goeldii

Goeldi's spiny-rat, *Proechimys goeldii*, is a spiny rat species from South America. It is found in Brazil.

Goff's pocket gopher
Goff's pocket gopher
Geomys pinetis goffi

Goff's southeastern pocket gopher (*Geomys pinetis goffi*) was a pocket gopher endemic to Brevard County, Florida, United States. The last sightings recorded were in 1955. They burrowed and lived mostly underground eating mainly underground...

Goitered Gazelle
Goitered Gazelle
Gazella subgutturosa

The goitered or black-tailed gazelle (*Gazella subgutturosa*) is a gazelle found in Georgia, Azerbaijan, Iran, parts of Iraq and Pakistan, Afghanistan, Tajikistan, Kyrgyzstan, Uzbekistan, Turkmenistan, Kazakhstan and in northwest China and...

Golan Heights blind mole-rat
Golan Heights blind mole-rat
Spalax golani

The Middle East blind mole-rat or Palestine mole-rat (*Nannospalax ehrenbergi*) is a species of rodent in the family Spalacidae.

Gold-and-white marmoset
Gold-and-white marmoset
Mico chrysoleucos

The gold-and-white marmoset (*Mico chrysoleucos*), also known as the golden-white tassel-ear marmoset, is a species of marmoset, a small monkey endemic to the Amazon rainforest in eastern Amazonas state, Brazil.

Golden angwantibo
Golden angwantibo
Arctocebus aureus

The golden angwantibo (*Arctocebus aureus*) is a strepsirrhine primate of the family Lorisidae. It shares the genus *Arctocebus* with the Calabar angwantibo (*Arctocebus calabarensis*) and together they are commonly called the golden pottos. The...

Golden Atlantic tree-rat
Golden Atlantic tree-rat
Phyllomys blainvillii

The golden Atlantic tree-rat, *Phyllomys blainvillii*, is a spiny rat species from South America. It is found in Brazil.

Golden Bamboo Lemur
Golden Bamboo Lemur
Haplemur aureus

The golden bamboo lemur (*Haplemur aureus*), bokombolomena or varibolomena in Malagasy, is a medium-sized bamboo lemur endemic to south-eastern Madagascar.

Golden Bandicoot
Golden Bandicoot
Isodon auratus

The golden bandicoot (*Isodon auratus*) is a short-nosed bandicoot found in northern Australia. It is the smallest of its genus. The golden bandicoot is now a threatened species. It was once found throughout much of northwestern Australia, with...

Golden bat
Golden bat
Mimon bennettii

The golden bat (*Mimon bennettii*) is a bat species from South America. It is found in Brazil, Colombia, French Guiana, Guyana, Suriname and Venezuela.

Golden Jackal
Golden Jackal
Canis aureus

The Golden jackal (*Canis aureus*) is a wolf-like canid. Despite its name, the Golden jackal is not closely related to the African Black-backed jackal or Side-striped jackal, which are part of the genus *Lupulella*, being instead closer to wolves and...

Golden Lion Tamarin
Golden Lion Tamarin
Leontopithecus rosalia

The Golden lion tamarin (*Leontopithecus rosalia*) is a small, endangered primate, endemic to the Atlantic coast of Brazil. It is a member of the family *Callitrichidae*. In spite of its name, this monkey isn't related to lions. Instead, this species...

Golden Monkey
Golden Monkey
Cercopithecus kandti

The golden monkey (*Cercopithecus kandti*) is a species of Old World monkey found in the Virunga volcanic mountains of Central Africa, including four national parks: Mgahinga, in south-west Uganda; Volcanoes, in north-west Rwanda; and Virunga and...

Golden mouse
Golden mouse
Ochrotomys nuttalli

The golden mouse (*Ochrotomys nuttalli*) is a species of New World mouse. It is usually 5-8 inches (12-25 cm) in body length, and has a soft pelage that ranges from golden-brownish to burnt orange in color. The genus name comes from the Greek...

Golden Oldfield mouse
Golden Oldfield mouse
Thomasomys aureus

The golden Oldfield mouse or golden thomomys (*Thomasomys aureus*) is a species of rodent in the family *Cricetidae*. It is found in Colombia, Ecuador, Peru, and Venezuela. Both the common and genus name commemorate the British zoologist Oldfield...

Golden Palm Civet
Golden Palm Civet
Paradoxurus zeylonensis

The Golden palm civet (*Paradoxurus zeylonensis*) is a viverrid that was described by Peter Simon Pallas in 1778. Its distribution is severely fragmented, and the extent and quality of its habitat in Sri Lanka's hill regions are declining.

Golden Snub-Nosed Monkey
Golden Snub-Nosed Monkey
Rhinopithecus roxellana

The Golden snub-nosed monkey (*Rhinopithecus roxellana*) is an Old World monkey endemic to a small area in temperate, mountainous forests of central and Southwest China. Snow occurs frequently within its range, and it can withstand colder

average...

Golden spiny mouse

Golden spiny mouse

Acomys russatus

The golden spiny mouse (*Acomys russatus*) gets its name from the reddish-orange spiny fur that covers its body from head to tail. This coarse, inflexible fur is thought to protect it from predation. Aside from the golden fur that covers its head...

Golden Takin

Golden Takin

Budorcas taxicolor bedfordi

The Golden takin (*Budorcas taxicolor bedfordi*) is a threatened subspecies of takin, native to China's. Takins have unique adaptations that help them stay warm and dry during cold Himalayan winters. Their large snout has sinus cavities that heat...

Golden viscacha rat

Golden viscacha rat

Pipanacoctomys aureus

The golden viscacha rat or golden vizcacha rat (*Pipanacoctomys aureus*) is the single species of the genus *Pipanacoctomys* of the rodent family Octodontidae. It has 92 chromosomes and has been regarded as tetraploid ($4x = 2n$). This octodontid and...

Golden-backed tree-rat

Golden-backed tree-rat

Mesembriomys macrurus

The golden-backed tree rat (*Mesembriomys macrurus*) is a species of rodent in the family Muridae, found only in Australia. It is present in the Charnley River-Artesian Range Wildlife Sanctuary in the Kimberley region of Western Australia. They...

Golden-backed uakari

Golden-backed uakari

Cacajao melanocephalus

The golden-backed uakari (*Cacajao melanocephalus*) or black-headed uakari, is a New World primate from the family Pitheciidae. It lives in the Amazon Rainforest, and is found in the countries of Brazil, Colombia, and Venezuela. It has black hair...

Golden-Bellied Capuchin

Golden-Bellied Capuchin

Sapajus xanthosternos

The golden-bellied capuchin (*Sapajus xanthosternos*), also known as the yellow-breasted or buff-headed capuchin, is a species of New World or neotropical monkey. It lives mainly in trees and are omnivorous, eating a wide variety of both plant and...

Golden-Bellied Mangabey

Golden-Bellied Mangabey

Cercocebus chrysogaster

The golden-bellied mangabey (*Cercocebus chrysogaster*) is a social Old World monkey found in swampy, humid forests south of the Congo River in the Democratic Republic of the Congo. It was previously considered a subspecies of the agile mangabey...

Golden-bellied treeshrew

Golden-bellied treeshrew

Tupaia chrysogaster

The golden-bellied treeshrew (*Tupaia chrysogaster*) is a treeshrew species within

the Tupaiidae. It is also called Mentawai treeshrew as it is endemic to the Indonesian Mentawai islands of Sipora, North and South Pagai. It lives in forests, and is...

Golden-brown mouse lemur
Golden-brown mouse lemur
Microcebus ravelobensis

The golden-brown mouse lemur (*Microcebus ravelobensis*), also known as the (Lac) Ravelobe mouse lemur, is part of the Cheirogaleidae family, and the smallest species of lemur. It is arboreal, nocturnal and usually social. It get its name from the...

Golden-Crowned Sifaka
Golden-Crowned Sifaka
Propithecus tattersalli

The golden-crowned sifaka or Tattersall's sifaka (*Propithecus tattersalli*) is a medium-sized lemur characterized by mostly white fur, prominent furry ears, and a golden-orange crown. It is one of the smallest sifakas (genus *Propithecus*),...

Golden-faced saki
Golden-faced saki
Pithecia chrysocephala

The golden-faced saki (*Pithecia chrysocephala*) is a species of saki monkey, a type of New World monkey. It is found in Brazil north of the Amazon, on both sides of the Rio Negro. This species was formerly considered a subspecies of the...

Golden-Headed Lion Tamarin
Golden-Headed Lion Tamarin
Leontopithecus chrysomelas

The golden-headed lion tamarin (*Leontopithecus chrysomelas*), also the golden-headed tamarin, is a lion tamarin endemic to Brazil. It is found only in the lowland and premontane tropical forest fragments in the state of Bahia, and therefore is...

Golden-Mantled Ground Squirrel
Golden-Mantled Ground Squirrel
Callospermophilus lateralis

The golden-mantled ground squirrel (*Callospermophilus lateralis*) is a ground squirrel native to western North America. It is distributed in the Rocky Mountains of British Columbia and Alberta, and through much of the western United States.

Golden-mantled howler
Golden-mantled howler
Alouatta palliata palliata

The golden-mantled howler (*Alouatta palliata palliata*) is a subspecies of the mantled howler, *A. palliata*. It ranges throughout much of Central America, in Guatemala, Honduras, Nicaragua and Costa Rica, and possibly Panama. The range limits...

Golden-mantled tamarin
Golden-mantled tamarin
Leontocebus tripartitus

The golden-mantled tamarin (*Leontocebus tripartitus*) is a tamarin species from South America. It is found in Ecuador and Peru, specifically in the upper Amazon (lowland), east of the Andes in Ecuador, and Northeast Peru; between the Rio Curaray...

Golden-Mantled Tree-Kangaroo
Golden-Mantled Tree-Kangaroo
Dendrolagus pulcherrimus

The golden-mantled tree-kangaroo (*Dendrolagus pulcherrimus*) is a species of tree-kangaroo endemic to the Northern New Guinea montane rain forests ecoregion of the island.

Golden-rumped elephant shrew

Golden-rumped elephant shrew

Rhynchocyon chrysopygus

The golden-rumped elephant shrew (*Rhynchocyon chrysopygus*) is a small African mammal. It is the largest species of the elephant shrew family along with its close relative the grey-faced sengi. It is classified as endangered.

Golden-tipped bat

Golden-tipped bat

Phoniscus papuensis

The golden-tipped bat (*Kerivoula papuensis*) is a species of Microchiropteran in the family Vespertilionidae. It is found in Papua New Guinea and in Australia, especially scattered along the eastern part of Australia. The species is considered...

Goldman's broad-clawed shrew

Goldman's broad-clawed shrew

Cryptotis goldmani

Goldman's broad-clawed shrew (*Cryptotis goldmani*) is a species of mammal in the family Soricidae. It is found in Guatemala and Mexico.

Goldman's diminutive woodrat

Goldman's diminutive woodrat

Nelsonia goldmani

Goldman's diminutive woodrat (*Nelsonia goldmani*) is a species of rodent in the family Cricetidae. It is found only in Mexico.

Goldman's nectar bat

Goldman's nectar bat

Lonchophylla mordax

Goldman's nectar bat (*Lonchophylla mordax*) is a bat species from South and Central America. It is found in Brazil, Colombia, Costa Rica, Ecuador and Panama.

Goldman's pocket gopher

Goldman's pocket gopher

Cratogeomys goldmani

Goldman's pocket gopher, (*Cratogeomys goldmani*) is a species of rodent in the pocket gopher family. It is distributed throughout northern Mexico.

Goldman's pocket mouse

Goldman's pocket mouse

Chaetodipus goldmani

Goldman's pocket mouse (*Chaetodipus goldmani*) is a species of rodent in the family Heteromyidae. It is endemic to Mexico, where it is threatened by the increasing conversion of its dry, scrubby habitat into agricultural land. As a result, the...

Goldman's spiny pocket mouse

Goldman's spiny pocket mouse

Heteromys goldmani

Goldman's spiny pocket mouse (*Heteromys goldmani*) is a species of rodent in the family Heteromyidae. It has been reported to be a subspecies of *Heteromys desmarestianus*. It is found in Guatemala and Mexico. Its natural habitat is subtropical or...

Goldman's water mouse

Goldman's water mouse
Rheomys raptor

Rheomys raptor, also known as the Goldman's water mouse or Goldman's Ichthyomyine, is a species of rodent in the family Cricetidae. This mouse is semiaquatic and its carnivorous diet includes invertebrates. The conservation status of the species...

Goliath shrew
Goliath shrew
Crocidura goliath

The goliath shrew (*Crocidura goliath*) is a species of mammal in the family Soricidae. It is found in Cameroon, Central African Republic, Republic of the Congo, Democratic Republic of the Congo, Equatorial Guinea, and Gabon. Its natural habitat...

Gomantong myotis
Gomantong myotis
Myotis gomantongensis

The Gomantong myotis (*Myotis gomantongensis*) is a species of bat in the family Vespertilionidae that is endemic to Sabah district of Malaysia.

Gongshan muntjac
Gongshan muntjac
Muntiacus gongshanensis

The Gongshan muntjac (*Muntiacus gongshanensis*) is a species of muntjac (a type of deer) living in the Gongshan mountains in northwestern Yunnan, southeast Tibet, Northeast India (especially in Arunachal Pradesh) and northern Myanmar.Ongoing...

Goodfellow's Tree-Kangaroo
Goodfellow's Tree-Kangaroo
Dendrolagus goodfellowi

Goodfellow's tree-kangaroo (*Dendrolagus goodfellowi*) also called the ornate tree-kangaroo, belongs to the family Macropodidae, which includes kangaroos, wallabies and their relatives, and the genus *Dendrolagus*, with eleven other species. The...

Goodfellow's tuco-tuco
Goodfellow's tuco-tuco
Ctenomys goodfellowi

Goodfellow's tuco-tuco (*Ctenomys goodfellowi*) is a species of rodent in the family Ctenomyidae. It is endemic to Bolivia, where it is found in the Chiquitano dry forest ecoregion, bordering on the cerrado. Its karyotype has $2n = 46$ and $FN = 68$

Goodman's Mouse Lemur
Goodman's Mouse Lemur
Microcebus lehilahytsara

Goodman's mouse lemur (*Microcebus lehilahytsara*) is a species of mouse lemur from the region near Andasibe in eastern Madagascar. The species is named in honor of primatologist Steven M. Goodman. "Lehilahytsara" is a combination of the Malagasy...

Goodwin's broad-clawed shrew
Goodwin's broad-clawed shrew
Cryptotis goodwini

Goodwin's broad-clawed shrew (*Cryptotis goodwini*) is a species of mammal in the shrew family, Soricidae. Body length and size of adults average 9.49 cm and 7.0 grams respectively making it a relatively larger shrew. What distinguishes this from...

Gorongoza gerbil

Gorongoza gerbil
Gerbilliscus inclusus

The Gorongoza gerbil (*Gerbilliscus inclusus*) is a species of rodent found in Mozambique, Tanzania, and Zimbabwe. Its natural habitat is moist savanna.

Gorontalo macaque
Gorontalo macaque
Macaca nigrescens

The Gorontalo macaque or Dumoga-bone macaque (*Macaca nigrescens*) is a species of primate in the family Cercopithecidae. It is endemic to the island of Sulawesi in Indonesia.

Gotel Mountain soft-furred mouse
Gotel Mountain soft-furred mouse
Praomys obscurus

The Gotel Mountain soft-furred mouse (*Praomys obscurus*) or Gotel Mountain praomys, is a species of rodent in the family Muridae. It is endemic to the Gotel Mountains in southeastern Nigeria. It occurs in fern-grassland, montane forest, along...

Gould's long-eared bat
Gould's long-eared bat
Nyctophilus gouldi

Gould's long-eared bat (*Nyctophilus gouldi*) is a microbat found in southern regions of Australia. It occurs in eastern Australia, from Queensland to Victoria, and in a smaller isolated range in the south-west of Western Australia.

Gould's wattled bat
Gould's wattled bat
Chalinolobus gouldii

Gould's wattled bat (*Chalinolobus gouldii*) is a species of Australian wattled bat named after the English naturalist John Gould.

Gounda mouse
Gounda mouse
Mus goundae

The Gounda mouse (*Mus goundae*) is a species of rodent in the family Muridae. It is found only in Central African Republic, around the Gounda River. Its natural habitat is moist savanna.

Goya tuco-tuco
Goya tuco-tuco
Ctenomys perrensi

The Goya tuco-tuco (*Ctenomys perrensi*) is a species of rodent in the family Ctenomyidae. It is endemic to Argentina.

Gracile Atlantic spiny rat
Gracile Atlantic spiny rat
Trinomys graciosus

Trinomys graciosus is a species in the mainly South American family Echimyidae, the spiny rats; it occurs in southeast Brazil from the south bank of the River Doce, Espirito Santo, southward to Teresopolis, Rio de Janeiro.

Gracile naked-tailed shrew
Gracile naked-tailed shrew
Crocidura maurisca

The gracile naked-tailed shrew (*Crocidura maurisca*) is a species of mammal in the family Soricidae. It is found in Burundi, Gabon, Kenya, and Uganda. Its natural habitat is swamps.

Gracile shrew mole
Gracile shrew mole
Uropsilus gracilis

The gracile shrew mole (*Uropsilus gracilis*) is a species of mammal in the family Talpidae. It is endemic to China; populations known from Myanmar likely represent other species (see below).The black-backed shrew mole (*U. atronates*) and the Snow...

Gracile shrew tenrec
Gracile shrew tenrec
Microgale gracilis

The gracile shrew tenrec (*Microgale gracilis*) is a species of mammal in the family Tenrecidae. It is endemic to Madagascar. Its natural habitat is subtropical or tropical moist montane forests.

Gracile tateril
Gracile tateril
Taterillus gracilis

The gracile tateril or slender gerbil (*Taterillus gracilis*) is a species of rodent found in Burkina Faso, Chad, Gambia, Ghana, Guinea, Ivory Coast, Mali, Niger, Nigeria, Senegal, Togo, and possibly Cameroon. Its natural habitats are dry savanna,...

Graells's tamarin
Graells's tamarin
Leontocebus nigricollis graellsii

Graells's tamarin, *Saguinus nigricollis graellsii*, is a subspecies of the black-mantled tamarin from the northwestern Amazon in southeastern Colombia, eastern Ecuador and northeastern Peru. It differs from other black-mantled tamarins in having a...

Grammomys brevirostris
Grammomys brevirostris
Grammomys brevirostris

Grammomys brevirostris is a species of rodent endemic to Kenya.

Granada hare
Granada hare
Lepus granatensis

The Granada hare (*Lepus granatensis*), also known as the Iberian hare, is a hare species that can be found on the Iberian Peninsula and on the island of Majorca.

Grand roundleaf bat
Grand roundleaf bat
Hipposideros grandis

The grand roundleaf bat (*Hipposideros grandis*), also known as the grand leaf-nosed bat, is a species of bat in the genus *Hipposideros*. It can be found in China, Myanmar, Thailand, and Vietnam.

Grandidier's free-tailed bat
Grandidier's free-tailed bat
Chaerephon leucogaster

Grandidier's free-tailed bat (*Chaerephon leucogaster*) is a species of bat in the family Molossidae. It is endemic to Madagascar. Its natural habitats are subtropical or tropical dry forests and dry savanna. It is threatened by habitat loss. It...

Grandidier's mongoose
Grandidier's mongoose

Galidictis grandidieri

Grandidier's mongoose (*Galidictis grandidieri*), also known as the giant-striped mongoose or Grandidier's vontsira, is a small carnivoran that lives only in a very small area of southwestern Madagascar, in areas of spiny forest vegetation. It is...

Grandidier's trident bat

Grandidier's trident bat

Paratriaenops auritus

Grandidier's trident bat (*Paratriaenops auritus*) is a species of bat in the family Hipposideridae endemic to Madagascar. It was formerly assigned to the genus *Triaienops*, but is now placed in the separate genus *Paratriaenops*.

Grandidier's tufted-tailed rat

Grandidier's tufted-tailed rat

Eliurus grandidieri

Grandidier's tufted-tailed rat (*Eliurus grandidieri*) is a species of rodent from the family Nesomyidae. Morphological evidence suggests that Grandidier's tufted-tailed rat is most closely related to Petter's tufted-tailed rat, *E. petterii*....

Grant's bushbaby

Grant's bushbaby

Paragalago granti

Grant's bushbaby (*Paragalago granti*), also known as Grant's lesser bushbaby or the Mozambique lesser bushbaby, is a species of primate in the family Galagidae. It is found in Malawi, Mozambique, Tanzania, and Zimbabwe. Its natural habitat is...

Grant's forest shrew

Grant's forest shrew

Sylvisorex granti

Grant's forest shrew (*Sylvisorex granti*) is a species of mammal in the family Soricidae found in the mountain forests of central Democratic Republic of the Congo, Kenya, Rwanda, Tanzania, and Uganda. Its natural habitat is subtropical or...

Grant's Gazelle

Grant's Gazelle

Nanger granti

Grant's gazelle (*Nanger granti*) is a species of gazelle distributed from northern Tanzania to South Sudan and Ethiopia, and from the Kenyan coast to Lake Victoria. Its Swahili name is swala granti. It was named for a 19th-century British...

Grant's golden mole

Grant's golden mole

Eremitalpa granti

Grant's golden mole (*Eremitalpa granti* ; colloquially also: dune shark) is a golden mole species. It is the only member of the genus *Eremitalpa*.

Grant's rock mouse

Grant's rock mouse

Aethomys granti

Grant's rock mouse or Grant's rock rat (*Aethomys granti*) is a species of rodent in the family Muridae found only in South Africa. It is sometimes included in the genus *Micaelamys*. Its natural habitats are subtropical or tropical dry shrubland and...

Grant's zebra

Grant's zebra

Equus quagga boehmi

Grant's zebra (*Equus quagga boehmi*) is the smallest of the seven subspecies of the plains zebra. This subspecies represents the zebra form of the Serengeti-Mara ecosystem.

Grasse's shrew

Grasse's shrew

Crocidura grassei

The Grasse's shrew (*Crocidura grassei*) is a species of mammal in the family Soricidae. It is found in Cameroon, Central African Republic, Republic of the Congo, Gabon, and Equatorial Guinea. Its natural habitat is subtropical or tropical moist...

Grassland mosaic-tailed rat

Grassland mosaic-tailed rat

Melomys burtoni

The grassland mosaic-tailed rat, or grassland melomys (*Melomys burtoni*), is a species of rodent in the family Muridae. It is found in Australia and Papua New Guinea. In Australia it is found along the northern coast from Kimberley to New South...

Grauer's large-headed shrew

Grauer's large-headed shrew

Paracrocidura graueri

Grauer's large-headed shrew (*Paracrocidura graueri*) is a species of mammal in the family Soricidae. It is endemic to Democratic Republic of the Congo. Its natural habitat is subtropical or tropical moist montane forests.

Gray Bat

Gray Bat

Myotis grisescens

The Gray bat (*Myotis grisescens*) is a species of microbat endemic to North America. It once flourished in caves all over the southeastern United States, but due to human disturbance, gray bat populations declined severely during the early and mid...

Gray birch mouse

Gray birch mouse

Sicista pseudonapaea

The gray birch mouse (*Sicista pseudonapaea*) is a species of rodent in the family Sminthidae. It is endemic to Kazakhstan, but possibly ranges into China. Its natural habitat is temperate forests.

Gray Brocket

Gray Brocket

Mazama gouazoubira

The Gray brocket (*Mazama gouazoubira*) is a species of brocket deer from South America. Unlike other species of brocket deer in its range, the Gray brocket has gray-brown fur without reddish tones.

Gray brush-furred rat

Gray brush-furred rat

Lophuromys aquilus

The gray brush-furred rat (*Lophuromys aquilus*) is a species of rodent in the family Muridae. It is found only in Democratic Republic of the Congo. Its natural habitats are subtropical or tropical swamps, subtropical or tropical moist montane...

Gray climbing mouse

Gray climbing mouse

Dendromus melanotis

The gray climbing mouse (*Dendromus melanotis*) is a species of rodent in the family Nesomyidae. It is found in Angola, Benin, Botswana, Democratic Republic of the Congo, Ethiopia, Guinea, Liberia, Malawi, Mozambique, Namibia, Nigeria, Rwanda, ...
Gray dorcopsis
Gray dorcopsis
Dorcopsis luctuosa

The gray dorcopsis or gray forest wallaby (*Dorcopsis luctuosa*) is a species of marsupial in the family Macropodidae. It is found in West Papua, Indonesia and Papua New Guinea.
Gray flying fox
Gray flying fox
Pteropus griseus

The gray flying fox (*Pteropus griseus*) is a species of flying fox in the family Pteropodidae. It is not to be confused with the grey flying fox (*Pteropus poliocephalus*). It is found in Indonesia, but not in the Philippines, despite occasional...
Gray Four-Eyed Opossum
Gray Four-Eyed Opossum
Philander opossum

The gray four-eyed opossum (*Philander opossum*) is an opossum species from Central and South America, ranging from southern Mexico to Peru, Bolivia and southwestern Brazil, at altitudes from sea level to 1600 m, but generally below 1,000 metres...
Gray Fox
Gray Fox
Urocyon cinereoargenteus

The Gray fox (*Urocyon cinereoargenteus*) is an omnivorous mammal of the family Canidae widely distributed throughout North America and Central America. It was once the most common fox in the eastern United States, and though still found there, ...
Gray leaf-eared mouse
Gray leaf-eared mouse
Graomys griseoflavus

The gray leaf-eared mouse (*Graomys griseoflavus*) is a rodent species from South America. It is found in Argentina, Bolivia, Brazil and Paraguay; its habitat includes the Gran Chaco. This is a variable and widely distributed species that can be...
Gray long-tongued bat
Gray long-tongued bat
Glossophaga leachii

The gray long-tongued bat (*Glossophaga leachii*) is a species of bat in the family Phyllostomidae. It is found in Costa Rica, El Salvador, Guatemala, Honduras, Mexico, and Nicaragua.
Gray Marmot
Gray Marmot
Marmota baibacina

The Gray marmot (*Marmota baibacina*) is a rodent that lives in Asia. It is one of the larger marmots in its genus. Gray marmots form social groups, live in burrows, and hibernate.
Gray Mouse Lemur
Gray Mouse Lemur
Microcebus murinus

The gray mouse lemur (*Microcebus murinus*), grey mouse lemur or lesser mouse lemur, is a small lemur, a type of strepsirrhine primate, found only on the island of Madagascar. Weighing 58 to 67 grams (2.0 to 2.4 oz), it is the largest of the mouse...

Gray sac-winged bat

Gray sac-winged bat

Balantiopteryx plicata

The gray sac-winged bat (*Balantiopteryx plicata*) is a species in the family Emballonuridae which comprises the 51 species of sac-winged bats. It is found in Mexico from Baja California Sur and Sonora to Guatemala, El Salvador, Honduras,...

Gray Seal

Gray Seal

Halichoerus grypus

The Gray seal (*Halichoerus grypus*) is found on both shores of the North Atlantic Ocean. In Latin *Halichoerus grypus* means "hook-nosed sea pig". It is a large seal of the family Phocidae, which are commonly referred to as "true seals" or "earless..."

Gray short-tailed bat

Gray short-tailed bat

Carollia subrufa

The gray short-tailed bat, or Hahn's short-tailed bat (*Carollia subrufa*), is a species of bat in the family Phyllostomidae native to Mexico and Central America.

Gray Short-Tailed Opossum

Gray Short-Tailed Opossum

Monodelphis domestica

The Gray short-tailed opossum (*Monodelphis domestica*) is a South American member of the family Didelphidae. Unlike most other marsupials, it does not have a true pouch. The Gray short-tailed opossum is used as a research model in science and is...

Gray Slender Loris

Gray Slender Loris

Loris lydekkerianus

The gray slender loris (*Loris lydekkerianus*) is a species of primate in the family Loridae. It is native to India and Sri Lanka and inhabits subtropical and tropical dry forests and subtropical or tropical moist lowland forests. It is threatened...

Gray slender opossum

Gray slender opossum

Marmosops incanus

The gray slender opossum (*Marmosops incanus*), is an opossum species from Brazil. This species is a semi-arboreal marsupial, moving on average 67.38% on the ground. They are solitary, nocturnal, and scansorial (tree climbers). They live off a diet...

Gray snub-nosed monkey

Gray snub-nosed monkey

Rhinopithecus brelichi

The gray snub-nosed monkey (*Rhinopithecus brelichi*), also known as Brelich's snub-nosed monkey, Guizhou snub-nosed monkey, and Guizhou golden monkey, is a species of primate in the family Cercopithecidae. It is endemic to China, where it is...

Gray spiny mouse

Gray spiny mouse

Acomys cineraceus

The gray spiny mouse (*Acomys cineraceus*) is a species of rodent in the family Muridae found in Ethiopia, Kenya, South Sudan, and Uganda. Its natural habitats are dry savanna, moist savanna, rocky areas, arable land, and rural gardens.

Gray tree rat

Gray tree rat

Lenothrix canus

{{Not to be confused with eastern grey squirrel, which is commonly referred to as a "tree rat" in its introduced range}}The gray tree rat (*Lenothrix canus*) is a species of rodent in the family Muridae and the only species in the monotypic genus...

Gray Whale

Gray Whale

Eschrichtius robustus

The Gray whale (*Eschrichtius robustus*) is a baleen whale that migrates between feeding and breeding grounds yearly. The common name of this whale comes from the gray patches and white mottling on its dark skin. Gray whales were once called devil...

Gray woolly monkey

Gray woolly monkey

Lagothrix cana

The gray woolly monkey (*Lagothrix lagothricha cana*) or Geoffroy's woolly monkey is a subspecies of the common woolly monkey from South America. It is found in Bolivia, Brazil and Peru. *L. l. cana* gets its common name, gray woolly monkey, from...

Gray-backed sportive lemur

Gray-backed sportive lemur

Lepilemur dorsalis

The gray-backed sportive lemur (*Lepilemur dorsalis*), also known as Gray's sportive lemur or back-striped sportive lemur, is a species of lemur in the family Lepilemuridae. It is endemic to Madagascar. It is threatened by habitat loss.

Gray-bellied caenolestid

Gray-bellied caenolestid

Caenolestes caniventer

The gray-bellied caenolestid (*Caenolestes caniventer*), or grey-bellied shrew opossum, is a shrew opossum found in humid, temperate forests and moist grasslands of western Ecuador and northwestern Peru. It was first described by American...

Gray-bellied mountain rat

Gray-bellied mountain rat

Limnomys bryophilus

The gray-bellied mountain rat, also known as the gray-bellied limnomys and the buffy-collared moss-mouse (*Limnomys bryophilus*), is a species of rodent in the family Muridae. It is found only in the Philippines, where it is common at high...

Gray-bellied night monkey

Gray-bellied night monkey

Aotus lemurinus

The gray-bellied night monkey (*Aotus lemurinus*), also called the grey-legged douroucoulis or lemurine owl monkey, is a small New World monkey of the family Aotidae. Native to tropical and subtropical forests of South America, the gray-bellied...

Gray-bellied pencil-tailed tree mouse

Gray-bellied pencil-tailed tree mouse

Chiropodomys muroides

The gray-bellied pencil-tailed tree mouse (*Chiropodomys muroides*) is a species of arboreal rodent in the family Muridae. It is endemic to Borneo where it is known from Gunung Kinabalu (Sabah, Malaysia) and from Long Petak in northern Kalimantan...

Gray-bellied pygmy mouse

Gray-bellied pygmy mouse

Mus triton

The gray-bellied pygmy mouse (*Mus triton*) is a species of rodent in the family Muridae. It is found in Angola, Burundi, Democratic Republic of the Congo, Kenya, Malawi, Mozambique, Rwanda, Tanzania, Uganda, and Zambia. Its natural habitats are...

Gray-bellied tree mouse

Gray-bellied tree mouse

Pogonomys sylvestris

The gray-bellied tree mouse (*Pogonomys sylvestris*) is a species of rodent in the family Muridae. It is found only in Papua New Guinea. It is consumed by the Kalam people of Papua New Guinea, and is hunted by women who look for it in its burrows.

Gray-cheeked flying squirrel

Gray-cheeked flying squirrel

Hylopetes lepidus

The arrow flying squirrel (*Hylopetes sagitta*) is a species of flying squirrel. It is endemic to Java and Bangka, Indonesia. The population is unknown as it has only been collected from a few localities. It is nocturnal and arboreal and may be...

Gray-collared chipmunk

Gray-collared chipmunk

Neotamias cinereicollis

The gray-collared chipmunk (*Neotamias cinereicollis*) is a species of rodent in the family Sciuridae. It is endemic to Arizona and New Mexico in the United States.

Gray-footed chipmunk

Gray-footed chipmunk

Neotamias canipes

The gray-footed chipmunk (*Neotamias canipes*) is a species of rodent in the family Sciuridae. It is endemic to New Mexico and in the Sierra Diablo and Guadalupe Mountains in the Trans-Pecos region of Texas in the United States. Its natural...

Gray-footed spiny rat

Gray-footed spiny rat

Proechimys poliopterus

The gray-footed spiny rat (*Proechimys poliopterus*) is a species of rodent in the family Echimyidae. It is found in Colombia and Venezuela.

Gray-handed night monkey

Gray-handed night monkey

Aotus griseimembra

The gray-handed night monkey (*Aotus griseimembra*) is a species of night monkey formerly considered a subspecies of Gray-bellied night monkey of the family Aotidae. Its range consists of parts of Colombia and Venezuela. The exact classification...

Gray-headed lemur

Gray-headed lemur

Eulemur cinereiceps

The gray-headed lemur (*Eulemur cinereiceps*), or gray-headed brown lemur, is a medium-sized primate, a cathemeral species of lemur in the family Lemnidae. Until a taxonomic revision in 2008, it was known as the white-collared brown lemur or...
Gray-headed thicket rat
Gray-headed thicket rat
Grammomys caniceps

The gray-headed thicket rat (*Grammomys caniceps*) is a species of rodent in the family Muridae. It is found in Kenya and Somalia. Its natural habitat is subtropical or tropical dry shrubland.
Gray-shanked douc
Gray-shanked douc
Pygathrix cinerea

The grey-shanked douc langur (*Pygathrix cinerea*) is a douc species native to the Vietnamese provinces of Quảng Nam, Quảng Ngãi, Bình Định, Kon Tum, and Gia Lai. The total population was estimated at 550 to 700 individuals in 2004. In 2016, Dr....
Gray-tailed narrow-headed rat
Gray-tailed narrow-headed rat
Stenocephalemys griseicauda

The gray-tailed narrow-headed rat (*Stenocephalemys griseicauda*) is a species of rodent in the family Muridae. It is found only in Ethiopia. Its natural habitats are subtropical or tropical moist montane forests, subtropical or tropical...
Gray-tailed vole
Gray-tailed vole
Microtus canicaudus

The gray-tailed vole (*Microtus canicaudus*) also known as the gray-tailed meadow vole or gray-tailed meadow mouse, is a rodent in the genus *Microtus* (small-eared "meadow voles") of the family Cricetidae. Voles are small mammals, and this species...
Gray's beaked whale
Gray's beaked whale
Mesoplodon grayi

Gray's beaked whale (*Mesoplodon grayi*), sometimes known as Haast's beaked whale, the scamperdown whale, or the southern beaked whale, is one of the better-known members of the genus *Mesoplodon*. This species is fairly gregarious and strands...
Grayish mouse opossum
Grayish mouse opossum
Tlacuatzin canescens

The grayish mouse opossum (*Tlacuatzin canescens*) is a species of opossum endemic to Mexico. It is the sole species in the genus *Tlacuatzin*.
Great Balkhan mouse-like hamster
Great Balkhan mouse-like hamster
Calomyscus mystax

The Great Balkhan mouse-like hamster (*Calomyscus mystax*) is a species of rodent in the family Calomyscidae. It is found in southwestern Turkmenistan, northern Iran and southern Azerbaijan. It was first discovered by the famed Soviet Zoologist and...
Great Basin pocket mouse
Great Basin pocket mouse
Perognathus parvus

The Great Basin pocket mouse (*Perognathus parvus*) is a species of rodent in the family Heteromyidae. It is found in British Columbia in Canada and the western United States.

Great bent-winged bat
Great bent-winged bat
Miniopterus tristis

The great bent-winged bat (*Miniopterus tristis*) is a species of vesper bat in the family *Miniopteridae*. It can be found in the following countries: Indonesia, Papua New Guinea, Philippines, Solomon Islands, and Vanuatu.

Great evening bat
Great evening bat
Ia io

The great evening bat (*Ia io*) is the largest bat in the vesper bat family (*Vespertilionidae*) and the only living species in the genus *Ia*. It is common to Eastern and Southeastern Asia (China, India, Laos, Nepal, Thailand and Vietnam), mainly...

Great Flying Fox
Great Flying Fox
Pteropus neohibernicus

The great flying fox (*Pteropus neohibernicus*), also known as the greater flying fox or Bismarck flying fox, is a species of megabat in the genus *Pteropus*, found throughout lowland areas of New Guinea and in the Bismarck Archipelago.

Conflicting...
Great fruit-eating bat
Great fruit-eating bat
Artibeus lituratus

The great fruit-eating bat (*Artibeus lituratus*) is a bat species in the family *Phyllostomidae* from South and Central America. It is found from Mexico to Brazil and Argentina, as well as in Antigua and Barbuda, Barbados, Grenada, Martinique,...

Great Gebril
Great Gebril
Rhombomys opimus

The great gerbil (*Rhombomys opimus*) is a large rodent found throughout much of Central Asia.

Great hopping mouse
Great hopping mouse
Notomys robustus

The great hopping mouse (*Notomys robustus*) is extinct. It is known only from skulls found in owl pellets in the Flinders Ranges. Some pellets also include bones of the introduced house mouse—indicating that it survived into historic times,...

Great jerboa
Great jerboa
Allactaga major

The great jerboa (*Allactaga major*) is a species of rodent in the family *Dipodidae*. It is found in Kazakhstan, Russia, Turkmenistan, Ukraine and Uzbekistan. It mainly lives in deserts.

Great Key Island giant rat
Great Key Island giant rat
Uromys siebersi

The Great Key Island giant rat, or Great Key Island uromys (*Uromys siebersi*) is a species of rodent in the family *Muridae*. It is known only from Great Key Island, Indonesia. The species name is after the ornithologist H. C. Siebers.

Great Lakes wolf
Great Lakes wolf

Canis lupus lycaon

The eastern wolf (*Canis lupus lycaon* or *Canis lycaon* or *Canis rufus lycaon*) also known as the timber wolf, Algonquin wolf or eastern timber wolf, is a canine of debated taxonomy native to the Great Lakes region and southeastern Canada. It is...
Great Plains wolf
Great Plains wolf
Canis lupus nubilus

The Great Plains wolf (*Canis lupus nubilus*), also known as the buffalo wolf or loafer, is an extinct subspecies of gray wolf that once extended throughout the Great Plains, from southern Manitoba and Saskatchewan in Canada southward to northern...

Great roundleaf bat
Great roundleaf bat
Hipposideros armiger

The great roundleaf bat, also known as the great Himalayan leaf-nosed bat (*Hipposideros armiger*) is a species of bat in the family Hipposideridae found in South Asia, Southeast Asia, and China. It gives birth to two young a year. It has been...

Great stripe-faced bat
Great stripe-faced bat
Vampyroides caraccioli

The great stripe-faced bat or stripe-faced vampire bat (*Vampyroides caraccioli*) is a bat species from South and Central America, where it is found from southern Mexico to Bolivia and northwestern Brazil, as well as on Trinidad. The great...

Great-tailed triok
Great-tailed triok
Dactylopsila megalura

The great-tailed triok (*Dactylopsila megalura*) is a species of marsupial in the family Petauridae. It is found in West Papua, Indonesia and Papua New Guinea. Its natural habitat is subtropical or tropical dry forests.

Greater Asiatic yellow bat
Greater Asiatic yellow bat
Scotophilus heathii

The greater Asiatic yellow bat (*Scotophilus heathii*) is a species of vesper bat. It is found in Afghanistan, Bangladesh, Cambodia, China, India, Indonesia, Laos, Myanmar, Nepal, Pakistan, Philippines, Sri Lanka, Thailand and Vietnam. Like many...

Greater bamboo bat
Greater bamboo bat
Tylonycteris robustula

The greater bamboo bat (*Tylonycteris robustula*) is a species of vesper bat in the family Vespertilionidae. It is found in Cambodia, China, India, Indonesia, Laos, Malaysia, Myanmar, the Philippines, Singapore and Thailand. It has two...

Greater Bamboo Lemur
Greater Bamboo Lemur
Prolemur simus

The greater bamboo lemur (*Prolemur simus*), also known as the broad-nosed bamboo lemur and the broad-nosed gentle lemur, is the largest bamboo lemur, at over five pounds or nearly 2.5 kilograms. It has greyish brown fur and white ear tufts, and...

Greater Bandicoot Rat
Greater Bandicoot Rat

Bandicota indica

The Greater bandicoot rat (*Bandicota indica*) is a large species of rodent from Asia. In Sinhala, the bandicoot rat is known as maha uru-meeya, and in Malayalam, it is known as panni-eli, thorappan. Both names directly translate to "pig-rat"....
Greater big-footed mouse
Greater big-footed mouse
Macrotarsomys ingens

The greater big-footed mouse or long-tailed big-footed mouse (*Macrotarsomys ingens*) is a nocturnal rodent found only in north west Madagascar. It was first described by F. Petter in 1959. It is listed as an endangered species as a result of...

Greater Bilby
Greater Bilby
Macrotis lagotis

The greater bilby (*Macrotis lagotis*), often referred to simply as the bilby since the lesser bilby (*Macrotis leucura*) became extinct in the 1950s, is an Australian species of nocturnal omnivorous animal in the order Peramelemorphia. Other...

Greater broad-nosed bat
Greater broad-nosed bat
Platyrrhinus vittatus

The greater broad-nosed bat (*Platyrrhinus vittatus*) is a species of bat in the family Phyllostomidae. It is found in Bolivia, Colombia, Costa Rica, Ecuador, Panama, Peru, and Venezuela.

Greater Bulldog Bat
Greater Bulldog Bat
Noctilio leporinus

The Greater bulldog bat (*Noctilio leporinus*) is a species of fishing bat native to Latin America. It uses echolocation to detect water ripples made by the fish upon which it preys, then uses the pouch between its legs to scoop the fish up and its...

Greater Cane Rat
Greater Cane Rat
Thryonomys swinderianus

The greater cane rat (*Thryonomys swinderianus*), also known as the grasscutter (in Ghana, Nigeria and other regions of West Africa), is one of two species of cane rats, a small family of African hystricognath rodents. It lives by reed-beds and...

Greater Chinese mole
Greater Chinese mole
Euroscaptor grandis

The greater Chinese mole (*Euroscaptor grandis*) is a species of mammal in the family Talpidae. It is found in China and potentially Myanmar.

Greater Congo shrew
Greater Congo shrew
Congosorex polli

The greater Congo shrew (*Congosorex polli*) is a species of mammal in the family Soricidae. It is endemic to the Democratic Republic of the Congo. Its natural habitat is subtropical or tropical moist lowland forests of Congo. It is currently...

Greater dog-like bat
Greater dog-like bat
Peropteryx kappleri

The greater dog-like bat (*Peropteryx kappleri*) is a bat species from Central America and South America. It is found from southern Mexico through Brazil and Peru.

Greater Dwarf Lemur
Greater Dwarf Lemur
Cheirogaleus major

The greater dwarf lemur (*Cheirogaleus major*), or the Geoffroy's dwarf lemur, is a lemur that is widely distributed over the primary and secondary forests near the eastern coast of Madagascar. They are also found in northern parts of Madagascar....

Greater dwarf shrew
Greater dwarf shrew
Suncus lixus

The greater dwarf shrew (*Suncus lixa*) is a species of mammal in the family Soricidae. It is found in Angola, Botswana, Democratic Republic of the Congo, Kenya, Malawi, Namibia, Tanzania, Zambia, and Zimbabwe. Its natural habitats are subtropical...

Greater Egyptian gerbil
Greater Egyptian gerbil
Gerbillus pyramidum

The greater Egyptian gerbil (*Gerbillus pyramidum*) is a small rodent in the family Muridae. It is native to northern Africa where it inhabits sandy deserts, semi-arid areas and oases. It is a common species, and the International Union for...

Greater Egyptian jerboa
Greater Egyptian jerboa
Jaculus orientalis

The greater Egyptian jerboa (*Jaculus orientalis*) is a species of rodent in the family Dipodidae. It is found in Algeria, Egypt, Libya, Morocco, Saudi Arabia, Tunisia, and is possibly extinct in the Negev Desert of Israel. Its natural habitats...

Greater fairy armadillo
Greater fairy armadillo
Calyptophractus retusus

The greater fairy armadillo (*Calyptophractus retusus*), also known as Burmeister's armadillo or the Chacoan fairy armadillo, is a species of armadillo in the family Chlamyphoridae. It is found in Argentina, Bolivia, and Paraguay. Its natural...

Greater False Vampire Bat
Greater False Vampire Bat
Lyroderma lyra

The Greater false vampire bat (*Lyroderma lyra*) is a species of bat native to Asia. It is also known as the Indian false vampire bat or Greater false-vampire.

Greater fat-tailed jerboa
Greater fat-tailed jerboa
Pygeretmus shitkovi

The greater fat-tailed jerboa (*Pygeretmus shitkovi*) is a species of rodent in the family Dipodidae. It is endemic to Kazakhstan. Its natural habitat is temperate desert.

Greater forest shrew
Greater forest shrew
Sylvisorex ollula

The greater forest shrew (*Sylvisorex ollula*) is a species of mammal in the family Soricidae found in Cameroon, the Central African Republic, the Republic of the Congo, Equatorial Guinea, Gabon, and Nigeria. Its natural habitat is subtropical or...

Greater ghost bat
Greater ghost bat
Diclidurus ingens

The greater ghost bat (*Diclidurus ingens*) is a bat species found in northwestern Brazil, southeastern Colombia, Guyana, Peru, and Venezuela.

Greater Glider
Greater Glider
Petauroides volans

The greater gliders are three species of large gliding marsupials in the genus *Petauroides*, all of which are found in eastern Australia. Until 2020 they were considered to be one species, *Petauroides volans*. In 2020 morphological and genetic...

Greater Grison
Greater Grison
Galictis vittata

The greater grison (*Galictis vittata*) is a species of mustelid native to Southern Mexico, Central America, and South America.

Greater guinea pig
Greater guinea pig
Cavia magna

The greater guinea pig (*Cavia magna*) is a species of rodent found in the coastal strip of Brazil and Uruguay, where it lives in moist grassland and marshes.

Greater hamster-rat
Greater hamster-rat
Beamys major

The greater hamster-rat, greater long-tailed pouched rat, or long-tailed pouched rat (*Beamys major*) is a species of rodent in the family Nesomyidae. It is found in Malawi, Tanzania, and Zambia. Its natural habitats are subtropical or tropical...

Greater hedgehog tenrec
Greater hedgehog tenrec
Setifer setosus

The greater hedgehog tenrec (*Setifer setosus*), also known as the large Madagascar hedgehog or sokina, is a species of mammal in the family Tenrecidae. It is endemic to Madagascar. Its natural habitats are subtropical or tropical dry forest,...

Greater Horseshoe Bat
Greater Horseshoe Bat
Rhinolophus ferrumequinum

The greater horseshoe bat (*Rhinolophus ferrumequinum*) is an insectivorous bat of the genus *Rhinolophus*. Its distribution covers Europe, Northern Africa, Central Asia and Eastern Asia. It is the largest of the horseshoe bats in Europe and is thus...

Greater Kudu
Greater Kudu
Tragelaphus strepsiceros

The Greater kudu (*Tragelaphus strepsiceros*) is a woodland antelope found in Africa. Despite occupying such widespread territory, Greater kudus are sparsely populated in most areas due to declining habitat, deforestation, and poaching.

Greater large-headed shrew
Greater large-headed shrew
Paracrocidura maxima

The greater large-headed shrew (*Paracrocidura maxima*) is a species of mammal in the family Soricidae. It is found in Burundi, Democratic Republic of the Congo, Rwanda, and Uganda. Its natural habitats are subtropical or tropical moist lowland...

Greater long-fingered bat
Greater long-fingered bat
Miniopterus inflatus

The greater long-fingered bat (*Miniopterus inflatus*) is a species in the family Miniopteridae. It is found in Cameroon, Central African Republic, Democratic Republic of the Congo, Equatorial Guinea, Ethiopia, Gabon, Guinea, Kenya, Liberia,...

Greater long-nosed armadillo
Greater long-nosed armadillo
Dasypus kappleri

The greater long-nosed armadillo (*Dasypus kappleri*) is a South American species of armadillo found in Colombia, Venezuela, Ecuador, Guyana, Suriname, French Guiana, Peru, Bolivia and Brazil. It is a solitary, nocturnal, terrestrial animal that...

Greater long-nosed bat
Greater long-nosed bat
Leptonycteris nivalis

The greater long-nosed bat or Mexican long-nosed bat (*Leptonycteris nivalis*) is a species of bat in the family Phyllostomidae. It is found in Mexico and the United States. It chiefly consumes pollen and nectar, particularly from agave plants and...

Greater long-tailed bat
Greater long-tailed bat
Choeroniscus periosus

The greater long-tailed bat (*Choeroniscus periosus*) is a species of bat in the family Phyllostomidae. It is found in Colombia and Ecuador. It is threatened by habitat loss.

Greater long-tailed hamster
Greater long-tailed hamster
Tscherskia triton

The greater long-tailed hamster (*Tscherskia triton*) is a rodent native to Siberia, the Korean Peninsula, and China. It is the only member of the genus *Tscherskia*.

Greater long-tailed shrew tenrec
Greater long-tailed shrew tenrec
Microgale principula

The greater long-tailed shrew tenrec (*Microgale principula*) is an afrotherian species of mammal in the family Tenrecidae. It is endemic to Madagascar, where its natural habitats are subtropical or tropical moist lowland forests or moist montane...

Greater Mindanao shrew
Greater Mindanao shrew
Crocidura grandis

The Greater Mindanao shrew (*Crocidura grandis*) is a species of mammal in the family Soricidae. It is endemic to the Philippines. Its natural habitat is subtropical or tropical dry forests. It is threatened by habitat loss.

Greater mole-rat
Greater mole-rat
Spalax microphthalmus

The greater blind mole-rat (*Spalax microphthalmus*) is a species of rodent in the family Spalacidae. It is found in Russia and Ukraine.

Greater monkey-faced bat
Greater monkey-faced bat
Pteralopex flanneryi

The greater monkey-faced bat or greater flying monkey (*Pteralopex flanneryi*) is a megabat endemic to Solomon Islands, Bougainville, in Papua New Guinea, and nearby small islands. It is listed as a critically endangered species and the population...

Greater Mouse-Deer
Greater Mouse-Deer
Tragulus napu

The greater mouse-deer, greater Malay chevrotain, or napu (*Tragulus napu*) is a species of even-toed ungulate in the family Tragulidae found in Sumatra, Borneo, and smaller Malaysian and Indonesian islands, and in southern Myanmar, southern...

Greater Mouse-Eared Bat
Greater Mouse-Eared Bat
Myotis myotis

The greater mouse-eared bat (*Myotis myotis*) is a European species of bat in the family Vespertilionidae.

Greater mouse-tailed bat
Greater mouse-tailed bat
Rhinopoma microphyllum

The greater mouse-tailed bat (*Rhinopoma microphyllum*) is a species of bat in the Rhinopomatidae family.

Greater musky fruit bat
Greater musky fruit bat
Ptenochirus jagori

The greater musky fruit bat (*Ptenochirus jagori*) is a species of megabat in the family Pteropodidae. It is endemic to the Philippines. It was named by Peters for Fedor Jagor.

Greater naked-tailed armadillo
Greater naked-tailed armadillo
Cabassous tatouay

The greater naked-tailed armadillo (*Cabassous tatouay*) is an armadillo species from South America.

Greater nectar bat
Greater nectar bat
Eonycteris major

The greater nectar bat or greater dawn bat (*Eonycteris major*) is a species of megabat within the genus *Eonycteris*. It is found in Brunei, Indonesia, Malaysia, and the Philippines. Its range is limited and includes Luzon to Maripipi in the...

Greater Noctule Bat
Greater Noctule Bat
Nyctalus lasiopterus

The Greater noctule bat (*Nyctalus lasiopterus*) is a rare carnivorous bat that is found only in Europe, West Asia, and North Africa. It is the largest and least

studied bat in Europe and is one of the few bat species to feed on passerine birds....

Greater Papuan pipistrelle
Greater Papuan pipistrelle
Pipistrellus collinus

The greater Papuan pipistrelle (*Pipistrellus collinus*) is a species of vesper bat found in Papua New Guinea and Irian Jaya.

Greater red musk shrew
Greater red musk shrew
Crocidura flavescens

The greater red musk shrew (*Crocidura flavescens*) is a species of mammal in the family Soricidae. It is found in Lesotho, Mozambique, South Africa, and Eswatini. Its natural habitats are moist savanna, temperate grassland, and rural gardens....

Greater round-eared bat
Greater round-eared bat
Tonatia bidens

The greater round-eared bat (*Tonatia bidens*) is a bat species from South America. It is found in northeastern and southern Brazil, northern Argentina, Paraguay and Bolivia. The species feeds on fruit as well as hunting small birds. Once caught,...

Greater roundleaf bat
Greater roundleaf bat
Hipposideros camerunensis

The greater roundleaf bat (*Hipposideros camerunensis*) is a species of bat in the family Hipposideridae found in Cameroon, the Democratic Republic of the Congo, and Kenya.

Greater sac-winged bat
Greater sac-winged bat
Saccopteryx bilineata

The greater sac-winged bat (*Saccopteryx bilineata*) is a bat of the family Emballonuridae native to Central and South America. They are the most common bats seen in the rainforest, as they often roost on the outside of large trees. They are...

Greater sheath-tailed bat
Greater sheath-tailed bat
Emballonura furax

The greater sheath-tailed bat or New Guinea sheath-tailed bat (*Emballonura furax*) is a species of sac-winged bat in the family Emballonuridae. It is endemic to New Guinea and some nearby islands. It is threatened by habitat loss.

Greater Short-Nosed Fruit Bat
Greater Short-Nosed Fruit Bat
Cynopterus sphinx

The greater short-nosed fruit bat (*Cynopterus sphinx*), or short-nosed Indian fruit bat, is a species of megabat in the family Pteropodidae found in South and Southeast Asia.

Greater short-tailed gerbil
Greater short-tailed gerbil
Dipodillus maghrebi

The greater short-tailed gerbil (*Dipodillus maghrebi*) is a rodent found mainly in Morocco.

Greater spear-nosed bat
Greater spear-nosed bat

Phyllostomus hastatus

The greater spear-nosed bat (*Phyllostomus hastatus*) is a bat species of the family Phyllostomidae from South and Central America. It is one of the larger bats of this region and is omnivorous.

Greater Spot-Nosed Monkey
Greater Spot-Nosed Monkey
Cercopithecus nictitans

The Greater spot-nosed monkey (*Cercopithecus nictitans*) is one of the smallest Old World monkeys. It is primarily arboreal and often associates with monkeys of other species.

Greater stick-nest rat
Greater stick-nest rat
Leporillus conditor

The greater stick-nest rat, also known as the house-building rat and wopilkara (*Leporillus conditor*) is a species of rodent in the family Muridae. They are about the size of a small rabbit and construct large nests of interwoven sticks. Once...

Greater stripe-backed shrew
Greater stripe-backed shrew
Sorex cylindricauda

The greater stripe-backed shrew or stripe-backed shrew (*Sorex cylindricauda*) is a species of mammal in the family Soricidae. It is endemic to China. It is threatened by habitat loss. The Greater stripe-backed shrew also shares 97% of its DNA...

Greater tree mouse
Greater tree mouse
Chiruromys forbesi

The greater tree mouse (*Chiruromys forbesi*) is a species of rodent in the family Muridae. It is found only in Papua New Guinea.

Greater tube-nosed bat
Greater tube-nosed bat
Murina leucogaster

The greater tube-nosed bat (*Murina leucogaster*) is a species of bat. An adult greater tube-nosed bat has a body length of 4.2-5.7 cm, a tail length of 3.6-4.1 cm, and a wing length of 3.7-4.4 cm. The species is found in India, Mongolia, China,...

Greater white-toothed shrew
Greater white-toothed shrew
Crocidura russula

The greater white-toothed shrew (*Crocidura russula*) is a small insectivorous mammal found in Europe and North Africa. It is the most common of the white-toothed shrews. This species is found along the Mediterranean, Netherlands, Belgium,...

Greater Wilfred's mouse
Greater Wilfred's mouse
Wilfredomys oenax

The greater Wilfred's mouse, *Wilfredomys oenax*, is a rodent species from South America. It is found in southern Brazil and Uruguay in subtropical lowland forest. It is arboreal to some degree. It is the only species in the genus *Wilfredomys*.

Greater yellow-shouldered bat
Greater yellow-shouldered bat
Sturnira magna

The greater yellow-shouldered bat (*Sturnira magna*) is a species of bat in the

family Phyllostomidae. It is found in Bolivia, Colombia, Ecuador, and Peru.
Greedy olalla rat
Greedy olalla rat
Olallamys edax

The greedy olalla rat (*Olallamys edax*) is a species of rodent in the family Echimyidae. It is found in Colombia and Venezuela. Its natural habitat is subtropical or tropical moist lowland forests. It is threatened by habitat loss. The species name...
Green acouchi
Green acouchi
Myoprocta pratti

The green acouchi (*Myoprocta pratti*) is a species of rodent in the acouchi genus, part of the family Dasyproctidae, from South America. Its distribution is in western Amazonia, west of the Rios Negro and Madeira, in northwestern Brazil, ...
Green bush squirrel
Green bush squirrel
Paraxerus poensis

The green bush squirrel (*Paraxerus poensis*) is a species of rodent in the family Sciuridae.
Green Monkey
Green Monkey
Chlorocebus sabaeus

The Green monkey (*Chlorocebus sabaeus*) is an Old World monkey which gets its name due to a green tinge in its golden fur. It lives in a wide range of wooded habitats and takes a wide variety of foods. These monkeys live in groups and have...
Green ringtail possum
Green ringtail possum
Pseudochirops archeri

The green ringtail possum (*Pseudochirops archeri*) is a species of ringtail possum found only in northern Australia. This makes it unique in its genus, all other members of which are found in New Guinea or nearby islands. The green ringtail...
Greenhall's dog-faced bat
Greenhall's dog-faced bat
Cynomops greenhalli

Greenhall's dog-faced bat (*Cynomops greenhalli*) is a South American bat species of the family Molossidae. It is found in Colombia, Peru, Ecuador, Venezuela, the Guianas, northeastern Brazil and Trinidad. It is an insect-eating bat, 40–97 mm in...
Greenish naked-backed fruit bat
Greenish naked-backed fruit bat
Dobsonia viridis

The greenish naked-backed fruit bat (*Dobsonia viridis*) is a species of megabat in the family Pteropodidae.
Greenish yellow bat
Greenish yellow bat
Scotophilus viridis

The eastern greenish yellow bat (*Scotophilus viridis*) is a species of vesper bat. It is found in Botswana, Central African Republic, Eswatini, Ethiopia, Kenya, Malawi, Mozambique, South Africa, Tanzania, Zambia, and Zimbabwe. Its natural...
Greenland Wolf

Greenland Wolf
Canis lupus orion

The Greenland wolf (*Canis lupus orion*) is a subspecies of Grey wolf that is native to Greenland. Historically, it was heavily persecuted, but today it is fully protected. The entire population of the Greenland wolf is very small, probably about...

Greenwood's shrew
Greenwood's shrew
Crocidura greenwoodi

The Greenwood's shrew (*Crocidura greenwoodi*) is a species of mammal in the family Soricidae. It is endemic to Somalia. Its natural habitats are subtropical or tropical moist lowland forest, dry savanna, subtropical or tropical dry shrubland, and...

Gregarious short-tailed rat
Gregarious short-tailed rat
Brachyuromys ramirohitra

The gregarious short-tailed rat (*Brachyuromys ramirohitra*) is a species of rodent in the family Nesomyidae. It is found only in Madagascar. Its natural habitats are subtropical or tropical dry forests and subtropical or tropical dry shrubland.

Gregory's wolf
Gregory's wolf
Canis lupus gregoryi

Gregory's wolf (*Canis rufus gregoryi*), also known as the Mississippi Valley wolf, was a subspecies of the red wolf. It was declared extinct in 1980. It once roamed the regions in and around the lower Mississippi River basin.

Gressitt's mosaic-tailed rat
Gressitt's mosaic-tailed rat
Paramelomys gressitti

Gressitt's mosaic-tailed rat (*Paramelomys gressitti*) is a species of rodent native to Papua New Guinea. Ongoing habitat degradation is a major threat. It is named after the collector, Judson Linsley Gressitt.

Grévy's Zebra
Grévy's Zebra
Equus grevyi

Grévy's zebras (*Equus grevyi*) are magnificent and elegant creatures. Although they belong to the horse family, these animals are actually more closely related to the African wild asses. Due to excessive and continuous hunting, they were once...

Grewcock's sportive lemur
Grewcock's sportive lemur
Lepilemur grewcockorum

Grewcock's sportive lemur (*Lepilemur grewcockorum*), or the Anjamangirana sportive lemur, is a sportive lemur endemic to Madagascar. It is a medium-sized sportive lemur with a total length of about 55 to 63 cm (22 to 25 in), of which 26–30 cm...

Grey dwarf hamster
Grey dwarf hamster
Cricetulus migratorius

The grey dwarf hamster, grey hamster or migratory hamster (*Cricetulus migratorius*) is a species of rodent in the family Cricetidae. Its range extends from Eastern Europe through the Middle East, Russia and Central Asia to Mongolia and western...

Grey long-eared bat
Grey long-eared bat

Plecotus austriacus

The grey long-eared bat (*Plecotus austriacus*) is a fairly large European bat. It has distinctive ears, long and with a distinctive fold. It hunts above woodland, often by day, and mostly for moths. In captivity, it has also been recorded to eat...

Grey red-backed vole

Grey red-backed vole

Myodes rufocanus

The grey red-backed vole or the grey-sided vole (*Myodes rufocanus*) is a species of vole. An adult grey red-backed vole weighs 20-50 grams. This species ranges across northern Eurasia, including northern China, the northern Korean Peninsula, and...

Grey Rhebok

Grey Rhebok

Pelea capreolus

The grey rhebok or grey rhebuck (*Pelea capreolus*), locally known as the reebok in Afrikaans, is a species of antelope native to South Africa, Lesotho, and Eswatini (Swaziland). The specific name *capreolus* is Latin for 'little goat', it is the...

Grey Wolf

Grey Wolf

Canis lupus

The wolf (*Canis lupus*), also known as the Gray wolf or Grey wolf, is a large canine native to Eurasia and North America. More than thirty subspecies of *Canis lupus* have been recognized, and gray wolves, as popularly understood, comprise wild...

Grey-bellied dunnart

Grey-bellied dunnart

Sminthopsis griseiventor

The grey-bellied dunnart (*Sminthopsis griseoventer*), alternately spelled gray-bellied dunnart, was described by the same people Kitchener, Stoddart and Henry along with the Kangaroo Island dunnart, Gilbert's dunnart and little long-tailed...

Grey-bellied squirrel

Grey-bellied squirrel

Callosciurus caniceps

The grey-bellied squirrel (*Callosciurus caniceps*) is a species of rodent in the family Sciuridae. It is found in forests, plantations and gardens in Peninsular Malaysia, Thailand, southern Myanmar, southern China (Yunnan) and possibly western...

Grey-Cheeked Mangabey

Grey-Cheeked Mangabey

Lophocebus albigena

The Grey-cheeked mangabey (*Lophocebus albigena*) is an Old World monkey from Central Africa. They have dark faces and look in shape overall like small, hairy baboons. Mangabeys live in groups and feed primarily on fruit.

Grey-crowned Central American squirrel monkey

Grey-crowned Central American squirrel monkey

Saimiri oerstedii citrinellus

The grey-crowned Central American squirrel monkey (*Saimiri oerstedii citrinellus*) is a subspecies of the Central American squirrel monkey. Its range is restricted to the Pacific coast of central Costa Rica. The northern end of its range is the...

Grey-faced sengi

Grey-faced sengi

Rhynchocyon udzungwensis

The grey-faced sengi (*Rhynchocyon udzungwensis*) is a species of elephant shrew that is endemic to the Udzungwa Mountains of south-central Tanzania. The discovery of the species was announced in January 2008; only 15 species of elephant shrew...
Grey-Headed Flying Fox
Grey-Headed Flying Fox
Pteropus poliocephalus

Grey-headed flying foxes (*Pteropus poliocephalus*) are large bats with dark brown eyes, black wings, dark grey head and body as well as a broad, reddish-brown collar. One of the most conspicuous features of this species is their fur, covering all...
Griffin's leaf-nosed bat
Griffin's leaf-nosed bat
Hipposideros griffini

Griffin's leaf-nosed bat (*Hipposideros griffini*) is a species of roundleaf bats. First seen in 2008, it was formally described in 2012. It is found only in Vietnam, and only at two places, Cát Bà Island in northern Vietnam, and Chư Mom Ray...
Griffith's long-fingered bat
Griffith's long-fingered bat
Miniopterus griffithsi

Griffith's long-fingered bat (*Miniopterus griffithsi*) is a bat in the genus *Miniopterus* which occurs in southern Madagascar. *M. griffithsi* was previously a part of the largest family of bats, the *Vespertilionidae*, which consist of 5 subfamilies....
Griselda's striped grass mouse
Griselda's striped grass mouse
Lemniscomys griselda

Griselda's lemniscomys or Griselda's striped grass mouse (*Lemniscomys griselda*) is a species of rodent in the family *Muridae*. It is found in Angola, Democratic Republic of the Congo, and Zambia. Its natural habitat is dry savanna.
Grivet
Grivet
Chlorocebus aethiops

The grivet (*Chlorocebus aethiops*) is an Old World monkey with long white tufts of hair along the sides of its face. Some authorities consider this and all of the members of the genus *Chlorocebus* to be a single species, *Cercopithecus aethiops*.
As...
Grizzled Giant Squirrel
Grizzled Giant Squirrel
Ratufa macroura

The grizzled giant squirrel (*Ratufa macroura*) is a large tree squirrel in the genus *Ratufa* found in the highlands of the Central and Uva provinces of Sri Lanka, and in patches of riparian forest along the Kaveri River and in the hill forests of...
Grizzled Mexican small-eared shrew
Grizzled Mexican small-eared shrew
Cryptotis obscura

The grizzled Mexican small-eared shrew (*Cryptotis obscura*) is a small mammal in the order *Eulipotyphla*. It is native in Mexican highlands. It can be found in dense, wet cloud forest, where it is found in the thick herbaceous undergrowth and leaf...
Grizzled Tree-Kangaroo

Grizzled Tree-Kangaroo
Dendrolagus inustus

The Grizzled tree-kangaroo (*Dendrolagus inustus*) is a very rare marsupial that lives in foothill forests of northern and western New Guinea and is indigenous to some of the offshore islands. It is a member of the macropod family and carries its...

Grizzly Bear
Grizzly Bear
Ursus arctos horribilis

The Grizzly bear (*Ursus arctos horribilis*) is a population or subspecies of the Brown bear that is found in North America. Native American tribes living among Brown bears often view them with a mixture of awe and fear. North American brown bears...

Grobben's gerbil
Grobben's gerbil
Gerbillus grobbeni

Grobben's gerbil (*Gerbillus grobbeni*) is a species of rodent, distributed mainly in Libya; Cyrenaica, Derna. Less than 250 individuals of this species are thought to persist in the wild. It is named after Austrian biologist Karl Grobбен.

Groove-toothed bat
Groove-toothed bat
Phoniscus atrox

The groove-toothed bat (*Phoniscus atrox*) is a species of bat in the family Vespertilionidae, the vesper bats. It is native to Indonesia, Malaysia, and Thailand. It is an uncommon species that depends on forests for survival, and it is threatened...

Groove-toothed flying squirrel
Groove-toothed flying squirrel
Aeretes melanopterus

The groove-toothed flying squirrel or North Chinese flying squirrel (*Aeretes melanopterus*) is a species of rodent in the family Sciuridae. It is monotypic within the genus *Aeretes*. It is endemic to China, and occurs in Sichuan, Gansu, Hebei, and...

Ground cuscus
Ground cuscus
Phalanger gymnotis

The ground cuscus (*Phalanger gymnotis*) is a marsupial from the order Diprotodontia and belongs within the family Phalangeridae, a diverse family consisting of the other cuscus species and the brushtail possums (*Trichosurus* spp.) and the...

Ground Pangolin
Ground Pangolin
Smutsia temminckii

The Ground pangolin (*Smutsia temminckii*) is an unusual animal that belongs to a group of pangolins which are among the most critically endangered and illegally trafficked animals in the world. They are armor-plated, distinguished from other...

Groundhog
Groundhog
Marmota monax

The groundhog (*Marmota monax*) is a lowland rodent of North America. It was first scientifically described by Carl Linnaeus in 1758. These adorable animals play an important role maintaining healthy soil in woodland and plain areas. They are...

Groves' dwarf lemur
Groves' dwarf lemur
Cheirogaleus grovesi

Groves' dwarf lemur (*Cheirogaleus grovesi*) is a species of dwarf lemur known from Ranomafana and Andringitra national parks and surrounding areas in southeastern Madagascar. The identification was based on comparison of mitochondrial and nuclear...

Guadalcanal monkey-faced bat
Guadalcanal monkey-faced bat
Pteralopex atrata

The Guadalcanal monkey-faced bat or Guadalcanal flying monkey (*Pteralopex atrata*) is a megabat endemic to Solomon Islands. It is listed as an endangered species. In 2013, Bat Conservation International listed this species as one of the 35...

Guadalcanal rat
Guadalcanal rat
Uromys porculus

The Guadalcanal rat (*Uromys porculus*) is a species of rodent in the family Muridae. It was found only in the Solomon Islands. Since this species was collected between 1886 and 1888 it has not been seen again. It is possibly extinct.

Guadalupe Fur Seal
Guadalupe Fur Seal
Arctocephalus townsendi

Guadalupe fur seals (*Arctocephalus townsendi*) are one of eight members of the fur seal genus *Arctocephalus*. They are the northernmost member of this genus. Sealers reduced the population of these fur seals to just a few dozen by the late 19th...

Guadeloupe big brown bat
Guadeloupe big brown bat
Eptesicus guadeloupensis

The Guadeloupe big brown bat (*Eptesicus guadeloupensis*) is a species of vesper bat. It is found only on the island of Guadeloupe. It is one of the 11 species of bat found on Guadeloupe, and one of the 3 that are endemic.

Guadeloupe big-eyed bat
Guadeloupe big-eyed bat
Chiroderma improvisum

The Guadeloupe big-eyed bat (*Chiroderma improvisum*) is a species of bat in the family Phyllostomidae. It is found in Guadeloupe and Montserrat. It is threatened by habitat loss mostly because of Hurricane Hugo, which destroyed 90% of the...

Guadeloupe Raccoon
Guadeloupe Raccoon
Procyon lotor minor

The Guadeloupe raccoon (*Procyon lotor minor*) is a common raccoon subspecies.

Guahiba gracile opossum
Guahiba gracile opossum
Cryptonanus guahybae

The Guahiba gracile opossum (*Cryptonanus guahybae*) is a species of opossum in the family Didelphidae. It is endemic to southern Brazil, where it is known only from three islands, Guahiba, São Lourenço, and Taquara, in the state of Rio Grande do...

Guaira spiny rat
Guaira spiny rat
Proechimys guairae

The Guaira spiny-rat (*Proechimys guairae*) is a species of rodent in the family Echimyidae. It is endemic to Venezuela. It is commonly referred to as 'casiragua' to avoid confusion with true rats (Muroidea)

Guajira mouse opossum

Guajira mouse opossum

Marmosa xerophila

The Guajira mouse opossum (*Marmosa xerophila*) is a species of opossum in the family Didelphidae. It is found in Colombia and Venezuela.

Guam flying fox

Guam flying fox

Pteropus tokudae

The Guam flying fox (*Pteropus tokudae*), also known as the little Marianas fruit bat, was a tiny megabat from Guam in the Marianas Islands in Micronesia that was confirmed extinct due to hunting or habitat changes. It was first recorded in 1931...

Guanaco

Guanaco

Lama guanicoe

The guanaco (*Lama guanicoe*) is a camelid native to South America. Described as an elegant animal, with a long, slender neck and fine legs by Charles Darwin, the guanaco is the biggest wild camelid family member in South America and is thought to...

Guatemalan Black Howler

Guatemalan Black Howler

Alouatta pigra

The Yucatán black howler, or Guatemalan black howler, (*Alouatta pigra*) is a species of howler monkey, a type of New World monkey, from Central America. It is found in Belize, Guatemala and Mexico, in and near the Yucatán Peninsula. It lives in...

Guatemalan broad-clawed shrew

Guatemalan broad-clawed shrew

Cryptotis griseoventris

The Guatemalan broad-clawed shrew (*Cryptotis griseoventris*) is a species of small-eared shrew in the family Soricidae. It is known from Guatemala and the Mexican state of Chiapas, where it has been found in montane forests of oak, pine and fir,...

Guatemalan deer mouse

Guatemalan deer mouse

Peromyscus guatemalensis

The Guatemalan deer mouse (*Peromyscus guatemalensis*) is a species of rodent in the family Cricetidae. It is found in Guatemala and Mexico.

Guatemalan myotis

Guatemalan myotis

Myotis cobanensis

The Guatemalan myotis (*Myotis cobanensis*) is a species of vesper bat. It is found only in Guatemala.

Guatemalan vole

Guatemalan vole

Microtus guatemalensis

The Guatemalan vole (*Microtus guatemalensis*) is a species of rodent in the family Cricetidae. It is found in Guatemala and Chiapas, Mexico, in montane pine-oak forest

and meadow at elevations between 2600 and 3100 m above sea level. It is...

Guayaquil squirrel
Guayaquil squirrel
Sciurus stramineus

The Guayaquil squirrel (*Sciurus stramineus*) is a tree squirrel endemic to Ecuador and Peru. It is a robust squirrel with a head-and-body length of 18 to 32 cm (7 to 13 in) and a similar length tail. The colour is variable; in Peru, a pale morph...

Guiana dolphin
Guiana dolphin
Sotalia guianensis

The Guiana dolphin (*Sotalia guianensis*), also known as the estuarine dolphin or costero, is a dolphin found in the coastal waters to the north and east of South America, and east of Central America. It is a member of the oceanic dolphin family...

Guianan bonneted bat
Guianan bonneted bat
Eumops maurus

The Guianan bonneted bat, or Thomas's mastiff bat, (*Eumops maurus*) is a species of free-tailed bat from South America. It is poorly understood, because it is difficult to capture, and may be one of the rarest Neotropical bats.

Guianan spear-nosed bat
Guianan spear-nosed bat
Phyllostomus latifolius

The Guianan spear-nosed bat (*Phyllostomus latifolius*) is a bat species from South America. It is found in Brazil, Colombia, Guyana, Suriname and Venezuela.

Guianan Squirrel Monkey
Guianan Squirrel Monkey
Saimiri sciureus

The Guianan squirrel monkey (*Saimiri sciureus*) is a species of squirrel monkey from South America. This species formerly applied to Humboldt's squirrel monkey and Collins' squirrel monkey, but genetic research in 2009 and 2015 revealed that these...

Guianan white-eared opossum
Guianan white-eared opossum
Didelphis imperfecta

The Guianan white-eared opossum (*Didelphis imperfecta*) is an opossum species from South America. It is found in Brazil, Suriname, French Guiana and Venezuela. Possessing the smallest distribution area of its genus, this species is endemic to the...

Guinea Baboon
Guinea Baboon
Papio papio

The Guinea baboon (*Papio papio*) is a baboon from the Old World monkey family. Some (older) classifications list only two species in the genus *Papio*, this one and the hamadryas baboon. In those classifications, all other *Papio* species are...

Guinea multimammate mouse
Guinea multimammate mouse
Mastomys erythroleucus

The Guinea multimammate mouse (*Mastomys erythroleucus*) is a species of rodent in the family Muridae found in Benin, Burkina Faso, Burundi, Cameroon, the Central African Republic, Chad, the Republic of the Congo, the Democratic Republic of

the...

Guinean gerbil

Guinean gerbil

Gerbilliscus guineae

The Guinean gerbil (*Gerbilliscus guineae*) is a species of rodent found in Burkina Faso, Gambia, Ghana, Guinea, Guinea-Bissau, Ivory Coast, Mali, Senegal, Sierra Leone, and possibly Liberia. Its natural habitats are subtropical or tropical dry...

Guinean grass rat

Guinean grass rat

Arvicanthis rufinus

The Guinean grass rat (*Arvicanthis rufinus*) is a species of rodent in the family Muridae. It is found in Benin, Ghana, Togo, possibly Cameroon, possibly Central African Republic, possibly Ivory Coast, possibly Guinea, possibly Liberia, and...

Guinean horseshoe bat

Guinean horseshoe bat

Rhinolophus guineensis

The Guinean horseshoe bat (*Rhinolophus guineensis*) is a species of bat in the family Rhinolophidae. It is found in Ivory Coast, Guinea, Liberia, Senegal, and Sierra Leone. Its natural habitats are subtropical or tropical dry forest, subtropical...

Gulf Coast Jaguarundi

Gulf Coast Jaguarundi

Herpailurus yagouaroundi cacomitli

The Gulf Coast jaguarundi is an endangered population of the jaguarundi (*Herpailurus yagouaroundi*) once ranging from southern Texas in the United States to eastern Mexico. In 2017, the International Union for Conservation of Nature no longer...

Gulf Coast kangaroo rat

Gulf Coast kangaroo rat

Dipodomys compactus

The Gulf Coast kangaroo rat (*Dipodomys compactus*) is a species of rodent in the family Heteromyidae found in Mexico and the state of Texas in the United States. Its appearance and ecology are very similar to those of its putative sister species,...

Gull Island vole

Gull Island vole

Microtus pennsylvanicus nesophilus

The Gull Island vole (*Microtus pennsylvanicus nesophilus*) is a subspecies of the meadow vole last collected in 1897. A ground-dwelling coastal beach grass herbivore endemic to Great Gull Island, New York, it disappeared after habitat destruction...

Gunning's golden mole

Gunning's golden mole

Neamblysomus gunningi

Gunning's golden mole (*Neamblysomus gunningi*) is a small mammal endemic to South Africa. It was listed in 2006 as an endangered species. Its decreasing numbers are due to habitat clearance or destruction and predation from domesticated cats and...

Gunnison's Prairie Dog

Gunnison's Prairie Dog

Cynomys gunnisoni

Gunnison's prairie dog (*Cynomys gunnisoni*) is one of five species of the prairie

dog. This species belongs to the squirrel family of rodents, and are predominantly related to the North American and Eurasian ground squirrels. Gunnison's prairie...
Günther's Dik-Dik
Günther's Dik-Dik
Madoqua guentheri

Günther's dik-dik (*Madoqua guentheri*) is a small elegant antelope that lives in arid zones of East Africa. The name dik-dik comes from a characteristic sound produced by females through their long, tubular snouts when they feel threatened.
Günther's vole
Günther's vole
Microtus guentheri

Günther's vole (*Microtus guentheri*) is a species of rodent in the family Cricetidae, also known by the name Levant vole. It is found in Bulgaria, Greece, Iran, Iraq, Israel, Jordan, Lebanon, North Macedonia, Serbia and Montenegro, Syria, Turkey, ...
Guramba shrew
Guramba shrew
Crocidura phaeura

The Guramba shrew (*Crocidura phaeura*) is a species of mammal in the family Soricidae. It is endemic to Ethiopia. Its natural habitats are subtropical or tropical dry forests.
Gursky's spectral tarsier
Gursky's spectral tarsier
Tarsius spectrumgurskyae

Gursky's spectral tarsier (*Tarsius spectrumgurskyae*) is a species of tarsier found in the island of Sulawesi in Indonesia. A recent taxonomic revision split this species off from the spectral tarsier and other tarsier species based on difference...
Guyanan red howler
Guyanan red howler
Alouatta macconnelli

The Guyanan red howler (*Alouatta macconnelli*) is a species of howler monkey, a type of New World monkey, native to Suriname, Guyana, Trinidad, French Guiana, Venezuela and Brazil.
Guyanan spiny rat
Guyanan spiny rat
Proechimys hoplomysoides

The Guyanan spiny-rat, *Proechimys hoplomysoides*, is a spiny rat species from South America. It is found in Brazil, Guyana and Venezuela. The species was first described by George Henry Hamilton Tate in 1939.
Guyenne spiny rat
Guyenne spiny rat
Proechimys guyannensis

The Guyenne spiny-rat, *Proechimys guyannensis*, or Cayenne spiny rat, is a spiny rat species from South America. It is found in Brazil, Colombia, French Guiana, Guyana, Surinam and Venezuela.
Habbema dasyure
Habbema dasyure
Murexia habbema

The Habbema dasyure (*Murexia habbema*) is a species of marsupial in the family Dasyuridae found in West Papua, Indonesia, and Papua New Guinea. Its natural

habitat is rocky areas. Some authorities place it in the genus *Micromurexia* along with...

Habromys schmidlyi
Habromys schmidlyi
Habromys schmidlyi

Habromys schmidlyi, sometimes known as Schmidly's deer mouse, is a species of rodent in the family *Cricetidae* found only in Mexico. Its natural habitat is cloud forest at elevation over 1,800 m. The name "Schmidly's deer mouse" is ambiguous, as it...

Hadromys loujacobsi
Hadromys loujacobsi
Hadromys loujacobs

Hadromys loujacobsi is an extinct species of rat, known only from three molars of early Pleistocene origin. They were found in the Upper Siwalik Group in the Punjab region of Pakistan.

Hagen's flying squirrel
Hagen's flying squirrel
Petinomys hageni

Hagen's flying squirrel (*Petinomys hageni*) is a species of rodent in the family *Sciuridae*. It is endemic to Indonesia.

Haggard's leaf-eared mouse
Haggard's leaf-eared mouse
Phyllotis haggardi

Haggard's leaf-eared mouse (*Phyllotis haggardi*) is a species of rodent in the family *Cricetidae*. It is found only in Ecuador.

Haig's tuco-tuco
Haig's tuco-tuco
Ctenomys haigi

Haig's tuco-tuco (*Ctenomys haigi*), known regionally as the Patagonian tuco-tuco, is a South American hystricognath rodent. Like other tuco-tucos it is subterranean and thus not often observed, although the "tuc-tuc" call of the males can be...

Hainald's rat
Hainald's rat
Rattus hainaldi

Hainald's rat (*Rattus hainaldi*) is a species of rodent in the family *Muridae*. It is found only on Flores Island in Indonesia, including on Mount Ranaka. Part of its habitat is protected within the Kelimutu National Park.

Hainan Black-Crested Gibbon
Hainan Black-Crested Gibbon
Nomascus hainanus

The Hainan black-crested gibbon (*Nomascus hainanus*) is a Critically Endangered species of gibbon. It was formerly considered a subspecies of the Eastern black crested gibbon from Hòa Bình and Cao Bằng provinces of Vietnam and Jingxi County in...

Hainan gymnure
Hainan gymnure
Neohylomys hainanensis

The Hainan gymnure (*Neohylomys hainanensis*) or Hainan moonrat is a species of mammal in the family *Erinaceidae*. Its natural habitat is subtropical or tropical dry forests. It was thought to be endemic to the island of Hainan, China where it is...

Hainan hare
Hainan hare
Lepus hainanus

The Hainan hare (*Lepus hainanus*) is a hare endemic to Hainan Island, China. The species was first described by Robert Swinhoe in 1870.

Hainan Island shrew
Hainan Island shrew
Crocidura wuchihensis

The Hainan Island shrew (*Crocidura wuchihensis*) is a species of mammal in the family Soricidae native to China and Vietnam. The IUCN has insufficient data to assess the level of population and its trend.

Hairless bat
Hairless bat
Cheiromeles torquatus

The hairless bat (*Cheiromeles torquatus*), also called the naked bulldog bat and greater naked bat, is a species of bat in the family Molossidae. The generic name *Cheiromeles* comes from the Greek word *cheir* (Greek for hand) and the species name...

Hairy Atlantic spiny rat
Hairy Atlantic spiny rat
Trinomys setosus

The hairy Atlantic spiny rat, *Trinomys setosus*, is a spiny rat species from South America. It is endemic to Brazil.

Hairy big-eared bat
Hairy big-eared bat
Micronycteris hirsuta

The hairy big-eared bat (*Micronycteris hirsuta*) is a bat species from South and Central America, as well as Trinidad and Tobago in the Caribbean.

Hairy big-eyed bat
Hairy big-eyed bat
Chiroderma villosum

The hairy big-eyed bat (*Chiroderma villosum*) is a bat species from South and Central America.

Hairy fruit-eating bat
Hairy fruit-eating bat
Artibeus hirsutus

The hairy fruit-eating bat (*Artibeus hirsutus*) is a species of bat in the family Phyllostomidae. It is endemic to Mexico. It is threatened by habitat loss.

Hairy harvest mouse
Hairy harvest mouse
Reithrodontomys hirsutus

The hairy harvest mouse (*Reithrodontomys hirsutus*) is a species of rodent in the family Cricetidae found only in Mexico.

Hairy little fruit bat
Hairy little fruit bat
Rhinophylla alethina

The hairy little fruit bat (*Rhinophylla alethina*) is a species of bat in the family Phyllostomidae found in Colombia and Ecuador. They are nocturnal creatures. They are listed as near-threatened by the IUCN.

Hairy long-nosed armadillo

Hairy long-nosed armadillo
Dasypus pilosus

The hairy long-nosed armadillo or woolly armadillo (*Dasypus pilosus*) is a species of armadillo in the family Dasypodidae. It is endemic to Peru. Its natural habitats are subtropical or tropical moist lowland forest and subtropical or tropical...

Hairy slit-faced bat
Hairy slit-faced bat
Nycteris hispida

The hairy slit-faced bat (*Nycteris hispida*) is a species of slit-faced bat widely distributed throughout forests and savannas in Africa. Two recognized subspecies exist: *N. h. hispida* and *N. h. pallida*. Various forest populations in western and...

Hairy yellow-shouldered bat
Hairy yellow-shouldered bat
Sturnira erythromos

The hairy yellow-shouldered bat (*Sturnira erythromos*) is a species of bat in the family Phyllostomidae native to South America. There are no recognised subspecies.

Hairy-eared cerrado mouse
Hairy-eared cerrado mouse
Thalpomys lasiotis

The hairy-eared cerrado mouse (*Thalpomys lasiotis*) or hairy-eared akodont, is a rodent species from South America. It is found in the cerrado grassland of Brazil.

Hairy-eared dwarf lemur
Hairy-eared dwarf lemur
Allocebus trichotis

The hairy-eared dwarf lemur (*Allocebus trichotis*), or hairy-eared mouse lemur, is one of the most scarcely known lemurs. *A. trichotis* is a nocturnal lemur that is endemic to Madagascar. It was originally named by Dr. Albert Günther in 1875 as...

Hairy-faced bat
Hairy-faced bat
Myotis annectans

The hairy-faced bat (*Myotis annectans*) is a species of vesper bat. It can be found in the following countries: India, Indonesia, Laos, Myanmar, and Thailand.

Hairy-footed dunnart
Hairy-footed dunnart
Sminthopsis hirtipes

The hairy-footed dunnart (*Sminthopsis hirtipes*) is a dunnart that has silver hairs on the soles of its hind feet accompanied by long hair on the side of its sole. It is an Australian marsupial similar to the Ooldea dunnart, with its upper body...

Hairy-footed flying squirrel
Hairy-footed flying squirrel
Belomys pearsonii

The hairy-footed flying squirrel (*Belomys pearsonii*) is a flying squirrel found in the mountains of the eastern Himalaya, Southeast Asia, southern China, and the island of Taiwan. It lives at elevations of 800–2,400 m (2,600–7,900 ft) above sea...

Hairy-footed gerbil
Hairy-footed gerbil
Gerbillurus paeba

The hairy-footed gerbil (*Gerbillurus paeba*) is a species of rodent found in

Angola, Botswana, Mozambique, Namibia, South Africa, and Zimbabwe. Its natural habitats are dry savanna, temperate shrubland, hot deserts, sandy shores, and urban areas....

Hairy-Fronted Muntjac
Hairy-Fronted Muntjac
Muntiacus crinifrons

The Hairy-fronted muntjac (*Muntiacus crinifrons*) is a species of deer currently found in East Asia. It is considered to be endangered, and was for a very long time one of the most poorly known deer in the world. It was also considered highly...

Hairy-legged myotis
Hairy-legged myotis
Myotis keaysi

The hairy-legged myotis (*Myotis keaysi*) is a species of mouse-eared bat. It is found from southern Tamaulipas in Mexico, through much of Central America and across northern South America as far east as Trinidad. Further south, it is found along...

Hairy-Legged Vampire Bat
Hairy-Legged Vampire Bat
Diphylla ecaudata

The hairy-legged vampire bat (*Diphylla ecaudata*) is one of three extant species of vampire bats. It mainly feeds on the blood of wild birds, but can also feed both on domestic birds and humans. This vampire bat lives mainly in tropical and...

Hairy-Nosed Otter
Hairy-Nosed Otter
Lutra sumatrana

The hairy-nosed otter (*Lutra sumatrana*) is a semiaquatic mammal endemic to Southeast Asia and one of the rarest and least known otter species. It is threatened by loss of natural resources and poaching.

Hairy-tailed antsangy
Hairy-tailed antsangy
Brachytarsomys villosa

The hairy-tailed antsangy (*Brachytarsomys villosa*) is a species of rodent in the family Nesomyidae. It is found only in Madagascar.

Hairy-tailed bat
Hairy-tailed bat
Lasiurus ebonus

The hairy-tailed bat (*Lasiurus ebonus*), is a bat species originally known only from its type locality, Ilha do Cardoso State Park in Brazil. A second specimen was collected in 2018, in Carlos Botelho State Park, approximately 100 kilometers away.

Hairy-tailed bolo mouse
Hairy-tailed bolo mouse
Necromys lasiurus

The hairy-tailed bolo mouse or hairy-tailed akodont (*Necromys lasiurus*) is a South American rodent species of the family Cricetidae. It is found in Argentina, Bolivia, Brazil, and Paraguay.

Hairy-tailed mole
Hairy-tailed mole
Parascalops breweri

The hairy-tailed mole (*Parascalops breweri*), also known as Brewer's mole, is a medium-sized North American mole. It is the only member of the genus *Parascalops*. The species epithet *breweri* refers to Thomas Mayo Brewer, an American naturalist.

Halcyon horseshoe bat
Halcyon horseshoe bat
Rhinolophus alcyone

The halcyon horseshoe bat (*Rhinolophus alcyone*) is a species of bat in the family Rhinolophidae. It is found in Cameroon, Republic of the Congo, Democratic Republic of the Congo, Ivory Coast, Equatorial Guinea, Ghana, Guinea, Kenya, Liberia,...

Halmahera blossom bat
Halmahera blossom bat
Syconycteris carolinae

The Halmahera blossom bat (*Syconycteris carolinae*) is a species of megabat in the family Pteropodidae. The bat is endemic to Halmahera and Bacan islands, of the northern Maluku Islands archipelago in Indonesia. It is an IUCN Red List Near...

Halmahera naked-backed fruit bat
Halmahera naked-backed fruit bat
Dobsonia crenulata

The Halmahera naked-backed fruit bat (*Dobsonia crenulata*) is a common and widespread species of megabat in the family Pteropodidae. It is endemic to Indonesia. It is assessed as least-concern by the IUCN as it has a wide range and seems to have...

Halmaheramys bokimekot
Halmaheramys bokimekot
Halmaheramys bokimekot

Halmaheramys bokimekot, or the spiny Boki Mekot rat, is a rodent found on the island of Halmahera in the Molucca archipelago, whose discovery was announced in 2013. This is the only locality where this particular species has been found. H....

Hamadryas Baboon
Hamadryas Baboon
Papio hamadryas

The strong and robust Hamadryas baboons (*Papio hamadryas*) belong to Old World monkeys. They are highly intelligent animals. Mature males of this species are identified by their silver manes and pink faces. In Ancient Egypt, they were considered...

Hamiguitan hairy-tailed rat
Hamiguitan hairy-tailed rat
Batomys hamiguitan

The Hamiguitan hairy-tailed rat (*Batomys hamiguitan*) is one of five species of rodent in the genus *Batomys*. It is in the diverse family Muridae. This species is found only in the Philippines. is a yellow-brown animal with a long furry tail,...

Hamilton's tomb bat
Hamilton's tomb bat
Taphozous hamiltoni

Hamilton's tomb bat (*Taphozous hamiltoni*) is a species of sac-winged bat in the family Emballonuridae. It is found in Chad, Kenya, Somalia, South Sudan, Tanzania, and Uganda. Its natural habitats are dry savanna and moist savanna. It is threatened...

Hamlyn's monkey
Hamlyn's monkey
Cercopithecus hamlyni

The Hamlyn's monkey (*Cercopithecus hamlyni*), also known as the owl-faced monkey, is a species of Old World monkey that inhabits the bamboo and primary rainforests of the Congo. This species is exceedingly rare and known only from a few...

Hammer-Headed Bat
Hammer-Headed Bat
Hypsignathus monstrosus

The Hammer-headed bat (*Hypsignathus monstrosus*) is a megabat that is found in Africa. It is the largest bat in continental Africa, with wingspans approaching 1 m (3.3 ft), and males almost twice as heavy as females. The Hammer-headed bat is...

Handley's nectar bat
Handley's nectar bat
Lonchophylla handleyi

Handley's nectar bat (*Lonchophylla handleyi*) is a species of bat in the family Phyllostomidae. It is found in Colombia, Ecuador, and Peru. It is threatened by habitat loss.

Handley's slender opossum
Handley's slender opossum
Marmosops handleyi

Handley's slender opossum (*Marmosops handleyi*) is a species of opossum in the family Didelphidae. It is endemic to Colombia.

Handley's tailless bat
Handley's tailless bat
Anoura cultrata

Handley's tailless bat (*Anoura cultrata*) is a species of bat in the family Phyllostomidae. It is found in Bolivia, Colombia, Costa Rica, Ecuador, Panama, Peru, and Venezuela.

Handleyomys alfaroi
Handleyomys alfaroi
Handleyomys alfaroi

Handleyomys alfaroi, also known as Alfaro's rice rat or Alfaro's oryzomys, is a species of rodent in the genus *Handleyomys* of family Cricetidae. It is found in Belize, Colombia, Costa Rica, Ecuador, El Salvador, Guatemala, Honduras, Mexico,...

Handleyomys chapmani
Handleyomys chapmani
Handleyomys chapmani

Handleyomys chapmani, also known as Chapman's oryzomys or Chapman's rice rat, is a species of rodent in the genus *Handleyomys* of family Cricetidae. It is found only in Mexico. It was previously placed in *Oryzomys* as *Oryzomys chapmani*, but has...

Handleyomys fuscatus
Handleyomys fuscatus
Handleyomys fuscatus

Handleyomys fuscatus, also known as the dusky-footed Handley's mouse or dusky-footed montane mouse, is a species of rodent in the tribe Oryzomyini of family Cricetidae. It was previously placed in the genus *Aepeomys*, but it is closely similar to...

Handleyomys intectus
Handleyomys intectus
Handleyomys intectus

The Colombian rice rat (*Handleyomys intectus*), also known as the white-footed Handley's mouse is a species of rodent in the family Cricetidae occurring only in Colombia. It was previously included in *Oryzomys*, but closely resembles the dusky...

Handleyomys melanotis
Handleyomys melanotis

Handleyomys melanotis

Handleyomys melanotis, also known as the black-eared oryzomys or black-eared rice rat, is a species of rodent in the genus *Handleyomys* of family Cricetidae. It is found in coastal lowland forest in western Mexico.

Handleyomys rhabdops
Handleyomys rhabdops
Handleyomys rhabdops

Handleyomys rhabdops, also known as the highland oryzomys or striped rice rat, is a species of rodent in the genus *Handleyomys* of family Cricetidae. It is nocturnal and is found in Guatemala and Mexico in montane forest at elevations from 1250 to...

Handleyomys rostratus
Handleyomys rostratus
Handleyomys rostratus

Handleyomys rostratus, also known as the long-nosed oryzomys, long-nosed rice rat, or rusty rice rat is a species of rodent in the genus *Handleyomys* of family Cricetidae. It is found in Belize, El Salvador, Guatemala, Honduras, Mexico, and...

Handleyomys saturatior
Handleyomys saturatior
Handleyomys saturatior

Handleyomys saturatior, also known as the cloud forest oryzomys or cloud forest rice rat, is a species of rodent in the genus *Handleyomys* of family Cricetidae. It is found in Belize, El Salvador, Guatemala, Honduras, Mexico, and Nicaragua in cloud...

Hapalomys gracilis
Hapalomys gracilis
Hapalomys gracilis

Hapalomys gracilis is a fossil rodent from the genus *Hapalomys* found in Longgupo in South China. The species designation *gracilis* is Latin for 'thin'. Only a first subchase of this species is known. This molar is smaller than that of all other...

Harbor Seal
Harbor Seal
Phoca vitulina

The Harbor seal (*Phoca vitulina*) is a true seal that lives along temperate and Arctic marine coastlines of the Northern Hemisphere. It is the most widely distributed species of pinniped (walruses, eared seals, and true seals). Harbor seals like...

Harbour Porpoise
Harbour Porpoise
Phocoena phocoena

The Harbour porpoise (*Phocoena phocoena*) is one of eight extant species of porpoise. It is one of the smallest species of cetacean. As its name implies, it stays close to coastal areas or river estuaries, and as such, is the most familiar...

Hardwicke's woolly bat
Hardwicke's woolly bat
Kerivoula hardwickii

Hardwicke's woolly bat (*Kerivoula hardwickii*) is a species of vesper bat in the family Vespertilionidae.

Haremma shrew
Haremma shrew

Crocidura harena

The Harena shrew (*Crocidura harena*) is a white-toothed shrew found only in one location in the Bale Mountains in southern Ethiopia. It occurs within an area of less than ten square kilometres, and is listed as a critically endangered species,...

Harlequin bat

Harlequin bat

Scotomanes ornatus

The harlequin bat (*Scotomanes ornatus*) is a species of bat in the family Vespertilionidae, the vesper bats. It is the only member of the monotypic genus *Scotomanes*. This bat is found in south-eastern Asia from India to China and Vietnam. This is a...

Harmless serotine

Harmless serotine

Eptesicus innoxius

The harmless serotine (*Eptesicus innoxius*) is a species of vesper bat. It has a restricted range in western Ecuador and northwestern Peru. An insectivorous species, it is a resident of tropical dry forest habitat, and is threatened by...

Harnessed bushbuck

Harnessed bushbuck

Tragelaphus scriptus

The bushbuck (*Tragelaphus scriptus*) is the nominate taxon of the bushbuck. It is a small to medium-sized antelope widespread in Africa. The Cape bushbuck is a southern and eastern subspecies which is recognised by some authors, which found...

Harp Seal

Harp Seal

Pagophilus groenlandicus

The Harp Seal (*Pagophilus groenlandicus*) is a species of earless seal, or true seal, native to the northernmost Atlantic Ocean and Arctic Ocean. In Greek, its scientific name translates to "ice-lover from Greenland," and its taxonomic synonym,...

Harpy fruit bat

Harpy fruit bat

Harpyionycteris whiteheadi

The harpy fruit bat (*Harpyionycteris whiteheadi*) is a species of megabat in the family Pteropodidae. It is endemic to the Philippines.

Harrington's gerbil

Harrington's gerbil

Taterillus harringtoni

Harrington's gerbil or Harrington's tateril (*Taterillus harringtoni*) is a species of rodent found in the Central African Republic, Ethiopia, Kenya, Somalia, South Sudan, Tanzania, and Uganda. Its natural habitats are dry savanna, subtropical or...

Harrington's rat

Harrington's rat

Desmomys harringtoni

Harrington's rat (*Desmomys harringtoni*) is a species of rodent in the family Muridae. It is found only in Ethiopia. Its natural habitats are subtropical or tropical moist montane forests and subtropical or tropical moist shrubland.

Harris's Antelope Squirrel

Harris's Antelope Squirrel

Ammospermophilus harrisi

Harris's antelope squirrel (*Ammospermophilus harrisi*) is a species of rodent in the family Sciuridae. It is found in Arizona and New Mexico in the United States, and in Sonora in Mexico. They are adapted to hot weather conditions including a...

Harrison's gerbil

Harrison's gerbil

Gerbillus mesopotamiae

Harrison's gerbil (*Gerbillus mesopotamiae*) is a gerbil, a small mammal in the rodent order. It is distributed mainly in the Tigris-Euphrates Valley in Iraq and western Iran. It is also known as the Mesopotamian gerbil.

Harrison's large-eared giant mastiff bat

Harrison's large-eared giant mastiff bat

Otomops harrisoni

Harrison's large-eared giant mastiff bat (*Otomops harrisoni*) is a species of bat found in Northeast Africa and the Arabian Peninsula. It was described as a new species in 2015. The IUCN evaluates it as a vulnerable species.

Harrison's tube-nosed bat

Harrison's tube-nosed bat

Murina harrisoni

Harrison's tube-nosed bat (*Murina harrisoni*) is a species of vesper bats (Vespertilionidae). Within the genus *Murina*, it belongs to the so-called 'cyclotis-group'. This species was only recently discovered and described. It is characterised by...

Hartebeest

Hartebeest

Alcelaphus buselaphus

The hartebeest (*Alcelaphus buselaphus*) is an African antelope. It is the only member of the genus *Alcelaphus*. Hartebeest are gregarious animals. They are very alert and non-aggressive. Hartebeest were formerly widespread in Africa, but...

Hartmann's Mountain Zebra

Hartmann's Mountain Zebra

Equus zebra hartmannae

Hartmann's mountain zebra (*Equus zebra hartmannae*) is a subspecies of the Mountain zebra found in Africa. It is easily distinguished from other similar zebra species by its dewlap and the lack of stripes on its belly.

Hartwig's soft-furred mouse

Hartwig's soft-furred mouse

Praomys hartwigi

Hartwig's soft-furred mouse or Hartwig's praomys (*Praomys hartwigi*) is a species of rodent in the family Muridae. It is found only in Cameroon. Its natural habitat is subtropical or tropical moist montane forest. It is threatened by habitat loss.

Harvey's duiker

Harvey's duiker

Cephalophus harveyi

The Harvey's red duiker (*Cephalophus harveyi*) is one of 19 species of duiker found in Tanzania and scattered through Kenya, southern Somalia and possibly central Ethiopia. Harvey's duikers are roughly 40 cm (15 in) tall at the shoulder and weigh...

Harwood's gerbil

Harwood's gerbil

Dipodillus harwoodi

Harwood's gerbil (*Dipodillus harwoodi*) is distributed mainly in Kenya.
Hastings River mouse
Hastings River mouse
Pseudomys oralis

The Hastings River mouse (*Pseudomys oralis*) is a species of Australian rodent in the subfamily Murinae of the family Muridae. It is found only in Australia.
Hatinh langur
Hatinh langur
Trachypithecus hatinhensis

The Hatinh langur (*Trachypithecus hatinhensis*) is a highly threatened Old World monkey found in limestone forests in Vietnam, primarily in the Quảng Bình Province. A recent survey discovered a small population living in the Quảng Trị Province....
Hatt's thicket rat
Hatt's thicket rat
Thamnomys major

Hatt's thicket rat (*Thamnomys major*), is a species of thicket rat indigenous to the northern slopes of Mount Karisimbi, an inactive volcano in North Kivu (Nord-Kivu) province of the Democratic Republic of the Congo (DRC). Since the species is...
Hatt's vesper rat
Hatt's vesper rat
Otonyctomys hatti

Hatt's vesper rat (*Otonyctomys hatti*), also known as Hatt's vesper mouse or Yucatán vesper rat, is a species of rodent in the family Cricetidae. It is the only species in the genus *Otonyctomys*. It is named for its discoverer, Robert T. Hatt.
Hausa mouse
Hausa mouse
Mus haussa

The Hausa mouse (*Mus haussa*) is a species of rodent in the family Muridae. It is found in Benin, Burkina Faso, Ivory Coast, Ghana, Mali, Mauritania, Niger, Nigeria, and Senegal. Its natural habitats are dry savanna, arable land, rural gardens, and...
Hausa genet
Hausa genet
Genetta thierryi

The Hausa genet (*Genetta thierryi*) is a genet species native to West African savannas. It is listed as Least Concern on the IUCN Red List. Hausa genets have been sighted in Senegal's wooded steppes, in moist woodlands in Guinea-Bissau, and in...
Hawaiian Hoary Bat
Hawaiian Hoary Bat
Lasiurus semotus

The Hawaiian hoary bat (*Aeorestes semotus*), also known as 'ōpe'ape'a, is a species of bat endemic to the islands of Hawai'i. Whereas the mainland hoary bat (*A. cinereus*) is found throughout North America, the Hawaiian hoary bat is distributed...
Hawaiian Monk Seal
Hawaiian Monk Seal
Neomonachus schauinslandi

The Hawaiian monk seal (*Neomonachus schauinslandi*) is an endangered species of earless seal native to the Hawaiian Islands. It is one of two extant monk seal

species; the other is the Mediterranean monk seal. A third species, the Caribbean monk...

Hawks' sportive lemur

Hawks' sportive lemur

Lepilemur tymerlachsoni

Hawks' sportive lemur (*Lepilemur tymerlachsoni*), also known as the Nosy Be sportive lemur or Nosy Be weasel lemur, is a sportive lemur endemic to Madagascar. It is a moderately large sportive lemur, with a total length of about 50 to 68 cm (20...

Hayman's dwarf epauletted fruit bat

Hayman's dwarf epauletted fruit bat

Micropteropus intermedius

Hayman's dwarf epauletted fruit bat or Hayman's epauletted fruit bat (*Micropteropus intermedius*) is a species of megabat in the family Pteropodidae. It is found in Angola and Democratic Republic of the Congo. Its natural habitats are subtropical...

Hazel Dormouse

Hazel Dormouse

Muscardinus avellanarius

The hazel dormouse or common dormouse (*Muscardinus avellanarius*) is a small mammal and the only living species in the genus *Muscardinus*.

Heart-nosed bat

Heart-nosed bat

Cardioderma cor

The heart-nosed bat (*Cardioderma cor*) is a species of bat in the family Megadermatidae. It is monotypic within the genus *Cardioderma*. It is found in eastern Sudan, north Tanzania, and south Zambia. In an experiment conducted in 2017, it was...

Heath mouse

Heath mouse

Pseudomys shortridgei

The heath mouse (*Pseudomys shortridgei*) is a species of mouse in the subfamily Murinae, the Old World rats and mice, found in Australia.

Heather shrew

Heather shrew

Crocidura erica

The heather shrew (*Crocidura erica*) is a species of mammal in the family Soricidae. It is endemic to Angola. It is threatened by habitat loss.

Heavenly hill rat

Heavenly hill rat

Bunomys coelestis

The heavenly hill rat (*Bunomys coelestis*) is a species of rodent in the family Muridae. It is found only in southwestern Sulawesi, Indonesia, where it has only been found on Mount Lompobatang. Its natural habitat is subtropical or tropical dry...

Heaviside's Dolphin

Heaviside's Dolphin

Cephalorhynchus heavisidii

Heaviside's dolphins (*Cephalorhynchus heavisidii*) are small stocky cetaceans native to the southwest coast of Africa. They spend time in small groups and usually remain nearshore in the mornings. They are very energetic and can dive up to 147...

Heavy-browed mouse opossum
Heavy-browed mouse opossum
Marmosa andersoni

The heavy-browed mouse opossum (*Marmosa andersoni*), or Anderson's mouse opossum, is a species of opossum in the family Didelphidae. It is endemic to a restricted range in southern Peru. This opossum inhabits forests; it is nocturnal and probably...

Heck's macaque
Heck's macaque
Macaca hecki

The Heck's macaque (*Macaca hecki*) is a macaque of Sulawesi, Indonesia. This Old World monkey is diurnal. It feeds on fruits.

Hector's beaked whale
Hector's beaked whale
Mesoplodon hectori

Hector's beaked whale (*Mesoplodon hectori*), is a small mesoplodont living in the Southern Hemisphere. This whale is named after Sir James Hector, a founder of the colonial museum in Wellington, New Zealand. The species has rarely been seen in...

Hector's Dolphin
Hector's Dolphin
Cephalorhynchus hectori

Hector's dolphin (*Cephalorhynchus hectori*) is one of four dolphin species belonging to the genus *Cephalorhynchus*. Hector's dolphin is the only cetacean endemic to New Zealand, and comprises two subspecies: *C. h. hectori*, the more numerous...

Heermann's kangaroo rat
Heermann's kangaroo rat
Dipodomys heermanni

Heermann's kangaroo rat (*Dipodomys heermanni*) is a species of rodent in the family Heteromyidae. Their long smooth pelage resembles typical kangaroo rats, with their dorsal side showing a mixed range of olive, black and orange colors. There are...

Helan Shan pika
Helan Shan pika
Ochotona argentata

The Helan Shan pika or silver pika (*Ochotona argentata*) is a species of mammal in the pika family, Ochotonidae. It is endemic to China where it is found in a small region of the Helan Mountains. It is listed as "Endangered" in the IUCN Red List...

Heller's broad-nosed bat
Heller's broad-nosed bat
Platyrrhinus helleri

Heller's broad-nosed bat (*Platyrrhinus helleri*) is a bat species from South and Central America.

Heller's pipistrelle
Heller's pipistrelle
Neoromicia helios

Heller's serotine (*Afronycteris helios*), formerly known as Heller's pipistrelle, is a species of vesper bat. It is only found in Africa, in countries such as Kenya, Somalia, Tanzania, and Uganda.

Hellwald's spiny rat
Hellwald's spiny rat
Maxomys hellwaldii

Hellwald's spiny rat (*Maxomys hellwaldii*) is a species of rodent in the family Muridae. It is found only in Indonesia.

Heptner's pygmy jerboa

Heptner's pygmy jerboa

Salpingotus heptneri

Heptner's pygmy jerboa (*Salpingotus heptneri*) is a species of rodent in the family Dipodidae. It is native to temperate deserts in Kazakhstan, Uzbekistan, and possibly Russia.

Herbert River ringtail possum

Herbert River ringtail possum

Pseudochirulus herbertensis

The Herbert River ringtail possum (*Pseudochirulus herbertensis*) is a species of possum found in northeastern Queensland, Australia. It is a dark brown to black species, sometimes with a white belly. The Herbert River ringtail possum was long...

Herbert's rock-wallaby

Herbert's rock-wallaby

Petrogale herberti

Herbert's rock-wallaby (*Petrogale herberti*) is a member of a group of seven very closely related rock-wallabies found in northeastern Queensland, Australia.

Herbert's is the most southerly and most widespread of the group. Herbert's rock-wallaby...

Herman's myotis

Herman's myotis

Myotis hermani

Herman's myotis (*Myotis hermani*) is a species of vesper bat. It is found only in Indonesia.

Hernández-Camacho's night monkey

Hernández-Camacho's night monkey

Aotus jorgehernandezi

Hernández-Camacho's night monkey (*Aotus jorgehernandezi*) is a species of night monkey of the family Aotidae. It was first described in 2007 by Thomas Defler and Marta Bueno. It has a gray neck and a white patch over each eye, separated by a...

Hero Shrew

Hero Shrew

Scutisorex somereni

The Hero shrew (*Scutisorex somereni*) is a large mammal native to the Congo Basin of Africa. It has corrugated interlocking vertebrae that are unique among mammals except for its sister species, Thor's hero shrew. This unique adaptation allows the...

Herpestes lemanensis

Herpestes lemanensis

Herpestes lemanensis

Herpestes lemanensis is an extinct mongoose species that was excavated in Tertiary depositions in France and described by Auguste Pomel in 1853. Its body size equals a large Indian civet, but its dentition resembles that of the Indian grey mongoose.

Hershkovitz's marmoset

Hershkovitz's marmoset

Mico intermedia

The Hershkovitz's marmoset (*Mico intermedia*), also known as the Aripuanã marmoset

is a marmoset species endemic to the south-central Amazon rainforest in Brazil. The common name is a reference to American zoologist Philip Hershkovitz.

Hershkovitz's titi

Hershkovitz's titi

Plecturocebus dubius

Hershkovitz's titi (*Plecturocebus dubius*) is a species of titi, a type of New World monkey, from South America. It is found in Bolivia, Brazil, and Peru. The common name is in reference to American zoologist Philip Hershkovitz, who described the...

Heude's Pig

Heude's Pig

Sus bucculentus

The Heude's pig (*Sus bucculentus*), also known as the Indochinese warty pig or Vietnam warty pig, was an alleged pig species found in Laos and Vietnam. It was virtually unknown and was feared extinct until the discovery of a skull from a recently...

Heuglin's gazelle

Heuglin's gazelle

Eudorcas tilonura

Heuglin's gazelle (*Eudorcas tilonura*), also known as the Eritrean gazelle, is a species of gazelle found east of the Nile River in Eritrea, Ethiopia and Sudan. It was considered a subspecies of the red-fronted gazelle (*E. rufifrons*) or...

Heuglin's striped grass mouse

Heuglin's striped grass mouse

Lemniscomys zebra

Heuglin's lemniscomys or Heuglin's striped grass mouse (*Lemniscomys zebra*) is a species of rodent in the family Muridae. It is found in Benin, Burkina Faso, Chad, Democratic Republic of the Congo, Ivory Coast, Gambia, Ghana, Guinea,...

Hewitt's red rock hare

Hewitt's red rock hare

Pronolagus saundersiae

The Hewitt's red rock hare (*Pronolagus saundersiae*) is a species of mammal in the family Leporidae. It had previously been classified as a subspecies of *Pronolagus rupestris*, but is now regarded as its own species due to differences in...

Highland brush mouse

Highland brush mouse

Abeomelomys sevia

The highland brush mouse (*Abeomelomys sevia*), also known as the Menzies' mouse, is a species of rodent in the family Muridae. It is endemic to Papua New Guinea, and is found in montane moss forests and in alpine zones over 2,000 m. It is the...

Highland Cattle

Highland Cattle

Bos primigenius taurus

The Highland is a Scottish breed of rustic cattle. It originated in the Scottish Highlands and the Outer Hebrides islands of Scotland and has long horns and a long shaggy coat. It is a hardy breed, able to withstand the intemperate conditions in...

Highland streaked tenrec

Highland streaked tenrec

Hemicentetes nigriceps

The highland streaked tenrec (*Hemicentetes nigriceps*) is an insectivore which

lives in the central upland regions of Madagascar. Its black and white striped body is covered with quills, which it will raise when agitated. The spines detach and...
Highland tuco-tuco
Highland tuco-tuco
Ctenomys opimus

The highland tuco-tuco (*Ctenomys opimus*) is a species of rodent in the family Ctenomyidae. It is found in high grassland in Argentina, Bolivia, Chile, and Peru where it lives in burrows.
Highland yellow-shouldered bat
Highland yellow-shouldered bat
Sturnira ludovici

The highland yellow-shouldered bat (*Sturnira ludovici*) is a species of bat in the family Phyllostomidae. It is found in Colombia, Costa Rica, Ecuador, El Salvador, Guatemala, Guyana, Honduras, Mexico, Nicaragua, Panama, and Venezuela.
Highlands punaré
Highlands punaré
Thrichomys inermis

The highlands punaré, *Thrichomys inermis*, is a caviomorph rodent of South America from the spiny rat family. It is endemic to gallery forest, savanna and rocky outcrop habitats in Bahia State within the Caatinga ecoregion of eastern Brazil at...
Highveld gerbil
Highveld gerbil
Gerbilliscus brantsii

The highveld gerbil (*Gerbilliscus brantsii*) is a species of rodent found in Angola, Botswana, Lesotho, Namibia, South Africa, Eswatini, Zambia, and Zimbabwe. Its natural habitats are dry savanna, temperate shrubland, subtropical or tropical dry...
Highveld golden mole
Highveld golden mole
Amblysomus septentrionalis

The highveld golden mole (*Amblysomus septentrionalis*) is a species of mammal in the golden mole family, Chrysochloridae. It is found in South Africa and Eswatini. Its natural habitats are temperate forests, subtropical or tropical dry forest,...
Hildebrandt's horseshoe bat
Hildebrandt's horseshoe bat
Rhinolophus hildebrandtii

Hildebrandt's horseshoe bat (*Rhinolophus hildebrandtii*) is a species of bat in the family Rhinolophidae found in Africa. Its natural habitats are dry savanna, moist savanna, caves, and subterranean habitats (other than caves).
Hildegard's broad-headed mouse
Hildegard's broad-headed mouse
Zelotomys hildegardae

Hildegard's broad-headed mouse (*Zelotomys hildegardae*) or Hildegard's zelotomys, is a species of rodent in the family Muridae. It is found in Central Africa.
Hildegard's shrew
Hildegard's shrew
Crocidura hildegardae

Hildegard's shrew (*Crocidura hildegardae*) is a species of mammal in the family Soricidae. It is found in Burundi, Cameroon, Central African Republic, Republic of

the Congo, Democratic Republic of the Congo, Ethiopia, Kenya, Rwanda, and...
Hildegarde's tomb bat
Hildegarde's tomb bat
Taphozous hildegardeae

Hildegarde's tomb bat (*Taphozous hildegardeae*) is a species of sac-winged bat in the family Emballonuridae. It is found near the coast in Kenya and Tanzania where it feeds in tropical dry forests and roosts in caves. It is a diurnal species and...
Hilgendorf's tube-nosed bat
Hilgendorf's tube-nosed bat
Murina hilgendorfi

Hilgendorf's tube-nosed bat (*Murina hilgendorfi*) is a species of vesper bat in the family Vespertilionidae. In Japan they are called 'tengu komori', after the mythical creature called the Tengu. It was formerly thought to be a subspecies of...
Hill's roundleaf bat
Hill's roundleaf bat
Hipposideros edwardshilli

Hill's roundleaf bat (*Hipposideros edwardshilli*) is a species of bat in the family Hipposideridae endemic to Papua New Guinea.
Hill's sheath-tailed bat
Hill's sheath-tailed bat
Taphozous hilli

Hill's sheath-tailed bat (*Taphozous hilli*) is a bat of the family Emballonuridae. They are found in the deserts of central Australia.
Hill's shrew
Hill's shrew
Crocidura hilliana

Hill's shrew (*Crocidura hilliana*) is a species of mammal in the family Soricidae. It is found in Thailand and Laos.
Hills' horseshoe bat
Hills' horseshoe bat
Rhinolophus hillorum

The Hills' horseshoe bat (*Rhinolophus hillorum*) is a species of bat in the family Rhinolophidae. It is found in Cameroon, Guinea, Liberia, and Nigeria. Its natural habitats are subtropical or tropical moist lowland forest, subtropical or...
Hilton Head white-tailed deer
Hilton Head white-tailed deer
Odocoileus virginianus hiltonensis

Hilton Head white-tailed deer (*Odocoileus virginianus hiltonensis*) are a subspecies of white-tailed deer indigenous to Hilton Head Island in South Carolina. The deer live in a mainly suburban environment and have developed (according to a study)...
Himalayan black bear
Himalayan black bear
Ursus thibetanus laniger

The Himalayan black bear (*Ursus thibetanus laniger*) is a subspecies of the Asian black bear found in the Himalayas of India, Bhutan, Nepal, China, and Pakistan.
Himalayan brown bear
Himalayan brown bear
Ursus arctos isabellinus

The Himalayan brown bear (*Ursus arctos isabellinus*), also known as the Himalayan red bear, isabelline bear or Dzu-Teh, is a subspecies of the brown bear and is known from northern Afghanistan, northern Pakistan, northern India, west China and...

Himalayan field mouse

Himalayan field mouse

Apodemus gurkha

The Himalayan field mouse (*Apodemus gurkha*) is a species of rodent in the family Muridae. It is endemic to Nepal.

Himalayan field rat

Himalayan field rat

Rattus nitidus

The Himalayan field rat (*Rattus nitidus*), sometimes known as the white-footed Indo-Chinese rat, is a species of rodent in the family Muridae. It has a wide range, being found in India, Bangladesh, Nepal, Bhutan, China, Myanmar, Laos, Thailand,...

Himalayan Goral

Himalayan Goral

Naemorhedus goral

The Himalayan goral (*Naemorhedus goral*) is a bovid species native to the Himalayas. It is listed as Near Threatened on the IUCN Red List because the population is thought to be declining significantly due to habitat loss and hunting for meat.

Himalayan Marmot

Himalayan Marmot

Marmota himalayana

The Himalayan marmot (*Marmota himalayana*) is a marmot species that inhabits alpine grasslands throughout the Himalayas and on the Tibetan Plateau. It is IUCN Red Listed as Least Concern because of its wide range and possibly large population.

Himalayan mole

Himalayan mole

Euroscaptor micrura

The Himalayan mole (*Euroscaptor micrurus*) or short-tailed mole is a species of mammal in the family Talpidae.

Himalayan pika

Himalayan pika

Ochotona himalayana

The Himalayan pika (*Ochotona himalayana*) is a species of small mammal in the pika family (*Ochotonidae*). It is found at high altitudes in remote areas of Ladakh, Uttarakhand and possibly also in Nepal & Tibet. The IUCN has listed this species as...

Himalayan Serow

Himalayan Serow

Capricornis sumatraensis thar

The Himalayan serow (*Capricornis sumatraensis thar*) is a subspecies of the Mainland serow. It is the official state animal of the Indian state of Mizoram.

Himalayan shrew

Himalayan shrew

Soriculus nigrescens

The Himalayan shrew (*Soriculus nigrescens*) is found in Bhutan, China, India, Myanmar, and Nepal, is currently the only species in the genus *Soriculus* within the

tribe Nectogalini, although the species *Chodsigoa* and *Episoriculus*, which occur in...

Himalayan striped squirrel
Himalayan striped squirrel
Tamias maclellandii

The Himalayan striped squirrel (*Tamias maclellandii*), also known as western striped squirrel, or Burmese striped squirrel, is a species of rodent in the family Sciuridae. It lives in a variety of forest from tropical to subtropical in Bhutan,...

Himalayan Tahr
Himalayan Tahr
Hemitragus jemlahicus

The Himalayan tahr (*Hemitragus jemlahicus*) is a large wild goat that lives on the mountain slopes and in the forests of the Himalayas. It is listed as Near Threatened on the IUCN Red List, as the population is declining because of hunting and...

Himalayan water shrew
Himalayan water shrew
Chimarrogale himalayica

The Himalayan water shrew (*Chimarrogale himalayica*) is a species of mammal in the family Soricidae. It is found in China, India, Japan, Laos, Myanmar, Taiwan, and Vietnam.

Himalayan whiskered bat
Himalayan whiskered bat
Myotis siligorensis

The Himalayan whiskered bat (*Myotis siligorensis*) is a species of vesper bat. It is found in Bangladesh, Bhutan, Cambodia, China, India, Indonesia, Laos, Malaysia, Nepal, and Vietnam.

Hinde's lesser house bat
Hinde's lesser house bat
Scotoecus hindei

Hinde's lesser house bat (*Scotoecus hindei*) is a species of vesper bat. It is found in Cameroon, Republic of the Congo, Kenya, Malawi, Mozambique, Nigeria, Somalia, South Sudan, Tanzania, and Zambia. Its natural habitat is dry savanna. Oldfield...

Hinde's rock rat
Hinde's rock rat
Aethomys hindei

Hinde's rock rat (*Aethomys hindei*) is a species of rodent in the family Muridae found in Burundi, Cameroon, the Central African Republic, the Democratic Republic of the Congo, Ethiopia, Kenya, Rwanda, Sudan, Tanzania, and Uganda. Its natural...

Hipposideros atrox
Hipposideros atrox
Hipposideros atrox

Hipposideros atrox, commonly known as the lesser bicolored leaf-nosed bat, is a species of bat found in Southeast Asia. Originally described as a subspecies in 1918, it was recognized as a full species in 2010. It uses echolocation to navigate...

Hipposideros bernardsigeei
Hipposideros bernardsigeei
Hipposideros bernardsigeei

Hipposideros bernardsigei is a hipposiderid species of bat known by fossil specimens, one of the many new taxa of microchiropterans discovered in the Riversleigh World Heritage Area.

Hipposideros besaoka
Hipposideros besaoka
Hipposideros besaoka

Hipposideros besaoka is an extinct bat from Madagascar in the genus *Hipposideros*. It is known from numerous jaws and teeth, which were collected in a cave at Anjohibe in 1996 and described as a new species in 2007. The site where *H. besaoka* was...

Hipposideros einnaythu
Hipposideros einnaythu
Hipposideros einnaythu

Hipposideros einnaythu, the house-dwelling leaf-nosed bat, is a species of bat found in Myanmar.

Hipposideros felix
Hipposideros felix
Hipposideros felix

Hipposideros felix is a species of bat known from Miocene fossil deposits at Li Mae Long in Thailand. The holotype is a tooth, the third molar, of a hipposiderid bat with affinities to the *Brachipposideros* group of fossil species found in...

Hipposideros rotalis
Hipposideros rotalis
Hipposideros rotalis

Hipposideros rotalis, the Laotian leaf-nosed bat or Laotian roundleaf bat, is a species of bat from the family Rhinolophidae which is thought to be endemic to Laos. Only a small number of individuals have been recorded, however the population...

Hipposideros winsburyorum
Hipposideros winsburyorum
Hipposideros winsburyorum

Hipposideros winsburyorum is a hipposiderid species of bat known by fossil specimens, one of the many new taxa of chiropterans discovered in the Riversleigh World Heritage Area. The species existed during the Pliocene.

Hirola
Hirola
Beatragus hunteri

The *hirola* is a critically endangered antelope that is found in a small area in Africa. It is medium-sized, tan to rufous-tawny in color with slightly lighter underparts, predominantly white inner ears, and a white tail that extends down to the...

Hispaniolan greater funnel-eared bat
Hispaniolan greater funnel-eared bat
Natalus major

The Hispaniolan greater funnel-eared bat (*Natalus major*) is a species of funnel-eared bat endemic to the island of Hispaniola (Dominican Republic and Haiti) in the Caribbean. First described in 1902, it has a complex taxonomic history, with some...

Hispaniolan hutia
Hispaniolan hutia
Plagiodontia aedium

The Hispaniolan hutia (*Plagiodontia aedium*) (Spanish: jutía) is a hutia species endemic to the island of Hispaniola (split between Haiti and the Dominican Republic). It is the only confirmed extant species of the genus *Plagiodontia*, and the...

Hispaniolan Solenodon
Hispaniolan Solenodon
Solenodon paradoxus

The Hispaniolan solenodon (*Solenodon paradoxus*), is a solenodon endemic to the Caribbean island of Hispaniola (shared by the Dominican Republic and Haiti). It was first described by Johann Friedrich von Brandt in 1833. A similar but smaller...

Hispid Cotton Rat
Hispid Cotton Rat
Sigmodon hispidus

The hispid cotton rat (*Sigmodon hispidus*) is a rodent species long thought to occur in parts of South America, Central America, and southern North America. However, recent taxonomic revisions, based on mitochondrial DNA sequence data, have split...

Hispid Hare
Hispid Hare
Caprolagus hispidus

The hispid hare (*Caprolagus hispidus*), also called Assam rabbit and bristly rabbit, is a leporid native to South Asia, whose historic range extended along the southern foothills of the Himalayas. Today, its habitat is highly fragmented with an...

Hispid hocicudo
Hispid hocicudo
Oxymycterus hispidus

The hispid hocicudo (*Oxymycterus hispidus*) is a rodent species from South America. It is found in Argentina and Brazil, in the Atlantic Forest.

Hispid pocket gopher
Hispid pocket gopher
Orthogeomys hispidus

The hispid pocket gopher (*Heterogeomys hispidus*) is a species of rodent in the family Geomyidae. It is found in Mexico, Belize, Guatemala, and Honduras. Some authors classify it in the genus *Orthogeomys*, but recent research has allowed this and...

Hispid pocket mouse
Hispid pocket mouse
Chaetodipus hispidus

The hispid pocket mouse (*Chaetodipus hispidus*) is a large pocket mouse native to the Great Plains region of North America. It is a member of the genus *Chaetodipus*.

Hoary bamboo rat
Hoary bamboo rat
Rhizomys pruinosus

The hoary bamboo rat (*Rhizomys pruinosus*) is a species of rodent in the family Spalacidae found in Southeast Asia (Cambodia, Laos, Malaysia, Myanmar, Thailand, and Vietnam), East Asia (China) and South Asia (India).

Hoary Bat
Hoary Bat
Lasiurus cinereus

The hoary bat (*Aeorestes cinereus*) is a species of bat in the vesper bat family, *Vespertilionidae*. It lives throughout most of North America (possibly including Hawaii, although this is disputed).

Hoary Fox

Hoary Fox

Lycalopex vetulus

The hoary fox or hoary zorro (*Lycalopex vetulus*), also known as raposinha-do-campo in Brazil (Portuguese for "meadow little fox"), is a species of zorro or "false" fox endemic to Brazil. Unlike many other foxes, it feeds primarily on small...

Hoary Marmot

Hoary Marmot

Marmota caligata

The Hoary marmot (*Marmota caligata*) is a species of marmot that lives the mountains of northwest North America. It is the largest North American ground squirrel and is often nicknamed "the whistler" for its high-pitched warning issued to alert...

Hoary wattled bat

Hoary wattled bat

Chalinolobus nigrogriseus

The hoary wattled bat (*Chalinolobus nigrogriseus*) is a species of vesper bat found in northern Australia and Papua New Guinea. Two subspecies are currently recognised:*C. n. nigrogriseus* (Gould, 1852)*C. n. rogersi* (Thomas, 1909)

Hodgson's bat

Hodgson's bat

Myotis formosus

Hodgson's bat (*Myotis formosus*), also called the copper-winged bat or black-and-orange myotis, is a species of vesper bat in the genus *Myotis*, the mouse-eared bats. Favouring mountain forests, it is found throughout Central, Southeast, and East...

Hodgson's brown-toothed shrew

Hodgson's brown-toothed shrew

Episoriculus caudatus

Hodgson's brown-toothed shrew (*Episoriculus caudatus*) is a species of mammal in the family *Soricidae*. It is found in China, India, and Myanmar.

Hodgson's giant flying squirrel

Hodgson's giant flying squirrel

Petaurista magnificus

Hodgson's giant flying squirrel (*Petaurista magnificus*) is a species of rodent in the family *Sciuridae*. This large flying squirrel lives in Himalayan forests in Asia. Like other flying squirrels, it is nocturnal and able to glide (not actually...

Hoffmann's pika

Hoffmann's pika

Ochotona hoffmanni

Hoffmann's pika (*Ochotona hoffmanni*) is a species of mammal in the pika family, *Ochotonidae*, that is endemic to Mongolia. It is currently listed as endangered by the International Union for Conservation of Nature (IUCN).

Hoffmann's rat

Hoffmann's rat

Rattus hoffmanni

Hoffmann's rat (*Rattus hoffmanni*) is a species of rodent in the family *Muridae*. It is found only in Indonesia, and is located throughout Sulawesi, as well as Malenge

island in the Togian Islands.
Hoffmann's Two-Toed Sloth
Hoffmann's Two-Toed Sloth
Choloepus hoffmanni

Hoffmann's two-toed sloth (*Choloepus hoffmanni*), also known as the northern two-toed sloth is a species of sloth from Central and South America. It is a solitary, largely nocturnal and arboreal animal, found in mature and secondary rainforests...
Hoffmanns's titi
Hoffmanns's titi
Plecturocebus hoffmannsi

Hoffmanns's titi (*Plecturocebus hoffmannsi*) is a species of titi, a type of New World monkey, endemic to Brazil. It was described as *Callicebus hoffmannsi* in 1908.
Hog Badger
Hog Badger
Arctonyx collaris

Hog badgers are three species of mustelid in the genus *Arctonyx*. They represent one of the two genera in the subfamily Melinae, alongside the true badgers (genus *Meles*).
Hog Deer
Hog Deer
Hyelaphus porcinus

The Indian hog deer (*Axis porcinus*) is a small deer native to the Indo-Gangetic Plain in Pakistan, northern India, Nepal, Bangladesh to mainland Southeast Asia. It also occurs in western Thailand, and is possibly extirpated from China (in...
Hokkaido red-backed vole
Hokkaido red-backed vole
Myodes rex

The Hokkaido red-backed vole (*Myodes rex*) is a species of rodent in the family Cricetidae. It is found at high altitudes on the island of Hokkaido in Japan and at lower altitudes on some smaller islands nearby. Its natural habitat is temperate...
Hokkaido wolf
Hokkaido wolf
Canis lupus hattai

The Hokkaido wolf (*Canis lupus hattai*), also known as the Ezo wolf (Japanese: 江戸オオカミ (蝦夷狼) 一, Hepburn: Ezo Ōkami) and in Russia as the Sakhalin wolf, is an extinct subspecies of gray wolf that once inhabited coastal north-east Asia. Its nearest...
Holland's sportive lemur
Holland's sportive lemur
Lepilemur hollandorum

Holland's sportive lemur (*Lepilemur hollandorum*), or the Mananara-Nord sportive lemur, is a sportive lemur that is endemic to Madagascar. It is one of 26 species in the genus *Lepilemur*. This lemur is found specifically in the Mananara-Nord...
Holochilus brasiliensis
Holochilus brasiliensis
Holochilus brasiliensis

Holochilus brasiliensis, also known as the Brazilian marsh rat or web-footed marsh rat, is a species of semiaquatic rodent from South America. It is found in northeastern Argentina, southern and eastern Brazil and in eastern Uruguay.
Holochilus chacarius
Holochilus chacarius

Holochilus chacarius

Holochilus chacarius, also known as the Chacoan marsh rat or Chaco marsh rat, is a semiaquatic rodent species in the family Cricetidae. It is found in Argentina and Paraguay within the Gran Chaco.

Hon Khoai squirrel

Hon Khoai squirrel

Callosciurus honkhoaiensis

The Hon Khoai squirrel (*Callosciurus honkhoaiensis*) is a species of squirrel in the family Sciuridae endemic to the small island group of Hon Khoai, which is located off the Indochinese Peninsula off the coast of south Vietnam. It can be...

Honduran fruit-eating bat

Honduran fruit-eating bat

Artibeus inopinatus

The Honduran fruit-eating bat (*Artibeus inopinatus*) is a species of bat in the family Phyllostomidae. It is found in El Salvador, Honduras, and Nicaragua.

Honduran small-eared shrew

Honduran small-eared shrew

Cryptotis hondurensis

The Honduran small-eared shrew (*Cryptotis hondurensis*) is a species of mammal in the family Soricidae. It is found in Honduras and possibly in El Salvador, Guatemala, and Nicaragua.

Honduran White Bat

Honduran White Bat

Ectophylla alba

The Honduran white bat (*Ectophylla alba*) is a unique species of bat, the only member of the genus *Ectophylla*. The genus and the species were both scientifically described for the first time in 1892. It constructs "tents" out of understory plant...

Honey Badger

Honey Badger

Mellivora capensis

The Honey badger (*Mellivora capensis*) is a mammal widely distributed in Africa, Southwest Asia, and the Indian subcontinent. Despite its name, the Honey badger does not closely resemble other badger species; instead, it bears more anatomical...

Honey Possum

Honey Possum

Tarsipes rostratus

The honey possum or noolbenger (*Tarsipes rostratus*), is a tiny species of marsupial that feeds on the nectar and pollen of a diverse range of flowering plants. Found in southwest Australia, it is an important pollinator for such plants as...

Hooded red-sided opossum

Hooded red-sided opossum

Monodelphis palliolata

The hooded red-sided opossum (*Monodelphis palliolata*) is a South American opossum species of the family Didelphidae. Until recently, it was viewed as a subspecies of *M. brevicaudata*. It is found in Colombia and Venezuela at altitudes from sea...

Hooded Seal

Hooded Seal

Cystophora cristata

The Hooded seal (*Cystophora cristata*) is a large phocid found only in the central and western North Atlantic. Hooded seal pups are known as "blue-backs" because their coats are blue-grey on the back with whitish bellies, though this coat is shed...

Hooded Skunk

Hooded Skunk

Mephitis macroura

The hooded skunk (*Mephitis macroura*) is a species of mammal in the family Mephitidae. *Mephitis* in Latin means "foul odor", *μακρός* (*makrós*) in Greek translates to "long" and *οὐρά* (*ourá*) translates to "tail".

Hooded spider monkey

Hooded spider monkey

Ateles geoffroyi grisescens

The hooded spider monkey (*Ateles geoffroyi grisescens*), is a subspecies of Geoffroy's spider monkey, a type of New World monkey, from Central America, native to Panama. It also might be found in a small portion of Colombia adjacent to Panama....

Hoogerwerf's rat

Hoogerwerf's rat

Rattus hoogerwerfi

Hoogerwerf's rat (*Rattus hoogerwerfi*) is a species of rodent in the family Muridae. It is named after zoologist Andries Hoogerwerf and is found only in western Sumatra, Indonesia, including Mount Leuser, where it is found only above 2000 m. It is...

Hoogstraal's gerbil

Hoogstraal's gerbil

Gerbillus hoogstraali

Hoogstraal's gerbil (*Gerbillus hoogstraali*) is distributed mainly in southwestern Morocco. There are thought to be less than 250 individuals in existence. The specific epithet honors American entomologist and parasitologist Harry Hoogstraal.

Hoogstraal's striped grass mouse

Hoogstraal's striped grass mouse

Lemniscomys hoogstraali

Hoogstraal's lemniscomys or Hoogstraal's striped grass mouse (*Lemniscomys hoogstraali*) is a species of rodent in the family Muridae. It is known only from the type specimen collected in 1961 at Paloich, north of Niayok, South Sudan. Its...

Hooper's mouse

Hooper's mouse

Peromyscus hooperi

Hooper's mouse (*Peromyscus hooperi*) is a species of rodent in the family Cricetidae. It is the only member of the *Peromyscus hooperi* species group, and is found only in Mexico. The species is named for Emmett Hooper, a researcher into the...

Hopi Chipmunk

Hopi Chipmunk

Neotamias rufus

The Hopi chipmunk, *Neotamias rufus*, is a small chipmunk found in Colorado, Utah and Arizona in the southwestern United States. It was previously grouped with the Colorado chipmunk, *T. quadrivittatus*. This species is listed as "Least Concern" on...

Hopkins's groove-toothed swamp rat

Hopkins's groove-toothed swamp rat
Pelomys hopkinsi

Hopkins's groove-toothed swamp rat (*Pelomys hopkinsi* Hayman, 1955) is a species of rodent in the family Muridae. It is found in Kenya, Rwanda, Uganda, possibly Burundi, and possibly Tanzania. Its natural habitat is swamps. It is threatened by...

Horn-skinned bat
Horn-skinned bat
Eptesicus floweri

The horn-skinned bat (*Eptesicus floweri*) is a species of vesper bat. It can be found in Mali and Sudan. It is found in subtropical or tropical dry shrubland, subtropical or tropical dry lowland grassland, and hot deserts.

Horse
Horse
Equus ferus caballus

The horse (*Equus ferus caballus*) is a domesticated, odd-toed, hoofed mammal. It belongs to the taxonomic family Equidae and is one of two extant subspecies of *Equus ferus*. The horse has evolved over the past 45 to 55 million years from a small...

Horse-tailed squirrel
Horse-tailed squirrel
Sundasciurus hippurus

The horse-tailed squirrel (*Sundasciurus hippurus*) is a species of rodent in the family Sciuridae. It is found throughout the islands of Borneo and Sumatra, as well as the southern half of the Malay Peninsula, which includes the entire nations of...

Horsfield's tarsier
Horsfield's tarsier
Cephalopachus bancanus

Horsfield's tarsier (*Cephalopachus bancanus*), also known as the western tarsier, is the only species of tarsier in the genus *Cephalopachus*. It occurs on Borneo, Sumatra and nearby islands and is, like other members of the group, entirely...

Horsfield's bat
Horsfield's bat
Myotis horsfieldii

Horsfield's bat (*Myotis horsfieldii*) is a species of vesper bat. It is found in China, India, Indonesia, Malaysia, Myanmar, the Philippines, Thailand, and Vietnam.

Horsfield's fruit bat
Horsfield's fruit bat
Cynopterus horsfieldii

Horsfield's fruit bat (*Cynopterus horsfieldii*) is a species of megabat native to South East Asia. It is named for Thomas Horsfield, an American naturalist who presented the type specimen to the British Museum.

Horsfield's shrew
Horsfield's shrew
Crocidura horsfieldii

Horsfield's shrew (*Crocidura horsfieldii*) is a species of mammal in the family Soricidae found in Cambodia, China, India, Japan, Laos, Nepal, Bhutan, Sri Lanka, Taiwan, Thailand, and Vietnam.

Horsfield's treeshrew
Horsfield's treeshrew
Tupaia javanica

Horsfield's treeshrew (*Tupaia javanica*), also called Javan treeshrew, is a treeshrew species within the Tupaiidae. It is endemic to the Indonesian islands of Sumatra, Bali, Java and Nias where it inhabits foremost primary forest. It was first...

Hose's Langur

Hose's Langur

Presbytis hosei

Hose's langur (*Presbytis hosei*) is a species of primate from Southeast Asia. It was first identified in Kutai National Park and Sangkulirang Peninsula, East Kalimantan, Indonesia, in 1985.

Hose's mongoose

Hose's mongoose

Herpestes brachyurus hosei

Hose's mongoose (*Herpestes brachyurus hosei*) is a subspecies of the short-tailed mongoose, but it is sometimes considered a separate species instead, *Herpestes hosei*. It is only known from a single specimen, an adult female taken in the Baram...

Hose's Palm Civet

Hose's Palm Civet

Diplogale hosei

Hose's palm civet (*Diplogale hosei*) is a viverrid species from the island of Borneo. It was named after the zoologist Charles Hose by Oldfield Thomas in 1892 who collected the first specimen in Sarawak in 1891. What little is known of the species...

Hose's pygmy flying squirrel

Hose's pygmy flying squirrel

Petaurillus hosei

The Hose's pygmy flying squirrel (*Petaurillus hosei*) is a species of rodent in the family Sciuridae. It was named for zoologist Charles Hose. It is endemic to Malaysia.

Hotson's jerboa

Hotson's jerboa

Allactaga hotsoni

Hotson's jerboa or the Iranian jerboa (*Allactaga hotsoni*), is a species of rodent in the family Dipodidae. It is found in Afghanistan, Iran, and Pakistan.

Hotson's mouse-like hamster

Hotson's mouse-like hamster

Calomyscus hotsoni

Hotson's mouse-like hamster (*Calomyscus hotsoni*) also known as Hotson's *calomyscus* or Hotson's brush-tailed mouse is a species of rodent in the family Calomyscidae. It is endemic to southwestern Pakistan and southeastern Iran (Musser and...

Hottentot golden mole

Hottentot golden mole

Amblysomus hottentotus

The Hottentot golden mole (*Amblysomus hottentotus*) is a species of mammal in the golden mole family, Chrysochloridae. It is found in South Africa, Eswatini, and possibly Lesotho. Its natural habitats are temperate forests, subtropical or...

Hourglass Dolphin

Hourglass Dolphin

Lagenorhynchus cruciger

The hourglass dolphin (*Lagenorhynchus cruciger*) is a small dolphin in the family Delphinidae that inhabits offshore Antarctic and sub-Antarctic waters. It is commonly seen from ships crossing the Drake Passage, but has a circumpolar...

House Mouse

House Mouse

Mus musculus

One of the most abundant and widespread mammals around the globe, the House mouse (*Mus musculus*) is found in almost any part of the world (except Antarctica) due to human travel. This rodent originated in Iran, Afghanistan, Pakistan, and northern...

Howell's forest shrew

Howell's forest shrew

Sylvisorex howelli

Howell's forest shrew (*Sylvisorex howelli*) is a species of mammal in the family Soricidae. It is endemic to Tanzania. Its natural habitat is subtropical or tropical moist montane forests.

Huanchaca mouse

Huanchaca mouse

Juscelinomys huanchacae

The Huanchaca mouse or Huanchaca akodont (*Juscelinomys huanchacae*) is a rodent species in the family Cricetidae. It is known from savannas in an area at an elevation of 700 metres (2,300 ft) in Serrania Huanchaca, Noel Kempff Mercado National...

Hubbard's sportive lemur

Hubbard's sportive lemur

Lepilemur hubbardorum

Hubbard's sportive lemur (*Lepilemur hubbardorum*), or the Zombitse sportive lemur, is a sportive lemur endemic to Madagascar. It has total length of about 51 to 59 cm (20 to 23 in), of which 23–25 cm (9.1–9.8 in) are tail. Hubbard's sportive...

Hubbs' beaked whale

Hubbs' beaked whale

Mesoplodon carlhubbsi

Hubbs' beaked whale (*Mesoplodon carlhubbsi*) was initially thought to be an Andrews' beaked whale when discovered by ichthyologist Carl Hubbs; however, it was named in his honor when it was discovered to be a new species. This species has the...

Hubert's multimammate mouse

Hubert's multimammate mouse

Mastomys huberti

Hubert's multimammate mouse, or Hubert's mastomys (*Mastomys huberti*) is a species of rodent in the family Muridae found in Burkina Faso, Gambia, Mali, Mauritania, Nigeria, and Senegal, and possibly Guinea, Guinea-Bissau, and Niger. Its natural...

Hudson Bay wolf

Hudson Bay wolf

Canis lupus hudsonicus

The Hudson Bay wolf (*Canis lupus hudsonicus*) is a subspecies of gray wolf native to northern Keewatin, including the northwestern coast of Hudson Bay in Canada. It was first classed as a distinct subspecies in 1941 by Edward Goldman, who...

Hudson's Oldfield mouse

Hudson's Oldfield mouse

Thomasomys hudsoni

Hudson's Oldfield mouse (*Thomasomys hudsoni*) is a species of rodent in the family Cricetidae. It is known only from the Andes in southern Ecuador, where it has been found at an elevation of 3100 m. It was formerly considered a subspecies of T....

Huemul

Huemul

Hippocamelus bisulcus

The south Andean deer (*Hippocamelus bisulcus*) is an endangered species of deer native to the mountains of Argentina and Chile. Along with the northern guanaco or taruca, it is one of the two mid-sized deer in the *Hippocamelus* genus and ranges...

Hugh's hedgehog

Hugh's hedgehog

Mesechinus hughii

Hugh's hedgehog (*Mesechinus hughii*), also sometimes referred to as the central Chinese hedgehog, is native to central China and Manchuria. It prefers open areas of dry steppe, but can be found in shrubland and forests. It is known to look for...

Humboldt big-eared brown bat

Humboldt big-eared brown bat

Histiotus humboldti

The Humboldt big-eared brown bat (*Histiotus humboldti*) is a species of vesper bat in the family Vespertilionidae. It is found in Colombia and Venezuela. *Histiotus humboldti* is distributed in the subtropical forests of the eastern foothills of the...

Humboldt marten

Humboldt marten

Martes americana humboldtensis

The Humboldt marten (*Martes caurina humboldtensis*) is an endangered, genetically distinct subspecies of the Pacific marten known from the old-growth coastal redwood forests, forests with dense shrub cover, areas with serpentine soils, and...

Humboldt's Flying Squirrel

Humboldt's Flying Squirrel

Glaucomys oregonensis

Humboldt's flying squirrel (*Glaucomys oregonensis*) is the only flying squirrel found in North America. It was named after the naturalist Alexander von Humboldt and California's Humboldt County, which is one of the areas inhabited by the squirrel.

Humboldt's Hog-Nosed Skunk

Humboldt's Hog-Nosed Skunk

Conepatus humboldtii

Humboldt's hog-nosed skunk (*Conepatus humboldtii*) is a type of hog-nosed skunk native to the open grassy areas in South America. Due to their secretive lifestyle and almost nocturnal habits, hog-nosed skunks are usually much less frequently seen...

Humboldt's squirrel monkey

Humboldt's squirrel monkey

Saimiri cassiquiarensis

Humboldt's squirrel monkey (*Saimiri cassiquiarensis*) is a species of squirrel monkey from Brazil, Colombia, Ecuador, Peru and Venezuela. It had previously been considered a subspecies of the common squirrel monkey, *S. sciureus*, but was elevated...

Hummelinck's vesper mouse

Hummelinck's vesper mouse

Calomys hummelincki

Hummelinck's vesper mouse (*Calomys hummelincki*) is a species of rodent in the family Cricetidae. It is found in Aruba, Brazil, Colombia, the Netherlands Antilles, and Venezuela.

Humpback Whale

Humpback Whale

Megaptera novaeangliae

The Humpback whale (*Megaptera novaeangliae*) is a species of baleen whale. It is a rorqual; a member of the family Balaenopteridae. The Humpback whale is known for breaching and other distinctive surface behaviors, making it popular with whale...

Hun shrew

Hun shrew

Crocidura attila

The Hun shrew (*Crocidura attila*) is a species of mammal in the family Soricidae. It is found in Cameroon, Central African Republic, Mali, Democratic Republic of the Congo, and Nigeria. Its natural habitats are subtropical or tropical moist...

Husson's yellow bat

Husson's yellow bat

Rhogeessa hussoni

Husson's yellow bat (*Rhogeessa (Rhogeessa) hussoni*) is a species of vesper bat found in Suriname and southern Brazil.

Hutan shrew

Hutan shrew

Crocidura hutanis

The Hutan shrew (*Crocidura hutanis*) is a species of mammal in the family Soricidae. It is known only from northern Sumatra in Indonesia.

Hutterer's brush-furred mouse

Hutterer's brush-furred mouse

Lophuromys huttereri

Hutterer's brush-furred mouse or Hutterer's brush-furred rat (*Lophuromys huttereri*) is a species of rodent in the family Muridae. It is found only in Democratic Republic of the Congo, where its natural habitat is subtropical or tropical moist...

Hutton's tube-nosed bat

Hutton's tube-nosed bat

Murina huttoni

Hutton's tube-nosed bat (*Murina huttoni*) is a species of vesper bat in the family Vespertilionidae. It can be found in the following countries: Bhutan, China, India, Laos, Malaysia, Myanmar, Nepal, Pakistan, Thailand, and Viet Nam. It lives...

Hydrochoerus gaylordi

Hydrochoerus gaylordi

Hydrochoerus gaylordi

Hydrochoerus gaylordi is an extinct species of capybara that lived in Grenada. This species was found in 1991 by Ronald Singer and his colleagues. The species was named after Joseph Gaylord as an acknowledgment of Singer.

Hylaeamys acritus

Hylaeamys acritus

Hylaeamys acritus

Hylaeamys acritus, formerly *Oryzomys acritus*, is an oryzomyine rodent of the family Cricetidae. The name is derived from the Greek word ἀκρίτος 'confused, doubtful',

because it could easily be confused with species such as *H. megacephalus* and...

Hylaeamys laticeps

Hylaeamys laticeps

Hylaeamys laticeps

Hylaeamys laticeps, also known as the Atlantic Forest oryzomys or the large-headed rice rat, is a species of rodent in the family Cricetidae. The rodent is endemic to the Atlantic Forest region of southeastern Brazil. The taxonomic history of...

Hylaeamys megacephalus

Hylaeamys megacephalus

Hylaeamys megacephalus

Hylaeamys megacephalus, also known as Azara's broad-headed oryzomys or the large-headed rice rat, is a species of rodent in the genus *Hylaeamys* of family Cricetidae, of which it is the type species. It is found mainly in lowland tropical...

Hylaeamys oniscus

Hylaeamys oniscus

Hylaeamys oniscus

Hylaeamys oniscus, the sowbug rice rat, is a rodent species in the family Cricetidae. It was formerly placed in the genus *Oryzomys* as *Oryzomys oniscus*. It occurs only in northeastern Brazil. It is close to *Hylaeamys laticeps*, which occurs further...

Hylaeamys perenensis

Hylaeamys perenensis

Hylaeamys perenensis

Hylaeamys perenensis, formerly *Oryzomys perenensis*, also known as the western Amazonian oryzomys, is an oryzomyine rodent of the family Cricetidae. It is found in western Amazonia (southeastern Colombia, eastern Ecuador, eastern Peru, northern...

Hylaeamys tatei

Hylaeamys tatei

Hylaeamys tatei

Hylaeamys tatei, also known as Tate's oryzomys or Tate's rice rat, is a South American rodent species of the family Cricetidae. It is known only from the eastern foothills of the Andes in central Ecuador, where it has been found at elevations...

Hylaeamys yunganus

Hylaeamys yunganus

Hylaeamys yunganus

Hylaeamys yunganus, also known as the Amazonian oryzomys or Yungas rice rat, is a species of rodent in the genus *Hylaeamys* of family Cricetidae. It is found in lowland tropical rainforest throughout Amazonia, in northeastern Bolivia, eastern...

Hylomyscus endorobae

Hylomyscus endorobae

Hylomyscus endorobae

Hylomyscus endorobae is a species of rodent of the genus *Hylomyscus* that is found only in select portions of the wet East African montane forests of the Kenyan Rift mountains of southwestern Kenya and Tanzania, and only at elevations above 2,000...

Hylomyscus vulcanorum

Hylomyscus vulcanorum

Hylomyscus vulcanorum

Hylomyscus vulcanorum is a species of rodent in the family Muridae.

Hyorhinomys stuempkei

Hyorhinomys stuempkei
Hyorhinomys stuempkei

The hog-nosed shrew rat or Sulawesi snouter (*Hyorhinomys stuempkei*) is a species of rodent in the family Muridae, more specifically in the subfamily Murinae, endemic to Sulawesi, Indonesia. This species was discovered in 2015 by Jacob A....

Hypogeomys australis
Hypogeomys australis
Hypogeomys australis

Hypogeomys australis is an extinct rodent from central and southeastern Madagascar. First described in 1903, it is larger than its close relative, the living *Hypogeomys antimena*, which occurs further west, but otherwise similar. Average length of...

Hypsiprymnodon karenblackae
Hypsiprymnodon karenblackae
Hypsiprymnodon karenblackae

Hypsiprymnodon karenblackae is a fossil species describing a small marsupial extant in Australia during the Early to Middle Miocene Epoch. The material was collected at the Australian Fossil Mammal Sites (Riversleigh). The taxon was published in...

Hypsugo anthonyi
Hypsugo anthonyi
Hypsugo anthonyi

Joffre's bat (*Mirostrellus joffrei*), also known as Joffre's pipistrelle, is a species of vesper bat in the family Vespertilionidae. It is the only member of the genus *Mirostrellus*. It is found in South and Southeast Asia. It is threatened by...

Iberian Lynx
Iberian Lynx
Lynx pardinus

The Iberian lynx (*Lynx pardinus*) is the rarest of the lynx species. It is the most threatened cat species, currently on the verge of extinction. By the turn of the 21st century, the Iberian lynx was on the verge of extinction but due to...

Iberian shrew
Iberian shrew
Sorex granarius

The Iberian shrew or Lagranja shrew (*Sorex granarius*) is a species of mammal in the family Soricidae. It is found in Portugal and Spain.

Iberian Wolf
Iberian Wolf
Canis lupus signatus

The Iberian wolf (*Canis lupus signatus*, or *Canis lupus lupus*, Spanish and Portuguese : Lobo ibérico) is a subspecies of Grey wolf that has been isolated from mixing with other wolf populations for over a century. The Iberian wolf forms the...

Idaho Ground Squirrel
Idaho Ground Squirrel
Urocitellus brunneus

The northern Idaho ground squirrel (*Urocitellus brunneus*) is a species of the largest genus of ground squirrels. This species and the Southern Idaho ground squirrel were previously considered conspecific, together called the Idaho ground squirrel.

Idaho pocket gopher
Idaho pocket gopher

Thomomys idahoensis

The Idaho pocket gopher (*Thomomys idahoensis*) is a species of rodent in the family Geomyidae. It is endemic to the Northwestern United States; found in Idaho, Montana, Utah and Wyoming. Its natural habitats are temperate forests, dry savanna, and...

Ihering's Atlantic spiny rat

Ihering's Atlantic spiny rat

Trinomys iheringi

Ihering's Atlantic spiny-rat or Ihering's spiny rat (*Trinomys iheringi*) is a spiny rat species from South America. It is endemic to Brazil, and was named in honor of Hermann von Ihering.

Ihering's three-striped opossum

Ihering's three-striped opossum

Monodelphis iheringi

Ihering's three-striped opossum (*Monodelphis iheringi*), is an opossum species found in Argentina and Brazil.

Ikonnikov's bat

Ikonnikov's bat

Myotis ikonnikovi

Ikonnikov's bat (*Myotis ikonnikovi*) is a species of vesper bat. An adult Ikonnikov's bat has a body length of 4.2-5.1 cm, a tail of 3.1-4.0 cm, and a wing length of 3.3-3.6 cm. It is found in eastern Siberia, the Ussuri region, Sakhalin,...

Ili Pika

Ili Pika

Ochotona iliensis

The Ili pika (*Ochotona iliensis*) is a species of mammal in the family Ochotonidae, endemic to northwest China. After its discovery in 1983, it was not documented again until 2002. The Ili pika population is declining and is currently considered...

Ilin Island cloudrunner

Ilin Island cloudrunner

Crateromys paulus

The Ilin Island cloudrunner (*Crateromys paulus*) is a critically endangered cloud rat from Ilin Island in the Philippines. It is called siyang by the Taubuwid Mangyan. It is a fluffy-coated, bushy-tailed rat and may have emerged from tree hollows...

Imaizumi's horseshoe bat

Imaizumi's horseshoe bat

Rhinolophus imaizumii

Imaizumi's horseshoe bat (*Rhinolophus imaizumii*) is a species of bat in the family Rhinolophidae. It is endemic to Japan. Its natural habitat is temperate forests. It is threatened by habitat loss. While it was formerly considered endangered by...

Imaizumi's red-backed vole

Imaizumi's red-backed vole

Myodes imaizumii

Imaizumi's red-backed vole (*Myodes imaizumii*) is a species of vole in the family Cricetidae. It is found in Japan and was initially designated as a subspecies of the Japanese red-backed vole, with studies of cranial and molar patterns...

Impala

Impala

Aepyceros melampus

The impala (*Aepyceros melampus*) is a medium-sized antelope that is found in Africa. It was first described to European audiences by German zoologist Hinrich Lichtenstein in 1812. Active mainly during the day, the impala may be gregarious or...

Inca Oldfield mouse
Inca Oldfield mouse
Thomasomys incanus

The Inca Oldfield mouse (*Thomasomys incanus*) is a species of rodent in the family Cricetidae. It is found only in Peru.

Incan caenolestid
Incan caenolestid
Lestoros inca

The Incan caenolestid (*Lestoros inca*), also known as the Incan shrew opossum or Peruvian caenolestid, is a caenolestid found in the southern Peruvian Andes. It was first described by English zoologist Oldfield Thomas in 1917. The head-and-body...

Incan hocicudo
Incan hocicudo
Oxymycterus inca

The Incan hocicudo (*Oxymycterus inca*) is a species of rodent in the family Cricetidae. It is found in Bolivia and Peru.

Incan little mastiff bat
Incan little mastiff bat
Mormopterus phrudus

The Incan little mastiff bat (*Mormopterus phrudus*) is a species of bat in the family Molossidae, the free-tailed bats. It belongs to the subgenus *Micronomus*.

Indian aurochs
Indian aurochs
Bos primigenius namadicus

The Indian aurochs (*Bos primigenius namadicus*) (Sindhi: انڊين جهنگلي ڏاند) is an extinct aurochs subspecies that is considered the wild ancestor of the domestic zebu cattle, which is mainly found in the Indian subcontinent and has been...

Indian boar
Indian boar
Sus scrofa cristatus

The Indian boar (*Sus scrofa cristatus*), also known as the Andamanese pig or Moupin pig, is a subspecies of wild boar native to India, Nepal, Burma, western Thailand and Sri Lanka. The Indian boar differs from its European counterpart by its large...

Indian Brown Mongoose
Indian Brown Mongoose
Herpestes fuscus

The Indian brown mongoose or brown mongoose (*Urva fusca*) is a carnivorous mammal from South Asia and Oceania. It is terrestrial and hunts prey at night.

Indian bush rat
Indian bush rat
Golunda ellioti

The Indian bush rat (*Golunda ellioti*) is a rodent species in the family Muridae. It is the only extant member of the genus *Golunda*, and is the only extant member of the tribe Arvicanthini found outside of Africa. The species is widely distributed...

Indian Desert Jird
Indian Desert Jird

Meriones hurrianae

The Indian desert jird or Indian desert gerbil (*Meriones hurrianae*) is a species of jird found mainly in the Thar Desert in India. Jirds are closely related to gerbils.

Indian Elephant

Indian Elephant

Elephas maximus indicus

The Indian elephant (*Elephas maximus indicus*) is one of three extant recognised subspecies of the Asian elephant. Since 1986, it has been listed as Endangered on the IUCN Red List as the wild population has declined by at least 50% since the...

Indian Flying Fox

Indian Flying Fox

Pteropus giganteus

The Indian flying fox (*Pteropus medius*) is a species of flying fox native to the Indian subcontinent. It is one of the largest bats in the world. This species is often regarded as vermin due to its destructive tendencies towards fruit farms, but...

Indian gerbil

Indian gerbil

Tatera indica

The Indian gerbil (*Tatera indica*) also known as "antelope rat", is a species of rodent in the family Muridae. It is found in southern Asia from Syria to Bangladesh. It is the only species in the genus *Tatera*. Members of the genus *Gerbilliscus* have...

Indian Giant Flying Squirrel

Indian Giant Flying Squirrel

Petaurista philippensis

The Indian giant flying squirrel (*Petaurista philippensis*) is a large rodent capable of gliding flight using a skin membrane stretched between the front and hind legs. It can be found only in the Southeast and South Asia, and southern and central...

Indian Giant Squirrel

Indian Giant Squirrel

Ratufa indica

The Indian giant squirrel (*Ratufa indica*) is a large multi-coloured tree squirrel species. It is a diurnal, arboreal, and mainly herbivorous squirrel. It is the state animal of the state of Maharashtra in western India and in there in local...

Indian Gray Mongoose

Indian Gray Mongoose

Herpestes edwardsii

The Indian grey mongoose (*Urva edwardsii*) is a mongoose species native to the Indian subcontinent and West Asia. It lives in burrows, hedgerows and thickets, among groves of trees, and is very bold and inquisitive but wary, seldom venturing far...

Indian hairy-footed gerbil

Indian hairy-footed gerbil

Gerbillus gleadowi

The Indian hairy-footed gerbil (*Gerbillus gleadowi*) is a species of rodent found mainly in Pakistan and northwestern India. They inhabit dry, sandy, and rocky country with sparse vegetation. Their burrows are often closed with sand. They feed...

Indian Hare
Indian Hare
Lepus nigricollis

The Indian hare (*Lepus nigricollis*) is a common species of hare. They are herbivores and live solitarily. Hairs nest in slight depressions called forms, and their young are able to fend for themselves shortly after birth.

Indian Hedgehog
Indian Hedgehog
Paraechinus micropus

The Indian hedgehog (*Paraechinus micropus*) is a small species of hedgehog native to South Asia. It can be compared to the Long-eared hedgehog (*Hemiechinus auritus*) which has a similar lifestyle and appearance.

Indian Leopard
Indian Leopard
Panthera pardus fusca

The Indian leopard (*Panthera pardus fusca*) is a leopard subspecies widely distributed on the Indian subcontinent. The Indian leopard is one of the big cats occurring on the Indian subcontinent, along with the Asiatic lion, Bengal tiger, Snow...

Indian Long-Eared Hedgehog
Indian Long-Eared Hedgehog
Hemiechinus collaris

The Indian long-eared hedgehog (*Hemiechinus collaris*) is a small mammal found in South Asia. It lives in burrows and comes out at night to search for food.

Indian Ocean Bottlenose Dolphin
Indian Ocean Bottlenose Dolphin
Tursiops aduncus

The Indo-Pacific bottlenose dolphin (*Tursiops aduncus*) is a species of bottlenose dolphin. This dolphin grows to 2.6 m (8.5 ft) long, and weighs up to 230 kg (510 lb). It lives in the waters around India, northern Australia, South China, the Red...

Indian Ocean humpback dolphin
Indian Ocean humpback dolphin
Sousa plumbea

The Indian Ocean humpback dolphin (*Sousa plumbea*) is a member of the Delphinidae family occupying coastal areas ranging from Southern Africa to Western Indochina. The Indo-Pacific humpback dolphin (*Sousa chinensis*) was formerly included within...

Indian Palm Squirrel
Indian Palm Squirrel
Funambulus palmarum

The Indian palm squirrel (*Funambulus palmarum*) is a species of rodent from India and Sri Lanka. In the late 19th century, this species was introduced to Madagascar, Réunion, Mayotte, Comoro Islands, Mauritius, Seychelles and Australia, where it...

Indian Pangolin
Indian Pangolin
Manis crassicaudata

The Indian pangolin (*Manis crassicaudata*) is a shy slow-moving mammal that feeds on ants and termites, digging them out of mounds and logs using its long claws, which are as long as its forelimbs. It is nocturnal and rests in deep burrows during...

Indian pipistrelle

Indian pipistrelle
Pipistrellus coromandra

The Indian pipistrelle (*Pipistrellus coromandra*) is a species of bat in the family Vespertilionidae found in Afghanistan, Bangladesh, Bhutan, Cambodia, India, Myanmar, Nepal, Pakistan, Sri Lanka, Thailand, and Vietnam.

Indian Porcupine
Indian Porcupine
Hystrix indica

The Indian crested porcupine (*Hystrix indica*) is a hystricomorph rodent species native to southern Asia and the Middle East. It is listed as Least Concern on the IUCN Red List. It belongs to the Old World porcupine family, Hystricidae.

Indian Rhinoceros
Indian Rhinoceros
Rhinoceros unicornis

The Indian rhinoceros (*Rhinoceros unicornis*) is the biggest of the three rhinos in Asia, and, along with the African white rhino, is the biggest of all rhino species. Its single black horn identifies it, along with its gray-brown hide and skin...

Indian roundleaf bat
Indian roundleaf bat
Hipposideros lankadiva

The Indian roundleaf bat (*Hipposideros lankadiva*), also known as the large Ceylon leaf-nosed bat or Kelaart's leaf-nosed bat is a species of bat in the family Hipposideridae. It is endemic to the Indian subcontinent, with marginal populations...

Indian Spotted Chevrotain
Indian Spotted Chevrotain
Moschiola indica

Indian spotted chevrotains (*Moschiola indica*) are even-toed ungulates found in South Asia. Chevrotains or mouse-deer are the smallest hoofed mammals in the world. Despite their common name of "mouse deer", they are not closely related to true deer.

Indian wild ass
Indian wild ass
Equus hemionus khar

The Indian wild ass (*Equus hemionus khar*), also called the Indian onager or, in the local Gujarati language, Ghudkhur and Khur, is a subspecies of the onager native to South Asia. It is currently listed as Near Threatened by IUCN. The previous...

Indian Wolf
Indian Wolf
Canis lupus pallipes

The Indian wolf (*Canis lupus pallipes*) is a subspecies of Grey wolf. It travels in smaller packs and is less vocal than other variants of the Grey wolf, and has a reputation for being cunning. The Indian wolf is one of the most endangered...

Indiana Bat
Indiana Bat
Myotis sodalis

The Indiana bat (*Myotis sodalis*) is a medium-sized mouse-eared bat native to North America. It lives primarily in Southern and Midwestern U.S. states and is listed as an endangered species. The Indiana bat is grey, black, or chestnut in color...

Indo-Pacific Finless Porpoise

Indo-Pacific Finless Porpoise
Neophocaena phocaenoides

The Indo-Pacific finless porpoise (*Neophocaena phocaenoides*) is an aquatic marine mammal found throughout most of the Indian Ocean, as well as the tropical and subtropical Pacific from Indonesia north to the Taiwan Strait. Finless porpoises are...

Indo-Pacific Humpback Dolphin
Indo-Pacific Humpback Dolphin
Sousa chinensis

The Indo-Pacific humpback dolphin (*Sousa chinensis*) is a species of humpback dolphin. This species is often referred to as the Chinese white dolphin in mainland China, Macao, Hong Kong, Taiwan, and Singapore as a common name. Some biologists...

Indochinese black langur
Indochinese black langur
Trachypithecus eburnus

The Indochinese black langur (*Trachypithecus eburnus*) is a poorly known lutung native to Laos and adjacent Vietnam. It was originally described as a subspecies of *T. auratus*, but was later found to be a member of the *T. francoisi* group, with some...

Indochinese flying squirrel
Indochinese flying squirrel
Hylopetes phayrei

The Indochinese flying squirrel (*Hylopetes phayrei*), also known as Phayre's flying squirrel, is a species of rodent in the family *Sciuridae*. It is found in China, Laos, Myanmar, Thailand, and Vietnam.

Indochinese hog deer
Indochinese hog deer
Hyelaphus annamiticus

The Indian hog deer (*Axis porcinus*) is a small deer native to the Indo-Gangetic Plain in Pakistan, northern India, Nepal, Bangladesh to mainland Southeast Asia. It also occurs in western Thailand, and is possibly extirpated from China (in...

Indochinese Leopard
Indochinese Leopard
Panthera pardus delacouri

The Indochinese leopard (*Panthera pardus delacouri*) is a leopard subspecies. In Indochina, leopards are rare outside protected areas and threatened by habitat loss as well as poaching for the illegal wildlife trade. The historical range of these...

Indochinese lutung
Indochinese lutung
Trachypithecus germaini

Germain's langur (*Trachypithecus germaini*) is a lutung native to Thailand, Burma, Cambodia, Laos and Vietnam. The Old World monkey was previously included in *Trachypithecus cristatus* and *Trachypithecus villosus*. Two subspecies were...

Indochinese serow
Indochinese serow
Capricornis sumatraensis maritimus

The Indochinese serow (*Capricornis sumatraensis maritimus*) is a vulnerable goat-antelope, a subspecies of the mainland serow, native to Cambodia, Laos, Myanmar, Thailand and Vietnam. In Cambodia the Indochinese serow, known as ស៊ីរ៉ែ /sat kɛh/,...

Indochinese short-tailed shrew
Indochinese short-tailed shrew
Blarinella griselda

The Indochinese short-tailed shrew (*Blarinella griselda*) is a species of mammal of the family Soricidae found in China and Vietnam. The species is a semifossorial red-toothed shrew with a stout body and short, slender tail. *Blarinella* is...

Indomalayan pencil-tailed tree mouse
Indomalayan pencil-tailed tree mouse
Chiropodomys gliroides

The Indomalayan pencil-tailed tree mouse or simply pencil-tailed tree mouse (*Chiropodomys gliroides*) is a species of arboreal rodent in the family Muridae. It is found in northeastern India, southern China (including Hainan), Myanmar, Thailand,...

Indonesian mountain weasel
Indonesian mountain weasel
Mustela lutreolina

The Indonesian mountain weasel (*Mustela lutreolina*) is a species of weasel that lives on the islands of Java and Sumatra in Indonesia at elevations over 1,000 metres (3,280 ft). They live in mountainous, tropical, and rainforest areas....

Indonesian short-nosed fruit bat
Indonesian short-nosed fruit bat
Cynopterus titthaecheilus

The Indonesian short-nosed fruit bat (*Cynopterus titthaecheilus*) is a species of megabat in the family Pteropodidae. It is endemic to Indonesia, and has three subspecies: *C. t. major* *C. t. terminus* *C. t. titthaecheilus*

Indonesian tomb bat
Indonesian tomb bat
Taphozous achates

The Indonesian tomb bat (*Taphozous achates*) is a species of sac-winged bat in the family Emballonuridae. It is found only in Indonesia.

Indri
Indri
Indri *indri*

The indri (*Indri indri*), also called the babakoto, is one of the largest living lemurs, with a head-and-body length of about 64–72 cm (25–28+1/2 in) and a weight of between 6 and 9.5 kg (13 and 21 lb). It has a black and white coat and maintains...

Indus River Dolphin
Indus River Dolphin
Platanista minor

The Indus River dolphin (*Platanista minor*) is a species of toothed whale native to Pakistan and northwestern India. This dolphin was the first discovered side-swimming cetacean. It is patchily distributed in five small, sub-populations that are...

Inland forest bat
Inland forest bat
Vespadelus baverstocki

The inland forest bat (*Vespadelus baverstocki*) is a vespertilionid bat that occurs in central and arid regions in Australia. They were first described in 1987, published in a review of poorly surveyed microbat populations. A tiny flying mammal,...

Inland hill rat
Inland hill rat
Bunomys penitus

The inland hill rat (*Bunomys penitus*) is a species of rodent in the family Muridae. It is found only in central and southern Sulawesi, Indonesia.

Inornate squirrel
Inornate squirrel
Callosciurus inornatus

The inornate squirrel (*Callosciurus inornatus*) is a species of rodent in the family Sciuridae. It is found in China, Laos, and Vietnam.

Inquisitive shrew mole
Inquisitive shrew mole
Uropsilus investigator

The inquisitive shrew mole (*Uropsilus investigator*) is a species of mammal in the family Talpidae. It is only known from Yunnan province of China, although its range is thought to extend over the border into Myanmar. A 2018 phylogenetic study...

Insular flying fox
Insular flying fox
Pteropus tonganus

The insular flying fox or Pacific flying fox (*Pteropus tonganus*) is a species of flying fox in the family Pteropodidae. It is geographically widespread, the most widespread flying fox in the Pacific: it is found in American Samoa, the Cook...

Insular horseshoe bat
Insular horseshoe bat
Rhinolophus keyensis

The insular horseshoe bat (*Rhinolophus keyensis*) is a species of bat in the family Rhinolophidae. It is endemic to Indonesia.

Insular mole
Insular mole
Mogera insularis

The insular mole (*Mogera insularis*) is a species of mammal in the family Talpidae. It is restricted to Hainan Island and Taiwan, where it is also known as the Formosan blind mole. The species was first described by Robert Swinhoe in 1863.

Insular myotis
Insular myotis
Myotis insularum

The insular myotis (*Myotis insularum*) is one of over 100 of species of vesper bat in the genus *Myotis*. It is found in possibly American Samoa and possibly Samoa.

Insular single leaf bat
Insular single leaf bat
Monophyllus plethodon

The insular single leaf bat or Lesser Antillean long-tongued bat (*Monophyllus plethodon*) is a species of bat in the family Phyllostomidae.

Insular vole
Insular vole
Microtus abbreviatus

The insular vole or St. Matthew Island vole (*Microtus abbreviatus*) is a species of rodent in the family Cricetidae. It occurs only on St. Matthew Island and the adjacent Hall Island, in Alaska. On these Bering Sea islands, insular voles live in...

Intelligent grass mouse
Intelligent grass mouse
Akodon iniscatus

The intelligent grass mouse or Patagonian akodont (*Akodon iniscatus*) is a species of rodent in the family Cricetidae. The species is found in Argentina and Chile.

Interior Alaskan wolf
Interior Alaskan wolf
Canis lupus pambasileus

The Interior Alaskan wolf (*Canis lupus pambasileus*), also known as the Yukon wolf, is a subspecies of gray wolf native to parts of British Columbia, the Northwest Territories, Interior Alaska and Yukon.

Intermediate horseshoe bat
Intermediate horseshoe bat
Rhinolophus affinis

The intermediate horseshoe bat (*Rhinolophus affinis*) is a bat species of the family Rhinolophidae ("nose crest") that is very widespread throughout much of the Indian subcontinent, southern and central China and Southeast Asia. It is listed by...

Intermediate long-fingered bat
Intermediate long-fingered bat
Miniopterus medius

The intermediate long-fingered bat (*Miniopterus medius*) is a species of vesper bat in the family Vespertilionidae. It can be found in the following countries: Native: Indonesia (Jawa, Kalimantan, Sulawesi), Malaysia and Thailand. Presence...

Intermediate roundleaf bat
Intermediate roundleaf bat
Hipposideros larvatus

The intermediate roundleaf bat (*Hipposideros larvatus*) is a species of bat in the family Hipposideridae. It is found in Bangladesh, Cambodia, China, India, Indonesia, Laos, Malaysia, Myanmar, Thailand, and Vietnam.

Intermediate slit-faced bat
Intermediate slit-faced bat
Nycteris intermedia

The intermediate slit-faced bat (*Nycteris intermedia*) is a species of slit-faced bat living in forest and savanna regions of west and central Africa. It is easily confused with *Nycteris arge* and *Nycteris nana*. *N. intermedia* is broadly...

Inyo shrew
Inyo shrew
Sorex tenellus

The Inyo shrew (*Sorex tenellus*) is a species of mammal in the family Soricidae.

Ipanema bat
Ipanema bat
Pygoderma bilabiatum

The Ipanema bat (*Pygoderma bilabiatum*) is a bat species of order Chiroptera and family Phyllostomidae. It is found in South America, specifically in northern Argentina, Bolivia, southeastern Brazil and Paraguay. It is monotypic within its genus.

Iranian jerboa
Iranian jerboa
Allactaga firouzi

The Iranian jerboa (*Allactaga firouzi*) is a rodent of the family Dipodidae and subfamily Allactaginae. They are characteristically known as hopping rodents and have only been found in a small population in Iran. Iranian jerboa are microhabitat...

Iranian shrew

Iranian shrew

Crocidura susiana

The Iranian shrew (*Crocidura susiana*) is a species of mammal in the family Soricidae. It is endemic to Iran. It is threatened by habitat loss.

Irenomys

Irenomys

Irenomys tarsalis

Irenomys tarsalis, also known as the Chilean climbing mouse, Chilean tree mouse, or long-footed irenomys, is a rodent found in Chile, from about 36° to 46°S, and in adjacent Argentina, mainly in forests. It is a large, long-tailed, soft-furred...

Iriomote Cat

Iriomote Cat

Prionailurus bengalensis iriomotensis

The Iriomote cat (*Prionailurus bengalensis iriomotensis*) is a subspecies of the Leopard cat that lives exclusively on the Japanese island of Iriomote. It is viewed by many biologists as a "living fossil," and is not much changed from its...

Irrawaddy Dolphin

Irrawaddy Dolphin

Orcaella brevirostris

The Irrawaddy dolphin (*Orcaella brevirostris*) is a euryhaline species of oceanic dolphin found in scattered subpopulations near sea coasts of the Bay of Bengal and Southeast Asia. It closely resembles the Australian snubfin dolphin and was not...

Irrawaddy squirrel

Irrawaddy squirrel

Callosciurus pygerythrus

The Irrawaddy squirrel (*Callosciurus pygerythrus*) or hoary-bellied Himalayan squirrel is a species of rodent in the family Sciuridae.

Isabel naked-tailed rat

Isabel naked-tailed rat

Solomys sapientis

The Isabel naked-tailed rat (*Solomys sapientis*) is a species of rodent in the family Muridae. It is found only on the island of Santa Isabel in the Solomon Islands.

Isabelle's ghost bat

Isabelle's ghost bat

Diclidurus isabella

Isabelle's ghost bat (*Diclidurus isabella*) is a bat species found in northwestern Brazil, Guyana, and Venezuela, and possibly Colombia. It was discovered in October 1916 by Emilie Snethlage, and described by Oldfield Thomas in 1920. While the...

Isalo serotine

Isalo serotine

Neoromicia malagasyensis

The Isalo serotine (*Laephotis malagasyensis*) is a vespertilionid bat of Madagascar in the genus *Laephotis*. It is known only from the vicinity of the Isalo National Park in the southwestern part of the island, where it has been caught in riverine...

Isarog shrew-rat
Isarog shrew-rat
Rhynchomys isarogensis

The Isarog shrew-rat or Mt Isarog shrew-rat (*Rhynchomys isarogensis*) is a species of rodent in the family Muridae. It is found only in the Philippines.

Isarog striped shrew-rat
Isarog striped shrew-rat
Chrotomys gonzalesi

The Isarog striped shrew-rat or Mount Isarog striped rat (*Chrotomys gonzalesi*) is a species of rodent in the family Muridae found only in the Philippines. Its natural habitat is subtropical or tropical dry forests.

Isla De La Juventud tree hutia
Isla De La Juventud tree hutia
Mysateles meridionalis

The Isla De La Juventud tree hutia or southern hutia (*Mysateles prehensilis meridionalis*) is a subspecies of rodent in the subfamily Capromyinae. It is endemic to lowland moist forests on Isla de la Juventud in Cuba. It is threatened by habitat...

Island Fox
Island Fox
Urocyon littoralis

The Island fox (*Urocyon littoralis*) is a small fox native to six of the eight Channel Islands of California. There are six subspecies, each unique to the island it lives on, reflecting its evolutionary history. They are generally docile, show...

Island mouse
Island mouse
Nesomys rufus

The island mouse or eastern red forest rat (*Nesomys rufus*) is a species of rodent in the family Nesomyidae. It is found only in Madagascar.

Island tube-nosed fruit bat
Island tube-nosed fruit bat
Nyctimene major

The island tube-nosed fruit bat (*Nyctimene major*) is a species of bat in the family Pteropodidae. It is found in Papua New Guinea and the Solomon islands.

Isothrix barbarabrownae
Isothrix barbarabrownae
Isothrix barbarabrownae

Isothrix barbarabrownae is a recently discovered species of rodents of the spiny rat family Echimyidae.

Issel's groove-toothed swamp rat
Issel's groove-toothed swamp rat
Pelomys isseli

Issel's groove-toothed swamp rat (*Pelomys isseli*) is a species of rodent in the family Muridae. It is found only in Uganda. Its natural habitats are arable land and rural gardens. It is threatened by habitat loss.

Isthmian mouse opossum
Isthmian mouse opossum
Marmosa isthmica

The Isthmian mouse opossum (*Marmosa isthmica*) is a species of opossum in the

family Didelphidae found in Colombia, Ecuador, and Panama. Foraging along branches and vines for fruit and insects, with the help of a prehensile tail, *M. isthmica* was...

Italian Wolf

Italian Wolf

Canis lupus italicus

The Italian wolf (*Canis lupus italicus* or *Canis lupus lupus*) is a subspecies of the Grey wolf. Although not universally recognized as a distinct subspecies, it nonetheless possesses a unique mtDNA haplotype and a distinct skull morphology.

The...

Ivory Coast rat

Ivory Coast rat

Dephomys eburneae

The Ivory Coast rat (*Dephomys eburneae*) is a species of rodent in the family Muridae. It is found in Ivory Coast, possibly Ghana, and Liberia. Its natural habitats are subtropical or tropical moist lowland forest and subtropical or tropical...

Ixtlán deer mouse

Ixtlán deer mouse

Habromys ixtlani

The Ixtlán deer mouse (*Habromys ixtlani*) is a species of rodent in the family Cricetidae. It is endemic to southwestern Mexico, and found in the Sierra de Juárez, a subrange of the Sierra Madre de Oaxaca in Oaxaca state. Although it was originally...

Ja slit-faced bat

Ja slit-faced bat

Nycteris major

The Ja slit-faced bat (*Nycteris major*) is a species of slit-faced bat that lives in the tropical and subtropical forests of Africa. Its habitat is fragmented and declining, and the species is therefore very rare. Logging and farming are the...

Jackass shrew

Jackass shrew

Crocidura arispa

The jackass shrew (*Crocidura arispa*) is a species of mammal belonging to the Soricidae family. It is endemic to Turkey. Its natural habitat is in rocky areas.

Jackson's fat mouse

Jackson's fat mouse

Steatomys jacksoni

The Jackson's fat mouse (*Steatomys jacksoni*) is a species of rodent in the family Nesomyidae. It is found in Ghana and Nigeria. Its natural habitats are arable land and...

Jackson's mongoose

Jackson's mongoose

Bdeogale jacksoni

Jackson's mongoose (*Bdeogale jacksoni*) is a mongoose species native to montane forests in Kenya, Uganda and Tanzania. It appears to be rare and has been classified as Near Threatened since 2008. Its long and dense fur is grizzled black and white....

Jackson's shrew

Jackson's shrew

Crocidura jacksoni

Jackson's shrew (*Crocidura jacksoni*) is a species of mammal in the family Soricidae. It is found in Burundi, Democratic Republic of the Congo, Kenya, Rwanda, South Sudan, and Uganda. Its natural habitats are subtropical or tropical moist lowland...

Jackson's soft-furred mouse

Jackson's soft-furred mouse

Praomys jacksoni

Jackson's soft-furred mouse or Jackson's praomys (*Praomys jacksoni*) is a species of rodent in the family Muridae. It is found in Angola, Burundi, Cameroon, Central African Republic, Republic of the Congo, Democratic Republic of the Congo,...

Jaguar

Jaguar

Panthera onca

The jaguar (*Panthera onca*) is a large cat species and the only living member of the genus *Panthera* native to the Americas. It is the largest cat species in the Americas and the third largest in the world. The jaguar has featured prominently in...

Jaguarundi

Jaguarundi

Puma yagouaroundi

The jaguarundi (*Herpailurus yagouaroundi*) is a wild cat native to the Americas. Secretive and alert, the jaguarundi hunts mainly during daytime and evening hours and feeds on various kinds of prey. It lives in a broad array of closed as well as...

Jalapan pine vole

Jalapan pine vole

Microtus quasiater

The Jalapan pine vole (*Microtus quasiater*) is a species of rodent in the family Cricetidae, found only in Mexico. The scientific name *quasiater* translates as "almost black", while the common name refers to the city of Jalapa, close to where the...

Jaliscan cotton rat

Jaliscan cotton rat

Sigmodon mascotensis

The Jaliscan cotton rat or Mexican cotton rat (*Sigmodon mascotensis*) is a species of rodent in the family Cricetidae. It is found only in Mexico. They commonly have brown fur with white fur on the belly. They are ground-dwelling and prefer open...

Jaliscan spiny pocket mouse

Jaliscan spiny pocket mouse

Heteromys spectabilis

The Jaliscan spiny pocket mouse (*Heteromys spectabilis*) is a species of heteromyid rodent endemic to Mexico. The specific name *spectabilis* means "striking", and refers to the animal's relatively bright coloration. It was formerly placed in the...

Jamaican coney

Jamaican coney

Geocapromys brownii

The Jamaican coney (*Geocapromys brownii*), also known as the Jamaican hutia and the Browns hutia, is a hutia endemic to Jamaica. It is in the order Rodentia and the subfamily Capromyinae. It is the only extant native land mammal on Jamaica.

Jamaican fig-eating bat

Jamaican fig-eating bat

Ariteus flavescens

The Jamaican fig-eating bat (*Ariteus flavescens*) is a species of bat in the family Phyllostomidae. It is the only living species in the genus *Ariteus*. The scientific name translates as "yellowish and warlike". There are no recognised subspecies.

Jamaican flower bat

Jamaican flower bat

Phyllonycteris aphylla

The Jamaican flower bat (*Phyllonycteris aphylla*) is a critically endangered species of bat in the family Phyllostomidae. It is endemic to Jamaica.

Jamaican Fruit Bat

Jamaican Fruit Bat

Artibeus jamaicensis

The Jamaican fruit bat (*Artibeus jamaicensis*) is a frugivorous bat from the Neotropics. One of their favored places for roosting is large leaves or "tents", which provide shelter from the weather and hide them from predators.

Jamaican greater funnel-eared bat

Jamaican greater funnel-eared bat

Natalus jamaicensis

The Jamaican greater funnel-eared bat (*Natalus jamaicensis*) is a species of funnel-eared bat found in Jamaica. It was first described as the subspecies *Natalus major jamaicensis*, later as a subspecies of *Natalus stramineus*, and now as its own...

Jamaican red bat

Jamaican red bat

Lasiurus degelidus

The Jamaican red bat (*Lasiurus degelidus*) is a species of vesper bat. It is endemic to Jamaica.

James' sportive lemur

James' sportive lemur

Lepilemur jamesorum

James' sportive lemur (*Lepilemur jamesorum*), or the Manombo sportive lemur, is a sportive lemur endemic to Madagascar. It is one of the larger sportive lemurs with a total length of about 60 to 67 cm (24 to 26 in), of which 28 to 32 cm (11 to...

James's gerbil

James's gerbil

Dipodillus jamesi

James's gerbil (*Dipodillus jamesi*) is a species of rodent endemic to the eastern coast of Tunisia.

Jameson's red rock hare

Jameson's red rock hare

Pronolagus randensis

Jameson's red rock hare (*Pronolagus randensis*) is a leporid found only on the continent of Africa, mostly in the southern portion of the continent. Zimbabwe and Namibia are thought to have the largest populations of the red rock hare and it is...

Japanese Badger

Japanese Badger

Meles anakuma

The Japanese badger (*Meles anakuma*) is a species of carnivoran of the family Mustelidae, the weasels and their kin. Endemic to Japan, it is found on Honshu,

Kyushu, Shikoku, and Shōdoshima. It shares the genus *Meles* with its close relatives, the...

Japanese black bear

Japanese black bear

Ursus thibetanus japonicus

The Japanese black bear (*Ursus thibetanus japonicus*) is a subspecies of the Asian black bear that lives on two main islands of Japan: Honshu and Shikoku. There are said to be 10,000 black bears in Japan. The population of black bears on Shikoku...

Japanese boar

Japanese boar

Sus scrofa leucomystax

The Japanese boar (*Sus scrofa leucomystax*), also known as the white-moustached pig, *inoshishi* (イノシシ), or *yama kujira* (山鯨, lit. "mountain whale"), is a subspecies of wild boar native to all of Japan, save for Hokkaido and the Ryukyu Islands.

Japanese dormouse

Japanese dormouse

Glirulus japonicus

The Japanese dormouse (*Glirulus japonicus*) is a species of rodent in the family Gliridae endemic to Japan. It is the only extant species within the genus *Glirulus*. Its natural habitat is temperate forests. In Japanese, it is called *yamane* (ヤマネ...

Japanese Dwarf Flying Squirrel

Japanese Dwarf Flying Squirrel

Pteromys momonga

The Japanese dwarf flying squirrel (*Pteromys momonga*; Japanese: ニホンモモンガ, Hepburn: Nihon momonga) is one of two species of Old World flying squirrels in the genus *Pteromys*. It is native to Japan where it inhabits sub-alpine forests and boreal...

Japanese Giant Flying Squirrel

Japanese Giant Flying Squirrel

Petaurista leucogenys

The Japanese giant flying squirrel (*Petaurista leucogenys*) is a large species of flying squirrel that lives in sub-alpine and boreal forests. It is active at night and able to glide (not actually fly like a bat) long distances between trees by...

Japanese grass vole

Japanese grass vole

Microtus montebelli

The Japanese grass vole (*Microtus montebelli*) is a species of rodent in the family Cricetidae. It is found only in Japan.

Japanese Hare

Japanese Hare

Lepus brachyurus

The Japanese hare (*Lepus brachyurus*) is a species of hare endemic to Japan. In Japanese, it is called the *Nousagi* (Japanese: 野ウサギ), meaning "field rabbit".

Japanese house bat

Japanese house bat

Pipistrellus abramus

The Japanese house bat or Japanese pipistrelle (*Pipistrellus abramus*) is a species of vesper bat. An adult has a body length of 3.6–4.8 cm (1.4–1.9 in), a tail of 2.9–4.0 cm (1.1–1.6 in), and a wing length of 3.2–3.6 cm (1.3–1.4 in). It

prefers...

Japanese long-eared bat
Japanese long-eared bat
Plecotus sacrimontis

The Japanese long-eared bat (*Plecotus sacrimontis*) is a species of vesper bat endemic to Japan, where it is found in Hokkaido, Honshu and Shikoku. It has distinctive, long ears, hence its Japanese name, the 'rabbit bat'. Formerly included as a...

Japanese Macaque
Japanese Macaque
Macaca fuscata

The Japanese macaque (*Macaca fuscata*) is a terrestrial Old World monkey species. Colloquially, they are referred to as "snow monkeys" because some live in areas where snow covers the ground for months each year - no other non-human primate is...

Japanese Marten
Japanese Marten
Martes melampus

The Japanese marten (*Martes melampus*) is a mammal in the marten genus most closely related to the sable. It is 0.5 m (1.5 ft) in length typically, not counting a 20-cm-long tail (7.9 in), and between 1,000 and 1,500 grams (2.2 and 3.3 lb) in...

Japanese mole
Japanese mole
Mogera wogura

The Japanese mole (*Mogera wogura*), also known as Temminck's mole, is a species of mole endemic to Japan. A solitary and diurnal species, it can live for up to 3.5 years in the wild. The Kobe mole (*M. kobeae*) was formerly described as a distinct...

Japanese mountain mole
Japanese mountain mole
Euroscaptor mizura

The Japanese mountain mole (*Oreoscaptor mizura*) is a species of Old World mole in the family Talpidae. It is endemic to Japan. It is the only member of the monotypic genus *Oreoscaptor*. Its natural habitats are temperate forests and temperate...

Japanese noctule
Japanese noctule
Nyctalus furvus

The Japanese noctule (*Nyctalus furvus*) is a species of bat belonging to the family Vespertilionidae. It is endemic to Japan.

Japanese Raccoon Dog
Japanese Raccoon Dog
Nyctereutes viverrinus

The Japanese raccoon dog (*Nyctereutes viverrinus*), also known as the tanuki is one of two species in the genus *Nyctereutes*, alongside the Common raccoon dog, of which it was formerly thought to be a subspecies. Currently, it is thought to...

Japanese red-backed vole
Japanese red-backed vole
Myodes andersoni

The Japanese red-backed vole, Wakayama red-backed vole, or Anderson's red-backed vole (*Myodes andersoni*) is a species of rodent in the family Cricetidae. It is found only on the island of Honshu in Japan. It was first described by the

British...

Japanese river otter

Japanese river otter

Lutra lutra whiteleyi

The Japanese otter (Japanese: ニホンカワウソ(日本川獺, Hepburn: Nihon-kawauso) (*Lutra nippon*) or Japanese river otter is an extinct species of otter formerly widespread in Japan. Dating back to the 1880s, it was even seen in Tokyo. The population suddenly...

Japanese Serow

Japanese Serow

Capricornis crispus

The Japanese serow (*Capricornis crispus*) is a goat-antelope, an even-toed ungulate mammal that lives in dense woodland in Japan. This animal is a national symbol of Japan and is protected in conservation areas. It is labeled a "living national...

Japanese short-tailed bat

Japanese short-tailed bat

Eptesicus japonensis

The Japanese short-tailed bat (*Eptesicus japonensis*) is a species of bat belonging to the family Vespertilionidae. It is endemic to Japan where it is found at the base of the northern Japanese Alps, the Chichibu Mountains and Oze National Park,...

Japanese shrew mole

Japanese shrew mole

Urotrichus talpoides

The Japanese shrew mole (*Urotrichus talpoides*) or himizu (ヒミズ) is a species of mammal in the family Talpidae. It is endemic to Japan and is found on Honshu, Shikoku, Kyushu, Awaji Island, Shodo Island, Oki Islands, Tsushima Island, Goto Islands,...

Japanese Squirrel

Japanese Squirrel

Sciurus lis

The Japanese squirrel (*Sciurus lis*) is a tree squirrel in the genus *Sciurus* endemic to Japan. It was described by Dutch zoologist Coenraad Jacob Temminck in 1844. The Japanese squirrel's range includes the islands of Honshū, Shikoku, and Kyūshū....

Japanese water shrew

Japanese water shrew

Chimarrogale platycephalus

The Japanese water shrew (*Chimarrogale platycephalus*), also called the flat-headed water shrew, is a species of mammal in the family Soricidae. It is endemic to Japanese Islands, Honshu and Kyushu. They are considered extinct in Shikoku.

Japanese Weasel

Japanese Weasel

Mustela itatsi

The Japanese weasel (*Mustela itatsi*) is a carnivorous mammal belonging to the genus *Mustela* in the family Mustelidae. The most closely related *Mustela* species is the Siberian weasel (*Mustela sibirica*). Its taxonomic species name, *itatsi*, is...

Japanese wolf

Japanese wolf

Canis lupus hodophilax

The Japanese wolf (Japanese: ニホンオオカミ (日本狼) , Hepburn: Nihon ōkami, or 山犬, yamainu ; *Canis lupus hodophilax*), also known as the Honshū wolf, is an extinct subspecies of the gray wolf that was once endemic to the islands of Honshū, Shikoku and...

Japen rat

Japen rat

Rattus jobiensis

The Japen rat (*Rattus jobiensis*) is a species of rodent in the family Muridae found only in Yapen, Biak-Supiori, and Owi islands of West Papua, Indonesia.

Jatna's tarsier

Jatna's tarsier

Tarsius supriatnai

The Jatna's tarsier (*Tarsius supriatnai*) is a species of tarsier found in Indonesia. Their call, or "songs" are known to be different from other similar tarsier species. The Jatna's tarsier has giant eyes and ears, and they also can rotate their...

Java Mouse-Deer

Java Mouse-Deer

Tragulus javanicus

The Java mouse-deer (*Tragulus javanicus*) is a species of even-toed ungulate in the family Tragulidae. When it reaches maturity it is about the size of a rabbit, making it the smallest living ungulate. It is found in forests in Java and perhaps...

Java pipistrelle

Java pipistrelle

Pipistrellus javanicus

The Java pipistrelle (*Pipistrellus javanicus*) is a species of pipistrelle bat found in South and Southeast Asia, including Afghanistan, Bangladesh, Brunei, Cambodia, China, India, Indonesia, Laos, Malaysia, Myanmar, Nepal, Pakistan, Philippines,...

Javan Ferret-Badger

Javan Ferret-Badger

Melogale orientalis

Javan ferret-badgers (*Melogale orientalis*) are mustelids found in Indonesia. These secretive animals live in or close to forested areas where they hunt a wide range of prey.

Javan ghost shrew

Javan ghost shrew

Crocidura umbra

The Javan ghost shrew (*Crocidura umbra*) is a species of mammal in the family Soricidae. It is endemic to Java where it is known only from Mount Gede and can be found from the type locality at 1,611 and 1,950 m elevation on Mt. Gede. Its natural...

Javan Leopard

Javan Leopard

Panthera pardus melas

The Javan leopard (*Panthera pardus melas*) is a leopard subspecies. Morphological research indicates that the Javan leopard is distinct from other Asian leopard subspecies and split off from other Asian leopard subspecies in the Middle Pleistocene...

Javan Lutung

Javan Lutung

Trachypithecus auratus

The Javan lutung (*Trachypithecus auratus*) is a species of Old World monkey. The Latin word *auratus* in its scientific name means "golden", and refers to a less common color variant. The Javan lungur has a specialised stomach to digest plant...

Javan mastiff bat

Javan mastiff bat

Otomops formosus

The Javan mastiff bat (*Otomops formosus*) is a species of bat in the family Molossidae endemic to Indonesia.

Javan Rhinoceros

Javan Rhinoceros

Rhinoceros sondaicus

The Javan rhinoceros (*Rhinoceros sondaicus*) is a very rare member of the family Rhinocerotidae and one of five extant rhinoceroses. Once the most widespread of Asian rhinoceroses, it is now critically endangered, with only one known population in...

Javan slit-faced bat

Javan slit-faced bat

Nycteris javanica

The Javan slit-faced bat (*Nycteris javanica*) is a species of slit-faced bats found on the Kangean Islands of Indonesia and Nusa Penida, Java. This species is also located in West Timor. This species' population is decreasing.

Javan Slow Loris

Javan Slow Loris

Nycticebus javanicus

The Javan slow loris (*Nycticebus javanicus*) is a venomous primate from Indonesia. It is most closely related to the Sunda slow loris and the Bengal slow loris. The Javan slow loris population is in sharp decline due to a number of threats and for...

Javan surili

Javan surili

Presbytis comata

The Javan surili (*Presbytis comata*) is an endangered species of Old World monkey endemic to the western half of Java, Indonesia, a biodiversity hotspot. Other common names by which it is known by include gray, grizzled or Sunda Island surili;...

Javan tailless fruit bat

Javan tailless fruit bat

Megaerops kusnotoi

The Javan Tailless Fruit Bat (*Megaerops kusnotoi*) is a species of megabat in the family Pteropodidae. It is endemic to Indonesia. The consumption of bushmeat, meat from undomesticated animals, is a big ecological problem in Indonesia, and a...

Javan thick-thumbed bat

Javan thick-thumbed bat

Glischropus javanus

The Javan thick-thumbed bat (*Glischropus javanus*) is a species of vesper bat. It can be found in Indonesia.

Javan Warty Pig

Javan Warty Pig

Sus verrucosus

The Javan warty pig (*Sus verrucosus*) is one of the rarest pig species. It is native to the Indonesian islands Java and Bawean and is the biggest suine on the island. It is considered extinct on Madura Island and has been listed as Endangered on...

Javanese flying squirrel
Javanese flying squirrel
Iomys horsfieldii

The Javanese flying squirrel (*Iomys horsfieldii*) is a species of rodent in the family *Sciuridae*. It is found in Indonesia, Malaysia, and Singapore.

Javanese shrew
Javanese shrew
Crocidura maxi

The Javanese shrew (*Crocidura maxi*) is a species of mammal in the family *Soricidae*. It is native to Indonesia and East Timor. It ranges throughout Java and the Lesser Sunda Islands. It has also been introduced to Ambon Island and the Aru Islands.

Jenkins's shrew
Jenkins's shrew
Crocidura jenkinsi

Jenkins's shrew (*Crocidura jenkinsi*) is a critically endangered species of mammal in the family *Soricidae*. It is endemic to South Andaman Island, India.

Jenkins's shrew tenrec
Jenkins's shrew tenrec
Microgale jenkinsae

Jenkins's shrew tenrec (*Microgale jenkinsae*) is a species of mammal in the family *Tenrecidae*. It is endemic to Madagascar. Its natural habitats are subtropical or tropical dry forests and subtropical or tropical dry shrubland. It is threatened by...

Jentink's dormouse
Jentink's dormouse
Graphiurus crassicaudatus

Jentink's dormouse (*Graphiurus crassicaudatus*) is a species of rodent in the family *Gliridae*. It is found in Cameroon, Ivory Coast, Ghana, Liberia, Nigeria, and Togo, and possibly Benin, Equatorial Guinea, and Sierra Leone. Its natural habitat...

Jentink's duiker
Jentink's duiker
Cephalophus jentinki

Jentink's duiker (*Cephalophus jentinki*), also known as gidi-gidi in Krio and kaikulowulei in Mende, is a forest-dwelling duiker found in the southern parts of Liberia, southwestern Côte d'Ivoire, and scattered enclaves in Sierra Leone. It is...

Jentink's flying squirrel
Jentink's flying squirrel
Hylopetes platyurus

The Jentink's flying squirrel (*Hylopetes platyurus*) is a species of squirrel native to Indonesia and Malaysia. They are nocturnal omnivores, and have an average body mass of 131.27 mm and an average body mass of 56.98 grams.

Jentink's squirrel
Jentink's squirrel
Sundasciurus jentinki

Jentink's squirrel (*Sundasciurus jentinki*) is a species of rodent in the family

Sciuridae. It is named in honor of the Dutch zoologist Fredericus Anna Jentink. It is found in Indonesia and Malaysia. Its natural habitat is subtropical or tropical...

Jico deer mouse

Jico deer mouse

Habromys simulatus

The Jico deer mouse (*Habromys simulatus*) is a species of rodent in the family Cricetidae found only in Mexico. Its natural habitat is subtropical or tropical moist lowland forests.

Joffre's pipistrelle

Joffre's pipistrelle

Hypsugo joffrei

Joffre's bat (*Mirostrellus joffrei*), also known as Joffre's pipistrelle, is a species of vesper bat in the family Vespertilionidae. It is the only member of the genus *Mirostrellus*. It is found in South and Southeast Asia. It is threatened by...

Johan's spiny mouse

Johan's spiny mouse

Acomys johannis

Johan's spiny mouse (*Acomys johannis*) is a species of rodent in the family Muridae. It is found in Benin, Burkina Faso, Cameroon, Chad, Ghana, Mali, Niger, Nigeria, and Togo. Its natural habitats are subtropical or tropical dry shrubland,...

Johnston's African dormouse

Johnston's African dormouse

Graphiurus johnstoni

The Johnston's African dormouse (*Graphiurus johnstoni*) is a species of rodent in the family Gliridae. It is found in Malawi, Zambia, and Zimbabwe. Its natural habitat is moist savanna.

Johnston's forest shrew

Johnston's forest shrew

Sylvisorex johnstoni

Johnston's forest shrew (*Sylvisorex johnstoni*) is a species of mammal in the family Soricidae found in Burundi, Cameroon, the Central African Republic, the Republic of the Congo, the Democratic Republic of the Congo, Equatorial Guinea, Gabon,...

Johnston's genet

Johnston's genet

Genetta johnstoni

Johnston's genet (*Genetta johnstoni*) is a genet species native to the Upper Guinean forests. As it is threatened by deforestation and conversion of rainforest to agriculturally and industrially used land, it is listed as Near Threatened on the...

Johnston's mangabey

Johnston's mangabey

Lophocebus johnstoni

Johnston's mangabey (*Lophocebus johnstoni*) is a species of crested mangabey in the family Cercopithecidae. It had been considered a subspecies of the gray-cheeked mangabey (*L. albigena*), but in 2007 was given the status of species by Colin...

Johnstone's mastiff bat

Johnstone's mastiff bat

Otomops johnstonei

Johnstone's mastiff bat (*Otomops johnstonei*) is a species of bat in the Molossidae family endemic to Indonesia.

Jolly's mouse lemur

Jolly's mouse lemur

Microcebus jollyae

Jolly's mouse lemur (*Microcebus jollyae*) is a species of mouse lemur from Mananjary and Kianjavato in Madagascar. The species is named in honor of primatologist Alison Jolly. Jolly's mouse lemur is uniformly reddish-brown with a small white...

Jones's roundleaf bat

Jones's roundleaf bat

Hipposideros jonesi

Jones's roundleaf bat (*Hipposideros jonesi*) is a species of bat in the family Hipposideridae. It is endemic to southern West Africa. Its natural habitats are subtropical or tropical dry forest, dry savanna, moist savanna, subtropical or tropical...

Jouvenet's shrew

Jouvenet's shrew

Crocidura jouvenetae

Jouvenet's shrew (*Crocidura jouvenetae*) is a species of mammal in the family Soricidae. It is found in Côte d'Ivoire, Guinea, Liberia, and Sierra Leone. This species is present in lowland tropical moist forest and possibly in montane forest.It...

Juan Fernández Fur Seal

Juan Fernández Fur Seal

Arctocephalus philippii

The Juan Fernández fur seal (*Arctocephalus philippii*) is the second smallest of the fur seals, second only to the Galápagos fur seal. There is still much that is unknown about this species. Scientists still do not know the average life span of...

Judean Mountains blind mole-rat

Judean Mountains blind mole-rat

Spalax judaei

The Middle East blind mole-rat or Palestine mole-rat (*Nannospalax ehrenbergi*) is a species of rodent in the family Spalacidae.

Jujuy tuco-tuco

Jujuy tuco-tuco

Ctenomys juris

The Jujuy tuco-tuco (*Ctenomys juris*) is a species of rodent in the family Ctenomyidae. It is known only from one location at an elevation of 500 m in southeastern Jujuy Province of northern Argentina.

Julia Creek Dunnart

Julia Creek Dunnart

Sminthopsis douglasi

The Julia Creek dunnart (*Sminthopsis douglasi*) is a marsupial with a buffy brown upperside and white underside. This dunnart has a body length of 100–135 mm with a tail of 60–105 mm to make a total length of 160–240 mm. Its weight is between 40...

Juliana's golden mole

Juliana's golden mole

Neamblysomus julianae

Juliana's golden mole (*Neamblysomus julianae*) is a golden mole endemic to South

Africa. It is listed as an endangered species due to habitat loss and a restricted range. Golden moles are an ancient group of mammals that live mostly below ground....

Juliomys anoblepas
Juliomys anoblepas
Juliomys anoblepas

Juliomys anoblepas is a rodent in the genus *Juliomys* of the subfamily Sigmodontinae known from a single broken skull. The specimen was collected by Peter Wilhelm Lund in the caves of Lagoa Santa, Minas Gerais, Brazil, in the first half of the...

Juliomys ossitenuis
Juliomys ossitenuis
Juliomys ossitenuis

Juliomys ossitenuis is a species of rodent in the family Cricetidae native to South America.

Juliomys rimofrons
Juliomys rimofrons
Juliomys rimofrons

Juliomys rimofrons, also known as the cleft-headed juliomys, is a South American rodent species in the family Cricetidae. It is found in southern Brazil. The species is arboreal and lives in montane forests. It is threatened because of the small...

Jungle Cat
Jungle Cat
Felis chaus

The Jungle cat (*Felis chaus*) is a medium-sized cat. Its name comes from the Caucasus Mountains, the place this cat was first discovered. In Asia, it is the Jungle cat, but in Africa, it is commonly named Reed cat or Swamp cat, due to its...

Jungle palm squirrel
Jungle palm squirrel
Funambulus tristriatus

The jungle palm squirrel, jungle striped squirrel, or Western Ghats squirrel (*Funambulus tristriatus*) is a species of rodent in the family Sciuridae endemic to India. Its natural habitats are subtropical or tropical dry forests and...

Jungle shrew
Jungle shrew
Suncus zeylanicus

The jungle shrew (*Suncus zeylanicus*) is a species of mammal in the family Soricidae endemic to Sri Lanka. Its natural habitat is subtropical or tropical dry forests. It is threatened by habitat loss.

Junín grass mouse
Junín grass mouse
Akodon juninensis

The Junín grass mouse (*Akodon juninensis*) is a species of rodent in the family Cricetidae. It is found only in Peru.

Junín red squirrel
Junín red squirrel
Sciurus pyrrhinus

The Junín red squirrel (*Sciurus pyrrhinus*) is a species of squirrel from Peru and Ecuador.

Junin slender opossum

Junin slender opossum
Marmosops juninensis

The Junin slender opossum (*Marmosops juninensis*) is a species of South American opossum in the family Didelphidae. It is known only from the valley of the Chanchamayu in Peru (a range that includes Junín Region) where it lives in Andean montane...

Juniper vole
Juniper vole
Neodon juldaschi

The juniper vole (*Microtus juldaschi*) is a species of rodent in the family Cricetidae. It is found in Afghanistan, China, Pakistan and Tajikistan. It was formerly classified in the genus *Neodon*, but genetic evidence indicates that it is...

Juruá red howler
Juruá red howler
Alouatta juara

The Juruá red howler (*Alouatta juara*) is a species of howler monkey, native to Peru and Brazil.

Kaapori capuchin
Kaapori capuchin
Cebus kaapori

The *Cebus kaapori* is a species of frugivorous, gracile capuchin endemic to the Brazilian Amazon. Their geographical home range is relatively small and is within the most densely populated region of the Amazon. With the strong human presence...

Kabul markhor
Kabul markhor
Capra falconeri jerdoni

The Kabul markhor (*Capra falconeri megaceros*) is a near-threatened goat-antelope native to Afghanistan and Pakistan.

Kachin red-backed vole
Kachin red-backed vole
Eothenomys cachinus

The Kachin red-backed vole (*Eothenomys cachinus*) is a species of rodent in the family Cricetidae. It is found in Kachin State in northern Burma. Although described as a species by Thomas in 1921, it was subsequently included as a subspecies of...

Kachin woolly bat
Kachin woolly bat
Kerivoula kachinensis

The Kachin woolly bat is a species of bat found in Southeast Asia.

Kafue lechwe
Kafue lechwe
Kobus leche kafuensis

The Kafue lechwe (*Kobus leche kafuensis*) is a subspecies of the southern lechwe. It is endemic to the Kafue Flats, Zambia. It is listed on the IUCN Red List as vulnerable.

Kafue mole-rat
Kafue mole-rat
Fukomys kafuensis

The Kafue mole-rat (*Fukomys kafuensis*) is a species of rodent in the family

Bathyergidae. It is endemic to Zambia. Its natural habitat is moist savanna.
Kahuzi swamp shrew
Kahuzi swamp shrew
Myosorex jejei

The Kahuzi swamp shrew (*Myosorex jejei*) is a species of mammal in the family Soricidae found in the Democratic Republic of the Congo. Its natural habitat is swamp. It is threatened by habitat loss.

Kaibab Squirrel
Kaibab Squirrel
Sciurus aberti kaibabensis

The Kaibab squirrel (*Sciurus aberti kaibabensis*) is an elusive isolated tassel-eared squirrel that lives in the Kaibab Plateau. It is now considered a subspecies of the Abert's squirrel (*Sciurus aberti*). This squirrel is not found anywhere else...

Kaiser's rock rat
Kaiser's rock rat
Aethomys kaiseri

Kaiser's rock rat (*Aethomys kaiseri*) is a species of rodent in the family Muridae found in Angola, Burundi, the Democratic Republic of the Congo, Kenya, Malawi, Rwanda, Tanzania, Uganda, and Zambia. Its natural habitats are subtropical or...
Kakadu dunnart
Kakadu dunnart
Sminthopsis bindi

The Kakadu dunnart (*Sminthopsis bindi*) is a dunnart first described in 1994 and whose closest relative is the Carpentarian dunnart. It typically has a body length of 50-85mm with a tail 60-105mm long, for a total length between 110-190mm. It...
Kakadu pebble-mound mouse
Kakadu pebble-mound mouse
Pseudomys calabyi

The Kakadu pebble-mound mouse (*Pseudomys calabyi*) is a rodent native to Australia. It is one of the pebble-mound mice.
Kalinowski's agouti
Kalinowski's agouti
Dasyprocta kalinowskii

Kalinowski's agouti (*Dasyprocta kalinowskii*) is a species of rodent in the family Dasyproctidae. It is endemic to southeast Peru. It occurs at elevations of up to 3,080 metres (10,100 ft) asl. It is threatened by habitat loss.
Kalinowski's mastiff bat
Kalinowski's mastiff bat
Mormopterus kalinowskii

Kalinowski's mastiff bat (*Mormopterus kalinowskii*) is a species of bat in the family Molossidae, the free-tailed bats. It is native to Peru and northern Chile. Relatively little is known about the species, but it is thought to be common in its...
Kalinowski's mouse opossum
Kalinowski's mouse opossum
Hyladelphys kalinowskii

Kalinowski's mouse opossum or the Peru gracile mouse opossum (*Hyladelphys kalinowskii*) is a species of opossum in the family Didelphidae. It is found in Brazil, French Guiana, Guyana, and Peru. Its natural habitat is subtropical or tropical...

Kalinowski's Oldfield mouse
Kalinowski's Oldfield mouse
Thomasomys kalinowskii

Kalinowski's Oldfield mouse (*Thomasomys kalinowskii*) is a species of rodent in the family Cricetidae. It is found only in Peru.

Kalko's round-eared bat
Kalko's round-eared bat
Lophostoma kalkoae

Kalko's round-eared bat (*Lophostoma kalkoae*) is a species of leaf-nosed bat endemic to Panama.

Kam dwarf hamster
Kam dwarf hamster
Cricetulus kamensis

The Kam dwarf hamster (*Cricetulus kamensis*) is a species of rodent in the family Cricetidae. It is found only in the mountains of western China where it inhabits grassland, shrubby marshes and steppes. Although it has a limited range, the...

Kamchatka brown bear
Kamchatka brown bear
Ursus arctos beringianus

The Kamchatka brown bear (*Ursus arctos beringianus*), also known as the "Far Eastern brown bear", or in Russian: Камчатский бурый медведь, romanized: Kamchatsky bury medved, is a subspecies of brown bear.

Kamchatka shrew
Kamchatka shrew
Sorex camtschatica

The Kamchatka shrew (*Sorex camtschatica*) is a species of mammal in the family Soricidae. It is endemic to Russia.

Kangaroo Island dunnart
Kangaroo Island dunnart
Sminthopsis aitkeni

The Kangaroo Island dunnart (*Sminthopsis aitkeni*) is a dark sooty-grey coloured dunnart species first described in 1969, with paler underparts of its body. It has an average body length of 170–198 mm, a snout to anus length of 80–93 mm, a tail...

Kano mole
Kano mole
Mogera kanoana

Kano's mole (*Mogera kanoana*) is a species of mole endemic to Taiwan. The specific name kanoana is derived from Tadao Kano, a Japanese naturalist who made the first record of these animals in 1940s. It is found in the mountainous center and the...

Kansu red deer
Kansu red deer
Cervus canadensis kansuensis

The Kansu red deer (*Cervus canadensis kansuensis*) is a subspecies of wapiti found in the Gansu province of China. This subspecies forms, along with the closely related Sichuan deer, and Tibetan red deer, the southernmost wapiti group.

Karimi's fat-tailed mouse opossum
Karimi's fat-tailed mouse opossum
Thylamys karimii

Karimi's fat-tailed mouse opossum (*Thylamys karimii*) is a species of opossum in the family Didelphidae. It is endemic to central and northeast Brazil, where it is

found in the cerrado and caatinga at elevations from 300 to 1100 m. This opossum...
Karoo rock elephant shrew
Karoo rock elephant shrew
Elephantulus pilicaudus

The Karoo rock elephant shrew or Karoo rock sengi (*Elephantulus pilicaudus*) is a type of shrew in the genus *Elephantulus*. Its known habitat is in South Africa (Northern Cape Province, Western Cape Province).

Kashmir cave bat
Kashmir cave bat
Myotis longipes

The Kashmir cave bat (*Myotis longipes*) is a species of vesper bat. It is endemic to the Western Himalayas of South Asia. It is found in the Western Himalayan broadleaf forests ecoregion, within Bhutan, India, Nepal, Pakistan, and Afghanistan.

Kashmir field mouse
Kashmir field mouse
Apodemus rusiges

The Kashmir field mouse (*Apodemus rusiges*) is a species of rodent in the family Muridae. It is found in India, Nepal, and Pakistan.

Kashmir flying squirrel
Kashmir flying squirrel
Eoglaucomys fimbriatus

The Kashmir flying squirrel (*Eoglaucomys fimbriatus*) is a species of rodent in the family Sciuridae. It is monotypic within the genus *Eoglaucomys*. It is found in India and Pakistan. Its natural habitat is subtropical or tropical dry forests. It...

Kashmir Gray Langur
Kashmir Gray Langur
Semnopithecus ajax

The Kashmir gray langur (*Semnopithecus ajax*) is a leaf-eating monkey found only in India. It lives at the highest elevation of any non-human primate in the world, ranging from 1,500 to 4,733 m (4,921 to 15,528 ft).

Kashmir musk deer
Kashmir musk deer
Moschus cupreus

The Kashmir musk deer (*Moschus cupreus*) is an endangered species of musk deer native to Afghanistan, India, and Pakistan. Recent studies have shown that the species is also native to western Nepal. This species was originally described as a...

Kashmir pygmy shrew
Kashmir pygmy shrew
Sorex planiceps

The Kashmir pygmy shrew (*Sorex planiceps*) is a species of mammal in the family Soricidae. It is found in India and Pakistan.

Kashmir stag
Kashmir stag
Cervus canadensis hanglu

The Kashmir stag (*Cervus hanglu hanglu*), also called hangul, is a subspecies of Central Asian red deer endemic to Kashmir and surrounding areas. It is found in dense riverine forests in the high valleys and mountains of Jammu and Kashmir and...

Kashmir white-toothed shrew
Kashmir white-toothed shrew
Crocidura pullata

The Kashmir white-toothed shrew (*Crocidura pullata*) is a species of mammal in the family Soricidae. It is found in India and Pakistan.

Katanglad shrew-mouse
Katanglad shrew-mouse
Crunomys suncoides

The Katanglad shrew-mouse (*Crunomys suncoides*), also known as the Kitanglad shrew-mouse is a species of rodent in the family Muridae. It is known only from one specimen taken at 2250 m on Mount Kitanglad, Bukidnon Province, Philippines.

Katinka's shrew
Katinka's shrew
Crocidura katinka

Katinka's shrew (*Crocidura katinka*) is a species of mammal in the family Soricidae. It has been recorded from Israel, Syria, and Palestine, but it may now be extinct in Israel. It may also be present in southwestern Iran.

Kazakhstan blind mole-rat
Kazakhstan blind mole-rat
Spalax uralensis

The Kazakhstan blind mole-rat or Ural blind mole-rat (*Spalax uralensis*) is a species of rodent in the family Spalacidae. It is endemic to Kazakhstan, along the Ural River basin and the flood plains of the Uil, Temir, and Emba rivers. Whereas the...

Kazbeg birch mouse
Kazbeg birch mouse
Sicista kazbegica

The Kazbeg birch mouse (*Sicista kazbegica*) is a species of rodent in the family Sminthidae. It is found in Georgia and Russia. Its natural habitat is temperate forests.

Keast's tube-nosed fruit bat
Keast's tube-nosed fruit bat
Nyctimene keasti

Keast's tube-nosed fruit bat (*Nyctimene keasti*) is a species of megabat in the family Pteropodidae found in Babar, Tanimbar, and the Kai Islands.

Keen's myotis
Keen's myotis
Myotis keenii

Keen's myotis (*Myotis keenii*) is a species of vesper bat. It is found in British Columbia in Canada and in Washington and Alaska in the United States. It is named after the Rev. John Henry Keen, who collected the specimen that formed the basis...

Kei flying fox
Kei flying fox
Pteropus keyensis

The Kei flying fox (*Pteropus keyensis*) is a species of megabat in the genus *Pteropus* found in the Kai Islands of Indonesia. It was formerly considered a subspecies of the black-bearded flying fox (*Pteropus melanopogon*). Very little is known...

Kei myotis
Kei myotis
Myotis stalkerii

The Kei myotis (*Myotis stalkerii*) is a species of vesper bat. It is found only in Indonesia.

Kelaart's long-clawed shrew

Kelaart's long-clawed shrew

Feroculus feroculus

Kelaart's long-clawed shrew (*Feroculus feroculus*) is a species of mammal in the family Soricidae. It is monotypic within the genus *Feroculus*. It is endemic to Sri Lanka and southern India. Its natural habitats are subtropical or tropical dry...

Kelaart's pipistrelle

Kelaart's pipistrelle

Pipistrellus ceylonicus

Kelaart's pipistrelle (*Pipistrellus ceylonicus*) is a species of vesper bat. It is found in southern and south-eastern Asia from Pakistan to Indonesia.

Kellen's dormouse

Kellen's dormouse

Graphiurus kelleni

Kellen's dormouse (*Graphiurus kelleni*) is a species of rodent in the family Gliridae. It is native to tropical Africa where its range extends from the Gambia and Senegal to Kenya and Tanzania. Its natural habitats are subtropical or tropical dry...

Kemp's gerbil

Kemp's gerbil

Gerbilliscus kempii

Kemp's gerbil (*Gerbilliscus kempii*) is a species of rodent. Mammal Species of the World considers *G. kempii* and *G. gambianus* to be synonyms, however the IUCN has assessed each taxon as were they different species. According to the IUCN in 2004,...

Kemp's grass mouse

Kemp's grass mouse

Deltamys kempii

Kemp's grass mouse (*Deltamys kempii*) is a species of South American rodent in the family Cricetidae. It is the only species in the genus *Deltamys*. It was formerly considered a member of the genus *Akodon*, but is now considered sufficiently...

Kemp's spiny mouse

Kemp's spiny mouse

Acomys kempii

Kemp's spiny mouse (*Acomys kempii*) is a species of rodent in the family Muridae. It is found in Kenya, Somalia, and Tanzania. Its natural habitats are dry savanna and rocky areas. It is one of only two mammals, the other being *Acomys percivali*,...

Kemp's thicket rat

Kemp's thicket rat

Thamnomys kempii

Kemp's thicket rat (*Thamnomys kempii*) is a species of rodent in the family Muridae. It is found in Burundi, Democratic Republic of the Congo, and Rwanda. Its natural habitat is subtropical or tropical moist montane forests. It is threatened by...

Kenai Peninsula wolf

Kenai Peninsula wolf

Canis lupus alces

The Kenai Peninsula wolf (*Canis lupus alces*), also known as the Kenai Peninsula

grey wolf, is an extinct subspecies of the gray wolf that lived on the Kenai Peninsula in southern Alaska.
Kenneth's white-toothed rat
Kenneth's white-toothed rat
Berylmys mackenziei

Kenneth's white-toothed rat (*Berylmys mackenziei*) is a species of rodent in the family Muridae. It is found in China, India, Myanmar, and Vietnam.
Kenya coast galago
Kenya coast galago
Paragalago cocos

The Kenya coast galago (*Paragalago cocos*) is a species of primate in the family Galagidae. It is found across Kenya and Tanzania, particularly around the northern coastal area of each of the countries, and has been located living from sea level...
Kenyan African mole-rat
Kenyan African mole-rat
Tachyoryctes ibeanus

The Kenyan African mole-rat or Kenyan mole-rat (*Tachyoryctes ibeanus*) is a species of rodent in the family Spalacidae. It is endemic to Kenya. Its natural habitats are dry savanna, moist savanna, arable land, pastureland, plantations, rural...
Kenyan big-eared free-tailed bat
Kenyan big-eared free-tailed bat
Tadarida lobata

The Kenyan big-eared free-tailed bat (*Tadarida lobata*) is a species of bat in the family Molossidae. It is found in Kenya and Zimbabwe. Its natural habitat is dry savanna.
Kenyan wattled bat
Kenyan wattled bat
Glauconycteris kenyacola

The Kenyan wattled bat (*Glauconycteris kenyacola*) is a species of vesper bat and a member of the family Vespertilionidae. It is found only in Kenya.
Kerala rat
Kerala rat
Rattus ranjinae

The Kerala rat (*Rattus ranjinae*) is a species of rodent in the family Muridae found only in Kerala, India. In Kerala, it is known only from fragmented locations in Alappuzha, Thrissur, and Thiruvananthapuram. Its natural habitats are...
Kerama deer
Kerama deer
Cervus nippon keramae

The Kerama deer (also Kerama Sika) is a endangered subspecies of the Sika deer native to the Kerama Islands.
Kerbis Peterhans's wood mouse
Kerbis Peterhans's wood mouse
Hylomyscus kerbispeterhansi

The Kerbis Peterhans's wood mouse (*Hylomyscus kerbispeterhansi*) is a species of rodent in the family Muridae. It is found in western Kenya, and presumably eastern Uganda on Mt. Elgon, and was first described as a new species to science in 2014....
Kerkennah Islands gerbil
Kerkennah Islands gerbil

Dipodillus zakariai

The Kerkennah Islands gerbil (*Dipodillus zakariai*) is a rodent from Kerkennah Islands, Tunisia.

Kermode bear

Kermode bear

Ursus americanus kermodei

The Kermode bear, sometimes called the spirit bear (*Ursus americanus kermodei*), is a subspecies of the American black bear and lives in the Central and North Coast regions of British Columbia, Canada. It is the official provincial mammal of...

Kerr's Atlantic tree-rat

Kerr's Atlantic tree-rat

Phyllomys kerri

Kerr's Atlantic tree-rat or Moojen's Atlantic tree rat (*Phyllomys kerri*), is a spiny rat species from South America. It is found in Brazil.

Key deer

Key deer

Odocoileus virginianus clavium

The Key deer (*Odocoileus virginianus clavium*) is an endangered deer that lives only in the Florida Keys. It is a subspecies of the white-tailed deer (*O. virginianus*). It is the smallest extant North American deer species.

Key Largo cotton mouse

Key Largo cotton mouse

Peromyscus gossypinus allapaticola

The Key Largo cotton mouse (*Peromyscus gossypinus allapaticola*) is a subspecies of the cotton mouse (*Peromyscus gossypinus*) and is found on Key Largo in the upper Florida Keys. It is a slightly larger mouse with a more reddish color than other...

Khajuria's leaf-nosed bat

Khajuria's leaf-nosed bat

Hipposideros durgadasi

The Khajuria's leaf-nosed bat, also known as Durga Das's leaf-nosed bat (*Hipposideros durgadasi*), is a species of bat in the family Hipposideridae. It is endemic to India. Its natural habitat is caves. It is threatened by habitat loss.

Khartoum gerbil

Khartoum gerbil

Dipodillus stigmonyx

The Khartoum gerbil (*Dipodillus stigmonyx*) is found mainly in Sudan.

Kiang

Kiang

Equus kiang

The kiang (*Equus kiang*) is a large graceful wild ass native to the Tibetan Plateau. It is closely related to the onager and has never been domesticated.

Kihaule's mouse shrew

Kihaule's mouse shrew

Myosorex kihaulei

Kihaule's mouse shrew (*Myosorex kihaulei*) is a species of mammal in the family Soricidae endemic to Tanzania where it is known only from the Udzungwa Mountains, at the western end of the Eastern Arc Mountains. Its natural habitats are...

Kilimanjaro mouse shrew

Kilimanjaro mouse shrew

Myosorex zinki

The Kilimanjaro mouse shrew (*Myosorex zinki*) is a species of mammal in the family Soricidae endemic to Tanzania. Its natural habitats are subtropical or tropical moist montane forests and swamps.

Kilimanjaro shrew

Kilimanjaro shrew

Crocidura monax

The Kilimanjaro shrew (*Crocidura monax*) is a species of mammal in the family Soricidae. It is found in Kenya and Tanzania.

Killer Whale

Killer Whale

Orcinus orca

The Killer whale (*Orcinus orca*) is a toothed whale belonging to the oceanic dolphin family. It is the largest member of this family. The sophisticated hunting techniques and vocal behaviors of Killer whales have been described as manifestations...

Kimberley rock rat

Kimberley rock rat

Zyzomys woodwardi

The Kimberley rock rat (*Zyzomys woodwardi*) is a species of rodent in the family Muridae. It is found only in Australia, specifically in the northern tropical part of the Northern Territory and adjacent Kimberley region of Western Australia, in...

Kinabalu shrew

Kinabalu shrew

Crocidura baluensis

The Kinabalu shrew (*Crocidura baluensis*) is a species in the family Soricidae. It is endemic to the mountain Mount Kinabalu on Borneo, and its sister peak, Mount Tambuyukon. Despite its resemblance in external morphology with the mountain shrew...

Kinabalu squirrel

Kinabalu squirrel

Callosciurus baluensis

The Kinabalu squirrel (*Callosciurus baluensis*) is a species of rodent in the family Sciuridae. It is endemic to highland forest in East Malaysia. Its name is a reference to Mount Kinabalu, though it is not restricted to this mountain. Its tail...

Kinda baboon

Kinda baboon

Papio cynocephalus kindae

The Kinda baboon (*Papio kindae*) is a species of baboon present in the miombo woodlands of Angola, the Democratic Republic of the Congo, Zambia, and possibly western Tanzania. It was once considered a subspecies of the yellow baboon (P....

King African mole-rat

King African mole-rat

Tachyoryctes rex

The King African mole-rat, King mole-rat, or Alpine mole-rat, (*Tachyoryctes rex*) is a burrowing rodent in the genus *Tachyoryctes* of family Spalacidae. It only occurs high on Mount Kenya, where it is common. Originally described as a separate...

King Colobus

King Colobus

Colobus polykomos

The king colobus (*Colobus polykomos*), also known as the western black-and-white colobus, is a species of Old World monkey, found in lowland and mountain rainforests in a region stretching from Senegal, through Guinea-Bissau, Guinea, Sierra Leone...

King genet

King genet

Genetta poensis

The king genet (*Genetta poensis*) is a small carnivoran native to the Republic of the Congo, Equatorial Guinea, Liberia, Ghana and Côte d'Ivoire. As it has not been recorded since 1946, it is listed as Data Deficient on the IUCN Red List. It...

King horseshoe bat

King horseshoe bat

Rhinolophus rex

The king horseshoe bat (*Rhinolophus rex*) is a species of bat in the family Rhinolophidae. It is endemic to the east coast of China.

King jird

King jird

Meriones rex

The king jird (*Meriones rex*) is a species of rodent in the family Muridae. It is found in Saudi Arabia and Yemen.

King rat

King rat

Uromys rex

The king rat (*Uromys rex*) is a large species of rodent in the family Muridae. It is endemic to the island of Guadalcanal in the Solomon Islands. Like the two other species of rodent in Guadalcanal, it is placed in the genus *Uromys*. It lives in...

Kinkajou

Kinkajou

Potos flavus

The kinkajou (*Potos flavus*) is a tropical rainforest mammal related to olingos, coatis, raccoons, and the ringtail and cacomistle. It is the only member of the genus *Potos* and is also known as the "honey bear". Kinkajous are arboreal, a lifestyle...

Kintampo rope squirrel

Kintampo rope squirrel

Funisciurus substriatus

The Kintampo rope squirrel (*Funisciurus substriatus*) is a species of rodent in the family Sciuridae. It is found in Benin, Burkina Faso, Ivory Coast, Ghana, Niger, and Togo. Its natural habitat is moist savanna.

Kipunji

Kipunji

Rungwecebus kipunji

The kipunji (*Rungwecebus kipunji*) is a species of Old World monkey that lives in Africa. It was discovered by researchers from the Wildlife Conservation Society, the University of Georgia, and Conservation International, in December 2003 and July...

Kirchner's viscacha rat

Kirchner's viscacha rat

Tympanoctomys kirchnerorum

Kirchner's viscacha rat or Kirchner's vizcacha rat (*Tympanoctomys kirchnerorum*) is

a species of rodent in the family Octodontidae described in 2014. It is one of two species in the genus Tympanoctomys. That species is endemic to Chubut Province...

Kirk's Dik-Dik

Kirk's Dik-Dik

Madoqua kirkii

Kirk's dik-dik (*Madoqua kirkii*) is a small antelope native to Eastern Africa and one of four species of dik-dik antelope. It is believed to have six subspecies and possibly a seventh existing in southwest Africa. Dik-diks are herbivores,...

Kit Fox

Kit Fox

Vulpes macrotis

The Kit fox (*Vulpes macrotis*) is a fox species that inhabits arid and semi-arid regions of North America. These foxes are the smallest of the four species of *Vulpes* occurring in North America and are among the smallest of the vulpines worldwide....

Kitti's hog-nosed bat

Kitti's hog-nosed bat

Craseonycteris thonglongyai

Kitti's hog-nosed bat (*Craseonycteris thonglongyai*), also known as the bumblebee bat, is a near-threatened species of bat and the only extant member of the family *Craseonycteridae*. It occurs in western Thailand and southeast Myanmar, where it...

Kivu climbing mouse

Kivu climbing mouse

Dendromus nyasae

The Kivu climbing mouse (*Dendromus nyasae*) is a species of rodent in the family *Nesomyidae*. It is found in Burundi, Democratic Republic of the Congo, Rwanda, Malawi, Tanzania, Uganda, and Zambia. Its natural habitats are subtropical or tropical...

Kivu giant pouched rat

Kivu giant pouched rat

Cricetomys kivuensis

The Kivu giant pouched rat (*Cricetomys kivuensis*) is a species of rodent in the family *Nesomyidae*. It is found in Southern Africa.

Kivu long-haired shrew

Kivu long-haired shrew

Crocidura lanosa

The Kivu long-haired shrew (*Crocidura lanosa*) is a species of mammal in the family *Soricidae*. It is found in Democratic Republic of the Congo and Rwanda. Its natural habitats are subtropical or tropical moist montane forests and swamps. It is...

Kivu shrew

Kivu shrew

Crocidura kivuana

The Kivu shrew (*Crocidura kivuana*) is a species of mammal in the family *Soricidae*. It is endemic to Democratic Republic of the Congo. Its natural habitat is swamps.

Klipspringer

Klipspringer

Oreotragus oreotragus

The klipspringer (*Oreotragus oreotragus*) is a small, sturdy antelope found in eastern and southern Africa. It was first described by German zoologist Eberhard August Wilhelm von Zimmermann in 1783. A gregarious animal, the klipspringer forms...

Kloss's gibbon
Kloss's gibbon
Hylobates klossii

Kloss's gibbon (*Hylobates klossii*), also known as the Mentawai gibbon, the bilou or dwarf siamang, is an endangered primate in the gibbon family, Hylobatidae. It is identifiable in that it is all black, resembling the siamang with its black fur,...

Kloss's mole
Kloss's mole
Euroscaptor klossi

Kloss's mole (*Euroscaptor klossi*) is a species of mammal in the family Talpidae. It is found in Laos and Thailand. It was named after zoologist C. Boden Kloss. The Malaysian mole (*E. malayanus*) was formerly considered a subspecies, but a 2008...

Kloss's squirrel
Kloss's squirrel
Callosciurus albescens

Kloss's squirrel or Kloss squirrel (*Callosciurus albescens*) is a species of rodent in the family Sciuridae. It is endemic to northern Sumatra in Indonesia. Population data is insufficient to assess its conservation status according to the IUCN....

Kluchor birch mouse
Kluchor birch mouse
Sicista kluchorica

The Kluchor birch mouse (*Sicista kluchorica*) is a species of rodent in the family Sminthidae. It is endemic to Russia. Its natural habitat is temperate forests.

Knox Jones's pocket gopher
Knox Jones's pocket gopher
Geomys knoxjonesi

Knox Jones's pocket gopher (*Geomys knoxjonesi*) is a species of pocket gopher found in Texas and New Mexico. This species is named for Dr. J. Knox Jones Jr. (1929–1992), a prolific mammalogist at Texas Tech University. It is a relatively small...

Koala
Koala
Phascolarctos cinereus

The koala or, inaccurately, koala bear (*Phascolarctos cinereus*) is an arboreal herbivorous marsupial native to Australia. It is the only extant representative of the family Phascolarctidae and its closest living relatives are the wombats. Because...

Kob
Kob
Kobus kob

The kob (*Kobus kob*) is an antelope found across Central Africa and parts of West Africa and East Africa. Together with the closely related reedbucks, waterbucks, lechwe, Nile lechwe, and puku, it forms the Reduncinae tribe. Found along the...

Kobayashi's bat
Kobayashi's bat
Eptesicus koyabashii

Kobayashi's bat (*Eptesicus kobayashii*) is a species of bat. An adult Kobayashi's bat has a body length of 6.0–6.3 cm, a tail length of 4.6–4.8 cm, and a wing length of 4.5–4.7 cm. The species is found only on the Korean Peninsula; it has been...

Kobe mole
Kobe mole
Mogera kobeae

The Japanese mole (*Mogera wogura*), also known as Temminck's mole, is a species of mole endemic to Japan. A solitary and diurnal species, it can live for up to 3.5 years in the wild. The Kobe mole (*M. kobeae*) was formerly described as a distinct...

Kock's mouse-eared bat
Kock's mouse-eared bat
Myotis dieteri

Kock's mouse-eared bat (*Myotis dieteri*) is a species of mouse-eared bat found in the Republic of the Congo.

Kodiak bear
Kodiak bear
Ursus arctos middendorffi

The Kodiak bear (*Ursus arctos middendorffi*), also known as the Kodiak brown bear, sometimes the Alaskan brown bear, inhabits the islands of the Kodiak Archipelago in southwest Alaska. It is the largest recognized subspecies or population of the...

Kodkod
Kodkod
Leopardus guigna

The kodkod (*Leopardus guigna*) (Spanish pronunciation:), also called güiña, is the smallest felid species native to the Americas. It lives primarily in central and southern Chile, as well as marginally in adjoining areas of Argentina. Its area...

Koepcke's hairy-nosed bat
Koepcke's hairy-nosed bat
Mimon koepckeae

Koepcke's hairy-nosed bat (*Mimon koepckeae*) is a species of bat endemic to Peru.

Koford's grass mouse
Koford's grass mouse
Akodon kofordi

Koford's grass mouse (*Akodon kofordi*) is a species of rodent in the family Cricetidae. It is found only in Peru. It is named after American biologist Carl B. Koford.

Kolan vole
Kolan vole
Caryomys inez

The Kolan vole, Inez's red-backed vole or Inez's vole (*Caryomys inez*) is a species of rodent in the family Cricetidae. It is found only in China. Two subspecies have been recognized, *Caryomys inez inez* from the northern part of its range and...

Kolar leaf-nosed bat
Kolar leaf-nosed bat
Hipposideros hypophyllus

The Kolar leaf-nosed bat (*Hipposideros hypophyllus*), or leafletted leaf-nosed bat is a species of bat in the family Hipposideridae. It is endemic to India. Its natural habitats are subtropical or tropical dry forests and caves. It is found in...

Kolombatovic's long-eared bat
Kolombatovic's long-eared bat
Plecotus kolombatovici

The Mediterranean long-eared bat (*Plecotus kolombatovici*), also known as Kolombatovic's long-eared bat, is a species of vesper bat ranging from Italy east through the Balkans east to Syria, and south to Jordan.

Komodo rat
Komodo rat
Komodomys rintjanus

The Komodo rat (*Komodomys rintjanus*) is a species of rodent in the family Muridae found only in the Lesser Sunda Islands of Indonesia, in Rintja, Padar, Lomblen, and Pantar islands. Its natural habitat is subtropical or tropical dry forests.It...

Kondana soft-furred rat
Kondana soft-furred rat
Millardia kondana

The Kondana soft-furred rat (*Millardia kondana*), also known as the Kondana rat or large metad, is an endangered species of rodent in the family Muridae. It is a nocturnal burrowing rat that is endemic to the Sinhgad plateau near Pune,...

Kongana shrew
Kongana shrew
Sylvisorex konganensis

The Kongana shrew (*Sylvisorex konganensis*) is a species of mammal in the family Soricidae endemic to Central African Republic. Its natural habitat is subtropical or tropical moist lowland forests.

Koopman's pencil-tailed tree mouse
Koopman's pencil-tailed tree mouse
Chiropodomys karlkoopmani

Koopman's pencil-tailed tree mouse (*Chiropodomys karlkoopmani*) is a species of arboreal rodent in the family Muridae. It is endemic to Pagai and Siberut islands in the Mentawai Islands, off the western coast of Sumatra in Indonesia.Its natural...

Koopman's rat
Koopman's rat
Rattus koopmani

Koopman's rat (*Rattus koopmani*) is a species of rodent in the family Muridae.It is found only on Peleng island, located in the Banggai Islands (Kepulauan Banggai) off the southeastern coast of Sulawesi, Indonesia.

Kordofan Giraffe
Kordofan Giraffe
Giraffa camelopardalis antiquorum

The Kordofan giraffe (*Giraffa camelopardalis antiquorum*) is a subspecies of giraffe. Its English name is a reference to Kordofan in Sudan.

Korean field mouse
Korean field mouse
Apodemus peninsulae

The Korean field mouse (*Apodemus peninsulae*), also known as the Korean wood mouse, is a species of mouse. It is distributed across Northeastern Asia, including the Russian Far East, northern China, the Korean Peninsula, Sakhalin, and Hokkaidō....

Korean fox
Korean fox
Vulpes vulpes peculiosa

The Korean fox (*Vulpes vulpes peculiosa*), also known as the Korean red fox, is a subspecies of red fox that lives in Korea, Russia andNortheast China. It has a body length of 66-68 cm, a tail length of 42-44 cm and a weight of 4.1-5.9 kg.

Korean hare
Korean hare
Lepus coreanus

The Korean hare (*Lepus coreanus*) is a species of hare native to the Korean Peninsula and adjoining parts of northeastern China. The Korean hare inhabits diverse habitats within its range, from remote mountain forests to cultivated land. Fur...

Korinch's rat
Korinch's rat
Rattus korinchi

Korinch's rat (*Rattus korinchi*) is a species of rodent in the family Muridae. It is found only in western Sumatra, Indonesia, and is only known from Mount Kerinci and Mount Talakmau, where it is endemic to high elevations above 2000 m. It is only...

Korrigum
Korrigum
Damaliscus lunatus korrigum

The korrigum (*Damaliscus lunatus korrigum*), also known as Senegal hartebeest, is a subspecies of the topi, a large African antelope.

Koslov's pika
Koslov's pika
Ochotona koslowi

Koslov's pika or Kozlov's pika (*Ochotona koslowi*) is a species of mammal in the family Ochotonidae. It is endemic to China. Its natural habitat is tundra. It is threatened by habitat loss. Kozlov's pika are herbivores, they are known as...

Kosrae flying fox
Kosrae flying fox
Pteropus ualanus

The Kosrae flying fox (*Pteropus ualanus*) is a species of megabat in the genus *Pteropus* found on the island of Kosrae, Micronesia.

Kouprey
Kouprey
Bos sauveli

The kouprey (*Bos sauveli*, Khmer: គោជ្រូក, Koŭprey, "forest ox"; also known as kouproh, "grey ox") is a little-known, forest-dwelling, wild bovine species native to Southeast Asia. A young male was sent to the Vincennes Zoo in 1937 where it was...

Kowari
Kowari
Dasyuroides byrnei

The kowari (*Dasyuroides byrnei*) is a small carnivorous marsupial native to central Australia. It was first described by a British-Australian evolutionary biologist, Sir Walter Baldwin Spencer in 1896. In an Australian Aboriginal language spoken...

Kozlov's pygmy jerboa
Kozlov's pygmy jerboa
Salpingotus kozlovi

Kozlov's pygmy jerboa (*Salpingotus kozlovi*) is a species of rodent in the family Dipodidae. It is found in northwestern China and southern and eastern Mongolia. Its natural habitat is temperate desert.

Kozlov's shrew
Kozlov's shrew
Sorex kozlovi

Kozlov's shrew (*Sorex kozlovi*) is a red-toothed shrew found only at the Mekong River, Tibet, China. This shrew is listed as a data deficient species.
Krebs's fat mouse

Krebs's fat mouse
Steatomys krebsii

The Krebs's fat mouse (*Steatomys krebsii*) is a species of rodent in the family Nesomyidae. It is found in Angola, Botswana, Lesotho, Namibia, South Africa, and Zambia. Its natural habitats are Mediterranean-type shrubby vegetation and subtropical...

Kri-kri

Kri-kri

Capra aegagrus cretica

The kri-kri (*Capra aegagrus cretica*), sometimes called the Cretan goat, Agrimi, or Cretan Ibex, is a feral goat inhabiting the Eastern Mediterranean, previously considered a subspecies of wild goat. The kri-kri is now found only on the island of...

Kuhl's pipistrelle

Kuhl's pipistrelle

Pipistrellus kuhlii

Kuhl's pipistrelle (*Pipistrellus kuhlii*) is a species of vesper bat that occurs in large areas of North Africa, Southern Europe and West Asia. It lives in temperate forests, subtropical or tropical dry shrubland, Mediterranean-type shrubby...

Kulina spiny rat

Kulina spiny rat

Proechimys kulinae

The Kulina spiny-rat, *Proechimys kulinae* or Javari spiny rat, is a spiny rat species from South America. It is found in Brazil and Peru.

Kultarr

Kultarr

Antechinomys laniger

The kultarr (*Antechinomys laniger*) (also called the "jerboa-marsupial" or marsupial jerboa) is a small insectivorous nocturnal marsupial inhabiting the arid interior of Australia. Preferred habitat includes stony deserts, shrubland, woodland,...

L'Hoest's Monkey

L'Hoest's Monkey

Cercopithecus lhoesti

L'Hoest's monkey (*Allochrocebus lhoesti*) or mountain monkey, is a guenon found in the upper eastern Congo basin. They mostly live in mountainous forest areas in small, female-dominated groups. They have a dark coat and can be distinguished by a...

La Plata Dolphin

La Plata Dolphin

Pontoporia blainvillei

The La Plata dolphin, franciscana or toninha (*Pontoporia blainvillei*) is a species of dolphin found in coastal Atlantic waters of southeastern South America. It is a member of the river dolphin group and the only one that lives in the ocean and...

La Touche's free-tailed bat

La Touche's free-tailed bat

Tadarida latouchei

The La Touche's free-tailed bat (*Tadarida latouchei*) is a species of bat in the family Molossidae. It is found in China, Japan, and Thailand.

Labrador Wolf

Labrador Wolf

Canis lupus labradorius

The Labrador wolf (*Canis lupus labradorius*) is a subspecies of Grey wolf. It is one of the least studied because it is very feared by people. It is closely related to the Newfoundland wolf.

Lac Alaotra bamboo lemur

Lac Alaotra bamboo lemur

Hapalemur alaotrensis

The Lac Alaotra bamboo lemur (*Hapalemur alaotrensis*), also known as the Lac Alaotra gentle lemur, Alaotran bamboo lemur, Alaotran gentle lemur, Alaotra reed lemur, or locally as the bandro, is a bamboo lemur. It is endemic to the reed beds in...

Lachaise's climbing mouse

Lachaise's climbing mouse

Dendromus lachaisei

Lachaise's climbing mouse (*Dendromus lachaisei*) is a species of climbing mouse described by Denys and Aniskine in 2012. First discovered on the slopes of Mount Nimba, it is now believed that the species is extant within Guinea, Liberia and Cote...

Lacustrine vole

Lacustrine vole

Microtus limnophilus

The lacustrine vole (*Microtus limnophilus*) is a species of rodent in the family Cricetidae. It is found in China and Mongolia.

Ladak pika

Ladak pika

Ochotona ladacensis

The Ladak pika (*Ochotona ladacensis*), also known as the Ladakh pika, is a species of mammal in the family Ochotonidae found in China, India, and Pakistan. Prior to identification as a separate species, specimens were thought to be of the plateau...

Ladew's Oldfield mouse

Ladew's Oldfield mouse

Thomasomys ladewi

Ladew's Oldfield mouse (*Thomasomys ladewi*) is a species of rodent in the family Cricetidae. It is found only in Bolivia.

Ladoga Ringed Seal

Ladoga Ringed Seal

Pusa hispida ladogensis

The Ladoga ringed seal (*Pusa hispida ladogensis*) is a freshwater subspecies of the Ringed seal which are found entirely in Lake Ladoga. It was isolated in freshwater lakes and separated from the Arctic ringed seal. This subspecies is related to...

Lady Burton's rope squirrel

Lady Burton's rope squirrel

Funisciurus isabella

The Lady Burton's rope squirrel (*Funisciurus isabella*) is a species of rodent in the family Sciuridae. It is found in Cameroon, Central African Republic, Republic of the Congo, Equatorial Guinea, and Gabon. Its natural habitats are tropical and...

Lagidium ahuacaense

Lagidium ahuacaense

Lagidium ahuacaense

Lagidium ahuacaense is a rodent in the mountain viscacha genus (*Lagidium*) that occurs in southern Ecuador. First observed in 2005 and formally described in 2009, it occurs more than 500 km (310 mi) north of the nearest previously known...

Lago Blanco tuco-tuco

Lago Blanco tuco-tuco

Ctenomys fodax

The Lago Blanco tuco-tuco (*Ctenomys fodax*) is a species of rodent in the family Ctenomyidae. It is known only from the Lago Blanco area in southern Argentina. Its karyotype has $2n = 28$ and $FN = 42$.

Lagos serotina

Lagos serotina

Eptesicus platyops

The *Lagos serotina* (*Eptesicus platyops*) is a species of vesper bat. It is found in Equatorial Guinea, Nigeria, and possibly Senegal.

Lagostomus crassus

Lagostomus crassus

Lagostomus crassus

Lagostomus crassus is an extinct species of viscacha that was native to Peru. It is known from a single skull found in southern Peru. The skull is not in any way fossilized, indicating that the animal it belonged to lived very recently. The skull...

Lake Baikal mountain vole

Lake Baikal mountain vole

Alticola olchonensis

The Lake Baikal mountain vole or Olkhon mountain vole (*Alticola olchonensis*) is a species of rodent in the family Cricetidae. It is found principally on the Olkhon and Ogoi islands on Lake Baikal, in southern Siberia. While it was originally...

Lake Chad gerbil

Lake Chad gerbil

Taterillus lacustris

The Lake Chad gerbil or Lake Chad tateril (*Taterillus lacustris*) is a species of rodent found in Cameroon and Nigeria. Its natural habitats are dry savanna, subtropical or tropical dry shrubland, and arable land.

Lake Mackay hare-wallaby

Lake Mackay hare-wallaby

Lagorchestes asomatus

The Lake Mackay hare-wallaby (*Lagorchestes asomatus*), also known as the central hare-wallaby or kuluwarri, is an extinct species of macropod formerly found in central Australia. Very little is known about it.

Lakeland Downs mouse

Lakeland Downs mouse

Leggadina lakedownensis

The Lakeland Downs mouse (*Leggadina lakedownensis*) is a species of rodent in the family Muridae. It is found only in Australia. Its natural habitat is dry savanna.

Lama dwarf hamster

Lama dwarf hamster

Cricetulus lama

The Lama dwarf hamster (*Cricetulus lama*) is a species of rodent in the family Cricetidae. It is found only in the mountains of western China where it inhabits grassland, shrubby marshes and steppes. Although it has a limited range, the...

Lami tuco-tuco
Lami tuco-tuco
Ctenomys lami

The Lami tuco-tuco (*Ctenomys lami*) is a species of rodent in the family Ctenomyidae. It is endemic to an area in the state of Rio Grande do Sul in southern Brazil, where it is found in the vicinity of sand dunes. The species is threatened by...

Laminate vlei rat
Laminate vlei rat
Otomys laminatus

The laminate vlei rat (*Otomys laminatus*) is a species of rodent in the family Muridae. It is found only in South Africa. Its natural habitats are subtropical or tropical high-altitude grassland and swamps.

Lamotte's roundleaf bat
Lamotte's roundleaf bat
Hipposideros lamottei

Lamotte's roundleaf bat (*Hipposideros lamottei*) is a species of bat found only at Mount Nimba on the border of Côte d'Ivoire, Guinea and Liberia. It is critically endangered.

Lamotte's shrew
Lamotte's shrew
Crocidura lamottei

Lamotte's shrew (*Crocidura lamottei*) is a species of mammal in the family Soricidae. It is found in Benin, Burkina Faso, Cameroon, Ivory Coast, Gambia, Ghana, Guinea, Guinea-Bissau, Liberia, Mali, Nigeria, Senegal, Sierra Leone, and Togo. Its...

Lamulate shrew
Lamulate shrew
Chodsigoa lamula

The lamulate shrew (*Chodsigoa lamula*) is a species of mammal in the family Soricidae. It is endemic to China.

Lander's horseshoe bat
Lander's horseshoe bat
Rhinolophus landeri

Lander's horseshoe bat (*Rhinolophus landeri*) is a species of bat in the family Rhinolophidae found in Africa. Its natural habitats are dry savanna, moist savanna, and caves.

Lang Bian white-bellied rat
Lang Bian white-bellied rat
Niviventer langbianis

The Lang Bian white-bellied rat (*Niviventer langbianis*) is a species of rodent in the family Muridae. It is found in India, Laos, Myanmar, Thailand, and Vietnam.

Lang's red colobus
Lang's red colobus
Piliocolobus langi

Lang's red colobus (*Piliocolobus langi*) is a species of red colobus monkey. Historically it had been treated as a subspecies of the Central African red colobus, (*P. foai*) but more recent taxonomies generally treat it as a separate...

Laotian Giant Flying Squirrel
Laotian Giant Flying Squirrel
Biswamoyopterus laoensis

The Laotian giant flying squirrel (*Biswamoyopterus laoensis*) is a species of flying squirrel native to Laos. It was the second described member in the genus *Biswamoyopterus*, after being first collected in September 2012 by scientists researching...

Laotian langur

Laotian langur

Trachypithecus laotum

The Laotian langur or white-browed black langur (*Trachypithecus laotum*) is a species of primate in the family Cercopithecidae. It is endemic to Laos. Its natural habitat is subtropical or tropical dry forests. Genetically the Indochinese black...

Laotian Rock Rat

Laotian Rock Rat

Laonastes aenigmamus

The Laotian rock rat or kha-nyou (*Laonastes aenigmamus*, Lao: ຂະຍຸ), sometimes called the "rat-squirrel", is a species of rodent found in the Khammouan region of Laos. The species was first described in a 2005 article by Paulina Jenkins and...

Lappet-eared free-tailed bat

Lappet-eared free-tailed bat

Chaerephon major

The lappet-eared free-tailed bat (*Chaerephon major*) is a species of bat in the family Molossidae. It is found in Benin, Burkina Faso, Democratic Republic of the Congo, Ivory Coast, Ghana, Guinea, Kenya, Liberia, Mali, Niger, Nigeria, Sudan,...

Lar Gibbon

Lar Gibbon

Hylobates lar

The Lar gibbon (*Hylobates lar*) is a captivating primate and a master of agility, being remarkably fast when swinging through the trees from branch to branch. It has long arms and hands, perfectly adapted to this means of locomotion. It is an...

Large Aden gerbil

Large Aden gerbil

Gerbillus poecilops

The Large Aden gerbil (*Gerbillus poecilops*) is a species of rodent in the family Muridae. It is found in Saudi Arabia and Yemen.

Large Asian roundleaf bat

Large Asian roundleaf bat

Hipposideros lekaguli

The large Asian roundleaf bat (*Hipposideros lekaguli*) is a species of bat in the family Hipposideridae. It is found in Malaysia, the Philippines, and Thailand. The specific name commemorates Thai zoologist and conservationist Boonsong Lekagul.

Large bamboo rat

Large bamboo rat

Rhizomys sumatrensis

The large bamboo rat, Sumatran rat, or Indomalayan rat (*Rhizomys sumatrensis*) is a species of rodent in the family Spalacidae found in Cambodia, China, Indonesia, Laos, Malaysia, Myanmar, Thailand, and Vietnam. It is one of four species of...

Large false serotine

Large false serotine

Hesperoptenus tomesi

The large false serotine (*Hesperoptenus tomesi*) is a species of vesper bat. It is

found only in Malaysia.

Large Flying Fox

Large Flying Fox

Pteropus vampyrus

The Large flying fox (*Pteropus vampyrus*) is a southeast Asian species of megabat. Despite its scientific name, it feeds exclusively on fruits, nectar, and flowers, like the other flying foxes of the genus *Pteropus*. It is noted for being one of...

Large forest bat

Large forest bat

Vespadelus darlingtoni

The large forest bat (*Vespadelus darlingtoni*) is a vespertilionid bat found in southeast Australia, Tasmania, and Lord Howe Island. They are classified as common.

Large fruit-eating bat

Large fruit-eating bat

Artibeus amplus

The large fruit-eating bat (*Artibeus amplus*) is a species of bat in the family Phyllostomidae. It is found in the South American neotropics, specifically in the countries of Colombia, Guyana, and Venezuela. The large fruit-eating bat is one of...

Large Indian Civet

Large Indian Civet

Viverra zibetha

The large Indian civet (*Viverra zibetha*) is a viverrid native to South and Southeast Asia. It is listed as Least Concern on the IUCN Red List. The global population is considered decreasing mainly because of trapping-driven declines in heavily...

Large Japanese Field Mouse

Large Japanese Field Mouse

Apodemus speciosus

Large Japanese field mice (*Apodemus speciosus*) are nocturnal rodents native to Japan. They are terrestrial and have excellent senses of hearing and smell.

Large Luzon forest rat

Large Luzon forest rat

Bullimus luzonicus

The Large Luzon Forest Rat, or Luzon Forest Rat (*Bullimus luzonicus*) is a species of rodent, the first of three to be described in the genus *Bullimus*. It is in the diverse family Muridae. It is found only in the Philippines. The rat has been...

Large Mindanao roundleaf bat

Large Mindanao roundleaf bat

Hipposideros coronatus

The large Mindanao roundleaf bat (*Hipposideros coronatus*) is a species of bat in the family Hipposideridae endemic to the Philippines.

Large Mindoro forest mouse

Large Mindoro forest mouse

Apomys gracilirostris

The large Mindoro forest mouse (*Apomys gracilirostris*) is a species of rodent in the family Muridae, from the genus *Apomys*. It is found only in the Philippines. Its natural habitat is subtropical or tropical moist montane forests. It is a large...

Large Mole

Large Mole

Mogera robusta

The Large mole (*Mogera robusta*) was formerly treated as a subspecies of the Japanese mole. It lives in a long burrow, seldom emerging on the surface of the ground during the day.

Large mosaic-tailed rat

Large mosaic-tailed rat

Mammelomys rattoides

The large mosaic-tailed rat (*Mammelomys rattoides*) is a species of rodent in the family Muridae. It is found in West Papua, Indonesia and Papua New Guinea.

Large myotis

Large myotis

Myotis chinensis

The large myotis (*Myotis chinensis*) is a species of vesper bat. It is found in central and southeast China, Hong Kong, Myanmar, Thailand, and northern Vietnam. It is also expected to occur in northern Laos, but not yet documented there.

Large New Guinea spiny rat

Large New Guinea spiny rat

Rattus praetor

The large New Guinea spiny rat (*Rattus praetor*) is a species of rodent in the family Muridae. It is found in Indonesia, Papua New Guinea, and the Solomon Islands.

Large pencil-tailed tree mouse

Large pencil-tailed tree mouse

Chiropodomys major

The large pencil-tailed tree mouse or greater pencil-tailed tree mouse (*Chiropodomys major*) is a species of arboreal rodent in the family Muridae. It is endemic to Borneo where it is only known from Sabah and Sarawak (Malaysia), although it is...

Large rufous horseshoe bat

Large rufous horseshoe bat

Rhinolophus rufus

The large rufous horseshoe bat (*Rhinolophus rufus*) is a species of bat in the family Rhinolophidae. It is endemic to the Philippines.

Large Samoan flying fox

Large Samoan flying fox

Pteropus coxi

The large Samoan flying fox (*Pteropus coxi*) is a species of fruit-eating megabat whose type specimen was originally collected in Samoa in 1856, but was not identified as a new species until 2009. The only known specimen was collected by an...

Large slit-faced bat

Large slit-faced bat

Nycteris grandis

The large slit-faced bat, *Nycteris grandis*, is a species of slit-faced bat with a broad distribution in forest and savanna habitats in West, Central, and East Africa. *N. marica* (Kershaw, 1923), is the available name for the southern savanna...

Large tree mouse

Large tree mouse

Pogonomys loriae

The large tree mouse (*Pogonomys loriae*) is a species of rodent in the family Muridae. It is found on the island of New Guinea (Indonesia and Papua New Guinea).

Large treeshrew
Large treeshrew
Tupaia tana

The large treeshrew (*Tupaia tana*) is a treeshrew species within the Tupaiidae. It is native to Sumatra and adjacent small islands, as well as in the lowlands and hills of Borneo.

Large vesper mouse
Large vesper mouse
Calomys callosus

The large vesper mouse (*Calomys callosus*) is a South American rodent species of the family Cricetidae. It is found in Argentina, Bolivia, Brazil and Paraguay. Its karyotype has $2n = 50$ and $FN = 66$. It was formerly synonymized with *C. expulsus*, but...

Large vlei rat
Large vlei rat
Otomys maximus

The large vlei rat (*Otomys maximus*) is a species of rodent in the family Muridae. It is found in Angola, Botswana, Democratic Republic of the Congo, Namibia, and Zambia. Its natural habitats are moist savanna, subtropical or tropical seasonally...

Large white-bellied rat
Large white-bellied rat
Niviventer excelsior

The large white-bellied rat (*Niviventer excelsior*), also known as the Sichuan niviventer, is a species of rodent in the family Muridae. It is found only in southwestern China. It occurs in Jiuzhaigou National Nature Reserve of Sichuan,...

Large-eared free-tailed bat
Large-eared free-tailed bat
Otomops martiensseni

The large-eared free-tailed bat (*Otomops martiensseni*) is a species of bat in the family Molossidae found in Angola, Central African Republic, Democratic Republic of the Congo, Ivory Coast, Djibouti, Ethiopia, Ghana, Kenya, Malawi, Rwanda, South...

Large-eared gray shrew
Large-eared gray shrew
Notiosorex evotis

The large-eared gray shrew (*Notiosorex evotis*) is a species of shrew.

Large-eared horseshoe bat
Large-eared horseshoe bat
Rhinolophus philippinensis

The large-eared horseshoe bat (*Rhinolophus philippinensis*) is a species of bat in the family Rhinolophidae. It is found in Australia, Indonesia, Malaysia, Papua New Guinea, and the Philippines.

Large-eared Oldfield mouse
Large-eared Oldfield mouse
Thomasomys macrotis

The large-eared Oldfield mouse (*Thomasomys macrotis*) is a species of rodent in the family Cricetidae. It is known only from a single locality in the Andes in north central Peru (which includes Rio Abiseo National Park), in montane forest at an...

Large-eared pied bat
Large-eared pied bat
Chalinolobus dwyeri

The large-eared pied bat (*Chalinolobus dwyeri*) is a species of vesper bat in the family Vespertilionidae. This vulnerable mammal can be found in Australia.

Large-Eared Pika

Large-Eared Pika

Ochotona macrotis

The Large-eared pika (*Ochotona macrotis*) is a small mountain-dwelling mammal found in Central Asia. Pikas nest among boulders and scree, and in the autumn they pull hay, soft twigs, and other stores of food under rocks to eat during the long,...

Large-eared sheath-tailed bat

Large-eared sheath-tailed bat

Emballonura diana

The large-eared sheath-tailed bat (*Emballonura diana*) is a species of sac-winged bat in the family Emballonuridae. It is found in Papua New Guinea and Solomon Islands. It is threatened by habitat loss.

Large-eared slit-faced bat

Large-eared slit-faced bat

Nycteris macrotis

The large-eared slit-faced bat, *Nycteris macrotis*, is a species of slit-faced bat which lives in forests and savannas throughout Africa. *Nycteris vinsoni* was once considered a synonym of *N. macrotis*, but it became recognized as a separate species...

Large-eared tenrec

Large-eared tenrec

Geogale aurita

The large-eared tenrec (*Geogale aurita*) is a species of mammal in the family Tenrecidae. It is the only species in the monotypic genus *Geogale*, and the only member of the subfamily Geogalinae. It is endemic to Madagascar where its natural...

Large-eared vole

Large-eared vole

Alticola macrotis

The large-eared vole (*Alticola macrotis*) is a species of rodent in the family Cricetidae. It is found in Mongolia and the Russian Federation.

Large-footed bat

Large-footed bat

Myotis adversus

The large-footed bat, large-footed mouse-eared bat, or large-footed myotis (*Myotis adversus*) is a species of vesper bat (family Vespertilionidae). It can be found in the following countries: Australia, Indonesia, Malaysia, Papua New Guinea,...

Large-headed capuchin

Large-headed capuchin

Sapajus macrocephalus

The large-headed capuchin (*Sapajus apella macrocephalus*) is a subspecies of the tufted capuchin monkey from South America. It is found in Bolivia, Brazil, Colombia, Ecuador and Peru. It was formerly thought to be its own species (S....

Large-scaled mosaic-tailed rat

Large-scaled mosaic-tailed rat

Mammelomys lanosus

The large-scaled mosaic-tailed rat (*Mammelomys lanosus*) is a species of rodent in the family Muridae. It is found in West Papua, Indonesia and Papua New Guinea.

Large-Spotted Civet
Large-Spotted Civet
Viverra megaspila

The Large-spotted civet (*Viverra megaspila*) is an endangered viverrid native to Southeast Asia.

Large-toothed hairy-tailed rat
Large-toothed hairy-tailed rat
Batomys dentatus

The large-toothed hairy-tailed rat (*Batomys dentatus*) is one of five species of rodent in the genus *Batomys*. It is in the diverse family Muridae. This species is found only in Philippines. Its natural habitat is subtropical or tropical dry forests.

Large-toothed shrew
Large-toothed shrew
Sorex macrodon

The large-toothed shrew or Mexican large-toothed shrew (*Sorex macrodon*) is one of 77 species within the genus *Sorex*. Registered on the IUCN Red List as vulnerable with a decreasing population, the Mexican large-toothed shrew has been recorded...

Lariang tarsier
Lariang tarsier
Tarsius lariang

The Lariang tarsier (*Tarsius lariang*) is a recently described tarsier occurring in the western part of the central core of Sulawesi. Six museum specimens of this species are known, two of which have been misidentified as the pygmy tarsier before...

Las Cajas water mouse
Las Cajas water mouse
Chibchanomys orcesi

The Las Cajas water mouse (*Chibchanomys orcesi*) is a species of rodent in the family Cricetidae. It is endemic to the Cajas National Park in Ecuador. Its natural habitat is the high grassland (páramo) of the park, where it lives near streams,...

Lasiurus atratus
Lasiurus atratus
Lasiurus atratus

Lasiurus atratus, the greater red bat, is a species of vesper bat. It is found in Guyana, Venezuela, French Guiana and Suriname.

Lataste's gerbil
Lataste's gerbil
Gerbillus latastei

Lataste's gerbil (*Gerbillus latastei*) is distributed mainly in Tunisia, Libya, and possibly Algeria. It is also known as the hairy-footed gerbil.

Latona's shrew
Latona's shrew
Crocidura latona

Latona's shrew (*Crocidura latona*) is a species of mammal in the family Soricidae. It is endemic to Democratic Republic of the Congo. Its natural habitat is subtropical or tropical moist lowland forest.

Laurie's moss mouse
Laurie's moss mouse
Pseudohydromys eleanorae

Laurie's moss mouse (*Pseudohydromys eleanorae*) is a species of mouse belonging to the family Muridae that is endemic to Papua New Guinea. It was first described in 2009.

LaVal's disk-winged bat

LaVal's disk-winged bat

Thyroptera lavalii

LaVal's disk-winged bat (*Thyroptera lavalii*) is a species of bat in the family Thyropteridae. It is native to Ecuador, Colombia, Peru, Venezuela and Brazil where it has been found near streams in tropical rainforest. The bat is insectivorous. It...

Lavaso dwarf lemur

Lavaso dwarf lemur

Cheirogaleus lavasoensis

The Lavaso dwarf lemur (*Cheirogaleus lavasoensis*) is a small, nocturnal strepsirrhine primate and a species of lemur that is endemic to three small, isolated patches of forest on the southern slopes of the Lavaso Mountains in southern...

Laxmann's shrew

Laxmann's shrew

Sorex caecutiens

Laxmann's shrew (*Sorex caecutiens*), or the masked shrew, is a species of shrew. Its range extends from northern Scandinavia and the Baltic to the Sea of Japan, including Hokkaidō, Sakhalin, and the Korean Peninsula. It favours mountain forests...

Layard's palm squirrel

Layard's palm squirrel

Funambulus layardi

Layard's palm squirrel or flame-striped jungle squirrel (*Funambulus layardi*) is a species of rodent in the family Sciuridae endemic to Sri Lanka. The validity of the subspecies *F. l. dravidianus* based on a single specimen from the southern tip...

Leach's single leaf bat

Leach's single leaf bat

Monophyllus redmani

Leach's single leaf bat (*Monophyllus redmani*), also known as Greater Antillean long-tongued bat, is a species of bat in the family Phyllostomidae. It is found in the southern Bahamas and in all the Greater Antilles (Cuba, Jamaica, Hispaniola...

Leadbeater's Possum

Leadbeater's Possum

Gymnobelideus leadbeateri

Leadbeater's possum (*Gymnobelideus leadbeateri*) is a critically endangered possum largely restricted to small pockets of alpine ash, mountain ash, and snow gum forests in the Central Highlands of Victoria, Australia, north-east of Melbourne. It...

Leaf Muntjac

Leaf Muntjac

Muntiacus putaoensis

The Leaf muntjac (*Muntiacus putaoensis*) is a small species of barking deer. It was documented in 1997 by biologist Alan Rabinowitz during his field study in the isolated Nogmung Township in Myanmar.

Least big-eared bat

Least big-eared bat

Neonycteris pusilla

The least big-eared bat (*Neonycteris pusilla*) is a South American bat species of the family Phyllostomidae, found in northwestern Brazil and eastern Colombia. It is monotypic within its genus.

Least chipmunk

Least chipmunk

Tamias minimus

The least chipmunk (*Neotamias minimus*) is the smallest species of chipmunk and the most widespread in North America.

Least dwarf shrew

Least dwarf shrew

Suncus infinitesimus

The least dwarf shrew (*Suncus infinitesimus*) is a species of mammal in the family Soricidae. It is found in Cameroon, Central African Republic, Democratic Republic of the Congo, Kenya, Nigeria, South Africa, Eswatini, Tanzania, and Uganda. Its...

Least forest mouse

Least forest mouse

Apomys musculus

The least forest mouse (*Apomys musculus*) is a species of rodent in the family Muridae. It is found only in the Philippines.

Least gerbil

Least gerbil

Gerbillus pusillus

The least gerbil (*Gerbillus pusillus*) is distributed mainly in South Sudan, southwestern Ethiopia, Kenya, and Tanzania.

Least groove-toothed swamp rat

Least groove-toothed swamp rat

Pelomys minor

The least groove-toothed swamp rat (*Pelomys minor*) is a species of rodent in the family Muridae. It is found in Angola, Democratic Republic of the Congo, Tanzania, and Zambia. Its natural habitat is moist savanna.

Least horseshoe bat

Least horseshoe bat

Rhinolophus pusillus

The least horseshoe bat (*Rhinolophus pusillus*) is a species of bat in the family Rhinolophidae. It is found in Cambodia, China, India, Indonesia, Laos, Malaysia, Myanmar, Nepal, Thailand and Vietnam. It is a food source of the parasite...

Least long-fingered bat

Least long-fingered bat

Miniopterus minor

The least long-fingered bat (*Miniopterus minor*) is a species of vesper bat in the family Miniopteridae. It can be found in the following countries: Republic of the Congo, Democratic Republic of the Congo, Kenya, São Tomé and Príncipe, and Tanzania.

Least pipistrelle

Least pipistrelle

Pipistrellus tenuis

The least pipistrelle (*Pipistrellus tenuis*) is a species of vesper bat.

Least pygmy squirrel

Least pygmy squirrel

Exilisciurus exilis

The least pygmy squirrel (*Exilisciurus exilis*), also known as the plain pygmy squirrel, is a species of rodent in the family Sciuridae. This plain olive-brown squirrel is endemic to forests, mostly below an altitude of 750 m (2,500 ft) but...
Least shrew tenrec
Least shrew tenrec
Microgale pusilla

The least shrew tenrec (*Microgale pusilla*) is a species of mammal in the family Tenrecidae. It is endemic to Madagascar. Its natural habitats are subtropical or tropical moist lowland forest, subtropical or tropical moist montane forest, swamps, ...
Least soft-furred mouse
Least soft-furred mouse
Praomys minor

The least soft-furred mouse or least praomys (*Praomys minor*) is a species of rodent in the family Muridae. It is found only in Democratic Republic of the Congo. Its natural habitat is subtropical or tropical moist lowland forest.
Least Weasel
Least Weasel
Mustela nivalis

The Least weasel (*Mustela nivalis*) lives deep in the Northeast Asian deciduous forest. It hunts day and night, aided by its sharp nails and long slender body. It is the world's smallest carnivore. According to Blackfoot legend, this species is...
Least woolly bat
Least woolly bat
Kerivoula minuta

The least woolly bat (*Kerivoula minuta*) is a species of vesper bat in the family Vespertilionidae. It is found in Malaysia and Thailand.
Least yellow bat
Least yellow bat
Rhogeessa mira

The least yellow bat (*Rhogeessa mira*) is a species of vesper bat. It is found only in Mexico.
Lechwe
Lechwe
Kobus leche

The lechwe (*Kobus leche*) is an antelope found in south-central Africa. These graceful animals live in marshy areas where they are important herbivores of aquatic plants and use knee-deep water as protection from predators.
Lelwel hartebeest
Lelwel hartebeest
Alcelaphus buselaphus lelwel

The Lelwel hartebeest (*Alcelaphus buselaphus lelwel*), also known as Jackson's hartebeest, is an antelope native to Central African Republic, Chad, the Democratic Republic of the Congo, Ethiopia, Kenya, Sudan, Tanzania, and Uganda. The Lelwel...
Lemming vole
Lemming vole
Alticola lemminus

The lemming vole (*Alticola lemminus*) is a species of rodent in the family Cricetidae. It is found only in the Russian Federation.

Lemur-like ringtail possum
Lemur-like ringtail possum
Hemibelideus lemuroides

The lemuroid ringtail possum (*Hemibelideus lemuroides*), also known as the lemur-like ringtail possum or the brushy-tailed ringtail, is a truly singular member of the ringtail possum group. It was once thought that they were gliding possums...

Lenis woolly bat
Lenis woolly bat
Kerivoula lenis

The lenis woolly bat (*Kerivoula lenis*) is a species of bat in the family *Vespertilionidae*. It is found in South and Southeast Asia.

Leopard
Leopard
Panthera pardus

The leopard (*Panthera pardus*) is one of the five extant species in the genus *Panthera*, a member of the cat family, *Felidae*. The leopard is distinguished by its well-camouflaged fur, opportunistic hunting behavior, broad diet, strength, and its...

Leopard Cat
Leopard Cat
Prionailurus bengalensis

The Leopard cat (*Prionailurus bengalensis*) is a small wild cat native to continental South, Southeast, and East Asia. Leopard cat subspecies differ widely in fur color, tail length, skull shape, and size of carnassials. Archaeological evidence...

Leopard Seal
Leopard Seal
Hydrurga leptonyx

The Leopard seal (*Hydrurga leptonyx*) is the second largest species of seal in the Antarctic (after the Southern elephant seal). Its only natural predator is the orca. It is the only species in the genus *Hydrurga*. Its closest relatives are the...

Leopardus vorohuensis
Leopardus vorohuensis
Leopardus vorohuensis

Leopardus vorohuensis is an extinct species of feline. Fossils of the species were found in the Vorohué Formation of Argentina and dated to the Uquian age of the early middle Pleistocene. It is the earliest known cat of the ocelot lineage of...

Leptomys arfakensis
Leptomys arfakensis
Leptomys arfakensis

Leptomys arfakensis is a species of rodent in the family of *Muridae*. It is found in the Arfak Mountains of West Papua, Indonesia.

Leptomys paulus
Leptomys paulus
Leptomys paulus

Leptomys paulus is a small rodent belonging to the family *Muridae* endemic to New Guinea.

Leschenault's rousette
Leschenault's rousette
Rousettus leschenaultii

Leschenault's rousette (*Rousettus leschenaultii*) is a species of fruit bat. The scientific name of the species was first published by Desmarest in 1820.

Lessa's tuco-tuco

Lessa's tuco-tuco

Ctenomys lessai

Lessa's Tuco-Tuco (*Ctenomys lessai*) is a species of tuco-tuco native to Bolivia. Only found near Lluthu Pampa, Cochabamba Department, at elevations of around 2,500 to 2,750 meters, the species measures about 255 millimeters in length and has...

Lesser Angolan epauletted fruit bat

Lesser Angolan epauletted fruit bat

Epomophorus grandis

The lesser Angolan epauletted fruit bat (*Epomophorus grandis*) is a species of megabat in the family Pteropodidae. It is found in Angola and Republic of the Congo. Its natural habitats are subtropical or tropical dry forest, subtropical or...

Lesser Antillean funnel-eared bat

Lesser Antillean funnel-eared bat

Natalus stramineus stramineus

The Lesser Antillean funnel-eared bat (*Natalus stramineus stramineus*) is a subspecies of the Mexican funnel-eared bat (*Natalus stramineus*) found to be endemic to the Lesser Antilles islands of the Caribbean. The subspecies was first described by...

Lesser Asiatic yellow bat

Lesser Asiatic yellow bat

Scotophilus kuhlii

The lesser Asiatic yellow bat (*Scotophilus kuhlii*) is a species of vesper bat. It is found in Bangladesh, India, Indonesia, Malaysia, Pakistan, the Philippines, Sri Lanka, and Taiwan.

Lesser bamboo bat

Lesser bamboo bat

Tylonycteris pachypus

The lesser bamboo bat or lesser flat-headed bat (*Tylonycteris pachypus*) is one of the smallest species of vesper bat, and is native to Southeast Asia.

Lesser bamboo rat

Lesser bamboo rat

Cannomys badius

The lesser bamboo rat (*Cannomys badius*) is a species of rodent in the family Spalacidae. It is monotypic within the genus *Cannomys*. It is found in Bangladesh, Bhutan, Cambodia, China, India, Myanmar, Nepal, and Thailand.

Lesser Bandicoot Rat

Lesser Bandicoot Rat

Bandicota bengalensis

The Lesser bandicoot rat (*Bandicota bengalensis*) is a giant rat of Southern Asia, not related to the true bandicoots which are marsupials. It is considered a pest in the cereal crops and gardens of India and Sri Lanka and emits piglike grunts...

Lesser Blind Mole-Rat

Lesser Blind Mole-Rat

Spalax leucodon

The Lesser blind mole-rat (*Nannospalax leucodon*) is a species of rodent found in Europe and a small part of Western Asia.

Lesser brown horseshoe bat
Lesser brown horseshoe bat
Rhinolophus stheno

The lesser brown horseshoe bat (*Rhinolophus stheno*) is a species of bat in the family Rhinolophidae. It is found in Indonesia, Laos, Malaysia, Thailand, and Vietnam.

Lesser bulldog bat
Lesser bulldog bat
Noctilio albiventris

The lesser bulldog bat (*Noctilio albiventris*) is an insectivorous and occasionally carnivorous bat of the (Neotropics), ranging through Central America and northern South America. Some unique characteristics of the bat include, large feet that...

Lesser cane rat
Lesser cane rat
Thryonomys gregorianus

The lesser cane rat (*Thryonomys gregorianus*) is a species of rodent in the family Thryomyidae. It is found in Cameroon, Chad, Democratic Republic of the Congo, Ethiopia, Kenya, Malawi, South Sudan, Tanzania, Uganda, Zambia, Zimbabwe, and...

Lesser Capybara
Lesser Capybara
Hydrochoerus isthmus

The lesser capybara (*Hydrochoerus isthmus*) is a large semiaquatic rodent of the family Caviidae found in eastern Panama, northwestern Colombia, and western Venezuela. The lesser capybara was described as a species in 1912, but was later...

Lesser Congo shrew
Lesser Congo shrew
Congosorex verheyeni

The lesser Congo shrew (*Congosorex verheyeni*) is a species of mammal in the family Soricidae found in Cameroon, the Central African Republic, the Republic of the Congo, and Gabon. Its natural habitat is subtropical or tropical moist lowland forest.

Lesser dog-like bat
Lesser dog-like bat
Peropteryx macrotis

The lesser dog-like bat (*Peropteryx macrotis*), also known as Peters' sac-winged bat, is a species of bat from South and Central America. First described in 1826, it was renamed in 1843 because the original scientific name was already in use for...

Lesser dwarf shrew
Lesser dwarf shrew
Suncus varilla

The lesser dwarf shrew (*Suncus varilla*) is a species of mammal in the family Soricidae. It is found in Botswana, Democratic Republic of the Congo, Lesotho, Malawi, Mozambique, South Africa, Tanzania, Zambia, and Zimbabwe.

Lesser Egyptian gerbil
Lesser Egyptian gerbil
Gerbillus gerbillus

The lesser Egyptian gerbil (*Gerbillus gerbillus*) is a small species of rodent in the family Muridae. It is native to North Africa and the Sinai Peninsula, where it lives in sandy habitats. It is a common species, and the International Union for...
Lesser Egyptian Jerboa

Lesser Egyptian Jerboa
Jaculus jaculus

The Lesser Egyptian jerboa (*Jaculus jaculus*) is a small rodent that lives in Africa and the Middle East. Jerboas look somewhat like miniature kangaroos and have some external similarities. Jerboas also move around in a similar manner to...

Lesser false vampire bat
Lesser false vampire bat
Megaderma spasma

The lesser false vampire bat (*Megaderma spasma*) is a bat found in South Asia and Southeast Asia from Sri Lanka and India in the west to Indonesia and the Philippines in the east. They live in caves and tree hollows. They are insectivorous. The...

Lesser fat-tailed jerboa
Lesser fat-tailed jerboa
Pygeretmus platyurus

The lesser fat-tailed jerboa (*Pygeretmus platyurus*) is a species of rodent in the family Dipodidae. It is endemic to West, Central, and Eastern Kazakhstan, as well as Northwestern Turkmenistan.

Lesser flying fox
Lesser flying fox
Pteropus mahaganus

The lesser flying fox or Sanborn's flying fox (*Pteropus mahaganus*) is a species of flying fox in the family Pteropodidae. It is found in Papua New Guinea and the Solomon Islands. The species is poorly known, but is believed to inhabit coastal...

Lesser forest shrew
Lesser forest shrew
Sylvisorex oriundus

The lesser forest shrew (*Sylvisorex oriundus*) is a species of mammal in the family Soricidae endemic to northeastern Democratic Republic of the Congo, where its type locality is Medje. Its natural habitat is subtropical or tropical moist lowland...

Lesser ghost bat
Lesser ghost bat
Diclidurus scutatus

The lesser ghost bat (*Diclidurus scutatus*) is a bat species found in South America. It is one of six bat species worldwide to have white fur.

Lesser gray-brown musk shrew
Lesser gray-brown musk shrew
Crocidura silacea

The lesser gray-brown musk shrew (*Crocidura silacea*) is a species of mammal in the family Soricidae. It is found in Angola, Malawi, Mozambique, South Africa, Swaziland, Tanzania, Uganda, Zambia, and Zimbabwe. Its natural habitats are dry...

Lesser great leaf-nosed bat
Lesser great leaf-nosed bat
Hipposideros turpis

The lesser great leaf-nosed bat or lesser roundleaf bat (*Hipposideros turpis*) is a species of bat in the family Hipposideridae. It is found in Japan. Its natural habitat is temperate forests. It is threatened by habitat loss.

Lesser Grison
Lesser Grison
Galictis cuja

The Lesser grison (*Galictis cuja*) is a species of mustelid from South America. Its common name 'grison' comes from French language and means "gray".

Lesser hairy-footed dunnart
Lesser hairy-footed dunnart
Sminthopsis youngsoni

The lesser hairy-footed dunnart (*Sminthopsis youngsoni*) is a small carnivorous Australian marsupial of the family *Dasyuridae*. It is a widespread and fairly common species, being found in many desert areas of Western Australia, Northern Territory...

Lesser hairy-winged bat
Lesser hairy-winged bat
Harpiocephalus harpia

The lesser hairy-winged bat (*Harpiocephalus harpia*) is a species of vesper bat in the family *Vespertilionidae*. It can be found in India, Indonesia, the Philippines, and Taiwan. They are known to eat various species of beetles. It is the only...

Lesser hamster-rat
Lesser hamster-rat
Beamys hindei

The lesser hamster-rat or long-tailed pouched rat (*Beamys hindei*) is a species of rodent in the family *Nesomyidae*. It is found in Kenya and Tanzania. Its natural habitat is subtropical or tropical moist lowland coastal forests. It is threatened...

Lesser Hedgehog Tenrec
Lesser Hedgehog Tenrec
Echinops telfairi

The lesser hedgehog tenrec (*Echinops telfairi*) is a species of mammal in the family *Tenrecidae*. It is the only species in the genus *Echinops* and is named in honour of Charles Telfair.

Lesser Horseshoe Bat
Lesser Horseshoe Bat
Rhinolophus hipposideros

The Lesser horseshoe bat (*Rhinolophus hipposideros*) is a small insectivorous bat found in Europe, Asia, and North Africa. It is closely related to its larger cousin, the Greater horseshoe bat. As with all horseshoe bats, the species gets its name...

Lesser iron-gray dwarf lemur
Lesser iron-gray dwarf lemur
Cheirogaleus minusculus

The lesser iron-gray dwarf lemur (*Cheirogaleus minusculus*), or large iron-gray dwarf lemur, is a small nocturnal lemur endemic to Madagascar. It is nocturnal and an arboreal quadruped, and is not very agile. It produces three types of...

Lesser Kudu
Lesser Kudu
Tragelaphus imberbis

The lesser kudu (*Tragelaphus imberbis*) is a forest antelope found in East Africa. It is placed in the genus *Tragelaphus* and family *Bovidae*. It was first scientifically described by the English zoologist Edward Blyth in 1869. The head-and-body...

Lesser large-footed bat
Lesser large-footed bat
Myotis hasseltii

The lesser large-footed bat (*Myotis hasseltii*) is a species of vesper bat. It can be found in Cambodia, India, Indonesia, Malaysia, Myanmar, Sri Lanka, Thailand, and Vietnam.

Lesser large-headed shrew

Lesser large-headed shrew

Paracrocidura schoutedeni

The lesser large-headed shrew (*Paracrocidura schoutedeni*) is a species of shrew in the family Soricidae. It is found in Cameroon, Central African Republic, Republic of the Congo, Democratic Republic of the Congo, Equatorial Guinea, and Gabon....

Lesser long-fingered bat

Lesser long-fingered bat

Miniopterus fraterculus

The lesser long-fingered bat (*Miniopterus fraterculus*), also known as the black clinging bat or lesser bent-winged bat, is a species of vesper bat in the family Vespertilionidae. It is found in western Southern Africa, south East Africa, and...

Lesser Long-Nosed Bat

Lesser Long-Nosed Bat

Leptonycteris yerbabuenae

The Lesser long-nosed bat (*Leptonycteris yerbabuenae*) is a medium-sized bat found in Central and North America. Enthusiasts for the bats often refer to Lesser long-nosed bats simply as leptos because they are the best-known members of the genus...

Lesser long-tailed shrew tenrec

Lesser long-tailed shrew tenrec

Microgale longicaudata

The lesser long-tailed shrew tenrec (*Microgale longicaudata*) is a species of mammal in the family Tenrecidae. It is active at all hours of the day and night, but each individual maintains its own pattern of rest and activity.

Lesser long-tongued bat

Lesser long-tongued bat

Choeroniscus minor

The lesser long-tongued bat (*Choeroniscus minor*), also called the lesser long-tailed bat, is a bat species from South America.

Lesser Mouse-Deer

Lesser Mouse-Deer

Tragulus kanchil

The lesser mouse-deer, lesser Malay chevrotain, or kanchil (*Tragulus kanchil*) is a species of even-toed ungulate in the family Tragulidae.

Lesser mouse-eared bat

Lesser mouse-eared bat

Myotis blythii

The lesser mouse-eared bat (*Myotis blythii*) is a species of insectivorous bat in the family Vespertilionidae.

Lesser mouse-tailed bat

Lesser mouse-tailed bat

Rhinopoma hardwickii

The lesser mouse-tailed bat (*Rhinopoma hardwickii*) is a species of microbat in the family Rhinopomatidae. Also referred to as Hardwicke's lesser mouse-tailed bat and long-tailed bat, it is named after Major General Thomas Hardwicke (1755-1835),...

Lesser musky fruit bat

Lesser musky fruit bat

Ptenochirus minor

The lesser musky fruit bat (*Ptenochirus minor*) is a species of megabat in the family Pteropodidae. It is endemic to the Philippines.

Lesser naked bat

Lesser naked bat

Cheiromeles parvidens

The lesser naked bat (*Cheiromeles parvidens*) is a species of bat in the family Molossidae, the free-tailed bats. It is native to Indonesia and the Philippines. This is a little-known species and its population status is unclear. It is known to...

Lesser naked-backed fruit bat

Lesser naked-backed fruit bat

Dobsonia minor

The lesser naked-backed fruit bat (*Dobsonia minor*) is a species of megabat in the family Pteropodidae. It is found in Indonesia and Papua New Guinea.

Lesser noctule

Lesser noctule

Nyctalus leisleri

The lesser noctule, Leisler's bat or the Irish bat (*Nyctalus leisleri*), is a species of insectivorous bat belonging to the vesper bat family, Vespertilionidae. The species was named to honour the naturalist Johann Philipp Achilles Leisler

Lesser Papuan pipistrelle

Lesser Papuan pipistrelle

Pipistrellus papuanus

The lesser Papuan pipistrelle (*Pipistrellus papuanus*) is a species of vesper bat. It can be found in Indonesia and Papua New Guinea.

Lesser pygmy flying squirrel

Lesser pygmy flying squirrel

Petaurillus emiliae

The lesser pygmy flying squirrel (*Petaurillus emiliae*) is a species of rodent in the family Sciuridae. It is endemic to Malaysia. It is listed as Data Deficient because it is known only from the type specimen collected in 1901. Adequate surveys...

Lesser ranee mouse

Lesser ranee mouse

Haeromys pusillus

The lesser ranee mouse (*Haeromys pusillus*) is a species of rodent in the family Muridae. It is found on the island of Borneo and the Palawan region (Philippines). Its natural habitat is subtropical or tropical dry forests.

Lesser red musk shrew

Lesser red musk shrew

Crocidura hirta

The lesser red musk shrew (*Crocidura hirta*) is a species of mammal in the family Soricidae. It is found in Angola, Botswana, Democratic Republic of the Congo, Malawi, Mozambique, Namibia, South Africa, Eswatini, Tanzania, Zambia, and Zimbabwe....

Lesser ricefield rat

Lesser ricefield rat

Rattus losea

The lesser ricefield rat (*Rattus losea*) is a rodent in the family Muridae. The species was first described by Robert Swinhoe in 1871. It is found in China, Laos,

Taiwan, Thailand, and Vietnam.
Lesser rock shrew
Lesser rock shrew
Crocidura serezhkyensis

The lesser rock shrew (*Crocidura serezhkyensis*) is a species of mammal in the family Soricidae. It is found in Azerbaijan, Kazakhstan, Tajikistan, and Turkey.
Lesser Ryukyu shrew
Lesser Ryukyu shrew
Crocidura watasei

The lesser Ryukyu shrew, or Watase's shrew (*Crocidura watasei*) is a common species of shrews that is endemic to Japan. It is often found living in bushes and grasslands along the river banks and in shrubs of lower elevations.
Lesser sac-winged bat
Lesser sac-winged bat
Saccopteryx leptura

The lesser sac-winged bat or lesser white-lined bat (*Saccopteryx leptura*) is a bat species of the family Emballonuridae from South and Middle America.
Lesser sheath-tailed bat
Lesser sheath-tailed bat
Emballonura monticola

The lesser sheath-tailed bat (*Emballonura monticola*) is a species of sac-winged bat in the family Emballonuridae. It is found in the Malay Peninsula (including Myanmar and Thailand), Borneo (both Indonesia and Malaysia), and many other parts of...
Lesser Short-Nosed Fruit Bat
Lesser Short-Nosed Fruit Bat
Cynopterus brachyotis

The Lesser short-nosed fruit bat (*Cynopterus brachyotis*) is a small bat that lives in Asia. These bats live in many types of habitats and are considered to be crop pests since they consume and damage fruit.
Lesser short-tailed gerbil
Lesser short-tailed gerbil
Dipodillus simoni

The lesser short-tailed gerbil (*Dipodillus simoni*) is distributed mainly from eastern Morocco to Egypt. It is also known as Simon's dipodil. After morphological and molecular studies in 2010 *Dipodillus* was ranged as a subgenus of *Gerbillus*, and...
Lesser small-toothed rat
Lesser small-toothed rat
Macruromys elegans

The lesser small-toothed rat or western small-toothed rat (*Macruromys elegans*) is a species of rodent in the family Muridae. It is found only in West Papua, Indonesia.
Lesser spear-nosed bat
Lesser spear-nosed bat
Phyllostomus elongatus

The lesser spear-nosed bat (*Phyllostomus elongatus*) is a bat species from South America. It is found in Bolivia, Brazil, Colombia, Ecuador, Guyana, Peru, Suriname and Venezuela.
Lesser Spot-Nosed Monkey
Lesser Spot-Nosed Monkey

Cercopithecus petaurista

The lesser spot-nosed monkey, lesser spot-nosed guenon, lesser white-nosed guenon, or lesser white-nosed monkey (*Cercopithecus petaurista*) is a species of primate in the family Cercopithecidae. It is found in Ivory Coast, Ghana, Guinea, Liberia,...

Lesser striped shrew

Lesser striped shrew

Sorex bedfordiae

The lesser striped shrew (*Sorex bedfordiae*) is a species of mammal in the family Soricidae. It is found in China, Myanmar, and Nepal.

Lesser Taiwanese shrew

Lesser Taiwanese shrew

Chodsigoa sodalis

The lesser Taiwanese shrew (*Chodsigoa sodalis*) is a rare species of shrew in the Soricomorpha order.

Lesser tree mouse

Lesser tree mouse

Chiruromys vates

The lesser tree mouse (*Chiruromys vates*) is a species of rodent in the family Muridae found only in Papua New Guinea.

Lesser tube-nosed bat

Lesser tube-nosed bat

Nyctimene minutus

The lesser tube-nosed bat (*Nyctimene minutus*) is a species of megabat in the family Pteropodidae. It is native to two of the Maluku Islands (Spice Islands) in northern Indonesia.

Lesser tube-nosed fruit bat

Lesser tube-nosed fruit bat

Paranyctimene raptor

The lesser tube-nosed fruit bat or unstriped tube-nosed bat (*Paranyctimene raptor*) is a species of megabat in the family Pteropodidae found in West Papua, Indonesia and Papua New Guinea.

Lesser tufted-tailed rat

Lesser tufted-tailed rat

Eliurus minor

The lesser tufted-tailed rat (*Eliurus minor*) is a species of rodent in the family Nesomyidae. It is found only in Madagascar.

Lesser white-toothed shrew

Lesser white-toothed shrew

Crocidura suaveolens

The lesser white-toothed shrew (*Crocidura suaveolens*) is a tiny shrew with a widespread distribution in Africa, Asia and Europe. Its preferred habitat is scrub and gardens and it feeds on insects, arachnids, worms, gastropods, newts and small...

Lesser Wilfred's mouse

Lesser Wilfred's mouse

Juliomys pictipes

The lesser Wilfred's mouse (*Juliomys pictipes*) is a species of South American rodents of the family Cricetidae. It was first described by Wilfred H. Osgood under the name *Thomasomys pictipes*, then into the genus *Wilfredomys*, and now known as...

Lesser woolly bat

Lesser woolly bat
Kerivoula lanosa

The lesser woolly bat (*Kerivoula lanosa*) is a species of vesper bat in the family Vespertilionidae. It is found in Botswana, Central African Republic, Democratic Republic of the Congo, Ivory Coast, Ethiopia, Gabon, Ghana, Guinea, Kenya, Liberia,...

Lesser woolly horseshoe bat
Lesser woolly horseshoe bat
Rhinolophus beddomei

The lesser woolly horseshoe bat (*Rhinolophus beddomei*), also called Beddome's horseshoe bat, is a species of bat in the family Rhinolophidae. It is found in India and Sri Lanka. Its natural habitats are subtropical or tropical moist lowland...

Lesser yellow bat
Lesser yellow bat
Scotophilus borbonicus

The lesser yellow bat (*Scotophilus borbonicus*) is a vesper bat found only on Madagascar and Réunion. On Réunion, it was considered common early in the 19th century, but was last sighted late in the 19th century. Only a single specimen attributed...

Lesser yellow-shouldered bat
Lesser yellow-shouldered bat
Sturnira nana

The lesser yellow-shouldered bat is a species of bat in the family Phyllostomidae. It is native to Peru and Ecuador. It is threatened by habitat loss.

Lesueur's hairy bat
Lesueur's hairy bat
Cistugo lesueurii

Lesueur's hairy bat (*Cistugo lesueurii*) also known as Lesueur's wing-gland bat is a species of bat in the family Cistugidae. It is found in dry savanna, Mediterranean-type shrubby vegetation, and hot deserts in Lesotho and South Africa. It is...

Lesula
Lesula
Cercopithecus lomamiensis

The lesula (*Cercopithecus lomamiensis*) is a species of Old World monkey in the guenon family, found in the Lomami Basin of the Congo. Though known to locals, it was unknown to the international scientific community until it was discovered in...

Levant mole
Levant mole
Talpa levantis

The Levant mole (*Talpa levantis*) is a species of mammal in the family Talpidae. It is found in Armenia, Azerbaijan, Georgia, Russia, and Turkey. There are two subspecies, *T. l. levantis* and *T. l. transcaucasica* (the Transcaucasian mole); the...

Lewis's tuco-tuco
Lewis's tuco-tuco
Ctenomys lewisi

Lewis's tuco-tuco (*Ctenomys lewisi*) is a species of rodent in the family Ctenomyidae. It is endemic to Bolivia.

Liberian Mongoose
Liberian Mongoose

Liberiictis kuhni

The Liberian mongoose (*Liberiictis kuhni*) is a mongoose species native to Africa. It was discovered in Liberia in 1958. Phylogenetic analysis shows it is closely related to other small, social mongooses and that the Banded mongoose is its closest...

Libyan jird

Libyan jird

Meriones libycus

The Libyan jird (*Meriones libycus*) is a species of rodent in the family Muridae. It is found in Mauritania, Morocco, Algeria, Tunisia, Libya, Egypt, Jordan, Syria, Saudi Arabia, Iran, Iraq, Afghanistan, Turkmenistan, Uzbekistan, Kazakhstan and...

Lichtenstein's Hartebeest

Lichtenstein's Hartebeest

Alcelaphus buselaphus lichtensteinii

Lichtenstein's hartebeest (*Alcelaphus buselaphus lichtensteinii*) is a subspecies of the hartebeest antelope. It is named in honor of the German zoologist Martin Lichtenstein.

Lichtenstein's jerboa

Lichtenstein's jerboa

Eremodipus lichtensteini

Lichtenstein's jerboa (*Eremodipus lichtensteini*) is a species of rodent in the family Dipodidae. It is monotypic within the genus *Eremodipus*. It is found in Kazakhstan, Turkmenistan, and Uzbekistan.

Liechtenstein's pine vole

Liechtenstein's pine vole

Microtus liechtensteini

Liechtenstein's pine vole (*Microtus liechtensteini*) is a species of rodent in the family Cricetidae. It is found in central and eastern Europe, from northern Italy through to Austria, Slovenia and Croatia.

Light-winged lesser house bat

Light-winged lesser house bat

Scotoecus albofuscus

The light-winged lesser house bat (*Scotoecus albofuscus*) is a species of vesper bat. It can be found in Benin, Democratic Republic of the Congo, Ivory Coast, Gambia, Ghana, Kenya, Malawi, Mozambique, Nigeria, Senegal, Sierra Leone, South Africa,...

Lima leaf-eared mouse

Lima leaf-eared mouse

Phyllotis limatus

The Lima leaf-eared mouse (*Phyllotis limatus*) or Lima pericote is a species of rodent in the family Cricetidae. It is found in a variety of habitats on the western slopes of the Andes from northern Chile to west-central Peru at elevations from...

Limestone rat

Limestone rat

Niviventer hinpoon

The limestone rat (*Niviventer hinpoon*) is a species of rodent in the family Muridae found only in the limestone karsts of Saraburi, Lopburi, Nakhon Sawan provinces, central Thailand. It is listed as an endangered species due to its highly...

Lindbergh's grass mouse

Lindbergh's grass mouse
Akodon lindberghi

Lindbergh's grass mouse (*Akodon lindberghi*) is a rodent species from South America. It is found in Brazil.

Linduan rousette
Linduan rousette
Rousettus linduensis

The Linduan rousette (*Rousettus linduensis*) is a species of megabat in the *Rousettus* genus of the family Pteropodidae. It is endemic to Indonesia and is known only from four specimens collected in the swamp forest of Lore Lindu National Park, in...

Lined pocket mouse
Lined pocket mouse
Chaetodipus lineatus

The lined pocket mouse (*Chaetodipus lineatus*) is a species of rodent in the family Heteromyidae. This pocket mouse is endemic to a small area of central Mexico.

Link rat
Link rat
Deomys ferrugineus

The link rat (*Deomys ferrugineus*) is a species of rodent in the family Muridae. It is also known by the common name Congo forest mouse. It is native to central Africa. It is 12-14.5 cm long with a 15-21 cm long tail. It weighs 40-70 g. It has...

Linnaeus's mouse opossum
Linnaeus's mouse opossum
Marmosa murina

Linnaeus's mouse opossum (*Marmosa murina*), also known as the common or murine mouse opossum, is a South American marsupial of the family Didelphidae.

Linnaeus's Two-Toed Sloth
Linnaeus's Two-Toed Sloth
Choloepus didactylus

Linnaeus's two-toed sloth (*Choloepus didactylus*), also known as the southern two-toed sloth, unau, or Linne's two-toed sloth is a species of sloth from South America, found in Venezuela, the Guyanas, Colombia, Ecuador, Peru, and Brazil north of...

Lion
Lion
Panthera leo

The lion (*Panthera leo*) is a large cat of the genus *Panthera* native to Africa and India. It is the most widely recognized animal symbol in human culture and it has been extensively depicted in sculptures and paintings, on national flags, and in...

Lion-Tailed Macaque
Lion-Tailed Macaque
Macaca silenus

The Lion-tailed macaque (*Macaca silenus*) is an Old World monkey. It is also known as the wanderoo and can be found only in the Western Ghats of South India.

Little bent-wing bat
Little bent-wing bat
Miniopterus australis

The little bent-wing bat or little long-fingered bat (*Miniopterus australis*) is a

species of vesper bat in the family Miniopteridae. It is found in Australia, Indonesia, the Philippines, and Vanuatu.

Little big-eared bat

Little big-eared bat

Micronycteris megalotis

The little big-eared bat (*Micronycteris megalotis*) is a bat species in the order Chiroptera and family Phyllostomidae. It is from South and Central America particularly Colombia, Venezuela, Guyana, French Guiana, Brazil, Peru, Ecuador, Bolivia,...

Little big-eyed bat

Little big-eyed bat

Chiroderma trinitatum

The little big-eyed bat (*Chiroderma trinitatum*) is a bat species from South and Central America.

Little black serotine

Little black serotine

Eptesicus andinus

The little black serotine (*Eptesicus andinus*) is a species of insectivorous vesper bat. It is found in Colombia, Ecuador, Peru, Venezuela, Bolivia and Brazil at elevations from 100 to 3300 m.

Little broad-nosed bat

Little broad-nosed bat

Scotorepens greyii

The little broad-nosed bat (*Scotorepens greyii*) (pronounced skoh'-toh-rep'-enz grey'-ee-ee') translates to Grey's darkness creeper. Sometimes called sometimes called Grey's broad-nosed after the third governor of South Australia Sir John Edward...

Little Brown Bat

Little Brown Bat

Myotis lucifugus

The Little brown bat (*Myotis lucifugus*) is an endangered species of mouse-eared microbat found in North America. It is similar in appearance to several other mouse-eared bats, including the Indiana bat, Northern long-eared bat, and Arizona...

Little collared fruit bat

Little collared fruit bat

Myonycteris torquata

The little collared fruit bat (*Myonycteris torquata*) is a species of megabat in the family Pteropodidae found in Angola, Cameroon, the Central African Republic, the Republic of the Congo, the Democratic Republic of the Congo, Ivory Coast,...

Little desert pocket mouse

Little desert pocket mouse

Chaetodipus arenarius

The little desert pocket mouse (*Chaetodipus arenarius*) is a species of small rodent in the family Heteromyidae. It is endemic to Baja California in Mexico.

Little forest bat

Little forest bat

Vespadelus vulturnus

The little forest bat (*Vespadelus vulturnus*) is a species of vesper bat in the family Vespertilionidae. It is found only in south-eastern Australia, including Tasmania. It is a tiny bat often weighing less than 4 g (0.14 oz) (males in some

areas...

Little free-tailed bat

Little free-tailed bat

Chaerephon pumilus

The little free-tailed bat (*Chaerephon pumilus*) is a species of the genus *Chaerephon* in the family *Molossidae*. It is widely distributed across Africa and islands around the continent.

Little goblin bat

Little goblin bat

Mormopterus minutus

The little goblin bat (*Mormopterus minutus*) is a species of bat in the family *Molossidae*, the free-tailed bats. It is endemic to Cuba. This bat is vulnerable due to habitat loss and degradation. During the day it can be found roosting in the palm...

Little golden-mantled flying fox

Little golden-mantled flying fox

Pteropus pumilus

The little golden-mantled flying fox (*Pteropus pumilus*) is a species of bat in the family *Pteropodidae*. It is found in Indonesia and the Philippines. Its natural habitat is subtropical or tropical dry forests. It is threatened by hunting and...

Little ground squirrel

Little ground squirrel

Spermophilus pygmaeus

The little ground squirrel or little souslik, (*Spermophilus pygmaeus*) is a species of rodent in the family *Sciuridae*. It is found from Eastern Europe to Central Asia. Its subspecies include *Spermophilus pygmaeus pygmaeus*, *Spermophilus pygmaeus*...

Little Indian field mouse

Little Indian field mouse

Mus booduga

The little Indian field mouse (*Mus booduga*) is a species of rodent in the family *Muridae*. It is found in Bangladesh, India, Myanmar, Nepal, and Sri Lanka.

Little Japanese horseshoe bat

Little Japanese horseshoe bat

Rhinolophus cornutus

The little Japanese horseshoe bat (*Rhinolophus cornutus*) is a species of bat in the family *Rhinolophidae*. It is found in Japan and possibly China. Its natural habitat is temperate forests. It is threatened by habitat loss. It has not yet been...

Little long-tailed dunnart

Little long-tailed dunnart

Sminthopsis dolichura

The Little long-tailed dunnart (*Sminthopsis dolichura*) is a dunnart that was, along with Gilbert's dunnart, described in 1984. The length from snout to tail is 150–200 mm of which head to anus is 65–50 mm and tail 85–105 mm long. Hind foot size...

Little margareta rat

Little margareta rat

Margaretamys parvus

The little margareta rat (*Margaretamys parvus*) is a species of rodent in the family *Muridae*. It is endemic to the island of Sulawesi in Indonesia.

Little native mouse

Little native mouse
Pseudomys delicatulus

The little native mouse (*Pseudomys delicatulus*), also known as the delicate mouse, is a species of rodent in the family Muridae. Kunwinjku of western Arnhem Land call this little creature Kijbuk (Reverend Peterson Nganjmirra, personal comment in...

Little Nepalese horseshoe bat
Little Nepalese horseshoe bat
Rhinolophus subbadius

The little Nepalese horseshoe bat (*Rhinolophus subbadius*) is a species of bat in the family Rhinolophidae. It is found in Bangladesh, China, India, Myanmar and Nepal, possibly in Bhutan.

Little pied bat
Little pied bat
Chalinolobus picatus

The little pied bat (*Chalinolobus picatus*) is a species of vesper bat in the family Vespertilionidae. It is found only in semi-arid woodlands in eastern Australia.

Little pocket mouse
Little pocket mouse
Perognathus longimembris

The little pocket mouse (*Perognathus longimembris*) is a species of rodent in the family Heteromyidae. It is found in Baja California and Sonora in Mexico and in Arizona, California, Idaho, Nevada, Oregon and Utah in the United States. Its...

Little red brocket
Little red brocket
Mazama rufina

The little red brocket or swamp brocket (*Mazama rufina*), also known as the Ecuador red brocket, is a small, little-studied deer native to the Andes of Colombia, Ecuador and northern Peru, where found in forest and páramo at altitudes between...

Little Red Flying Fox
Little Red Flying Fox
Pteropus scapulatus

The little red flying-fox (*Pteropus scapulatus*) is a megachiropteran bat native to northern and eastern Australia. The species weighs about half a kilogram, one US pound, and is the smallest species of *Pteropus* in mainland Australia. P....

Little Red Kaluta
Little Red Kaluta
Dasykaluta rosamondae

The Little red kaluta (*Dasykaluta rosamondae*) is a small marsupial that lives in the arid regions of northwest Australia. These animals are active at night and live for about four years in captivity.

Little soft-furred rat
Little soft-furred rat
Rattus mollicomulus

The little soft-furred rat (*Rattus mollicomulus*) is a species of rodent in the family Muridae. It is found only on the upper slopes of Mount Lampobattang in Bantaeng Regency, South Sulawesi, Indonesia. It gets its name from its soft, silky fur....

Little tube-nosed bat
Little tube-nosed bat
Murina aurata

The little tube-nosed bat (*Murina aurata*) is a species of bat. An adult little tube-nosed bat has a body length of 4.0-4.6 cm, a tail length of 2.8-3.6 cm, and a wing length of 3.0-3.3 cm. The species is found across South and East Asia, from...
Little white-shouldered bat
Little white-shouldered bat
Ametrida centurio

The little white-shouldered bat (*Ametrida centurio*) is a species of bat from South and Central America. It is the only species within its genus, the name of which translates as "reaper" or "destroyer".
Little wood mouse
Little wood mouse
Hylomyscus parvus

The lesser *hylomyscus* or little wood mouse (*Hylomyscus parvus*) is a species of rodent in the family Muridae. It is found in Cameroon, Central African Republic, Republic of the Congo, Democratic Republic of the Congo, Gabon, and possibly...
Little woolly mouse opossum
Little woolly mouse opossum
Marmosa phaea

The little woolly mouse opossum (*Marmosa phaea*) is a nocturnal, arboreal and mainly solitary South American marsupial of the family Didelphidae. It is native to the western slopes of the Andes in Colombia, Ecuador and Peru, where it lives at...
Little yellow bat
Little yellow bat
Rhogeessa parvula

The little yellow bat (*Rhogeessa parvula*) is a species of vesper bat found only in Mexico.
Little yellow-shouldered bat
Little yellow-shouldered bat
Sturnira lilium

The little yellow-shouldered bat (*Sturnira lilium*) is a bat species from South and Central America. This species is a frugivore and an effective seed disperser. It roosts alone in tree cavities, on branches, vines, and under palm leaves, usually...
Littledale's whistling rat
Littledale's whistling rat
Parotomys littledalei

Littledale's whistling rat (*Parotomys littledalei*) is one of two species of murid rodent in the genus *Parotomys*, the other being Brants's whistling rat (*Parotomys brantsii*). It is found in Namibia and South Africa. Its natural habitats are...
Livingstone's Fruit Bat
Livingstone's Fruit Bat
Pteropus livingstonii

Livingstone's fruit bat (*Pteropus livingstonii*) is an Old World fruit bat found only in the Union of the Comoros in the western Indian Ocean. It is the largest and rarest bat of all Comorian species.
Livingstone's house bat
Livingstone's house bat
Scotophilus livingstonii

Livingstone's yellow bat or Livingstone's house bat (*Scotophilus livingstonii*) is a species of bat found in Africa.

Llama
Llama
Lama glama

The llama(*Lama glama*) is a domesticated South American camelid. It is widely used as a meat and pack animal by Andean cultures since the Pre-Columbian era. The ancestors of llamas are thought to have originated from the Great Plains of North...

Llanos long-nosed armadillo
Llanos long-nosed armadillo
Dasypus sabanicola

The Llanos long-nosed armadillo (*Dasypus sabanicola*) or northern long-nosed armadillo is a species of armadillo in the family Dasypodidae. It is endemic to Colombia and Venezuela, where its habitat is the intermittently flooded grassland of the...

Lodgepole chipmunk
Lodgepole chipmunk
Neotamias speciosus

The Lodgepole chipmunk (*Neotamias speciosus*) is a species of rodent in the family Sciuridae. It is found in the U.S. state of California at elevations from 1,500 to 3,000 metres (4,900 to 9,800 ft). The Lodgepole chipmunk has a variety of common...

Lomami red colobus
Lomami red colobus
Piliocolobus parmentieri

The Lomami red colobus (*Piliocolobus parmentieri*) is a type of red colobus monkey from central Africa. Historically it had been treated as a subspecies of the Central African red colobus, (*P. foai*) but more recent taxonomies generally treat it...

Lombok flying fox
Lombok flying fox
Pteropus lombocensis

The Lombok flying fox (*Pteropus lombocensis*) is a species of megabat in the genus *Pteropus*. It is endemic to Indonesia. This species has been listed on Appendix II of CITES since 1990, along with most others in the genus *Pteropus*. It was...

Lonchophylla fornicata
Lonchophylla fornicata
Lonchophylla fornicata

Lonchophylla fornicata is a species of bat found in South America.

Lonchophylla inexpectata
Lonchophylla inexpectata
Lonchophylla inexpectata

Lonchophylla inexpectata is a species of leaf-nosed bat found in Brazil.

Lonchophylla orienticollina
Lonchophylla orienticollina
Lonchophylla orienticollina

Lonchophylla orienticollina is a species of bat found in Colombia, Venezuela, and Ecuador.

Long-Beaked Common Dolphin
Long-Beaked Common Dolphin
Delphinus capensis

The common dolphin (*Delphinus delphis*) is the most abundant cetacean in the world,

with a global population of about six million. Despite this fact and its vernacular name, the common dolphin is not thought of as the archetypal dolphin, with...

Long-clawed ground squirrel
Long-clawed ground squirrel
Spermophilopsis leptodactylus

The long-clawed ground squirrel (*Spermophilopsis leptodactylus*) is a squirrel species native to grasslands and deserts in northeastern Iran, Tajikistan, Turkmenistan, northwestern Afghanistan, Uzbekistan and Kazakhstan. It is the only member of...

Long-clawed mole vole
Long-clawed mole vole
Prometheomys schaposchnikowi

The long-clawed mole vole (*Prometheomys schaposchnikowi*) is a species of rodent in the family Cricetidae. It is the only species in the genus *Prometheomys*. It is found in Georgia and Turkey.

Long-clawed shrew
Long-clawed shrew
Sorex unguiculatus

The long-clawed shrew (*Sorex unguiculatus*) is a species of shrew. An adult long-clawed shrew has a weight of less than 20 g and a body length of 54–97 mm, with a tail of 40–53 mm. It is distributed through the uplands of northeastern Asia,...

Long-eared chipmunk
Long-eared chipmunk
Neotamias quadrimaculatus

The long-eared chipmunk (*Neotamias quadrimaculatus*), also called the Sacramento chipmunk or the four-banded chipmunk, is a species of rodent in the squirrel family, Sciuridae. It is endemic to the central and northern Sierra Nevada of California...

Long-eared flying mouse
Long-eared flying mouse
Idiurus macrotis

The long-eared flying mouse (*Idiurus macrotis*) or long-eared scaly-tailed flying squirrel, is a species of flying mouse from western and central Africa. It is not actually a squirrel, nor a mouse, though it is a rodent. Not much is known about...

Long-eared gymnure
Long-eared gymnure
Hylomys megalotis

The long-eared gymnure (*Hylomys megalotis*) is a eulipotyphlan that is found in Laos. This specific type of gymnure has long ears and a long skull compared to that of others. It is also recognized for its broad forefeet, stout claws, and naked...

Long-Eared Hedgehog
Long-Eared Hedgehog
Hemiechinus auritus

The long-eared hedgehog (*Hemiechinus auritus*) is a species of hedgehog native to Central Asian countries and some countries of the Middle East. The long-eared hedgehog lives in burrows that it either makes or finds and is distinguished by its...

Long-Eared Jerboa
Long-Eared Jerboa
Euchoreutes naso

The long-eared jerboa (*Euchoreutes naso*) is a nocturnal mouse-like rodent with a

long tail, long hind legs for jumping, and exceptionally large ears. It is distinct enough that authorities consider it to be the only member of both its genus,...

Long-eared myotis

Long-eared myotis

Myotis evotis

The long-eared myotis (*Myotis evotis*) is a species of vesper bat in the suborder Microchiroptera. It can be found in western Canada, the western United States, and Baja California in Mexico.

Long-fingered bat

Long-fingered bat

Myotis capaccinii

The long-fingered bat (*Myotis capaccinii*) is a carnivorous species of vesper bat. It is native to coastal areas around the Mediterranean Sea, as well as a few patches of land in western Iran. Due to the fact that its population is in decline, it...

Long-fingered triok

Long-fingered triok

Dactylopsila palpator

The long-fingered triok (*Dactylopsila palpator*) is a species of marsupial in the family Petauridae. It is found in West Papua, Indonesia and Papua New Guinea. It is known as blc in the Kalam language of Papua New Guinea.

Long-Finned Pilot Whale

Long-Finned Pilot Whale

Globicephala melas

Long-finned pilot whales (*Globicephala melas*) are large oceanic dolphins. Pilot whales get their name from the original belief that there was a "pilot" or lead individual in their groups. They have also earned the nickname of "pothead whale" in...

Long-footed potoroo

Long-footed potoroo

Potorous longipes

The long-footed potoroo (*Potorous longipes*) is a small marsupial found in southeastern Australia, restricted to an area around the coastal border between New South Wales and Victoria. It was first recorded in 1967 when an adult male was caught...

Long-footed rat

Long-footed rat

Tarsomys apoensis

The long-footed rat (*Tarsomys apoensis*) is a species of rodent in the family Muridae. It is found only in the highlands of Mindanao, Philippines, including Mount Apo, Mount Kitanglad, and Mount Malindang.

Long-footed shrew

Long-footed shrew

Crocidura crenata

The long-footed shrew (*Crocidura crenata*) is a species of mammal in the family Soricidae. It is found in Cameroon, Central African Republic, Republic of the Congo, Democratic Republic of the Congo, Equatorial Guinea, and Gabon. Its natural...

Long-footed treeshrew

Long-footed treeshrew

Tupaia longipes

The long-footed treeshrew (*Tupaia longipes*) is a treeshrew species within the Tupaiidae. It is endemic to Borneo and threatened due to deforestation and degradation of habitat.

Long-footed water rat

Long-footed water rat

Leptomys elegans

The long-footed water rat (*Leptomys elegans*) is a species of rodent in the family Muridae. It is found in the mountains of southern Papua New Guinea. Its natural habitat is subtropical or tropical dry forest.

Long-furred Atlantic tree-rat

Long-furred Atlantic tree-rat

Phyllomys medius

The long-furred Atlantic tree-rat or Thomas's Atlantic tree-rat (*Phyllomys medius*), is a spiny rat species from South America and is found in Brazil.

Long-Haired Rat

Long-Haired Rat

Rattus villosissimus

The Long-haired rat (*Rattus villosissimus*) is a rodent native to Australia. This species is well known for its population irruptions over vast areas of Australia due to which it received an alternative common name - the plague rat.

Long-haired rousette

Long-haired rousette

Rousettus lanosus

The long-haired fruit bat (*Stenonycteris lanosus*), also known as the long-haired rousette, is a species of megabat in the family Pteropodidae. It is the only member of the genus *Stenonycteris*. It was formerly classified in the genus *Rousettus*...

Long-headed hill rat

Long-headed hill rat

Bunomys prolatus

The long-headed hill rat (*Bunomys prolatus*) is a species of rodent in the family Muridae. It is found only in Sulawesi, Indonesia, where it is only known from Mount Tambusisi. Its natural habitat is subtropical or tropical dry forests. It is...

Long-legged bat

Long-legged bat

Macrophyllum macrophyllum

The long-legged bat (*Macrophyllum macrophyllum*) is a member of the Phyllostomidae family in the order Chiroptera. Both males and females of this species are generally small, with wingspans reaching 80mm with an average weight ranging between 6...

Long-legged myotis

Long-legged myotis

Myotis volans

The long-legged myotis (*Myotis volans*) is a species of vesper bat that can be found in western Canada, Mexico, and the western United States.

Long-Nosed Bandicoot

Long-Nosed Bandicoot

Perameles nasuta

The long-nosed bandicoot (*Perameles nasuta*), a marsupial, is a species of bandicoot found in eastern Australia, from north Queensland along the east coast to Victoria. Around 40 centimetres (16 in) long, it is sandy- or grey-brown with a long...

Long-nosed caenolestid
Long-nosed caenolestid
Rhyncholestes raphanurus

The long-nosed caenolestid (*Rhyncholestes raphanurus*), also known as the Chilean shrew opossum or long-nosed shrew opossum, is a shrew opossum that occurs in temperate forests of Argentina and southern Chile. It was first described by American...

Long-nosed dasyure
Long-nosed dasyure
Murexia naso

The long-nosed dasyure (*Murexia naso*) is a species of marsupial in the family Dasyuridae. It is found in Indonesia and Papua New Guinea. Its natural habitat is subtropical or tropical dry forests.

Long-nosed echymipera
Long-nosed echymipera
Echymipera rufescens

The long-nosed echymipera (*Echymipera rufescens*), or long-nosed spiny bandicoot, is a species of marsupial in the family Peramelidae. It is found in Australia, Indonesia, and Papua New Guinea. Its natural habitat is subtropical or tropical dry...

Long-nosed hociendo
Long-nosed hociendo
Oxymycterus nasutus

The long-nosed hociendo (*Oxymycterus nasutus*) is a South American rodent species found in southeastern Brazil and Uruguay.

Long-nosed Luzon forest mouse
Long-nosed Luzon forest mouse
Apomys sacobianus

The long-nosed Luzon forest mouse (*Apomys sacobianus*), also known as the Pinatubo volcano mouse is a species of rodent in the family Muridae. It is found only in the Philippines. Its natural habitat is subtropical or tropical dry forests. It is...

Long-nosed mole
Long-nosed mole
Euroscaptor longirostris

The long-nosed mole (*Euroscaptor longirostris*) is a species of mammal in the family Talpidae. It is endemic to China, where it has a widespread distribution. Orlov's mole (*E. orlovi*) and Kuznetsov's mole (*E. kuznetsovi*), both of which range...

Long-nosed mongoose
Long-nosed mongoose
Herpestes naso

The long-nosed mongoose (*Xenogale naso*) is a mongoose native to Central African wetlands and rainforests. It has been listed as Least Concern on the IUCN Red List since 1996.

Long-nosed mosaic-tailed rat
Long-nosed mosaic-tailed rat
Paramelomys levipes

The long-nosed mosaic-tailed rat or Papuan lowland paramelomys (*Paramelomys levipes*) is a species of rodent in the family Muridae. It is found only in Papua New Guinea.

Long-nosed paramelomys

Long-nosed paramelomys
Paramelomys naso

The long-nosed paramelomys (*Paramelomys naso*) is a species of rodent of the family Muridae endemic to New Guinea. It is found in the lowlands of the south of the country.

Long-Nosed Potoroo
Long-Nosed Potoroo
Potorous tridactylus

The long-nosed potoroo (*Potorous tridactylus*) is a species of potoroo. These small marsupials are part of the rat-kangaroo family. The long-nosed potoroo contains two subspecies, *P. t. tridactylus* from mainland Australia, and *P. t. apicalis* from...

Long-nosed short-tailed opossum
Long-nosed short-tailed opossum
Monodelphis scalops

The long-nosed short-tailed opossum (*Monodelphis scalops*), is an opossum species in Argentina and Brazil.

Long-snouted bat
Long-snouted bat
Platalina genovensium

The long-snouted bat (*Platalina genovensium*) is a species of bat in the family Phyllostomidae. It is monotypic within the genus *Platalina*. It is endemic to northern Peru and northern Chile. It feeds almost exclusively on the nectar and fruit of...

Long-tailed armored tree-rat
Long-tailed armored tree-rat
Makalata macrura

The long-tailed armored tree-rat, *Makalata macrura*, is a spiny rat species from South America. It is found in Brazil, with a population in Ecuador which is referable either to this species or to *Makalata didelphoides*. Initially considered a large...

Long-tailed birch mouse
Long-tailed birch mouse
Sicista caudata

The long-tailed birch mouse (*Sicista caudata*) is a species of birch mouse found in Northeast Asia. It has been reported from the Ussuri region of Manchuria, Sakhalin Island and Primorsky Krai in Russia, and northern North Korea. It is listed as...

Long-tailed brown-toothed shrew
Long-tailed brown-toothed shrew
Episoriculus leucops

The long-tailed brown-toothed shrew (*Episoriculus leucops*) is a species of mammal in the family Soricidae. It is found in China, India, Myanmar, Nepal, and Vietnam.

Long-Tailed Chinchilla
Long-Tailed Chinchilla
Chinchilla lanigera

The long-tailed chinchilla (*Chinchilla lanigera*), also called the Chilean, coastal, common, or lesser chinchilla, is one of two species of rodent from the genus *Chinchilla* : the other species being *C. chinchilla*. Both species are endangered in...

Long-tailed dunnart
Long-tailed dunnart
Sminthopsis longicaudata

The long-tailed dunnart (*Sminthopsis longicaudata*) is an Australian dunnart that, like the little long-tailed dunnart, has a tail longer than its body. It is also one of the larger dunnarts at a length from snout to tail of 260–306 mm of which...
Long-tailed dwarf hamster
Long-tailed dwarf hamster
Cricetulus longicaudatus

The long-tailed dwarf hamster (*Cricetulus longicaudatus*) is a species of rodent in the family Cricetidae. It is found in China, Kazakhstan, Mongolia, and Russia.
Long-tailed forest shrew
Long-tailed forest shrew
Myosorex longicaudatus

The long-tailed forest shrew, or long-tailed mouse shrew, (*Myosorex longicaudatus*) is a species of mammal in the family Soricidae. It is endemic to South Africa, where its natural habitats are Mediterranean-type shrubby vegetation and swamps.
Long-tailed fruit bat
Long-tailed fruit bat
Notopteris macdonaldi

The long-tailed fruit bat, long-tailed blossom bat, or Fijian blossom bat (*Notopteris macdonaldi*) is a species of megabat in the family Pteropodidae. It is found in Fiji and Vanuatu. They roost as large colonies in caves and forage in a range of...
Long-tailed giant rat
Long-tailed giant rat
Leopoldamys sabanus

The long-tailed giant rat (*Leopoldamys sabanus*) is a species of rodent in the family Muridae. It is found in Bangladesh, Cambodia, Indonesia, Laos, Malaysia, Thailand, and Vietnam.
Long-Tailed Goral
Long-Tailed Goral
Naemorhedus caudatus

The long-tailed goral or Amur goral (*Naemorhedus caudatus*) is a species of ungulate of the family Bovidae found in the mountains of eastern and northern Asia, including Russia, China, and Korea. A population of this species exists in the Korean...
Long-tailed ground squirrel
Long-tailed ground squirrel
Urocitellus undulatus

The long-tailed ground squirrel or Eversmann's souslik (*Urocitellus undulatus*) is a species of rodent in the squirrel family Sciuridae. It is found in China, Kazakhstan, Mongolia, and Russia.
Long-tailed house bat
Long-tailed house bat
Eptesicus hottentotus

The long-tailed house bat or long-tailed serotine bat (*Eptesicus hottentotus*) is a species of vesper bat found in Kenya, Lesotho, Malawi, Mozambique, Namibia, South Africa, Zambia, and Zimbabwe. It is found in dry savanna, moist savanna,...
Long-Tailed Macaque
Long-Tailed Macaque
Macaca fascicularis

The Long-tailed macaque (*Macaca fascicularis*) is a cercopithecine primate native to

Southeast Asia. This species has a long history alongside humans. It has been alternately seen as an agricultural pest, a sacred animal, and, more recently, the...

Long-Tailed Marmot
Long-Tailed Marmot
Marmota caudata

The Long-tailed marmot (*Marmota caudata*) is a large ground squirrel that lives in mountainous regions in the central parts of Asia. As suggested by its name, this is a relatively long-tailed species of marmot.

Long-tailed mole
Long-tailed mole
Scaptonyx fusicaudus

The long-tailed mole (*Scaptonyx fusicauda*) is a species of mole in the family Talpidae. It is found in China, Vietnam and Myanmar.

Long-tailed mountain rat
Long-tailed mountain rat
Niviventer rapit

The long-tailed mountain rat (*Niviventer rapit*) is a species of rodent in the family Muridae. It is endemic to Borneo and found in Indonesia and Malaysia. Recorded at elevations of 940–3,360 m (3,080–11,020 ft) above sea level, it is a poorly...

Long-tailed mountain shrew
Long-tailed mountain shrew
Episoriculus macrurus

The long-tailed mountain shrew (*Episoriculus macrurus*) is a species of mammal in the family Soricidae. It is found in China, Myanmar, Nepal, and Vietnam.

Long-tailed mouse
Long-tailed mouse
Pseudomys higginsi

The long-tailed mouse (*Pseudomys higginsi*) is a native Australian mammal in the Order Rodentia and the Family Muridae. It is found only on the island of Tasmania. The long-tailed mouse is an omnivore that feeds on insects and a range of plants....

Long-tailed musk shrew
Long-tailed musk shrew
Crocidura dolichura

The long-tailed musk shrew (*Crocidura dolichura*) is a species of mammal in the family Soricidae. It is found in Burundi, Cameroon, Central African Republic, Republic of the Congo, Democratic Republic of the Congo, Gabon, Nigeria, and Uganda.

Long-Tailed Pangolin
Long-Tailed Pangolin
Phataginus tetradactyla

The Long-tailed pangolin (*Phataginus tetradactyla*), also called the African black-bellied pangolin, or Ipi, is a diurnal, arboreal pangolin species belonging to the family Manidae, in the order Pholidota. The common names for this species stem...

Long-tailed planigale
Long-tailed planigale
Planigale ingrami

The long-tailed planigale (*Planigale ingrami*), also known as Ingram's planigale or the northern planigale, is the smallest of all marsupials, and one of the smallest

of all mammals. It is rarely seen but is a quite common inhabitant of the...

Long-tailed pocket mouse

Long-tailed pocket mouse

Chaetodipus formosus

The long-tailed pocket mouse (*Chaetodipus formosus*) is a species of rodent in the family Heteromyidae. It is found in Arizona, California, Nevada and Utah in the United States and Baja California in Mexico.

Long-tailed porcupine

Long-tailed porcupine

Trichys fasciculata

The long-tailed porcupine (*Trichys fasciculata*) is a species of rodent in the family Hystricidae. It is monotypic within the genus *Trichys*, and is found in Brunei, Indonesia, and Malaysia.

Long-tailed pygmy possum

Long-tailed pygmy possum

Cercartetus caudatus

The long-tailed pygmy possum (*Cercartetus caudatus*) is a diprotodont marsupial found in the rainforests of northern Australia and New Guinea (Western New Guinea (Indonesia) and Papua New Guinea). Living at altitudes of above 1,500 m, it eats...

Long-tailed shrew

Long-tailed shrew

Sorex dispar

The long-tailed shrew or rock shrew (*Sorex dispar*) is a small North American shrew found in Atlantic Canada and the Northeastern United States. This shrew is slate grey in color with a pointed snout, a long tail, and lighter underparts. It is...

Long-tailed shrew rat

Long-tailed shrew rat

Tateomys macrocercus

The long-tailed shrew rat (*Tateomys macrocercus*) is a species of rodent in the family Muridae. It is found only in Central Sulawesi, Indonesia, where it is known only from Mount Nokilalaki in Sigi Regency.

Long-tailed spiny rat

Long-tailed spiny rat

Proechimys longicaudatus

The long-tailed spiny rat, *Proechimys longicaudatus*, is a spiny rat species from South America. It is found in Bolivia, Brazil and Paraguay.

Long-tailed Talaud mosaic-tailed rat

Long-tailed Talaud mosaic-tailed rat

Melomys talaudium

The long-tailed Talaud mosaic-tailed rat or the long-tailed Talaud melomys (*Melomys talaudium*) is a species of rodent in the family Muridae. It is endemic to Karakelong and Salebabu in the Talaud Islands in Indonesia where it occurs in forest...

Long-tailed vole

Long-tailed vole

Microtus longicaudus

The long-tailed vole (*Microtus longicaudus*), in some areas known as the San Bernardino long-tailed vole, is a small vole found in western North America. They have short ears and a long tail. Their fur is gray brown with light gray underparts....

Long-Tailed Weasel

Long-Tailed Weasel
Mustela frenata

The Long-tailed weasel is an endearing-looking species of mustelid common throughout America and southern Canada. Looking like a cute, curious, lively kitten, it is one of nature's most ferocious and relentless predators and is known as "nature's..."

Long-tongued arboreal mouse
Long-tongued arboreal mouse
Rhagomys longilingua

The long-tongued arboreal mouse (*Rhagomys longilingua*) is a South American rodent species of the family Cricetidae. It is found in a variety of habitats, including dense forest, in Bolivia and Peru at elevations from 450 to 2,100 metres (1,480...

Long-tongued fruit bat
Long-tongued fruit bat
Macroglossus sobrinus

The long-tongued fruit bat (*Macroglossus sobrinus*) is a species of megabat. It is nectarivorous, feeding on nectar from primarily banana flowers. It is found in several countries in South and Southeast Asia.

Long-Tongued Nectar Bat
Long-Tongued Nectar Bat
Macroglossus minimus

The long-tongued nectar bat (*Macroglossus minimus*), also known as the northern blossom bat, honey nectar bat, least blossom-bat, dagger-toothed long-nosed fruit bat, and lesser long-tongued fruit bat, is a species of megabat. *M. minimus* is one...

Long-toothed pipistrelle
Long-toothed pipistrelle
Hypsugo dolichodon

The long-toothed pipistrelle (*Hypsugo dolichodon*) is a species of bat of the genus *Hypsugo*. It is a small bat, with a length of 35.2–38.4 mm (1.38–1.51 in) of forearm, and 5.9–7 mm (0.23–0.27 in) of foot. It feeds on insects and has especially...

Long-winged tomb bat
Long-winged tomb bat
Taphozous longimanus

The long-winged tomb bat (*Taphozous longimanus*) is a species of sac-winged bat in the family Emballonuridae. It is found in Bangladesh, Cambodia, India, Indonesia, Malaysia, Myanmar, Singapore, Sri Lanka, and Thailand.

Lophuromys eisenrauti
Lophuromys eisenrauti
Lophuromys eisenrauti

Lophuromys eisenrauti, also known as Eisenraut's brush-furred rat or Mount Lefo brush-furred mouse, is a species of rodent in the family Muridae. It is only found in Mount Lefo, the western area of Cameroon.

Lord Derby's scaly-tailed squirrel
Lord Derby's scaly-tailed squirrel
Anomalurus derbianus

Lord Derby's scaly-tailed squirrel (*Anomalurus derbianus*) is an anomalurid rodent native to Africa. It was named after Edward Smith-Stanley, 13th Earl of Derby.

Lord Howe long-eared bat
Lord Howe long-eared bat

Nyctophilus howensis

The Lord Howe long-eared bat (*Nyctophilus howensis*) was a vespertilionid bat known only by a single specimen, a skull found on Lord Howe Island in 1972. A mammalian insectivorous species resembling the long-eared *Nyctophilus*, with an elongated...

Lorentz's mosaic-tailed rat
Lorentz's mosaic-tailed rat
Paramelomys lorentzii

Lorentz's mosaic-tailed rat (*Paramelomys lorentzii*) is a species of rodent in the family Muridae. It is found in West Papua, Indonesia and Papua New Guinea. The rat is named after Hendrikus Albertus Lorentz, a Dutch explorer who passed through...

Loring's rat
Loring's rat
Thallomys loringi

Loring's rat (*Thallomys loringi*) is a species of rodent in the family Muridae. It is found in Kenya and Tanzania. Its natural habitats are subtropical or tropical dry forests, dry savanna, and subtropical or tropical dry shrubland. The species is...

Lorrain dormouse
Lorrain dormouse
Graphiurus lorrainae

The Lorrain dormouse (*Graphiurus lorrainae*) is a species of rodent in the family Gliridae. It is found in Cameroon, Democratic Republic of the Congo, Ghana, Guinea-Bissau, Liberia, Nigeria, Sierra Leone, Tanzania, and Uganda. Its natural...

Louis's yellow-shouldered bat
Louis's yellow-shouldered bat
Sturnira luisi

Louis's yellow-shouldered bat (*Sturnira luisi*) is a species of bat in the family Phyllostomidae. It is found in Colombia, Costa Rica, Ecuador, Panama, and Peru.

Louise's spiny mouse
Louise's spiny mouse
Acomys louisae

Louise's spiny mouse (*Acomys louisae*) is a species of rodent in the family Muridae. It is found in Djibouti, Ethiopia, Kenya, and Somalia. Its natural habitats are dry savanna, subtropical or tropical dry lowland grassland, and rocky areas.

Louisiana black bear
Louisiana black bear
Ursus americanus luteolus

The Louisiana black bear (*Ursus americanus luteolus*), one of 16 subspecies of the American black bear, is found in parts of Louisiana, mainly along the Mississippi River Valley and the Atchafalaya River Basin. It was classified as 'threatened'...

Lovat's climbing mouse
Lovat's climbing mouse
Dendromus lovati

Lovat's climbing mouse (*Dendromus lovati*) is a species of rodent in the family Nesomyidae. It is found only in Ethiopia. Its natural habitat is subtropical or tropical high-altitude grassland. It is threatened by habitat loss.

Lovely-haired rat
Lovely-haired rat
Taeromys callitrichus

The lovely-haired rat (*Taeromys callitrichus*) is a species of rodent in the family Muridae. It is found only in Sulawesi, Indonesia.

Low's squirrel
Low's squirrel
Sundasciurus lowii

Low's squirrel (*Sundasciurus lowii*) is a species of rodent in the family Sciuridae. It is found in Borneo and nearby islets (Indonesia and Malaysia). Its former subspecies *S. l. robinsoni* from Sumatra and Malay Peninsula and *S. l. natunensis*...

Lowe's gerbil
Lowe's gerbil
Dipodillus lowei

Lowe's gerbil (*Dipodillus lowei*) is distributed mainly in Sudan; Jebel Marra. Less than 250 individuals of this species are thought to persist in the wild.

Lowe's mona monkey
Lowe's mona monkey
Cercopithecus lowei

Lowe's mona monkey (*Cercopithecus lowei*) is an Old World monkey in the family Cercopithecidae found from the Ivory Coast to Ghana. It was previously classified as a subspecies of Campbell's mona monkey *C. campbelli*.

Lowe's shrew
Lowe's shrew
Chodsigoa parca

The Lowe's shrew (*Chodsigoa parca*) is a species of mammal in the family Soricidae. It is found in China, Myanmar, Thailand, and Vietnam.

Lowland Anoa
Lowland Anoa
Bubalus depressicornis

Anoa, also known as dwarf buffalo and sapiutan, is a subgenus of *Bubalus* comprising two species endemic to the island of Sulawesi in Indonesia: the mountain anoa (*Bubalus quarlesi*) and the lowland anoa (*Bubalus depressicornis*). Both live in...

Lowland brush mouse
Lowland brush mouse
Pogonomelomys bruijnii

The lowland brush mouse (*Pogonomelomys bruijnii*) is a species of rodent in the family Muridae. It is found in Indonesia on the island of Sulawesi and on the Vogelkop Peninsula in Papua Province, Indonesia.

Lowland long-nosed squirrel
Lowland long-nosed squirrel
Hyosciurus ileile

The lowland long-nosed squirrel (*Hyosciurus ileile*) is a species of rodent in the family Sciuridae. It is endemic to Sulawesi, Indonesia. Its natural habitat is subtropical or tropical dry lowland grassland.

Lowland mosaic-tailed rat
Lowland mosaic-tailed rat
Paramelomys platyops

The lowland mosaic-tailed rat (*Paramelomys platyops*) is a species of rodent in the family Muridae. It is found in Indonesia and Papua New Guinea.

Lowland ringtail possum
Lowland ringtail possum
Pseudochirulus canescens

The lowland ringtail possum (*Pseudochirulus canescens*) is a species of marsupial

in the family Pseudocheiridae. It is found in Indonesia, the Solomon Islands, and Papua New Guinea.

Lowland Streaked Tenrec

Lowland Streaked Tenrec

Hemicentetes semispinosus

The lowland streaked tenrec (*Hemicentetes semispinosus*) is a small tenrec found in Madagascar. It belongs to the family Tenrecidae in the order Afrosoricida, and more specifically to the subfamily of the spiny tenrecs Tenrecinae. Its natural...

Lowlands tree-kangaroo

Lowlands tree-kangaroo

Dendrolagus spadix

The lowlands tree-kangaroo or lowland tree-kangaroo (*Dendrolagus spadix*) is a species of tree-kangaroo in the family Macropodidae. This marsupial is endemic to the Southern New Guinea lowland rainforests ecoregion in southwestern Papua New...

Loxodonta exoptata

Loxodonta exoptata

Loxodonta exoptata

Loxodonta exoptata is an extinct species of elephant in the genus *Loxodonta*, from Africa. A 2009 study suggested that *Loxodonta exoptata* gave rise to *L. atlantica*, which gave rise to *L. africana*. The molars of *L. exoptata* are distinguished from...

Loyalty bent-winged bat

Loyalty bent-winged bat

Miniopterus robustior

The Loyalty bent-winged bat (*Miniopterus robustior*) is a species of vesper bat in the family Miniopteridae. It is found only in New Caledonia.

Lubang forest mouse

Lubang forest mouse

Apomys lubangensis

The Lubang forest mouse (*Apomys lubangensis*) is a forest mouse endemic to Lubang Island in the Philippines.

Lucifer titi

Lucifer titi

Cheracebus lucifer

The *Lucifer titi* (*Cheracebus lucifer*) is a species of titi, a type of New World monkey, from South America. It is found in Brazil, Colombia, Ecuador, and Peru. It was described as *Callicebus lucifer* in 1914. The *Lucifer titi* has previously been...

Lucina's shrew

Lucina's shrew

Crocidura lucina

Lucina's shrew (*Crocidura lucina*) is a species of mammal in the family Soricidae. It is endemic to Ethiopia. The mammal's natural habitats are subtropical or tropical high-altitude grassland and swamps.

Ludia's shrew

Ludia's shrew

Crocidura ludia

Ludia's shrew (*Crocidura ludia*) is a species of mammal in the family Soricidae. It is found in Cameroon, Central African Republic, Republic of the Congo, and Democratic Republic of the Congo. Its natural habitat is subtropical or tropical moist...

Luis Manuel's tailless bat

Luis Manuel's tailless bat
Anoura luismanueli

Luis Manuel's tailless bat (*Anoura luismanueli*) is a species of bat in the family Phyllostomidae. It is endemic to Venezuela and the eastern slope of the Cordillera Oriental of Colombia.

Lukolela swamp rat
Lukolela swamp rat
Praomys lukolelae

The Lukolela swamp rat (*Congomys lukolelae*) is a species of rodent in the family Muridae. It is found only in Democratic Republic of the Congo. Its natural habitat is subtropical or tropical moist lowland forest. It is threatened by habitat loss.

Lumholtz's Tree-Kangaroo
Lumholtz's Tree-Kangaroo
Dendrolagus lumholtzi

Lumholtz's tree-kangaroo (*Dendrolagus lumholtzi*) is a heavy-bodied tree-kangaroo found in rain forests of the Atherton Tableland Region of Queensland. Its status is classified as near threatened by the IUCN, and authorities consider it as rare....

Lund's Atlantic tree-rat
Lund's Atlantic tree-rat
Phyllomys lundii

Lund's Atlantic tree-rat, (*Phyllomys lundii*), is a spiny rat species from South America. It is found in Brazil.

Lunda rope squirrel
Lunda rope squirrel
Funisciurus bayonii

The Lunda rope squirrel (*Funisciurus bayonii*) is a species of rodent in the family Sciuridae. It is found in Angola and Democratic Republic of the Congo. Its natural habitat is moist savanna.

Lundomys
Lundomys
Lundomys molitor

Lundomys molitor, also known as Lund's amphibious rat or the greater marsh rat, is a semiaquatic rat species from southeastern South America. Its distribution is now restricted to Uruguay and nearby Rio Grande do Sul, Brazil, but it previously...

Lusitanian pine vole
Lusitanian pine vole
Microtus lusitanicus

The Lusitanian pine vole (*Microtus lusitanicus*) is a species of rodent in the family Cricetidae. It is endemic to the northwestern half of the Iberian Peninsula where it occurs in central and northern Portugal, northwest Spain, and in the extreme...

Lutrine opossum
Lutrine opossum
Lutreolina crassicaudata

Lutreolina is a genus of opossum found in South America. Both extant species in this genus are known as lutrine opossums. They have an otter-like body plan and occasionally semiaquatic tendencies, hence the genus name *Lutreolina*, which is Latin...

Luzon Aurora forest mouse
Luzon Aurora forest mouse
Apomys aurorae

The Luzon Aurora forest mouse (*Apomys aurorae*) is a forest mouse endemic to Aurora in Luzon, Philippines.

Luzon broad-toothed rat

Luzon broad-toothed rat

Abditomys latidens

The Luzon broad-toothed rat (*Abditomys latidens*) is a species of rodent in the family Muridae. It is endemic to central and northern Luzon in the Philippines. It is the only member of the genus *Abditomys*.

Luzon Cordillera forest mouse

Luzon Cordillera forest mouse

Apomys abrae

The Luzon Cordillera forest mouse (*Apomys abrae*) is a species of rodent in the family Muridae found only in the Philippines.

Luzon fruit bat

Luzon fruit bat

Otopteropus cartilagonodus

The Luzon fruit bat (*Otopteropus cartilagonodus*) is a species of megabat in the family Pteropodidae. It is monotypic within the genus *Otopteropus*. It is endemic to the Philippines. Its natural habitat is subtropical or tropical dry forest. It is...

Luzon giant forest mouse

Luzon giant forest mouse

Apomys magnus

The Luzon giant forest mouse (*Apomys magnus*) is a forest mouse endemic to Luzon, Philippines.

Luzon hairy-tailed rat

Luzon hairy-tailed rat

Batomys granti

The Luzon forest rat or Luzon hairy-tailed rat (*Batomys granti*) is one of five species of rodent in the genus *Batomys*. It is in the diverse family Muridae. This species is found only in the Philippines. This species was the first of the five to...

Luzon montane forest mouse

Luzon montane forest mouse

Apomys datae

The Luzon montane forest mouse (*Apomys datae*) is a species of rodent in the family Muridae, from the genus *Apomys*. It occurs only in the Philippines, where it has been found on the large northern island Luzon (in the Cordillera Central and on the...

Luzon short-nosed rat

Luzon short-nosed rat

Tryphomys adustus

The Luzon short-nosed rat (*Tryphomys adustus*) is a species of rodent in the family Muridae. It is the only species in the genus *Tryphomys*. It is found only in the Philippines, and is known only from Benguet, Laguna, and Tarlac provinces.

Luzon shrew

Luzon shrew

Crocidura grayi

The Luzon shrew (*Crocidura grayi*) is a species of mammal in the family Soricidae. It is endemic to the Philippines. It is threatened by habitat loss.

Luzon striped rat

Luzon striped rat
Chrotomys whiteheadi

The Luzon striped rat (*Chrotomys whiteheadi*) is a species of rodent in the family Muridae. It is found only in the Philippines. Its natural habitat is subtropical or tropical dry forest.

Luzon Zambales forest mouse
Luzon Zambales forest mouse
Apomys zambalensis

The Luzon Zambales forest mouse (*Apomys zambalensis*) is a forest mouse endemic to Zambales in Luzon, Philippines.

Lyle's Flying Fox
Lyle's Flying Fox
Pteropus lylei

Lyle's flying fox (*Pteropus lylei*) is a species of flying fox in the family Pteropodidae. It is found in Cambodia, Thailand and Vietnam, with an outlying population in Yunnan, China. It faces persecution from farmers and it is killed for...

Lynx issiodorensis
Lynx issiodorensis
Lynx issiodorensis

Lynx issiodorensis, sometimes called the Issoire lynx, is an extinct species of lynx that inhabited Europe during the late Pliocene to Pleistocene epochs, and may have originated in Africa during the late Pliocene. It is named after the town of...

Macaca anderssoni
Macaca anderssoni
Macaca anderssoni

Macaca anderssoni is a prehistoric species of macaque from the Pleistocene of China. It weighed between 7.5 kilograms (17 lb) and 10.5 kilograms (23 lb).

Macaca jiangchuanensis
Macaca jiangchuanensis
Macaca jiangchuanensis

Macaca jiangchuanensis is a prehistoric macaque from the early Pleistocene of China.

Macaca libyca
Macaca libyca
Macaca libyca

Macaca libyca is a prehistoric macaque from the Late Miocene of Wadi Natrun, Egypt.

Macaca majori
Macaca majori
Macaca majori

Macaca majori, commonly known as the dwarf macaque, is a prehistoric macaque from the Early Pleistocene of Sardinia, Italy.

MacArthur's mouse lemur
MacArthur's mouse lemur
Microcebus macarthurii

MacArthur's mouse lemur (*Microcebus macarthurii*), or the Anjiahely mouse lemur, is a species of mouse lemur known only from Makira Natural Park in northeastern Madagascar. The species is named in honor of the founder of the MacArthur Foundation,...

MacArthur's shrew
MacArthur's shrew
Crocidura macarthuri

MacArthur's shrew (*Crocidura macarthuri*) is a species of mammal in the family Soricidae. It is found in Kenya and Somalia. Its natural habitat is dry savanna. It is threatened by habitat loss.

MacConnell's bat
MacConnell's bat
Mesophylla macconnelli

MacConnell's bat, *Mesophylla macconnelli*, is a bat species from South and Central America. It is monotypic within the genus *Mesophylla*.

MacConnell's climbing mouse
MacConnell's climbing mouse
Rhipidomys macconnelli

MacConnell's climbing mouse (*Rhipidomys macconnelli*) is a nocturnal and arboreal rodent species from South America. It is found in Brazil, Guyana and Venezuela, where it inhabits rainforest at elevations from 300 to 2800 m.

Macedonian mouse
Macedonian mouse
Mus macedonicus

The Macedonian mouse (*Mus macedonicus*) is a species of rodent in the family Muridae and order Rodentia. This rodent lives in the area from eastern Georgia and western Bulgaria to Israel. It is considered part of a Palearctic group along with...

MacFarlane's bear
MacFarlane's bear
Ursus inopinatus

MacFarlane's bear was a grizzly bear killed in Canada's Northwest Territories in 1864. Inuit hunters shot and killed an enormous yellow-furred bear and gave the skin and skull to the Fort Anderson post manager and amateur naturalist Roderick...

Machado's butterfly bat
Machado's butterfly bat
Glauconycteris machadoi

Machado's butterfly bat (*Glauconycteris machadoi*) is a species of vesper bat in the family Vespertilionidae. It is found only in Angola. Its natural habitat is subtropical or tropical moist lowland forests.

Macinnes's mouse-tailed bat
Macinnes's mouse-tailed bat
Rhinopoma macinnesi

The Macinnes's mouse-tailed bat (*Rhinopoma macinnesi*) is a species of bat in the Rhinopomatidae family. It is found in Ethiopia, Kenya, Somalia, and Uganda. Its natural habitats are subtropical or tropical dry shrubland, hot deserts, and...

Mackenzie River wolf
Mackenzie River wolf
Canis lupus mackenzii

The Mackenzie River wolf or Mackenzie Arctic Wolf (*Canis lupus mackenzii*) is a subspecies of gray wolf which is found in Canada's southern portion of Northwest Territories. Not much has been published on *Canis lupus mackenzii* but one of the most...

Mackilligin's gerbil
Mackilligin's gerbil

Dipodillus mackilligini

Mackilligin's gerbil (*Dipodillus mackilligini*) also known as Mackilligin's dipodil, is a species of rodent in the family Muridae. It occurs in the southeastern deserts of Egypt and Sudan, around the area of the Halaib Triangle. It has been...

Macclaud's horseshoe bat

Macclaud's horseshoe bat

Rhinolophus macclaudi

Macclaud's horseshoe bat (*Rhinolophus macclaudi*) is a species of bat in the family Rhinolophidae. It is endemic to Guinea. Its natural habitats are moist savanna, caves, and subterranean habitats (other than caves). It is one of five African...

Macleay's dorcopsis

Macleay's dorcopsis

Dorcopsulus macleayi

Macleay's dorcopsis (*Dorcopsulus macleayi*), also known as the Papuan dorcopsis or the Papuan forest wallaby, is a species of marsupial in the family Macropodidae. It is endemic to Papua New Guinea, where its natural habitat is subtropical or...

Macleay's mustached bat

Macleay's mustached bat

Pteronotus macleayii

Macleay's mustached bat (*Pteronotus macleayii*) is a species of bat in the family Mormoopidae. It is found in Cuba and Jamaica, and is threatened by habitat loss. The species is named for William Sharp Macleay, who collected the type specimen.

MacMillan's shrew

MacMillan's shrew

Crocidura macmillani

MacMillan's shrew (*Crocidura macmillani*) is a species of mammal in the family Soricidae. It is endemic to the Ethiopian Highlands. Its natural habitats are subtropical or tropical moist montane forests, and moist savanna. It is threatened by...

Macmillan's thicket rat

Macmillan's thicket rat

Grammomys macmillani

Macmillan's thicket rat (*Grammomys macmillani*) is a species of rodent in the family Muridae. It is found in Central African Republic, Democratic Republic of the Congo, Ethiopia, Kenya, South Sudan, Tanzania, and Uganda. Its natural habitats are...

Macroderma godthelpi

Macroderma godthelpi

Macroderma godthelpi

Macroderma godthelpi is a species of bat known from fossil material found in Australia, one of the larger carnivorous megadermatid family of the order Chiroptera. They resembled the modern species *Macroderma gigas*, known as a false vampire or...

Macroderma koppa

Macroderma koppa

Macroderma koppa

Macroderma koppa is a species of bat known from fossil material found in Australia, one of the larger carnivorous megadermatid family of the order Chiroptera. They resembled the modern species *Macroderma gigas*, known as a false vampire or ghost...

Macroderma malugara

Macroderma malugara
Macroderma malugara

Macroderma malugara is a species of bat known from fossil material found in Australia. The name describes a 'good killer' in the local language, and was similar in size and probably habits of the modern Macroderma gigas (known as the ghost bat)....

Macroscelides flavicaudatus
Macroscelides flavicaudatus
Macroscelides flavicaudatus

Macroscelides flavicaudatus (common name: Namib round-eared sengi or Namib round-eared elephant shrew) is a species of elephant shrew in the family Macroscelididae found in the central Namib desert and southern Namibia.

Macroscelides micus
Macroscelides micus
Macroscelides micus

Macroscelides micus (common name Etendeka round-eared sengi or Etendeka round-eared elephant shrew) is a species of elephant shrew in the family Macroscelididae. It is only found in gravel plains in the Etendeka formation of north-west Namibia....

Macroscelides proboscideus
Macroscelides proboscideus
Macroscelides proboscideus

The round-eared elephant shrew (Macroscelides proboscideus) or round-eared sengi (called the Karoo round-eared elephant shrew to distinguish it from its sister species; formerly misleadingly named the "short-eared elephant shrew"), is a species...

Madagascan Flying Fox
Madagascan Flying Fox
Pteropus rufus

The Madagascan flying fox, Madagascar flying-fox, or Madagascar fruit bat (Pteropus rufus) is a species of megabat in the genus Pteropus. It is endemic to Madagascar. Its natural habitat is subtropical or tropical moist lowland forests. It is...

Madagascan fruit bat
Madagascan fruit bat
Eidolon dupreanum

The Madagascan fruit bat (Eidolon dupreanum) is a species of bat in the family Pteropodidae. It is endemic to Madagascar and is listed as "Vulnerable" by the IUCN because it is hunted as bushmeat.

Madagascan large free-tailed bat
Madagascan large free-tailed bat
Tadarida fulminans

The Madagascan large free-tailed bat (Tadarida fulminans) is a species of bat in the family Molossidae. It is found in Democratic Republic of the Congo, Kenya, Madagascar, Malawi, Rwanda, South Africa, Tanzania, Zambia, and Zimbabwe. Its natural...

Madagascan pygmy shrew
Madagascan pygmy shrew
Suncus madagascariensis

The Madagascan pygmy shrew (Suncus madagascariensis) is a species of mammal in the family Soricidae. It is the only known Malagasy shrew.

Madagascan rousette
Madagascan rousette

Rousettus madagascariensis

The Madagascan rousette (*Rousettus madagascariensis*) or Madagascar rousette, is a species of megabat in the family Pteropodidae endemic to Madagascar. Its natural habitat is subtropical or tropical dry forests. It is threatened by habitat loss.
Madagascar free-tailed bat
Madagascar free-tailed bat
Otomops madagascariensis

The Madagascar free-tailed bat or Malagasy giant mastiff bat (*Otomops madagascariensis*) is a species of free-tailed bat formerly included as a subspecies of the large-eared free-tailed bat, but was later considered to be a distinct, endemic...
Madagascar sheath-tailed bat
Madagascar sheath-tailed bat
Coleura kibomalandy

The Madagascar sheath-tailed bat (*Coleura kibomalandy*) is a species of sac-winged bat found in Madagascar.
Madagascar sucker-footed bat
Madagascar sucker-footed bat
Myzopoda aurita

The Madagascar sucker-footed bat, Old World sucker-footed bat, or simply sucker-footed bat (*Myzopoda aurita*) is a species of bat in the family Myzopodidae endemic to Madagascar, especially in the eastern part of the forests. The genus was...
Madame Berthe's Mouse Lemur
Madame Berthe's Mouse Lemur
Microcebus berthae

Madame Berthe's mouse lemur (*Microcebus berthae*) or Berthe's mouse lemur is the smallest of the mouse lemurs and the smallest primate in the world; the average body length is 9.2 cm (3.6 in) and seasonal weight is around 30 g (1.1 oz)....
Madeira pipistrelle
Madeira pipistrelle
Pipistrellus maderensis

The Madeira pipistrelle (*Pipistrellus maderensis*) is a species of vesper bat. It is endemic to Azores, Madeira and the Canary Islands.
Madidi titi
Madidi titi
Plecturocebus aureipalatii

The Madidi titi, also known as the GoldenPalace.com monkey or the golden palace monkey, is a titi, a kind of New World monkey, discovered in western Bolivia's Madidi National Park in 2004. Its scientific name is *Plecturocebus aureipalatii*, the...
Madras treeshrew
Madras treeshrew
Anathana ellioti

The Madras treeshrew (*Anathana ellioti*), also known as the Indian treeshrew, is a species of treeshrew in the monotypic genus *Anathana* found in the hill forests of central and southern India. The genus name is derived from the Tamil name of...
Madura horseshoe bat
Madura horseshoe bat
Rhinolophus madurensis

The Madura horseshoe bat (*Rhinolophus madurensis*) is a species of bat from the

family Rhinolophidae. Current taxonomy treats the Madura horseshoe bat as a species separate of the Sulawesi horseshoe bat and not including parvus, but Csorba et al....

Maduran leaf-nosed bat
Maduran leaf-nosed bat
Hipposideros madurae

The Maduran leaf-nosed bat (*Hipposideros madurae*) is a species of bat in the family Hipposideridae endemic to Indonesia. This species is known from the eastern half of Java and on Madura Island. It is found below 1,000 metres (3,300 ft). It is an...

Maendeleo horseshoe bat
Maendeleo horseshoe bat
Rhinolophus maendeleo

The Maendeleo horseshoe bat (*Rhinolophus maendeleo*) is a recently discovered species of bats in the family Rhinolophidae. It inhabits caves of the Coastal Lowland forests of Tanzania. Its closest relative is the Adam's horseshoe bat. It was...

Magdalena rat
Magdalena rat
Xenomys nelsoni

The Magdalena rat (*Xenomys nelsoni*) is a species of rodent in the family Cricetidae which is found only in a small region of western Mexico. It is the only species in the genus *Xenomys*. The common name comes from the village where the first...

Magdalena spiny rat
Magdalena spiny rat
Proechimys magdalenae

The Magdalena spiny rat (*Proechimys magdalenae*) is a species of rodent in the family Echimyidae. It is endemic to Colombia.

Magellanic tuco-tuco
Magellanic tuco-tuco
Ctenomys magellanicus

The Magellanic tuco-tuco (*Ctenomys magellanicus*) is a species of rodent in the family Ctenomyidae. It is found in Argentina and Chile. Its natural habitat is subtropical or tropical dry lowland grassland. It is also known as the cururo by the Ona...

Maggie Taylor's roundleaf bat
Maggie Taylor's roundleaf bat
Hipposideros maggietaylorae

The Maggie Taylor's roundleaf bat (*Hipposideros maggietaylorae*) is a species of bat in the family Hipposideridae. It is found in West Papua (Indonesia) and Papua New Guinea.

Maghreb garden dormouse
Maghreb garden dormouse
Eliomys munbyanus

The Maghreb garden dormouse (*Eliomys munbyanus*) is a nocturnal species of rodent in the family Gliridae. It is found in Algeria, Libya, Morocco, Tunisia and Western Sahara. Its natural habitats vary from humid forests to semi-deserts. It is a...

Mahogany Glider
Mahogany Glider
Petaurus gracilis

The mahogany glider (*Petaurus gracilis*) is an endangered gliding possum native to a small region of coastal Queensland in Australia.

Mahomet mouse

Mahomet mouse

Mus mahomet

The Mahomet mouse (*Mus mahomet*) is a species of rodent in the family Muridae. It is found in Ethiopia, Kenya, and Uganda. Its natural habitats are subtropical or tropical moist montane forests and subtropical or tropical high-altitude shrubland. It...

Mainland drill

Mainland drill

Mandrillus leucophaeus leucophaeus

The mainland drill (*Mandrillus leucophaeus leucophaeus*) is a subspecies of the endangered drill. It is distinguished by ringed yellow and black coloring on its crown, and is otherwise similar to the Bioko drill.

Mainland serow

Mainland serow

Capricornis sumatraensis

The mainland serow (*Capricornis sumatraensis*) is a serow species native to the Himalayas, Southeast Asia and China.

Major's long-fingered bat

Major's long-fingered bat

Miniopterus majori

Major's long-fingered bat (*Miniopterus majori*) is a species of vesper bat in the family Vespertilionidae. It is found only in Madagascar. It is similar to *M. schreibersi* of Africa, differing by having a shorter forearm, slightly longer digits and...

Major's long-tailed tenrec

Major's long-tailed tenrec

Microgale majori

Major's long-tailed tenrec (*Microgale majori*) is a species of mammal in the family Tenrecidae. It is endemic to Madagascar. Its natural habitat is the eastern humid forest of the island, as well as some western forests, where it has been seen at...

Major's pine vole

Major's pine vole

Microtus majori

Major's pine vole (*Microtus majori*) is a species of rodent in the family Cricetidae found in Caucasus region and its vicinities (Russia, Georgia, Azerbaijan, Armenia, Turkey, Iran).

Major's tufted-tailed rat

Major's tufted-tailed rat

Eliurus majori

Major's tufted-tailed rat (*Eliurus majori*) is a species of rodent in the family Nesomyidae. It is found only in Madagascar. Its natural habitat is subtropical or tropical dry forests. It is threatened by habitat loss. The species was named in...

Majorcan hare

Majorcan hare

Lepus granatensis solisi

The Majorcan hare (*Lepus granatensis solisi*) is a subspecies of the Granada hare, from Majorca, Spain. It was described only in 1992. Most sources assume it is an extinct species.

Makira flying fox
Makira flying fox
Pteropus cognatus

The Makira flying fox (*Pteropus cognatus*) is a species of megabat in the genus *Pteropus*, found in the Solomon Islands. The species is currently decreasing and is endangered due to threats from logging and hunting. In 2013, Bat Conservation...

Makira roundleaf bat
Makira roundleaf bat
Hipposideros demissus

The Makira roundleaf bat (*Hipposideros demissus*) is a species of bat in the family *Hipposideridae* endemic to the Solomon Islands.

Makwassie musk shrew
Makwassie musk shrew
Crocidura maquassiensis

The Makwassie musk shrew (*Crocidura maquassiensis*) is a species of mammal in the family *Soricidae*. It is found in South Africa, Eswatini, and Zimbabwe. Its natural habitat is rocky areas.

Malabar large-spotted civet
Malabar large-spotted civet
Viverra civettina

The Malabar large-spotted civet (*Viverra civettina*), also known as the Malabar civet, is a viverrid endemic to the Western Ghats of India. It is listed as Critically Endangered on the IUCN Red List as the population is estimated to number fewer...

Malabar spiny dormouse
Malabar spiny dormouse
Platacanthomys lasiurus

The Malabar spiny dormouse (*Platacanthomys lasiurus*) is a species of muroid rodent endemic to the Western Ghats of India. It is the only extant species in the genus *Platacanthomys* and although resembling a dormouse, it is not closely related....

Malagasy Civet
Malagasy Civet
Fossa fossana

The Malagasy or striped civet (*Fossa fossana*), also known as the fanaloka (Malagasy,) or jabady, is an euplerid endemic to Madagascar. The Malagasy civet is a small mammal, about 47 centimetres (19 in) long excluding the tail (which is only...

Malagasy Giant Rat
Malagasy Giant Rat
Hypogeomys antimena

The Malagasy giant rat (*Hypogeomys antimena*) is a rodent found only in Madagascar. Habitat loss, limited range, and other factors made it an endangered species. It is the only extant species in the genus *Hypogeomys*; another species, *Hypogeomys*...

Malagasy mountain mouse
Malagasy mountain mouse
Monticolomys koopmani

The Malagasy mountain mouse or Koopman's montane voalavo (*Monticolomys koopmani*) is a rodent within the subfamily *Nesomyinae* of the family *Nesomyidae*. It is monotypic within the genus *Monticolomys*, and is closely related to the big-footed mouse...

Malagasy mouse-eared bat
Malagasy mouse-eared bat

Myotis goudoti

The Malagasy mouse-eared bat (*Myotis goudotii*) is a species of bat in the family Vespertilionidae that is endemic to Madagascar.

Malagasy serotine

Malagasy serotine

Neoromicia matroka

The Malagasy serotine (*Laephotis matroka*), is a species of vesper bat. It is found only in Madagascar. It was formerly classified in *Neoromicia* before phylogenetic analysis found it to belong to *Laephotis*.

Malagasy slit-faced bat

Malagasy slit-faced bat

Nycteris madagascariensis

The Malagasy slit-faced bat (*Nycteris madagascariensis*) is a species of slit-faced bat in Madagascar. Very little is known about this species.

Malagasy white-bellied free-tailed bat

Malagasy white-bellied free-tailed bat

Mops leucostigma

The Malagasy white-bellied free-tailed bat (*Mops leucostigma*) is a species of bat in the family Molossidae. It is endemic to Madagascar.

Malaita tube-nosed fruit bat

Malaita tube-nosed fruit bat

Nyctimene malaitensis

The Malaita tube-nosed fruit bat (*Nyctimene malaitensis*) is a species of bat in the family Pteropodidae. It is endemic only to the islands of Malaita and Makira (formerly known as San Cristóbal) in the Solomon Islands. The species occurs in...

Malawi bushbaby

Malawi bushbaby

Paragalago nyasae

The Malawi bushbaby (*Paragalago nyasae*) is a species of primate in the family Galagidae. It lives in southern Malawi and the neighboring region in Mozambique. The IUCN considers it to be part of the species *Paragalago granti*, Grant's bushbaby.

Malayan Civet

Malayan Civet

Viverra zangalunga

The Malayan civet (*Viverra zangalunga*) is a member of the viverrid family. It is listed as "Least Concern" by IUCN as it is relatively widely distributed, appears to be tolerant of degraded habitats, and occurs in a number of protected areas.

Malayan field rat

Malayan field rat

Rattus tiomanicus

The Malayan field rat, Malaysian field rat or Malaysian wood rat, (*Rattus tiomanicus*) is a species of rodent in the family Muridae. It is nocturnal and mainly arboreal and is found in Malaysia, Thailand, Indonesia and the Philippines. It is a...

Malayan free-tailed bat

Malayan free-tailed bat

Mops mops

The Malayan free-tailed bat (*Mops mops*) is a species of bat in the family Molossidae. It is found in Indonesia and Malaysia.

Malayan horseshoe bat
Malayan horseshoe bat
Rhinolophus malayanus

The Malayan horseshoe bat (*Rhinolophus malayanus*) is a species of bat in the family Rhinolophidae. It is found in Cambodia, Laos, Malaysia, Myanmar, Thailand, and Vietnam.

Malayan mountain spiny rat
Malayan mountain spiny rat
Maxomys inas

The Malayan mountain spiny rat (*Maxomys inas*) is a species of rodent in the family Muridae. It is found only in Malaysia.

Malayan Porcupine
Malayan Porcupine
Hystrix brachyura

The Malayan porcupine or Himalayan porcupine (*Hystrix brachyura*) is a species of rodent in the family Hystricidae. Three subspecies are extant in South and Southeast Asia.

Malayan pygmy shrew
Malayan pygmy shrew
Suncus malayanus

The Malayan pygmy shrew (*Suncus malayanus*) is a species of mammal in the family Soricidae. It can be found in Malaysia and Thailand. Its natural habitat is subtropical or tropical dry forests.

Malayan roundleaf bat
Malayan roundleaf bat
Hipposideros nequam

The Malayan roundleaf bat (*Hipposideros nequam*) is a horseshoe bat found only in Malaysia. It is listed as a data deficient species.

Malayan shrew
Malayan shrew
Crocidura malayana

The Malayan shrew (*Crocidura malayana*) is a species of mammal in the family Soricidae. It is endemic to Malaysia. It is threatened by habitat loss.

Malayan slit-faced bat
Malayan slit-faced bat
Nycteris tragata

The Malayan slit-faced bat (*Nycteris tragata*) is a species of slit-faced bat which lives in Indonesia, Malaysia, Myanmar, and Thailand.

Malayan tailless leaf-nosed bat
Malayan tailless leaf-nosed bat
Coelops robinsoni

The Malayan tailless leaf-nosed bat (*Coelops robinsoni*) is a species of bat in the family Hipposideridae. It is a very small bat which has long and soft fur. The fur coloration is brown to blackish on the dorsal surface and ashy on the ventral...

Malayan Tapir
Malayan Tapir
Tapirus indicus

The Malayan tapir (*Tapirus indicus*) (otherwise known as Asian tapir) is the only living species of Old World tapir. There are only 4 tapir species around the globe, and this animal is the largest of them. Meanwhile, the Asian tapir is the most...

Malayan tree rat
Malayan tree rat
Pithecheir parvus

The Malayan tree rat (*Pithecheir parvus*) is a species of rodent in the family Muridae. It is found only in Malaysia.

Malayan water shrew
Malayan water shrew
Chimarrogale hantu

The Malayan water shrew (*Chimarrogale hantu*), also known as the hantu water shrew, is a red-toothed shrew recorded only from the Malaysian state of Selangor. It was listed as a critically endangered, but is now considered near threatened. It gets...

Malayan Weasel
Malayan Weasel
Mustela nudipes

The Malayan weasel (*Mustela nudipes*) is a small mammal from Asia. It is very poorly known but assumed to occur at low densities and behave elusively based on low detection rates.

Malbrouck
Malbrouck
Chlorocebus cynosuros

The malbrouck (*Chlorocebus cynosuros*) is an Old World primate from Africa that belongs to the genus *Chlorocebus*. The species is sometimes classified as a subspecies of the vervet monkey (*C. pygerythrus*), or of the widespread grivet (*C. aethiops*).

Maluku myotis
Maluku myotis
Myotis moluccarum

The Maluku myotis (*Myotis moluccarum*), also commonly known as the Arafura large-footed bat, is a species of mouse-eared bat. Native to Indonesia, and possibly New Guinea and northern Australia, it one of several species known as "fishing bats"...

Mamfe shrew
Mamfe shrew
Crocidura virgata

The Mamfe shrew (*Crocidura virgata*) is a species of mammal in the family Soricidae. It was discovered by Sanderson in 1940, and can be found in Nigeria and Cameroon.

Mamore arboreal rice rat
Mamore arboreal rice rat
Oecomys mamorae

The mamore arboreal rice rat, (*Oecomys mamorae*), also known as the Mamore oecomys is an arboreal species of rodent in the genus *Oecomys* of family Cricetidae. Its distribution extends over much of Bolivia and into nearby Brazil and Paraguay....

Manado fruit bat
Manado fruit bat
Boneia bidens

The Manado fruit bat (*Boneia bidens*) is a species of megabat in the family Pteropodidae. It is the only member of the genus *Boneia*. It was formerly classified in the genus *Rousettus*, but phylogenetic evidence supports them being distinct and...

Manavi long-fingered bat
Manavi long-fingered bat

Miniopterus manavi

The Manavi long-fingered bat (*Miniopterus manavi*) is a bat in the genus *Miniopterus* that occurs in east-central Madagascar. First described in 1906, this species was later included in the mainland African *M. minor*. A 1995 revision united...

Manchurian hare

Manchurian hare

Lepus mandshuricus

The Manchurian hare (*Lepus mandshuricus*) is a species of hare found in northeastern China and Russia, the Amur River basin, and the higher mountains of northern North Korea. It lives in forests and the IUCN has assessed its conservation status...

Manchurian sika deer

Manchurian sika deer

Cervus nippon mantchuricus

The Manchurian sika deer or Dybowski's sika deer (*Cervus nippon mantchuricus* or *Cervus nippon dybowskii*) is a subspecies of deer, the largest of the 14 subspecies of sika deer. It was first described by Robert Swinhoe in 1864.

Manchurian wapiti

Manchurian wapiti

Cervus canadensis xanthopygus

The Manchurian wapiti (*Cervus canadensis xanthopygus*) is a subspecies of the wapiti native to East Asia.

Mandarin vole

Mandarin vole

Lasiopodomys mandarinus

The mandarin vole (*Lasiopodomys mandarinus*) is a species of vole found in central China as well as the southern and central Korean Peninsula. An adult mandarin vole is typically of about 115 millimeters in length, in addition to about 28 mm of...

Mandelli's mouse-eared bat

Mandelli's mouse-eared bat

Myotis sicarius

Mandelli's mouse-eared bat (*Myotis sicarius*) is a species of vesper bat. It can be found in India and Nepal. It is found in subtropical or tropical moist montane forests. It is threatened by habitat loss. The name honours the ornithologist Louis...

Mandrill

Mandrill

Mandrillus sphinx

The mandrill (*Mandrillus sphinx*) is a large Old World monkey native to west-central Africa. It is one of the most colorful mammals in the world and its closest living relative is the drill. Both species were traditionally thought to be baboons,...

Maned Rat

Maned Rat

Lophiomys imhausi

The Maned rat (*Lophiomys imhausi*) is a long-haired and bushy-tailed African rodent that superficially resembles a porcupine. It is the world's only poisonous rodent, that borrows toxins from plants to fend off predators.

Maned Sloth

Maned Sloth

Bradypus torquatus

The maned sloth (*Bradypus torquatus*) is a three-toed sloth that lives only in Brazil. It is one of four species of three-toed sloth.

Maned Wolf

Maned Wolf

Chrysocyon brachyurus

The Maned wolf (*Chrysocyon brachyurus*) is a large canine of South America. Its markings resemble those of foxes, but it is neither a fox nor a wolf. It is the only species in the genus *Chrysocyon* (meaning "golden dog"). It is the largest canine...

Maneless Zebra

Maneless Zebra

Equus quagga borensis

The Maneless zebra (*Equus quagga borensis*) is a subspecies of the Plains zebra spread over the northern parts of eastern Africa.

Manenguba shrew

Manenguba shrew

Crocidura manengubae

The Manenguba shrew (*Crocidura manengubae*) is a species of mammal in the family Soricidae. It is endemic to Cameroon. Its natural habitat is subtropical or tropical moist montane forests.

Manicore marmoset

Manicore marmoset

Mico manicorensis

The Marca's marmoset (*Mico marcai*) is a species of marmoset that is endemic to the Amazon, in the Aripuanã-Manicoré interfluvium in Brazil. Its body is light grey, with orange legs, a black tail, a pinkish face, and naked ears. It is about 9...

Manipur bush rat

Manipur bush rat

Hadromys humei

The Manipur bush rat (*Hadromys humei*), also known as Hume's rat or Hume's hadromys, is a species of rodent in the family Muridae. It is found in northeastern India, and is listed as endangered.

Manipur white-toothed rat

Manipur white-toothed rat

Berylmys manipulus

The Manipur white-toothed rat (*Berylmys manipulus*) is a species of rodent in the family Muridae. The Manipur white-toothed rat is found in northeastern India, northern and central Myanmar, and Yunnan province of China (west of the Salween River)....

Manitoba wolf

Manitoba wolf

Canis lupus griseoalbus

The Manitoba wolf (*Canis lupus griseoalbus*), also known as the grey-white wolf, is an extinct subspecies of gray wolf that roamed in the southern Northwest Territories, northern Alberta, Saskatchewan, and south-central Manitoba. This wolf is...

Manitoban Elk

Manitoban Elk

Cervus canadensis manitobensis

The Manitoban elk (*Cervus canadensis manitobensis*) is a subspecies of elk found in

the United States and Canada. The subspecies was driven into near extinction by 1900 but has recovered since then.
Mantiqueira Atlantic tree-rat
Mantiqueira Atlantic tree-rat
Phyllomys mantiqueirensis

The Mantiqueira Atlantic Tree-rat (*Phyllomys mantiqueirensis*) is a spiny rat species from South America. It is found in Brazil.
Mantled Guereza
Mantled Guereza
Colobus guereza

The mantled guereza (*Colobus guereza*), also known simply as the guereza, the eastern black-and-white colobus, or the Abyssinian black-and-white colobus, is a black-and-white colobus, a type of Old World monkey. It is native to much of west...
Mantled Howler Monkey
Mantled Howler Monkey
Alouatta palliata

The mantled howler (*Alouatta palliata*) is a species of howler monkey, a type of New World monkey, from Central and South America. It is one of the monkey species most often seen and heard in the wild in Central America. It takes its "mantled"...
Mantled mastiff bat
Mantled mastiff bat
Otomops secundus

The mantled mastiff bat (*Otomops secundus*) is a species of bat in the family Molossidae endemic to Papua New Guinea.
Manu short-tailed bat
Manu short-tailed bat
Carollia manu

The Manu short-tailed bat (*Carollia manu*) is a bat species found in Peru and Bolivia.
Manus Island mosaic-tailed rat
Manus Island mosaic-tailed rat
Melomys matambuai

The Manus Island mosaic-tailed rat or Manus melomys (*Melomys matambuai*) is a species of rodent in the family Muridae. It is endemic to the island of Manus in Papua New Guinea where it occurs in forest habitats and is largely arboreal. The...
Manusela mosaic-tailed rat
Manusela mosaic-tailed rat
Melomys fraterculus

The Manusela mosaic-tailed rat (*Melomys fraterculus*) is a species of rodent in the family Muridae. It is found only in Indonesia. It lives in the rainforests on the island of Seram in Indonesia. Specimens of it weigh 66.5g and have a head and...
Manzano Mountain cottontail
Manzano Mountain cottontail
Sylvilagus cognatus

The robust cottontail or Holzner's cottontail (*Sylvilagus holzneri*) is a species of cottontail rabbit native to high-altitude regions of the southwestern United States and western Mexico.
Marajó short-tailed opossum
Marajó short-tailed opossum
Monodelphis maraxina

The Marajó short-tailed opossum (*Monodelphis maraxina*) is an opossum species from South America. It is endemic to Brazil, in the Amazon River delta region within Maranhão state.

Maranhão red-handed howler

Maranhão red-handed howler

Alouatta ululata

The Maranhão red-handed howler (*Alouatta ululata*) is an endangered species of howler monkey endemic to forests (for example Babaçu forests) in the northeastern Brazilian states of Ceará, Maranhão and Piauí. It was previously thought to be a...

Marañón white-fronted capuchin

Marañón white-fronted capuchin

Cebus yuracus

The Marañón white-fronted capuchin or Peruvian white-fronted capuchin (*Cebus yuracus*) is a species of gracile capuchin monkey from the upper Amazon Basin. It had been regarded as synonymous with the shock-headed capuchin (*C. cuscinus*), which...

Marbled Cat

Marbled Cat

Pardofelis marmorata

The Marbled cat (*Pardofelis marmorata*) is a small wild cat that lives in mountainous areas of Asia. The Marbled cat is closely related to the Asian golden cat and the Bay cat, all of which diverged from other felids about 9.4 million years ago.

Marbled Polecat

Marbled Polecat

Vormela peregusna

The marbled polecat (*Vormela peregusna*) is a small mammal belonging to the monotypic genus *Vormela* within the mustelid subfamily *Ictonychinae*. *Vormela* is from the German word *Würmlein*, which means "little worm". The specific name *peregusna* comes...

Marca's marmoset

Marca's marmoset

Mico marcai

The Marca's marmoset (*Mico marcai*) is a species of marmoset that is endemic to the Amazon, in the Aripuanã-Manicoré interfluvium in Brazil. Its body is light grey, with orange legs, a black tail, a pinkish face, and naked ears. It is about 9...

Marco Polo sheep

Marco Polo sheep

Ovis ammon polii

The Marco Polo sheep (*Ovis ammon polii*) is a subspecies of argali sheep, named after Marco Polo. Their habitat are the mountainous regions of Central Asia. Marco Polo sheep are distinguishable mostly by their large size and spiraling horns....

Mareeba rock-wallaby

Mareeba rock-wallaby

Petrogale mareeba

The Mareeba rock-wallaby (*Petrogale mareeba*) is a rare species of rock-wallaby found around Mareeba in northeastern Queensland, Australia.

Maremma boar

Maremma boar

Sus scrofa majori

The Maremman boar (*Sus scrofa majori*) is a subspecies of wild boar native to Maremma in Central Italy. It is much smaller than the Central European boar and the two subspecies have been interbred on farms. Genetic studies (Larson et al. 2005)...

Margarita Island kangaroo rat

Margarita Island kangaroo rat

Dipodomys merriami margaritae

The Margarita Island kangaroo rat (*Dipodomys merriami margaritae*) is a subspecies of rodent in the family Heteromyidae. It is endemic to Mexico, where it is found only on Isla Santa Margarita of Magdalena Bay, Baja California Sur. The natural...

Margay

Margay

Leopardus wiedii

The margay (*Leopardus wiedii*) is a small wild cat native to Central and South America. Until the 1990s, margays were hunted illegally for the wildlife trade, which resulted in a large population decrease.

Margot Marsh's mouse lemur

Margot Marsh's mouse lemur

Microcebus margotmarshae

Margot Marsh's mouse lemur or the Antafondro mouse lemur (*Microcebus margotmarshae*) is a species of mouse lemur endemic to Madagascar. Its holotype was first collected on 21 May 2006, proposed in 2006 by Andriantompohavana et al., and was...

Mariana Fruit Bat

Mariana Fruit Bat

Pteropus mariannus

The Mariana fruit bat (*Pteropus mariannus*) is a medium-sized megabat, also known as the Mariana flying fox and fanihi in Chamorro. Habitat loss has driven it to endangered status, and it is listed as threatened by the US Fish and Wildlife Service.

Marie's vole

Marie's vole

Volemys musseri

Marie's vole (*Volemys musseri*) is a species of rodent in the family Cricetidae. It is found only in China. *Volemys musseri* is one of two species in the genus *Volemys* along with the Szechuan vole (*Volemys millicens*).

Marine Otter

Marine Otter

Lontra felina

The Marine otter (*Lontra felina*) is a rare and relatively unknown South American mammal. It is the smallest of the New World otters and the smallest marine mammal. This charismatic marine species displays charm and dexterity. It lives along South...

Marinkelle's sword-nosed bat

Marinkelle's sword-nosed bat

Lonchorhina marinkellei

Marinkelle's sword-nosed bat (*Lonchorhina marinkellei*) is a bat species from South America. It is found in Colombia. In 2013, Bat Conservation International listed this species as one of the 35 species of its worldwide priority list of...

Maritime shrew

Maritime shrew

Sorex maritimensis

The maritime shrew (*Sorex maritimensis*) is a species of mammal in the family Soricidae. It is found in New Brunswick and Nova Scotia in Canada.
Maritime striped squirrel
Maritime striped squirrel
Tamias maritimus

The maritime striped squirrel or eastern striped squirrel (*Tamias maritimus*) is a species of rodent in the family Sciuridae. It is found in southern and eastern China (including Hainan), Taiwan, and Laos and Vietnam east of Mekong. This species...
Markhor
Markhor
Capra falconeri

The markhor (*Capra falconeri*) is a large *Capra* species. It is the national animal of Pakistan, where it is also known as the screw horn or "screw-horned goat", *mārkhor* (مارخور) in Urdu from Classical Persian.
Marley's golden mole
Marley's golden mole
Amblysomus marleyi

Marley's golden mole (*Amblysomus marleyi*) is a species of burrowing mammal in the golden mole family, Chrysochloridae. It is found in South Africa and possibly Eswatini. It has been separated from *Amblysomus hottentotus* by Bronner (1995b, 1996,...
Marmoset rat
Marmoset rat
Hapalomys longicaudatus

The marmoset rat (*Hapalomys longicaudatus*), also known as the greater marmoset rat, is a species of rodent in the family Muridae. It is found in Malaysia, Myanmar, and Thailand. Its natural habitat is subtropical or tropical dry forests. It is...
Marohita mouse lemur
Marohita mouse lemur
Microcebus marohita

The Marohita mouse lemur (*Microcebus marohita*) is a species of mouse lemur known only from the Marohita Forest in eastern Madagascar, near the village of Marolambo. Specimens were first collected in December 2003, and its discovery was announced...
Maroon Leaf Monkey
Maroon Leaf Monkey
Presbytis rubicunda

The maroon langur, maroon leaf monkey, or red leaf monkey (*Presbytis rubicunda*) is a member of the family Cercopithecidae. It is found on the southeast Asian island of Borneo and the nearby smaller Karimata. *P. rubicunda* mostly live in forests...
Marovaza house bat
Marovaza house bat
Scotophilus marovaza

The Marovaza yellow bat or Marovaza house bat (*Scotophilus marovaza*) is a species of bat found in Madagascar.
Marsh Deer
Marsh Deer
Blastocerus dichotomus

The marsh deer (*Blastocerus dichotomus*) is the largest deer species from South

America reaching a length of 2 m (6.6 ft) and a shoulder height of 1.2 m (3.9 ft). It is found in Argentina, Bolivia, Peru, Brazil, Uruguay and Paraguay. Formerly...
Marsh Mongoose
Marsh Mongoose
Atilax paludinosus

The marsh mongoose (*Atilax paludinosus*), also known as the water mongoose or the vansire, is a medium-sized mongoose native to sub-Saharan Africa that inhabits foremost freshwater wetlands. It has been listed as Least Concern on the IUCN Red...
Marsh Rabbit
Marsh Rabbit
Sylvilagus palustris

The marsh rabbit (*Sylvilagus palustris*) is a small cottontail rabbit found in marshes and swamps of coastal regions of the Eastern and Southern United States. It is a strong swimmer and found only near regions of water. It is similar in...
Marsh Rice Rat
Marsh Rice Rat
Oryzomys palustris

The marsh rice rat (*Oryzomys palustris*) is a semiaquatic North American rodent in the family Cricetidae. It usually occurs in wetland habitats, such as swamps and salt marshes. It is found mostly in the eastern and southern United States, from...
Marsh shrew
Marsh shrew
Sorex bendirii

The marsh shrew (*Sorex bendirii*), also known as the Pacific water shrew, Bendire's water shrew, Bendire's shrew and Jesus shrew is the largest North American member of the genus *Sorex* (long-tailed shrews). Primarily covered in dark-brown fur, it...
Marshall's horseshoe bat
Marshall's horseshoe bat
Rhinolophus marshalli

Marshall's horseshoe bat (*Rhinolophus marshalli*) is a species of bat in the family Rhinolophidae. It is found in Laos, Malaysia, Thailand, and Vietnam.
Marsican brown bear
Marsican brown bear
Ursus arctos marsicanus / *arctos*

The Marsican brown bear (*Ursus arctos arctos*, formerly *Ursus arctos marsicanus*), also known as the "Apennine brown bear", and *orso bruno marsicano* in Italian, is a critically endangered population or subspecies of the Eurasian brown bear, with a...
Marsupial Mole
Marsupial Mole
Genus *Notoryctes*

The northern marsupial mole or kakarratul (*Notoryctes caurinus*) is a marsupial in the family Notoryctidae, an endemic animal of arid regions of Central Australia. It lives in the loose sand of dunes and river plains in the desert, spending...
Martins's tamarin
Martins's tamarin
Saguinus martinsi

Martins's tamarin (*Saguinus martinsi*) or Martin's ochraceous bare-face tamarin, is a species of tamarin endemic to Brazil.
Masai Giraffe

Masai Giraffe
Giraffa camelopardalis tippelskirchii

The Masai giraffe (*Giraffa camelopardalis tippelskirchi*) is a subspecies of giraffe native to East Africa. The Masai giraffe is the tallest land animal on Earth and is currently the national animal of Tanzania.

Mashona mole-rat
Mashona mole-rat
Fukomys darlingi

The Mashona mole-rat (*Fukomys darlingi*) is a species of rodent in the family Bathyergidae. It is found in Mozambique and Zimbabwe. Its natural habitats are subtropical or tropical dry shrubland, subtropical or tropical dry lowland grassland, and...

Masked flying fox
Masked flying fox
Pteropus personatus

The masked flying fox, Moluccan masked flying fox or masked fruit bat (*Pteropus personatus*), is a species of flying fox in the family Pteropodidae. It is endemic to Indonesia. It is part of a species complex of closely related species. The...

Masked mouse-tailed dormouse
Masked mouse-tailed dormouse
Myomimus personatus

The masked mouse-tailed dormouse (*Myomimus personatus*) is a species of rodent in the family Gliridae. It is found in Iran and Turkmenistan.

Masked Palm Civet
Masked Palm Civet
Paguma larvata

The masked palm civet (*Paguma larvata*), also called the gem-faced civet, is a palm civet species native to the Indian subcontinent and Southeast Asia. It has been listed as least concern on the IUCN Red List since 2008 as it occurs in many...

Masked ringtail possum
Masked ringtail possum
Pseudochirulus larvatus

The masked ringtail possum (*Pseudochirulus larvatus*) is a marsupial possum of the family Pseudocheiridae. It is found in northeastern New Guinea in the Star Mountains, the eastern Central Cordillera, the Huon Peninsula and the northern coast...

Masked white-tailed rat
Masked white-tailed rat
Uromys hadrourus

The masked white-tailed rat (*Uromys hadrourus*) is a species of rodent in the family Muridae. It is found only in Australia, in north-east Queensland. Descriptive English common names for this species include Masked White-tailed Rat and Thornton...

Masoala fork-marked lemur
Masoala fork-marked lemur
Phaner furcifer

The Masoala fork-marked lemur (*Phaner furcifer*), also known as the eastern fork-marked lemur or Masoala fork-crowned lemur, is a species of lemur found in the coastal forests of northeastern Madagascar. It is a small nocturnal animal with large...

Master leaf-eared mouse

Master leaf-eared mouse
Phyllotis magister

The master leaf-eared mouse (*Phyllotis magister*) is a species of rodent in the family Cricetidae. It is found in Chile and Peru.

Matabeleland mole-rat
Matabeleland mole-rat
Cryptomys nimrodi

The Matabeleland mole-rat, *Cryptomys nimrodi*, is a species of mole-rat found in Zimbabwe.

Matapalo broad-nosed bat
Matapalo broad-nosed bat
Platyrrhinus matapalensis

The Matapalo broad-nosed bat, *Platyrrhinus matapalensis*, is a species of leaf-nosed bat described in 2005. It is found in South America.

Mato Grosso dog-faced bat
Mato Grosso dog-faced bat
Neoplatymops mattogrossensis

The Mato Grosso dog-faced bat (*Neoplatymops mattogrossensis*), is a bat species from South America. It is found in Brazil, Colombia, Guyana and Venezuela.

Matschie's Tree-Kangaroo
Matschie's Tree-Kangaroo
Dendrolagus matschiei

Matschie's tree-kangaroo (*Dendrolagus matschiei*), also known as the Huon tree-kangaroo is a tree-kangaroo native to the Huon Peninsula of northeastern New Guinea island, within the nation of Papua New Guinea. Under the IUCN classification,...

Matses's big-eared bat
Matses's big-eared bat
Micronycteris matses

Matses's big-eared bat (*Micronycteris matses*) is a bat species from South America. It is found in Brazil.

Matthey's mouse
Matthey's mouse
Mus mattheyi

Matthey's mouse (*Mus mattheyi*) is a species of rodent in the family Muridae. Matthey's mouse was first described by Francis Petter (1923-2012) in 1969 and was named after Robert Matthey (1900-1982).It is found in Burkina Faso, Ivory Coast,...

Maués marmoset
Maués marmoset
Mico mauesi

The Maués marmoset (*Mico mauesi*) is a marmoset endemic to Brazil. It is found only on the west bank of the Maués Açu River, in the Amazonas state.

Māui dolphin
Māui dolphin
Cephalorhynchus hectori maui

Māui dolphin, Maui's dolphin, or popoto (*Cephalorhynchus hectori maui*) is a subspecies of the Hector's dolphin (*Cephalorhynchus hectori*)—New Zealand's only endemic cetacean. Māui dolphins are only found off the west coast of New Zealand's North...

Maule tuco-tuco
Maule tuco-tuco

Ctenomys maulinus

The Maule tuco-tuco (*Ctenomys maulinus*) is a species of rodent in the family Ctenomyidae. It is found in Argentina and Chile where it occupies several different types of habitats. It is a common species and the IUCN has assessed its conservation...

Mauritanian Flying Fox

Mauritanian Flying Fox

Pteropus niger

The Mauritian flying fox (*Pteropus niger*), also known as Greater Mascarene flying fox or Mauritius fruit bat is a large megabat species endemic to Mauritius and La Réunion.

Mauritanian gerbil

Mauritanian gerbil

Gerbillus mauritaniae

The Mauritanian gerbil (*Gerbillus mauritaniae*) is distributed mainly in northern Mauritania. Some authorities place it in a separate genus, *Monodia*. It is also sometimes considered a subspecies of the Sudan gerbil.

Mauritanian shrew

Mauritanian shrew

Crocidura lusitania

The Mauritanian shrew (*Crocidura lusitania*) is a species of mammal in the family Soricidae. It is found in Algeria, Burkina Faso, Eritrea, Ethiopia, Gambia, Guinea, Guinea-Bissau, Mali, Mauritania, Morocco, Niger, Nigeria, Senegal, Sierra Leone....

Mauritian tomb bat

Mauritian tomb bat

Taphozous mauritanus

The Mauritian tomb bat (*Taphozous mauritanus*) is a species of sac-winged bat in the family Emballonuridae that is found in central and southern Africa and Madagascar. It was discovered in 1818 by Étienne Geoffroy Saint-Hilaire, and is...

Maximowicz's vole

Maximowicz's vole

Microtus maximowiczii

Maximowicz's vole (*Microtus maximowiczii*) is a species of rodent in the family Cricetidae. It is found in northeastern China, Mongolia, and eastern Russia.

Maxomys baeodon

Maxomys baeodon

Maxomys baeodon

Maxomys baeodon, also known as the small Bornean maxomys or small spiny rat, is a species of rodent in the family Muridae. It is known only from Sarawak and Sabah in the Malaysian part of Borneo.

Maxomys tajuddinii

Maxomys tajuddinii

Maxomys tajuddinii

Maxomys tajuddinii is a species of rodent in the genus *Maxomys*. It is found in Indonesia and Malaysia. It is a rodent of medium size, with a tail length of 106.9-122.3 mm, feet of 27.62-30.04 mm and a weight of up to 70 g.

Maxwell's duiker

Maxwell's duiker

Philantomba maxwellii

The Maxwell's duiker (*Philantomba maxwellii*) is a small antelope found in western

Africa.
Maya mouse
Maya mouse
Peromyscus mayensis

The Maya mouse (*Peromyscus mayensis*) is a species of rodent in the family Cricetidae. It is found only in Guatemala.

Mayor's mouse
Mayor's mouse
Mus mayori

Mus mayori is a species of rodent in the genus *Mus*, the mice. Its common names include Mayor's mouse, highland rat, and spiny mouse. It is endemic to Sri Lanka. This mouse lives in tropical and subtropical forest types and wet grassland habitat....

Mazama pocket gopher
Mazama pocket gopher
Thomomys mazama

The *Mazama pocket gopher* (*Thomomys mazama*) is a smooth-toothed pocket gopher restricted to the Pacific Northwest. The herbivorous species ranges from coastal Washington, through Oregon, and into north-central California. Four subspecies of the...

McIlhenny's four-eyed opossum
McIlhenny's four-eyed opossum
Philander mcilhennyi

McIlhenny's four-eyed opossum (*Philander mcilhennyi*) is a South American species of Opossum. Found in Brazil and Peru, it is almost entirely black, except for white spots above each eye. This species is named for John Stauffer "Jack" McIlhenny...

Meadow Jumping Mouse
Meadow Jumping Mouse
Zapus hudsonius

The meadow jumping mouse (*Zapus hudsonius*) is the most widely distributed mouse in the family Zapodidae. Its range extends from the Atlantic coast in the east to the Great Plains west, and from the arctic tree lines in Canada and Alaska to the...

Mearns coyote
Mearns coyote
Canis latrans mearnsi

The coyote (*Canis latrans*) is a species of canine native to North America. It is smaller than its close relative, the wolf, and slightly smaller than the closely related eastern wolf and red wolf. It fills much of the same ecological niche as...

Mearns's grasshopper mouse
Mearns's grasshopper mouse
Onychomys arenicola

Mearns's grasshopper mouse or the Chihuahuan grasshopper mouse (*Onychomys arenicola*) is a grasshopper mouse found in southwestern New Mexico, West Texas, and north-central Mexico. They are similar to *Onychomys torridus*, but differ in karyotype...

Mearns's pouched mouse
Mearns's pouched mouse
Saccostomus mearnsi

Mearns's pouched mouse (*Saccostomus mearnsi*) is a species of rodent in the family Nesomyidae. It is found in Ethiopia, Kenya, Somalia, Tanzania, and Uganda. Its natural habitats are dry savanna, subtropical or tropical dry shrubland, hot deserts,...

Mearns's squirrel
Mearns's squirrel
Tamiasciurus mearnsi

Mearns's squirrel (*Tamiasciurus douglasii mearnsi*) is a subspecies of the Douglas squirrel endemic to Mexico. It is endangered and occurs in low densities, and is threatened by habitat loss. It is possibly also threatened by competition from the...

Mechow's mole-rat
Mechow's mole-rat
Fukomys mechowii

Mechow's mole-rat (*Fukomys mechowii*) is a species of rodent in the family Bathyergidae. It is found in Angola, Democratic Republic of the Congo, Zambia, and possibly Malawi. Its natural habitats are moist savanna, subtropical or tropical dry...

Mechuka giant flying squirrel
Mechuka giant flying squirrel
Petaurista mechukaensis

The Mechuka giant flying squirrel (*Petaurista mechukaensis*) is a species of rodent in the family Sciuridae. First described in 2007 from East Himalayan forests at altitudes of 1,500–2,500 m (4,900–8,200 ft) in the region of Mechuka of...

Medellín small-eared shrew
Medellín small-eared shrew
Cryptotis medellinia

The Medellín small-eared shrew (*Cryptotis medellinia*) is a species of mammal in the family Soricidae. It is endemic to Colombia, where it is known from the northern parts of the Cordillera Occidental and Cordillera Central at elevations from...

Mediterranean horseshoe bat
Mediterranean horseshoe bat
Rhinolophus euryale

The Mediterranean horseshoe bat (*Rhinolophus euryale*) is a species of insectivorous bat in the family Rhinolophidae. It is found in the Mediterranean region and balkan peninsula, as well as parts of Italy.

Mediterranean Monk Seal
Mediterranean Monk Seal
Monachus monachus

The Mediterranean monk seal (*Monachus monachus*) is a monk seal belonging to the family Phocidae. As of 2015, it is estimated that fewer than 700 individuals survive in three or four isolated subpopulations in the Mediterranean, (especially) in...

Mediterranean pine vole
Mediterranean pine vole
Microtus duodecimcostatus

The Mediterranean pine vole (*Microtus duodecimcostatus*) is a species of rodent in the family Cricetidae. It is found in France, Andorra, Portugal, and Spain where it lives in a network of shallow tunnels.

Mediterranean water shrew
Mediterranean water shrew
Neomys anomalus

The Mediterranean, Southern or Miller's water shrew (*Neomys anomalus*) is a species of mammal in the family Soricidae.

Medium-tailed brush-furred rat
Medium-tailed brush-furred rat
Lophuromys medicaudatus

The medium-tailed brush-furred rat (*Lophuromys medicaudatus*) is a species of rodent in the family Muridae. It has been recorded from the Democratic Republic of the Congo, Rwanda, and Uganda. It lives in rainforest and swamps. It is a rare...

Medje free-tailed bat
Medje free-tailed bat
Mops congicus

The Medje free-tailed bat (*Mops congicus*) is a species of bat in the family Molossidae. It is found in Cameroon, Democratic Republic of the Congo, and Uganda. Its natural habitats are subtropical or tropical dry forests and subtropical or...

Meerkat
Meerkat
Suricata suricatta

The meerkat (*Suricata suricatta*) or suricate is a small mongoose found in southern Africa. They live in rock crevices in stony, often calcareous areas, and in large burrow systems in plains. They use a broad variety of calls to communicate with...

Megaoryzomys
Megaoryzomys
Megaoryzomys curioi

Megaoryzomys curioi, also known as the Galápagos giant rat, is an extinct species of sigmodontine rodent, known only from Santa Cruz Island in the Galápagos Islands. It likely met its demise when European settlers introduced invasive species to...

Mehely's horseshoe bat
Mehely's horseshoe bat
Rhinolophus mehelyi

Mehely's horseshoe bat (*Rhinolophus mehelyi*) is a species of insectivorous bat in the family Rhinolophidae found in Southern Europe and parts of the Middle East. It is distributed in a narrow band around the Mediterranean Sea from North-Western...

Melanomys caliginosus
Melanomys caliginosus
Melanomys caliginosus

Melanomys caliginosus, also known as the dusky melanomys or dusky rice rat, is a species of rodent in the genus *Melanomys* of family Cricetidae. It is found from Central America, in Honduras, Nicaragua, Costa Rica, and Panama, into South America,...

Melanomys robustulus
Melanomys robustulus
Melanomys robustulus

Melanomys robustulus, also known as the robust melanomys or robust dark rice rat, is a species of rodent in the genus *Melanomys* of family Cricetidae. It is found on the eastern slope of the Andes in Ecuador.

Melanomys zunigae
Melanomys zunigae
Melanomys zunigae

Melanomys zunigae, also known as Zuniga's melanomys or Zuniga's dark rice rat, is a species of rodent in the genus *Melanomys* of family Cricetidae. Known only from a small region of coastal Peru, it is listed as "critically endangered" by the IUCN...

Melck's house bat

Melck's house bat
Neoromicia melckorum

Melck's house bat (*Neoromicia melckorum*) is a species of vesper bat. It is found in Democratic Republic of the Congo, Kenya, Malawi, Mozambique, South Africa, Tanzania, Zambia, Zimbabwe and Madagascar. Its natural habitats are dry savanna and...

Melissa's yellow-eared bat
Melissa's yellow-eared bat
Vampyressa melissa

Melissa's yellow-eared bat (*Vampyressa melissa*) is a species of bat in the family Phyllostomidae. It is found in southern Colombia, Ecuador and Peru.

Meller's mongoose
Meller's mongoose
Rhynchogale melleri

Meller's mongoose (*Rhynchogale melleri*) is a species of mongoose found in Africa. It occurs in Democratic Republic of the Congo, Eswatini, Malawi, Mozambique, South Africa, Tanzania, Zambia and Zimbabwe. It is the only member of the genus...

Melon-Headed Whale
Melon-Headed Whale
Peponocephala electra

The Melon-headed whale (*Peponocephala electra*) belongs to the oceanic dolphin family. It is closely related to the Pygmy killer whale and Pilot whale. Collectively these dolphin species are known by the common name blackfish. Melon-headed whales...

Menchu's little yellow bat
Menchu's little yellow bat
Rhogeessa menchuae

Menchu's little yellow bat (*Rhogeessa menchuae*) is a species of vesper bat found in Central America. It was described as a new species in 2012.

Mendoza tuco-tuco
Mendoza tuco-tuco
Ctenomys mendocinus

The Mendoza tuco-tuco (*Ctenomys mendocinus*) is a species of rodent in the family Ctenomyidae.

Mentawai langur
Mentawai langur
Presbytis potenziani

The Mentawai langur (*Presbytis potenziani*) is a species of primate in the family Cercopithecidae. It is endemic to the Mentawai Islands in Indonesia. Its natural habitat is subtropical or tropical dry forests. The Siberut langur (*P. siberu*) was...

Mentawai long-tailed giant rat
Mentawai long-tailed giant rat
Leopoldamys siporanus

The Mentawai long-tailed giant rat (*Leopoldamys siporanus*) is a species of rodent in the family Muridae. It is endemic to the Mentawai Islands archipelago off the west coast of Sumatra, in Indonesia. Its natural habitat is subtropical or tropical...

Mentawai rat
Mentawai rat
Rattus lugens

The Mentawai rat (*Rattus lugens*) is a species of rodent in the family Muridae. It is found only in the Mentawai Islands of Indonesia, on the islands of Siberut, Sipora, Pagai Utara (North Pagai island), and Pagai Selatan (South Pagai island).
Mentawai squirrel
Mentawai squirrel
Callosciurus melanogaster

The Mentawai squirrel (*Callosciurus melanogaster*) is a species of rodent in the family Sciuridae. It is one of 20 or so species endemic to the Mentawai Islands off the west coast of Sumatra. There are three subspecies: *C. m. melanogaster*, *C. m.*
Mentawai three-striped squirrel
Mentawai three-striped squirrel
Lariscus obscurus

The Mentawai three-striped squirrel (*Lariscus obscurus*) is a species of rodent in the family Sciuridae. It is endemic to Indonesia. Its natural habitat is subtropical or tropical dry forests. It is threatened by habitat loss.
Mentawi flying squirrel
Mentawi flying squirrel
Iomys sipora

The Mentawi flying squirrel (*Iomys sipora*) is a species of rodent in the family Sciuridae. It is endemic to Indonesia where it is only known from the Mentawai Islands (Sipura and North Pagai). Its natural habitat is lowland tropical primary...
Menzbier's marmot
Menzbier's marmot
Marmota menzbieri

The Menzbier's marmot (*Marmota menzbieri*) is a species of rodent in the family Sciuridae from Central Asia. Its name commemorates Russian zoologist Mikhail Aleksandrovich Menzbier.
Menzies' echymipera
Menzies' echymipera
Echymipera echinista

Menzies' echymipera (*Echymipera echinista*), or Menzies' spiny bandicoot or Fly River bandicoot, is a species of marsupial in the family Peramelidae. It is endemic to Papua New Guinea.
Mérida brocket
Mérida brocket
Mazama bricenii

The Mérida brocket (*Mazama bricenii*), also known as the Meroia brocket or rufous brocket, is a small species of deer. It is found in forest and páramo at altitudes of 1,000–3,500 metres (3,300–11,500 ft) in the Andes of northern Colombia and...
Merida small-eared shrew
Merida small-eared shrew
Cryptotis meridensis

The Merida small-eared shrew (*Cryptotis meridensis*) is a species of shrew that is endemic to Venezuela.
Meridional serotine
Meridional serotine
Eptesicus isabellinus

The meridional serotine (*Eptesicus isabellinus*) is a species of bat native to the Iberian Peninsula, Morocco, Algeria, Tunisia and Libya.

Merriam's Chipmunk
Merriam's Chipmunk
Neotamias merriami

Merriam's chipmunk (*Neotamias merriami*) is a species of rodent in the family Sciuridae. It is found in central and southern California in the United States and a small area in northern Baja California, Mexico. The dental formula for *Tamias*...

Merriam's elk
Merriam's elk
Cervus canadensis merriami

The Merriam's elk (*Cervus canadensis merriami*) is an extinct subspecies of elk once found in the arid lands of the southwestern United States (in Arizona, New Mexico and Texas), as well as in Mexico. Uncontrolled hunting and cattle grazing since...

Merriam's ground squirrel
Merriam's ground squirrel
Urocitellus canus

Merriam's ground squirrel (*Urocitellus canus*) is a species of rodent in the family Sciuridae. It occurs in the western United States in Idaho, Nevada, and Oregon.

Merriam's Kangaroo Rat
Merriam's Kangaroo Rat
Dipodomys merriami

Merriam's kangaroo rat (*Dipodomys merriami*) is a species of rodent in the family Heteromyidae. The species name commemorates Clinton Hart Merriam.

Merriam's pocket gopher
Merriam's pocket gopher
Cratogeomys merriami

Merriam's pocket gopher (*Cratogeomys merriami*) is a species of rodent in the family Geomyidae. It is endemic to Mexico, where it is found in the area of the Valley of Mexico and the Valley of Toluca at elevations from 1800 to 4000 m. Its favored...

Merriam's pocket mouse
Merriam's pocket mouse
Perognathus merriami

Merriam's pocket mouse (*Perognathus merriami*) is a species of rodent in the family Heteromyidae. It is found in northeast Mexico and New Mexico, Oklahoma and Texas in the United States. Its habitat is shortgrass prairie, desert areas with scrub...

Merriam's shrew
Merriam's shrew
Sorex merriami

Merriam's shrew (*Sorex merriami*) is a species of mammal in the family Soricidae. It is endemic to the western United States and extreme southern British Columbia in Canada.

Merriam's small-eared shrew
Merriam's small-eared shrew
Cryptotis merriami

Merriam's small-eared shrew (*Cryptotis merriami*) is a species of mammal in the family Soricidae. It is found in Chiapas, Guatemala, El Salvador, Honduras, Nicaragua and Costa Rica.

Mesquite mouse
Mesquite mouse
Peromyscus merriami

The mesquite mouse (*Peromyscus merriami*) is a species of rodent in the family Cricetidae found in Mexico and in Arizona in the United States.

Mexican agouti

Mexican agouti

Dasyprocta mexicana

The Mexican agouti (*Dasyprocta mexicana*), also known as the Mexican black agouti, is a species of rodent in the family Dasyproctidae. It is native to lowland evergreen forest and second growth in southern Mexico (Veracruz, Oaxaca, Chiapas and...

Mexican big-eared bat

Mexican big-eared bat

Corynorhinus mexicanus

The Mexican big-eared bat (*Corynorhinus mexicanus*) is a species of vesper bat endemic to Mexico. They are nocturnal and insectivorous. Their very large ears are located across their foreheads, and when captured, the bats are observed to curl...

Mexican Bobcat

Mexican Bobcat

Lynx rufus escuinapae

The Mexican bobcat (*Lynx rufus escuinapae* syn. *Lynx rufus oaxacensis*) is a population of the bobcat in Mexico. As of 2017, it is uncertain whether or not this is a valid subspecies.

Mexican cottontail

Mexican cottontail

Sylvilagus cunicularius

The Mexican cottontail (*Sylvilagus cunicularius*) is a species of cottontail rabbit in the family Leporidae. It is endemic to Mexico where its natural habitats are temperate forests, subtropical or tropical dry forests and pastureland.

Mexican deer mouse

Mexican deer mouse

Peromyscus mexicanus

The Mexican deer mouse (*Peromyscus mexicanus*) is a species of forest-dwelling rodent in the family Cricetidae. It is found in southern Mexico and throughout much of Central America.

Mexican dog-faced bat

Mexican dog-faced bat

Cynomops mexicanus

The Mexican dog-faced bat (*Cynomops mexicanus*) is a bat species of the family Molossidae from Central America. It is found from Nayarit in Mexico to Costa Rica at elevations up to 1500 m. It was formerly considered a subspecies of *C. greenhalli*....

Mexican fox squirrel

Mexican fox squirrel

Sciurus nayaritensis

The Mexican fox squirrel (*Sciurus nayaritensis*) is a species of tree squirrel found throughout the Sierra Madre Occidental of Mexico as far south as Jalisco – and northward into the Chiricahua Mountains of southeastern Arizona, U.S. This species,...

Mexican Free-Tailed Bat

Mexican Free-Tailed Bat

Tadarida brasiliensis

The Mexican free-tailed bat (*Tadarida brasiliensis*) is a medium-sized bat native to the Americas and is regarded as one of the most abundant mammals in North America. It has been claimed to have the fastest horizontal speed (as opposed to stoop...
Mexican funnel-eared bat
Mexican funnel-eared bat
Natalus stramineus

The Mexican funnel-eared bat (*Natalus stramineus*) is a bat species found in Central America and the Caribbean.
Mexican Gray Squirrel
Mexican Gray Squirrel
Sciurus aureogaster

The Mexican gray squirrel (or red-bellied squirrel) (*Sciurus aureogaster*) is a tree squirrel in the genus *Sciurus* native to Guatemala and eastern and southern Mexico. It has been introduced to the Florida Keys. The alternate name should not be...
Mexican Gray Wolf
Mexican Gray Wolf
Canis lupus baileyi

The Mexican gray wolf (*Canis lupus baileyi*) is a subspecies of the Gray wolf, commonly known as "El lobo". Once there were thousands of these wolves but in the U.S. many had been killed by the mid-1970s, and just a handful existed in zoos. In...
Mexican greater funnel-eared bat
Mexican greater funnel-eared bat
Natalus mexicanus

The Mexican greater funnel-eared bat (*Natalus mexicanus*) is a species of bat found in Central America. While initially and currently described as a species, from 1959–2006 it was considered a subspecies of the Mexican funnel-eared bat, *Natalus*...
Mexican ground squirrel
Mexican ground squirrel
Ictidomys mexicanus

The Mexican ground squirrel (*Ictidomys mexicanus*) is a species of rodent in the family *Sciuridae*. It is found in Mexico and the United States. One of its closest relatives is the thirteen-lined ground squirrel (*Ictidomys tridecemlineatus*).
Mexican Hairy Dwarf Porcupine
Mexican Hairy Dwarf Porcupine
Coendou mexicanus

The Mexican hairy dwarf porcupine (*Coendou mexicanus*) is a species of rodent found in Central and North America. Its closest relatives are the Andean porcupine (*Coendou quichua*) and the Stump-tailed porcupine (*Coendou rufescens*).
Mexican harvest mouse
Mexican harvest mouse
Reithrodontomys mexicanus

The Mexican harvest mouse (*Reithrodontomys mexicanus*) is a species of rodent in the family *Cricetidae*. It is found in Colombia, Costa Rica, Ecuador, El Salvador, Guatemala, Honduras, Mexico, Nicaragua, and Panama in a variety of habitats at...
Mexican howler
Mexican howler
Alouatta palliata mexicana

The Mexican howler (*Alouatta palliata mexicana*) is a subspecies of the mantled

howler, *A. palliata*. This subspecies is found predominantly in forests between south eastern Mexico and north eastern Peru. Typical of its species, the Mexican howler...

Mexican long-tailed shrew

Mexican long-tailed shrew

Sorex oreopolus

The Mexican long-tailed shrew (*Sorex oreopolus*) is a species of mammal in the family Soricidae. It is endemic to Mexico.

Mexican long-tongued bat

Mexican long-tongued bat

Choeronycteris mexicana

The Mexican long-tongued bat (*Choeronycteris mexicana*) is a species of bat in the family Phyllostomidae. It is monotypic within the genus *Choeronycteris*. The species is found in El Salvador, Guatemala, Honduras, Mexico, and the United States.

Mexican mouse opossum

Mexican mouse opossum

Marmosa mexicana

The Mexican mouse opossum (*Marmosa mexicana*) is a species of North American opossum in the family Didelphidae.

Mexican Prairie Dog

Mexican Prairie Dog

Cynomys mexicanus

The Mexican prairie dog (*Cynomys mexicanus*) is a burrowing rodent native to Mexico. They are closely related to squirrels, chipmunks, and marmots. Mexican prairie dogs originated about 230,000 years ago from a peripherally isolated population of...

Mexican pronghorn

Mexican pronghorn

Antilocapra americana mexicana

The Mexican pronghorn (*Antilocapra americana mexicana*) is a pronghorn native to Mexico. It was found in the United States (in Arizona), but is considered extirpated there.

Mexican shrew

Mexican shrew

Megasorex gigas

The Mexican shrew (*Megasorex gigas*) is a species of mammal from the subfamily Soricinae in the family Soricidae. It is monotypic within the genus *Megasorex* and is endemic to Mexico.

Mexican small-eared shrew

Mexican small-eared shrew

Cryptotis mexicana

The Mexican small-eared shrew (*Cryptotis mexicana*) is a species of mammal in the family Soricidae. It is endemic to Mexico.

Mexican Spider Monkey

Mexican Spider Monkey

Ateles geoffroyi vellerosus

The Mexican spider monkey (*Ateles geoffroyi vellerosus*) is a subspecies of Geoffroy's spider monkey, and is one of the largest types of New World monkey. It is considered to be endangered according to the IUCN Red List since 2020, mostly due to...

Mexican spiny pocket mouse

Mexican spiny pocket mouse

Heteromys irroratus

The Mexican spiny pocket mouse (*Heteromys irroratus*) is a species of rodent in the family Heteromyidae. It is native to Mexico and Texas in the United States where it is found in dry, scrubby habitats. The IUCN has assessed it as being of "least...

Mexican volcano mouse
Mexican volcano mouse
Neotomodon alstoni

The Mexican volcano mouse (*Neotomodon alstoni*) is a species of rodent in the family Cricetidae endemic to high elevation areas of the Trans-Mexican Volcanic Belt.

Mexican vole
Mexican vole
Microtus mexicanus

The Mexican vole (*Microtus mexicanus*) is a species of vole. The Hualapai Mexican vole (*M. m. hualapaiensis*) was formerly recognized as a subspecies and was a protected taxon under the United States Endangered Species Act until 2017. Several...

Mexican water mouse
Mexican water mouse
Rheomys mexicanus

The Mexican water mouse, Mexican fishing mouse or Goodwin's water mouse (*Rheomys mexicanus*), is a species of semiaquatic rodent in the family Cricetidae. It has a restricted range in the state of Oaxaca in southern Mexico, Threatened by...

Miahuatlán cotton rat
Miahuatlán cotton rat
Sigmodon planifrons

The Miahuatlán cotton rat (*Sigmodon planifrons*) was formerly considered a rodent species in the family Cricetidae. It is found only on the Pacific slope of the Sierra de Miahuatlán in the Mexican state of Oaxaca, where it lives in deciduous...

Mianzini African mole-rat
Mianzini African mole-rat
Tachyoryctes annectens

The Mianzini mole-rat or Mianzini African mole-rat (*Tachyoryctes annectens*) is a species of rodent in the family Spalacidae endemic to Kenya. Its natural habitats are dry savanna, pastureland, and rural gardens. Some taxonomic authorities...

Michoacan deer mouse
Michoacan deer mouse
Osgoodomys banderanus

The Michoacan deer mouse (*Osgoodomys banderanus*) is a species of rodent in the family Cricetidae. It is the only species in the genus *Osgoodomys*, and is endemic to Mexico.

Michoacan pocket gopher
Michoacan pocket gopher
Zygogeomys trichopus

The Michoacan pocket gopher (*Zygogeomys trichopus*) is a species of rodent in the family Geomyidae. It is monotypic within the genus *Zygogeomys*. It is endemic to Mexico where its natural habitat is temperate, high-altitude forests. Its numbers...

Micklem's mole-rat
Micklem's mole-rat
Fukomys micklemi

The Micklem's mole-rat (*Fukomys micklemi*) is a species of rodent in the family Bathyergidae. It is found in Zambia. It is a subterranean species of chisel tooth diggers. It is a social species, and this has been suggested to account for its more...

Microcebus manitatra
Microcebus manitatra
Microcebus manitatra

The Bemanasy mouse lemur, *Microcebus manitatra*, is a species of mouse lemur described in 2016 from Madagascar. It was discovered by a team of researchers at the German Primate Center. It was initially discovered among closely related species such...

Microgale grandidieri
Microgale grandidieri
Microgale grandidieri

Microgale grandidieri, also known as Grandidier's shrew tenrec, is a species of shrew tenrec occurring in the dry forests of western and southwestern Madagascar. Populations of this species were formerly included in *Microgale brevicaudata* ; M....

Microgale macpheeii
Microgale macpheeii
Microgale macpheeii

Microgale macpheeii is an extinct shrew tenrec from southeastern Madagascar. It is known only from two partial skulls found in Andrahomana cave, which radiocarbon dating of associated rodent remains suggests are about 3000 years old. It is the...

Micromys erythrotis
Micromys erythrotis
Micromys erythrotis

The Indochinese harvest mouse (*Micromys erythrotis*) is a species of rodent in the genus *Micromys* that is native to Asia.

Micronomus
Micronomus
Micronomus norfolkensis

Micronomus norfolkensis is a species of molossid bat, a family of flying mammals. The bat is endemic to Australia, where it occurs from southeastern Queensland to eastern New South Wales. They are the sole species of genus *Micronomus* and referred...

Micronycteris giovanniae
Micronycteris giovanniae
Micronycteris giovanniae

Micronycteris giovanniae is a species of leaf-nosed bat found in Ecuador.

Microryzomys altissimus
Microryzomys altissimus
Microryzomys altissimus

Microryzomys altissimus, also known as the Páramo colilargo or highland small rice rat, is a species of rodent in the genus *Microryzomys* of family Cricetidae. It is found in Colombia, Ecuador, and Peru, but the Colombian segment may be a separate...

Microryzomys minutus
Microryzomys minutus
Microryzomys minutus

Microryzomys minutus, also known as the montane colilargo or the forest small rice rat, is a species of rodent in the genus *Microryzomys* of family Cricetidae. It is found in Bolivia, Colombia, Ecuador, Peru, and Venezuela, but these populations...
Midas free-tailed bat
Midas free-tailed bat
Mops midas

The Midas free-tailed bat (*Mops midas*) is a species of bat scientifically classified in the order Chiroptera and the family Molossidae. It is distributed from western Africa to Saudi Arabia and further south. Its natural habitats are dry...
Midday jird
Midday jird
Meriones meridianus

The midday jird, or midday gerbil (*Meriones meridianus*), is a species of rodent in the family Muridae and native to sandy deserts in Afghanistan, China, Iran, Azerbaijan, Kazakhstan, Kyrgyzstan, Mongolia, Russia, Tajikistan, Turkmenistan, and...
Middendorff's vole
Middendorff's vole
Microtus middendorffi

Middendorff's vole (*Microtus middendorffi*) is a species of rodent in the family Cricetidae. It is found only in Russia, most commonly north Siberia. The common name commemorates Alexander Theodor von Middendorff (1815–1894), a German-Russian...
Middle East blind mole-rat
Middle East blind mole-rat
Spalax ehrenbergi

The Middle East blind mole-rat or Palestine mole-rat (*Nannospalax ehrenbergi*) is a species of rodent in the family Spalacidae.
Millard's rat
Millard's rat
Dacnomys millardi

Millard's rat (*Dacnomys millardi*) is a species of rodent in the family Muridae. It is the only species in the genus *Dacnomys*. It is found in China, India, Laos, and Nepal.
Miller's long-tongued bat
Miller's long-tongued bat
Glossophaga longirostris

Miller's long-tongued bat (*Glossophaga longirostris*) is a bat species from northern South America and the Windward Islands of the Lesser Antilles. It is found in northern Brazil, Venezuela, Colombia, Guyana, Trinidad and Tobago, Grenada, the...
Miller's mastiff bat
Miller's mastiff bat
Molossus pretiosus

Miller's mastiff bat (*Molossus pretiosus*) is a species of bat in the family Molossidae. It is found in Brazil, Colombia, Costa Rica, Guyana, Mexico, Nicaragua, and Venezuela.
Miller's saki
Miller's saki
Pithecia milleri

Miller's saki (*Pithecia milleri*), also known as Miller's monk saki, is a species

of saki monkey, a type of New World monkey. Its range includes parts of southwestern Colombia, northeastern Ecuador, and perhaps adjacent areas in Peru. This...

Miller's striped mouse
Miller's striped mouse
Hybomys planifrons

Miller's striped mouse or the Liberian forest hybomys (*Hybomys planifrons*) is a species of rodent in the family Muridae. It is found in Ivory Coast, Guinea, Liberia, and Sierra Leone. Its natural habitats are subtropical or tropical dry forest,...

Millet's leopoldamys
Millet's leopoldamys
Leopoldamys milleti

Millet's leopoldamys (*Leopoldamys milleti*) is a species of rodent from the family Muridae. It lives in the Langbian highlands of southern Vietnam, although its distribution limits are currently unresolved. It is listed as Least Concern on the...

Milne-Edwards' sportive lemur
Milne-Edwards' sportive lemur
Lepilemur edwardsi

Milne-Edwards' sportive lemur (*Lepilemur edwardsi*), or Milne-Edwards' weasel lemur, is a species of lemur in the family Lepilemuridae. It is endemic to Madagascar. Its natural habitat is subtropical or tropical dry forests. It is threatened by...

Milne-Edwards's sifaka
Milne-Edwards's sifaka
Propithecus edwardsi

Milne-Edwards's sifaka (*Propithecus edwardsi*), or Milne-Edwards's simpona, is a large arboreal, diurnal lemur endemic to the eastern coastal rainforest of Madagascar. Milne-Edwards's sifaka is characterized by a black body with a light-colored...

Milton's titi
Milton's titi
Plecturocebus miltoni

Milton's titi (*Plecturocebus miltoni*) is a species of titi, a type of New World monkey, from southern Amazon rainforest, Brazil. It was named after the Brazilian primatologist Milton Thiago de Mello. Milton's titi was discovered in 2011 by Julio...

Mimic tree rat
Mimic tree rat
Xenuromys barbatus

The mimic tree rat, rock-dwelling giant rat, or rock-dwelling rat (*Xenuromys barbatus*) is a species of rodent in the family Muridae found in West Papua, Indonesia and Papua New Guinea.

Minahassa pipistrelle
Minahassa pipistrelle
Pipistrellus minahassae

The Minahassa pipistrelle (*Pipistrellus minahassae*) is a species of vesper bat. It can be found in Indonesia.

Minahassa ranee mouse
Minahassa ranee mouse
Haeromys minahassae

The Minahassa ranee mouse (*Haeromys minahassae*) is a species of rodent in the family Muridae. It is found only on the island of Sulawesi (Indonesia).

Minca spiny rat

Minca spiny rat

Proechimys mincae

The Minca spiny rat (*Proechimys mincae*) is a species of rodent in the family Echimyidae. It is endemic to Colombia.

Mindanao flying squirrel

Mindanao flying squirrel

Petinomys mindanensis

The Mindanao flying squirrel (*Petinomys mindanensis*) is a common species of squirrel that is endemic to the Philippines.

Mindanao hairy-tailed rat

Mindanao hairy-tailed rat

Batomys salomonseni

The Mindanao hairy-tailed rat (*Batomys salomonseni*) is one of five species of rodent in the genus *Batomys*. It is in the diverse family Muridae. This species is found only in the Philippines.

Mindanao lowland forest mouse

Mindanao lowland forest mouse

Apomys littoralis

The Mindanao lowland forest mouse (*Apomys littoralis*) is a species of rodent in the family Muridae. It is found only in the Philippines.

Mindanao montane forest mouse

Mindanao montane forest mouse

Apomys insignis

The Mindanao montane forest mouse (*Apomys insignis*) is a species of rodent in the family Muridae. It is found only in the Philippines.

Mindanao mountain rat

Mindanao mountain rat

Limnomys sibuanus

The Mindanao mountain rat or long-tailed moss mouse (*Limnomys sibuanus*), is a species of rodent in the family Muridae. It is found only in the Philippines, where it is present at high altitudes in the Kitanglad Mountain Range on the island of...

Mindanao pygmy fruit bat

Mindanao pygmy fruit bat

Alionycteris paucidentata

The Mindanao pygmy fruit bat (*Alionycteris paucidentata*) is a species of megabat in the family Pteropodidae. It is monotypic within the genus *Alionycteris*. It is endemic to the Philippines. Its natural habitat is subtropical or tropical dry...

Mindanao shrew

Mindanao shrew

Crocidura beatus

The Mindanao shrew (*Crocidura beatus*) is a species of mammal in the family Soricidae. It is endemic to the Philippines. Its natural habitat is subtropical or tropical dry forests. It is threatened by habitat loss.

Mindanao shrew-rat

Mindanao shrew-rat

Crunomys melanius

The Mindanao shrew-rat (*Crunomys melanius*) is a species of rodent in the family Muridae. It is found only in the Philippines.

Mindanao squirrel

Mindanao squirrel

Sundasciurus mindanensis

The Mindanao squirrel (*Sundasciurus mindanensis*) is a species of rodent in the family Sciuridae. It is endemic to the Philippines.

Mindanao treeshrew

Mindanao treeshrew

Tupaia everetti

The Mindanao treeshrew (*Tupaia everetti*), also called the Philippine tree shrew, is a species of treeshrew endemic to the Mindanao region in the Philippines. It was formerly considered the only member of the genus *Urogale*, but that genus was...

Mindomys

Mindomys

Mindomys hammondi

Mindomys is a genus of sigmodontine rodents in the family Cricetidae. It includes two species known only from Ecuador, Hammond's rice rat (*Mindomys hammondi*) and the Kutukú rat (*Mindomys kutuku*).:16

Mindoro black rat

Mindoro black rat

Rattus mindorensis

The Mindoro black rat (*Rattus mindorensis*) is a species of rodent in the family Muridae. It is found only in hilly and forested areas of Mindoro island, the Philippines.

Mindoro climbing rat

Mindoro climbing rat

Anonymomys mindorensis

The Mindoro climbing rat or Mindoro rat (*Anonymomys mindorensis*) is a species of rodent in the family Muridae. It is found only in the Philippines, and is known only from Ilong Peak in the Halcon Mountains. It is the only species in the genus...

Mindoro shrew

Mindoro shrew

Crocidura mindorus

The Mindoro shrew (*Crocidura mindorus*) is a species of mammal in the family Soricidae. It is endemic to the Philippines. It is threatened by habitat loss.

Mindoro stripe-faced fruit bat

Mindoro stripe-faced fruit bat

Styloctenium mindorensis

The Mindoro stripe-faced fruit bat (*Styloctenium mindorensis*), nicknamed the "flying fox" for its foxlike face (although it is not a flying fox bat), is a species of large Philippine megabat. It is endemic to the island of Mindoro. The Mindoro...

Mindoro striped rat

Mindoro striped rat

Chrotomys mindorensis

The Mindoro striped rat (*Chrotomys mindorensis*) is a species of rodent in the family Muridae. It is found only in the Philippines.

Miniopterus aelleni

Miniopterus aelleni

Miniopterus aelleni

Miniopterus aelleni is a bat in the genus *Miniopterus* that occurs on Anjouan in the Comoros and in northern and western Madagascar. It is a small brown bat; its forearm length is 35 to 41 mm (1.4 to 1.6 in). The long tragus (a projection in the...

Miniopterus brachytragos

Miniopterus brachytragos

Miniopterus brachytragos

Miniopterus brachytragos is a bat in the genus *Miniopterus* that occurs in northern and western Madagascar. Populations of this species have historically been included in *Miniopterus manavi*, but molecular data published in 2008 and 2009 indicate...

Miniopterus griveaudi

Miniopterus griveaudi

Miniopterus griveaudi

Miniopterus griveaudi is a bat in the genus *Miniopterus* found on Grande Comore and Anjouan in the Comoros and in northern and western Madagascar. First described in 1959 from Grande Comore as a subspecies of the mainland African *M. minor*, it was...

Miniopterus macrocneme

Miniopterus macrocneme

Miniopterus macrocneme

Miniopterus macrocneme is a bat in the genus *Miniopterus*. It is found primarily in the Loyalty Islands of New Caledonia, though it is also found in New Guinea. It has been considered a subspecies of *Miniopterus pusillus* in recent years. It lives...

Miniopterus maghrebensis

Miniopterus maghrebensis

Miniopterus maghrebensis

Miniopterus maghrebensis, also known as the Maghrebian bent-wing bat, is a species of bat found in North Africa.

Miniopterus mahafaliensis

Miniopterus mahafaliensis

Miniopterus mahafaliensis

Miniopterus mahafaliensis is a bat in the genus *Miniopterus* that occurs in southwestern Madagascar. Populations of this species have historically been included in *Miniopterus manavi*, but molecular data published in 2008 and 2009 indicate that...

Miniopterus newtoni

Miniopterus newtoni

Miniopterus newtoni

Miniopterus newtoni is a species of bat that is endemic to São Tomé and Príncipe.

Miniopterus pallidus

Miniopterus pallidus

Miniopterus pallidus

Miniopterus pallidus, the pale bent-wing bat, is a species of bat found in Middle-East. Until 2010, it was considered as a sub-species of *M. schreibersii*.

Miniopterus sororculus

Miniopterus sororculus

Miniopterus sororculus

Miniopterus sororculus is a species of bat endemic to the highland forests of Madagascar.

Miniopterus tao

Miniopterus tao

Miniopterus tao

Miniopterus tao is a fossil bat in the genus Miniopterus from the Pleistocene of Zhoukoudian in China. It is known from a number of mandibles (lower jaws), which were initially identified as the living species Miniopterus schreibersii in 1963...
Miniopterus zapfei
Miniopterus zapfei
Miniopterus zapfei

Miniopterus zapfei is a fossil bat in the genus Miniopterus from the middle Miocene of France. First described in 2002, it is known only from the site of La Grive M, where it occurs with another fossil Miniopterus species, the smaller and more...
Minor epauletted fruit bat
Minor epauletted fruit bat
Epomophorus minor

The minor epauletted fruit bat (Epomophorus minor) is a species of megabat in the family Pteropodidae. It is found in Zambia, Tanzania, Mozambique and Kenya.
Minor red bat
Minor red bat
Lasiurus minor

The minor red bat (Lasiurus minor) is a species of bat from the family Vespertilioninae.
Minute fruit bat
Minute fruit bat
Cynopterus minutus

The minute fruit bat (Cynopterus minutus) is a species of megabat within the family Pteropodidae. It is found in Sumatra, Java, Borneo and Sulawesi.
Miophocaena
Miophocaena
Miophocaena nishinoi

Miophocaena nishinoi is an extinct species of porpoise from the Late Miocene Koetoi Formation of Japan, dating to around 6.4–5.5 million years ago (mya), represented by a partial skull. the genus name derives from Ancient Greek mio for the...
Mira climbing rat
Mira climbing rat
Tylomys mirae

The Mira climbing rat (Tylomys mirae) is a species of rodent in the family Cricetidae. It is found in Colombia and Ecuador.
Mirza's eastern moss rat
Mirza's eastern moss rat
Mirzamys norahae

Mirza's eastern moss rat (Mirzamys norahae) is a species of rodent endemic to New Guinea.
Mirza's western moss rat
Mirza's western moss rat
Mirzamys louiseae

Mirza's western moss rat (Mirzamys louiseae) is a species of rodent endemic to Papua New Guinea.
Mishmi Takin
Mishmi Takin
Budorcas taxicolor taxicolor

The Mishmi takin (Budorcas taxicolor taxicolor) is an endangered goat-antelope

native to Asia. It is a subspecies of takin.
Misonne's soft-furred mouse
Misonne's soft-furred mouse
Praomys misonnei

Misonne's soft-furred mouse or Misonne's praomys (*Praomys misonnei*) is a species of rodent in the family Muridae. It is found in Democratic Republic of the Congo, Kenya, and Uganda. Its natural habitats are subtropical or tropical moist lowland...
Miss Ryley's soft-furred rat
Miss Ryley's soft-furred rat
Millardia kathleenae

Miss Ryley's soft-furred rat (*Millardia kathleenae*) is a species of rodent in the family Muridae. It is found only in central Myanmar, and was also recorded on Mount Popa.
Miss Waldron's Red Colobus
Miss Waldron's Red Colobus
Piliocolobus waldronae

Miss Waldron's red colobus (*Piliocolobus waldronae*) is a species of the red colobus native to West Africa. It has not been officially sighted since 1978 and was considered extinct in 2000. However, new evidence suggests that a very small number...
Mistratoan yellow-shouldered bat
Mistratoan yellow-shouldered bat
Sturnira mistratensis

The Mistratoan yellow-shouldered bat (*Sturnira mistratensis*), is a species of leaf-nosed bat indigenous to the Cordillera Occidental, in the Mistrato municipality in Risaralda, Colombia. Since existing information derives exclusively from the...
Mitchell's hopping mouse
Mitchell's hopping mouse
Notomys mitchellii

Mitchell's hopping mouse (*Notomys mitchellii*) also known as the pankot, is the largest extant member of the genus *Notomys*, weighing between 40 and 60 g (1.4 and 2.1 oz). *N. mitchellii* is a bipedal rodent with large back legs, similar to a jerboa...
Mitred horseshoe bat
Mitred horseshoe bat
Rhinolophus mitratus

The mitred horseshoe bat (*Rhinolophus mitratus*) is a species of bat in the family Rhinolophidae. It is endemic to India. Little is known about the species, because it is known only from the holotype, the specimen used to describe the species to...
Mittendorf's striped grass mouse
Mittendorf's striped grass mouse
Lemniscomys mittendorfi

Mittendorf's lemniscomys or Mittendorf's striped grass mouse (*Lemniscomys mittendorfi*) is a species of rodent in the family Muridae. It is endemic to Cameroon where it is found at high elevations on a single mountain. Its natural habitat is...
Mittermeier's mouse lemur
Mittermeier's mouse lemur
Microcebus mittermeieri

Mittermeier's mouse lemur (*Microcebus mittermeieri*) is a species of mouse lemur

known only from Anjanaharibe-Sud Special Reserve in Madagascar. The species is named in honor of primatologist Russell Mittermeier, who is President of Conservation...

Mittermeier's sportive lemur
Mittermeier's sportive lemur
Lepilemur mittermeieri

Mittermeier's sportive lemur (*Lepilemur mittermeieri*) is a sportive lemur endemic to the Ampasindava Peninsula in Madagascar.

Mo's spiny rat
Mo's spiny rat
Maxomys moi

Mo's spiny rat (*Maxomys moi*) is a species of rodent in the family Muridae. It is found in Laos and Vietnam.

Mogollon mountain wolf
Mogollon mountain wolf
Canis lupus mogollonensis

The Mogollon mountain wolf (*Canis lupus mogollonensis*) is an extinct subspecies of gray wolf whose range once included Arizona and New Mexico. It is darker than its more northern cousins, and has a highly arched frontal bone.

Mohave Ground Squirrel
Mohave Ground Squirrel
Xerospermophilus mohavensis

The Mohave ground squirrel (*Xerospermophilus mohavensis*) is a species of ground squirrel found only in the Mojave Desert in California. The squirrel was first described in 1886 by Frank Stephens of San Diego. It is listed as a threatened species...

Mohol Bushbaby
Mohol Bushbaby
Galago moholi

The Mohol bushbaby (*Galago moholi*) is a medium-sized primate native to southern Africa. It is physically very similar to the Senegal bushbaby and was formerly considered to be its southern variety.

Molaccan prehensile-tailed rat
Molaccan prehensile-tailed rat
Rattus morotaiensis

The Molaccan prehensile-tailed rat (*Rattus morotaiensis*) is a species of rodent in the family Muridae. It is found only in the Halmahera Islands of Indonesia, including Morotai, Halmahera, and Batjan islands.

Mole-like rice tenrec
Mole-like rice tenrec
Oryzorictes hova

The mole-like rice tenrec (*Oryzorictes hova*), also known as the fossorial tenrec or hova rice tenrec, is a species of mammal in the tenrec family. Like all other tenrecs, it is endemic to Madagascar. Its natural habitats are subtropical or...

Molina's grass mouse
Molina's grass mouse
Akodon molinae

Molina's grass mouse (*Akodon molinae*) is a species of rodent in the family Cricetidae. It is found only in Argentina.

Molina's Hog-Nosed Skunk
Molina's Hog-Nosed Skunk

Conepatus chinga

Molina's hog-nosed skunk (*Conepatus chinga*) is a skunk species from South America. Like all skunk species, Molina's hog-nosed skunk is known for its ability to spray a liquid with a strong, unpleasant scent from its anal glands.

Moloney's mimic bat
Moloney's mimic bat
Mimetillus moloneyi

Moloney's mimic bat (*Mimetillus moloneyi*) is a species of vesper bat. It can be found in Benin, Cameroon, Central African Republic, Republic of the Congo, Democratic Republic of the Congo, Ivory Coast, Equatorial Guinea, Ethiopia, Gabon, Ghana,...

Molossus alvarezi
Molossus alvarezi
Molossus alvarezi

Molossus alvarezi, or Alvarez's mastiff bat, is a species of bat in the family Molossidae, native to the Yucatán Peninsula. It lives within a relative homogenous environment within perennial forests, low forests, and a band of xeric vegetation.

Moluccan flying fox
Moluccan flying fox
Pteropus chrysoproctus

The Moluccan flying fox (*Pteropus chrysoproctus*), also known as the Ambon flying fox, is a species of megabat in the genus *Pteropus*. It is found in the low-lying forests (less than 250 m above sea level) of Seram Island (including Manusela...

Mona Monkey
Mona Monkey
Cercopithecus mona

The mona monkey (*Cercopithecus mona*) is an Old World monkey that lives in western Africa between Ghana and Cameroon. The mona monkey can also be found on the island of Grenada as it was transported to the island aboard slave ships headed to the...

Monard's African climbing mouse
Monard's African climbing mouse
Dendromus leucostomus

Monard's African climbing mouse (*Dendromus leucostomus*) is a species of rodent in the family Nesomyidae. It is endemic to east-central Angola, but only recorded in the type locality of Caluquembe. The IUCN red list of threatened species lists...

Monard's dormouse
Monard's dormouse
Graphiurus monardi

Monard's dormouse (*Graphiurus monardi*) is a species of rodent in the family Gliridae. It is found in Angola, Democratic Republic of the Congo, and possibly Zambia. Its natural habitat is moist Central Zambezian miombo woodlands savanna.

Moncton's mosaic-tailed rat
Moncton's mosaic-tailed rat
Paramelomys moncktoni

Moncton's mosaic-tailed rat (*Paramelomys moncktoni*, previously known as *Melomys moncktoni*) is a species of rodent in the family Muridae. It is found only in Papua New Guinea. The species lives in lowlands in the north-east and south-east portion...

Mondolfi's four-eyed opossum
Mondolfi's four-eyed opossum
Philander mondolfii

Mondolfi's four-eyed opossum (*Philander mondolfii*) is a South American species of opossum found in Colombia and Venezuela, first described in 2006. It inhabits foothills of the Cordillera de Mérida and those on the eastern side of the Cordillera...

Mongalla free-tailed bat
Mongalla free-tailed bat
Mops demonstrator

The Mongalla free-tailed bat (*Mops demonstrator*) is a species of bat in the family Molossidae. It is found in Burkina Faso, Cameroon, Democratic Republic of the Congo, Ivory Coast, Ghana, Kenya, Mali, Sudan, and Uganda. Its natural habitats are...

Mongalla gazelle
Mongalla gazelle
Eudorcas albonotata

The Mongalla gazelle (*Eudorcas albonotata*) is a species of gazelle found in the floodplain and savanna of South Sudan. It was first described by British zoologist Walter Rothschild in 1903. The taxonomic status of the Mongalla gazelle is widely...

Mongolian five-toed jerboa
Mongolian five-toed jerboa
Allactaga sibirica

The Mongolian five-toed jerboa or Siberian jerboa (*Allactaga sibirica*) is a species of rodent in the family Dipodidae. It is found in China, Kazakhstan, Mongolia, and Turkmenistan.

Mongolian gazelle
Mongolian gazelle
Procapra gutturosa

The Mongolian gazelle (*Procapra gutturosa*) is a medium-sized antelope native to Central Asian steppes.

Mongolian Gerbil
Mongolian Gerbil
Meriones unguiculatus

The Mongolian gerbil (*Meriones unguiculatus*) is a small burrowing rodent often used in science and kept as a small house pet. Their use in science dates back to the latter half of the 19th century but these small animals became popular as pets in...

Mongolian hamster
Mongolian hamster
Allocricetulus curtatus

The Mongolian hamster (*Allocricetulus curtatus*) is a species of rodent in the family Cricetidae. It is one of two members of the genus *Allocricetulus*, and is found in China and Mongolia. The Mongolian hamster is a wild hamster and cannot be...

Mongolian silver vole
Mongolian silver vole
Alticola semicanus

The Mongolian silver vole (*Alticola semicanus*) is a species of rodent in the family Cricetidae. It is found only in Mongolia.

Mongolian three-toed jerboa
Mongolian three-toed jerboa
Stylodipus sungorus

The Mongolian three-toed jerboa (*Stylodipus sungorus*) is a species of rodent in the family Dipodidae. It is found in Mongolia and possibly China.

Mongolian vole

Mongolian vole

Microtus mongolicus

The Mongolian vole (*Microtus mongolicus*) is a species of rodent in the family Cricetidae. It is found in China, Mongolia, and Russia.

Mongolian wild ass

Mongolian wild ass

Equus hemionus hemionus

The Mongolian wild ass (*Equus hemionus hemionus*), also known as Mongolian khulan, is the nominate subspecies of the onager. It is found in southern Mongolia and northern China. It was previously found in eastern Kazakhstan and southern Siberia...

Mongoose Lemur

Mongoose Lemur

Eulemur mongoz

The mongoose lemur (*Eulemur mongoz*) is a small primate in the family Lemnidae, native to Madagascar and introduced to the Comoros Islands. These arboreal animals have pointed faces, long, bushy tails, dark-brown upper parts, pale bellies, and...

Monito Del Monte

Monito Del Monte

Dromiciops gliroides

The "monkey of the mountains", the Monito del monte (*Dromiciops gliroides*), is not a monkey. It is a South American marsupial and is often referred to as a "living fossil" because it is the only living member of Microbiotheria, an otherwise...

Monjon

Monjon

Petrogale burbridgei

The monjon is a common name for the species *Petrogale burbridgei*, the smallest of the rock-wallabies (*Petrogale*), found in the north-west Australia. They are restricted to a small area of the Kimberley region and on nearby islands in the...

Monk Saki

Monk Saki

Pithecia monachus

The monk saki (*Pithecia monachus*) also known as Geoffroy's monk saki, is a species of saki monkey, a type of New World monkey, from South America. It is found in forested areas of northwestern Brazil and northeastern Peru.

Mono's short-tailed bat

Mono's short-tailed bat

Carollia monohernandezi

Mono's short-tailed bat (*Carollia monohernandezi*) is a leaf-nosed bat species from South America found in Colombia and Panama. Specimens have been collected in habitats ranging from dry to tropical rain forest at elevations ranging from 30 to...

Monodelphis saci

Monodelphis saci

Monodelphis saci

The Saci short-tailed opossum (*Monodelphis saci*), also known as gnome opossum, is a species of opossum endemic to the rainforests of Brazil.

Montagne d'Ambre dwarf lemur
Montagne d'Ambre dwarf lemur
Cheirogaleus andysabini

The Montagne d'Ambre dwarf lemur or Andy Sabin's dwarf lemur (*Cheirogaleus andysabini*) is a species of dwarf lemur known only from Amber Mountain National Park and the surrounding area in northern Madagascar. It was identified in 2005, but not...

Montane African climbing mouse
Montane African climbing mouse
Dendromus insignis

The montane African climbing mouse or remarkable climbing mouse (*Dendromus insignis*) is a species of rodent in the family Nesomyidae. It is found in Democratic Republic of the Congo, Kenya, Rwanda, Tanzania, and Uganda.

Montane Atlantic Forest rat
Montane Atlantic Forest rat
Delomys collinus

The montane Atlantic Forest rat (*Delomys collinus*) is a South American rodent species of the family Cricetidae. It is endemic to part of the Atlantic Forest region of southeastern Brazil, where it is found at elevations from 1000 to 2700 m in...

Montane bamboo rat
Montane bamboo rat
Dactylomys peruanus

The montane bamboo rat or Peruvian bamboo rat (*Dactylomys peruanus*), is a species of rodent in the family Echimyidae. It is found in Bolivia and Peru. Its natural habitat is subtropical or tropical moist lowland forests.

Montane cotton rat
Montane cotton rat
Sigmodon zanzonensis

The montane cotton rat (*Sigmodon zanzonensis*) is a rodent species in the family Cricetidae. It is found in the highlands of Chiapas, Mexico and Guatemala. The eastern and southern limits of its range have not yet been determined.

Montane fish-eating rat
Montane fish-eating rat
Neusticomys monticolus

The montane fish-eating rat (*Neusticomys monticolus*) is a species of semiaquatic rodent in the family Cricetidae. It inhabits the Andes Mountains of Colombia and Ecuador.

Montane grass mouse
Montane grass mouse
Akodon montensis

The montane grass mouse (*Akodon montensis*) is a rodent species from South America. It is found in Argentina, Brazil, Paraguay and Uruguay.

Montane Guinea Pig
Montane Guinea Pig
Cavia tschudii

The Montane guinea pig (*Cavia tschudii*) is a species of caviid rodent found in the Andes in South America. It is the likely main ancestor of *Cavia porcellus*, the Domestic cavy or Domestic guinea pig, which appears to be a hybrid that includes...

Montane long-nosed squirrel
Montane long-nosed squirrel

Hyosciurus heinrichi

The montane long-nosed squirrel (*Hyosciurus heinrichi*) is a species of rodent in the family *Sciuridae*. It is endemic to central Sulawesi, Indonesia. Its natural habitat is subtropical or tropical dry lowland grassland. It was discovered during a...

Montane monkey-faced bat

Montane monkey-faced bat

Pteralopex pulchra

The montane monkey-faced bat or montane flying monkey (*Pteralopex pulchra*) is a megabat endemic to the Solomon Islands. It is listed as a critically endangered species. Due to its imperiled status, it is identified by the Alliance for Zero...

Montane mouse shrew

Montane mouse shrew

Myosorex blarina

The montane mouse shrew (*Myosorex blarina*) is a species of mammal in the family *Soricidae* endemic to Uganda. Its natural habitats are subtropical or tropical moist montane forests and swamps. It is threatened by habitat loss.

Montane myotis

Montane myotis

Myotis oxyotus

The montane myotis (*Myotis oxyotus*) is a species of vesper bat. It is found in Bolivia, Colombia, Costa Rica, Ecuador, Panama, Peru, and Venezuela.

Montane Oldfield mouse

Montane Oldfield mouse

Thomasomys oreas

The montane Oldfield mouse (*Thomasomys oreas*) is a species of rodent in the family *Cricetidae*. It is found only in Bolivia.

Montane shaggy rat

Montane shaggy rat

Dasymys montanus

The montane shaggy rat (*Dasymys montanus*) is a species of rodent in the family *Muridae*. It is found in Uganda and possibly Democratic Republic of the Congo. Its natural habitats are subtropical or tropical moist montane forest, subtropical or...

Montane shrew

Montane shrew

Sorex monticolus

The montane shrew (*Sorex monticolus*) is a species of mammal in the family *Soricidae* commonly known as the dusky shrew. *Monticolus* is derived from the Latin root word *mons* meaning mountain. It is found in Alaska, western Canada, the western...

Montane shrew tenrec

Montane shrew tenrec

Microgale monticola

The montane shrew tenrec (*Microgale monticola*) is a species of mammal in the family *Tenrecidae*. It is endemic to Madagascar. Its natural habitat is subtropical or tropical moist montane forests.

Montane Sumatran white-bellied rat

Montane Sumatran white-bellied rat

Niviventer fraterculus

The montane Sumatran white-bellied rat (*Niviventer fraterus*), also known as the Montane Sumatran niviventer, is a species of rodent in the family Muridae. It is found in the montane forests along the mountains of western Sumatra, Indonesia.

Montane vole

Montane vole

Microtus montanus

The montane vole (*Microtus montanus*) is a species of vole native to the western United States and Canada.

Montane water vole

Montane water vole

Arvicola scherman

The montane water vole (*Arvicola scherman*) is a species of vole found throughout Europe, from the mountains of northern Spain through central Europe and eastwards through to central Romania. Initially regarded as a species, it was reassigned as...

Montane white-toothed shrew

Montane white-toothed shrew

Crocidura montis

The montane white-toothed shrew (*Crocidura montis*) is a species of mammal in the family Soricidae. It is found in Kenya, South Sudan, Tanzania, and Uganda. Its natural habitats are subtropical or tropical moist montane forest and subtropical or...

Montane wood mouse

Montane wood mouse

Hylomyscus denniae

The montane hylomyscus or montane wood mouse (*Hylomyscus denniae*) is a species of rodent in the family Muridae. A long-coated species with brownish-grey upper parts and whitish-grey underparts, it occurs in the uplands of tropical Central Africa...

Monte gerbil mouse

Monte gerbil mouse

Eligmodontia moreni

The Monte gerbil mouse or Monte laucha (*Eligmodontia moreni*) is a species of rodent in the family Cricetidae. It is found only in Argentina.

Moojen's Atlantic spiny rat

Moojen's Atlantic spiny rat

Trinomys moojeni

Moojen's Atlantic spiny-rat, *Trinomys moojeni*, is a spiny rat species from South America. It is found in Brazil. It is named after twentieth-century Brazilian zoologist João Moojen.

Moon forest shrew

Moon forest shrew

Sylvisorex lunaris

The moon forest shrew (*Sylvisorex lunaris*) is a species of mammal in the family Soricidae. It is found in Burundi, Rwanda, and Uganda. Its natural habitat is subtropical or tropical moist montane forests.

Moon striped mouse

Moon striped mouse

Hybomys lunaris

The moon striped mouse or Ruwenzori hybomys (*Hybomys lunaris*) is a species of rodent in the family Muridae. It is found in Democratic Republic of the Congo, Rwanda, and Uganda. Its natural habitat is subtropical or tropical moist montane...

Moon-toothed degu
Moon-toothed degu
Octodon lunatus

The moon-toothed degu (*Octodon lunatus*) is a species of rodent in the family Octodontidae. It is endemic to Chile, occurring in mountainous areas along the Pacific coast in the central part of the country.

Moonrat
Moonrat
Echinosorex gymnura

The moonrat (*Echinosorex gymnura*) is a southeast Asian species of mammal in the family Erinaceidae (hedgehogs and gymnures). It is the only species in the genus *Echinosorex*. The moonrat is a fairly small, primarily carnivorous animal which,...

Moonshine shrew
Moonshine shrew
Crocidura luna

The moonshine shrew (*Crocidura luna*) is a species of mammal in the family Soricidae. It is found in Angola, Burundi, Democratic Republic of the Congo, Kenya, Malawi, Mozambique, Rwanda, Tanzania, Uganda, Zambia, and Zimbabwe. Its natural habitat...

Moor macaque
Moor macaque
Macaca maura

The Moor macaque (*Macaca maura*) is a macaque monkey with brown/black body fur with a pale rump patch and pink bare skin on the rump. It has ischial callosities, which are oval-shaped. It is about 50-58.5 cm long, and eats figs, bamboo seeds,...

Moore's woolly lemur
Moore's woolly lemur
Avahi mooreorum

Moore's woolly lemur (*Avahi mooreorum*), or the Masoala woolly lemur, is a woolly lemur endemic to Madagascar. It has brown-grey fur and a reddish tail. It is named in honor of the Gordon and Betty Moore Foundation of San Francisco.

Moose
Moose
Alces alces

The moose (in North America) or elk (in Eurasia) (*Alces alces*) is a member of the New World deer subfamily. It is the largest and heaviest extant species in the deer family. Hunting and other human activities have caused a reduction in the size...

Mops bakarii
Mops bakarii
Mops bakarii

Mops bakarii is a species of mastiff bat, described in 2008. It is endemic to Pemba Island, which lies in the Indian Ocean off the coast of Tanzania.

Morgan's gerbil mouse
Morgan's gerbil mouse
Eligmodontia morgani

Morgan's gerbil mouse (*Eligmodontia morgani*), also known as the western Patagonian laucha, is a South American species of rodent in the family Cricetidae. It is named for J. P. Morgan, one of the sponsors of the expedition that first identified...

Mormopterus francoismoutoui
Mormopterus francoismoutoui
Mormopterus francoismoutoui

Mormopterus francoismoutoui is a species of free-tailed bat that is endemic to the island of Réunion, which is east of Madagascar.

Moroccan jird

Moroccan jird

Meriones grandis

The Moroccan jird (Meriones grandis) is a species of rodent from the family Muridae. The species was first discovered by Ángel Cabrera in 1907. It is endemic to Morocco, northern Algeria, and Tunisia. It had been included as part of Shaw's jird...

Morris's bat

Morris's bat

Myotis morrisi

Morris's bat (Myotis morrisi) is a species of vesper bat in the family Vespertilionidae. It is found in Ethiopia and Nigeria. Its natural habitats are dry savanna, moist savanna, caves, and subterranean habitats (other than caves).

Morro Bay kangaroo rat

Morro Bay kangaroo rat

Dipodomys heermanni morroensis

The Morro Bay kangaroo rat, Dipodomys heermanni morroensis, is endemic to San Luis Obispo County, California. This subspecies of "Heermann" kangaroo rat today lives only in a restricted 2 km area south of Morro Bay in San Luis Obispo County,...

Moss-forest blossom bat

Moss-forest blossom bat

Syconycteris hobbit

The moss-forest blossom bat (Syconycteris hobbit) is a species of megabat in the family Pteropodidae found in New Guinea. Its natural habitat is subtropical or tropical dry forests. It is threatened by habitat loss. Originally discovered in New...

Moss-forest rat

Moss-forest rat

Rattus niobe

The moss-forest rat (Rattus niobe) is a species of rodent in the family Muridae. It is found in Indonesia and Papua New Guinea.

Mossy forest shrew

Mossy forest shrew

Crocidura musseri

The mossy forest shrew (Crocidura musseri) is a species of shrew native to Indonesia.

Mottle-faced tamarin

Mottle-faced tamarin

Saguinus inustus

The mottle-faced tamarin (Saguinus inustus) is a species of tamarin from South America. It is found in Brazil and Colombia.

Mottled tuco-tuco

Mottled tuco-tuco

Ctenomys latro

The mottled tuco-tuco (Ctenomys latro) is a species of rodent in the family Ctenomyidae. It is endemic to Argentina. Its natural habitat is subtropical or tropical dry lowland grassland.

Mottled-tailed shrew mouse

Mottled-tailed shrew mouse
Pseudohydromys fuscus

The mottled-tailed shrew mouse (*Pseudohydromys fuscus*) is a species of rodent in the family Muridae. It is found in West Papua, Indonesia and Papua New Guinea.

Mouflon

Mouflon

Ovis orientalis

The mouflon (*Ovis gmelini*) is a wild sheep native to the Caspian region. It is thought to be the ancestor of all modern domestic sheep breeds.

Mount Apo forest mouse

Mount Apo forest mouse

Apomys hylocetes

The Mount Apo forest mouse (*Apomys hylocetes*) is a species of rodent in the family Muridae. It is found only in the Philippines.

Mount Banahaw forest mouse

Mount Banahaw forest mouse

Apomys banahao

The Mount Banahaw forest mouse (*Apomys banahao*) is a forest mouse endemic to the Mount Banahaw area in the Philippines.

Mount Cameroon brush-furred rat

Mount Cameroon brush-furred rat

Lophuromys roseveari

The Mount Cameroon brush-furred rat or Rosevear's brush-furred mouse (*Lophuromys roseveari*) is a species of rodent in the family Muridae. It is found only in Cameroon. Its natural habitats are subtropical or tropical moist montane forests,...

Mount Cameroon forest shrew

Mount Cameroon forest shrew

Sylvisorex morio

The Mount Cameroon forest shrew (*Sylvisorex morio*) is a species of mammal in the family Soricidae endemic to Cameroon. Its natural habitat is subtropical or tropical moist montane forests.

Mount Carmel blind mole-rat

Mount Carmel blind mole-rat

Spalax carmeli

The Middle East blind mole-rat or Palestine mole-rat (*Nannospalax ehrenbergi*) is a species of rodent in the family Spalacidae.

Mount Claro rock-wallaby

Mount Claro rock-wallaby

Petrogale sharmani

The Mount Claro rock-wallaby (*Petrogale sharmani*), also known as Sharman's rock-wallaby, is a species of rock-wallaby found in northeastern Queensland, Australia. It is a member of a group of seven very closely related species also including...

Mount Data shrew-rat

Mount Data shrew-rat

Rhynchomys soricoides

The Mount Data shrew-rat (*Rhynchomys soricoides*) is a species of rodent in the family Muridae. It is found only in the Philippines.

Mount Elgon vlei rat

Mount Elgon vlei rat

Otomys jacksoni

The Mount Elgon vlei rat (*Otomys jacksoni*) is a species of rodent in the family Muridae. It is found in Kenya and Uganda. Its natural habitat is subtropical or tropical high-altitude grassland. It is threatened by habitat loss.

Mount Gaoligong flying squirrel
Mount Gaoligong flying squirrel
Biswamoyopterus gaoligongensis

The Mount Gaoligong flying squirrel (*Biswamoyopterus gaoligongensis*) is a species of large, arboreal flying squirrel endemic to highland forests in southwestern China. It is the third known member of the highly elusive genus *Biswamoyopterus*. It...

Mount Gargues pipistrelle
Mount Gargues pipistrelle
Pipistrellus aero

The Mount Gargues pipistrelle (*Pipistrellus aero*) is a species of vesper bat found in Kenya and believed to be widely distributed across highlands in Ethiopia. It typically lives in subtropical or tropical forests.

Mount Graham red squirrel
Mount Graham red squirrel
Tamiasciurus hudsonicus grahamensis

The Mount Graham red squirrel (*Tamiasciurus fremonti grahamensis*) is an endangered subspecies of the southwestern red squirrel (*Tamiasciurus fremonti*) native to the Pinaleno Mountains of Arizona. It is smaller than most other subspecies of red...

Mount Irid forest mouse
Mount Irid forest mouse
Apomys iridensis

The Mount Irid forest mouse (*Apomys iridensis*) is a forest mouse endemic to Mount Irid in Luzon, Philippines.

Mount Isarog shrew-mouse
Mount Isarog shrew-mouse
Archboldomys luzonensis

The Mount Isarog shrew-mouse (*Archboldomys luzonensis*) is a species of rodent in the family Muridae found only in the Philippines. Its natural habitat is subtropical or tropical dry forest. It is threatened by habitat loss due to logging...

Mount Kahuzi climbing mouse
Mount Kahuzi climbing mouse
Dendromus kahuziensis

The Mount Kahuzi climbing mouse (*Dendromus kahuziensis*) is a rodent found only in the Democratic Republic of the Congo. It is listed as a critically endangered species due to illegal logging; it is also threatened by fire. Only two specimens...

Mount Kenya mole shrew
Mount Kenya mole shrew
Surdisorex polulus

The Mount Kenya mole shrew (*Surdisorex polulus*) is a species of mammal in the family Soricidae endemic to Mount Kenya in Kenya. Its natural habitat is tropical high-altitude bamboo and grassland, particularly the bamboo-belt of Mount Kenya and...

Mount Lyell shrew
Mount Lyell shrew
Sorex lyelli

The Mount Lyell shrew (*Sorex lyelli*) is a species of mammal in the family Soricidae. It is named for Mount Lyell in Yosemite National Park, the area where the shrew has been most commonly found.

Mount Mabu horseshoe bat
Mount Mabu horseshoe bat
Rhinolophus mabuensis

The Mount Mabu horseshoe bat (*Rhinolophus mabuensis*) is a species of horseshoe bat that is endemic to the Southeast African nation of Mozambique.

Mount Mangan forest mouse
Mount Mangan forest mouse
Apomys manganensis

The Mount Mangan forest mouse (*Apomys manganensis*) is a forest mouse endemic to Mount Mangan in Luzon, Philippines.

Mount Missim long-eared bat
Mount Missim long-eared bat
Nyctophilus shirleyae

The Mount Missim long-eared bat (*Nyctophilus shirleyae*) is a species of slit-faced bat found in Papua New Guinea.

Mount Oku hylomyscus
Mount Oku hylomyscus
Hylomyscus grandis

The Mount Oku hylomyscus (*Hylomyscus grandis*) is a species of rodent in the family Muridae. Found only on Mount Oku, Cameroon, in tropical Central Africa, its natural habitat is tropical moist montane forests. It has a very small range and is...

Mount Oku rat
Mount Oku rat
Lamottemys okuensis

The Mount Oku rat (*Lamottemys okuensis*) is a species of rodent in the family Muridae. The genus *Lamottemys* is monotypic, and this is the only species. It is found only in Cameroon where its natural habitat is subtropical or tropical moist...

Mount Pirri isthmus rat
Mount Pirri isthmus rat
Isthmomys pirrensis

The Mount Pirri isthmus rat (*Isthmomys pirrensis*) is a species of rodent in the family Cricetidae. Its natural habitat is subtropical or tropical moist lowland forests. It is found only in Panama. At the time of its discovery, it was considered...

Mount Popa pipistrelle
Mount Popa pipistrelle
Pipistrellus paterculus

The Mount Popa pipistrelle (*Pipistrellus paterculus*) is a species of vesper bat. It is found in China, India, Myanmar, Thailand, and Vietnam.

Mount Tapulao forest mouse
Mount Tapulao forest mouse
Apomys brownorum

The Mount Tapulao forest mouse (*Apomys brownorum*) is a forest mouse endemic to the Mount Tapulao area in the Philippines. It is named after the American zoologist Barbara Elaine Russell Brown.
Mountain Anoa

Mountain Anoa
Bubalus quarlesi

Anoa, also known as dwarf buffalo and sapiutan, is a subgenus of *Bubalus* comprising two species endemic to the island of Sulawesi in Indonesia: the mountain anoa (*Bubalus quarlesi*) and the lowland anoa (*Bubalus depressicornis*). Both live in...

Mountain Beaver
Mountain Beaver
Aplodontia rufa

The Mountain beaver (*Aplodontia rufa*) is a rodent from North America. It should not be confused with true North American and Eurasian beavers, to which it is not closely related; instead its closest relatives are squirrels.

Mountain brushtail possum
Mountain brushtail possum
Trichosurus cunninghami

The mountain brushtail possum, or southern bobuck (*Trichosurus cunninghami*), is a nocturnal, semi-arboreal marsupial of the family *Phalangeridae* native to southeastern Australia. It was not described as a separate species until 2002.

Mountain Cottontail
Mountain Cottontail
Sylvilagus nuttallii

The mountain cottontail or Nuttall's cottontail (*Sylvilagus nuttallii*) is a species of mammal in the family *Leporidae*. It is found in Canada and the United States.

Mountain cuscus
Mountain cuscus
Phalanger carmelitae

The mountain cuscus (*Phalanger carmelitae*) is a species of marsupial in the family *Phalangeridae* found in West Papua, Indonesia and Papua New Guinea.

Mountain degu
Mountain degu
Octodontomys gliroides

The mountain degu (*Octodontomys gliroides*) is a species of rodent in the family *Octodontidae*. It is the only species in the genus *Octodontomys*. It is found in the foothills of the Andes in Argentina, Bolivia and Chile.

Mountain Gazelle
Mountain Gazelle
Gazella gazella

The mountain gazelle or the Palestine mountain gazelle (*Gazella gazella*) is a species of gazelle widely but unevenly distributed.

Mountain giant Sunda rat
Mountain giant Sunda rat
Sundamys infraluteus

The mountain giant Sunda rat, *Sundamys infraluteus* (also known as the giant mountain rat or giant rat of Sumatra) is a large rat, around 480 to 640 millimeters (19 to 25 inches) in total length (230 to 290 millimeters excluding the tail).

It...
Mountain Goat
Mountain Goat
Oreamnos americanus

The Mountain goat (*Oreamnos americanus*) is a hoofed mammal endemic to mountainous

areas of western North America. A subalpine to alpine species, it is a sure-footed climber commonly seen on cliffs and ice. Despite its vernacular name the Mountain...

Mountain gorilla

Mountain gorilla

Gorilla beringei beringei

The mountain gorilla (*Gorilla beringei beringei*) is one of the two subspecies of the eastern gorilla. It is listed as endangered by the IUCN as of 2018. There are two populations: One is found in the Virunga volcanic mountains of Central Africa,...

Mountain ground squirrel

Mountain ground squirrel

Xerus princeps

The mountain ground squirrel (*Geosciurus princeps*) is a rodent that is native to southwestern Angola, western Namibia, and western South Africa. It is also known as the Kaoko ground squirrel or the Damara ground squirrel. It is the closest...

Mountain Hare

Mountain Hare

Lepus timidus

Mountain hares (*Lepus timidus*) are largely adapted to polar and mountainous habitats. They are shy but swift animals. They nest in slight depressions called forms, and their young are able to fend for themselves shortly after birth.

Mountain mosaic-tailed rat

Mountain mosaic-tailed rat

Paramelomys rubex

The mountain mosaic-tailed rat (*Paramelomys rubex*) is a species of rodent in the family Muridae. It is found in West Papua, Indonesia and Papua New Guinea.

Mountain noctule

Mountain noctule

Nyctalus montanus

The mountain noctule (*Nyctalus montanus*) is a species of bat found in Afghanistan, India, Pakistan, and Nepal.

Mountain Nyala

Mountain Nyala

Tragelaphus buxtoni

The mountain nyala (*Tragelaphus buxtoni*) or balbok is an antelope found in high altitude woodland in a small part of central Ethiopia. It is a monotypic species (without any identified subspecies) first described by English naturalist

Richard...

Mountain paca

Mountain paca

Cuniculus taczanowskii

The mountain paca (*Cuniculus taczanowskii*) is a small burrow-dwelling rodent whose habitats are high altitude South American forests. Pacas are nocturnal, sedentary, and solitary animals with territorial tendencies. It eats mostly fruits and...

Mountain pademelon

Mountain pademelon

Thylogale lanatus

The mountain pademelon (*Thylogale lanatus*) is one of seven species of the genus *Thylogale*. It is found only in Papua New Guinea.

Mountain pocket gopher

Mountain pocket gopher
Thomomys monticola

The mountain pocket gopher (*Thomomys monticola*) is a species of rodent in the family Geomyidae. It is found in California and Nevada, and is endemic there, in the Western United States. The Sierra Nevada are part of its range.

Mountain Pygmy Possum
Mountain Pygmy Possum
Burramys parvus

The mountain pygmy possum (*Burramys parvus*); also simply known as the Burramys, is a small, mouse-sized (weighs 45 grams (1.6 oz)) nocturnal marsupial of Australia found in dense alpine rock screes and boulder fields, mainly southern Victoria...

Mountain Reedbuck
Mountain Reedbuck
Redunca fulvorufula

The mountain reedbuck (*Redunca fulvorufula*) is an antelope found in mountainous areas of much of sub-Saharan Africa.

Mountain reindeer
Mountain reindeer
Rangifer tarandus tarandus

The mountain reindeer (*Rangifer tarandus tarandus*), also called the Norwegian reindeer, northern reindeer, Common reindeer or mountain caribou, is a mid-sized to large subspecies of the reindeer that is native to the western Scandinavian...

Mountain spiny pocket mouse
Mountain spiny pocket mouse
Heteromys oresterus

The mountain spiny pocket mouse (*Heteromys oresterus*) is a species of rodent in the family Heteromyidae. It is endemic to Costa Rica. Its natural habitat is subtropical or tropical moist lowland forests. It is threatened by habitat loss.

Mountain spiny rat
Mountain spiny rat
Maxomys alticola

The mountain spiny rat (*Maxomys alticola*) is a species of rodent in the family Muridae. It is found only in Malaysia.

Mountain Tapir
Mountain Tapir
Tapirus pinchaque

The mountain tapir, also known as the Andean tapir or woolly tapir (*Tapirus pinchaque*) is the smallest of the four widely recognized species of tapir. It is the only one to live outside of tropical rainforests in the wild. It is most easily...

Mountain treeshrew
Mountain treeshrew
Tupaia montana

The mountain treeshrew (*Tupaia montana*) is a treeshrew species within the Tupaiidae. It is endemic to Borneo and inhabits montane forests in Sarawak and Sabah. The first specimen was described by Oldfield Thomas and was part of a zoological...

Mountain tube-nosed fruit bat
Mountain tube-nosed fruit bat
Nyctimene ceratans

The mountain tube-nosed fruit bat (*Nyctimene ceratonyx*) is a species of megabat in the family Pteropodidae. It is endemic to New Guinea island within Papua New Guinea (nation) and Western New Guinea of Indonesia; and to New Britain Island of Papua...
Mountain viscacha rat
Mountain viscacha rat
Octomys mimax

The mountain viscacha rat or mountain vizcacha rat (*Octomys mimax*), historically viscacha rat or vizcacha rat, is a species of rodent in the family Octodontidae. It is endemic to Argentina. It is the only living species within the genus *Octomys*....
Mountain water rat
Mountain water rat
Baiyankamys habbema

The mountain water rat (*Baiyankamys habbema*) is a semiaquatic species of rodent in the family Muridae. It is found in West Papua, Indonesia and Papua New Guinea. Its natural habitat is rivers.
Mountain Weasel
Mountain Weasel
Mustela altaica

The mountain weasel (*Mustela altaica*), also known as the pale weasel, Altai weasel or solongoi, primarily lives in high-altitude environments, as well as rocky tundra and grassy woodlands. This weasel rests in rock crevices, tree trunks, and...
Mountain Zebra
Mountain Zebra
Equus zebra

Mountain zebras (*Equus zebra*) are African mammals with distinctive black-and-white striped coats. Their dazzling stripes make them among the most recognizable mammals. Mountain zebras are so-called due to their amazing climbing ability, allowing...
Moupin pika
Moupin pika
Ochotona thibetana

The Moupin pika (*Ochotona thibetana*), also known as Ribetischer Pika, Moupin-Pika, Pika del Tibet, and Manipuri pika, is a species of mammal in the pika family, Ochotonidae. It has many subspecies, some of which may be distinct species. Its...
Mouse bandicoot
Mouse bandicoot
Microperoryctes murina

The mouse bandicoot (*Microperoryctes murina*) is a species of marsupial in the family Peramelidae. It is endemic to West Papua, Indonesia. Its natural habitat is subtropical or tropical dry forests. Type locality: Indonesia, Prov. of Papua (=...
Mouse-tailed Atlantic spiny rat
Mouse-tailed Atlantic spiny rat
Trinomys myosuros

The mouse-tailed Atlantic spiny-rat, *Trinomys myosuros*, is a spiny rat species from South America. It is found in Brazil.
Mouselike pipistrelle
Mouselike pipistrelle
Hypsugo musciculus

The mouselike pipistrelle (*Hypsugo musciculus*) is a species of vesper bat in the family Vespertilionidae. It is found in Cameroon, Republic of the Congo, Democratic Republic of the Congo, and Gabon. Its natural habitats are subtropical or

tropical...
Moustached guenon
Moustached guenon
Cercopithecus cephus

The moustached guenon or moustached monkey (*Cercopithecus cephus*) is a species of primate in the family Cercopithecidae. It is found in Angola, Cameroon, Central African Republic, Republic of the Congo, Democratic Republic of the Congo,...

Moustached Tamarin
Moustached Tamarin
Saguinus mystax

The moustached tamarin (*Saguinus mystax*) is a New World monkey and a species of tamarin. The moustached tamarin is named for the lack of coloring in the facial hair surrounding their mouth, appearing similar to a moustache. As with all New World...

Mozambican horseshoe bat
Mozambican horseshoe bat
Rhinolophus mossambicus

The Mozambican horseshoe bat (*Rhinolophus mossambicus*) is a species of horseshoe bat found in Africa.

Mozambique thicket rat
Mozambique thicket rat
Grammomys cometes

The Mozambique thicket rat (*Grammomys cometes*) is an African species of rodent in the mouse family, Muridae. It is found in Mozambique, South Africa, and Zimbabwe. Its natural habitat is subtropical or tropical dry forests.

Muennink's spiny rat
Muennink's spiny rat
Tokudaia muenninki

Muennink's spiny rat (*Tokudaia muenninki*) or Okinawa spiny rat (Japanese: オキナワトグネズミ, romanized: Okinawa togenezumi or 沖縄棘鼠) is a species of rodent in the family Muridae. Endemic to Okinawa Island, Japan, its natural habitat is subtropical moist...

Muenster yellow-toothed cavy
Muenster yellow-toothed cavy
Galea monasteriensis

The Muenster yellow-toothed cavy (*Galea monasteriensis*) is a species of rodent in the family Caviidae. It is known only from one location in Valle Hermoso in the Bolivian Andes, at an elevation of 2557 m. Specimens from this location were shipped...

Muisk vole
Muisk vole
Microtus mujanensis

The Muisk vole (*Microtus mujanensis*) is a species of rodent in the family Cricetidae. It is found only in Russia.

Mule Deer
Mule Deer
Odocoileus hemionus

The Mule deer (*Odocoileus hemionus*) is one of the most iconic and beloved wildlife in the American West. It is so-named because of its large ears that look like a mule's ears.

Muli pika

Muli pika
Ochotona muliensis

The Muli pika (*Ochotona muliensis*) is a species of mammal in the family Ochotonidae. It is endemic to China. Its natural habitat is temperate grassland. It is threatened by habitat loss. It is a rarely found, one of the six pika species endemic...

Mullah spiny mouse
Mullah spiny mouse
Acomys mullah

The Mullah spiny mouse (*Acomys mullah*) is a species of rodent in the family Muridae found in Djibouti, Eritrea, Ethiopia, and Somalia. Its natural habitats are subtropical or tropical dry shrubland and rocky areas.

Müller's giant Sunda rat
Müller's giant Sunda rat
Sundamys muelleri

Müller's giant Sunda rat (*Sundamys muelleri*) is a species of rodent in the family Muridae. It is found in Indonesia, Malaysia, Myanmar, the Philippines, and Thailand.

Müller's gibbon
Müller's gibbon
Hylobates muelleri

Müller's gibbon (*Hylobates muelleri*), also known as the southern grey gibbon, is a primate in the gibbon family, Hylobatidae.

Muna-Buton macaque
Muna-Buton macaque
Macaca ochreata brunnescens

The Muna-Buton macaque (*Macaca ochreata brunnescens*) is one of seven Sulawesi macaques in Indonesia. It is only found on two small islands off the south-east coast: the islands of Buton and Muna. This Old World monkey is diurnal and arboreal,...

Munduruku marmoset
Munduruku marmoset
Mico munduruku

The Munduruku marmoset (*Mico munduruku*) is a marmoset endemic to Brazil. It is found only in the southern Amazon, in an area of approximately 120,000 km², from the right bank of the Jamanxim River, below the mouth of the Rio Novo, to the mouth...

Mura's saddleback tamarin
Mura's saddleback tamarin
Leontocebus fuscicollis mura

Mura's saddleback tamarin (*Leontocebus fuscicollis mura*) is a subspecies of monkey that was first seen by scientists in 2007 in the Brazilian state of Amazonas. The monkey, which is mostly gray and brown, weighs 213 grams (7.5 oz), is 240...

Murina recondita
Murina recondita
Murina recondita

Murina recondita is a species of bat found in Taiwan.

Murree vole
Murree vole
Hyperacrius wynnei

The Murree vole (*Hyperacrius wynnei*) is a species of rodent in the family

Cricetidae. It is endemic to the Himalayas in Pakistan and India. It was named by William Thomas Blanford after its collector, geologist Arthur Beavor Wynne.

Mus lepidoides
Mus lepidoides
Mus lepidoides

Mus lepidoides is a species of rodent in the family Muridae.

Mus musculus domesticus
Mus musculus domesticus
Mus musculus domesticus

Mus musculus domesticus, the Western European house mouse, is a subspecies of the house mouse (*Mus musculus*). Some laboratory mouse strains, such as C57BL/6, are domesticated from *M. m. domesticus*.

Mus nitidulus
Mus nitidulus
Mus nitidulus

Mus nitidulus is a medium-sized rodent of the genus *Mus* that appears in central Myanmar.

Muskox
Muskox
Ovibos moschatus

The muskox (*Ovibos moschatus*) is a hoofed mammal native to the Arctic. It is noted for its thick coat and for the strong odor emitted by males during the seasonal rut, from which its name derives. This musky odor has the effect of attracting...

Muskrat
Muskrat
Ondatra zibethicus

The muskrat (*Ondatra zibethicus*) is a medium-sized semiaquatic rodent native to North America. It lives in wetlands over a wide range of climates and habitats. It has important effects on the ecology of wetlands and is a resource of food and fur...

Musky Rat-Kangaroo
Musky Rat-Kangaroo
Hypsiprymnodon moschatus

The musky rat-kangaroo (*Hypsiprymnodon moschatus*) is a small marsupial found only in the rainforests of northeastern Australia. First described in the later 19th century, the only other species are known from fossil specimens. They are similar...

Musschenbroek's spiny rat
Musschenbroek's spiny rat
Maxomys musschenbroekii

Musschenbroek's spiny rat (*Maxomys musschenbroekii*) is a species of rodent in the family Muridae. It is found only in Indonesia.

Musser's shrew mouse
Musser's shrew mouse
Pseudohydromys musseri

Musser's shrew mouse (*Pseudohydromys musseri*) is a species of rodent in the family Muridae. It is found only in Papua New Guinea. Its natural habitat is subtropical or tropical moist montane forests.

Musso's fish-eating rat
Musso's fish-eating rat
Neusticomys mussoi

Musso's fish-eating rat (*Neusticomys mussoi*) is a species of rodent in the family Cricetidae endemic to western Venezuela, where it has been found at altitudes of 1000 to 1200 m. It is semiaquatic and feeds on freshwater invertebrates.

Mustela Mopbie
Mustela Mopbie
Mustela mopbie

Mustela mopbie is a small species of weasel that occurs in southwestern China. It was discovered in 2025. This species has a slender body, which enables it to enter narrow crevices to catch prey. The researchers who discovered *Mustela mopbie*...

Mutable sun squirrel
Mutable sun squirrel
Heliosciurus mutabilis

The mutable sun squirrel (*Heliosciurus mutabilis*) is a species of rodent in the family Sciuridae. It is found in Malawi, Mozambique, Tanzania, Zambia, and Zimbabwe. Its natural habitats are subtropical or tropical moist lowland forest,...

Muton's soft-furred mouse
Muton's soft-furred mouse
Praomys mutoni

Muton's soft-furred mouse or riverine *praomys* (*Praomys mutoni*) is a species of rodent in the family Muridae. It is found only in Democratic Republic of the Congo. Its natural habitats are subtropical or tropical swampland and rivers.

Myanmar Snub-Nosed Monkey
Myanmar Snub-Nosed Monkey
Rhinopithecus strykeri

The Myanmar snub-nosed monkey (*Rhinopithecus strykeri*) is a critically endangered species of colobine monkey discovered in 2010 in northern Burma (Myanmar). It was formally described as a novel species of primate in 2011 based on its fur, beard,...

Myosorex varius
Myosorex varius
Myosorex varius

The forest shrew (*Myosorex varius*) is a species of shrew in the mouse shrew family, Soricidae. It is found in Lesotho, South Africa, and Eswatini. Its natural habitats include temperate forests, dry savanna, Mediterranean-type shrubby...

Myotis diminutus
Myotis diminutus
Myotis diminutus

Myotis diminutus is a species of mouse-eared bat found in Ecuador and Colombia. It was recently described as a new species in 2011.

Myotis fimbriatus
Myotis fimbriatus
Myotis fimbriatus

The fringed long-footed myotis (*Myotis fimbriatus*) is a species of vesper bat in the family Vespertilionidae. It is found in China, Taiwan and Hong Kong. The fringed long-footed myotis is described as having short, thick, brown fur with pale...

Myotis flavus
Myotis flavus
Myotis flavus

Hodgson's bat (*Myotis formosus*), also called the copper-winged bat or black-and-orange myotis, is a species of vesper bat in the genus *Myotis*, the mouse-eared

bats. Favouring mountain forests, it is found throughout Central, Southeast, and East...

Myotis izecksohni
Myotis izecksohni
Myotis izecksohni

Myotis izecksohni is a species of mouse-eared bat that is endemic to Brazil and Argentina.

Myotis laval
Myotis laval
Myotis laval

LaVal's myotis (Myotis laval) is a species of bat found in Brazil and Paraguay.

Myotis macropus
Myotis macropus
Myotis macropus

The southern myotis (Myotis macropus), also known as large-footed myotis, is a species of vesper bat (Vespertilionidae) in genus Myotis. The southern myotis is one of only two Australian "fishing" bats and feeds by trawling its specially adapted...

Myotis midastactus
Myotis midastactus
Myotis midastactus

Myotis midastactus is a golden-coloured species of vesper bats from South America, where it occurs in Bolivia and Paraguay.

Myotis nyctor
Myotis nyctor
Myotis nyctor

The Barbados myotis (Myotis nyctor) is a species of bat found in the Lesser Antilles. It was previously considered a subspecies of Schwartz's myotis, Myotis martiniquensis, but was elevated to species rank in 2012.

Myotis phanluongi
Myotis phanluongi
Myotis phanluongi

Myotis phanluongi is a species of mouse-eared bat. It was described by Borisenko et al. in 2008. It is known from Vietnam. Very small Myotis species with forearm length about 34-36 mm, closely related to Myotis siligorensis. Described from the Hon...

Myotis taiwanensis
Myotis taiwanensis
Myotis taiwanensis

The fringed long-footed myotis (Myotis fimbriatus) is a species of vesper bat in the family Vespertilionidae. It is found in China, Taiwan and Hong Kong. The fringed long-footed myotis is described as having short, thick, brown fur with pale...

Myotis vivesi
Myotis vivesi
Myotis vivesi

Myotis vivesi, the fish-eating bat or fish-eating myotis, is a species of bat that lives around the Gulf of California, and feeds on fish and crustaceans. It is the largest species of the genus Myotis in the Americas, and has exceptionally large...

Mzab gundi
Mzab gundi
Massoutiera mzabi

The Mzab gundi (*Massoutiera mzabi*) is a species of rodent in the family Ctenodactylidae. It is monotypic within the genus *Massoutiera*. It is found in Algeria, Chad, Mali, Niger, and possibly Libya. The Mzab gundi can live in a variety of...

Nabarlek

Nabarlek

Petrogale concinna

Nabarleks (*Petrogale concinna*), are a tiny species of macropod found in northern Australia. They are a shy and nocturnal animal that resides in rocky hollows and forages in the surrounding area. Their diet is grasses, sedges, and ferns found in...

Nagtglas's African dormouse

Nagtglas's African dormouse

Graphiurus nagtglasii

The Nagtglas's African dormouse (*Graphiurus nagtglasii*) is a species of rodent in the family Gliridae. It is found in Cameroon, Central African Republic, Ghana, Liberia, Nigeria, Sierra Leone, and possibly Gabon. Its natural habitat is...

Nairobi grass rat

Nairobi grass rat

Arvicanthis nairobae

The Nairobi grass rat (*Arvicanthis nairobae*) is a species of rodent in the family Muridae. It is found in Kenya, Tanzania, and possibly Ethiopia. Its natural habitat is dry savanna.

Naked Mole-Rat

Naked Mole-Rat

Heterocephalus glaber

The Naked mole-rat (*Heterocephalus glaber*) is a burrowing rodent native to parts of East Africa. It is closely related to the blind mole-rats. The Naked mole-rat exhibits a highly unusual set of physical and behavioral traits that allow it to thrive in...

Naked-eared deer mouse

Naked-eared deer mouse

Peromyscus gymnotis

The naked-eared deer mouse (*Peromyscus gymnotis*) is a species of rodent in the family Cricetidae. It is found in El Salvador, Guatemala, Honduras, Mexico, and Nicaragua.

Naked-nosed shrew tenrec

Naked-nosed shrew tenrec

Microgale gymnorhyncha

The naked-nosed shrew tenrec (*Microgale gymnorhyncha*) is a species of mammal in the family Tenrecidae. It is endemic to Madagascar. Its natural habitats are subtropical or tropical moist lowland forest and subtropical or tropical moist montane...

Naked-rumped pouched bat

Naked-rumped pouched bat

Saccolaimus saccolaimus

The naked-rumped pouched bat (*Saccolaimus saccolaimus*), also known as pouched tomb bat, is a species of sac-winged bat in the family Emballonuridae. It is found in Australia, Bangladesh, India, Indonesia, Malaysia, Papua New Guinea, the...

Naked-Rumped Tomb Bat

Naked-Rumped Tomb Bat

Taphozous nudiventris

The Naked-rumped tomb bat (*Taphozous nudiventris*) is a species of sac-winged bat found in northern Africa, the Middle East, and southeastern Asia. It is often associated with human settlements, but is not tolerant of much disturbance at its...

Naked-soled conyrat

Naked-soled conyrat

Reithrodon typicus

The naked-soled conyrat (*Reithrodon typicus*) is a species of rodent in the family Cricetidae. It is an herbivore of grasslands in northern Argentina, southern Brazil, and Uruguay. Its karyotype has $2n = 28$.

Namaqua dune mole-rat

Namaqua dune mole-rat

Bathyergus janetta

The Namaqua dune mole-rat (*Bathyergus janetta*) is a species of rodent in the family Bathyergidae. It is found in Namibia and South Africa. Its natural habitats are subtropical or tropical dry shrubland, caves, and sandy shores. The IUCN assessment...

Namaqua rock rat

Namaqua rock rat

Aethomys namaquensis

The Namaqua rock rat (*Aethomys namaquensis*) is a species of rodent in the family Muridae. It is sometimes included in the genus *Micaelamys*. It is found in Angola, Botswana, Lesotho, Malawi, Mozambique, Namibia, South Africa, Eswatini, Zambia, and...

Namaqua slender mongoose

Namaqua slender mongoose

Galerella sanguinea swalius

The Namaqua slender mongoose (*Herpestes sanguineus swalius*), also known as the Namibian slender mongoose, is a subspecies of the common slender mongoose. It is endemic to Namibia.

Namdapha Flying Squirrel

Namdapha Flying Squirrel

Biswamoyopterus biswasi

The Namdapha flying squirrel (*Biswamoyopterus biswasi*) is a rare species of flying squirrel native to India. It was the sole member in the genus *Biswamoyopterus* until the description of the Laotian giant flying squirrel (*Biswamoyopterus laoensis*)...

Namib brush-tailed gerbil

Namib brush-tailed gerbil

Gerbillurus setzeri

The Namib brush-tailed gerbil or Setzer's hairy-footed gerbil (*Gerbillurus setzeri*) is a species of rodent endemic to Angola and Namibia. Its natural habitats are sandy and gravelly plains. It stays in its burrow by day, emerging at night to...

Namib long-eared bat

Namib long-eared bat

Laephotis namibensis

The Namib long-eared bat (*Laephotis namibensis*) is a species of vesper bat in the family Vespertilionidae found in Namibia. It is found in these habitats: dry savanna and temperate desert.

Nancy Ma's night monkey

Nancy Ma's night monkey

Aotus nancymae

Nancy Ma's night monkey (*Aotus nancymae*) is a night monkey species from South America. It is found in Brazil and Peru. It is known in medical research as a model organism for studying the Duffy antigen. Nancy Ma's night monkeys have also been...
Napo saki
Napo saki
Pithecia napensis

The Napo saki (*Pithecia napensis*), also known as the Napo monk saki, is a species of saki monkey, a type of New World monkey. Its range includes parts of eastern Ecuador and northern Peru. The name is derived from the Napo River in its locality....
Napo spiny rat
Napo spiny rat
Proechimys quadruplicatus

The Napo spiny rat, *Proechimys quadruplicatus*, is a spiny rat species from South America. It is found in Brazil, Colombia, Ecuador, Peru and Venezuela.
Narrow-faced kangaroo rat
Narrow-faced kangaroo rat
Dipodomys venustus

The narrow-faced kangaroo rat (*Dipodomys venustus*) is a species of rodent in the family Heteromyidae. It is endemic to California in the United States. The dental formula of *Dipodomys venustus* is $1.0.1.31.0.1.3 \times 2 = 20$, which is the dental...
Narrow-headed slender opossum
Narrow-headed slender opossum
Marmosops cracens

The narrow-headed slender opossum (*Marmosops cracens*) is a species of opossum in the family Didelphidae. It is endemic to Venezuela.
Narrow-headed vole
Narrow-headed vole
Microtus gregalis

The narrow-headed vole (*Microtus gregalis*) is a species of rodent in the family Cricetidae. Ranging over northern and central Asia and also into Alaska, it is the only species in the subgenus *Stenocranius*.
Narrow-nosed harvest mouse
Narrow-nosed harvest mouse
Reithrodontomys tenuirostris

The narrow-nosed harvest mouse (*Reithrodontomys tenuirostris*) is a species of rodent in the family Cricetidae. It is found in Guatemala and Mexico.
Narrow-nosed planigale
Narrow-nosed planigale
Planigale tenuirostris

The narrow-nosed planigale (*Planigale tenuirostris*) is a species of very small marsupial carnivore of the family Dasyuridae.
Narrow-skulled pocket mouse
Narrow-skulled pocket mouse
Chaetodipus artus

The narrow-skulled pocket mouse (*Chaetodipus artus*) is a species of rodent in the family Heteromyidae. It is endemic to western Mexico, living west of the Sierra Madre Occidental crest.
Narrow-striped marsupial shrew
Narrow-striped marsupial shrew

Phascolosorex dorsalis

The narrow-striped dasyure or narrow-striped marsupial shrew (*Phascolosorex dorsalis*) is a species of marsupial in the family Dasyuridae found in West Papua, Indonesia and Papua New Guinea. Its natural habitat is subtropical or tropical dry...

Narrow-striped mongoose

Narrow-striped mongoose

Mungotictis decemlineata

The narrow-striped mongoose or narrow-striped vongsira (*Mungotictis decemlineata*) is a member of the family Eupleridae endemic to Madagascar. It inhabits the Madagascar dry deciduous forests in western and southwestern Madagascar, where it lives...

Narrow-tailed white-bellied rat

Narrow-tailed white-bellied rat

Niviventer lepturus

The narrow-tailed white-bellied rat (*Niviventer lepturus*) is a species of rodent in the family Muridae. It is found only in Indonesia.

Narrow-winged pipistrelle

Narrow-winged pipistrelle

Pipistrellus stenopterus

The narrow-winged pipistrelle (*Pipistrellus stenopterus*) is a species of vesper bat. It can be found in Brunei Darussalam, Indonesia, and Malaysia.

Narwhal

Narwhal

Monodon monoceros

The narwhal, also known as a narwhale (*Monodon monoceros*), is a medium-sized toothed whale with a large "tusk" from a protruding canine tooth. It is one of two living species of whale in the family Monodontidae, along with the Beluga whale....

Nasolo's shrew tenrec

Nasolo's shrew tenrec

Microgale nasoloi

Nasolo's shrew tenrec (*Microgale nasoloi*) is a species of mammal in the family Tenrecidae. It is endemic to Madagascar. Its natural habitats are subtropical or tropical dry forests and subtropical or tropical moist montane forests. It is...

Nasuetia meridensis

Nasuetia meridensis

Nasuetia meridensis

The eastern mountain coati or eastern dwarf coati (*Nasuetia meridensis*) is a small procyonid found in cloud forest and páramo at elevations of 2,000–4,000 m (6,600–13,100 ft) in the Andes of western Venezuela. Until 2009, it was included as a...

Nasuetia olivacea

Nasuetia olivacea

Nasuetia olivacea

The western mountain coati or western dwarf coati (*Nasuetia olivacea*) is a small procyonid, found in cloud forest and páramo at altitudes of 1,300–4,250 metres (4,270–13,940 ft) in the Andes of Colombia and Ecuador. A population discovered in...

Natal free-tailed bat

Natal free-tailed bat

Mormopterus acetabulosus

The Natal free-tailed bat (*Mormopterus acetabulosus*) is a species of bat in the family Molossidae, the free-tailed bats. It is endemic to the island of Mauritius. It is known from fewer than five locations in its range, but it is common at a few...

Natal long-fingered bat

Natal long-fingered bat

Miniopterus natalensis

The Natal long-fingered bat (*Miniopterus natalensis*) is a species of vesper bat in the family Miniopteridae. It can be found in the following countries: Angola, Botswana, Democratic Republic of the Congo, possibly Ethiopia, Kenya, Lesotho,...

Natal Multimammate Mouse

Natal Multimammate Mouse

Mastomys natalensis

The Natal multimammate mouse (*Mastomys natalensis*) is a species of rodent found only in Africa. The Natal multimammate rat has been used as a laboratory animal since 1939 and has great value for researchers.

Natal red rock hare

Natal red rock hare

Pronolagus crassicaudatus

The Natal red rock hare or greater red rock hare (*Pronolagus crassicaudatus*) is a species of mammal in the family Leporidae (rabbits and hares). It has a slightly grizzled, grayish brown head and reddish brown upperparts. The dense fur is thick...

Nathusius's pipistrelle

Nathusius's pipistrelle

Pipistrellus nathusii

Nathusius' pipistrelle (*Pipistrellus nathusii*) is a small bat in the genus *Pipistrellus*. It is very similar to the common pipistrelle and has been overlooked in many areas until recently but it is widely distributed across Europe. It was...

Natterer's bat

Natterer's bat

Myotis nattereri

Natterer's bat (*Myotis nattereri*) is a European vespertilionid bat with pale wings. It has brown fur tending to greyish-white on its underside. It is found across most of the continent of Europe, parts of the Near East and North Africa. It feeds...

Natterer's tuco-tuco

Natterer's tuco-tuco

Ctenomys nattereri

The Bolivian tuco-tuco (*Ctenomys boliviensis*) is a species of rodent in the family Ctenomyidae. It is found in Argentina, Bolivia, and Paraguay.

Natuna Island surili

Natuna Island surili

Presbytis natunae

The Natuna Island surili (*Presbytis natunae*) is a species of primate in the family Cercopithecidae.

Navivasha African mole-rat

Navivasha African mole-rat

Tachyoryctes naivashae

The Naivasha African mole-rat (*Tachyoryctes naivashae*) is a species of rodent in the family Spalacidae. It is endemic to Kenya. Its natural habitats are subtropical

or tropical moist montane forests, moist savanna, arable land, and...

Nayarit mouse

Nayarit mouse

Peromyscus simulus

The Nayarit mouse or Sinaloan deer mouse (*Peromyscus simulus*) is a species of cricetid rodent endemic to Mexico. It was considered a subspecies of brush mouse until 1977.

Neacomys dubosti

Neacomys dubosti

Neacomys dubosti

Neacomys dubosti, also known as Dubost's neacomys or Dubost's bristly mouse, is a species of South American rodent in the genus *Neacomys* of family Cricetidae. It is found in French Guiana, southeastern Suriname and nearby Amapá, Brazil. It was...

Neacomys guianae

Neacomys guianae

Neacomys guianae

Neacomys guianae, also known as the Guianan neacomys or Guiana bristly mouse, is a nocturnal rodent species from South America. It is found in lowland tropical rainforests in northeastern Brazil, French Guiana, Guyana, Suriname and Venezuela. Its...

Neacomys minutus

Neacomys minutus

Neacomys minutus

Neacomys minutus, also known as the minute neacomys, the small bristly mouse, or the minute spiny mouse, is a rodent species from South America in the genus *Neacomys*. It is found in Brazil.

Neacomys musseri

Neacomys musseri

Neacomys musseri

Neacomys musseri, also known as Musser's neacomys or Musser's bristly mouse, is a rodent species from South America. It is found in far western Brazil and southeastern Peru.

Neacomys paracou

Neacomys paracou

Neacomys paracou

Neacomys paracou, also known as the Paracou neacomys or Paracou bristly mouse, is a rodent species from South America in the genus *Neacomys*. It is found in northern Brazil, French Guiana, Guyana, Suriname and southeastern Venezuela.

Neacomys pictus

Neacomys pictus

Neacomys pictus

Neacomys pictus, also known as the painted neacomys or painted bristly mouse, is a species of rodent in the genus *Neacomys* of family Cricetidae. It is found only in Panama.

Neacomys spinosus

Neacomys spinosus

Neacomys spinosus

Neacomys spinosus, also known as the common neacomys, common bristly mouse, or bristly mouse, is a nocturnal rodent species from South America in the genus *Neacomys*. It is found in Bolivia, Brazil, Colombia, Ecuador and Peru, where it often lives...

Neacomys tenuipes
Neacomys tenuipes
Neacomys tenuipes

Neacomys tenuipes, also known as the narrow-footed neacomys or narrow-footed bristly mouse, is found along the northern Andes from northwestern Venezuela through Colombia into Ecuador, in rainforest at elevations from 400 to 1750 m. Populations...

Neave's mouse
Neave's mouse
Mus neavei

Neave's mouse (*Mus neavei*) is a species of rodent in the family Muridae. It is found in Democratic Republic of the Congo, Mozambique, South Africa, Tanzania, Zambia, and Zimbabwe. Its natural habitat is dry savanna.

Neblina slender opossum
Neblina slender opossum
Marmosops neblina

The neblina slender opossum (*Marmosops neblina*) is an opossum species from South America. It is found in Brazil, Ecuador, Venezuela, and perhaps Peru.

Neblina uakari
Neblina uakari
Cacajao hosomi

The *Neblina uakari* (*Cacajao hosomi*) or black-headed uakari, is a newly described species of monkey from the far northwest Brazilian Amazon and adjacent southern Venezuela. It was found by Jean-Phillipe Boubli of the University of Auckland and...

Necklace pipistrelle
Necklace pipistrelle
Arielulus torquatus

The necklace sprite or necklace pipistrelle (*Thainycteris torquatus*) is a species of vesper bat endemic to Taiwan.

Nectomys apicalis
Nectomys apicalis
Nectomys apicalis

Nectomys apicalis, also known as the western Amazonian nectomys, is a semiaquatic species of rodent in the genus *Nectomys* of family Cricetidae. It is found east of the Andes in Ecuador, Peru, and Bolivia, east into western Brazil; further to the...

Nectomys magdalenae
Nectomys magdalenae
Nectomys grandis

Nectomys grandis, also known as the Magdalena-Cauca water rat, is a nocturnal, semiaquatic species of rodent in the genus *Nectomys* of family Cricetidae. It is found in western and northern Colombia at altitudes from sea level to 2000 m, including...

Nectomys palmipes
Nectomys palmipes
Nectomys palmipes

Nectomys palmipes, also known as the Trinidad nectomys or Trinidad water rat, is a species of semiaquatic rodent in genus *Nectomys* of family Cricetidae. It is found on the island of Trinidad and on the nearby mainland of Venezuela.

Nectomys rattus

Nectomys rattus
Nectomys rattus

Nectomys rattus, the small-footed bristly mouse, Amazonian nectomys, Amazonian mouse, or common water rat is a species of rodent in the genus Nectomys of family Cricetidae. It is found in Brazil, Colombia, French Guiana, Guyana, Suriname, and...

Nectomys squamipes
Nectomys squamipes
Nectomys squamipes

Nectomys squamipes, also known as the Atlantic Forest nectomys, South American water rat, or scaly-footed water rat, is a semiaquatic insectivorous rodent species. It is from Argentina, Brazil, and Paraguay – found primarily near forest rivers...

Negev shrew
Negev shrew
Crocidura ramona

The Negev shrew (*Crocidura ramona*), also known as the Ramon's shrew (Hebrew: חֲדָף הַרְמוֹן, ḥadaf haramon) is a species of mammal in the family Soricidae. So far, it is only known from Israel. It is found in three regions: Mizpe Ramon and Sede...

Negros shrew
Negros shrew
Crocidura negrina

The Negros shrew (*Crocidura negrina*) is a white-toothed shrew found only on the island of Negros in the Philippines. It is locally called the katsurí and is listed as a critically endangered species due to habitat loss and a restricted range.

Nehring's blind mole-rat
Nehring's blind mole-rat
Spalax nehringi

The Anatolian blind mole-rat or Nehring's blind mole-rat (*Nannospalax xanthodon*) is a species of rodent in the family Spalacidae. It is found in Armenia, Georgia, and Turkey.

Neill's long-tailed giant rat
Neill's long-tailed giant rat
Leopoldamys neilli

Neill's long-tailed giant rat (*Leopoldamys neilli*) is a species of rodent in the family Muridae. It is found in Thailand, Laos, Vietnam, and possibly Myanmar. Its natural habitats are subtropical or tropical dry shrubland and rocky areas....

Nelson's collared lemming
Nelson's collared lemming
Dicrostonyx nelsoni

Nelson's collared lemming (*Dicrostonyx nelsoni*) is a species of rodent in the family Cricetidae. It is found in western and southwestern Alaska in the United States.

Nelson's giant deer mouse
Nelson's giant deer mouse
Megadontomys nelsoni

Nelson's giant deer mouse (*Megadontomys nelsoni*) is a species of rodent in the family Cricetidae. It is found only in Mexico.

Nelson's kangaroo rat
Nelson's kangaroo rat
Dipodomys nelsoni

Nelson's kangaroo rat (*Dipodomys nelsoni*) is a species of rodent in the family Heteromyidae which is endemic to the central plateau of Mexico.

Nelson's pocket mouse

Nelson's pocket mouse

Chaetodipus nelsoni

Nelson's pocket mouse (*Chaetodipus nelsoni*) is a species of rodent in the family Heteromyidae. It is found in Mexico and in New Mexico and Texas in United States. It is named in honor of the American naturalist Edward William Nelson.

Nelson's small-eared shrew

Nelson's small-eared shrew

Cryptotis nelsoni

Nelson's small-eared shrew (*Cryptotis nelsoni*) is a species of mammal in the family Soricidae. It is endemic to eastern Mexico. The species was discovered by Edward William Nelson and Edward Alphonso Goldman in 1894, who collected a number of...

Nelson's spiny pocket mouse

Nelson's spiny pocket mouse

Heteromys nelsoni

Nelson's spiny pocket mouse (*Heteromys nelsoni*) is a species of rodent in the family Heteromyidae. It is found in Mexico and Guatemala. Its natural habitat is subtropical or tropical moist lowland forests. It is threatened by habitat loss.

Nendo tube-nosed fruit bat

Nendo tube-nosed fruit bat

Nyctimene sanctacrucis

The Nendo or Santa Cruz tube-nosed fruit bat (*Nyctimene sanctacrucis*) is a megabat from the Santa Cruz Group of the Solomon Islands, near the eastern limit of the distribution of tube-nosed fruit bats. It has tube-like nostrils and a wingspan of...

Neoromicia robertsi

Neoromicia robertsi

Neoromicia robertsi

Neoromicia robertsi is a species of vesper bat found in Madagascar. It is a recently described species, as it was first described in 2012.

Neotropical pygmy squirrel

Neotropical pygmy squirrel

Sciurillus pusillus

The Neotropical pygmy squirrel (*Sciurillus pusillus*) is a South American species of tree squirrel, being the only living species in the genus *Sciurillus* and the subfamily *Sciurillinae*. Genetic analysis has shown it to be the sister group to all...

Neotropical River Otter

Neotropical River Otter

Lontra longicaudis

The neotropical otter or neotropical river otter (*Lontra longicaudis*) is an otter species found in Middle America, South America, and the island of Trinidad. It is physically similar to the northern and southern river otter, which occur directly...

Nepal Gray Langur

Nepal Gray Langur

Semnopithecus schistaceus

The Nepal gray langur (*Semnopithecus schistaceus*) is a species of Old World monkey found only in Asia. Langurs mostly walk quadrupedally and spend half of their time on the ground and the other half in trees.

Nepal myotis

Nepal myotis

Myotis nipalensis

Myotis nipalensis commonly known as Nepal myotis is a vesper bat of genus *Myotis*.

Nephelomys albigularis

Nephelomys albigularis

Nephelomys albigularis

Nephelomys albigularis, also known as the white-throated oryzomys or Tomes's rice rat, is a species of rodent in the genus *Nephelomys* of family Cricetidae. Described in 1860, it was the first *Nephelomys* species to be discovered. It was originally...

Nephelomys auriventer

Nephelomys auriventer

Nephelomys auriventer

Nephelomys auriventer, also known as the golden-bellied oryzomys or Ecuadorian rice rat, is a species of rodent in the genus *Nephelomys* of family Cricetidae. Oldfield Thomas originally described it, in 1899, as a species of *Oryzomys*, *Oryzomys*...

Nephelomys caracolus

Nephelomys caracolus

Nephelomys caracolus

Nephelomys caracolus, also known as the Costa Central oryzomys or caracol rice rat, is a species of rodent in the genus *Nephelomys* of family Cricetidae. It is found in cloud forest in the Cordillera de la Costa Central of Aragua, Miranda, and the...

Nephelomys childi

Nephelomys childi

Nephelomys childi

Nephelomys childi is a species of rodent in the genus *Nephelomys* of family Cricetidae. The type locality is at Bogotá, Colombia, and the type locality of its junior synonym, *oconnelli*, is at a place known as Buenavista, about 50 miles (80 km)...

Nephelomys devius

Nephelomys devius

Nephelomys devius

Nephelomys devius, also known as the Talamancan oryzomys, Boquete rice rat, Chiriqui rice rat, or montane rice rat, is a species of rodent in the genus *Nephelomys* of family Cricetidae. It is found in cloud forest in the highlands of Costa Rica...

Nephelomys keaysi

Nephelomys keaysi

Nephelomys keaysi

Nephelomys keaysi, also known as Keays's oryzomys or Keays's rice rat, is a species of rodent in the genus *Nephelomys* of family Cricetidae. It is found from southeastern Peru to northern Bolivia on the eastern slope of the Andes in Yungas humid...

Nephelomys levipes

Nephelomys levipes

Nephelomys levipes

Nephelomys levipes, also known as the nimble-footed oryzomys or light-footed rice rat, is a species of rodent in the genus *Nephelomys* of family Cricetidae. It is

found on the eastern slope of the Andes from southeastern Peru into west-central...
Nephelomys maculiventer
Nephelomys maculiventer
Nephelomys maculiventer

Nephelomys maculiventer is a species of rodent in the genus *Nephelomys* of family Cricetidae. The type locality is in Colombia, at "Sierra El Libano, alt. 6000 ft, Santa Marta District". It was originally described on the basis of 47 specimens,...
Nephelomys meridensis
Nephelomys meridensis
Nephelomys meridensis

Nephelomys meridensis, also known as the Mérida oryzomys, is a species of rodent in the genus *Nephelomys* of family Cricetidae. It is found in cloud forest in the Sierra Nevada de Mérida of western Venezuela at elevations from 1100 to 4000 m. It...
Nephelomys moerex
Nephelomys moerex
Nephelomys moerex

Nephelomys moerex is a species of rodent in the genus *Nephelomys* of family Cricetidae. The type locality is at Mindo in western Ecuador, where it has been recorded together with three other rodents of the oryzomyine group, *Sigmodontomys*...
Nephelomys nimbosus
Nephelomys nimbosus
Nephelomys nimbosus

Nephelomys nimbosus is a species of rodent in the genus *Nephelomys* of family Cricetidae. Its type locality is at San Antonio on the northeastern slope of the Tungurahua in the Andes of Ecuador, at an altitude of about 6,700 feet (2,000 m). The...
Nephelomys pectoralis
Nephelomys pectoralis
Nephelomys pectoralis

Nephelomys pectoralis is a species of rodent in the genus *Nephelomys* of family Cricetidae. Its type locality is 40 miles (64 km) west of the city of Popayán, Cauca Department, Colombia, at an altitude of 10,340 feet (3,150 m). American zoologist...
Nephelomys pirrensis
Nephelomys pirrensis
Nephelomys pirrensis

Nephelomys pirrensis, also known as the Mount Pirre rice rat, is a species of rodent in the genus *Nephelomys* of family Cricetidae. Its type locality is at Mount Pirri or Pirre in eastern Panama, at an altitude of 4,500 feet (1,400 m), and it has...
Neriad horseshoe bat
Neriad horseshoe bat
Rhinolophus nereis

The neriad horseshoe bat (*Rhinolophus nereis*) is a species of bat in the family Rhinolophidae. It is endemic to Siantan Island of the Anambas Islands, and to the North Natuna Islands, west of Borneo in the South China Sea within Indonesia....
Nesomys narindaensis
Nesomys narindaensis
Nesomys narindaensis

Nesomys narindaensis is an extinct rodent that lived in northwestern Madagascar. It is known from subfossil skull bones and isolated molars found in several sites during field work that started in 2001. First described in 2010, it is placed in...
Nesoryzomys fernandinae
Nesoryzomys fernandinae
Nesoryzomys fernandinae

Nesoryzomys fernandinae, also known as the Fernandina nesoryzomys, Fernandina rice rat, or Fernandina Galápagos mouse, is a species of rodent in the genus *Nesoryzomys* of family Cricetidae. It is found only on Fernandina in the Galápagos Islands...
Nesoryzomys indefessus narboroughi
Nesoryzomys indefessus narboroughi
Nesoryzomys indefessus narboroughi

Nesoryzomys indefessus narboroughi is a subspecies of *Nesoryzomys indefessus*, a rodent in the genus *Nesoryzomys*, from Fernandina Island in the Galápagos Islands. The only other subspecies occurred on Indefatigable Island, but is now extinct. Some...
Nesoryzomys swarthi
Nesoryzomys swarthi
Nesoryzomys swarthi

Nesoryzomys swarthi, also known as the Santiago nesoryzomys or Santiago Galápagos mouse, is a species of rodent in the genus *Nesoryzomys* of family Cricetidae. It is found only on Santiago in the Galápagos Islands. Its natural habitat is...
Neumann's grass rat
Neumann's grass rat
Arvicanthis neumanni

Neumann's grass rat (*Arvicanthis neumanni*) is a species of rodent in the family Muridae. It is found in Ethiopia, Somalia, Sudan, Tanzania, and possibly Kenya. Its natural habitats are dry savanna and subtropical or tropical dry shrubland.
Neuquén grass mouse
Neuquén grass mouse
Akodon neocenus

The Neuquén grass mouse (*Akodon neocenus*) is a species of rodent in the family Cricetidae. It is found only in Argentina. As of 2017, the IUCN synonymizes it with *Akodon dolores*.
New Britain naked-backed fruit bat
New Britain naked-backed fruit bat
Dobsonia praedatrix

The New Britain naked-backed fruit bat (*Dobsonia praedatrix*) is a species of megabat in the family Pteropodidae. It is endemic to New Britain island in northern Papua New Guinea.
New Britain water rat
New Britain water rat
Hydromys neobritannicus

The New Britain water rat (*Hydromys neobritannicus*) is a species of semiaquatic rodent in the family Muridae. It is found only on the island of New Britain in Papua New Guinea. It is threatened by habitat loss.
New Caledonia blossom bat
New Caledonia blossom bat
Notopteris neocaledonica

The New Caledonia blossom bat (*Notopteris neocaledonica*) is an uncommon species of megabat in the family Pteropodidae. The species lives in caves in northern New

Caledonia, and forms colonies of up to 300.
New Caledonia flying fox
New Caledonia flying fox
Pteropus vetulus

The New Caledonia flying fox (*Pteropus vetulus*) is a species of flying fox in the family Pteropodidae. It is endemic to New Caledonia. The habitat of the species is highly fragmented, and possibly decreasing due to deforestation. Other threats...
New Caledonia wattled bat
New Caledonia wattled bat
Chalinolobus neocaledonicus

The New Caledonian wattled bat (*Chalinolobus neocaledonicus*) is a species of vesper bat, family Vespertilionidae. It is found only in New Caledonia.
New Caledonian long-eared bat
New Caledonian long-eared bat
Nyctophilus nebulosus

The New Caledonian long-eared bat, species *Nyctophilus nebulosus*, is a vespertilionid bat found in New Caledonia. They are only recorded at Mount Koghis, near Nouméa, and the population is decreasing.
New England Cottontail
New England Cottontail
Sylvilagus transitionalis

The New England cottontail (*Sylvilagus transitionalis*), also called the gray rabbit, brush rabbit, wood hare, wood rabbit, or cooney, is a species of cottontail rabbit represented by fragmented populations in areas of New England, specifically...
New Georgian monkey-faced bat
New Georgian monkey-faced bat
Pteralopex taki

The New Georgian monkey-faced bat or New Georgian flying monkey (*Pteralopex taki*) is a recently described species of megabat endemic to the Solomon Islands, more specifically New Georgia and Vangunu Islands. It is presumably extinct on...
New Guinea big-eared bat
New Guinea big-eared bat
Pharotis imogene

The New Guinea big-eared bat or Papuan big-eared bat, (*Pharotis imogene*), is a vespertilionid bat endemic to Papua New Guinea. It is listed as a critically endangered species due to ongoing habitat loss. It is the only known member of the genus...
New Guinea free-tailed bat
New Guinea free-tailed bat
Austronomus kuboriensis

The New Guinea free-tailed bat (*Austronomus kuboriensis*), sometimes designated the New Guinea mastiff bat, is an extant species of free-tailed bat that inhabits the Chimbu highlands of Papua New Guinea. Although Koopman describes *A. kuboriensis*...
New Guinea long-eared bat
New Guinea long-eared bat
Nyctophilus microtis

The New Guinea long-eared bat, species *Nyctophilus microtis*, is a small flying mammal. It is found only in Papua New Guinea.
New Guinea waterside rat
New Guinea waterside rat

Parahydromys asper

The New Guinea waterside rat (*Parahydromys asper*) is the only member of the genus *Parahydromys*. It is considered part of the New Guinea Old Endemics, meaning its ancestors were part of the first wave of murine rodents to colonize the island. It...

New Guinean jumping mouse

New Guinean jumping mouse

Lorentzimys nouhuysi

The New Guinean jumping mouse (*Lorentzimys nouhuysi*) is a species of rodent in the family Muridae. It is found in West Papua, Indonesia and Papua New Guinea.

New Guinean planigale

New Guinean planigale

Planigale novaeguineae

The New Guinean planigale (*Planigale novaeguineae*), also known as the Papuan planigale, is a species of small marsupial carnivore native to the Trans-Fly savanna and grasslands of New Guinea.

New Guinean quoll

New Guinean quoll

Dasyurus albopunctatus

The New Guinean quoll (*Dasyurus albopunctatus*), also known as the New Guinea quoll or New Guinea native cat, is a carnivorous marsupial mammal native to New Guinea. It is the second-largest surviving marsupial carnivore of New Guinea.

New Guinean rat

New Guinean rat

Rattus novaeguineae

The New Guinean rat (*Rattus novaeguineae*) is a species of rodent in the family Muridae. It is found only in parts of central Papua New Guinea.

New Holland Mouse

New Holland Mouse

Pseudomys novaehollandiae

The New Holland mouse (*Pseudomys novaehollandiae*) is a species of rodent in the family Muridae. It was first described by George Waterhouse in 1843. It vanished from view for over a century before its rediscovery in Ku-ring-gai Chase National...

New Ireland forest rat

New Ireland forest rat

Rattus sanila

The New Ireland forest rat (*Rattus sanila*) is a large rodent in the family Muridae. It is endemic to New Ireland, in the Bismarck Archipelago, Papua New Guinea.

New Mexico shrew

New Mexico shrew

Sorex neomexicanus

The New Mexico shrew (*Sorex neomexicanus*) is a species of mammal in the family Soricidae. It is found only in New Mexico in the Capitan and Sandia-Manzano Mountains. Its total length is 103 to 121 millimeters. Its tail length is 39 to 54...

New Zealand Fur Seal

New Zealand Fur Seal

Arctocephalus forsteri

Arctocephalus forsteri (common names include the Australasian fur seal, South Australian fur seal, New Zealand fur seal, Antipodean fur seal, or long-nosed fur seal) is a species of fur seal found mainly around southern Australia and New Zealand....

New Zealand Greater Short-Tailed Bat

New Zealand Greater Short-Tailed Bat

Mystacina robusta

The New Zealand greater short-tailed bat (*Mystacina robusta*) is one of two species of New Zealand short-tailed bats, unique to New Zealand. Larger than the New Zealand lesser short-tailed bat, there have been no confirmed sightings of this...

New Zealand Lesser Short-Tailed Bat

New Zealand Lesser Short-Tailed Bat

Mystacina tuberculata

The New Zealand lesser short-tailed bat (*Mystacina tuberculata*) is a small omnivorous species of bat. It is one of two extant and three overall mammal species unique to New Zealand. Lesser short-tailed bats have unique adaptations that...

New Zealand long-tailed bat

New Zealand long-tailed bat

Chalinolobus tuberculatus

The New Zealand long-tailed bat (*Chalinolobus tuberculatus*), also known as the long-tailed wattled bat or pekapeka-tou-roa (Māori), is one of 15 species of bats in the genus *Chalinolobus* variously known as "pied bats", "wattled bats" or...

New Zealand sea lion

New Zealand sea lion

Phocarctos hookeri

The New Zealand sea lion (*Phocarctos hookeri*), once known as Hooker's sea lion, and as pakake or whakahao (male) and kaki (female) in Māori, is a species of sea lion that is endemic to New Zealand and primarily breeds on New Zealand's...

Newfoundland Black Bear

Newfoundland Black Bear

Ursus americanus hamiltoni

The Newfoundland black bear (*Ursus americanus hamiltoni*) is a subspecies of the American black bear that is found in Canada. It has one of the longest hibernation periods of any bear in North America.

Newfoundland Pine Marten

Newfoundland Pine Marten

Martes americana atrata

The Newfoundland pine marten (*Martes americana atrata*) is a subspecies of the American marten. It inhabits only the island of Newfoundland and is one of only 14 species of land mammals native to the island. It has been geographically and...

Newfoundland wolf

Newfoundland wolf

Canis lupus beothucus

The Newfoundland wolf (*Canis lupus beothucus*) was a subspecies of grey wolf that was native to Newfoundland. As a food source, the species would prey and rely on the Newfoundland Caribou. During the autumn and early winter, some wolves would...

Niagara free-tailed bat

Niagara free-tailed bat

Mops niangarae

The Niagara free-tailed bat or Niagara mops bat (*Mops niangarae*) is a species of bat in the family Molossidae known only from its holotype. It is endemic to

Democratic Republic of the Congo. Its natural habitat is either moist tropical forest...

Nicaraguan harvest mouse
Nicaraguan harvest mouse
Reithrodontomys paradoxus

The Nicaraguan harvest mouse (*Reithrodontomys paradoxus*) is a species of rodent in the family Cricetidae. It is found in Costa Rica and Nicaragua.

Nicaraguan pocket gopher
Nicaraguan pocket gopher
Orthogeomys matagalpae

Cherrie's pocket gopher (*Heterogeomys cherriei*) is a species of rodent in the family Geomyidae. It is found in Honduras, Nicaragua, and Costa Rica. It is threatened by habitat loss. Some authors classify it in the genus *Orthogeomys*, but recent...

Nicaraguan spider monkey
Nicaraguan spider monkey
Ateles geoffroyi geoffroyi

The Nicaraguan spider monkey (*Ateles geoffroyi geoffroyi*), is a subspecies of Geoffroy's spider monkey, a type of New World monkey, from Central America. It is native to Nicaragua and parts of Costa Rica closest to Nicaragua plus the Guanacaste...

Niceforo's big-eared bat
Niceforo's big-eared bat
Trinycteris nicefori

Niceforo's big-eared bat (*Trinycteris nicefori*) is a bat species from South and Central America, ranging from Chiapas to Bolivia and northeastern Brazil. Its habitat is primary and secondary forest at altitudes from sea level to 1000 m. It is...

Nicobar flying fox
Nicobar flying fox
Pteropus faunulus

The Nicobar flying fox (*Pteropus faunulus*) is a species of flying fox in the family Pteropodidae. It is endemic to India. Its natural habitats are subtropical or tropical moist lowland forest and subtropical or tropical swamps. It is threatened...

Nicobar long-tailed macaque
Nicobar long-tailed macaque
Macaca fascicularis umbrosa

The Nicobar long-tailed macaque (*Macaca fascicularis umbrosa*, popularly known as the Nicobar monkey) is a subspecies of the crab-eating macaque (*M. fascicularis*), endemic to the Nicobar Islands in the Bay of Bengal. This primate is found on...

Nicobar shrew
Nicobar shrew
Crocidura nicobarica

The Nicobar shrew or Nicobar white-tailed shrew (*Crocidura nicobarica*) is a critically endangered species of mammal in the family Soricidae. It is endemic to the Great Nicobar Island of India.

Nicobar treeshrew
Nicobar treeshrew
Tupaia nicobarica

The Nicobar treeshrew (*Tupaia nicobarica*) is a treeshrew species within the

Tupaiidae. It is endemic to the Nicobar Islands where it inhabits the islands' rain forests. It is threatened by habitat loss. Although previously listed as an endangered...

Niger Delta red colobus
Niger Delta red colobus
Piliocolobus epieni

The Niger Delta red colobus (*Piliocolobus epieni*) is a critically endangered species of colobus monkey endemic to the western part of the Niger Delta. It is threatened by hunting and habitat loss.

Nigeria-Cameroon chimpanzee
Nigeria-Cameroon chimpanzee
Pan troglodytes ellioti

The Nigeria-Cameroon chimpanzee (*Pan troglodytes ellioti*) is a subspecies of the common chimpanzee which inhabits the rainforest along the border of Nigeria and Cameroon. Male Nigeria-Cameroon chimpanzees can weigh up to 70 kilos with a body...

Nigerian free-tailed bat
Nigerian free-tailed bat
Chaerephon nigeriae

The Nigerian free-tailed bat (*Chaerephon nigeriae*) is a species of bat in the family Molossidae. It is native to two widely separated parts of Africa, and is sometimes considered to represent two separate species.

Nigerian gerbil
Nigerian gerbil
Gerbillus nigeriae

The Nigerian gerbil (*Gerbillus nigeriae*) is distributed mainly in northern Nigeria and Burkina Faso.

Nigerian mole-rat
Nigerian mole-rat
Fukomys foxi

The Nigerian mole-rat (*Fukomys foxi*) is a species of rodent in the family Bathyergidae. It is found in northern Nigeria and Cameroon. A colonial, subterranean species, its natural habitats are tropical dry lowland grassland, riverside woodland...

Nigerian shrew
Nigerian shrew
Crocidura nigeriae

The Nigerian shrew (*Crocidura nigeriae*) is a species of mammal in the family Soricidae. The animal is found in Benin, Burkina Faso, Cameroon, Ivory Coast, Nigeria, Togo, and there are claims of it also being found in Ghana. Its natural habitat...

Nikolaus's mouse
Nikolaus's mouse
Megadendromus nikolausi

Nikolaus's mouse (*Megadendromus nikolausi*) is a species of rodent in the family Nesomyidae. It is the only species in the genus *Megadendromus*. It is found only in Ethiopia. Its natural habitat is subtropical or tropical high-altitude shrubland....

Nile Lechwe
Nile Lechwe
Kobus megaceros

The Nile lechwe or Mrs Gray's lechwe (*Kobus megaceros*) is an endangered species of

antelope found in swamps and grasslands in Sudan and Ethiopia.

Nilgai

Nilgai

Boselaphus tragocamelus

The nilgai (*Boselaphus tragocamelus*) is the largest Asian antelope and is native to the Indian subcontinent. It is the sole member of the genus *Boselaphus* and was described by Peter Simon Pallas in 1766. The nilgai has been associated with Indian...

Nilgiri Langur

Nilgiri Langur

Semnopithecus johnii

The Nilgiri langur (*Semnopithecus johnii*) is a langur (a type of Old World monkey). This primate has glossy black fur on its body and golden brown fur on its head. It is similar in size and long-tailed like the gray langurs. Females have a white...

Nilgiri long-tailed tree mouse

Nilgiri long-tailed tree mouse

Vandeleuria nilagirica

Nilgiri vandeleuria (*Vandeleuria nilagirica*), also known as the Nilgiri long-tailed tree mouse and the Indian long-tailed tree mouse, is a species of rodent in the family Muridae. There has been some dispute as to whether this specimen is...

Nilgiri Marten

Nilgiri Marten

Martes gwatkinsii

The Nilgiri marten (*Martes gwatkinsii*) is the only marten species native to southern India. It lives in the hills of the Nilgiris and parts of the Western Ghats. With only around a thousand members left it is listed as Vulnerable on the IUCN Red...

Nilgiri striped squirrel

Nilgiri striped squirrel

Funambulus sublineatus

The Nilgiri striped squirrel (*Funambulus sublineatus*) is a threatened species of rodent, a small squirrel (Sciuridae) from rainforests in the southern Western Ghats, including the Nilgiris, in Peninsular India. It formerly included *Funambulus*...

Nilgiri Tahr

Nilgiri Tahr

Nilgiritragus hylocrius

The Nilgiri tahr (*Nilgiritragus hylocrius*) is an ungulate that is endemic to India. It is the state animal of Tamil Nadu. Despite its local name, it is more closely related to the sheep of the genus *Ovis* than the ibex and wild goats of the genus...

Nillu rat

Nillu rat

Rattus montanus

The Nillu rat or Sri Lankan mountain rat (*Rattus montanus*) is a species of rodent in the family Muridae. It is found only in Sri Lanka where it is known as நில்லு எலி (Nillu Eli) in Tamil and නෙලු මීයා (Nelu Meeya) in Sinhalese.

Nimba otter shrew

Nimba otter shrew

Micropotamogale lamottei

The Nimba otter shrew (*Micropotamogale lamottei*) is a dwarf otter shrew and

belongs to the mammal family Potamogalidae. Otter shrews are shrew-like afrotherian mammals found in sub-Saharan Africa. They are most closely related to the tenrecs of...

Nimba shrew

Nimba shrew

Crocidura nimbae

The Nimba shrew (*Crocidura nimbae*) is a species of mammal in the family Soricidae. It is found in Ivory Coast, Guinea, Liberia, and Sierra Leone. Its natural habitat is subtropical or tropical moist lowland forests. It is threatened by habitat...

Nimble-footed mouse

Nimble-footed mouse

Peromyscus levipes

The nimble-footed mouse (*Peromyscus levipes*) is a species of rodent in the family Cricetidae. It is found only in Mexico.

Nine-Banded Armadillo

Nine-Banded Armadillo

Dasypus novemcinctus

The Nine-banded armadillo (*Dasypus novemcinctus*) is a mammal found in North, Central, and South America. This makes it the most widespread of the armadillos. Its ancestors originated in South America and remained there until the formation of the...

Ningbing false antechinus

Ningbing false antechinus

Pseudantechinus ningbing

The Ningbing false antechinus (*Pseudantechinus ningbing*), also known as the Ningbing pseudantechinus, is a small species of carnivorous marsupial found in north-western Australia. It is locally common throughout the Kimberley region of Western...

Niobe ground squirrel

Niobe ground squirrel

Lariscus niobe

The Niobe ground squirrel (*Lariscus niobe*) is a species of rodent in the family Sciuridae. It is endemic to the mountains of western Sumatra, and eastern Java, in Indonesia. Its natural habitat is subtropical or tropical dry forests. It is...

Niobe's shrew

Niobe's shrew

Crocidura niobe

Niobe's shrew (*Crocidura niobe*) is a species of mammal in the family Soricidae. It is native to the Albertine Rift montane forests.

Noack's roundleaf bat

Noack's roundleaf bat

Hipposideros ruber

Noack's roundleaf bat (*Hipposideros ruber*) is a species of bat in the family Hipposideridae. It is found throughout tropical Africa. Its natural habitats are subtropical or tropical moist lowland forests, moist savanna, caves, and subterranean...

Noble mouse-like hamster

Noble mouse-like hamster

Calomyscus grandis

The noble mouse-like hamster or the noble calomyscus, (*Calomyscus grandis*) is a species of mouse-like hamster from Iran. It is the largest species of *Calomyscus*

and was initially described as a subspecies of *Calomyscus bailwardi*. The animal is...

Nolthenius's long-tailed climbing mouse

Nolthenius's long-tailed climbing mouse

Vandeleuria nolthenii

Nolthenius's long-tailed climbing mouse (*Vandeleuria nolthenii*), also known as Sri Lanka highland tree mouse or Podi-gas-miya (Sinhala: *ශ්‍රී ලංකා ගස් මීයා*), is a species of rodent in the family Muridae. The species is endemic to the highlands...

Nonsense rat

Nonsense rat

Rattus burrus

The nonsense rat, Nicobar Archipelago rat, or Miller's Nicobar rat (*Rattus burrus*) is endemic to the Nicobar Islands in India. It lives on Great Nicobar, Little Nicobar, and Trinket islands. On Car Nicobar Island, *Rattus palmarum* and *Rattus*...

North African boar

North African boar

Sus scrofa algira

The Barbary wild boar (*Sus scrofa algira*) is a smaller subspecies of wild boar native to North Africa including Morocco, Algeria and Tunisia. It is a prey animal for past and present predators of the Atlas Mountains, such as striped hyenas,...

North African elephant

North African elephant

Loxodonta africana pharaoensis

The North African elephant (*Loxodonta africana pharaoensis*) is an extinct possible subspecies of the African bush elephant (*Loxodonta africana*), or possibly a separate elephant species, that existed in North Africa, north of the Sahara, until...

North African elephant shrew

North African elephant shrew

Petrosaltator rozeti

The North African elephant shrew (*Petrosaltator rozeti*) or North African sengi is a species of elephant shrew in the family Macroscelididae. It is found in Algeria, Libya, Morocco, and Tunisia, and (since the extinction of the North African...

North African gerbil

North African gerbil

Dipodillus campestris

The North African gerbil (*Dipodillus campestris*) is a species of rodent in the family Muridae. It is found in North Africa where its natural habitats are arable land and rocky areas of the Maghreb, and hot Saharan deserts.

North African hedgehog

North African hedgehog

Atelerix algirus

The North African hedgehog (*Atelerix algirus*) is a mammal species in the family Erinaceidae. Little is known about this hedgehog, even though the most common breed of domesticated hedgehogs is a result of crossing a Four-toed hedgehog with a...

North African white-toothed shrew

North African white-toothed shrew

Crocidura pachyura

The North African white-toothed shrew (*Crocidura pachyura* or *Crocidura ichnusae*) is a species of mammal in the family Soricidae. It is found on the islands of Ibiza, Sardinia, and Pantelleria in the Mediterranean Sea, in northern Algeria, and

in...

North American brown lemming
North American brown lemming
Lemmus trimucronatus

The Canadian lemming or Nearctic brown lemming (*Lemmus trimucronatus*) is a small North American lemming.

North American Cougar
North American Cougar
Puma concolor couguar

The North American cougar (*Puma concolor couguar*) is a cougar subspecies in North America. It is the biggest cat in North America. It was once common in eastern North America and is still prevalent in the western half of the continent. It thus...

North American Least Shrew
North American Least Shrew
Cryptotis parva

The North American least shrew (*Cryptotis parva*) is one of the smallest mammals. It can be found in wet and dry habitats and can be active at any time of the day or night. This tiny shrew rarely lives more than a year in the wild and has many...

North American Porcupine
North American Porcupine
Erethizon dorsatum

The North American porcupine (*Erethizon dorsatum*) is the second largest rodent in North America, after the North American beaver. In addition, the range of this species is the northernmost of all porcupines. Its ancestors crossed the Atlantic...

North American River Otter
North American River Otter
Lontra canadensis

The North American river otter (*Lontra canadensis*) is a semiaquatic mammal that lives along its waterways and coasts. It is the only river otter that is found north of Mexico. These animals live in burrows and like most predators, prey upon the...

North Atlantic Right Whale
North Atlantic Right Whale
Eubalaena glacialis

The North Atlantic right whale (*Eubalaena glacialis*) is a rare baleen whale, one of three right whale species belonging to the genus *Eubalaena*, all of which were formerly classified as a single species. Because of their docile nature, their slow...

North China sika deer
North China sika deer
Cervus nippon mandarinus

The North China sika deer or Mandarin sika deer (*Cervus nippon mandarinus*) is one of the many subspecies of sika deer. It is a large subspecies with some of the most prominent spots of all subspecies, which is permanent throughout the year. It...

North Pacific Right Whale
North Pacific Right Whale
Eubalaena japonica

The North Pacific right whale (*Eubalaena japonica*) is a very large, thickset baleen whale species that is extremely rare and endangered. The Northeast Pacific

population, which summers in the southeastern Bering Sea and Gulf of Alaska, may have...

North Sulawesi Babirusa
North Sulawesi Babirusa
Babirusa celebensis

The North Sulawesi babirusa (*Babirusa celebensis*) is a pig-like animal native to Sulawesi and some nearby islands (Lembeh, Buton and Muna) in Indonesia. It has two pairs of large tusks composed of enlarged canine teeth. The upper canines...

Northeast African Cheetah
Northeast African Cheetah
Acinonyx jubatus soemmeringii

The Northeast African cheetah (*Acinonyx jubatus soemmeringii*) is a cheetah subspecies. It was first described under the scientific name *Cynailurus soemmeringii* by the Austrian zoologist Leopold Fitzinger in 1855 on the basis of a specimen from...

Northeast African mole-rat
Northeast African mole-rat
Tachyoryctes splendens

The northeast African mole-rat (*Tachyoryctes splendens*) is a species of rodent in the family Spalacidae and is found in Ethiopia, Somalia, and northwest Kenya. Its natural habitats are subtropical or tropical moist montane forests, dry savanna,...

Northeastern coyote
Northeastern coyote
Canis latrans thomasi

The coyote (*Canis latrans*) is a species of canine native to North America. It is smaller than its close relative, the wolf, and slightly smaller than the closely related eastern wolf and red wolf. It fills much of the same ecological niche as...

Northern Amazon red squirrel
Northern Amazon red squirrel
Sciurus igniventris

The northern Amazon red squirrel, *Sciurus igniventris*, is a squirrel species from South America. It occurs in Brazil, Colombia, Ecuador, Peru and Venezuela.

Northern Baja deer mouse
Northern Baja deer mouse
Peromyscus fraterculus

The Northern Baja deer mouse (*Peromyscus fraterculus*) is a species of deer mouse native to Southern California and the Baja California peninsula as well as several islands in the Gulf of California. *P. fraterculus* was previously considered a...

Northern Bat
Northern Bat
Eptesicus nilssonii

The Northern bat (*Eptesicus nilssonii*) is the most abundant species of bat in northern Eurasia. They are nocturnal and fast-flying, adapted to hunting airborne insects using echolocation.

Northern bettong
Northern bettong
Bettongia tropica

The northern bettong (*Bettongia tropica*) is a small potoroid marsupial which is restricted to some areas of mixed open Eucalyptus woodlands and Allocasuarina forests bordering rainforests in far northeastern Queensland, Australia. They are known...

Northern birch mouse
Northern birch mouse
Sicista betulina

The northern birch mouse (*Sicista betulina*) is a small rodent about 5 to 8 cm long (without the tail), weighing 5 to 13 g. It lives in northern Europe and Asia in forest and marsh zones. It hibernates in burrows. It eats shoots, grains, berries,...

Northern bog lemming
Northern bog lemming
Synaptomys borealis

The northern bog lemming (*Synaptomys borealis*) is a small North American lemming. It is one of two species in the genus *Synaptomys*, the other being the southern bog lemming.

Northern Bottlenose Whale
Northern Bottlenose Whale
Hyperoodon ampullatus

The northern bottlenose whale (*Hyperoodon ampullatus*) is a species of beaked whale in the ziphiid family, being one of two members of the genus *Hyperoodon*. The northern bottlenose whale was hunted heavily by Norway and Britain in the 19th and...

Northern broad-nosed bat
Northern broad-nosed bat
Scotorepens sanborni

The northern broad-nosed bat (*Scotorepens sanborni*) is a species of the vespertilionid family of microbats. It can be found in northern Australia, Timor-Leste, and Papua New Guinea.

Northern brown bandicoot
Northern brown bandicoot
Isodon macrourus

The northern brown bandicoot (*Isodon macrourus*), a marsupial species, is a bandicoot found only on the northern and eastern coasts of Australia and nearby islands, mainly Papua New Guinea. It is not, however, found far inland.

Northern brown howler
Northern brown howler
Alouatta guariba guariba

The northern brown howler (*Alouatta guariba guariba*) is the type subspecies of the brown howler, native to Brazil. It is listed as critically endangered, with fewer than 250 individuals restricted to the vicinity of the Jequitinhonha River. The...

Northern brush-tailed phascogale
Northern brush-tailed phascogale
Phascogale pirata

The northern brush-tailed phascogale (*Phascogale pirata*), also known as the northern brush-tailed wambenger or northern brush-tailed mousesack is a species of marsupial in the family Dasyuridae. It is endemic to northern Australia.

Northern brushtail possum
Northern brushtail possum
Trichosurus arnhemensis

The northern brushtail possum (*Trichosurus arnhemensis*) is a nocturnal marsupial inhabiting northern Australia. The northern brushtail possum is sometimes considered a species, however more often than not is considered a subspecies of the common...

Northern buffed-cheeked gibbon
Northern buffed-cheeked gibbon
Nomascus annamensis

The northern buffed-cheeked gibbon (*Nomascus annamensis*) is a newly discovered species of crested gibbon which is found in Vietnam, Cambodia, and Laos. Its habitat is in the humid subtropical and seasonal tropical forests of these countries.

Northern caenolestid
Northern caenolestid
Caenolestes convelatus

The northern caenolestid (*Caenolestes convelatus*), also known as the blackish shrew opossum, is a shrew opossum found in Colombia and Ecuador. It is listed as Vulnerable by the IUCN.

Northern cave bat
Northern cave bat
Vespadelus caurinus

The northern cave bat (*Vespadelus caurinus*) is a vespertilionid bat that occurs in most of Australia.

Northern Chinese boar
Northern Chinese boar
Sus scrofa moupinensis

The northern Chinese boar (*Sus scrofa moupinensis*) is a subspecies of wild boar native to China and Vietnam. The subspecies was described by Alphonse Milne-Edwards in 1871. It also occurs in Sichuan. It is likely to be the ancestor of domestic...

Northern Collared Lemming
Northern Collared Lemming
Dicrostonyx groenlandicus

The Northern collared lemming (*Dicrostonyx groenlandicus*) is a small rodent found in Arctic North America and Wrangel Island. At one time, it was considered to be a subspecies of the Arctic lemming (*Dicrostonyx torquatus*). Some sources believe...

Northern common cuscus
Northern common cuscus
Phalanger orientalis

The northern common cuscus (*Phalanger orientalis*), also known as the grey cuscus, is a species of marsupial in the family Phalangeridae native to northern New Guinea and adjacent smaller islands, but is now also found in the Bismarck...

Northern fin whale
Northern fin whale
Balaenoptera physalus physalus

The northern fin whale (*Balaenoptera physalus physalus*) is a subspecies of fin whale that lives in the North Atlantic Ocean and North Pacific Ocean. It has been proposed that the northern Pacific population represents a separate subspecies, B....

Northern Flying Squirrel
Northern Flying Squirrel
Glaucomys sabrinus

The Northern flying squirrel (*Glaucomys sabrinus*) is one of three species of the genus *Glaucomys*, the only flying squirrels found in North America. They are found in forests across North America. They are proficient gliders but uncoordinated...

Northern free-tailed bat
Northern free-tailed bat

Chaerephon johorensis

The northern free-tailed bat (*Chaerephon johorensis*) is a species of bat in the family Molossidae. It is found in Indonesia and Malaysia.

Northern freetail bat

Northern freetail bat

Chaerephon jobensis

The northern freetail bat (*Chaerephon jobensis*) is a species of bat found in Yapen, Seram Island, Western New Guinea and Northern Australia.

Northern Fur Seal

Northern Fur Seal

Callorhinus ursinus

The northern fur seal (*Callorhinus ursinus*) is an eared seal found along the north Pacific Ocean, the Bering Sea, and the Sea of Okhotsk. It is the largest member of the fur seal subfamily (Arctocephalinae) and the only living species in the...

Northern ghost bat

Northern ghost bat

Diclidurus albus

The northern ghost bat (*Diclidurus albus*) is a bat species from South America, Trinidad, and Central America. It is a relatively rare, completely white, insectivorous bat, with an unusual sac at the base of its tail.

Northern giant mouse lemur

Northern giant mouse lemur

Mirza zaza

The northern giant mouse lemur (*Mirza zaza*), or northern dwarf lemur, is a species of lemur discovered in 2005. They are part of the primate order, and classified in the family Cheirogaleidae. Previously, both populations of giant mouse lemurs...

Northern glider

Northern glider

Petaurus abiri

The northern glider (*Petaurus abidi*) is a species of marsupial in the family Petauridae. It is endemic to Papua New Guinea, becoming known to science in 1981 after being discovered in the Torricelli Mountains. This species has been found in...

Northern gracile opossum

Northern gracile opossum

Gracilinanus marica

The northern gracile opossum (*Gracilinanus marica*) is a species of opossum in the family Didelphidae. It is found in Colombia and Venezuela. Its natural habitat is subtropical or tropical moist lowland forests.

Northern grass mouse

Northern grass mouse

Necromys urichi

The northern grass mouse, or northern akodont, (*Necromys urichi*) is a species of rodent in the family Cricetidae. It is found in Brazil, Colombia, Venezuela and Trinidad and Tobago.

Northern grasshopper mouse

Northern grasshopper mouse

Onychomys leucogaster

The northern grasshopper mouse (*Onychomys leucogaster*) is a North American carnivorous rodent of the family Cricetidae. It ranges over much of the western

part of the continent, from southern Saskatchewan and central Washington to Tamaulipas in...
Northern Greater Galago
Northern Greater Galago
Otolemur garnettii

The Northern greater galago (*Otolemur garnettii*) is an arboreal primate found only in Africa. They are quadrupedal and can hop short distances from tree to tree.
Northern groove-toothed shrew mouse
Northern groove-toothed shrew mouse
Microhydromys richardsoni

The northern groove-toothed shrew mouse (*Microhydromys richardsoni*) is a species of rodent in the family Muridae found in West Papua, Indonesia and Papua New Guinea.
Northern Hairy-Nosed Wombat
Northern Hairy-Nosed Wombat
Lasiorhinus krefftii

The Northern hairy-nosed wombat (*Lasiorhinus krefftii*) is one of three extant species of Australian marsupials known as wombats. It is one of the rarest land mammals in the world and is critically endangered. Its historical range extended across...
Northern hopping mouse
Northern hopping mouse
Notomys aquilo

The northern hopping mouse (*Notomys aquilo*) is a species of rodent in the family Muridae. It is found only in coastal northern Australia, from Arnhem Land to the Cobourg Peninsula. This mouse weighs 25 to 30 grams and is brown above and white...
Northern leaf-nosed bat
Northern leaf-nosed bat
Hipposideros stenotis

The northern leaf-nosed bat (*Hipposideros stenotis*) is a micro-bat of the family Hipposideridae, known as "leaf-nosed" bats. The species is endemic to northern regions of Australia. They are highly manoeuvrable in flight, and use echolocation to...
Northern Lion
Northern Lion
Panthera leo leo

The Northern lion (*Panthera leo leo*) is a lion subspecies, which is present in West Africa, northern Central Africa and India. In West and Central Africa, it is restricted to fragmented and isolated populations with a declining trajectory. The...
Northern little yellow-eared bat
Northern little yellow-eared bat
Vampyressa thuyne

The northern little yellow-eared bat (*Vampyressa thuyne*) is a bat species from South and Central America, where it is found from southern Mexico to Bolivia, the Guianas and western Brazil.
Northern Long-Eared Bat
Northern Long-Eared Bat
Myotis septentrionalis

The Northern long-eared bat (*Myotis septentrionalis*) is a small bat native to North America. It is distinguishable by its long ears when comparing it to other bats in

its genus. The geographic range of this species includes 37 states.

Northern Luzon Giant Cloud Rat

Northern Luzon Giant Cloud Rat

Phloeomys pallidus

The Northern Luzon giant cloud rat (*Phloeomys pallidus*) is a large rodent only found in the Philippines. It is also known as the Northern Luzon slender-tailed cloud rat, and as bu-ot in Filipino.

Northern Luzon shrew-rat

Northern Luzon shrew-rat

Crunomys fallax

The Northern Luzon shrew-rat (*Crunomys fallax*) is a species of rodent in the family Muridae. It is found only in the Philippines. Its natural habitat is subtropical or tropical dry forest. It is threatened by habitat loss.

Northern Lynx

Northern Lynx

Lynx lynx lynx

The Northern lynx (*Lynx lynx lynx*) is a subspecies of the Eurasian lynx (*Lynx lynx*).

Northern mole vole

Northern mole vole

Ellobius talpinus

The northern mole vole (*Ellobius talpinus*) is a species of rodent in the family Cricetidae. It is distributed over large parts of Eastern Europe and Asia.

Northern Muriqui

Northern Muriqui

Brachyteles hypoxanthus

The northern muriqui (*Brachyteles hypoxanthus*) is one of two species of muriqui. They are also known as the woolly spider monkey as they exhibit the woollen pelt of woolly monkeys and the long prehensile tail of spider monkeys. They are the...

Northern nail-tail wallaby

Northern nail-tail wallaby

Onychogalea unguifera

The northern or sandy nail-tail wallaby (*Onychogalea unguifera*) is a species of macropod found across northern Australia on arid and sparsely wooded plains. The largest species of the genus *Onychogalea*, it is a solitary and nocturnal herbivorous...

Northern Naked-Tailed Armadillo

Northern Naked-Tailed Armadillo

Cabassous centralis

The Northern naked-tailed armadillo (*Cabassous centralis*) is one of only two species of armadillo found outside of South America, the other being the more widely distributed Nine-banded armadillo. Unlike other armadillos with which this species...

Northern needle-clawed bushbaby

Northern needle-clawed bushbaby

Euoticus pallidus

The northern needle-clawed bushbaby (*Euoticus pallidus*) is a species of strepsirrhine primate in the family Galagidae. It is found in the coastal region of Cameroon and Nigeria, and on the island of Bioko, Equatorial Guinea in lower-elevation...

Northern Olingo

Northern Olingo
Bassaricyon gabbii

The Northern olingo (*Bassaricyon gabbii*) is a tree-dwelling member of the raccoon family. It was the first species of olingo that was described scientifically. The part of the scientific name of this animal honors William More Gabb (1839-1878),...

Northern Palawan tree squirrel
Northern Palawan tree squirrel
Sundasciurus juvencus

The Northern Palawan tree squirrel (*Sundasciurus juvencus*) is a species of rodent in the family Sciuridae. It is endemic to the Philippines. Its natural habitat is subtropical or tropical dry forests. It is threatened by habitat loss.

Northern Palm Squirrel
Northern Palm Squirrel
Funambulus pennantii

The Northern palm squirrel (*Funambulus pennantii*) is a species of rodent found in Asia. It is semi-arboreal and lives in tropical and subtropical forests and many other rural and urban habitats.

Northern Pig-Tailed Macaque
Northern Pig-Tailed Macaque
Macaca leonina

The Northern pig-tailed macaque (*Macaca leonina*) is a species of macaque in the family Cercopithecidae. It was considered a subspecies of the Southern pig-tailed macaque but is now classified as individual species.

Northern Pika
Northern Pika
Ochotona hyperborea

The northern pika (*Ochotona hyperborea*) is a species of pika found across mountainous regions of northern Asia, from the Ural Mountains to northern Japan and south through Mongolia, Manchuria and northern Korea. An adult northern pika has a body...

Northern pipistrelle
Northern pipistrelle
Pipistrellus westralis

The northern pipistrelle (*Pipistrellus westralis*), also known as Koopman's or the mangrove pipistrelle, is a species of vesper bat found only in Australia. It is one of Australia's smallest bat species. On average, it weighs 3 g (0.11 oz).

Northern Plains Gray Langur
Northern Plains Gray Langur
Semnopithecus entellus

The Northern plains gray langur (*Semnopithecus entellus*) is a species of primate found in Asia. Langurs mostly walk quadrupedally and spend half of their time on the ground and the other half in trees.

Northern Pocket Gopher
Northern Pocket Gopher
Thomomys talpoides

The Northern pocket gopher (*Thomomys talpoides*) is a small burrowing rodent from North America. These animals are commonly known for their extensive tunneling activities and their ability to destroy farms and gardens.

Northern pudu
Northern pudu
Pudu mephistophiles

The northern pudu (*Pudu mephistophiles*, Mapudungun püdü or püdü) is a species of South American deer native to the Andes of Colombia, Venezuela, Peru and Ecuador. It is the world's smallest deer and is classified as Data Deficient in the IUCN Red...

Northern pygmy mouse
Northern pygmy mouse
Baiomys taylori

The northern pygmy mouse (*Baiomys taylori*) is a species of rodent in the family Cricetidae. It is known as ratón-pigmeo norteño in the Spanish-speaking areas of its range. It is found in Mexico and the United States. Parasites of the northern...

Northern Quoll
Northern Quoll
Dasyurus hallucatus

The northern quoll (*Dasyurus hallucatus*), also known as the northern native cat, the North Australian native cat or the satanellus is a carnivorous marsupial native to Australia.

Northern red-backed vole
Northern red-backed vole
Myodes rutilus

The northern red-backed vole (*Myodes rutilus*) is a small slender vole found in Alaska, northern Canada, Scandinavia and northern Russia.

Northern red-sided opossum
Northern red-sided opossum
Monodelphis brevicaudata

The northern red-sided opossum or the Guianan short-tailed opossum, *Monodelphis brevicaudata*, is an opossum species from South America. It is found in Bolivia, Brazil, French Guiana, Guyana, Suriname and Venezuela.

Northern Right Whale Dolphin
Northern Right Whale Dolphin
Lissodelphis borealis

Northern right whale dolphins (*Lissodelphis borealis*) are small, slender species of cetacean that live in the cold and temperate waters of the North Pacific Ocean.

They lack a dorsal fin and appear superficially porpoise-like. Northern right...
Northern rock mouse
Northern rock mouse
Peromyscus nasutus

The northern rock mouse (*Peromyscus nasutus*) is a species of rodent in the family Cricetidae. It is found in Mexico and the United States.

Northern Rocky Mountain wolf
Northern Rocky Mountain wolf
Canis lupus irremotus

The northern Rocky Mountain wolf (*Canis lupus irremotus*), also known as the northern Rocky Mountain timber wolf, is a subspecies of gray wolf native to the northern Rocky Mountains. It is a light-colored, medium to large-sized subspecies with a...

Northern rufous mouse lemur
Northern rufous mouse lemur
Microcebus tavaratra

The northern rufous mouse lemur (*Microcebus tavaratra*), northern brown mouse lemur, or Tavaratra mouse lemur is found in Northern Madagascar from the Ankarana

National Park in the West to the river Manambato in Northeast and up to the Irodo...

Northern Short-Tailed Shrew
Northern Short-Tailed Shrew
Blarina brevicauda

The Northern short-tailed shrew (*Blarina brevicauda*) is the largest shrew in its genus and occurs in the northeastern region of North America. It is a highly active, and voracious insectivore. It is notable in that it is one of the few venomous...

Northern shrew tenrec
Northern shrew tenrec
Microgale jobihely

The northern shrew tenrec (*Microgale jobihely*) is a species of mammal in the family Tenrecidae. It is endemic to Madagascar, where it has a restricted disjoint range in two locations 485 km apart in the north and east of the island. In the...

Northern smooth-tailed treeshrew
Northern smooth-tailed treeshrew
Dendrogale murina

The northern smooth-tailed treeshrew (*Dendrogale murina*) is a species of treeshrew in the family Tupaiidae found in Cambodia, Thailand, and Vietnam. Their diet primarily consists of invertebrates but rarely includes fruits and plants.

Northern Sportive Lemur
Northern Sportive Lemur
Lepilemur septentrionalis

The northern sportive lemur (*Lepilemur septentrionalis*), also known as the Sahafary sportive lemur or northern weasel lemur, is a species of lemur in the family Lepilemuridae. It is endemic to Madagascar. As a result of severe ecological and...

Northern Sulawesi echiothrix
Northern Sulawesi echiothrix
Echiothrix leucura

The Northern Sulawesi echiothrix or Sulawesi spiny rat (*Echiothrix leucura*) is a species of rodent in the family Muridae. It is endemic to northeastern Sulawesi, Indonesia.

Northern Sumatran Rhinoceros
Northern Sumatran Rhinoceros
Dicerorhinus sumatrensis lasiotis

The Northern Sumatran rhinoceros (*Dicerorhinus sumatrensis lasiotis*) was the most widespread subspecies of Sumatran rhinoceros. It was also the only known subspecies native to mainland Asia. While it has been officially declared extinct on...

Northern sword-nosed bat
Northern sword-nosed bat
Lonchorhina inusitata

The northern sword-nosed bat (*Lonchorhina inusitata*) is a bat species from South America. It is found in Brazil, French Guiana, Suriname and Venezuela.

Northern Tamandua
Northern Tamandua
Tamandua mexicana

The northern tamandua (*Tamandua mexicana*) is a species of tamandua, an anteater in the family Myrmecophagidae. They live in tropical and subtropical forests from southern Mexico, through Central America, and to the edge of the northern Andes.

Northern three-striped opossum
Northern three-striped opossum
Monodelphis americana

The northern three-striped opossum (*Monodelphis americana*) is an opossum species from South America. It is endemic to Atlantic Forest ecoregions of coastal Brazil.

Northern three-toed jerboa
Northern three-toed jerboa
Dipus sagitta

The northern three-toed jerboa (*Dipus sagitta*) is a species of rodent in the family Dipodidae. It is the only extant species within the genus *Dipus*. It ranges across Iran, Uzbekistan, Turkmenistan, Kazakhstan, Russia, China and Mongolia. A common...

Northern Treeshrew
Northern Treeshrew
Tupaia belangeri

The northern treeshrew (*Tupaia belangeri*) is a treeshrew species native to Southeast Asia. In 1841, the German zoologist Johann Andreas Wagner first used the specific name *Cladobates belangeri* for treeshrews that had been collected in Pegu during...

Northern viscacha
Northern viscacha
Lagidium peruanum

The northern viscacha (*Lagidium peruanum*) is a species of viscacha, a rodent in the family Chinchillidae. It is known from Peru and Chile, at elevations from 300 to 5000 m, and may also be present in Bolivia.

Northern voalavo
Northern voalavo
Voalavo gymnocaudus

The northern voalavo (*Voalavo gymnocaudus*), also known as the naked-tailed voalavo or simply the voalavo, is a rodent in the family Nesomyidae found in the highlands of northern Madagascar. Discovered in 1994 and formally described in 1998, it...

Northern water rat
Northern water rat
Paraleptomys rufilatus

The northern water rat (*Paraleptomys rufilatus*) is an endangered species of rodent in the family Muridae found in the highlands of New Guinea.

Northern white rhinoceros
Northern white rhinoceros
Ceratotherium simum cottoni

The northern white rhinoceros (*Ceratotherium simum cottoni*), or northern square-lipped rhinoceros, is one of two subspecies of the white rhinoceros (the other being the southern white rhinoceros). Formerly found in several countries in East and...

Northern White-Breasted Hedgehog
Northern White-Breasted Hedgehog
Erinaceus roumanicus

Northern white-breasted hedgehogs (*Erinaceus roumanicus*) are small mammals easily recognized by their spines. They nocturnal secretive animals and roll into a tight spiny ball when feeling threatened.

Northern yellow bat
Northern yellow bat

Lasiurus intermedius

The northern yellow bat (*Dasypterus intermedius*) is a non-migratory bat in the family Vespertilionidae, typically active year-round except during abnormally frigid winter weather, during which they will induce torpor.

Northwest African Cheetah

Northwest African Cheetah

Acinonyx jubatus hecki

The Northwest African cheetah (*Acinonyx jubatus hecki*) is a cheetah subspecies native to the Sahara and the Sahel. It was described by German zoologist Max Hilzheimer in 1913 under the scientific name *Acinonyx hecki*.

Northwestern deer mouse

Northwestern deer mouse

Peromyscus keeni

The northwestern deer mouse or Keen's mouse (*Peromyscus keeni*) is a species of rodent in the family Cricetidae. It is found in British Columbia in Canada and in Alaska and Washington in the United States. It was named after the Rev. John Henry...

Northwestern fat mouse

Northwestern fat mouse

Steatomys caurinus

The northwestern fat mouse (*Steatomys caurinus*) is a species of rodent in the family Nesomyidae. It is native to tropical West Africa where it occurs on grassland and crop land.

Northwestern Wolf

Northwestern Wolf

Canis lupus occidentalis

The Northwestern wolf (*Canis lupus occidentalis*) is a subspecies of Gray wolf in western North America. Arguably it is the largest grey wolf subspecies in the world.

Norway Lemming

Norway Lemming

Lemmus lemmus

The Norway lemming, also known as the Norwegian lemming (*Lemmus lemmus*) is a common species of lemming found in northern Fennoscandia, where it is the only vertebrate species endemic to the region. The Norway lemming dwells in tundra and fells,...

Norwegian red deer

Norwegian red deer

Cervus elaphus atlanticus

The Norwegian red deer (*Cervus elaphus atlanticus*) is a small subspecies of red deer native to Norway. Today it is farmed on a commercial basis since the 1980s.

Notiomys

Notiomys

Notiomys edwardsii

Notiomys edwardsii, also known as Edwards's long-clawed mouse, Edward's long-clawed akodont, or Milne-Edwards' long-clawed mouse, is a rodent in the tribe Abrotrichini from southern Argentina. It is the only species in the genus *Notiomys*,...

Notiosorex cockrumi

Notiosorex cockrumi

Notiosorex cockrumi

Notiosorex cockrumi, also called Cockrum's gray shrew or Cockrum's desert shrew, is a tiny species of shrews named in 2003. This red-toothed shrew, which is as light as a penny, is the first new mammal species from Arizona since 1977. Its range...

Nubian Giraffe

Nubian Giraffe

Giraffa camelopardalis camelopardalis

The Nubian giraffe (*Giraffa camelopardalis camelopardalis*) is the nominate subspecies of giraffe. It used to be widespread everywhere on Northeast Africa. It is currently extinct in the wild of the Democratic Republic of Congo, Egypt and Eritrea.

Nubian Ibex

Nubian Ibex

Capra nubiana

The Nubian ibex (*Capra nubiana*) is a rare desert-dwelling goat species found in mountainous areas of Africa, and the Middle East. It was historically considered to be a subspecies of the Alpine ibex (*C. ibex*), but is now considered a distinct...

Nubian Wild Ass

Nubian Wild Ass

Equus africanus africanus

The Nubian wild ass (*Equus africanus africanus*) is the nominate subspecies of African wild ass, and one of the ancestors of the domestic donkey, which was domesticated about 6,000 years ago. It is presumed to be extinct, though two populations...

Nubra pika

Nubra pika

Ochotona nubrica

The Nubra pika (*Ochotona nubrica*) is a species of mammal of the pika family, Ochotonidae. It is found in Bhutan, China, India, Nepal, and Pakistan.

Nullarbor dwarf bettong

Nullarbor dwarf bettong

Bettongia pusilla

The Nullarbor dwarf bettong, *Bettongia pusilla*, was a potoroine marsupial that occurred in Australia. The animal is only known from skeletons found in caves of the Nullarbor Plain and is now classified as recently extinct.

Numbat

Numbat

Myrmecobius fasciatus

The numbat (*Myrmecobius fasciatus*) is an insectivorous marsupial that was once widespread across southern Australia, but is now restricted to several small colonies in Western Australia. It is therefore considered an endangered species and...

Nusatenggara short-nosed fruit bat

Nusatenggara short-nosed fruit bat

Cynopterus nusatenggara

The Nusatenggara short-nosed fruit bat (*Cynopterus nusatenggara*) is a species of megabat within the family Pteropodidae found in Indonesia. It has three subspecies: *C. n. nusatenggara* *C. n. sinagai* *C. n. wetarensis*

Nut-colored yellow bat

Nut-colored yellow bat

Scotophilus nux

The nut-colored yellow bat (*Scotophilus nux*) is a species of vesper bat. It can be

found in Cameroon, Democratic Republic of the Congo, Ivory Coast, Equatorial Guinea, Ghana, Guinea, Kenya, Liberia, Nigeria, Rwanda, Sierra Leone, and Uganda. It...

Nyala

Nyala

Tragelaphus angasii

The nyala (*Tragelaphus angasii*) is a spiral-horned antelope native to southern Africa. It was first described in 1849 by George French Angas. They are very shy cautious creatures. They inhabit thickets within dense and dry savanna woodlands and...

Nycticebus bancanus

Nycticebus bancanus

Nycticebus bancanus

The Bangka slow loris (*Nycticebus bancanus*) is a strepsirrhine primate and a species of slow loris that is native to southwestern Borneo and the island of Bangka. Originally considered a subspecies or synonym of the Bornean slow loris...

Nycticebus borneanus

Nycticebus borneanus

Nycticebus borneanus

Nycticebus borneanus, the Bornean slow loris, is a strepsirrhine primate and a species of slow loris that is native to central south Borneo in Indonesia. Formerly considered a subspecies or synonym of *N. menagensis*, it was promoted to full...

Nycticebus kayan

Nycticebus kayan

Nycticebus kayan

The Kayan River slow loris (*Nycticebus kayan*) is a strepsirrhine primate and a species of slow loris that is native to the northern and central highland region of the island of Borneo. The species was originally thought to be a part of the...

Nyctophilus arnhemensis

Nyctophilus arnhemensis

Nyctophilus arnhemensis

Nyctophilus arnhemensis, known as the northern or Arnhem long-eared bat, is a species of Chiroptera (bats) native to northern regions of Australia. The distribution range is from north-western Queensland to northern Western Australia.

Nyctophilus corbeni

Nyctophilus corbeni

Nyctophilus corbeni

The south-eastern long-eared bat or Corben's long-eared bat (*Nyctophilus corbeni*), is a species of bat found in Australia. It occurs in the woodlands of the Murray Darling Basin and adjacent areas.

Nyctophilus daedalus

Nyctophilus daedalus

Nyctophilus daedalus

Nyctophilus daedalus is a species of bat in the family Vespertilionidae, a flying mammal endemic to northern Australia. They are also referred to as the pallid long-eared bat or northern long-eared bat.

Nyctophilus geoffroyi

Nyctophilus geoffroyi

Nyctophilus geoffroyi

Nyctophilus geoffroyi is a vespertilionid bat, a flying nocturnal mammal found in Australia. The species is relatively common. They have been referred to as the

lesser long-eared bat.
Nyctophilus major
Nyctophilus major
Nyctophilus major

Nyctophilus major, referred to as a western long-eared bat, is a species found in forests and woodlands of Southwest Australia.

Nyika climbing mouse
Nyika climbing mouse
Dendromus nyikae

The Nyika climbing mouse (*Dendromus nyikae*) is a species of rodent in the family Nesomyidae. It is found in Angola, Democratic Republic of the Congo, Malawi, Mozambique, South Africa, Tanzania, Zambia, and Zimbabwe. Its natural habitat is moist...

Nyika rock rat
Nyika rock rat
Aethomys nyikae

The Nyika rock rat (*Aethomys nyikae*) is a species of rodent in the family Muridae found in Angola, the Democratic Republic of the Congo, Malawi, and Zambia. Its natural habitat is subtropical or tropical dry forest.

Nyiro shrew
Nyiro shrew
Crocidura macowi

The Nyiro shrew (*Crocidura macowi*) is a species of mammal in the family Soricidae. It is endemic to Kenya.

O'Connell's spiny rat
O'Connell's spiny rat
Proechimys oconnelli

O'Connell's spiny-rat (*Proechimys oconnelli*) is a species of rodent in the family Echimyidae. It is endemic to Colombia.

Oaxaca giant deer mouse
Oaxaca giant deer mouse
Megadontomys cryophilus

The Oaxaca giant deer mouse (*Megadontomys cryophilus*) is a species of rodent in the family Cricetidae. It is found only in Mexico.

Oaxacan broad-clawed shrew
Oaxacan broad-clawed shrew
Cryptotis peregrina

The Oaxacan broad-clawed shrew (*Cryptotis peregrina*) is a species of mammal in the family Soricidae. It is found in Oaxaca in Mexico.

Oaxacan pocket gopher
Oaxacan pocket gopher
Orthogeomys cuniculus

The giant pocket gopher (*Orthogeomys grandis*) is a species of rodent in the family Geomyidae. It is found in Mexico, Guatemala, El Salvador and Honduras. It is the type species of the genus *Orthogeomys* ; some zoologists also include in this...

Obi mosaic-tailed rat
Obi mosaic-tailed rat
Melomys obiensis

The Obi mosaic-tailed rat (*Melomys obiensis*) is a species of rodent in the family Muridae. It is found only in Indonesia.

Occidental gerbil
Occidental gerbil
Gerbillus occiduus

The occidental gerbil (*Gerbillus occiduus*) is distributed mainly southwestern Morocco. Less than 250 individuals of this species are thought to persist in the wild.

Ocelot
Ocelot
Leopardus pardalis

The ocelot (*Leopardus pardalis*) is a medium-sized wild cat that was first described by Carl Linnaeus in 1758. The association of the ocelot with humans dates back to the Aztec and Incan civilizations; it has occasionally been kept as a pet.

Ochotona spanglei
Ochotona spanglei
Ochotona spanglei

Ochotona spanglei is an extinct species of pika (mammal in the family Ochotonidae), known from Late Miocene - Early Pliocene fossil from Oregon (USA). Fossils were also found in Nebraska referred to as *Ochotona cf. spanglei*. *Ochotona spanglei* is...

Ochre bush squirrel
Ochre bush squirrel
Paraxerus ochraceus

The ochre bush squirrel (*Paraxerus ochraceus*) is a species of rodent in the family Sciuridae found in Kenya, Somalia, and Tanzania. Its natural habitat is dry savanna.

Ochre mole-rat
Ochre mole-rat
Fukomys ochraceocinereus

The Ochre mole-rat (*Fukomys ochraceocinereus*) is a species of rodent in the family Bathyergidae. It is found in Central African Republic, Democratic Republic of the Congo, South Sudan, and Uganda. Its natural habitats are moist savanna,...

Oecomys auyantepui
Oecomys auyantepui
Oecomys auyantepui

Oecomys auyantepui, also known as the Guianan oecomys and north Amazonian arboreal rice rat, is a species of rodent in the genus *Oecomys* from South America. It is found in Guyana, Suriname, French Guiana, and nearby regions of Venezuela and...

Oecomys bicolor
Oecomys bicolor
Oecomys bicolor

Oecomys bicolor, also known as the white-bellied oecomys or bicolored arboreal rice rat, is a species of rodent in the genus *Oecomys* of family Cricetidae. It has a wide distribution in the Amazon biome, occurring in northwestern Brazil, northern...

Oecomys catherinae
Oecomys catherinae
Oecomys catherinae

Oecomys catherinae, also known as the Atlantic Forest oecomys, is a species of rodent in the genus *Oecomys* from eastern Brazil. It lives in a variety of habitats including in the Atlantic Forest, Cerrado and Caatinga ecoregions. It is related to...

Oecomys cleberi

Oecomys cleberi
Oecomys cleberi

Oecomys cleberi, also known as Cleber's oecomys or Cleber's arboreal rice rat, is a species of rodent in the genus Oecomys of family Cricetidae. Known only from the Federal District of Brazil, its taxonomic status relative to O. concolor and O....

Oecomys concolor
Oecomys concolor
Oecomys concolor

Oecomys concolor, also known as the unicolored oecomys, unicolored rice rat, or unicolored arboreal rice rat, is a species of rodent in the genus Oecomys of family Cricetidae. It is found in tropical rainforest in the Amazon biome, but its range...

Oecomys flavicans
Oecomys flavicans
Oecomys flavicans

Oecomys flavicans, also known as the tawny oecomys or yellow arboreal rice rat, is a species of rodent in the genus Oecomys of family Cricetidae. It is found in the mountains of northwestern Venezuela and nearby Colombia.

Oecomys paricola
Oecomys paricola
Oecomys paricola

Oecomys paricola, also known as the Brazilian oecomys, Brazilian arboreal rice rat, or South Amazonian arboreal rice rat, is a species of rodent in the genus Oecomys of family Cricetidae. It is found in central Brazil south of the Amazon, where...

Oecomys phaeotis
Oecomys phaeotis
Oecomys phaeotis

Oecomys phaeotis, also known as the dusky arboreal rice rat or dusky oecomys, is a species of rodent in the genus Oecomys of family Cricetidae. It is found on the eastern slopes of the Andes of Peru, at 1500 to 2000 m altitude.

Oecomys rex
Oecomys rex
Oecomys rex

Oecomys rex, also known as the regal oecomys or king arboreal rice rat, is a species of rodent in the genus Oecomys of family Cricetidae. It is found in Guyana, Suriname, French Guiana, and nearby parts of Venezuela and Brazil.

Oecomys roberti
Oecomys roberti
Oecomys roberti

Oecomys roberti, also known as Robert's oecomys or Robert's arboreal rice rat, is a rodent species from South America in the genus Oecomys. It has a broad distribution in the Amazon biome, being found in Bolivia, Brazil, French Guiana, Guyana,...

Oecomys rutilus
Oecomys rutilus
Oecomys rutilus

Oecomys rutilus, also known as the reddish oecomys or red arboreal rice rat, is a species of rodent in the genus Oecomys of family Cricetidae. It is found in Guyana, Suriname, French Guiana, and nearby regions of Brazil and Venezuela.

Oecomys speciosus
Oecomys speciosus
Oecomys speciosus

Oecomys speciosus, also known as the savannah oecomys, arboreal rice rat, or Venezuelan arboreal rice rat, is a species of rodent in the genus *Oecomys* of family Cricetidae. It ranges over northeastern Colombia and much of Venezuela, including the...

Oecomys superans
Oecomys superans
Oecomys superans

Oecomys superans, also known as the large oecomys or foothill arboreal rice rat, is a species of rodent in the genus *Oecomys* of family Cricetidae. It is found along the eastern slope of the Andes in southern Colombia, Ecuador, and Peru and east...

Oecomys sydandersoni
Oecomys sydandersoni
Oecomys sydandersoni

Oecomys sydandersoni is an arboreal species of rodent in the genus *Oecomys*. It lives in forest patches in a small area in eastern Bolivia. It is a medium-sized species, weighing about 45 g (1.6 oz), with mostly grayish and brownish fur and short...

Oecomys trinitatis
Oecomys trinitatis
Oecomys trinitatis

Oecomys trinitatis, also known as the long-furred oecomys, long-furred rice rat, Trinidad arboreal rice rat, or big arboreal rice rat, is a species of rodent in the genus *Oecomys* of family Cricetidae. As currently constituted, it has a wide...

Ogilby's duiker
Ogilby's duiker
Cephalophus ogilbyi

Ogilby's duiker (*Cephalophus ogilbyi*) is a small antelope found in Sierra Leone, Liberia, Ghana, southeastern Nigeria, Bioko Island and possibly Gabon. No subspecies are recognized. The two former subspecies, the white-legged duiker *Cephalophus*...

Ogilvie Mountains collared lemming
Ogilvie Mountains collared lemming
Dicrostonyx nunatakensis

The Ogilvie Mountains collared lemming (*Dicrostonyx nunatakensis*) is a species of rodent in the family Cricetidae. It is found only in Yukon Territory, Canada. Its natural habitat is tundra.

Ognev's long-eared bat
Ognev's long-eared bat
Plecotus ognevi

Ognev's long-eared bat (*Plecotus ognevi*) is a species of bat found in Asia. By 2006, it was recognized as a separate species from the *P. auritus* species complex.

Ohiya rat
Ohiya rat
Srilankamys ohiensis

The Ohiya rat (*Srilankamys ohiensis*), or Sri Lanka bi-colored rat, is a species of rodent in the family Muridae. It is the only species in the genus *Srilankamys*. It is found only in Sri Lanka where it is known locally as මී කෑ දෙපැහැ මීයා in...

Okapi
Okapi
Okapia johnstoni

The okapi (*Okapia johnstoni*) is an artiodactyl mammal that is found only in central Africa. It was not discovered until 1901. *Okapia johnstoni*, its taxonomic name, honors its native Central African name, as well as the man who 'discovered' it,...

Okinawa flying fox

Okinawa flying fox

Pteropus loochoensis

The Okinawa flying fox (*Pteropus loochoensis*) is a species of megabat in the genus *Pteropus*. It is endemic to possibly Japan. It was previously listed as extinct by the IUCN, but because the two known specimens are taxonomically uncertain and of...

Oku mouse shrew

Oku mouse shrew

Myosorex okuensis

The Oku mouse shrew (*Myosorex okuensis*) is a species of mammal in the family Soricidae endemic to Cameroon. Its natural habitat is subtropical or tropical moist montane forests. It is threatened by habitat loss.

Oldfield mouse

Oldfield mouse

Peromyscus polionotus

The oldfield mouse or beach mouse (*Peromyscus polionotus*) is a nocturnal species of rodent in the family Cricetidae and primarily eats seeds. It lives in holes throughout the southeastern United States in beaches and sandy fields. Predators to...

Oldfield white-bellied rat

Oldfield white-bellied rat

Niviventer culturatus

The Oldfield white-bellied rat or soft-furred Taiwan niviventer (*Niviventer culturatus*) is a species of phat rat in the family Muridae. It is found only in Taiwan. It has also been considered a subspecies of *Niviventer niviventer* and included in...

Oligoryzomys andinus

Oligoryzomys andinus

Oligoryzomys andinus

Oligoryzomys andinus, also known as the Andean colilargo or Andean pygmy rice rat, is a species of rodent in genus *Oligoryzomys* of family Cricetidae. It is found in the Andes of southern Peru and western Bolivia, but may in fact include more than...

Oligoryzomys arenalis

Oligoryzomys arenalis

Oligoryzomys arenalis

Oligoryzomys arenalis, also known as the sandy colilargo or sandy pygmy rice rat, is a species of rodent in the genus *Oligoryzomys* of family Cricetidae. It is found in the Andes of Peru at 400 to 2850 m altitude, but may include more than one...

Oligoryzomys brendae

Oligoryzomys brendae

Oligoryzomys brendae

Oligoryzomys brendae, also known as Brenda's colilargo, is a South American species of rodent in the genus *Oligoryzomys*. It is found only in Tucumán Province in northwestern Argentina, but its taxonomic status requires revision.

Oligoryzomys chacoensis

Oligoryzomys chacoensis

Oligoryzomys chacoensis

Oligoryzomys chacoensis, also known as the Chacoan colilargo or Chacoan pygmy rice rat, is a rodent species from South America. It is found in the Gran Chaco region of southeastern Bolivia, southwestern Brazil, Paraguay, and northeastern...

Oligoryzomys destructor
Oligoryzomys destructor
Oligoryzomys destructor

Oligoryzomys destructor, also known as Tschudi's colilargo or the destructive pygmy rice rat, is a species of rodent in the genus *Oligoryzomys* of family Cricetidae. It is found along the eastern Andes from southern Colombia, through Ecuador,...

Oligoryzomys flavescens
Oligoryzomys flavescens
Oligoryzomys flavescens

Oligoryzomys flavescens, also known as the flavescens colilargo or yellow pygmy rice rat is a species of rodent in the genus *Oligoryzomys* of family Cricetidae. It is found in southern South America, occurring in southern Brazil, Paraguay,...

Oligoryzomys fornesi
Oligoryzomys fornesi
Oligoryzomys fornesi

Oligoryzomys fornesi, also known as Fornes' colilargo, is a species of rodent in the genus *Oligoryzomys* of family Cricetidae. It is found from northeastern Brazil via Paraguay into northeastern Argentina. Its karyotype has $2n = 62$ and $FNa = 64$.

Oligoryzomys fulvescens
Oligoryzomys fulvescens
Oligoryzomys fulvescens

Oligoryzomys fulvescens, also known as the fulvous colilargo, fulvous pygmy rice rat, or northern pygmy rice rat, is a species of rodent in the genus *Oligoryzomys* of family Cricetidae. It is found from southern Mexico through Central America into...

Oligoryzomys griseolus
Oligoryzomys griseolus
Oligoryzomys griseolus

Oligoryzomys griseolus, also known as the grizzled colilargo or the grayish pygmy rice rat, is a species of rodent in the genus *Oligoryzomys* of family Cricetidae. It is found in the Andes of Venezuela and nearby Colombia. Its karyotype has $2n = \dots$

Oligoryzomys longicaudatus
Oligoryzomys longicaudatus
Oligoryzomys longicaudatus

Oligoryzomys longicaudatus, also known as the long-tailed colilargo or long-tailed pygmy rice rat, is a species of rodent in the genus *Oligoryzomys* of the family Cricetidae. It is found in the southern Andes of Chile and Argentina, with an...

Oligoryzomys magellanicus
Oligoryzomys magellanicus
Oligoryzomys magellanicus

Oligoryzomys magellanicus, also known as the Patagonian colilargo and the Magellanic pygmy rice rat, is a species of rodent in the genus *Oligoryzomys* of the family Cricetidae. It is found in the southernmost parts of Argentina and Chile,...

Oligoryzomys microtis
Oligoryzomys microtis
Oligoryzomys microtis

Oligoryzomys microtis, also known as the small-eared colilargo or small-eared pygmy rice rat, is a species of rodent in the genus *Oligoryzomys* of family Cricetidae. It

is found in western Brazil, eastern Peru, Bolivia, and northern Paraguay.

Oligoryzomys moojeni
Oligoryzomys moojeni
Oligoryzomys moojeni

Oligoryzomys moojeni is a species of rodent from South America in the genus Oligoryzomys of family Cricetidae. It is known only from central Brazil, where it is found in the cerrado and gallery forests. It is named after twentieth-century...

Oligoryzomys nigripes
Oligoryzomys nigripes
Oligoryzomys nigripes

Oligoryzomys nigripes, also known as the black-footed colilargo or the black-footed pygmy rice rat, is a rodent in the genus Oligoryzomys of family Cricetidae.

Oligoryzomys nigripes is a species that has been further divided into different sister...

Oligoryzomys rupestris
Oligoryzomys rupestris
Oligoryzomys rupestris

Oligoryzomys rupestris is a species of rodent in the genus Oligoryzomys of family Cricetidae. It is known only from eastern Brazil, where it has been found in several localities in the campos rupestres montane savanna ecoregion. This is a small...

Oligoryzomys stramineus
Oligoryzomys stramineus
Oligoryzomys stramineus

Oligoryzomys stramineus, also known as the straw-colored colilargo or straw-colored pygmy rice rat, is a species of rodent in the genus Oligoryzomys of the family Cricetidae. It occurs only in the cerrado and caatinga ecoregions of northeastern...

Oligoryzomys vegetus
Oligoryzomys vegetus
Oligoryzomys vegetus

Oligoryzomys vegetus, also known as the sprightly colilargo or sprightly pygmy rice rat, is a species of rodent in the genus Oligoryzomys of family Cricetidae. It is found only in the mountains of Costa Rica and western Panama.

Olinguito
Olinguito
Bassaricyon neblina

The olinguito (Bassaricyon neblina) is a mammal of the raccoon family Procyonidae that lives in montane forests in the Andes of western Colombia and Ecuador. It was classified as belonging to a new species in 2013. The specific name neblina is...

Olive Baboon
Olive Baboon
Papio anubis

The Olive baboon (Papio anubis) is a member of the Old World monkeys. The species is the most wide-ranging of all baboons, being native to 25 countries throughout Africa.

Olive colobus
Olive colobus
Procolobus verus

The olive colobus monkey (Procolobus verus), also known as the green colobus or Van Beneden's colobus, is a species of primate in the family Cercopithecidae. Its

English name refers to its dull olive upperparts. It is the smallest example of all...

Olive montane mouse
Olive montane mouse
Aepeomys lugens

The olive montane mouse (*Aepeomys lugens*) is a species of rodent in the family Cricetidae. It is found in Ecuador and Venezuela.

Olive-backed pocket mouse
Olive-backed pocket mouse
Perognathus fasciatus

The olive-backed pocket mouse (*Perognathus fasciatus*) is a species of rodent in the family Heteromyidae. It is found in the central Great Plains of Canada and the United States where it is widespread and relatively common; the IUCN considers it...

Oliver's warty pig
Oliver's warty pig
Sus oliveri

Oliver's warty pig or Mindoro warty pig (*Sus oliveri*) is a small species in the pig genus (*Sus*) which can only be found on the island of Mindoro in the central Philippines. This species previously treated to be a subspecies of *S. philippensis*,...

Ollala brothers's titi
Ollala brothers's titi
Plecturocebus olallae

The Ollala brothers's titi (*Plecturocebus olallae*) is a species of titi, a type of New World monkey, endemic to Bolivia. There are between 110 and 150 individuals in the wild.

Olrog's chaco mouse
Olrog's chaco mouse
Andalgalomys olrogi

Olrog's chaco mouse (*Andalgalomys olrogi*) is a species of South American rodent in the family Cricetidae, endemic to Argentina. The natural habitat of the species is hot deserts. Its karyotype has $2n = 60$. The species is named after...

Olrog's four-eyed opossum
Olrog's four-eyed opossum
Philander olrogi

Olrog's four-eyed opossum (*Philander olrogi*) is a South American species of opossum found in eastern Bolivia, first described in 2008 based on specimens collected in 1974. It inhabits the lowland Amazon rainforest, with an elevation range of 150...

Olympic Marmot
Olympic Marmot
Marmota olympus

The Olympic marmot (*Marmota olympus*) is a rodent in the squirrel family, Sciuridae; it occurs only in the U.S. state of Washington, on the middle elevations of the Olympic Peninsula. The closest relatives of this species are the hoary marmot and...

Olympic shrew
Olympic shrew
Sorex rohweri

The Olympic shrew (*Sorex rohweri*) is a rare species of shrew that lives in only 13

spots in northwest Washington state and, a recent discovery, in Burns Bog, located in Delta, BC. Not much is known about this species. It was only described in...
Omilteme cottontail
Omilteme cottontail
Sylvilagus insonus

The Omilteme cottontail (*Sylvilagus insonus*) is a cottontail rabbit found only in the state of Guerrero, Mexico in the mountain range of Sierra Madre del Sur. Belonging to the family Leporidae, it is one of fourteen species in the genus...
Omura's Whale
Omura's Whale
Balaenoptera omurai

Omura's whale or the dwarf fin whale (*Balaenoptera omurai*) is a species of rorqual about which very little is known. Before its formal description, it was referred to as a small, dwarf or pygmy form of Bryde's whale by various sources. The...
Onager
Onager
Equus hemionus

The onager (*Equus hemionus*) is a species of the family Equidae that is found in Asia. The onager was described and given its binomial name by German zoologist Peter Simon Pallas in 1775. Unlike most horses and donkeys, onagers have never been...
Oncilla
Oncilla
Leopardus tigrinus

The oncilla (*Leopardus tigrinus*), also known as the northern tiger cat, little spotted cat, and tigrillo, is a small spotted cat ranging from Central America to central Brazil. It is listed as Vulnerable on the IUCN Red List, and the population...
One-striped opossum
One-striped opossum
Monodelphis unistriata

The one-striped opossum (*Monodelphis unistriata*) is a possibly extinct opossum species from South America. It is known only from two specimens found in Brazil in 1821 and Argentina in 1899.
One-toothed shrew mouse
One-toothed shrew mouse
Pseudohydromys ellermani

The one-toothed shrew mouse (*Pseudohydromys ellermani*) is a species of rodent in the family Muridae. It is found in West Papua, Indonesia and Papua New Guinea, at elevations from 1,200 to 3,000 m. It is most closely related to *Pseudohydromys*...
Ontong Java flying fox
Ontong Java flying fox
Pteropus howensis

The Ontong Java flying fox (*Pteropus howensis*) is a species of flying fox in the family Pteropodidae. It is endemic to the Ontong Java Atoll in the Solomon Islands. Its natural habitats are subtropical or tropical dry forests and subtropical or...
Ooldea dunnart
Ooldea dunnart
Sminthopsis ooldea

The Ooldea dunnart (*Sminthopsis ooldea*), also called Troughton's dunnart after the person who found the species, is an Australian marsupial similar to the hairy-

footed dunnart. It is greyish-yellow on its upper body and white on the underside...
Opdenbosch's mangabey
Opdenbosch's mangabey
Lophocebus opdenboschi

Opdenbosch's mangabey (*Lophocebus opdenboschi*) is a species of crested mangabey in the family Cercopithecidae. It has also been treated as a subspecies of *Lophocebus aterrimus*. It is found in the Democratic Republic of the Congo (formerly...
Opossum rat
Opossum rat
Rattus marmosurus

The opossum rat (*Rattus marmosurus*) is a species of rodent in the family Muridae. It is found only in northern and central Sulawesi, Indonesia.
Orange leaf-nosed bat
Orange leaf-nosed bat
Rhinonicteris aurantia

The orange leaf-nosed bat (*Rhinonicteris aurantia*) is a bat in the family Hipposideridae. It is the only living species in the genus *Rhinonicteris* which is endemic to Australia, occurring in the far north and north-west of the continent. They...
Orange nectar bat
Orange nectar bat
Lonchophylla robusta

The orange nectar bat (*Lonchophylla robusta*) is a species of bat in the family Phyllostomidae. It is found in Colombia, Costa Rica, Ecuador, Nicaragua, Panama, Peru, and Venezuela. Orange nectar bats in Costa Rica were observed utilising a unique...
Orange-bellied Himalayan squirrel
Orange-bellied Himalayan squirrel
Dremomys lokriah

The orange-bellied Himalayan squirrel (*Dremomys lokriah*) is a species of rodent in the family Sciuridae. It is found in Bangladesh, China, India, Myanmar, Nepal and Bhutan.
Orange-brown Atlantic tree-rat
Orange-brown Atlantic tree-rat
Phyllomys brasiliensis

The orange-brown Atlantic tree-rat or red-nosed tree-rat (*Phyllomys brasiliensis*), is a spiny rat species from South America. It is found in Brazil.
Orbiculus leaf-nosed bat
Orbiculus leaf-nosed bat
Hipposideros orbiculus

The orbiculus leaf-nosed bat (*Hipposideros orbiculus*), also known as the orbiculus roundleaf bat and small disc roundleaf bat, is a species of bat from the family Hipposideridae. The species has been found on the island of Sumatra in Indonesia...
Orcés's long-tongued bat
Orcés's long-tongued bat
Lonchophylla orcesi

Orcés's long-tongued bat (*Lonchophylla orcesi*) is a species of leaf-nosed bat found in Ecuador.
Ord's Kangaroo Rat
Ord's Kangaroo Rat

Dipodomys ordii

Ord's kangaroo rat (*Dipodomys ordii*) is a small rodent found only in western North America. Their common name comes from their bipedal form. They hop in a manner similar to the much larger kangaroo.

Oreoryzomys

Oreoryzomys

Oreoryzomys balneator

Oreoryzomys balneator, also known as the Peruvian rice rat or Ecuadoran oryzomys, is a species of rodent in the tribe Oryzomyini of family Cricetidae. It is found in Ecuador and northern Peru in cloud forest at elevations from 1500 to 1800 m. It...

Oribi

Oribi

Ourebia ourebi

The oribi (; *Ourebia ourebi*) is a small antelope found in eastern, southern and western Africa. The sole member of its genus, it was described by the German zoologist Eberhard August Wilhelm von Zimmermann in 1783. Eight subspecies are...

Oriental Basin pocket gopher

Oriental Basin pocket gopher

Cratogeomys fulvescens

The Oriental Basin pocket gopher (*Cratogeomys fulvescens*) is a species of pocket gopher which is endemic to Mexico. It was first described in 1895 by Clinton Hart Merriam. It was considered to be a subspecies of Merriam's pocket gopher...

Oriental shrew

Oriental shrew

Crocidura orientalis

The Oriental shrew (*Crocidura orientalis*) is a species of mammal in the family Soricidae. It is endemic to Indonesia. It is threatened by habitat loss.

Oriental Small-Clawed Otter

Oriental Small-Clawed Otter

Aonyx cinerea

The Oriental small-clawed otter (*Aonyx cinereus*), also known as the Asian small-clawed otter is an otter species native to South and Southeast Asia. It has short claws that do not extend beyond the pads of its webbed digits. It is the smallest...

Orinoco agouti

Orinoco agouti

Dasyprocta guamara

The Orinoco agouti (*Dasyprocta guamara*) is a species of rodent in the family Dasyproctidae. It is endemic to Delta Amacuro in Venezuela, where it is found in areas with rainforest or mangrove.

Orinoco sword-nosed bat

Orinoco sword-nosed bat

Lonchorhina orinocensis

The Orinoco sword-nosed bat (*Lonchorhina orinocensis*) is a species of bat in the family Phyllostomidae. It is found in Colombia and Venezuela.

Orizaba deer mouse

Orizaba deer mouse

Peromyscus beatae

The Orizaba deer mouse (*Peromyscus beatae*) is a small species of rodent in the family Cricetidae, native to El Salvador, Guatemala, Honduras, and Mexico. It is

found in thorn scrub and favors rocky areas with brush and scattered trees. They are...

Orizaba long-tailed shrew
Orizaba long-tailed shrew
Sorex orizabae

The Orizaba long-tailed shrew (*Sorex orizabae*) is a species of mammal in the family Soricidae. It is found in the states of Federal District, México, Michoacán, Morelos, Puebla, Tlaxcala, and Veracruz in Mexico. It is named after Pico de...

Ornate cuscus
Ornate cuscus
Phalanger ornatus

The ornate cuscus or Molluccan cuscus (*Phalanger ornatus*) is a species of marsupial in the family Phalangeridae. It is endemic to Indonesia, where it is found on the North Maluku islands of Halmahera, Bacan and Morotai, at elevations from sea...

Ornate flying fox
Ornate flying fox
Pteropus ornatus

The ornate flying fox (*Pteropus ornatus*) is a species of flying fox in the family Pteropodidae. It is endemic to New Caledonia. Its natural habitat is subtropical or tropical dry forests. It is threatened due to habitat destruction and hunting,...

Ornate Shrew
Ornate Shrew
Sorex ornatus

The Ornate shrew (*Sorex ornatus*) is a small mammal found only in western North America. These animals have long, pointed snouts, small ears, which are often not visible, and scent glands located on the sides of their bodies. As their eyesight is...

Ornate spider monkey
Ornate spider monkey
Ateles geoffroyi ornatus

The ornate spider monkey (*Ateles geoffroyi ornatus*), is a subspecies of Geoffroy's spider monkey, a type of New World monkey, from Central America, native to Costa Rica and Panama. Other common names for this subspecies include the brilliant...

Ornate titi
Ornate titi
Plecturocebus ornatus

The ornate titi (*Plecturocebus ornatus*) is a species of titi and is the smallest member of the family Pitheciidae, which also includes uakaris and saki monkeys. As it is a member of this family, it is classified as a type of New World monkey. It...

Orycteropus abundulafus
Orycteropus abundulafus
Orycteropus abundulafus

Orycteropus abundulafus is an extinct species of aardvark. Its fossil was found in northern Chad. It lived in the Mio-Pliocene epochs.

Oryzomys albiventer
Oryzomys albiventer
Oryzomys albiventer

Oryzomys albiventer is a rodent in the genus *Oryzomys* of family Cricetidae from interior western Mexico, in the states of Jalisco, Guanajuato, and Michoacán. First

described in 1901 as a separate species, it was later lumped under *O. couesi* and...
Oryzomys antillarum
Oryzomys antillarum
Oryzomys antillarum

Oryzomys antillarum, also known as the Jamaican rice rat, is an extinct rodent of Jamaica. A member of the genus *Oryzomys* within the family Cricetidae, it is similar to *O. couesi* of mainland Central America, from where it may have dispersed to...
Oryzomys couesi
Oryzomys couesi
Oryzomys couesi

Oryzomys couesi, also known as Coues's rice rat, is a semiaquatic rodent in the family Cricetidae occurring from southernmost Texas through Mexico and Central America into northwestern Colombia. It is usually found in wet habitats, such as...
Oryzomys dimidiatus
Oryzomys dimidiatus
Oryzomys dimidiatus

Oryzomys dimidiatus, also known as the Nicaraguan oryzomys, Thomas's rice rat, or the Nicaraguan rice rat, is a rodent in the family Cricetidae. It is known from only three specimens, all collected in southeastern Nicaragua since 1904. Placed in...
Oryzomys gorgasi
Oryzomys gorgasi
Oryzomys gorgasi

Oryzomys gorgasi, also known as Gorgas's oryzomys or Gorgas's rice rat, is a rodent in the genus *Oryzomys* of family Cricetidae. First recorded in 1967, it is known from only a few localities, including a freshwater swamp in the lowlands of...
Oryzomys nelsoni
Oryzomys nelsoni
Oryzomys nelson

Oryzomys nelsoni is an extinct rodent of María Madre Island, Nayarit, Mexico. Within the genus *Oryzomys* of the family Cricetidae, it may have been most closely related to the mainland species *O. albiventer*. Since its first description in 1898,...
Oryzomys peninsulae
Oryzomys peninsulae
Oryzomys peninsulae

Oryzomys peninsulae, also known as the Lower California rice rat, is a species of rodent from western Mexico. Restricted to the southern tip of the Baja California peninsula, it is a member of the genus *Oryzomys* of family Cricetidae. Only about...
Osgood's horseshoe bat
Osgood's horseshoe bat
Rhinolophus osgoodi

Osgood's horseshoe bat (*Rhinolophus osgoodi*) is a species of bat in the family Rhinolophidae. It is endemic to China. Its name refers to Wilfred Hudson Osgood.
Osgood's leaf-eared mouse
Osgood's leaf-eared mouse
Phyllotis osgoodi

Osgood's leaf-eared mouse (*Phyllotis osgoodi*) is a species of rodent in the family Cricetidae. It is found only in the Altiplano of northeastern Chile. The species is named after American zoologist Wilfred Hudson Osgood.
Osgood's mouse

Osgood's mouse
Peromyscus gratus

Osgood's mouse (*Peromyscus gratus*) is a species of rodent in the family Cricetidae. It is found only in Mexico. Its name references Wilfred Hudson Osgood.
Osgood's rat
Osgood's rat
Rattus osgoodi

Osgood's rat (*Rattus osgoodi*) is a species of rodent in the family Muridae. It is found only in Lâm Đồng Province in the Central Highlands (Tây Nguyên) of southern Vietnam. Its name references Wilfred Hudson Osgood.
Osgood's short-tailed opossum
Osgood's short-tailed opossum
Monodelphis osgoodi

Osgood's short-tailed opossum (*Monodelphis osgoodi*) is a species of opossum in the family Didelphidae. It is found in Bolivia and Peru. Its natural habitats are subtropical or tropical moist lowland forest and subtropical or tropical dry lowland...

Osman Hill's mangabey
Osman Hill's mangabey
Lophocebus osmani

The Osman Hill's mangabey (*Lophocebus osmani*), also known as the rusty-mantled mangabey, is a species of crested mangabey in the family Cercopithecidae with a restricted distribution in West Africa.

Otter Civet
Otter Civet
Cynogale bennettii

The Otter civet (*Cynogale bennettii*) is a semiaquatic viverrid native to Southeast Asia. It is believed to be undergoing severe population decline and is classified as an endangered species by the IUCN Red List.

Otto's sportive lemur
Otto's sportive lemur
Lepilemur otto

Otto's sportive lemur (*Lepilemur otto*), or the Ambodimahabibo sportive lemur, is a sportive lemur endemic to Madagascar. Its coloration is primarily grey-brown. It is known only from Ambodimahabibo, between the Mahajamba River in the west and...

Oubangui mouse
Oubangui mouse
Mus oubanguii

The Oubangui mouse (*Mus oubanguii*) is a species of rodent in the family Muridae. It is found only in the Central African Republic. Its natural habitat is dry savanna.

Oustalet's red colobus
Oustalet's red colobus
Piliocolobus oustaleti

Oustalet's red colobus (*Piliocolobus oustaleti*) is a species of red colobus monkey. It lives in various types of forest in southern South Sudan, southern Central African Republic, northern Democratic Republic of the Congo and northeastern...

Owl's spiny rat
Owl's spiny rat
Carterodon sulcidens

Owl's spiny rat, *Carterodon sulcidens*, is a rodent species from South America in the family Echimyidae. It is found in Brazil. It is the only species in the genus *Carterodon*. The Owl's spiny rat has evolved from a single species that has had...
Owston's palm civet
Owston's palm civet
Chrotogale owstoni

Owston's palm civet (*Chrotogale owstoni*) is a civet native to Vietnam, Laos and southern China. It is listed as Endangered by IUCN because of an ongoing population decline, estimated to be more than 50% over the last three generations, inferred...
Oxymycterus hucucha
Oxymycterus hucucha
Oxymycterus hucucha

Oxymycterus hucucha, also known as the Quechuan Hociudo, is a species of rodent in the genus *Oxymycterus* of family Cricetidae from South America. It is found only in a small region of the Andes in central Bolivia, where it lives in cloud forest...
Oyapock's fish-eating rat
Oyapock's fish-eating rat
Neusticomys oyapocki

Oyapock's fish-eating rat (*Neusticomys oyapocki*) is a species of rodent in the family Cricetidae. It is found in French Guiana and Brazil.
Ozark big-eared bat
Ozark big-eared bat
Corynorhinus townsendii ingens

The Ozark big-eared bat is an endangered species found only in a small number of caves in Arkansas, Oklahoma and Missouri, the southern central United States. Also known as the western big-eared bat, the long-eared bat, and the lump-nosed bat,...
Ozimops beccarii
Ozimops beccarii
Ozimops beccarii

Beccari's free-tailed bat (*Ozimops beccarii*) is a species of bat in the free-tailed bat family Molossidae found to Indonesia and Papua New Guinea. This bat can be found in several habitat types, including savanna, tropical moist forest, and...
Ozimops cobourgianus
Ozimops cobourgianus
Ozimops cobourgianus

Ozimops cobourgianus is a species of molossid bat, insectivorous flying mammals known as freetail bats, which are found in north and west coastal regions of Australia. First described in 1959, the group were later recognised as species...
Ozimops halli
Ozimops halli
Ozimops halli

Ozimops halli, also referred to as the Cape York free-tailed bat, is a species of molossid bat found at the Cape York Peninsula in Australia.
Ozimops kitcheneri
Ozimops kitcheneri
Ozimops kitcheneri

Ozimops kitcheneri, the south-western free-tailed bat, is a species of molossid bat found in Southwest Australia. A small flying mammal, it forages in forests and woodlands for insects.
Ozimops loriae
Ozimops loriae

Ozimops loriae

Ozimops loriae is a species of bat found in Australia and Papua New Guinea. The common names include Loria's mastiff bat and little northern free-tailed bat. The species was formerly described as Mormopterus loriae, and assigned to an Australian...

Ozimops lumsdenae

Ozimops lumsdenae

Ozimops lumsdenae

Ozimops lumsdenae is a species of molossid bat found in Australia, the largest of the genus Ozimops.

Ozimops petersi

Ozimops petersi

Ozimops petersi

Ozimops petersi, the inland free-tailed bat is a species of bat found in Australia. It is notable for being able to tolerate the most extreme body temperature range of any known mammal.

Ozimops planiceps

Ozimops planiceps

Ozimops planiceps

Ozimops planiceps is a small bat in the family Molossidae, native to Australia and Indonesia.

Ozimops ridei

Ozimops ridei

Ozimops ridei

Ozimops ridei is a species of molossid bat found in eastern Australia.

Paca

Paca

Cuniculus paca

The lowland paca (Cuniculus paca), also known as the spotted paca, is a large rodent found in tropical and sub-tropical America, from east-central Mexico to northern Argentina, and has been introduced to Cuba and Algeria. The animal is called...

Pacarana

Pacarana

Dinomys branickii

The pacarana (Dinomys branickii) is a rare and slow-moving hystricognath rodent indigenous to South America. Native Tupi people call it the pacarana (false paca) because it is superficially similar to the paca, a different rodent which is not in...

Pacific degu

Pacific degu

Octodon pacificus

The Pacific degu (Octodon pacificus), also known as the Mocha Island degu, is a species of rodent in the family Octodontidae. It is endemic to Mocha Island in Chile. Its natural habitat is subtropical or tropical moist lowland forests. It is...

Pacific jumping mouse

Pacific jumping mouse

Zapus trinotatus

The Pacific jumping mouse (Zapus trinotatus) is a species of rodent in the family

Zapodidae. It is found in Canada and the United States. Its natural habitats are temperate grassland and swamps.

Pacific pocket mouse

Pacific pocket mouse

Perognathus longimembris pacificus

The Pacific pocket mouse, *Perognathus longimembris pacificus*, is endemic to California. It lives in sandy coastal soils of the coastal sage scrub ecoregion. It eats seeds and some insects. It was believed to be extinct until 1993, when a small...

Pacific sheath-tailed bat

Pacific sheath-tailed bat

Emballonura semicaudata

The Pacific sheath-tailed bat or Polynesian sheath-tailed bat (*Emballonura semicaudata*) is a species of sac-winged bat in the family Emballonuridae found in American Samoa, Fiji, Guam, Micronesia, Palau, Samoa (where it is called pe'a vai,...

Pacific shrew

Pacific shrew

Sorex pacificus

The Pacific shrew (*Sorex pacificus*) is a species of mammal in the family Soricidae. It is endemic to western Oregon in the United States. The Pacific Shrew is native to western Oregon, more specifically from the Siltcoos lake to the coast going...

Pacific spiny rat

Pacific spiny rat

Proechimys decumanus

The Pacific spiny rat (*Proechimys decumanus*) is a species of rodent in the family Echimyidae. It is found in Ecuador and Peru.

Pacific White-Sided Dolphin

Pacific White-Sided Dolphin

Lagenorhynchus obliquidens

The Pacific white-sided dolphin (*Lagenorhynchus obliquidens*), also known as the hookfin porpoise, is a very active dolphin found in the cool or temperate waters of the North Pacific Ocean.

Pagai Island macaque

Pagai Island macaque

Macaca pagensis

The Pagai Island macaque (*Macaca pagensis*), also known as the Pagai macaque or Bokkoi, is an Old World monkey endemic to the Mentawai Islands off the west coast of Sumatra. It is listed as critically endangered on the IUCN Red List due to its...

Pagai spiny rat

Pagai spiny rat

Maxomys pagensis

The Pagai spiny rat (*Maxomys pagensis*) is a species of rodent in the family Muridae. It is found only in the Mentawai Islands of Indonesia, on South Pagai, North Pagai, Sipora, and Siberut islands.

Painted Bat

Painted Bat

Kerivoula picta

The Painted bat (*Kerivoula picta*) is a vesper bat found in Asia. It is also known

as "butterfly bat" (projapoti badur), "rongin chamchika" (coloured bat), or "komola-badami chamchika" (orange-brown bat) in Bengali.

Painted big-eared mouse

Painted big-eared mouse

Auliscomys pictus

The painted big-eared mouse (*Auliscomys pictus*) is a species of rodent in the family Cricetidae. It is found in Bolivia and Peru, and possibly Chile.

Painted ringtail possum

Painted ringtail possum

Pseudochirulus forbesi

The painted ringtail possum or moss-forest ringtail possum (*Pseudochirulus forbesi*) is a species of marsupial in the family Pseudocheiridae. It is found in Papua New Guinea.

Painted spiny pocket mouse

Painted spiny pocket mouse

Heteromys pictus

The painted spiny pocket mouse (*Heteromys pictus*) is a species of rodent in the family Heteromyidae. It is found in Mexico and the northern tip of Guatemala. It was formerly placed in the genus *Liomys*, which is now recognized to be paraphyletic...

Painted tree-rat

Painted tree-rat

Callistomys pictus

The painted tree-rat (*Callistomys pictus*) is a species of spiny rat from Brazil, restricted to north-eastern Bahia in eastern Brazil. It is the only species in the genus *Callistomys*.

Painted treeshrew

Painted treeshrew

Tupaia picta

The painted treeshrew (*Tupaia picta*) is a treeshrew species of the family Tupaiidae. The first specimen was described by Oldfield Thomas and was part of a zoological collection from northern Borneo obtained by the British Museum of Natural History.

Palaeomephitis

Palaeomephitis

Palaeomephitis steinheimensis

Palaeomephitis steinheimensis is an extinct species of mephitid of the Miocene epoch in Europe. It is the earliest known species of the family Mephitidae.

Palawan Bearded Pig

Palawan Bearded Pig

Sus ahoenobarbus

The Palawan bearded pig (*Sus ahoenobarbus*) is a pig species found only in Southeast Asia. It is the largest wild pig in the Philippines. Until recently, it was considered a subspecies of the Bornean bearded pig (*Sus barbatus*).

Palawan Binturong

Palawan Binturong

Arctictis binturong whitei

The Palawan binturong (*Arctictis binturong whitei*) is a subspecies of the binturong. This is a medium-sized feliform mammal native to the Philippines.

Palawan flying squirrel

Palawan flying squirrel

Hylopetes nigripes

The Palawan flying squirrel or tapilak (*Hylopetes nigripes*) is a species of rodent in the family Sciuridae. It is endemic to the Philippines. Its natural habitat is subtropical or tropical dry forests. Its population is decreasing as time goes on....

Palawan fruit bat

Palawan fruit bat

Acerodon leucotis

The Palawan fruit bat (*Acerodon leucotis*), also known as the Palawan flying fox, is a species of megabat found in forests of Palawan, Balabac and Busuanga in the Philippines. It is listed as vulnerable by the IUCN and is declining due to hunting...

Palawan montane squirrel

Palawan montane squirrel

Sundasciurus rabori

The Palawan montane squirrel (*Sundasciurus rabori*) is a species of rodent in the family Sciuridae. It is endemic to the Philippines. Its natural habitat is subtropical or tropical dry forests. It is threatened by habitat loss.

Palawan moss shrew

Palawan moss shrew

Palawanosorex muscorum

The Palawan moss shrew (*Palawanosorex muscorum*) is a species of shrew found on Mount Mantalingajan in the Philippines. The shrew is one of two other unique species only found on Mount Mantalingajan.

Palawan pencil-tailed tree mouse

Palawan pencil-tailed tree mouse

Chiropodomys calamianensis

The Palawan pencil-tailed tree mouse (*Chiropodomys calamianensis*) is a species of arboreal rodent in the family Muridae. It is known from lowland forest near sea level, coconut groves, and bamboo thickets. It is endemic to the Palawan Faunal...

Palawan shrew

Palawan shrew

Crocidura palawanensis

The Palawan shrew (*Crocidura palawanensis*) is a species of mammal in the family Soricidae. It is endemic to the Philippines and known from Palawan and Balabac Islands, from sea level to 1,300 m (4,300 ft) asl. It occurs in old growth and scrubby...

Palawan soft-furred mountain rat

Palawan soft-furred mountain rat

Palawanomys furvus

The Palawan soft-furred mountain rat (*Palawanomys furvus*) is a species of rodent in the family Muridae. It is the only species in the genus *Palawanomys*. It is found only in Palawan, Philippines, and has been recorded on Mount Mantalingajan.

Palawan spiny rat

Palawan spiny rat

Maxomys panglima

The Palawan spiny rat (*Maxomys panglima*) is a species of rodent in the family Muridae. It is found only in the Philippines.

Palawan stink badger

Palawan stink badger

Mydaus marchei

The Palawan stink badger (*Mydaus marchei*), pantot or tuldo in the Batak language, is a carnivorous of the western Philippines named for its resemblance to badgers, its powerful smell, and the largest island to which it is native, Palawan. Like...
Palawan treeshrew
Palawan treeshrew
Tupaia palawanensis

The Palawan treeshrew (*Tupaia palawanensis*) is a treeshrew species endemic to the Palawan Island, Philippines, where it occurs from sea level to an elevation of 1,400 m (4,600 ft). The population is considered steady. Formerly, it was considered...
Pale field rat
Pale field rat
Rattus tunneyi

The pale field rat (*Rattus tunneyi*) is a small rat endemic to Australia. It is a nocturnal and herbivorous rodent that resides throughout the day in shallow burrows made in loose sand. Once widespread, the range has become greatly reduced and it...
Pale fork-marked lemur
Pale fork-marked lemur
Phaner pallescens

The pale fork-marked lemur (*Phaner pallescens*), or western fork-marked lemur, is known from western Madagascar; south of the Fiherenana River to the region of Soalala. It is listed on CITES Appendix I.
Pale Fox
Pale Fox
Vulpes pallida

The pale fox (*Vulpes pallida*) is a species of fox found in the band of African Sahel from Senegal in the west to Sudan in the east. It is one of the least studied of all canid species, in part due to its remote habitat and its sandy coat that...
Pale gerbil
Pale gerbil
Gerbillus perpallidus

The pale gerbil (*Gerbillus perpallidus*) is endemic to Egypt and is distributed mainly in the northwestern part of the country. It is also known as the pallid gerbil. The pale gerbil has pale orange fur, with white underparts, white forelimbs and...
Pale gray shrew
Pale gray shrew
Crocidura pergrisea

The pale gray shrew (*Crocidura pergrisea*) is a species of mammal in the family Soricidae. It is endemic to Pakistan. It is threatened by habitat loss.
Pale kangaroo mouse
Pale kangaroo mouse
Microdipodops pallidus

The pale kangaroo mouse or Soda Spring Valley kangaroo mouse (*Microdipodops pallidus*) is a species of rodent in the family Heteromyidae. It is endemic to California and Nevada in the United States.
Pale leaf-eared mouse
Pale leaf-eared mouse
Graomys domorum

The pale leaf-eared mouse (*Graomys domorum*) is a species of rodent in the family

Cricetidae. It is found in Argentina and Bolivia.

Pale pygmy jerboa

Pale pygmy jerboa

Salpingotus pallidus

The pale pygmy jerboa or pallid pygmy jerboa (*Salpingotus pallidus*) is a species of rodent in the family Dipodidae endemic to Kazakhstan.

Pale shrew tenrec

Pale shrew tenrec

Microgale fotsifotsy

The pale shrew tenrec (*Microgale fotsifotsy*), also known as the pale-footed shrew tenrec is a species of mammal in the family Tenrecidae. It is endemic to Madagascar. Its natural habitats are subtropical or tropical moist lowland forest and...

Pale spear-nosed bat

Pale spear-nosed bat

Phyllostomus discolor

The pale spear-nosed bat (*Phyllostomus discolor*) is a species of phyllostomid bat from South and Central America.

Pale-faced bat

Pale-faced bat

Phylloderma stenops

The pale-faced bat (*Phylloderma stenops*) is a bat species from South and Central America.

Pale-Throated Three-Toed Sloth

Pale-Throated Three-Toed Sloth

Bradypus tridactylus

The pale-throated sloth (*Bradypus tridactylus*), occasionally known as the Ai, is a species of three-toed sloth that inhabits tropical rainforests in northern South America. It is similar in appearance to, and often confused with, the...

Pallas's Cat

Pallas's Cat

Otocolobus manul

The Pallas's cat (*Otocolobus manul*) is a small wild cat well camouflaged and adapted to the cold continental climate in its native range. It has rounded rather than vertical slit pupils, a unique feature among small cats. It inhabits rocky...

Pallas's long-tongued bat

Pallas's long-tongued bat

Glossophaga soricina

Pallas's long-tongued bat (*Glossophaga soricina*) is a South and Central American bat with a fast metabolism that feeds on nectar.

Pallas's pika

Pallas's pika

Ochotona pallasii

Pallas's pika (*Ochotona pallasii*), also known as the Mongolian pika, is a species of small mammals in the pika family, Ochotonidae. It is found mainly in the mountains of western Mongolia.

Pallas's Squirrel

Pallas's Squirrel

Callosciurus erythraeus

Pallas's squirrel (*Callosciurus erythraeus*) is a medium-sized rodent that lives in

forests of Asia. They live in trees, eat a wide range of food, and cache acorns in the autumn.

Pallas's tube-nosed bat

Pallas's tube-nosed bat

Nyctimene cephalotes

Pallas's tube-nosed bat (*Nyctimene cephalotes*), also known as the Torresian tube-nosed bat or northern tube-nosed bat, is a species of megabat in the *Nyctimene* genus found in Indonesia. Its range may extend to New Guinea, but sightings may be...

Pallid Atlantic Forest rat

Pallid Atlantic Forest rat

Delomys sublineatus

The pallid Atlantic Forest rat (*Delomys sublineatus*) is a rodent species from South America. It is found in Brazil.

Pallid Atlantic tree-rat

Pallid Atlantic tree-rat

Phyllomys lamarum

The pallid Atlantic tree-rat (*Phyllomys lamarum*) is a species of rodent in the family Echimyidae. It is a species of spiny rat. It is endemic to Brazil, where it is found from Paraiba to Minas Gerais.

Pallid Bat

Pallid Bat

Antrozous pallidus

The Pallid bat (*Antrozous pallidus*) is the sole species of its genus and is closely related to Van Gelder's bat (*Bauerus dubiaquercus*). The Pallid bat was first described in 1856. It is the state bat of California.

Pallid beach mouse

Pallid beach mouse

Peromyscus polionotus decoloratus

The pallid beach mouse or Ponce de Leon beach mouse (*Peromyscus polionotus decoloratus*) was known from two locations in Florida, Ponce Park, Volusia County and Bulow, Flagler County. No individuals have been seen since 1959. The average pallid...

Pallid large-footed myotis

Pallid large-footed myotis

Myotis macrotarsus

The pallid large-footed myotis or Philippine large-footed myotis (*Myotis macrotarsus*) is a species of vesper bat. It can be found in the following countries: Malaysia and Philippines. It is found in caves and arable land.

Palm rat

Palm rat

Rattus palmarum

The palm rat (*Rattus palmarum*) is a species of rodent in the family Muridae. It is found in the Nicobar Islands, on Car Nicobar and Great Nicobar islands. The palm rat's natural habitats are subtropical or tropical dry forest and subtropical or...

Palmer's chipmunk

Palmer's chipmunk

Neotamias palmeri

Palmer's chipmunk (*Neotamias palmeri*) is a species of rodent in the family Sciuridae, endemic to Nevada. Its natural habitat is temperate forests. It is threatened by habitat loss.

Pampas Cat
Pampas Cat
Leopardus colocola

The Pampas cat (*Leopardus colocola*) is a small wild cat native to South America. It is named after the Pampas, fertile South American low grasslands. It is listed as Near Threatened on the IUCN Red List as the loss of its native habitat may cause...

Pampas Deer
Pampas Deer
Ozotoceros bezoarticus

The Pampas deer (*Ozotoceros bezoarticus*) is a species of deer that live in the grasslands of South America at low elevations. They are known as veado-campeiro in Portuguese and as venado or gama in Spanish. Their habitat includes water and hills,...

Pampas Fox
Pampas Fox
Lycalopex gymnocercus

The Pampas fox (*Lycalopex gymnocercus*), also known as grey pampean fox, Pampas zorro, Azara's fox, or Azara's zorro (in Spanish also called aguará chaí, anglicized as aguarachay, in Portuguese also called graxaim), is a medium-sized zorro, or...

Panama slender opossum
Panama slender opossum
Marmosops invictus

The Panama slender opossum (*Marmosops invictus*), also known as the slaty slender mouse opossum, is a species of opossum in the family Didelphidae. It is endemic to Panama, where it has been found in tropical rainforest habitats, including...

Panamanian climbing rat
Panamanian climbing rat
Tylomys panamensis

The Panamanian climbing rat (*Tylomys panamensis*) is a species of rodent in the family Cricetidae. It is endemic to Panama.

Panamanian Night Monkey
Panamanian Night Monkey
Aotus zonalis

The Panamanian night monkey or Chocoan night monkey (*Aotus zonalis*) is a species of night monkey formerly considered a subspecies of the gray-bellied night monkey of the family Aotidae. Its range consists of Panama and the Chocó region of...

Panamanian spiny pocket mouse
Panamanian spiny pocket mouse
Heteromys adspersus

The Panamanian spiny pocket mouse (*Heteromys adspersus*), also known as Peter's spiny pocket mouse, is a species of heteromyid rodent endemic to Panama. It is very closely related to Salvin's spiny pocket mouse, and has been placed in the same...

Panamanian White-Faced Capuchin
Panamanian White-Faced Capuchin
Cebus imitator

The Panamanian white-faced capuchin (*Cebus imitator*) is a medium-sized New World monkey native to the forests of Central America. It is a highly intelligent monkey and has been trained to assist paraplegic persons. It is noted for its tool use,...

Panamint chipmunk
Panamint chipmunk

Neotamias panamintinus

The Panamint chipmunk (*Neotamias panamintinus*) is a species of rodent in the squirrel family, *Sciuridae*. It is endemic to desert mountain areas of southeast California and southwest Nevada in the United States. It is considered a species of least...

Panamint kangaroo rat

Panamint kangaroo rat

Dipodomys panamintinus

The Panamint kangaroo rat (*Dipodomys panamintinus*) is a species of rodent in the family *Heteromyidae*. It is endemic to the Mojave Desert in eastern California and western Nevada, in the United States.

Panay cloudbunner

Panay cloudbunner

Crateromys heaneyi

The Panay cloudbunner (*Crateromys heaneyi*) is the second-largest cloud rat, a squirrel-like rodent that is found on the island of Panay in the Philippines. It is the most endangered rodent species in Panay, and one of the only few known cloud...

Panay shrew

Panay shrew

Crocidura panayensis

The Panay shrew (*Crocidura panayensis*) is a species of shrew from the Philippines.

Panniet naked-backed fruit bat

Panniet naked-backed fruit bat

Dobsonia pannietensis

The Panniet naked-backed fruit bat (*Dobsonia pannietensis*), also known as the De Vis's Bare-backed Fruit Bat and Panaeati Bare-backed Fruit Bat, is a species of megabat in the family *Pteropodidae*. It roosts in groups, within caves and tree hollows.

Pantanal Cat

Pantanal Cat

Leopardus colocola braccatus

The Pantanal cat (*Leopardus colocola braccatus*) is a Pampas cat subspecies, a small wild cat native to South America. It is named after the Pantanal wetlands in central South America

Panthera leo fossilis

Panthera leo fossilis

Panthera leo fossilis

Panthera leo fossilis is a fossil cat of the genus *Panthera*, which was first excavated near Mauer in Germany, and lived during the Upper Pleistocene. Bone fragments of *P. l. fossilis* were also excavated near Pakefield in the United Kingdom, which...

Panthera onca augusta

Panthera onca augusta

Panthera onca augusta

Panthera onca augusta, commonly known as the Pleistocene jaguar or simply the giant jaguar, is an extinct subspecies of the jaguar that was endemic to North America during the Pleistocene epoch (1.8 mya–11,000 years ago).

Panthera pardus spelaea

Panthera pardus spelaea

Panthera pardus spelaea

Panthera pardus spelaea, sometimes called the European Ice Age leopard or Late Pleistocene leopard, is a fossil leopard subspecies, which roamed Europe in the Late Pleistocene. The youngest known bone fragments date to about 32,000 to 26,000...

Panthera tigris acutidens
Panthera tigris acutidens
Panthera tigris acutidens

Panthera tigris acutidens or Wanhshien tiger is an extinct tiger subspecies, which was scientifically described in 1928 based on fossils excavated near Wanhshien in southern China's Sichuan Province. Otto Zdansky named it *Felis acutidens*. After the...

Panthera tigris soloensis
Panthera tigris soloensis
Panthera tigris soloensis

Panthera tigris soloensis, known as the Ngandong tiger, is an extinct subspecies of the modern tiger species. It inhabited the Sundaland region of Indonesia during the Pleistocene epoch.

Panthera tigris trinilensis
Panthera tigris trinilensis
Panthera tigris trinilensis

Panthera tigris trinilensis, known as the Trinil tiger, is an extinct tiger subspecies dating from about 1.2 million years ago that was found at the locality of Trinil, Java, Indonesia. The fossil remains are now stored in the Dubois Collection...

Panthera youngi
Panthera youngi
Panthera youngi

Panthera youngi is a fossil cat species that was described in 1934; fossil remains of this cat were excavated in a *Sinanthropus* formation in Choukoutien, northeastern China. Upper and lower jaws excavated in Japan's Yamaguchi Prefecture were also...

Pantropical Spotted Dolphin
Pantropical Spotted Dolphin
Stenella attenuata

The pantropical spotted dolphin (*Stenella attenuata*) is a species of dolphin found in all the world's temperate and tropical oceans. The species was beginning to come under threat due to the killing of millions of individuals in tuna purse...

Papillose woolly bat
Papillose woolly bat
Kerivoula papillosa

The papillose woolly bat (*Kerivoula papillosa*) is a species of vesper bat in the family *Vespertilionidae*. It is found in Brunei, India, Indonesia, Malaysia, and Vietnam.

Papua grassland mosaic-tailed rat
Papua grassland mosaic-tailed rat
Melomys lutillus

The Papua grassland mosaic-tailed rat (*Melomys lutillus*), also known as the grassland melomys, is a species of rodent in the family *Muridae*. It is endemic to the island of New Guinea where it is present from sea level to altitudes of about 2,200...

Papuan bandicoot
Papuan bandicoot
Microperoryctes papuensis

The Papuan bandicoot (*Microperoryctes papuensis*) is a species of marsupial in the family Peramelidae. It is endemic to Papua New Guinea. Its natural habitat is subtropical or tropical dry forests. *M. papuensis* is a small bandicoot with a soft...

Papuan sheath-tailed bat
Papuan sheath-tailed bat
Saccolaimus mixtus

The Papuan sheath-tailed bat (*Saccolaimus mixtus*) is a species of bat in the family Emballonuridae which occurs at the Cape York Peninsula and New Guinea. The poorly known species hunts in open forests for night flying insects.

Para dog-faced bat
Para dog-faced bat
Cynomops paranus

The Para dog-faced bat (*Cynomops paranus*), also called the brown dog-faced bat, is a South American bat species of the family Molossidae. It is found in Panama, Colombia, Ecuador, Peru, Venezuela, Guyana, Surinam, French Guiana, Brazil, and...

Pará spiny tree-rat
Pará spiny tree-rat
Mesomys stimulax

The Pará spiny tree rat, *Mesomys stimulax*, or Surinam spiny tree rat, is a spiny rat species from South America. It is found in Brazil.

Paradox vole
Paradox vole
Microtus paradoxus

The paradox vole (*Microtus paradoxus*) is a species of rodent in the family Cricetidae found in southern Turkmenistan.

Paraguaian hairy dwarf porcupine
Paraguaian hairy dwarf porcupine
Coendou spinosus

The Paraguaian hairy dwarf porcupine, *Coendou spinosus*, is a South American porcupine species from the family Erethizontidae. It is found in Argentina, Brazil, Paraguay and Uruguay. They have a short tail and gray brown quills and feed on fruits,...

Paraguana moustached bat
Paraguana moustached bat
Pteronotus paraguayanensis

The Paraguana moustached bat (*Pteronotus paraguayanensis*) occurs only on the Paraguaná Peninsula of Venezuela. The entire population uses three caves, one of which is subject to human vandalism. Their total range is less than 400 km² (150 sq mi)....

Paraguaná spiny pocket mouse
Paraguaná spiny pocket mouse
Heteromys oasicus

The Paraguaná spiny pocket mouse (*Heteromys oasicus*) is a South American species of rodent in the family Heteromyidae. It is known from two localities at elevations above 200 m, Cerro Santa Ana and the Fila de Monte Cano, within the Paraguaná...

Paraguayan bolo mouse
Paraguayan bolo mouse
Necromys lenguarum

The Paraguayan bolo mouse or Paraguayan akodont (*Necromys lenguarum*) is a species

of rodent in the family Cricetidae. According to the IUCN, it is present in Bolivia, Paraguay, and Peru, and possibly also in Argentina and Brazil. It is found at...

Paraguayan fat-tailed mouse opossum

Paraguayan fat-tailed mouse opossum

Thylamys macrurus

The Paraguayan fat-tailed mouse opossum (*Thylamys macrurus*) is a species of opossum in the family Didelphidae. It is found in forested areas of Brazil and Paraguay. It is known only from a few specimens. For two listed specimens, one had a...

Paraguayan punaré

Paraguayan punaré

Thrichomys pachyurus

The Paraguayan punaré, *Thrichomys pachyurus*, is a caviomorph rodent of South America from the spiny rat family. With its skull averaging 55 mm long, it is the largest species in the genus *Thrichomys*. It is found in savannas and forest edges in...

Paramo hocicudo

Paramo hocicudo

Oxymycterus paramensis

The Paramo hocicudo (*Oxymycterus paramensis*) is a species of rodent in the family Cricetidae. It is found in Argentina, Bolivia, and Peru. The Argentine hocicudo is sometimes considered conspecific.

Paramo Oldfield mouse

Paramo Oldfield mouse

Thomasomys paramorum

The paramo Oldfield mouse (*Thomasomys paramorum*) is a species of rodent in the family Cricetidae. It is found only in Ecuador.

Paramushir shrew

Paramushir shrew

Sorex leucogaster

The Paramushir shrew (*Sorex leucogaster*) is a species of mammal in the family Soricidae. It is endemic to Russia. Its natural habitat is temperate forests. It is threatened by habitat loss.

Paraná grass mouse

Paraná grass mouse

Akodon paranaensis

The Paraná grass mouse (*Akodon paranaensis*) is a South American rodent species of the family Cricetidae. It is found in northeastern Argentina and southeastern Brazil.

Paratriaenops furculus

Paratriaenops furculus

Paratriaenops furculus

Paratriaenops furculus, also known as Trouessart's trident bat, is a species of bat in the family Hipposideridae. It is endemic to Madagascar. It was formerly assigned to the genus *Triaenops*, but is now placed in the separate genus *Paratriaenops*....

Paratriaenops pauliani

Paratriaenops pauliani

Paratriaenops pauliani

Paratriaenops pauliani is a species of bat in the family Hipposideridae. It is endemic to Aldabra Atoll of the western Seychelles, where it was found on Picard

Island. It was formerly considered to be part of the species *Triaenops furculus*, known...

Pardine genet

Pardine genet

Genetta pardina

The pardine genet (*Genetta pardina*), also known as the West African large spotted genet, is a genet species living in West Africa. As it is widely distributed and common, it is listed as Least Concern on the IUCN Red List.

Pariente's fork-marked lemur

Pariente's fork-marked lemur

Phaner parienti

Pariente's fork-marked lemur (*Phaner parienti*), or the Sambirano fork-marked lemur is a species of lemur endemic to the Sambirano region of north-western Madagascar.

This lemur has light brown to gray upperparts, a prominent facial fork and...

Parissi's slit-faced bat

Parissi's slit-faced bat

Nycteris parisii

Parissi's slit-faced bat (*Nycteris parisii*) is a species of slit-faced bat that may live in dry savanna regions of East Africa, near the Duba Valley and Benue River along with large rivers. It is only known from three specimens. One was found in...

Parma Wallaby

Parma Wallaby

Macropus parma

The parma wallaby (*Notamacropus parma*) is the smallest member of the marsupial genus *Notamacropus*. It inhabits wet sclerophyll (hard-leaved) forests of northern New South Wales, Australia. Adult parma wallabies typically weigh 4–5 kg, with the...

Parnell's mustached bat

Parnell's mustached bat

Pteronotus parnellii

Parnell's mustached bat (*Pteronotus parnellii*) is an insectivorous bat native to North, Central, the Caribbean, and South America. This bat species ranges from southern Sonora, Mexico, south to Brazil. It has a wider historical range; fossil...

Parti-coloured bat

Parti-coloured bat

Vespertilio murinus

The parti-coloured bat or rearmouse (*Vespertilio murinus*) is a species of vesper bat that lives in temperate Eurasia, from Western and Southern Europe, eastwards over the Caucasus and Iran into Mongolia, north-east China, Korea, Afghanistan and...

Particolored flying squirrel

Particolored flying squirrel

Hylopetes alboniger

The particolored flying squirrel (*Hylopetes alboniger*) is a species of rodent in the family *Sciuridae*. It is found in Bangladesh, Bhutan, Cambodia, China, India, Laos, Myanmar, Nepal, Thailand, and Vietnam. Its natural habitat is subtropical or...

Patagonian bonneted bat

Patagonian bonneted bat

Eumops patagonicus

The Patagonian bonneted bat (*Eumops patagonicus*), also called the Patagonian dwarf bonneted bat, is a species of free-tailed bat found in Argentina, Bolivia and Paraguay.

Patagonian chinchilla mouse
Patagonian chinchilla mouse
Euneomys chinchilloides

The Patagonian chinchilla mouse (*Euneomys chinchilloides*) is a species of rodent in the family Cricetidae. It was first described by George Robert Waterhouse in 1839. It is found in Tierra del Fuego and neighboring areas of southernmost...

Patagonian Mara
Patagonian Mara
Dolichotis patagonum

The Patagonian mara (*Dolichotis patagonum*) is a relatively large rodent found in open and semiopen habitats of South America. A population of this species has also been recorded in the northern United Arab Emirates, possibly as a result of...

Patagonian opossum
Patagonian opossum
Lestodelphys halli

The Patagonian opossum (*Lestodelphys halli*) is the sole species in genus *Lestodelphys*.

Patagonian weasel
Patagonian weasel
Lyncodon patagonicus

The Patagonian weasel (*Lyncodon patagonicus*) is a small mustelid that is the only member of the genus *Lyncodon*. Its geographic range is the Pampas of western Argentina and sections of Chile. An early mention of the animal is in the Journal of...

Patas Monkey
Patas Monkey
Erythrocebus patas

The common patas monkey (*Erythrocebus patas*), also known as the wadi monkey or hussar monkey, is a ground-dwelling monkey distributed over semi-arid areas of West Africa, and into East Africa.

Patricia's disk-winged bat
Patricia's disk-winged bat
Thyroptera wynneae

Patricia's disk-winged bat (*Thyroptera wynneae*) is a species of disk-winged bat found in South America.

Patrizi's trident leaf-nosed bat
Patrizi's trident leaf-nosed bat
Asellia patrizii

Patrizi's trident leaf-nosed bat (*Asellia patrizii*) is a species of bat in the family Hipposideridae. It is found in Eritrea, Ethiopia, and Saudi Arabia. Its natural habitats are subtropical or tropical dry shrubland and caves. It is threatened...

Patton's spiny rat
Patton's spiny rat
Proechimys pattoni

Patton's spiny-rat, *Proechimys pattoni*, is a spiny rat species from South America. It is found in Brazil and Peru.

Paucident planigale
Paucident planigale
Planigale gilesi

The paucident planigale (*Planigale gilesi*), also known as Giles' planigale, is a very small species of carnivorous marsupial of the family Dasyuridae.

Paucidentomys
Paucidentomys
Paucidentomys vermidax

Paucidentomys is a genus of rodents of a type commonly known as shrew-rats which was discovered in 2012 in a remote rainforest on the Indonesian island, Sulawesi. The genus is monotypic, consisting of the species *Paucidentomys vermidax*, or the...

Pavel's Seram mosaic-tailed rat
Pavel's Seram mosaic-tailed rat
Melomys paveli

The Pavel's Seram mosaic-tailed rat (*Melomys paveli*) is a species of rodent in the family Muridae. It is found only on the south coast of the island of Seram in Indonesia. At one time it was thought to be a subspecies of the black-tailed...

Peale's Dolphin
Peale's Dolphin
Lagenorhynchus australis

Peale's dolphin (*Lagenorhynchus australis*) is a small dolphin found in the waters around Tierra del Fuego at the foot of South America. It is also commonly known as the black-chinned dolphin or even Peale's black-chinned dolphin. However, since...

Peale's free-tailed bat
Peale's free-tailed bat
Nyctinomops aurispinosus

Peale's free-tailed bat (*Nyctinomops aurispinosus*) is a bat species from South and Central America.

Pearson's chaco mouse
Pearson's chaco mouse
Andalgalomys pearsoni

Pearson's chaco mouse (*Andalgalomys pearsoni*) is a species of rodent in the family Cricetidae. This mouse is found in the Gran Chaco ecoregion of southeastern Bolivia and western Paraguay at elevations up to 400 m. The species is named after...

Pearson's horseshoe bat
Pearson's horseshoe bat
Rhinolophus pearsonii

Pearson's horseshoe bat (*Rhinolophus pearsonii*) is a species of bat in the family Rhinolophidae. It is found in Bangladesh, Bhutan, China, India, Laos, Malaysia, Myanmar, Nepal, Thailand and Vietnam. The species is named after John Thomas

Pearson....
Pearson's long-clawed shrew
Pearson's long-clawed shrew
Solisorex pearsoni

Pearson's long-clawed shrew (*Solisorex pearsoni*) is a species of mammal in the family Soricidae. It is monotypic within the genus *Solisorex*. It is endemic to Sri Lanka. Its natural habitats are subtropical or tropical dry forests and subtropical...

Pearson's tuco-tuco
Pearson's tuco-tuco
Ctenomys pearsoni

Pearson's tuco-tuco (*Ctenomys pearsoni*) is a species of rodent in the family Ctenomyidae. It is endemic to Uruguay, where it is found at elevations below 200 m. This tuco-tuco constructs burrows with multiple openings (an average of 13)...

Peary Caribou

Peary Caribou

Rangifer tarandus pearyi

The Peary caribou (*Rangifer tarandus pearyi*) is a subspecies of caribou found in the High Arctic islands of Nunavut and the Northwest Territories in Canada. Peary caribou were named after the American explorer Robert Peary. They are called tuktu...

Pel's flying squirrel

Pel's flying squirrel

Anomalurus pelii

Pel's flying squirrel or Pel's scaly-tailed squirrel (*Anomalurus pelii*) is a species of rodent in the family Anomaluridae. It is found in Liberia, Ivory Coast, and Ghana, where it lives in lowland tropical rainforests.

Pel's pouched bat

Pel's pouched bat

Saccolaimus peli

Pel's pouched bat (*Saccolaimus peli*) is a species of sac-winged bat in the family Emballonuridae. It is found in Angola, Cameroon, Republic of the Congo, Democratic Republic of the Congo, Ivory Coast, Equatorial Guinea, Gabon, Ghana, Guinea,...

Peleng leaf-nosed bat

Peleng leaf-nosed bat

Hipposideros pelingensis

The Peleng leaf-nosed bat (*Hipposideros pelingensis*) is a species of bat native to Sulawesi and other adjacent Indonesian islands. It has been recorded in Marus National Park and Lambu Sango National Reserve.

Peleng rat

Peleng rat

Rattus pelurus

The Peleng rat (*Rattus pelurus*) is a species of rodent in the family Muridae. It is found only on Peleng Island off the southeastern coast of Sulawesi, Indonesia.

Peleng tarsier

Peleng tarsier

Tarsius pelengensis

The Peleng tarsier (*Tarsius pelengensis*), or the Peleng Island tarsier, is a nocturnal primate found on the island of Peleng, just east of Sulawesi, Indonesia.

Pelew flying fox

Pelew flying fox

Pteropus pelewensis

The Pelew flying fox (*Pteropus pelewensis*) is a species of megabat in the genus *Pteropus* found in the Palau Islands. A subspecies found on Yap, the Yap flying fox, is considered as a separate species by some authorities. The species is listed as...

Pemba flying fox

Pemba flying fox

Pteropus voeltzkowi

The Pemba flying fox (*Pteropus voeltzkowi*) is a species of flying fox in the family Pteropodidae. It is endemic to the island of Pemba on the coast of Tanzania.

Pen-Tailed Treeshrew
Pen-Tailed Treeshrew
Ptilocercus lowii

The Pen-tailed treeshrew (*Ptilocercus lowii*) is a small mammal native to Asia. These animals are considered very close relatives of primates, with the colugos being closer to primates.

Peninsular horseshoe bat
Peninsular horseshoe bat
Rhinolophus robinsoni

The Peninsular horseshoe bat (*Rhinolophus robinsoni*) is a species of horseshoe bat found in Malaysia and Thailand.

Peninsular myotis
Peninsular myotis
Myotis peninsularis

The peninsular myotis (*Myotis peninsularis*) is a species of vesper bat. It is endemic to northwestern Mexico, found only within Baja California Sur state on the southern Baja California Peninsula. Its habitats include the southern Peninsular...

Peninsular shrew
Peninsular shrew
Crocidura negligens

The peninsular shrew (*Crocidura negligens*) is a species of mammal in the family Soricidae. It is found in the Malay Peninsula and Thailand (south of the Kra Isthmus). It is also found on the adjacent islands of Koh Samui, Pulau Tioman, and Pulau...

Pennant's colobus
Pennant's colobus
Procolobus pennantii

Pennant's colobus or Pennant's red colobus (*Piliocolobus pennantii*) is a species of tree-dwelling primate in the family Cercopithecidae. It is endemic to tropical Central Africa. Three subspecies have traditionally been recognised but its...

Peracchi's nectar bat
Peracchi's nectar bat
Lonchophylla peracchii

Peracchi's nectar bat (*Lonchophylla peracchii*) is a species of nectar-feeding bat in the family Phyllostomidae. It was first described from the Atlantic Forest in southeastern Brazil.

Perameles allinghamensis
Perameles allinghamensis
Perameles allinghamensis

Perameles allinghamensis, the Bluff Downs bandicoot, is a small extinct bandicoot that lived in Australia 4 million years ago in the Pliocene period. It was discovered at the Bluff Downs fossil site in northern Queensland. Its diet probably...

Perameles bowensis
Perameles bowensis
Perameles bowensis

Perameles bowensis is an extinct Pliocene-aged species of bandicoot. Fossils have been found in the Wellington Caves of New South Wales. The bandicoot was about 20 centimeters long. It is believed to have gone extinct in the Late Pliocene. It is...

Percival's spiny mouse
Percival's spiny mouse

Acomys percivali

Percival's spiny mouse (*Acomys percivali*) is a species of rodent in the family Muridae. It is found in Kenya, Somalia, South Sudan, and Uganda. Its natural habitats are dry savanna, subtropical or tropical dry shrubland, and rocky areas. It is...

Percival's trident bat

Percival's trident bat

Cloeotis percivali

Percival's trident bat (*Cloeotis percivali*) is a species of bat in the family Hipposideridae. It is monotypic within the genus *Cloeotis*. It is found in Botswana, Democratic Republic of the Congo, Eswatini, Kenya, Mozambique, South Africa,...

Perdido Key beach mouse

Perdido Key beach mouse

Peromyscus polionotus trissyllepsis

The Perdido Key beach mouse (*Peromyscus polionotus trissyllepsis*) is an endangered subspecies of the oldfield mouse. It is found on Perdido Key, Florida. The small white and gray mouse, weighing only 13–16 g (0.46–0.56 oz), blends in well with...

Père David's Deer

Père David's Deer

Elaphurus davidianus

The Père David's deer (*Elaphurus davidianus*) is a species of deer, the only extant member of the genus *Elaphurus*. Based on genetic comparisons, Père David's deer is closely related to Eld's deer. Père David's deer were hunted almost to extinction...

Père David's mole

Père David's mole

Talpa davidiana

Père David's mole (*Talpa davidiana*) is a mole found in the Middle East, ranging from south-central Turkey east to Kurdistan Province, Iran, although it could also range south into Syria. It is listed as a data deficient species. The species is...

Père David's rock squirrel

Père David's rock squirrel

Sciurotamias davidianus

The Père David's rock squirrel (*Sciurotamias davidianus*), also known as the Chinese rock squirrel, is a species of rodent in the family Sciuridae. It is endemic to China, where it is found widely in rocky habitats in the eastern and central...

Père David's vole

Père David's vole

Eothenomys melanogaster

Père David's vole (*Eothenomys melanogaster*) is a species of rodent in the family Cricetidae. It is found in China, Myanmar, Taiwan, and Thailand. This species is a member of the melanogaster group, one of the two main groups of *Eothenomys* voles....

Perny's long-nosed squirrel

Perny's long-nosed squirrel

Dremomys pernyi

Perny's long-nosed squirrel (*Dremomys pernyi*) is a species of rodent in the family Sciuridae. It is found in Northeast India, southern and central China, northern Myanmar, northern Vietnam, and Taiwan.

Peromyscus nesodytes

Peromyscus nesodytes
Peromyscus nesodytes

The giant island deer mouse (*Peromyscus nesodytes*) became extinct approximately 8000 years BP and lived during the late Pleistocene on California's Channel Islands.

Peromyscus sagax
Peromyscus sagax
Peromyscus sagax

Peromyscus sagax (La Palma deer mouse or Michoacán deer mouse), is a species of mouse native to an area around La Palma, Los Reyes, Michoacán in Mexico.

Peromyscus schmidlyi
Peromyscus schmidlyi
Peromyscus schmidlyi

Schmidly's deer mouse (*Peromyscus schmidlyi*), is a recently described species of deer mouse from the mountains of western Mexico. It is part of the highly complex and well-studied *Peromyscus boylii* species complex. The uniqueness of *Peromyscus*...

Perote ground squirrel
Perote ground squirrel
Xerospermophilus perotensis

The Perote ground squirrel (*Xerospermophilus perotensis*) is a species of rodent in the family Sciuridae. It is endemic to Mexico and is currently at risk of extinction. Perote ground squirrels live within the same area as rock squirrels...

Perote mouse
Perote mouse
Peromyscus bullatus

The Perote mouse (*Peromyscus bullatus*), or Perote deer mouse, is a species of rodent in the family Cricetidae. It is found only in Mexico.

Perrier's sifaka
Perrier's sifaka
Propithecus perrieri

Perrier's sifaka (*Propithecus perrieri*) is a lemur endemic to Madagascar. It was formerly considered to be a subspecies of diademed sifaka. It has a very small range in northeastern Madagascar where its habitat is dry deciduous or semihumid...

Perrin's beaked whale
Perrin's beaked whale
Mesoplodon perrini

Perrin's beaked whale (*Mesoplodon perrini*) is part of the toothed whale suborder and is one of over 90 known cetaceans in existence today. Beaked whales are part of the family Ziphiidae, which are the second most diverse group out of all marine...

Persian Fallow Deer
Persian Fallow Deer
Dama dama mesopotamica

The Persian fallow deer (*Dama mesopotamica*) is a deer species once native to all of the Middle East, but currently only living in Iran and Israel. It was reintroduced in Israel. It has been listed as endangered on the IUCN Red List since 2008....

Persian jird
Persian jird
Meriones persicus

The Persian jird (*Meriones persicus*) is a species of rodent in the family Muridae

found in Afghanistan, Armenia, Azerbaijan, Iran, Iraq, Pakistan, Turkey, and Turkmenistan. The Persian jird is about 6 inches (15 cm) long and has a brush-tipped...

Persian Leopard

Persian Leopard

Panthera pardus tulliana

The Persian leopard is a leopard subspecies native to Western and Central Asia. Since 2016, it has been listed as Endangered on the IUCN Red List, as the wild population is estimated at less than 1000 mature individuals.

Persian Onager

Persian Onager

Equus hemionus onager

The Persian onager (*Equus hemionus onager*), also called the Persian wild ass or Persian zebra, is a subspecies of onager (Asiatic wild ass) native to Iran. It is listed as Endangered, with no more than 600 individuals left in the wild and only...

Persian vole

Persian vole

Microtus irani

The Persian vole (*Microtus irani*) is a species of rodent in the family Cricetidae. It is only found in Iran.

Peruvian climbing mouse

Peruvian climbing mouse

Rhipidomys modicus

The Peruvian climbing mouse or lesser Peruvian rhipidomys (*Rhipidomys modicus*), is a South American species of rodent in the family Cricetidae. It is endemic to the eastern slopes of the Andes in central Peru, where it is found at altitudes from...

Peruvian cotton rat

Peruvian cotton rat

Sigmodon peruanus

The Peruvian cotton rat (*Sigmodon peruanus*) is a species of rodent in the family Cricetidae. It is found in Ecuador and Peru.

Peruvian fish-eating rat

Peruvian fish-eating rat

Neusticomys peruviansis

The Peruvian fish-eating rat (*Neusticomys peruviansis*) is a species of semiaquatic rodent in the family Cricetidae. It is found only in eastern Peru, where it is known from locations at elevations from 200 to 400 m.

Peruvian night monkey

Peruvian night monkey

Aotus miconax

The Peruvian night monkey (*Aotus miconax*), also known as the Andean night monkey, is a nocturnal New World monkey endemic to northern Peru. Adults weigh around 1 kg (2.2 lb) and measure up to 50 cm (20 in) in length. Its colour is grey to light...

Peruvian Oldfield mouse

Peruvian Oldfield mouse

Thomasomys eleusis

The Peruvian Oldfield mouse (*Thomasomys eleusis*) is a species of rodent in the family Cricetidae. It is found only in Peru.

Peruvian small-eared shrew

Peruvian small-eared shrew

Cryptotis peruviansis

The Peruvian small-eared shrew (*Cryptotis peruviansis*) is a species of mammal in the family Soricidae. It is known only from northern Peru, where it has been found in shrubby "elfin" cloud forest in the Andes at elevations from 2050 to 3150 m....

Peruvian Spider Monkey

Peruvian Spider Monkey

Ateles chamek

The Peruvian spider monkey (*Ateles chamek*) also known as the black-faced black spider monkey, is a species of spider monkey that lives in Peru, as well as in Brazil and in Bolivia. At 60 centimetres (2 feet) long, they are relatively large among...

Peruvian tree-rat

Peruvian tree-rat

Makalata rhipidura

The Peruvian tree rat (*Makalata rhipidura*) is a species of rodent in the family Echimyidae. It is found in northeastern Peru and adjacent Ecuador, where it is found in the Amazon rainforest. It is nocturnal and arboreal. The etymology of the...

Peruvian tuco-tuco

Peruvian tuco-tuco

Ctenomys peruanus

The Peruvian tuco-tuco (*Ctenomys peruanus*) is a species of rodent in the family Ctenomyidae. It is endemic to Peru and Bolivia.

Peruvian vesper mouse

Peruvian vesper mouse

Calomys sorellus

The Peruvian vesper mouse (*Calomys sorellus*) is a species of rodent in the family Cricetidae. It is found only in Peru.

Peters's climbing rat

Peters's climbing rat

Tylomys nudicaudus

Peters's climbing rat (*Tylomys nudicaudus*) is a species of rodent in the family Cricetidae. It is found in Belize, El Salvador, Guatemala, Honduras, Mexico, and Nicaragua.

Peters's disk-winged bat

Peters's disk-winged bat

Thyroptera discifera

Peters's disk-winged bat (*Thyroptera discifera*) is a bat species mainly found in South and Central America. They belong to the family Thyropteridae which are a small group of disk-winged bats. These bats have a distinguishing feature of...

Peters's duiker

Peters's duiker

Cephalophus callipygus

Peters's duiker (*Cephalophus callipygus*) is a small antelope found in Gabon, Equatorial Guinea, southern Cameroon, and northern Republic of the Congo. Peters's duikers typically weigh about 40 lb (18 kg), and are about 20 in (50 cm) at the...

Peters's dwarf epauletted fruit bat

Peters's dwarf epauletted fruit bat

Micropteropus pusillus

Peters's dwarf epauletted fruit bat (*Micropteropus pusillus*) is a species of

megabat in the family Pteropodidae. It is found in Angola, Benin, Burkina Faso, Burundi, Cameroon, Central African Republic, Chad, Republic of the Congo, Democratic...

Peters's epauletted fruit bat

Peters's epauletted fruit bat

Epomophorus crypturus

The Peters's epauletted fruit bat (*Epomophorus crypturus*) is a species of megabat in the family Pteropodidae. It is found in Angola, Botswana, Democratic Republic of the Congo, Eswatini, Malawi, Mozambique, Namibia, Tanzania, Zambia, and...

Peters's flat-headed bat

Peters's flat-headed bat

Platymops setiger

Peters's flat-headed bat (*Platymops setiger*) is a species of bat in the family Molossidae and the monotypic genus *Platymops*. It is found in Ethiopia, Kenya, and South Sudan. Its natural habitats are dry savanna and rocky areas.

Peters's fruit bat

Peters's fruit bat

Cynopterus luzoniensis

Peters's fruit bat (*Cynopterus luzoniensis*) is a species of megabat within the family Pteropodidae. It is found in Sulawesi, Philippines, and adjacent small islands.

Peters's mouse

Peters's mouse

Mus setulosus

Peters' mouse (*Mus setulosus*) is a species of rodent in the family Muridae. It is found in Benin, Cameroon, Central African Republic, Republic of the Congo, Democratic Republic of the Congo, Ivory Coast, Equatorial Guinea, Ethiopia, Gabon, Ghana,...

Peters's musk shrew

Peters's musk shrew

Crocidura gracilipes

The Peters's musk shrew (*Crocidura gracilipes*) is a species of mammal in the family Soricidae. It is endemic to Tanzania.

Peters's myotis

Peters's myotis

Myotis ater

Peters's myotis or the small black myotis (*Myotis ater*) is a species of insectivorous vesper bat. It is found in Indonesia, Malaysia, Thailand, Vietnam, and the Philippines; its exact distribution is uncertain as it is difficult to distinguish...

Peters's pipistrelle

Peters's pipistrelle

Falsistrellus petersi

The Peters's pipistrelle (*Hypsugo petersi*) is a species of vesper bat in the family Vespertilionidae. It is found in Indonesia, Malaysia, and the Philippines. Formerly classified in the genus *Falsistrellus*, phylogenetic evidence supports it...

Peters's sheath-tailed bat

Peters's sheath-tailed bat

Paremballonura atrata

Peters's sheath-tailed bat (*Paremballonura atrata*) is a species of sac-winged bat

in the family Emballonuridae. It is found only in Madagascar. It is threatened by habitat loss.

Peters's squirrel

Peters's squirrel

Sciurus oculatus

Peters's squirrel (*Sciurus oculatus*) is a tree squirrel in the genus *Sciurus* endemic to Mexico. It was first described by the German naturalist and explorer Wilhelm Peters in 1863. Three subspecies are recognised. It is a common species, and the...

Peters's striped mouse

Peters's striped mouse

Hybomys univittatus

Peters's striped mouse or Peters's hybomys (*Hybomys univittatus*) is a species of rodent in the family Muridae. It is found in Angola, Burundi, Cameroon, Central African Republic, Republic of the Congo, Democratic Republic of the Congo, Equatorial...

Peters's trumpet-eared bat

Peters's trumpet-eared bat

Phoniscus jagorii

Peters' trumpet-eared bat (*Phoniscus jagorii*) is a species of bat in the family Vespertilionidae, the vesper bats. It is native to Indonesia, Laos, Malaysia, Vietnam, and the Philippines. It was named by Peters for Fedor Jagor. It is also known...

Peters's tube-nosed bat

Peters's tube-nosed bat

Harpiola grisea

Peters's tube-nosed bat (*Harpiola grisea*) is a species of vesper bat in the family Vespertilionidae, found in the Indian Subcontinent, mainly in the Western Himalayas. They have tube-shaped nostrils (hence the name) which assist them with their...

Peters's wrinkle-lipped bat

Peters's wrinkle-lipped bat

Mormopterus jugularis

Peters's wrinkle-lipped bat (*Mormopterus jugularis*), also called Peters's goblin bat, is a species of bat in the family Molossidae, the free-tailed bats. It is endemic to Madagascar, where it is widespread and in some areas abundant. It commonly...

Peterson's chinchilla mouse

Peterson's chinchilla mouse

Euneomys petersoni

Peterson's chinchilla mouse (*Euneomys petersoni*) is a species of rodent in the family Cricetidae. It is found in west central and southern Argentina and nearby areas of Chile. The species is named after American paleontologist Olaf A. Peterson...

Peterson's free-tailed bat

Peterson's free-tailed bat

Mops petersoni

Peterson's free-tailed bat (*Mops petersoni*) is a species of bat in the family Molossidae. It is found in Cameroon and Ghana, and its natural habitats are subtropical or tropical dry forest and subtropical or tropical moist lowland forest. It is...

Peterson's long-fingered bat

Peterson's long-fingered bat
Miniopterus petersoni

Peterson's long-fingered bat (*Miniopterus petersoni*) is a bat in the genus *Miniopterus* which occurs in southeast Madagascar. It was described by Steven M. Goodman et al. in 2008. While *M. petersoni* is similar to *M. sororculus*, the two species...

Petter's big-footed mouse
Petter's big-footed mouse
Macrotarsomys petteri

Petter's big-footed mouse (*Macrotarsomys petteri*), is a Madagascan rodent in the genus *Macrotarsomys*. With a head and body length of 150 mm (5.9 in) and body mass of 105 g (3.7 oz), It is the largest species of its genus. The upperparts are...

Petter's gerbil
Petter's gerbil
Taterillus petteri

Petter's gerbil or Petter's tateril (*Taterillus petteri*) is a species of rodent found in Burkina Faso, Mali, and Niger. Its natural habitats are dry savanna, subtropical or tropical dry shrubland, arable land, and rural gardens.

Petter's soft-furred mouse
Petter's soft-furred mouse
Praomys petteri

Petter's soft-furred mouse, or Petter's praomys (*Praomys petteri*) is a species of rodent in the family Muridae. It is found in Cameroon, Central African Republic, and Republic of the Congo. Its natural habitat is subtropical or tropical moist...

Petter's sportive lemur
Petter's sportive lemur
Lepilemur petteri

Petter's sportive lemur (*Lepilemur petteri*) is a sportive lemur endemic to Madagascar. It is one of 26 species in the genus *Lepilemur*. It is one of the smaller sportive lemurs with a total length of about 49 to 54 cm (19 to 21 in), of which...

Petter's tufted-tailed rat
Petter's tufted-tailed rat
Eliurus petteri

Petter's tufted-tailed rat (*Eliurus petteri*) is a rodent in the genus *Eliurus* found in lowland eastern Madagascar. First described in 1994, it is most closely related to the smaller *Eliurus grandidieri*. Virtually nothing is known of its natural...

Peyrieras's woolly lemur
Peyrieras's woolly lemur
Avahi peyrierasi

Peyrieras's woolly lemur or Peyrieras's avahi (*Avahi peyrierasi*) is a species of woolly lemur native to southeastern Madagascar. It weighs about 1 kg.

Pfeiffer's red bat
Pfeiffer's red bat
Lasiurus pfeifferi

Pfeiffer's red bat (*Lasiurus pfeifferi*) is a species of bat from the family Vespertilioninae and is endemic to Cuba. It is listed as Near Threatened by the IUCN Red List due to a significant population decline, caused by human population density...

Phayre's Leaf Monkey

Phayre's Leaf Monkey
Trachypithecus phayrei

Phayre's leaf monkey (*Trachypithecus phayrei*) is a species of Old World monkey native to South and Southeast Asia. It is listed as Endangered on the IUCN Red List and is threatened by hunting and loss of habitat.

Phayre's squirrel
Phayre's squirrel
Callosciurus phayrei

Phayre's squirrel (*Callosciurus phayrei*) is a species of rodent in the family Sciuridae. It is found in forests in China (Yunnan only) and Myanmar.

Philip Myers's akodont
Philip Myers's akodont
Akodon philipmyersi

Philip Myers's akodont (*Akodon philipmyersi*) is a recently described species of grass mouse from Misiones Province, Argentina. Like other grass mice, *A. philipmyersi* is a small, non-descript, greyish-brown mouse with prominent ears. The species...

Philippine dawn bat
Philippine dawn bat
Eonycteris robusta

The Philippine dawn bat (*Eonycteris robusta*) is a species of megabat in the family Pteropodidae found in the Philippines.

Philippine Deer
Philippine Deer
Rusa marianna

The Philippine deer (*Rusa marianna*), also known as the Philippine sambar or Philippine brown deer, is a vulnerable deer species endemic to the Philippines. It was first described from introduced populations in the Mariana Islands, hence the...

Philippine Flying Lemur
Philippine Flying Lemur
Cynocephalus volans

The Philippine flying lemur or Philippine colugo (*Cynocephalus volans*), known locally as kagwang, is one of two species of colugo or "flying lemurs." It is monotypic of its genus. Although called a flying lemur, it cannot fly and is not a lemur....

Philippine forest horseshoe bat
Philippine forest horseshoe bat
Rhinolophus inops

The Philippine forest horseshoe bat (*Rhinolophus inops*) is a species of bat in the family Rhinolophidae. It is endemic to the Philippines.

Philippine forest rat
Philippine forest rat
Rattus everetti

The Philippine forest rat is a species of rodent in the family Muridae. It is found only in the Philippines, and is located throughout the archipelago. The scientific name commemorates British colonial administrator and zoological collector...

Philippine forest roundleaf bat
Philippine forest roundleaf bat
Hipposideros obscurus

The Philippine forest roundleaf bat (*Hipposideros obscurus*) is a species of bat in the family Hipposideridae. It is endemic to the Philippines.

Philippine gray flying fox

Philippine gray flying fox

Pteropus speciosus

The Philippine gray flying fox (*Pteropus speciosus*) is a species of flying fox in the family Pteropodidae. It is found in Indonesia and the Philippines. Its natural habitat is subtropical or tropical dry forests. It is threatened by habitat loss.

Philippine long-fingered bat

Philippine long-fingered bat

Miniopterus paululus

The Philippine long-fingered bat (*Miniopterus paululus*) is a species of bat from the family Miniopteridae. The Philippine long-fingered bat is native to Indonesia, the Philippines, Malaysia, and Timor-Leste. Due to the dubious taxonomic status...

Philippine Long-Tailed Macaque

Philippine Long-Tailed Macaque

Macaca fascicularis philippensis

The Philippine long-tailed macaque (*Macaca fascicularis philippensis*) is a subspecies of the Long-tailed macaque. It is known in various Philippine languages as matching /matsing or the more general term unggoy ("monkey").

Philippine Mouse-Deer

Philippine Mouse-Deer

Tragulus nigricans

The Philippine mouse-deer (*Tragulus nigricans*), also known as the Balabac chevrotain or pilandok (in Filipino), is a small, nocturnal ruminant, which is endemic to Balabac and nearby smaller islands (Bugsuk and Ramos) southwest of Palawan in the...

Philippine Naked-Backed Fruit Bat

Philippine Naked-Backed Fruit Bat

Dobsonia chapmani

The Philippine naked-backed fruit bat (*Dobsonia chapmani*) is a megabat that mostly lives on Negros Island. The wings of these bats meet along the midline of their bodies, making them very agile fliers. They roosted in caves and were so abundant...

Philippine Pangolin

Philippine Pangolin

Manis culionensis

The Philippine pangolin or Palawan pangolin (*Manis culionensis*), also locally known as balintong, is a pangolin species endemic to the Palawan province of the Philippines. Its habitat includes primary and secondary forests, as well as...

Philippine porcupine

Philippine porcupine

Hystrix pumila

The Philippine porcupine or Palawan porcupine (*Hystrix pumila*) is a species of rodent in the family Hystricidae (Old World porcupines) endemic to the island of Palawan in the Philippines. It is known locally as durian or landak. Apparently, its...

Philippine pygmy roundleaf bat

Philippine pygmy roundleaf bat

Hipposideros pygmaeus

The Philippine pygmy roundleaf bat (*Hipposideros pygmaeus*), also called the

Philippine pygmy leaf-nosed bat, is a species of bat in the family Hipposideridae. It is endemic to the Philippines, where it has been recorded on Bohol, Luzon...
Philippine pygmy squirrel
Philippine pygmy squirrel
Exilisciurus concinnus

The Philippine pygmy squirrel (Exilisciurus concinnus) is a species of rodent in the family Sciuridae. It is endemic to the Philippines.
Philippine slow loris
Philippine slow loris
Nycticebus menagensis

The Philippine slow loris (Nycticebus menagensis) is a strepsirrhine primate and a species of slow loris that is native to the north and east coastal areas of the island of Borneo, as well as the Sulu Archipelago in the Philippines. The species...
Philippine Tarsier
Philippine Tarsier
Carlito syrichta

Known as the "world's smallest monkey" because of its similarity in appearance to that primate, tarsiers, along with lemurs, tree shrews, and lorises, actually are members of a more primitive suborder of Prosimii or prosimian. They are amongst...
Philippine tree squirrel
Philippine tree squirrel
Sundasciurus philippinensis

The Philippine tree squirrel (Sundasciurus philippinensis) is a species of rodent in the family Sciuridae which is endemic to the Philippines.
Philippine tube-nosed fruit bat
Philippine tube-nosed fruit bat
Nyctimene rabori

The Philippine tube-nosed fruit bat (Nyctimene rabori) locally known in Tagalog as Bayakan is a species of bat in the family Pteropodidae. It is endemic to the Philippines and known from the islands of Cebu, Negros and Sibuyan. It occurs in and...
Philippine Warty Pig
Philippine Warty Pig
Sus philippensis

The Philippine warty pig (Sus philippensis) is one of four known species in the pig genus (Sus) endemic to the Philippines. It has multiple native common names, but it is most widely known as baboy damo ("bush pig") in Tagalog.
Phillips's gerbil
Phillips's gerbil
Gerbilliscus phillipsi

Phillips's gerbil (Gerbilliscus phillipsi) is a species of rodent found in Ethiopia, Kenya, and Somalia. Its natural habitat is subtropical or tropical dry shrubland.
Phillips's kangaroo rat
Phillips's kangaroo rat
Dipodomys phillipsii

Phillips's kangaroo rat (Dipodomys phillipsii) is a species of rodent in the family Heteromyidae. It is endemic to Mexico. Its natural habitat is hot deserts. The species is named after John Phillips, an official of a Mexican mining company who...

Phillips's mouse
Phillips's mouse
Mus phillipsi

Phillips's mouse (*Mus phillipsi*) is a species of rodent in the family Muridae. It is found only in India. Its natural habitats are subtropical or tropical dry forests, subtropical or tropical dry lowland grassland, and hot deserts. It is threatened...

Phillips's small-eared shrew
Phillips's small-eared shrew
Cryptotis phillipsii

Phillips' small-eared shrew or Phillips' short-eared shrew (*Cryptotis phillipsii*) is a species of mammal in the family Soricidae found in Mexico.

Phou Khao Khouay leaf-nosed bat
Phou Khao Khouay leaf-nosed bat
Hipposideros khaokhouayensis

The Phou Khao Khouay leaf-nosed bat (*Hipposideros khaokhouayensis*) is a species of bat found in Laos and Vietnam. It was described as a new species in 2006. It is considered vulnerable to extinction by the IUCN.

Phyllomys sulinus
Phyllomys sulinus
Phyllomys sulinus

Phyllomys sulinus, the southern Atlantic tree-rat, is a species of spiny rat from South America, described in 2008. It is found in the subtropical region of southern Brazil, from the states of São Paulo to Rio Grande do Sul. The etymology of the...

Pichi
Pichi
Zaedyus pichiy

The pichi (*Zaedyus pichiy*) is the only living member of the genus *Zaedyus*, [2] and the only armadillo to hibernate. Fossil remains from the Cerro Azul Formation indicate this species had already evolved during the late Miocene epoch.

Pichincha Oldfield mouse
Pichincha Oldfield mouse
Thomasomys vulcani

The Pichincha Oldfield mouse (*Thomasomys vulcani*) is a species of rodent in the family Cricetidae. It is present in the Cordillera Occidental of the Andes of Ecuador, where its habitats include shrubby páramo and montane forest. It is nocturnal...

Piebald shrew
Piebald shrew
Diplomesodon pulchellum

The piebald shrew (*Diplomesodon pulchellum*) is a shrew found in the Turan Lowland east of the Caspian Sea in Iran, Turkmenistan and Uzbekistan. It grows to 2–23/4 inches in length, and usually hunts for insects and lizards at night. It is the...

Pied bat
Pied bat
Niumbaha superba

The pied butterfly bat (*Glauconycteris superba*), also known as the pied bat or badger bat, is a rare species of vesper bat in the family Vespertilionidae.

Pied Tamarin
Pied Tamarin
Saguinus bicolor

The Pied tamarin (*Saguinus bicolor*) is a Critically Endangered primate species found in a restricted area of the Brazilian Amazon Rainforest. It was named the mascot of Manaus, Brazil in 2005.

Pig-tailed langur

Pig-tailed langur

Simias concolor

The pig-tailed langur (*Simias concolor*), monotypic in genus *Simias*, is a large Old World monkey, endemic to several small islands off the coast of Sumatra in Indonesia. Its face is black, its fur is blackish-brown and it has a relatively short...

Pikumche pericote

Pikumche pericote

Loxodontomys pikumche

The Pikumche pericote (*Loxodontomys pikumche*) is a species of rodent in the family Cricetidae. It is known only from central Chile, where its range includes xeric shrublands (the matorral).

Pilar tuco-tuco

Pilar tuco-tuco

Ctenomys pilarensis

The Pilar tuco-tuco (*Ctenomys pilarensis*) is a species of rodent in the family Ctenomyidae. It is found in Ñeembucú and Misiones Departments in southern Paraguay, east of the city of Pilar. The species lives in areas with sandy soil. It is...

Pilbara ningauí

Pilbara ningauí

Ningauí timealeyi

The Pilbara ningauí (*Ningauí timealeyi*), sometimes known as Ealey's ningauí, is a tiny species of marsupial carnivore found in Australia.

Pileated Gibbon

Pileated Gibbon

Hylobates pileatus

The Pileated gibbon (*Hylobates pileatus*) is an endangered species of primate from Southeast Asia. Part of its scientific name means "forest walker". Gibbons live in trees and unlike most of the great apes, they frequently form long-term pair bonds.

Pilliga mouse

Pilliga mouse

Pseudomys pilligaensis

Pseudomys pilligaensis, commonly known as the Pilliga mouse or poolkoo, is a species of rodent in the family Muridae. Its distribution comprises the Pilliga forest region of New South Wales, Australia, specimens have also been trapped in the...

Pine Marten

Pine Marten

Martes martes

The European pine marten (*Martes martes*) is native to Northern Europe and belongs to the mustelid family, other members of which are mink, otter, badger, weasel, and wolverine. It is an agile animal, adapted to living in the forests. It is known...

Pinheiro's slender opossum

Pinheiro's slender opossum

Marmosops pinheiroi

Pinheiro's slender opossum (*Marmosops pinheiroi*) is an opossum species from South

America. It is found in Brazil, French Guiana, Guyana, Suriname and Venezuela.
Pink Fairy Armadillo
Pink Fairy Armadillo
Chlamyphorus truncatus

The Pink fairy armadillo (*Chlamyphorus truncatus*) is the smallest species of armadillo first described by Richard Harlan in 1825. It is the only armadillo that has a dorsal shell that is almost completely separate from the body. This solitary, ...
Pinyon Mouse
Pinyon Mouse
Peromyscus truei

The pinyon mouse (*Peromyscus truei*) is native to the southwestern United States and Baja California in Mexico. These medium-sized mice are often distinguished by their relatively large ears. The range of this species extends from southern Oregon...
Pipistrellus hanaki
Pipistrellus hanaki
Pipistrellus hanaki

Hanak's dwarf bat or Hanak's pipistrelle (*Pipistrellus hanaki*) is a species of bat only found in Cyrenaica, Libya and Crete, Greece.
Pipistrellus raceyi
Pipistrellus raceyi
Pipistrellus raceyi

Pipistrellus raceyi, also known as Racey's pipistrelle, is a bat from Madagascar, in the genus *Pipistrellus*. Although unidentified species of *Pipistrellus* had been previously reported from Madagascar since the 1990s, *P. raceyi* was not formally...
Pitman's shrew
Pitman's shrew
Crocidura pitmani

Pitman's shrew (*Crocidura pitmani*) is a species of mammal in the family Soricidae. It is endemic to Zambia.
Pittier's crab-eating rat
Pittier's crab-eating rat
Ichthyomys pittieri

Pittier's crab-eating rat (*Ichthyomys pittieri*) is a species of semiaquatic rodent in the family Cricetidae. It is endemic to Venezuela. The natural habitats of this species are rivers and swamps. Its karyotype has $2n = 92$ and $FNa = 98$. This was...
Piute ground squirrel
Piute ground squirrel
Urocitellus mollis

The Piute ground squirrel (*Urocitellus mollis*) is a species of rodent in the family Sciuridae. It is endemic to the Great Basin region of the western United States, where it is found in parts of California, Idaho, Nevada, Oregon, Utah, and...
Plain brush-tailed rat
Plain brush-tailed rat
Isothrix pagurus

The Plain brush-tailed rat, *Isothrix pagurus*, is a spiny rat species from South America. It is found in Brazil.
Plains bison
Plains bison

Bison bison bison

The Plains bison (*Bison bison bison*) is one of two subspecies/ecotypes of the American bison, the other being the wood bison (*B. b. athabasca*). A natural population of Plains bison survives in Yellowstone National Park (the Yellowstone Park...

Plains coyote

Plains coyote

Canis latrans latrans

The coyote (*Canis latrans*) is a species of canine native to North America. It is smaller than its close relative, the wolf, and slightly smaller than the closely related eastern wolf and red wolf. It fills much of the same ecological niche as...

Plains harvest mouse

Plains harvest mouse

Reithrodontomys montanus

The plains harvest mouse (*Reithrodontomys montanus*) is a species of rodent in the family Cricetidae. It is found in northern Mexico and the central United States.

Plains Pocket Gopher

Plains Pocket Gopher

Geomys bursarius

The Plains pocket gopher (*Geomys bursarius*) is a burrowing rodent from North America. Pocket gophers are so named in reference to their externally located, fur-lined cheek pouches. They are the most highly fossorial rodents found in North America.

Plains pocket mouse

Plains pocket mouse

Perognathus flavescens

The plains pocket mouse (*Perognathus flavescens*) is a heteromyid rodent of North America. It ranges from southwestern Minnesota and southeastern North Dakota to northern Texas east of the Rockies, and from northern Utah and Colorado to northern...

Plains Rat

Plains Rat

Pseudomys australis

The plains rat (*Pseudomys australis*), also known as the palyoora, plains mouse and eastern mouse, is a conilurine rodent native to arid and semi-arid Australia. Referred to as the palyoora or yarlle by Indigenous groups, the plains rat was once...

Plains viscacha

Plains viscacha

Lagostomus maximus

The plains viscacha or plains vizcacha (*Lagostomus maximus*) is a species of viscacha, a rodent in the family Chinchillidae. It is the only living species within the genus *Lagostomus*. It is found in Argentina, Bolivia, and Paraguay. The plains...

Plains viscacha rat

Plains viscacha rat

Tympanoctomys barrerae

The plains viscacha rat, plains vizcacha rat, red viscacha rat, or red vizcacha rat (*Tympanoctomys barrerae*) is a species of rodent in the family Octodontidae native to Argentina. It is one of three species in the genus *Tympanoctomys*.

Plains Zebra

Plains Zebra
Equus quagga

The Plains zebra (*Equus quagga*, formerly *Equus burchellii*) is the most common and geographically widespread species of zebra. It is the most recognizable grazing animal on the African continent. Their closest relatives are donkeys and horses.
The...

Plantain Squirrel
Plantain Squirrel
Callosciurus notatus

The Plantain squirrel (*Callosciurus notatus*) is a species of rodent from Southeast Asia. They live in a wide range of habitats and fruit farmers consider them to be pests.

Plateau mouse
Plateau mouse
Peromyscus melanophrys

The plateau mouse (*Peromyscus melanophrys*) is a species of rodent in the family Cricetidae. It is found only in Mexico.

Plateau Pika
Plateau Pika
Ochotona curzoniae

The Plateau pika (*Ochotona curzoniae*) is a small mountain-dwelling mammal. It prefers to live in high elevations, mostly in the Tibetan Plateau, which is where the common name originates from.

Plateau vole
Plateau vole
Lasiopodomys fuscus

The plateau vole (*Lasiopodomys fuscus*) is a species of rodent in the family Cricetidae. It is found only in China. Its natural habitat is temperate grassland.

Platypus
Platypus
Ornithorhynchus anatinus

The platypus (*Ornithorhynchus anatinus*), sometimes referred to as the duck-billed platypus, is a semiaquatic, egg-laying mammal endemic to eastern Australia, including Tasmania. Together with the four species of echidna, it is one of the five...

Platyrrhinus albericoi
Platyrrhinus albericoi
Platyrrhinus albericoi

Platyrrhinus albericoi is a species of leaf-nosed bat found in South America.

Platyrrhinus ismaeli
Platyrrhinus ismaeli
Platyrrhinus ismaeli

Platyrrhinus ismaeli is a species of bat found in South America.

Pleasant bolo mouse
Pleasant bolo mouse
Necromys amoenus

The pleasant bolo mouse, or pleasant akodont, (*Necromys amoenus*) is a species of rodent in the family Cricetidae. It is found on grassland at high altitudes in Bolivia and Peru.

Pleasant gerbil

Pleasant gerbil
Gerbillus amoenus

The pleasant gerbil (*Gerbillus amoenus*) is a species of rodent found mainly in Libya and Egypt, and possibly Mauritania to Tunisia. This species is about 6 cm in body length, with a brown agouti-style coat, a white belly and a very long tail. It...

Pleistocene New Zealand sea lion
Pleistocene New Zealand sea lion
Neophoca palatina

Neophoca palatina, the Pleistocene New Zealand sea lion, is an extinct species of pinniped known from a nearly-complete adult male skull found at Ōhope Beach on the North Island in 1937. It was found in a stratum from the late Castlecliffian...

Plush-coated ringtail possum
Plush-coated ringtail possum
Pseudocheirops corinnae

The plush-coated ringtail possum or golden ringtail possum (*Pseudocheirops corinnae*) is a species of marsupial in the family Pseudocheiridae. It is found in Indonesia, Papua New Guinea, and the Solomon Islands. Its natural habitat is subtropical...

Pocketed free-tailed bat
Pocketed free-tailed bat
Nyctinomops femorosaccus

The pocketed free-tailed bat (*Nyctinomops femorosaccus*) is a species of bat in the family Molossidae found in Mexico and in Arizona, California, New Mexico, and Texas in the United States. They resemble the Brazilian free-tailed bat ("Tadarida...

Pocock's highland rat
Pocock's highland rat
Rattus pococki

Pocock's highland rat (*Rattus pococki*) is a species of rat of the family Muridae. It is native to New Guinea.

Podogymnura truei
Podogymnura truei
Podogymnura truei

Podogymnura truei, also known as the Mindanao gymnure, Mindanao moonrat, or Mindanao wood shrew, is a mammal of the family Erinaceidae. It is endemic to the Mindanao islands of the Philippines. Erinaceidae is a family of small mammals that...

Podolsk mole-rat
Podolsk mole-rat
Spalax zemni

The Podolsk blind mole-rat or Podolian mole-rat (*Spalax zemni*) is a species of rodent in the family Spalacidae. It is endemic to western and central Ukraine.

Pohle's fruit bat
Pohle's fruit bat
Casinycteris ophiodon

Pohle's fruit bat (*Casinycteris ophiodon*) is a near threatened species of megabat found in the subtropical or tropical forests of Cameroon, the Republic of Congo, Côte d'Ivoire, Ghana and Liberia.

Polar Bear
Polar Bear
Ursus maritimus

The Polar bear (*Ursus maritimus*) is a hypercarnivorous bear found within the Arctic Circle, encompassing the Arctic Ocean, its surrounding seas and surrounding land masses. It is the largest extant bear species, as well as the largest extant land...

Polia's shrew

Polia's shrew

Crocidura polia

Polia's shrew (*Crocidura polia*) is a species of mammal in the family Soricidae. It is endemic to Democratic Republic of the Congo.

Polynesian Rat

Polynesian Rat

Rattus exulans

The Polynesian rat (*Rattus exulans*) is the third most widespread species of rat in the world behind the Brown rat and Black rat. The Polynesian rat originated in Southeast Asia, and like its relatives, has become widespread, migrating to most of...

Pomona roundleaf bat

Pomona roundleaf bat

Hipposideros pomona

The Pomona roundleaf bat, Pomona leaf-nosed bat, or Andersen's leaf-nosed bat (*Hipposideros pomona*) is a species of bat in the family Hipposideridae that is endemic to India.

Poncelet's giant rat

Poncelet's giant rat

Solomys ponceleti

Poncelet's giant rat or Poncelet's naked-tailed rat (*Solomys ponceleti*) is a species of rodent in the family Muridae. It is found in Papua New Guinea and Solomon Islands.

Pond bat

Pond bat

Myotis dasycneme

The pond bat (*Myotis dasycneme*) is a species of vesper bat. It is found in Eurasia from France to Russia and Kazakhstan.

Popayán Oldfield mouse

Popayán Oldfield mouse

Thomasomys popayanus

The Popayán Oldfield mouse (*Thomasomys popayanus*), also called the "Popayán thomasomys", is a species of rodent in the family Cricetidae. It is present in the Andes of southwestern and central western Colombia, at elevations from 1800 to 3200 m....

Porcupine caribou

Porcupine caribou

Rangifer tarandus granti

The Porcupine caribou or Grant's caribou (*Rangifer tarandus granti*) is a subspecies of the reindeer or caribou found in Alaska, United States, and Yukon and the Northwest Territories, Canada. It resembles the subspecies known as the...

Portenko's shrew

Portenko's shrew

Sorex portenkoi

Portenko's shrew (*Sorex portenkoi*) is a species of mammal in the family Soricidae

that is endemic to Russia.
Porteous's tuco-tuco
Porteous's tuco-tuco
Ctenomys porteousi

The Mendoza tuco-tuco (*Ctenomys mendocinus*) is a species of rodent in the family Ctenomyidae.
Porter's rock rat
Porter's rock rat
Aconaemys porteri

Porter's rock rat (*Aconaemys porteri*) is a species of rat in the family Octodontidae. It is found in Argentina and Chile at altitudes between 900 and 2,000 meters above sea level.
Portuguese ibex
Portuguese ibex
Capra pyrenaica lusitanica

The Portuguese ibex (*Capra pyrenaica lusitanica*) is an extinct subspecies of Iberian ibex that inhabited the north mountainous zones of Portugal, Galicia, Asturias and western Cantabria. In size and colouration it was much like the Spanish...
Pouched gerbil
Pouched gerbil
Desmodilliscus braueri

The pouched gerbil (*Desmodilliscus braueri*) is a species of rodent in the family Muridae. It is the only species in the genus *Desmodilliscus* and the subtribe *Desmodilliscina*. It is found across western Africa from Mauritania east to Sudan. Its...
Pousargues African fat mouse
Pousargues African fat mouse
Steatomys opimus

Pousargues's African fat mouse (*Steatomys opimus*) is an extant species of rodent indigenous to the Central African Republic, Cameroon, and the Democratic Republic of the Congo. Given the distribution of the species throughout central Africa, the...
Pousargues's mongoose
Pousargues's mongoose
Dologale dybowskii

Pousargues's mongoose (*Dologale dybowskii*), also known as the African tropical savannah mongoose, is a mongoose native to Central Africa. It is listed as data deficient on the IUCN Red List as little is known about its distribution and...
Prairie shrew
Prairie shrew
Sorex haydeni

The prairie shrew (*Sorex haydeni*) is a small shrew found in the Canadian prairies and midwestern United States. At one time, this species was considered to be a subspecies of the similar cinereus shrew, *S. cinereus*. It is brown in color with...
Prairie Vole
Prairie Vole
Microtus ochrogaster

The prairie vole (*Microtus ochrogaster*) is a small vole found in central North America. The vole has long, coarse grayish-brown fur on the upper portion of the body and yellowish fur on the lower portion of the body. It has short ears and a...

Praomys coetzei
Praomys coetzei
Praomys coetzei

Praomys coetzei is a species of mouse in the family Muridae.
Pratt's roundleaf bat
Pratt's roundleaf bat
Hipposideros pratti

Pratt's roundleaf bat (Hipposideros pratti) is a species of bat in the family Hipposideridae. It is found in China, Malaysia, Myanmar, Thailand, and Vietnam.
Pratt's vole
Pratt's vole
Eothenomys chinensis

Pratt's vole (Eothenomys chinensis) is a species of rodent in the family Cricetidae. It is endemic to Mount Emei, Sichuan, China. It was named in 1891 for Antwerp Edgar Pratt.
Preble's meadow jumping mouse
Preble's meadow jumping mouse
Zapus hudsonius preblei

Preble's meadow jumping mouse (Zapus hudsonius preblei) is a subspecies of meadow jumping mouse, endemic to the upland habitats of Colorado and Wyoming in North America. It is found nowhere else in the world. It is listed as Threatened under the...
Preble's shrew
Preble's shrew
Sorex preblei

Preble's shrew (Sorex preblei) is a small shrew distributed across the Great Basin of the United States and southern British Columbia in Canada. It belongs to the order Eulipotyphla, family Soricidae and genus Sorex.
Prehensile-tailed hutia
Prehensile-tailed hutia
Mysateles prehensilis

The prehensile-tailed hutia (Mysateles prehensilis) is a species of hutia in the subfamily Capromyinae endemic to Cuba. It is the only member of the genus Mysateles. It is an arboreal foliovore, found in both primary and secondary forest.
Prehensile-tailed rat
Prehensile-tailed rat
Pogonomys mollipilosus

The prehensile-tailed rat (Pogonomys mollipilosus) is a species of rodent that can be found in New Guinea and Australia; it occurs in the tropical rainforest of Queensland.
Prehesile-Tailed Porcupine
Prehesile-Tailed Porcupine
Coendou prehensilis

The Brazilian porcupine is a porcupine found in Brazil, Argentina, Colombia, Venezuela, Guyana, French Guiana, Peru, Paraguay, Suriname, Bolivia and Trinidad, with a single record from Ecuador. It inhabits tropical forests at elevations up to...
Pretty-Faced Wallaby
Pretty-Faced Wallaby
Macropus parryi

The whiptail wallaby (*Notamacropus parryi*), also known as the pretty-faced wallaby, is a species of wallaby found in eastern Australia. It is locally common from Cooktown in Queensland to near Grafton in New South Wales.

Preuss's monkey

Preuss's monkey

Allochrocebus preussi

The Preuss's monkey (*Allochrocebus preussi*), also known as Preuss's guenon, is a diurnal primate that lives terrestrially in mountainous (up to 2500 m) forests of eastern Nigeria, western Cameroon and Bioko in Equatorial Guinea. It was formerly...

Preuss's red colobus

Preuss's red colobus

Procolobus preussi

Preuss's red colobus (*Piliocolobus preussi*) is a red colobus primate species endemic to the Cross-Sanaga Rivers ecoregion. An important population occurs in Korup National Park, Southwest Province, Cameroon, but the species' distribution is...

Prevost's Squirrel

Prevost's Squirrel

Callosciurus prevostii

Prevost's squirrel (*Callosciurus prevostii*) is a colorful rodent found in forests of Southeast Asia. This species is declining in some regions because of habitat loss and hunting, it can live in somewhat disturbed habitats and often visits...

Pribilof Island shrew

Pribilof Island shrew

Sorex pribilofensis

The Pribilof Island shrew (*Sorex pribilofensis*) is a species of mammal in the family Soricidae. It is found only on Alaska's Pribilof Islands.

Primordial tapecua

Primordial tapecua

Tapecomys primus

The primordial tapecua (*Tapecomys primus*) is a species of medium-sized rodent in the family Cricetidae. The type locality is in southeastern Bolivia. It is the only known species of the genus *Tapecomys*. Two specimens were found in 1991 in a...

Prince Bernhard's titi

Prince Bernhard's titi

Plecturocebus bernhardi

Prince Bernhard's titi monkey (*Plecturocebus bernhardi*) is a species of titi, within the subfamily Callicebinae. It is named for Prince Bernhard of the Netherlands, a naturalist who created the Order of the Golden Ark honoring conservationists...

Prince Demidoff's bushbaby

Prince Demidoff's bushbaby

Galagoides demidovii

Prince Demidoff's bushbaby (*Galagoides demidovii*), also known as Prince Demidoff's galago, is a species of primate in the family Galagidae. It is native to parts of tropical West and Central Africa.

Prince of Wales flying squirrel

Prince of Wales flying squirrel

Glaucomys sabrinus griseifrons

The Prince of Wales flying squirrel, *Glaucomys sabrinus griseifrons*, is a

subspecies of the northern flying squirrel. *Glaucomys sabrinus griseifrons* is endemic to Prince of Wales Island in Alaska and has a unique coloration compared to other...

Principal gerbil

Principal gerbil

Gerbillus principulus

The principal gerbil (*Gerbillus principulus*) is a species of gerbil distributed mainly in Sudan; Jebel Meidob; El Malha. Fewer than 250 individuals of this species are thought to persist in the wild.

Pristine mustached bat

Pristine mustached bat

Pteronotus pristinus

The pristine mustached bat (*Pteronotus (Phyllodia) pristinus*) is an extinct Late Quaternary species of bat in the endemic Neotropical family Mormoopidae. It was distributed in Cuba and possibly Florida (United States).

Proboscis bat

Proboscis bat

Rhynchonycteris naso

The proboscis bat (*Rhynchonycteris naso*) is a bat species from South and Central America. Other common names include Long-nosed proboscis bat, sharp-nosed bat, Brazilian long-nosed bat and river bat in English, and murciélago narizón in Spanish....

Proboscis Monkey

Proboscis Monkey

Nasalis larvatus

The Proboscis monkey (*Nasalis larvatus*) is one of the largest monkey species native to Asia. It has a characteristic long nose that distinguishes this animal from other monkeys. As a matter of fact, another name of this primate is 'Monyet...

Pronghorn

Pronghorn

Antilocapra americana

The pronghorn (*Antilocapra americana*) is a species of even-toed mammal native to North America. Though not an antelope, it is known colloquially in North America as the American antelope, prong buck, pronghorn antelope, and prairie antelope,...

Proserpine rock-wallaby

Proserpine rock-wallaby

Petrogale persephone

The Proserpine rock-wallaby (*Petrogale persephone*) is a species of rock-wallaby restricted to a small area in Conway National Park, Dryander National Park, Gloucester Island National Park, and around the town of Airlie Beach, all in Whitsunday...

Protocetus

Protocetus

Protocetus atavus

Protocetus atavus ("first whale") is an extinct species of primitive cetacean from Egypt. It lived during the middle Eocene period 45 million years ago. The first discovered protocetid, *Protocetus atavus* was described by Fraas 1904 based on a...

Przewalski's Gazelle

Przewalski's Gazelle

Procapra przewalskii

In the wild, Przewalski's gazelle (*Procapra przewalskii*) is found only in China.

Once widespread, its range has declined to six populations near Qinghai Lake. This gazelle was named after Nikolai Przhevalsky, a Russian explorer who collected a...
Przewalski's gerbil
Przewalski's gerbil
Brachiones przewalskii

Przewalski's gerbil or Przewalski's jird (*Brachiones przewalskii*) is a species of rodent in the family Muridae. It is the only species in the genus *Brachiones*, and is found only in China.
Przewalski's Horse
Przewalski's Horse
Equus ferus

Przewalski's horse (*Equus ferus przewalskii* or *Equus przewalskii*), otherwise known as P-horse, is named after Nikolai Przewalski (pronounced "shuh-val-skee"), a Russian explorer. In 1870, he was the first to discover a subspecies of this horse....
Przewalski's steppe lemming
Przewalski's steppe lemming
Eolagurus przewalskii

Przewalski's steppe lemming (*Eolagurus przewalskii*) is a species of rodent in the family Cricetidae. It is found in China and Mongolia.
Pseudomys vandycki
Pseudomys vandycki
Pseudomys vandycki

Pseudomys vandycki is a species of the murid family, mice and rats, that is known from fossils discovered in Queensland, Australia.
Pseudoryzomys
Pseudoryzomys
Pseudoryzomys simplex

Pseudoryzomys simplex, also known as the Brazilian false rice rat or false oryzomys, is a species of rodent in the family Cricetidae from south-central South America. It is found in lowland palm savanna and thorn scrub habitats. It is a...
Pterophocaena
Pterophocaena
Pterophocaena nishinoi

Pterophocaena nishinoi is an extinct species of porpoise discovered in the Late Miocene Wakkanai Formation of Japan dating to 9.3–9.2 million years ago (mya), and may represent an intermediate phase between porpoises and dolphins. It is one of...
Pteropus pelagicus
Pteropus pelagicus
Pteropus pelagicus

Pteropus pelagicus is a species of fruit bat in the family Pteropodidae. It includes two subspecies that were formerly recognized as full species– *Pteropus insularis* (Chuuk flying fox) and *Pteropus phaeocephalus* (Mortlock flying fox). It is...
Pu Hoat muntjac
Pu Hoat muntjac
Muntiacus puhoatensis

The Pu Hoat muntjac (*Muntiacus puhoatensis*) is a species of muntjac only known from Pu Hoat region in Vietnam, which is bordering Laos. It is sometimes considered to be conspecific with Roosevelt's muntjac, and its habitat and behavior are...
Puebla deer mouse

Puebla deer mouse
Peromyscus mekisturus

The Puebla deer mouse (*Peromyscus mekisturus*) is a species of rodent in the family Cricetidae. It is found only in Mexico.

Puerto Rican flower bat
Puerto Rican flower bat
Phyllonycteris major

The Puerto Rican flower bat (*Phyllonycteris major*) is an extinct species of bat from the family Phyllostomidae (leaf-nosed bats). It was native to Puerto Rico and is known only from subfossil skeletal material.

Puerto Rican long-nosed bat
Puerto Rican long-nosed bat
Monophyllus plethodon frater

The Puerto Rican long-nosed bat (*Monophyllus plethodon frater*) is known only from a skull fragment excavated in the large Cathedral Cave near Morovis, Puerto Rico, by Dr. H. E. Anthony prior to 1917. This species was never observed or documented...

Puku
Puku
Kobus vardonii

The puku (*Kobus vardonii*) is a medium-sized antelope found in wet grasslands in southern Democratic Republic of Congo, Namibia, Tanzania, Zambia and more concentrated in the okavango delta in Botswana. Nearly one-third of all puku are found in...

Puma pumoides
Puma pumoides
Puma pumoides

Puma pumoides is an extinct prehistoric cat that was described in 1956 by Alfredo Castellanos using the scientific name *Felis pumoides*. Castellanos excavated its fossil remains in the Reartes Valley located in the province of Córdoba, Argentina,...

Puna mouse
Puna mouse
Punomys lemminus

The puna mouse (*Punomys lemminus*) is a species of rodent in the family Cricetidae. It is endemic to southern Peru, where it is found in puna habitat at elevations of 4400 to 4900 m in the Cordillera Occidental.

Pundt's tuco-tuco
Pundt's tuco-tuco
Ctenomys pundti

Pundt's tuco-tuco (*Ctenomys pundti*) is a species of rodent in the family Ctenomyidae. It is endemic to the Pampas of southern Córdoba and San Luis Provinces in central Argentina. The species is named after Argentine landowner and collector...

Pungent pipistrelle
Pungent pipistrelle
Falsistrellus mordax

The pungent pipistrelle (*Hypsugo mordax*) is a species of vesper bat in the family Vespertilionidae. It is found in Indonesia and possibly the Philippines. Formerly classified in the genus *Falsistrellus*, phylogenetic evidence supports it belonging...

Puno grass mouse
Puno grass mouse
Akodon subfuscus

The Puno grass mouse (*Akodon subfuscus*) is a species of rodent in the family Cricetidae. It is found in Bolivia and Peru.

Punta de Vacas chinchilla rat
Punta de Vacas chinchilla rat
Abrocoma vaccarum

The Punta de Vacas chinchilla rat (*Abrocoma vaccarum*), also known as the Mendoza chinchilla rat, is a species of chinchilla rats in the family Abrocomidae. Found only in Argentina, this species was once thought to be part of *Abrocoma cinerea*....

Puntilla tuco-tuco
Puntilla tuco-tuco
Ctenomys coludo

The Puntilla tuco-tuco (*Ctenomys coludo*) is a species of rodent in the family Ctenomyidae. It is endemic to central Argentina. The common name of the species comes from the municipality of La Puntilla at the type locality. It was first described...

Purple-Faced Langur
Purple-Faced Langur
Semnopithecus vetulus

The purple-faced langur (*Semnopithecus vetulus*), also known as the purple-faced leaf monkey, is a species of Old World monkey that is endemic to Sri Lanka. The animal is a long-tailed arboreal species, identified by a mostly brown appearance,...

Purple-necked rock-wallaby
Purple-necked rock-wallaby
Petrogale purpureicollis

The purple-necked rock-wallaby (*Petrogale purpureicollis*) is a species of rock-wallaby first described in 1924 by Albert Sherbourne Le Souef, then director of the Taronga Zoo in Sydney, Australia, who noted a purple colouration around the neck...

Purus red howler
Purus red howler
Alouatta puruensis

The Purús red howler (*Alouatta puruensis*) is a species of howler monkey native to Brazil, Peru and north of Bolivia.

Pygmy bamboo bat
Pygmy bamboo bat
Tylonycteris pygmaeus

The pygmy bamboo bat (*Tylonycteris pygmaeus*) is a species of vesper bat in the family Vespertilionidae. It is found in Southwest China and was discovered in 2007. The species is around 4 cm long and weighs between 2.6 and 3.5 grams.

Pygmy beaked whale
Pygmy beaked whale
Mesoplodon peruvianus

The pygmy beaked whale (*Mesoplodon peruvianus*), also known as the bandolero beaked whale, Peruvian beaked whale and lesser beaked whale, is the smallest of the mesoplodonts and one of the newest discoveries. There were at least two dozen...

Pygmy blue whale
Pygmy blue whale
Balaenoptera musculus brevicauda

The pygmy blue whale (*Balaenoptera musculus brevicauda*) is a subspecies of the blue whale (*Balaenoptera musculus*) found in the Indian Ocean and the southern Pacific Ocean. Reaching lengths of up to 24 metres (79 ft) it is smaller than the other...

Pygmy brocket
Pygmy brocket
Mazama nana

The pygmy brocket (*Mazama nana*) is a brocket deer species from South America. It is found in southern Brazil, Argentina and Paraguay. It is a small deer with short legs, weighing 15 to 20 kilograms (33 to 44 lb). It is reddish-brown in...

Pygmy brown-toothed shrew
Pygmy brown-toothed shrew
Chodsigoa parva

The pygmy brown-toothed shrew (*Chodsigoa parva*) is a species of shrew in the order Eulipotyphla. It is distributed in China. *C. parva* was initially thought to be the same as *Chodsigoa lamula*, but it was found to be a separate species.

Pygmy fruit bat
Pygmy fruit bat
Aethalops alecto

The pygmy fruit bat (*Aethalops alecto*), also known as the grey fruit bat, is a species of megabat.

Pygmy fruit-eating bat
Pygmy fruit-eating bat
Dermanura phaeotis

The pygmy fruit-eating bat (*Dermanura phaeotis*) is a bat of the family Phyllostomidae. The specific name *phaeotis* is of Greek derivation, coming from the word *phaios* meaning dusky, referring to their dusky gray coloration.

Pygmy gerbil
Pygmy gerbil
Gerbillus henleyi

The pygmy gerbil (*Gerbillus henleyi*) is distributed mainly in Algeria to Israel and the Arabian Peninsula. It is also known as Henley's gerbil or pygmy dipodil. *Gerbillus henleyi* is a long-tailed small gerbil with a back coat that is dark...

Pygmy Hippo
Pygmy Hippo
Choeropsis liberiensis

The Pygmy hippo (*Choeropsis liberiensis* or *Hexaprotodon liberiensis*) is a small hippopotamid native to the forests and swamps of West Africa. It is one of only two extant species in the family Hippopotamidae, the other being its much larger...

Pygmy Hog
Pygmy Hog
Porcula salvania

The Pygmy hog (*Porcula salvania*) is a suid found in the Himalayas. It is the smallest and the rarest wild pig in the world and is now listed as Endangered on the IUCN Red List.

Pygmy killer whale
Pygmy killer whale
Feresa attenuata

The pygmy killer whale (*Feresa attenuata*) is a poorly known and rarely seen oceanic dolphin. It derives its common name from sharing some physical

characteristics with the orca also known as the killer whale. It is the smallest cetacean species...

Pygmy long-eared bat

Pygmy long-eared bat

Nyctophilus walkeri

The pygmy long-eared bat (*Nyctophilus walkeri*) is a vespertilionid bat, found in the north of the Australian continent. An insectivorous flying hunter, they are one of the tiniest mammals in Australia, weighing only a few grams and one or two...

Pygmy Marmoset

Pygmy Marmoset

Cebuella pygmaea

The pygmy marmoset, genus *Cebuella*, is a small genus of New World monkey native to rainforests of the western Amazon Basin in South America. It is notable for being the smallest monkey and one of the smallest primates in the world, at just over...

Pygmy mouse lemur

Pygmy mouse lemur

Microcebus myoxinus

The pygmy mouse lemur (*Microcebus myoxinus*), also known as Peters' mouse lemur or dormouse lemur, is a primate weighing only 43–55 g (1.5–1.9 oz); it is the second smallest of the mouse lemurs. Its dorsal side is a rufous-brown colour, and...

Pygmy Rabbit

Pygmy Rabbit

Brachylagus idahoensis

The pygmy rabbit (*Brachylagus idahoensis*) is a rabbit species native to the United States. It is also the only native rabbit species in North America to dig its own burrow. The pygmy rabbit differs significantly from species within either the...

Pygmy Right Whale

Pygmy Right Whale

Caperea marginata

The pygmy right whale (*Caperea marginata*) may be a member of the cetotheres, a family of baleen whales, which until 2012 were thought to be extinct; *C. marginata* has otherwise been considered the sole member of the family Neobalaenidae. First...

Pygmy ringtail possum

Pygmy ringtail possum

Pseudochirulus mayeri

The pygmy ringtail possum (*Pseudochirulus mayeri*) is a species of marsupial in the family Pseudocheiridae. It is found in the montane forest regions of Papua New Guinea and West Papua, Indonesia. They are "widespread along the Central...

Pygmy rock mouse

Pygmy rock mouse

Petromyscus collinus

The pygmy rock mouse (*Petromyscus collinus*) is a species of rodent in the family Nesomyidae. It is found in Angola, Namibia, and South Africa. Its natural habitat is subtropical or tropical dry shrubland.

Pygmy round-eared bat

Pygmy round-eared bat

Lophostoma brasiliense

The pygmy round-eared bat (*Lophostoma brasiliense*) is a bat species from South and Central America.

Pygmy scaly-tailed flying squirrel

Pygmy scaly-tailed flying squirrel

Idiurus zenkeri

The pygmy scaly-tailed flying squirrel (*Idiurus zenkeri*) is a species of rodent in the family Anomaluridae. It is found in Cameroon, Central African Republic, Republic of the Congo, Democratic Republic of the Congo, Equatorial Guinea, and...

Pygmy short-tailed opossum

Pygmy short-tailed opossum

Monodelphis kunsii

The pygmy short-tailed opossum, scientifically named *Monodelphis kunsii*, is an opossum species from South America. *M. kunsii* is a marsupial from the family Didelphidae of the order Didelphimorphia. Although it is a marsupial, it lacks the...

Pygmy shrew tenrec

Pygmy shrew tenrec

Microgale parvula

The pygmy shrew tenrec (*Microgale parvula*) is a species of placental mammal in the family Tenrecidae. It is endemic to Madagascar. Its natural habitats are subtropical or tropical moist lowland forests and subtropical or tropical moist montane...

Pygmy Slow Loris

Pygmy Slow Loris

Nycticebus pygmaeus

The pygmy slow loris (*Xanthonycticebus pygmaeus*) is a species of slow loris found east of the Mekong River in Vietnam, Laos, eastern Cambodia, and China. It occurs in a variety of forest habitats, including tropical dry forests, semi-evergreen,...

Pygmy sperm whale

Pygmy sperm whale

Kogia breviceps

The pygmy sperm whale (*Kogia breviceps*) is one of two extant species in the family Kogiidae in the sperm whale superfamily. They are not often sighted at sea, and most of what is known about them comes from the examination of stranded specimens.

Pygmy Spotted Skunk

Pygmy Spotted Skunk

Spilogale pygmaea

The Pygmy spotted skunk (*Spilogale pygmaea*) is a rare species of mammal that can be found only in Mexico. These animals are noted for their scent glands, which they use to deter predators.

Pygmy Tarsier

Pygmy Tarsier

Tarsius pumilus

The pygmy tarsier (*Tarsius pumilus*), also known as the mountain tarsier or the lesser spectral tarsier, is a nocturnal primate found in central Sulawesi, Indonesia, in an area with lower vegetative species diversity than the lowland tropical...

Pygmy three-toed sloth

Pygmy three-toed sloth

Bradypus pygmaeus

The pygmy three-toed sloth (*Bradypus pygmaeus*), also known as the monk sloth or dwarf sloth, is a sloth endemic to Isla Escudo de Veraguas, a small island off the Caribbean coast of Panama. The species was first described by Robert P. Anderson...

Pygmy treeshrew

Pygmy treeshrew

Tupaia minor

The pygmy treeshrew (*Tupaia minor*) is a treeshrew species within the family Tupaiidae. It is native to Thailand, Malaysia and Indonesia. The generic name is derived from the Malay word tupai meaning squirrel or small animals that resemble squirrels.

Pyrenean Chamois

Pyrenean Chamois

Rupicapra pyrenaica

The Pyrenean chamois (*Rupicapra pyrenaica*) is a goat-antelope that lives in the Pyrenees, Cantabrian Mountains and Apennine Mountains. It is one of the two species of the genus *Rupicapra*, the other being the chamois, *Rupicapra rupicapra*.

Pyrenean Desman

Pyrenean Desman

Galemys pyrenaicus

The Pyrenean desman also called Iberian desman (*Galemys pyrenaicus*) is a small semiaquatic, globally threatened mammal related to moles and shrews, and, along with the Russian desman (*Desmana moschata*) , is one of the two extant members of the...

Pyrenean ibex

Pyrenean ibex

Capra pyrenaica pyrenaica

The Pyrenean ibex (*Capra pyrenaica pyrenaica*), Aragonese and Spanish common name bucardo, Basque common name bukardo, Catalan common name herc and French common name bouquetin, was one of the four subspecies of the Iberian ibex or Iberian wild...

Qazvin vole

Qazvin vole

Microtus qazvinensis

The Qazvin vole (*Microtus qazvinensis*) is a species of rodent in the family Cricetidae. It is found in Iran and is normally given as either part of the *Microtus* or *Sumeriomys* subgenus. It is a close relative of *M. guentheri* distinguished by...

Qinling panda

Qinling panda

Ailuropoda melanoleuca qinlingensis

The Qinling panda (*Ailuropoda melanoleuca qinlingensis*) is a subspecies of the giant panda, discovered in the 1960s but not recognized as a subspecies until 2005. Besides the nominate subspecies, it is the first giant panda subspecies to be...

Quaestor horticudo

Quaestor horticudo

Oxymycterus quaestor

The quaestor horticudo (*Oxymycterus quaestor*) is a species of rodent in the family Cricetidae. It is found in southeastern Brazil and northeastern Argentina, where it lives in forest and moist and dry scrub.

Quagga

Quagga

Equus quagga quagga

The quagga (*Equus quagga quagga*) is a subspecies of the Plains zebra that was endemic to South Africa until it was hunted to extinction in the late 19th century by European settler-colonists. Some were taken to zoos in Europe, but breeding...

Quechuan mouse opossum

Quechuan mouse opossum
Marmosa macrotarsus

The Quechuan mouse opossum (*Marmosa macrotarsus*) is a South American opossum species of the family Didelphidae. It is known from two areas of montane forest on the eastern slopes of the Andes in Peru, at altitudes from 300 to 2700 m. The type...

Queen Charlotte Islands caribou
Queen Charlotte Islands caribou
Rangifer tarandus dawsoni

The Queen Charlotte Islands caribou or Dawson's caribou (*Rangifer tarandus dawsoni*) is an extinct subspecies of the reindeer (or the caribou in North America) that once lived on Graham Island, the largest of the Haida Gwaii islands in British...

Quokka
Quokka
Setonix brachyurus

The quokka (*Setonix brachyurus*), is a small macropod about the size of a domestic cat. In 1696, Dutch explorer Willem de Vlamingh mistook these animals for giant rats, and renamed the Wadjemup island 't Eylandt 't Rottenest, which means "the rat..."

Raccoon
Raccoon
Procyon lotor

The raccoon (*Procyon lotor*), sometimes called the common raccoon to distinguish it from other species, is a mammal native to North America. It is the largest of the procyonid family. Three of the raccoon's most distinctive features are its...

Raccoon Dog
Raccoon Dog
Nyctereutes procyonoides

The Common raccoon dog (*Nyctereutes procyonoides*) is a canid indigenous to mainland East Asia and northern Vietnam. Despite its name, its closest relatives are the true foxes, and not the American raccoons. Among the Canidae, the Raccoon dog...

Radde's shrew
Radde's shrew
Sorex raddei

Radde's shrew (*Sorex raddei*) is a species of mammal in the family Soricidae. It is found in Armenia, Azerbaijan, Georgia, Russia, and Turkey.

Raffray's bandicoot
Raffray's bandicoot
Peroryctes raffrayana

Raffray's bandicoot (*Peroryctes raffrayana*) is a species of marsupial in the family Peroryctidae. It is found in Indonesia and Papua New Guinea. Its natural habitat is subtropical or tropical dry forests.

Raffray's sheath-tailed bat
Raffray's sheath-tailed bat
Emballonura raffrayana

Raffray's sheath-tailed bat (*Emballonura raffrayana*) is a species of sac-winged bat in the family Emballonuridae. It is found in eastern Indonesia (including Western New Guinea), Papua New Guinea, and Solomon Islands.

Rafinesque's big-eared bat
Rafinesque's big-eared bat

Corynorhinus rafinesquii

Rafinesque's big-eared bat (*Corynorhinus rafinesquii*), sometimes known as the southeastern big-eared bat, is a species of vesper bat native to the southeastern United States.

Rahm's brush-furred rat

Rahm's brush-furred rat

Lophuromys rahmi

Rahm's brush-furred rat (*Lophuromys rahmi*) is a species of rodent in the family Muridae. It is found in Democratic Republic of the Congo, Rwanda, and Uganda. Its natural habitat is subtropical or tropical moist montane forests. It is threatened by...

Railer bat

Railer bat

Mops thersites

The Railer bat (*Mops thersites*) is a species of bat in the family Molossidae. It is found in Cameroon, Central African Republic, Republic of the Congo, Democratic Republic of the Congo, Ivory Coast, Equatorial Guinea, Gabon, Ghana, Guinea,...

Rain forest shrew

Rain forest shrew

Sylvisorex pluvialis

The rain forest shrew (*Sylvisorex pluvialis*) is a species of mammal in the family Soricidae endemic to Cameroon. Its natural habitat is subtropical or tropical moist lowland forest. It is known only from its type locality, and from Kongana,...

Rainey's shrew

Rainey's shrew

Crocidura raineyi

Rainey's shrew (*Crocidura raineyi*) is a species of mammal in the family Soricidae. It is endemic to Kenya. Its natural habitat is subtropical or tropical moist lowland forests. It is threatened by habitat loss.

Rajah spiny rat

Rajah spiny rat

Maxomys rajah

The Rajah spiny rat (*Maxomys rajah*) also known as the brown spiny rat is endemic to Thailand and Malaysia, Sumatra, Borneo, and adjacent islands (Payne et al., 1985).

Rakali

Rakali

Hydromys chrysogaster

The rakali (*Hydromys chrysogaster*) is an Australian native rodent first described in 1804. Having adapted to and colonized a unique niche of a semiaquatic and nocturnal lifestyle, this rakali lives in burrows on the banks of rivers, lakes and...

Ramanantsoavana's woolly lemur

Ramanantsoavana's woolly lemur

Avahi ramanantsoavanai

Ramanantsoavana's woolly lemur (*Avahi ramanantsoavanai*), also known as Ramanantsoavana's avahi or the Manombo woolly lemur, is a species of woolly lemur native to southeastern Madagascar. It weighs about 1 kg. It was originally considered a...

Randrianasolo's sportive lemur

Randrianasolo's sportive lemur

Lepilemur randrianasoloi

Randrianasolo's sportive lemur (*Lepilemur randrianasoloi*), or the Bemaraha sportive lemur, is a sportive lemur endemic to Madagascar. It has a total length of about 49 to 56 cm (19 to 22 in), of which 21–26 cm (8.3–10.2 in) are tail....

Ranee mouse

Ranee mouse

Haeromys margarettae

The ranee mouse (*Haeromys margarettae*) is a species of rodent in the family Muridae. It is restricted to the island of Borneo, in the provinces of Sarawak (Malaysia) and Sabah (Malaysia). Its natural habitat is subtropical or tropical dry forests.

Ratanaworabhan's fruit bat

Ratanaworabhan's fruit bat

Megaerops niphanae

The Ratanaworabhan's fruit bat (*Megaerops niphanae*) is a species of megabat in the family Pteropodidae. It is found in Bhutan, India, Thailand, and Vietnam.

Recife broad-nosed bat

Recife broad-nosed bat

Platyrrhinus recifinus

The Recife broad-nosed bat, *Platyrrhinus recifinus*, is a species of bat from South America. It is named for the city of Recife in Brazil, where it was first recorded by Oldfield Thomas in 1901. Previously considered "Vulnerable", the Recife...

Reclusive ringtail possum

Reclusive ringtail possum

Pseudocheirops coronatus

The reclusive ringtail possum or Monk ringtail possum (*Pseudocheirops coronatus*) is a species of marsupial in the family Pseudocheiridae. It is endemic to the Arfak Mountains in the Vogelkop Peninsula of West Papua, Indonesia.

Red acouchi

Red acouchi

Myoprocta acouchy

The red acouchi, *Myoprocta acouchy*, is a species of rodent in the acouchi genus, part of the family Dasyproctidae, from South America. It is found in the Guyanese subregion of the Amazon biome, including Guyana, Suriname, French Guiana, and...

Red and white giant flying squirrel

Red and white giant flying squirrel

Petaurista alborufus

The red and white giant flying squirrel (*Petaurista alborufus*) is a species of rodent in the family Sciuridae. It is a very large, dark rufous-red, buff and white flying squirrel found in forests at altitudes of 800–3,500 m (2,600–11,500 ft) in...

Red Brocket

Red Brocket

Mazama americana

The red brocket (*Mazama americana*) is a species of brocket deer from forests in South America, ranging from northern Argentina to Colombia and the Guianas. It also occurs on the Caribbean island of Trinidad (it also occurred on the island of...

Red bush squirrel

Red bush squirrel

Paraxerus palliatus

The red bush squirrel or red-bellied coast squirrel (*Paraxerus palliatus*) is a species of rodent in the family Sciuridae, which is found in bush and forest along the eastern seaboard of Africa. It is closely related to *P. lucifer*, *P.*

vexillarius...

Red climbing mouse

Red climbing mouse

Vernaya fulva

The red climbing mouse (*Vernaya fulva*), also known as Vernay's climbing mouse, is a species of rodent in the family Muridae. It is named after explorer and antiques dealer, Arthur Vernay, who sponsored the expedition during which the animal was...

Red Deer

Red Deer

Cervus elaphus

The Red deer (*Cervus elaphus*) is one of the largest deer species. It is closely related and slightly larger American elk or wapiti, native to North America and eastern parts of Asia, had been regarded as a subspecies of red deer, but recently it...

Red forest duiker

Red forest duiker

Cephalophus natalensis

The red forest duiker, Natal duiker, or Natal red duiker (*Cephalophus natalensis*) is a small antelope found in central to southern Africa. It is one of 22 extant species form the subfamily Cephalophinae. While the red forest duiker is very...

Red Fox

Red Fox

Vulpes vulpes

The Red fox (*Vulpes vulpes*) is the largest of the true foxes and one of the most widely distributed members of the order Carnivora, being present across the entire Northern Hemisphere. Its range has increased alongside human expansion, having...

Red fruit bat

Red fruit bat

Stenoderma rufum

The red fruit bat or red fig-eating bat (*Stenoderma rufum*) is a species of bat in the family Phyllostomidae, in the monotypic genus *Stenoderma*. It is found in Puerto Rico and the U.S. Virgin Islands.

Red gazelle

Red gazelle

Eudorcas rufina

The red gazelle (*Eudorcas rufina*) is an extinct species or subspecies of gazelle. It was formerly considered a member of the genus *Gazella* within the subgenus *Eudorcas* before *Eudorcas* was promoted to a full genus. It is thought to have lived in...

Red Giant Flying Squirrel

Red Giant Flying Squirrel

Petaurista petaurista

The red giant flying squirrel or common giant flying squirrel (*Petaurista petaurista*) is a species of rodent in the family Sciuridae (squirrels). It is found in a wide variety of forest-types, plantations and more open habitats with scattered...

Red goral

Red goral

Naemorhedus baileyi

The red goral (*Naemorhedus baileyi*) is a species of even-toed ungulate in the subfamily Caprinae in the family Bovidae. It is found in India, Tibet and Myanmar. Its natural habitats are seasonal mountainous areas 1,000 to 2,000 meters above sea...

Red hartebeest

Red hartebeest

Alcelaphus buselaphus caama

The red hartebeest (*Alcelaphus buselaphus caama*) or Cape hartebeest is a subspecies of the hartebeest found in Southern Africa. More than 130,000 individuals live in the wild. The red hartebeest is closely related to the tsessebe and the...

Red hocicudo

Red hocicudo

Oxymycterus rufus

The red hocicudo (*Oxymycterus rufus*) is a rodent species from South America. It is found in Argentina, Brazil and Uruguay.

Red Howler Monkey

Red Howler Monkey

Alouatta seniculus

The Colombian red howler or Venezuelan red howler (*Alouatta seniculus*) is a South American species of howler monkey, a type of New World monkey, found in the western Amazon Basin in Venezuela, Colombia, Ecuador, Peru and Brazil. The population...

Red Kangaroo

Red Kangaroo

Macropus rufus

The Red kangaroo (*Osphranter rufus*) is the largest of all kangaroos, the largest terrestrial mammal native to Australia, and the largest extant marsupial. It is an iconic symbol of Australia and like all Australian wildlife, the Red kangaroo is...

Red Lemur

Red Lemur

Eulemur rufus

The red lemur (*Eulemur rufus*), also known as the rufous brown lemur or northern red-fronted lemur, is a species of lemur from Madagascar. Until 2001, the species *E. rufus* was considered a subspecies of the common brown lemur, *E. fulvus*, after...

Red mouse opossum

Red mouse opossum

Marmosa rubra

The red mouse opossum (*Marmosa rubra*) is a South American marsupial of the family Didelphidae. Its range includes eastern Ecuador and Peru.

Red myotis

Red myotis

Myotis ruber

The red myotis (*Myotis ruber*) is a vesper bat species found in Argentina, Brazil, Paraguay and Uruguay.

Red Panda

Red Panda

Ailurus fulgens

The Red panda (*Ailurus fulgens*) is a small mammal native to the eastern Himalayas and southwestern China. It is not closely related to the Giant panda, which is a bear, though both possess elongated wrist bones or "false thumbs" used for

grasping...
Red River Hog
Red River Hog
Potamochoerus porcus

The Red river hog (*Potamochoerus porcus*) is a wild member of the pig family that lives in Africa. It is rarely seen away from rainforests, and generally prefers areas near rivers or swamps.

Red rock rat
Red rock rat
Aethomys chrysophilus

The red rock rat, or red veld rat, (*Aethomys chrysophilus*) is a species of rodent in the family Muridae native to southern Africa.

Red Ruffed Lemur
Red Ruffed Lemur
Varecia rubra

The red ruffed lemur (*Varecia rubra*) is one of two species in the genus *Varecia*, the ruffed lemurs; the other is the black-and-white ruffed lemur (*Varecia variegata*). Like all lemurs, it is native to Madagascar. It occurs only in the...

Red serow
Red serow
Capricornis rubidus

The red serow (*Capricornis rubidus*) is a goat-antelope thought to be native to southern Bangladesh and northern Myanmar. It has been sometimes been considered a subspecies of *C. sumatraensis*. In the northeastern part of India, the red serow...

Red Slender Loris
Red Slender Loris
Loris tardigradus

The red slender loris (*Loris tardigradus*) is a small, nocturnal strepsirrhine primate native to the rainforests of Sri Lanka. This is No. 6 of the 10 focal species and No. 22 of the 100 EDGE mammal species worldwide considered the most...

Red spiny rat
Red spiny rat
Maxomys surifer

The red spiny rat (*Maxomys surifer*) is a rodent species in the family Muridae that inhabits forests in Myanmar, Thailand, Malaysia, Cambodia, Laos, Vietnam and on Java, Borneo and Sumatra islands. In China, it has been recorded only in Yunnan...

Red Squirrel
Red Squirrel
Sciurus vulgaris

The Red squirrel (*Sciurus vulgaris*) is a species of tree squirrel in the genus *Sciurus* common throughout Europe and Asia. It lives in trees and its long tail helps the squirrel to balance and steer when jumping from tree to tree and running along...

Red three-striped opossum
Red three-striped opossum
Monodelphis umbristriata

The red three-striped opossum (*Monodelphis umbristriata*) is an opossum species from South America. It is found in Brazil.

Red tree rat
Red tree rat
Pithecheir melanurus

The red tree rat (*Pithecheirus melanurus*) is a species of rodent in the family Muridae. It is found only in western Java, Indonesia.

Red tree vole

Red tree vole

Arborimus longicaudus

The red tree vole (*Arborimus longicaudus*) is a rodent of the Pacific Northwest, found in the US states of Oregon and California. They were formerly known as *Phenacomys longicaudus* and have also been called the red tree mouse.

Red wolf

Red wolf

Canis lupus rufus

The red wolf (*Canis rufus*) is a canine native to the southeastern United States. Its size is intermediate between the coyote (*Canis latrans*) and gray wolf (*Canis lupus*). The red wolf's taxonomic classification as being a separate species, a...

Red-backed bearded saki

Red-backed bearded saki

Chiropotes chiropotes

The red-backed bearded saki (*Chiropotes chiropotes*) is a New-World monkey, from South America. It is a species of bearded saki.

Red-Bellied Lemur

Red-Bellied Lemur

Eulemur rubriventer

The red-bellied lemur (*Eulemur rubriventer*) is a medium-sized strepsirrhine primate with a luxuriant chestnut brown coat. This lemur is endemic to eastern Madagascan rainforests and is distinguished by patches of white skin below the eyes,...

Red-bellied marsupial shrew

Red-bellied marsupial shrew

Phascolosorex doriae

The red-bellied dasyure or red-bellied marsupial shrew (*Phascolosorex doriae*) is a species of marsupial in the family Dasyuridae endemic to West Papua, Indonesia. Its natural habitat is subtropical or tropical dry forests.

Red-bellied mosaic-tailed rat

Red-bellied mosaic-tailed rat

Protochromys fellowsi

The red-bellied mosaic-tailed rat (*Protochromys fellowsi*) is a species of rodent in the family Muridae. It is the only species in the genus *Protochromys*. It is found only in Papua New Guinea.

Red-bellied squirrel

Red-bellied squirrel

Rubrisciurus rubriventer

The red-bellied squirrel or Sulawesi giant squirrel (*Rubrisciurus rubriventer*) is a species of squirrel. Until recently, it was described as a species in the genus *Callosciurus*, but since the 1990s it is generally placed in its own genus...

Red-bellied titi

Red-bellied titi

Plecturocebus moloch

The red-bellied titi or dusky titi (*Plecturocebus moloch*) is a species of titi, a type of New World monkey, endemic to Brazil. It lives in forests and thickets. It has a rounded head and a thick, soft coat and frequently adopts a characteristic...

Red-brown pipistrelle
Red-brown pipistrelle
Hypsugo kitcheneri

The red-brown pipistrelle (*Hypsugo kitcheneri*) is a species of vesper bat in the family Vespertilionidae. It is found in Indonesia and possibly Malaysia.

Red-capped tamarin
Red-capped tamarin
Saguinus pileatus

The red-capped tamarin, *Saguinus mystax pileatus*, is subspecies of moustached tamarin from South America. They are found in Brazil, on the eastern margin of the Tefé and Coari rivers. Previously recognised as a separate species, *Saguinus pilatus*,...

Red-cheeked dunnart
Red-cheeked dunnart
Sminthopsis virginiae

The red-cheeked dunnart (*Sminthopsis virginiae*) is so called because of the distinctive red hair on its cheek. It is an Australasian marsupial. Its total length is 167–270 mm; its average body length is 80–135 mm with a tail of 87–135 mm. Ear...

Red-cheeked flying squirrel
Red-cheeked flying squirrel
Hylopetes spadiceus

The red-cheeked flying squirrel (*Hylopetes spadiceus*) is a species of rodent in the family Sciuridae. It is found in Indonesia, Malaysia, Myanmar, Singapore, Thailand, and Vietnam.

Red-cheeked ground squirrel
Red-cheeked ground squirrel
Spermophilus erythrogenys

The red-cheeked ground squirrel (*Spermophilus erythrogenys*) is a species of rodent in the family Sciuridae. It is commonly referred to as the red-cheeked ground souslik and there are several recognized subspecies. It is found in central Asia....

Red-cheeked rope squirrel
Red-cheeked rope squirrel
Funisciurus leucogenys

The red-cheeked rope squirrel (*Funisciurus leucogenys*) is a species of rodent in the family Sciuridae. It is found in Benin, Cameroon, Central African Republic, Equatorial Guinea, Ghana, Nigeria, and Togo. Its natural habitats are subtropical or...

Red-Crested Tree-Rat
Red-Crested Tree-Rat
Santamartamys rufodorsalis

The Red-crested tree-rat (*Santamartamys rufodorsalis*) is a species of tree-rat found in South America. It is nocturnal and is believed to feed on plant matter. It is a very rare and poorly known small rodent known only from three specimens.

Red-eared guenon
Red-eared guenon
Cercopithecus erythrotis

The red-eared guenon (*Cercopithecus erythrotis*), also called red-eared monkey, or russet-eared guenon is a primate species in the family Cercopithecidae. It is native to subtropical and tropical moist lowland forests in Cameroon, Equatorial...

Red-Faced Spider Monkey
Red-Faced Spider Monkey
Ateles paniscus

The red-faced spider monkey (*Ateles paniscus*) also known as the Guiana spider monkey or red-faced black spider monkey, is a species of spider monkey found in the rain forests in northern South America. The species faces issues with hunting and...

Red-Flanked Duiker
Red-Flanked Duiker
Cephalophus rufilatus

The red-flanked duiker (*Cephalophus rufilatus*) is a species of small antelope found in western and central Africa in countries as far apart as Senegal and Sudan. Red-flanked duikers grow to almost 15 in (35 cm) in height and weigh up to 31 lb...

Red-Fronted Gazelle
Red-Fronted Gazelle
Eudorcas rufifrons

The Red-fronted gazelle (*Eudorcas rufifrons*) is widely but unevenly distributed gazelle across the middle of Africa. It was first described by British zoologist John Edward Gray.

Red-Fronted Lemur
Red-Fronted Lemur
Eulemur rufifrons

The red-fronted lemur (*Eulemur rufifrons*), also known as the red-fronted brown lemur or southern red-fronted brown lemur, is a species of lemur from Madagascar. Until 2001, it was considered a subspecies of the common brown lemur, *E. fulvus*. In...

Red-handed howler
Red-handed howler
Alouatta belzebul

The red-handed howler (*Alouatta belzebul*) is a vulnerable species of howler monkey, a type of New World monkey. It is endemic to Brazil, found in the southeastern Amazon and disjunctly in the Atlantic Forest between Rio Grande do Norte and Sergipe.

Red-Handed Tamarin
Red-Handed Tamarin
Saguinus midas

Red-handed tamarins (*Saguinus midas*) are New World monkeys named for the contrasting reddish-orange hair on their feet and hands.

Red-headed titi
Red-headed titi
Cheracebus regulus

The red-headed titi (*Cheracebus regulus*) is a species of titi, a type of New World monkey, endemic to Brazil. It was originally described as *Callicebus regulus* in 1927.

Red-hipped squirrel
Red-hipped squirrel
Dremomys pyrrhomerus

The red-hipped squirrel (*Dremomys pyrrhomerus*) is a species of rodent in the family Sciuridae. It is found in both China and Vietnam.

Red-Legged Pademelon
Red-Legged Pademelon
Thylogale stigmatica

The red-legged pademelon (*Thylogale stigmatica*) is a species of small macropod found on the northeastern coast of Australia and in New Guinea. In Australia it has a scattered distribution from the tip of Cape York Peninsula in Queensland to...
Red-legged sun squirrel
Red-legged sun squirrel
Heliosciurus rufobrachium

The red-legged sun squirrel (*Heliosciurus rufobrachium*) is a species of rodent in the family *Sciuridae*, also commonly known as the isabelline red-legged sun squirrel. It is native to tropical western and central Africa where its range extends...
Red-Necked Pademelon
Red-Necked Pademelon
Thylogale thetis

The red-necked pademelon (*Thylogale thetis*) is a forest-dwelling marsupial living in the eastern coastal region of Australia.
Red-Necked Wallaby
Red-Necked Wallaby
Macropus rufogriseus

The Red-necked wallaby (*Notamacropus rufogriseus*) is a medium-sized macropod marsupial, common in the more temperate and fertile parts of eastern Australia, including Tasmania. Its name comes from the reddish fur on its shoulders and nape.
Red-nosed mouse
Red-nosed mouse
Wiedomys pyrrhorhinos

The red-nosed mouse (*Wiedomys pyrrhorhinos*) is an arboreal rodent species endemic to Brazil. It is found in caatinga and cerrado habitat in southeast Brazil.
Red-Rumped Agouti
Red-Rumped Agouti
Dasyprocta leporina

Red-rumped agoutis (*Dasyprocta leporina*) are rodents from South America. They are related to guinea pigs and look quite similar, but are larger and have longer legs adapted for fast running.
Red-Shanked Douc
Red-Shanked Douc
Pygathrix nemaeus

The Red-shanked douc (*Pygathrix nemaeus*) is a diurnal Old World monkey. They are the most colorful monkeys among all species of primates. Red-shanked doucs live in fission-fusion, multilevel societies, and communicate with each other using...
Red-tailed chipmunk
Red-tailed chipmunk
Neotamias ruficaudus

The red-tailed chipmunk (*Neotamias ruficaudus*) is a species of rodent in the family *Sciuridae*. It is found in Alberta and British Columbia in Canada and Montana, Idaho and Washington in the United States.
Red-Tailed Monkey
Red-Tailed Monkey
Cercopithecus ascanius

The red-tailed monkey, also known as the black-cheeked white-nosed monkey, red-tailed guenon, redtail monkey, or Schmidt's guenon (*Cercopithecus ascanius*) is a species of primate in the family *Cercopithecidae*. It is found in Angola,

Cameroon,...

Red-tailed phascogale

Red-tailed phascogale

Phascogale calura

The red-tailed phascogale (*Phascogale calura*), also known as the red-tailed wambenger or red-tailed mousesack, is a small carnivorous marsupial found in Central and Western Australia. It is closely related to the brush-tailed phascogale...

Red-tailed sportive lemur

Red-tailed sportive lemur

Lepilemur ruficaudatus

The red-tailed sportive lemur (*Lepilemur ruficaudatus*), or red-tailed weasel lemur, is native to Madagascar like all lemurs. It is a nocturnal species feeding largely on leaves, though they also eat some fruit. Individuals weigh around 800 g...

Red-Tailed Squirrel

Red-Tailed Squirrel

Sciurus granatensis

The red-tailed squirrel (*Sciurus granatensis*) is a species of tree squirrel distributed from southern Central America to northern South America.

Red-throated squirrel

Red-throated squirrel

Dremomys gularis

The red-throated squirrel (*Dremomys gularis*) is a rodent of the family *Sciuridae*. This species of rodent is distributed in parts of southeastern Asia, in areas of the Red River Valley of northern Vietnam and southern central Yunnan in China. The...

Reddish tuco-tuco

Reddish tuco-tuco

Ctenomys frater

The reddish tuco-tuco (*Ctenomys frater*) is a species of rodent in the family *Ctenomyidae*. Five subspecies have been recognized, some formerly designated as separate species. It is found in Argentina and Bolivia at altitudes from 600 to 4,500....

Reddish-gray mouse lemur

Reddish-gray mouse lemur

Microcebus griseorufus

The reddish-gray mouse lemur (*Microcebus griseorufus*) also known as the gray-brown mouse lemur or rufous-gray mouse lemur, is found in Western Madagascar in the region around Beza Mahafaly Reserve, north to Lamboharana.

Reddish-gray musk shrew

Reddish-gray musk shrew

Crocidura cyanea

The reddish-gray musk shrew (*Crocidura cyanea*) is a species of mammal in the family *Soricidae*. It is found in Angola, Botswana, Democratic Republic of the Congo, Eswatini, Lesotho, Malawi, Mozambique, Namibia, Nigeria, South Africa, Tanzania,...

Reed vole

Reed vole

Microtus fortis

The reed vole (*Microtus fortis*) is a species of vole. It is found in northern and

central Eurasia, including northern China and the Korean Peninsula. This species is somewhat larger and longer-tailed than most other voles.

Reeves's Muntjac

Reeves's Muntjac

Muntiacus reevesi

Reeves's muntjac (*Muntiacus reevesi*) is a muntjac species native to East Asia. It takes its name from John Reeves, a naturalist employed by the British East India Company in the 19th century.

Reig's grass mouse

Reig's grass mouse

Akodon reigi

Reig's grass mouse (*Akodon reigi*) is a South American rodent species found in Brazil and Uruguay. It is named after Argentine biologist Osvaldo Reig (1929–1992).

Reig's montane mouse

Reig's montane mouse

Aepeomys reigi

Reig's montane mouse (*Aepeomys reigi*) is a species of rodent in the family Cricetidae. It is found only in Venezuela. It is named after Argentine biologist Osvaldo Reig (1929–1992).

Reig's opossum

Reig's opossum

Monodelphis reigi

Reig's opossum (*Monodelphis reigi*) is a South American opossum species of the family Didelphidae, discovered in 2004. It is named after Argentine biologist Osvaldo Reig (1929–1992). It was initially found in montane forest in Canaima National...

Reig's tuco-tuco

Reig's tuco-tuco

Ctenomys osvaldoreigi

Reig's tuco-tuco (*Ctenomys osvaldoreigi*) is a species of rodent in the family Ctenomyidae. It is endemic to central Argentina, where it is known only from a grassland location in Cordoba Province at an elevation above 2000 m in the Sierras...

Reindeer

Reindeer

Rangifer tarandus

The reindeer (*Rangifer tarandus*), also known as the caribou in North America, is a species of deer with circumpolar distribution. Reindeer occur in both migratory and sedentary populations, and their herd sizes vary greatly in different regions....

Remy's pygmy shrew

Remy's pygmy shrew

Suncus remyi

Remy's pygmy shrew (*Suncus remyi*) is a species of mammal in the family Soricidae. It is found in Cameroon, the Central African Republic, the Republic of the Congo, and Gabon. Its natural habitat is subtropical or tropical moist lowland forests....

Rendall's serotine

Rendall's serotine

Neoromicia rendalli

Rendall's serotine (*Pseudoromicia rendalli*) is a species of vesper bat. It is found in Benin, Botswana, Burkina Faso, Cameroon, Central African Republic, Chad, Republic of the Congo, Democratic Republic of the Congo, Gambia, Ghana, Kenya,

Malawi,...
Rennell flying fox
Rennell flying fox
Pteropus rennelli

The Rennell flying fox (*Pteropus rennelli*) is a species of flying fox found in the Solomon Islands. It is an endangered species risking extinction.

Reticulated Giraffe
Reticulated Giraffe
Giraffa camelopardalis reticulata

The Reticulated giraffe (*Giraffa camelopardalis reticulata*) is a subspecies of giraffe native to the Horn of Africa. The Reticulated giraffe was described and given its binomial name by British zoologist William Edward de Winton in 1899, however...

Rhabdomys dilectus
Rhabdomys dilectus
Rhabdomys dilectus

The mesic four-striped grass rat (*Rhabdomys dilectus*) is a species of rodent in the family Muridae.

Rhesus Macaque
Rhesus Macaque
Macaca mulatta

The Rhesus macaque (*Macaca mulatta*) is a species of Old World monkey. It lives both in trees and on the ground. It mainly eats fruit, but will also include 100 different plant species in its diet. It spends most of its day feeding and resting,...

Rhim Gazelle
Rhim Gazelle
Gazella leptoceros

The Rhim gazelle (*Gazella leptoceros*) is an endangered species of antelope well adapted to desert life. It is considered an endangered species.

Rhinoceros sondaicus annamiticus
Rhinoceros sondaicus annamiticus
Rhinoceros sondaicus annamiticus

The Vietnamese Javan Rhino (*Rhinoceros sondaicus annamiticus*) also known as the Indo-Chinese Javan Rhino is a possibly extinct subspecies of the Javan rhinoceros that formerly lived in Laos, Cambodia, Thailand and Malaysia, and may still...

Rhinolophus belligerator
Rhinolophus belligerator
Rhinolophus belligerator

The Poso horseshoe bat or belligerent horseshoe bat (*Rhinolophus belligerator*) is an endangered species of horseshoe bat found only on Sulawesi Island, which is part of Indonesia.

Rhinolophus hilli
Rhinolophus hilli
Rhinolophus hilli

Rhinolophus hilli, Hill's horseshoe bat, is a species of bat in the family Rhinolophidae. It is endemic to Rwanda. Its natural habitats are subtropical or tropical moist montane forests, caves, and subterranean habitats (other than caves). In...

Rhinolophus luctoides
Rhinolophus luctoides

Rhinolophus luctoides

The Selangor woolly horseshoe bat or luctus-like horseshoe bat (Rhinolophus luctoides) is a bat species of the family Rhinolophidae endemic to Malaysia. Populations of this species were previously classified within the woolly horseshoe bat (R....

Rhinolophus mcintyreii

Rhinolophus mcintyreii

Rhinolophus mcintyreii

McIntyre's horseshoe bat (Rhinolophus mcintyreii) is a species of horseshoe bat that is endemic to Papua New Guinea.

Rhinolophus microglobosus

Rhinolophus microglobosus

Rhinolophus microglobosus

Rhinolophus microglobosus is a species of horseshoe bat found in Southeast Asia.

Rhinolophus monticolus

Rhinolophus monticolus

Rhinolophus monticolus

Rhinolophus monticolus or Mountain horseshoe bat is a species of bat in the family Rhinolophidae. It is found from Greater Mekong in Southeast Asia.

Rhinolophus proconsulis

Rhinolophus proconsulis

Rhinolophus proconsulis

The Bornean woolly horseshoe bat or Proconsul's horseshoe bat (Rhinolophus proconsulis) is an endangered species of horseshoe bat found in Malaysia. Though it was discovered in 1959, it was not recognized as a distinct species until 2013.

Rhinolophus sedulus

Rhinolophus sedulus

Rhinolophus sedulus

The lesser woolly horseshoe bat (Rhinolophus sedulus) is a species of bat in the family Rhinolophidae. It is found in Brunei, Indonesia, and Malaysia. It is assessed as near-threatened by the IUCN.

Rhinolophus tatar

Rhinolophus tatar

Rhinolophus tatar

The Sulawesi broad-eared horseshoe bat or Tatar horseshoe bat (Rhinolophus tatar) is a species of horseshoe bat found in Sulawesi, Indonesia.

Rhinolophus xinanzhongguoensis

Rhinolophus xinanzhongguoensis

Rhinolophus xinanzhongguoensis

Rhinolophus xinanzhongguoensis, the wedge-sellaed horseshoe bat or southwestern China horseshoe bat, is a species of horseshoe bat from China.

Rhoads's Oldfield mouse

Rhoads's Oldfield mouse

Thomasomys rhoadsi

Rhoads's Oldfield mouse (Thomasomys rhoadsi) is a species of rodent in the family Cricetidae. It is found only in Ecuador.

Rhynchomys labo

Rhynchomys labo

Rhynchomys labo

The Labo shrew rat (*Rhynchomys labo*) is a species of shrewlike rat in the subfamily Murinae. It was discovered at elevations above 1250 m on Mt. Labo of the Bicol Peninsula of Luzon in the Philippines and described in 2019.

Rhynchomys mingan

Rhynchomys mingan

Rhynchomys mingan

The Mingan shrew rat (*Rhynchomys mingan*) is a species of shrewlike rat in the subfamily Murinae. It was discovered at elevations above 1450 m on Mt. Mingan of the central Sierra Madre of Luzon in the Philippines and described in 2019.

Riama mosaic-tailed rat

Riama mosaic-tailed rat

Melomys howi

The Riama mosaic-tailed rat (*Melomys howi*), is a species of rodent in the family Muridae. It is found only on the island of Riama in Indonesia, a small island to the west of Selaru in the Tanimbar Islands. Very little is known about this...

Ribbon Seal

Ribbon Seal

Histiophoca fasciata

The ribbon seal (*Histiophoca fasciata*) is a medium-sized pinniped from the true seal family (Phocidae). A seasonally ice-bound species, it is found in the Arctic and Subarctic regions of the North Pacific Ocean, notably in the Bering Sea and...

Ribboned rope squirrel

Ribboned rope squirrel

Funisciurus lemniscatus

The ribboned rope squirrel (*Funisciurus lemniscatus*) is a species of rodent in the family Sciuridae. It is found in Cameroon, Republic of the Congo, Democratic Republic of the Congo, and Equatorial Guinea. Its natural habitat is subtropical or...

Ricefield Rat

Ricefield Rat

Rattus argentiventer

The Ricefield rat (*Rattus argentiventer*) is a medium-sized species of rat found throughout Southeast Asia. They live in groups and are usually active during the nighttime hours.

Richardson's collared lemming

Richardson's collared lemming

Dicrostonyx richardsoni

The Richardson's collared lemming (*Dicrostonyx richardsoni*) is a small North American lemming. At one time, they were considered to be a subspecies of the Arctic lemming, *Dicrostonyx torquatus*. Some sources believe they are a subspecies of the...

Richardson's Ground Squirrel

Richardson's Ground Squirrel

Urocitellus richardsonii

Richardson's ground squirrel (*Urocitellus richardsonii*), also known as the dakrat or flickertail, is a North American ground squirrel in the genus *Urocitellus*. Like a number of other ground squirrels, they are sometimes called prairie dogs or...

Richmond's squirrel

Richmond's squirrel

Sciurus richmondi

Richmond's squirrel (*Sciurus richmondi*) is a poorly known tree squirrel in the

genus *Sciurus* endemic to Nicaragua, which is likely a synonym of the red-tailed squirrel (*Sciurus granatensis*). It is locally known as the ardilla del rama.
Rickart's dyak fruit bat
Rickart's dyak fruit bat
Dyacopterus rickarti

Rickart's dyak fruit bat (*Dyacopterus rickarti*) is a species of megabat in the family Pteropodidae found in Luzon and Mindanao islands, in the Philippines.
Rickett's big-footed bat
Rickett's big-footed bat
Myotis pilosus

Rickett's big-footed bat (*Myotis pilosus*) is a species of vesper bat. It can be found in southern and eastern China, Vietnam, and Laos. This species has often been called *Myotis ricketti*, but the older *M. pilosus* has priority. The erroneous...
Ridley's bat
Ridley's bat
Myotis ridleyi

Ridley's bat (*Myotis ridleyi*) is a species of vesper bat. It is found in Indonesia and Malaysia.
Ridley's leaf-nosed bat
Ridley's leaf-nosed bat
Hipposideros ridleyi

Ridley's leaf-nosed bat, Ridley's roundleaf bat, or Singapore roundleaf horseshoe bat (*Hipposideros ridleyi*) is a species of bat in the family Hipposideridae. It is found in Brunei, Malaysia and Singapore. Its natural habitat is subtropical or...
Ring-tailed ground squirrel
Ring-tailed ground squirrel
Notocitellus annulatus

The ring-tailed ground squirrel (*Notocitellus annulatus*) is a species of rodent in the family Sciuridae. It is endemic to the Pacific coast region of central Mexico. It is a common species and feeds mainly on fruits and nuts. The IUCN has...
Ring-Tailed Lemur
Ring-Tailed Lemur
Lemur catta

The Ring-tailed lemur (*Lemur catta*) is a large strepsirrhine primate and the most recognized lemur. It belongs to Lemnidae, one of five lemur families, and is the only member of the Lemur genus. Like all lemurs it is endemic to the island of...
Ring-Tailed Vontsira
Ring-Tailed Vontsira
Galidia elegans

The Ring-tailed vontsira (*Galidia elegans*) is a carnivorous native to Madagascar. Its local common name is vontsira mena, meaning 'red vontsira' in Malagasy.
Ringed Seal
Ringed Seal
Pusa hispida

The Ringed seal (*Pusa hispida*) is a small earless seal that lives in the Arctic and sub-Arctic regions. It is the most abundant and wide-ranging ice seal in the Northern Hemisphere. Ringed seals are one of the primary prey of Polar bears and...
Ringtail
Ringtail
Bassariscus astutus

The ringtail (*Bassariscus astutus*) is a member of the raccoon family native to arid regions of North America. It is the state mammal of Arizona.

Rio Acari marmoset
Rio Acari marmoset
Mico acariensis

The Rio Acari marmoset (*Mico acariensis*) is a marmoset species endemic to Brazil. It was first described in 2000.

Rio Beni titi
Rio Beni titi
Plecturocebus modestus

Rio Beni titi (*Plecturocebus modestus*) is a species of titi, a type of New World monkey, endemic to Bolivia.

Río Cesar white-fronted capuchin
Río Cesar white-fronted capuchin
Cebus cesarae

The Río Cesar white-fronted capuchin (*Cebus cesarae*) is a species of gracile capuchin monkey from the Río Cesar Valley in northern Colombia. It had previously been considered a subspecies of the white-fronted capuchin (*C. albifrons*).
Genetic...

Rio de Janeiro arboreal rat
Rio de Janeiro arboreal rat
Phaenomys ferrugineus

The Rio de Janeiro arboreal rat (*Phaenomys ferrugineus*) is a rodent species from South America. It is found in Brazil. It is the only species in the genus *Phaenomys*.

Rio Grande ground squirrel
Rio Grande ground squirrel
Ictidomys parvidens

The Rio Grande ground squirrel (*Ictidomys parvidens*) is a species of squirrel in the family *Sciuridae*. It is found in the south-western United States (Texas and New Mexico) and in north-eastern Mexico.

Rio Guaporé mouse
Rio Guaporé mouse
Juscelinomys guaporensis

The Rio Guaporé mouse or Guaporé akodont (*Juscelinomys guaporensis*) was formerly considered a rodent species in the family *Cricetidae*. It is known only from a small savanna in eastern Bolivia near the Rio Guaporé. However, in 2012 it was...

Rio Mayo titi
Rio Mayo titi
Plecturocebus oenanthe

The Rio Mayo titi (*Plecturocebus oenanthe*) is a species of titi, a type of New World monkey, endemic to Peru. The Rio Mayo titi, was thought previously to have a small range of origin in the Alto Mayo valley, but research has proven that the...

Rio Negro brush-tailed rat
Rio Negro brush-tailed rat
Isothrix negrensis

The Rio Negro brush-tailed rat or dark brush-tailed tree rat (*Isothrix negrensis*), is a spiny rat species from South America. It is found in Brazil.

Rio Negro tuco-tuco
Rio Negro tuco-tuco
Ctenomys rionegrensis

The Rio Negro tuco-tuco (*Ctenomys rionegrensis*) is a species of rodent in the family Ctenomyidae. It is found in a small fragmented range in Entre Ríos Province in northeastern Argentina and in the Río Negro Department of western Uruguay. It is...

Rio Purus titi

Rio Purus titi

Cheracebus purinus

The collared titi (*Cheracebus torquatus*) is a species of titi, a type of New World monkey. It is endemic to northern Brazil.

Rio Tapajós saki

Rio Tapajós saki

Pithecia irrorata

The Rio Tapajós saki or Gray's bald-faced saki (*Pithecia irrorata*) is a species of saki monkey, a type of New World monkey, from South America. It is found in parts of western Brazil, southeastern Colombia, southeastern Peru and possibly...

Riparian myotis

Riparian myotis

Myotis riparius

The riparian myotis (*Myotis riparius*), is a vespertilionid bat species from South and Central America. It is a medium-sized bat compared to other South American myotis.

Risso's Dolphin

Risso's Dolphin

Grampus griseus

Risso's dolphin (*Grampus griseus*) is one of the most mysterious and little known-species of dolphin with a rather unusual appearance. It is named after Antoine Risso, whose study of the animal formed the basis of the recognized description by...

Riverine Rabbit

Riverine Rabbit

Bunolagus monticularis

The Riverine rabbit (*Bunolagus monticularis*) is the only member of its genus because of the unique traits that separate it from the other leporids. It is one of the most endangered mammals in the world.

Roach's mouse-tailed dormouse

Roach's mouse-tailed dormouse

Myomimus roachi

The Roach's mouse-tailed dormouse or ground dormouse, also known simply as the mouse-tailed dormouse, (*Myomimus roachi*) is a species of rodent in the family Gliridae. It is found in Bulgaria, Turkey, and possibly the far east of Greece.

Roan Antelope

Roan Antelope

Hippotragus equinus

The roan antelope (*Hippotragus equinus*) is a savanna antelope found in western, central, and southern Africa. The roan antelope is one of the largest species of antelope; only eland, bongo and large male greater kudu can exceed them in weight....

Robbins's tateril

Robbins's tateril

Taterillus arenarius

Robbins's tateril (*Taterillus arenarius*), or the Sahel gerbil, is a species of rodent found across Mauritania and possibly Mali and Niger. Its natural habitats are dry savanna and subtropical or tropical dry shrubland.

Robbins's yellow bat

Robbins's yellow bat

Scotophilus nucella

Robbins's yellow bat (*Scotophilus nucella*) is a species of vesper bat. It is found in Ivory Coast, Ghana, and Uganda.

Robert's hocicudo

Robert's hocicudo

Oxymycterus roberti

Robert's hocicudo (*Oxymycterus roberti*) is a rodent species from South America. It is found in Brazil and Paraguay.

Robert's shaggy rat

Robert's shaggy rat

Dasymys robertsii

Robert's shaggy rat (*Dasymys robertsii*) is a species of rodent in the genus *Dasymys* that lives in South Africa. It was described in 2004.

Robert's snow vole

Robert's snow vole

Chionomys roberti

Robert's snow vole (*Chionomys roberti*) is a species of rodent in the family Cricetidae. It is found in Azerbaijan, Georgia, the Russian Federation, and Turkey. Its natural habitats are temperate forests and temperate grassland.

Roberto's spiny rat

Roberto's spiny rat

Proechimys roberti

Roberto's spiny-rat, *Proechimys roberti*, or Para spiny rat, is a spiny rat species from South America. It is found in Brazil. Morphological, karyological, and mitochondrial DNA (cytochrome b) data indicate that *Proechimys oris* is likely a junior...

Roberts' lechwe

Roberts' lechwe

Kobus leche robertsi

Roberts' lechwe (*Kobus leche robertsi*) or Kawambwa lechwe is an extinct subspecies of lechwe. It was found around Kawambwa, Zambia.

Roberts's flat-headed bat

Roberts's flat-headed bat

Sauromys petrophilus

Roberts's flat-headed bat (*Sauromys petrophilus*) is a species of free-tailed bat native to southern Africa. It is the only species in the genus *Sauromys*. The scientific name translates as "rock loving lizard-mouse", while the common name honours...

Robinson's mouse opossum

Robinson's mouse opossum

Marmosa robinsoni

Robinson's mouse opossum (*Marmosa robinsoni*) is a species of opossum in the family Didelphidae. It is found in Belize, Colombia, Ecuador, Grenada, Honduras, Panama, Peru, Trinidad and Tobago, and Venezuela. Robinson's mouse opossums move along...

Roborovski Hamster

Roborovski Hamster

Phodopus roborovskii

The Roborovski hamster (*Phodopus roborovskii*) is the smallest of three species of hamster in the genus *Phodopus*. It lives in the deserts of Central Asia and its average lifespan is only 2-3 years. These tiny creatures are known for their speed...

Robust cottontail

Robust cottontail

Sylvilagus robustus

The robust cottontail or Holzner's cottontail (*Sylvilagus holzneri*) is a species of cottontail rabbit native to high-altitude regions of the southwestern United States and western Mexico.

Robust golden mole

Robust golden mole

Amblysomus robustus

The robust golden mole (*Amblysomus robustus*) is a species of mammal in the golden mole family, *Chrysochloridae*. It is endemic to parts of Mpumalanga province in South Africa. Its natural habitats are temperate forests, subtropical or tropical...

Robust tuco-tuco

Robust tuco-tuco

Ctenomys tuconax

The robust tuco-tuco (*Ctenomys tuconax*) is a species of rodent in the family *Ctenomyidae*. It is a burrowing rodent and is endemic to the Tucumán Province of Argentina.

Robust yellow bat

Robust yellow bat

Scotophilus robustus

The robust yellow bat (*Scotophilus robustus*) is a species of vesper bat. It is found only in Madagascar.

Rock Cavy

Rock Cavy

Kerodon rupestris

The rock cavy is a rodent species endemic to eastern Brazil, which has also been introduced to the Atlantic island of Fernando de Noronha. The rock cavy is called *mocó* in Brazilian Portuguese, and *koriko pexerumen* in the Xukuru language of...

Rock dormouse

Rock dormouse

Graphiurus platyops

The rock dormouse or flat-headed African dormouse (*Graphiurus platyops*) is a species of rodent in the family *Gliridae*. It is found in Botswana, Mozambique, South Africa, Eswatini, Zambia, and Zimbabwe where it lives among rocks in upland areas....

Rock Hyrax

Rock Hyrax

Procavia capensis

The Rock hyrax (*Procavia capensis*) is a medium-sized terrestrial mammal native to Africa and the Middle East. It is also commonly referred to in South Africa as the *dassie*. These cute animals are the only extant terrestrial afrotherians in the...

Rock pocket mouse

Rock pocket mouse

Chaetodipus intermedius

The rock pocket mouse (*Chaetodipus intermedius*) is one of 19 species of pocket mice in the genus *Chaetodipus*. It is sometimes grouped in the genus *Perognathus*.

Rock Squirrel

Rock Squirrel

Otospermophilus variegatus

The rock squirrel (*Otospermophilus variegatus*) is a species of rodent in the family *Sciuridae* and is native to Mexico and the Southwestern United States, including southern Nevada, Utah, Colorado, Arizona, New Mexico, West Texas, and the...

Rock vole

Rock vole

Microtus chrotorrhinus

The rock vole (*Microtus chrotorrhinus*) is a medium-sized vole found in eastern North America. It is also called the yellow-nosed vole.

Rock-haunting ringtail possum

Rock-haunting ringtail possum

Petropseudes dahli

The rock-haunting ringtail possum (*Petropseudes dahli*), also known as the rock ringtail possum, is a species of Australian possum. It is found in rocky escarpments in the Kimberley, Arnhem Land and Gulf of Carpentaria across Western Australia...

Rock-loving mouse

Rock-loving mouse

Mus saxicola

The rock-loving mouse (*Mus saxicola*), also known as the brown spiny mouse, is a species of rodent in the family *Muridae*. It is found in India, Nepal, and Pakistan.

Rocky Mountain elk

Rocky Mountain elk

Cervus canadensis nelsoni

The Rocky Mountain elk (*Cervus canadensis nelsoni*) is a subspecies of elk found in the Rocky Mountains and adjacent ranges of Western North America.

Rodrigues Flying Fox

Rodrigues Flying Fox

Pteropus rodricensis

The Rodrigues flying fox or Rodrigues fruit bat (*Pteropus rodricensis*) is a species of bat in the family *Pteropodidae*, the flying foxes or fruit bats. It is endemic to Rodrigues, an island in the Indian Ocean belonging to Mauritius. Its natural...

Rodriguez's harvest mouse

Rodriguez's harvest mouse

Reithrodontomys rodriguezi

Rodriguez's harvest mouse (*Reithrodontomys rodriguezi*) is a species of rodent in the family *Cricetidae*. It is found only in Costa Rica.

Roe Deer

Roe Deer

Capreolus capreolus

The Roe deer (*Capreolus capreolus*) is a small species of deer. The male of the species is sometimes referred to as a roebuck. In the English language, this deer was originally simply called a 'roe', but over time the word 'roe' has become a...
Rohu's bat

Rohu's bat
Philetor brachypterus

Rohu's bat (*Philetor brachypterus*) is a species of vesper bat. It belongs to the monotypic genus *Philetor*. It is found in Brunei, Indonesia, Malaysia, Nepal, Papua New Guinea, and the Philippines.

Roig's pericote
Roig's pericote
Andalgalomys roigi

Roig's pericote or Roig's chaco mouse (*Andalgalomys roigi*) is a species of rodent in the family *Cricetidae*. Its karyotype has $2n = 60$ and $FN = 120$. It may be a subspecies of *Olrog's chaco mouse*.

Roig's tuco-tuco
Roig's tuco-tuco
Ctenomys roigi

Roig's tuco-tuco (*Ctenomys roigi*) is a species of rodent in the family *Ctenomyidae*. It is endemic to a small region near the Paraná River in Corrientes Province, northeastern Argentina, where lives on sand dunes, and near rivers.

Development is...

Roloway Monkey
Roloway Monkey
Cercopithecus roloway

The Roloway monkey (*Cercopithecus roloway*) is an endangered Old World monkey that can be found only in West Africa. It was previously considered a subspecies of the Diana monkey (*C. diana*). It is still difficult to gather data on wild Roloway...

Roman mole
Roman mole
Talpa romana

The Roman mole (*Talpa romana*) is a species of mammal in the family *Talpidae*. It is found in southern Italy. It was last recorded on Sicily in 1885. There is also an unconfirmed report about an isolated subpopulation in the Var region of southern...

Romanian hamster
Romanian hamster
Mesocricetus newtoni

The Romanian hamster or Dobrudja hamster (*Mesocricetus newtoni*) is a species of rodent in the family *Cricetidae*. It is found in Bulgaria and Romania.

Ronald's opossum
Ronald's opossum
Monodelphis ronaldei

Ronald's opossum (*Monodelphis ronaldei*) is a South American opossum species of the family *Didelphidae*. It was discovered in 2004, and most closely resembles *M. adusta*. It is known only from Manú National Park, Peru, where it inhabits the Amazon...

Rondo dwarf galago
Rondo dwarf galago
Paragalago rondoensis

The Rondo dwarf galago (*Paragalago rondoensis*) or Rondo bushbaby is a species of primate in the family *Galagidae*. The dwarf galagos are the smallest members of the genus *Galagoides*. It weighs less than 100 grams, making it the smallest known...

Rondon's marmoset
Rondon's marmoset
Mico rondoni

Rondon's marmoset (*Mico rondoni*), also known as the Rondônia marmoset, is a small species of monkey from the family Callitrichidae found in the south-western Amazon in Brazil. It is endemic to the state of Rondônia, and its range bordered by the...
Roosevelt elk
Roosevelt elk
Cervus canadensis roosevelti

The Roosevelt elk (*Cervus canadensis roosevelti*), also known commonly as the Olympic elk and Roosevelt's wapiti, is the largest of the four surviving subspecies of elk (*Cervus canadensis*) in North America by body mass (although by antler size,...
Roosevelt's muntjac
Roosevelt's muntjac
Muntiacus rooseveltorum

A single specimen of the Roosevelt's muntjac or Roosevelt's barking deer (*Muntiacus rooseveltorum*) was presented to the Field Museum in 1929 following the Kelley-Roosevelts expedition organized by Theodore (Jnr) and Kermit Roosevelt. The...
Roosevelt's shrew
Roosevelt's shrew
Crocidura roosevelti

Roosevelt's shrew (*Crocidura roosevelti*) is a species of mammal in the family Soricidae. It is found in Angola, Cameroon, Central African Republic, Democratic Republic of the Congo, Rwanda, Tanzania, and Uganda. Its natural habitat is moist...
Roosmalen's dwarf porcupine
Roosmalen's dwarf porcupine
Coendou roosmalenorum

Roosmalen's dwarf porcupine, *Coendou roosmalenorum*, is a porcupine species from the New World porcupine family likely endemic to northern Brazil. Only three specimens were known at the time, and only one had a collection locality. It is named for...
Roosmalens' dwarf marmoset
Roosmalens' dwarf marmoset
Mico humilis

The Roosmalens' dwarf marmoset (*Mico humilis*), also known as the black-crowned dwarf marmoset, is a small New World monkey native to the Amazon Rainforest, on the east bank of the lower Madeira River, and the west bank of the Aripuanã River, in...
Roraima mouse
Roraima mouse
Podoxymys roraimae

The Roraima mouse (*Podoxymys roraimae*) is a species of rodent in the family Cricetidae. It is the only species in the genus *Podoxymys*. It is found only in Guyana.
Rory Cooper's false antechinus
Rory Cooper's false antechinus
Pseudantechinus roryi

Rory Cooper's false antechinus (*Pseudantechinus roryi*), also known as the tan false antechinus and the tan pseudantechinus, is a recently named species of small carnivorous marsupial which inhabits rocky outcrops in Western Australia. Nothing is...
Rosalinda gerbil
Rosalinda gerbil

Gerbillus rosalinda

The Rosalinda gerbil (*Gerbillus rosalinda*) is distributed mainly central Sudan.
Rosalinda's Oldfield mouse
Rosalinda's Oldfield mouse
Thomasomys rosalinda

Rosalinda's Oldfield mouse (*Thomasomys rosalinda*) is a species of rodent in the family Cricetidae. It is found only in Peru.
Rosenberg's fruit-eating bat
Rosenberg's fruit-eating bat
Dermanura rosenbergi

Rosenberg's fruit-eating bat (*Dermanura rosenbergi*, often misspelled *rosenbergii*) is a species of bat in the family Phyllostomidae. It is found in humid tropical forests in the El Chocó region on the coast of western Colombia and northwestern...
Rosevear's serotine
Rosevear's serotine
Neoromicia roseveari

Rosevear's serotine (*Pseudoromicia roseveari*) is a species of vesper bat that lives in Guinea and Liberia. It was described as a new species in 2013. It is listed as endangered by the IUCN.
Rosevear's striped grass mouse
Rosevear's striped grass mouse
Lemniscomys roseveari

Rosevear's lemniscomys or Rosevear's striped grass mouse (*Lemniscomys roseveari*) is a species of rodent in the family Muridae. It is found in Zambia and possibly Angola, where its natural habitat is cryptosephalum dry forest. The species is...
Ross Seal
Ross Seal
Ommatophoca rossii

The Ross seal (*Ommatophoca rossii*) is a true seal (family Phocidae) with a range confined entirely to the pack ice of Antarctica. It is the only species of the genus *Ommatophoca*. First described during the Ross expedition in 1841, it is the...
Rossel Island melomys
Rossel Island melomys
Melomys arcium

The Rossel Island melomys (*Melomys arcium*) is a species of rodent from the family Muridae. It is endemic to Rossel Island in Papua New Guinea, but Flannery (1955) presumed that "it did exist previously on other islands of the group, but has...
Rothschild's cuscus
Rothschild's cuscus
Phalanger rothschildi

Rothschild's cuscus (*Phalanger rothschildi*), also called the Obi Island cuscus, is a species of marsupial in the family Phalangeridae. It is endemic to the islands of Obi, Bisa and Obilatu in the Obi Islands of Maluku province, Indonesia.
Rothschild's Giraffe
Rothschild's Giraffe
Giraffa camelopardalis rothschildi

Rothschild's giraffe (*Giraffa camelopardalis rothschildi*) is a subspecies of the giraffe. It is one of the most endangered distinct populations of giraffe, with 1,399 mature individuals estimated in the wild in 2018.
Rothschild's porcupine

Rothschild's porcupine
Coendou rothschildi

The Andean porcupine (Coendou quichua) or Quichua porcupine is a species of rodent in the family Erethizontidae. It is found in the Andes of northern Ecuador and Colombia as well as in Panama. This porcupine is little known, but is probably...

Rothschild's rock-wallaby
Rothschild's rock-wallaby
Petrogale rothschildi

Rothschild's rock-wallaby (Petrogale rothschildi) - sometimes known as the Roebourne rock-wallaby, is a species of macropod found in Western Australia, in the Pilbara district and the Dampier Archipelago. It is not currently considered to be...

Rothschild's woolly rat
Rothschild's woolly rat
Mallomys rothschildi

Rothschild's woolly rat (Mallomys rothschildi) is a species of rodent in the family Muridae. It is found on the island of New Guinea: both in the West Papua region of Indonesia and Papua New Guinea. The species has been known to eat karuka nuts...

Rothschild's zokor
Rothschild's zokor
Eospalax rothschildi

The Rothschild's zokor (Eospalax rothschildi) is a species of rodent in the family Spalacidae. It is endemic to China.

Rough-haired golden mole
Rough-haired golden mole
Chrysospalax villosus

The rough-haired golden mole (Chrysospalax villosus) is a species of mammal that live mostly below ground. They have shiny coats of dense fur and a streamlined, formless appearance. They have no visible eyes or ears; in fact, they are blind

....
Rough-Toothed Dolphin
Rough-Toothed Dolphin
Steno bredanensis

The rough-toothed dolphin (Steno bredanensis) is a species of dolphin that can be found in deep warm and tropical waters around the world. The species was first described by Georges Cuvier in 1823. The genus name Steno, of which this species is...

Round-eared tube-nosed bat
Round-eared tube-nosed bat
Murina cyclotis

The round-eared tube-nosed bat (Murina cyclotis), is a species of bat in the family Vespertilionidae from Central and Southeast Asia.

Round-eared tube-nosed fruit bat
Round-eared tube-nosed fruit bat
Nyctimene cyclotis

The round-eared tube-nosed fruit bat (Nyctimene cyclotis) is a species of bat in the family Pteropodidae. It is possibly conspecific with Nyctimene certans, although the taxonomy remains unresolved. The possible synonymy of the species was...

Round-Tailed Ground Squirrel

Round-Tailed Ground Squirrel
Xerospermophilus tereticaudus

The round-tailed ground squirrel (*Xerospermophilus tereticaudus*), known as "Ardillón cola redonda" in Spanish, live in the desert of the Southwestern United States and Northwestern Mexico. They are called "ground squirrels" because they burrow...

Round-Tailed Muskrat
Round-Tailed Muskrat
Neofiber alleni

The Round-tailed muskrat (*Neofiber alleni*) is a semi-aquatic species of rodent, sometimes called the Florida water rat. It is found only in the southeastern United States, where its natural habitat is swamps.

Royal Antelope
Royal Antelope
Neotragus pygmaeus

The royal antelope (*Neotragus pygmaeus*) is a West African antelope, recognized as the world's smallest antelope. It was first described by Swedish zoologist Carl Linnaeus in 1758. It stands up to merely 25 centimetres (10 in) at the shoulder and...

Royal Vole
Royal Vole
Myodes regulus

The Royal vole (*Craseomys regulus*) is a small rodent that occurs only on the Korean Peninsula. It lives underground in a burrow and comes out at night to feed on various vegetation.

Royle's mountain vole
Royle's mountain vole
Alticola roylei

Royle's mountain vole (*Alticola roylei*) is a species of rodent in the family Cricetidae. It is found in China, Nepal, Pakistan and India.

Royle's pika
Royle's pika
Ochotona roylei

Royle's pika (*Ochotona roylei*), also called the Himalayan mouse hare or hui shutu, is a species of pika. It is found in Bhutan, China, India, Nepal, and Pakistan.

Ruatan Island agouti
Ruatan Island agouti
Dasyprocta ruatanica

The Ruatan Island agouti (*Dasyprocta ruatanica*), also called the Roatán Island agouti, is a species of agouti in the family Dasyproctidae.

Rudd's African mole-rat
Rudd's African mole-rat
Tachyoryctes ruddi

Rudd's mole-rat (*Tachyoryctes ruddi*) is a species of rodent in the family Spalacidae. It is found in Kenya and Uganda. Its natural habitats are subtropical or tropical moist lowland forest, subtropical or tropical moist montane forest,...

Rudd's mouse
Rudd's mouse
Uranomys ruddi

Rudd's mouse or the white-bellied brush-furred rat (*Uranomys ruddi*) is the only

member of the genus *Uranomys*. This animal is closely related to the spiny mice, brush-furred mice, and the link rat.

Ruddy Mongoose

Ruddy Mongoose

Herpestes smithii

The Ruddy mongoose (*Urva smithii*) is a small carnivorous mammal native to hill forests in South Asia. Mongooses are carnivores and among other prey, they hunt various rodents and insects and therefore are often kept to control pests.

Ruddy treeshrew

Ruddy treeshrew

Tupaia splendidula

The ruddy treeshrew (*Tupaia splendidula*) is a treeshrew species in the family Tupaiidae. It is endemic to Borneo, the Natuna Islands and the Anambas Islands.

Rufous dog-faced bat

Rufous dog-faced bat

Molossops neglectus

The rufous dog-faced bat (*Molossops neglectus*), is a bat species from South America. It is found in Argentina, Brazil, Colombia, Guyana, Peru and Suriname.

Rufous elephant shrew

Rufous elephant shrew

Elephantulus rufescens

The rufous elephant shrew, rufous sengi or East African long-eared elephant-shrew (*Galegeeska rufescens*) is a species of elephant shrew in the family Macroscelididae. Found in Ethiopia, Kenya, Somalia, South Sudan, Tanzania and Uganda, its...

Rufous Hare-Wallaby

Rufous Hare-Wallaby

Lagorchestes hirsutus

The Rufous hare-wallaby (*Lagorchestes hirsutus*) is a small macropod from Australia. It was once widespread in the central and western deserts but a number of threats, caused the last wild population of this species on mainland Australia to go...

Rufous horseshoe bat

Rufous horseshoe bat

Rhinolophus rouxii

The rufous horseshoe bat (*Rhinolophus rouxii*) is a species of bat in the family Rhinolophidae. It is found in China, India, Myanmar, Nepal, Sri Lanka, and Vietnam. It has been identified as a vector of severe acute respiratory syndrome...

Rufous mouse opossum

Rufous mouse opossum

Marmosa lepida

The rufous mouse opossum (*Marmosa lepida*) or little rufous mouse opossum is an opossum species from South America. The species has been found in Bolivia, Surinam, French Guinea, Brazil, Colombia, Ecuador, Guyana, Peru and Surinam in lowland...

Rufous mouse-eared bat

Rufous mouse-eared bat

Myotis bocagii

The rufous mouse-eared bat (*Myotis bocagii*) is a species of vesper bat. It can be found in the following countries: Angola, Benin, Burundi, Cameroon, Central African Republic, Republic of the Congo, Democratic Republic of the Congo, Ivory Coast,...

Rufous rat-kangaroo

Rufous rat-kangaroo

Aepyprymnus rufescens

The rufous rat-kangaroo (*Aepyprymnus rufescens*), more commonly known as the rufous bettong, is a small marsupial species of the family Potoroidae found in Australia. It is not classified as threatened. The rufous bettong is about the size of a...

Rufous soft-furred spiny rat

Rufous soft-furred spiny rat

Diplomys labilis

The rufous soft-furred spiny-rat (*Diplomys labilis*), known as ratón espinoso meaning "spiny" or "thorny" rat in Spanish, is a species of rodent in the family Echimyidae. It is found in Colombia, Ecuador, and Panama. In November 2015, it was...

Rufous trident bat

Rufous trident bat

Triaenops persicus

The rufous trident bat, Persian trident bat, or triple nose-leaf bat (*Triaenops persicus*) is a species of bat in the genus *Triaenops*. It occurs in southwestern Pakistan, southern Iran, the United Arab Emirates, Oman, and Yemen. In the last...

Rufous-bellied bolo mouse

Rufous-bellied bolo mouse

Necromys lactens

The rufous-bellied bolo mouse or white-chinned akodont, (*Necromys lactens*) is a species of rodent in the family Cricetidae. It is found in Argentina and Bolivia where it inhabits the dry valleys of the eastern Andes. Its conservation status is...

Rümmler's brush mouse

Rümmler's brush mouse

Coccymys ruemmleri

Rümmler's brush mouse or Rümmler's mouse (*Coccymys ruemmleri*) is a species of rodent in the family Muridae. It is found in Indonesia and Papua New Guinea. Its natural habitats are subtropical or tropical moist montane forests and subtropical or...

Rumpi mouse shrew

Rumpi mouse shrew

Myosorex rumpii

The Rumpi mouse shrew (*Myosorex rumpii*) is a Myosoricinae shrew found only on the Rumpi Hills, Cameroon. It is listed as a critically endangered species due to habitat loss and a restricted range.

Rungwe dwarf galago

Rungwe dwarf galago

Galagoides sp.

The Rungwe dwarf galago is a newly identified species of eastern dwarf galago. Specimens were first collected in the 1930s, but were identified as different species. A formal description of the species is presently being made.

Rupicolous gerbil

Rupicolous gerbil

Dipodillus rupicola

The rupicolous gerbil (*Dipodillus rupicola*) is a species of rodent in the family Muridae. It is found only in Mali. Its natural habitats are dry savanna and rocky areas.

Rupp's mouse

Rupp's mouse

Stenocephalemys ruppi

Rupp's mouse or Rupp's stenocephalemys (*Stenocephalemys ruppi*) is a species of rodent in the family Muridae. It is found only in Ethiopia. Its natural habitat is subtropical or tropical high-altitude shrubland. It is threatened by habitat loss.
Rüppell's broad-nosed bat
Rüppell's broad-nosed bat
Scoteanax rueppellii

Rüppell's or the greater broad-nosed bat (*Scoteanax rueppellii*) is a species of vespertilionid microbat found in eastern Australia.
Rüppell's Fox
Rüppell's Fox
Vulpes rueppellii

Rüppell's fox (*Vulpes rueppellii*), also called Rüppell's sand fox, is a fox species living in desert and semi-desert regions of North Africa, the Middle East, and southwestern Asia. It has been listed as Least Concern on the IUCN Red List since...
Rüppell's horseshoe bat
Rüppell's horseshoe bat
Rhinolophus fumigatus

Rüppell's horseshoe bat (*Rhinolophus fumigatus*) is a species of bat in the family Rhinolophidae found in Africa. Its natural habitats are subtropical or tropical dry forests, dry savanna, moist savanna, caves and other subterranean habitats....
Rüppell's pipistrelle
Rüppell's pipistrelle
Pipistrellus rueppellii

Rüppell's bat (*Vansonia rueppellii*), also known as Rüppell's pipistrelle, is a species of vesper bat found in Africa and Asian republics such as Iraq and Israel. It is the only member of the genus Vansonia. It is found in dry and moist savanna, ...
Rusa Deer
Rusa Deer
Rusa timorensis

The Javan rusa or Sunda sambar (*Rusa timorensis*) is a deer native to Indonesia and East Timor. Introduced populations exist in a wide variety of locations in the Southern Hemisphere.
Ruschi's rat
Ruschi's rat
Arawayaomys ruschii

Ruschi's rat or Ruschi's spiny mouse (*Arawayaomys ruschii*) is a rodent species found in Argentina and Brazil. Some cranial features suggest it may be an archaic relative of the paramo Oldfield mouse (*Thomasomys paramorum*). The upper parts are...
Russet free-tailed bat
Russet free-tailed bat
Chaerephon russatus

The russet free-tailed bat (*Chaerephon russatus*) is a species of bat in the family Molossidae. It is found in Cameroon, Democratic Republic of the Congo, Ghana, and Kenya. Its natural habitat is subtropical or tropical moist lowland forests. It...
Russet ground squirrel
Russet ground squirrel
Spermophilus major

The russet ground squirrel (*Spermophilus major*) is a species of rodent in the family Sciuridae. It is also known as the hibernating large ground squirrel and the large-toothed souslik. It is found in west central Asia where its natural habitat...

Russian Desman
Russian Desman
Desmana moschata

The Russian desman (*Desmana moschata*) (Russian: выхухоль vykhukhol') is a small semiaquatic mammal that inhabits the Volga, Don and Ural River basins in Russia, Ukraine and Kazakhstan. It constructs burrows into the banks of ponds and...

Rusty pipistrelle
Rusty pipistrelle
Pipistrellus rusticus

The rusty pipistrelle (*Pipistrellus rusticus*) is a species of vesper bat. It is found in Angola, Botswana, Burkina Faso, Central African Republic, Chad, Ethiopia, Ghana, Kenya, Malawi, Mozambique, Namibia, Nigeria, Senegal, South Africa, Sudan,...

Rusty-bellied brush-furred rat
Rusty-bellied brush-furred rat
Lophuromys sikapusi

The rusty-bellied brush-furred rat (*Lophuromys sikapusi*) is a species of rodent in the family Muridae. It is found in Angola, Benin, Cameroon, Central African Republic, Republic of the Congo, Democratic Republic of the Congo, Ivory Coast,...

Rusty-sided Atlantic tree-rat
Rusty-sided Atlantic tree-rat
Phyllomys pattoni

The rusty-sided Atlantic tree-rat, *Phyllomys pattoni*, is a spiny rat species from South America. It is found in Brazil.

Rusty-Spotted Cat
Rusty-Spotted Cat
Prionailurus rubiginosus

The Rusty-spotted cat (*Prionailurus rubiginosus*) is one of the cat family's smallest members, of which historical records are known only from India and Sri Lanka. In 2012, it was also recorded in the western Terai of Nepal. Since 2016, the global...

Rusty-Spotted Genet
Rusty-Spotted Genet
Genetta maculata

The Rusty-spotted genet (*Genetta maculata*) is a small African carnivoran. These cat-like animals can move their eyes within their sockets to a limited extent, and move their heads to focus on moving objects. They can also move their pinnae by...

Ruwenzori colobus
Ruwenzori colobus
Colobus angolensis ruwenzori

The Ruwenzori colobus (*Colobus angolensis ruwenzorii*), also known as Ruwenzori black-and-white colobus, is a subspecies of the Angola colobus. This primate is distributed from the Afromontane forests of the Ruwenzori Mountains across the...

Ruwenzori duiker
Ruwenzori duiker
Cephalophus rubidus

The Ruwenzori duiker or Ruwenzori red duiker (*Cephalophus rubidus*) is a stocky but small antelope found only in the Ruwenzori Mountains between Uganda and, probably, the Democratic Republic of Congo. They may be a subspecies of the black-fronted...
Ruwenzori horseshoe bat
Ruwenzori horseshoe bat
Rhinolophus ruwenzorii

The Ruwenzori horseshoe bat (*Rhinolophus ruwenzorii*) is a species of bat in the family Rhinolophidae. It is found in Democratic Republic of the Congo, Rwanda, and Uganda. Its natural habitats are subtropical or tropical moist lowland forest,...
Ruwenzori otter shrew
Ruwenzori otter shrew
Micropotamogale ruwenzorii

The Ruwenzori otter shrew (*Micropotamogale ruwenzorii*) is a species of semiaquatic dwarf otter shrew of the family Potamogalidae. It is found in Democratic Republic of the Congo, Rwanda, and Uganda. Otter shrews are shrew-like afrotherian...
Ruwenzori shrew
Ruwenzori shrew
Ruwenzorisorex suncoides

The Ruwenzori shrew (*Ruwenzorisorex suncoides*) is a species of mammal in the family Soricidae. It is monotypic within the genus *Ruwenzorisorex*. It is found in Burundi, Democratic Republic of the Congo, Rwanda, and Uganda. It is semiaquatic,...
Ruwenzori sun squirrel
Ruwenzori sun squirrel
Heliosciurus ruwenzorii

The Ruwenzori sun squirrel (*Heliosciurus ruwenzorii*) is a species of rodent in the family Sciuridae. It is found in Burundi, Democratic Republic of the Congo, Rwanda, and Uganda. Its natural habitats are subtropical or tropical moist montane...
Ruwenzori thicket rat
Ruwenzori thicket rat
Grammomys ibeanus

The Ruwenzori thicket rat (*Grammomys ibeanus*) is a species of rodent in the family Muridae. It is found in Kenya, Malawi, South Sudan, Tanzania, Uganda, and Zambia. Its natural habitats are subtropical or tropical moist montane forests and...
Ruwenzori vlei rat
Ruwenzori vlei rat
Otomys dartmouthi

The Ruwenzori vlei rat (*Otomys dartmouthi*) is a species of rodent in the family Muridae. It is found only in Uganda. Its natural habitat is subtropical or tropical high-altitude grassland.
Rwandan shaggy rat
Rwandan shaggy rat
Dasymys rwandae

The Rwandan shaggy rat (*Dasymys rwandae*) is a species of shaggy marsh rat endemic to north-western Rwanda, close to the Virunga Mountains.
Ryukyu Flying Fox
Ryukyu Flying Fox
Pteropus dasymallus

The Ryukyu flying fox or Ryukyu fruit bat (*Pteropus dasymallus*) is a species of megabat in the family Pteropodidae. It is found in Japan, Taiwan, and the Batanes and Babuyan Islands of the Philippines. Its natural habitats are subtropical or...

Ryukyu long-tailed giant rat
Ryukyu long-tailed giant rat
Diplothrix legata

The Ryukyu long-tailed giant rat or Ryukyu rat (*Diplothrix legata*) is a species of rodent in the family Muridae. It is the only extant species in the genus *Diplothrix*. It is found only in the Ryukyu Islands of Japan. Its natural habitat is...

Ryukyu mouse
Ryukyu mouse
Mus caroli

The Ryukyu mouse (*Mus caroli*) is a species of rodent in the family Muridae. It is found in Cambodia, China, Indonesia, Japan, Laos, Malaysia, Taiwan, Thailand, and Vietnam.

Ryukyu shrew
Ryukyu shrew
Crocidura orii

The Ryukyu shrew (*Crocidura orii*), also known as Orii's shrew is a species of mammal in the family Soricidae. It is endemic to Japan. It is threatened by habitat loss.

Ryukyu spiny rat
Ryukyu spiny rat
Tokudaia osimensis

The Ryukyu spiny rat (*Tokudaia osimensis*) is a species of rodent in the family Muridae. Endemic to Amami Ōshima island in the Amami Islands of the Ryukyu archipelago of Japan, its natural habitat is subtropical moist broadleaf forest. The...

Ryukyu tube-nosed bat
Ryukyu tube-nosed bat
Murina ryukyuana

The Ryukyu tube-nosed bat (*Murina ryukyuana*) is a species of vesper bat found only in Japan.

Sable
Sable
Martes zibellina

The sable (*Martes zibellina*) is a species of marten, a small omnivorous mammal primarily living in the forest environments of Russia. The name sable appears to be of Slavic origin and entered most Western European languages via the early medieval...

Sable Antelope
Sable Antelope
Hippotragus niger

The Sable antelope (*Hippotragus niger*) is an antelope which inhabits wooded savanna in East and Southern Africa. Local names for this species include swartwitpens (Afrikaans), kgama or phalafala (Sotho), mBarapi or palahala (Swahili), kukurugu,...

Sado mole
Sado mole
Mogera tokudae

The Sado mole or Tokuda's mole (*Mogera tokudae*) is a species of mammal in the family Talpidae. It is endemic to Sado Island, Japan.
Sage's rock rat

Sage's rock rat
Aconaemys sagei

Sage's rock rat (*Aconaemys sagei*) is a species of rodent in the family Octodontidae. It is found in Argentina and possibly Chile.

Sagebrush vole
Sagebrush vole
Lemmiscus curtatus

The sagebrush vole (*Lemmiscus curtatus*) is a tiny vole found in western North America. This is the only member of genus *Lemmiscus*. They are somewhat similar in appearance to lemmings. They have chunky bodies with short legs and a very short tail...

Sahamalaza sportive lemur
Sahamalaza sportive lemur
Lepilemur sahamalazensis

The Sahamalaza sportive lemur (*Lepilemur sahamalazensis*) is a species of sportive lemur endemic to northern Madagascar.

Saharan shrew
Saharan shrew
Crocidura tarfayensis

The Saharan shrew (*Crocidura tarfayensis*) is a species of mammal in the family Soricidae. It is found in Mauritania and Morocco. Its natural habitats are rocky areas and sandy shores. It is threatened by habitat loss.

Saharan Striped Polecat
Saharan Striped Polecat
Ictonyx libycus

The Saharan striped polecat (*Ictonyx libycus*) is a carnivorous mammal native to Africa. It is a nocturnal hunter known to spray a foul, skunk-like anal emission when threatened.

Sahelian tiny shrew
Sahelian tiny shrew
Crocidura pasha

The Sahelian tiny shrew (*Crocidura pasha*) is a species of mammal in the family Soricidae. It is found in Ethiopia and Sudan. Its natural habitat is dry Sahel savanna.

Sahyadris forest rat
Sahyadris forest rat
Rattus satarae

The Sahyadris forest rat (*Rattus satarae*) is a species of rat belonging to the family Muridae. It is native to the northern Western Ghats in India where it is split between three regions, Satara in Maharashtra, the Nilgiri mountains in Tamil...

Saiga Antelope
Saiga Antelope
Saiga tatarica

The Saiga antelope (*Saiga tatarica*) is a critically endangered antelope that during antiquity inhabited a vast area of the Eurasian steppe. During the Pleistocene, they also occurred in Beringian North America and the British Isles. It is...

Saimaa ringed seal
Saimaa ringed seal
Pusa hispida saimensis

The Saimaa ringed seal (*Pusa hispida saimensis*, Finnish: *Saimaannorppa*) is a subspecies of ringed seal (*Pusa hispida*). They are among the most endangered seals in the world, having a total population of only about 400 individuals. The only...

Saimiri annectens
Saimiri annectens
Saimiri annectens

Saimiri annectens, originally described as *Laventiana annectens* and later as *Neosaimiri annectens*, is an extinct species of New World monkey in the genus *Saimiri* (squirrel monkeys) from the Middle Miocene (Laventan in the South American land...

Saimiri fieldsi
Saimiri fieldsi
Saimiri fieldsi

Saimiri fieldsi is an extinct species of New World monkey in the genus *Saimiri* (squirrel monkeys) from the Middle Miocene (Laventan in the South American land mammal ages; 13.8 to 11.8 Ma). Its remains have been found at the...

Saint Lawrence Island shrew
Saint Lawrence Island shrew
Sorex jacksoni

The Saint Lawrence Island shrew (*Sorex jacksoni*) is a species of mammal in the family Soricidae. It is found only on Alaska's St. Lawrence Island.

Takeji horseshoe bat
Takeji horseshoe bat
Rhinolophus takejiensis

The *Takeji horseshoe bat* (*Rhinolophus takejiensis*) is a species of bat in the family Rhinolophidae. It is endemic to Zambia. Its natural habitats are subtropical or tropical dry forest, subtropical or tropical moist lowland forest, and moist...

Sakhalin vole
Sakhalin vole
Microtus sachalinensis

The *Sakhalin vole* (*Microtus sachalinensis*) is a species of rodent in the family Cricetidae. It is found only in Russia.

Salenski's shrew
Salenski's shrew
Chodsigoa salenskii

Salenski's shrew (*Chodsigoa salenskii*) is a red-toothed shrew found only in northern Sichuan, China, where it is known from Wolong National Nature Reserve. It is listed as a critically endangered species due to habitat loss and a restricted range.

Salim Ali's fruit bat
Salim Ali's fruit bat
Latidens salimalii

Salim Ali's fruit bat (*Latidens salimalii*) is a rare megabat species in the monotypic genus *Latidens*. It was first collected by Angus Hutton, a planter and naturalist in the High Wavy Mountains in the Western Ghats of Theni district, Tamil Nadu...

Saline red bat
Saline red bat
Lasiurus salinae

The *saline red bat* (*Lasiurus salinae*) is a species of bat from the family Vespertilionidae. It was formerly included as a subspecies or a synonym of the

eastern red bat and the desert red bat, but is distinct. The species is present in...

Salokko rat

Salokko rat

Taeromys arcuatus

The Salokko rat (*Taeromys arcuatus*) is a species of rodent in the family Muridae. It is found only in Sulawesi, Indonesia. The Salokko rat has been found in the Mekongga Mountains of southeastern Sulawesi, and Lore Lindu National Park, Central...

Salt Marsh Harvest Mouse

Salt Marsh Harvest Mouse

Reithrodontomys raviventris

The salt marsh harvest mouse (*Reithrodontomys raviventris*), also known as the red-bellied harvest mouse, is an endangered rodent endemic to the San Francisco Bay Area salt marshes in California. The two distinct subspecies are both endangered...

Salt's Dik-Dik

Salt's Dik-Dik

Madoqua saltiana

Salt's dik-dik (*Madoqua saltiana*) is a small antelope that lives in dry areas of Africa. It is named after Henry Salt, who was the first European to acknowledge the species in Abyssinia in the early 19th century.

Salta tuco-tuco

Salta tuco-tuco

Ctenomys saltarius

The Salta tuco-tuco (*Ctenomys saltarius*) is a species of rodent in the family Ctenomyidae. It is endemic to Argentina and Bolivia.

Salvin's big-eyed bat

Salvin's big-eyed bat

Chiroderma salvini

Salvin's big-eyed bat (*Chiroderma salvini*) is a species of bat in the family Phyllostomidae. It is found in Bolivia, Colombia, Costa Rica, Ecuador, El Salvador, Guatemala, Honduras, Mexico, Panama, Peru, and Venezuela.

Salvin's spiny pocket mouse

Salvin's spiny pocket mouse

Heteromys salvini

Salvin's spiny pocket mouse (*Heteromys salvini*) is a small to medium-sized rodent in the family Heteromyidae. It was formerly placed in the genus *Liomys*, which is now recognized to be paraphyletic and has been subsumed into *Heteromys*.

Samar squirrel

Samar squirrel

Sundasciurus samarensis

The Samar squirrel (*Sundasciurus samarensis*) is a species of rodent in the family Sciuridae. It was historically known as lalagsing. It is endemic to the Philippines, where it has been recorded from Samar and Leyte Islands. Its natural habitat...

Sambar

Sambar

Rusa unicolor

The sambar (*Rusa unicolor*) is a large deer native to the Indian subcontinent and Southeast Asia. The name "sambar" is also sometimes used to refer to the Philippine deer called the "Philippine sambar", and the Javan rusa called the "Sunda sambar".

Sambirano mouse lemur
Sambirano mouse lemur
Microcebus sambiranensis

The Sambirano mouse lemur (*Microcebus sambiranensis*) is a small, recently discovered primate and like the other mouse lemurs can only be found on the island of Madagascar. The dorsal side is both cinnamon and rufous-cinnamon and is grey...

Sambirano woolly lemur
Sambirano woolly lemur
Avahi unicolor

The Sambirano woolly lemur (*Avahi unicolor*), also known as the Sambirano avahi or unicolor woolly lemur, is a species of woolly lemur native to western Madagascar.

Samoa flying fox
Samoa flying fox
Pteropus samoensis

The Samoa flying fox or Samoan flying fox (*Pteropus samoensis*) is a species of flying fox in the family Pteropodidae. It is found in American Samoa, Fiji, and Samoa (where it is known as pe'a and pe'a vao). Its natural habitat is subtropical or...

San Bernardino kangaroo rat
San Bernardino kangaroo rat
Dipodomys merriami parvus

The San Bernardino kangaroo rat (*Dipodomys merriami parvus*) is a species of rodent in the family Heteromyidae. It is one of 19 recognized subspecies of Merriam's kangaroo rat (*Dipodomys merriami*) that are spread throughout the arid regions of...

San Cristobal shrew
San Cristobal shrew
Sorex stizodon

The San Cristobal shrew (*Sorex stizodon*) is a species of mammal in the family Soricidae. It is endemic to Mexico.

San Diego pocket mouse
San Diego pocket mouse
Chaetodipus fallax

The San Diego pocket mouse (*Chaetodipus fallax*) is a rodent species in the family Heteromyidae. It occupies the northern region of Baja California near San Diego extending into Mexico.

San Esteban Island mouse
San Esteban Island mouse
Peromyscus stephani

The San Esteban Island mouse (*Peromyscus stephani*) is a species of rodent in the family Cricetidae. It is endemic to Mexico, where it is known only from San Esteban Island in the northern Gulf of California.

San Felipe hutia
San Felipe hutia
Mesocapromys sanfelipensis

The San Felipe hutia (*Mesocapromys sanfelipensis*), also known as the little earth hutia is a species of rodent. It is found in Cuba, and is listed on the 2019 IUCN Red List as Critically Endangered or possibly extinct.

San Joaquin Antelope Squirrel
San Joaquin Antelope Squirrel
Ammospermophilus nelsoni

The San Joaquin antelope squirrel or Nelson's antelope squirrel (*Ammospermophilus nelsoni*), is a species of antelope squirrel, in the San Joaquin Valley of the U.S. state of California.

San Joaquin kit fox

San Joaquin kit fox

Vulpes macrotis mutica

The endangered San Joaquin kit fox (*Vulpes macrotis mutica*) was formerly very common in the San Joaquin Valley and through much of Central California. Its 1990 population was estimated to be 7,000. This subspecies is still endangered, after...

San Joaquin pocket mouse

San Joaquin pocket mouse

Perognathus inornatus

The San Joaquin pocket mouse or Salinas pocket mouse (*Perognathus inornatus*) is a species of rodent in the family Heteromyidae. It is endemic to California in the United States where it lives in desert and semi-desert habitats.

San José brush rabbit

San José brush rabbit

Sylvilagus mansuetus

The San José brush rabbit (*Sylvilagus bachmani mansuetus*) is a critically endangered subspecies of the brush rabbit, in the family Leporidae.

San José Island kangaroo rat

San José Island kangaroo rat

Dipodomys merriami insularis

The San José Island kangaroo rat (*Dipodomys merriami insularis*) is a subspecies of rodent in the family Heteromyidae. It is endemic to Mexico, where it is found only on San José Island off the east coast of Baja California Sur. (*Dipodomys*...

San Juan tuco-tuco

San Juan tuco-tuco

Ctenomys johannis

The San Juan tuco-tuco (*Ctenomys johannis*) is a species of rodent in the family Ctenomyidae. It is endemic to west central Argentina, where it is known only from southern San Juan Province.

San Lorenzo mouse

San Lorenzo mouse

Peromyscus interparietalis

The San Lorenzo mouse (*Peromyscus interparietalis*) is a species of rodent in the family Cricetidae. It is endemic to Mexico, where it is found only from the islands of San Lorenzo Norte, San Lorenzo Sur, and Salsipuedes off the east coast of...

San Luis tuco-tuco

San Luis tuco-tuco

Ctenomys pontifex

The San Luis tuco-tuco (*Ctenomys pontifex*) is a species of rodent in the family Ctenomyidae. It is endemic to Argentina.

San Quintin kangaroo rat

San Quintin kangaroo rat

Dipodomys gravipes

The San Quintin kangaroo rat (*Dipodomys gravipes*) is a species of rodent in the family Heteromyidae. It is endemic to Mexico, where it is known only from western Baja California. Its natural habitat includes arid lowlands with sparse...

Sanborn's big-eared bat

Sanborn's big-eared bat
Micronycteris sanborni

Sanborn's big-eared bat (*Micronycteris sanborni*) is a bat species from South America. It is found in Bolivia and Brazil.

Sanborn's bonneted bat
Sanborn's bonneted bat
Eumops hansae

Sanborn's bonneted bat (*Eumops hansae*), is a bat species from South and Central America. It was first described from a specimen collected at Colonia Hansa, Brazil, and is sometimes also known as the Hansa bonneted bat.

Sanborn's squirrel
Sanborn's squirrel
Sciurus sanborni

Sanborn's squirrel (*Sciurus sanborni*) is or was a little known tree squirrel described in 1944 from the skin and skeleton of a single female specimen collected in Peru in 1941. Subsequently, over the proceeding years only a handful of either...

Sand Cat
Sand Cat
Felis margarita

The Sand cat (*Felis margarita*) is a small wild cat that inhabits sandy and stony deserts far from water sources. The first Sand cat known to science was discovered in the Algerian Sahara and described in 1858. In the early 1970s, these adorable...

Sand gerbil
Sand gerbil
Gerbillus syrticus

The sand gerbil (*Gerbillus syrticus*) is distributed mainly in northeastern Libya. Less than 250 individuals are thought to be left in existence. It is sometimes considered a subspecies of the pygmy gerbil.

Sand-colored soft-furred rat
Sand-colored soft-furred rat
Millardia gleadowi

The sand-colored soft-furred rat (*Millardia gleadowi*) is a species of rodent in the family Muridae. It is found in India and Pakistan.

Sandhill dunnart
Sandhill dunnart
Sminthopsis psammophila

The sandhill dunnart (*Sminthopsis psammophila*) is a species of small carnivorous Australian marsupial of the family Dasyuridae. It is known from four scattered arid areas of Australia: near Lake Amadeus in Northern Territory, the central Eyre...

Sandstone false antechinus
Sandstone false antechinus
Pseudantechinus bilarni

The sandstone false antechinus, *Pseudantechinus bilarni*, also known as the sandstone pseudantechinus, the sandstone antechinus, the sandstone dibbler, Harney's antechinus and the Northern dibbler, is a species of small carnivorous marsupial,...

Sandy inland mouse
Sandy inland mouse
Pseudomys hermannsburgensis

The sandy inland mouse (*Pseudomys hermannsburgensis*) is a species of rodent in the family Muridae. Also known as the Hermannsburg (Mission) false-mouse or Hermannsburg mouse, it is endemic to Australia and found widely yet sparsely through arid...

Sandy mole-rat
Sandy mole-rat
Spalax arenarius

The sandy blind mole-rat (*Spalax arenarius*) is an endangered species of rodent in the family Spalacidae. It is endemic to Ukraine.

Sanford's brown lemur
Sanford's brown lemur
Eulemur sanfordi

Sanford's brown lemur (*Eulemur sanfordi*), or Sanford's lemur, is a species of strepsirrhine primate in the family Lemnidae. Sanford's brown lemur was previously considered a subspecies of the common brown lemur (*Eulemur fulvus*) but was raised...

Sangai
Sangai
Rucervus eldii eldii

The sangai is an endemic and endangered subspecies of Eld's deer found only in Manipur, India. It is also the state animal of Manipur. Its common English name is Manipur brow-antlered deer or Eld's deer and the scientific name is *Rucervus eldii*...

Sanghir squirrel
Sanghir squirrel
Prosciurillus rosenbergii

The Sanghir squirrel (*Prosciurillus rosenbergii*) is a species of rodent in the family Sciuridae. It is endemic to the volcanic Sangir Islands (where they are the only squirrel) in Indonesia, located just north of the equator in the Celebes Sea,...

Sangihe tarsier
Sangihe tarsier
Tarsius sangirensis

The Sangihe tarsier (*Tarsius sangirensis*), also known as Sangihe Island tarsier, is a small primate found on Sangir Island, which is located about 200 kilometers north-east of the island of Sulawesi in Indonesia. In 2008 a population of the...

Sanje mangabey
Sanje mangabey
Cercocebus sanjei

The Sanje mangabey (*Cercocebus sanjei*) is a highly endangered Old World monkey of the white-eyelid mangabey group from the Eastern Arc Mountains in Tanzania. They are about 50-65 centimetres (20-26 in) in length, excluding the tail, and their...

Santa Catarina's guinea pig
Santa Catarina's guinea pig
Cavia intermedia

Santa Catarina's guinea pig (*Cavia intermedia*) or Moleques do Sul cavy is a rare guinea pig species of southeastern South America.

Santa Cruz mouse
Santa Cruz mouse
Peromyscus sejugis

The Santa Cruz mouse (*Peromyscus sejugis*) is a species of rodent in the family

Cricetidae. It is endemic to Mexico, where it is found only on two small islands in the southern Gulf of California. Feral cats on Santa Cruz Island are a threat.
Santa Marta porcupine
Santa Marta porcupine
Coendou sanctamartae

The Santa Marta porcupine (*Coendou sanctamartae*) is a rodent in the family Erethizontidae. It is known from dry forests on the lower slopes of the Sierra Nevada de Santa Marta and Serranía del Perijá mountains of northern Colombia, at altitudes...
Santa Marta white-fronted capuchin
Santa Marta white-fronted capuchin
Cebus malitiosus

The Santa Marta white-fronted capuchin (*Cebus malitiosus*) is a species of gracile capuchin monkey from Colombia. It was formerly considered a subspecies of the *Cebus albifrons* or a synonym of the Colombian white-faced capuchin (*C. capucinus*),...
Santander dwarf squirrel
Santander dwarf squirrel
Microsciurus santanderensis

The Santander dwarf squirrel (*Microsciurus santanderensis*) is a small tree squirrel endemic to Colombia.
Santarem marmoset
Santarem marmoset
Mico humeralifer

The Santarem marmoset (*Mico humeralifer*), also known as the black and white tassel-ear marmoset, is a marmoset endemic to the Brazilian states of Amazonas and Pará.
São Paulo grass mouse
São Paulo grass mouse
Akodon sanctipaulensis

The São Paulo grass mouse (*Akodon sanctipaulensis*) is a rodent species from South America. It is found in Brazil.
São Tomé collared fruit bat
São Tomé collared fruit bat
Myonycteris brachycephala

The São Tomé collared fruit bat (*Myonycteris brachycephala*) is a species of megabat in the family Pteropodidae. It is endemic to São Tomé and Príncipe. Its natural habitats are subtropical or tropical moist lowland forest and subtropical or...
São Tomé free-tailed bat
São Tomé free-tailed bat
Chaerephon tomensis

The São Tomé free-tailed bat (*Chaerephon tomensis*) is a species of bat in the family Molossidae. It is endemic to São Tomé and Príncipe. Its natural habitats are dry savanna, moist savanna, and plantations. It is threatened by habitat loss. Only...
São Tomé leaf-nosed bat
São Tomé leaf-nosed bat
Macronycteris thomensis

The São Tomé leaf-nosed bat (*Macronycteris thomensis*) is a species of bat in the family Hipposideridae. It is endemic to the island of São Tomé, in the Gulf of Guinea off the western coast of Africa. The bat's natural habitats are subtropical

or...

São Tomé shrew

São Tomé shrew

Crocidura thomensis

The São Tomé shrew (*Crocidura thomensis*) is a white-toothed shrew about 3.0 in (7.6 cm) long found only on São Tomé Island, São Tomé and Príncipe. It is listed as a critically endangered species due to habitat loss and a restricted range. It

was...

Saola

Saola

Pseudoryx nghetinhensis

The saola (*Pseudoryx nghetinhensis*) is one of the world's rarest large mammals, a forest-dwelling bovine native to the Annamite Range in Vietnam and Laos. It was described in 1993 following the discovery of remains in Vũ Quang Nature Reserve by a...

Sarawak surili

Sarawak surili

Presbytis chrysomelas

The Sarawak surili (*Presbytis chrysomelas*) is a species of primate in the family Cercopithecidae. It is endemic to the southeast Asian island of Borneo, where it is distributed north of the Kapuas River in Kalimantan, Indonesia, the Malaysia...

Sarcophilus laniarius

Sarcophilus laniarius

Sarcophilus laniarius

Sarcophilus laniarius is an extinct species of large Tasmanian devil. Richard Owen originally called the specimen on which the genus was based *Dasyurus laniarius*. Pleistocene fossil deposits in limestone caves at Naracoorte, South Australia include...

Sardinian long-eared bat

Sardinian long-eared bat

Plecotus sardus

The Sardinian long-eared bat (*Plecotus sardus*) is a species of bat endemic to Sardinia, Italy. This species was discovered in 2002 in the caves of central Sardinia, the type locality being a cave in Lanaitto's Valley in the Oliena District. It...

Sardinian Wildcat

Sardinian Wildcat

Felis lybica sarda

The Sardinian wildcat is an isolated population of feral cats (*Felis catus*) on the island of Sardinia, introduced during the Roman Empire. It has historically been misidentified as a Eurasian lynx (*Lynx lynx*) and an African wildcat (*Felis*...

Satéré marmoset

Satéré marmoset

Mico saterei

The Satéré marmoset (*Mico saterei*) is a marmoset species endemic to Brazil.

Saudi gazelle

Saudi gazelle

Gazella saudia

The Saudi gazelle (*Gazella saudia*) is a *Gazella* species that was once native to the Arabian Peninsula. It was declared extinct in the wild in 1996, as it was last observed in the wild in 1970. The Saudi gazelle was officially declared extinct...

Saunder's vlei rat
Saunder's vlei rat
Otomys saundersiae

Saunder's vlei rat (*Otomys saundersiae*) is a species of rodent in the family Muridae. It is found only in South Africa. Its natural habitats are subtropical or tropical dry shrubland, Mediterranean-type shrubby vegetation, and subtropical or...

Saussure's shrew
Saussure's shrew
Sorex saussurei

Saussure's shrew (*Sorex saussurei*) is a species of mammal in the family Soricidae, that is found in Mexico. There is also a disjointed population of shrews in Guatemala that is provisionally assigned to this species, but may represent a distinct...

Savanna dwarf shrew
Savanna dwarf shrew
Crocidura nanilla

The savanna dwarf shrew (*Crocidura nanilla*) is a species of mammal in the family Soricidae. It is found in Guinea, Kenya, Mauritania, Senegal, Tanzania, and Uganda. Its natural habitats are dry savanna and moist savanna.

Savanna gerbil
Savanna gerbil
Gerbilliscus validus

The savanna gerbil (*Gerbilliscus validus*) is a species of rodent found in Angola, Burundi, Democratic Republic of the Congo, Ethiopia, Kenya, Mali, Rwanda, Sudan, Tanzania, Uganda, Zambia, and Zimbabwe. Its natural habitats are dry savanna,...

Savanna path shrew
Savanna path shrew
Crocidura viaria

The savanna path shrew (*Crocidura viaria*) is a species of mammal in the family Soricidae. It is found in Benin, Burkina Faso, Cameroon, Chad, Eritrea, Ethiopia, Ghana, Kenya, Mali, Mauritania, Morocco, Niger, Nigeria, Senegal, Somalia, Sudan,...

Savanna shrew
Savanna shrew
Crocidura fulvastra

The savanna shrew (*Crocidura fulvastra*) is a species of mammal in the family Soricidae. It is found in Benin, Burkina Faso, Cameroon, Central African Republic, Chad, Democratic Republic of the Congo, Djibouti, Eritrea, Ethiopia, Kenya, Mali,...

Savanna swamp shrew
Savanna swamp shrew
Crocidura longipes

The Savanna swamp shrew (*Crocidura longipes*) is a species of mammal in the family Soricidae. It is endemic to Nigeria. Its natural habitat is swamps. It is threatened by habitat loss.

Savi's pine vole
Savi's pine vole
Microtus savii

Savi's pine vole (*Microtus savii*) is a species of rodent in the family Cricetidae. It is found in France and Italy.

Savi's pipistrelle
Savi's pipistrelle
Hypsugo savii

Savi's pipistrelle (*Hypsugo savii* sometimes classified as *Pipistrellus savii*) is a species of vesper bat found across North West Africa, the Mediterranean region and the Middle East. It feeds at night on flying insects. In the summer it roosts...

Savile's bandicoot rat
Savile's bandicoot rat
Bandicota savilei

Savile's bandicoot rat (*Bandicota savilei*) is a species of rodent in the family Muridae found in Myanmar, Thailand, and Vietnam.

Saxatilomys
Saxatilomys
Saxatilomys paulinae

Saxatilomys paulinae is a species of murid rodent native to central Laos and Vietnam, separated to a monotypic genus *Saxatilomys*. It was first discovered in the Khammouan Limestone National Biodiversity Conservation Area in Khammouan Province,...

Scaglia's tuco-tuco
Scaglia's tuco-tuco
Ctenomys scagliai

Scaglia's tuco-tuco (*Ctenomys scagliai*) is a species of rodent in the family Ctenomyidae. It is endemic to a locality in Tucumán Province, northern Argentina. The species is named after Argentine naturalist Galileo Juan Scaglia (1915–1989). Its...

Scaly-footed small-eared shrew
Scaly-footed small-eared shrew
Cryptotis squamipes

The scaly-footed small-eared shrew (*Cryptotis squamipes*) is a species of mammal in the family Soricidae. It is found in Colombia and Ecuador.

Scaly-tailed possum
Scaly-tailed possum
Wyulda squamicaudata

The scaly-tailed possum (*Wyulda squamicaudata*) is found in northwestern Australia, where it is restricted to the Kimberley. They are light grey over most of the body and have relatively short ears and muzzle. The mostly hairless tail is able to...

Schaller's mouse shrew
Schaller's mouse shrew
Myosorex schalleri

Schaller's mouse shrew (*Myosorex schalleri*) is a species of mammal in the family Soricidae endemic to the Democratic Republic of Congo. Its natural habitat is subtropical or tropical moist montane forests.

Schaub's myotis
Schaub's myotis
Myotis schaubi

Schaub's myotis (*Myotis schaubi*) is a species of vesper bat. It is found in Armenia, Azerbaijan and Iran and known from fossils from Hungary. It is related to *Myotis nattereri*.

Schelkovnikov's pine vole
Schelkovnikov's pine vole
Microtus schelkovnikovi

Schelkovnikov's pine vole (*Microtus schelkovnikovi*) is a species of rodent in the family Cricetidae. It is found in Azerbaijan and Iran. It has recently been considered the sole species in the subgenus Hyrcanicola.

Schidlovsky's vole

Schidlovsky's vole

Microtus schidlovskii

Schidlovsky's vole (*Microtus schidlovskii*) is a species of rodent in the family Cricetidae. It is normally found in northwestern Armenia, and was long considered a subspecies of the social vole until relisted as a species by Golenishchev in 2002.

Schlieffen's bat

Schlieffen's bat

Nycticeinops schlieffeni

Schlieffen's serotine (*Nycticeinops schlieffeni*), also known as Schlieffen's bat or Schlieffen's twilight bat, is a species of vesper bat found in Africa. It has been placed in numerous genera since its first description in 1859, but...

Schmidts's big-eared bat

Schmidts's big-eared bat

Micronycteris schmidtorum

Schmidts's big-eared bat (*Micronycteris schmidtorum*) is a bat species from South and Central America.

Schneider's leaf-nosed bat

Schneider's leaf-nosed bat

Hipposideros speoris

Schneider's leaf-nosed bat or Schneider's roundleaf bat (*Hipposideros speoris*) is a species of bat in the family Hipposideridae. It is endemic to South Asia. Its natural habitats are subtropical or tropical dry forests, caves, and urban areas.

Schreber's yellow bat

Schreber's yellow bat

Scotophilus nigrita

Schreber's yellow bat (*Scotophilus nigrita*) or the giant house bat, is a species of vesper bat. It is found in Benin, Democratic Republic of the Congo, Ivory Coast, Ghana, Kenya, Malawi, Mozambique, Nigeria, Senegal, Tanzania, Togo, and...

Schultz's round-eared bat

Schultz's round-eared bat

Lophostoma schulzi

Schultz's round-eared bat (*Lophostoma schulzi*) is a species of bat in the family Phyllostomidae. It is found in Brazil, French Guiana, Guyana, and Suriname. It is threatened by habitat loss.

Schwartz's myotis

Schwartz's myotis

Myotis martiniquensis

Schwartz's myotis (*Myotis martiniquensis*) is a species of vesper bat. It is found in Barbados and Martinique.

Scimitar Oryx

Scimitar Oryx

Oryx dammah

The Scimitar oryx (*Oryx dammah*) is a *Oryx* species that was once widespread across North Africa. In 2000, it was declared extinct in the wild on the IUCN Red List. Scimitar oryx inhabited semideserts and deserts and were adapted to live in extreme...

Sciurocheirus makandensis
Sciurocheirus makandensis
Sciurocheirus makandensis

Sciurocheirus makandensis (Makandé squirrel galago) is a species of squirrel galago native to Gabon, Africa.

Sciurus ingrami
Sciurus ingrami
Sciurus ingrami

Sciurus ingrami, usually called Ingram's squirrel in English, is a squirrel found in South America. It is known as serelepe in southeastern Brazil. It is found in the Atlantic Forest Biome of Brazil and Misiones Province, Argentina. It is regarded...

Sclater's golden mole
Sclater's golden mole
Chlorotalpa sclateri

Sclater's golden mole (Chlorotalpa sclateri) is a species of mammal in the family Chrysochloridae. It is found in Lesotho and South Africa. Its natural habitats are subtropical or tropical dry shrubland, subtropical or tropical high-altitude...

Sclater's guenon
Sclater's guenon
Cercopithecus sclateri

Sclater's guenon (Cercopithecus sclateri), also known as Sclater's monkey and the Nigerian monkey, is an Old World monkey that was first described by Reginald Innes Pocock in 1904 and named after Philip Sclater. It is an arboreal and diurnal...

Sclater's mouse shrew
Sclater's mouse shrew
Myosorex sclateri

Sclater's mouse shrew (Myosorex sclateri) is a species of mammal in the family Soricidae endemic to South Africa. Its natural habitats are subtropical or tropical moist lowland forests and swamps. It is threatened by habitat loss.

Sclater's shrew
Sclater's shrew
Sorex sclateri

Sclater's shrew (Sorex sclateri) is a species of mammal in the family Soricidae. It is endemic to Mexico.

Scolomys melanops
Scolomys melanops
Scolomys melanops

Scolomys melanops, also known as the short-nosed scolomys, South American spiny mouse, Ecuadorian spiny mouse, or gray spiny mouse, is a species of rodent in the genus Scolomys of family Cricetidae. It is a forest mouse and was thought to be...

Scolomys ucayalensis
Scolomys ucayalensis
Scolomys ucayalensis

Scolomys ucayalensis, also known as the long-nosed scolomys or Ucayali spiny mouse is a nocturnal rodent species from South America. It is part of the genus Scolomys within the tribe Oryzomyini. It is found in Brazil, Colombia, Ecuador and Peru...

Scotophilus tandrefana
Scotophilus tandrefana
Scotophilus tandrefana

The Malagasy yellow bat (*Scotophilus tandrefana*), sometimes known as the western yellow bat, is a species of vesper bat endemic to Madagascar.

Scott's mouse-eared bat

Scott's mouse-eared bat

Myotis scotti

Scott's mouse-eared bat (*Myotis scotti*) is a species of vesper bat. It is found only in Ethiopia, in subtropical or tropical moist montane forests and shrubland.

It is threatened by habitat loss.

Scott's sportive lemur

Scott's sportive lemur

Lepilemur scottorum

Scott's sportive lemur (*Lepilemur scottorum*), or the Masoala sportive lemur, is a sportive lemur endemic to Madagascar. This is one of 26 species in the genus *Lepilemur*. It has greyish brown fur and a black-tipped tail. It is named in honor of...

Scottish red deer

Scottish red deer

Cervus elaphus scoticus

The Scottish red deer (*Cervus elaphus scoticus*) is a subspecies of red deer, which is native to Great Britain. Like the red deer of Ireland, it migrated from continental Europe sometime in the Stone Age. The Scottish red deer is farmed for meat,...

Scottish Wildcat

Scottish Wildcat

Felis silvestris silvestris

The Scottish wildcat (*Felis silvestris silvestris*) is a European wildcat population in Scotland. It was once widely distributed across Great Britain, but the population has declined drastically since the turn of the 20th century. It is now...

Screaming Hairy Armadillo

Screaming Hairy Armadillo

Chaetophractus vellerosus

The screaming hairy armadillo (*Chaetophractus vellerosus*) is a species of armadillo also known as the small screaming armadillo, crying armadillo or the small hairy armadillo. It is a burrowing armadillo found in the central and southern parts...

Scrub Hare

Scrub Hare

Lepus saxatilis

The scrub hare (*Lepus saxatilis*) is one of two species of hares found in southern Namibia, Mozambique, South Africa, Eswatini and Lesotho. Although it is listed as a least concern species, the population has been declining and is expected to...

Scully's tube-nosed bat

Scully's tube-nosed bat

Murina tubinaris

Scully's tube-nosed bat (*Murina tubinaris*) is a species of vesper bat in the family Vespertilionidae. It can be found in the following countries: India, Laos, Myanmar, Pakistan, Thailand, and Viet Nam. Members of the ashy-gray tube-nosed bat...

Sculptor squirrel

Sculptor squirrel

Glyphotes simus

The sculptor squirrel or Bornean pygmy squirrel (*Glyphotes simus*) is a species of rodent in the family Sciuridae. It was described by Michael Rogers Oldfield Thomas in 1898. It is monotypic within the genus *Glyphotes*. It is endemic to Malaysia...

Sea Mink

Sea Mink

Neovison macrodon

The sea mink (*Neogale macrodon*) is a recently extinct species of mink that lived on the eastern coast of North America around the Gulf of Maine on the New England seaboard. It was most closely related to the American mink (*Neogale vison*), with...

Sea Otter

Sea Otter

Enhydra lutris

The Sea otter (*Enhydra lutris*) is a marine mammal native to the coasts of the northern and eastern North Pacific Ocean. They are the heaviest members of the weasel family, but among the smallest marine mammals. Unlike most marine mammals, the sea...

Seal's sportive lemur

Seal's sportive lemur

Lepilemur seali

Seal's sportive lemur (*Lepilemur seali*), or the Anjanaharibe-Sud sportive lemur, is a sportive lemur endemic to Madagascar. It is a large sportive lemur with a total length of about 57 to 64 cm (22 to 25 in), of which 25 to 28 cm (9.8 to...

Seba's short-tailed bat

Seba's short-tailed bat

Carollia perspicillata

Seba's short-tailed bat (*Carollia perspicillata*) is a common and widespread bat species in the family Phyllostomidae. They are found in Central America, the northern parts of South America, and in the Antilles islands.

Sechuran Fox

Sechuran Fox

Lycalopex sechurae

The Sechuran fox (*Lycalopex sechurae*) is a small South American species of canid. It is closely related to other South American "false" foxes or zorro and is one of ten extant species of canid endemic to South America.

Secretive dwarf squirrel

Secretive dwarf squirrel

Prosciurillus abstrusus

The secretive dwarf squirrel (*Prosciurillus abstrusus*) is a species of rodent in the family Sciuridae. It is endemic to Sulawesi, Indonesia, where it is found on the Mengkoka Mountains in southeastern Sulawesi. Its natural habitat is montane...

Sei Whale

Sei Whale

Balaenoptera borealis

The Sei whale (*Balaenoptera borealis*) is a baleen whale, the third-largest rorqual after the Blue whale and the Fin whale. It inhabits most oceans and adjoining seas and prefers deep offshore waters. The whale's name comes from the Norwegian word...

Selangor pygmy flying squirrel

Selangor pygmy flying squirrel

Petaurillus kinlochii

The Selangor pygmy flying squirrel (*Petaurillus kinlochii*) is a species of rodent in the family Sciuridae. It is endemic to Malaysia where it has only been found in the State of Selangor.

Selangor silvered langur

Selangor silvered langur

Trachypithecus selangorensis

The Selangor silvered langur (*Trachypithecus selangorensis*) is a species of leaf monkey found on the west coast of Peninsular Malaysia. It had been previously considered a form of silvery lutung. Roos and colleagues elevated this population to a...

Selinda veld rat

Selinda veld rat

Aethomys silindensis

The Selinda veld rat or Silinda rock rat (*Aethomys silindensis*) is a species of rodent in the family Muridae found in possibly Mozambique and Zimbabwe. Its natural habitat is subtropical or tropical dry forests.

Selous' zebra

Selous' zebra

Equus quagga selousi

Selous' zebra (*Equus quagga selousi*) is a subspecies of the plains zebra distributed across southeastern Africa. It is found mostly in Mozambique.

Selous's mongoose

Selous's mongoose

Paracynictis selousi

Selous's mongoose (*Paracynictis selousi*) is a mongoose species native to Southern Africa. It is the only member of the genus *Paracynictis*.

Seminole bat

Seminole bat

Lasiurus seminolus

The Seminole bat (*Lasiurus seminolus*) is a species of bat in the family Vespertilionidae.

Semliki red colobus

Semliki red colobus

Piliocolobus semlikiensis

The Semliki red colobus (*Piliocolobus semlikiensis*) is a type of red colobus monkey from central Africa. Historically it has been treated as a subspecies of the Central African red colobus (*P. foai*) but more recent taxonomies generally treat it...

Semon's leaf-nosed bat

Semon's leaf-nosed bat

Hipposideros semoni

Semon's leaf-nosed bat or Semon's roundleaf bat (*Hipposideros semoni*) is a species of bat in the family Hipposideridae. It is found in Australia and Papua New Guinea.

Senegal Bushbaby

Senegal Bushbaby

Galago senegalensis

The Senegal bushbaby (*Galago senegalensis*) is a small, nocturnal primate, a member of the galago family Galagidae. It is thought that the name "bush baby" may come either from the animals' cries or from their appearance.

Senegal gerbil

Senegal gerbil

Taterillus pygargus

The Senegal gerbil or Senegal tateril (*Taterillus pygargus*) is a species of rodent found in Niger, Senegal, possibly Gambia, possibly Mali, and possibly Mauritania. Its natural habitats are dry savanna, subtropical or tropical dry shrubland,...

Senegal one-striped grass mouse

Senegal one-striped grass mouse

Lemniscomys linulus

Lemniscomys linulus, commonly known as the Senegal grass mouse or Senegal one-striped grass mouse, is a species of rodent in the family Muridae. It is found in Ivory Coast, Guinea, Mali, and Senegal and its natural habitat is dry savanna. At one...

Senegalese wolf

Senegalese wolf

Canis lupaster anthus

The Senegalese wolf (*Canis lupaster anthus*), also known as the grey jackal, slender jackal or anthus, is the nominate subspecies of the African golden wolf native to Senegal.

Senkaku mole

Senkaku mole

Mogera uchidai

The Senkaku mole (*Mogera uchidai*), also known as the Ryukyu mole, is a species of mammal in the family Talpidae. It was formerly classified as being the only species in the genus *Nesosaptor*. It is endemic to the Uotsuri-jima (Japanese: 魚釣島;...

Sepia short-tailed opossum

Sepia short-tailed opossum

Monodelphis adusta

The sepia short-tailed opossum (*Monodelphis adusta*) is a species of opossum in the family Didelphidae found in Colombia, Ecuador, Panama, Peru and Venezuela. Its habitat consists of different types of forests up to 2200 meters above sea level, as...

Seram bandicoot

Seram bandicoot

Rhynchomeles prattorum

The Seram bandicoot (*Rhynchomeles prattorum*), also known as the Seram Island long-nosed bandicoot, is a member of the order Peramelemorphia that is endemic to the island of Seram in Indonesia. It is the only species in the genus *Rhynchomeles*.

Seram long-tailed mosaic-tailed rat

Seram long-tailed mosaic-tailed rat

Melomys fulgens

The Seram long-tailed mosaic-tailed rat (*Melomys fulgens*), also known as the Seram long-tailed melomys, is a species of rodent in the family Muridae. It is found only on the south coast of the island of Seram in Indonesia. At one time it was...

Seri's sheath-tailed bat

Seri's sheath-tailed bat

Emballonura serii

Seri's sheath-tailed bat (*Emballonura serii*) is a species of sac-winged bat in the family Emballonuridae. It is found in the Bismarck Archipelago (Papua New Guinea) and Yapen Island (Indonesia). Its natural habitat is caves.

Seri's tree-kangaroo

Seri's tree-kangaroo

Dendrolagus stellarum

Seri's tree-kangaroo (*Dendrolagus stellarum*) is a species of tree-kangaroo native and endemic to montane forests of west-central New Guinea. Seri's tree-kangaroo was originally considered to be a subspecies of *Dendrolagus dorianus*, but was...

Serotine Bat

Serotine Bat

Eptesicus serotinus

The Serotine bat (*Eptesicus serotinus*) is a fairly large Eurasian bat with quite large ears. It has a wingspan of around 37 cm (15 in) and often hunts in woodland. Its name serotine is derived from the Latin *serotinus*, which means 'evening',...

Serra do Mar grass mouse

Serra do Mar grass mouse

Akodon serrensis

The Serra do Mar grass mouse or Cerrado grass mouse (*Akodon serrensis*) is a rodent species from South America. It is found in Brazil. Pardiñas et al. (2016) transferred this species from the genus *Akodon* to the separate akodontine genus *Castoria*....

Serval

Serval

Leptailurus serval

The serval (*Leptailurus serval*) is a wild cat native to Africa. It is rare in North Africa and the Sahel, but widespread in sub-Saharan countries, except rainforest regions. Across its range, it occurs in protected areas, and hunting it is...

Servaline Genet

Servaline Genet

Genetta servalina

The Servaline genet (*Genetta servalina*) is a slender cat-like animal from Central Africa where it is widely distributed and considered common. Genets have large eyes which they can move within their sockets to a limited extent. They also move...

Servant mouse

Servant mouse

Mus famulus

The servant mouse (*Mus famulus*) or Bonhote's mouse, is a species of rodent in the family Muridae. It is found only in the Western Ghats of South India, where it is restricted to Eravikulam National Park, Avalanche, Kalapatti, and Coonoor.

Setirostris

Setirostris

Setirostris eleryi

Setirostris eleryi is a species of small insectivorous bat found in inland eastern Australia. It is the sole species of the molossid genus *Setirostris*, a name that refers to the coarse bristles on their faces. Earlier common names have referred...

Setzer's mouse-tailed dormouse

Setzer's mouse-tailed dormouse

Myomimus setzeri

The Setzer's mouse-tailed dormouse (*Myomimus setzeri*) is a species of rodent in the family Gliridae. It is endemic to Iran. Very little information is available for this genus *Myomimus* member, with just only 10 specimens known. It is found...

Setzer's pygmy mouse

Setzer's pygmy mouse

Mus setzeri

Setzer's pygmy mouse (*Mus setzeri*) is a species of rodent in the family Muridae. It is found in Botswana, Namibia, and Zambia. Its natural habitats are dry savanna, rivers, freshwater lakes, and intermittent freshwater lakes.

Seurat's spiny mouse

Seurat's spiny mouse

Acomys seurati

Seurat's spiny mouse (*Acomys seurati*) is a species of rodent in the family Muridae. It is found only in Algeria. Its natural habitats are rocky areas and hot deserts.

Seven-banded armadillo

Seven-banded armadillo

Dasypus septemcinctus

The seven-banded armadillo (*Dasypus septemcinctus*), also known as the Brazilian lesser long-nosed armadillo, is a species of armadillo from South America found in Paraguay, Argentina, Bolivia and Brazil. It is a solitary nocturnal, terrestrial...

Severtzov's birch mouse

Severtzov's birch mouse

Sicista severtzovi

The Severtzov's birch mouse, or dark birch mouse (*Sicista severtzovi*) is a species of rodent in the family Sminthidae. It is endemic to East-European steppes (E Ukraine and S-W Russia).

Severtzov's jerboa

Severtzov's jerboa

Allactaga severtzovi

Severtzov's jerboa (*Allactaga severtzovi*) is an herbivorous species of rodent in the family Dipodidae. It is found in Kazakhstan, Tajikistan, Turkmenistan, and Uzbekistan.

Seychelles fruit bat

Seychelles fruit bat

Pteropus seychellensis

The Seychelles fruit bat or Seychelles flying fox (*Pteropus seychellensis*) is a megabat found on the granitic islands of Seychelles, and on the Comoros and Mafia Island. It is a significant component of the ecosystems for the islands, dispersing...

Seychelles sheath-tailed bat

Seychelles sheath-tailed bat

Coleura seychellensis

The Seychelles sheath-tailed bat (*Coleura seychellensis*) is a sac-winged bat found in the central granitic islands of the Seychelles. It is an insectivorous bat, feeding primarily in forest clearings at night and roosting in communal roosts by...

Shadowy broad-nosed bat

Shadowy broad-nosed bat

Platyrrhinus umbratus

The shadowy broad-nosed bat (*Platyrrhinus umbratus*) is a species of bat in the family Phyllostomidae. It is found in Colombia and Venezuela.

Shaggy bat

Shaggy bat

Centronycteris maximiliani

The shaggy bat (*Centronycteris maximiliani*) is a bat species from South America. It appears to be a slow flier and has a rather regular pattern of foraging in its

home range, a feature shared with other emballonurids. It is also aerial insectivore.

Shamel's horseshoe bat
Shamel's horseshoe bat
Rhinolophus shameli

Shamel's horseshoe bat (*Rhinolophus shameli*) is a species of bat in the family Rhinolophidae. It is found in Cambodia, Laos, Myanmar, Thailand and Vietnam. The eponym for the species name "shameli " was American mammalogist H. Harold Shamel.
Shansei vole
Shansei vole
Myodes shanseius

The Shansei vole (*Myodes shanseius*) is a species of rodent in the family Cricetidae. It is found only in north-central China where its habitat is forests.
Shanxi sika deer
Shanxi sika deer
Cervus nippon grassianus

The Shanxi sika deer (*Cervus nippon grassianus*) is a possibly extinct subspecies of the sika deer. It is also one of its largest, being 105–110 cm tall at the shoulders and weighing 100 kg. The color is tawny or grayish dark, and is brown on the...
Shark Bay Mouse
Shark Bay Mouse
Pseudomys fieldi

Gould's mouse (*Pseudomys gouldii*), also known as the Shark Bay mouse and djoongari in the Pintupi and Luritja languages, is a species of rodent in the murid family. Once ranging throughout Australia from Western Australia to New South Wales, its...
Sharpe's Grysbok
Sharpe's Grysbok
Raphicerus sharpei

Sharpe's or northern grysbok (*Raphicerus sharpei*) is a small, shy, solitary antelope that is found from tropical to south-eastern Africa.
Shaw Mayer's brush mouse
Shaw Mayer's brush mouse
Pogonomelomys mayeri

Shaw Mayer's brush mouse (*Pogonomelomys mayeri*) is a species of rodent in the family Muridae. It is found in West Papua, Indonesia and Papua New Guinea.
Shaw Mayer's water rat
Shaw Mayer's water rat
Baiyankamys shawmayeri

Shaw Mayer's water rat (*Baiyankamys shawmayeri*) is a semiaquatic species of rodent in the family Muridae. It is found in the mountains of Papua New Guinea.
Shaw's jird
Shaw's jird
Meriones shawi

Shaw's jird (*Meriones shawi*) is a species of rodent in the family Muridae. It is found in Algeria, Egypt, Libya, Morocco, and Tunisia. Its natural habitats are arable land, pastureland, and rural gardens. They typically live 1–2 years. Shaw's jird...
Sheath-tailed mouse
Sheath-tailed mouse
Mus fragilicauda

The sheath-tailed mouse (*Mus fragilicauda*) is a mouse found in two locations in central Thailand and in Laos. They were discovered and documented in 2002. It is the only known *Mus* species to lose its tail integument when handled. It is sometimes...

Shepherd's Beaked Whale
Shepherd's Beaked Whale
Tasmacetus shepherdi

Shepherd's beaked whale (*Tasmacetus shepherdi*) was first known to science in 1937. It was named by W. R. B. Oliver after George Shepherd, curator of the Wanganui Museum, who collected the type specimen near Ohawe on the south Taranaki coast of...

Sherman's Fox Squirrel
Sherman's Fox Squirrel
Sciurus niger shermani

Sherman's fox squirrel (*Sciurus niger shermani*) is a subspecies of the Fox squirrel found in the United States. It nests in trees and has lost much of its habitat to farming and development.

Shield-faced roundleaf bat
Shield-faced roundleaf bat
Hipposideros lylei

The shield-faced roundleaf bat (*Hipposideros lylei*) is a species of bat in the family Hipposideridae. It is found in China, Laos, Malaysia, Myanmar, Thailand and Vietnam.

Shield-nosed leaf-nosed bat
Shield-nosed leaf-nosed bat
Hipposideros scutinares

The shield-nosed leaf-nosed bat or shield-nosed roundleaf bat (*Hipposideros scutinares*) is a bat from Laos and Vietnam.

Shining thicket rat
Shining thicket rat
Grammomys poensis

The shining thicket rat (*Grammomys poensis*) is a species of rodent in the family Muridae.

Shinto shrew
Shinto shrew
Sorex shinto

The Shinto shrew (*Sorex shinto*) is a species of shrew of the genus *Sorex* that lives only on the islands of Japan. It is a mole-like mammal with a pointed snout, very small ears, and a relatively long tail. Like most shrews, it is tiny, has poor...

Shiny guinea pig
Shiny guinea pig
Cavia fulgida

The shiny guinea pig (*Cavia fulgida*) is a guinea pig species of southeastern South America. The rodent is endemic to Brazil. It is native to the Atlantic Forest ecoregions.

Shipton's mountain cavy
Shipton's mountain cavy
Microcavia shiptoni

Shipton's mountain cavy (*Microcavia shiptoni*) is a species of rodent in the family

Caviidae. It is endemic to Argentina.
Shock-headed capuchin
Shock-headed capuchin
Cebus cuscinus

The shock-headed capuchin (*Cebus cuscinus*) is a species of gracile capuchin monkey from Bolivia and Peru. It was previously classified as a subspecies of the Humboldt's white-fronted capuchin (*C. albifrons*), but in 2013 Mittermeier and Rylands...
Short-Beaked Common Dolphin
Short-Beaked Common Dolphin
Delphinus delphis

The common dolphin (*Delphinus delphis*) is the most abundant cetacean in the world, with a global population of about six million. Despite this fact and its vernacular name, the common dolphin is not thought of as the archetypal dolphin, with...
Short-Beaked Echidna
Short-Beaked Echidna
Tachyglossus aculeatus

The Short-beaked echidna (*Tachyglossus aculeatus*) is one of four living species of echidna. It is covered in fur and spines and has a distinctive snout and a specialized tongue, which it uses to catch its insect prey at a great speed. The...
Short-eared bat
Short-eared bat
Cyttarops alecto

The short-eared bat (*Cyttarops alecto*) is a bat species from South and Central America. It is found in Brazil, Costa Rica, Guyana and Nicaragua. It is monotypic within its genus.
Short-Eared Dog
Short-Eared Dog
Atelocynus microtis

The short-eared dog (*Atelocynus microtis*), also known as the short-eared zorro or small-eared dog, is a unique and elusive canid species endemic to the Amazonian basin. This is the only species assigned to the genus *Atelocynus*.
Short-eared possum
Short-eared possum
Trichosurus caninus

The short-eared possum (*Trichosurus caninus*) is a species of marsupial in the family Phalangeridae endemic to Australia. Found north of Sydney, the species was once classed as a mountain brushtail possum, which is its closest relative. In the...
Short-eared rock-wallaby
Short-eared rock-wallaby
Petrogale brachyotis

The short-eared rock-wallaby (*Petrogale brachyotis*) is a species of rock-wallaby found in northern Australia, in the northernmost parts of the Northern Territory and Western Australia. It is much larger than its three closest relatives, the...
Short-faced mole
Short-faced mole
Scaptochirus moschatus

The short-faced mole (*Scaptochirus moschatus*) is a species of mammal in the family Talpidae. It is monotypic within the genus *Scaptochirus*. It is endemic to China.
Short-Finned Pilot Whale

Short-Finned Pilot Whale
Globicephala macrorhynchus

The short-finned pilot whale (*Globicephala macrorhynchus*) is one of the two species of cetaceans in the genus *Globicephala*, which it shares with the long-finned pilot whale (*G. melas*). It is part of the oceanic dolphin family (*Delphinidae*).It...

Short-footed Luzon tree rat
Short-footed Luzon tree rat
Carpomys melanurus

The short-footed Luzon tree rat or greater dwarf cloud rat (*Carpomys melanurus*) is a species of rodent in the family *Muridae*. It is found only in the Philippines, specifically in northern Luzon. Its natural habitat is tropical moist montane...

Short-furred Atlantic tree-rat
Short-furred Atlantic tree-rat
Phyllomys unicolor

The short-furred Atlantic tree-rat or Wagner's Atlantic tree rat (*Phyllomys unicolor*), is a spiny rat species from South America. It is found in Brazil.

Short-furred dasyure
Short-furred dasyure
Murexia longicaudata

The short-furred dasyure (*Murexia longicaudata*), also known as the short-haired marsupial mouse, is a member of the order *Dasyuromorphia*. It was once recognised as the only species in the genus *Murexia*, but now five species are recognised. It...

Short-haired water rat
Short-haired water rat
Paraleptomys wilhelmina

The short-haired water rat (*Paraleptomys wilhelmina*) is a species of rodent in the family *Muridae*.It is found in West Papua, Indonesia and Papua New Guinea. There are two currently undescribed species allied with this taxon.

Short-headed broad-nosed bat
Short-headed broad-nosed bat
Platyrrhinus brachycephalus

The short-headed broad-nosed bat (*Platyrrhinus brachycephalus*) is a bat species from South America. It is found in Bolivia, northwestern Brazil, Colombia, Ecuador, French Guiana, Guyana, Peru, Suriname and Venezuela.

Short-headed roundleaf bat
Short-headed roundleaf bat
Hipposideros breviceps

The short-headed roundleaf bat (*Hipposideros breviceps*) is a species of bat in the family *Hipposideridae* endemic to Indonesia. It is known only from the type specimen.

Short-nosed harvest mouse
Short-nosed harvest mouse
Reithrodontomys brevirostris

The short-nosed harvest mouse (*Reithrodontomys brevirostris*) is a species of rodent in the family *Cricetidae* found in Costa Rica and Nicaragua.

Short-palated fruit bat
Short-palated fruit bat
Casinycteris argynnis

The short-palated fruit bat (*Casinycteris argynnis*) is a species of frugivorous

megabat in the family Pteropodidae. It is found in Cameroon, Central African Republic, and Democratic Republic of the Congo. Its natural habitat is subtropical or...

Short-snouted elephant shrew
Short-snouted elephant shrew
Elephantulus brachyrhynchus

The short-snouted elephant shrew or short-snouted sengi (*Elephantulus brachyrhynchus*) is a species of elephant shrew in the family Macroscelididae. It is found over a wide area of Africa. Its natural habitats are dry savanna and subtropical or...

Short-tailed bandicoot rat
Short-tailed bandicoot rat
Nesokia indica

The short-tailed bandicoot rat (*Nesokia indica*) is a species of rodent in the family Muridae. Other common names include short-tailed mole-rat, Indian bandicoot, bandicoot-rat, flat-tooth rat and short-tailed nesokia.

Short-tailed brush-furred rat
Short-tailed brush-furred rat
Lophuromys brevicaudus

The short-tailed brush-furred rat (*Lophuromys brevicaudus*) is a species of rodent in the family Muridae. It is found only in Ethiopia. Its natural habitats are subtropical or tropical high-altitude shrubland and subtropical or tropical...

Short-Tailed Chinchilla
Short-Tailed Chinchilla
Chinchilla chinchilla

The short-tailed chinchilla (*Chinchilla chinchilla*; formerly *C. brevicaudata*)—also called the Bolivian, Peruvian, or royal chinchilla—is an endangered species of South American rodent, and one of two species in the genus *Chinchilla*. Their...

Short-tailed gymnure
Short-tailed gymnure
Hylomys suillus

The short-tailed gymnure (*Hylomys suillus*) is a small mammal from the family of the Erinaceidae. The scientific name of the species is first published by Salomon Müller in 1840.

Short-tailed mongoose
Short-tailed mongoose
Herpestes brachyurus

The short-tailed mongoose (*Urva brachyura*) is a mongoose species native to Peninsular Malaysia, Sumatra and Borneo. It inhabits evergreen forest and rural gardens from sea level to an elevation of 1,500 m (4,900 ft). It is listed as Near...

Short-tailed roundleaf bat
Short-tailed roundleaf bat
Hipposideros curtus

The short-tailed roundleaf bat (*Hipposideros curtus*) is a species of bat in the family Hipposideridae. It is found in Cameroon and Equatorial Guinea. Its natural habitats are subtropical or tropical moist lowland forests and caves. It is...

Short-tailed shrew tenrec
Short-tailed shrew tenrec
Microgale brevicaudata

The short-tailed shrew tenrec (*Microgale brevicaudata*) is a species of mammal in

the family Tenrecidae. It is endemic to Madagascar. Its natural habitats are subtropical or tropical dry forest and subtropical or tropical moist lowland forest. It...

Short-tailed spiny rat
Short-tailed spiny rat
Proechimys brevicauda

The short-tailed spiny-rat, *Proechimys brevicauda*, or Huallaga spiny rat, is a spiny rat species from South America. It is found in Bolivia, Brazil, Colombia, Ecuador and Peru.

Short-tailed Talaud mosaic-tailed rat
Short-tailed Talaud mosaic-tailed rat
Melomys caurinus

The short-tailed Talaud mosaic-tailed rat or the short-tailed Talaud melomys (*Melomys caurinus*) is a species of rodent in the family Muridae. It is endemic to Karakelong and Salebabu in the Talaud Islands in Indonesia where it occurs in forest...

Shortridge's horseshoe bat
Shortridge's horseshoe bat
Rhinolophus shortridgei

Shortridge's horseshoe bat (*Rhinolophus shortridgei*) is a species of Horseshoe bat native to Northeast India, northern Myanmar, and southern China. It was first described in 1918 by Knud Andersen, and was considered a subspecies of Blyth's...

Shortridge's langur
Shortridge's langur
Trachypithecus shortridgei

The Shortridge's langur (*Trachypithecus shortridgei*) is a lutung native to Burma and China.

Shortridge's long-fingered bat
Shortridge's long-fingered bat
Miniopterus shortridgei

Shortridge's long-fingered bat (*Miniopterus shortridgei*) is a bat in the genus *Miniopterus* which occurs throughout Indonesia. It was described by Eleanor Mary Ord Laurie and John Edwards Hill in 1957.

Shortridge's mouse
Shortridge's mouse
Mus shortridgei

Shortridge's mouse (*Mus shortridgei*) is a species of rodent in the family Muridae. It is found in Cambodia, Laos, Myanmar, Thailand, and Vietnam.

Shortridge's multimammate mouse
Shortridge's multimammate mouse
Mastomys shortridgei

Shortridge's multimammate mouse (*Mastomys shortridgei*) is a rodent species in the family Muridae. It is native to Angola, Botswana and Namibia. Its natural habitats are moist savanna, subtropical or tropical seasonally wet or flooded lowland...

Shortridge's rat
Shortridge's rat
Thallomys shortridgei

Shortridge's rat (*Thallomys shortridgei*) is a species of rodent in the family Muridae. It is found only in South Africa. Its natural habitat is subtropical or tropical dry shrubland.

Shortridge's rock mouse

Shortridge's rock mouse
Petromyscus shortridgei

Shortridge's rock mouse (*Petromyscus shortridgei*) is a species of rodent in the family Nesomyidae. It is found in Angola and Namibia. Its natural habitat is subtropical or tropical dry shrubland.

Shrew gymnure
Shrew gymnure
Neotetracus sinensis

The shrew gymnure, or shrew hedgehog (*Neotetracus sinensis*), is a species of mammal in the family Erinaceidae and is the only extant species in the genus *Neotetracus*. It is found in China, Myanmar, and Vietnam.

Shrew-faced squirrel
Shrew-faced squirrel
Rhinosciurus laticaudatus

The shrew-faced squirrel (*Rhinosciurus laticaudatus*), also known as the long-nosed squirrel, is a species of rodent in the family Sciuridae. It is monotypic within the genus *Rhinosciurus*. It is found in forests in Peninsular Malaysia (possibly...

Shrew-toothed shrew tenrec
Shrew-toothed shrew tenrec
Microgale soricoides

The shrew-toothed shrew tenrec (*Microgale soricoides*) is a species of mammal in the family Tenrecidae. It is endemic to Madagascar. Its natural habitats are subtropical or tropical moist lowland forest and subtropical or tropical moist montane...

Siamang Gibbon
Siamang Gibbon
Symphalangus syndactylus

The Siamang gibbon (*Symphalangus syndactylus*) is an arboreal, black-furred gibbon native to the forests of Southeast Asia. The largest of the gibbons, the siamang can be twice the size of other gibbons. It is the only species in the genus...

Siau Island tarsier
Siau Island tarsier
Tarsius tumpara

The Siau Island tarsier (*Tarsius tumpara*) is a species of tarsier from the tiny volcanic island of Siau in Indonesia. The *T. tumpara* species is one of 14 species and 7 subspecies in the tarsier family called Tarsiidae. They belong to the...

Siberian brown lemming
Siberian brown lemming
Lemmus sibiricus

The West Siberian lemming or Western Siberian brown lemming (*Lemmus sibiricus*) is a true lemming species found in the Russian Federation. Like other lemmings, it belongs to the Cricetidae family of rodents. It is endemic to Russia, where it has...

Siberian Chipmunk
Siberian Chipmunk
Eutamias sibiricus

The Siberian chipmunk (*Eutamias sibiricus*), also called common chipmunk is native to northern Asia from central Russia to China, Korea, and Hokkaidō in northern Japan. It was imported from South Korea and introduced in Europe as a pet in the 1960s.

Siberian Flying Squirrel

Siberian Flying Squirrel
Pteromys volans

The Siberian flying squirrel (*Pteromys volans*) is an Old World flying squirrel with a range from the Baltic Sea in the west to the Pacific Coast in the east. It is the only species of flying squirrel found in Europe. It is considered vulnerable...

Siberian Ibex
Siberian Ibex
Capra sibirica

The Siberian ibex (*Capra sibirica*), also known as the Altai ibex, Central Asia(n) ibex, Gobi ibex, Himalayan ibex, Mongolian ibex or Tian Shan ibex, is a species of ibex that lives in central Asia. It has traditionally been treated as a...

Siberian large-toothed shrew
Siberian large-toothed shrew
Sorex daphaenodon

The Siberian large-toothed shrew (*Sorex daphaenodon*) is a species of shrew. An adult Siberian large-toothed shrew has a weight of 4.6-6.0 grams and a body length of 5.5-6.4 centimeters, with a tail of 2.4-3.75 centimeters. This species is found...

Siberian Lynx
Siberian Lynx
Lynx lynx wrangeli

The Siberian lynx (*Lynx lynx wrangeli*) is a subspecies of Eurasian lynx living in the Russian Far East. It is the second most common subspecies of the Eurasian lynx.

Siberian Musk Deer
Siberian Musk Deer
Moschus moschiferus

The Siberian musk deer (*Moschus moschiferus*) is a musk deer found in the mountain forests of Northeast Asia. The small shape of these animals allows them to hide from predators through tiny openings in the rocky terrain and also allows them to...

Siberian Roe Deer
Siberian Roe Deer
Capreolus pygargus

The Siberian roe deer, eastern roe deer, or Asian roe (*Capreolus pygargus*), is a species of roe deer found in northeastern Asia. In addition to Siberia and Mongolia, it is found in Kazakhstan, the Tian Shan Mountains of Kyrgyzstan, eastern...

Siberian shrew
Siberian shrew
Crocidura sibirica

The Siberian shrew (*Crocidura sibirica*) is a species of mammal in the family Soricidae. It is found in Russia, possibly China, and possibly Mongolia.

Siberian Weasel
Siberian Weasel
Mustela sibirica

Siberian weasels (*Mustela sibirica*) are medium-sized weasels native to Asia. They live in various forests and open areas. They build their nests inside fallen logs and often line their nesting chambers with bird feathers and rodent hair.

Siberian zokor
Siberian zokor

Myospalax myospalax

The Siberian zokor (*Myospalax myospalax*) is a species of rodent in the family Spalacidae. It is found in Kazakhstan and Russia.

Siberut flying squirrel

Siberut flying squirrel

Petinomys lugens

The Siberut flying squirrel (*Petinomys lugens*) is a species of rodent in the family Sciuridae. It is endemic to Indonesia. Its natural habitat is subtropical or tropical dry forests. It is threatened by habitat loss.

Siberut macaque

Siberut macaque

Macaca siberu

The Siberut macaque (*Macaca siberu*) is a vulnerable species of macaque, which is endemic to Siberut Island in Indonesia. It was formerly considered conspecific with the Pagai Island macaque (*M. pagensis*) which is overall paler in color, but...

Sibree's dwarf lemur

Sibree's dwarf lemur

Cheirogaleus sibreei

Sibree's dwarf lemur (*Cheirogaleus sibreei*) is a small nocturnal lemur endemic to Madagascar. The name of this dwarf lemur commemorates the English missionary and naturalist James Sibree (1836–1929). In 2010, a research team confirmed that they...

Sibuyan shrew

Sibuyan shrew

Crocidura ninoyi

The Sibuyan shrew (*Crocidura ninoyi*) is a species of shrew from the Philippines.

Sibuyan striped shrew-rat

Sibuyan striped shrew-rat

Chrotomys sibuyanensis

Sibuyan striped shrew rat (*Chrotomys sibuyanensis*) is a species of rodent in the family Muridae. The holotype was collected in 1992; however, it wasn't formally described until 2005.

Sichuan deer

Sichuan deer

Cervus canadensis macneilli

The Sichuan deer (*Cervus canadensis macneilli*), also known as MacNeill's deer, is a subspecies of the wapiti native to Western China.

Sichuan field mouse

Sichuan field mouse

Apodemus latronum

The Sichuan field mouse (*Apodemus latronum*) is a species of rodent in the family Muridae. It is found in Qinghai and Sichuan provinces of China, and in India and Burma.

Sichuan sika deer

Sichuan sika deer

Cervus nippon sichuanicus

The Sichuan sika deer (*Cervus nippon sichuanicus*) is one of the many subspecies of sika deer. It was discovered in 1978 and was proven to be a distinct subspecies. It is found in the mountains of northern Sichuan and southern Gansu, which holds...

Sichuan takin

Sichuan takin

Budorcas taxicolor tibetana

The Sichuan takin or Tibetan takin (*Budorcas taxicolor tibetana*) is a subspecies of takin (goat-antelope). *Budorcas* from Greek *bous* ("ox" or "cow") and *dorkas* ("gazelle"); *taxicolor* from Latin *taxus* ("badger") and *color* ("hue") referring to...

Sicilian shrew

Sicilian shrew

Crocidura sicula

The Sicilian shrew (*Crocidura sicula*) is a species of mammal in the family Soricidae. It is found in Sicily (Italy) and Gozo (Malta). Its natural habitat is temperate shrubland.

Sicilian wolf

Sicilian wolf

Canis lupus cristaldii

The Sicilian wolf (*Canis lupus cristaldii*) (Sicilian: *lupu sicilianu*) is an extinct subspecies of the gray wolf that was endemic to Sicily. It was paler than the mainland Italian wolf and comparable in size to the extant Arabian wolf and extinct...

Sicista concolor

Sicista concolor

Sicista concolor

Sicista concolor, the Chinese birch mouse, is a species of rodent in the family Sminthidae. It is native to China, India and Pakistan.

Side-Striped Jackal

Side-Striped Jackal

Canis adustus

The Side-striped jackal (*Lupulella adusta*) is a canine native to Africa. Unlike the smaller and related Black-backed jackal, which dwells in open plains, the Side-striped jackal prefers to live in woodland and scrub areas.

Sierra de Perijá white-fronted capuchin

Sierra de Perijá white-fronted capuchin

Cebus leucocephalus

The Sierra de Perijá white-fronted capuchin (*Cebus leucocephalus*) is a species of gracile capuchin monkey from Colombia and Venezuela. It had formerly been regarded as a subspecies of the Humboldt's white-fronted capuchin but was reclassified by...

Sierra del Tontal chinchilla rat

Sierra del Tontal chinchilla rat

Abrocoma shistacea

The Sierra del Tontal chinchilla rat (*Abrocoma shistacea*) is a species of chinchilla rat. Found only in Argentina, this species was once thought to be part of *Abrocoma cinerea*. In 2002, Braun and Mares from the University of Oklahoma confirmed...

Sierra Leone free-tailed bat

Sierra Leone free-tailed bat

Mops brachypterus

The Sierra Leone free-tailed bat (*Mops brachypterus*) is a species of bat in the family Molossidae. It is found in Cameroon, Ivory Coast, Equatorial Guinea, Gabon, Ghana, Guinea, Kenya, Liberia, Mozambique, Nigeria, Sierra Leone, Tanzania, Togo,...

Sierra Madre forest mouse

Sierra Madre forest mouse

Apomys sierrae

The Sierra Madre forest mouse (*Apomys sierrae*) is a forest mouse endemic to the Sierra Madre Range of eastern Luzon, Philippines.

Sierra Madre ground squirrel

Sierra Madre ground squirrel

Callospermophilus madrensis

The Sierra Madre ground squirrel (*Callospermophilus madrensis*) is a species of rodent in the squirrel family. It is endemic to the Sierra Madre Occidental, in northern Mexico. Its natural habitat is subtropical or tropical dry lowland grassland.

Sierra Madre shrew mouse

Sierra Madre shrew mouse

Soricomys musseri

The Sierra Madre shrew mouse (*Soricomys musseri*) is a species of mammal from the Philippines.

Sierra Nevada Bighorn Sheep

Sierra Nevada Bighorn Sheep

Ovis canadensis sierrae

The Sierra Nevada bighorn sheep (*Ovis canadensis sierrae*) is a subspecies of Bighorn sheep found only in the Sierra Nevada mountains of California. It was listed as a federally endangered subspecies in 2000. In 2016, over 600 Sierra bighorn...

Sierra Nevada Red Fox

Sierra Nevada Red Fox

Vulpes vulpes necator

The Sierra Nevada red fox (*Vulpes vulpes necator*) is a subspecies of Red fox. It is likely one of the most endangered mammals in North America. The High Sierra fox shares most of its physical characteristics with the Red fox, though it is...

Sierra Tontal tuco-tuco

Sierra Tontal tuco-tuco

Ctenomys tulduco

The Sierra Tontal tuco-tuco (*Ctenomys tulduco*) is a species of rodent in the family Ctenomyidae. It is known only from a small locality in the Sierra del Tontal, San Juan Province, west central Argentina.

Sigmodontomys alfari

Sigmodontomys alfari

Sigmodontomys alfari

Sigmodontomys alfari, also known as the short-tailed sigmodontomys, Alfaro's rice water rat, Cana rice rat, or Allen's rice rat, is a species of rodent in the subfamily Sigmodontinae of family Cricetidae. It is found from Honduras through...

Sika Deer

Sika Deer

Cervus nippon

The Sika deer (*Cervus nippon*) is a species of deer native to much of East Asia and introduced to other parts of the world. Its name comes from the Japanese word shika which means "deer". In Japan, the species is known as the nihonjika (Japan...

Sikkim mountain vole

Sikkim mountain vole

Neodon sikimensis

The Sikkim mountain vole (*Neodon sikimensis*) is a species of rodent in the family

Cricetidae. It is found in Bhutan, India, Nepal and China.

Sikkim rat

Sikkim rat

Rattus andamanensis

The Sikkim rat (*Rattus andamanensis*) is a species of rodent in the family Muridae. It is found in Bhutan, Cambodia, China, India, Laos, Myanmar, Nepal, Thailand, and Vietnam. The rat's coloring is brownish upperparts and a white underside. *Rattus...*

Silent dormouse

Silent dormouse

Graphiurus surdus

The silent dormouse (*Graphiurus surdus*) is a species of rodent in the family Gliridae. It is found in Cameroon, Democratic Republic of the Congo, Equatorial Guinea, and Gabon. Its natural habitat is subtropical or tropical moist lowland forests.

Silent grass mouse

Silent grass mouse

Akodon surdus

The silent grass mouse or slate-bellied akodont (*Akodon surdus*) is a species of rodent in the family Cricetidae. It is endemic to Peru.

Silky Anteater

Silky Anteater

Cyclopes didactylus

The silky anteater, also known as the pygmy anteater, has traditionally been considered a single species of anteater, *Cyclopes didactylus*, in the genus *Cyclopes*, the only living genus in the family Cyclopedidae. Found in southern Mexico, and...

Silky cuscus

Silky cuscus

Phalanger sericeus

The silky cuscus (*Phalanger sericeus*) is a species of marsupial in the family Phalangeridae. It is found in Indonesia and Papua New Guinea.

Silky mouse

Silky mouse

Pseudomys apodemoides

The silky mouse (*Pseudomys apodemoides*) is a species of rodent in the family Muridae. It is found only in Australia.

Silky Oldfield mouse

Silky Oldfield mouse

Thomasomys bombycinus

The silky Oldfield mouse (*Thomasomys bombycinus*) is a species of rodent in the family Cricetidae. It is found only in Colombia.

Silky pocket mouse

Silky pocket mouse

Perognathus flavus

The silky pocket mouse (*Perognathus flavus*) is a species of rodent in the family Heteromyidae. It is found in northern and central Mexico and the southwest region of the United States. It is a species of least concern, according to the IUCN,...

Silky short-tailed bat

Silky short-tailed bat

Carollia brevicauda

The silky short-tailed bat (*Carollia brevicauda*) is a bat species from South and Central America. It is found in Bolivia, Brazil, Colombia, Ecuador, French Guiana, Guyana, Panama, Peru, Suriname, Mexico and Venezuela.

Silky Sifaka

Silky Sifaka

Propithecus candidus

The silky sifaka (*Propithecus candidus*) is a large lemur characterized by long, silky, white fur. It has a very restricted range in northeastern Madagascar, where it is known locally as the simpona. It is one of the rarest mammals on Earth.

The...

Silky tuco-tuco

Silky tuco-tuco

Ctenomys sericeus

The silky tuco-tuco (*Ctenomys sericeus*) is a species of rodent in the family Ctenomyidae. It is endemic to Argentina.

Silver Dik-Dik

Silver Dik-Dik

Madoqua piacentinii

The Silver dik-dik (*Madoqua piacentinii*) is a small antelope found in Africa. It is the smallest species of dik-dik. These antelope are named for the alarm calls of the females which they make through their long, tubular snouts when they feel...

Silver fruit-eating bat

Silver fruit-eating bat

Dermanura glauca

The silver fruit-eating bat (*Dermanura glauca*) is a South American bat species of the family Phyllostomidae.

Silver monkey

Silver monkey

Cercopithecus doggetti

The silver monkey (*Cercopithecus doggetti*) is a species of Old World monkey found primarily in East Africa. Its range includes Burundi, Tanzania, Rwanda, Uganda, and the Democratic Republic of the Congo. The silver monkey was previously...

Silver mountain vole

Silver mountain vole

Alticola argentatus

The silver mountain vole (*Alticola argentatus*) is a species of rodent in the family Cricetidae. They are distinguished by their silver-grey pelage, long vibrissae, rootless hypsodont molars and angular skull shape. Like many mammals of the...

Silver-Backed Chevrotain

Silver-Backed Chevrotain

Tragulus versicolor

The Silver-backed chevrotain (*Tragulus versicolor*) is a small mysterious deer known only from Vietnam. It was first described in 1910. Facing multiple threats in the wild, the Silver-backed chevrotain was feared to have gone extinct and in 2019,...

Silver-Haired Bat

Silver-Haired Bat

Lasionycteris noctivagans

The silver-haired bat (*Lasionycteris noctivagans*) is a solitary migratory species of vesper bat in the family Vespertilionidae and the only member of the genus

Lasionycteris.
Silver-headed antechinus
Silver-headed antechinus
Antechinus argentus

The silver-headed antechinus (*Antechinus argentus*) is a species of dasyurid marsupial of the genus *Antechinus*. Described in 2013, the species is known only from so far is known to occur in just three locations: Blackdown Tableland National Park,...

Silver-tipped myotis
Silver-tipped myotis
Myotis albescens

The silver-tipped myotis (*Myotis albescens*) is a species of mouse-eared bat found in a range of lowland habitats in the Americas. It is part of the vesper bat genus *Myotis*, which includes many common species across the world. Genetic analysis...

Silvered bat
Silvered bat
Glauconycteris argentata

The silvered bat (*Glauconycteris argentata*) is a species of vesper bat in the family *Vespertilionidae*. It is found in Angola, Burundi, Cameroon, Republic of the Congo, Democratic Republic of the Congo, Equatorial Guinea, Kenya, Malawi, Rwanda,...

Silvery flying fox
Silvery flying fox
Pteropus argentatus

The silvery flying fox (*Pteropus argentatus*), also known as Ambon flying fox, or silvery fruit bat, is a megabat in the genus *Pteropus* from eastern Indonesia. It is known from a single, damaged, and immature specimen that was thought to have...

Silvery Gibbon
Silvery Gibbon
Hylobates moloch

The silvery gibbon (*Hylobates moloch*), also known as the Javan gibbon, is a primate in the gibbon family *Hylobatidae*. It is endemic to the Indonesian island of Java, where it inhabits undisturbed rainforests up to an altitude of 2,450 m...

Silvery greater galago
Silvery greater galago
Otolemur monteiri

The silvery greater galago (*Otolemur monteiri*) is a nocturnal primate from the galago family. It is usually found in *Brachystegia* woodland, from Angola to Tanzania, western Kenya and Rwanda. The species was separated from the brown greater galago...

Silvery Lutung
Silvery Lutung
Trachypithecus cristatus

The silvery lutung (*Trachypithecus cristatus*), also known as the silvered leaf monkey or the silvery langur, is an Old World monkey. It is arboreal, living in coastal, mangrove, and riverine forests in Peninsular Malaysia, Sumatra, and Borneo. It...

Silvery Marmoset
Silvery Marmoset
Mico argentatus

The silvery marmoset (*Mico argentatus*) is a New World monkey that lives in the

eastern Amazon Rainforest in Brazil. The fur of the silvery marmoset is colored whitish silver-grey except for a dark tail. Remarkable are its naked, flesh-colored...

Silvery mole-rat

Silvery mole-rat

Heliophobius argenteocinereus

The silvery mole-rat, silvery blesmol, or silky mole-rat (*Heliophobius argenteocinereus*) is a species of mole-rat of East Africa which occurs in southern Kenya, Tanzania, southeastern Democratic Republic of Congo, Mozambique and Malawi. Solitary...

Silvery woolly monkey

Silvery woolly monkey

Lagothrix poeppigii

The silvery woolly monkey (*Lagothrix lagothricha poeppigii*), also known as Poeppig's woolly monkey or the red woolly monkey, is a subspecies of the common woolly monkey from South America. Named after the German zoologist Eduard Friedrich...

Simalur rat

Simalur rat

Rattus simalurensis

The Simalur rat (*Rattus simalurensis*) is a species of rodent in the family Muridae. It is found only in Indonesia, on Simalur island and the nearby islands of Sumat, Lasia, and Babi

Simmons' mouse lemur

Simmons' mouse lemur

Microcebus simmonsi

Simmons' mouse lemur (*Microcebus simmonsi*) is a species of mouse lemur known only from Betampona Special Reserve, Analalava Special Community Reserve near Foulpointe, and Zahamena National Park in Madagascar. The species is named in honor of Dr....

Simon's spiny rat

Simon's spiny rat

Proechimys simonsi

Simons's spiny rat, *Proechimys simonsi*, is a spiny rat species from South America. It is found in Bolivia, Brazil, Colombia, Ecuador and Peru. It was named for American scientific collector Perry O. Simons.

Sinaloan mastiff bat

Sinaloan mastiff bat

Molossus sinaloae

The Sinaloan mastiff bat (*Molossus sinaloae*) is a species of bat in the family Molossidae, native to Mexico, Central America and northern South America.

Sinaloan pocket mouse

Sinaloan pocket mouse

Chaetodipus pernix

The Sinaloan pocket mouse (*Chaetodipus pernix*) is one of 17 species of pocket mice in the genus *Chaetodipus*. Two subspecies of *C. pernix* are recognized, *C. p. pernix* and *C. p. rostratus*, all are endemic to Mexico. This small mouse lives among...

Sind bat

Sind bat

Rhyneptesicus nasutus

The Sind bat (*Rhyneptesicus nasutus*) is a species of vesper bat. It is found in

Afghanistan, Iran, Iraq, Oman, Pakistan, Saudi Arabia, and Yemen. It is the only member of the genus *Rhyneptesicus*.

Sindh Ibex

Sindh Ibex

Capra aegagrus blythi

The Sindh ibex (*Capra aegagrus blythi*) is a vulnerable subspecies of the Wild goat found in Asia. It is also known as the Turkman wild goat.

Singapore whiskered bat

Singapore whiskered bat

Vespertilio oreias

The Singapore whiskered bat (*Vespertilio oreias*) is or was a possible species of vesper bat endemic to Singapore. No specimens have been found since its original scientific description in 1840 by Dutch zoologist Coenraad Temminck. Modern...

Singing vole

Singing vole

Microtus miurus

The singing vole (*Microtus miurus*), is a medium-sized vole found in northwestern North America, including Alaska and northwestern Canada.

Single-striped grass mouse

Single-striped grass mouse

Lemniscomys rosalia

The single-striped grass mouse or single-striped lemniscomys (*Lemniscomys rosalia*) is a species of rodent in the family Muridae. It is found in Angola, Botswana, Kenya, Malawi, Mozambique, Namibia, South Africa, Eswatini, Tanzania, Zambia, and...

Sinnamary brush-tailed rat

Sinnamary brush-tailed rat

Isothrix sinnamariensis

The Sinnamary brush-tailed rat (*Isothrix sinnamariensis*), is a species of rodent in the family Echimyidae. It is found in French Guiana, Guyana, and possibly Suriname. Its natural habitat is subtropical or tropical moist lowland forests. It is...

Sipora flying squirrel

Sipora flying squirrel

Hylopetes sipora

The Sipora flying squirrel (*Hylopetes sipora*) is a species of rodent in the family Sciuridae. It is endemic to Indonesia. Its natural habitat is subtropical or tropical dry forests. It is threatened by habitat loss.

Sir David Attenborough's myotis

Sir David Attenborough's myotis

Myotis attenboroughi

Sir David Attenborough's myotis (*Myotis attenboroughi*) is a species of small bat in the family Vespertilionidae that is endemic to the Caribbean island of Tobago. Its presence on nearby Trinidad is as yet uncertain. It is the only mammal species...

Sir David's Long-Beaked Echidna

Sir David's Long-Beaked Echidna

Zaglossus attenboroughi

Sir David's long-beaked echidna (*Zaglossus attenboroughi*) is one of the three species from the genus *Zaglossus* that occurs in New Guinea. It is named in honor of Sir David Attenborough, the eminent naturalist.

Siskiyou chipmunk
Siskiyou chipmunk
Neotamias siskiyou

The Siskiyou chipmunk (*Neotamias siskiyou*) is a species of rodent in the family Sciuridae. It is endemic to northern California and central Oregon in the United States.

Sitatunga
Sitatunga
Tragelaphus spekii

The sitatunga or marshbuck (*Tragelaphus spekii*) is a swamp-dwelling antelope found throughout central Africa, centering on the Democratic Republic of the Congo, the Republic of the Congo, Cameroon, parts of Southern Sudan, Equatorial Guinea,...

Sitka deer
Sitka deer
Odocoileus hemionus sitkensis

The Sitka deer or Sitka black-tailed deer (*Odocoileus hemionus sitkensis*), is a subspecies of mule deer (*Odocoileus hemionus*), similar to the Columbian black-tailed subspecies (*O. h. columbianus*). Their name originates from Sitka, Alaska, and...

Six-Banded Armadillo
Six-Banded Armadillo
Euphractus sexcinctus

The Six-banded armadillo (*Euphractus sexcinctus*) is a species of mammal found in South America. It was first described by Swedish zoologist Carl Linnaeus in 1758. Six-banded armadillos are efficient diggers and due to their poor eyesight, rely on...

Skomer vole
Skomer vole
Myodes glareolus skomerensis

The Skomer vole (*Myodes glareolus skomerensis*) is a subspecies of bank vole endemic to the island of Skomer, off the west coast of Wales. The bank vole was probably introduced by humans at some time after the last glaciation. It is one of four...

Skywalker hoolock gibbon
Skywalker hoolock gibbon
Hoolock tianxing

The Skywalker hoolock gibbon or Gaoligong hoolock gibbon (*Hoolock tianxing*) is an arboreal primate in the gibbon family, Hylobatidae. It is one of three species of hoolock gibbon and was first described in January 2017 in the American Journal of...

Slender broad-nosed bat
Slender broad-nosed bat
Platyrrhinus angustirostris

The slender broad-nosed bat (*Platyrrhinus angustirostris*) is a species of bat in the family Phyllostomidae. As a phyllostomid bat, it is characterized by a narrow uropatagium which is fringed with hair; a white dorsal stripe; large inner upper...

Slender harvest mouse
Slender harvest mouse
Reithrodontomys gracilis

The slender harvest mouse(*Reithrodontomys gracilis*) is a species of rodent in the family Cricetidae. A small mouse-like rodent distributed throughout a portion

Central America.
Slender Mongoose
Slender Mongoose
Galereella sanguinea

The common slender mongoose (*Herpestes sanguineus*), also known as the black-tipped mongoose or the black-tailed mongoose, is a very common mongoose species native to sub-Saharan Africa.

Slender Oldfield mouse
Slender Oldfield mouse
Thomasomys gracilis

The slender Oldfield mouse (*Thomasomys gracilis*) is a species of rodent in the family Cricetidae. It is found in Ecuador and Peru.

Slender rat
Slender rat
Rattus verecundus

The slender rat (*Rattus verecundus*) is a species of rodent in the family Muridae. It is found in West Papua, Indonesia and Papua New Guinea.

Slender shrew
Slender shrew
Sorex gracillimus

The slender shrew (*Sorex gracillimus*) is a species of shrew. An adult slender shrew has a weight of 1.5-5.3 grams and a body length of 4.7-6.0 centimeters, with a tail of 4-5 centimeters; this makes it one of the smaller shrews found in its...

Slender squirrel
Slender squirrel
Sundasciurus tenuis

The slender squirrel (*Sundasciurus tenuis*) is a species of rodent in the family Sciuridae. It is arboreal and found in Indonesia, Singapore, Malaysia, and Thailand. The body is brown on the upper parts and light grey on the underparts. The body...

Slender treeshrew
Slender treeshrew
Tupaia gracilis

The slender treeshrew (*Tupaia gracilis*) is a treeshrew species within the Tupaiidae. It is native to Borneo and inhabits foremost lowland old forest.

Slender tube-nosed bat
Slender tube-nosed bat
Murina gracilis

The slender tube-nosed bat (*Murina gracilis*) is a species of vesper bat in the family Vespertilionidae found only in Taiwan.

Slender yellow bat
Slender yellow bat
Rhogeessa gracilis

The slender yellow bat (*Rhogeessa gracilis*) is a species of vesper bat. It is found only in Mexico.

Slender-tailed dunnart
Slender-tailed dunnart
Sminthopsis murina

The slender-tailed dunnart (*Sminthopsis murina*), also known as the common dunnart in Australia, is a dasyurid marsupial. It has an average body length of 7 to 12

centimeters (2.8–4.7 in) with a tail length of 5.5 to 13 centimetres (2.2–5.1 in)....

Slender-tailed squirrel
Slender-tailed squirrel
Protoxerus aubinnii

The slender-tailed squirrel (*Protoxerus aubinnii*) is a species of rodent in the family *Sciuridae* found in Ivory Coast, Ghana, Liberia, and Sierra Leone. Its natural habitat is subtropical or tropical moist lowland forests.

Slevin's mouse
Slevin's mouse
Peromyscus slevini

Slevin's mouse (*Peromyscus slevini*), also known as the Catalina deer mouse, is a species of rodent in the family *Cricetidae*. It is endemic to Isla Santa Catalina off the east coast of Baja California Sur, an island with an area of about 40 km²...

Sloggett's vlei rat
Sloggett's vlei rat
Myotomys sloggetti

Sloggett's vlei rat or ice rat (*Myotomys sloggetti*) is a species of rodent in the family *Muridae*. It is found in southern Lesotho and South Africa where its natural habitats are subtropical or tropical high-altitude grassland, swamps, and rocky...

Sloth Bear
Sloth Bear
Melursus ursinus

The Sloth bear (*Melursus ursinus*) is a myrmecophagous bear species native to the Indian subcontinent. It is listed as vulnerable on the IUCN Red List, mainly because of habitat loss and degradation. When their territories are encroached upon by...

Small Asian Mongoose
Small Asian Mongoose
Herpestes javanicus

The Javan mongoose (*Urva javanica*) is a mongoose species native to Southeast Asia.
Small Asian sheath-tailed bat
Small Asian sheath-tailed bat
Emballonura alecto

The small Asian sheath-tailed bat (*Emballonura alecto*) is a species of sac-winged bat in the family *Emballonuridae*. It is found in Borneo (Brunei, Indonesia, Malaysia), Sulawesi, and the Philippines.

Small bent-winged bat
Small bent-winged bat
Miniopterus pusillus

The small bent-winged bat (*Miniopterus pusillus*) is a species of vesper bat in the family *Miniopteridae*. A novel version of the Coronavirus has been identified in this species. It can be found in the following countries: Bangladesh, Bhutan,...

Small big-eared brown bat
Small big-eared brown bat
Histiotus montanus

The small big-eared brown bat (*Histiotus montanus*) is a species of vesper bat in the family *Vespertilionidae*. It can be found in the following countries: Argentina, Bolivia, Brazil, Chile, Colombia, Ecuador, Peru, Uruguay, and Venezuela.
Small dorcopsis

Small dorcopsis
Dorcopsulus vanheurni

The lesser forest wallaby or small dorcopsis (*Dorcopsulus vanheurni*) is a species of marsupial in the family Macropodidae. It is found in the mountainous interior of West Papua, Indonesia and Papua New Guinea. Its natural habitat is subtropical...

Small five-toed jerboa
Small five-toed jerboa
Allactaga elater

The small five-toed jerboa (*Allactaga elater*) is a rodent of the family Dipodidae and genus *Allactaga*, that has five digits.

Small flying fox
Small flying fox
Pteropus hypomelanus

The small flying fox, island flying fox or variable flying fox (*Pteropus hypomelanus*) is a species of flying fox in the family Pteropodidae. It is found in Australia, Cambodia, Indonesia, Malaysia, the Maldives, Myanmar, Papua New Guinea, the...

Small hocicudo
Small hocicudo
Oxymycterus hiska

The small hocicudo (*Oxymycterus hiska*) is a little known species of rodent in the family Cricetidae. It is found in Bolivia and Peru on the eastern slopes of the Andes.

Small Indian Civet
Small Indian Civet
Viverricula indica

The small Indian civet (*Viverricula indica*) is a civet native to South and Southeast Asia. It is listed as Least Concern on the IUCN Red List because of its widespread distribution, widespread habitat use and healthy populations living in...

Small Japanese field mouse
Small Japanese field mouse
Apodemus argenteus

The small Japanese field mouse (*Apodemus argenteus*) is a species of rodent in the family Muridae. It is endemic to Japan.

Small Japanese mole
Small Japanese mole
Mogera imaizumii

The small Japanese mole (*Mogera imaizumii*) is a species of mammal in the family Talpidae. It is endemic to Japan. Even though they are extinct in central Tokyo, they are found in the grounds of the Imperial Palace.

Small Luzon forest mouse
Small Luzon forest mouse
Apomys microdon

The small Luzon forest mouse (*Apomys microdon*) is a species of rodent in the family Muridae. It is found only in the Philippines.

Small mouse-tailed bat
Small mouse-tailed bat
Rhinopoma muscatellum

The small mouse-tailed bat (*Rhinopoma muscatellum*) is a species of bat in the

Rhinopomatidae family. It is found in Afghanistan, Iran, Oman, and possibly Ethiopia. The small mouse-tailed bat ranges from the Sistan Basin in Iran well into the...

Small pencil-tailed tree mouse

Small pencil-tailed tree mouse

Chiropodomys pusillus

The small pencil-tailed tree mouse or lesser pencil-tailed tree mouse (*Chiropodomys pusillus*) is a species of arboreal rodent in the family Muridae. It is endemic to Borneo where it is only known from Sabah and Sarawak (Malaysia) and from...

Small red brocket

Small red brocket

Mazama bororo

The small red brocket (*Mazama bororo*) is a small species of deer in the family Cervidae. It is endemic to Atlantic Forest in Paraná, Santa Catarina and São Paulo in southeastern Brazil. This species, which only was scientifically described in...

Small rufous horseshoe bat

Small rufous horseshoe bat

Rhinolophus subrufus

The small rufous horseshoe bat (*Rhinolophus subrufus*) is a species of bat in the family Rhinolophidae. It is endemic to the Philippines on the island of Camiguin, Catanduanes, Luzon, Mindanao and Mindoro at elevations from sea level to 1,000m.

Small Samoan flying fox

Small Samoan flying fox

Pteropus allenorum

The small Samoan flying fox (*Pteropus allenorum*) is a species of fruit-eating megabat whose type specimen was originally collected in Samoa in 1856, but was not identified as a new species until 2009. Its wingspan was at least two feet, and it...

Small sun squirrel

Small sun squirrel

Heliosciurus punctatus

The small sun squirrel (*Heliosciurus punctatus*) is a species of rodent in the family Sciuridae. It is found in Ivory Coast, Ghana, Liberia, Sierra Leone, and possibly Guinea. Its natural habitat is subtropical or tropical moist lowland forests.

Small vesper mouse

Small vesper mouse

Calomys laucha

The small vesper mouse (*Calomys laucha*) is a rodent species from South America. It is found in Argentina, Bolivia, Brazil, Paraguay and Uruguay. It is one of the hosts of hantavirus, causing hantavirus pulmonary syndrome

Small white-toothed rat

Small white-toothed rat

Berylmys berdmorei

The small white-toothed rat (*Berylmys berdmorei*) is a species of rodent in the family Muridae. It is found in Cambodia, Laos, Myanmar, Thailand, and Vietnam.

Small white-winged flying fox

Small white-winged flying fox

Desmalopex microleucopterus

The small white-winged flying fox (*Desmalopex microleucopterus*) is a species of megabat in the family Pteropodidae. It is known from Mindoro Island, in the

Philippines. Only 13 specimens have ever been found; 1 in 1998 and 12 found in 2006.

Small woolly bat

Small woolly bat

Kerivoula intermedia

The small woolly bat (*Kerivoula intermedia*) is a species of vesper bat in the family Vespertilionidae. It is found only in Malaysia and is little known. It is only slightly larger than the Kitti's hog-nosed bat, one of the smallest mammals in the...

Small-eared dormouse

Small-eared dormouse

Graphiurus microtis

The small-eared dormouse (*Graphiurus microtis*) is a species of rodent in the family Gliridae. It is found in Angola, Botswana, Eritrea, Ethiopia, Kenya, Lesotho, Malawi, Mozambique, Namibia, South Africa, Sudan, Eswatini, Tanzania, Zambia, and...

Small-eared rat

Small-eared rat

Taeromys microbullatus

The small-eared rat (*Taeromys microbullatus*), is an extant species of Old World rodent that inhabits the southeast of Sulawesi, an Indonesian island. The rat is found in the Mekongga Mountains of southeastern Sulawesi. Following the...

Small-footed shrew

Small-footed shrew

Crocidura parvipes

The small-footed shrew (*Crocidura parvipes*) is a species of mammal in the family Soricidae. It is found in Angola, Burundi, Cameroon, Central African Republic, Chad, Democratic Republic of the Congo, Ethiopia, Kenya, Nigeria, Rwanda, South...

Small-toothed fruit bat

Small-toothed fruit bat

Neopteryx frosti

The small-toothed fruit bat or small-toothed Flying fox (*Neopteryx frosti*) is a species of megabat in the family Pteropodidae. It is monotypic within the genus *Neopteryx*.

Small-toothed harvest mouse

Small-toothed harvest mouse

Reithrodontomys microdon

The small-toothed harvest mouse (*Reithrodontomys microdon*) is a species of rodent in the family Cricetidae. It is found in Guatemala and Mexico.

Small-toothed long-eared bat

Small-toothed long-eared bat

Nyctophilus microdon

The small-toothed long-eared bat, species *Nyctophilus microdon*, is a vespertilionid bat. This flying mammal is found only in Papua New Guinea.

Small-toothed mole

Small-toothed mole

Euroscaptor parvidens

The small-toothed mole (*Euroscaptor parvidens*) is a species of mammal in the family Talpidae. It is found in Di Linh, in southern Vietnam, and Rakho near the border of Vietnam and China, and in southern Yunnan, China, near the Vietnamese border....

Small-Toothed Palm Civet
Small-Toothed Palm Civet
Arctogalidia trivirgata

The small-toothed palm civet (*Arctogalidia trivirgata*) is a small feliform mammal native to dense forests of Southeast Asia. The scientific name of this species can be translated as the 'three-striped bear-weasel'.

Small-toothed sportive lemur
Small-toothed sportive lemur
Lepilemur microdon

The small-toothed sportive lemur (*Lepilemur microdon*), or small-toothed weasel lemur, is a primate species in the family Lepilemuridae that—like all extant lemurs—is endemic to Madagascar. The species lives in dense rainforest in southeastern...

Smaller horseshoe bat
Smaller horseshoe bat
Rhinolophus megaphyllus

The smaller horseshoe bat (*Rhinolophus megaphyllus*) is a species of bat in the family Rhinolophidae. It is found in Australia and Papua New Guinea.

Smith Island cottontail
Smith Island cottontail
Sylvilagus floridanus hitchensi

Smith Island cottontail, Smith's Island cottontail or Hitchen's cottontail (*Sylvilagus floridanus hitchensi*), was a subspecies of the Eastern cottontail rabbit that lived mainly on two islands on the tip of the Delmarva Peninsula in Virginia. It...

Smith's bush squirrel
Smith's bush squirrel
Paraxerus cepapi

Smith's bush squirrel (*Paraxerus cepapi*), also known as the yellow-footed squirrel or tree squirrel, is an African bush squirrel which is native to woodlands of the southern Afrotropics.

Smith's red rock hare
Smith's red rock hare
Pronolagus rupestris

Smith's red rockhare, Smith's red rock hare or Smith's red rock rabbit (*Pronolagus rupestris*) is a species of mammal in the family Leporidae (rabbits and hares), and is the smallest member of the genus *Pronolagus*. The upperparts and gular collar...

Smith's shrew
Smith's shrew
Chodsigoa smithii

The Smith's shrew (*Chodsigoa smithii*) is a species of mammal in the family Soricidae. It is endemic to Southwest China, living primarily in mountainous broad-leaved forests at elevations of 900–3000 meters.

Smith's vole
Smith's vole
Myodes smithii

Smith's vole (*Myodes smithii*) is a species of rodent in the family Cricetidae. It is also known as Smith's red-backed vole and is found only in Japan. This vole is named after Richard Gordon Smith, (1858–1918) who, after falling out with his...

Smith's woolly bat
Smith's woolly bat
Kerivoula smithii

Smith's woolly bat (*Kerivoula smithii*) is a species of vesper bat in the family Vespertilionidae. It is found in Cameroon, Democratic Republic of the Congo, Kenya, Nigeria, and Uganda. Its natural habitats are subtropical or tropical dry forests,...

Smith's zokor

Smith's zokor

Eospalax smithii

Smith's zokor (*Eospalax smithii*) is a species of rodent in the family Spalacidae, endemic to China.

Smithers's horseshoe bat

Smithers's horseshoe bat

Rhinolophus smithersi

Smithers's horseshoe bat (*Rhinolophus smithersi*) is a species of horseshoe bat found in South Africa and Zimbabwe. It was described as a new species in 2012.

Smoke-bellied rat

Smoke-bellied rat

Niviventer eha

The smoke-bellied rat (*Niviventer eha*) is a species of rodent in the family Muridae. It is found in China, India, Myanmar, and Nepal.

Smoky bat

Smoky bat

Amorphochilus schnablii

The smoky bat (*Amorphochilus schnablii*) is a species of bat in the family Furipteridae. It is monotypic within the genus *Amorphochilus*. Its natural habitat is rocky shores. It is threatened by habitat loss. This bat is also called the thumbless...

Smoky flying squirrel

Smoky flying squirrel

Pteromyscus pulverulentus

The smoky flying squirrel (*Pteromyscus pulverulentus*) is a species of rodent in the family Sciuridae. It is monotypic within the genus *Pteromyscus*. It is found in parts of Brunei, Indonesia, Peninsular Malaysia, and southern Thailand, but its...

Smoky grass mouse

Smoky grass mouse

Akodon fumeus

The smoky grass mouse (*Akodon fumeus*) is a species of rodent in the family Cricetidae. It is found in Argentina, Bolivia and Peru.

Smoky mouse

Smoky mouse

Pseudomys fumeus

The smoky mouse (*Pseudomys fumeus*) is a species of rodent in the family Muridae native to southeastern Australia. It was first described in 1934 and its species name is Latin for "smoky". As its name suggests, it is a grey-furred mouse, darker...

Smoky pocket gopher

Smoky pocket gopher

Cratogeomys fumosus

The smoky pocket gopher (*Cratogeomys fumosus*) is a species of rodent in the family Geomyidae. It is endemic to Mexico (Querétaro). Its natural habitat is subtropical or tropical dry lowland grassland. Molecular phylogenetics has revealed that...

Smoky shrew
Smoky shrew
Sorex fumeus

The smoky shrew (*Sorex fumeus*) is a medium-sized North American shrew found in eastern Canada and the northeastern United States and extends further south along the Appalachian Mountains.

Smoky white-toothed shrew
Smoky white-toothed shrew
Crocidura fumosa

The smoky white-toothed shrew (*Crocidura fumosa*) is a species of mammal in the family Soricidae. It is endemic to Kenya. Its natural habitat is subtropical or tropical moist montane forests. It is threatened by habitat loss.

Smooth-Coated Otter
Smooth-Coated Otter
Lutrogale perspicillata

The Smooth-coated otter (*Lutrogale perspicillata*) is an otter species found in most of the Indian subcontinent and Southeast Asia. As its name indicates, its fur is smooth and shorter than that of other otter species.

Snow Leopard
Snow Leopard
Uncia uncia

The Snow leopard (*Panthera uncia*) is a felid native to the mountain ranges of Central and South Asia. The Snow leopard is adapted to living in a cold, mountainous environment and is capable of killing most animals in its range.

Snow Mountains grassland mosaic-tailed rat
Snow Mountains grassland mosaic-tailed rat
Melomys frigidicola

The Snow Mountains grassland mosaic-tailed rat (*Melomys frigidicola*), also known as the Snow Mountains grassland melomys, is a species of rodent in the family Muridae. It is endemic to the mountainous west part of the island of New Guinea, where...

Snow sheep
Snow sheep
Ovis nivicola

The snow sheep (*Ovis nivicola*), or Siberian bighorn sheep, is a species of sheep from the mountainous areas in the northeast of Siberia. One subspecies, the Putorana snow sheep (*Ovis nivicola borealis*), lives isolated from the other forms in the...

Snow-footed Oldfield mouse
Snow-footed Oldfield mouse
Thomasomys niveipes

The snow-footed Oldfield mouse (*Thomasomys niveipes*) is a species of rodent in the family Cricetidae. It is found only in Colombia.

Snowshoe Hare
Snowshoe Hare
Lepus americanus

The Snowshoe hare (*Lepus americanus*) is a secretive forest-dwelling mammal found only in North America. It has the name "snowshoe" because of the large size of its hind feet. The animal's feet prevent it from sinking into the snow when it hops...

Social pipistrelle
Social pipistrelle

Arielulus societatis

The social sprite (*Arielulus societatis*), also known as the social pipistrelle, is a species of vesper bat found only in Malaysia on the Malay Peninsula.

Social tuco-tuco

Social tuco-tuco

Ctenomys sociabilis

The social tuco-tuco (*Ctenomys sociabilis*) is a species of rodent in the family Ctenomyidae. It is endemic to Argentina. Its natural habitat is subtropical or tropical dry lowland grassland.

Social vole

Social vole

Microtus socialis

The social vole (*Microtus socialis*) is a species of rodent in the family Cricetidae. It is found in China, Iran, Kazakhstan, Syria, Turkey, and Ukraine.

Sody's tree rat

Sody's tree rat

Kadarsanomys sodyi

Sody's tree rat (*Kadarsanomys sodyi*) is a rodent species in the family Muridae that has been recorded only in bamboo forest in Gunung Gede Pangrango National Park, West Java, Indonesia. It was first recorded during surveys carried out between...

Sody's yellow house bat

Sody's yellow house bat

Scotophilus collinus

Sody's yellow bat or Sody's yellow house bat (*Scotophilus collinus*) is a species of vesper bat. It is native to Island Southeast Asia, where it is found in Indonesia, Malaysia, and Timor-Leste. This species was described in 1936.

Soemmerring's Gazelle

Soemmerring's Gazelle

Nanger soemmerringii

Soemmerring's gazelle (*Nanger soemmerringii*) is a gazelle species native to the Horn of Africa. It was described and given its binomen by German physician Philipp Jakob Cretzschmar in 1828. The species is currently very understudied due to its...

Soft grass mouse

Soft grass mouse

Akodon mollis

The soft grass mouse is a species of rodent in the family Cricetidae. It is found in Ecuador and Peru.

Soft-furred Oldfield mouse

Soft-furred Oldfield mouse

Thomasomys laniger

The soft-furred Oldfield mouse (*Thomasomys laniger*) is a species of rodent in the family Cricetidae. It is found in Colombia and Venezuela.

Soft-furred rat

Soft-furred rat

Millardia meltada

The soft-furred rat (*Millardia meltada*), or soft-furred metad, is a species of rodent in the family Muridae native to South Asia.

Soft-spined Atlantic spiny rat

Soft-spined Atlantic spiny rat

Trinomys dimidiatus

The soft-spined Atlantic spiny-rat, *Trinomys dimidiatus*, is a spiny rat species from South America. It is endemic to Brazil.

Sokolov's dwarf hamster

Sokolov's dwarf hamster

Cricetulus sokolovi

Sokolov's dwarf hamster (*Cricetulus sokolovi*) is a species of rodent in the hamster and vole family Cricetidae. Previously listed as conspecific with Chinese striped hamster, it has been listed as a separate species since 1988. It has a...

Solomon's naked-backed fruit bat

Solomon's naked-backed fruit bat

Dobsonia inermis

The Solomon's naked-backed fruit bat (*Dobsonia inermis*) is a species of megabat in the family Pteropodidae. It is endemic to the Solomon Islands.

Solomons flying fox

Solomons flying fox

Pteropus rayneri

The Solomons flying fox (*Pteropus rayneri*) is a species of flying fox in the family Pteropodidae. It is endemic to the Solomon Islands archipelago. The population abundance has been marked by periodic die-offs of generally unknown cause,...

Solomons mastiff bat

Solomons mastiff bat

Chaerephon solomonis

The Solomons mastiff bat (*Chaerephon solomonis*) is a species of bat in the family Molossidae. It is endemic to the Solomon Islands.

Somali bushbaby

Somali bushbaby

Galago gallarum

The Somali bushbaby (*Galago gallarum*), or the Somali lesser galago, as it is also known, is a species of nocturnal, arboreal primate in the family Galagidae. It is found in Ethiopia, Kenya, and Somalia. It is threatened by habitat loss.

Somali dwarf shrew

Somali dwarf shrew

Crocidura nana

The Somali dwarf shrew (*Crocidura nana*) is a species of mammal in the family Soricidae. It is found in Ethiopia and Somalia. Its natural habitat is subtropical or tropical dry lowland grassland. Type Locality: Somalia, Dollo

Somali elephant shrew

Somali elephant shrew

Galegeeska revoilii

The Somali elephant shrew or Somali sengi (*Galegeeska revoilii*) is a species of elephant shrew in the family Macroscelididae.

Somali golden mole

Somali golden mole

Calcochloris tytonis

The Somali golden mole (*Calcochloris tytonis*) is a golden mole endemic to Somalia. In 1964, Dr. Alberto Simonetta of the University of Florence discovered the mole's jaw and ear bone fragments in a barn owl pellet in Giohar, Somalia. The Somali...

Somali hedgehog

Somali hedgehog
Atelerix sclateri

The Somali hedgehog (*Atelerix sclateri*) is a species of mammal in the family Erinaceidae. It is endemic to Somalia and Somaliland. The Somali hedgehog is nocturnal.

Somali pygmy gerbil
Somali pygmy gerbil
Microdillus peeli

The Somali pygmy gerbil (*Microdillus peeli*) is a species of rodent in the family Muridae. It is the only species in the genus *Microdillus*. It is found only in Somalia. Its natural habitat is subtropical or tropical dry shrubland.

Somali serotine
Somali serotine
Neoromicia somalica

The Somali serotine (*Neoromicia somalica*) is a species of vesper bat. It is found in Benin, Botswana, Burkina Faso, Cameroon, Central African Republic, Chad, Republic of the Congo, Democratic Republic of the Congo, Ivory Coast, Eritrea, Ethiopia,...

Somali shrew
Somali shrew
Crocidura somalica

The Somali shrew (*Crocidura somalica*) is a species of mammal in the family Soricidae. It is found in Ethiopia, Mali, Somalia, and Sudan. Its natural habitat is dry savanna.

Somali wild ass
Somali wild ass
Equus africanus somaliensis

The Somali wild ass (*Equus africanus somaliensis*) is a subspecies of the African wild ass. It is found in Somalia, the Southern Red Sea region of Eritrea, and the Afar Region of Ethiopia. The legs of the Somali wild ass are striped, resembling...

Somali Wild Dog
Somali Wild Dog
Lycaon pictus somalicus

The Somali wild dog (*Lycaon pictus somalicus*) is a subspecies of the African wild dog. It is smaller than the East African wild dog subspecies, has shorter and coarser fur, and has a weaker dentition. Its colour closely approaches that of the...

Somalia gerbil
Somalia gerbil
Gerbillus dunni

Dunn's gerbil (*Gerbillus dunni*) is a rodent species distributed mainly in Eritrea, Ethiopia, Somaliland and Djibouti. It is an animal of dry grassland and rocky plains. It is also sometimes called the "Somalia gerbil", not to be confused with...

Somalian gerbil
Somalian gerbil
Dipodillus somalicus

The Somalian gerbil (*Dipodillus somalicus*) is found only in Somalia.

Somalian slender mongoose
Somalian slender mongoose
Galerella ochracea

The Somali slender mongoose (*Herpestes ochraceus*) is a small mammal found in Somalia and adjacent regions. It is a small to medium-sized carnivore, averaging about 0.6 kg (1.3 lb) in weight. It was first described by John Edward Gray in 1848,...

Somali trident bat
Somali trident bat
Asellia italosomalia

The Somali trident bat (*Asellia italosomalia*) is a species of bat found in the Horn of Africa.

Sombre bat
Sombre bat
Eptesicus tatei

The sombre bat (*Eptesicus tatei*) is a species of vesper bat. It is found only in India. Its natural habitat is subtropical or tropical moist montane forests.

Sommer's Sulawesi rat
Sommer's Sulawesi rat
Sommeromys macrorhinos

Sommer's Sulawesi rat (*Sommeromys macrorhinos*) is a species of rodent in the family Muridae from Sulawesi. It is the only species in the monotypic genus *Sommeromys* and was described by Musser & Durden in 2002.

Sonnerat's shrew
Sonnerat's shrew
Diplomesodon sonnerati

Sonnerat's shrew (*Diplomesodon sonnerati* or *Crociodura sonnerati*) is a species of shrew that was first described by Pierre Sonnerat from Pondicherry somewhere in 1813.

Sonoma chipmunk
Sonoma chipmunk
Neotamias sonomae

The Sonoma chipmunk (*Neotamias sonomae*) is a small ground-dwelling rodent in the squirrel family Sciuridae. It is endemic to northwestern California in the United States. Members of *Neotamias* are characterized by having 2 premolars, transitional...

Sonoran harvest mouse
Sonoran harvest mouse
Reithrodontomys burti

The Sonoran harvest mouse (*Reithrodontomys burti*) is a species of rodent in the family Cricetidae. It is found only in Mexico.

Sonoran pronghorn
Sonoran pronghorn
Antilocapra americana sonoriensis

The Sonoran pronghorn (*Antilocapra americana sonoriensis*) is an endangered subspecies of pronghorn. Numerous and ongoing threats exist to the survival of the Sonoran pronghorn at present.

Sooretamys
Sooretamys
Sooretamys angouya

Sooretamys angouya, also known as the rat-headed rice rat, and Paraguayan rice rat, is a rodent species from South America. It is found in northeastern Argentina, southern Brazil and Paraguay in forested areas within the Atlantic Forest and...

Sooty dunnart
Sooty dunnart
Sminthopsis fuliginosus

The sooty dunnart (*Sminthopsis fuliginosus*) is a species of dunnart found in Western Australia. It is one of the least-known of the dunnarts, with the IUCN classifying it as data deficient. It was formerly believed to be a subspecies of the...

Sooty Mangabey
Sooty Mangabey
Cercocebus atys

The Sooty mangabey (*Cercocebus atys*) is an Old World monkey found in forests of Africa. These rare and endangered monkeys live in groups and communicate with each other using a wide repertoire of vocalizations, including distinct alarm calls for...

Sooty mustached bat
Sooty mustached bat
Pteronotus quadridens

The sooty mustached bat (*Pteronotus quadridens*) is a species of bat in the family Mormoopidae. It is found in Cuba, the Dominican Republic, Haiti, Jamaica, and Puerto Rico.

Sooty roundleaf bat
Sooty roundleaf bat
Hipposideros fuliginosus

The sooty roundleaf bat (*Hipposideros fuliginosus*) is a species of bat in the family Hipposideridae. It is found in Cameroon, Democratic Republic of the Congo, Ivory Coast, Gabon, Ghana, Guinea, Liberia, Nigeria, Sierra Leone, and Uganda. Its...

Soprano Pipistrelle
Soprano Pipistrelle
Pipistrellus pygmaeus

The Soprano pipistrelle (*Pipistrellus pygmaeus*) is a small species of bat that lives in Europe. It is also known as the 55 kHz pipistrelle and the Brown pipistrelle.

Sorensen's leaf-nosed bat
Sorensen's leaf-nosed bat
Hipposideros sorenseni

Sorensen's leaf-nosed bat (*Hipposideros sorenseni*) is a species of bat in the family Hipposideridae that is endemic to Indonesia.

Sorex ornatus sinuosus
Sorex ornatus sinuosus
Sorex ornatus sinuosus

Sorex ornatus sinuosus, the Suisun shrew or Suisun ornate shrew, is a subspecies of the ornate shrew that occurs in the tidal marshes of the northern shores of San Pablo and Suisun bays (northern arms of the San Francisco Bay, as far east as...

Soriano's yellow-shouldered bat
Soriano's yellow-shouldered bat
Sturnira soriano

Soriano's yellow-shouldered bat (*Sturnira soriano*), is an extant species of leaf-nosed bat indigenous to Bolivia and Venezuela, although its precise distribution is uncertain. Following the postulation of the species in 2005, *S. soriano*...
Soricine brucie

Soricine brucie
Bucepattersonius soricinus

The soricine brucie (*Bucepattersonius soricinus*) is a rodent species from South America. It is found in Brazil.

Soricomys kalinga
Soricomys kalinga
Soricomys kalinga

The Kalinga shrew mouse (*Soricomys kalinga*) is a rodent of the genus *Soricomys* found in the northern province of Kalinga, island of Luzon, in the northern Philippines.

South African Giraffe
South African Giraffe
Giraffa camelopardalis giraffa

The South African giraffe (*Giraffa camelopardalis giraffa*) is a subspecies of giraffe. South African giraffes are uncommon in captivity. As of 2010, there are around 45 South African giraffes breeding in zoos. Approximately 12,000 privately owned...

South African pouched mouse
South African pouched mouse
Saccostomus campestris

The South African pouched mouse or southern African pouched mouse (*Saccostomus campestris*) is a species of rodent in the family Nesomyidae, which is viewed as actually representing a complex of at least three undescribed species. It is found in...

South African Springhare
South African Springhare
Pedetes capensis

The South African springhare (*Pedetes capensis*) (Afrikaans: springhaas) is a medium-sized terrestrial and burrowing rodent. Despite the name, it is not a hare. It is one of two extant species in the genus *Pedetes*, and is native to southern...

South American Coati
South American Coati
Nasua nasua

The South American coati (*Nasua nasua*) is a member of the raccoon family (Procyonidae), found in South America. Coatis are also known as coatimundis. The name "coatimundi" comes from the Tupian languages of Brazil, where it means "lone coati"....

South American Cougar
South American Cougar
Puma concolor concolor

The South American cougar (*Puma concolor concolor*) is a cougar subspecies. The cougar holds historical cultural significance amongst many South American indigenous people. People in the Andes regard it as being either a snatcher of souls or as a...

South American Fur Seal
South American Fur Seal
Arctocephalus australis

The South American fur seal (*Arctocephalus australis*) breeds on the coasts of Peru, Chile, the Malvinas Islands, Argentina, Uruguay and Brazil. The total population is around 250,000. However, population counts are sparse and outdated. Although...

South American Gray Fox
South American Gray Fox
Lycalopex griseus

The South American gray fox (*Lycalopex griseus*), also known as the Patagonian fox, the chilla or the gray zorro, is a species of *Lycalopex*, the "false" foxes. It is endemic to the southern part of South America.

South American hoary bat
South American hoary bat
Lasiurus villosissimus

The South American hoary bat (*Aeorestes villosissimus*) is a species of vesper bat found in South America.

South American Sea Lion
South American Sea Lion
Otaria flavescens

The South American sea lion (*Otaria flavescens*, formerly *Otaria byronia*), also called the southern sea lion and the Patagonian sea lion, is a sea lion found on the western and southeastern coasts of South America. It is the only member of the...

South China field mouse
South China field mouse
Apodemus draco

The South China field mouse (*Apodemus draco*) is a species of rodent in the family Muridae. It is found in China, India, and Myanmar.

South China sika deer
South China sika deer
Cervus nippon kopschi

The South China sika deer (*Cervus nippon kopschi*) is one of many subspecies of sika deer. Standing 85 cm or 33 in tall at the shoulders, it is a small subspecies that is only a little larger than its Japanese counterpart. The back is brown with...

South-Central Black Rhinoceros
South-Central Black Rhinoceros
Diceros bicornis minor

The South-central black rhinoceros (*Diceros bicornis minor*) is a subspecies of the Black rhinoceros. Although it is the most numerous of the Black rhino subspecies, it is nevertheless designated as critically endangered on the IUCN's red list.

South-Western Black Rhinoceros
South-Western Black Rhinoceros
Diceros bicornis occidentalis

The South-western black rhinoceros (*Diceros bicornis occidentalis*) is a subspecies of the Black rhinoceros. This subspecies is often mistaken for either the extinct Southern black rhinoceros (*D. b. bicornis*) or the southeastern subspecies (*D. b....*

Southeast African Cheetah
Southeast African Cheetah
Acinonyx jubatus jubatus

The Southeast African cheetah (*Acinonyx jubatus jubatus*) is the nominate cheetah subspecies. It was first described by German naturalist Johann Christian Daniel von Schreber in his book *Die Säugethiere in Abbildungen nach der Natur mit...*

Southeast Asian long-fingered bat
Southeast Asian long-fingered bat
Miniopterus fuscus

The Southeast Asian long-fingered bat (*Miniopterus fuscus*) is a species of vesper bat in the family Vespertilionidae. It is endemic to Japan. It has been assessed as endangered by the IUCN.

Southeast Asian shrew
Southeast Asian shrew
Crocidura fuliginosa

The Southeast Asian shrew (*Crocidura fuliginosa*) is a species of mammal in the family Soricidae. It is found in Cambodia, India, China, Laos, Malaysia, Myanmar, Thailand, and Vietnam.

Southeastern four-eyed opossum
Southeastern four-eyed opossum
Philander frenatus

The southeastern four-eyed opossum (*Philander frenatus*) is an opossum species native to South America. It is found in Atlantic Forest ecoregions, in Brazil, Paraguay and Argentina.

Southeastern myotis
Southeastern myotis
Myotis austroriparius

The southeastern myotis (*Myotis austroriparius*) is a small bat found throughout the Gulf Coastal Plain and the Lower Mississippi Alluvial Plain of the southeastern United States.

Southeastern pocket gopher
Southeastern pocket gopher
Geomys pinetis

The southeastern pocket gopher (*Geomys pinetis*) is a species of pocket gopher that is native to the southeastern United States. It occurs in Alabama, Georgia, and Florida, where it is the only pocket gopher.

Southeastern shrew
Southeastern shrew
Sorex longirostris

The southeastern shrew (*Sorex longirostris*) is a species of mammal in the family Soricidae. It is found in the southeastern United States.

Southeastern Spanish ibex
Southeastern Spanish ibex
Capra pyrenaica hispanica

The Southeastern Spanish ibex or Beceite ibex (*Capra pyrenaica hispanica*) is a goat that is endemic to Spain. The Southeastern Spanish ibex inhabits the Sierra Nevada, Sierra de las Nieves Natural Park, Sierra de Cazorla, Sierra de Grazalema,...

Southeastern xanthurus rat
Southeastern xanthurus rat
Rattus salocco

The southeastern xanthurus rat (*Rattus salocco*), is a rodent of the genus *Rattus*.

Southern African Hedgehog
Southern African Hedgehog
Atelerix frontalis

The Southern African hedgehog (*Atelerix frontalis*) is a species of mammal in the family Erinaceidae. It is found in Angola, Botswana, Lesotho, Namibia, South Africa, Tanzania and Zimbabwe.

Southern African spiny mouse

Southern African spiny mouse
Acomys spinosissimus

The southern African spiny mouse (*Acomys spinosissimus*) is a species of rodent in the family Muridae. It is found in Botswana, Democratic Republic of the Congo, Malawi, Mozambique, South Africa, Tanzania, Zambia, and Zimbabwe. Its natural...

Southern African vlei rat
Southern African vlei rat
Otomys irroratus

The Southern African vlei rat (*Otomys irroratus*) is a species of rodent in the vlei rat genus, *Otomys*, of the family Muridae in the order Rodentia. This is the type species of the genus. It is native to the grasslands and swamps of southern...

Southern African wildcat
Southern African wildcat
Felis lybica cafra

The Southern African wildcat (*Felis lybica cafra*) is an African wildcat subspecies. In 2007, it was tentatively recognized as a distinct subspecies on the basis of genetic analysis. It is also known in English as the 'bush cat'.

Southern Amazon red squirrel
Southern Amazon red squirrel
Sciurus spadiceus

The southern Amazon red squirrel (*Sciurus spadiceus*), is a squirrel species from South America where it inhabits forests in much of north-western South America east of the Andes. Three subspecies are currently recognised. It is a dark red...

Southern bent-wing bat
Southern bent-wing bat
Miniopterus orianae bassanii

The southern bent-wing bat (*Miniopterus orianae bassanii*) is one of two subspecies of the Australasian bent-wing bat. Its population size has declined rapidly since the 1950s, and it is classified as critically endangered by the Australian...

Southern big-eared brown bat
Southern big-eared brown bat
Histiotus magellanicus

The southern big-eared brown bat (*Histiotus magellanicus*) is a species of bat from the family Vespertilionidae. Although current taxonomy treats the southern big-eared brown bat as a separate species, it is often treated as a subspecies of the...

Southern big-eared mouse
Southern big-eared mouse
Loxodontomys micropus

The southern big-eared mouse (*Loxodontomys micropus*), also known as the southern pericote, is a species of rodent in the family Cricetidae. It is found in Argentina and Chile, and is one of only two species in its genus. The type specimen was...

Southern birch mouse
Southern birch mouse
Sicista subtilis

The southern birch mouse (*Sicista subtilis*) is a species of birch mouse in the family Sminthidae. It is native to southern Russia, Kazakhstan, and potentially northern Mongolia and China.

Southern black rhinoceros
Southern black rhinoceros
Diceros bicornis bicornis

The southern black rhinoceros or Cape rhinoceros (*Diceros bicornis bicornis*) is an extinct subspecies of the black rhinoceros that was once abundant in South Africa from the Cape Province to Transvaal, southern Namibia, and possibly also Lesotho...
Southern Bog Lemming
Southern Bog Lemming
Synaptomys cooperi

Southern bog lemmings (*Synaptomys cooperi*) are small North American rodents. They are active year-round and cache grasses and sedge stems in underground chambers.
Southern Bottlenose Whale
Southern Bottlenose Whale
Hyperoodon planifrons

The southern bottlenose whale (*Hyperoodon planifrons*) is a species of whale, in the Ziphiid family, one of two members of the genus *Hyperoodon*. Seldom observed, the southern bottlenose whale is resident in Antarctic waters. The species was first...
Southern Brown Bandicoot
Southern Brown Bandicoot
Isoodon obesulus

The southern brown bandicoot (*Isoodon obesulus*) is a short-nosed bandicoot, a type of marsupial, found mostly in southern Australia. A similar species in South Western Australia, known as the quenda (from the Noongar word 'kwinda') was...
Southern brown howler
Southern brown howler
Alouatta guariba clamitans

The southern brown howler (*Alouatta guariba clamitans*) is a monkey subspecies of brown howler native to southeastern Brazil (Minas Gerais to Rio Grande do Sul) and far northeastern Argentina (Misiones). Gregorin, 2006, considered the southern...
Southern climbing mouse
Southern climbing mouse
Rhipidomys austrinus

The southern climbing mouse (*Rhipidomys austrinus*) is a species of rodent in the family Cricetidae. It is found in Argentina and Bolivia in forested valleys and on slopes on the eastern side of the Andes Mountains.
Southern common cuscus
Southern common cuscus
Phalanger mimicus

The southern common cuscus (*Phalanger mimicus*), also known as Australian cuscus, grey cuscus, grey phalanger, and to-ili, is an arboreal marsupial endemic to Australia (Cape York), southern New Guinea and possibly the Aru Islands. Until...
Southern cotton rat
Southern cotton rat
Sigmodon hirsutus

The southern cotton rat (*Sigmodon hirsutus*) is a rodent species in the family Cricetidae. It is found from southern Chiapas in Mexico through Central America, except for Belize, and as far east as northern Colombia and Venezuela. It lives in...
Southern dog-faced bat
Southern dog-faced bat
Cynomops planirostris

The southern dog-faced bat (*Cynomops planirostris*), is a Central and South

American bat species of the family Molossidae. It is found in northern Argentina, Bolivia, Brazil, Colombia, French Guiana, Guyana, Panama, Peru, Paraguay, Suriname and...

Southern fin whale

Southern fin whale

Balaenoptera physalus quoyi

The southern fin whale (*Balaenoptera physalus quoyi*) is a subspecies of fin whale that lives in the Southern Ocean. At least one other subspecies of fin whale, the northern fin whale (*B. p. physalus*), exists in the northern hemisphere.

Southern Flying Squirrel

Southern Flying Squirrel

Glaucomys volans

The southern flying squirrel or the assapan (*Glaucomys volans*) is one of three species of the genus *Glaucomys* and one of three flying squirrel species found in North America. It is found in deciduous and mixed woods in the eastern half of North...

Southern forest bat

Southern forest bat

Vespadelus regulus

The southern forest bat (*Vespadelus regulus*) is a vespertilionid bat found in Australia.

Southern giant pouched rat

Southern giant pouched rat

Cricetomys ansorgei

The southern giant pouched rat (*Cricetomys ansorgei*) is a species of rodent in the family Nesomyidae. It is distributed in the savannah of East and Southern Africa.

Southern giant slender-tailed cloud rat

Southern giant slender-tailed cloud rat

Phloeomys cumingi

The southern giant slender-tailed cloud rat or southern Luzon giant cloud rat (*Phloeomys cumingi*), also known as bugkon in Filipino, is a vulnerable species of rodent in the family Muridae, found only in southern Luzon in the Philippines. It is...

Southern Grasshopper Mouse

Southern Grasshopper Mouse

Onychomys torridus

The Southern grasshopper mouse (*Onychomys torridus*) is a species of predatory rodent native to North America. Notable for its resistance to venom, it routinely preys on the highly venomous Arizona bark scorpion.

Southern groove-toothed moss mouse

Southern groove-toothed moss mouse

Microhydromys argenteus

The southern groove toothed moss-mouse (*Microhydromys argenteus*) is a species of rodent in the family Muridae found in Southern Papua New Guinea. As opposed to *M. richardsoni*, *argenteus* sp can be differentiated most prominently by its...

Southern Hairy-Nosed Wombat

Southern Hairy-Nosed Wombat

Lasiorhinus latifrons

The southern hairy-nosed wombat (*Lasiorhinus latifrons*) is one of three extant species of wombats. It is found in scattered areas of semiarid scrub and mallee from the eastern Nullarbor Plain to the New South Wales border area. It is the...

Southern Idaho ground squirrel
Southern Idaho ground squirrel
Urocitellus endemicus

The southern Idaho ground squirrel (*Urocitellus endemicus*) is a species of the largest genus of ground squirrels. This species and the Northern Idaho ground squirrel were previously considered conspecific, together called the Idaho ground squirrel.

Southern lesser bamboo lemur
Southern lesser bamboo lemur
Haplemur meridionalis

The southern lesser bamboo lemur (*Haplemur meridionalis*), also known as the southern bamboo lemur, rusty-gray lesser bamboo lemur, and southern gentle lemur, is a species of bamboo lemur endemic to southern Madagascar.

Southern Lion
Southern Lion
Panthera leo melanochaita

The Southern lion is a lion subspecies from Southern and East Africa. In this region, lion populations were referred to by several regional names, including Katanga lion, Transvaal lion, Kalahari lion, Southeast African lion, Southwest African...

Southern little yellow-eared bat
Southern little yellow-eared bat
Vampyressa pusilla

The southern little yellow-eared bat (*Vampyressa pusilla*) is a frugivorous bat species from South America. It is found in Brazil, Argentina and Paraguay.

Southern long-nosed armadillo
Southern long-nosed armadillo
Dasypus hybridus

The southern long-nosed armadillo (*Dasypus hybridus*) is a species of armadillo native to South America.

Southern long-nosed bat
Southern long-nosed bat
Leptonycteris curasoae

The southern long-nosed bat (*Leptonycteris curasoae*) is a South American species of bat in the family Phyllostomidae.

Southern mole vole
Southern mole vole
Ellobius fuscocapillus

The southern mole vole (*Ellobius fuscocapillus*) is a species of rodent in the family Cricetidae. It is found in Afghanistan, Iran, Pakistan, and Turkmenistan.

Southern mountain cavy
Southern mountain cavy
Microcavia australis

The southern mountain cavy (*Microcavia australis*) is a species of South American rodent in the family Caviidae.

Southern multimammate mouse
Southern multimammate mouse
Mastomys coucha

The southern multimammate mouse or southern African mastomys (*Mastomys coucha*) is a species of rodent in the family Muridae which is endemic to southern Africa

(Angola, Botswana, Lesotho, Mozambique, Namibia, South Africa, and Zimbabwe). It is...

Southern Muriqui

Southern Muriqui

Brachyteles arachnoides

The southern muriqui (*Brachyteles arachnoides*) is a muriqui (woolly spider monkey) species endemic to Brazil.

Southern myotis

Southern myotis

Myotis aelleni

The southern myotis (*Myotis aelleni*) is a species of vesper bat. It is found only in Argentina. Based on qualitative and quantitative analyses of type specimens, Novaes, Wilson, Ruedi and Moratelli treat this taxon as a junior synonym of *Myotis*...

Southern naked-tailed armadillo

Southern naked-tailed armadillo

Cabassous unicinctus

The southern naked-tailed armadillo (*Cabassous unicinctus*) is a species of small armadillo from South America.

Southern needle-clawed bushbaby

Southern needle-clawed bushbaby

Euoticus elegantulus

The southern needle-clawed bushbaby (*Euoticus elegantulus*) is a species of strepsirrhine primate in the family Galagidae. Found in Cameroon, Central African Republic, Republic of the Congo, and possibly Democratic Republic of the Congo, its...

Southern ningau

Southern ningau

Ningau yvonneae

Southern ningau refers to *Ningau yvonneae*, a tiny marsupial carnivore related to the Dasyuridae family. Similar in appearance to *Ningau ridei*, found throughout central Australia, this species occurs in spinifex on semi-arid sandplains across...

Southern Palawan tree squirrel

Southern Palawan tree squirrel

Sundasciurus steerii

The Southern Palawan tree squirrel (*Sundasciurus steerii*) (local name "bising") is a species of rodent in the family Sciuridae. It is endemic to the Philippines. Its natural habitat is subtropical or tropical dry forests.

Southern Pig-Tailed Macaque

Southern Pig-Tailed Macaque

Macaca nemestrina

The southern pig-tailed macaque (*Macaca nemestrina*), also known as the Sundaland pig-tailed macaque and Sunda pig-tailed macaque, is a medium-sized macaque that lives in southern Thailand, Malaysia, and Indonesia. It is known locally as berok.

Southern pocket gopher

Southern pocket gopher

Thomomys umbrinus

The southern pocket gopher (*Thomomys umbrinus*) is a species of rodent in the family Geomyidae. It is found in Mexico and the United States, usually in high altitude grassland and shrubland. It feeds on plant material and has an extensive

burrow...
Southern Pudu
Southern Pudu
Pudu puda

Southern pygmy mouse
Southern pygmy mouse
Baiomys musculus

The southern pygmy mouse (*Baiomys musculus*) is a species of rodent in the family Cricetidae. It is found in El Salvador, Guatemala, Honduras, Mexico, and Nicaragua.
Southern Red Muntjac
Southern Red Muntjac
Muntiacus muntjak

The Southern red muntjac (*Muntiacus muntjak*) is a deer species native to Southeast Asia. It was formerly known as the Indian muntjac or the Common muntjac before the species was taxonomically revised to represent only populations of Sunda and...
Southern red-backed vole
Southern red-backed vole
Myodes gapperi

The southern red-backed vole or Gapper's red-backed vole (*Myodes gapperi*) is a small slender vole found in Canada and the northern United States. It is closely related to the western red-backed vole (*Myodes californius*), which lives to the...
Southern red-sided opossum
Southern red-sided opossum
Monodelphis sorex

The southern red-sided opossum (*Monodelphis sorex*) is an opossum species from South America. It is found in Argentina, Brazil and Paraguay. This opossum has small rounded ears, a long pointed nose and a tail which is half as long as its body.
The...
Southern Reedbuck
Southern Reedbuck
Redunca arundinum

The Southern reedbuck (*Redunca arundinum*) is an elegant antelope from Africa. It was first described by Pieter Boddaert, a Dutch physician and naturalist, in 1785. It gets its name from two Latin words: *reduncas* (meaning bent backward and curved,...
Southern Right Whale
Southern Right Whale
Eubalaena australis

The Southern right whale (*Eubalaena australis*) is a baleen whale, one of three species classified as right whales. These whales are known to be rather active on the water surface and curious towards human vessels. They like to interact with...
Southern Right Whale Dolphin
Southern Right Whale Dolphin
Lissodelphis peronii

The southern right whale dolphin (*Lissodelphis peronii*) is a small and slender species of cetacean, found in cool waters of the Southern Hemisphere. It is one of two species of right whale dolphin (genus *Lissodelphis*). This genus is...
Southern River Otter
Southern River Otter
Lontra provocax

The southern river otter (*Lontra provocax*) is a species of otter that lives in Chile and Argentina. Although called a "river otter", it inhabits both marine and freshwater environments. It sometimes is considered a subspecies of *Lontra*...

Southern Rocky Mountain wolf

Southern Rocky Mountain wolf

Canis lupus youngi

The southern Rocky Mountain wolf (*Canis lupus youngi*) is an extinct subspecies of gray wolf which was once distributed over southeastern Idaho, southwestern Wyoming, northeastern Nevada, Utah, western and central Colorado, northwestern Arizona...

Southern Short-Tailed Shrew

Southern Short-Tailed Shrew

Blarina carolinensis

The southern short-tailed shrew (*Blarina carolinensis*) is a gray, short-tailed shrew that inhabits the eastern United States.

Southern small-toothed moss mouse

Southern small-toothed moss mouse

Pseudohydromys pumehanae

The southern small-toothed moss mouse (*Pseudohydromys pumehanae*) is a species of rodent in the family Muridae found on the slopes of Mount Bosavi, Papua New Guinea.

Southern spiny pocket mouse

Southern spiny pocket mouse

Heteromys australis

The southern spiny pocket mouse (*Heteromys australis*) is a species of rodent in the family Heteromyidae. It is found in Colombia, Ecuador, and Panama.

Southern Spotted Skunk

Southern Spotted Skunk

Spilogale angustifrons

The Southern spotted skunk (*Spilogale angustifrons*) is a species of mammal found in Central America and in southern Mexico. At one time this skunk was considered to be a subspecies of the Eastern spotted skunk (*Spilogale putorius*).

Southern Tamandua

Southern Tamandua

Tamandua tetradactyla

The southern tamandua (*Tamandua tetradactyla*), also called the collared anteater or lesser anteater, is a species of anteater from South America and the island of Trinidad in the Caribbean. It is a solitary animal found in many habitats, from...

Southern Three-Banded Armadillo

Southern Three-Banded Armadillo

Tolypeutes matacus

The southern three-banded armadillo (*Tolypeutes matacus*), also known as La Plata three-banded armadillo or Azara's domed armadillo, is an armadillo species from South America. It is found in parts of southwestern Brazil, northern Argentina,...

Southern three-striped opossum

Southern three-striped opossum

Monodelphis theresa

The southern three-striped opossum (*Monodelphis theresa*) is an opossum species from South America. It is found in the Atlantic Forest region of Brazil.

Southern Tiger Cat

Southern Tiger Cat

Leopardus guttulus

The Southern tiger cat (*Leopardus guttulus*) is a small wild cat species native to South America. It was long considered to be a subspecies of the oncilla and was recognized as a distinct species in 2013. The Southern tiger cat is in fact closely...

Southern Tree Hyrax
Southern Tree Hyrax
Dendrohyrax arboreus

The Southern tree hyrax (*Dendrohyrax arboreus*) is a small mammal found in Africa. These animals are superficially similar to marmots, or over-large pikas, but are much more closely related to elephants and sirenians.

Southern tuco-tuco
Southern tuco-tuco
Ctenomys australis

The southern tuco-tuco (*Ctenomys australis*) is a species of rodent in the family Ctenomyidae. It is endemic to Argentina.

Southern Viscacha
Southern Viscacha
Lagidium viscacia

The southern viscacha (*Lagidium viscacia*) is a species of viscacha, a rodent in the family Chinchillidae found in Argentina, Bolivia, Chile, and Peru. It is a colonial animal living in small groups in rocky mountain areas. It has long ears and...

Southern vole
Southern vole
Microtus levis

The East European vole (*Microtus mystacinus*) is a species of vole (rodent) in the family Cricetidae.

Southern White Rhinoceros
Southern White Rhinoceros
Ceratotherium simum simum

The Southern white rhinoceros (*Ceratotherium simum simum*) is one of the two subspecies of the white rhinoceros. It is the most common and widespread subspecies of rhinoceros. Southern white rhinos are one of the largest and heaviest land animals...

Southern White-Breasted Hedgehog
Southern White-Breasted Hedgehog
Erinaceus concolor

The Southern white-breasted hedgehog (*Erinaceus concolor*) is a spiny mammal found in Europe and Asia. These hedgehogs are becoming very popular as pets in the United States and Iran.

Southern woolly lemur
Southern woolly lemur
Avahi meridionalis

The southern woolly lemur (*Avahi meridionalis*), or southern avahi, has been recently recognized as a separate species of woolly lemur in 2006 by Zaramody et al. It is a nocturnal and pair-living species. Groups can range from 2 (the parental...

Southern yellow bat
Southern yellow bat
Lasiurus ega

The southern yellow bat (*Dasypterus ega*) is a species of vesper bat that belongs

to suborder microchiroptera (microbat) in the family Vespertilionidae. It is native to South, North and Central America, from the Rio Grande Valley of Texas in the...
Southwest China vole
Southwest China vole
Eothenomys custos

The Southwest China vole (*Eothenomys custos*) is a species of rodent in the family Cricetidae. It is found only in Yunnan and Sichuan, China. It occurs in Cangshan Erhai Nature Reserve.
Southwestern myotis
Southwestern myotis
Myotis auriculus

The southwestern myotis (*Myotis auriculus*) is a species of vesper bat. It is found in Guatemala, Mexico, and in Arizona and New Mexico in the United States.
Southwestern water vole
Southwestern water vole
Arvicola sapidus

The southwestern water vole or southern water vole (*Arvicola sapidus*) is a large amphibious vole native to most of France and southwestwards through Spain and Portugal. It is listed on the IUCN Red List as vulnerable. Although historically...
Sowell's short-tailed bat
Sowell's short-tailed bat
Carollia sowelli

Sowell's short-tailed bat (*Carollia sowelli*) is a common bat species in the family Phyllostomidae. It is found from San Luis Potosi (Mexico) through Central America to west Panama. The species is named after American philanthropist James N. Sowell.
Sowerby's beaked whale
Sowerby's beaked whale
Mesoplodon bidens

Sowerby's beaked whale (*Mesoplodon bidens*), also known as the North Atlantic or North Sea beaked whale, is a species of toothed whale. It was the first mesoplodont whale to be described. James Sowerby, an English naturalist and artist, first...
Spade-Toothed Whale
Spade-Toothed Whale
Mesoplodon traversii

The Spade-toothed whale (*Mesoplodon traversii*) is the rarest species of beaked whale. As of a find in 2024, only six specimens have been confirmed, all in the South Pacific, five in New Zealand, and one in Chile. This cetacean is probably the...
Spanish Ibex
Spanish Ibex
Capra pyrenaica

The Iberian ibex (*Capra pyrenaica*), also known as the Spanish ibex, Spanish wild goat, or Iberian wild goat, is a species of ibex endemic to the Iberian Peninsula. Four subspecies have been described; two are now extinct. The Portuguese...
Spanish mole
Spanish mole
Talpa occidentalis

The Spanish mole (*Talpa occidentalis*) is a species of mammal in the family Talpidae. It is found in Portugal and Spain.
Spanish red deer
Spanish red deer

Cervus elaphus hispanicus

The Spanish red deer (*Cervus elaphus hispanicus*), is a subspecies of the red deer native to Spain. The Spanish red deer is a polygynous subspecies, which means the males have two or more mates; during mating season, males show a dark ventral...

Speckled dasyure

Speckled dasyure

Neophascogale lorentzi

The speckled dasyure (*Neophascogale lorentzii*), also known as the long-clawed marsupial mouse, is a member of the order Dasyuromorphia. It is an inhabitant of Papua, Indonesia and Papua New Guinea. It is the only member of the genus...

Speckled Ground Squirrel

Speckled Ground Squirrel

Spermophilus suslicus

The speckled ground squirrel or spotted souslik (*Spermophilus suslicus*) is a species of rodent in the family Sciuridae from Eastern Europe. *Spermophilus suslicus* consists of three subspecies: *S. s. boristhenicus*, *S. s. guttatus*, and *S. s.*

Speckled spiny tree-rat

Speckled spiny tree-rat

Pattonomys semivillosus

The speckled spiny tree-rat (*Pattonomys semivillosus*) is a species of rodent in the family Echimyidae. It is found in northern Colombia and Venezuela. It can be found in gallery forest and dry forest. Its diet includes fruit and seeds.

Spectacled Bear

Spectacled Bear

Tremarctos ornatus

The Spectacled bear (*Tremarctos ornatus*) is a species of bear native to the Andes Mountains in western South America. It is the only living species of bear native to South America, and the last remaining short-faced bear (subfamily Tremarctinae)....

Spectacled dormouse

Spectacled dormouse

Graphiurus ocularis

The spectacled dormouse or namtap (*Graphiurus ocularis*) is a species of rodent in the family Gliridae, and one of four dormouse species endemic to South Africa. It has a bushy, squirrel-like tail, a fairly short muzzle, small ears and soft fur....

Spectacled Flying Fox

Spectacled Flying Fox

Pteropus conspicillatus

The spectacled flying fox (*Pteropus conspicillatus*), also known as the spectacled fruit bat, is a megabat that lives in Australia's north-eastern regions of Queensland. It is also found in New Guinea and on the offshore islands including...

Spectacled hare-wallaby

Spectacled hare-wallaby

Lagorchestes conspicillatus

The spectacled hare-wallaby (*Lagorchestes conspicillatus*) is a species of macropod found in Australia and New Guinea. In Australia, a small sub-population is found on Barrow Island, while the mainland type is widespread, though in decline,...

Spectacled Porpoise

Spectacled Porpoise

Phocoena dioptrica

The spectacled porpoise (*Phocoena dioptrica*) is a small to midsize porpoise indigenous to the Southern Ocean. It is one of the most poorly studied cetaceans, partly due to its remote range in the southern ocean. What little is known about this...

Spectacled slender opossum
Spectacled slender opossum
Marmosops ocellatus

The Spectacled slender opossum (*Marmosops ocellatus*), also known as little spotted slender opossum is a species in the marsupial genus *Marmosops* from South America.

Spectral Bat
Spectral Bat
Vampyrum spectrum

The spectral bat (*Vampyrum spectrum*), also called the great false vampire bat or Linnaeus's false vampire bat, is a large, carnivorous leaf-nosed bat found in Mexico, Central America, and South America. It is the only member of the genus *Vampyrum*...

Spectral Tarsier
Spectral Tarsier
Tarsius spectrum

The spectral tarsier (*Tarsius spectrum*, also called *Tarsius tarsier*) is a species of tarsier found on the island of Selayar in Indonesia. It is apparently less specialized than the Philippine tarsier or Horsfield's tarsier; for example, it lacks...

Speke's Gazelle
Speke's Gazelle
Gazella spekei

Speke's gazelle (*Gazella spekei*) is the smallest of the gazelle species. It is confined to the Horn of Africa, where it inhabits stony brush, grass steppes, and semi deserts. This species has been sometimes regarded as a subspecies of the *Dorcas*...

Speke's pectinator
Speke's pectinator
Pectinator spekei

Speke's pectinator (*Pectinator spekei*) or Speke's gundi, is a species of rodent in the family *Ctenodactylidae*. It is monotypic within the genus *Pectinator*. It is found in Djibouti, Eritrea, Ethiopia, and Somalia. Its natural habitats are...

Speothos pacivorus
Speothos pacivorus
Speothos pacivorus

The Pleistocene bush dog (*Speothos pacivorus*) is an extinct canid species in the genus *Speothos* from the Late Pleistocene. It was a relative of the extant bush dog. When compared to the bush dog, *S. pacivorus* had an overall larger body size, a...

Sperm Whale
Sperm Whale
Physeter macrocephalus

The Sperm whale or cachalot (*Physeter macrocephalus*) is the largest of the toothed whales and the largest toothed predator. It is the only living member of the genus *Physeter* and one of three extant species in the sperm whale family, along with...

Spermophilus brevicauda
Spermophilus brevicauda
Spermophilus brevicauda

Spermophilus brevicauda (Brandt's ground squirrel) is a species of rodent in the family Sciuridae. It is found in eastern Kazakhstan and the northern half of the Xinjiang region of China. A generic revision of the genus was undertaken in 2007...

Spermophilus pallidicauda
Spermophilus pallidicauda
Spermophilus pallidicauda

The pallid ground squirrel (Spermophilus pallidicauda) is a species of rodent in the family Sciuridae. It is found in Mongolia and the adjacent Nei Mongol autonomous region. A generic revision of the genus was undertaken in 2007 which helped...

Spermophilus ralli
Spermophilus ralli
Spermophilus ralli

The relict ground squirrel (Spermophilus relictus) is a species of rodent in the family Sciuridae. It is found in Kazakhstan, Kyrgyzstan and Uzbekistan. It is thought to be synonymous with Spermophilus ralli, whose name was formerly used for the...

Spermophilus relictus
Spermophilus relictus
Spermophilus relictus

The relict ground squirrel (Spermophilus relictus) is a species of rodent in the family Sciuridae. It is found in Kazakhstan, Kyrgyzstan and Uzbekistan. It is thought to be synonymous with Spermophilus ralli, whose name was formerly used for the...

Spider Monkey
Spider Monkey
Genus Ateles

Spider monkeys are New World monkeys belonging to the genus Ateles, part of the subfamily Atelinae, family Atelidae. Like other atelines, they are found in tropical forests of Central and South America, from southern Mexico to Brazil. The genus...

Spiked Atlantic spiny rat
Spiked Atlantic spiny rat
Trinomys paratus

The spiked Atlantic spiny-rat or Espirito Santo spiny rat (Trinomys paratus), is a spiny rat species from South America. It is found in Brazil.

Spinifex Hopping Mouse
Spinifex Hopping Mouse
Notomys alexis

The Spinifex hopping mouse (Notomys alexis) is a tiny semi-fossorial, burrowing rodent. It spends a great deal of energy not just foraging for food, but also transporting it back to their burrows.

Spinner Dolphin
Spinner Dolphin
Stenella longirostris

The spinner dolphin (Stenella longirostris) is a small dolphin found in off-shore tropical waters around the world. It is famous for its acrobatic displays in which it rotates around its longitudinal axis as it leaps through the air. It is a...

Spiny Ceram rat
Spiny Ceram rat
Rattus feliceus

The spiny Ceram rat (*Rattus feliceus*) is a species of rodent in the family Muridae. It is found only in Seram, Indonesia, which it is named after.
Spiny long-footed rat
Spiny long-footed rat
Tarsomys echinatus

The spiny long-footed rat (*Tarsomys echinatus*), also known as the Mindanao spiny rat, is a species of rodent in the family Muridae. It is found only in Mindanao, Philippines, in Bukidnon and South Cotabato provinces. It has been recorded in...
Spiny pocket mouse
Spiny pocket mouse
Chaetodipus spinatus

The spiny pocket mouse (*Chaetodipus spinatus*) is a species of rodent in the family Heteromyidae and order Rodentia. It is found in Baja California in Mexico and in Arizona, California and Nevada.
Spix's disk-winged bat
Spix's disk-winged bat
Thyroptera tricolor

Spix's disk-winged bat (*Thyroptera tricolor*), is a species of bat in the family Thyropteridae. It is native to the Americas from Mexico to Brazil. The most prominent anatomical feature of this bat, which distinguishes it from other species, is...
Spix's night monkey
Spix's night monkey
Aotus vociferans

Spix's night monkey (*Aotus vociferans*), also known as the Colombian gray night monkey, noisy night monkey and Spix's owl monkey, is a night monkey species from South America. It is found in Brazil, Colombia, Ecuador and Peru. The Spix's night...
Spix's red-handed howler
Spix's red-handed howler
Alouatta discolor

Spix's red-handed howler (*Alouatta discolor*) is a species of Howler monkey native to the southeastern Amazon in Brazil. It is threatened by deforestation and hunting.
Spix's white-fronted capuchin
Spix's white-fronted capuchin
Cebus unicolor

Spix's white-fronted capuchin (*Cebus unicolor*) is a species of gracile capuchin monkey. It had previously been classified as a subspecies of the Humboldt's white-fronted capuchin (*C. albifrons*). Following genetic studies by Boubli, et al,...
Spix's yellow-toothed cavy
Spix's yellow-toothed cavy
Galea spixii

Spix's yellow-toothed cavy (*Galea spixii*) is a rodent, a cavy species from South America. It is found in Bolivia east of the Andes and much of south central to northeastern Brazil. The species is found in open savanna and semiarid habitats, such...
Splendid climbing mouse
Splendid climbing mouse
Rhipidomys nitela

The splendid climbing mouse (*Rhipidomys nitela*) is a rodent species from South

America. It is found in Brazil, French Guiana, Guyana, Suriname and Venezuela.
Spotted bat
Spotted bat
Euderma maculatum

The spotted bat (*Euderma maculatum*) is a species of vesper bat and the only species of the genus *Euderma*.
Spotted bolo mouse
Spotted bolo mouse
Necromys punctulatus

The spotted bolo mouse or Ecuadorian akodont (*Necromys punctulatus*) is a species of rodent in the family Cricetidae. It is known from Ecuador and may also occur in Colombia. Little is known of its status and range.
Spotted free-tailed bat
Spotted free-tailed bat
Chaerephon bivittatus

The spotted free-tailed bat (*Chaerephon bivittatus*) is a species of bat in the family Molossidae. It is found in Burundi, Djibouti, Eritrea, Ethiopia, Kenya, Mozambique, Rwanda, South Sudan, Tanzania, Uganda, Zambia, and Zimbabwe. Its natural...
Spotted giant flying squirrel
Spotted giant flying squirrel
Petaurista elegans

The spotted giant flying squirrel (*Petaurista elegans*), also known as the lesser giant flying squirrel, is a species of rodent in the family Sciuridae. It is found in hill and mountain forests at altitudes of 200–4,000 m (660–13,120 ft) in...
Spotted Ground Squirrel
Spotted Ground Squirrel
Xerospermophilus spilosoma

The spotted ground squirrel (*Xerospermophilus spilosoma*) is a species of ground squirrel in the rodent family Sciuridae. It is found throughout Mexico and the central and western United States. Characterized by a white spotted back, the spotted...
Spotted Hyena
Spotted Hyena
Crocuta crocuta

A highly misunderstood animal, the Spotted hyena (*Crocuta crocuta*) has the reputation of being cowardly and sly, but it is actually a fascinating and intelligent creature with a very interesting social system. It looks like a dog but is closer...
Spotted linsang
Spotted linsang
Prionodon pardicolor

The spotted linsang (*Prionodon pardicolor*) is a linsang, a tree-dwelling carnivorous mammal, native to much of Southeast Asia. It is widely, though usually sparsely, recorded, and listed as Least Concern on the IUCN Red List.
Spotted Seal
Spotted Seal
Phoca largha

The spotted seal (*Phoca largha*), also known as the larga seal or largha seal, is a member of the family Phocidae, and is considered a "true seal". It inhabits ice floes and waters of the north Pacific Ocean and adjacent seas. It is primarily...

Spotted-Necked Otter
Spotted-Necked Otter
Hydricotis maculicollis

The spotted-necked otter (*Hydricotis maculicollis*), or speckle-throated otter, is an otter native to sub-Saharan Africa.

Spotted-winged fruit bat
Spotted-winged fruit bat
Balionycteris maculata

The spotted-winged fruit bat, (*Balionycteris maculata*) is the smallest megabat in the world, and the only species in the genus *Balionycteris*. It inhabits forests in Indonesia and Malaysia.

Springbok
Springbok
Antidorcas marsupialis

The springbok (*Antidorcas marsupialis*) is a medium-sized antelope found mainly in southern and southwestern Africa. The sole member of the genus *Antidorcas*, this bovid was first described by the German zoologist Eberhard August Wilhelm von...

Spurred roundleaf bat
Spurred roundleaf bat
Hipposideros calcaratus

The spurred roundleaf bat (*Hipposideros calcaratus*) is a species of bat in the family *Hipposideridae*. It is found in Papua New Guinea and the Solomon Islands.

Spurrell's free-tailed bat
Spurrell's free-tailed bat
Mops spurrelli

Spurrell's free-tailed bat (*Mops spurrelli*) is a species of bat in the family *Molossidae* named after Herbert George Flaxman Spurrell. It is found in Cameroon, Central African Republic, Ivory Coast, Equatorial Guinea, Ghana, Liberia, Sierra...

Spurrell's woolly bat
Spurrell's woolly bat
Kerivoula phalaena

Spurrell's woolly bat (*Kerivoula phalaena*) is a species of vesper bat in the family *Vespertilionidae* named after Herbert George Flaxman Spurrell. It is found in Cameroon, Democratic Republic of the Congo, Ivory Coast, Ghana, Guinea, Liberia, and...

Spy hociendo
Spy hociendo
Oxymycterus delator

The spy hociendo (*Oxymycterus delator*) is a species of rodent in the family *Cricetidae*. It is found only in Paraguay.

Squirrel Glider
Squirrel Glider
Petaurus norfolcensis

The squirrel glider (*Petaurus norfolcensis*) is a nocturnal gliding possum. The squirrel glider is one of the wrist-winged gliders of the genus *Petaurus*.

Squirrel-toothed rat
Squirrel-toothed rat
Anisomys imitator

The squirrel-toothed rat (*Anisomys imitator*), also known as the New Guinea giant rat, powerful-toothed rat, uneven-toothed rat, or narrow-toothed giant rat, is a

species of rodent in the family Muridae. It is the only species in the genus...
Sri Lankan axis deer
Sri Lankan axis deer
Axis axis ceylonensis

The Sri Lankan axis deer (*Axis axis ceylonensis*) or Ceylon spotted deer is a subspecies of axis deer (*Axis axis*) that inhabits only Sri Lanka. The name chital is not used in Sri Lanka. Its validity is disputed, and some maintain that the axis...
Sri Lankan Elephant
Sri Lankan Elephant
Elephas maximus maximus

The Sri Lankan elephant (*Elephas maximus maximus*) is one of three recognised subspecies of the Asian elephant. Its population is now largely restricted to the dry zone in the north, east and southeast of Sri Lanka. In the past, Sri Lankan...
Sri Lankan jackal
Sri Lankan jackal
Canis aureus naria

The Sri Lankan jackal (*Canis aureus naria*), also known as the Southern Indian jackal is a subspecies of golden jackal native to southern India and Sri Lanka. On the Asian mainland, the Sri Lankan jackal occurs in the whole southern part of the...
Sri Lankan Leopard
Sri Lankan Leopard
Panthera pardus kotiya

The Sri Lankan leopard (*Panthera pardus kotiya*) is a leopard subspecies native to Sri Lanka. It was first described in 1956 by Sri Lankan zoologist Paules Edward Pieris Deraniyagala. It is colloquially known as kotiyā and chiruthai.
Sri Lankan lion
Sri Lankan lion
Panthera leo sinhaleyus

The Sri Lankan lion (*Panthera leo sinhaleyus*), also known as the Ceylonese lion, is an extinct prehistoric subspecies of lion, excavated in Sri Lanka. It is believed to have become extinct prior to the arrival of culturally modern humans, C....
Sri Lankan long-tailed shrew
Sri Lankan long-tailed shrew
Crocidura miya

The Sri Lankan long-tailed shrew (*Crocidura miya*) is a species of mammal in the family Soricidae. It is endemic to Sri Lanka where it is known, in Sinhala, as ශ්‍රී ලංකා කුණු හික් මීයා. It is threatened by habitat loss.
Sri Lankan sambar deer
Sri Lankan sambar deer
Rusa unicolor unicolor

The Sri Lankan sambar or Indian sambar (*Rusa unicolor unicolor*) - it is also spelt "sambhur" or "sambur" - ගෝනා (gōṇā) in Sinhala, is a subspecies of the sambar that lives in India and Sri Lanka. This subspecies is one of the largest sambar...
Sri Lankan shrew
Sri Lankan shrew
Suncus fellowesgordoni

The Sri Lankan shrew (*Suncus fellowesgordoni*), also called Gordon's pygmy shrew, is a species of mammal in the family Soricidae. It is endemic to Sri Lanka. It is

threatened by habitat loss. It is known as හික් මීයා in Sinhala. It was named...
Sri Lankan sloth bear
Sri Lankan sloth bear
Melursus ursinus inornatus

The Sri Lankan sloth bear (*Melursus ursinus inornatus*) is a subspecies of the sloth bear which is found mainly in lowland dry forests in the island of Sri Lanka.
Sri Lankan Spotted Chevrotain
Sri Lankan Spotted Chevrotain
Moschiola meminna

Sri Lankan spotted chevrotain (*Moschiola meminna*) is an even-toed ungulate, member of the chevrotain family. These animals are the smallest hoofed mammals in the world.
St Kilda house mouse
St Kilda house mouse
Mus musculus muralis

The St Kilda house mouse (*Mus musculus muralis*) is an extinct subspecies of the house mouse found only on the islands of the St Kilda archipelago of northwest Scotland. They were first described, alongside the St Kilda field mouse, by natural...
St. Aignan's trumpet-eared bat
St. Aignan's trumpet-eared bat
Kerivoula agnella

St. Aignan's trumpet-eared bat (*Kerivoula agnella*) is a species of vesper bat in the family Vespertilionidae. It is found only in Papua New Guinea, and is only endemic to the islands, not the mainland. The species has been recorded on the islands...
Star-Nosed Mole
Star-Nosed Mole
Condylura cristata

The Star-nosed mole (*Condylura cristata*) is a small mole found in moist, low areas in the northern parts of North America. It is the only member of the tribe having a touch organ with more than 25,000-minute sensory receptors, known as Eimer's...
Steadfast tube-nosed fruit bat
Steadfast tube-nosed fruit bat
Paranyctimene tenax

The steadfast tube-nosed fruit bat (*Paranyctimene tenax*) is a species of megabat in the family Pteropodidae found in West Papua, Indonesia, and Papua New Guinea. The species was first proposed as a subgeneric arrangement of *Nyctimene*, published...
Steenbok
Steenbok
Raphicerus campestris

The steenbok (*Raphicerus campestris*) is a common small antelope of southern and eastern Africa. It is sometimes known as the steinbuck or steinbok.
Steere's spiny rat
Steere's spiny rat
Proechimys steerei

Steere's spiny rat, *Proechimys steerei*, is a spiny rat species from South America. It is found in Bolivia, Brazil and Peru.
Stein's cuscus
Stein's cuscus
Phalanger vestitus

Stein's cuscus (*Phalanger vestitus*) is a species of marsupial in the family Phalangeridae. It is found in Indonesia and Papua New Guinea.

Stein's paramelomys
Stein's paramelomys
Paramelomys steini

Stein's paramelomys, (*Paramelomys steini*), is a species of rodent native to Indonesia. Official observation of this species was only obtained on the Wayland Mountain on the island of New Guinea (Papua Province, Indonesia). It has only been...

Stein's rat
Stein's rat
Rattus steini

Stein's rat (*Rattus steini*), also known as the small spiny rat, is a species of rodent in the family Muridae. It is found in West Papua, Indonesia and Papua New Guinea.

Steinbach's tuco-tuco
Steinbach's tuco-tuco
Ctenomys steinbachi

Steinbach's tuco-tuco (*Ctenomys steinbachi*) is a species of rodent in the family Ctenomyidae. It is endemic to Bolivia. The species is named after zoological collector Dr. José Steinbach (1856-1929).

Stejneger's beaked whale
Stejneger's beaked whale
Mesoplodon stejnegeri

Stejneger's beaked whale (*Mesoplodon stejnegeri*), also known as the Bering Sea beaked whale or the saber-toothed whale, is a relatively unknown member of the genus *Mesoplodon* inhabiting the North Pacific Ocean. Leonhard Hess Stejneger collected...

Stella wood mouse
Stella wood mouse
Hylomyscus stella

The Stella *hylomyscus* or Stella wood mouse (*Hylomyscus stella*) is a species of rodent in the family Muridae. It is found in Burundi, Cameroon, Central African Republic, Republic of the Congo, Democratic Republic of the Congo, Equatorial Guinea,...

Steller Sea Lion
Steller Sea Lion
Eumetopias jubatus

The Steller sea lion (*Eumetopias jubatus*) is a near-threatened species of sea lion in the northern Pacific. It is the sole member of the genus *Eumetopias* and the largest of the eared seals (Otariidae). Among pinnipeds, it is inferior in size only...

Stephen Nash's titi
Stephen Nash's titi
Plecturocebus stephennashi

Stephen Nash's titi (*Plecturocebus stephennashi*), also known as just Nash's titi or Stephen Nash's monkey, is a species of titi, a type of New World monkey, endemic to the eastern bank of the Purus River in Brazil. It was discovered by Marc van...

Stephens's kangaroo rat
Stephens's kangaroo rat
Dipodomys stephensi

Stephens's kangaroo rat (*Dipodomys stephensi*) is a species of rodent in the family Heteromyidae. It is endemic to the Southern California region of the United States, primarily in western Riverside County. The species is named after American...

Steppe brown bear

Steppe brown bear

Ursus arctos priscus

The steppe brown bear (*Ursus arctos priscus*) is a disputed extinct subspecies of brown bear that lived in Eurasia during either the Pleistocene or the early Holocene epochs, but its geological age is uncertain. Fossils of the bear have been...

Steppe field mouse

Steppe field mouse

Apodemus witherbyi

The steppe field mouse (*Apodemus witherbyi*) is a species of rodent in the family Muridae found in Georgia, Armenia, Azerbaijan, Israel, Turkey, Jordan, Iran, Turkmenistan, Pakistan, and probably Afghanistan, Iraq, Lebanon, and Syria. The Mount...

Steppe Lemming

Steppe Lemming

Lagurus lagurus

The Steppe lemming (*Lagurus lagurus*) is a small rodent from Eastern Europe and parts of Asia. It is similar in appearance to the Norway lemming (*Lemmus lemmus*).

Steppe mouse

Steppe mouse

Mus spicilegus

The steppe mouse or mound-building mouse (*Mus spicilegus*) is a species of rodent in the family Muridae. It is found in grassland and other open areas in Austria, Bosnia and Herzegovina, Bulgaria, Croatia, Czech Republic, Hungary, North...

Steppe pika

Steppe pika

Ochotona pusilla

The steppe pika (*Ochotona pusilla*) is a small mammal of the pika family, Ochotonidae. It is found in the steppes of southern Russia and northern Kazakhstan.

Steppe Polecat

Steppe Polecat

Mustela eversmanii

The steppe polecat (*Mustela eversmanii*), also known as the white or masked polecat, is a species of mustelid native to Central and Eastern Europe and Central Asia. It is listed as Least Concern on the IUCN Red List because of its wide...

Steppe wolf

Steppe wolf

Canis lupus campestris

The steppe wolf (*Canis lupus campestris*), also known as the Caspian Sea wolf, is a subspecies of grey wolf native to the Caspian steppes, the steppe regions of the Caucasus, the lower Volga region, southern Kazakhstan north to the middle of the...

Stickeen brown bear

Stickeen brown bear

Ursus arctos stikeenensis

The Stickeen brown bear (*Ursus arctos stikeenensis*), also known as Stikine brown bear, is a large, North American brown bear that is most commonly dark brown in

color but can also range from blonde to black, featuring a distinctive hump on its...

Stiff-spine spiny rat

Stiff-spine spiny rat

Proechimys echinothrix

The stiff-spine spiny-rat, *Proechimys echinothrix*, or Tefe spiny rat, is a spiny rat species from South America. It is found in Brazil and Colombia.

Stirton's deer mouse

Stirton's deer mouse

Peromyscus stirtoni

Stirton's deer mouse (*Peromyscus stirtoni*) is a species of rodent in the family Cricetidae. It is found in El Salvador, Guatemala, Honduras, and Nicaragua. *P. stirtoni* is widely distributed and is presumed to have a large population and a...

Stock's vampire bat

Stock's vampire bat

Desmodus stocki

Desmodus stocki, or Stock's vampire bat, is an extinct species of vampire bat native to Pleistocene Florida, farther north than any living vampire bats. It weighed about 50% more than the common vampire bat (*Desmodus rotundus*), and was also more...

Stolička's mountain vole

Stolička's mountain vole

Alticola stoliczkanus

Stolička's or Stoliczka's mountain vole (*Alticola stoliczkanus*) is a species of rodent in the family Cricetidae. It is found in China, Pakistan, India and Nepal.

Stoliczka's trident bat

Stoliczka's trident bat

Aselliscus stoliczkanus

Stoliczka's trident bat (*Aselliscus stoliczkanus*) is a species of bat in the family Hipposideridae. It is found in China, Laos, Malaysia, Myanmar, Thailand, and Vietnam. In northern Thailand, it is found in Doi Pha Hom Pok National Park.

Stolzmann's crab-eating rat

Stolzmann's crab-eating rat

Ichthyomys stolzmanni

Stolzmann's crab-eating rat (*Ichthyomys stolzmanni*) is a species of rodent in the family Cricetidae. It is found in Ecuador and Peru. The species is believed to be nocturnal and semiaquatic, and has been found at an elevation range of 900 to...

Stone dormouse

Stone dormouse

Graphiurus rupicola

The stone dormouse (*Graphiurus rupicola*) is a species of rodent in the family Gliridae. It is found in Namibia, South Africa, and possibly Angola. Its natural habitat is rocky areas in the Karoo. Though it has a limited range, it is a fairly...

Stone Sheep

Stone Sheep

Ovis dalli stonei

The Stone sheep (*Ovis dalli stonei*) is the more southern subspecies of Thinhorn sheep, *Ovis dalli*.

Storey's African mole-rat

Storey's African mole-rat

Tachyoryctes storeyi

Storey's African mole-rat (*Tachyoryctes storeyi*) is a species of rodent in the family Spalacidae endemic to Kenya. Its natural habitat is subtropical or tropical dry lowland grassland. Some taxonomic authorities consider it to be conspecific with...

Strand's birch mouse

Strand's birch mouse

Sicista strandi

Strand's birch mouse (*Sicista strandi*) is a species of rodent in the family Sminthidae. It is endemic to Russia.

Strange big-eared brown bat

Strange big-eared brown bat

Histiotus alienus

The strange big-eared brown bat (*Histiotus alienus*), is a bat species from South America. It is found in Brazil.

Strap-Toothed Beaked Whale

Strap-Toothed Beaked Whale

Mesoplodon layardii

The Strap-toothed beaked whale (*Mesoplodon layardii*) is one of the largest members of toothed whales. They are the most poorly known group of large mammals.

Straw-Coloured Fruit Bat

Straw-Coloured Fruit Bat

Eidolon helvum

The straw-coloured fruit bat (*Eidolon helvum*) is a large fruit bat that is the most widely distributed of all the African megabats. It is quite common throughout its area ranging from the southwestern Arabian Peninsula, across forest and savanna...

Streaked dwarf porcupine

Streaked dwarf porcupine

Coendou ichillus

The streaked dwarf porcupine, *Coendou ichillus*, is a porcupine species in the family Erethizontidae. It is known from the lowlands (below an altitude of 400 m) of eastern Ecuador, and may be present in Peru as well. It appears to be nocturnal and...

Stripe-faced dunnart

Stripe-faced dunnart

Sminthopsis macroura

The striped-faced dunnart (*Sminthopsis macroura*) is a small, Australian, nocturnal, "marsupial mouse," part of the family Dasyuridae. The species' distribution occurs throughout much of inland central and northern Australia, occupying a range of...

Stripe-headed round-eared bat

Stripe-headed round-eared bat

Tonatia saurophila

The stripe-headed round-eared bat (*Tonatia saurophila*) is a species of bat from family Phyllostomidae. It can be found on forests in Central and South America.

Stripe-Necked Mongoose

Stripe-Necked Mongoose

Herpestes vitticollis

The Stripe-necked mongoose (*Urva vitticollis*) is a small terrestrial carnivorous mammal from South Asia. It was described by an English zoologist in 1835 and has

two recognized subspecies.
Striped Atlantic Forest rat
Striped Atlantic Forest rat
Delomys dorsalis

The striped Atlantic Forest rat (*Delomys dorsalis*) is a rodent species from South America. It is found in Argentina and Brazil.

Striped bandicoot
Striped bandicoot
Microperoryctes longicauda

The striped bandicoot (*Microperoryctes longicauda*) is a species of marsupial in the family Peramelidae. It is found in West Papua, Indonesia and Papua New Guinea. Its natural habitat is subtropical or tropical dry forests. The Striped bandicoot...

Striped bush squirrel
Striped bush squirrel
Paraxerus flavovittis

The striped bush squirrel (*Paraxerus flavovittis*) is a species of rodent in the family Sciuridae found in Kenya, Malawi, Mozambique, and Tanzania. Its natural habitats are moist savanna and plantations.

Striped Dolphin
Striped Dolphin
Stenella coeruleoalba

The striped dolphin (*Stenella coeruleoalba*) is an extensively studied dolphin found in temperate and tropical waters of all the world's oceans. It is a member of the oceanic dolphin family, Delphinidae.

Striped Field Mouse
Striped Field Mouse
Apodemus agrarius

The striped field mouse (*Apodemus agrarius*) is a rodent in the family Muridae. The range of this species stretches from Eastern Europe to Eastern Asia.

Striped Ground Squirrel
Striped Ground Squirrel
Xerus erythropus

The striped ground squirrel (*Euxerus erythropus*) is a species of squirrel native to Africa. It was first described by Geoffroy in 1803, but the original publication may be unavailable, so that the binomial authority is today more often cited as...

Striped hairy-nosed bat
Striped hairy-nosed bat
Mimon crenulatum

The striped hairy-nosed bat (*Mimon crenulatum*) is a bat species from South and Central America. It is found in Brazil, Colombia, Ecuador, French Guiana, Guyana, Suriname and Venezuela.

Striped Hog-Nosed Skunk
Striped Hog-Nosed Skunk
Conepatus semistriatus

The Striped hog-nosed skunk (*Conepatus semistriatus*) is a skunk species from Central and South America. It is considered a generalist species because it can thrive in, and withstand disturbed environmental conditions. Striped hog-nosed skunks can...

Striped Hyena
Striped Hyena

Hyaena hyaena

Hyenas are the most common large carnivore in Africa. Hyenas are neither members of the dog nor the cat family. They are unique, and so have their own family: Hyaenidae, which has four members: Striped hyenas, the "laughing" Spotted hyenas, Brown...

Striped leaf-nosed bat

Striped leaf-nosed bat

Macronycteris vittatus

The striped leaf-nosed bat (*Macronycteris vittatus*) is a species of bat native to eastern and southern Africa. It was formerly considered part of *M. commersoni*, which is now viewed as being restricted to Madagascar. Both *commersoni* and it were...

Striped Polecat

Striped Polecat

Ictonyx striatus

The striped polecat (*Ictonyx striatus*) - also called the African polecat, zoril, zorille, zorilla, Cape polecat, and African skunk - is a member of the family Mustelidae that resembles a skunk (of the family Mephitidae). The name "zorilla" comes...

Striped Possum

Striped Possum

Dactylopsila trivirgata

The striped possum or common striped possum (*Dactylopsila trivirgata*) is a member of the marsupial family Petauridae. It is found mainly in New Guinea. The species is black with three white stripes running head to tail, and its head has white...

Striped Skunk

Striped Skunk

Mephitis mephitis

The Striped skunk (*Mephitis mephitis*) is a skunk of the genus *Mephitis* that occurs across much of North America. These animals possess highly developed musk-filled scent glands to ward off predators. They have a long history of association with...

Striped treeshrew

Striped treeshrew

Tupaia dorsalis

The striped treeshrew (*Tupaia dorsalis*) is a treeshrew species within the Tupaiidae. It is endemic to Borneo and known only from a few individuals in Sabah, Sarawak, Brunei and Kalimantan.

Strong tuco-tuco

Strong tuco-tuco

Ctenomys validus

The strong tuco-tuco (*Ctenomys validus*) is a species of rodent in the family Ctenomyidae. It is endemic to Argentina.

Strong-tailed Oldfield mouse

Strong-tailed Oldfield mouse

Thomasomys ischyrius

The strong-tailed Oldfield mouse (*Thomasomys ischyrius*) is a species of rodent in the family Cricetidae. It is found only in Peru.

Stuhlmann's golden mole

Stuhlmann's golden mole

Chrysochloris stuhlmanni

Stuhlmann's golden mole (*Chrysochloris stuhlmanni*) is a species of mammal in the family Chrysochloridae. It is found in Cameroon, Democratic Republic of the Congo, Kenya, Tanzania, and Uganda. Its natural habitats are subtropical or tropical...

Stump-Tailed Macaque

Stump-Tailed Macaque

Macaca arctoides

The stump-tailed macaque (*Macaca arctoides*), also called the bear macaque, is a species of macaque native to South Asia and Southeast Asia. In India, it occurs south of the Brahmaputra River, in the northeastern part of the country. Its range in...

Stump-tailed porcupine

Stump-tailed porcupine

Coendou rufescens

The stump-tailed porcupine (*Coendou rufescens*) is a species of rodent in the family Erethizontidae. It is found mainly in Colombia, with a few records from Ecuador. This species was formerly sometimes assigned to *Echinoprocta*, a genus no longer...

Sturdee's pipistrelle

Sturdee's pipistrelle

Pipistrellus sturdeeii

Sturdee's pipistrelle (*Pipistrellus sturdeeii*), also known as the Bonin pipistrelle bat, is a bat that was thought to have lived in Japan before officially becoming extinct in 2000. In 2020 the IUCN changed its official status to "extinct".

Sturnira bakeri

Sturnira bakeri

Sturnira bakeri

Sturnira bakeri is a species of bat found in South-America.

Sturnira hondurensis

Sturnira hondurensis

Sturnira hondurensis

Sturnira hondurensis is a species of bat found in Central America. Previously, it was considered a subspecies of the highland yellow-shouldered bat, but it has been considered distinct since 2010.

Sturnira koopmanhilli

Sturnira koopmanhilli

Sturnira koopmanhilli

Sturnira koopmanhilli is a species of leaf-nosed bat found in South America.

Sturnira perla

Sturnira perla

Sturnira perla

Sturnira perla is a species of yellow-shouldered bat found in Ecuador.

Subalpine woolly rat

Subalpine woolly rat

Mallomys istapantap

The subalpine woolly rat (*Mallomys istapantap*) is a species of rodent in the family Muridae. It is found in West Papua, Indonesia and Papua New Guinea.

Subantarctic Fur Seal

Subantarctic Fur Seal

Arctocephalus tropicalis

The subantarctic fur seal (*Arctocephalus tropicalis*) is found in the southern

parts of the Indian, Pacific, and Atlantic Oceans. It was first described by Gray in 1872 from a specimen recovered in northern Australia—hence the inappropriate...
Submyotodon moupinensis
Submyotodon moupinensis
Submyotodon moupinensis

The Moupin broad-muzzled bat (*Submyotodon moupinensis*) is a bat in the family Vespertilionidae endemic to southern China.
Subtropical antechinus
Subtropical antechinus
Antechinus subtropicus

The subtropical antechinus (*Antechinus subtropicus*) is a species of small carnivorous marsupial of the family Dasyuridae. It was previously thought to be conspecific with the brown antechinus (*Antechinus stuartii*). The subtropical antechinus is...
Sucre spiny rat
Sucre spiny rat
Proechimys urichi

The Sucre spiny-rat (*Proechimys urichi*) is a species of rodent in the family Echimyidae. It is endemic to Venezuela.
Sudan gerbil
Sudan gerbil
Gerbillus nancillus

The Sudan gerbil (*Gerbillus nancillus*) is distributed mainly in central Sudan.
Sudanian grass rat
Sudanian grass rat
Arvicanthis ansorgei

The Sudanian grass rat (*Arvicanthis ansorgei*) is a species of rodent in the family Muridae. It is found in Burkina Faso, Guinea-Bissau, Mali, Niger, Senegal, possibly Benin, possibly Ivory Coast, possibly Ghana, possibly Guinea, and possibly...
Sugar Glider
Sugar Glider
Petaurus breviceps

The Sugar glider (*Petaurus breviceps*) is a small arboreal gliding possum that belongs to the marsupial infraclass. It is so called due to loving sweet food such as sugar and honey, while the word 'glider' refers to their gliding habit when moving...
Sula rat
Sula rat
Rattus elaphinus

The Sula rat (*Rattus elaphinus*) is a species of rodent in the family Muridae. It is found only in Indonesia, on Taliabu and Mangole islands in the Sula Archipelago. On Sanana island, only the introduced *Rattus tanezumi* and *Rattus exulans* are found.
Sulawesi Bear Cuscus
Sulawesi Bear Cuscus
Ailurops ursinus

The Sulawesi bear cuscus (*Ailurops ursinus*) is a species of arboreal marsupial found in Southeast Asia.
Sulawesi dwarf cuscus
Sulawesi dwarf cuscus
Strigocuscus celebensis

The Sulawesi dwarf cuscus (*Strigocuscus celebensis*) is a species of arboreal marsupial in the family Phalangeridae endemic to Sulawesi and nearby islands in Indonesia. It inhabits tropical moist lowland forest and is nocturnal, folivorous and...

Sulawesi flying fox
Sulawesi flying fox
Acerodon celebensis

The Sulawesi flying fox or Sulawesi fruit bat (*Acerodon celebensis*) is a species of megabat endemic to Indonesia. It is classified as "Vulnerable" by the IUCN due to unsustainable levels of hunting.

Sulawesi forest rat
Sulawesi forest rat
Taeromys punicans

The Sulawesi forest rat (*Taeromys punicans*) is a species of rodent in the family Muridae. It is found only in central Sulawesi, Indonesia. It is known only from Pinedapa, Poso Pesisir, Poso Regency.

Sulawesi free-tailed bat
Sulawesi free-tailed bat
Mops sarasinorum

The Sulawesi free-tailed bat (*Mops sarasinorum*) is a species of bat in the family Molossidae. It is found in Indonesia and the Philippines.

Sulawesi giant rat
Sulawesi giant rat
Paruromys dominator

The Sulawesi giant rat (*Paruromys dominator*) is a species of rodent in the family Muridae. It is monotypic in the genus *Paruromys*. The species is endemic to Sulawesi in Indonesia, where it inhabits forests at elevations from sea level to the...

Sulawesi harpy fruit bat
Sulawesi harpy fruit bat
Harpyionycteris celebensis

The Sulawesi harpy fruit bat (*Harpyionycteris celebensis*) is a species of megabat in the family Pteropodidae. It is endemic to Indonesia.

Sulawesi horseshoe bat
Sulawesi horseshoe bat
Rhinolophus celebensis

The Sulawesi horseshoe bat (*Rhinolophus celebensis*) is a species of bat in the family Rhinolophidae. It is endemic to Indonesia.

Sulawesi montane rat
Sulawesi montane rat
Taeromys hamatus

The Sulawesi montane rat (*Taeromys hamatus*) is a species of rodent in the family Muridae. It is found only in central Sulawesi, Indonesia, on Mount Lohi, Mount Kanino, and Mount Nokilalaki.

Sulawesi naked-backed fruit bat
Sulawesi naked-backed fruit bat
Dobsonia exoleta

The Sulawesi naked-backed fruit bat (*Dobsonia exoleta*) is a species of megabat in the family Pteropodidae. It is endemic to Indonesia.

Sulawesi palm civet
Sulawesi palm civet
Macrogalidia musschenbroekii

The Sulawesi palm civet (*Macrogalidia musschenbroekii*), also known as Sulawesi civet, musang and brown palm civet is a little-known palm civet endemic to Sulawesi. It is listed as Vulnerable on the IUCN Red List due to population decline...

Sulawesi root rat
Sulawesi root rat
Gracilimus radix

Gracilimus radix, the Sulawesi root rat, is a species of rat in the family Muridae. It is a monotypic species, as it the only member of the genus *Gracilimus*. It is only known from the Mamasa Regency in Sulawesi, Indonesia, where it is endemic to...

Sulawesi rousette
Sulawesi rousette
Rousettus celebensis

The Sulawesi rousette or Sulawesi fruit bat (*Pilonycteris celebensis*) is a species of megabat in the family Pteropodidae endemic to Sulawesi, an island in Indonesia. It is presently the only member of the genus *Pilonycteris*.

Sulawesi shrew
Sulawesi shrew
Crocidura lea

The Sulawesi shrew (*Crocidura lea*) is a species of mammal in the family Soricidae. It is endemic to the central and northern provinces of Sulawesi in Indonesia. It is a fairly common species and the International Union for Conservation of Nature...

Sulawesi soft-furred rat
Sulawesi soft-furred rat
Eropeplus canus

The Sulawesi soft-furred rat (*Eropeplus canus*) is a species of rodent in the family Muridae. It is the only species in the genus *Eropeplus*. It is found only in Indonesia. Its natural habitat is subtropical or tropical dry forest. It is threatened...

Sulawesi stripe-faced fruit bat
Sulawesi stripe-faced fruit bat
Styloctenium wallacei

The Wallace's or Sulawesi stripe-faced fruit bat (*Styloctenium wallacei*) is a species of megabat in the family Pteropodidae. It is endemic to Sulawesi and the nearby Togian Islands of Indonesia. Cave paintings resembling these bats have been...

Sulawesi tiny shrew
Sulawesi tiny shrew
Crocidura levicula

The Sulawesi tiny shrew (*Crocidura levicula*) is a species of mammal in the family Soricidae. It is endemic to the island of Sulawesi in Indonesia.

Sulawesi white-handed shrew
Sulawesi white-handed shrew
Crocidura rhoditis

The Sulawesi white-handed shrew or Temboan shrew (*Crocidura rhoditis*) is a species of mammal in the family Soricidae. It is endemic to the island of Sulawesi in Indonesia. It is a fairly common species and the population seems stable so the...

Sulawesi yellow bat
Sulawesi yellow bat
Scotophilus celebensis

The Sulawesi yellow bat (*Scotophilus celebensis*) is a species of vesper bat. It is found only in Indonesia.

Sulawesian shrew rat

Sulawesian shrew rat

Melasmothrix naso

The Sulawesian shrew rat (*Melasmothrix naso*) is a species of rodent in the family Muridae. It is the only species in the genus *Melasmothrix*. It is found only in central Sulawesi, Indonesia, and is known from the localities of Rano Rano and Mount...

Sumatran clouded leopard

Sumatran clouded leopard

Neofelis diardi diardi

The Sumatran clouded leopard (*Neofelis diardi diardi*) is a subspecies of the Sunda clouded leopard and is native to the Indonesian islands of Sumatra and Batu. It differs in molecular, craniomandibular and dental characteristics from the Bornean...

Sumatran dhole

Sumatran dhole

Cuon alpinus sumatrensis

The dhole (; *Cuon alpinus*) is a canid native to Central, South, East, and Southeast Asia. Other English names for the species include Asian wild dog, Asiatic wild dog, Indian wild dog, whistling dog, red dog, and mountain wolf. It is genetically...

Sumatran Elephant

Sumatran Elephant

Elephas maximus sumatranus

The Sumatran elephant (*Elephas maximus sumatranus*) is one of three recognized subspecies of the Asian elephant. In 2011, IUCN upgraded the conservation status of the Sumatran elephant from endangered to critically endangered in its Red List as...

Sumatran flying squirrel

Sumatran flying squirrel

Hylopetes winstoni

The Sumatran flying squirrel (*Hylopetes winstoni*) is a flying squirrel only found on the island of Sumatra. It is listed as data deficient on the IUCN red list. Originally discovered in 1949, it is known only from a single specimen. It is a...

Sumatran giant shrew

Sumatran giant shrew

Crocidura lepidura

The Sumatran giant shrew (*Crocidura lepidura*) is a shrew of the genus *Crocidura*. It is native to the Indonesian island of Sumatra, where it is found both in the rainforests to the west of the island and in the hillsides in the south and east....

Sumatran lar gibbon

Sumatran lar gibbon

Hylobates lar vestitus

The Sumatran lar gibbon (*Hylobates lar vestitus*), also known as the Sumatran white-handed gibbon, is a subspecies of the lar gibbon, a primate in the gibbon family Hylobatidae. It is native to the island of Sumatra, Indonesia. It shares the tree...

Sumatran long-tailed shrew

Sumatran long-tailed shrew
Crocidura paradoxura

The Sumatran long-tailed shrew (*Crocidura paradoxura*) is a species of mammal in the family Soricidae. It is endemic to Indonesia. It is threatened by habitat loss.
Sumatran mastiff bat
Sumatran mastiff bat
Mormopterus doriae

The Sumatran mastiff bat (*Mormopterus doriae*) is a species of bat in the family Molossidae, the free-tailed bats. It is only known from Sumatra in Indonesia. It was described in 1907 and has not been recorded since.
Sumatran muntjac
Sumatran muntjac
Muntiacus montanus

The Sumatran muntjac (*Muntiacus muntjak montanus*) is a subspecies of Indian muntjac in the deer family which can be the size of a large dog. It was discovered in 1914, but had not been sighted since 1930 until one was snared and freed from a...
Sumatran Orangutan
Sumatran Orangutan
Pongo abelii

The Sumatran orangutan (*Pongo abelii*) is one of the three species of orangutans. Critically Endangered, and found only in the north of the Indonesian island of Sumatra, it is rarer than the Bornean orangutan but more common than the recently...
Sumatran porcupine
Sumatran porcupine
Hystrix sumatrae

The Sumatran porcupine (*Hystrix sumatrae*) is a species of rodent in the family Hystricidae. It is endemic to the Indonesian island of Sumatra, where it is hunted for food.
Sumatran Rhinoceros
Sumatran Rhinoceros
Dicerorhinus sumatrensis

The Sumatran rhinoceros (*Dicerorhinus sumatrensis*) is a rare member of the family Rhinocerotidae and one of five extant species of rhinoceros. It is now critically endangered, with only five substantial populations in the wild. Sumatran rhinos,...
Sumatran Serow
Sumatran Serow
Capricornis sumatraensis sumatraensis

The Sumatran serow (*Capricornis sumatraensis sumatraensis*) is a subspecies of the Mainland serow. It is also known as the Southern serow and was previously considered its own species.
Sumatran shrewlike mouse
Sumatran shrewlike mouse
Mus crociduroides

The Sumatran shrewlike mouse (*Mus crociduroides*) is a species of mouse. It is endemic to Indonesia. This mouse is known only from the mountains of western Sumatra, where it lives in rainforest habitat. It occurs in Kerinci Seblat National Park.
Sumatran spiny rat
Sumatran spiny rat

Maxomys hylomyoides

The Sumatran spiny rat (*Maxomys hylomyoides*) is a species of rodent in the family Muridae. It is found only in Indonesia.

Sumatran Striped Rabbit

Sumatran Striped Rabbit

Nesolagus netscheri

The Sumatran striped rabbit (*Nesolagus netscheri*) is a rare species of rabbit found only in forests in western Sumatra, Indonesia, and surrounding areas. It was last seen in 1972 and went unreported until an individual rabbit was photographed in...

Sumatran surili

Sumatran surili

Presbytis melalophos

The black-crested Sumatran langur (*Presbytis melalophos*) is a species of primate in the family Cercopithecidae. It is endemic to Sumatra in Indonesia. Its natural habitat is subtropical or tropical dry forests. It is threatened by habitat...

Sumatran Tiger

Sumatran Tiger

Panthera tigris sondaica

The Sumatran tiger is a population of Sunda Island tiger (*Panthera tigris sondaica*) on the Indonesian island of Sumatra. This is the only surviving tiger population in the Sunda Islands, where the Bali and Javan tigers are extinct. Sequences from...

Sumatran water shrew

Sumatran water shrew

Chimarrogale sumatrana

The Sumatran water shrew (*Chimarrogale sumatrana*) is a red-toothed shrew found only in the Padang highlands of western Sumatra, Indonesia. Its natural habitats are streams in montane forests. The species is only known from a holotype, which is...

Sumba roundleaf bat

Sumba roundleaf bat

Hipposideros sumbae

The Sumba roundleaf bat (*Hipposideros sumbae*) is a species of bat in the family Hipposideridae. It lives in Indonesia and East Timor. It is present on the islands of Sumba, Rote, Sumbawa, Flores, Semau, and Savu.

Sumichrast's harvest mouse

Sumichrast's harvest mouse

Reithrodontomys sumichrasti

Sumichrast's harvest mouse (*Reithrodontomys sumichrasti*) is a species of rodent in the family Cricetidae. It is found in Costa Rica, El Salvador, Guatemala, Honduras, Mexico, Nicaragua, and Panama.

Sumichrast's vesper rat

Sumichrast's vesper rat

Nyctomys sumichrasti

Sumichrast's vesper rat (*Nyctomys sumichrasti*) is a rodent of the family Cricetidae found from southern Mexico to Panama. It is named for the collector of the first specimen, and its closest relative is probably Hatt's vesper rat, a similar, but...

Summit rat

Summit rat

Rattus baluensis

The summit rat (*Rattus baluensis*) is a species of rodent in the family Muridae. It is found only on Mount Kinabalu and Mount Tambuyukon, Malaysia, and has been recorded at altitudes of 2,040 to 2,477 m on Mt. Tambuyukon and 2,670 to 3,426 m on...

Sun Bear

Sun Bear

Helarctos malayanus

The Sun bear (*Helarctos malayanus*) is found in the tropical forests of Southeast Asia. The animal gets its name from the characteristic orange-to-cream-colored chest patch. It is also known as the "honey bear", for its love of honeycombs and...

Sun-tailed monkey

Sun-tailed monkey

Allochrocebus solatus

The sun-tailed monkey (*Allochrocebus solatus*) from Gabon is one of the least studied primates in its habitat. It was discovered as a new species in 1988, and is classified as a guenon, which is a member of the genus *Cercopithecus*, but was...

Sunburned rat

Sunburned rat

Rattus adustus

The sunburned rat (*Rattus adustus*) is a species of rat from Enggano Island in Indonesia. It is only known from the holotype and has not been recorded since its description in 1940.

Sunda Clouded Leopard

Sunda Clouded Leopard

Neofelis diardi

The Sunda clouded leopard (*Neofelis diardi*) is a medium-sized wild cat native to Borneo and Sumatra. In 2006, it was classified as a species, distinct from the Clouded leopard in mainland Southeast Asia.

Sunda flying fox

Sunda flying fox

Acerodon mackloti

The Sunda flying fox or Sunda fruit bat (*Acerodon mackloti*) is a species of bat in the family Pteropodidae. It is endemic to Indonesia.

Sunda Flying Lemur

Sunda Flying Lemur

Galeopterus variegatus

The Sunda flying lemur (*Galeopterus variegatus*), also known as Sunda colugo, Malayan flying lemur and Malayan colugo, is a colugo species. It is native throughout Southeast Asia ranging from southern Myanmar, Thailand, southern Vietnam, Malaysia...

Sunda Leopard Cat

Sunda Leopard Cat

Prionailurus javanensis

The Sunda leopard cat (*Prionailurus javanensis*) is a small wild cat species native to the Sundaland islands of Java, Bali, Borneo, Sumatra and the Philippines that is considered distinct from the leopard cat occurring in mainland South and...

Sunda long-eared bat

Sunda long-eared bat

Nyctophilus heran

The Sunda long-eared bat (*Nyctophilus heran*) is a species of vesper bat. It is

found only in Indonesia.

Sunda Pangolin

Sunda Pangolin

Manis javanica

The Sunda pangolin (*Manis javanica*), also known as the Malayan or Javan pangolin, is a species of pangolin. It is found throughout Southeast Asia, including Myanmar, Thailand, Cambodia, Laos, Malaysia, Singapore, Vietnam, and the islands of...

Sunda porcupine

Sunda porcupine

Hystrix javanica

The Sunda porcupine (*Hystrix javanica*) is a species of rodent in the family Hystricidae. It is endemic to Indonesia. Due to the popularity of the hunting and consumption of the Sunda porcupine as an aphrodisiac, the Ministry of Environment and...

Sunda shrew

Sunda shrew

Crocidura monticola

The Sunda shrew (*Crocidura monticola*) is a species of mammal in the family Soricidae found in Indonesia and Malaysia. Populations from the Malay Peninsula, Sumatra and Borneo are now considered to belong to *Crocidura neglecta*. The widespread...

Sunda Slow Loris

Sunda Slow Loris

Nycticebus coucang

The Sunda slow loris (*Nycticebus coucang*) or greater slow loris is a strepsirrhine primate and a species of slow loris native to Indonesia, West Malaysia, southern Thailand and Singapore. It measures 27 to 38 cm (11 to 15 in) from head to tail...

Sunda Stink Badger

Sunda Stink Badger

Mydaus javanensis

The Sunda stink badger (*Mydaus javanensis*) is a mammal native to Indonesia and Malaysia. Despite the common name, stink badgers are not closely related to true badgers, and are, instead, Old World relatives of the skunks.

Sundaic mountain leopoldamys

Sundaic mountain leopoldamys

Leopoldamys ciliatus

The Sundaic mountain leopoldamys (*Leopoldamys ciliatus*) is a species of rodent from the family Muridae. It was formerly considered a subspecies of Edwards's long-tailed giant rat, and it has one synonym: *Leopoldamys setiger*. The Sundaic mountain...

Sundevall's jird

Sundevall's jird

Meriones crassus

Sundevall's jird (*Meriones crassus*) is a species of rodent in the family of Muridae. It is found in Afghanistan, Algeria, Egypt, Iran, Iraq, Israel, Jordan, Kuwait, Libya, Pakistan, Morocco, Niger, Palestine, Oman, Saudi Arabia, Sudan, Syria,...

Sundevall's roundleaf bat

Sundevall's roundleaf bat

Hipposideros caffer

Sundevall's roundleaf bat (*Hipposideros caffer*), also called Sundevall's leaf-

nosed bat, is a species of bat in the family Hipposideridae. These bats are very similar in appearance to the closely related Noack's roundleaf bat, and the two have in...

Suni

Suni

Nesotragus moschatus

The Suni (*Nesotragus moschatus*) is a small antelope. It occurs in dense underbrush from central Kenya to KwaZulu-Natal in South Africa. Suni are around 30 to 43 centimetres (12 to 17 inches) high at the shoulder and weigh 4.5 to 5.4 kilograms...

Superagüi lion tamarin

Superagüi lion tamarin

Leontopithecus caissara

The black-faced lion tamarin or Superagüi lion tamarin (*Leontopithecus caissara*) is a small New World monkey of the family Callitrichidae. It is endangered and endemic to coastal forests in southeastern Brazil. There are several conservation...

Surat serotine

Surat serotine

Eptesicus dimissus

The Surat helmeted bat (*Cassidix dimissus*) is a species of vesper bat. It ranges from Thailand and Laos west to Nepal.

Surdisorex schlitteri

Surdisorex schlitteri

Surdisorex schlitteri

Represented by a single specimen, the Mount Elgon mole shrew (*Surdisorex schlitteri*) is a species of African mole shrew known to date only from the ericaceous zone of Mt. Elgon in Western Kenya. In 1984, a specimen of the genus *Surdisorex* was...

Suriname lowland forest cottontail

Suriname lowland forest cottontail

Sylvilagus parentum

The Suriname tapeti (*Sylvilagus parentum*) or Suriname lowland forest cottontail is a South American species of cottontail rabbit described in 2017. It is known from the lowlands of western Suriname, and was described from specimens collected by...

Svalbard reindeer

Svalbard reindeer

Rangifer tarandus platyrhynchus

The Svalbard reindeer (*Rangifer tarandus platyrhynchus*) is a small subspecies of reindeer found on the Svalbard archipelago of Norway. They are the only large grazing mammal in the European High Arctic, and this makes them exceptional for...

Swamp antechinus

Swamp antechinus

Antechinus minimus

The swamp antechinus (*Antechinus minimus*), also known as the little Tasmanian marsupial mouse, is a species of shrew-like marsupial of the family Dasyuridae and as such is related to dunnarts, quolls and the Tasmanian devil.

Swamp musk shrew

Swamp musk shrew

Crocidura mariquensis

The swamp musk shrew (*Crocidura mariquensis*), or musk shrew, is a species of mammal in the family Soricidae. It occurs in Angola, Botswana, Democratic Republic

of the Congo, Mozambique, Namibia, South Africa, Eswatini, Zambia and Zimbabwe.
Its...
Swamp Rabbit
Swamp Rabbit
Sylvilagus aquaticus

The Swamp rabbit (*Sylvilagus aquaticus*) is a large cottontail rabbit found in the southern United States. As its name implies it has a strong preference for wet areas, and it will readily take to the water and swim.
Swamp Wallaby
Swamp Wallaby
Wallabia bicolor

The swamp wallaby (*Wallabia bicolor*) is a small macropod marsupial of eastern Australia. This wallaby is also commonly known as the black wallaby, with other names including black-tailed wallaby, fern wallaby, black pademelon, stinker (in...
Swarthy gerbil
Swarthy gerbil
Gerbillus aquilus

The swarthy gerbil (*Gerbillus aquilus*) is distributed mainly in eastern Iran, southern Afghanistan, and western Pakistan.
Swayne's hartebeest
Swayne's hartebeest
Alcelaphus buselaphus swaynei

Swayne's hartebeest (*Alcelaphus buselaphus swaynei*) is an endangered antelope native to Ethiopia. Two of the largest remaining populations are located in Senkelle Swayne's Hartebeest Sanctuary, Nechisar National Park and Maze National Park. It...
Swift Fox
Swift Fox
Vulpes velox

The Swift fox (*Vulpes velox*) is a small fox around the size of a domestic cat found in the western grasslands of North America. It is closely related to the Kit fox. The Swift fox became nearly extinct in the 1930s as a result of predator control...
Swift fruit bat
Swift fruit bat
Thoopterus nigrescens

The swift fruit bat (*Thoopterus nigrescens*) is a species of megabat in the family Pteropodidae.
Swinhoe's striped squirrel
Swinhoe's striped squirrel
Tamiops swinhoei

Swinhoe's striped squirrel is a small species of rodent in the family Sciuridae known scientifically as *Tamiops swinhoei*. This species is found mostly in China and Southeast Asia. Their diet consists of mostly seeds, fruits, nuts and ginger...
Swinny's horseshoe bat
Swinny's horseshoe bat
Rhinolophus swinnyi

Swinny's horseshoe bat (*Rhinolophus swinnyi*) is a species of bat in the family Rhinolophidae. In English, *R. swinnyi* is commonly referred to as Swinny's horseshoe bat. In Afrikaans, it is commonly referred to as Swinny se saalneusvlermuis.
This...
Swynnerton's bush squirrel

Swynnerton's bush squirrel
Paraxerus vexillarius

Swynnerton's bush squirrel (*Paraxerus vexillarius*), also known as Svynnerton's bush squirrel, is a species of rodent in the family Sciuridae endemic to Tanzania. Its natural habitat is subtropical or tropical moist montane forests. It is...

Sykes' monkey
Sykes' monkey
Cercopithecus albogularis

Sykes' monkey (*Cercopithecus albogularis*), also known as the white-throated monkey or Samango monkey, is an Old World monkey found between Ethiopia and South Africa, including south and east Democratic Republic of Congo. It is named after...

Sylvilagus palustris hefneri
Sylvilagus palustris hefneri
Sylvilagus palustris hefneri

Sylvilagus palustris hefneri, also known as the Lower Keys marsh rabbit, is an endangered subspecies of marsh rabbit named after Playboy founder Hugh Hefner.

Syrian brown bear
Syrian brown bear
Ursus arctos syriacus

The Syrian brown bear (*Ursus arctos syriacus* or *Ursus arctos arctos*) is a relatively small subspecies of brown bear native to the Middle East and the Caucasus mountain range.

Syrian elephant
Syrian elephant
Elephas maximus asurus

The Syrian elephant or Western Asiatic elephant (*Elephas maximus asurus*) was the westernmost population of the Asian elephant (*Elephas maximus*), which became extinct in ancient times. Skeletal remains of *E. m. asurus* have been recorded from the...

Syrian Hamster
Syrian Hamster
Mesocricetus auratus

The Syrian hamster (*Mesocricetus auratus*) is a small rodent belonging to the hamster subfamily, Cricetinae. In 1930, Professor Aharoni captured a female Syrian hamster (otherwise called Golden or Teddy bear hamster) with her twelve pups in Aleppo...

Syrian wild ass
Syrian wild ass
Equus hemionus hemippus

The Syrian wild ass (*Equus hemionus hemippus*), less commonly known as a hemippe, an achdari, or a Mesopotamian or Syrian onager, is an extinct subspecies of onager native to the Arabian peninsula and surrounding areas. It ranged across...

Szechuan vole
Szechuan vole
Volemys millicens

The Szechuan vole (*Volemys millicens*) is a species of rodent in the family Cricetidae. It is found only in northwestern Sichuan, China. It is one of two species in the genus *Volemys* along with Marie's vole (*Volemys musseri*).

Szechwan myotis
Szechwan myotis
Myotis altarium

The Szechwan myotis (*Myotis altarium*) is a species of vesper bat. It is found in China and Thailand.

Tacarcuna bat

Tacarcuna bat

Lasiurus castaneus

The Tacarcuna bat (*Lasiurus castaneus*) is a species of vesper bat. It is found in Costa Rica, Panama and possibly Colombia.

Tacoma pocket gopher

Tacoma pocket gopher

Thomomys mazama tacomensis

The Tacoma pocket gopher (*Thomomys mazama tacomensis*), was a subspecies of the Mazama pocket gopher that was restricted to a few isolated populations in the southern Puget Sound area and on the Olympic Peninsula in Washington. The animal became...

Taczanowski's Oldfield mouse

Taczanowski's Oldfield mouse

Thomasomys taczanowskii

Taczanowski's Oldfield mouse (*Thomasomys taczanowskii*) is a species of rodent in the family Cricetidae. It is found only in Peru.

Taiga shrew

Taiga shrew

Sorex isodon

The taiga shrew (*Sorex isodon*), also known as the even-toothed shrew can achieve a body length of about 67 millimeters, with a tail of about 43 millimeters. This shrew is very similar to the long-clawed shrew. This species inhabits forested...

Taiga vole

Taiga vole

Microtus xanthognathus

The taiga vole (*Microtus xanthognathus*) is a large vole found in northwestern North America, including Alaska and northwestern Canada. The name "taiga vole" comes from its living in the boreal taiga zone. It is also sometimes called the...

Tailed tailless bat

Tailed tailless bat

Anoura caudifer

The tailed tailless bat (*Anoura caudifer*) is a species of leaf-nosed bat from South America.

Tailless fruit bat

Tailless fruit bat

Megaerops ecaudatus

The tailless fruit bat (*Megaerops ecaudatus*) is a species of fruit bat in the family Pteropodidae.

Tailless Tenrec

Tailless Tenrec

Tenrec ecaudatus

The tailless tenrec (*Tenrec ecaudatus*), also known as the common tenrec, is a species of mammal in the family Tenrecidae. It is the only member of the genus *Tenrec*. Native to Madagascar, it is also found in the Comoros, Mauritius, Réunion, and...

Taita shrew

Taita shrew

Suncus aequatorius

The Taita shrew (*Suncus aequatorius*) is an extant species of white-toothed shrew from two localities in the Taita Hills mountain range in the Taita-Taveta District of southwestern Kenya. Given the continuing decline in the quality of this...

Taiva shrew tenrec
Taiva shrew tenrec
Microgale taiva

The taiva shrew tenrec (*Microgale taiva*) is a species of mammal in the family Tenrecidae. It is endemic to Madagascar. Its natural habitats are subtropical or tropical moist lowland forest and subtropical or tropical moist montane forest. It is...

Taiwan big-eared bat
Taiwan big-eared bat
Plecotus taivanus

The Taiwan long-eared bat (*Plecotus taivanus*), also known as the Taiwan big-eared bat, is a species of vesper bat in the family Vespertilionidae. It is found only in Taiwan.

Taiwan broad-muzzled myotis
Taiwan broad-muzzled myotis
Submyotodon latirostris

The Taiwan broad-muzzled bat or Taiwan broad-muzzled myotis (*Submyotodon latirostris*) is a species of vesper bat found in Taiwan.

Taiwan field mouse
Taiwan field mouse
Apodemus semotus

The Taiwan field mouse, also called Formosan wood mouse (*Apodemus semotus*), is a species of rodent in the family Muridae. It is found only in Taiwan. The Taiwan field mouse is primarily distributed in the montane region between 1,400 and 3,000 m....

Taiwan Serow
Taiwan Serow
Capricornis swinhoei

The Taiwan serow (*Capricornis swinhoei*) is a small ruminant mammal that is native to East Asia. Like their smaller relatives, the gorals, serows are slightly larger and slower-moving, and somewhat less agile, than gorals; however, they can still...

Taiwan tube-nosed bat
Taiwan tube-nosed bat
Murina puta

The Taiwan tube-nosed bat (*Murina puta*) is a species of vesper bat in the family Vespertilionidae. It is found only in Taiwan. It is a close relative of Hutton's tube-nosed bat, and might even be the same species.

Taiwan vole
Taiwan vole
Microtus kikuchii

The Taiwan vole (*Microtus kikuchii*, now *Alexandromys kikuchii*) is a species of rodent in the family Cricetidae. It is an endemic species of Taiwan. This species is herbivorous with a preference for the Yushan cane (*Yushania niitakayamensis*).

Taiwanese brown-toothed shrew
Taiwanese brown-toothed shrew
Episoriculus fumidus

The Taiwanese brown-toothed shrew (*Episoriculus fumidus*) is a species of shrew in

the tribe Nectogalini. It is found only in Taiwan. It prefers dense ground cover in forests and subalpine shrublands in high mountains of central Taiwan.

Taiwanese gray shrew

Taiwanese gray shrew

Crocidura tanakae

The Taiwanese gray shrew (*Crocidura tanakae*) is a species of mammal in the family Soricidae. Previously believed to be endemic to Taiwan, it is now also known to occur in Vietnam.

Taiwanese mole shrew

Taiwanese mole shrew

Anourosorex yamashinai

The Taiwanese mole shrew (*Anourosorex yamashinai*) is one of four species of red-toothed shrews in the genus *Anourosorex*. This species is endemic to Taiwan.

Takin

Takin

Budorcas taxicolor

The takin (*Budorcas taxicolor*) is a large species of ungulate found in the eastern Himalayas. It is the national animal of Bhutan. The specific name 'taxicolor' comes from Latin and roughly can be translated as 'badger-like coloration'.

Talamancan small-eared shrew

Talamancan small-eared shrew

Cryptotis gracilis

The Talamancan small-eared shrew (*Cryptotis gracilis*) is a species of mammal in the family Soricidae. It is found in Costa Rica and Panama.

Talamancan yellow-shouldered bat

Talamancan yellow-shouldered bat

Sturnira mordax

The Talamancan yellow-shouldered bat (*Sturnira mordax*) is a species of bat in the family Phyllostomidae. It is found only in Costa Rica and Panama, and there are no subspecies.

Talas tuco-tuco

Talas tuco-tuco

Ctenomys talarum

The Talas tuco-tuco (*Ctenomys talarum*) is a species of tuco-tuco endemic to eastern Argentina.

Talaud bear cuscus

Talaud bear cuscus

Ailurops melanotis

The Talaud bear cuscus (*Ailurops melanotis*) is a species of marsupial in the family Phalangeridae. It is endemic to Salibabu Island in the Talaud Islands, Indonesia. Its natural habitat is subtropical or tropical dry forests. The species is...

Talaud flying fox

Talaud flying fox

Acerodon humilis

The Talaud flying fox or Talaud fruit bat (*Acerodon humilis*) is a species of bat in the family Pteropodidae. It is endemic to the islands of Salebabu and Karekaleng in the Talaud Archipelago of Indonesia. Its natural habitat is subtropical or...

Talazac's shrew tenrec

Talazac's shrew tenrec

Nesogale talazaci

Talazac's shrew tenrec (*Nesogale talazaci*) is a species of mammal in the family Tenrecidae. It is endemic to Madagascar. Its natural habitats are subtropical or tropical moist lowland forest and subtropical or tropical moist montane forest. It...

Talpa martinorum
Talpa martinorum
Talpa martinorum

Martino's mole (*Talpa martinorum*) is a species of small blind mole in the family Talpidae found in eastern Thrace along the western Black Sea coast in south-eastern Bulgaria and north-eastern Turkey, mainly in the Strandzha mountain range. In...

Tamá small-eared shrew
Tamá small-eared shrew
Cryptotis tamensis

The Tamá small-eared shrew (*Cryptotis tamensis*) is a species of mammal in the family Soricidae. It is known from the Cordillera Oriental of Colombia and the Páramo de Tamá of western Venezuela, where it has been found primarily in cloud forest...

Tamaraw
Tamaraw
Bubalus mindorensis

The tamaraw (*Bubalus mindorensis*) is a small hoofed mammal native to the Philippines. It is the only endemic Philippine bovine. It is believed, however, to have once also thrived on the larger island of Luzon. The tamaraw was originally found all...

Tamarisk jird
Tamarisk jird
Meriones tamariscinus

The tamarisk jird (*Meriones tamariscinus*) is a species of rodent in the family Muridae. It is found in China, Mongolia, Kazakhstan, Kyrgyzstan, Russia, Turkmenistan, and Uzbekistan. It prefers habitats with grass or shrub cover. Tamarisk jirds...

Tammar Wallaby
Tammar Wallaby
Macropus eugenii

The tammar wallaby (*Notamacropus eugenii*), also known as the dama wallaby or darma wallaby, is a small macropod native to South and Western Australia. Though its geographical range has been severely reduced since European colonisation, the...

Tana River mangabey
Tana River mangabey
Cercocebus galeritus

The Tana River mangabey (*Cercocebus galeritus*) is a highly endangered species of primate in the family Cercopithecidae. Some authorities have included the taxa *agilis* and *sanjei* as subspecies of this species, while others award these full...

Tana River red colobus
Tana River red colobus
Procolobus rufomitrat

The Tana River red colobus (*Piliocolobus rufomitrat*), also called the eastern red colobus, is a highly endangered species of primate in the family Cercopithecidae. It is endemic to a narrow zone of gallery forest near the Tana River in...

Tanala tufted-tailed rat

Tanala tufted-tailed rat
Eliurus tanala

The Tanala tufted-tailed rat (*Eliurus tanala*) is a species of rodent in the family Nesomyidae. It is found only in Madagascar.

Tanezumi rat
Tanezumi rat
Rattus tanezumi

The Tanezumi rat (*Rattus tanezumi*), also known as the Asian rat or Asian house rat, is a species of rodent in the family Muridae. It is closely related to the black rat (*Rattus rattus*). It is widespread in eastern, southern and south-eastern...

Tantalus monkey
Tantalus monkey
Chlorocebus tantalus

The tantalus monkey (*Chlorocebus tantalus*) is an Old World monkey from Africa that ranges from Ghana to Sudan. It was originally described as a subspecies of the grivet (*Chlorocebus aethiops*). All species in *Chlorocebus* were formerly in the...

Tanyuromys aphantus
Tanyuromys aphantus
Tanyuromys aphantus

Tanyuromys aphantus, known as the long-tailed sigmodontomys, Harris's rice water rat, or the long-tailed rice rat, is a species of rodent in the family Cricetidae. It is known from Costa Rica, Panama, and Ecuador. In 2012, it was reassigned to...

Tanyuromys thomasei
Tanyuromys thomasei
Tanyuromys thomasei

Thomas Lee's rice water rat (*Tanyuromys thomasei*) is a species of rodent in the family Cricetidae. It is endemic to Ecuador, where it is found in middle-elevation forests on Pacific-facing Andean slopes. It is known from both intact and...

Tanzanian shaggy rat
Tanzanian shaggy rat
Dasymys sua

The Tanzanian shaggy rat (*Dasymys sua*) is a species of shaggy marsh rat endemic to eastern Tanzania, near the Uluguru Mountains.

Tanzanian shrew
Tanzanian shrew
Crocibura tanzaniana

The Tanzanian shrew (*Crocibura tanzaniana*) is a species of mammal in the family Soricidae. It is endemic to Tanzania. Its natural habitat is subtropical or tropical moist montane forests. It is threatened by habitat loss.

Tanzanian vleisig rat
Tanzanian vleisig rat
Otomys lacustris

The Tanzanian vleisig rat (*Otomys lacustris*) is a species of rodent in the family Muridae. It is found in Kenya, Malawi, and Tanzania. Its natural habitats are subtropical or tropical high-altitude grassland and swamps. It is threatened by habitat loss.

Tanzanian woolly bat
Tanzanian woolly bat
Kerivoula africana

The Tanzanian woolly bat or Dobson's painted bat (*Kerivoula africana*) is a species of vesper bat in the family Vespertilionidae. It is found only in Tanzania. Locally, it is known as the "popo" or "tunge".

Tapanuli Orangutan

Tapanuli Orangutan

Pongo tapanuliensis

The Tapanuli orangutan (*Pongo tapanuliensis*) is a species of orangutan found in the island of Sumatra in Indonesia. It is one of three known species of orangutan, alongside the Sumatran orangutan, found farther northwest on the island, and the...

Tapeti

Tapeti

Sylvilagus brasiliensis

The tapeti (*Sylvilagus brasiliensis*) is a species of cottontail rabbit. As traditionally defined, its range extends from southern Mexico to northern Argentina, but this includes several distinctive population that have since been split into...

Tapirus mesopotamicus

Tapirus mesopotamicus

Tapirus mesopotamicus

Tapirus mesopotamicus is an extinct species of tapir that lived in South America during the Pleistocene. It is considered a possible ancestor of all extant South American tapirs.

Tapirus oliverasi

Tapirus oliverasi

Tapirus oliverasi

Tapirus oliverasi is a likely invalid extinct species of tapir from South America. *T. oliverasi* was described based on dentary remains from the Early Pleistocene deposits at the Libertad Formation in Uruguay that were of larger size than that of...

Tapulao shrew-rat

Tapulao shrew-rat

Rhynchomys tapulao

The Tapulao shrew-rat (*Rhynchomys tapulao*) is a rodent in the subfamily Murinae. It was described in 2007.

Tarabul's gerbil

Tarabul's gerbil

Gerbillus tarabuli

Tarabul's gerbil (*Gerbillus tarabuli*) is a species of small rodent which is found in arid regions of north western Africa.

Tarabundí vole

Tarabundí vole

Microtus oaxacensis

The Tarabundí vole (*Microtus oaxacensis*) is a species of rodent in the family Cricetidae. It is found only in Mexico.

Tarai gray langur

Tarai gray langur

Semnopithecus hector

The Tarai gray langur (*Semnopithecus hector*) is an Old World monkey, and was formerly considered a subspecies of the northern plains gray langur. The species is listed as Near Threatened, as there are probably not many more than 10,000 mature...

Tarbagan marmot
Tarbagan marmot
Marmota sibirica

The Tarbagan marmot (*Marmota sibirica*) is a species of rodent in the family Sciuridae. It is found in China (Inner Mongolia and Heilongjiang), northern and western Mongolia, and Russia (southwest Siberia, Tuva, Transbaikalia). In the Mongolian...

Tarella shrew
Tarella shrew
Crocidura tarella

The tarella shrew (*Crocidura tarella*) is a species of mammal in the family Soricidae. It is found in Democratic Republic of the Congo and Uganda. Its natural habitats are subtropical or tropical moist lowland forest and subtropical or tropical...

Target rat
Target rat
Stochomys longicaudatus

The target rat (*Stochomys longicaudatus*) is a species of rodent in the family Muridae. It is the only species in the genus *Stochomys*. Its natural habitat is subtropical or tropical moist lowland forests.

Tarija akodont
Tarija akodont
Akodon pervalens

The Tarija akodont (*Akodon pervalens*), or Tarija grass mouse, is a species in the family Cricetidae native to Bolivia, and possibly Argentina.

Tarpan
Tarpan
Equus ferus ferus

The term tarpan (*Equus ferus ferus*) refers to free-ranging horses of the Russian steppe from the 18th to the 20th century. It is generally unknown whether those horses represented genuine wild horses, feral domestic horses or hybrids.The last...

Tarsius fuscus
Tarsius fuscus
Tarsius fuscus

The Makassar tarsier (*Tarsius fuscus*) is a species of tarsier. Its range is in Indonesia in the southwestern peninsula of the island of Sulawesi, near Makassar. At one point the taxon was downgraded to a junior synonym of the spectral tarsier...

Taruca
Taruca
Hippocamelus antisensis

The taruca (*Hippocamelus antisensis*) is a medium-sized deer species that lives in the high regions of South America. Its common name, taruca, means "deer" in both, the Quechua and Aymara languages. It is closely related to the Southern guemal.

Tasmanian Devil
Tasmanian Devil
Sarcophilus harrisii

The Tasmanian devil is a carnivorous marsupial of the family Dasyuridae. The size of a small dog, the Tasmanian devil became the largest carnivorous marsupial in the world, following the extinction of the thylacine in 1936. It is related to...

Tasmanian Pademelon
Tasmanian Pademelon
Thylogale billiardierii

The Tasmanian pademelon (*Thylogale billiardierii*), also known as the rufous-bellied pademelon or red-bellied pademelon, is the sole species of pademelon found in Tasmania, and was formerly found throughout southeastern Australia. This pademelon...

Tasmanian Pygmy Possum
Tasmanian Pygmy Possum
Cercartetus lepidus

The Tasmanian pygmy possum (*Cercartetus lepidus*) is the world's smallest possum. It was first described by Oldfield Thomas in 1888, after he identified that a museum specimen labelled as an Eastern pygmy possum in fact represented a species then...

Tate's fat-tailed mouse opossum
Tate's fat-tailed mouse opossum
Thylamys tatei

Tate's fat-tailed mouse opossum (*Thylamys tatei*) is a species of opossum in the family Didelphidae, named after American zoologist George Henry Hamilton Tate. It is found at elevations of 300 to 3,000 m along the coast of central Peru. The...

Tate's shrew rat
Tate's shrew rat
Tateomys rhinogradoides

Tate's shrew rat (*Tateomys rhinogradoides*) is a species of rodent in the family Muridae. It is found only in central Sulawesi, Indonesia, where it has been recorded on Mount Latimodjong (Mount Rantemario), Mount Tokala, and Mount Nokilalaki. The...

Tate's three-striped dasyure
Tate's three-striped dasyure
Myoictis wavicus

The Tate's three-striped dasyure (*Myoictis wavicus*) is a member of the order Dasyuromorphia. This marsupial carnivore lives in New Guinea. This species inhabits primary montane forest. Like its congeners, it is thought to be diurnal and largely...

Tate's triok
Tate's triok
Dactylopsila tatei

Tate's triok, also known as the Fergusson Island striped possum (*Dactylopsila tatei*), is a species of marsupial in the family Petauridae. It is endemic to Papua New Guinea. Its natural habitat is subtropical or tropical dry forests.

Tate's woolly mouse opossum
Tate's woolly mouse opossum
Marmosa paraguayana

Tate's woolly mouse opossum (*Marmosa paraguayana*) is an omnivorous, arboreal South American marsupial of the family Didelphidae, named after American zoologist George Henry Hamilton Tate. It is native to Atlantic coastal forests of Brazil,...

Tatra Chamois
Tatra Chamois
Rupicapra rupicapra tatra

The Tatra chamois (*Rupicapra rupicapra tatra*) is a subspecies of the chamois. It is found only in the Tatra Mountains in Slovakia and Poland.

Tatra pine vole
Tatra pine vole

Microtus tatricus

The Tatra pine vole (*Microtus tatricus*), also called the Tatra vole or Tatra ground vole, is a species of vole endemic to the Carpathian mountain range in Slovakia, Poland, Ukraine and Romania. Two subspecies have been recognised. *M. t. tatricus*...

Taurine cattle

Taurine cattle

Bos taurus taurus

Cattle (*Bos taurus*) are large domesticated bovines. They are most widespread species of the genus *Bos*. Adult females are referred to as cows and adult males are referred to as bulls. Cattle are commonly raised as livestock for meat (beef or veal,...

Taurus ground squirrel

Taurus ground squirrel

Spermophilus taurensis

The Taurus ground squirrel (*Spermophilus taurensis*) is a species of rodent in the family *Sciuridae*. It is endemic to the Taurus Mountains of Turkey. It was first identified as a distinct species to the Asia Minor ground squirrel in 2007.

Tawitawi forest rat

Tawitawi forest rat

Rattus tawitawiensis

The Tawitawi forest rat or Tawitawi Island rat (*Rattus tawitawiensis*) is a species of rodent in the family *Muridae*. It is found only in Tawi-Tawi, Philippines.

Tawny deer mouse

Tawny deer mouse

Peromyscus perfulvus

The tawny deer mouse or marsh mouse (*Peromyscus perfulvus*) is a species of rodent in the family *Cricetidae*. It is found only in Mexico.

Tawny tuco-tuco

Tawny tuco-tuco

Ctenomys fulvus

The tawny tuco-tuco (*Ctenomys fulvus*) is a species of burrowing rodent in the family *Ctenomyidae*. It is found in the desert regions of northern Chile and adjoining areas of Argentina.

Tawny-bellied cotton rat

Tawny-bellied cotton rat

Sigmodon fulviventer

The tawny-bellied cotton rat (*Sigmodon fulviventer*) is a species of rodent in the family *Cricetidae*. It is found in Mexico and in the US states of Arizona and New Mexico.

Tayra

Tayra

Eira barbara

The tayra (*Eira barbara*) is a forest-dwelling animal from the weasel family, native to the Americas. Tayras are also known as the toluco or perico ligero in Central America, motete in Honduras, irara in Brazil, san hol or viejo de monte in the...

Tehuantepec Jackrabbit

Tehuantepec Jackrabbit

Lepus flavigularis

The Tehuantepec jackrabbit (*Lepus flavigularis*) is a medium-sized hare found only

in Mexico. It exists in three separated populations and is the most endangered of the hares.

Telefomin cuscus

Telefomin cuscus

Phalanger matanim

The Telefomin cuscus (Phalanger matanim) is a critically endangered possum found on New Guinea. It is named after the Telefol ethnic group, who hunted the animal long before it was identified scientifically by the Australian zoologist Tim...

Telefomin roundleaf bat

Telefomin roundleaf bat

Hipposideros corynophyllus

The Telefomin roundleaf bat (Hipposideros corynophyllus) is a species of bat in the family Hipposideridae. It is found in West Papua (Indonesia) and Papua New Guinea.

Telford's shrew

Telford's shrew

Crocidura telfordi

Telford's shrew (Crocidura telfordi) is a species of mammal in the family Soricidae. It is endemic to Tanzania. Its natural habitat is subtropical or tropical moist montane forests. It is threatened by habitat loss.

Temchuk's bolo mouse

Temchuk's bolo mouse

Necomys temchuki

Temchuk's bolo mouse (Necomys temchuki) is a species of rodent in the family Cricetidae. It is found only in Argentina.

Temminck's flying fox

Temminck's flying fox

Pteropus temminckii

Temminck's flying fox (Pteropus temminckii) is a species of flying fox in the family Pteropodidae. It is found in Indonesia. The species was classified as "Vulnerable" in 2008 by the IUCN due to threats from habitat destruction and hunting.

Temminck's flying squirrel

Temminck's flying squirrel

Petinomys setosus

Temminck's flying squirrel (Petinomys setosus) is a species of rodent in the family Sciuridae. It is found in Indonesia, Malaysia, Myanmar, and Thailand.

Temminck's mouse

Temminck's mouse

Mus musculoides

Temminck's mouse (Mus musculoides) is a species of the genus Mus and of the subgenus Nannomys. It is found throughout West Africa, Central Africa, and East Africa. An adult generally weighs 0.006 kg.

Temminck's mysterious bat

Temminck's mysterious bat

Nycticeius aenobarbus

Temminck's mysterious bat (Nycticeius aenobarbus) is a species of bat of the family Vespertilionidae. As the name suggests, there is very little known information about this bat. The species is listed as Data Deficient on the IUCN Red List, and...

Temminck's striped mouse

Temminck's striped mouse
Hybomys trivirgatus

Temminck's striped mouse or West African hybomys (*Hybomys trivirgatus*) is a species of rodent in the family Muridae. It is found in Ivory Coast, Ghana, Guinea, Liberia, Nigeria, and Sierra Leone. Its natural habitat is subtropical or tropical...

Temminck's trident bat
Temminck's trident bat
Aselliscus tricuspidatus

Temminck's trident bat (*Aselliscus tricuspidatus*) is a species of bat in the family Hipposideridae. It is found in Indonesia, Papua New Guinea, the Solomon Islands, and Vanuatu.

Temotu flying fox
Temotu flying fox
Pteropus nitendiensis

The Temotu flying fox (*Pteropus nitendiensis*) is a species of flying fox in the family Pteropodidae. It is endemic to the Solomon Islands. It is threatened by habitat destruction due to subsistence agricultural practices, as well as natural...

Tenasserim lutung
Tenasserim lutung
Trachypithecus barbei

The Tenasserim lutung (*Trachypithecus barbei*) is a species of lutung. It is found in Myanmar and Thailand. It is named after the Tenasserim Hills. Phylogenetic evidence indicates that the Tenasserim lutung is an ancestor of the far more widespread...

Tenasserim white-bellied rat
Tenasserim white-bellied rat
Niviventer tenaster

The Tenasserim white-bellied rat (*Niviventer tenaster*) is a species of rodent in the family Muridae. It is named after the Tenasserim Hills and is found above 1,000 m in forested limestone mountainous areas. Its distribution includes India,...

Tenkile
Tenkile
Dendrolagus scottae

The tenkile (*Dendrolagus scottae*), also known as Scott's tree-kangaroo, is a species of tree-kangaroo in the family Macropodidae. It is endemic to a very small area of the Torricelli Mountains of Papua New Guinea. Its natural habitat is...

Tent-making bat
Tent-making bat
Uroderma bilobatum

The tent-making bat (*Uroderma bilobatum*) is an American leaf-nosed bat (Phyllostomidae) found in lowland forests of Central and South America. This medium-sized bat has a gray coat with a pale white stripe running down the middle of the back....

Tete veld aethomys
Tete veld aethomys
Aethomys ineptus

The Tete veld aethomys or Tete veld rat (*Aethomys ineptus*) is a species of rodent in the family Muridae. It is found in South Africa and Eswatini. Its natural habitats are temperate forests, shrubland, and grassland. The common name refers to

the...

Texas antelope squirrel

Texas antelope squirrel

Ammospermophilus interpres

The Texas antelope squirrel (*Ammospermophilus interpres*) is a species of rodent in the family *Sciuridae*. It is found in Mexico and in both Texas and New Mexico within the United States.

Texas kangaroo rat

Texas kangaroo rat

Dipodomys elator

The Texas kangaroo rat (*Dipodomys elator*) is a rodent of the family *Heteromyidae*. It is found in Texas and Oklahoma in the United States, where it often lives in association with brush species, like mesquite and lotebush, growing in areas with...

Texas mouse

Texas mouse

Peromyscus attwateri

The Texas mouse (*Peromyscus attwateri*) is a species of rodent in the family *Cricetidae*. It is found in Arkansas, Kansas, Missouri, Oklahoma, and Texas in the United States. This species is named in honor of Henry Philemon Attwater.

Texas pocket gopher

Texas pocket gopher

Geomys personatus

The Texas pocket gopher (*Geomys personatus*) is a species of rodent in the family *Geomyidae*. It is found in Tamaulipas in Mexico and in Texas in the United States.

Texas wolf

Texas wolf

Canis lupus monstrabilis

The Texas wolf (*Canis lupus monstrabilis*) is an extinct subspecies of gray wolf, distinct from the Texas red wolf (*Canis lupus rufus*), whose range once included southern and western Texas and northeastern Mexico.

Thaeler's pocket gopher

Thaeler's pocket gopher

Orthogeomys thaeleri

The Darien pocket gopher (*Heterogeomys dariensis*) is a species of rodent in the family *Geomyidae*. It occurs in Panama and Colombia. Some authors classify it in the genus *Orthogeomys*, but recent research has allowed this and its related...

Thai horseshoe bat

Thai horseshoe bat

Rhinolophus siamensis

The Thai horseshoe bat (*Rhinolophus siamensis*), sometimes called the Thai leaf-nosed bat, is a species of bat from the family *Rhinolophidae*. It is frequently listed as a subspecies of the Big-eared horseshoe bat, but this may be a result of the...

Thailand roundleaf bat

Thailand roundleaf bat

Hipposideros halophyllus

The Thailand roundleaf bat (*Hipposideros halophyllus*) is a species of bat in the family *Hipposideridae*. It is endemic to Thailand.

Thalia's shrew

Thalia's shrew

Crocidura thalia

Thalia's shrew (*Crocidura thalia*) is a species of mammal in the family Soricidae. It is endemic to Ethiopia. The mammal's natural habitats are subtropical or tropical moist montane forests, moist savanna, and subtropical or tropical...

Theobald's tomb bat
Theobald's tomb bat
Taphozous theobaldi

Theobald's tomb bat (*Taphozous theobaldi*) is a species of sac-winged bat in the family Emballonuridae. It is found in India, Indonesia, Myanmar, Thailand, and Vietnam.

Therese's shrew
Therese's shrew
Crocidura theresae

Therese's shrew (*Crocidura theresae*) is a species of mammal in the family Soricidae. It is found in Ivory Coast, Ghana, Guinea, Liberia, and Sierra Leone. Its natural habitats are subtropical or tropical moist lowland forest and moist savanna.

Theropithecus brumpti
Theropithecus brumpti
Theropithecus brumpti

Theropithecus brumpti was a large terrestrial monkey that lived in the mid to late Pliocene. It is an extinct species of papionin. This fossil primate is mostly known from skulls and mandibles found in Pliocene deposits excavated in the Shungura...

Theropithecus darti
Theropithecus darti
Theropithecus darti

Theropithecus darti is an extinct species of *Theropithecus* from the middle to late Pliocene of Africa.

Theropithecus oswaldi
Theropithecus oswaldi
Theropithecus oswaldi

Theropithecus oswaldi is an extinct species of *Theropithecus* from the early to middle Pleistocene of Kenya, Ethiopia, Tanzania, South Africa, Spain, Morocco and Algeria. It appears to have been a specialised grazer. The species went extinct in...

Thespian grass mouse
Thespian grass mouse
Akodon mimus

The thespian grass mouse or hocicudo-like akodont (*Akodon mimus*) is a species of rodent in the family Cricetidae. It is found in Bolivia and Peru.

Thick-eared bat
Thick-eared bat
Eptesicus pachyotis

The thick-eared bat (*Eptesicus pachyotis*) is a species of vesper bat native to China, India, Myanmar, Bangladesh and Thailand. Very little is known about the status and ecology of this bat, except that it may prefer to inhabit tropical moist...

Thick-spined porcupine
Thick-spined porcupine
Hystrix crassispinis

The thick-spined porcupine (*Hystrix crassispinis*) is a species of rodent in the family Hystricidae. It is endemic to the island of Borneo and found in Brunei, Indonesia, and Malaysia.

Thick-tailed pygmy jerboa

Thick-tailed pygmy jerboa

Salpingotus crassicauda

The thick-tailed pygmy jerboa (*Salpingotus crassicauda*) is a species of rodent in the family Dipodidae. It is found in China, Kazakhstan, and Mongolia. Its natural habitats are temperate grassland and temperate desert. It is threatened by...

Thick-tailed shrew

Thick-tailed shrew

Crocidura brunnea

The thick-tailed shrew (*Crocidura brunnea*) is a species of mammal in the family Soricidae. It is found on the islands of Java and Bali in Indonesia.

Thick-tailed three-toed jerboa

Thick-tailed three-toed jerboa

Stylodipus telum

The thick-tailed three-toed jerboa (*Stylodipus telum*) is a species of rodent in the family Dipodidae. It is found in China, Kazakhstan, Russia, Turkmenistan, Ukraine, and Uzbekistan. Its typical habitat is steppe, desert and mountain grassland...

Thick-thumbed myotis

Thick-thumbed myotis

Myotis rosseti

The thick-thumbed myotis (*Myotis rosseti*) is a species of vesper bat. It can be found in Cambodia, Laos, and Thailand.

Thin mouse shrew

Thin mouse shrew

Myosorex tenuis

The thin mouse shrew (*Myosorex tenuis*) is a species of mammal in the family Soricidae found in South Africa and possibly Mozambique.

Thin sand rat

Thin sand rat

Psammomys vexillaris

The thin sand rat or lesser sand rat (*Psammomys vexillaris*) is a species of rodent in the family Muridae. It has also been previously named the pale sand rat based on work published by Oldfield Thomas in 1925. It is found in Algeria, Libya, and...

Thinhorn Sheep

Thinhorn Sheep

Ovis dalli

The Thinhorn sheep (*Ovis dalli*) is a wild sheep native to North America. These sheep live in dry alpine areas where they browse grasses and sedges and acquire minerals to supplement their diet from mineral licks.

Thirteen-Lined Ground Squirrel

Thirteen-Lined Ground Squirrel

Ictidomys tridecemlineatus

The Thirteen-lined ground squirrel (*Ictidomys tridecemlineatus*) is a ground squirrel that is widely distributed over grasslands and prairies of North America. It is well known for standing upright to survey its domain, diving down into its burrow...

Thollon's red colobus

Thollon's red colobus
Procolobus tholloni

Thollon's red colobus (*Piliocolobus tholloni*), also known as the Tshuapa red colobus, is a species of red colobus monkey from the Democratic Republic of the Congo and lower Republic of the Congo. It is found south of Congo River and west of...

Thomas's big-eared brown bat
Thomas's big-eared brown bat
Histiotus laephotis

Thomas's big-eared brown bat (*Histiotus laephotis*) is a species of vesper bat found in South America.

Thomas's broad-nosed bat
Thomas's broad-nosed bat
Platyrrhinus dorsalis

Thomas's broad-nosed bat (*Platyrrhinus dorsalis*) is a species of bat in the family Phyllostomidae. It is found in Bolivia, Colombia, Ecuador, Panama, and Peru.

Thomas's bushbaby
Thomas's bushbaby
Galagoides thomasi

Thomas's bushbaby (*Galagoides thomasi*) is a species of primate in the family Galagidae. It is found in Angola, Burundi, Cameroon, Democratic Republic of the Congo, Equatorial Guinea, Gabon, Kenya, Nigeria, Rwanda, Tanzania, Uganda, and Zambia.

Thomas's Ethiopian brush-furred rat
Thomas's Ethiopian brush-furred rat
Lophuromys brunneus

Thomas's Ethiopian brush-furred rat (*Lophuromys brunneus*), also called the brown brush-furred rat or the brown brush-furred mouse, is a species of brush-furred mouse from Southern Ethiopia.

Thomas's flying squirrel
Thomas's flying squirrel
Aeromys thomasi

Thomas's flying squirrel (*Aeromys thomasi*) is a species of rodent in the family Sciuridae. It is one of two species in the genus *Aeromys*. It is found in Indonesia and Malaysia.

Thomas's fruit-eating bat
Thomas's fruit-eating bat
Dermanura watsoni

Thomas's fruit-eating bat (*Dermanura watsoni*), sometimes also popularly called Watson's fruit-eating bat, is a species of bat in the family Phyllostomidae. It is found from southern Mexico, through Central America to Colombia. Its South American...

Thomas's giant deer mouse
Thomas's giant deer mouse
Megadontomys thomasi

Thomas's giant deer mouse (*Megadontomys thomasi*) is a species of rodent in the family Cricetidae. It is found only in Mexico.

Thomas's horseshoe bat
Thomas's horseshoe bat
Rhinolophus thomasi

Thomas's horseshoe bat (*Rhinolophus thomasi*) is a species of bat in the family Rhinolophidae. It is found in China, Laos, Myanmar, Thailand, and Vietnam.
Thomas's langur
Thomas's langur
Presbytis thomasi

The Thomas's langur (*Presbytis thomasi*) is a species of primate in the family Cercopithecidae. It is endemic to northern Sumatra, Indonesia. Its natural habitat is subtropical or tropical dry forests. It is threatened by habitat loss. Its native...
Thomas's mosaic-tailed rat
Thomas's mosaic-tailed rat
Paramelomys mollis

Thomas's mosaic-tailed rat (*Paramelomys mollis*) is a species of rodent in the family Muridae. It is found in West Papua, Indonesia and Papua New Guinea.
Thomas's nectar bat
Thomas's nectar bat
Lonchophylla thomasi

Thomas's nectar bat (*Lonchophylla thomasi*) is a bat species from South and Central America. Thomas's nectar bat pollinates *marcgravia*.
Thomas's Oldfield mouse
Thomas's Oldfield mouse
Thomasomys pyrrhonotus

Thomas's Oldfield mouse (*Thomasomys pyrrhonotus*) is a species of rodent in the family Cricetidae. It is found in Ecuador and Peru.
Thomas's pika
Thomas's pika
Ochotona thomasi

Thomas's pika (*Ochotona thomasi*), also known as the Thomas-pika, is a species of small mammal in the pika family, Ochotonidae. The fur on its upper body is reddish brown in summer, and mouse grey in winter. It is a generalist herbivore...
Thomas's pine vole
Thomas's pine vole
Microtus thomasi

Thomas's pine vole (*Microtus thomasi*) is a species of rodent in the family Cricetidae. It is found in Bosnia and Herzegovina, Greece, Montenegro, North Macedonia, Serbia, and possibly Albania.
Thomas's pygmy jerboa
Thomas's pygmy jerboa
Salpingotus thomasi

Thomas's pygmy jerboa (*Salpingotus thomasi*) is a species of rodent in the family Dipodidae. It is endemic to Afghanistan. Its natural habitat is temperate desert. This species is known only from the type specimen and lately is synonymized with...
Thomas's pygmy mouse
Thomas's pygmy mouse
Mus sorella

The Thomas's pygmy mouse (*Mus sorella*) is a species of rodent in the family Muridae. It is found in Angola, Cameroon, Republic of the Congo, Democratic Republic of the Congo, Gabon, Kenya, Rwanda, South Sudan, Tanzania, and Uganda. Its natural...
Thomas's rock rat
Thomas's rock rat

Aethomys thomasi

Thomas's rock rat (*Aethomys thomasi*) is a species of rodent in the family Muridae found only in Angola. Its natural habitat is subtropical or tropical dry shrubland.

Thomas's rope squirrel
Thomas's rope squirrel
Funisciurus anerythrus

Thomas's rope squirrel or redless tree squirrel (*Funisciurus anerythrus*) is a species of rodent in the family Sciuridae. It is found in Benin, Cameroon, Central African Republic, Democratic Republic of the Congo, Equatorial Guinea, Gabon, and...

Thomas's sac-winged bat
Thomas's sac-winged bat
Balantiopteryx io

Thomas's sac-winged bat (*Balantiopteryx io*) is a species of sac-winged bat in the family Emballonuridae. It is found in Belize, Guatemala, and Mexico.

Thomas's shaggy bat
Thomas's shaggy bat
Centronycteris centralis

The Thomas's shaggy bat (*Centronycteris centralis*) is a bat species from South America. It was previously included in the shaggy bat but Simmons and Handley (1998) showed that the species were distinct.

Thomas's shrew tenrec
Thomas's shrew tenrec
Microgale thomasi

Thomas's shrew tenrec (*Microgale thomasi*) is a species of mammal in the family Tenrecidae. It is endemic to Madagascar. Its natural habitat is subtropical or tropical moist lowland forests. It is threatened by habitat loss.

Thomas's small-eared shrew
Thomas's small-eared shrew
Cryptotis thomasi

Thomas's small-eared shrew (*Cryptotis thomasi*) is a species of mammal in the family Soricidae. It is found in Colombia, Ecuador, and Peru.

Thomas's water mouse
Thomas's water mouse
Rheomys thomasi

Thomas's water mouse (*Rheomys thomasi*) is a species of rodent in the family Cricetidae found in El Salvador, Guatemala, and Mexico at altitudes of 400 to 2700 m. It lives near forest streams and is semiaquatic; its carnivorous diet includes both...

Thomas's yellow bat
Thomas's yellow bat
Rhogeessa io

Thomas's yellow bat (*Rhogeessa io*) is a species of bat from the family Vespertilionidae.

Thomas's yellow-shouldered bat
Thomas's yellow-shouldered bat
Sturnira thomasi

The Thomas's yellow-shouldered bat (*Sturnira thomasi*) is a species of bat in the family Phyllostomidae. It is endemic to Guadeloupe. It is threatened by habitat

loss.

Thomasomys ucucha

Thomasomys ucucha

Thomasomys ucucha

Thomasomys ucucha, also known as the *ucucha thomomys*, is a rodent in the genus *Thomasomys* of the family *Cricetidae*. It is known only from high altitude forest and grassland habitats in the Cordillera Oriental of Ecuador. Seven other species of...

Thomson's Gazelle

Thomson's Gazelle

Eudorcas thomsonii

The Thomson's gazelle (*Eudorcas thomsonii*) is named after Joseph Thomson, a Scottish explorer, and is sometimes referred to as a "tommie". Thomson's gazelles are endemic to eastern Africa, where the overall population of this species has...

Thongaree's disc-nosed bat

Thongaree's disc-nosed bat

Eudiscoderma thongareeae

Thongaree's disc-nosed bat (*Eudiscoderma thongareeae*) is a critically endangered species of bat found in Thailand. It is the only member of the genus *Eudiscoderma*.

Thor's hero shrew

Thor's hero shrew

Scutisorex thori

Thor's hero shrew (*Scutisorex thori*) is an extant species of shrew native to the Democratic Republic of the Congo. It and its sister species, the hero shrew (*Scutisorex somereni*), are the only mammal species known to have interlocking vertebrae.

Thornicroft's Giraffe

Thornicroft's Giraffe

Giraffa camelopardalis thornicrofti

Thornicroft's giraffe (*Giraffa camelopardalis thornicrofti*) is a subspecies of giraffe that lives only in Zambia's South Luangwa Valley. This subspecies lives only in the wild, with no captive populations. It was originally named after Harry...

Thorold's deer

Thorold's deer

Cervus albirostris

Thorold's deer (*Cervus albirostris*) is a threatened species of deer found at high altitudes in the eastern Tibetan Plateau. It is also known as the White-lipped deer for the white patches around its muzzle. This deer was first scientifically...

Three-striped dasyure

Three-striped dasyure

Myoictis melas

The three-striped dasyure (*Myoictis melas*) is a member of the order *Dasyuromorphia*. This marsupial carnivore lives in West Papua, Indonesia and Papua New Guinea.

Three-striped ground squirrel

Three-striped ground squirrel

Lariscus insignis

The three-striped ground squirrel (*Lariscus insignis*) is a species of rodent in the family *Sciuridae*. It is found in Indonesia, Malaysia, and Thailand.

Three-striped night monkey

Three-striped night monkey

Aotus trivirgatus

The three-striped night monkey (*Aotus trivirgatus*), also known as northern night monkey or northern owl monkey, is one of several species of owl monkeys currently recognised. It is found in Venezuela and north-central Brazil. Until 1983, all the...

Thrichomys laurentius

Thrichomys laurentius

Thrichomys laurentius

Thrichomys laurentius, the Sao Lourenço punaré, is a South American caviomorph rodent of the spiny rat family. It was formerly considered a subspecies of *T. apereoides*. It is found in the northern Caatinga ecoregion of northeastern Brazil, a...

Thumbless bat

Thumbless bat

Furipterus horrens

The thumbless bat (*Furipterus horrens*) is a species of insectivorous bat in the family Furipteridae, in the monotypic genus *Furipterus*. It is found in Costa Rica, Brazil, Venezuela; Colombia; Ecuador; Suriname; French Guiana; Guyana; Panama;...

Tian Shan wapiti

Tian Shan wapiti

Cervus canadensis songaricus

The Tian Shan wapiti or Tian Shan maral (*Cervus canadensis songaricus*), is a subspecies of *C. canadensis*. It is also called the Tian Shan elk in North American English.

Tiang

Tiang

Damaliscus lunatus tiang

The tiang (*Damaliscus lunatus tiang*) is a subspecies of the topi, an African antelope. Depending on which topi populations you want to call tiang, they may be found in southern Chad, the northern Central African Republic, and southwestern Sudan...

Tibetan Antelope

Tibetan Antelope

Pantholops hodgsonii

The Tibetan antelope or chiru (*Pantholops hodgsonii*) is a medium-sized bovid native to the northeastern Tibetan plateau. Fewer than 150,000 mature individuals are left in the wild, but the population is currently thought to be increasing. In the...

Tibetan blue bear

Tibetan blue bear

Ursus arctos pruinosus

The Tibetan bear or Tibetan blue bear (*Ursus arctos pruinosus*) is a subspecies of the brown bear (*Ursus arctos*) found in the eastern Tibetan Plateau. It is also known as the Himalayan blue bear, Himalayan snow bear, Tibetan brown bear, or the...

Tibetan dwarf hamster

Tibetan dwarf hamster

Cricetulus alticola

The Tibetan dwarf hamster (*Cricetulus alticola*) is a species of rodent in the family Cricetidae. It is found not only in Tibet and China, but also in India and Nepal in mountainous regions at altitudes of up to about 5,200 m (17,100 ft).

Tibetan Fox

Tibetan Fox
Vulpes ferrilata

The Tibetan fox (*Vulpes ferrilata*) is a species of true fox native to Asia. It lives a solitary life and is widespread in the Tibetan Plateau's steppes and semi-deserts.

Tibetan Macaque
Tibetan Macaque
Macaca thibetana

The Tibetan macaque (*Macaca thibetana*) is a macaque species found in China. It is the largest species of macaque and one of the largest monkeys found in Asia. Only the Proboscis monkey and the larger species of Gray langur are bigger in size...

Tibetan red deer
Tibetan red deer
Cervus canadensis wallichi

The Tibetan red deer (*Cervus canadensis wallichi*) also known as shou, is a subspecies of elk (wapiti) native to the southern Tibetan highlands and Bhutan. Once believed to be near-extinct, its population has increased to over 8,300, the majority...

Tibetan shrew
Tibetan shrew
Sorex thibetanus

The Tibetan shrew (*Sorex thibetanus*) is a species of mammal in the family Soricidae. It is found in western China.

Tibetan wolf
Tibetan wolf
Canis lupus chanco

The Himalayan wolf (*Canis lupus chanco*) is a canine of debated taxonomy. It is distinguished by its genetic markers, with mitochondrial DNA indicating that it is genetically basal to the Holarctic grey wolf, genetically the same wolf as the...

Tickell's bat
Tickell's bat
Hesperoptenus tickelli

Tickell's bat (*Hesperoptenus tickelli*) is a species of vesper bat. It can be found in the following countries: Bangladesh Bhutan, Cambodia, possibly China, India, Myanmar, Nepal, Sri Lanka and Thailand.

Tien Shan birch mouse
Tien Shan birch mouse
Sicista tianshanica

The Tien Shan birch mouse (*Sicista tianshanica*) is a species of rodent in the family Sminthidae. It is found in China, Kazakhstan and Kyrgyzstan.

Tien Shan dhole
Tien Shan dhole
Cuon alpinus hesperius

The Tian Shan dhole (*Cuon alpinus hesperius*), also known as the Siberian dhole, Western Asiatic dhole, or northern dhole is a subspecies of dhole native to the Altai and Tian Shan mountain ranges, and possibly Pamir and Kashmir.

Tien Shan red-backed vole
Tien Shan red-backed vole
Myodes centralis

The Tien Shan red-backed vole (*Myodes centralis*) is a species of rodent in the

family Cricetidae. It is found in China and Kyrgyzstan.

Tien Shan shrew

Tien Shan shrew

Sorex asper

The Tien Shan shrew (*Sorex asper*) is a species of mammal in the family Soricidae. It is found in China and Kazakhstan.

Tien Shan vole

Tien Shan vole

Microtus ilaeus

The Tien Shan vole (*Microtus ilaeus*) is a species of rodent in the family Cricetidae. It is found in Afghanistan, Kazakhstan, Kyrgyzstan, Tajikistan, and Turkmenistan.

Tiger

Tiger

Panthera tigris

The tiger (*Panthera tigris*) is the largest living cat species and a member of the genus *Panthera*. It is among the most recognizable and popular of the world's charismatic megafauna. It featured prominently in the ancient mythology and folklore of...

Tiger Quoll

Tiger Quoll

Dasyurus maculatus

The Tiger quoll (*Dasyurus maculatus*) is a carnivorous marsupial native to Australia. It is the world's second-largest extant carnivorous marsupial, behind the Tasmanian devil. The Tiger quoll is one the most violent animals, found in the...

Tilda's yellow-shouldered bat

Tilda's yellow-shouldered bat

Sturnira tildae

Tilda's yellow-shouldered bat (*Sturnira tildae*) is a bat species from South America. It is found in Bolivia, Brazil, Colombia, Ecuador, French Guiana, Guyana, Peru, Suriname, Venezuela, and Trinidad and Tobago.

Timor rat

Timor rat

Rattus timorensis

The Timor rat (*Rattus timorensis*) is a species of rodent in the family Muridae found in Indonesian West Timor, where it lives in the teak forests. It is known from a specimen collected near the summit of Mount Mutis.

Timor roundleaf bat

Timor roundleaf bat

Hipposideros crumeniferus

The Timor roundleaf bat (*Hipposideros crumeniferus*) is a species of bat in the family Hipposideridae. It is endemic to Indonesia.

Timor shrew

Timor shrew

Crocidura tenuis

The Timor shrew or thin shrew (*Crocidura tenuis*) is a species of mammal in the family Soricidae. It is endemic to Timor. It is threatened by habitat loss.

Timor warty pig

Timor warty pig

Sus celebensis timoriensis

The Timor warty pig (*Sus celebensis timoriensis*) is a subspecies of *Sus celebensis*, or Celebes warty pig. Though described as a separate species, it is a feral form of the Celebes warty pig found in the Lesser Sunda Islands.

Timorese horseshoe bat

Timorese horseshoe bat

Rhinolophus montanus

The Timorese horseshoe bat (*Rhinolophus montanus*) is a species of bat native to Timor-Leste.

Tinfields rock rat

Tinfields rock rat

Aethomys stannarius

Tinfields rock rat (*Aethomys stannarius*) is a species of rodent in the family Muridae found in Cameroon and Nigeria. Its natural habitats are dry savanna, subtropical or tropical dry shrubland, and subtropical or tropical dry lowland grassland.

Tiny fat mouse

Tiny fat mouse

Steatomys parvus

The tiny fat mouse (*Steatomys parvus*) is a species of rodent in the family Nesomyidae. It is found in Angola, Botswana, Ethiopia, Kenya, Mozambique, Namibia, South Sudan, Tanzania, Uganda, Zambia, and Zimbabwe. Its natural habitats are subtropical...

Tiny pipistrelle

Tiny pipistrelle

Pipistrellus nanulus

The tiny pipistrelle (*Pipistrellus nanulus*) is a species of vesper bat. It can be found in Benin, Burkina Faso, Cameroon, Central African Republic, Democratic Republic of the Congo, Ivory Coast, Equatorial Guinea, Gabon, Ghana, Guinea, Kenya,...

Tiny serotine

Tiny serotine

Neoromicia guineensis

The tiny serotine (*Neoromicia guineensis*) is a species of vesper bat.

Tiny tuco-tuco

Tiny tuco-tuco

Ctenomys minutus

The tiny tuco-tuco (*Ctenomys minutus*) is a tuco-tuco species from South America. It is found in Brazil and Bolivia.

Tiny yellow bat

Tiny yellow bat

Rhogeessa minutilla

The tiny yellow bat (*Rhogeessa minutilla*) is a species of vesper bat. It is found in Colombia and Venezuela.

Tipton kangaroo rat

Tipton kangaroo rat

Dipodomys nitratoides nitratoides

The Tipton kangaroo rat (*Dipodomys nitratoides nitratoides*), is a subspecies of the San Joaquin kangaroo rat, a rodent in the family Heteromyidae.

Titania's woolly bat

Titania's woolly bat

Kerivoula titania

Titania's woolly bat (*Kerivoula titania*) is a species of bat found in Southeast Asia.

Toad mouse

Toad mouse

Mus bufo

The toad mouse (*Mus bufo*) is a species of rodent in the family Muridae. It is found in Burundi, Democratic Republic of the Congo, Rwanda, and Uganda. Its natural habitats are subtropical or tropical moist montane forests and arable land.

Tocantins vesper mouse

Tocantins vesper mouse

Calomys tocantinsi

The Tocantins vesper mouse (*Calomys tocantinsi*) is a South American rodent species of the family Cricetidae. It is found in Brazil. Its karyotype has $2n = 46$ and $FNa = 66$.

Togian Babirusa

Togian Babirusa

Babyrousa togeanensis

The Togian babirusa (*Babyrousa togeanensis*) is the largest species of babirusa. It is found in Indonesia and was considered a subspecies of the Buru babirusa (*Babyrousa babyrussa*) until 2002.

Togo mouse

Togo mouse

Leimacomys buettneri

The Togo mouse (*Leimacomys buettneri*), also known as Büttner's African forest mouse or the groove-toothed forest mouse, is a unique muroid rodent known from only two specimens taken from near the type locality of Bismarckburg, near Yege, Togo,...

Tokunoshima spiny rat

Tokunoshima spiny rat

Tokudaia tokunoshimensis

The Tokunoshima spiny rat (*Tokudaia tokunoshimensis*) is a rodent found only on the island of Tokunoshima in the Ryukyu Islands of Japan. Due to its small habitat, it is considered endangered. It is commonly found in the secondary and primary...

Tolai hare

Tolai hare

Lepus tolai

The tolai hare (*Lepus tolai*) is a species of hare native to Central Asia, Mongolia, and Northern and Central China. It inhabits semi-desert, steppes, rocky habitats, and forest meadows. It is relatively common, even in areas with heavy human...

Toltec cotton rat

Toltec cotton rat

Sigmodon toltecus

The Toltec cotton rat (*Sigmodon toltecus*) is a rodent species in the family Cricetidae. It is found in eastern Mexico from the Rio Grande to the Yucatán Peninsula, as well as in Belize and northern Guatemala. It prefers moist grassland habitat....

Toltec fruit-eating bat

Toltec fruit-eating bat

Dermanura tolteca

The Toltec fruit-eating bat (*Dermanura tolteca*) is a species of bat in the family Phyllostomidae. It is also sometimes called the "lowland fruit eating bat."

Tome's spiny rat

Tome's spiny rat

Proechimys semispinosus

Tome's spiny rat (*Proechimys semispinosus*), also known as Tomes' spiny rat or the Central American spiny rat, is a species of spiny rat distributed from Honduras to Ecuador. The IUCN has assessed its conservation status as being of "least concern".

Tomes's sword-nosed bat

Tomes's sword-nosed bat

Lonchorhina aurita

Tomes's sword-nosed bat (*Lonchorhina aurita*), also known as the common sword-nosed bat is a bat species from South and Central America. It is also found in the Bahamas, as is known only from one specimen collected on the island of New...

Tondano rat

Tondano rat

Taeromys taerae

The Tondano rat (*Taeromys taerae*) is a species of rodent in the family Muridae. It is found only in Indonesia. *Taeromys taerae* is found in the highlands of northeastern Sulawesi. The rat is a morphological relative of *T. hamatus*, which is found...

Tonkean Macaque

Tonkean Macaque

Macaca tonkeana

The Tonkean black macaque or Tonkean macaque (*Macaca tonkeana*) is a species of primate in the family Cercopithecidae. It is endemic to central Sulawesi and the nearby Togian Islands in Indonesia. It is threatened by habitat loss. Widespread...

Tonkin Snub-Nosed Monkey

Tonkin Snub-Nosed Monkey

Rhinopithecus avunculus

The Tonkin snub-nosed monkey or Dollman's snub-nosed monkey (*Rhinopithecus avunculus*) is a slender-bodied arboreal Old World monkey, endemic to northern Vietnam. It has black and white fur, a pink nose and lips, and blue patches around the eyes....

Tonkin weasel

Tonkin weasel

Mustela tonkinensis

The Tonkin weasel (*Mustela tonkinensis*) is a species of weasel described by Björkegren in 1941. It is known only from a singular specimen collected from an undisclosed location in Northern Vietnam. Originally believed to be a form of either the...

Topi

Topi

Damaliscus lunatus jimela

Damaliscus lunatus jimela is a subspecies of topi, and is usually just called a topi. It is a highly social and fast type of antelope found in the savannas, semi-deserts, and floodplains of sub-Saharan Africa.

Toque Macaque

Toque Macaque

Macaca sinica

Tora hartebeest
Tora hartebeest
Alcelaphus buselaphus tora

The tora hartebeest (*Alcelaphus buselaphus tora*) is an extremely endangered antelope, native to Eritrea and Ethiopia. It has possibly been extirpated from Sudan. One of the most critically endangered large mammals in the world, it is threatened...

Townsend's Big-Eared Bat
Townsend's Big-Eared Bat
Corynorhinus townsendii

The Townsend's big-eared bat (*Corynorhinus townsendii*) is a species of vesper bat found in North America.

Townsend's Chipmunk
Townsend's Chipmunk
Neotamias townsendii

Townsend's chipmunk (*Neotamias townsendii*) is a species of rodent in the squirrel family, *Sciuridae*. It lives in the forests of the Pacific Northwest of North America, from extreme southwestern British Columbia through western Washington and...

Townsend's ground squirrel
Townsend's ground squirrel
Urocitellus townsendii

Townsend's ground squirrel (*Urocitellus townsendii*) is a species of rodent in the family *Sciuridae*. It is found in high desert shrublands in several areas of the United States.

Townsend's mole
Townsend's mole
Scapanus townsendii

The Townsend's mole (*Scapanus townsendii*) is a fossorial mammal in the family *Talpidae*, and is the largest North American mole. It was named after the American naturalist John Kirk Townsend. The name was selected at the request of Thomas Nuttall...

Townsend's pocket gopher
Townsend's pocket gopher
Thomomys townsendii

Townsend's pocket gopher (*Thomomys townsendii*) is a species of pocket gopher endemic to the northwestern United States.

Townsend's vole
Townsend's vole
Microtus townsendii

Townsend's vole (*Microtus townsendii*) is a species of rodent in the family *Cricetidae*, the sister species of *M. canicaudus*. It is found in temperate grasslands of British Columbia in Canada and in the states of Washington and Oregon in the...

Tranier's tateril
Tranier's tateril
Taterillus tranieri

Tranier's tateril (*Taterillus tranieri*) is a species of rodent found in Mali and Mauritania. Its natural habitats are dry savanna, subtropical or tropical dry shrubland, and arable land.
Transandinomys bolivaris

Transandinomys bolivaris
Transandinomys bolivaris

Transandinomys bolivaris, also known as the long-whiskered rice rat, is a rodent in the family Cricetidae. It is found in humid forest from northeastern Honduras to western Ecuador, up to 1,800 m (5,900 ft) above sea level. Since it was first...

Transandinomys talamancae
Transandinomys talamancae
Transandinomys talamancae

Transandinomys talamancae is a rodent in the family Cricetidae that occurs from Costa Rica to southwestern Ecuador and northern Venezuela. Its habitat consists of lowland forests up to 1,500 m (5,000 ft) above sea level. With a body mass of 38 to...

Transbaikal zokor
Transbaikal zokor
Myospalax psilurus

The *Transbaikal zokor* (*Myospalax psilurus*) is a species of rodent in the family Spalacidae. It is found in China, Mongolia, and Russia.

Transcaspian vole
Transcaspian vole
Microtus transcaspicus

The *Transcaspian vole* (*Microtus transcaspicus*) is a species of rodent in the family Cricetidae. It is found in Afghanistan, Iran, and Turkmenistan.

Transcaucasian mole vole
Transcaucasian mole vole
Ellobius lutescens

The *Transcaucasian mole vole* (*Ellobius lutescens*) is a species of rodent in the family Cricetidae. It is found in Armenia, Azerbaijan, Georgia, Iran, and Turkey.

Transcaucasian water shrew
Transcaucasian water shrew
Neomys teres

The *Transcaucasian water shrew* (*Neomys teres*) is a species of mammal in the family Soricidae. It is found in Armenia, Azerbaijan, Georgia, and possibly Iran and Turkey.

Transitional colilargo
Transitional colilargo
Microakodontomys transitorius

The *transitional colilargo* (*Microakodontomys transitorius*) also known as the intermediate lesser grass mouse, is a species of rodent in the tribe Oryzomyini known only from two individuals found in the Federal District of Brazil.

Although...
Transvolcanic deer mouse
Transvolcanic deer mouse
Peromyscus hylocetes

The *transvolcanic deer mouse* (*Peromyscus hylocetes*) is a species of rodent in the genus *Peromyscus* of the family Cricetidae found only in Mexico.

Travancore flying squirrel
Travancore flying squirrel
Petinomys fuscocapillus

Travancore flying squirrel (*Petinomys fuscocapillus*), also known as the small flying squirrel, is a flying squirrel found in South India and Sri Lanka.

Travancore flying squirrels were thought to be extinct but were rediscovered in 1989 after a...

Tree bat

Tree bat

Ardops nichollsi

The tree bat (*Ardops nichollsi*) is a species of bat in the family Phyllostomidae and the only species in the genus *Ardops*. It is found in Dominica, Guadeloupe, Martinique, Montserrat, Netherlands Antilles, Saint Lucia, Saba and Saint Vincent and...

Tree Pangolin

Tree Pangolin

Phataginus tricuspis

The Tree pangolin (*Phataginus tricuspis*) is a species of pangolins ("scaly anteaters"), and is native to equatorial Africa. It is the most common of the African forest pangolins.

Trefoil horseshoe bat

Trefoil horseshoe bat

Rhinolophus trifolius

The trefoil horseshoe bat (*Rhinolophus trifolius*) is a species of bat in the family Rhinolophidae. It is found in Brunei, India, Indonesia, Malaysia, Myanmar, Singapore, and Thailand. In Borneo locally common up to 1,800m, including mangroves.

Trefoil-toothed giant rat

Trefoil-toothed giant rat

Lenomys meyeri

The trefoil-toothed giant rat (*Lenomys meyeri*) is a species of rodent in the family Muridae. It is found only in Sulawesi, Indonesia, where it is located throughout the island.

Tres Marias Island mouse

Tres Marias Island mouse

Peromyscus madrensis

The Tres Marias island mouse or Tres Marias deer mouse (*Peromyscus madrensis*) is a species of rodent in the family Cricetidae. It is found only on the Islas Marias off the west coast of Mexico (part of the state of Nayarit). When last assessed,...

Tres Marias rabbit

Tres Marias rabbit

Sylvilagus graysoni

The Tres Marias cottontail or Tres Marias rabbit (*Sylvilagus graysoni*) is a species of mammal in the family Leporidae.

Tres Marias Raccoon

Tres Marias Raccoon

Procyon lotor insularis

The Tres Marias raccoon (*Procyon lotor insularis*) is sometimes historically considered a separate species, however, now it is regarded to be a subspecies of the Common raccoon, introduced to the Islas Marias in the recent past. It is slightly...

Trevor's free-tailed bat

Trevor's free-tailed bat

Mops trevori

The Trevor's free-tailed bat (*Mops trevori*) is a species of bat in the family Molossidae. It is found in Central African Republic, Democratic Republic of the

Congo, Ivory Coast, Ghana, Guinea, Nigeria, South Sudan, and Uganda. Its natural...
Triaenops goodmani
Triaenops goodmani
Triaenops goodmani

Triaenops goodmani is an extinct bat from Madagascar in the genus *Triaenops*. It is known from three lower jaws collected in a cave at Anjohibe in 1996, and described as a new species in 2007. The material is at most 10,000 years old. A bat...

Tricolored Bat
Tricolored Bat
Perimyotis subflavus

The Tricolored bat (*Perimyotis subflavus*) is a species of microbat native to eastern North America. Its common name "tricolored bat" derives from the coloration of the hairs on its back, which have three distinct color bands. It is the smallest...

Tricolored big-eared bat
Tricolored big-eared bat
Glyphonictis sylvestris

The tricolored big-eared bat (*Glyphonictis sylvestris*) is a bat species from South and Central America.

Trident bat
Trident bat
Asellia tridens

The trident bat or trident leaf-nosed bat (*Asellia tridens*) is a species of bat in the family Hipposideridae. It is widely distributed in the Middle East, South and Central Asia, and North, East, and Central Africa. Its natural habitats are...

Trigonictis macrodon
Trigonictis macrodon
Trigonictis macrodon

Trigonictis macrodon is an extinct species of mammal related to the living grison (genus *Galictis*). It lived in North America during the Pliocene to Pleistocene epochs, from ~4.1–1.6 Ma. (AEO), existing for approximately 2.5 million years.

Fossil...
Trinidad dog-like bat
Trinidad dog-like bat
Peropteryx trinitatis

The Trinidad dog-like bat (*Peropteryx trinitatis*) is a species of bat from the family Emballonuridae. It is native to Aruba, French Guiana, Grenada, Trinidad and Tobago, and Venezuela. The bat is considered to be rare everywhere in its...

Trinidad spiny pocket mouse
Trinidad spiny pocket mouse
Heteromys anomalus

The Trinidad spiny pocket mouse (*Heteromys anomalus*) is a species of rodent in the family Heteromyidae. It is found in Colombia, Trinidad and Tobago, and Venezuela.

Trinidad spiny rat
Trinidad spiny rat
Proechimys trinitatis

The Trinidad spiny rat (*Proechimys trinitatis*) is a species of rodent in the family Echimyidae. It is found in Trinidad and Tobago and northern Venezuela.

Trinidadian funnel-eared bat
Trinidadian funnel-eared bat
Natalus tumidirostris

The Trinidadian funnel-eared bat (*Natalus tumidirostris*) is a species of bat in the family Natalidae. It is endemic to Colombia, Venezuela, Guyana, Suriname, French Guiana, Trinidad and Tobago and Netherlands Antilles.

Tristram's jird

Tristram's jird

Meriones tristrami

Meriones tristrami, known as Tristram's jird, is a species of rodent that lives in the Middle East. It is named after the Reverend Henry Baker Tristram who collected the first specimens. It is up to 155 mm (6.1 in) long, and lives in burrows in...

Tropical antechinus

Tropical antechinus

Antechinus adustus

The tropical antechinus (*Antechinus adustus*), also known as the rusty antechinus, is a species of small marsupial carnivore, particularly closely related to the brown antechinus (*Antechinus stuartii*).It is found in a small area of tropical vine...

Tropical big-eared brown bat

Tropical big-eared brown bat

Histiotus velatus

The tropical big-eared brown bat (*Histiotus velatus*), is a bat species from South America. It is found in Bolivia, Peru, Argentina, Brazil and Paraguay

Tropical bottlenose whale

Tropical bottlenose whale

Indopacetus pacificus

The tropical bottlenose whale (*Indopacetus pacificus*), also known as the Indo-Pacific beaked whale or Longman's beaked whale, was considered to be the world's rarest cetacean until recently, but the spade-toothed whale now holds that position....

Tropical ground squirrel

Tropical ground squirrel

Notocitellus adocetus

The tropical ground squirrel (*Notocitellus adocetus*) is a species of rodent in the family Sciuridae. At one time the species was originally described as *Spermophilus adocetus*, but the genus *Spermophilus* was revised and subdivided in 2009 and it...

Tropical pocket gopher

Tropical pocket gopher

Geomys tropicalis

The tropical pocket gopher (*Geomys tropicalis*) is a species of rodent in the family Geomyidae. It is endemic to Mexico.Its natural habitat is hot deserts. It is threatened by habitat loss.

Tropical small-eared shrew

Tropical small-eared shrew

Cryptotis tropicalis

The tropical small-eared shrew (*Cryptotis tropicalis*) is a very small mammal of the family Soricidae. The species is found in the eastern highlands of Chiapas, Mexico, and parts of Belize and Guatemala. Until recently, it was considered a...

Tropical vlei rat

Tropical vlei rat

Otomys tropicalis

The tropical vlei rat (*Otomys tropicalis*) is a species of rodent in the family

Muridae. It is found in Burundi, Democratic Republic of the Congo, Kenya, Rwanda, South Sudan, and Uganda. Its natural habitats are subtropical or tropical seasonally...

Troughton's sheath-tailed bat
Troughton's sheath-tailed bat
Taphozous troughtoni

Troughton's sheath-tailed bat (*Taphozous troughtoni*) is a species of sheath-tailed bat in the family Emballonuridae, found only in Australia.

Trowbridge's shrew
Trowbridge's shrew
Sorex trowbridgii

Trowbridge's shrew (*Sorex trowbridgii*) is a species of mammal in the family Soricidae. It is found in southern British Columbia in Canada and in Washington, Oregon, and California in the United States.

True's beaked whale
True's beaked whale
Mesoplodon mirus

True's beaked whale (*Mesoplodon mirus*) is a medium-sized whale in the genus *Mesoplodon*. It is native to the northern Atlantic Ocean. The common name is in reference to Frederick W. True, a curator at the United States National Museum (now the...

True's shrew mole
True's shrew mole
Dymecodon pilirostris

True's shrew mole (*Dymecodon pilirostris*) is a species of mammal in the family Talpidae. It is endemic to Japan (Honshu, Shikoku and Kyushu) and is a common species above 1000 meters in grassland, shrubland and forest. It is the only species in...

True's vole
True's vole
Hyperacrius fertilis

True's vole (*Hyperacrius fertilis*) is a species of rodent in the family Cricetidae. It is found in India and Pakistan.

Trujillo
Trujillo
Scotophilus trujilloi

Trujillo or Truxillo may refer to:

Truong Son muntjac
Truong Son muntjac
Muntiacus truongsongensis

The Truong Son muntjac or Annamite muntjac (*Muntiacus truongsongensis*) is a species of muntjac deer. It is one of the smallest muntjac species, at about 15 kg (33 lb), half the size of the Indian muntjac (or common muntjac). It was discovered in...

Tschudi's slender opossum
Tschudi's slender opossum
Marmosops impavidus

Tschudi's slender opossum (*Marmosops impavidus*) is an opossum species from South America, named after Swiss naturalist Johann Jakob von Tschudi. It is found in Bolivia, Brazil, Colombia, Ecuador, Paraguay, Peru and Venezuela.

Tschudi's yellow-shouldered bat
Tschudi's yellow-shouldered bat

Sturnira oporaphilum

Tschudi's yellow-shouldered bat (*Sturnira oporaphilum*) is a species of leaf-nosed bat indigenous to Argentina, Ecuador, and Peru, with its range also encompassing Bolivia.

Tsing-ling pika

Tsing-ling pika

Ochotona huangensis

The Tsing-ling pika (*Ochotona syrx*) is a species of pika endemic to the mountains in Central China. It inhabits mountainous forests and shrublands. It is a poorly known species. It is a rarely found, one of the six pika species endemic to...

Tsingy tufted-tailed rat

Tsingy tufted-tailed rat

Eliurus antsingy

The Tsingy tufted-tailed rat (*Eliurus antsingy*) is a species of rodent in the family Nesomyidae. It is endemic to western and northern Madagascar, and has been observed mainly in dry forest. As a result of deforestation, the habitat of E....

Tsolov's mouse-like hamster

Tsolov's mouse-like hamster

Calomyscus tsolovi

Tsolov's mouse-like hamster or Syrian calomyscus (*Calomyscus tsolovi*) is a species of rodent in the family Calomyscidae. It is endemic to Syria.

Tube-lipped nectar bat

Tube-lipped nectar bat

Anoura fistulata

The tube-lipped nectar bat (*Anoura fistulata*) is a bat from Ecuador. It was described in 2005. It has a remarkably long tongue, which it uses to drink nectar. It additionally consumes pollen and insects.

Tucuman tuco-tuco

Tucuman tuco-tuco

Ctenomys tucumanus

The Tucuman tuco-tuco (*Ctenomys tucumanus*) is a species of rodent in the family Ctenomyidae. It is endemic to Argentina.

Tucuxi

Tucuxi

Sotalia fluviatilis

The tucuxi (*Sotalia fluviatilis*) is a freshwater dolphin found in the rivers of the Amazon basin. Despite being found in geographic locations similar to those of 'true' river dolphins such as the boto, the tucuxi is not closely related to them...

Tuft-tailed spiny tree-rat

Tuft-tailed spiny tree-rat

Lonchothrix emiliae

The tuft-tailed spiny tree rat (*Lonchothrix emiliae*) is a spiny rat species from South America. It is known from Brazil south of the Amazon River, where it has been found in grassland and gallery forest. It is the only species in the genus...

Tufted Deer

Tufted Deer

Elaphodus cephalophus

The tufted deer (*Elaphodus cephalophus*) is a small species of deer characterized

by a prominent tuft of black hair on its forehead and fang-like canines for the males. It is a close relative of the muntjac, living somewhat further north over a...

Tufted Gray Langur
Tufted Gray Langur
Semnopithecus priam

The tufted gray langur (*Semnopithecus priam*), also known as Madras gray langur, and Coromandel sacred langur, is an Old World monkey, one of the species of langurs. This, like other gray langurs, is mainly a leaf-eating monkey. It is found in...

Tufted ground squirrel
Tufted ground squirrel
Rheithrosciurus macrotis

The tufted ground squirrel or groove-toothed squirrel (*Rheithrosciurus macrotis*) is a species of rodent in the family Sciuridae. It is the sole species in the genus *Rheithrosciurus*. It is found only on the island of Borneo. Confirmed elements of...

Tufted pygmy squirrel
Tufted pygmy squirrel
Exilisciurus whiteheadi

The tufted pygmy squirrel (*Exilisciurus whiteheadi*) is a species of rodent in the family Sciuridae. It is endemic to highland forest in Borneo. The common name of this tiny squirrel refers to its distinctive ear-tufts. Its diet consists mainly of...

Tufted-tailed spiny tree-rat
Tufted-tailed spiny tree-rat
Mesomys occultus

The tufted-tailed spiny tree-rat or furtive spiny tree-rat (*Mesomys occultus*) is a spiny rat species from South America. It is known from Amazonas, northwestern Brazil, where it is found in tropical rainforest. The species is arboreal.

Tule elk
Tule elk
Cervus canadensis nannodes

The tule elk (*Cervus canadensis nannodes*) is a subspecies of elk found only in California, ranging from the grasslands and marshlands of the Central Valley to the grassy hills on the coast. The subspecies name derives from the tule, a species of...

Tule shrew
Tule shrew
Sorex ornatus juncensis

The tule shrew (*Sorex ornatus juncensis*) is a possibly extinct subspecies of the ornate shrew (*Sorex ornatus*). It was confined to the Baja California peninsula in Mexico.

Tullberg's soft-furred mouse
Tullberg's soft-furred mouse
Praomys tullbergi

Tullberg's soft-furred mouse or Tullberg's praomys (*Praomys tullbergi*) is a species of rodent in the family Muridae in Africa. It is found in Angola, Benin, Cameroon, Republic of the Congo, Democratic Republic of the Congo, Ivory Coast,...

Tumbala climbing rat
Tumbala climbing rat
Tylomys tumbalensis

The Tumbala climbing rat (*Tylomys tumbalensis*) is a species of rodent in the family Cricetidae. It is found in Mexico, where it is known only from one locality in Tumbalá, Chiapas. The species is threatened by deforestation.

Tundra shrew

Tundra shrew

Sorex tundrensis

The tundra shrew (*Sorex tundrensis*) is a small shrew found in Alaska, the northern Yukon Territory, the MacKenzie Delta region of the Northwest Territories, extreme northwestern British Columbia and eastern Russia. At one time, this animal was...

Tundra vole

Tundra vole

Microtus oeconomus

The tundra vole (*Microtus oeconomus*) or root vole is a medium-sized vole found in Northern and Central Europe, Asia, and northwestern North America, including Alaska and northwestern Canada. In the western part of the Netherlands, the tundra...

Tundra wolf

Tundra wolf

Canis lupus albus

The tundra wolf (*Canis lupus albus*), also known as the Turukhan wolf, is a subspecies of grey wolf native to Eurasia's tundra and forest-tundra zones from Finland to the Kamchatka Peninsula. It was first described in 1792 by Robert Kerr, who...

Tupaia miocenica

Tupaia miocenica

Tupaia miocenica

Tupaia miocenica is a fossil treeshrew from the Miocene of Thailand. Known only from a single tooth, an upper first or second molar, it is among the few known fossil treeshrews. With a length of 3.57 mm, the tooth is large for a treeshrew. At the...

Turbo shrew

Turbo shrew

Crocidura turba

The turbo shrew (*Crocidura turba*) is a species of mammal in the family Soricidae. It is found in Angola, Burundi, Cameroon, Central African Republic, Republic of the Congo, Democratic Republic of the Congo, Equatorial Guinea, Gabon, Kenya,...

Turkestan Lynx

Turkestan Lynx

Lynx lynx isabellinus

The Turkestan lynx (*Lynx lynx isabellinus*) is a subspecies of Eurasian lynx native to Central Asia. It is also known as Central Asian lynx, Tibetan lynx, or Himalayan lynx. It is widespread from west in Central Asia, from South Asia to China and...

Turkestan rat

Turkestan rat

Rattus pyctoris

The Turkestan rat (*Rattus pyctoris*) is a species of rodent in the family Muridae.

Turkestan red pika

Turkestan red pika

Ochotona rutila

The Turkestan red pika (*Ochotona rutila*) is a species of mammal in the family Ochotonidae. The summer fur at its back is bright rufous and the ventral fur is white or ochraceous. The winter dorsal fur is pale brown and the ventral fur is

white...

Turkestani long-eared bat
Turkestani long-eared bat
Otonycteris leucophaea

The Turkestani long-eared bat, *Otonycteris leucophaea*, is a species of bat found in Asia. Though it was initially described in 1873 as a species, for many years it was considered synonymous with the desert long-eared bat, *Otonycteris hemprichii*....

Turkish hamster
Turkish hamster
Mesocricetus brandti

The Turkish hamster (*Mesocricetus brandti*), also referred to as Brandt's hamster, Azerbaijani hamster, or avurtlak, is a species of hamster native to Turkey, Armenia and other surrounding nations. The Turkish hamster, first catalogued in 1878,...

Turkmen wild goat
Turkmen wild goat
Capra aegagrus turcmenica

The Turkmen wild goat or bearded ibex (*Capra aegagrus turcmenica*) is a vulnerable sub-species of wild goat native to Iran and Turkmenistan. Turkmen wild goats are present in scattered populations in the central Kopet Dag along the border...

Turkmenian fox
Turkmenian fox
Vulpes vulpes flavescens

The Turkmenian fox (*Vulpes vulpes flavescens*), also known as the Persian fox, is an Asiatic subspecies of red fox distinguished by its very small size and primitive cranial features. It inhabits the Middle Asian plains and approximately south of...

Turkmenian Kulan
Turkmenian Kulan
Equus hemionus kulan

The Turkmenian kulan (*Equus hemionus kulan*) is a subspecies of onager (Asiatic wild ass) native to Central Asia. It is one of the largest subspecies of the onagers. The Turkmenian kulan was declared Endangered in 2016.

Tursiops truncatus gillii
Tursiops truncatus gillii
Tursiops truncatus gillii

The common bottlenose dolphin or Atlantic bottlenose dolphin (*Tursiops truncatus*) is a wide-ranging marine mammal of the family Delphinidae. The common bottlenose dolphin is a very familiar dolphin species due to the wide exposure it receives in...

Turuchan pika
Turuchan pika
Ochotona turuchanensis

The Turuchan pika (*Ochotona turuchanensis*) is a species of pika found in isolated regions in the Central Siberian Plateau. It is a small (16-19 cm) rock dwelling species that is active during the day due to the low temperature at night. It was...

Tuva silver vole
Tuva silver vole
Alticola tuvinicus

The Tuva silver vole (*Alticola tuvinicus*) is a species of rodent in the family Cricetidae. It is found in Mongolia and Russian Federation.
Tweedy's crab-eating rat

Tweedy's crab-eating rat
Ichthyomys tweedii

Tweedy's crab-eating rat (*Ichthyomys tweedii*) is a species of rodent in the family Cricetidae. It is found in two disjoint regions in western Ecuador and central Panama. The species is found near fast-flowing streams in primary and secondary...

Tyler's mouse opossum
Tyler's mouse opossum
Marmosa tyleriana

Tyler's mouse opossum (*Marmosa tyleriana*) is a South American marsupial of the family Didelphidae. It lives in rainforests of the Guiana Highlands of southern Venezuela at elevations between 1300 and 2200 m. The species has only been found on...

Typical striped grass mouse
Typical striped grass mouse
Lemniscomys striatus

The typical striped grass mouse (*Lemniscomys striatus*) is a small rodent of the suborder Myomorpha in the family Muridae.

Typical vlei rat
Typical vlei rat
Otomys typus

The typical vlei rat (*Otomys typus*) is a species of vlei rat in the rodent family Muridae. It is found only in Ethiopia. Its natural habitats are subtropical or tropical moist montane forests, subtropical or tropical high-altitude shrubland, and...

Tyrrhenian mole
Tyrrhenian mole
Talpa tyrrhenica

The Tyrrhenian mole (*Talpa tyrrhenica*) is an extinct species of mammal in the family Talpidae. It was endemic to the Mediterranean islands of Corsica and Sardinia during the Pleistocene epoch. First described in 1945 by Dorothea Bate, this...

Udine shrew
Udine shrew
Sorex arunchi

The Udine shrew (*Sorex arunchi*) is a species of mammal in the family Soricidae. It is found in the Udine province of north-east Italy and in western Slovenia.

Udzungwa red colobus
Udzungwa red colobus
Procolobus gordonorum

The Udzungwa red colobus (*Piliocolobus gordonorum*), also known as the Uzungwa red colobus or Iringa red colobus, is a species of primate in the family Cercopithecidae. It is endemic to riverine and montane forest in the Udzungwa Mountains in...

Uganda mangabey
Uganda mangabey
Lophocebus ugandae

The Uganda mangabey (*Lophocebus ugandae*) is a species of Old World monkey found only in Uganda and in the Minziro Forest Reserve, just over the border in Tanzania. This crested mangabey was previously thought to just be a population of the...

Ugandan Kob
Ugandan Kob

Kobus kob thomasi

The Ugandan kob (*Kobus kob thomasi*) is a subspecies of the kob. It appears on the coat of arms of Uganda, along with a Grey crowned crane, representing the abundant wildlife present in the country.

Ugandan lowland shrew
Ugandan lowland shrew
Crocidura selina

The Ugandan lowland shrew (*Crocidura selina*) is a species of mammal in the family Soricidae. It is found in Kenya and Uganda. Its natural habitats are subtropical or tropical swamps and subtropical or tropical moist montane forest. It is...

Ugandan musk shrew
Ugandan musk shrew
Crocidura mutesae

The Ugandan musk shrew (*Crocidura mutesae*) is a species of mammal in the family Soricidae. It is found in Uganda, the Democratic Republic of Congo and the Central African Republic. Its range, population size and habits are poorly known.

Ugandan red colobus
Ugandan red colobus
Piliocolobus tephrosceles

The Ugandan red colobus (*Piliocolobus tephrosceles*) or ashy red colobus is an endangered species of red colobus monkey, recognised as a distinct species since 2001. There is disagreement however over taxonomy with many considering the

Ugandan...
Uinta Chipmunk
Uinta Chipmunk
Neotamias umbrinus

The Uinta chipmunk or hidden forest chipmunk (*Neotamias umbrinus*), is a species of chipmunk in the family Sciuridae. It is endemic to the United States. Formerly known as *Tamias umbrinus*, phylogenetic studies have shown it to be sufficiently...

Uinta Ground Squirrel
Uinta Ground Squirrel
Urocitellus armatus

The Uinta ground squirrel (*Urocitellus armatus*), commonly called a "chisler" and Potgut in northern Utah, is a species of rodent native to the western United States.

Ultimate shrew
Ultimate shrew
Crocidura ultima

The ultimate shrew (*Crocidura ultima*) is a species of mammal in the family Soricidae. It is endemic to Kenya. Its natural habitat is subtropical or tropical moist montane forests. It is threatened by habitat loss.

Uluguru bushbaby
Uluguru bushbaby
Paragalago orinus

The Uluguru bushbaby (*Paragalago orinus*), also known as the mountain dwarf galago or the Amani dwarf galago, is a species of primate in the family Galagidae. Like all galagos, it is a strepsirrhine primate. It is endemic to the Eastern Arc...

Umboi tube-nosed fruit bat
Umboi tube-nosed fruit bat
Nyctimene vizcaccia

The Umboi tube-nosed fruit bat (*Nyctimene vizcaccia*) is a species of megabat in the family Pteropodidae. It is found in Papua New Guinea and the Solomon Islands.
Unadorned rock-wallaby
Unadorned rock-wallaby
Petrogale inornata

The unadorned rock-wallaby (*Petrogale inornata*) is a member of a group of closely related rock-wallabies found in northeastern Queensland, Australia. It is paler than most of its relatives and even plainer, hence its common name.The unadorned...
Unalaska collared lemming
Unalaska collared lemming
Dicrostonyx unalascensis

The Unalaska collared lemming (*Dicrostonyx unalascensis*) is a species of rodent in the family Cricetidae.This species is found on two islands, Umnak and Unalaska, in the Aleutian Archipelago of Alaska in the United States.Its natural habitat is...
Underwood's bonneted bat
Underwood's bonneted bat
Eumops underwoodi

Underwood's bonneted bat (*Eumops underwoodi*) is a species of bat in the family Molossidae found in Belize, Costa Rica, El Salvador, Guatemala, Honduras, Mexico, Nicaragua and the southwestern United States.
Underwood's long-tongued bat
Underwood's long-tongued bat
Hylonycteris underwoodi

Underwood's long-tongued bat (*Hylonycteris underwoodi*) is a species of bat in the family Phyllostomidae. It is monotypic within the genus *Hylonycteris*. It is found in Belize, Guatemala, Mexico, Nicaragua, and Panama. *Hylonycteris underwoodi* feed...
Underwood's pocket gopher
Underwood's pocket gopher
Orthogeomys underwoodi

Underwood's pocket gopher (*Heterogeomys underwoodi*) is a species of rodent in the family Geomyidae. It is endemic to Costa Rica. Some authors classify it in the genus *Orthogeomys*, but recent research has allowed this and its related species to...
Underwood's water mouse
Underwood's water mouse
Rheomys underwoodi

Underwood's water mouse (*Rheomys underwoodi*) is a species of rodent in the family Cricetidae. It is found in Costa Rican and western Panamanian cloud forest at altitudes from 1500 to 2000 m. This mouse lives near streams in highland forests and...
Unduavi gracile opossum
Unduavi gracile opossum
Cryptomys unduaviensis

The Unduavi gracile opossum (*Cryptomys unduaviensis*) is a species of opossum in the family Didelphidae. It is native to northern Bolivia, where it has been found in seasonally flooded grassland. Some of the specimens recognized by Voss et al....
Unexpected cotton rat
Unexpected cotton rat
Sigmodon inopinatus

The unexpected cotton rat (*Sigmodon inopinatus*) is a species of rodent in the

family Cricetidae. It is found only in Ecuador at elevations of 3500 to 4000 m, where it has been found in association with streams and marshes. It is also known as...

Ungava collared lemming
Ungava collared lemming
Dicrostonyx hudsonius

The Ungava collared lemming or Labrador collared lemming (*Dicrostonyx hudsonius*) is a small North American lemming. This species has a short, chunky body covered with brownish-grey fur, with a thin dark stripe along the back and a yellow line...

Unicolored Oldfield mouse
Unicolored Oldfield mouse
Thomasomys monochromos

The unicolored Oldfield mouse (*Thomasomys monochromos*) is a species of rodent in the family Cricetidae. It is found only in Colombia.

Unstriped ground squirrel
Unstriped ground squirrel
Xerus rutilus

The unstriped ground squirrel (*Xerus rutilus*) is a species of rodent (order Rodentia) in the family Sciuridae. It is the only member of the genus *Xerus* not to have a stripe. It is found in Djibouti, Eritrea, Ethiopia, Kenya, Somalia, Sudan,...

Upemba lechwe
Upemba lechwe
Kobus anseli

The Upemba lechwe (*Kobus lechwe anseli*) is a subspecies of antelope found only in the Upemba wetlands in the Democratic Republic of Congo. It was described in 2005, after analysis of 35 museum specimens collected in 1926 and 1947–8. Some...

Upemba shrew
Upemba shrew
Crocidura zimmeri

The Upemba shrew (*Crocidura zimmeri*) is a species of shrew in the family Soricidae. It is endemic to Democratic Republic of the Congo. Its natural habitat is swamps. It is threatened by habitat loss.

Upper Galilee Mountains blind mole-rat
Upper Galilee Mountains blind mole-rat
Spalax galili

The Middle East blind mole-rat or Palestine mole-rat (*Nannospalax ehrenbergi*) is a species of rodent in the family Spalacidae.

Ural field mouse
Ural field mouse
Apodemus uralensis

The Ural field mouse (*Apodemus uralensis*) is a species of rodent in the family Muridae. It is also known as the pygmy field mouse. It is found in Armenia, Austria, Azerbaijan, Belarus, Bulgaria, China, Croatia, Czech Republic, Estonia, Georgia,...

Ural mouse-like hamster
Ural mouse-like hamster
Calomyscus urartensis

The Ural mouse-like hamster or Azerbaijani mouse-like hamster (*Calomyscus urartensis*) is a medium-sized mouse-like hamster from Nakhichevan, Azerbaijan and northwestern Azerbaijan Province, Iran. It is among the better studied

representatives of...

Urial

Urial

Ovis orientalis

The urial (00R-ee-əl; *Ovis vignei*), also known as the arkars or shapo, is a wild sheep native to Central and South Asia. It is listed as Vulnerable on the IUCN Red List.

Uromys vika

Uromys vika

Uromys vika

The Vangunu giant rat (*Uromys vika*), locally known as the vika, is a giant arboreal species of rodent in the family Muridae. The rat was discovered in the island of Vangunu in the Solomon Islands in 2015, after years of searching based on local...

Ursine colobus

Ursine colobus

Colobus vellerosus

The ursine colobus (*Colobus vellerosus*, white-thighed colobus, Geoffroy's black-and-white colobus, white-thighed black-and-white colobus) is a species of primate in the family Cercopithecidae.

Ursine howler

Ursine howler

Alouatta arctoidea

The ursine howler (*Alouatta arctoidea*) is a species of howler monkey native to Venezuela and possibly Colombia. It is sometimes considered a subspecies of the Venezuelan red howler and classified as *Alouatta seniculus arctoidea*.

Ursine tree-kangaroo

Ursine tree-kangaroo

Dendrolagus ursinus

The ursine tree-kangaroo (*Dendrolagus ursinus*) is a species of marsupial in the family Macropodidae. It is endemic to the Vogelkop and possibly the Fakfak Peninsulas, West Papua, Indonesia. Other common names for this species include the black...

Ursus americanus carlottae

Ursus americanus carlottae

Ursus americanus carlottae

The Haida Gwaii black bear (*Ursus americanus carlottae*), also known as the Queen Charlotte Islands black bear, is a morphologically distinct subspecies of the American black bear. The most significant morphological differences are its large...

Ursus deningeri

Ursus deningeri

Ursus deningeri

Ursus deningeri (Deninger's bear) is an extinct species of bear, endemic to Eurasia during the Pleistocene for approximately 1.7 million years, from 1.8 to 0.1 million years ago. The range of this bear has been found to encompass both Europe and...

Ursus dolinensis

Ursus dolinensis

Ursus dolinensis

Ursus dolinensis (the Gran Dolina bear) is an extinct mammalian carnivore species of the Ursidae family. Its fossilized remains were unearthed from the lowest layers

of the stratigraphic sequence at the archaeological and paleontological site of...

Ursus etruscus
Ursus etruscus
Ursus etruscus

Ursus etruscus (the Etruscan bear) is an extinct species of bear, endemic to Europe, Asia and North Africa during the Pliocene through Pleistocene, living from ~5.3 million to 100,000 years ago.

Ursus ingressus
Ursus ingressus
Ursus ingressus

Ursus ingressus (the Gamssulzen Cave bear) is an extinct species of the family Ursidae that lived in Central Europe during the Late Pleistocene. It is named after the Gamssulzen Cave in Austria, where the holotype of this species was found.

Ursus maritimus tyrannus
Ursus maritimus tyrannus
Ursus maritimus tyrannus

Ursus maritimus tyrannus (meaning tyrant polar bear) is an extinct subspecies of polar bear, known from a single fragmentary ulna found in the gravels of the Thames at Kew Bridge, London. It was named by the Finnish paleontologist Björn Kurtén in...

Ursus minimus
Ursus minimus
Ursus minimus

Ursus minimus (the Auvergne bear) is an extinct species of bear, endemic to Europe during the Pliocene and Pleistocene, living from 5.3 to 1.8 Mya, existing for about 3.5 million years. *U. minimus* appears to have given rise to *Ursus etruscus*. The...

Ursus rossicus
Ursus rossicus
Ursus rossicus

Ursus rossicus (the Pleistocene small cave bear) is an extinct species of bear that lived in the steppe regions of northern Eurasia and Siberia during the Pleistocene.

Usambara shrew
Usambara shrew
Crocidura usambarae

The Usambara shrew (*Crocidura usambarae*) is a species of mammal in the family Soricidae. It is endemic to Tanzania. Its natural habitat is subtropical or tropical moist montane forests. It is threatened by habitat loss.

Uspallata chinchilla rat
Uspallata chinchilla rat
Abrocoma uspallata

The Uspallata chinchilla rat (*Abrocoma uspallata*) is a species of chinchilla rat in the family Abrocomidae native to Argentina. This species was identified in 2002, by Braun and Mares from the University of Oklahoma. Only a single specimen has...

Ussuri Black Bear
Ussuri Black Bear
Ursus thibetanus ussuricus

The Ussuri black bear (*Ursus thibetanus ussuricus*) is a large subspecies of the Asiatic black bear. It is named after the Ussuri River.

Ussuri brown bear
Ussuri brown bear
Ursus arctos lasiotus

The Ussuri brown bear (*Ursus arctos lasiotus*), also known as the Ezo brown bear or the black grizzly bear, is a subspecies of the brown bear or a population of the Eurasian brown bear (*U. a. arctos*). One of the largest brown bears, a very large...

Ussuri Dhole

Ussuri Dhole

Cuon alpinus alpinus

The Ussuri dhole (*Cuon alpinus alpinus*), also known as the Indian dhole, Eastern Asiatic dhole, Chinese dhole or southern dhole, is the nominate subspecies of the dhole native to East Asia. The Ussuri dhole is also native to China, however it is...

Ussuri shrew

Ussuri shrew

Sorex mirabilis

The Ussuri shrew (*Sorex mirabilis*), also known as the giant shrew, is a species of shrew found in Northeast Asia. An adult Ussuri shrew has a total length including the tail of 137 to 170 mm (5.4 to 6.7 in). It is found in valleys and on the...

Ussuri tube-nosed bat

Ussuri tube-nosed bat

Murina ussuriensis

The Ussuri tube-nosed bat (*Murina ussuriensis*) is a species of vesper bat in the family Vespertilionidae. It is threatened by habitat loss. It is the only species of bat that hibernates in snowbanks.

Ussuri white-toothed shrew

Ussuri white-toothed shrew

Crocidura lasiura

The Ussuri white-toothed shrew (*Crocidura lasiura*) is a species of musk shrew found on the mainland Northeast Asia. It is common and widespread, and is one of the largest shrews found in the region, with adult weight of 14-25 grams. It should...

Uta Hick's bearded saki

Uta Hick's bearded saki

Chiropotes utahicki

Uta Hick's bearded saki (*Chiropotes utahicki*) is an endangered species of bearded saki, a type of New World monkey. It is endemic to Brazil, where restricted to the Amazon between the Xingu and Tocantins Rivers. It was formerly treated as a...

Utah Prairie Dog

Utah Prairie Dog

Cynomys parvidens

The Utah prairie dog (*Cynomys parvidens*) is the smallest species of prairie dog, a member of the squirrel family of rodents. It is a protected species as it faces various threats the most dangerous being habitat loss.

Uzungwe vlei rat

Uzungwe vlei rat

Otomys uzungwensis

The Uzungwe vlei rat (*Otomys uzungwensis*) is a species of rodent in the family Muridae. It is found in Malawi, Tanzania, and Zambia. Its natural habitats are subtropical or tropical high-altitude grassland and swamps. It is threatened by habitat loss.

Vagrant shrew

Vagrant shrew

Sorex vagrans

The vagrant shrew (*Sorex vagrans*), also known as the wandering shrew, is a medium-sized North American shrew. At one time, the montane shrew and the Orizaba long-tailed shrew were considered to belong to the same species.

Val's gundi

Val's gundi

Ctenodactylus vali

Val's gundi (*Ctenodactylus vali*) is a species of rodent in the family Ctenodactylidae. It is known from two widely separated areas of North Africa.

Valais shrew

Valais shrew

Sorex antinorii

The Valais shrew (*Sorex antinorii*) is a species of mammal in the family Soricidae.

Vampyriscus nymphaea

Vampyriscus nymphaea

Vampyriscus nymphaea

Vampyriscus nymphaea is a species of bat in the family Phyllostomidae, the leaf-nosed bats. It is known commonly as the striped yellow-eared bat. It is native to Central and South America, where it occurs in Colombia, Costa Rica, Ecuador,...

Van Deusen's rat

Van Deusen's rat

Rattus vandeuseni

Van Deusen's rat (*Rattus vandeuseni*) is a species of rodent in the family Muridae. It is endemic to the mountains of southeast Papua New Guinea.

Van Gelder's bat

Van Gelder's bat

Bauerus dubiaquercus

Van Gelder's bat or Van Gelder's big-eared bat (*Bauerus dubiaquercus*) is a species of vesper bat in the family Vespertilionidae. It is found in Belize, Costa Rica, Honduras, and Mexico. The species is monotypic within its genus. It is part of the...

Van Sung's shrew

Van Sung's shrew

Chodsigoa caovansunga

Van Sung's shrew (*Chodsigoa caovansunga*), also known as Cao Van Sung mountain shrew is a species of shrew in the Soricomorpha order. Specimens of *Chodsigoa caovansunga* have been found in Vietnam. *Chodsigoa caovansunga* was named after Cao Van...

Van Zyl's golden mole

Van Zyl's golden mole

Cryptochloris zyli

Van Zyl's golden mole (*Cryptochloris zyli*) is a golden mole endemic to the Western Cape Province, South Africa. It is listed as an endangered species due to habitat loss. Golden moles are an ancient group of mammals who live mostly below ground....

Vancouver Island marmot

Vancouver Island marmot

Marmota vancouverensis

The Vancouver Island marmot (*Marmota vancouverensis*) naturally occurs only in the high mountains of Vancouver Island, in British Columbia. This particular marmot species is large compared to some other marmots, and most other rodents. Marmots

as...

Vancouver Island Wolf
Vancouver Island Wolf
Canis lupus crassodon

The Vancouver Coastal Sea wolf or Vancouver Coastal Island wolf (*Canis lupus crassodon*) is a subspecies of grey wolf, found along the Pacific Coast of Canada. These are one of the shyest and most elusive creatures rarely seen by humans. They are...

Vanikoro flying fox
Vanikoro flying fox
Pteropus tuberculatus

The Vanikoro flying fox (*Pteropus tuberculatus*), also known locally as basapine, is a species of bat in the family Pteropodidae. It has only been found in the Vanikoro island group located in the southern Solomon Islands. The species as a whole...

Vanuatu flying fox
Vanuatu flying fox
Pteropus anetianus

The Vanuatu flying fox or white flying fox (*Pteropus anetianus*) is a species of flying fox in the family Pteropodidae. It is endemic to Vanuatu. It is most closely related to the Samoa flying fox.

Vanzolini's bald-faced saki
Vanzolini's bald-faced saki
Pithecia vanzolinii

Vanzolini's bald-faced saki (*Pithecia vanzolinii*) is a species of saki monkey, a type of New World monkey. Its range is in western Brazil. The monkey is named after Brazilian zoologist Paulo Vanzolini. This species was originally described by...

Vaquita
Vaquita
Phocoena sinus

Vaquitas (*Phocoena sinus*) are record-holders among all cetaceans. Thus, these animals are the smallest cetaceans, meanwhile being the smallest porpoises; they have the smallest range; and finally, vaquitas are the most critically endangered...

Variable grass mouse
Variable grass mouse
Akodon varius

The variable grass mouse (*Akodon varius*) is a species of rodent in the family Cricetidae. It is known only from Bolivia, where it is found at elevations from 400 to 3000 m.

Variable pocket gopher
Variable pocket gopher
Orthogeomys heterodus

The variable pocket gopher (*Heterogeomys heterodus*) is a species of rodent in the family Geomyidae. It is endemic to Costa Rica, usually being found in grasslands and tropical forests at higher altitudes, up to 8,000 feet. It is threatened by...

Varied white-fronted capuchin
Varied white-fronted capuchin
Cebus versicolor

The varied white-fronted capuchin (*Cebus versicolor*) is a species of gracile capuchin monkey from Colombia. It had been classified as a subspecies of the white-

fronted capuchin (*C. albifrons*) Genetic analysis by Jean Boubli in 2012 revealed it...

Variegated butterfly bat
Variegated butterfly bat
Glauconycteris variegata

The variegated butterfly bat (*Glauconycteris variegata*) is a species of vesper bat. It is sometimes also called the leaf-winged bat, or simply the butterfly bat. It is not currently endangered, but may be threatened by habitat loss in some parts...

Variegated Squirrel
Variegated Squirrel
Sciurus variegatoides

The variegated squirrel (*Sciurus variegatoides*) is a tree squirrel in the genus *Sciurus* found in Costa Rica, El Salvador, Guatemala, Honduras, southern Mexico, Nicaragua, and Panama. Fifteen subspecies are recognised. It is a common squirrel and...

Variegated wolf
Variegated wolf
Canis lupaster soudanicus

The variegated wolf (*Canis lupaster soudanicus*), also known as the Nubian wolf, is a subspecies of African golden wolf native to Sudan and Somalia. It is smaller and more lightly built than the Egyptian wolf, standing 38 cm (15 in) at the...

Veldkamp's dwarf epauletted fruit bat
Veldkamp's dwarf epauletted fruit bat
Nanonycteris veldkampii

Veldkamp's dwarf epauletted fruit bat (*Nanonycteris veldkampii*) is a species of bat in the family Pteropodidae. It is monotypic within the genus *Nanonycteris*. It is found in Benin, Cameroon, Central African Republic, Ivory Coast, Ghana, Guinea,...

Velvet climbing mouse
Velvet climbing mouse
Dendroprionomys rousseti

The velvet climbing mouse (*Dendroprionomys rousseti*) is a species of rodent in the family Nesomyidae. It is found only in Republic of the Congo.

Velvety free-tailed bat
Velvety free-tailed bat
Molossus molossus

The velvety free-tailed bat or Pallas's mastiff bat (*Molossus molossus*), is a bat species in the family Molossidae.

Velvety fruit-eating bat
Velvety fruit-eating bat
Enchisthenes hartii

The velvety fruit-eating bat (*Enchisthenes hartii*), also known as Hart's little fruit bat, is a species of bat in the family Phyllostomidae. It is monotypic within the genus *Enchisthenes*. It is found in Central America, Mexico, the United...

Velvety myotis
Velvety myotis
Myotis simus

The velvety myotis (*Myotis simus*), is a species of vesper bat from South America.

Venezuelan climbing mouse
Venezuelan climbing mouse

Rhipidomys venezuelae

The Venezuelan climbing mouse (*Rhipidomys venezuelae*) is a species of rodent in the family Cricetidae. It is found in Colombia, Tobago, and Venezuela.

Venezuelan fish-eating rat

Venezuelan fish-eating rat

Neusticomys venezuelae

The Venezuelan fish-eating rat (*Neusticomys venezuelae*) is a species of a Rodent in the Cricetidae family. It is found in Guyana and Venezuela.

Venezuelan lowland rabbit

Venezuelan lowland rabbit

Sylvilagus varynaensis

The Venezuelan lowland rabbit (*Sylvilagus varynaensis*), also known as the Barinas wild rabbit, is a cottontail rabbit species found in western Venezuela.

Veracruz shrew

Veracruz shrew

Sorex veraecrucis

The Veracruz shrew (*Sorex veraecrucis*) is a species of mammal in the family Soricidae. It is found in Mexico.

Verapaz shrew

Verapaz shrew

Sorex veraepacis

The Verapaz shrew (*Sorex veraepacis*) is a species of mammal in the family Soricidae. It is found in Guatemala and Mexico.

Verhagen

Verhagen

Lophuromys verhageni

Verhagen is a Dutch-language toponymic surname. It is a contraction of Van der Hagen, meaning "from/of the haag. A haag was a bushland, hedged lot, or (private) hunting ground. The name could also specifically refer to an origin in The Hague...

Verheyen's multimammate mouse

Verheyen's multimammate mouse

Mastomys kollmannspergeri

Verheyen's multimammate mouse (*Mastomys kollmannspergeri*) is a species of rodent in the family Muridae found in Cameroon, Chad, Niger, Nigeria, and Sudan. Its natural habitats are dry savanna, intermittent rivers, intermittent freshwater lakes,...

Verhoeven's giant tree rat

Verhoeven's giant tree rat

Papagomys theodorverhoeveni

Verhoeven's giant tree rat (*Papagomys theodorverhoeveni*) is an extinct rat of subfamily Murinae that lived on Flores in Indonesia. It was judged to be extinct in 1996. However, experts believe that it died out before 1500 AD. The species is...

Vernay's climbing mouse

Vernay's climbing mouse

Dendromus vernayi

Vernay's climbing mouse (*Dendromus vernayi*) is a species of rodent in the family Nesomyidae. It is found only in Angola. Its natural habitat is moist savanna.

Verreaux's mouse

Verreaux's mouse

Myomyscus verreauxii

Verreaux's mouse or Verreaux's white-footed rat (*Myomyscus verreauxii*) is a species of rodent in the family Muridae. It is the only member of the genus *Myomyscus*. Other species that had been previously assigned to *Myomyscus* are now considered to...

Verreaux's Sifaka

Verreaux's Sifaka

Propithecus verreauxi

Verreaux's sifaka (*Propithecus verreauxi*), or the white sifaka, is a medium-sized primate in one of the lemur families, the Indriidae. It lives in Madagascar and can be found in a variety of habitats from rainforest to western Madagascar dry...

Verschuren's swamp rat

Verschuren's swamp rat

Praomys verschureni

Verschuren's swamp rat (*Congomys verschureni*) is a species of rodent in the family Muridae. It is found only in Democratic Republic of the Congo. Its natural habitat is subtropical or tropical moist lowland forests. It is threatened by habitat loss.

Vervet Monkey

Vervet Monkey

Chlorocebus pygerythrus

The Vervet monkey (*Chlorocebus pygerythrus*) is an Old World monkey native to Africa. The term "vervet" is also used to refer to all the members of the genus *Chlorocebus*. These adorable monkeys live in social groups and involve their communication...

Vicuna

Vicuna

Vicugna vicugna

The vicuña (*Lama vicugna*) or vicuna is one of the two wild South American camelids, which live in the high alpine areas of the Andes. Vicunas are relatives of the llama, and are now believed to be the wild ancestor of domesticated alpacas, which...

Vieira's long-tongued bat

Vieira's long-tongued bat

Xeronycteris vieirai

Vieira's long-tongued bat (*Xeronycteris vieirai*) is a species of bat from northeastern Brazil, discovered in 2005 by Gregorin and Ditchfield. It is the only species in the genus *Xeronycteris*.

Vieira's titi

Vieira's titi

Plecturocebus vieirai

Vieira's titi monkey (*Plecturocebus vieirai*) is a species of titi, a type of New World monkey, from central-northern Brazil.

Vietnam ferret-badger

Vietnam ferret-badger

Melogale cucphuongensis

The Vietnam ferret-badger (*Melogale cucphuongensis*) is a member of the family Mustelidae native to Vietnam. It was described in 2011 and is known from only two specimens.

Vietnamese sika deer

Vietnamese sika deer

Cervus nippon pseudaxis

The Vietnamese sika deer (*Cervus nippon pseudaxis*) also known as the indochinese sika deer is one of the many subspecies of the sika deer. It is one of the smaller subspecies, due to the tropical environment they live in. They were previously...

Villa's gray shrew

Villa's gray shrew

Notiosorex villa

Villa's gray shrew (*Notiosorex villai*) is a shrew native to northeastern Mexico, where it is called musaraña. It is assessed vulnerable by the IUCN due to its small extent of occurrence and degradation of habitat.

Vincent's bush squirrel

Vincent's bush squirrel

Paraxerus vincenti

Vincent's bush squirrel (*Paraxerus vincenti*) is a species of rodent in the family Sciuridae, named in honour of Jack Vincent. It is endemic to Mozambique. Its natural habitat is subtropical or tropical moist montane forests, and it is threatened...

Vinogradov's jerboa

Vinogradov's jerboa

Allactaga vinogradovi

Vinogradov's jerboa (*Allactaga vinogradovi*) is a species of rodent in the family Dipodidae. It is found in Kazakhstan, Kyrgyzstan, Tajikistan, and Uzbekistan.

Vinogradov's jird

Vinogradov's jird

Meriones vinogradovi

Vinogradov's jird (*Meriones vinogradovi*) is a species of rodent in the family Muridae. It is found in Armenia, Azerbaijan, Iran, Syria, and Turkey.

Vinson's slit-faced bat

Vinson's slit-faced bat

Nycteris vinsoni

Vinson's slit-faced bat (*Nycteris vinsoni*) is a species of slit-faced bat known only from two specimens. Both specimens were smoked out of a baobab tree in a national park in southern Mozambique. Virtually nothing is known about this species. N....

Vipos tuco-tuco

Vipos tuco-tuco

Ctenomys viperinus

The Vipos tuco-tuco (*Ctenomys viperinus*) is a species of rodent in the family Ctenomyidae. It is endemic to northern Tucumán Province in northern Argentina. The common name is a reference to the city of Vipos in the area.

Virginia Big-Eared Bat

Virginia Big-Eared Bat

Corynorhinus townsendii virginianus

The Virginia big-eared bat (*Corynorhinus townsendii virginianus*) is one of two endangered subspecies of the Townsend's big-eared bat. It is the state bat of Virginia.

Virginia Opossum

Virginia Opossum

Didelphis virginiana

The Virginia opossum (*Didelphis virginiana*) is the only marsupial found in North America north of Mexico. In the United States, it is typically referred to simply as a possum. Virginia opossums are familiar to many North Americans as they are...

Visagie's golden mole
Visagie's golden mole
Chrysochloris visagiei

Visagie's golden mole (*Chrysochloris visagiei*) is a small, insectivorous mammal of the family Chrysochloridae, the golden moles, endemic to South Africa.

Visayan Leopard Cat
Visayan Leopard Cat
Prionailurus javanensis sumatranus

The Visayan leopard cat is a Sunda leopard cat (*Prionailurus javanensis sumatranus*) population. It has been listed as vulnerable on the IUCN Red List in 2008 under its former scientific name *P. bengalensis rabori* as its range is estimated to be...

Visayan Spotted Deer
Visayan Spotted Deer
Rusa alfredi

The Visayan spotted deer (*Rusa alfredi*), also known as the Visayan deer, the Philippine spotted deer or Prince Alfred's deer, is a nocturnal and endangered species of deer located primarily in the rainforests of the Visayan islands of Panay and...

Visayan Warty Pig
Visayan Warty Pig
Sus cebifrons

The Visayan warty pig (*Sus cebifrons*) is a critically endangered species in the pig genus (*Sus*). It is known by many names in the region (depending on the island and linguistic group) with most translating into 'wild pig': baboy ilahas ('wild...

Visored bat
Visored bat
Sphaeronycteris toxophyllum

The visored bat, (*Sphaeronycteris toxophyllum*), is a bat species from tropical South America. It is the only species in the genus *Sphaeronycteris*. Although visored bats have some unique characteristics, they are thought to be most closely...

Voalavoanala
Voalavoanala
Gymnuromys roberti

The voalavoanala (*Gymnuromys roberti*) is a species of rodent in the family Nesomyidae. It is the only species in the genus *Gymnuromys*. It is found only in Madagascar. Its natural habitat is tropical dry forests.

Vogelkop mountain rat
Vogelkop mountain rat
Rattus arfakiensis

The Vogelkop mountain rat, *Rattus arfakiensis*, is a species of rat native to Indonesia. It is found only in the Bird's Head Peninsula (possibly throughout the Arfak Mountains) of Papua Province, Indonesia.

Vogelkop ringtail possum
Vogelkop ringtail possum
Pseudochirulus schlegeli

The Vogelkop ringtail possum (*Pseudochirulus schlegeli*) is a species of marsupial in the family Pseudocheiridae. It is endemic to the Vogelkop Peninsula, West Papua, Indonesia. It is threatened by habitat loss.

Voi shrew
Voi shrew

Crocidura voi

The Voi shrew (*Crocidura voi*) is a species of mammal in the family Soricidae. It is found in Benin, Burkina Faso, Cameroon, Central African Republic, Chad, Ethiopia, Ghana, Kenya, Mali, Niger, Nigeria, Somalia, Sudan, and Togo. Its natural...

Volcano harvest mouse

Volcano harvest mouse

Reithrodontomys chrysopsis

The volcano harvest mouse (*Reithrodontomys chrysopsis*) is a species of rodent in the family Cricetidae. It is found only in Mexico.

Volcano mouse

Volcano mouse

Mus vulcani

The volcano mouse or Javan shrew-like mouse (*Mus vulcani*) is a species of rodent in the family Muridae endemic to Indonesia.

Volcano Rabbit

Volcano Rabbit

Romerolagus diazi

The volcano rabbit (*Romerolagus diazi*), also known as teporingo or zacatuche, is a small rabbit that resides in the mountains of Mexico. It is the world's second-smallest rabbit, second only to the pygmy rabbit. It has small rounded ears, short...

Volcano shrew

Volcano shrew

Sylvisorex vulcanorum

The volcano shrew (*Sylvisorex vulcanorum*) is a species of mammal in the family Soricidae found in the high-altitude rainforest of Burundi, eastern Democratic Republic of the Congo, Rwanda, and Uganda. Its natural habitats are subtropical or...

Von der Decken's sifaka

Von der Decken's sifaka

Propithecus deckenii

Von der Decken's sifaka (*Propithecus deckenii*) is a sifaka endemic to Madagascar. It has a length of 92 to 107 centimeters, of which 42-48 centimeters are tail. Von der Decken's Sifaka lives in western Madagascar. It lives in dry deciduous...

Voracious shrew

Voracious shrew

Crocidura vorax

The voracious shrew (*Crocidura vorax*) is a common and widespread species of shrew native to China, India, Laos, Thailand, and Vietnam.

Vordermann's flying squirrel

Vordermann's flying squirrel

Petinomys vordermanni

Vorderman's flying squirrel (*Petinomys vordermanni*) is a species of rodent in the family Sciuridae. It is found in Indonesia, Malaysia, and Myanmar. It was described in 1890 by the Dutch zoologist Fredericus Anna Jentink, who named it after its...

Vordermann's pipistrelle

Vordermann's pipistrelle

Hypsugo vordermanni

The Vorderman's pipistrelle (*Hypsugo vordermanni*) is a species of vesper bat found

in Brunei Darussalam, Indonesia, and Malaysia. It was described in 1890 by the Dutch zoologist Fredericus Anna Jentink, who named it after its discoverer, the...
Vulpes riffautae
Vulpes riffautae
Vulpes riffautae

Vulpes riffautae is an extinct species of fox from the late Miocene of Chad (approximately 7 ma). Fossils of *V. riffautae* potentially represent the earliest record of the dog family, Canidae, in the Old World. *V. riffautae* was intermediate in...
Vulpes skinneri
Vulpes skinneri
Vulpes skinneri

Vulpes skinneri is a species of extinct fox in the genus *Vulpes* from the early Pleistocene, identified based on fossil remains dated to about 2 million years ago. The species is known from a single partial skeleton discovered in the Malapa Fossil...
Vulpes vulpes kurdistanica
Vulpes vulpes kurdistanica
Vulpes vulpes kurdistanica

The Trans-Caucasian fox (*Vulpes vulpes kurdistanica*) is a subspecies of the red fox, found in the northeast part of Turkey.
Wagner's bonneted bat
Wagner's bonneted bat
Eumops glaucinus

Wagner's bonneted bat or Wagner's mastiff bat (*Eumops glaucinus*), is a species of bat in the family Molossidae. It is found in the Americas from Argentina and Peru north to Mexico, and Cuba. Populations in Florida in the United States are now...
Wagner's gerbil
Wagner's gerbil
Dipodillus dasyurus

Wagner's gerbil (*Dipodillus dasyurus*) is a gerbil that is native mainly to the Nile Delta, Israel, the Sinai, Syria, Iraq and the Arabian Peninsula. It also referred to as the rough-tailed dipodil or Wadi Hof gerbil. They are solo, burrowing...
Wagner's mustached bat
Wagner's mustached bat
Pteronotus personatus

Wagner's mustached bat (*Pteronotus personatus*) is a bat species from South and Central America. It is one of the few New World bats species known to perform Doppler shift compensation behavior.
Wahlberg's Epauletted Fruit Bat
Wahlberg's Epauletted Fruit Bat
Epomophorus wahlbergi

Wahlberg's epauletted fruit bat (*Epomophorus wahlbergi*) is a species of megabat in the family Pteropodidae. It is commonly found across southern Africa.
Waigeou cuscus
Waigeou cuscus
Spilocuscus papuensis

The Waigeou cuscus or Waigeou spotted cuscus (*Spilocuscus papuensis*) is a species of marsupial in the family Phalangeridae. It is endemic to the island of Waigeo in Indonesia, and consequently the spelling Waigeo cuscus is often used instead of...

Waiomys
Waiomys
Waiomys mamasae

Waiomys is a genus of rodents from the family Muridae. It is endemic to Sulawesi, Indonesia. The genus is monotypic, consisting of the species Waiomys mamasae (Sulawesi water rat). It is known only from Mount Gandangdewata, Mamasa Regency, West...

Walia Ibex
Walia Ibex
Capra walie

The walia ibex (Capra walie, Amharic: ዋሊያ wālyā) is an endangered species of ibex. It is sometimes considered a subspecies of the Alpine ibex. Threats against the species include habitat loss, poaching, and restricted range; only about 500...

Wall-roosting mouse-eared bat
Wall-roosting mouse-eared bat
Myotis muricola

The wall-roosting mouse-eared bat, or Nepalese whiskered myotis (Myotis muricola) is a species of vesper bat whose type locality is Nepal.

Wallace's dasyure
Wallace's dasyure
Myoictis wallacei

Wallace's dasyure (Myoictis wallacei) or Wallace's three-striped dasyure is a member of the order Dasyuromorphia. This marsupial carnivore is found in Indonesia and Papua New Guinea. Specifically, it is found on the Aru Islands of the Maluku...

Wallace's large spiny rat
Wallace's large spiny rat
Halmaheramys wallacei

The Wallace's large spiny rat (Halmaheramys wallacei) is a rodent found on the Obi Islands of Bisa and Obi in the Molucca archipelago of Indonesia. The genus Halmaheramys was originally thought to include only one species, endemic to Halmahera,...

Wallace's tarsier
Wallace's tarsier
Tarsius wallacei

Wallace's tarsier, Tarsius wallacei is a species of Sulawesi tarsier (all tarsiers from the genus tarsius are from Sulawesi and its surrounding islands). It is found in the forests of what is referred to geologically as the neck of Sulawesi where...

Walrus
Walrus
Odobenus rosmarus

The walrus (Odobenus rosmarus) is a large flippered marine mammal with a discontinuous distribution about the North Pole in the Arctic Ocean and subarctic seas of the Northern Hemisphere. The walrus is the only living species in the family...

Walston's tube-nosed bat
Walston's tube-nosed bat
Murina walstoni

Walston's tube-nosed bat (Murina walstoni) is a species in the vesper bat family Vespertilionidae, found in the Greater Mekong region of Southeast Asia, specifically the Đắk Lắk Province of Vietnam and the Koh Kong and Ratanakiri provinces of...

Walter Verheyen's mouse
Walter Verheyen's mouse
Hylomyscus walterverheyeni

Walter Verheyen's mouse (*Hylomyscus walterverheyeni*) is a species of rodent of the genus *Hylomyscus* that is found in the Central African lowland and mountain forests.

Walter's duiker
Walter's duiker
Philantomba walteri

Walter's duiker (*Philantomba walteri*) is a species of duiker found in Togo, Benin and Nigeria. It was described in 2010. Its name commemorates Professor Walter Verheyen, who was the first to obtain a specimen of this species of duiker from Togo...

Wandering Oldfield mouse
Wandering Oldfield mouse
Thomasomys erro

The wandering Oldfield mouse (*Thomasomys erro*), also called the wandering thomasomys, is a species of rodent in the family Cricetidae. It has been found in both the Cordillera Oriental and Cordillera Occidental of the Andes in central and north...

Wandering small-eared shrew
Wandering small-eared shrew
Cryptotis montivaga

The wandering small-eared shrew (*Cryptotis montivaga*) is a species of shrew in the family Soricidae. It is endemic to Ecuador.

Ward's field mouse
Ward's field mouse
Apodemus pallipes

Ward's field mouse (*Apodemus pallipes*) is a species of rodent in the family Muridae. It is found in Kyrgyzstan, Tajikistan, Afghanistan, India, Iran, Nepal, and Pakistan.

Ward's red-backed vole
Ward's red-backed vole
Eothenomys wardi

The Ward's red-backed vole (*Eothenomys wardi*) is a species of rodent in the family Cricetidae. It is found in China, specifically the north-western area of Yunnan Province. It is sometimes considered a subspecies of *Eothenomys chinensis*, but...

Washington ground squirrel
Washington ground squirrel
Urocitellus washingtoni

The Washington ground squirrel (*Urocitellus washingtoni*) is a squirrel that lives in the Pacific Northwest states of Washington and Oregon, United States.

Water Buffalo
Water Buffalo
Bubalus bubalis

The Water buffalo (*Bubalus bubalis*) is a large bovid from the Indian subcontinent and Southeast Asia. Today, it is also found in Europe, Australia, North America, South America, and some African countries. Two extant types of water buffalo are...

Water Chevrotain
Water Chevrotain
Hyemoschus aquaticus

The water chevrotain (*Hyemoschus aquaticus*), also known as the fanged deer, is a small ruminant found in tropical Africa. It is the largest of the 10 species of chevrotains, basal even-toed ungulates which are similar to deer, but are barely...

Water Deer

Water Deer

Hydropotes inermis

The Water deer (*Hydropotes inermis*) is a small deer native to China and Korea. It was first described to the Western world by Robert Swinhoe in 1870. Water deer have developed long canine teeth which protrude from the upper jaw; it is due to...

Water Opossum

Water Opossum

Chironectes minimus

The water opossum (*Chironectes minimus*), also locally known as the yapok, is a marsupial of the family Didelphidae. It is the only living member of its genus, *Chironectes*. This semiaquatic creature is found in and near freshwater streams and...

Water vole

Water vole

Microtus richardsoni

The water vole (*Microtus richardsoni*) is the largest North American vole. It is found in the northwestern United States and southern parts of western Canada. This animal has been historically considered a member of genus *Arvicola*, but molecular...

Waterbuck

Waterbuck

Kobus ellipsiprymnus

The waterbuck (*Kobus ellipsiprymnus*) is a large antelope found in sub-Saharan Africa. It was first described by Irish naturalist William Ogilby in 1833. The waterbuck is predominantly a grazer and is mostly found on grassland.

Waterhouse's leaf-nosed bat

Waterhouse's leaf-nosed bat

Macrotus waterhousii

Waterhouse's leaf-nosed bat (*Macrotus waterhousii*) is a species of big-eared bat in the family Phyllostomidae. It is found in the Cayman Islands, Cuba, the Dominican Republic, Guatemala, Haiti, Jamaica, and Mexico, with a range from Sonora to...

Waterhouse's swamp rat

Waterhouse's swamp rat

Scapteromys tumidus

Waterhouse's swamp rat (*Scapteromys tumidus*) is a semiaquatic rodent species from South America. It is found in southern Brazil, Uruguay and northern Argentina, where it lives in freshwater and salt marshes, as well as open grassland of the...

Waters's gerbil

Waters's gerbil

Gerbillus watersi

Waters's gerbil (*Gerbillus watersi*) is distributed mainly in Sudan, Somalia, and Djibouti.

Watson's climbing rat

Watson's climbing rat

Tylomys watsoni

Watson's climbing rat (*Tylomys watsoni*) is a species of rodent in the family

Cricetidae. It is found in Costa Rica and Panama.

Watts's pipistrelle

Watts's pipistrelle

Pipistrellus wattsi

Watt's pipistrelle (*Pipistrellus wattsi*) is a species of vesper bat. It is found only in Papua New Guinea.

Watts's spiny rat

Watts's spiny rat

Maxomys wattsi

Watts's spiny rat (*Maxomys wattsi*) is a species of rodent in the family Muridae. It is endemic to Sulawesi, Indonesia.

Weasel sportive lemur

Weasel sportive lemur

Lepilemur mustelinus

The weasel sportive lemur (*Lepilemur mustelinus*), also known as the greater sportive lemur, weasel lemur, or greater weasel lemur, is a species of lemur native to northeastern Madagascar. Its habitat includes rainforests and tropical...

Web-footed tenrec

Web-footed tenrec

Microgale mergulus

The web-footed tenrec, Malagasy otter shrew, or aquatic tenrec (*Microgale mergulus*) is the only known semiaquatic tenrec (the related African otter shrews have similar habits), and is found in eastern Madagascar, especially in and around...

Webb's tufted-tailed rat

Webb's tufted-tailed rat

Eliurus webbi

Webb's tufted-tailed rat (*Eliurus webbi*) is a species of rodent in the family Nesomyidae. It is found only in Madagascar. Its natural habitat is subtropical or tropical dry forests. It is threatened by habitat loss.

Weber's dwarf squirrel

Weber's dwarf squirrel

Prosciurillus weberi

Weber's dwarf squirrel (*Prosciurillus weberi*) is a species of rodent in the family Sciuridae. It is endemic to northern and western Sulawesi, Indonesia. Its natural habitat is subtropical or tropical dry forests.

Weddell Seal

Weddell Seal

Leptonychotes weddellii

The Weddell seal (*Leptonychotes weddellii*) is a relatively large and abundant true seal with a circumpolar distribution surrounding Antarctica. The Weddell seal was discovered and named in the 1820s during expeditions led by British sealing...

Wedge-capped capuchin

Wedge-capped capuchin

Cebus olivaceus

The wedge-capped capuchin or Guianan weeper capuchin (*Cebus olivaceus*) is a capuchin monkey from South America. It is found in northern Brazil, Guyana and Venezuela. The genus *Cebus* is divided into several different species. However, taxonomists...

Welwitsch's bat

Welwitsch's bat

Myotis welwitschii

Welwitsch's bat also known as Welwitsch's mouse-eared bat or Welwitsch's myotis (*Myotis welwitschii*) is a species of vesper bat native to Africa.

West African Giraffe

West African Giraffe

Giraffa camelopardalis peralta

The West African giraffe (*Giraffa camelopardalis peralta*) is a subspecies of the giraffe distinguished by its light-colored spots, which is found in the Sahel regions of West Africa. In the 19th century, it ranged from Senegal to Lake Chad, yet...

West African long-tailed shrew

West African long-tailed shrew

Crocidura muricauda

The West African long-tailed shrew (*Crocidura muricauda*) is a species of mammal in the family Soricidae. It can be found in Ivory Coast, Ghana, Guinea, Liberia, and Sierra Leone. Its natural habitat is subtropical or tropical moist lowland...

West African oyan

West African oyan

Poiana leightoni

The West African oyan (*Poiana leightoni*), also known as the West African linsang, is a linsang species native to the Upper Guinean forests in West Africa. It is one of the least known small carnivores in Africa.

West African Potto

West African Potto

Perodicticus potto

The West African potto (*Perodicticus potto*) is a nocturnal strepsirrhine primate found in tropical West Africa. It is also known as Bosman's potto, after Willem Bosman, who described the species in 1704. In some English-speaking parts of Africa,...

West African pygmy shrew

West African pygmy shrew

Crocidura obscurior

The West African pygmy shrew or obscure white-toothed shrew (*Crocidura obscurior*) is a species of mammal in the family Soricidae. It is found in Ivory Coast, Ghana, Guinea, Liberia, and Sierra Leone. Its natural habitat is subtropical or...

West African shaggy rat

West African shaggy rat

Dasymys rufulus

The West African shaggy rat (*Dasymys rufulus*) is a species of rodent in the family Muridae. It is found in Benin, Cameroon, Ivory Coast, Gambia, Ghana, Guinea, Guinea-Bissau, Liberia, Mali, Nigeria, Senegal, Sierra Leone, and Togo. Its natural...

West African wild dog

West African wild dog

Lycaon pictus manguensis

The African wild dog (*Lycaon pictus*), also called the African painted dog and the African hunting dog, is a wild canine which is a native species to sub-Saharan Africa. It is the largest wild canine in Africa, and the only extant member of the...

West Caucasian Tur

West Caucasian Tur

Capra caucasica

The West Caucasian tur (*Capra caucasica*) is a mountain-dwelling goat-antelope native to the western half of the Caucasus Mountains range, in Georgia and European Russia. It is listed as Endangered on the IUCN Red List, as the wild population is...

West Indian Manatee
West Indian Manatee
Trichechus manatus

The West Indian manatee (*Trichechus manatus*) is the largest member of the aquatic mammal order Sirenia (which also includes the dugong and the extinct Steller's sea cow). Manatees are obligate herbivores, have developed vocal communication...

Western barbastelle
Western barbastelle
Barbastella barbastellus

The western barbastelle (*Barbastella barbastellus*), also known as the barbastelle or barbastelle bat, is a European bat in the genus *Barbastella*. This species is found from Portugal to Azerbaijan and from Sweden to Canary Islands, where a...

Western barred bandicoot
Western barred bandicoot
Perameles bougainville

The Western barred bandicoot (*Perameles bougainville*), also known as the Marl, is a small species of bandicoot; now extinct across most of its former range, the western barred bandicoot only survives on offshore islands and in fenced sanctuaries...

Western bent-winged bat
Western bent-winged bat
Miniopterus magnater

The western bent-winged bat (*Miniopterus magnater*) is a species of vesper bat in the family Vespertilionidae. It can be found in the following countries: China, India, Indonesia, Laos, Malaysia, Myanmar, Papua New Guinea, Thailand, Timor-Leste,...

Western black rhinoceros
Western black rhinoceros
Diceros bicornis longipes

The western black rhinoceros (*Diceros bicornis longipes*) or West African black rhinoceros is an extinct subspecies of the black rhinoceros. It was declared extinct by the IUCN in 2011. The western black rhinoceros was believed to have been...

Western broad-nosed bat
Western broad-nosed bat
Scotorepens balstoni

The western or inland broad-nosed bat (*Scotorepens balstoni*) is a species of vespertilionid bats. They are endemic to Australia and widespread throughout the inland, especially in arid and semi-arid regions. This insectivorous microbat,...

Western broad-toothed field mouse
Western broad-toothed field mouse
Apodemus epimelas

The western broad-toothed field mouse (*Apodemus epimelas*) is a species of rodent in the genus *Apodemus* from southeastern Europe. It is related to *A. mystacinus*, which occurs further to the east. It is found in Albania, Bosnia and Herzegovina,...

Western brush wallaby

Western brush wallaby
Macropus irma

The western brush wallaby (*Notamacropus irma*), also known as the black-gloved wallaby, is a species of wallaby found in the southwestern coastal region of Western Australia. The wallaby's main threat is predation by the introduced red fox...

Western chestnut mouse
Western chestnut mouse
Pseudomys nanus

The western chestnut mouse (*Pseudomys nanus*) is a species of rodent in the family Muridae. It is native to northern Australia and various close islands, with the vast majority found in Queensland and the Northern Territory.

Western chimpanzee
Western chimpanzee
Pan troglodytes verus

The western chimpanzee, or West African chimpanzee, (*Pan troglodytes verus*) is a Critically Endangered subspecies of the common chimpanzee. It inhabits western Africa, mainly in Côte d'Ivoire, Guinea, Liberia, Mali, and Sierra Leone but with...

Western dwarf squirrel
Western dwarf squirrel
Microsciurus mimulus

The western dwarf squirrel (*Microsciurus mimulus*) is a small tree squirrel in the genus *Microsciurus* and tribe Sciurini found in Colombia, Ecuador, and Panama. The table below lists the three recognized subspecies of *Microsciurus mimulus*, along...

Western false pipistrelle
Western false pipistrelle
Falsistrellus mackenziei

The western false pipistrelle, species *Falsistrellus mackenziei*, is a vespertilionid bat that occurs in Southwest Australia. The population is declining due to loss of its habitat, old growth in tall eucalypt forest which has largely been clear...

Western gerbil
Western gerbil
Gerbillus hesperinus

The western gerbil (*Gerbillus hesperinus*) is distributed mainly in northern Morocco. It is listed as Endangered by the IUCN.

Western Gorilla
Western Gorilla
Gorilla gorilla

The Western gorilla (*Gorilla gorilla*) is a great ape found in Africa, one of two species of the hominid genus *Gorilla*. Two subspecies are recognized and both are listed Critically Endangered.

Western Gray Squirrel
Western Gray Squirrel
Sciurus griseus

The western gray squirrel (*Sciurus griseus*) is an arboreal rodent found along the western coast of the United States and Mexico. It is a tree squirrel. In some places, this species has also been known as the silver-gray squirrel, the California...

Western Grey Kangaroo

Western Grey Kangaroo
Macropus fuliginosus

The western grey kangaroo (*Macropus fuliginosus*), also referred to as a western grey giant kangaroo, black-faced kangaroo, mallee kangaroo, and sooty kangaroo, is a large and very common kangaroo found across almost the entire southern part of...

Western hartebeest
Western hartebeest
Alcelaphus buselaphus major

The western hartebeest (*Alcelaphus buselaphus major*) is an antelope native to the medium to tall grassland plains of Benin, Burkina Faso, Cameroon, the Central African Republic, Chad, Côte d'Ivoire, Ghana, Guinea, Guinea-Bissau, Mali, Niger,...

Western harvest mouse
Western harvest mouse
Reithrodontomys megalotis

The western harvest mouse (*Reithrodontomys megalotis*) is a small neotomine mouse native to most of the western United States. Many authorities consider the endangered salt marsh harvest mouse to be a subspecies, but the two are now usually...

Western heather vole
Western heather vole
Phenacomys intermedius

The western heather vole (*Phenacomys intermedius*) is a small vole found in western North America. Until recently, the eastern heather vole, (*Phenacomys ungava*), was considered to be a subspecies. These animals are similar in appearance to the...

Western Hoolock Gibbon
Western Hoolock Gibbon
Hoolock hoolock

Western hoolock gibbons (*Hoolock hoolock*) are large primates from the gibbon family, Hylobatidae. They live in family groups and form long-term pair bonds. Their primary mode of locomotion is swinging from branch to branch. They can also leap and...

Western jumping mouse
Western jumping mouse
Zapus princeps

The western jumping mouse (*Zapus princeps*), is a species of rodent in the family Zapodidae. It is found in Canada and the United States. Western jumping mice evolved during the Pleistocene, possibly from the fossil species *Zapus burtoni*, which is...

Western lesser bamboo lemur
Western lesser bamboo lemur
Haplemur occidentalis

The western lesser bamboo lemur (*Haplemur occidentalis*), also known as the northern bamboo lemur, western gentle lemur, or Sambirano lesser bamboo lemur, is a species of bamboo lemur endemic to Madagascar.

Western Long-Beaked Echidna
Western Long-Beaked Echidna
Zaglossus bruijnii

The western long-beaked echidna (*Zaglossus bruijnii*) is one of the four extant echidnas and one of three species of *Zaglossus* that occur in New Guinea. Originally described as *Tachyglossus bruijnii*, this is the type species of *Zaglossus*.

Western long-tongued bat

Western long-tongued bat
Glossophaga morenoi

The western long-tongued bat (*Glossophaga morenoi*) is a species of bat in the family Phyllostomidae. It is endemic to Mexico.

Western lowland gorilla
Western lowland gorilla
Gorilla gorilla gorilla

The western lowland gorilla (*Gorilla gorilla gorilla*) is one of two Critically Endangered subspecies of the western gorilla (*Gorilla gorilla*) that lives in montane, primary and secondary forest and lowland swampland in central Africa in Angola,...

Western lowland olingo
Western lowland olingo
Bassaricyon medius

The western lowland olingo (*Bassaricyon medius*) is a species of olingo from Central and South America, where it is known from Panama and from Colombia and Ecuador west of the Andes.

Western mastiff bat
Western mastiff bat
Eumops perotis

The western mastiff bat (*Eumops perotis*), also known as the western bonneted bat, the greater mastiff bat, or the greater bonneted bat, is a member of the free-tailed bat family, Molossidae. It is found in the Western United States, Mexico and...

Western Moose
Western Moose
Alces alces andersoni

The Western moose (*Alces alces andersoni*) is a subspecies of moose. It is the second largest North American subspecies of moose, second to the Alaskan moose. This subspecies is prey to timber wolves, bears.

Western mouse
Western mouse
Pseudomys occidentalis

The western mouse or walyadji refers to *Pseudomys occidentalis*, a species of rodent in the family Muridae. Once widespread across a larger range, it has become restricted to around ten reserves of remnant bushland in Southwest Australia and...

Western naked-backed fruit bat
Western naked-backed fruit bat
Dobsonia peronii

The western naked-backed fruit bat (*Dobsonia peronii*) is a species of megabat in the family Pteropodidae. It is endemic to Indonesia. Its natural habitat is subtropical or tropical dry forests. It is threatened by habitat loss.

Western nectar bat
Western nectar bat
Lonchophylla hesperia

The western nectar bat (*Lonchophylla hesperia*) is a species of bat in the family Phyllostomidae. It is found in Ecuador and Peru. It is threatened by habitat loss.

Western nesomys
Western nesomys
Nesomys lambertoni

The western nesomys (*Nesomys lambertoni*), also known as the lowland red forest rat, is a species of rat endemic to Madagascar.
Western New Guinea mountain rat
Western New Guinea mountain rat
Rattus arrogans

The western New Guinea mountain rat (*Rattus arrogans*) is a species of rodent in the family Muridae. It is widespread in the mountains of central and western New Guinea.
Western pebble-mound mouse
Western pebble-mound mouse
Pseudomys chapmani

Western pebble-mound mouse or Ngadji, species *Pseudomys chapmani*, is a burrowing and mound building rodent in the family Muridae. They occur in the Pilbara, a remote region in the northwest of Australia.
Western purple-faced langur
Western purple-faced langur
Semnopithecus vetulus nestor

The western purple-faced langur (*Semnopithecus vetulus nestor*), also known as the north lowland wetzone purple-faced langur, is a subspecies of purple-faced langur endemic to Sri Lanka. It lives in the wet zone in western Sri Lanka around the...
Western Pygmy Possum
Western Pygmy Possum
Cercartetus concinnus

The western pygmy possum (*Cercartetus concinnus*), also known as the southwestern pygmy possum or the mundarda, is a small marsupial found in Australia. Genetic studies indicate its closest relative is probably the eastern pygmy possum, from...
Western Quoll
Western Quoll
Dasyurus geoffroii

The western quoll (*Dasyurus geoffroii*) is Western Australia's largest endemic mammalian carnivore. One of the many marsupial mammals native to Australia, it is also known as the chuditch. The species is currently classed as near-threatened.
Western Red Colobus
Western Red Colobus
Piliocolobus badius

The western red colobus (*Piliocolobus badius*), also known as the bay red colobus, rust red colobus or Upper Guinea red colobus, is a species of Old World monkey in West African forests from Senegal to Ghana. All other species of red colobuses...
Western red-backed vole
Western red-backed vole
Myodes californicus

The western red-backed vole (*Myodes californicus*) is a species of vole in the family Cricetidae. It is found in California and Oregon in the United States and lives mainly in coniferous forest. The body color is chestnut brown, or brown mixed...
Western ringtail possum
Western ringtail possum
Pseudocheirus occidentalis

The western ringtail possum or ngwayir refers to a species of possum, *Pseudocheirus occidentalis*, found in a small area of Southwest Australia. They are a cat-sized marsupial with a stocky build, dark greyish-brown fur, pale underparts and a

long...

Western rock elephant shrew
Western rock elephant shrew
Elephantulus rupestris

The western rock elephant shrew or western rock sengi (*Elephantulus rupestris*) is a species of elephant shrew in the family Macroscelididae. It is found in Namibia, South Africa, possibly Angola, and possibly Botswana. Its natural habitats are...

Western round-eared bat
Western round-eared bat
Lophostoma occidentale

The western round-eared bat (*Lophostoma occidentale*) is a bat species found only on the Pacific coast of northwestern Ecuador.

Western Saharan spiny mouse
Western Saharan spiny mouse
Acomys airensis

The Western Saharan spiny mouse or Air spiny mouse (*Acomys airensis*) is a species of small, insectivorous rodent in the family Muridae found arid regions of western Africa.

Western sheath-tailed bat
Western sheath-tailed bat
Paremballonura tiavato

The western sheath-tailed bat (*Paremballonura tiavato*) is a species of sac-winged bat (family Emballonuridae) found in Madagascar.

Western shrew mouse
Western shrew mouse
Pseudohydromys occidentalis

The western shrew mouse (*Pseudohydromys occidentalis*) is a species of rodent in the family Muridae. It is found in West Papua, Indonesia and Papua New Guinea.

Western small-footed bat
Western small-footed bat
Myotis ciliolabrum

The western small-footed bat (*Myotis ciliolabrum*), also known as the western small-footed myotis, is a species of vesper bat native to North America.

Western Spanish ibex
Western Spanish ibex
Capra pyrenaica victoriae

The Western Spanish ibex or Gredos ibex (*Capra pyrenaica victoriae*) is a subspecies of Iberian ibex native to Spain, in the Sierra de Gredos. It was later introduced to other sites in Spain (Las Batuecas, La Pedriza, Riaño) and to northern...

Western Spotted Skunk
Western Spotted Skunk
Spilogale gracilis

The western spotted skunk (*Spilogale gracilis*) is a spotted skunk of western North America.

Western sucker-footed bat
Western sucker-footed bat
Myzopoda schliemanni

The western sucker-footed bat (*Myzopoda schliemanni*) is a Malagasy bat. Little is known about its habits, but they are assumed to be similar to those of the

Madagascar sucker-footed bat.
Western tree hyrax
Western tree hyrax
Dendrohyrax dorsalis

The western tree hyrax (*Dendrohyrax dorsalis*), also called the western tree dassie or Beecroft's tree hyrax, is a species of tree hyrax within the family Procaviidae. It can be distinguished from other hyraxes by short coarse fur, presence of...

Western vlei rat
Western vlei rat
Otomys occidentalis

The western vlei rat (*Otomys occidentalis*) is a species of rodent in the family Muridae. It is found in Cameroon and Nigeria. Its natural habitats are subtropical or tropical high-altitude grassland and swamps. It is threatened by habitat loss.

Western water rat
Western water rat
Hydromys hussoni

The western water rat (*Hydromys hussoni*) is a semiaquatic species of rodent in the family Muridae. It is found in West Papua, Indonesia and Papua New Guinea. It is threatened by habitat loss.

Western white-eared giant rat
Western white-eared giant rat
Hyomys dammermani

The western white-eared giant rat (*Hyomys dammermani*) is a species of rodent in the family Muridae. It is found in West Papua, Indonesia and Papua New Guinea.

Western woolly lemur
Western woolly lemur
Avahi occidentalis

The western woolly lemur or western avahi (*Avahi occidentalis*) is a species of woolly lemur native to western Madagascar, where they live in dry deciduous forests. These nocturnal animals weigh 0.7-0.9 kg. It is a folivorous species. The western...

Western yellow bat
Western yellow bat
Lasiurus xanthinus

The western yellow bat (*Dasypterus xanthinus*) is a species of vesper bat. It is found in Mexico and the southwestern United States. This species roosts in trees such as *Populus fremontii*, *Platanus wrightii*, and *Quercus arizonica*. If available,...

Wetzel's climbing mouse
Wetzel's climbing mouse
Rhipidomys wetzeli

Wetzel's climbing mouse (*Rhipidomys wetzeli*) is a species of rodent in the family Cricetidae. It is endemic to Venezuela.

Weyland ringtail possum
Weyland ringtail possum
Pseudochirulus caroli

The Weyland ringtail possum (*Pseudochirulus caroli*) is a species of marsupial in the family Pseudocheiridae. It is endemic to the western Central Cordillera of Papua Province, Indonesia. *P. caroli* is also "known from four localities west of the...

Weyns's duiker

Weyns's duiker
Cephalophus weynsi

Weyns's duiker (*Cephalophus weynsi*) is a tiny antelope found in the Democratic Republic of Congo, Uganda and western Kenya. It is sometimes spelled "Weyn's", "Weyns", or "Weyns'" duiker. Weyns's duikers average about 33 lb (15 kg) in weight when...

Whiskered bat
Whiskered bat
Myotis mystacinus

The whiskered bat (*Myotis mystacinus*) is a small European bat with long fur. Although uncommon, *M. mystacinus* is often found around human habitation and around water; it is similar to Brandt's bat (*Myotis brandtii*), from which it was...

Whiskered flying squirrel
Whiskered flying squirrel
Petinomys genibarbis

The whiskered flying squirrel (*Petinomys genibarbis*) is a species of flying squirrel. It is found in Indonesia and Malaysia.

Whitaker's shrew
Whitaker's shrew
Crocidura whitakeri

Whitaker's shrew (*Crocidura whitakeri*) is a species of mammal in the family Soricidae. It is found in Western Sahara, Algeria, Morocco, Tunisia. Its natural habitats are subtropical or tropical dry shrubland, rocky and sandy coasts. It is a...

White marmoset
White marmoset
Mico leucippe

The white marmoset (*Mico leucippe*), or golden-white bare-ear marmoset, is a vulnerable species of marmoset, a small monkey endemic to the Amazon rainforest in Pará, Brazil.

White Rhinoceros
White Rhinoceros
Ceratotherium simum

The White rhino (*Ceratotherium simum*) is the second-largest mammal in the world. An alternative name for this species is the "Square-lipped rhinoceros", referring to the hairless, square upper lip of the animal. The word "rhino" derives from an...

White-ankled mouse
White-ankled mouse
Peromyscus pectoralis

The white-ankled mouse (*Peromyscus pectoralis*) is a species of rodent in the family Cricetidae. It is found in Mexico and in New Mexico, Oklahoma and Texas in the United States. The white-ankled mouse is commonly found in coexistence with the...

White-Beaked Dolphin
White-Beaked Dolphin
Lagenorhynchus albirostris

The white-beaked dolphin (*Lagenorhynchus albirostris*) is a marine mammal belonging to the family Delphinidae (oceanic dolphins) in the suborder Odontoceti (toothed whales).

White-bellied big-eared bat
White-bellied big-eared bat

Micronycteris minuta

The white-bellied big-eared bat (*Micronycteris minuta*) is a bat species from South and Central America, as well as Trinidad and Tobago in the Caribbean.

White-bellied duiker

White-bellied duiker

Cephalophus leucogaster

The white-bellied duiker (*Cephalophus leucogaster*) is a duiker found in central Africa. Little is known on the ecology of the species, and only some information on habitat and diet is available. Duikers are a fairly small species of antelope that...

White-bellied fat-tailed mouse opossum

White-bellied fat-tailed mouse opossum

Thylamys pallidior

The white-bellied fat-tailed mouse opossum (*Thylamys pallidior*) is a species of opossum in the family Didelphidae. It is found in Argentina, Bolivia, Chile and Peru.

White-bellied free-tailed bat

White-bellied free-tailed bat

Mops niveiventer

The white-bellied free-tailed bat (*Mops niveiventer*) is a species of bat in the family Molossidae. It is found in Angola, Democratic Republic of the Congo, and Zambia. Its natural habitats are subtropical or tropical moist lowland forests and...

White-bellied lesser house bat

White-bellied lesser house bat

Scotoecus albigula

The white-bellied lesser house bat (*Scotoecus albigula*) is a species of vesper bat. It can be found in Angola, Kenya, Malawi, Mozambique, Somalia, Uganda, and Zambia.

White-bellied Luzon tree rat

White-bellied Luzon tree rat

Carpomys phaeurus

The white-bellied Luzon tree rat (*Carpomys phaeurus*) is a species of rodent in the family Muridae. It is found only in Philippines. Its natural habitat is subtropical or tropical dry forests.

White-bellied mosaic-tailed rat

White-bellied mosaic-tailed rat

Melomys leucogaster

The white-bellied mosaic-tailed rat (*Melomys leucogaster*) is a species of rodent in the family Muridae. It is found in West Papua, Indonesia and Papua New Guinea.

White-bellied moss mouse

White-bellied moss mouse

Pseudohydromys sandrae

The white-bellied moss mouse, (*Pseudohydromys sandrae*), is a species of mouse endemic to Papua New Guinea. It was first described in 2009. The type specimen was an adult male found between 800 and 850 m, in Southern Highlands Province of Papua...

White-Bellied Musk Deer

White-Bellied Musk Deer

Moschus leucogaster

The white-bellied musk deer or Himalayan musk deer (*Moschus leucogaster*) is a musk deer species occurring in the Himalayas of Nepal, Bhutan, India, Pakistan and China. It is listed as endangered on the IUCN Red List because of overexploitation...

White-bellied nesomys

White-bellied nesomys

Nesomys audeberti

The white-bellied nesomys (*Nesomys audeberti*), also known as the lowland red forest rat, is a species of rat endemic to Madagascar.

White-bellied rat

White-bellied rat

Niviventer niviventer

The white-bellied rat (*Niviventer niviventer*) is a species of rodent in the family Muridae. It is found in India, Nepal, Bhutan and Pakistan.

White-bellied slender opossum

White-bellied slender opossum

Marmosops noctivagus

The white-bellied slender opossum (*Marmosops noctivagus*) is a species of opossum from South America. It is found in Bolivia, Brazil, Ecuador and Peru. Recent research suggests that *M. dorothea* is a synonym of *M. noctivagus*.

White-bellied spider monkey

White-bellied spider monkey

Ateles belzebuth

The white-bellied spider monkey (*Ateles belzebuth*), also known as the white-fronted or long-haired spider monkey, is an endangered species of spider monkey, a type of New World monkey. It is found in the north-western Amazon in Colombia,...

White-bellied woolly mouse opossum

White-bellied woolly mouse opossum

Marmosa constantiae

The white-bellied woolly mouse opossum (*Marmosa constantiae*) is a small pouchless marsupial of the family Didelphidae. It was formerly assigned to the genus *Micoureus*, which was made a subgenus of *Marmosa* in 2009. The specific epithet was given...

White-bellied yellow bat

White-bellied yellow bat

Scotophilus leucogaster

The white-bellied yellow bat (*Scotophilus leucogaster*) or white-bellied house bat, is a species of vesper bat in the genus *Scotophilus*, the house bats. It can be found in Angola, Benin, Botswana, Burkina Faso, Cameroon, Central African Republic,...

White-Cheeked Gibbon

White-Cheeked Gibbon

Genus *Nomascus*

The northern white-cheeked gibbon (*Nomascus leucogenys*) is a Critically Endangered species of gibbon native to South East Asia. It is closely related to the southern white-cheeked gibbon (*Nomascus siki*), with which it was previously considered...

White-cheeked macaque

White-cheeked macaque

Macaca leucogenys

The white-cheeked macaque (*Macaca leucogenys*) is a species of macaque found only in Mêdog County in southeastern Tibet and Arunachal Pradesh in northeastern India.

The white-cheeked macaque lives in forest habitats, from tropical forests to...

White-Cheeked Spider Monkey

White-Cheeked Spider Monkey

Ateles marginatus

The white-cheeked spider monkey (*Ateles marginatus*) is a species of spider monkey, a type of New World monkey, endemic to Brazil. It moves around the forest canopy in small family groups of two to four, part of larger groups of a few dozen...

White-coated titi

White-coated titi

Plecturocebus pallescens

The white-coated titi (*Plecturocebus pallescens*) is a species of titi, a type of New World monkey, from South America. It is found in Bolivia, Brazil, and Paraguay.

White-collared fruit bat

White-collared fruit bat

Megaerops wetmorei

The white-collared fruit bat (*Megaerops wetmorei*) is a species of megabat found in Southeast Asia.

White-eared cotton rat

White-eared cotton rat

Sigmodon leucotis

The white-eared cotton rat (*Sigmodon leucotis*) is a species of rodent in the family Cricetidae. It is found only in Mexico.

White-Eared Opossum

White-Eared Opossum

Didelphis albiventris

The White-eared opossum (*Didelphis albiventris*) is a marsupial found in South America. It is a terrestrial and, sometimes, arboreal animal that lives in a wide range of habitats.

White-eared pocket mouse

White-eared pocket mouse

Perognathus alticollis

The white-eared pocket mouse (*Perognathus alticollis*) is a species of rodent in the family Heteromyidae. It is endemic to the San Bernardino Mountains and the Tehachapi Mountains of southern California in the United States. There are two...

White-eared titi

White-eared titi

Plecturocebus donacophilus

The white-eared titi, *Plecturocebus donacophilus*, also known as the Bolivian titi or Bolivian gray titi, is a species of titi, a type of New World monkey, from eastern Bolivia and an area of western Brazil. The species has a range that extends...

White-Faced Saki

White-Faced Saki

Pithecia pithecia

The white-faced saki (*Pithecia pithecia*), called the Guianan saki and the golden-faced saki, is a species of the New World saki monkey. They can be found in Brazil, French Guiana, Guyana, Suriname and Venezuela. This species lives in the...

White-faced spiny tree-rat

White-faced spiny tree-rat

Echymys chrysurus

The white-faced spiny tree-rat, *Echimys chrysurus*, is a spiny rat species from South America. It is found in Brazil, French Guiana, Guyana and Suriname. The etymology of the species name derives from the two ancient greek words χρυσός (khrusós),...

White-footed climbing mouse

White-footed climbing mouse

Rhipidomys leucodactylus

The white-footed climbing mouse (*Rhipidomys leucodactylus*) is a species of South American rodent found in Bolivia, Brazil, Ecuador, French Guiana, Guyana, Peru, Suriname and Venezuela. It is the type species of the genus and the type location...

White-footed dunnart

White-footed dunnart

Sminthopsis leucopus

The white-footed dunnart (*Sminthopsis leucopus*) is a marsupial that occurs on Tasmania and mainland Australia. It occurs along the coast and in inner Gippsland and Alpine areas up to 400 metres near Narbethong. In southern New South Wales, the...

White-footed fox

White-footed fox

Vulpes vulpes pusilla

The white-footed fox (*Vulpes vulpes pusilla*), also known as the desert fox, is a small, Asiatic subspecies of red fox which occurs throughout most of northwestern Indian subcontinent, Pakistan's desert districts from Rawalpindi to Rajasthan and...

White-Footed Mouse

White-Footed Mouse

Peromyscus leucopus

The white-footed mouse (*Peromyscus leucopus*) is a rodent native to North America from Ontario, Quebec, Labrador, and the Maritime Provinces (excluding the island of Newfoundland) to the southwestern United States and Mexico. In the Maritimes,...

White-footed saki

White-footed saki

Pithecia albicans

The white-footed saki, buffy saki or white saki (*Pithecia albicans*) is a species of saki monkey, a type of New World monkey, endemic to western Brazil south of the Amazon.

White-footed sportive lemur

White-footed sportive lemur

Lepilemur leucopus

The white-footed sportive lemur, white-footed weasel lemur, or dry-bush weasel lemur (*Lepilemur leucopus*) is a species of lemur in the family Lepilemuridae, the sportive lemurs. It is similar in appearance to other lemurs in the family, with a...

White-footed tamarin

White-footed tamarin

Saguinus leucopus

The white-footed tamarin (*Saguinus leucopus*) is a tamarin species endemic to Colombia. It is a silvery brown colour with pale streaks and russet underparts, and is very similar in appearance to the cotton-top tamarin, from which it is separated...

White-footed vole

White-footed vole
Arborimus albipes

The white-footed vole (*Arborimus albipes*) is a species of rodent in the family Cricetidae. It is found only in the United States. Its natural habitat is temperate forests.

White-Fronted Brown Lemur
White-Fronted Brown Lemur
Eulemur albifrons

The white-headed lemur (*Eulemur albifrons*), also known as the white-headed brown lemur, white-fronted brown lemur, or white-fronted lemur, is a species of primate in the family Lemnidae. It is only found in north-eastern Madagascar. It is...

White-Fronted Capuchin
White-Fronted Capuchin
Cebus albifrons

Humboldt's white-fronted capuchin (*Cebus albifrons*) is a species of gracile capuchin monkey. It is found in Colombia, Venezuela, Brazil, and potentially the island of Trinidad.

White-fronted surili
White-fronted surili
Presbytis frontata

The white-fronted surili (*Presbytis frontata*) is a species of primate in the family Cercopithecidae, the Old World monkeys. It is endemic to the large international island of Borneo, in Indonesia, Malaysia, and possibly Brunei. Its body is mainly...

White-Headed Capuchin
White-Headed Capuchin
Cebus capucinus

The Colombian white-faced capuchin (*Cebus capucinus*), also known as the Colombian white-headed capuchin or Colombian white-throated capuchin, is a medium-sized New World monkey of the family Cebidae, subfamily Cebinae. It is native to the...

White-headed langur
White-headed langur
Trachypithecus poliocephalus

The white-headed langur (*Trachypithecus leucocephalus*) is a critically endangered species of langur endemic to Guangxi, China.

White-headed marmoset
White-headed marmoset
Callithrix geoffroyi

The white-headed marmoset (*Callithrix geoffroyi*), also known as the tufted-ear marmoset, Geoffroy's marmoset, or Geoffrey's marmoset, is a marmoset endemic to forests in eastern Brazil, where it is native to Bahia, Espírito Santo, and Minas...

White-legged duiker
White-legged duiker
Cephalophus crusalbum

The white-legged duiker (*Cephalophus crusalbum*) is a medium-sized antelope species from the subfamily of duikers (Cephalophinae) within the family of bovids (Bovidae). It is native to Gabon and the Republic of the Congo. It was described as...

White-lined broad-nosed bat
White-lined broad-nosed bat

Platyrrhinus lineatus

The white-lined broad-nosed bat (*Platyrrhinus lineatus*) is a bat species from South America. It is found in southern and eastern Brazil, Paraguay, Uruguay, northern Argentina, Bolivia, Peru, Ecuador, Colombia, French Guiana and Suriname.

White-Lipped Peccary

White-Lipped Peccary

Tayassu pecari

The white-lipped peccary (*Tayassu pecari*) is a species of peccary found in Central and South America and the only member of the genus *Tayassu*. Multiple subspecies have been identified. White-lipped peccaries are similar in appearance to pigs,...

White-lipped tamarin

White-lipped tamarin

Saguinus labiatus

The white-lipped tamarin (*Saguinus labiatus*), also known as the red-bellied tamarin, is a tamarin which lives in the Amazon area of Brazil and Bolivia. The red belly of these New World monkeys is its most remarkable outward characteristic....

White-mantled tamarin

White-mantled tamarin

Leontocebus weddelli melanoleucus

The white-mantled tamarin, *Saguinus weddelli melanoleucus*, is a subspecies of Weddell's saddle-back tamarin, a tamarin monkey from South America. It is found in Brazil, between Rio Jurua and Rio Tarauacá.

White-Nosed Coati

White-Nosed Coati

Nasua narica

The White-nosed coati (*Nasua narica*) is a species of coati and a member of the family Procyonidae (raccoons and their relatives). Local Spanish names for this animal include pizote, Mexican Bin-dog', antoon, and tejón, depending upon the region.

White-nosed saki

White-nosed saki

Chiropotes albinasus

The white-nosed saki (*Chiropotes albinasus*) is a species of bearded saki, a type of New World monkey, endemic to the south-central Amazon rainforest in Brazil. Both its scientific and common name were caused by the authors working from dead...

White-Sided Jackrabbit

White-Sided Jackrabbit

Lepus callotis

The White-sided jackrabbit (*Lepus callotis*) is a jackrabbit that lives in a limited range in North America. It is considered threatened in New Mexico because its numbers are declining in recent years.

White-spined Atlantic spiny rat

White-spined Atlantic spiny rat

Trinomys albispinus

The white-spined Atlantic spiny rat (*Trinomys albispinus*) is a spiny rat species endemic to Brazil.

White-striped dorcopsis

White-striped dorcopsis

Dorcopsis hageni

The white-striped dorcopsis or greater forest wallaby (*Dorcopsis hageni*) is a

species of marsupial in the family Macropodidae. It is found in the northern part of West Papua, Indonesia and Papua New Guinea. It is a common species in suitable...

White-striped free-tailed bat
White-striped free-tailed bat
Austronomus australis

The white-striped free-tailed bat (*Austronomus australis*) is a species of bat in the family Molossidae. Its echolocation calls are audible to humans, which is a characteristic found in only a few microbat species. The species was formerly...

White-Tailed Antelope Squirrel
White-Tailed Antelope Squirrel
Ammospermophilus leucurus

The white-tailed antelope squirrel (*Ammospermophilus leucurus*) is a diurnal species of ground squirrel, scientifically classified in the order Rodentia and family Sciuridae, found in arid regions of the southwestern United States and the Baja...

White-tailed antsangy
White-tailed antsangy
Brachytarsomys albicauda

The white-tailed antsangy (*Brachytarsomys albicauda*) is a species of rodent in the family Nesomyidae. It is found only in Madagascar.

White-Tailed Deer
White-Tailed Deer
Odocoileus virginianus

The White-tailed deer (*Odocoileus virginianus*) is a medium-sized deer native to the Americas where it is the most widely distributed wild ungulate. Texas is home to the most White-tailed deer of any U.S. state or Canadian province, with an...

White-tailed dunnart
White-tailed dunnart
Sminthopsis granulipes

The white-tailed dunnart (*Sminthopsis granulipes*), also known as the ash-grey dunnart, is a dunnart native to Australia.

White-Tailed Jackrabbit
White-Tailed Jackrabbit
Lepus townsendii

The White-tailed jackrabbit (*Lepus townsendii*) is a large hare found in western North America. Whitetails are often seen in urban and suburban parks in Western Canada. People often come across bunnies alone during the day in spring and mistakenly...

White-tailed mole
White-tailed mole
Parascaptor leucura

The white-tailed mole (*Parascaptor leucura*) is a species of mammal in the family Talpidae. It is found in Bangladesh, China, India, and Myanmar. It is the only species in the genus *Parascaptor*.

White-Tailed Mongoose
White-Tailed Mongoose
Ichneumia albicauda

The white-tailed mongoose (*Ichneumia albicauda*) is a species in the mongoose family Herpestidae. It is the only member of the genus *Ichneumia*.

White-tailed mountain vole

White-tailed mountain vole
Alticola albicauda

The white-tailed mountain vole (*Alticola albicauda*) is a species of vole in the family Cricetidae. It is found in India and Pakistan.

White-tailed olalla rat
White-tailed olalla rat
Olallamys albicauda

The white-tailed olalla rat (*Olallamys albicauda*) is a species of rodent in the family Echimyidae. It is a medium-sized rat with a reddish-brown dorsum, a whitish venter, and a tail characterized by a whitish tip. It is endemic to Colombia. Its...

White-Tailed Prairie Dog
White-Tailed Prairie Dog
Cynomys leucurus

The white-tailed prairie dog (*Cynomys leucurus*) is found in western Wyoming and western Colorado with small areas in eastern Utah and southern Montana. The largest populations are in Wyoming where they are known colloquially as "chisellers".

This...
White-tailed rat
White-tailed rat
Mystromys albicaudatus

The white-tailed rat (*Mystromys albicaudatus*) also known as the white-tailed mouse, is the only member of the subfamily Mystromyinae in the family Nesomyidae. This species is sometimes placed in the subfamily Cricetinae due to similarities in...

White-tailed titi
White-tailed titi
Plecturocebus discolor

The white-tailed titi, *Plecturocebus discolor*, is a species of titi, a type of New World monkey, from South America. It is found in Colombia, Ecuador and Peru. It was described in 1848 as *Callicebus discolor*.

White-thighed surili
White-thighed surili
Presbytis siamensis

The white-thighed surili (*Presbytis siamensis*) is a species of primate from the family of old world monkeys (Cercopithecidae). This species lives arboreal amongst the sub-montane forests. It is endemic to the Thai-Malay Peninsula, the Riau...

White-throated grass mouse
White-throated grass mouse
Akodon simulator

The white-throated or grey-bellied grass mouse (*Akodon simulator*) is a species of rodent in the family Cricetidae. It is found in Argentina and Bolivia.

White-throated guenon
White-throated guenon
Cercopithecus erythrogaster

The white-throated guenon (*Cercopithecus erythrogaster*), also known as the red-bellied monkey and the red-bellied guenon, is a diurnal primate that lives on trees of rainforests or tropical areas of Nigeria and Benin. The white-throated guenon is...

White-throated round-eared bat
White-throated round-eared bat
Lophostoma silvicolium

The white-throated round-eared bat (*Lophostoma silvicolum*) is a South and Central American bat species found from Honduras to Bolivia, Paraguay and Brazil. It creates roosts inside the nests of the termite, *Nasutitermes corniger*. It thrives on a...

White-tipped Oldfield mouse
White-tipped Oldfield mouse
Thomasomys caudivarius

The white-tipped Oldfield mouse (*Thomasomys caudivarius*) is a species of rodent in the family Cricetidae. It is found in the Andes from central Ecuador to northern Peru, at elevations from 2500 to 3350 m, where it lives in montane forest.

White-tipped tufted-tailed rat
White-tipped tufted-tailed rat
Eliurus penicillatus

The white-tipped tufted-tailed rat (*Eliurus penicillatus*) is a rodent endemic to Madagascar. It is known from only two specimens, one collected from Ampitambe forest in 1895 or 1896 and the second in 2000. It is listed by the International Union...

White-toothed brush mouse
White-toothed brush mouse
Coccymys albidens

The white-toothed brush mouse (*Coccymys albidens*), also known as the white-toothed melomys or white-toothed mouse, is a species of rodent in the family Muridae. It is found only in West Papua, Indonesia. Its natural habitats are subtropical or...

White-toothed tuco-tuco
White-toothed tuco-tuco
Ctenomys leucodon

The white-toothed tuco-tuco (*Ctenomys leucodon*) is a species of rodent in the family Ctenomyidae. It is found in Bolivia and Peru.

White-winged dog-like bat
White-winged dog-like bat
Peropteryx leucoptera

The white-winged dog-like bat (*Peropteryx leucoptera*) is a bat species from South America. It is found in northern Brazil, southeastern Colombia, Ecuador, French Guiana, Guyana, eastern Peru, Surinam and Venezuela.

White-winged flying fox
White-winged flying fox
Desmalopex leucopterus

The white-winged flying fox (*Desmalopex leucopterus*), also known as the mottled-winged flying fox is a species of bat in the family Pteropodidae. They are endemic to the Philippines. Their natural habitat is subtropical or tropical dry forests....

White-winged serotine
White-winged serotine
Neoromicia tenuipinnis

The white-winged serotine (*Pseudoromicia tenuipinnis*) is a species of vesper bat. It can be found in the following countries: Angola, Burundi, Cameroon, Republic of the Congo, Democratic Republic of the Congo, Ivory Coast, Equatorial Guinea,...

White-Winged Vampire Bat
White-Winged Vampire Bat
Diaemus youngi

The white-winged vampire bat (*Diaemus youngi*), a species of vampire bat, is the only member of the genus *Diaemus*. They are found from Mexico to northern Argentina and are present on the islands of Trinidad and Margarita.

Whitehead's spiny rat

Whitehead's spiny rat

Maxomys whiteheadi

Whitehead's spiny rat (*Maxomys whiteheadi*) is a species of rodent in the family Muridae. It is found in Indonesia, Malaysia, and Thailand.

Whitehead's woolly bat

Whitehead's woolly bat

Kerivoula whiteheadi

Whitehead's woolly bat (*Kerivoula whiteheadi*) is a species of vesper bat in the family Vespertilionidae. It is found in Brunei, Indonesia, Malaysia, the Philippines, and Thailand.

Whitish dwarf squirrel

Whitish dwarf squirrel

Prosciurillus leucomus

The whitish dwarf squirrel (*Prosciurillus leucomus*) is a species of rodent in the family Sciuridae. It is endemic to Indonesia, where it is found on Sulawesi, Buton Island, Muna Island, Kabeana Island, and adjacent islands to the southeast of...

Wide-toothed hutia

Wide-toothed hutia

Plagiodontia araeum

Hyperplagiodontia, rarely called the wide-toothed hutia, is an extinct genus of hutia which contains a single species, *Hyperplagiodontia araeum*. The species was originally described as a member of the genus *Plagiodontia* along with the extant...

Wied's marmoset

Wied's marmoset

Callithrix kuhlii

Wied's marmoset (*Callithrix kuhlii*), also known as Wied's black-tufted-ear marmoset, is a New World monkey that lives in tropical and subtropical forests of eastern Brazil. Unlike other marmosets, Wied's marmoset lives in groups consisting of 4...

Wild Bactrian Camel

Wild Bactrian Camel

Camelus ferus

The Wild Bactrian camel is a critically endangered species of camel living in parts of northern China and southern Mongolia. Until recently, wild Bactrian camels were thought to have descended from domesticated Bactrian camels that became feral...

Wild Boar

Wild Boar

Sus scrofa

The Wild boar (*Sus scrofa*) is a suid native to much of Eurasia and North Africa. The species is now one of the widest-ranging mammals in the world, as well as the most widespread suiform. It has been assessed as the least concern on the IUCN Red...

Wild Goat

Wild Goat

Capra aegagrus

The Wild goat (*Capra aegagrus*) is a nearly threatened goat species native to Asia. It is one of the ancestors of the domestic goat.

Wild Yak
Wild Yak
Bos mutus

The Wild yak (*Bos mutus*) is a large, wild cattle native to the Himalayas. It is the ancestor of the domestic yak (*Bos grunniens*). Wild yaks are among the largest bovids and are second only to the gaur in shoulder height. They are also the largest...

Wildcat
Wildcat
Felis silvestris

The wildcat is a species complex comprising two small wild cat species: the European wildcat (*Felis silvestris*) and the African wildcat (*F. lybica*). The association of African wildcats and humans appears to have developed along with the...

Williams' jerboa
Williams' jerboa
Allactaga williamsi

Williams's jerboa (*Allactaga williamsi*) is a species of jerboas native to Afghanistan, Armenia, Azerbaijan, Iran and Turkey.

Williamson's mouse-deer
Williamson's mouse-deer
Tragulus williamsoni

Williamson's mouse-deer (*Tragulus williamsoni*) is a species of even-toed ungulate in the family Tragulidae. It is found in Thailand, and possibly in China. The species is named after the collector Walter James Franklin Williamson.

Wilson's spiny mouse
Wilson's spiny mouse
Acomys wilsoni

Wilson's spiny mouse (*Acomys wilsoni*) is a species of rodent in the family Muridae. It is found in Ethiopia, Kenya, Somalia, South Sudan, Tanzania, and Uganda. Its natural habitats are dry savanna, subtropical or tropical dry shrubland, and rocky...

Wimmer's shrew
Wimmer's shrew
Crocidura wimmeri

Wimmer's shrew (*Crocidura wimmeri*) is a white-toothed shrew found only in Côte d'Ivoire. It is listed as a critically endangered species due to habitat loss and a restricted range. Shrew species assignment is a challenge due to their strong...

Winkelmann's mouse
Winkelmann's mouse
Peromyscus winkelmanni

Winkelmann's mouse (*Peromyscus winkelmanni*) is a species of rodent in the family Cricetidae found only in Mexico, and is named for John R. Winkelmann, who collected the first specimens.

Wisent
Wisent
Bison bonasus

The wisent (*Bison bonasus*) or the European bison is a European species of bison. These animals were hunted to extinction in the wild in the early 20th century. Some of them were kept alive in captivity, and have since been reintroduced into...
Woermann's bat

Woermann's bat
Megaloglossus woermanni

Woermann's bat or Woermann's fruit bat (*Megaloglossus woermanni*) is a species of megabat in the family Pteropodidae. It is monotypic within the genus *Megaloglossus*. It is found in Angola, Benin, Cameroon, Central African Republic, Republic of...

Wolf's Mona Monkey
Wolf's Mona Monkey
Cercopithecus wolfi

Wolf's mona monkey (*Cercopithecus wolfi*) is a colorful Old World monkey from central Africa. It was first described from a living specimen in the Zoological Garden at Dresden. It was brought in 1887 by Dr Ludwig Wolf from somewhere in central...

Wolffsohn's leaf-eared mouse
Wolffsohn's leaf-eared mouse
Phyllotis wolffsohni

Wolffsohn's leaf-eared mouse (*Phyllotis wolffsohni*) is a species of rodent in the family Cricetidae. It is found only in Bolivia.

Wolffsohn's viscacha
Wolffsohn's viscacha
Lagidium wolffsohni

Wolffsohn's viscacha (*Lagidium wolffsohni*) is a rare species of rodent in the family Chinchillidae. This species occurs in southwestern Argentina and adjacent Chile.

Wollaston's roundleaf bat
Wollaston's roundleaf bat
Hipposideros wollastoni

Wollaston's roundleaf bat (*Hipposideros wollastoni*) is a species of bat in the family Hipposideridae. It is found in West Papua, Indonesia and Papua New Guinea. It was named after the explorer Sandy Wollaston.

Wolverine
Wolverine
Gulo gulo

The wolverine (*Gulo gulo*) is the largest land-dwelling species of the family Mustelidae. It is a muscular carnivore and a solitary animal. The wolverine has a reputation for ferocity and strength out of proportion to its size, with the documented...

Wondiwoi Tree-Kangaroo
Wondiwoi Tree-Kangaroo
Dendrolagus mayri

The Wondiwoi tree-kangaroo (*Dendrolagus mayri*) is an elusive and rare bear-like mammal that was considered extinct until rediscovery in 2018. It is a species of tree-kangaroo, a group of long-tailed animals native to Australia and New Guinea that...

Wongai ningau
Wongai ningau
Ningau ridei

The Wongai ningau, a common name of *Ningau ridei*, is a tiny carnivorous marsupial native to the arid open grasslands of inland Australia. Their diet is mainly small insects, and occasionally larger prey such as spiders, grasshoppers and...

Wood bison
Wood bison

Bison bison athabasca

The wood bison (*Bison bison athabasca*) or mountain bison (often called the wood buffalo or mountain buffalo), is a distinct northern subspecies or ecotype of the American bison. Its original range included much of the boreal forest regions of...

Wood lemming

Wood lemming

Myopus schisticolor

The wood lemming (*Myopus schisticolor*) is a species of rodent in the family Cricetidae. It belongs to the rodent subfamily Arvicolinae, so is a relative of the voles, lemmings, and muskrats. It is found in the taiga biome of China, Estonia,...

Wood Mouse

Wood Mouse

Apodemus sylvaticus

The Wood mouse (*Apodemus sylvaticus*) is a murid rodent native to Europe and northwestern Africa. It is found across most of Europe and is a very common and widespread species, is commensal with people, and is sometimes considered a pest.

The Wood...

Wood sprite gracile opossum

Wood sprite gracile opossum

Gracilinanus dryas

The wood sprite gracile opossum (*Gracilinanus dryas*) is a mammal. It is a species of opossum in the family Didelphidae. It is found in Colombia and Venezuela. Its natural habitat is subtropical or tropical moist lowland forests. It is threatened...

Wood's slit-faced bat

Wood's slit-faced bat

Nycteris woodi

Wood's slit-faced bat (*Nycteris woodi*) is a species of slit-faced bat that lives in the dry savanna regions of Southern Africa. Its numbers are declining due to habitat loss from logging and farming, pesticide use, and the decline of baobab trees...

Woodford's fruit bat

Woodford's fruit bat

Melonycteris woodfordi

Woodford's fruit bat (*Melonycteris woodfordi*), also known as the orange fruit bat, is a species of megabat in the family Pteropodidae. It is endemic to the Solomon Islands.

Woodland Dormouse

Woodland Dormouse

Graphiurus murinus

The woodland dormouse (*Graphiurus murinus*) is a species of rodent in the family Gliridae. It is native to southern and eastern Africa and is also known as the African dormouse, African dwarf dormouse, African pygmy dormouse, or colloquially as...

Woodland Jumping Mouse

Woodland Jumping Mouse

Napaeozapus insignis

The woodland jumping mouse (*Napaeozapus insignis*) is a species of jumping mouse found in North America. It can jump up to 3 m (9.8 ft) using its extremely strong feet and long tail.

Woodland Oldfield mouse

Woodland Oldfield mouse
Thomasomys hylophilus

The Woodland Oldfield mouse (*Thomasomys hylophilus*) is a rare species of rodent found in forests of South America.

Woodland thicket rat
Woodland thicket rat
Grammomys dolichurus

The woodland thicket rat (*Grammomys dolichurus*) is a species of rodent in the family Muridae. It is found in Angola, Burundi, Democratic Republic of the Congo, Kenya, Malawi, Mozambique, Rwanda, South Africa, South Sudan, Eswatini, Tanzania,...

Woodland Vole
Woodland Vole
Microtus pinetorum

The Woodland vole (*Microtus pinetorum*) is a small rodent found in North America. It is a relative of lemmings and hamsters but with a stouter body and a longer, hairy tail. It is also known as the Pine vole.

Woodlark cuscus
Woodlark cuscus
Phalanger lullulae

The Woodlark cuscus (*Phalanger lullulae*) is a species of marsupial in the family Phalangeridae endemic to Papua New Guinea, specifically on Madau and Woodlark Island, a part of the Milne Bay Province of Papua New Guinea. It happens to be the...

Woolley's false antechinus
Woolley's false antechinus
Pseudantechinus woolleyae

Woolley's false antechinus (*Pseudantechinus woolleyae*), also known as Woolley's pseudantechinus, is a species of small carnivorous marsupial belonging to the family Dasyuridae. It is found in the Australian state of Western Australia, primarily...

Woolley's moss mouse
Woolley's moss mouse
Pseudohydromys patriciae

Woolley's moss mouse, (*Pseudohydromys patriciae*), is a species of mouse endemic to Papua New Guinea. It was first described in 2009.

Woolley's three-striped dasyure
Woolley's three-striped dasyure
Myoictis leucura

Woolley's three-striped dasyure (*Myoictis leucura*) is a member of the order Dasyuromorphia. This marsupial carnivore lives in Papua New Guinea. This species inhabits primary montane forest. Like its congeners, it appears to be diurnal and largely...

Woolly dormouse
Woolly dormouse
Dryomys laniger

The woolly dormouse (*Dryomys laniger*) is a species of rodent in the family Gliridae endemic to Turkey.

Woolly Flying Squirrel
Woolly Flying Squirrel
Eupetaurus cinereus

Eupetaurus is a genus of rodent in the family Sciuridae. Members of this genus are known as woolly flying squirrels. They are large to very large flying squirrels found in the highest rocky cliffs near the treeline of the Himalayas and Tibetan...
Woolly giant rat
Woolly giant rat
Kunsia tomentosus

The woolly giant rat (*Kunsia tomentosus*) is a species of large burrowing rodent native to South America. No subspecies are currently recognised. It is the only member of the genus *Kunsia*.
Woolly hare
Woolly hare
Lepus oiostolus

The woolly hare (*Lepus oiostolus*) is a species of mammal in the family Leporidae. It is found in western and central China, northern India, and Nepal, where its typical habitat is montane grassland. It has a wide range and is present in some...
Woolly horseshoe bat
Woolly horseshoe bat
Rhinolophus luctus

The great woolly horseshoe bat (*Rhinolophus luctus*) is a species of bat in the family Rhinolophidae. It is endemic to Indonesia. The northern woolly horseshoe bat (*R. perniger*), Malaysian woolly horseshoe bat (*R. morio*), and Andersen's woolly...
Woolly mouse opossum
Woolly mouse opossum
Marmosa demerarae

The woolly mouse opossum or long-furred woolly mouse opossum (*Marmosa demerarae*), known locally as the *cuíca*, is a South American marsupial of the family Didelphidae. Its range includes central Colombia, Venezuela, French Guiana, Guyana, ...
Woolly-headed spiny tree-rat
Woolly-headed spiny tree-rat
Mesomys leniceps

The woolly-headed spiny tree-rat (*Mesomys leniceps*) is a species of rodent in the family Echimyidae. It is endemic to Peru. The etymology of the species name corresponds to the Latin word *leniceps* constructed from *lēnis* meaning calm, gentle, and...
Woosnam's broad-headed mouse
Woosnam's broad-headed mouse
Zelotomys woosnami

Woosnam's broad-headed mouse or Woosnam's zelotomys (*Zelotomys woosnami*) is a species of rodent in the family Muridae. It is found in Angola, Botswana, Namibia, and South Africa. Its natural habitat is dry savanna.
Woosnam's brush-furred rat
Woosnam's brush-furred rat
Lophuromys woosnami

The Woosnam's brush-furred rat (*Lophuromys woosnami*) is a species of rodent in the family Muridae. It is found in Burundi, the Democratic Republic of the Congo, Rwanda, and Uganda. Its natural habitat types include mountain forest clearings and...
Woylie
Woylie

Bettongia penicillata

The woylie or brush-tailed bettong (*Bettongia penicillata*) is an extremely rare, small marsupial, belonging to the genus *Bettongia*, that is endemic to Australia. There are two subspecies: *B. p. ogilbyi*, and the now extinct *B. p. penicillata*.

Wrangel Island lemming
Wrangel Island lemming
Lemmus portenkoi

The East Siberian lemming, Eastern Siberian brown lemming, or Ognev's lemming (*Lemmus paulus*) is a species of lemming endemic to Russia, where it has a disjunct distribution throughout parts of Siberia east of the Verkhoyansk Range.

Wrangel lemming
Wrangel lemming
Dicrostonyx vinogradovi

The northern collared lemming or Nearctic collared lemming (*Dicrostonyx groenlandicus*), sometimes called the Peary Land collared lemming in Canada, is a small lemming found in Arctic North America and Wrangel Island. At one time, it was...

Wright's sportive lemur
Wright's sportive lemur
Lepilemur wrightae

Wright's sportive lemur (*Lepilemur wrightae*), or the Kalambatrietra sportive lemur, is a sportive lemur endemic to Madagascar. It is one of the larger sportive lemurs with a total length of about 52 to 64 cm (20 to 25 in), of which 24–27 cm...

Wrinkle-faced bat
Wrinkle-faced bat
Centurio senex

The wrinkle-faced bat (*Centurio senex*) is a species of bat in the family Phyllostomidae and the only identified member of the genus *Centurio*. This bat is found in various countries in and around Central America. It eats fruit but is not...

Wrinkle-lipped free-tailed bat
Wrinkle-lipped free-tailed bat
Chaerephon plicatus

The wrinkle-lipped free-tailed bat (*Chaerephon plicatus*) is a species of bat in the family Molossidae. It is found in Bangladesh, Bhutan, Cambodia, China, Cocos (Keeling) Islands, India, Indonesia, Laos, Malaysia, Myanmar, Nepal, the Philippines,...

Wroughton's free-tailed bat
Wroughton's free-tailed bat
Otomops wroughtoni

Wroughton's free-tailed bat (*Otomops wroughtoni*) is a free-tailed bat formerly considered to be confined to the Western Ghats area of India, though it has also recently been discovered in northeast India and in a remote part of Cambodia. It is...

Wyoming ground squirrel
Wyoming ground squirrel
Urocitellus elegans

The Wyoming ground squirrel (*Urocitellus elegans*) is a species of rodent in the family Sciuridae. It is endemic to the Northwestern United States.

Wyoming pocket gopher
Wyoming pocket gopher

Thomomys clusius

The Wyoming pocket gopher (*Thomomys clusius*) is a species of gopher that is endemic to the United States. Between 1915 and 1979, it was generally considered to be a subspecies of the northern pocket gopher.

Xanthippe's shrew

Xanthippe's shrew

Crocidura xantippe

Xanthippe's shrew (*Crocidura xantippe*) is a species of mammal in the family Soricidae. It is found in Kenya and Tanzania. Its natural habitats are dry savanna and subtropical or tropical dry shrubland.

Yakushima macaque

Yakushima macaque

Macaca fuscata yakui

The Yakushima macaque (*Macaca fuscata yakui*) is a subspecies of Japanese macaque that is indigenous to Yakushima Island (Kagoshima Prefecture).It is also known as the Yaku macaque.

Yalden's rat

Yalden's rat

Desmomys yaldeni

Yalden's rat (*Desmomys yaldeni*) is a species of rodent in the family Muridae, found only in Ethiopia, where its natural habitat is the montane forests. It is threatened by habitat loss.The rat got its name in honour of Derek Yalden, a zoologist.

Yamdena mosaic-tailed rat

Yamdena mosaic-tailed rat

Melomys cooperae

The Yamdena mosaic-tailed rat (*Melomys cooperae*) is a species of rat belonging to the family Muridae. It was first described in 1995 from specimens found on Yamdena Island in Indonesia. No other examples of this species have been found; as a...

Yanbaru whiskered bat

Yanbaru whiskered bat

Myotis yanbarensis

The Yanbaru whiskered bat (*Myotis yanbarensis*) is a species of vesper bat in the genus *Myotis*.It is known only from three islands of the Ryukyu Archipelago, south of Japan, Okinawa (where the Yanbaru forest is situated), Amami Ōshima, and...

Yangtze Finless Porpoise

Yangtze Finless Porpoise

Neophocaena asiaeorientalis

The Yangtze finless porpoise (*Neophocaena asiaeorientalis*) is a species of toothed whale found only in the Yangtze River in China. It is the country's only known freshwater cetacean following the possible extinction of the baiji. The Yangtze...

Yankari shrew

Yankari shrew

Crocidura yankariensis

The Yankari shrew (*Crocidura yankariensis*) is a species of mammal in the family Soricidae. It is found in Cameroon, Central African Republic, Chad, Ethiopia, Kenya, Nigeria, Somalia, and Sudan. Its natural habitat is dry savanna.

Yarkand deer

Yarkand deer

Cervus elaphus yarkandensis

The Yarkand deer (*Cervus hanglu yarkandensis*), also known as the Theenivs deer, Tarim deer, or Lop Nor stag, is a subspecies of the Central Asian red deer that is native to the province of Xinjiang, China. It is similar in ecology to the related...

Yarkand hare

Yarkand hare

Lepus yarkandensis

The Yarkand hare (*Lepus yarkandensis*) is a species of mammal in the family Leporidae. It has soft, straight, sandy brown dorsal pelage which has grayish-black stripes, and completely white ventral pelage. Endemic to China, the Yarkand hare is...

Yates's big-eared bat

Yates's big-eared bat

Micronycteris yatesi

Yates's big-eared bat (*Micronycteris yatesi*) is a species of leaf-nosed bat found in Bolivia.

Yellow Baboon

Yellow Baboon

Papio cynocephalus

The yellow baboon (*Papio cynocephalus*) is a baboon in the family of Old World monkeys. The species epithet means "dog-head" in Greek, due to the dog-like shape of the muzzle and head. Yellow baboons have slim bodies with long arms and legs, and...

Yellow golden mole

Yellow golden mole

Calcochloris obtusirostris

The yellow golden mole (*Calcochloris obtusirostris*) is a species of mammal in the family Chrysochloridae. It is found in Mozambique, South Africa (KwaZulu-natal and the limpopo), and Zimbabwe. The yellow golden mole natural habitats are...

Yellow ground squirrel

Yellow ground squirrel

Spermophilus fulvus

The yellow ground squirrel (*Spermophilus fulvus*) is a large and sturdy ground squirrel species native to Afghanistan, China, Iran, Kazakhstan, Turkmenistan, Uzbekistan and Russia. It inhabits sandy steppes with *Artemisia*, glasswort and...

Yellow isthmus rat

Yellow isthmus rat

Isthmomys flavidus

The yellow isthmus rat (*Isthmomys flavidus*) is a species of rodent in the family Cricetidae. It is found only in Panama. It was discovered by W. W. Brown Jr. on the southern slope of Volcan de Chiriqui (8° 49' N, 82° 32' W). He found it common...

Yellow Mongoose

Yellow Mongoose

Cynictis penicillata

The yellow mongoose (*Cynictis penicillata*), sometimes referred to as the red meerkat, is a member of the mongoose family. It averages about 0.45 kg (1 lb) in weight and about 510 mm (20 in) in length. It lives in open country, semi-desert...

Yellow serotine

Yellow serotine

Neoromicia flavescens

The yellow serotine (*Neoromicia flavescens*) is a species of vesper bat. It is found

in Angola, Burundi, Cameroon, Malawi, and Mozambique. Its natural habitats are subtropical or tropical dry forests, subtropical or tropical moist lowland forests,...

Yellow steppe lemming

Yellow steppe lemming

Eolagurus luteus

The yellow steppe lemming (*Eolagurus luteus*) is a species of rodent in the family Cricetidae. It is found in China, Kazakhstan, and Mongolia. Its natural habitat is temperate desert.

Yellow-Backed Duiker

Yellow-Backed Duiker

Cephalophus silvicultor

The yellow-backed duiker (*Cephalophus silvicultor*) is a forest dwelling antelope in the order Artiodactyla from the family Bovidae. Yellow-backed duikers are the most widely distributed of all duikers. They are found mainly in Central and...

Yellow-bellied brush-furred rat

Yellow-bellied brush-furred rat

Lophuromys luteogaster

The yellow-bellied brush-furred rat (*Lophuromys luteogaster*) is a species of rodent in the family Muridae. It has been recorded from the Democratic Republic of the Congo, Rwanda, and Uganda. It lives in rainforest and swamps. Although the...

Yellow-bellied climbing mouse

Yellow-bellied climbing mouse

Rhipidomys ochrogaster

The yellow-bellied climbing mouse (*Rhipidomys ochrogaster*) is a species of arboreal rodent in the family Cricetidae. It is known only from southeastern Peru, where it has been found in cloud forest at an elevation of 1830 m. This species was...

Yellow-Bellied Glider

Yellow-Bellied Glider

Petaurus australis

The Yellow-bellied glider is the least studied and least known of its family. It is amongst the largest gliding possums of Australia, the loudest, and the best glider. They have a distinctive very long thickly furred mostly black tail and for...

Yellow-Bellied Marmot

Yellow-Bellied Marmot

Marmota flaviventris

The Yellow-bellied marmot (*Marmota flaviventris*) is a large, stout-bodied ground squirrel in the marmot genus. It is one of fourteen species of marmots and is native to mountainous regions of North America. They live in burrows in colonies and...

Yellow-bellied sheath-tailed bat

Yellow-bellied sheath-tailed bat

Saccolaimus flaviventris

The yellow-bellied sheath-tailed bat (*Saccolaimus flaviventris*), also known as the yellow-bellied sheathtail or yellow-bellied pouched bat, is a microbat species of the family Emballonuridae found extensively in Australia and less commonly in...

Yellow-Bellied Weasel

Yellow-Bellied Weasel

Mustela kathiah

The Yellow-bellied weasel (*Mustela kathiah*) is a small carnivorous that lives in

forests in central and eastern Asia. Despite their size, these animals are active predators and feed on small mammals.

Yellow-cheeked chipmunk
Yellow-cheeked chipmunk
Neotamias ochrogenys

The yellow-cheeked chipmunk (*Neotamias ochrogenys*), also known as the redwood chipmunk, is a species of rodent in the squirrel family, *Sciuridae*. It is endemic to areas near the coast of northern California in the United States where it inhabits...

Yellow-Cheeked Gibbon
Yellow-Cheeked Gibbon
Nomascus gabriellae

The Yellow-cheeked gibbon (*Nomascus gabriellae*) is a species of ape that lives in forests of Southeast Asia. Little is known about this species in the wild, but it is thought that it has a life span of approximately 46 years. It was discovered...

Yellow-crowned brush-tailed rat
Yellow-crowned brush-tailed rat
Isothrix bistrata

The yellow-crowned brush-tailed rat, *Isothrix bistrata*, is a South American brush-tailed rat species from the family *Echimyidae*. It is found in Bolivia, Brazil, Colombia, Ecuador, Peru, and Venezuela. They are nocturnal and arboreal animals found...

Yellow-faced horseshoe bat
Yellow-faced horseshoe bat
Rhinolophus virgo

The yellow-faced horseshoe bat (*Rhinolophus virgo*) is a species of bat in the family *Rhinolophidae*. It is endemic to the Philippines.

Yellow-faced pocket gopher
Yellow-faced pocket gopher
Cratogeomys castanops

The yellow-faced pocket gopher (*Cratogeomys castanops*) is a species of pocket gopher that is native to shortgrass prairies in the south-western United States and northern Mexico. It is the species that lives north of the Southern Coahuila...

Yellow-Footed Antechinus
Yellow-Footed Antechinus
Antechinus flavipes

The Yellow-footed antechinus (*Antechinus flavipes*) is a shrew-like marsupial found in Australia. These animals are also sometimes called 'broad-footed marsupial mice', 'pouched mice', or 'Antechinus shrews'. However, the majority of those common...

Yellow-Footed Rock-Wallaby
Yellow-Footed Rock-Wallaby
Petrogale xanthopus

The yellow-footed rock-wallaby (*Petrogale xanthopus*), formerly known as the ring-tailed rock-wallaby, is a member of the macropod family (the marsupial family that includes the kangaroos, wallabies, tree-kangaroos, and wallaroos).

Yellow-haired hill rat
Yellow-haired hill rat
Bunomys chrysocomus

The yellow-haired hill rat (*Bunomys chrysocomus*) is a species of rodent in the family *Muridae*. It is found only in Sulawesi, Indonesia, including Lore Lindu

National Park.
Yellow-lipped bat
Yellow-lipped bat
Vespadelus douglasorum

The yellow-lipped cave bat (*Vespadelus douglasorum*) is a vespertilionid bat which only occurs in the Kimberley region of northwest Australia. The bat was first captured at Tunnel Creek in 1958 and a description published nearly twenty years...

Yellow-Necked Mouse
Yellow-Necked Mouse
Apodemus flavicollis

The yellow-necked mouse (*Apodemus flavicollis*), also called yellow-necked field mouse, yellow-necked wood mouse, and South China field mouse, is closely related to the wood mouse, with which it was long confused. It was only recognised as a...

Yellow-nosed cotton rat
Yellow-nosed cotton rat
Sigmodon ochrognathus

The yellow-nosed cotton rat (*Sigmodon ochrognathus*) is a species of rodent in the family Cricetidae. It is native to Mexico and to the states of Arizona, New Mexico, and Texas in the United States, where it inhabits mountain grassland, scrub,...

Yellow-Pine Chipmunk
Yellow-Pine Chipmunk
Neotamias amoenus

The yellow-pine chipmunk (*Neotamias amoenus*) is a species of order Rodentia in the family Sciuridae. It is found in western North America: parts of Canada and the United States.

Yellow-rumped leaf-eared mouse
Yellow-rumped leaf-eared mouse
Phyllotis xanthopygus

The yellow-rumped leaf-eared mouse (*Phyllotis xanthopygus*) otherwise known as the Patagonian leaf-eared mouse is a species of rodent in the family Cricetidae and order Rodentia. It is the most widespread member of the genus.

Yellow-sided opossum
Yellow-sided opossum
Monodelphis dimidiata

The yellow-sided opossum (*Monodelphis dimidiata*) is an opossum species from South America. It is found in Argentina, Brazil and Uruguay. They have grey or black fur on their dorsal side with yellowish fur on the lateral side that continues down...

Yellow-spotted brush-furred rat
Yellow-spotted brush-furred rat
Lophuromys flavopunctatus

The yellow-spotted brush-furred rat (*Lophuromys flavopunctatus*) is a species of rodent in the family Muridae found in Angola, the Democratic Republic of the Congo, Ethiopia, Kenya, Malawi, Mozambique, South Sudan, Tanzania, and Zambia. Its...

Yellow-Spotted Rock Hyrax
Yellow-Spotted Rock Hyrax
Heterohyrax brucei

The Yellow-spotted rock hyrax (*Heterohyrax brucei*) is a mammal found only in Africa. The name "hyrax" comes from the Greek word "ὑραξ" and means shrew-mouse. This species has several common names including Bush hyrax, Yellow-spotted hyrax, Bush...

Yellow-striped chevrotain

Yellow-striped chevrotain
Moschiola kathygre

The yellow-striped chevrotain (*Moschiola kathygre*) is a species of chevrotain described in 2005. It is found in the wet zones of Sri Lanka. It was recognized as a species distinct from *Moschiola meminna* based on the phylogenetic species concept.

Yellow-tailed rat
Yellow-tailed rat
Rattus xanthurus

The yellow-tailed rat (*Rattus xanthurus*) is a species of rodent in the family Muridae. It is found only in northeastern Sulawesi, Indonesia.

Yellow-Tailed Woolly Monkey
Yellow-Tailed Woolly Monkey
Oreonax flavicauda

The Yellow-tailed woolly monkey (*Lagothrix flavicauda*) is a rare New World monkey native to Peru. It was first described by Alexander von Humboldt in 1812 based on a skin found 10 years earlier. Humboldt had never seen a live animal of this...

Yellow-throated big-eared bat
Yellow-throated big-eared bat
Lamproncycteris brachyotis

The yellow-throated big-eared bat or orange-throated bat (*Lamproncycteris brachyotis*) is a species of bat from South and Central America, where it ranges from southern Mexico to Brazil. It is monotypic within the genus *Lamproncycteris*. A frugivore...

Yellow-Throated Marten
Yellow-Throated Marten
Martes flavigula

The yellow-throated marten (*Martes flavigula*) is a marten species native to Asia. It is listed as Least Concern on the IUCN Red List due to its wide distribution, evidently relatively stable population, occurrence in a number of protected areas,...

Yellow-throated squirrel
Yellow-throated squirrel
Sciurus gilvigularis

The yellow-throated squirrel (*Sciurus gilvigularis*) is a tree squirrel in the genus *Sciurus* endemic to South America. It is found in Brazil, Guyana and Venezuela.

Yellow-winged bat
Yellow-winged bat
Lavia frons

The yellow-winged bat (*Lavia frons*) is one of five species of false vampire bat (family Megadermatidae) from Africa.

Yellowish myotis
Yellowish myotis
Myotis levis

The yellowish myotis (*Myotis levis*), is a vesper bat species from South America. It is found in Argentina, Brazil, Paraguay, and Uruguay. Dinelli's myotis (*M. dinellii*) was formerly considered a subspecies of *M. levis*, but was split as a...

Yemeni mouse
Yemeni mouse
Myomyscus yemeni

The Yemeni mouse (*Ochromyscus yemeni*) is a species of rodent in the family Muridae. It is found in Saudi Arabia and Yemen. It is the only modern member of the tribe Praomyini to be found outside of Africa. It was initially described as a...
Yemeni mouse-tailed bat
Yemeni mouse-tailed bat
Rhinopoma hadramauticum

The Yemeni mouse-tailed bat (*Rhinopoma hadramauticum*) is an endangered species of bat found in Yemen. It is only known from one roost, and its population is estimated at 150 individuals.
Yemeni trident leaf-nosed bat
Yemeni trident leaf-nosed bat
Triaenops parvus

The Yemeni trident leaf-nosed bat (*Triaenops parvus*) is a species of bat found in the Middle East.
Yepes's mulita
Yepes's mulita
Dasypus mazzai

Yepes's mulita or the Yungas lesser long-nosed armadillo (*Dasypus mazzai*) is a species of armadillo in the family Dasypodidae. It is endemic to Argentina and Bolivia. Its natural habitat is subtropical dry forests. The species was renamed D....
Yezo sika deer
Yezo sika deer
Cervus nippon yesoensis

The Yezo sika deer (*Cervus nippon yesoensis*, Japanese: エゾシカ / 蝦夷鹿, romanized: yezoshika, Ainu: ユク yuk) is one of the many subspecies of the sika deer. The sika that inhabit the island of Hokkaido are indigenous, although it is not known whether...
Yolanda's tuco-tuco
Yolanda's tuco-tuco
Ctenomys yolandae

Yolanda's tuco-tuco (*Ctenomys yolandae*) is a species of rodent in the family Ctenomyidae. The species is endemic to Santa Fe Province, northeast Argentina, where it lives near the Paraná and San Javier rivers. Its karyotype has $2n = 50$ and $FN = ...$
Yonenaga's Atlantic spiny rat
Yonenaga's Atlantic spiny rat
Trinomys yonenagae

Yonenaga's Atlantic spiny-rat (*Trinomys yonenagae*) or torch-tail spiny rat is a spiny rat species endemic to Brazil. Locally, it is known as rabo de facho. Named for Yatiyo Yonenaga-Yassuda, a cytogenetics researcher, it is considered an...
Yucatan brown brocket
Yucatan brown brocket
Mazama pandora

The Yucatan brown brocket (*Odocoileus pandora*) is a small species of deer native to the Yucatán Peninsula of Mexico, Belize and Guatemala. While it is found in humid tropical forest like most other brocket deer, the Yucatan brown brocket also...
Yucatan deer mouse
Yucatan deer mouse
Peromyscus yucatanicus

The Yucatan deer mouse (*Peromyscus yucatanicus*) is a species of rodent in the family Cricetidae. The species is found in Mexico and Guatemala; an example habitat is the Petenes mangroves ecoregion of the Yucatan.

Yucatan small-eared shrew

Yucatan small-eared shrew

Cryptotis mayensis

The Yucatan small-eared shrew (*Cryptotis mayensis*) is a species of mammal in the family Soricidae. It is mainly known from lowlands of Guatemala, Belize and Mexico's Yucatán Peninsula, where it has been found in dry scrubland and tropical dry...

Yucatan spider monkey

Yucatan spider monkey

Ateles geoffroyi yucatanensis

The Yucatan spider monkey (*Ateles geoffroyi yucatanensis*), also known by its mayan name "Ma'ax", is a disputed subspecies of Geoffroy's spider monkey, and is one of the largest types of New World monkey. It inhabits Mexico, Guatemala and Belize....

Yucatan squirrel

Yucatan squirrel

Sciurus yucatanensis

The Yucatan squirrel (, *Sciurus yucatanensis*), originally named the Yucatan gray squirrel, also once named the Campeche squirrel, is a tree squirrel in the genus *Sciurus* found in the Yucatán Peninsula and adjacent areas. It is native to northern...

Yucatan yellow bat

Yucatan yellow bat

Rhogeessa aeneus

The Yucatan yellow bat (*Rhogeessa aeneus*) is a species of bat found in the Yucatán Peninsula in Mexico, and possibly also in Belize and Guatemala. These small insectivorous bats forage on flying insects (most likely mosquitoes) at dawn and dusk.

Yulungshan vole

Yulungshan vole

Eothenomys proditor

The Yulungshan vole, Yulong Chinese vole, Yulongxuen Chinese vole, or Yulongxuen red-backed vole (*Eothenomys proditor*) is a species of rodent in the family Cricetidae, endemic to Jade Dragon Snow Mountain in the Sichuan-Yunnan border region of...

Yuma myotis

Yuma myotis

Myotis yumanensis

The Yuma myotis (*Myotis yumanensis*) is a species of vesper bat native to western North America.

Yunnan bush rat

Yunnan bush rat

Hadromys yunnanensis

The Yunnan bush rat (*Hadromys yunnanensis*) is a species of rodent from the family Muridae. It has just recently been released from synonymy with the Manipur bush rat, and so there is very little information about it. It was recognized as a...

Yunnan hare

Yunnan hare

Lepus comus

The Yunnan hare (*Lepus comus*) is a medium-sized species of mammal in the family Leporidae. It has soft, flat, and long dorsal pelage which is grayish brown or dark gray in color, and whitish ventral pelage. It was considered endemic to China...

Yunnan lar gibbon

Yunnan lar gibbon

Hylobates lar yunnanensis

The Yunnan lar gibbon (*Hylobates lar yunnanensis*), also known as the Yunnan white-handed gibbon, is a subspecies of the lar gibbon, a primate in the gibbon family, Hylobatidae. This Chinese subspecies is thought to be extinct.

Yunnan red-backed vole

Yunnan red-backed vole

Eothenomys miletus

The Yunnan red-backed vole (*Eothenomys miletus*) is a species of rodent in the family Cricetidae. It is found in Southwest China, specifically Yunnan Province. It is the largest member of the genus *Eothenomys*, with a higher cranium, a soft, thick...

Zacatecan deer mouse

Zacatecan deer mouse

Peromyscus difficilis

The Zacatecan deer mouse or southern rock mouse (*Peromyscus difficilis*) is a species of rodent in the family Cricetidae. It is found only in Mexico, and is not considered endangered.

Zacatecas harvest mouse

Zacatecas harvest mouse

Reithrodontomys zacatecae

The Zacatecas harvest mouse (*Reithrodontomys zacatecae*) is a species of rodent in the family Cricetidae. It is found only in Mexico.

Zacatecas shrew

Zacatecas shrew

Sorex emarginatus

The Zacatecas shrew (*Sorex emarginatus*) is a species of mammal in the family Soricidae. It is endemic to Mexico.

Zaglossus hacketti

Zaglossus hacketti

Zaglossus hacketti

Murrayglossus is a genus in the family Tachyglossidae. It contains a single species, *Murrayglossus hacketti*, the giant echidna, an extinct species of echidna from Western Australia that is dated to the Pleistocene. It is known only from a few...

Zaglossus robustus

Zaglossus robustus

Zaglossus robustus

Zaglossus robustus is an extinct species of long-beaked echidna known from the middle Miocene (about 13 or 14 million years ago) of Gulgong, New South Wales, Australia. It may belong in the genus *Megalibgwilia*. The supposed fossil platypus...

Zagros Mountains mouse-like hamster

Zagros Mountains mouse-like hamster

Calomyscus bailwardi

The Zagros Mountains mouse-like hamster (*Calomyscus bailwardi*) is a relatively

little-known rodent which was the first species of mouse-like hamster to be described. The species is distributed throughout much of southern Iran, particularly in...

Zaisan mole vole
Zaisan mole vole
Ellobius tancrei

The Zaisan mole vole (*Ellobius tancrei*), or eastern mole vole, is a species of rodent in the family Cricetidae. It is found in central Asia.

Zambian mole-rat
Zambian mole-rat
Fukomys amatus

The Zambian mole-rat (*Fukomys amatus*) is a species of rodent in the family Bathyergidae. It is found in Zambia and the Democratic Republic of the Congo. It was formerly considered a subspecies of *Cryptomys hottentotus*.

Zanj sun squirrel
Zanj sun squirrel
Heliosciurus undulatus

The Zanj sun squirrel (*Heliosciurus undulatus*) is a species of rodent in the family Sciuridae found in eastern Africa.

Zanzibar bushbaby
Zanzibar bushbaby
Paragalago zanzibaricus

The Zanzibar bushbaby, Matundu dwarf galago, Udzungwa bushbaby, or Zanzibar galago (*Paragalago zanzibaricus*) is a primate of the family Galagidae. An adult typically weighs 150 grams (5.3 oz), its head-body length is 14 to 15 centimetres (5.5 to...

Zanzibar Red Colobus
Zanzibar Red Colobus
Piliocolobus kirkii

The Zanzibar red colobus (*Piliocolobus kirkii*) is a species of red colobus monkey endemic to Unguja, the main island of the Zanzibar Archipelago, off the coast of Tanzania. It is also known as Kirk's red colobus after Sir John Kirk, the British...

Zanzibar servaline genet
Zanzibar servaline genet
Genetta servalina archeri

The Zanzibar servaline genet (*Genetta servalina archeri*) is a recently discovered subspecies of servaline genet endemic to Unguja Island in the Tanzanian archipelago of Zanzibar. Its conservation status is uncertain.

Zaphir's shrew
Zaphir's shrew
Crocidura zaphiri

Zaphir's shrew (*Crocidura zaphiri*) is a species of mammal in the family Soricidae. It is found in Ethiopia and Kenya. Its natural habitats are subtropical or tropical dry forest and subtropical or tropical moist lowland forest.

Zarudny's jird
Zarudny's jird
Meriones zarudnyi

Zarudny's jird (*Meriones zarudnyi*) is a species of rodent in the family Muridae. It is found in Afghanistan, Iran, and Turkmenistan.

Zarudny's rock shrew
Zarudny's rock shrew

Crocidura zarudnyi

Zarudny's rock shrew (*Crocidura zarudnyi*) is a species of mammal in the family Soricidae. It is found in Afghanistan, Iran, and Pakistan.

Zebra Duiker

Zebra Duiker

Cephalophus zebra

The Zebra duiker (*Cephalophus zebra*) is a small antelope found in Africa. It is believed to be one of the earliest duiker species to have evolved. The species is locally known as the marking deer in Liberia.

Zebu

Zebu

Bos taurus indicus

The zebu (; *Bos indicus* or *Bos taurus indicus*), sometimes known in the plural as indicine cattle or humped cattle, is a species or subspecies of domestic cattle originating in the Indian sub-continent. Zebu are characterised by a fatty hump on...

Zempoaltepec

Zempoaltepec

Peromyscus melanocarpus

The zempoaltepec (*Peromyscus melanocarpus*) is a species of rodent in the family Cricetidae. It is found only in Mexico.

Zempoaltepec deer mouse

Zempoaltepec deer mouse

Habromys lepturus

The Zempoaltepec deer mouse (*Habromys lepturus*) also known as the slender-tailed deer mouse, is a species of rodent in the family Cricetidae. It is endemic to southeastern Mexico, at one locality on Cerro Zempoaltepec above 2,500 metres (8,200 ft),...

Zempoaltépec vole

Zempoaltépec vole

Microtus umbrosus

The Zempoaltépec vole (*Microtus umbrosus*) is a species of rodent in the family Cricetidae. The name *Microtus* is from the Greek word *mikros* meaning small and *otus* meaning ear. The name *umbrosus* could be from the Latin *umbros* meaning shady. It is...

Zena's brush-furred rat

Zena's brush-furred rat

Lophuromys zena

According to UniProt, Zena's brush-furred rat (*Lophuromys zena*) is a species of rat. However, the IUCN lists this as a synonym for *Lophuromys flavopunctatus* (yellow-spotted brush-furred rat).

Zenati myotis

Zenati myotis

Myotis zenatius

The Zenati myotis (*Myotis zenatius*) is a rare species of mouse-eared bat that is restricted to North Africa. It is very rare in Morocco, being restricted to just four localities ranging from the Rif to the Atlas Mountains. Aside from Morocco, it is...

Zenker's fruit bat

Zenker's fruit bat

Scotonycteris zenkeri

Zenker's fruit bat or Tear-drop bat (*Scotonycteris zenkeri*) is a species of megabat in the family Pteropodidae. It is found in Cameroon, Central African Republic, Republic of the Congo, Democratic Republic of the Congo, Ivory Coast, Equatorial...

Ziama horseshoe bat
Ziama horseshoe bat
Rhinolophus ziama

The Ziama horseshoe bat (*Rhinolophus ziama*) is a species of bat in the family Rhinolophidae. It was first described in 2002. It is found in Guinea, Sierra Leone, and Liberia. Its natural habitats are subtropical or tropical moist lowland forests...

Ziegler's water rat
Ziegler's water rat
Hydromys ziegleri

Ziegler's water rat (*Hydromys ziegleri*), discovered in the mid-2000s, is a semiaquatic species of rodent native to the mountains Papua New Guinea of which little is known.

Zulu golden mole
Zulu golden mole
Amblysomus hottentotus iris

The Zulu golden mole (*Amblysomus hottentotus iris*) is a golden mole native to Transvaal, South Africa.

Zulu serotine
Zulu serotine
Neoromicia zuluensis

The Zulu serotine (*Neoromicia zuluensis*), also called the Zulu pipistrelle, aloe bat, or aloe serotine, is a species of vesper bat found in Angola, Botswana, Democratic Republic of the Congo, Ethiopia, Kenya, Malawi, Namibia, South Africa, South...

Zygodontomys brevicauda
Zygodontomys brevicauda
Zygodontomys brevicauda

Zygodontomys brevicauda, also known as the short-tailed zygodont, short-tailed cane mouse, or common cane mouse, is a species of rodent in the genus *Zygodontomys* of tribe Oryzomyini.

Zygodontomys brunneus
Zygodontomys brunneus
Zygodontomys brunneus

Zygodontomys brunneus, also known as the brown zygodont, brown cane mouse, or Colombian cane mouse, is a rodent species in the genus *Zygodontomys* of tribe Oryzomyini. It is found only in Colombia.