

Fish

3497 species

[Home](#)

Fish

[Use Filters](#)

Sort By:

Popularity

A to Z

Group By:

None

Family

Population Status



Abactochromis

Abactochromis labrosus

Abactochromis is a monotypic genus of cichlid fish containing the single species Abactochromis labrosus (formerly known as Melanochromis labrosus). The genus name combines the Latin for "driven away from" ("ab" - away from, "actus" - driven) with...



Abalistes stellatus

Abalistes stellatus

Abalistes stellatus is a member of the triggerfish family that occurs along the coasts of the Indian Ocean, the Red Sea, the Persian Gulf, and along the western edge of the Pacific Ocean.



Abbottina rivularis

Abbottina rivularis

Abbottina rivularis, also known as the Chinese false gudgeon or the Amur false gudgeon, is a species of ray-finned fish in the family Cyprinidae, the carps and minnows. It is native to China, Korea, and Japan, but it has been introduced to the...



Ablabys taenianotus

Ablabys taenianotus

Ablabys taenianotus, the cockatoo waspfish, is a small fish species that belongs to the scorpionfishes family Scorpaenidae.



Abramites hypselonotus

Abramites hypselonotus

Abramites hypselonotus – known as the marbled headstander or the high-backed headstander – is a member of the family Anostomidae of the order Characiformes. Under normal fish classifying rules, the marbled headstander would be classified as a...



Abudefduf troschelii

Abudefduf troschelii

Abudefduf troschelii, the Pacific sergeant major or Panama sergeant major, is a species of damselfish belonging to the family Pomacentridae that can be identified by the pronounced black stripes on the lateral sides of the fish. Its specific name...



Abyssal grenadier

Coryphaenoides armatus

The abyssal grenadier, Coryphaenoides armatus, is an abyssal fish of the genus Coryphaenoides, found in all the world's oceans, at depths between 800 and 4,000 metres (2,600 and 13,100 ft). Its adult length is 20 to 40 centimetres (8 to 16 in),...



Abyssal spiderfish

Bathypterois longipes

The abyssal spiderfish, Bathypterois longipes, is a species of deepsea tripod fish, a demersal fish living on the bottom of the Atlantic Ocean, even on the abyssal plain. It is one of the deepest-dwelling fish known. This solitary fish grows about...



Abyssobrotula galathea

Abyssobrotula galathea

Abyssobrotula galathea is a species of cusk eel in the family Ophidiidae. It is the deepest-living fish known; one specimen, trawled from a depth of 8,370 m (27,460 ft) in the Puerto Rico Trench in 1970, holds the record for the deepest fish...



Acadian redfish

Sebastes fasciatus

The Acadian redfish (Sebastes fasciatus), also known as the Atlantic redfish, Acadian rockfish, or Labrador redfish, is a species of marine ray-finned fish belonging to the subfamily Sebastinae, the rockfishes, part of the family Scorpaenidae. It...



Acanthemblemaria medusa

Acanthemblemaria medusa

Acanthemblemaria medusa, the medusa blenny, is a species of chaenopsid blenny found in coral reefs around Lesser Antilles, in the western central Atlantic ocean. It can reach a maximum length of 4 cm (1.6 in) TL.



Acanthicus adonis

Acanthicus adonis

Acanthicus adonis, the adonis pleco or polka dot lyre-tail pleco, is a large species of armored catfish. It was originally described from the lower Tocantins River in Brazil, but individuals resembling the species have also been recorded from...



Acanthicus hystrix

Acanthicus hystrix

Acanthicus hystrix, the lyre-tail pleco, is a species of armored catfish native to the Amazon, Tocantins–Araguaia and Orinoco basins. It is typically found at various depths on a rock or rock–gravel bottom in the main channel of rivers in places...



Acanthobrama telavivensis

Acanthobrama telavivensis

Acanthobrama telavivensis, commonly known as the Yarkon bream or Yarkon bleak, is a species of freshwater ray-finned fish of the family Cyprinidae found only in Israel, in the Yarkon River system.



Acanthogobius flavimanus

Acanthogobius flavimanus

Acanthogobius flavimanus is a species of fish in the goby family known by the common name yellowfin goby. Other common names include mahaze, Japanese river goby, Oriental goby, and spotted goby. It is native to Asia, where its range includes...



Acanthopagrus australis

Acanthopagrus australis

Acanthopagrus australis, the yellowfin bream, also known as sea bream, surf bream, silver bream or eastern black bream, is a species of marine and freshwater fish of the porgy family, Sparidae. It is a deep-bodied fish, occasionally confused with...



Acanthopagrus berda

Acanthopagrus berda

Acanthopagrus berda, the goldsilk seabream, sly bream, picnic seabream, black sea bream, black porgy, picky bream, silver bream or river bream, is a species of ray-finned fish belonging to the family Sparidae, the sea breams and porgies. This...



Acanthopagrus butcheri

Acanthopagrus butcheri

The Black bream (Acanthopagrus butcheri), also commonly known as the southern black bream, southern bream and blue-nosed bream, is a species of anadromous ray-finned fish of the porgy family Sparidae. A deep-bodied fish, it is occasionally...



Acanthopagrus latus

Acanthopagrus latus

Acanthopagrus latus, the yellowfin seabream, grey bream, Houttuyn's yellowfin seabream, Japanese bream or yellow-finned black porgy, is a species of marine ray-finned fish belonging to the family Sparidae, the seabreams and porgies. This fish is...



Acanthopagrus schlegelii

Acanthopagrus schlegelii

Acanthopagrus schlegelii, the blackhead seabream, black porgy or black seabream, is a species of marine ray-finned fish belonging to the family Sparidae, the seabreams and porgies. This species is found in the Western Pacific Ocean. The blackhead...



Acanthurus achilles

Acanthurus achilles

Acanthurus achilles, the Achilles tang, redtail surgeonfish or redspot surgeonfish, is a marine ray-finned fish belonging to the family Acanthuridae, the surgeonfishes, unicornfishes and tangs. This fish is found in the Pacific Ocean.



Acanthurus bariene

Acanthurus bariene

Acanthurus bariene, the bariene surgeonfish, black-spot surgeonfish, or eye-spot surgeonfish, is a species of marine ray-finned fish belonging to the family Acanthuridae, the surgeonfishes, unicornfishes and tangs. This species is found in the...



Acanthurus blochii

Acanthurus blochii

Acanthurus blochii, the ringtail surgeonfish or dark surgeonfish, is a marine ray-finned fish belonging to the family Acanthuridae, the surgeonfishes, unicornfishes and tangs. This fish is found in the Indo-Pacific.



Acanthurus coeruleus

Acanthurus coeruleus

Acanthurus coeruleus is a species of marine ray-finned fish belonging to the family Acanthuridae, the surgeonfishes, unicornfishes and tangs, found in the Atlantic Ocean. It can grow up to 39 centimetres (15 in) long. Common names include...



Acanthurus dussumieri

Acanthurus dussumieri

Acanthurus dussumieri, commonly known as Dussumier's surgeonfish, the eye-stripe surgeonfish or the ornate surgeonfish, is a ray-finned fish from the Indo-Pacific. It occasionally makes its way into the aquarium trade. It grows to a maximum...



Acanthurus gahhm

Acanthurus gahhm

Acanthurus gahhm, the black surgeonfish, monk surgeonfish or lined surgeonfish, is a species of marine ray-finned fish belonging to the family Acanthuridae, the surgeonfishes, unicornfishes or tangs. This fish is found in the Western Indian Ocean.



Acanthurus guttatus

Acanthurus guttatus

Acanthurus guttatus, the whitespotted surgeonfish, spotted surgeonfish, mustard surgeonfish, mustard tang or spotband surgeonfish, is a species of marine ray-finned fish belonging to the family Acanthuridae, the surgeonfishes, unicornfishes or...



Acanthurus japonicus

Acanthurus japonicus

Acanthurus japonicus, the Japan surgeonfish, white-faced surgeonfish, gold rim tang, powder brown tang and white-nose surgeonfish, is a species of marine ray-finned fish belonging to the family Acanthuridae, the surgeonfishes, unicornfishes or...



Acanthurus leucosternon

Acanthurus leucosternon

Acanthurus leucosternon the blue surgeonfish, powder blue tang and powder-blue surgeonfish, is a species of marine ray-finned fish belonging to the family Acanthuridae, the surgeonfishes, unicornfishes and tangs. This species is found in the...



Acanthurus lineatus

Acanthurus lineatus

Acanthurus lineatus, the lined surgeonfish, blue banded surgeonfish, blue-lined surgeonfish, clown surgeonfish, pyjama tang, striped surgeonfish, and zebra surgeonfish, is a species of marine ray-finned fish belonging to the family Acanthuridae,...



Acanthurus maculiceps

Acanthurus maculiceps

Acanthurus maculiceps, the white-freckled surgeonfish, yellow-freckled surgeonfish, pale-lined surgeonfish, spotted-faced surgeonfish or earbar surgeonfish, is a species of marine ray-finned fish belonging to the family Acanthuridae, the...



Acanthurus mata

Acanthurus mata

Acanthurus mata, the pale surgeonfish, blue-lined surgeonfish, mata surgeonfish, striped surgeonfish, tailring surgeon, white-tail lancet or yellowmask surgeonfish, is a species of marine ray-finned fish belonging to the family Acanthuridae, the...



Acanthurus monroviae

Acanthurus monroviae

Acanthurus monroviae, the Monrovia doctorfish, is a species of marine ray-finned fish belonging to the family Acanthuridae, the surgeonfishes, unicornfishes and tangs. It is found in the eastern Atlantic Ocean.



Acanthurus nigricans

Acanthurus nigricans

Acanthurus nigricans, the goldrim surgeonfish, velvet surgeon, whitecheek surgeonfish, yellow-banded surgeonfish or yellowrimmed surgeonfish, is a species of marine ray-finned fish belonging to the family Acanthuridae, the surgeonfishes,...



Acanthurus nigricauda

Acanthurus nigricauda

Acanthurus nigricauda, the epaulette surgeonfish, black-barred surgeonfish, eye-line surgeonfish, shoulderbar surgeonfish, white-tail surgeonfish or blackstreak surgeonfish, is a species of marine ray-finned fish belonging to the family...



Acanthurus nigrofuscus

Acanthurus nigrofuscus

Acanthurus nigrofuscus, the brown surgeonfish, blackspot surgeonfish, brown tang, dusky surgeon, lavender tang or spot-cheeked surgeonfish, is a species of marine ray-finned fish belonging to the family Acanthuridae, which includes the...



Acanthurus nigroris

Acanthurus nigroris

Acanthurus nigroris, the bluelined surgeonfish, is a species of marine ray-finned fish belonging to the family Acanthuridae, which includes the surgeonfishes, unicornfishes and tangs. This species and A. nigros have been regarded as synonymous...



Acanthurus olivaceus

Acanthurus olivaceus

Acanthurus olivaceus, the orange-band surgeonfish, the orange-shoulder surgeonfish or the orangebar tang, is a species of marine ray-finned fish belonging to the family Acanthuridae, this family includes the surgeonfishes, unicornfishes and...



Acanthurus pyroferus

Acanthurus pyroferus

Acanthurus pyroferus, chocolate surgeonfish, mimic surgeonfish, orange-gilled surgeonfish, Pacific mimic surgeon, and yellowspot surgeon, is a species of marine ray-finned fish belonging to the family Acanthuridae, which includes the...



Acanthurus thompsoni

Acanthurus thompsoni

Acanthurus thompsoni, the night surgeonfish, chocolate surgeonfish, Thompson's surgeonfish, Thompson's tang or whitetail surgeonfish, is a species of marine ray-finned fish belonging to the family Acanthuridae which includes the...



Acanthurus tractus

Acanthurus tractus

Acanthurus tractus, the five-band surgeonfish, ocean surgeon, or ocean surgeonfish, is a species of ray-finned fish in the family Acanthuridae found in the western Atlantic Ocean, Florida, the Bahamas, the Caribbean Sea and the Gulf of Mexico....



Acanthurus triostegus

Acanthurus triostegus

Acanthurus triostegus, the convict surgeonfish, convict surgeon, convict tang or fiveband surgeonfish, is a species of marine ray-finned fish belonging to the family Acanthuridae which includes the surgeonfishes, unicornfishes and tangs. This...



Acanthurus xanthopterus

Acanthurus xanthopterus

Acanthurus xanthopterus, the yellowfin surgeonfish yellowmask surgeonfish, ringtail surgeonfish or Cuvier's surgeonfish, is a species of marine ray-finned fish belonging to the family Acanthuridae which includes the surgeonfishes, unicornfishes...



Acantopsis dialuzona

Acantopsis dialuzona

Acantopsis dialuzona is a loach native to the swift, clear streams and rivers of mainland and archipelagic Southeast Asia, Indonesia, Peninsular Malaysia, Thailand, but Southeast Asia outside its range, including as Acantopsis. A. choirorhynchos...



Acheilognathus melanogaster

Acheilognathus melanogaster

Acheilognathus melanogaster is a species of brackish, freshwater ray-finned fish in the genus Acheilognathus. It is endemic to Japan.



Achirus lineatus

Achirus lineatus

The lined sole (Achirus lineatus) is a ray-finned flatfish found in the Western Atlantic. Its common length is 17 cm. Often considered a trash fish in commercial trawling, it is of little or no economic value.



Achoerodus viridis

Achoerodus viridis

The eastern blue groper (Achoerodus viridis) is a species of wrasse native to southeastern Australia from Hervey Bay in southern Queensland to Wilsons Promontory in Victoria. They occur in coastal waters, preferring rocky areas at a depth of...



Acipenser oxyrinchus

Acipenser oxyrinchus

Acipenser oxyrinchus is a species of sturgeon.



Adolfo's catfish

Corydoras adolfoi

Adolfo's catfish (Corydoras adolfoi) is a tropical freshwater fish belonging to the Corydoradinae sub-family of the family Callichthyidae. It originates in inland waters in South America, and is found in the Rio Negro basin and the Rio Uapes in...



Adrianichthys kruyti

Adrianichthys kruyti

Adrianichthys kruyti (the duck-billed buntingi or duckbilled buntingi) is a critically endangered species of ricefish that is endemic to Lake Poso, Sulawesi, Indonesia. It was discovered by Max Wilhelm Carl Weber on his Siboga Expedition to...



Adriatic sturgeon

Acipenser naccarii

The Adriatic sturgeon (Acipenser naccarii) is a species of fish in the family Acipenseridae. It is native to the Adriatic Sea and large rivers which flow in it of Albania, Greece, Italy, Montenegro, Croatia, Bosnia and Herzegovina and Slovenia....



Aeoliscus strigatus

Aeoliscus strigatus

Aeoliscus strigatus, also known as the razorfish, jointed razorfish or coral shrimpfish, is a member of the family Centriscidae of the order Syngnathiformes. This unique fish adopts a head-down tail-up position as an adaptation for hiding among...



African angelshark

Squatina africana

The African angelshark (Squatina africana) is an angelshark of the family Squatinidae.



African arowana

Heterotis niloticus

The African arowana or Nile arowana (Heterotis niloticus) is a species of bonytongue. Despite being called an "arowana", the African arowana is more closely related to arapaimas, the only other members in the subfamily Arapaiminae, than the South...



African banded barb

Enteromius fasciolatus

The African banded barb, Angola barb, blue-barred barb or fire barb (Enteromius fasciolatus) is a ray-finned fish species in the family Cyprinidae. It is found mainly in Angola and Zambia, extending into neighboring Botswana, the Democratic...



African brown knifefish

Xenomystus nigri

The African brown knifefish (Xenomystus nigri), is the only species in the genus Xenomystus of the family Notopteridae. This fish is found in the Chad, Nile, Congo, Ogowe and Niger basins, as well as coastal river basins in Sierra Leone, Liberia,...



African butter catfish

Schilbe mystus

The African butter catfish (Schilbe mystus) is a species of fish in the family Schilbeidae. It is native to many major river systems in Africa. Other common names for the fish include butter fish, butter barbel, African glass catfish, lubangu,...



African jewelfish

Hemichromis bimaculatus

The African jewelfish (Hemichromis bimaculatus), also known as jewel cichlid or jewelfish, is from the family Cichlidae.



African leaffish

Polycentropsis abbreviata

the African leaffish (Polycentropsis abbreviata) is a species of African leaffish native to fresh waters of western Africa. Although placed in the family Nandidae by FishBase and by the 5th Edition of Fishes of the World many recent authorities...



African longfin eel

Anguilla mossambica

The African longfin eel (*Anguilla mossambica*), also known simply as the longfin eel, is an eel in the family Anguillidae. It was described by Wilhelm Peters in 1852, originally under the genus *Muraena*. It is a tropical eel known from freshwaters...



African moon tetra

Bathyaethiops caudomaculatus

The African moon tetra (Bathyaethiops caudomaculatus) is a species of African tetra, found in slow flowing rivers of the middle and lower course of the Congo River. The fish is often kept in aquaria but little is known about its distribution...



African pompano

Alectis ciliaris

The African pompano (Alectis ciliaris), also known as the pennant-fish or threadfin trevally, is a widely distributed species of tropical marine fish in the jack family, Carangidae. The species is found in tropical waters worldwide, with adults...



African sawtail catshark

Galeus polli

The African sawtail catshark (Galeus polli) is a species of catshark, part of the family Scyliorhinidae. Demersal in nature, it is found at depths of 160–720 m (520–2,360 ft) off the western African coast from Morocco to South Africa. This...



African threadfish

Alectis alexandrina

The African threadfish (Alectis alexandrina), also known as the Alexandria pompano, is a species of large marine fish in the jack family, Carangidae. The species is distributed along the coast of tropical Africa in the eastern Atlantic Ocean,...



Agamyxis pectinifrons

Agamyxis pectinifrons

Agamyxis pectinifrons, the spotted talking catfish, spotted raphael catfish or whitebarred catfish, is a species of thorny catfish found in the Amazon basin where it has been recorded from Bolivia, Brazil, Colombia and Peru. This species grows to...



Agonomalus jordani

Agonomalus jordani

Agonomalus jordani is a fish in the family Agonidae. It was described by David Starr Jordan and Edwin Chapin Starks in 1904. It is a marine, temperate water-dwelling fish which is known from the northwestern Pacific Ocean, including Japan, the...



Ailia coila

Ailia coila

Ailia coila, also known as the Gangetic ailia is a species of catfish in the family Ailiidae native to India, Bangladesh, Nepal and Pakistan. This species grows to a length of 30 centimetres (12 in) TL. Locally this fish is known as "kajoli in...



Alabama bass

Micropterus henshalli

Micropterus henshalli, the Alabama bass, is a medium-sized freshwater ray finned fish, a black bass from the genus Micropterus which is part of the sunfish family Centrarchidae. This species is endemic to the southeastern United States where it...



Alabama cavefish

Speoplatyrhinus poulsoni

The Alabama cavefish (Speoplatyrhinus poulsoni) is a critically endangered species of amblyopsid cavefish found only in underground pools in Key Cave, located in northwestern Alabama, United States in the Key Cave National Wildlife Refuge. It was...



Alabama shad

Alosa alabamae

The Alabama shad (Alosa alabamae) is an anadromous species of clupeid fish endemic to the United States where it breeds in medium to large flowing rivers from the Mississippi River drainage to the Suwannee River, Florida, as well as some other...



Alabama sturgeon

Scaphirhynchus suttkusi

The Alabama sturgeon (Scaphirhynchus suttkusi) is a species of sturgeon native to the United States of America and now only believed to exist in 130 miles (210 km) of the lower Alabama River. The fish has a distinctive yellowish-orange color....



Alaska blackfish

Dallia pectoralis

The Alaska blackfish (Dallia pectoralis) is a species of freshwater fish in the esocid family (Esocidae) of order Esociformes. It inhabits Arctic regions of Alaska as well as Siberia and the Bering Sea islands.



Alaska plaice

Pleuronectes quadrituberculatus

Alaska plaice (*Pleuronectes quadrituberculatus*) is a saltwater fish that live in the North Pacific Ocean. Alaska plaice are right-eye flounders which live on the sandy bottoms of the continental shelf, up to 600 metres deep. Their geographic...



Alaska pollock

Gadus chalcogrammus

The Alaska pollock or walleye pollock (*Gadus chalcogrammus*) is a marine fish species of the cod genus *Gadus* and family Gadidae. It is a semi-pelagic schooling fish widely distributed in the North Pacific, with largest concentrations found in the...



Alaska whitefish

Coregonus nelsonii

The Alaska whitefish (*Coregonus nelsonii*) is a species of whitefish in the family Salmonidae. It is found in parts of northwestern North America, where it occurs only in small and large rivers, and rarely in lakes. The maximum length recorded for...



Albacore

Thunnus alalunga

The albacore (*Thunnus alalunga*), known also as the longfin tuna, is a species of tuna of the order Scombriformes. It is found in temperate and tropical waters across the globe in the epipelagic and mesopelagic zones. There are six distinct stocks...



Albatrossia

Albatrossia pectoralis

Albatrossia pectoralis, the giant grenadier or giant rattail, is a very large rattail, and the only member of the genus *Albatrossia*. It is found in the north Pacific from northern Japan to the Okhotsk and Bering Seas, east to the Gulf of Alaska,...



Alburnoides bipunctatus

Alburnoides bipunctatus

Alburnoides bipunctatus, known vernacularly as the schneider, spirlin, bleak, riffle minnow, and others, is a species of small (9-cm average length) freshwater fish in the family Cyprinidae. It is found in Afghanistan, Armenia, Austria...



Alburnus arborella

Alburnus arborella

Alburnus arborella is a species of ray-finned fish in the carp family Cyprinidae. It is a freshwater fish occurring in lakes and streams in Europe. It is distributed in Croatia, Italy, Slovenia, and Switzerland. This species is up to 10...



Alburnus tarichi

Alburnus tarichi

Alburnus tarichi, known as the tarek, pearl mullet, Van fish or Van shah kuli, is a species of cyprinid fish, found only in Turkey, where it is the only fish known to inhabit Lake Van. It is endemic to the Lake Van basin. It is locally known in...



Alcolapia alcalica

Alcolapia alcalica

Alcolapia alcalica, the common natron tilapia or soda cichlid, is an endangered species of fish in the family Cichlidae. It is endemic to the hypersaline, warm Lake Natron in Ngorongoro District of Arusha Region, its drainage and the Shombole...



Alcolapia grahami

Alcolapia grahami

Alcolapia grahami, the Lake Magadi tilapia or Graham's cichlid, is a vulnerable species of fish in the family Cichlidae. It is specialised to live in hot, alkaline waters in springs and lagoons around hypersaline lakes.



Alcolapia latilabris

Alcolapia latilabris

Alcolapia latilabris, the wide-lipped Natron tilapia, is a species of small fish in the family Cichlidae. It is endemic to the hypersaline, warm Lake Natron in Tanzania. It lives near springs in the southern part of the lake. It reaches up to 6.2...



Alcolapia ndalalani

Alcolapia ndalalani

Alcolapia ndalalani, the narrow-mouthed Natron tilapia, is a species of small fish in the family Cichlidae. It is endemic to the hypersaline, warm Lake Natron in Tanzania. Here it lives in creeks and springs at the southern shores of the lake. It...



Alepisaurus ferox

Alepisaurus ferox

Alepisaurus ferox, also known as the long snouted lancetfish, longnose lancetfish, or cannibal fish, is a species of lancetfish found in the ocean depths down to 1,830 m (6,000 ft). This species grows to 215 cm (85 in) in total length and a...



Alestes baremoze

Alestes baremoze

Alestes baremoze, the pebbly fish or silversides, is a species of characin fish from the freshwater systems of northern and western Africa. It has some importance as a commercially exploited food fish.



Alewife

Alosa pseudoharengus

The alewife (Alosa pseudoharengus; pl.: alewives) is an anadromous species of herring found in North America. It is one of the "typical" North American shads, attributed to the subgenus Pomolobus of the genus Alosa. As an adult it is a marine...



Alfonsino

Beryx decadactylus

The alfonsino (*Beryx decadactylus*), also known as the alfonsin, longfinned beryx, red bream, or imperador, is a species of deepwater berycid fish of the order Beryciformes. It can be found in temperate and subtropical ocean waters nearly...



Allard's clownfish

Amphiprion allardi

Allard's clownfish or Allard's anemonefish (*Amphiprion allardi*) is a marine fish belonging to the family Pomacentridae, the clownfishes and damselfishes, from the western Indian Ocean off the coast of East Africa and the Mascarenes.



Alligator gar

Atractosteus spatula

The alligator gar (*Atractosteus spatula*) is a euryhaline ray-finned fish in the clade Ginglymodi of the infraclass Holostei, being most closely related to the bowfins. It is the largest species in the gar family (Lepisosteidae), and is among the...



Alligator pipefish

Syngnathoides biaculeatus

The alligator pipefish or double-ended pipefish (*Syngnathoides biaculeatus*) is a species of fish in the family Syngnathidae and is the only species in the monotypic genus *Syngnathoides*. It is found in shallow water in the tropical and subtropical...



Allis shad

Alosa alosa

The allis shad (*Alosa alosa*) is a widespread Northeast Atlantic species of fish in the herring family, Clupeidae. It is an anadromous fish which migrates into fresh water to spawn. It is found in the eastern Atlantic Ocean, the western Baltic Sea...



Alosa agone

Alosa agone

Alosa agone is a species of ray-finned fish in the genus Alosa. It is an endangered species.



Alosa braschnikowi

Alosa braschnikowi

Alosa braschnikowi, the Caspian marine shad or Brazhnikov's shad, is one of the clupeid fish species endemic to the Caspian Sea. This is a relatively large shad, typically 30 cm and up to 50 cm length. It also has a more "herring-like" body shape...



Alosa caspia

Alosa caspia

Alosa caspia is a species of clupeid fish, one of the species of shad (genus Alosa) endemic to the Caspian Sea basin. FishBase treats separately three subspecies from the Caspian: Caspian shad, Alosa caspia caspia (Eichwald, 1838) Enzeli shad,...



Alosa maeotica

Alosa maeotica

Alosa maeotica, known as the Black Sea shad or Azov shad, is a species of clupeid fish endemic to the Sea of Azov and the western part of the Black Sea basin. It is found in Bulgaria, Georgia, Moldova, Romania, Russia, Turkey, and Ukraine.This is...



Alpine bullhead

Cottus poecilopus

The alpine bullhead or Siberian bullhead (Cottus poecilopus) is a species of freshwater fish in the family Cottidae of sculpins. It is found in China, the Czech Republic, Denmark, Finland, Germany, North Korea, Moldova, Norway, Poland, Romania,...



Altolamprologus calvus

Altolamprologus calvus

Altolamprologus calvus is a cichlid endemic to the southern shoreline of Lake Tanganyika in eastern Africa. The species has an extremely laterally compressed body and a high dorsal fin. Males may grow to 13 cm (5 inches), while females are...



Altolamprologus compressiceps

Altolamprologus compressiceps

Altolamprologus compressiceps is a species of fish in the family Cichlidae. It is endemic to the shallow rocky areas of Lake Tanganyika. It is not considered threatened by the IUCN.



Aluterus scriptus

Aluterus scriptus

Aluterus scriptus, commonly known as scrawled filefish, broomtail filefish or scribbled leatherjacket, is a marine fish belonging to the family Monacanthidae.



Alvord cutthroat trout

Oncorhynchus clarki alvordensis

The Alvord cutthroat trout, *Oncorhynchus clarkii alvordensis*, was a subspecies of cutthroat trout. It was known only from Trout Creek in Oregon and Virgin Creek in Nevada, although it may have lived in several of the larger Alvord Basin streams...



Amatitlania sajica

Amatitlania sajica

Amatitlania sajica, the T-bar cichlid or Sajica cichlid, is a Central American species of cichlid found in freshwater streams and lakes on the Pacific slope of Costa Rica. The fish is tan colored with seven indistinct bars on the body. The third...



Amatitlania septemfasciata

Amatitlania septemfasciata

Amatitlania septemfasciata (sometimes called the Seven-stripe Cichlid or Cutter's Cichlid) is a fish species in the cichlid family. It is found on the Atlantic slope of Costa Rica, between the San Juan River drainage and the Banano River. Several...



Amatitlania siquia

Amatitlania siquia

Amatitlania siquia', is a species of cichlid native to Central America. These cichlid can be identified by black melanin-based stripes in males and an additional bright orange spot on the abdomen for females. The size and pigmentation of the spots...



Amazon leafyfish

Monocirrhus polyacanthus

The Amazon leaffish (*Monocirrhus polyacanthus*), or South American leaffish, is a species of fish belonging to the family Polycentridae. It is found in the Amazon basin in Bolivia, Brazil, Colombia, Peru and Venezuela, where it inhabits edges of...



Amazon molly

Poecilia formosa

The Amazon molly (Poecilia formosa) is a freshwater fish native to warm, fresh waters between Tuxpan River in northeastern Mexico and the Rio Grande and the Nueces River in the southern parts of the U.S. state of Texas. It reproduces through...



Amazonsprattus

Amazonsprattus scintilla

Amazonsprattus scintilla, the Rio Negro pygmy anchovy, is a fresh-water anchovy that is endemic to the Amazon River system in South America.



Amblygaster clupeioides

Amblygaster clupeioides

Amblygaster clupeioides, the bleeker smoothbelly sardinella, blue pilchard, sharp-nosed pilchard, or sharpnose sardine, is a reef-associated marine species of sardinellas in the herring family Clupeidae. It is one of the three species of genus...



Amblygaster sirm

Amblygaster sirm

Amblygaster sirm, the spotted sardinella, also known as the northern pilchard, spotted pilchard, spotted sardine, and trenched sardine, is a reef-associated marine species of sardinellas in the herring family Clupeidae.



Amblyglyphidodon aureus

Amblyglyphidodon aureus

Amblyglyphidodon aureus also known as the golden damselfish is a species of marine fish in the family Pomacentridae, the damselfishes and clownfishes. It is native to the central Indo-Pacific.



Amblyglyphidodon curacao

Amblyglyphidodon curacao

Amblyglyphidodon curacao; also known as the staghorn damselfish , the clouded damselfish or the black-snouted sergeant major is a species of marine fish in the family Pomacentridae, the damselfishes and the clownfishes. It is widespread...



Amblygobius phalaena

Amblygobius phalaena

Amblygobius phalaena, the Sleeper Banded goby, white-barred goby, is a species of goby native to tropical reefs of the western Pacific Ocean and through the central Indo-Pacific area at depths of from 2 to 20 metres (6.6 to 65.6 ft). This species...



Ameca shiner

Notropis amecae

The Ameca shiner (Notropis amecae) is a species of cyprinid fish in the family Cyprinidae. The Ameca shiner was described in 1986 from upper parts of the Ameca River drainage in Jalisco, Mexico. Although already feared extinct by 1969, and listed...



Ameiurus platycephalus

Ameiurus platycephalus

Ameiurus platycephalus, the flat bullhead, is a species of North American freshwater catfish native to the waters of the southeastern United States from Virginia to Georgia. This species grows to a maximum length of 29 cm (11 in) TL though it is...



American butterfish

Peprilus triacanthus

The American butterfish (Peprilus triacanthus), also known as the Atlantic butterfish, is a butterfish of the family Stromateidae.



American conger

Conger oceanicus

The American conger (Conger oceanicus) is a species of eel in the family Congridae. Other common names for this fish include conger, dog eel, poison eel and sea eel. It is a marine fish with a widespread distribution in the Western Atlantic from...



American eel

Anguilla rostrata

The American eel (*Anguilla rostrata*) is a facultative catadromous fish found on the eastern coast of North America. Freshwater eels are fish belonging to the elopomorph superorder, a group of phylogenetically ancient teleosts. The American eel...



American gizzard shad

Dorosoma cepedianum

The American gizzard shad (*Dorosoma cepedianum*), also known as the mud shad, is a member of the herring family of fish and is native to large swaths of fresh and brackish waters in the United States of America, as well as portions of Quebec....



American paddlefish

Polyodon spathula

The American paddlefish (*Polyodon spathula*, also known as a Mississippi paddlefish, spoon-billed cat, or spoonbill) is a species of ray-finned fish. It is the last living species of paddlefish (*Polyodontidae*). This family is most closely related...



American pickerel

Esox americanus

The American pickerels are two subspecies of *Esox americanus*, a medium-sized species of North American freshwater predatory fish belonging to the pike family (genus *Esox* in family *Esocidae* of order *Esociformes*): Redfin pickerel, sometimes called...



American plaice

Hippoglossoides platessoides

The American plaice, American sole or long rough dab (*Hippoglossoides platessoides*) is a North Atlantic flatfish that belongs, along with other right-eyed flounders, to the family *Pleuronectidae*. In the northwest Atlantic (*H. p. platessoides*) it...



American shad

Alosa sapidissima

The American shad (Alosa sapidissima) is a species of anadromous clupeid fish naturally distributed on the North American coast of the North Atlantic, from Newfoundland to Florida, and as an introduced species on the North Pacific coast. The...



Ammodytes americanus

Ammodytes americanus

Ammodytes americanus, also known as American sand lance, American sand eel, and sand launce, is a small fish in the family Ammodytidae. First described by James Ellsworth De Kay in 1842, it is widespread in the western North Atlantic. Like all...



Amphilophus citrinellus

Amphilophus citrinellus

Amphilophus citrinellus is a large cichlid fish endemic to the San Juan River and adjacent watersheds in Costa Rica and Nicaragua. In the aquarium trade A. citrinellus is often sold under the trade name of Midas cichlid. A. citrinellus are...



Amphilophus labiatus

Amphilophus labiatus

Amphilophus labiatus is a large cichlid fish endemic to Lake Managua and Lake Nicaragua in Central America. It is also known by the common name red devil cichlid, which it shares with another closely related cichlid, A. citrinellus.



Amphilophus zaliosus

Amphilophus zaliosus

Amphilophus zaliosus is a species of cichlid that inhabits Lake Apoyo in Nicaragua. It is known in the aquarium trade as the arrow cichlid. It is an elongate species in the Midas cichlid species complex. The arrow cichlid shares its habitat with...



Amphiprion akallopisos

Amphiprion akallopisos

The nosestripe clownfish or nosestripe anemonefish, skunk clownfish, Amphiprion akallopisos, is an anemonefish (also called clownfish) that lives in association with sea anemones. A. akallopisos is found in the Indian Ocean. It resides in shallow...



Amphiprion akindynos

Amphiprion akindynos

Amphiprion akindynos, the Barrier Reef anemonefish, is a species of anemonefish that is principally found in the Great Barrier Reef of Australia, but also in nearby locations in the Western Pacific. The species name 'akindynos' is Greek, meaning...



Amphiprion barberi

Amphiprion barberi

Amphiprion barberi, also known as fiji anemonefish a species of anemonefish that is found in the western Pacific Ocean. It was previously considered a geographic color variation of other anemonefish, initially Amphiprion rubrocinctus from 1972...



Amphiprion chagosensis

Amphiprion chagosensis

Amphiprion chagosensis, the Chagos anemonefish, is a marine fish belonging to the family Pomacentridae, the clownfishes and damselfishes. It is named for the Chagos Archipelago in the Indian Ocean and it is endemic to the archipelago. The...



Amphiprion chrysogaster

Amphiprion chrysogaster

Amphiprion chrysogaster, the Mauritian anemonefish, is a marine fish belonging to the family Pomacentridae, the clownfishes and damselfishes. It is endemic to Mauritius and probably Réunion.



Amphiprion fuscocaudatus

Amphiprion fuscocaudatus

Amphiprion fuscocaudatus (the Seychelles anemonefish) is a marine fish belonging to the family Pomacentridae, the clownfishes and damselfishes.



Amphiprion latezonatus

Amphiprion latezonatus

Amphiprion latezonatus, also known as the wide-band anemonefish, is a species of anemonefish found in subtropical waters off the east coast of Australia. Like all anemonefishes, it forms a symbiotic mutualism with sea anemones and is unaffected...



Amphiprion latifasciatus

Amphiprion latifasciatus

Amphiprion latifasciatus (Madagascar anemonefish) is a marine fish belonging to the family Pomacentridae, the clownfishes and damselfishes.



Amphiprion leucokranos

Amphiprion leucokranos

Amphiprion leucokranos (White bonnet anemonefish) is a naturally occurring hybrid anemonefish found in the western central Pacific Ocean. Like all anemonefishes it forms a symbiotic mutualism with sea anemones and is unaffected by the stinging...



Amphiprion mccullochi

Amphiprion mccullochi

Amphiprion mccullochi, also known as whitesnout anemonefish or McCulloch's anemonefish, is a species of anemonefish found in subtropical waters at Lord Howe Island and Norfolk Island.. It was named for Allan McCulloch, a former Curator of Fishes...



Amphiprion nigripes

Amphiprion nigripes

Amphiprion nigripes (Maldive anemonefish or blackfinned anemonefish), is a marine fish belonging to the family Pomacentridae, which includes clownfishes and damselfishes.



Amphiprion omanensis

Amphiprion omanensis

Amphiprion omanensis (Oman anemonefish) is a marine fish belonging to the family Pomacentridae, the clownfishes and damselfishes.



Amphiprion pacificus

Amphiprion pacificus

Amphiprion pacificus, is a species of anemonefish that is found in the western Pacific Ocean. Like all anemonefishes it forms a symbiotic mutualism with sea anemones and is unaffected by the stinging tentacles of the host anemone. It is a...



Amphiprion thiellei

Amphiprion thiellei

Amphiprion thiellei (Thielle's anemonefish) is thought to be a naturally occurring hybrid anemonefish found in the vicinity of Cebu, Philippines. Like all anemonefishes it forms a symbiotic mutualism with sea anemones and is unaffected by the...



Amu Darya sturgeon

Pseudoscaphirhynchus kaufmanni

The Amu Darya sturgeon or false shovelnose sturgeon (Pseudoscaphirhynchus kaufmanni) is a critically endangered species of fish in the family Acipenseridae. It is found in Turkmenistan, Uzbekistan, Tajikistan and perhaps Afghanistan. It inhabits...



Amur bitterling

Rhodeus sericeus

The Amur bitterling (Rhodeus sericeus) is a small fish of the carp family. It is sometimes just called "bitterling", which dates back to the time when the European bitterling (Rhodeus amarus) was still considered conspecific with R. sericeus, and...



Amur catfish

Silurus asotus

Silurus asotus, commonly called the Japanese common catfish, Amur catfish, Far Eastern catfish, and Chinese catfish, is a catfish species belonging to the Siluridae family. It lives in widespread distribution in freshwater habitats across East...



Amur pike

Esox reichertii

The Amur pike (*Esox reichertii*), also known as the blackspotted pike, is a pike native to the Amur River system in east Asia, as well as freshwater habitat on the island of Sakhalin. Closely related to the northern pike, it reaches a length of...



Anabas cobojius

Anabas cobojius

Anabas cobojius, the Gangetic koi, popularly known as Koi in Bengali is a species of climbing gourami native to Bangladesh and India, where it occurs in many types of standing water bodies. This species reaches a total length of 30 cm (12 in) and...



Anabas testudineus

Anabas testudineus

The climbing perch (Anabas testudineus) is a species of amphibious freshwater fish in the family Anabantidae (the climbing gouramis). A labyrinth fish native to Far Eastern Asia, the fish inhabits freshwater systems from Pakistan, India,...



Anableps anableps

Anableps anableps

Anableps anableps, the largescale four-eyes, is a species of four-eyed fish found in fresh and brackish waters of northern South America and Trinidad. This species grows to a length of 30 centimetres (12 in) TL. This fish can occasionally be...



Anaocypris hispanica

Anaocypris hispanica

Anaocypris hispanica, the Spanish minnowcarp, is a small species of ray-finned fish species in the family Cyprinidae. It is the only living member of the genus Anaocypris. It is endemic to the Iberian Peninsula and is found in the basin of the...



Anarchias seychellensis

Anarchias seychellensis

Anarchias seychellensis is a moray eel found in coral reefs in the Pacific and Indian Oceans. It was first named by J. L. B. Smith in 1962, and is commonly known as the Seychelles moray or the marbled reef-eel.



Anarhichas minor

Anarhichas minor

Anarhichas minor, the spotted wolffish, spotted sea cat or leopardfish, is a large marine fish of the family Anarhichadidae. This bottom-dwelling species is found across the North Atlantic and adjacent parts of the Arctic Ocean from north of...



Anarhichas orientalis

Anarhichas orientalis

Anarhichas orientalis, the Bering wolffish, is a marine fish in the family Anarhichadidae, the "wolffish".



Anchoa hepsetus

Anchoa hepsetus

Anchoa hepsetus, commonly known as the broad-striped anchovy, is a species of ray-finned fish in the family Engraulidae. It grows to be about 15 cm (6 in) long. The broad-striped anchovy is found in the west Atlantic along the North American...



Anchoa mitchilli

Anchoa mitchilli

Anchoa mitchilli is a species of ray-finned fish in the family Engraulidae, the anchovies. Its common names include bay anchovy and common anchovy. It is native to the western Atlantic Ocean and Gulf of Mexico. It is one of the most common fish...



Ancistrus cirrhosus

Ancistrus cirrhosus

Ancistrus cirrhosus, the Jumbie teta, is a species of armored catfish found in the Paraná River basin of Argentina and Uruguay. This species grows to a length of 8.9 centimetres (3.5 in) SL.



Ancistrus hoplogenyys

Ancistrus hoplogenyys

Ancistrus hoplogenyys is a species of armored catfish native to Argentina, Brazil, Guyana, Paraguay, Peru, and Uruguay, where it occurs in the basins of the Amazon River, the Essequibo River, and the Paraguay River. It has also been reported from...



Ancistrus temminckii

Ancistrus temminckii

Ancistrus temminckii is a species of armoured suckermouth catfish in the family Loricariidae of order Siluriformes. It is found in Suriname in the Saramacca River, Suriname River and Maroni River basins. Males grow to about 10 centimetres (3.9...



Andinoacara latifrons

Andinoacara latifrons

Andinoacara latifrons, the platinum acara, is a species of fish in the family Cichlidae in the order Perciformes, native to the Magdalena, Atrato, Sinú, and San Juan River basins in Colombia. The scientific species name is indicative to its...



Andinoacara stalsbergi

Andinoacara stalsbergi

Andinoacara stalsbergi is a species of South American freshwater fish in the family Cichlidae. It was previously included in A. rivulatus, but was described as a separate species in 2009. The specific name honours the Norwegian aquarist Alf...



Anguilla bengalensis

Anguilla bengalensis

The mottled eel (Anguilla bengalensis), also known as the African mottled eel, the Indian longfin eel, the Indian mottled eel, the long-finned eel or the river eel, is a demersal, catadromous eel in the family Anguillidae. It was described by...



Anguilla bengalensis bengalensis

Anguilla bengalensis bengalensis

Anguilla bengalensis bengalensis, the Indian mottled eel, is a subspecies of eel in the genus Anguilla of the family Anguillidae. It is found throughout the Indian subcontinent and neighbouring regions including the East Indies. Showing the...



Anguilla bicolor

Anguilla bicolor

Anguilla bicolor is a species of eel in the genus Anguilla of the family Anguillidae, consisting of two subspecies.



Angular roughshark

Oxynotus centrina

Sharks portal The angular roughshark (*Oxynotus centrina*) is a rough shark of the family *Oxynotidae*.



Anisotremus virginicus

Anisotremus virginicus

Anisotremus virginicus, the porkfish, also known as the Atlantic porkfish sweetlips, dogfish or paragrater grunt, is a species of marine ray-finned fish, a grunt belonging to the family Haemulidae. It is native to the western Atlantic Ocean.



Anomalochromis

Anomalochromis thomasi

Anomalochromis is a genus of fish in the family Cichlidae, containing the single species Anomalochromis thomasi, the African butterfly cichlid. It is a small cichlid growing to a length of 6–8 centimetres (2.4–3.1 in). The natural habitat of A....



Anoplogaster cornuta

Anoplogaster cornuta

Anoplogaster cornuta, the common fangtooth, is a species of deep sea fish found in temperate and tropical oceans worldwide. It is found at depths of from 2 to 5,000 metres (10 to 16,400 ft) with the adults usually found from 500 to 5,000 metres...



Anostomus anostomus

Anostomus anostomus

Anostomus anostomus, also known as the striped headstander, striped anostomus, and striped anastomus, is a species of fish.



Antarctic silverfish

Pleuragramma antarctica

The Antarctic silverfish (Pleuragramma antarcticum), or Antarctic herring, is a species of marine ray-finned fish belonging to the family Nototheniidae, the notothen or cod icefishes. It is native to the Southern Ocean and the only truly pelagic...



Antarctic toothfish

Dissostichus mawsoni

The Antarctic toothfish (Dissostichus mawsoni), also known as the Antarctic cod, is a large, black or brown fish found in very cold (subzero) waters of the Southern Ocean near Antarctica. It is the largest fish in the Southern Ocean, feeding on...



Antennatus coccineus

Antennatus coccineus

Antennatus coccineus, the scarlet or freckled frogfish, is a species of frogfish originally classified as Chironectes coccineus and Antennarius coccineus. It lives within tropical waters and has a central distribution being around...



Anthias anthias

Anthias anthias

Anthias anthias, the swallowtail sea perch or marine goldfish, is a species of marine ray-finned fish from the grouper and sea bass family Serranidae. It is native to the eastern Atlantic Ocean and the Mediterranean Sea where it is associated...



Apache trout

Oncorhynchus apache

The Apache trout or Arizona trout, *Oncorhynchus apache*, is a species of freshwater fish in the salmon family (family Salmonidae) of order Salmoniformes. It is one of the Pacific trouts.



Aphanius danfordii

Aphanius danfordii

Aphanius danfordii, the Kızılırmak toothcarp or Sultan Sazlığı toothcarp, is a species of killifish belonging to the family Cyprinodontidae. It is endemic to the Kızılırmak River and the upper Seyhan River drainage systems and is now restricted...



Aphanius transgrediens

Aphanius transgrediens

Aphanius transgrediens, the Acı Göl toothcarp or Acipinar killifish, is a species of freshwater fish in the family Cyprinodontidae. It is endemic to the springs of Lake Acıgöl in Turkey. It is threatened by a reduction in rainfall from climate...



Aphareus furca

Aphareus furca

Aphareus furca, the small toothed jobfish, blue smalltooth jobfish, fork-tailed snapper or snapper jobbyfish is a species of marine ray-finned fish, a snapper belonging to the family Lutjanidae. It is found in the Indo-Pacific region.



Aphyosemion australe

Aphyosemion australe

Aphyosemion australe, the lyretail panchax, golden panchax or Cape Lopez lyretail, is a species of freshwater fish belonging to the family Aplocheilidae. It is found around Cape Lopez and in surrounding areas in Gabon.



Aphyosemion bivittatum

Aphyosemion bivittatum

Aphyosemion bivittatum (twostripe lyretail, two-banded killi, red lyretail) is a species of freshwater fish belonging to the family Aplocheilidae. It is found in rivers in southeastern Nigeria and southwestern Cameroon. It was originally...



Apistogramma agassizii

Apistogramma agassizii

Apistogramma agassizii, commonly known as Agassiz's dwarf cichlid, is a species of cichlid found in the Marañón and Ucayali River in Peru, some tributaries of the Amazon River, as well as downstream to the estuary in the Atlantic. It is named...



Apistogramma borellii

Apistogramma borellii

Apistogramma borellii, common names umbrella cichlid, umbrella apisto and yellow dwarf cichlid, is a species of fish in the genus Apistogramma. It is a South American dwarf cichlid, found in the Paraguay River and Paraná River. They have been...



Apistogramma cacatuoides

Apistogramma cacatuoides

Apistogramma cacatuoides or the cockatoo dwarf cichlid is a South American cichlid and the Apistogramma species most commonly bred in captivity.



Apistogramma hongsloui

Apistogramma hongsloui

Apistogramma hongsloui is a species of dwarf cichlid fish, native to the Orinoco basin in South America. They are often kept in the aquariums and prefer to have soft, acidic water.



Apistogramma macmasteri

Apistogramma macmasteri

Apistogramma macmasteri is a dwarf cichlid in the tribe Geophagini, one of the tribes of the subfamily of American cichlids, the Cichlinae. It is a freshwater fish that lives in the rivers Guaytiquía and Metica in the Meta River system. The Meta...



Apistogramma nijsseni

Apistogramma nijsseni

Apistogramma nijsseni is a species of cichlid fish, endemic to highly restricted local black water habitats in the Quebrada Carahuayte, a small stream in the Ucayali River drainage, southern Peru. The male reaches a maximum length of 8 cm (3 in),...



Apistogramma panduro

Apistogramma panduro

Apistogramma panduro (A-183), also commonly referred to as Apistogramma pandurini, is a small species of cichlid freshwater fish from the Amazon river basin in Peru. It is closely related to A. nijsseni, so much so that they are placed in the A....



Apistogramma viejita

Apistogramma viejita

Apistogramma viejita is a dwarf cichlid in the subfamily Cichlinae, in the tribe Geophagini. It is a benthopelagic freshwater fish that lives in the Meta River and the Orinoco River in Colombia. They grow up to 4.6 cm in standard length. The males...



Apocryptes

Apocryptes bato

Apocryptes bato is a species of mudskipper native to India, Bangladesh and Myanmar where it can be found in tropical rivers, estuaries and coastal waters of the Indian Ocean. This species can reach a length of 26 centimetres (10 in) TL. It is of...



Apogon imberbis

Apogon imberbis

Apogon imberbis, the cardinalfish, the Mediterranean cardinalfish or king of the mullets, is a species of ray-finned fish, a cardinalfish belonging to the family Apogonidae. It is widely distributed in the Mediterranean and along the warm...



Apogon maculatus

Apogon maculatus

Apogon maculatus, commonly known as flamefish, is a species of cardinalfish from the western Atlantic. Apogon maculatus is a nocturnal fish, and it usually hides in shadowy areas. This fish can be viewed with a red light in a dark room if needed...



Apolemichthys trimaculatus

Apolemichthys trimaculatus

Apolemichthys trimaculatus, the threespot angelfish or flagfin angelfish, is a demersal marine ray-finned fish, a marine angelfish belonging to the family Pomacanthidae. It has a wide Indo-Pacific distribution.



Apollo sharkminnow

Luciosoma spilopleura

The Apollo sharkminnow (*Luciosoma spilopleura*) is a species of cyprinid fish found in Thailand, Sumatra and Borneo.



Aquitanian pike

Esox aquitanicus

The Aquitanian pike (Esox aquitanicus) is a species of pike, a predatory fish native from the Charente to the Adour river drainages in the French Great South-West region. It was scientifically described in 2014, but had formerly been regarded as...



Arabian carpetshark

Chiloscyllium arabicum

The Arabian carpetshark (Chiloscyllium arabicum) is a species of carpet shark in the family Hemiscylliidae, inhabiting coral reefs and other shallow coastal habitats from the Persian Gulf to India. Reaching 78 cm (31 in) long, this shark is...



Arabian toothcarp

Aphaniops dispar

The Arabian toothcarp (Aphaniops dispar), known also as the Arabian pupfish or mother-of-Pearl fish is a species of killifish belonging to the family Cyprinodontidae. It can be found from the shores of the Red Sea south to Ethiopia, the Gulf of...



Aral barbel

Luciobarbus brachycephalus

The Aral barbel (Luciobarbus brachycephalus) is a species of ray-finned fish in the genus Luciobarbus. It is found in the Aral basin, Chu drainage and southern and western Caspian Sea. For spawning, it migrates up to larger tributaries of the...



Arapaima gigas

Arapaima gigas

Arapaima gigas, also known as pirarucu or simply arapaima, is a species of arapaima native to the basin of the Amazon River. Once believed to be the sole species in the genus, it is among the largest freshwater fish. The species is an obligate...



Arapaima leptosoma

Arapaima leptosoma

Arapaima leptosoma is a species of freshwater fish endemic to Brazil. It is a member of the arapaimas, a genus of air-breathing fish that contains some of the world's largest freshwater fish. Like other members of the genus Arapaima, this fish...



Arc-eye hawkfish

Paracirrhites arcatus

The arc-eye hawkfish (Paracirrhites arcatus), the ringeye hawkfish, horseshoe hawkfish or whiteline hawkfish, is a species of marine ray-finned fish, a hawkfish belonging to the family Cirrhitidae. It is found in shallow waters in the tropical...



Archocentrus

Archocentrus centrarchus

Archocentrus is a genus of cichlid fish from Central America. It currently contains a single species, the flier cichlid (Archocentrus centrarchus), which is found in stagnant and slow-moving freshwater habitats such as lakes, ponds, ditches,...



Archosargus probatocephalus

Archosargus probatocephalus

Archosargus probatocephalus, the sheepshead, sheepshead seabream or convict fish, is a species of marine ray-finned fish belonging to the family Sparidae, the seabreams and porgies. This species is found in the Western Atlantic Ocean. The...



Arctic char

Salvelinus alpinus

The Arctic char or Arctic charr (Salvelinus alpinus) is a cold-water fish in the family Salmonidae, native to alpine lakes, as well as Arctic and subarctic coastal waters in the Holarctic.



Arctic cisco

Coregonus autumnalis

Arctic cisco (Coregonus autumnalis), also known as omul Russian: Омуль, is an anadromous species of freshwater whitefish that inhabits the Arctic parts of Siberia, Alaska and Canada. It has a close freshwater relative in several lakes of Ireland,...



Arctic flounder

Liopsetta glacialis

The Arctic flounder (Liopsetta glacialis), also known as the Christmas flounder, eelback flounder and Polar plaice, is a flatfish of the family Pleuronectidae. It is a demersal fish that lives on coastal mud bottoms in salt, brackish and fresh...



Arctic grayling

Thymallus arcticus

The Arctic grayling (Thymallus arcticus) is a species of freshwater fish in the salmon family Salmonidae. T. arcticus is widespread throughout the Arctic and Pacific drainages in Canada, Alaska, and Siberia, as well as the upper Missouri River...



Arctic skate

Amblyraja hyperborea

The Arctic skate (Amblyraja hyperborea) is a species of fish in the family Rajidae. It lives near the seabed between 140 and 2,500 m deep in the Arctic Ocean and waters around Canada and northern and north-western Europe, in the northern Pacific...



Arctogadus

Arctogadus glacialis

Arctogadus glacialis, known also with ambiguous common names Arctic cod and polar cod, is an Arctic species of fish in the cod family Gadidae, related to the true cod (genus *Gadus*). *Arctogadus glacialis* is found in icy water. They grow to about...



Areolate grouper

Epinephelus areolatus

The areolate grouper (*Epinephelus areolatus*), also known as the yellowspotted rockcod, areolate rockcod, green-spotted rock-cod, squaretail grouper or squaretail rock-cod, is a species of marine ray-finned fish, a grouper from the subfamily...



Argentine anchoita

Engraulis anchoita

The Argentine anchoita (*Engraulis anchoita*) or Argentine anchovy is an anchovy of the genus *Engraulis*, found in and around waters of Argentina, Uruguay and southern Brazil.



Argentine hake

Merluccius hubbsi

The Argentine hake (Merluccius hubbsi) is a merluccid hake of the genus Merluccius, found in the southwestern Atlantic Ocean, along the coast of Argentina, and Uruguay. This fish was described by an Argentine ichthyologist, Tomás Marini in...



Argyrolepecus aculeatus

Argyrolepecus aculeatus

Argyrolepecus aculeatus, the lovely hatchetfish or Atlantic silver hatchetfish, is a species of fish in the family Sternoptychidae. It may exceed 70 millimetres (2.8 in) standard length (SL). It lives in the mesopelagic zone of all oceans and...



Argyrolepecus affinis

Argyrolepecus affinis

Argyrolepecus affinis is a species of ray-finned fish in the family Sternoptychidae, found in the tropical and subtropical Atlantic, Indian and Pacific Oceans. Common names for this fish include Pacific hatchetfish, deepsea hatchetfish and...



Argyrolepecus gigas

Argyrolepecus gigas

Argyrolepecus gigas, the giant hatchetfish or greater silver hatchetfish, is a marine fish of the genus Argyrolepecus. It is found in every ocean except the north Pacific in the mesopelagic zone of tropical and subtropical waters. "Giant" in...



Argyrolepecus hemigymnus

Argyrolepecus hemigymnus

Argyrolepecus hemigymnus, the half-naked hatchetfish, short silver hatchetfish or spurred hatchetfish, is a deep-sea hatchetfish of the genus Argyrolepecus found mesopelagically in the Atlantic, Indian, and Pacific Oceans as well as in the...



Argyrosomus japonicus

Argyrosomus japonicus

Argyrosomus japonicus is a silvery to bronze-green colored fish, a member of the family Sciaenidae, which may grow up to 2 m (6 ft 7 in) in length. It is known as Japanese meagre (FAO), mullo way or jewfish in the eastern states of Australia,...



Argyrosomus regius

Argyrosomus regius

Argyrosomus regius, also known as the meagre, croaker, jewfish, shade-fish, sowa, kir, corvina, salmon-bass or stone bass, is a species of fish of the family Sciaenidae. This large fish has a pearly-silver to brownish coloration and a...



Aristotle's catfish

Silurus aristotelis

Aristotle's catfish (Silurus aristotelis) is a species of fish in the family Siluridae. It is endemic to Greece, where it occurs in the Acheloos River drainage. Its natural habitat is freshwater lakes. It is threatened by habitat loss. This...



Arius manillensis

Arius manillensis

Arius manillensis is a species of marine catfish endemic to the island of Luzon, Philippines. It is commonly known as the Manila sea catfish or kanduli. It is fished commercially.



Arnoldichthys

Arnoldichthys spilopterus

Arnoldichthys spilopterus, the Niger tetra, is a characin endemic to Nigeria. It is the only member of its genus.



Arothron caeruleopunctatus

Arothron caeruleopunctatus

Arothron caeruleopunctatus, also known as the blue-spotted pufferfish, is a demersal marine fish belonging to the family Tetraodontidae.



Arothron meleagris

Arothron meleagris

Arothron meleagris, commonly known as the guineafowl puffer or golden puffer, is a pufferfish from the Indo-Pacific, and Eastern Pacific. It is occasionally harvested for the aquarium trade. It reaches 50 cm in length. Guineafowl puffers have...



Arothron reticularis

Arothron reticularis

Arothron reticularis, variously known as the reticulated pufferfish, reticulated blowfish or reticulated toadfish, is a ray-finned fish in the family Tetraodontidae. It is native to the tropical and sub-tropical Indo-Pacific region where its...



Arothron stellatus

Arothron stellatus

Arothron stellatus, also known as the stellate pufferfish, starry puffer, or starry toadfish, is a demersal marine fish belonging to the family Tetraodontidae. It is found in shallow water in the Indo-Pacific region.



Arripis trutta

Arripis trutta

Arripis trutta, known as the Australian salmon in Australia and as kahawai in New Zealand, is a South Pacific marine fish and one of the four extant species within the genus Arripis, native to the cooler waters around the southeastern Australian...



Arrow goby

Clevelandia ios

The arrow goby (Clevelandia ios) is a species of goby native to marine and brackish waters of the Pacific coast of North America from British Columbia to Baja California. This species grows to a length of 6.4 centimetres (2.5 in) SL, though most...



Arrowhead dogfish

Deania profundorum

Sharks portalThe arrowhead dogfish (Deania profundorum) is a small little known deepwater dogfish of the family Centrophoridae.



Arrowtooth flounder

Atheresthes stomias

The arrowtooth flounder (Atheresthes stomas) is a fish in the family Pleuronectidae. It can be caught from the Bering Sea to Santa Rosa Island, California. At present, it is the most common fish in the Gulf of Alaska. Data is insufficient for...



Arroyo chub

Gila orcuttii

The arroyo chub (Gila orcuttii) is a species of cyprinid fish found only in the coastal streams of southern California, United States. The shape of the arroyo chub is somewhat chunky, with a deep body and thick, caudal peduncle. The eyes are...



Arulius barb

Dawkinsia arulius

The Arulius barb (Dawkinsia arulius) is a tropical cyprinid fish native to the Kaveri River basin of south east India. Other common names include Tamiraparani barb, Silas barb and longfin barb.



Asian arowana

Scleropages formosus

The Asian arowana (Scleropages formosus) comprises several phenotypic varieties of freshwater fish distributed geographically across Southeast Asia. While most consider the different varieties to belong to a single species, work by Pouyaud et al....



Asian sheepshead wrasse

Semicossyphus reticulatus

The Asian sheephead wrasse (*Semicossyphus reticulatus*) is one of the largest species of wrasse. Native to the western Pacific Ocean, it inhabits rocky reef areas around the Korean Peninsula, China, Japan, and the Ogasawara Islands. It can reach...



Asian swamp eel

Monopterus albus

The Asian swamp eel (*Monopterus albus*), also known as rice eel, ricefield eel, rice paddy eel or white rice-field eel, is a commercially important, air-breathing species of fish in the family Synbranchidae. It occurs in East and Southeast Asia,...



Asoka barb

Systemus asoka

The Asoka barb (Systemus asoka) is a species of cyprinid fish endemic to Sri Lanka where it is found in the upper reaches of the Sitawaka River and its tributaries, and Kelani near Kitulgala. This fish can reach a length of 17 centimetres (6.7...



Asp

Leuciscus aspius

The asp (Leuciscus aspius) is a European freshwater fish of the Cyprinidae family. It is sometimes considered by taxonomic authorities to be one of two members of the genus Aspius. It is protected under Appendix III of the Bern Convention and...



Aspidoras fuscoguttatus

Aspidoras fuscoguttatus

Aspidoras fuscoguttatus or Brown-point Shield Skin Longirostris is a tropical freshwater fish belonging to the Corydoradinae sub-family of the family Callichthyidae. It originates in inland waters in South America, and is found in the upper...



Aspiolucius esocinus

Aspiolucius esocinus

Aspiolucius esocinus or the pike asp is a species of cyprinid fish native to rivers, such as Amu Darya and Syr Darya, and occasionally lakes in Tajikistan, Kyrgyzstan, Turkmenistan, and Uzbekistan, where it was a common fish until 1980s. After...



Astatotilapia burtoni

Astatotilapia burtoni

Astatotilapia burtoni is a species of fish in the family Cichlidae. It is found in Lake Tanganyika and its surrounding waterways, including parts of Burundi, Rwanda, Tanzania, and Zambia. Its natural habitats are rivers, intermittent rivers,....



Astatotilapia flavijosephi

Astatotilapia flavijosephi

Astatotilapia flavijosephi, the Jordan mouthbrooder, is a vulnerable species of freshwater fish in the family Cichlidae (cichlids). It is found in the central Jordan River system, including Lake Tiberias (Kinneret), in Israel, Jordan and Syria,...



Asterophysus batrachus

Asterophysus batrachus

Asterophysus batrachus, the gulper catfish or ogre catfish, is a species of catfish (order Siluriformes) of the family Auchenipteridae. It is native to the Rio Negro and Orinoco basins in Brazil, Colombia and Venezuela, where mostly found in...



Astronesthes niger

Astronesthes niger

Astronesthes niger, commonly known as snaggletooth, is a species of small, deep sea fish in the family Stomiidae. It occurs in the tropical and subtropical Atlantic Ocean, the Caribbean Sea and the Gulf of Mexico, as well as the Indian Ocean and...



Astronotus crassipinnis

Astronotus crassipinnis

Astronotus crassipinnis is a South American fish in the cichlid family from the southern Amazon basin and the Paraná–Paraguay basins. It is not as well-known or common in the aquarium trade as its relative, the more northerly distributed oscar...



Astroscopus guttatus

Astroscopus guttatus

Astroscopus guttatus (northern stargazer) is a fish that can reach lengths of 22 inches (56 cm) and is found on the Atlantic shores between the states of North Carolina and New York in the United States. The northern stargazer can be found up to...



Astyanax jordani

Astyanax jordani

Astyanax jordani is a freshwater fish of the characin family (family Characidae) of order Characiformes, native to Mexico. It is sometimes called the cave tetra, or by its local Spanish name tetra ciego. A blind cave fish, A. jordani is very...



Atlantic blue marlin

Makaira nigricans

The Atlantic blue marlin (*Makaira nigricans*) is a species of marlin endemic to the Atlantic Ocean. It is closely related to, and usually considered conspecific with, the Indo-Pacific blue marlin, then simply called blue marlin. Some authorities...



Atlantic bluefin tuna

Thunnus thynnus

The Atlantic bluefin tuna (*Thunnus thynnus*) is a species of tuna in the family Scombridae. It is variously known as the northern bluefin tuna (mainly when including Pacific bluefin as a subspecies), giant bluefin tuna, and formerly as the...



Atlantic bonito

Sarda sarda

The Atlantic bonito (*Sarda sarda*) is a large mackerel-like fish of the family Scombridae. It is common in shallow waters of the Atlantic Ocean, the Mediterranean Sea, and the Black Sea, where it is an important commercial and game fish.



Atlantic bumper

Chloroscombrus chrysurus

The Atlantic bumper (*Chloroscombrus chrysurus*) is a game fish in the family, Carangidae. It was first described by the "father of modern taxonomy", Carl Linnaeus in the book, *Systema Naturae*. Atlantic bumper are known to eat smaller fish,...



Atlantic chub mackerel

Scomber colias

Atlantic chub mackerel (*Scomber colias*), also known as Tinker mackerel, is a pelagic schooling species of mackerel found in the Atlantic Ocean, the Mediterranean Sea, and the Black Sea. It was originally thought to be a subspecies of the chub...



Atlantic cod

Gadus morhua

The Atlantic cod (pl.: cod; Gadus morhua) is a fish of the family Gadidae, widely consumed by humans. It is also commercially known as cod or codling. Dry cod may be prepared as unsalted stockfish, and as cured salt cod or clipfish. In the western...



Atlantic croaker

Micropogonias undulatus

The Atlantic croaker (Micropogonias undulatus) is a species of marine ray-finned fish belonging to the family Sciaenidae and is closely related to the black drum (Pogonias cromis), the silver perch (Bairdiella chrysoura), the spot croaker...



Atlantic flyingfish

Cheilopogon melanurus

The Atlantic flyingfish (Cheilopogon melanurus) is a flying fish in the family Exocoetidae. It was first described by the French zoologist, Achille Valenciennes in a 22-volume work entitled Histoire naturelle des poissons (Natural History of...



Atlantic goliath grouper

Epinephelus itajara

The Atlantic goliath grouper or itajara (Epinephelus itajara), also known as the jewfish, is a saltwater fish of the grouper family and one of the largest species of bony fish. The species can be found in the West Atlantic ranging from...



Atlantic halibut

Hippoglossus hippoglossus

The Atlantic halibut (Hippoglossus hippoglossus) is a flatfish of the family Pleuronectidae. They are demersal fish living on or near sand, gravel or clay bottoms at depths of between 50 and 2,000 m (200 and 6,600 ft). The halibut is among the...



Atlantic herring

Clupea harengus

Atlantic herring (Clupea harengus) is a herring in the family Clupeidae. It is one of the most abundant fish species in the world. Atlantic herrings can be found on both sides of the Atlantic Ocean, congregating in large schools. They can grow up...



Atlantic horse mackerel

Trachurus trachurus

The Atlantic horse mackerel (Trachurus trachurus), also known as the European horse mackerel or common scad, is a species of jack mackerel in the family Carangidae, which includes the jacks, pompanos and trevallies. It is found in the eastern...



Atlantic lizardfish

Synodus saurus

The Atlantic lizardfish (Synodus saurus), is a species of lizardfish that lives primarily in the eastern Atlantic Ocean.



Atlantic mackerel

Scomber scombrus

The Atlantic mackerel (*Scomber scombrus*), also known as Boston mackerel, Norwegian mackerel, Scottish mackerel or just mackerel, is a species of mackerel found in the temperate waters of the Mediterranean Sea, the Black Sea, and the northern...



Atlantic mudskipper

Periophthalmus barbarus

The Atlantic mudskipper (*Periophthalmus barbarus*) is a species of mudskipper native to fresh, marine, and brackish waters of the tropical Atlantic coasts of Africa, including most offshore islands, through the Indian Ocean and into the western...



Atlantic pomfret

Brama brama

The Atlantic pomfret (*Brama brama*), also known as Ray's bream, is a species of marine ray-finned fish, a pomfret of the family Bramidae. It is found in the Atlantic, Indian, and South Pacific Oceans, at depths down to 1,000 m (3,300 ft). Its...



Atlantic sailfish

Istiophorus albicans

The Atlantic sailfish (*Istiophorus albicans*) is a species of marine fish in the family Istiophoridae of the order Istiophoriformes. It is found in the Atlantic Ocean and the Caribbean Sea, except for large areas of the central North Atlantic and...



Atlantic salmon

Salmo salar

The Atlantic salmon (*Salmo salar*) is a species of ray-finned fish in the family Salmonidae. It is the third largest of the Salmonidae, behind Siberian taimen and Pacific Chinook salmon, growing up to a meter in length. Atlantic salmon are found...



Atlantic saury

Scomberesox saurus

The Atlantic saury (Scomberesox saurus) is a fish of the family Scomberesocidae found in the North Atlantic Ocean from the Gulf of Saint Lawrence, Canada south to Bermuda and North Carolina in the western Atlantic and from Iceland to Morocco in...



Atlantic sharpnose shark

Rhizoprionodon terraenovae

The Atlantic sharpnose shark (Rhizoprionodon terraenovae) is a species of requiem shark in the family Carcharhinidae. It is found in the subtropical waters of the north-western Atlantic Ocean, between latitudes 43°N and 18°N.



Atlantic silverside

Menidia menidia

The Atlantic silverside (Menidia menidia), also known as spearing in the northeastern United States, is a small species of fish from the West Atlantic, ranging from the Gulf of St. Lawrence in Canada to northeastern Florida in the USA. It is one...



Atlantic sixgill shark

Hexanchus vitulus

The Atlantic sixgill shark (Hexanchus vitulus) is a rare species of hexanchid shark found in the Atlantic Ocean at depths that are greater than 300 meters. These depths are known as mesopelagic and bathypelagic in tropical and temperate waters...



Atlantic Spanish mackerel

Scomberomorus maculatus

The Atlantic Spanish mackerel (Scomberomorus maculatus) is a migratory species of mackerel that swims to the northern Gulf of Mexico in spring, returns to southern Florida in the eastern Gulf, and to Mexico in the western Gulf in the fall.



Atlantic stargazer

Uranoscopus scaber

The Atlantic stargazer (Uranoscopus scaber) is a marine, subtropical fish of family Uranoscopidae. Its body is suited for living on the sea floor, and is one of few fish capable of bioelectrogenesis, or the ability to generate an electric charge.



Atlantic stingray

Dasyatis sabinus

The Atlantic stingray (Hypanus sabinus) is a species of stingray in the family Dasyatidae, common along the Atlantic coast of North America from Chesapeake Bay to Mexico, including brackish and freshwater habitats. It may be distinguished from...



Atlantic sturgeon

Acipenser oxyrinchus oxyrinchus

The Atlantic sturgeon (Acipenser oxyrinchus oxyrinchus) is a member of the family Acipenseridae, and, along with other sturgeon, it is sometimes considered a living fossil. The Atlantic sturgeon is one of two subspecies of A. oxyrinchus, the...



Atlantic tarpon

Megalops atlanticus

The Atlantic tarpon (Megalops atlanticus) is a ray-finned fish that inhabits coastal waters, estuaries, lagoons, and rivers. It is also known as the silver king. It is found in the Atlantic Ocean, typically in tropical and subtropical regions,...



Atlantic thread herring

Opisthonema oglinum

The Atlantic thread herring (Opisthonema oglinum) is a herring-like fish in the family Clupeidae. It has a dark blue-gray back, silvery sides, a white belly, and a small head. It grows up to 38 cm in length. It can be found in shallow waters and...



Atlantic threadfin

Polydactylus octonemus

The Atlantic threadfin (*Polydactylus octonemus*) is a species of ray-finned fish a threadfin from the family Polynemidae which is native to subtropical and temperate waters of the western Atlantic Ocean and the Gulf of Mexico.



Atlantic torpedo

Tetronarce nobiliana

The Atlantic torpedo (*Tetronarce nobiliana*) is a species of electric ray in the family Torpedinidae. It is found in the Atlantic Ocean, from Nova Scotia to Brazil in the west and from Scotland to West Africa and off southern Africa in the east,...



Atlantic tripletail

Lobotes surinamensis

The Atlantic tripletail (*Lobotes surinamensis*), or tripletail, is a species of marine ray-finned fish belonging to the family Lobotidae. This fish is found in tropical and subtropical waters around the world except for the eastern Pacific Ocean.



Atlantic trumpetfish

Aulostomus strigosus

The Atlantic trumpetfish (*Aulostomus strigosus*), also referred to as Atlantic coronetfish, is a species of trumpetfish in the family Aulostomidae. It is a tropical marine fish found in shallow coastal waters in the eastern Atlantic Ocean from...



Atlantic weasel shark

Paragaleus pectoralis

The Atlantic weasel shark (Paragaleus pectoralis) is a weasel shark of the family Hemigaleidae, found in the eastern Atlantic Ocean. Also known as the little tiger shark, it is the only species of the genus Paragaleus to have been recorded off...



Atlantic whitefish

Coregonus huntsmani

The Atlantic whitefish (Coregonus huntsmani) is a coregonine fish inhabiting some freshwater lakes within Nova Scotia, Canada. It is known to survive only in the Petite Rivière watershed as landlocked populations. Earlier it was also found in the...



Atlantic wolffish

Anarhichas lupus

The Atlantic wolffish (Anarhichas lupus), also known as the seawolf, Atlantic catfish, ocean catfish, devil fish, wolf eel (the common name for its Pacific relative), woof or sea cat, is a marine fish of the wolffish family Anarhichadidae, native...



Atlantic wreckfish

Polyprion americanus

The Atlantic wreckfish (Polyprion americanus), also known as the stone bass or bass groper (among other names), is a marine, bathydemersal, and oceanodromous ray-finned fish in the family Polyprionidae. It has a worldwide, if disjunct,...



Atopodontus adriaensi

Atopodontus adriaensi

Atopodontus adriaensi is a species of upside-down catfish endemic to Gabon where it occurs in rapids of the Ivindo, Okano, Ngounié and the Nyanga Rivers. This species grows to a length of 10.0 centimetres (3.9 in) SL. This species is the only...



Aulonocara jacobfreibergi

Aulonocara jacobfreibergi

Aulonocara jacobfreibergi, the eureka red peacock, is a species of fish in the family Cichlidae. It is also known as Freiberg's peacock or fairy cichlid. There are several colour morphs including red or yellow. They generally can grow to around...



Aulonocara nyassae

Aulonocara nyassae

Aulonocara nyassae, known as the emperor cichlid, is a species of haplochromine Cichlid that is endemic to Lake Malawi in Africa. It has been recorded from the southeastern arm of the lake and may be present in the southwestern arm. This species...



Aulostomus maculatus

Aulostomus maculatus

Aulostomus maculatus, the West Atlantic trumpetfish, is a long-bodied fish with an upturned mouth. It often swims vertically while trying to blend with vertical coral, such as sea rods, sea pens, and pipe sponges.



Auriglobus modestus

Auriglobus modestus

Auriglobus modestus, the bronze puffer or golden puffer, is a ray-finned fish in the family Tetraodontidae. It is found in the middle parts of rivers to their estuaries in southeast Asia, including Cambodia, Laos, Thailand, Vietnam, Malaysia and...



Aurora trout

Salvelinus fontinalis timagamiensis

The aurora trout, *Salvelinus fontinalis timagamiensis*, is a variant or subspecies of the brook trout native to two lakes in the Temagami District of Ontario, Canada. The existence of the fish was brought to the attention of the angling world by...



Australasian flying fish

Cheilopogon pinnatibarbatus melanocercus

The Australasian flying fish, Cheilopogon pinnatibarbatus melanocercus, is a subspecies of flyingfish of the family Exocoetidae, found off New South Wales of Australia, and around New Zealand, in surface waters. The Australasian flying fish feeds...



Australasian snapper

Pagrus auratus

The Australasian snapper (Pagrus auratus) or silver seabream is a species of porgie found in coastal waters of Australia, Philippines, Indonesia, mainland China, Taiwan, Japan and New Zealand. Its distribution areas in the Northern and Southern...



Australian angelshark

Squatina australis

The Australian angelshark (Squatina australis) is a species of angelshark, family Squatinidae, found in the subtropical waters of southern Australia from Western Australia to New South Wales between latitudes 18°S and 41°S, at depths down to 255...



Australian barracuda

Sphyraena novaehollandiae

The Australian barracuda, arrow barracuda, Australian sea pike, sea pike, snook, or shortfin barracuda, Sphyraena novaehollandiae, is a barracuda of the genus Sphyraena which occurs in the south-western Pacific Ocean.



Australian bass

Macquaria novemaculeata

The Australian bass (Macquaria novemaculeata) is a small- to medium-sized species of primarily freshwater (but estuarine spawning) fish found in coastal rivers and streams along the east coast of Australia. A member of the genus Macquaria...



Australian blacktip shark

Carcharhinus tilstoni

The Australian blacktip shark (Carcharhinus tilstoni) is a species of requiem shark, in the family Carcharhinidae, endemic to northern and eastern Australia. Favoring the upper and middle parts of the water column, it can be found from the...



Australian bonito

Sarda australis

The Australian bonito, horse mackerel or little bonito, Sarda australis is a fish of the family Scombridae and is found in eastern Australia and New Zealand. They swim at depths reaching depths of approximately 30 m (98 ft), in open water. Its...



Australian clownfish

Amphiprion rubrocinctus

Amphiprion rubrocinctus, also known as the Australian clownfish or red anemonefish, is a species of anemonefish that is endemic to north west Australia. Like all anemonefishes it forms a symbiotic mutualism with sea anemones and is unaffected by...



Australian ghostshark

Callorhinchus milii

The Australian ghostshark (Callorhinchus milii) is a cartilaginous fish (Chondrichthyes) belonging to the subclass Holocephali (chimaera). Sharks, rays and skates are the other members of the cartilaginous fish group and are grouped under the...



Australian grayling

Prototroctes maraena

The Australian grayling (Prototroctes maraena) is a primarily freshwater fish found in coastal rivers in south-eastern mainland Australia and Tasmania. In past decades it has also been known as the cucumber mullet or cucumber herring, for its...



Australian herring

Arripis georgianus

The Australian herring (Arripis georgianus), also known as the ruff, tommy ruff, or Australian ruff, is one of four Australasian fish species within the genus Arripis. It closely resembles its sister species, the Australian salmon, although it...



Australian marbled catshark

Atelomycterus macleayi

Sharks portalThe Australian marbled catshark (Atelomycterus macleayi) is a catshark of the family Scyliorhinidae, found in the eastern Indian Ocean, endemic to Western Australia between latitudes 12 and 21°S, from the surface to 245 m (800 ft)...



Australian sharpnose shark

Rhizoprionodon taylori

Sharks portalThe Australian sharpnose shark (Rhizoprionodon taylori) is a requiem shark, belonging to the family Carcharhinidae. It is found in the tropical waters of the western Pacific Ocean off Papua New Guinea and northern Australia, between...



Australian smelt

Retropinna semoni

The Australian smelt (Retropinna semoni) is a small, pelagic silvery freshwater fish found in large numbers in waters of the south eastern Australian mainland.



Australian spotted mackerel

Scomberomorus munroi

The Australian spotted mackerel (Scomberomorus munroi) is a species of fish in the family Scombridae. Common fork length ranges between 50 and 80 cm. Specimens have been recorded at up to 104 cm in length, and weighing up to 10.2 kg. It is found...



Australian swellshark

Cephaloscyllium laticeps

The Australian swellshark (Cephaloscyllium laticeps) or draughtboard shark, is a species of catshark, and part of the family Scyliorhinidae, endemic to southern Australia. This bottom-dwelling species can be found on the continental shelf down to...



Australian weasel shark

Hemigaleus australiensis

The Australian weasel shark (Hemigaleus australiensis) is an uncommon species of ground shark in the family Hemigaleidae. It inhabits shallow waters off northern Australia to a depth of 170 m (560 ft); smaller sharks frequent sand and seagrass...



Australoheros facetus

Australoheros facetus

Australoheros facetus, formerly Cichlasoma facetum, the chameleon cichlid or chanchito, is a species of cichlid from the subfamily Cichlasomatinae which is native to northern Argentina, Paraguay, Uruguay and southern Brazil.



Awaous guamensis

Awaous guamensis

Awaous guamensis is a species of goby native to the Pacific islands from the Marianas to Vanuatu, New Caledonia and Fiji where it can be found in fresh, brackish and marine waters. Recent work based upon morphological and genetic differences has...



Awaous stamineus

Awaous stamineus

Awaous stamineus, commonly known as 'O'opu nakea, is a species of goby native to the Hawaiian Islands. It has been previously considered a synonym of Awaous guamensis but recent work based upon morphological and genetic differences has recognized...



Ayumodoki

Parabotia curtus

The ayumodoki or kissing loach (Parabotia curtus) is a species of ray-finned fish in the family Botiidae. It is found in lakes and streams on Honshu, the largest island in Japan. Spawning grounds for kissing loach are ditches and small reservoirs...



Badis badis

Badis badis

Badis badis, also known as the blue perch or blue badis, is a small species of Asian freshwater fish in the family Badidae of the order Anabantiformes. It is found in ponds, rivers, ditches and swamps in northern India, eastern Pakistan,...



Bagarius bagarius

Bagarius bagarius

Bagarius bagarius, also known as the giant devil catfish or goonch (Bengali: বাঘাইর), is a species of catfish in the genus Bagarius. It is generally reported as being found in large and medium rivers in South Asia, and is likely synonymous with...



Bagarius rutilus

Bagarius rutilus

Bagarius rutilus is a species of sisorid catfish native to Laos, Vietnam and Yunnan in China where it is found in the Nam Sam, Nam Ma and Red River basins. This species grows to a length of 100 centimetres (39 in) SL.



Bagarius suchus

Bagarius suchus

The crocodile catfish (Bagarius suchus) is a species of sisorid catfish native to Laos, Cambodia and Thailand, where it occurs in the Mekong and Chao Phraya river basins. This species reaches a maximum length of 70 centimetres (28 in) SL.



Bagarius yarrelli

Bagarius yarrelli

Bagarius yarrelli, also known as the giant devil catfish or goonch, is a very large species of catfish in the genus Bagarius found in rivers in the Indian subcontinent. The species reaches up to 2 m (6.6 ft) in length. It may be synonymous with...



Bahamas sawshark

Pristiophorus schroederi

Sharks portalThe Bahamas sawshark, Pristiophorus schroederi, is a sawshark of the family Pristiophoridae, found in the western Central Atlantic Ocean from the Bahamas and Cuba at depths of between 400 and 1,000 m. These sharks are at least 80 cm...



Bairdiella chrysoura

Bairdiella chrysoura

Bairdiella chrysoura, the American silver perch, silver croaker or goldtail croaker, is a species of marine ray-finned fish belonging to the family Sciaenidae, the drums and croakers. This species is widespread along the eastern coast of North...



Baja California rainbow trout

Oncorhynchus mykiss nelsoni

The Baja California rainbow trout or San Pedro Martir trout or Nelson's trout (*Oncorhynchus mykiss nelsoni*) is a localized subspecies of the rainbow trout (*Oncorhynchus mykiss*), a freshwater fish in the family Salmonidae.Baja California rainbow...



Bala shark

Balantiocheilos melanopterus

The bala shark (*Balantiocheilos melanopterus*) also known as the tricolor shark, tricolor sharkminnow, silver shark, or shark minnow, is a fish of the family Cyprinidae, and is one of the two species in the genus *Balantiocheilos*. This species is...



Balao halfbeak

Hemiramphus balao

The Balao halfbeak (*Hemiramphus balao*), occasionally called the Balao for short, is an ocean-going species of fish in the family Hemiramphidae. It was first described by the French naturalist Charles Alexandre Lesueur in 1821. They are used as...



Bald notothen

Pagothenia borchgrevinki

The bald notothen (*Pagothenia borchgrevinki*), also known as the bald rockcod, is a species of marine ray-finned fish belonging to the family Nototheniidae, the notothen or cod icefishes. It is native to the Southern Ocean.



Baldchin groper

Choerodon rubescens

The baldchin groper (Choerodon rubescens) is a species of wrasse endemic to Western Australia, where it occurs on coral and weed-grown rocky reefs. This species can reach a length of 90 cm (35 in).



Balkhash perch

Perca schrenkii

The Balkhash perch (Perca schrenkii) is a species of perch endemic to the Lake Balkhash and Lake Alakol watershed system, which lies mainly in Kazakhstan. It is similar to the other two species of perch, and grows to a comparable size, but has a...



Ballan wrasse

Labrus bergylta

The ballan wrasse (Labrus bergylta) is a species of marine ray finned fish from the family Labridae, the wrasses. It is found in the eastern Atlantic Ocean, where it inhabits rocky areas. Like many wrasse species, it is a protogynous...



Ballerus ballerus

Ballerus ballerus

Ballerus ballerus, also known as the zope or the blue bream, is a species of cyprinid fish native to Eurasia.



Balloon shark

Cephaloscyllium sufflans

The balloon shark (Cephaloscyllium sufflans) is a species of catshark, and part of the family Scyliorhinidae, endemic to the southwestern Indian Ocean off South Africa and Mozambique. Benthic in nature, it is found over sandy and muddy flats at...



Ballyhoo

Hemiramphus brasiliensis

The ballyhoo halfbeak or ballyhoo (Hemiramphus brasiliensis) is a baitfish of the halfbeak family (Hemiramphidae). It is similar to the Balao halfbeak (H. balao) in most features. Ballyhoo are frequently used as cut bait and for trolling purposes...



Balon's ruffe

Gymnocephalus baloni

Balon's ruffe (Gymnocephalus baloni), also known as the Danube ruffe, is a species of freshwater ray-finned fish, a ruffe, from the family Percidae which is native to the drainages of the Danube from delta to Germany and the Dnieper from delta to...



Banded angelfish

Apolemichthys arcuatus

The banded angelfish (Apolemichthys arcuatus), also known as the bandit angelfish and three spine angelfish, is a distinctive species ray-finned fish belonging to the family Pomacanthidae. It is endemic to deeper reefs in Hawaii and the Johnston...



Banded archerfish

Toxotes jaculatrix

The banded archerfish (Toxotes jaculatrix) is a brackish water perciform fish of the archerfish genus Toxotes. It is silvery in colour and has a dorsal fin towards the posterior end. It has distinctive, semi-triangular markings along its sides....



Banded butterflyfish

Chaetodon striatus

The banded butterflyfish (Chaetodon striatus) is a species of marine ray-finned fish, a butterflyfish found in the tropical western Atlantic Ocean from Brazil to Bermuda. Common names include the banded butterflyfish, the butterbun, the...



Banded cichlid

Heros severus

Heros severus (previously known as Cichlasoma severum), is a species of tropical freshwater cichlid native to the upper Orinoco and upper Rio Negro basins in South America. It has historically been confused with several other species in the...



Banded corydoras

Scleromystax barbatus

The banded corydoras or bearded catfish (Scleromystax barbatus) is a subtropical freshwater fish belonging to the subfamily Corydoradinae of the family Callichthyidae. It originates in coastal drainages in South America from Rio de Janeiro to...



Banded houndshark

Triakis scyllium

The banded houndshark (Triakis scyllium) is a species of houndshark in the family Triakidae, common in the northwestern Pacific Ocean from the southern Russian Far East to Taiwan. Found on or near the bottom, it favors shallow coastal habitats...



Banded killifish

Fundulus diaphanus

The banded killifish (Fundulus diaphanus) is a North American species of temperate freshwater killifish belonging to the genus Fundulus of the family Fundulidae. Its natural geographic range extends from Newfoundland to South Carolina, and west...



Banded knifefish

Gymnotus carapo

The banded knifefish (Gymnotus carapo) is a species of gymniform knifefish native to a wide range of freshwater habitats in South America. It is the most widespread species of Gymnotus, but it has frequently been confused with several relatives,...



Banded kōkopu

Galaxias fasciatus

The banded kōkopu (Galaxias fasciatus) is a galaxiid of the genus Galaxias, found only in New Zealand, including the Chatham and Stewart/Rakiura islands. It commonly grows to 20–25 cm, but has been recorded growing to around 30 cm. Juvenile...



Banded pipefish

Dunckerocampus dactyliophorus

The banded pipefish or ringed pipefish (Dunckerocampus dactyliophorus) is a species of fish in the Syngnathidae (seahorses and pipefish) family.



Banded pygmy sunfish

Elassoma zonatum

The banded pygmy sunfish, Elassoma zonatum, is a species of pygmy sunfish endemic to the United States, where it is found from Indiana and Illinois to Texas to the Atlantic coast. It prefers densely vegetated bodies of slow-moving water. This...



Banded sculpin

Cottus carolinae

The banded sculpin (Cottus carolinae) is a freshwater fish dwelling mostly in small to moderate sized streams in areas of swift current. Young and juvenile C. carolinae can mainly be found in pools, riffles, and other shallow habitats while...



Banded sunfish

Enneacanthus obesus

The banded sunfish (Enneacanthus obesus) is a freshwater fish of the family, Centrarchidae. They can grow to 2–3 inches long. They are native to North America.



Bandtail puffer

Sphoeroides spengleri

The bandtail puffer (Sphoeroides spengleri) is a species in the family Tetraodontidae, or pufferfishes. It can grow to a length of about 30 cm and is common in the Caribbean and observed from Massachusetts, USA in the north to Santa Catarina,...



Bandula barb

Pethia bandula

The bandula barb (Pethia bandula) is a species of cyprinid endemic to Sri Lanka where it is only known from near Galapitamada in the Warakapola Divisional Secretariat. As this critically endangered species only was known from a single unprotected...



Banggai cardinalfish

Pterapogon kauderni

The Banggai cardinalfish (Pterapogon kauderni) is a small tropical cardinalfish in the family Apogonidae. It is the only member of its genus. This attractive fish is popular in the aquarium trade. It is among the relatively few marine fish to...



Bantam sunfish

Lepomis symmetricus

The bantam sunfish (Lepomis symmetricus) is a species of freshwater fish in the genus Lepomis common throughout Louisiana, in extreme southeastern Texas, in southern Arkansas, and in a few places in western Kentucky and western Tennessee.



Bar jack

Caranx ruber

The bar jack (Caranx ruber), also known as the carbonero, red jack, blue-striped cavalla or passing jack, is a common species of inshore marine fish classified in the jack family, Carangidae. The bar jack is distributed through the western...



Barbel steed

Hemibarbus labeo

The barbel steed (*Hemibarbus labeo*) is a species of small freshwater fish in the family Cyprinidae. It is found throughout the Amur basin in eastern Asia to northern Vietnam, Japan and islands of Hainan and Taiwan.



Barbeled houndshark

Leptocharias smithii

The barbeled houndshark (*Leptocharias smithii*) is a species of ground shark and the only member of the family Leptochariidae. This demersal species is found in the coastal waters of the eastern Atlantic Ocean from Mauritania to Angola, at depths...



Barbelthroat carpetshark

Cirrhoscyllium expolitus

Sharks portalThe barbelthroat carpetshark (Cirrhoscyllium expolitus) is a carpetshark of the family Parascylliidae found in the South China Sea between Luzon in the Philippines and China, between latitudes 23°N and 10°N, at depths between 180...



Barbodes sirang

Barbodes sirang

Barbodes sirang, known locally as the sirang, is a possibly extinct species of cyprinid fish endemic to Lake Lanao in Mindanao, the Philippines. This species can reach a length of 8.6 centimetres (3.4 in) TL.



Barbodes tumba

Barbodes tumba

Barbodes tumba is a species of cyprinid fish endemic to Mindanao, Philippines where it is found in Lake Lanao and associated water systems. This species can reach a length of 12.9 centimetres (5.1 in) SL.



Barbodes wynaadensis

Barbodes wynaadensis

Barbodes wynaadensis is a species of cyprinid fish. It is endemic to the Wyanad Plateau and its surroundings in the southern Western Ghats, India. It occurs in fast-flowing rivers and streams with rock substrates. This species can reach a length...



Barbour's seahorse

Hippocampus barbouri

Barbour's seahorse (Hippocampus barbouri) is a species of fish of the family Syngnathidae.



Barca snakehead

Channa barca

The Barca snakehead (Channa barca) is a rare species of snakehead. It is endemic to the upper Brahmaputra river basin in northeastern India and Bangladesh. Records from Nepal are of doubtful validity. Overall it has been assessed as data...



Barfin flounder

Verasper moseri

The barfin flounder (*Verasper moseri*) is a flatfish of the family Pleuronectidae. It is a demersal fish that lives on sandy, muddy bottoms at depths of up to 900 metres (3,000 ft). It can reach up to 70 centimetres (28 in) in length and can weigh...



Barilius barila

Barilius barila

Barilius barila (Bengali: বারালি) is a tropical fish in genus Barilius of the family Cyprinidae. It is commercially very important due to its superior taste. It is found in the Asian countries of India, Nepal, Bangladesh and Myanmar. Normally...



Barilius bendelisis

Barilius bendelisis

Barilius bendelisis is a fish in genus Barilius of the family Cyprinidae. It is found in Pakistan, India, Nepal, Bangladesh, Sri Lanka, Bhutan and Myanmar.



Barndoor skate

Dipturus laevis

The barndoor skate (Dipturus laevis) is a species of marine cartilaginous fish in the skate family Rajidae of the order Rajiformes. It is native to the northwestern Atlantic Ocean, and is found from the Grand Banks of Newfoundland and the...



Barramundi

Lates calcarifer

The barramundi (Lates calcarifer), Asian sea bass, or giant sea perch, or dangri is a species of catadromous fish in the family Latidae of the order Perciformes. The species is widely distributed in the Indo-West Pacific, spanning the waters of...



Barred bichir

Polypterus delhezi

The barred bichir, armoured bichir, bandback bichir, or banded bichir (Polypterus delhezi) is an elongated fish found in the Congo River, specifically in the upper and middle portions. This species is one of the more commonly available in...



Barred hogfish

Bodianus scrofa

The barred hogfish (Bodianus scrofa) is a species of wrasse native to the eastern Atlantic Ocean, where it occurs around the Macaronesian island groups of the Azores, Madeira, the Canary Islands and Cape Verde. This species occurs on rocky reefs...



Barred moray

Echidna polyzona

The barred moray (Echidna polyzona), also known as the banded moray, the dark-banded eel, the girdled moray, the girdled reef eel, the many banded moray eel, the ringed moray, the ringed reef moray, the striped moray and the zebra eel,) is a...



Barred mudskipper

Periophthalmus argentilineatus

The barred mudskipper (Periophthalmus argentilineatus) or silverlined mudskipper, is a species of mudskippers native to marine, fresh and brackish waters from the African coast of the Indian Ocean, to the Marianas and Samoa in the western Pacific...



Barred sand bass

Paralabrax nebulifer

The barred sand bass (Paralabrax nebulifer) is a species of fish in the family Serranidae, the sea basses and groupers. It is native to California and Baja California, where it lives in the coastal waters of the eastern Pacific Ocean.



Barred surfperch

Amphistichus argenteus

The barred surfperch (Amphistichus argenteus) is a species of surfperch native to inshore waters from northern California, United States to southern Baja California, Mexico. This species can reach a length of 43 centimetres (17 in) TL though most...



Barred thicklip

Hemigymnus fasciatus

The barred thicklip wrasse (Hemigymnus fasciatus) is a species of fish belonging to the wrasse family, native from the Indo-Pacific.



Bartail flathead

Platycephalus indicus

The bartail flathead (Platycephalus indicus), also known as the Indian flathead, gobi or Indo-Pacific flathead, is a species of largely marine ray-finned fish belonging to the family Platycephalidae, the flatheads. This species is found in the...



Baryancistrus xanthellus

Baryancistrus xanthellus

Baryancistrus xanthellus is a species of armored catfish endemic to Brazil. It is a benthic fresh water fish that lives in the Volta Grande and Xingu River, as well as the area immediately above Belo Monte falls and the Iriri River. It reaches a...



Basa

Pangasius bocourti

Basa (Pangasius bocourti) is a species of catfish in the family Pangasiidae. Basa are native to the Mekong and Chao Phraya basins in Mainland Southeast Asia. These fish are important as a food source, and also on the international market. They...



Basking shark

Cetorhinus maximus

The basking shark (Cetorhinus maximus) is the second-largest living shark and fish, after the whale shark. It is one of three plankton-eating shark species, along with the whale shark and megamouth shark. Typically, basking sharks reach 7.9 m (26...



Bastard sturgeon

Acipenser nudiiventris

The bastard sturgeon, also known as the fringebarbel sturgeon, ship sturgeon, spiny sturgeon, or thorn sturgeon (Acipenser nudiiventris), is a species of fish in the family Acipenseridae. These fish are typically found along the benthos of...



Bat ray

Myliobatis californica

The bat ray (Myliobatis californica) is an eagle ray found in muddy or sandy sloughs, estuaries and bays, kelp beds and rocky-bottomed shoreline in the eastern Pacific Ocean, between the Oregon coast and the Gulf of California. It is also found...



Bathypterois grallator

Bathypterois grallator

The tripod fish or tripod spiderfish, Bathypterois grallator, is a deep-sea benthic fish in the family Ipnopidae found at lower latitudes. It is now relatively well known from photographs and submersible observations, and seems to prefer to perch...



Bathysaurus mollis

Bathysaurus mollis

Bathysaurus mollis is a species of fish in the family Bathysauridae, commonly called bathysaur or highfin lizardfish.



Bay pipefish

Syngnathus leptorhynchus

The bay pipefish (Syngnathus californiensis) is a pipefish native to the eelgrass beds of the Eastern Pacific (Southern Baja California to Gulf of Alaska), where its sinuous shape and green color allow it to blend in with the waving blades of...



Bayad

Bagrus bajad

The bayad (Bagrus bajad), is a species of bagrid catfish from Africa.



Bear Lake sculpin

Cottus extensus

The Bear Lake sculpin (Cottus extensus) occasionally referred to incorrectly as a "bullhead", is a species of freshwater sculpin endemic to Bear Lake on the Utah-Idaho border. It is one of only four sculpins native to Utah, and the only extant...



Beardless barb

Cyclocheilichthys apogon

The beardless barb (Cyclocheilichthys apogon) is a species of freshwater fish in the family Cyprinidae. It is widespread in Southeast Asia. It grows to 25 cm (9.8 in) total length.



Beaufortia kweichowensis

Beaufortia kweichowensis

Beaufortia kweichowensis is a species of gastromyzontid loach native to rivers in China. The common names for this popular aquarium species are Chinese hillstream loach, Hong Kong pleco, butterfly hillstream loach, and Chinese sucker fish.



Bedotia madagascariensis

Bedotia madagascariensis

Bedotia madagascariensis (zona) is a species of fish in the family Bedotiidae. It is endemic to Madagascar, where found in rivers and lakes between the Ivoloina River and the Manambolo Creek. It is commonly seen in the aquarium trade, where it...



Beige catshark

Parmaturus bigus

Sharks portalThe beige catshark (*Parmaturus bigus*) is a cat shark of the family Scyliorhinidae. The first recorded specimen was a female recorded off the coast of Queensland, Australia around Lord Howe Island. Its length was 72 cm.Recently, a...



Belontia hasselti

Belontia hasselti

The Malay combtail (*Belontia hasselti*) is a species of gourami native to southeast Asia where it is found in peat swamps of the Malay Peninsula and the Greater Sunda Islands. This species can reach a length of 20 centimetres (7.9 in) SL. It is a...



Belontia signata

Belontia signata

Belontia signata, the Ceylonese combtail, is a species of gourami endemic to Sri Lanka. This species inhabits shallow, slow-flowing clear-water streams. It can reach a length of 18 centimetres (7.1 in) TL though most are only around 10...



Belted sandfish

Serranus subligarius

The belted sandfish (*Serranus subligarius*), also known as the dwarf sea bass or stubby sea bass, is a species of marine ray-finned fish, a sea bass from the subfamily Serraninae, classified as part of the family Serranidae which includes the...



Beluga

Huso huso

The beluga, also known as the beluga sturgeon or great sturgeon (Huso huso), is a species of anadromous fish in the sturgeon family (Acipenseridae) of order Acipenseriformes. It is found primarily in the Caspian and Black Sea basins, and formerly...



Bengal danio

Devario devario

The Bengal danio or Sind danio (Devario devario) is a subtropical fish belonging to the minnow family (Cyprinidae). Originating in Pakistan, India, Nepal, Bangladesh, and Afghanistan, this fish is sometimes kept in community tanks by fish-keeping...



Bennett's stingray

Hemitrygon bennetti

The Bennett's stingray or frilltailed stingray (Hemitrygon bennetti, often misspelled benetti or bennettii) is a little-known species of stingray in the family Dasyatidae, with a wide but ill-defined distribution in the Indian and Pacific Oceans....



Bentfin devil ray

Mobula thurstoni

The bentfin devil ray (Mobula thurstoni), also known as the lesser devil ray, smoothtail devil ray, smoothtail mobula or Thurston's devil ray, is a species of ray in the family Mobulidae. It is found worldwide in tropical, subtropical and warm...



Benthoosema glaciale

Benthoosema glaciale

Benthoosema glaciale, or glacier lantern fish, is the most common species of lanternfish and important part of the midwater ecosystem of northern North Atlantic. It feeds on small crustaceans, including copepods, krill and amphipods, along with...



Bering cisco

Coregonus laurettae

The Bering cisco or Lauretta whitefish (Coregonus laurettae) is a freshwater whitefish found in Alaska and part of Russia. It is often considered to be the same species as the more common Arctic cisco (Coregonus autumnalis).



Beringraja rhina

Beringraja rhina

Beringraja rhina, commonly known as the longnose skate, is a species of skate in the family Rajidae from the northeast Pacific. It is found from the eastern Bering Sea and Alaska to Baja California and the Gulf of California. The longnose skate...



Betta albimarginata

Betta albimarginata

Betta albimarginata is a species of betta fish, endemic to the island of Borneo where it is only found in the Indonesian province of Kalimantan Timur. It inhabits the shallows (5 to 10 cm (2.0 to 3.9 in)) of forest streams amongst vegetation and...



Betta bellica

Betta bellica

Betta bellica, the slim betta or slender betta, is a species of betta native to eastern Sumatra in Indonesia and Peninsular Malaysia. It is an inhabitant of swamp forests and peat swamps. This species grows to a length of 10 cm (3.9 in). It can...



Betta burdigala

Betta burdigala

Betta burdigala is a species of gourami endemic to Bangka Island in Indonesia. It is an inhabitant of peat swamps. This species grows to a length of 2.5 cm (0.98 in).



Betta channoides

Betta channoides

Betta channoides is a species of betta endemic to the island of Borneo in Indonesia, where it is only found in the province of Kalimantan Timur. It is an inhabitant of the shallows of acidic brown-water forest streams. This species grows to a...



Betta coccina

Betta coccina

Betta coccina is a species of betta native to Peninsular Malaysia and Sumatra, Indonesia. It grows to a length of 5.6 cm (2.2 in). It can be found in the aquarium trade.



Betta dimidiata

Betta dimidiata

Betta dimidiata is a species of gourami endemic to Kalimantan Barat in Indonesia. It is an inhabitant of swamp forest and forest streams, and can grow to a length of 4.6 cm (1.8 in).



Betta enisae

Betta enisae

Betta enisae is a species of gourami endemic to the Kapuas River basin of Indonesia. It is an inhabitant of forest streams with slightly acidic waters, and can mostly be found in the shallows amongst the leaf litter. This species grows to a...



Betta gladiator

Betta gladiator

Betta gladiator is a species of gourami endemic to Sabah, Malaysia. It is an inhabitant of the acidic waters of heath forest. It is known as a particularly territorial and aggressive species, especially in captivity. This species grows to a...



Betta livida

Betta livida

Betta livida is a species of gourami endemic to Malaysia, where it is only found in the state of Selangor. An inhabitant of peat swamps, it grows to a length of 3.6 cm (1.4 in).



Betta miniopinna

Betta miniopinna

Betta miniopinna is a species of gourami endemic to Bintan Island in the Riau Archipelago of Indonesia. The species reaches 2.4 cm (0.9 inches) in standard length and is believed to be a facultative air-breather.



Betta pallifina

Betta pallifina

Betta pallifina is a species of gourami endemic to the island of Borneo.



Betta persephone

Betta persephone

Betta persephone is a species of labyrinth fish endemic to Malaysia where it is only known from Johor. This betta's specific epithet derives from the Greek goddess Persephone, queen of the Underworld, a reference to the largely blackish...



Betta raja

Betta raja

Betta raja is a species of betta endemic to the island of Sumatra in Indonesia. It is an inhabitant of swamps in lowland forests. This species grows to a length of 6.4 centimetres (2.5 in) SL. It is used as bait by local fishermen and has also...



Betta simorum

Betta simorum

Betta simorum is a species of gourami endemic to Indonesia. This species grows to a length of 6.4 centimetres (2.5 in) SL. This species can also be found in the aquarium trade. The species was named simorum and uses the Latin genitive plural "of...



Betta smaragdina

Betta smaragdina

Betta smaragdina, commonly known as the emerald green betta, blue betta or Mekong fighting fish (Thai: ปลากัดเขียว or ปลากัดอีสาน) is a species of gourami native to Southeast Asia. The species gets its green and blue colors due to refraction and...



Betta spilotogena

Betta spilotogena

Betta spilotogena is a species of gourami endemic to Indonesia where it is only known from the islands of Bintan and Singkep. It inhabits swamps and nearby streams preferring areas with plentiful vegetation and 0.5 metres (1.6 ft) or less in...



Betta unimaculata

Betta unimaculata

Betta unimaculata, the Howong betta, is a species of labyrinth fish endemic to the island of Borneo where it is only known from the eastern portion of the island. This species grows to a length of 12.6 centimetres (5.0 in) TL, one of the largest...



Bicolor angelfish

Centropyge bicolor

The bicolor angelfish (Centropyge bicolor) is a marine species of fish, easily recognizable by its yellow tail, yellow front half of their body, and blue rear with blue patterns above and around the eye. Other names of this angelfish include:...



Bicolored foxface

Siganus uspi

The bicolored foxface (Siganus uspi), also known as the Uspi rabbitfish, is a species of marine ray-finned fish, a rabbitfish belonging to the family Siganidae. It is found at coral reefs in Fiji in the Pacific Ocean. It occasionally makes its...



Bidenichthys beeblebroxi

Bidenichthys beeblebroxi

Bidenichthys beeblebroxi is a species of common reef fish of the family Bythitidae, and one of three species in the genus Bidenichthys. The species is found in the coastal waters off North Island and northern South Island, New Zealand. It is a...



Bidyanus bidyanus

Bidyanus bidyanus

The silver perch (Bidyanus bidyanus) is a medium-sized freshwater fish of the family Terapontidae endemic to the Murray-Darling river system in south-eastern Australia.



Big skate

Beringraja binocularata

The big skate (Beringraja binocularata) is the largest species of skate (family Rajidae) in the waters off North America. They are found along the Pacific Coast from Alaska to Baja California, typically from the intertidal zone to a depth of 120 m...



Big-belly seahorse

Hippocampus abdominalis

The big-belly seahorse (Hippocampus abdominalis) or pot-bellied seahorse is one of the largest seahorse species in the world, with a length of up to 35 cm (14 in), and is the largest in Australia. Seahorses are members of the family Syngnathidae,...



Big-scale sand smelt

Atherina boyeri

The big-scale sand smelt (Atherina boyeri) is a species of fish in the family Atherinidae. It is a euryhaline amphidromous fish, up to 20 cm in length.



Big-scaled redfin

Tribolodon hakonensis

The big-scaled redfin (Tribolodon hakonensis), also known as the Japanese dace and ugui, is a medium-sized Asian fish. First described by Albert Günther in 1877 as Leuciscus hakonensis, it is the type specimen of the genus Tribolodon, having been...



Bigeye chub

Hybopsis amblops

Bigeye chub (Hybopsis amblops) is a species of freshwater fish in the carp family (Cyprinidae).



Bigeye houndshark

Iago omanensis

The bigeye houndshark (Iago omanensis) is a species of houndshark, belonging to the family Triakidae. It is found in the deep waters of the continental shelves in the western Indian Ocean, from the Red Sea to southwestern India, between latitudes...



Bigeye sand tiger

Odontaspis noronhai

The bigeye sand tiger (Odontaspis noronhai) is an extremely rare species of mackerel shark in the family Odontaspidae, with a possible worldwide distribution. A large, bulky species reaching at least 3.6 m (12 ft) in length, the bigeye sand...



Bigeye scad

Selar crumenophthalmus

The bigeye scad (Selar crumenophthalmus) is a species of oceanic fish found in tropical regions around the globe. Other common names include purse-eyed scad, goggle-eyed scad, akule, chicharro, charrito ojón, jacks, matang baka, mushimas and...



Bigeye snapper

Lutjanus lutjanus

The bigeye snapper (Lutjanus lutjanus), also known as the bigeye seaperch, red sea lined snapper, golden striped snapper, rosy snapper, yellow snapper, or simply snapper, is a species of marine ray-finned fish, a snapper belonging to the family...



Bigeye thresher

Alopias superciliosus

The bigeye thresher (Alopias superciliosus) is a species of thresher shark, family Alopidae, found in temperate and tropical oceans worldwide. Like the other thresher sharks, nearly half its total length consists of the elongated upper lobe of...



Bigeye trevally

Caranx sexfasciatus

The bigeye trevally (*Caranx sexfasciatus*), also known as the bigeye jack, great trevally, six-banded trevally and dusky jack, is a species of widespread large marine fish classified in the jack family Carangidae. The bigeye trevally is...



Bigeye tuna

Thunnus obesus

The bigeye tuna (Thunnus obesus) is a species of true tuna of the genus Thunnus, belonging to the wider mackerel family Scombridae. In Hawaiian, it is one of two species known as 'ahi, the other being the yellowfin tuna. Bigeye tuna are found in...



Bigeyed sixgill shark

Hexanchus nakamurai

The bigeyed sixgill shark (Hexanchus nakamurai) is a cow shark of the family Hexanchidae. Its dorsal surface has a brownish-gray color, and is sharply separated from the light coloring of its ventral surface. The eyes are a fluorescent green...



Bighead carp

Hypophthalmichthys nobilis

The bighead carp (Hypophthalmichthys nobilis) is a species of cyprinid freshwater fish native to East Asia, and is one of several Asian carps introduced into North America. It is one of the most intensively exploited fishes in fish farming, with...



Bigmouth buffalo

Ictiobus cyprinellus

The bigmouth buffalo (Ictiobus cyprinellus) is a fish native to North America that is in decline. It is the largest North American species in the Catostomidae or "sucker" family, and is one of the longest-lived and latest-maturing freshwater...



Bignose shark

Carcharhinus altimus

The bignose shark (Carcharhinus altimus) is a species of requiem shark, in the family Carcharhinidae. Distributed worldwide in tropical and subtropical waters, this migratory shark frequents deep waters around the edges of the continental shelf....



Bigtooth river stingray

Potamotrygon henlei

The bigtooth river stingray or Tocantins River ray (Potamotrygon henlei) is a species of freshwater fish in the family Potamotrygonidae. It is endemic to the lower Tocantins basin and Araguaia basin in Brazil, and prefers muddy bottoms. It is...



Binni

Mesopotamichthys sharpeyi

The binni (*Mesopotamichthys sharpeyi*) is a species of cyprinid fish endemic to the Tigris–Euphrates Basin in the Middle East. This fish mostly inhabits lakes and marshes, especially in densely vegetated places where it also lays its eggs, but...



Biotodoma cupido

Biotodoma cupido

Biotodoma cupido, commonly known as the green-streaked eartheater or cupid cichlid, is a species of cichlid native to the Amazon and Essequibo basins in Bolivia, Brazil, Colombia, Guayana, and Peru. It is sometimes seen in the aquarium trade. It...



Birdbeak dogfish

Deania calcea

Sharks portalThe birdbeak dogfish (*Deania calcea*) is a dogfish shark of the family Centrophoridae found in the Pacific Ocean around Honshū, Japan, southern Australia, New Zealand, and Chile, and in the Atlantic Ocean from Iceland south to the...



Biwa trout

Oncorhynchus rhodurus

The Biwa trout (*Oncorhynchus rhodurus*) is an anadromous salmonid fish of the genus *Oncorhynchus*, endemic to Lake Biwa in Shiga Prefecture, Japan, but also introduced to Lake Ashi and Lake Chūzenji. While called trout, the fish is most closely...



Black and white snapper

Macolor niger

The black and white snapper (*Macolor niger*), the black and white seaperch or black snapper, is a species of marine ray-finned fish, a snapper belonging to the family Lutjanidae. It is native to the Indian Ocean and the western Pacific Ocean.



Black arowana

Osteoglossum ferreirai

The black arowana (*Osteoglossum ferreirai*) is a South American freshwater bony fish of the family Osteoglossidae. Black arowanas are sometimes kept in aquariums, but they are predatory and require a very large tank. It is generally common, but...



Black buffalo

Ictiobus niger

The black buffalo (*Ictiobus niger*) is a North American species of freshwater fish in the Catostomidae or sucker family. With a lifespan greater than 100 years it is among the longest-lived of freshwater fish species. Found in the Mississippi...



Black bullhead

Ameiurus melas

The black bullhead or black bullhead catfish (*Ameiurus melas*) is a species of bullhead catfish. Like other bullhead catfish, it has the ability to thrive in waters that are low in oxygen, brackish, turbid and/or very warm. It also has barbels...



Black carp

Mylopharyngodon piceus

The black carp (Mylopharyngodon piceus) or Chinese black roach is a species of cyprinid freshwater fish and the sole extant species of the genus Mylopharyngodon. It is native to lakes and rivers in East Asia, ranging from the Amur Basin across...



Black crappie

Pomoxis nigromaculatus

The black crappie (Pomoxis nigromaculatus) is a freshwater fish found in North America, one of the two types of crappies. It is very similar to the white crappie (P. annularis) in size, shape, and habits, except that it is darker, with a pattern...



Black dogfish

Centroscyllium fabricii

The black dogfish (Centroscyllium fabricii) is a species of dogfish shark in the family Etmopteridae. It is common over the outer continental shelf and continental slope at depths of 180–2,250 m (590–7,380 ft). Females generally inhabit deeper...



Black drum

Pogonias cromis

The black drum (Pogonias cromis), also known as the drum or drummer, is a species of marine ray-finned fish belonging to the family Sciaenidae, the drums and croakers. It is found in the western North Atlantic Ocean off the eastern coast of North...



Black ghost knifefish

Apteronotus albifrons

The black ghost knifefish (Apteronotus albifrons) is a tropical fish belonging to the ghost knifefish family (Apteronotidae). They originate in freshwater habitats in South America where they range from Venezuela to the Paraguay–Paraná River....



Black goby

Gobius niger

The black goby (Gobius niger) is a species of ray-finned fish found in the Eastern Atlantic and Mediterranean Sea and Black Sea. It inhabits estuaries, lagoons, and inshore water over seagrass and algae. It feeds on a variety of invertebrates and...



Black lancer

Bagrichthys macracanthus

The black lancer (Bagrichthys macracanthus) is a species of bagrid catfish found in Vietnam, Cambodia, Indonesia, Laos, Malaysia and Thailand. It grows to a length of 25.0 cm. It is the only species of the genus Bagrichthys that is traded for the...



Black margate

Anisotremus surinamensis

The black margate (Anisotremus surinamensis), also known as black bream, black thicklip, dogfish, lippe, Mexican bull, pompon, Spanish grunt, surf bream, sweetlips or thicklip grunt, is a species of marine ray-finned fish, a grunt belonging to...



Black marlin

Istiompax indica

The black marlin (Istiompax indica) is a species of marlin found in tropical and subtropical areas of the Indian and Pacific Oceans. Reaching lengths of over 4.5 m (15 ft), it is one of the largest marlins and also one of the largest bony fish....



Black musselcracker

Cymatoceps nasutus

The black musselcracker (Cymatoceps nasutus) is a species of marine ray-finned fish belonging to the family Sparidae, which includes the seabreams and porgies. This species is the only species in the monospecific genus Cymatoceps. It is endemic...



Black neon tetra

Hyphessobrycon herbertaxelrodi

The black neon tetra (Hyphessobrycon herbertaxelrodi) is a freshwater fish of the characin family (Characidae) of the order Characiformes. It is native to the Paraguay basin of southern Brazil. They are often found in the aquarium trade.



Black phantom tetra

Hyphessobrycon megalopterus

The black phantom tetra (Hyphessobrycon megalopterus), or simply phantom tetra, is a small freshwater fish of the characin family (Characidae) of order Characiformes. It is native to the upper Paraguay basin and upper Madeira basin (including...



Black redhorse

Moxostoma duquesnei

The black redhorse (Moxostoma duquesnei, also duquesnii, duquesnei) is a species of freshwater fish endemic to Ontario and the eastern half of the United States, where it lives in streams and small to medium rivers. A bottom-feeder, it feeds on...



Black rockfish

Sebastes melanops

The black rockfish (Sebastes melanops), also known variously as the black seaperch, black bass, black rock cod, sea bass, black snapper and Pacific Ocean perch, is a species of marine ray-finned fish belonging to the subfamily Sebastinae, the...



Black ruby barb

Pethia nigrofasciata

The black ruby barb (Pethia nigrofasciata, formerly Puntius nigrofasciatus) or purplehead barb is a tropical cyprinid fish endemic to Sri Lanka, where it occurs in forested streams from the Kelani basin to the Nilwala basin. They are found in...



Black scabbardfish

Aphanopus carbo

The black scabbardfish (Aphanopus carbo) is a bathypelagic cutlassfish of the family Trichiuridae found in the Atlantic Ocean between latitudes 69°N and 27°N at depths between 180 and 1,700 m (600 and 5,600 ft). Its length is up to 110 cm (3.6...



Black scorpionfish

Scorpaena porcus

The black scorpionfish (*Scorpaena porcus*), also known as the European scorpionfish or small-scaled scorpionfish, is a venomous scorpionfish, common in marine subtropical waters. It is widespread in the Eastern Atlantic Ocean from the British...



Black sea bass

Centropristis striata

The black sea bass (*Centropristis striata*) is a species of marine ray-finned fish, a sea bass from the subfamily Serraninae which is part of the family Serranidae, which also includes the groupers and anthias. It is found in the western Atlantic...



Black Sea salmon

Salmo labrax

The Black Sea salmon (Salmo labrax) is a fairly small species of salmon, at about 20 inches (510 mm) long on average and rarely reaching over 30 in (760 mm). It inhabits the northern Black Sea coasts and inflowing rivers. There are anadromous....



Black Sea sprat

Clupeonella cultriventris

The Black Sea sprat, Clupeonella cultriventris, is a small fish of the herring family, Clupeidae. It is found in the Black Sea and Sea of Azov and rivers of its basins: Danube, Dnister, Dnipro (Ukraine), Southern Bug, Don, Kuban. It has...



Black seabream

SpondylIOSoma cantharus

The black seabream (SpondylIOSoma cantharus) is a protogynous species of Sparidae fishes, recognisable by their oval, compressed body and jaws, which contain 4–6 rows of slender teeth that are larger at the front. They are silvery in colour with...



Black sharkminnow

Labeo chrysophekadion

The black sharkminnow (Labeo chrysophekadion), also known as the black shark or black labeo, is a species of freshwater fish in the carp family. It is found in the Mekong and Chao Phraya river basins, Malay Peninsula, Sumatra, Java and Borneo. It...



Black snakehead

Channa melasoma

The black snakehead (Channa melasoma) is a species of snakehead native to Thailand, Indonesia, Malaysia, Philippines, and Singapore. This commercially important species reaches a length of 30 cm (12 in). The Black Snakehead inhabits large to...



Black swallower

Chiasmodon niger

The black swallower, Chiasmodon niger, is a species of deep sea fish in the family Chiasmodontidae. It is known for its ability to swallow fish larger than itself. It has a worldwide distribution in tropical and subtropical waters, in the...



Black tetra

Gymnocorymbus ternetzi

The black tetra (Gymnocorymbus ternetzi), also known as the black skirt tetra, petticoat tetra, high-fin black skirt tetra, black widow tetra and blackamoor, is a freshwater fish of the characin family (Characidae). It is native to the Paraguay...



Black triggerfish

Melichthys niger

The black triggerfish or black durgon (Melichthys niger), called Humuhumu'ele'ele in Hawaiian, is a blimp-shaped triggerfish with bright white lines running along its dorsal and anal fins. From distance, it appears to be completely black....



Black-banded trevally

Seriolina nigrofasciata

The black-banded trevally (*Seriolina nigrofasciata*) is a species of carangid native to the Indian Ocean, the western Pacific Ocean, and the Atlantic coast of southeastern South Africa. This species inhabits reefs and rocky bottoms at depths from...



Black-barred danio

Danio absconditus

The black-barred danio (Danio absconditus) is a species of Danio discovered in Myanmar by Tin Win in 2005 and described in 2015 by Sven Oscar Kullander and Ralf Britz.



Black-faced blenny

Tripterygion delaisi

The black-faced blenny (Tripterygion delaisi) is a small benthic fish from the family Tripterygiidae (triplefin-blennies). It occurs at depths of 3 to 40 metres (9.8 to 131.2 ft) and lives on the substrate under large rocks, cliffs or other...



Black-ray goby

Stonogobiops nematodes

Stonogobiops nematodes, the Filament-finned prawn-goby, the Antenna goby, the high-fin goby, the red-banded goby, the high-fin red-banded goby, the striped goby, the barber-pole goby, or the black-ray Goby, is a species of marine goby native to...



Black-striped pipefish

Syngnathus abaster

The black-striped pipefish (Syngnathus abaster) is a species of fish in the family Syngnathidae. It is found in the eastern Atlantic from the southern Gulf of Biscay to Gibraltar, also in the Mediterranean and Black Seas. As the introduced...



Blackback butterflyfish

Chaetodon melannotus

The black-backed butterflyfish or blackback butterflyfish (*Chaetodon melannotus*) is a species of butterflyfish (family Chaetodontidae). It is widespread through the Indo-Pacific area from the Red Sea and East Africa to Samoa, to southern Japan...



Blackbanded sunfish

Enneacanthus chaetodon

The blackbanded sunfish (*Enneacanthus chaetodon*) is a freshwater fish species of the sunfish family (Centrarchidae). They are found in the United States ranging from New Jersey to central Florida.> The defining feature of this Black-banded...



Blackbelly lanternshark

Etmopterus lucifer

Sharks portalThe blackbelly lanternshark or lucifer shark, *Etmopterus lucifer*, is a shark of the family Etmopteridae, found around the world in tropical to temperate seas, at depths between 150 and 1,250 meters. Its length is up to 47...



Blackbelly triggerfish

Rhinecanthus verrucosus

The blackbelly triggerfish (*Rhinecanthus verrucosus*) is a ray-finned fish in the family Balistidae found in the Indo-Pacific. It occasionally makes its way into the aquarium trade. It is sometimes known as the blackpatch triggerfish.



Blackcap basslet

Gramma melacara

The blackcap basslet (*Gramma melacara*), or blackcap gramma, is a species of fish inhabiting reefs in the tropical western Atlantic Ocean. It prefers vertical surfaces with crevices in which it can hide. It can be found at depths of from 10 to 180...



Blackcheek moray eel

Gymnothorax breedeni

The blackcheek moray eel or masked moray (Gymnothorax breedeni) is a species of marine fish in the family Muraenidae.



Blackchin tilapia

Sarotherodon melanotheron

The blackchin tilapia (Sarotherodon melanotheron) is a species of cichlid native to coastal west Africa. It is a paternal mouthbrooder which has been introduced to Asia and North America. In the Philippines, it is also informally called gloria or...



Blackfin barracuda

Sphyraena qenie

The blackfin barracuda (Sphyraena qenie), also known as the chevron barracuda, is a species of barracuda that ranges from the Red Sea and East Africa to the Indo and Western Pacific and as far as French Polynesia. The blackfin barracuda reaches a...



Blackfin icefish

Chaenocephalus aceratus

The blackfin icefish (Chaenocephalus aceratus), also known as the Scotia Sea icefish, is a species of crocodile icefish belonging to the family Channichthyidae. The blackfin icefish belongs to Notothenioidei, a suborder of fishes that accounts...



Blackfin scad

Alepes melanoptera

The blackfin scad (Alepes melanoptera) (also known as the shortfinned trevally and Chinese trevally) is a species of tropical marine fish of the jack family Carangidae. The species inhabits inshore waters throughout the Indo-Pacific region,...



Blackfin seabass

Lateolabrax latus

The blackfin seabass (Lateolabrax latus) is a Perciforme fish in the family lateolabracidae, found primarily in the shallow waters of the Pacific coast of Asia, in Japan and in South Korea. There are only two species in the genus Lateolabrax,...



Blackfin snapper

Lutjanus buccanella

The blackfin snapper (Lutjanus buccanella), also known as the blackspot snapper, blackfin red snapper, gun-mouth backfin, gun-mouth snapper, redfish and wrenchman is a species of marine ray-finned fish, a snapper belonging to the family...



Blackfin tuna

Thunnus atlanticus

The blackfin tuna (Thunnus atlanticus) is a species of tuna in the family Scombridae. It is occasionally referred to as the Bermuda tuna, blackfinned albacore, or deep bodied tunny. They are the smallest tuna species in the genus Thunnus,...



Blackline rasbora

Rasbora borapetensis

The blackline rasbora (Rasbora borapetensis) is a fish of the family Cyprinidae found in Asia in the Mekong, Chao Phraya, and Mae Klong basins, and also the northern Malay Peninsula. In the aquarium trade, it is known by a variety of other names,...



Blackmouth catshark

Galeus melastomus

The blackmouth catshark (Galeus melastomus) is a species of catshark, and part of the family Scyliorhinidae, common in the northeastern Atlantic Ocean from Iceland to Senegal, including the Mediterranean Sea. It is typically found over the...



Blacknose shark

Carcharhinus acronotus

The blacknose shark (Carcharhinus acronotus) is a species of requiem shark, belonging to the family Carcharhinidae, common in the tropical and subtropical waters of the western Atlantic Ocean. This species generally inhabits coastal seagrass,...



Blacknose shiner

Notropis heterolepis

The blacknose shiner (Notropis heterolepis) is a species of fish belonging to the family Leuciscidae.



Blacksaddle filefish

Paraluteres prionurus

The blacksaddle filefish, Paraluteres prionurus, is a filefish of the family Monacanthidae. It reaches a length of a maximum 11 cm. The blacksaddle filefish are found in pairs or in small schools and inhabit reefs across the Indian and Pacific...



Blacksail snake mackerel

Thyrsitoides marleyi

The blacksail snake mackerel (Thyrsitoides marleyi), known also as the black snoek, is a species of snake mackerel found in the Indo-Pacific from shallow water to a depth of at least 400 m (1,300 ft) where they appear to prefer slopes on...



Blackside hawkfish

Paracirrhites forsteri

The black-side hawkfish (Paracirrhites forsteri), freckled hawkfish or Forster's hawkfish, is a species of marine ray-finned fish, a hawkfish belonging to the family Cirrhitidae. It is from the Indo-Pacific. It is occasionally found in the...



Blackspot climbing perch

Ctenopharynx intermedius

The blackspot climbing perch (Ctenopharynx intermedius) is a species of fish in the family Cichlidae. It is native to Lake Malawi and the upper Shire River.



Blackspot seabream

Pagellus bogaraveo

The blackspot seabream (Pagellus bogaraveo), also known as the red seabream and as the besugo, is a species of marine ray-finned fish belonging to the family Sparidae, which includes the seabreams and porgies. This is a widespread species in the...



Blackspot shark

Carcharhinus sealei

The blackspot shark (Carcharhinus sealei) is a small species of requiem shark in the family Carcharhinidae found in the tropical Indo-West Pacific Ocean between latitudes 24°N and 30°S, from the surface to a depth around 40 m (130 ft). Its length...



Blackspot tuskfish

Choerodon schoenleinii

The blackspot tuskfish (Choerodon schoenleinii) is a wrasse native to the Indian Ocean and the western Pacific Ocean from Mauritius to Indonesia and Australia north to the Ryukyus. This species occurs on reefs, preferring areas with sandy...



Blackspotted puffer

Arothron nigropunctatus

The blackspotted puffer (Arothron nigropunctatus), also known as the dog-faced puffer, is a tropical marine fish belonging to the family Tetraodontidae.



Blackstripe topminnow

Fundulus notatus

The blackstripe topminnow, Fundulus notatus, is a small freshwater fish in the family Fundulidae, found in central North America.



Blacktail butterflyfish

Chaetodon austriacus

The blacktail butterflyfish (Chaetodon austriacus), also known as black-tailed butterflyfish or exquisite butterflyfish, is a species marine ray-finned fish, a butterflyfish belonging to the family Chaetodontidae. It is native to the western...



Blacktail shiner

Cyprinella venusta

The blacktail shiner (Cyprinella venusta) is a small freshwater fish in the family Cyprinidae native to the United States.



Blackthroat seaperch

Doederleinia berycoides

The blackthroat seaperch (Doederleinia berycoides), also known as the rosy seabass, is a species of fish in the family Acropomatidae, the temperate ocean-basses or lanternbellies. It is the only species in the monotypic genus Doederleinia. It is...



Blacktip grouper

Epinephelus fasciatus

The blacktip grouper (Epinephelus fasciatus), also known as the redbanded grouper, blacktipped cod, black-tipped rockcod, footballer cod, red-barred cod, red-barred rockcod, scarlet rock-cod or weathered rock-cod, is a species of marine...



Blacktip reef shark

Carcharhinus melanopterus

The blacktip reef shark (Carcharhinus melanopterus) is a species of requiem shark, in the family Carcharhinidae, which can be easily identified by the prominent black tips on its fins (especially on the first dorsal fin and its caudal fin). Among...



Blacktip shark

Carcharhinus limbatus

The blacktip shark (*Carcharhinus limbatus*) is a species of requiem shark, and part of the family Carcharhinidae. It is common to coastal tropical and subtropical waters around the world, including brackish habitats. Genetic analyses have revealed...



Blacktip tope

Hypogaleus hyugaensis

Sharks portalThe blacktip tope (*Hypogaleus hyugaensis*), also known as pencil shark or blacktip topeshark, is a houndshark of the family Triakidae, and the only member of the genus *Hypogaleus*. It is found in the deep waters of the continental...



Blacktip trevally

Caranx heberi

The blacktip trevally (Caranx heberi), also known as the blacktip kingfish or yellowtail kingfish, is a species of large marine fish classified in the jack family Carangidae. The blacktip trevally is distributed throughout the tropical to...



Bleeding heart tetra

Hyphessobrycon erythrostigma

The bleeding heart tetra, Hyphessobrycon erythrostigma, is a freshwater tropical fish native to the Upper Amazon River basin. It grows to the size of 2.5 in (64 mm) and lives for around five years.



Blicca bjoerkna

Blicca bjoerkna

Blicca bjoerkna, alternatively called the white bream or the silver bream, is a European species of freshwater fish in the family Cyprinidae.



Blind cave eel

Ophisternon candidum

The blind cave eel (Ophisternon candidum) is a species of cavefish in the family Synbranchidae. It is the longest cavefish in Australia (up to 40 cm (16 in)) and one of the only three vertebrates in Australia that is restricted to underground...



Blind electric ray

Typhlonarke aysoni

The blind electric ray (*Typhlonarke aysoni*) is a little-known species of sleeper ray in the family Narkidae, endemic to New Zealand. It is found on the bottom, typically at a depth of 300–400 m (980–1,310 ft). Reaching 38 cm (15 in) in length....



Blind shark

Brachaelurus waddi

The blind shark (*Brachaelurus waddi*) is one of two species of carpet sharks in the family Brachaeluridae, along with the bluegrey carpetshark (*Brachaelurus colcloughi*). Found along the coast of eastern Australia, this nocturnal, bottom-dwelling...



Blind swamp eel

Ophisternon infernale

The blind swamp eel (Ophisternon infernale) is a species of fish in the family Synbranchidae. It is endemic to Mexico where it lives in cave systems and is known in Spanish as the anguila ciega. The International Union for Conservation of Nature...



Blob sculpin

Psychrolutes phrictus

The blob sculpin (Psychrolutes phrictus) is a species of deep-sea fish of the family Psychrolutidae. It feeds mainly on crustaceans, molluscs, and sea pens. It lives off the continental shelves in very deep water (839–2800 m) in the North Pacific...



Blonde ray

Raja brachyura

The blonde ray or blonde skate (Raja brachyura) is a species of ray fish in the family Rajidae.



Bloodfin tetra

Aphyocharax anisitsi

The bloodfin tetra (Aphyocharax anisitsi) is a species of characin from the Paraná River basin in South America. The bloodfin is a relatively large tetra, growing to 5.5 cm. Its notable feature (as the name suggests) is the blood-red colouration...



Blotched foxface

Siganus unimaculatus

The blotched foxface (*Siganus unimaculatus*), also called the blackblotch foxface or one-spot foxface, is a species of marine ray-finned fish, a rabbitfish belonging to the family Siganidae. It is found at reefs and lagoons in the central...



Blotched snakehead

Channa maculata

The blotched snakehead (Channa maculata) is a species of snakehead. It is one of four species of the genus Channa native to China. It is also to native northern Vietnam and Taiwan, but has been widely introduced to other countries, where it is an...



Blotcheye soldierfish

Myripristis berndti

The blotcheye soldierfish (Myripristis berndti) is a species of soldierfish belonging to the family Holocentridae.



Blotchy swellshark

Cephaloscyllium umbratile

The blotchy swellshark, or Japanese swellshark, (Cephaloscyllium umbratile) is a common species of catshark, belonging to the family Scyliorhinidae. The Blotchy swellshark is found at depths of 90–200 m (300–660 ft) in the northwestern Pacific...



Bludger

Carangoides gymnostethus

The bludger (Carangoides gymnostethus), also known as the bludger trevally, nakedbreast trevally or Bleeker's jackfish, is a widespread species of large marine fish in the jack family, Carangidae. The bludger inhabits the tropical and subtropical...



Blue acara

Andinoacara pulcher

The blue acara (Andinoacara pulcher) is a colorful freshwater fish in the cichlid family. This fish can be found in various freshwater habitats, ranging from standing water to flowing streams, in Venezuela and Trinidad. They can reach lengths of...



Blue and gold snapper

Lutjanus viridis

The blue-and-gold snapper (Lutjanus viridis) is a species of marine ray-finned fish, a snapper belonging to the family Lutjanidae. It is native to the eastern Pacific Ocean where it is important to local subsistence fisheries.



Blue blanquillo

Malacanthus latovittatus

The blue blanquillo, Malacanthus latovittatus, also known as the banded blanquillo, striped blanquillo, false whiting, sand tilefish or eye of the sea, is a species of marine ray-finned fish, a tilefish belonging to the family Malacanthidae. It...



Blue catfish

Ictalurus furcatus

The blue catfish (*Ictalurus furcatus*) is a large species of North American catfish, reaching a length of 65 in (170 cm) and a weight of 165 lb (75 kg). The continent's largest, it can live to 20 years, with a typical fish being between 25–46 in...



Blue cod

Parapercis colias

The New Zealand blue cod (*Parapercis colias*) is a temperate marine fish of the family Pinguipedidae. It is also known by its Māori names, rāwaru, pākirikiri and patutuki, and by its other names in English, Boston blue cod, New Zealand cod or sand...



Blue danio

Danio kerri

The blue danio, Kerr's danio or turquoise danio (*Danio kerri*), is a tropical fish belonging to the genus *Danio* in family Cyprinidae.



Blue grenadier

Macruronus novaezelandiae

The blue grenadier (also known as hoki, blue hake, New Zealand whiptail, or whiptail hake, *Macruronus novaezelandiae*) is a merluccid hake of the family Merlucciidae found around southern Australia and New Zealand, as well as off both the Atlantic...



Blue jack mackerel

Trachurus picturatus

The blue jack mackerel (Trachurus picturatus) is a species of mackerel-like fish in the family Carangidae. Their maximum reported length is 60 cm, with a common length of 25 cm. They are coastal fish found at depths to 370 m off the Bay of Biscay...



Blue ling

Molva dypterygia

The blue ling (Molva dypterygia) is a member of the cod family from the North Atlantic. It is usually 70 to 110 cm long, but the maximum length is 155 cm. Blue ling feed on fish (flatfishes, gobies, rocklings) and crustaceans and benthic...



Blue lyretail

Fundulopanchax gardneri

The blue lyretail (Fundulopanchax gardneri), also known as the Gardner's killi and formerly as the steel-blue aphyosemion. It is a species of killifish endemic to freshwater habitats in Nigeria and Cameroon.



Blue mackerel

Scomber australasicus

The blue mackerel (Scomber australasicus), also called Japanese mackerel, Pacific mackerel, slimy mackerel or spotted chub mackerel, is a fish of the family Scombridae, found in tropical and subtropical waters of the Pacific Ocean from Japan...



Blue maomao

Scorpiis violacea

The blue maomao (Scorpiis violacea), also known as the violet sweep, blue sweep or hardbelly, is a species of marine ray-finned fish, a member of the subfamily Scorpioninae, part of the sea chub family Kyphosidae. It is native to the southwestern...



Blue mbuna

Labeotropheus fuelleborni

The blue mbuna (*Labeotropheus fuelleborni*) is a species of cichlid found in Lake Malawi where it inhabits areas with rocky substrates. This species can reach a length of 30 centimetres (12 in) SL. This species is important to local commercial...



Blue moki

Latridopsis ciliaris

Blue moki (*Latridopsis ciliaris*) is a species of marine ray finned fish belonging to the family Latridae, the trumpeters. It is native to the southwestern Pacific Ocean around New Zealand and occasionally off southeastern Australia at depths of...



Blue panchax

Aplocheilichthys panchax

The blue panchax or whitespot (*Aplocheilichthys panchax*) is a common freshwater fish found in a large variety of habitats due to its high adaptability. This species is native to southern Asia from Pakistan, India to Indonesia. It has been discovered...



Blue parrotfish

Scarus coeruleus

The blue parrotfish (*Scarus coeruleus*) is a member of the parrotfish genus *Scarus*. It is found on coral reefs in shallow water in the tropical and subtropical parts of the western Atlantic Ocean and the Caribbean Sea. They usually forage in a...



Blue rockfish

Sebastes mystinus

The blue rockfish (*Sebastes mystinus*) or blue seaperch, is a species of marine ray-finned fish belonging to the subfamily Sebastinae, the rockfishes, part of the family Scorpaenidae. It is found in the northeastern Pacific Ocean, ranging from...



Blue runner

Caranx crysos

The blue runner (Caranx crysos), also known as the bluestripe jack, Egyptian scad, hardtail jack or hardnose, is a common species of moderately large marine fish classified in the jack family, Carangidae. The blue runner is distributed across the...



Blue shark

Prionace glauca

The blue shark (Prionace glauca), also known as the great blue shark, is a species of requiem shark, in the family Carcharhinidae, which inhabits deep waters in the world's temperate and tropical oceans. Averaging around 3.1 m (10 ft) and...



Blue stingray

Dasyatis chrysonota

The blue stingray (Dasyatis chrysonota) is a species of whiptail stingray of the family Dasyatidae often found in the coastal waters of southern Africa.



Blue sucker

Cycleptus elongatus

The blue sucker (Cycleptus elongatus) is a long-lived freshwater species of fish in the sucker family that is of conservation concern. The species has an average weight of 2–3 kilograms (4.4–6.6 lb) and an average length of 76 centimetres (30...



Blue trevally

Carangoides ferdau

The blue trevally (Carangoides ferdau), also known as the banded trevally, barred trevally, Ferdau's trevally or Forskaal's jackfish, is a common, widespread species of pelagic marine fish classified in the jack family, Carangidae. The blue...



Blue tuskfish

Choerodon cyanodus

The blue tuskfish, Choerodon cyanodus, is a species of wrasse native to the Indian and western Pacific Oceans, where it is known to occur around Australia, but has been claimed to occur more widely. It inhabits reefs. This species can reach a...



Blue warehou

Seriolella brama

The blue warehou (*Seriolella brama*) or common warehou is a medusafish of the family Centrolophidae found off southern Australia and around New Zealand, at depths of between 5m and 400m. Its length is up to about 75 cm. The blue warehou is not to...



Blue weed whiting

Haletta semifasciata

The blue weed whiting (*Haletta semifasciata*) is a species of ray-finned fish, a weed whiting from the family Odacidae, which is endemic to Australia where it is only found along the southern coast. It is found in brackish and marine waters in...



Blue whiting

Micromesistius poutassou

The blue whiting (*Micromesistius poutassou*) one of the two species in the genus *Micromesistius* in the family Gadidae, which also contains cod, haddock, whiting, and pollock. It is common in the northeast Atlantic Ocean from Morocco to Iceland and...



Blue-eyed plec

Panaque cochliodon

The blue-eyed plec (*Panaque cochliodon*) is a herbivorous freshwater armored catfish endemic to Colombia where it occurs in the Cauca and Magdalena River basins. It is distinguished from many loricariids by being dark grey to black and having...



Blue-spotted jawfish

Opistognathus rosenblatti

The blue-spotted jawfish (*Opistognathus rosenblatti*) is a species of jawfish native to the Gulf of California. It is an inhabitant of reefs where it is found in large colonies at depths of around 12 metres (39 ft). This species hides in its...



Blue-spotted spinefoot

Siganus corallinus

The blue-spotted spinefoot (Siganus corallinus), the coral rabbitfish, coral spinefoot, ocellated spinefoot or orange spinefoot is a species of marine ray-finned fish, a rabbitfish belonging to the family Siganidae. It is found in the...



Blueback herring

Alosa aestivalis

The blueback herring, blueback shad, or summer shad (Alosa aestivalis) is an anadromous species of herring from the east coast of North America, with a range from Nova Scotia to Florida. Blueback herring form schools and are believed to migrate...



Bluebarred pygmy sunfish

Elassoma okatie

The bluebarred pygmy sunfish, Elassoma okatie, is a species of pygmy sunfish endemic to South Carolina, United States, where it prefers waters with dense vegetation growth in the Edisto and Savannah River drainages. This species can reach 3.4 cm...



Bluecheek butterflyfish

Chaetodon semilarvatus

The blue-cheeked butterflyfish (Chaetodon semilarvatus) is a species of marine ray-finned fish, a butterflyfish belonging to the family Chaetodontidae. It is found in the northwestern Indian Ocean.



Bluefin gurnard

Chelidonichthys kumu

The bluefin gurnard or Pacific red gurnard (Chelidonichthys kumu) is a species of marine ray-finned fishes belonging to the family Triglidae, the gurnards and sea robins. Its Māori names are kumukumu and pūwahaiau. It is found in the western...



Bluefin trevally

Caranx melampygus

The bluefin trevally (Caranx melampygus), also known as the bluefin jack, bluefin kingfish, bluefanned crevalle, blue ulua, omilu, and spotted trevally, is a species of large, widely distributed marine fish classified in the jack family,...



Bluefish

Pomatomus saltatrix

The bluefish (Pomatomus saltatrix) is the only extant species of the family Pomatomidae. It is a marine pelagic fish found around the world in temperate and subtropical waters, except for the northern Pacific Ocean. Bluefish are known as tailor...



Bluegill

Lepomis macrochirus

The bluegill (Lepomis macrochirus), sometimes referred to as "bream," "brim," "sunny," or, as is common in Texas, "copper nose", is a species of North American freshwater fish, native to and commonly found in streams, rivers, lakes, ponds and...



Bluegrey carpetshark

Brachaelurus colcloughi

The bluegrey carpetshark (Brachaelurus colcloughi) or Colclough's shark, is an uncommon species of carpet shark endemic to shallow inshore waters off northeastern Australia. It is one of the two extant members of the family Brachaeluridae. The...



Bluehead chub

Nocomis leptcephalus

The bluehead chub (Nocomis leptcephalus) is a cyprinid native to North America. Its name is due to its appearance, as breeding males have a blue head. Adult bluehead chubs are, on average, between 70 and 160 mm (2.8 and 6.3 in) in length. They...



Bluelashed butterflyfish

Chaetodon bennetti

The bluelashed butterflyfish (Chaetodon bennetti), also known as the eclipse butterflyfish, archer butterflyfish or Bennett's butterflyfish, is a species of marine ray-finned fish, a butterflyfish belonging to the family Chaetodontidae. It is...



Bluering angelfish

Pomacanthus annularis

The bluering angelfish (Pomacanthus annularis), also known as the Annularis Angelfish and the Blue King Angelfish, is a species of marine ray-finned fish, a marine angelfish belonging to the family Pomacanthidae. It is member of the genus...



Blueside wrasse

Cirrhilabrus cyanopleura

The blueside wrasse, Cirrhilabrus cyanopleura, is a species of wrasse native to the eastern Indian Ocean and the western Pacific Ocean. It is found on reefs in schools swimming 1 to 2 m (3.3 to 6.6 ft) above the bottom. It occurs at depths from 2...



Bluespine unicornfish

Naso unicornis

The bluespine unicornfish (*Naso unicornis*), also known as the short-nose unicornfish, is a species of marine ray-finned fish belonging to the family Acanthuridae, the surgeonfishes, unicornfishes and tangs. This species is found in the...



Bluespot mullet

Crenimugil seheli

The bluespot mullet (Crenimugil seheli) is a member of the ray-finned fish family Mugilidae found worldwide in coastal temperate and tropical waters, and in some species in fresh water. Crenimugil seheli have served as an important source of food...



Bluespotted cornetfish

Fistularia commersonii

The bluespotted cornetfish (Fistularia commersonii), also known as smooth cornetfish or smooth flutemouth, is a marine fish which belongs to the family Fistulariidae. This very long and slender reef-dweller belongs to the same order as the...



Bluespotted ribbontail ray

Taeniura lymma

The bluespotted ribbontail ray (Taeniura lymma) is a species of stingray in the family Dasyatidae. Found from the intertidal zone to a depth of 30 m (100 ft), this species is common throughout the tropical Indian and western Pacific Oceans in...



Bluespotted sunfish

Enneacanthus gloriosus

The bluespotted sunfish (Enneacanthus gloriosus) is a species of fish in the family Centrarchidae, the sunfishes. It is native to the southeastern and eastern United States, its distribution extending as far north as New Jersey, Pennsylvania, and...



Bluestreak cleaner wrasse

Labroides dimidiatus

The bluestreak cleaner wrasse, *Labroides dimidiatus*, is one of several species of cleaner wrasses found on coral reefs from Eastern Africa and the Red Sea to French Polynesia. Like other cleaner wrasses, it eats parasites and dead tissue off...



Bluestriped fangblenny

Plagiotremus rhinorhynchos

Plagiotremus rhinorhynchos, commonly called the bluestriped fangblenny, is a species of combtooth blenny found in coral reefs in the Pacific and Indian ocean. This species reaches a length of 12 centimetres (4.7 in) SL. It is also known as the...



Bluestriped grunt

Haemulon sciurus

The bluestriped grunt (*Haemulon sciurus*), also known as the boar grunt, golden grunt, humpback grunt, redmouth grunt, or yellow grunt, is a species of marine ray-finned fish, a grunt belonging to the family Haemulidae. It is found in the western...



Blunthead puffer

Sphoeroides pachygaster

The blunthead puffer, *Sphoeroides pachygaster*, is a pufferfish of the family Tetraodontidae, found circumglobally in tropical and temperate seas, at depths between 50 and 500 m. First recorded in the Mediterranean Sea off the Spanish coast in...



Bluntnose minnow

Pimephales notatus

The bluntnose minnow (*Pimephales notatus*) is a species of temperate freshwater fish belonging to the genus *Pimephales* of the cyprinid family. Its natural geographic range extends from the Great Lakes south along the Mississippi River basin to...



Bluntnose sixgill shark

Hexanchus griseus

The bluntnose sixgill shark (Hexanchus griseus), often simply called the cow shark, is the largest hexanchoid shark, growing to 20 ft (6.1 m) in length. It is found in tropical and temperate waters worldwide and its diet is widely varied by...



Bluntnose stingray

Hypanus say

The bluntnose stingray or Say's stingray (Hypanus say, often misspelled sayi) is a species of stingray in the family Dasyatidae, native to the coastal waters of the western Atlantic Ocean from the U.S. state of Massachusetts to Venezuela. It is a...



Bocaccio rockfish

Sebastes paucispinis

The bocaccio rockfish (Sebastes paucispinis) is a species of marine ray-finned fish belonging to the subfamily Sebastinae, the rockfishes, part of the family Scorpaenidae. It is found in the northeast Pacific Ocean.



Bodianus mesothorax

Bodianus mesothorax

Bodianus mesothorax, the split-level hogfish, blackbelt hogfish, black-belt hogfish, coral hogfish, eclipse hogfish, eclipse pigfish, mesothorax hogfish or yellow-spotted hogfish, is a species of wrasse native to the western Pacific Ocean and the...



Boeseman's rainbowfish

Melanotaenia boesemani

Boeseman's rainbowfish (Melanotaenia boesemani), also known as the Boesemani rainbowfish, is a species of fish in the family Melanotaeniidae.



Bombay duck

Harpadon nehereus

Harpadon nehereus, (common names include Bombay duck, bummalu, bombil, bombili, boomla, lote, loitta) is a species of lizardfish. Adults may reach a maximum length of 40 cm (16 in), but the usual size is around 25 cm (10 in).



Bonefish

Albula vulpes

The bonefish (Albula vulpes) is the type species of the bonefish family (Albulidae), the only family in order Albuliformes.



Bonnethead

Sphyrna tiburo

The bonnethead (Sphyrna tiburo), also called a bonnet shark or shovelhead, is a small member of the hammerhead shark genus Sphyrna, and part of the family Sphyrnidae. It is an abundant species in the littoral zone of the North Atlantic and the...



Bonneville cisco

Prosopium gemmifer

The Bonneville cisco (*Prosopium gemmifer*) is a species of cisco endemic to Bear Lake along the Utah-Idaho border, United States. It is one of three species of whitefish endemic to Bear Lake, the others being the Bear Lake whitefish and the...



Bonneville cutthroat trout

Oncorhynchus clarkii utah

The Bonneville cutthroat trout (Oncorhynchus clarkii utah) is a subspecies of cutthroat trout native to tributaries of the Great Salt Lake and Sevier Lake. Most of the fish's current and historic range is in Utah, but they are also found in...



Bony-eared assfish

Acanthonus armatus

The bony-eared assfish (Acanthonus armatus) is a bathypelagic species of cusk-eel found in tropical and sub-tropical oceans at depths of from 1,171 to 4,415 metres (3,842 to 14,485 ft). It has been found as far north as Queen Charlotte Sound off...



Bonytail chub

Gila elegans

The bonytail chub or bonytail (Gila elegans) is a cyprinid freshwater fish native to the Colorado River basin of Arizona, California, Colorado, Nevada, New Mexico, Utah and Wyoming in the southwestern United States; it has been extirpated from...



Boops boops

Boops boops

Boops boops (; from Ancient Greek βόωψ, literally 'ox-eyed'), commonly called the bogue, is a species of seabream native to the eastern Atlantic.



Boraras brigittae

Boraras brigittae

Boraras brigittae is a species of ray-finned fish in the genus Boraras, also known as chili rasboras or mosquito rasboras, are a very small species of fishes native to the swamps of South West Borneo, Indonesia. It is found natively in blackwater...



Boraras merah

Boraras merah

Boraras merah is a species of ray-finned fish in the genus Boraras, also known as Phoenix rasbora. It is between 15 and 20mm long with a red base colour to the body.



Boraras naevus

Boraras naevus

Boraras naevus is a fish in the family Cyprinidae endemic to Thailand. Its locations include swampy areas north of Surat Thani, the Tapi River drainage (located on the Gulf of Thailand), and the Andaman Sea slope of the Malay Peninsula near Trang.



Boreogadus saida

Boreogadus saida

Boreogadus saida, known as the polar cod or as the Arctic cod, is a fish of the cod family Gadidae, related to the true cod (genus Gadus). Another fish species for which both the common names Arctic cod and polar cod are used is Arctogadus...



Borna snakehead

Channa amphibeus

The Borna snakehead, or Chel snakehead, Channa amphibeus, is an extremely rare species of snakehead. C. amphibeus is native only to the Chel River (in the drainage of the Brahmaputra), of the Bhutan foothills, in northern Bengal, India.As with...



Borneo shark

Carcharhinus borneensis

The Borneo shark (Carcharhinus borneensis) is a species of requiem shark, and part of the family Carcharhinidae. Extremely rare, it is known only from inshore waters around Mukah in northwestern Borneo, though it may once have been more widely...



Borriqueta porgy

Boridia grossidens

The Borriqueta porgy (Boridia grossidens) is a species of marine ray-finned fish, a grunt belonging to the family Haemulidae. It is native to the Atlantic Coast of South America.



Bothus podas

Bothus podas

Bothus podas, also known as the wide-eyed flounder, is a flounder in the genus Bothus, native to the Mediterranean Sea and the Atlantic Coast of Africa.During the reproductive season, males court and mate successively with females in their...



Botia dario

Botia dario

Botia dario (Bengal loach or queen loach) is a species of fish in the loach family Botiidae found in the Brahmaputra and Ganges basins in Bangladesh, Bhutan and northern India. The species is overall widespread. In Bangladesh, B. dario is...



Bowfin

Amia calva

The bowfin (Amia calva) is a bony fish, native to North America. Common names include mudfish, mud pike, dogfish, grindle, grinnel, swamp trout, and choupique. It is regarded as a relict, being one of only two surviving species of the...



Brachygobius doriae

Brachygobius doriae

Brachygobius doriae, the bumblebee goby, is a species of goby native to fresh and brackish waters of Indonesia, Brunei and Singapore, Malaysia, Cambodia, Thailand and Vietnam. This species can reach a length of 4.2 centimetres (1.7 in) TL. It is...



Brama australis

Brama australis

Brama australis, the Southern Ray's bream or Pacific pomfret, is a species of marine ray-finned fish from the family Bramidae, the pomfrets. It is found in the southern Oceans.



Bramble shark

Echinorhinus brucus

The bramble shark (Echinorhinus brucus) is one of the two species of sharks in the family Echinorhinidae. Aside from the eastern Pacific Ocean, it is found in tropical and temperate waters worldwide. This rarely encountered shark swims close to...



Branchiostegus japonicus

Branchiostegus japonicus

Branchiostegus japonicus, the horsehead tilefish, Japanese horsehead tilefish, red amadai or the red tilefish, is a species of marine ray-finned fish, a tilefish belonging to the family Malacanthidae. It is native to the western Pacific Ocean.



Brassy minnow

Hybognathus hankinsoni

The brassy minnow (Hybognathus hankinsoni) is a species of fish belonging to the family Cyprinidae. The family Cyprinidae consists of mainly freshwater minnows and carps. The fish gets its scientific name (Hybognathus hankinsoni) from the Greek...



Brassy trevally

Caranx papuensis

The brassy trevally, Caranx papuensis (also known as the brassy kingfish, Papuan trevally, tea-leaf trevally, and green back trevally) is a species of large marine fish classified in the jack family, Carangidae.



Brazilian flathead

Percophis brasiliensis

The Brazilian flathead (Percophis brasiliensis) is a species of duckbill which is the only species in the genus Percophis, the type genus of the monotypic subfamily Percophinae of the duckbill family Percophidae. It occurs in the south western...



Brazilian large-eyed stingray

Dasyatis marianae

The Brazilian large-eyed stingray, *Hypanus marianae*, is a species of stingray in the family Dasyatidae. Endemic to northeastern Brazil, adults of this species inhabit shallow coral and sandstone reefs while the young are also found near beaches...



Brazilian sharpnose shark

Rhizoprionodon lalandii

The Brazilian sharpnose shark (*Rhizoprionodon lalandii*) is a requiem shark of the family Carcharhinidae. It is found in the tropical waters of the western Atlantic Ocean between latitudes 13° N and 33° S, at depths between 3 and 70 m; it has been...



Brazilian snapper

Lutjanus alexandrei

The Brazilian snapper, (Lutjanus alexandrei) is a species of snapper native to the tropical waters of the Atlantic off the coast of Brazil.



Brevitrygon walga

Brevitrygon walga

Brevitrygon walga, the dwarf whipray or mangrove whipray, is a small stingray, a cartilaginous fish in the family Dasyatidae. It is a demersal fish and is found over the continental and insular shelf of the west central Pacific Ocean where it is...



Bridgelip sucker

Catostomus columbianus

The bridgelip sucker (Catostomus columbianus) is a fish in the family Catostomidae that occupies the Columbia River system. Like all sucker fish, they live only in fresh water. Bridgelip suckers have a tendency to live in deeper waters during the...



Brill

Scophthalmus rhombus

The brill (Scophthalmus rhombus) is a species of flatfish in the turbot family (Scophthalmidae) of the order Pleuronectiformes. Brill can be found in the northeast Atlantic, Black Sea, Baltic Sea, and Mediterranean, primarily in deeper offshore...



Brilliant rasbora

Rasbora einthovenii

The brilliant rasbora (Rasbora einthovenii) is a species of ray-finned fish in the genus Rasbora. It is found in the Malay Peninsula and in Borneo. Adults can grow up to 5 cm (2 inches)



Brindled madtom

Noturus miurus

The brindled madtom (Noturus miurus) is a small catfish of the family Ictaluridae that is native to the eastern United States.



Broad stingray

Bathytoshia lata

The Broad stingray (Bathytoshia lata), also known as the Brown stingray or Hawaiian stingray, is a species of stingray in the family Dasyatidae. They range across the Indo-Pacific from southern Africa to Hawaii, and are the predominant species of...



Broad whitefish

Coregonus nasus

The broad whitefish (Coregonus nasus) is a freshwater whitefish species. Dark silvery in colour, and like a herring in its shape. Its distinctive features includes: a convex head, short gill rakers, and a mild overbite. It is found in the...



Broadbarred king mackerel

Scomberomorus semifasciatus

The broadbarred king mackerel or grey mackerel (Scomberomorus semifasciatus) is a species of fish in the family Scombridae found in tropical waters of the western Pacific, along the northern coast of Australia and the southern coast of Papua New...



Broadfin shark

Lamiopsis temminckii

Sharks portalThe broadfin shark (*Lamiopsis temminckii*) is a tropical whaler shark, characterized by the broad shape of its pectoral fins. They are classified as requiem sharks of the family Carcharhinidae. This is one of two species in the genus...



Broadhead catfish

Clarias macrocephalus

The broadhead catfish (*Clarias macrocephalus*) is an economically important air-breathing catfish that is a native of Southeast Asia. The fish is farmed in ponds for use in human consumption in countries such as Thailand (Thai: ปลาอุกอุย) and the...



Broadnose sevengill shark

Notorynchus cepedianus

The broadnose sevengill shark (*Notorynchus cepedianus*) is the only extant member of the genus *Notorynchus*, in the family Hexanchidae. It is recognizable because of its seven gill slits, while most shark species have five gill slits, with the...



Broadnosed pipefish

Syngnathus typhle

The broadnosed pipefish or deepnosed pipefish (*Syngnathus typhle*) is a fish of the family Syngnathidae (seahorses and pipefishes). It is native to the Eastern Atlantic from Vardø in Norway, Baltic Sea (north to the Gulf of Finland) and the...



Bronze corydoras

Corydoras aeneus

The bronze corydoras (Corydoras aeneus), also known as the green corydoras, bronze catfish, lightspot corydoras or wavy catfish, is a species of freshwater fish in the armored catfish family, Callichthyidae, often kept as an aquarium fish. It is...



Bronze featherback

Notopterus notopterus

The bronze featherback (Notopterus notopterus; Assamese: কাঁহুলি kandhuli, Bengali: ফলি, Bengali: কাংলা, Thai: ปลาซลาด, ปลาฉลาด, ปลาดอง, Vietnamese: Cá thát lát, Burmese: ငါးဖတ်) is a ray-finned fish in the family Notopteridae found in South and...



Brook silverside

Labidesthes sicculus

Labidesthes sicculus, also known as the Brook silverside is a North American species of Neotropical silverside. The brook silverside lives in slow moving rivers and lakes from the Great Lakes to the Mississippi Basin and Gulf Coastal Plains. The...



Brook trout

Salvelinus fontinalis

The brook trout (Salvelinus fontinalis) is a species of freshwater fish in the char genus Salvelinus of the salmon family Salmonidae. It is native to Eastern North America in the United States and Canada, but has been introduced elsewhere in...



Broomtail wrasse

Cheilinus lunulatus

The broomtail wrasse (Cheilinus lunulatus) is a species of wrasse native to the Red Sea and Indian Ocean.



Brotula barbata

Brotula barbata

Brotula barbata, commonly known as the bearded brotula, Atlantic bearded brotula, or sugarcod, is a species of cusk-eel in the genus Brotula. It lives in the Atlantic Ocean, in depths of up to 300 meters. Its coloring ranges from olive-brown to...



Brown bullhead

Ameiurus nebulosus

The brown bullhead (Ameiurus nebulosus) is a fish of the family Ictaluridae that is widely distributed in North America. It is a species of bullhead catfish and is similar to the black bullhead (Ameiurus melas) and yellow bullhead (Ameiurus...



Brown catshark

Apristurus brunneus

The brown catshark (Apristurus brunneus) is commonly found in the Pacific Ocean, ranging from the northern Pacific waters off the coast of British Columbia and south to the Baja California peninsula in Mexico. They may live as far south as...



Brown garden eel

Heteroconger longissimus

The brown garden eel (Heteroconger longissimus), also known simply as the garden eel, is an eel in the family Congridae (conger/garden eels). It was described by Albert Günther in 1870. It is a tropical, marine eel which is known from the eastern...



Brown ghost knifefish

Apteronotus leptorhynchus

The brown ghost knifefish (Apteronotus leptorhynchus) is a species of weakly electric knifefish in the family Apteronotidae. Only the brown ghost knifefish, a vertebrate, has been proven to have negligible brain aging thus far. In the current...



Brown meagre

Sciaena umbra

The brown meagre or corb (Sciaena umbra) is a species of marine ray-finned fish belonging to the family Sciaenidae, the drums and croakers. This species found in, the eastern Atlantic, Mediterranean Sea and Black Sea occurring in shallow waters...



Brown rockfish

Sebastes auriculatus

The brown rockfish (Sebastes auriculatus), whose other names include brown seaperch, chocolate bass, brown bass and brown bomber, is a species of marine ray-finned fish belonging to the subfamily Sebastinae, the rockfishes, part of the family...



Brown shyshark

Haploblepharus fuscus

The brown shyshark or plain happy (Haploblepharus fuscus) is a species of catshark, part of the family Scyliorhinidae. It is endemic to the shallow, coastal waters of South Africa from west of Cape Agulhas to KwaZulu-Natal. This benthic species...



Brown trout

Salmo trutta

The brown trout (*Salmo trutta*) is a species of salmonid ray-finned fish and the most widely distributed species of the genus *Salmo*, endemic to most of Europe, West Asia and parts of North Africa, and has been widely introduced globally as a game...



Brown wrasse

Labrus merula

The brown wrasse (*Labrus merula*) is a species of wrasse native to the Eastern Atlantic from Portugal to Morocco, including the Azores, as well as in the Mediterranean Sea. This species can reach 45 cm (18 in) in standard length, though most do...



Brown-marbled grouper

Epinephelus fuscoguttatus

The brown-marbled grouper or tiger grouper (*Epinephelus fuscoguttatus*) is a benthic marine fish which belongs to the family Serranidae or also known as the groupers.



Brownbanded bamboo shark

Chiloscyllium punctatum

The brownbanded bamboo shark (*Chiloscyllium punctatum*), is a bamboo shark in the family Hemiscylliidae that can be found in the Indo-West Pacific from Japan to northern Australia, between latitudes 34° N and 26° S. It is regularly bred in public...



Brownspeckled grouper

Epinephelus chlorostigma

The brownspeckled grouper (*Epinephelus chlorostigma*), also known as the brown speckled reef cod, brown-speckled rockcod, coral grouper or honeycomb cod, is a species of marine ray-finned fish, a grouper from the subfamily Epinephelinae which is part...



Brycinus longipinnis

Brycinus longipinnis

Primarily known as long-fin tetra, Brycinus longipinnis is also called the African long-finned tetra among other common names.



Bucktooth tetra

Exodon paradoxus

The bucktooth tetra (Exodon paradoxus) is the only member of the genus Exodon and is a freshwater fish of the characin family (family Characidae) of order Characiformes. It is native to the Amazon Basin and Guyana. Though first described in 1845,...



Buenos Aires tetra

Hyphessobrycon anisitsi

The Buenos Aires tetra (Psalidodon anisitsi) is a tropical fish from South America. It was first observed in the wild in 1907, by Carl H. Eigenmann.



Bufoceratias shaoi

Bufoceratias shaoi

Bufoceratias shaoi is a species of double angler, a type of anglerfish. The fish is bathypelagic and has been found at depths ranging from 0 to 1,200 metres (0 to 3,937 ft). It has been found in the western Indian Ocean and western Pacific Ocean....



Bufoceratias wedli

Bufoceratias wedli

Bufoceratias wedli is a deepsea anglerfish found in the mesopelagic to bathypelagic regions of the ocean, ranging from a depth of 300 to 1750 m. It is a double angler with two lures on its back, the anterior lure being the smaller.



Bulatmai barbel

Luciobarbus capito

The Bulatmai barbel (Luciobarbus capito) is a species of ray-finned fish in the genus Luciobarbus from the Aral and Caspian basins, including rivers that flow into these.



Bull ray

Aetomylaeus bovinus

Aetomylaeus bovinus, also known as the bull ray, duckbill ray, or duckbill eagle ray, is a species of large stingray of the family Myliobatidae found around the coasts of Europe and Africa.



Bull shark

Carcharhinus leucas

The bull shark (Carcharhinus leucas), also known as the Zambezi shark (informally zambi) in Africa and Lake Nicaragua shark in Nicaragua, is a species of requiem shark commonly found worldwide in warm, shallow waters along coasts and in rivers....



Bull trout

Salvelinus confluentus

The bull trout (*Salvelinus confluentus*) is a char of the family Salmonidae native to northwestern North America. Historically, *S. confluentus* has been known as the "Dolly Varden" (*S. malma*), but was reclassified as a separate species in 1980....



Bullet tuna

Auxis rochei

The bullet tuna, *Auxis rochei*, is a species of tuna, in the family Scombridae, found circumglobally in tropical oceans, including the Mediterranean Sea, in open surface waters to depths of 50 m (164 ft). The population of bullet tuna in the...



Bullhead minnow

Pimephales vigilax

The bullhead minnow (Pimephales vigilax) is a species of freshwater demersal fish, native to the Mississippi River system in the United States, to tributaries of Lake Huron and Lake Michigan and to northern Mexico.



Bullneck seahorse

Hippocampus minotaur

The bullneck seahorse (Hippocampus Minotaur) is a pygmy seahorse in the genus Hippocampus. This seahorse has never been found in the wild, and little is known about its natural habitat. The only known specimens were collected on the coast of...



Bullnose ray

Myliobatis freminvillii

The bullnose ray (Myliobatis freminvillei) is an eagle ray, which is widely distributed in the western Atlantic. It is found at depth above 10 m (33 ft) in coastal waters from Cape Cod down to Argentina, but is absent from parts of the western...



Bumblebee fish

Brachygobius xanthonus

The bumblebee fish (Brachygobius xanthonus) is a species of fresh and brackish water goby native to Thailand and Indonesia. This species can reach a length of 3.8 centimetres (1.5 in) SL and is found in lower parts of rivers, coastal areas,...



Burbot

Lota lota

The burbot (Lota lota), also known as bubbot, mariah, loche, cusk, freshwater cod, freshwater ling, freshwater cusk, the lawyer, coney-fish, lingcod, or eelpout, is a species of coldwater ray-finned fish native to the subarctic regions of the...



Burmese bamboo shark

Chiloscyllium burmensis

The Burmese bamboo shark, Chiloscyllium burmensis, is an extremely rare bamboo shark in the family Hemiscylliidae. The first specimen was caught 1963 off the coast from Rangoon in Myanmar (known as Burma at the time) in a depth of 29 – 33 m. This...



Burmese border loach

Botia kubotai

The Burmese Border loach, angelicus loach or polka dot loach, Botia kubotai, is a recently described species that has quickly become a popular tropical fish for freshwater aquariums. In 2002, fish collectors working in western Thailand began to...



Burmese trout

Raiamas guttatus

The Burmese trout (Raiamas guttatus) is a species of ray-finned fish in the genus Raiamas. They are found in the Irrawaddy, Mekong, Chao Phraya, Salween River basins and also in the northern Malay Peninsula.



Burnt-tailed barb

Balantiocheilos ambusticauda

The burnt-tailed barb (Balantiocheilos ambusticauda), also known as Siamese bala-shark, is a possibly extinct freshwater fish species from the family Cyprinidae. It is or was endemic to the Mae Klong and Chao Phraya River basins in Thailand.



Bushymouth catfish

Ancistrus dolichopterus

Ancistrus dolichopterus, one of the Ancistrus species called the bushymouth catfish, is a species of armored catfish native to Brazil. It is found in middle Amazon basin and in the basins of the Rio Negro, the lower Trombetas, the Tefé, the...



Butterfly kingfish

Gasterochisma melampus

The butterfly kingfish (Gasterochisma melampus) is an ocean-dwelling ray-finned bony fish in the mackerel family, Scombridae – a family which it shares with the tunas, mackerels, Spanish mackerels, and bonitos. Unlike the 50 species from those...



Butterfly perch

Caesioperca lepidoptera

The butterfly perch (Caesioperca lepidoptera) is a marine species of ray-finned fish in the subfamily Anthiinae, part of the sea bass family Serranidae. This fish is found in the eastern Indian Ocean, and the southwestern Pacific Ocean, including...



Butterfly splitfin

Ameca splendens

The butterfly splitfin or butterfly goodeid (Ameca splendens) is a bony fish from the monotypic genus Ameca of the splitfin family (Goodeidae). It was formerly found throughout the Ameca River drainage in Mexico; the type locality is Rio...



Cabezon

Scorpaenichthys marmoratus

The cabezon (Scorpaenichthys marmoratus) is a large species of sculpin native to the Pacific coast of North America. Although the genus name translates literally as "scorpion fish", true scorpionfish (such as lionfish) belong to the related...



Cactus roach

Rutilus virgo

The cactus roach (*Rutilus virgo*) is a species of fish in the family Cyprinidae, native to the basin of the Danube River upstream of the Iron Gate, very numerous in the Sava basin. It is also recorded in the Ukrainian Zakarpattian region. This...



Caecobarbus

Caecobarbus geertsi

Caecobarbus geertsi, the African blind barb or Congo blind barb (known as Nzonzi a mpofu in the local Kikongo language, meaning "blind barb"), is a species of cyprinid fish. This threatened cavefish is only known from Democratic Republic of the...



Caesio caerulaurea

Caesio caerulaurea

Caesio caerulaurea, the blue and gold fusilier, blue fusilier, gold-band fusilier or scissor-tailed fusilier, is a species of marine fish in the family Caesionidae. It is widespread throughout the tropical waters of the Indo-Pacific area,...



Caesio cuning

Caesio cuning

Caesio cuning, the redbelly yellowtail fusilier, yellowtail fusilier, red-bellied fusilier or robust fusilier, is a species of marine ray-finned fish, a fusilier belonging to the family Caesionidae. It is native to the Indian and Western Pacific...



Caesio teres

Caesio teres

Caesio teres, the yellow and blueback fusilier, beautiful fusilier, blue and gold fusilier or yellow-tail fusilier, is a species of marine, pelagic ray-finned fish belonging to the family Caesionidae. It occurs in the Indian and Western Pacific...



California corbina

Menticirrhus undulatus

The California corbina (*Menticirrhus undulatus*), or California kingcroaker, is marine demersal fish in the croaker family. It can often be found along sandy beaches and in shallow bays. This species travels in small groups along the surf zone in...



California flying fish

Cheilopogon pinnatibarbatus californicus

The California flying fish, Cheilopogon pinnatibarbatus californicus, is a subspecies of Bennett's flying fish. Prior to the 1970s, the California flying fish was known as a distinct species, with the scientific classification Cypselurus...



California halibut

Paralichthys californicus

The California halibut or California flounder (Paralichthys californicus) is a large-tooth flounder native to the waters of the Pacific Coast of North America from the Quillayute River in Washington to Magdalena Bay in Baja California. It feeds...



California killifish

Fundulus parvipinnis

The California killifish (Fundulus parvipinnis) is a type of killifish (Fundulidae) found along the coast of southern California and Baja California. Like the other members of the family, California killifish are small, no more than about 11...



California lizardfish

Synodus lucioceps

The California lizardfish (Synodus lucioceps) is a species of lizardfish primarily inhabiting the Californian coast. It has a long, brown body, which offers it camouflage in its habitat, the sandy bottom of the ocean. From its hiding spot on the...



California moray

Gymnothorax mordax

The California moray (Gymnothorax mordax) is a moray eel of the family Muraenidae, found in the eastern Pacific from just north of Santa Barbara to Santa Maria Bay in Baja California. They are the only species of moray eel found off California,...



California roach

Hesperoleucus symmetricus

The California roach (Hesperoleucus symmetricus) is a cyprinid fish native to western North America and abundant in the intermittent streams throughout central California. Once considered the sole member of its genus, it has recently been split...



California sheephead

Semicossyphus pulcher

The California sheephead (Semicossyphus pulcher) is a species of wrasse native to the eastern Pacific Ocean. Its range is from Monterey Bay, California, to the Gulf of California, Mexico. It can live for up to 20 years in favorable conditions and...



Californian anchovy

Engraulis mordax

The Californian anchovy or northern anchovy (*Engraulis mordax*) is a species of anchovy found in the Pacific Ocean, ranging from Mexico to British Columbia.



Callichthys callichthys

Callichthys callichthys

Callichthys callichthys, the cascarudo, armored catfish, bubblenest catfish, hassar, or mailed catfish, is a subtropical freshwater fish belonging to the subfamily Callichthyinae of the family Callichthyidae.



Callorhinchus callorhynchus

Callorhinchus callorhynchus

Callorhinchus callorhynchus, the American elephantfish or cockfish, is a species of fish in the family Callorhinchidae found in southern Brazil, Peru, Chile, Argentina, and Uruguay. Its natural habitat is open seas. Being a holocephalan from the...



Calophysus macropterus

Calophysus macropterus

Calophysus macropterus (also known as the Piracatinga, Vulture Catfish, or Zamurito) is a species of catfish (order Siluriformes) of the monotypic genus Calophysus of the family Pimelodidae. This fish has also been placed in its own family,...



Calotomus carolinus

Calotomus carolinus

Calotomus carolinus, commonly known as Carolines parrotfish, is a species of marine ray-finned fish, a parrotfish, in the family Scaridae. It is also known as the starry-eye parrotfish, stareye parrotfish, bucktooth parrotfish, Christmas...



Calotomus japonicus

Calotomus japonicus

Calotomus japonicus, commonly known as the Japanese parrotfish, is a species of parrotfish commonly found throughout the northwest Pacific. The species has been classified as Least Concern by the IUCN.



Cambodian logsucker

Garra cambodgiensis

The Cambodian logsucker (Garra cambodgiensis), also known as stonelapping minnow or false Siamese algae eater, is a species of ray-finned fish in the genus Garra. It lives in Southeast Asia.



Camouflage grouper

Epinephelus polyphekadion

The camouflage grouper (*Epinephelus polyphekadion*), also known as the blue-tailed cod, camouflage rockcod, small-toothed rockcod, smooth flowery rock-cod, snout-spot grouper or snout-spot rock-cod, is a species of marine ray-finned fish, a...



Canara pearlspot

Etroplus canarensis

The Canara pearlspot (*Etroplus canarensis*), also known as banded chromide or Canara pearlspot cichlid, is an endangered species of cichlid endemic to South Karnataka in India.



Canary damsel

Similiparma lurida

The Canary damsel (*Similiparma lurida*), or Cape Verde gregory (which is also a common name of *Stegastes imbricatus*) is a species of marine fish of the family Pomacentridae. It lives primary in shallow, subtropical waters of the eastern Atlantic...



Canary rockfish

Sebastes pinniger

The canary rockfish (*Sebastes pinniger*), also known as the orange rockfish, is a species of marine ray-finned fish belonging to the subfamily Sebastinae, the rockfishes, part of the family Scorpaenidae. It is native to the waters of the Pacific...



Candiru

Vandellia cirrhosa

Candiru (*Vandellia cirrhosa*), also known as cañero, toothpick fish, or vampire fish, is a species of parasitic freshwater catfish in the family Trichomycteridae native to the Amazon Basin where it is found in the countries of Bolivia, Brazil,...



Canterbury mudfish

Neochanna burrowsius

The Canterbury mudfish (Neochanna burrowsius), also known as the kowaro, is found only on the Canterbury Plains in New Zealand. Like other Neochanna species, it is a small, tubular and flexible fish which lacks scales. They are able to survive...



Cantherhines macrocerus

Cantherhines macrocerus

Cantherhines macrocerus, commonly known as the whitespotted filefish, American whitespotted filefish, or whitespotted limefish, is a marine fish found along the coast of Florida extending southward into the Caribbean. This species is distinct and...



Canthigaster jactator

Canthigaster jactator

Canthigaster jactator is a member of the pufferfish family that is endemic to the Hawaiian islands. It occasionally makes its way into the aquarium trade. It grows to a length of 9 centimeters (3.5 in).



Canthigaster rostrata

Canthigaster rostrata

Canthigaster rostrata, commonly known as the Caribbean sharp-nose puffer, is a pufferfish from the Western Central Atlantic. The Caribbean sharp-nose puffer is a small fish with a maximum length of 12 cm or approximately 4.7 inches. It can be...



Canthigaster solandri

Canthigaster solandri

Canthigaster solandri, commonly known as the spotted sharpnose puffer, is a ray-finned species of fish in the pufferfish family. It grows to a length of 11.5 centimeters (4.5 inches) in total length. It lives in the tropical Indo-Pacific: from...



Cape elephantfish

Callorhinchus capensis

The Cape elephantfish, josef, or St Joseph shark (Callorhinchus capensis) is a species of fish in the family Callorhinchidae.



Cape Fear shiner

Notropis mekistocholas

The Cape Fear shiner (Notropis mekistocholas) is a North American species of freshwater fish in the family Cyprinidae. It is endemic to the central part of the state of North Carolina in the southeastern United States, and is only found in the...



Cape galaxias

Galaxias zebratus

The Cape galaxias (Galaxias zebratus) is a species of freshwater fish of the family Galaxiidae. It is a small fish, rarely larger than 6 cm, that inhabits clear streams, rivers and ponds of South Africa. It shares the same habitat as imported...



Cape gurnard

Chelidonichthys capensis

The Cape gurnard (Chelidonichthys capensis) is a species of marine ray-finned fishes belonging to the family Triglidae, the gurnards and sea robins. This species is found in the southeastern Atlantic Ocean and the southwestern Indian Ocean. This...



Cape horse mackerel

Trachurus capensis

The Cape horse mackerel (Trachurus capensis) is a mackerel-like species in the family Carangidae. It is a pelagic species of the south eastern Atlantic Ocean which is a target of fisheries, mainly as bycatch.



Cape kurper

Sandelia capensis

The Cape kurper (Sandelia capensis) is a species of fish in the family Anabantidae, the climbing gouramis or climbing perches. It is endemic to South Africa.



Cape whitefish

Pseudobarbus capensis.[1]

The Cape whitefish or Berg-breede River whitefish (Cheilobarbus capensis) is a ray-finned fish species in the family Cyprinidae. It was formerly placed with the South African redfins in Pseudobarbus. It is tetraploid. Its closest living relative...



Capelin

Mallotus villosus

The capelin or caplin (Mallotus villosus) is a small forage fish of the smelt family found in the North Atlantic, North Pacific and Arctic oceans. In summer, it grazes on dense swarms of plankton at the edge of the ice shelf. Larger capelin also...



Capoeta damascina

Capoeta damascina

Capoeta damascina, the Levantine scraper or Mesopotamian barb, is a species of cyprinid fish from the Near East region. It is reported from Iraq, Israel, Jordan, Lebanon, Syria, Iran and Turkey. It is a bottom feeding fish, up to 50 cm (1.6 ft)...



Caranx lugubris

Caranx lugubris

Caranx lugubris, the black jack, black trevally, black kingfish, coal fish or black ulua, is a species of large ocean fish in the jack family Carangidae. The species has a circumtropical distribution, found in oceanic, offshore waters of the...



Carassioides acuminatus

Carassioides acuminatus

Carassioides acuminatus or black fish is a species of cyprinid fish in the subfamily Cyprininae. It is found in central and northern Vietnam and the Pearl River and Hainan in China. It may have been introduced into Laos. It occurs in slow flowing...



Carcharhinus humani

Carcharhinus humani

Carcharhinus humani, also known as the Human's whaler shark, is a species of requiem shark, in the family Carcharhinidae. It inhabits the western Indian Ocean near the Socotra Islands, off Kuwait, Mozambique, and South Africa.



Carcharhinus tjutjot

Carcharhinus tjutjot

The Indonesian whaler shark (Carcharhinus tjutjot), is a species of requiem shark belonging to the family Carcharhinidae. Until recently, it was thought to be a junior synonym of the whitecheek shark (C. dussumieri). The Indonesian whaler shark,...



Cardinal tetra

Paracheirodon axelrodi

The cardinal tetra (Paracheirodon axelrodi) is a freshwater fish of the family Characidae of order Characiformes. It is native to the upper Orinoco and Negro Rivers in South America. Growing to about 3 cm (1.2 in) total length, the cardinal tetra...



Caribbean reef shark

Carcharhinus perezii

The Caribbean reef shark (Carcharhinus perezii) is a species of requiem shark, belonging to the family Carcharhinidae. It is found in the tropical waters of the western Atlantic Ocean from Florida to Brazil, and is the most commonly encountered...



Caribbean roughshark

Oxynotus caribbaeus

Sharks portalThe Caribbean roughshark (Oxynotus caribbaeus) is a rough shark of the family Oxynotidae, found on the upper continental slopes of the Caribbean Sea, at depths between 400 and 450 m (1,310 and 1,480 ft). It reaches a length around...



Caribbean sharpnose shark

Rhizoprionodon porosus

The Caribbean sharpnose shark (*Rhizoprionodon porosus*) is a requiem shark, and part of the family Carcharhinidae.



Carinotetraodon imitator

Carinotetraodon imitator

Carinotetraodon imitator, commonly known as the dwarf Malabar pufferfish, is a freshwater pufferfish found in the Western Ghats of India, with little information available on its complete distribution. It is one of the smallest pufferfish in the...



Carinotetraodon irrubesco

Carinotetraodon irrubesco

Carinotetraodon irrubesco, known commonly as the red-tail dwarf puffer, is a freshwater pufferfish found only in the lower Banyuasin basin in South Sumatra and the Sambas River in West Kalimantan.



Carolina hammerhead

Sphyrna gilberti

The Carolina hammerhead (Sphyrna gilberti) is a species of hammerhead shark, and part of the family Sphyrnidae, found in the western Atlantic Ocean. Their pupping grounds are in nearshore waters off the southeastern U.S. with the highest...



Carolina pygmy sunfish

Elassoma boehlkei

The Carolina pygmy sunfish, *Elassoma boehlkei*, is a species of pygmy sunfish endemic to the United States, where it is only known from the Waccamaw and Santee River drainages in the Carolinas. This species prefers waters with dense vegetation. It...



Carpenter's chimaera

Chimaera lignaria

The carpenter's chimaera (Chimaera lignaria), also known as the giant chimaera or the giant purple chimaera, is a species of fish in the family Chimaeridae.



Carpenter's flasher wrasse

Paracheilinus carpenteri

The Carpenter's flasher wrasse, Paracheilinus carpenteri, is a species of wrasse native to the western Pacific Ocean. It can be found on reefs at depths from 27 to 45 m (89 to 148 ft). This species can reach 8 cm (3.1 in) in standard length. It...



Caspian kutum

Rutilus kutum

The Caspian kutum (Rutilus kutum) or Caspian white fish is a member of the family Cyprinidae from brackish water habitats of the Caspian Sea and from its freshwater tributaries. It is typically a medium-sized fish, reaching 45–55 cm in length,...



Caspian roach

Rutilus caspicus

The Caspian roach (Rutilus caspicus) is a species of roach fish living in the Caspian Sea. The Caspian roach can be distinguished from other roaches by its laterally compressed body, silvery grey iris, rounded snout and grey pectoral pelvic and...



Catla

Labeo catla

Catla (Labeo catla), (Bengali: কাতলা, romanized: kātālā) also known as the major South Asian carp, is an economically important South Asian freshwater fish in the carp family Cyprinidae. It is native to rivers and lakes in northern India,...



Catostomus discobolus

Catostomus discobolus

The bluehead sucker (Catostomus discobolus) is endemic to the intermountain Western US. A total of 23 members of the genus Catostomus has been found in North America. C. discobolus and C. yarrowi are two sister species that have very similar...



Catostomus latipinnis

Catostomus latipinnis

Catostomus latipinnis (flannemouth sucker) is a North American fish identified by its enlarged lower lips. It belongs to the genus Catostomus, commonly known as suckers. Historically, the flannemouth sucker ranged in the Colorado River Basin,...



Caulophryne jordani

Caulophryne jordani

Caulophryne jordani, commonly known as the fanfin angler, is a species of fanfin, a type of anglerfish. The fish is primarily found in the bathyal zone at depths ranging from 100 to 1,510 metres (330 to 4,950 ft). It has many unusual tentacles...



Caulophryne polynema

Caulophryne polynema

Caulophryne polynema is a species of fish in the family Caulophrynidae, the fanfins. It is known commonly as the hairy fanfin. It is native to the Atlantic and eastern Pacific Oceans. It occurs off the eastern coast of North America across to...



Celebes rainbowfish

Marosatherina ladigesii

The Celebes rainbowfish (Marosatherina ladigesii) is a species of sailfin silverside endemic to Sulawesi (formerly known as Celebes) in Indonesia. It is the only known member of its genus.



Central mudminnow

Umbra limi

The central mudminnow (Umbra limi) is a small fish in the family Umbridae of the order Esociformes. It is found in central and eastern North America in productive waters. It is fairly tolerant of low oxygen concentrations and, as a result, it is...



Central stoneroller

Campostoma anomalum

The central stoneroller, or Ohio stoneroller (Campostoma anomalum), is a fish in the family Cyprinidae endemic to North America.



Centrolophus

Centrolophus niger

The black ruff (*Centrolophus niger*) is a medusafish, the only member of the genus *Centrolophus*. It is a pelagic fish found in all tropical and temperate oceans at depths of 0 to 1,000 m (0 to 3,280 ft). Its length is typically up to 60 cm (24...



Centromochlus perugiaie

Centromochlus perugiaie

Centromochlus perugiaie is a species of benthopelagic fish, a member of the Auchenipteridae (driftwood catfish) family. They are widely known as honeycomb catfish or oil catfish because of their striking skin pattern. Honeycomb catfish are under 3...



Centropomus parallelus

Centropomus parallelus

Centropomus parallelus is a species of fish in the family Centropomidae, the snooks and robalos. It is known by several common names, including fat snook, smallscale fat snook, little snook, and chucumite. It is native to the western Atlantic...



Centropyge eibli

Centropyge eibli

Centropyge eibli, the blacktail angelfish, red stripe angelfish, orangelined angelfish, or Eibl dwarf angel is a species of marine ray-finned fish, a marine angelfish belonging to the family Pomacanthidae. It is found near reefs in the Indo-Pacific.



Centropyge fisheri

Centropyge fisheri

Centropyge fisheri, the orange angelfish, whitetail angelfish, damsel angelfish, yellowtail angelfish, Hawaiian flame angelfish, Fisher's angelfish, Fisher's dwarf angelfish or Fisher's pygmy angelfish, is a species of marine ray-finned fish, a...



Centropyge heraldi

Centropyge heraldi

Centropyge heraldi, the yellow angelfish or Herald's angelfish is a species of marine ray-finned fish, a marine angelfish belonging to the family Pomacanthidae. It comes from the Pacific Ocean and sometimes makes its way into the aquarium trade.



Centropyge potteri

Centropyge potteri

Centropyge potteri, commonly known as the russet angelfish, Potter's angelfish or Potter's pygmy angelfish, is a species of marine ray-finned fish, a marine angelfish belonging to the family Pomacanthidae. It is found in the central Pacific Ocean.



Centropyge tibicen

Centropyge tibicen

Centropyge tibicen, the keyhole angelfish, black angelfish, whitespot angelfish or puller angelfish, is a species of marine ray-finned fish, a marine angelfish belonging to the family Pomacanthidae. It is found in the Indo-Pacific region.



Cephalopholis argus

Cephalopholis argus

Cephalopholis argus, the peacock hind, roi, bluespotted grouper, and celestial grouper, is a species of marine ray-finned fish, a member of the subfamily Epinephelinae, the groupers, and part of the family Serranidae, which also includes the...



Cephalopholis boenak

Cephalopholis boenak

Cephalopholis boenak, the chocolate hind, brownbarred rockcod, brown-banded cod or brown-banded rockcod, is a species of marine ray-finned fish, a grouper from the subfamily Epinephelinae which is in the family Serranidae which also includes the...



Cephalopholis formosa

Cephalopholis formosa

Cephalopholis formosa, the Bluelined hind or bluelined rockcod is a species of marine ray-finned fish, a grouper from the subfamily Epinephelinae which is in the family Serranidae which also includes the anthias and sea basses. It is found in the...



Cephalopholis fulva

Cephalopholis fulva

Cephalopholis fulva, the coney or the butterfish, is a species of marine ray-finned fish, a grouper from the subfamily Epinephelinae which is in the family Serranidae which also includes the anthias and sea basses. It is found in the western...



Cephalopholis spiloparaea

Cephalopholis spiloparaea

Cephalopholis spiloparaea, known commonly as the strawberry hind strawberry rock cod, strawberry cod or orange rock cod, is a species of marine ray-finned fish, a grouper from the subfamily Epinephelinae which is in the family Serranidae which...



Cephalopholis taeniops

Cephalopholis taeniops

Cephalopholis taeniops, the African hind, bluespotted sea bass or spotted grouper, is a species of marine ray-finned fish, a grouper from the subfamily Epinephelinae which is in the family Serranidae which also includes the anthias and sea...



Cephalopholis urodeta

Cephalopholis urodeta

Cephalopholis urodeta, the darkfin hind, flagtail rockcod, chevron rockcod, duskyfin hind, duskyfin rockcod, flagtail grouper or V-tail grouper, is a species of marine ray-finned fish, a grouper from the subfamily Epinephelinae which is in the...



Cepola haastii

Cepola haastii

Cepola haastii is a species of marine ray-finned fish belonging to the family Cepolidae, the bandfishes. It is found on the inner continental shelf around New Zealand. Its length is between 15 and 25 cm. This species is known as the red bandfish,...



Cepola macrophthalma

Cepola macrophthalma

Cepola macrophthalma is a species of marine ray-finned fish belonging to the family Cepolidae, the bandfishes. It is found in the eastern Atlantic and Mediterranean from Senegal north to the British Isles. This species is known as the red...



Cero

Scomberomorus regalis

The cero (*Scomberomorus regalis*), also known as the pintado, kingfish, cero mackerel, cerite or painted mackerel, is a ray-finned bony fish in the family Scombridae, better known as the mackerel family. More specifically, this fish is a member of...



Cetoscarus bicolor

Cetoscarus bicolor

Cetoscarus bicolor, also known as the bicolour parrotfish or bumphead parrotfish, is a species of fish belonging to the family Scaridae. It is found only in the Red Sea.



Cetoscarus ocellatus

Cetoscarus ocellatus

Cetoscarus ocellatus, common name Spotted parrotfish, is a species of parrotfish belonging to the family Scaridae.



Ceylon killifish

Aplocheilus dayi

The Ceylon killifish (Sinhala: උඩ හඳයා, romanized: uda handaya; Aplocheilus dayi) is a species of killifish endemic to Sri Lanka. This species grows to a length of 9 cm (3.5 in). Males and females have a black dot at the rear end of the base of...



Ceylon snakehead

Channa orientalis

The Ceylon snakehead (Channa orientalis) is a species of snakehead found in freshwater habitats, typically shaded streams, in southwestern Sri Lanka (although occasionally claimed to occur in other countries, this is misidentifications of...



Chaca bankanensis

Chaca bankanensis

Chaca bankanensis is a species of angler catfish found in the Sundaland region, where it occurs in Brunei, Indonesia, Malaysia, and Thailand. It is found in peat swamps and streams. This species grows to a length of 20.0 cm (7.9 in). It is found...



Chaca chaca

Chaca chaca

Chaca chaca is a species of angler catfish found in the Ganges-Brahmaputra River system of India and the Ayeyarwady River of Myanmar, where it is found in rivers, canals and ponds of grassland, scrubland, deciduous forest and rainforest habitats....



Chaetodermis

Chaetodermis penicilligerus

Chaetodermis penicilligerus, also known as the prickly leather-jacket or tasselled leather-jacket, is a demersal marine fish belonging to the family Monacanthidae. This fish is in the monotypic genus Chaetodermis. It occasionally makes its way...



Chaetodon daedalma

Chaetodon daedalma

Chaetodon daedalma or the wrought-iron butterflyfish is a species of marine ray-finned fish, a butterflyfish belonging to the family Chaetodontidae. It is native to the western Pacific ocean near central and southern Japan.



Chaetodon falcula

Chaetodon falcula

Chaetodon falcula, the blackwedged butterflyfish or falcula butterflyfish, is a species of marine ray-finned fish a butterflyfish belonging to the family Chaetodontidae. It is found in the Indian Ocean.



Chaetodon larvatus

Chaetodon larvatus

Chaetodon larvatus, commonly known as the hooded butterflyfish or orangeface butterflyfish (and with many other common names), is a species of marine ray-finned fish, a butterflyfish belonging to the family Chaetodontidae. It is found in the Red...



Chaetodon melapterus

Chaetodon melapterus

Chaetodon melapterus, the Arabian butterflyfish, blackfin butterflyfish, or black-finned melon butterflyfish, is a species of marine ray-finned fish, a butterflyfish belonging to the family Chaetodontidae. It is found in the north western Indian...



Chaetodon plebeius

Chaetodon plebeius

Chaetodon plebeius, the blueblotch butterflyfish, bluespot butterflyfish, bluedash butterflyfish or grey-blotched butterflyfish, is a species of marine ray-finned fish, a butterflyfish belonging to the family Chaetodontidae. It is native to the...



Chaetodon ulietensis

Chaetodon ulietensis

Chaetodon ulietensis, the Pacific double-saddle butterflyfish or false falcula butterflyfish, is a species of butterflyfish (family Chaetodontidae). It flourishes in coral-rich environments in the central Indo-Pacific region. Their range extends...



Chaetodontoplus duboulayi

Chaetodontoplus duboulayi

Chaetodontoplus duboulayi, the scribbled angelfish, is a species of marine ray-finned fish, a marine angelfish belonging to the family Pomacanthidae. This species is from the southwestern Pacific Ocean.



Chain catshark

Scyliorhinus retifer

The chain catshark or chain dogfish (Scyliorhinus retifer) is a small, reticulated catshark that is biofluorescent. The species is common in the Northwest Atlantic, Gulf of Mexico, and Caribbean. It is harmless and rarely encountered by humans....



Chain moray

Echidna catenata

Echidna catenata, commonly known as the chain moray, is a moray eel found in shallow parts of the western Atlantic Ocean and from islands elsewhere in the Atlantic. It occasionally makes its way into the aquarium trade. It grows to a maximum...



Chain pickerel

Esox niger

The chain pickerel (Esox niger) is a species of freshwater fish in the pike family (family Esocidae) of order Esociformes. The chain pickerel and the American pickerel (E. americanus) belong to the Esox genus of pike.



Chameleon goby

Tridentiger trigonocephalus

The chameleon goby (Tridentiger trigonocephalus) is a species of ray-finned fish native to marine and brackish waters along the coasts of eastern Asia. It has also spread to other parts of the world where it is found in waters with varying...



Champscephalus esox

Champscephalus esox

Champscephalus esox, the pike icefish or northern icefish, is a species of marine ray-finned fish belonging to the family Channichthyidae, the crocodile icefishes.



Channa bleheri

Channa bleheri

Channa bleheri (the rainbow snakehead) is a species of dwarf snakehead that is endemic to the Brahmaputra River basin in the Indian states of Assam and Arunachal Pradesh. It is among the most colorful species of snakehead.



Channa burmanica

Channa burmanica

Channa burmanica is a species of snakehead which is endemic to northern Burma. It is a very small species of snakehead included in the informal group of dwarf snakeheads. Its colouring closely resembles that of Channa gachua and Channa bleheri....



Channa gachua

Channa gachua

Channa gachua, the dwarf snakehead, is a species of fish in the family Channidae. The name "dwarf snakehead" is also used for several other species of small snakeheads. C. gachua is native to freshwater habitats in southern Asia, where it has a...



Channa harcourtbutleri

Channa harcourtbutleri

Channa harcourtbutleri, the Burmese snakehead, is a species of snakehead endemic to Inle Lake and surroundings in Myanmar. Locally called nga ohn-ma, among aquarists it is considered one of the dwarf snakeheads, but no significant import for...



Channa marulius

Channa marulius

Channa marulius (bullseye snakehead or great snakehead) is a large species of snakehead native to South Asia. Populations in Southeast Asia are now regarded as separate species.



Channa micropeltes

Channa micropeltes

Channa micropeltes, giant snakehead, giant mudfish or toman harimau, is among the largest species in the family Channidae, capable of growing to 1.5 m (4.9 ft) in length and a weight of 20 kg (44 lb). It is native to the fresh waters of Southeast...



Channa pardalis

Channa pardalis

Channa pardalis is a species of dwarf snakehead in the genus Channa. It was first described in 2016 from Khasi Hills, Meghalaya of northeastern India. Prior to its scientific description, it was known as Channa sp. "True Blue" or Channa sp....



Channa pleurophthalma

Channa pleurophthalma

Channa pleurophthalma, the ocellated snakehead is a species of Southeast Asian freshwater fish in the snakehead family.



Channa pulchra

Channa pulchra

Channa pulchra is a species of snakehead fish in the family Channidae which is native to Myanmar. It was first described in 2007 by R. Britz from a specimen collected from the Kyeintali Chaung (stream) basin in Rakhine Yoma, western Myanmar. The...



Channa punctata

Channa punctata

Channa punctata, the spotted snakehead, is a species of snakehead. It is found in the Indian Subcontinent and nearby areas, ranging across Afghanistan, Pakistan, India, Sri Lanka, Nepal, Bangladesh, Myanmar and Tibet. Its natural habitats are...



Channa stewartii

Channa stewartii

Channa stewartii is a species of dwarf snakehead in the family Channidae, which is native to Nepal and the Indian states of Arunachal Pradesh, Assam, Manipur, Meghalaya, Nagaland and Tripura. It may also be found in Bangladesh. This freshwater...



Channa striata

Channa striata

Channa striata, the striped snakehead, is a species of snakehead fish. It is also known as the common snakehead, chevron snakehead, or snakehead murrel and generally referred simply as mudfish. It is native to South and Southeast Asia, and has...



Channel catfish

Ictalurus punctatus

The channel catfish (*Ictalurus punctatus*) is North America's most numerous catfish species. It is the official fish of Kansas, Missouri, Nebraska, and Tennessee, and is informally referred to as a "channel cat". In the United States, they are the...



Chauliodus danae

Chauliodus danae

Chauliodus danae, or dana viperfish, is a species of viperfish in the family Stomiidae. The Dana Viperfish is mostly found in the bathyal zone. The species was first caught and recognised on the Dana expeditions 1920-1922 and named after the...



Cheat minnow

Pararhinichthys bowersi

The cheat minnow (Pararhinichthys bowersi) is a demersal, freshwater fish endemic to the United States, most commonly found in the Ohio River basin. It is the only species in the genus Pararhinichthys. Its taxonomic status is debatable and...



Checker barb

Oliotius oligolepis

The checker barb (Oliotius oligolepis) is a species of cyprinid fish endemic to creeks, rivers, and lakes in Sumatra, Indonesia. It has also been established in the wild in Colombia. The adult males have red fins with black tips. It will grow up...



Checkerboard wrasse

Halichoeres hortulanus

The checkerboard wrasse (Halichoeres hortulanus) is a fish belonging to the wrasse family. It is native to the area including the Indian Ocean to central Pacific Ocean.



Checkered puffer

Sphoeroides testudineus

The checkered puffer (Sphoeroides testudineus) is a species in the family Tetraodontidae, or pufferfishes. Normal sizes: 4 to 7 in. (10 to 18 cm) Travel / Living Depth: 3– 35 ft. (1-12m). Location and Distribution: Caribbean, The Bahamas, Florida,...



Cheilopogon nigricans

Cheilopogon nigricans

Cheilopogon nigricans, the blacksail flyingfish, also known as African flyingfish, or leaping flyingfish, is a flying fish in the family Exocoetidae. It is an oceanodromous, plankton-eating marine fish which has commercial value.



Chelmon marginalis

Chelmon marginalis

Chelmon marginalis, the margined coralfish or Western beaked butterflyfish, is a species of marine ray-finned fish, a butterflyfish in the family Chaetodontidae. It is a reef fish which is endemic to Australia.



Cherokee darter

Etheostoma scotti

The Cherokee darter (Etheostoma scotti) is a rare species of freshwater ray-finned fish, a darter from the subfamily Etheostomatinae, part of the family Percidae, which also contains the perches, ruffes and pikeperches. It is endemic to Georgia...



Cherry barb

Puntius titteya

The cherry barb (Puntius titteya) is a tropical freshwater fish belonging to the family Cyprinidae. It is endemic to Sri Lanka, and introduced populations have become established in Mexico and Colombia. The cherry barb was named Puntius titteya...



Cherubfish

Centropyge argi

The cherubfish (Centropyge argi), also known as the pygmy angelfish, is a species of marine ray-finned fish, a marine angelfish belonging to the family Pomacanthidae. It is found in the western Atlantic Ocean.



Chevron butterflyfish

Chaetodon trifascialis

The chevron butterflyfish (Chaetodon trifascialis), also known as chevroned butterflyfish, triangulate butterflyfish or V-lined butterflyfish, is a species of marine ray-finned fish, a butterflyfish belonging to the family Chaetodontidae. It has...



Chilean devil ray

Mobula tarapacana

The Chilean devil ray (*Mobula tarapacana*), also known as the box ray, greater Guinean mobula, sicklefin devil ray or the spiny mobula, is a species of ray in the family Mobulidae. It is often observed worldwide, basking just below the surface in...



Chilean jack mackerel

Trachurus murphyi

The Chilean jack mackerel (Trachurus murphyi), sometimes called the Inca scad or Peruvian jack mackerel, is a species of jack mackerel in the genus Trachurus of the family Carangidae. Since the 1970s, it has become one of the world's more...



Chilodus punctatus

Chilodus punctatus

Chilodus punctatus, known as the spotted headstander or pearl headstander, is a headstander, a type of fish, in the genus Chilodus. The spotted headstander has a body that extends grayish and green colors over his body and sets off the colors...



Chilomycterus antillarum

Chilomycterus antillarum

Chilomycterus antillarum, the web burrfish, spiny box puffer, bridled burrfish or striped burrfish, is a species of fish in the family Diodontidae native to the Western Atlantic Ocean, southern Florida, and the Bahamas to Brazil. The species can...



Chilomycterus reticulatus

Chilomycterus reticulatus

Chilomycterus reticulatus, the spotfin burrfish, spotted burrfish, Pacific burrfish, spotfin porcupinefish or few-spined porcupinefish, is a species of fish in the genus Chilomycterus, which is part of the porcupinefish family Diodontidae.



Chilotilapia

Chilotilapia rhoadesii

Chilotilapia is a genus of freshwater fish in the cichlid family. It contains the sole species Chilotilapia rhoadesii, the Malawi bream, which is endemic to Lake Malawi in East Africa, where it prefers areas with muddy substrates from the...



Chimaera bahamaensis

Chimaera bahamaensis

Chimaera bahamaensis, commonly known as the Bahamas ghost shark, is a species of fish in the family Chimaeridae. It is found in North Atlantic Ocean around the Bahamas, specifically it has been found east of Andros Island. Chimaera bahamaensis is...



China rockfish

Sebastes nebulosus

The China rockfish (Sebastes nebulosus), the yellowstripe rockfish or yellowspotted rockfish, is a species of marine ray-finned fish belonging to the subfamily Sebastinae, the rockfishes, part of the family Scorpaenidae. It is native to the...



Chinamanfish

Symphorus nematophorus

The Chinamanfish (Symphorus nematophorus), Chinaman snapper, galloper or thread-finned sea perch, is species of marine ray-finned fish, a snapper belonging to the family Lutjanidae. It is found in the Western Pacific Ocean.



Chinese bahaba

Bahaba taipingensis

The Chinese bahaba (Bahaba taipingensis), also known as the giant yellow croaker, is a critically endangered species of marine and brackish water fish in the family Sciaenidae. It is a large fish, reaching lengths up to 2 m (6 ft 7 in) and...



Chinese high-fin banded shark

Myxocyprinus asiaticus

The Chinese high-fin banded shark (Myxocyprinus asiaticus) is a popular freshwater aquarium fish that belongs to the family Catostomidae. It grows to about 1.35 m (4 ft 5 in) long and is unsuitable for most home aquariums. In addition to Chinese...



Chinese mackerel

Scomberomorus sinensis

The Chinese mackerel (Scomberomorus sinensis), also known as the Chinese seerfish, is a ray-finned bony fish in the family Scombridae, better known as the mackerel family. More specifically, this fish is a member of the tribe Scomberomorini, the...



Chinese rice fish

Oryzias sinensis

The Chinese rice fish (Oryzias sinensis) is a species of fish in the genus Oryzias. This freshwater fish occurs in swamps, stagnant parts of streams, rice fields and marshes, and is up to 3.1 cm (1.2 in) long. It was formerly considered a...



Chinese sleeper

Perccottus glenii

The Chinese sleeper (Perccottus glenii), also known as the Amur sleeper, is a species of freshwater sleeper native to the Amur River basin in eastern Asia with introduced populations in other regions of Eurasia. It is currently the only known...



Chinese sturgeon

Acipenser sinensis

The Chinese sturgeon (Acipenser sinensis; Chinese: 中華鰻; pinyin: zhōnghuá xún) is a critically endangered member of the family Acipenseridae in the order Acipenseriformes. Historically, this anadromous fish was found in China, Japan, and the...



Chinese trumpetfish

Aulostomus chinensis

The Chinese trumpetfish, Aulostomus chinensis, is a demersal marine fish belonging to the family Aulostomidae.



Chinook salmon

Oncorhynchus tshawytscha

The Chinook salmon (*Oncorhynchus tshawytscha*) is the largest and most valuable species of Pacific salmon. Its common name is derived from the Chinookan peoples. Other vernacular names for the species include king salmon, Quinnot salmon, Tsumen,...



Chionobathyscus

Chionobathyscus dewitti

Chionobathyscus dewitti is a species of marine ray-finned fish belonging to the family Channichthyidae, the crocodile icefishes. This species is found in the Southern Ocean.



Chionodraco hamatus

Chionodraco hamatus

Chionodraco hamatus is a species of marine ray-finned fish belonging to the family Channichthyidae, the crocodile icefishes. It is found in the Southern Ocean.



Chionodraco rastrospinosus

Chionodraco rastrospinosus

The ocellated icefish (Chionodraco rastrospinosus) is a fish of the family Channichthyidae. It lives in the cold waters off Antarctica and is known for having transparent haemoglobin-free blood.C. rastrospinosus live in the Southern Ocean up to a...



Chirostoma bartoni

Chirostoma bartoni

Chirostoma bartoni, the Alberca silverside, is a species of neotropical silverside endemic to the Lerma River basin of Mexico. Typical adult specimens are approximately 7.1 cm in length. The Alberca silverside was found only in the Alberca...



Chirostomias

Chirostomias pliopterus

Chirostomias pliopterus is a species of barbeled dragonfish found in the Atlantic Ocean. This species grows to a length of 20.5 centimetres (8.1 in) SL. This species is the only described member of its genus. This scaleless dragonfish was also...



Chiselmouth

Acrocheilus alutaceus

The chiselmouth (Acrocheilus alutaceus) is an unusual cyprinid fish of western North America. It is named for the sharp hard plate on its lower jaw, which is used to scrape rocks for algae. It is the sole member of the monotypic genus Acrocheilus...



Chitala chitala

Chitala chitala

Chitala chitala (Assamese: চিতল sitawl, Bengali: চিতল, chitol) is a knifefish from Bangladesh, India, Nepal, and Pakistan, found in the Brahmaputra, Indus, Ganges and Mahanadi River basins. It is sometimes known as the Indian featherback or...



Chitala hypselonotus

Chitala hypselonotus

Chitala hypselonotus is a species of knifefish found in freshwater habitats in Borneo and Sumatra in southeast Asia.



Chlorurus microrhinos

Chlorurus microrhinos

Chlorurus microrhinos, the blunt-head parrotfish or steephead parrotfish, is a species of marine ray-finned fish, a parrotfish from the family Scaridae. It is found in the Indo-Pacific region.



Chlorurus sordidus

Chlorurus sordidus

Chlorurus sordidus, known commonly as the daisy parrotfish or bullethead parrotfish, is a species of marine fish in the family Scaridae.



Chlorurus spilurus

Chlorurus spilurus

Chlorurus spilurus, known commonly as the Pacific daisy parrotfish or Pacific bullethead parrotfish and in Hawaiian called uhu, is a species of marine fish in the family Scaridae. The Pacific daisy parrotfish is widespread throughout the...



Chocolate gourami

Sphaerichthys osphromenoides

The chocolate gourami (Sphaerichthys osphromenoides) is a species of gourami native to the Malay Peninsula, Sumatra, and Borneo. Chocolate gouramis reach a length of 6 cm (2.4 in). These fish have a chocolate colour with golden bands running down...



Chologaster cornuta

Chologaster cornuta

Chologaster cornuta, commonly named swampfish, ricefish, or riceditch killifish, is a freshwater fish of the family Amblyopsidae. It is the only living species of the genus Chologaster. It only lives in US rivers in the Atlantic Coastal Plain...



Christmas wrasse

Thalassoma trilobatum

The Christmas wrasse (Thalassoma trilobatum), also known as the ladder wrasse, green-barred wrasse or green-blocked wrasse, is a species of ray-finned fish, a wrasse from the family Labridae which is native to the Indian Ocean and the western...



Chromis atripectoralis

Chromis atripectoralis

The black-axil chromis (Chromis atripectoralis), also known as the blackfin chromis or blue-green puller, is a damselfish from the tropical Indo-Pacific. This fish can reach almost 12 centimetres (4.7 in) in length. It inhabits lagoons and reefs...



Chromis chromis

Chromis chromis

Chromis chromis, the damselfish or Mediterranean chromis, is a small species of ray-finned fish of the family Pomacentridae from the Eastern Atlantic and Mediterranean.



Chromis cyanea

Chromis cyanea

Chromis cyanea (blue chromis) is a damselfish in the subfamily Pomacentrinae, found in Bermuda, southern Florida and the Caribbean Sea. It is collected for the aquarium trade.



Chromis notata

Chromis notata

The pearl-spot chromis or spottedfin puller (Chromis notata) is a damselfish of the family Pomacentridae, found in the northwest Pacific from southern Korea, the coast of Jeju Island, southern Japan, the Ryukyu Islands, Taiwan, and China, at...



Chromis punctipinnis

Chromis punctipinnis

The blacksmith (Chromis punctipinnis), also known as the blacksmith chromis and blacksmith damselfish, is a species of fish in the damselfish family. It is native to the subtropical northeastern Pacific Ocean, where its range is from Monterey Bay,...



Chromis viridis

Chromis viridis

Chromis viridis (green chromis) is a species of damselfish. Individuals tend to be iridescent apple-green and light blue, and reach a maximal length of 10 cm. It is sometimes called the "blue-green chromis", but that may also refer to Chromis...



Chrysiptera cyanea

Chrysiptera cyanea

Chrysiptera cyanea is a species of damselfish found in the wide Indo-West Pacific but not known in the Red Sea. A few individuals were observed in the Mediterranean Sea in 2013 off Slovenia, a likely aquarium release. Common names include blue...



Chrysiptera hemicyanea

Chrysiptera hemicyanea

Chrysiptera hemicyanea, known commonly as the azure damselfish, azure demoiselle, half-blue demoiselle, and yellow-dipped damsel, is a species of damselfish.



Chrysiptera parasema

Chrysiptera parasema

Chrysiptera parasema, also known as yellowtail damselfish, yellowtail blue damsel, goldtail demoiselle and other variations, is a saltwater species of fish from the Indo-Pacific. It was described by Henry Weed Fowler in 1918.



Chrysiptera springeri

Chrysiptera springeri

Chrysiptera springeri, commonly known as Springer's demoiselle, is a species of damselfish in the family Pomacentridae.



Chrysiptera talboti

Chrysiptera talboti

Chrysiptera talboti, known commonly as Talbot's damselfish and Talbot's demoiselle, is a species of damselfish. It is a marine fish from the eastern Indian Ocean and western Pacific.



Chrysiptera taupou

Chrysiptera taupou

Chrysiptera taupou, known commonly as the southseas devil, southseas demoiselle, and Fiji damsel, is a species of damselfish. It is native to the western Pacific Ocean from the Coral Sea to Samoa.



Chub mackerel

Scomber japonicus

The chub mackerel, Pacific mackerel, or Pacific chub mackerel (Scomber japonicus) is a species of fish in the tuna and mackerel family, Scombridae. This species of mackerel closely resembles the Atlantic chub mackerel.



Chum salmon

Oncorhynchus keta

The chum salmon (Oncorhynchus keta), also known as dog salmon or keta salmon, is a species of anadromous salmonid fish from the genus Oncorhynchus (Pacific salmon) native to the coastal rivers of the North Pacific and the Beringian Arctic, and is...



Cichla intermedia

Cichla intermedia

Cichla intermedia, the royal peacock bass, is a large species of cichlid found in the Orinoco River basin in Venezuela and Colombia.



Cichla monoculus

Cichla monoculus

Cichla monoculus, sometimes known as the tucanare peacock bass ("peacock bass" is also used for some of its relatives) or toukounaré, is a very large species of cichlid, and a prized game fish. It is native to the Amazon basin in South America,...



Cichla ocellaris

Cichla ocellaris

Cichla ocellaris, sometimes known as the butterfly peacock bass ("peacock bass" is also used for some of its relatives), is a very large species of cichlid from South America, and a prized game fish. It reaches 74 cm (29 in) in length. It is...



Cichla orinocensis

Cichla orinocensis

Cichla orinocensis, sometimes known as the Orinoco peacock bass, tucunaré or tucunaré miri, is a very large species of cichlid. This peacock bass is native to the Rio Negro and Orinoco River basins in northern South America. It mainly occurs in...



Cichla temensis

Cichla temensis

Cichla temensis, the speckled pavon, speckled peacock bass, painted pavon, or three-barred peacock bass, is a very large South American cichlid, and a prized food and game fish. Reaching up to 1 m (3.3 ft) in length and 13 kg (29 lb) in weight....



Cichlasoma bimaculatum

Cichlasoma bimaculatum

The black acara (Cichlasoma bimaculatum) is an omnivorous, freshwater, tropical fish. The species originates from South America, specifically in the Caroni River in the Orinoco basin in Venezuela, Essequibo River in Guyana, and the Rio Branco...



Cigar wrasse

Cheilio inermis

The cigar wrasse, Cheilio inermis, is a species of wrasse native to the Indo-Pacific. It is mainly found on tropical reefs at depths to 30 m (98 ft) in the Indo-Pacific region, Red Sea included. They inhabit seagrass beds and algae-covered flats,...



Cinnamon clownfish

Amphiprion melanopus

Amphiprion melanopus, also known as the cinnamon clownfish, fire clownfish, red and black anemonefish, black-backed anemonefish or dusky anemonefish is a widely distributed anemonefish chiefly found in the western and southern parts of the...



Cirrhinus molitorella

Cirrhinus molitorella

Cirrhinus molitorella (mud carp or dace) is a species of ray-finned fish in the genus Cirrhinus found mainly in southern China and Vietnam.



Cirrhinus mrigala

Cirrhinus mrigala

Cirrhinus mrigala is a species of ray-finned fish in the genus Cirrhinus. It is found in northern India, Pakistan, Bangladesh and Nepal. This species and the Mrigal carp (Cirrhinus cirrhosus) are both considered distinct.



Citharichthys stigmaeus

Citharichthys stigmaeus

Citharichthys stigmaeus, the speckled sanddab, is a species of flounder in the genus Citharichthys. It is native to the eastern Pacific Ocean, ranging from Alaska in the north to Baja California, Mexico in the south. It is a much smaller cousin of...



Clanwilliam yellowfish

Labeobarbus seeberi

The Clanwilliam yellowfish (Labeobarbus seeberi) is a ray-finned fish species in the family Cyprinidae. It has long been placed in Barbus, the "wastebin genus" for barbs, by default; however, the species is increasingly being restored to related...



Clarias anguillaris

Clarias anguillaris

Clarias anguillaris is a species of African airbreathing catfish also known as the mudfish. This species is of minor importance in commercial fisheries. It grows to a length of 100 cm (39.4 inches) TL.



Clarias batu

Clarias batu

Clarias batu (sometimes known under the common name Tioman walking catfish) is a species of clariid catfish. It is only known from Tioman Island off the eastern coast of Peninsular Malaysia. The eel-like catfish found on this island were...



Clarias brachysoma

Clarias brachysoma

Clarias brachysoma (walking catfish) is a species of fish in the family Clariidae. It is endemic to Sri Lanka. Its natural habitats are rain forest streams and lowland rivers. It is threatened by habitat loss. It grows to a length of 40 cm. During...



Clarias cavernicola

Clarias cavernicola

The golden cave catfish (Clarias cavernicola) is a critically endangered species of airbreathing catfish. This cavefish is only known to live in the Aigamas cave, Otjozondjupa region, Namibia. It has also been reported from the nearby Dragon's...



Clarias gariepinus

Clarias gariepinus

Clarias gariepinus or African sharptooth catfish is a species of catfish of the family Clariidae, the airbreathing catfishes.



Clarias nieuhofii

Clarias nieuhofii

Clarias nieuhofii, the slender walking catfish, is a species of clariid catfish. It has a wide distribution in Southeast Asia including southern Vietnam, Cambodia, Thailand, Malay Peninsula, parts of the Greater Sunda Islands (Indonesia and...



Clarion angelfish

Holacanthus clarionensis

The Clarion angelfish (Holacanthus clarionensis) is a species of marine ray-finned fish, a marine angelfish belonging to the family Pomacanthidae. It is found in the eastern Pacific Ocean, almost exclusively near islands off the Pacific coast of...



Clark's anemonefish

Amphiprion clarkii

Clark's anemonefish (*Amphiprion clarkii*), also known as the yellowtail clownfish, is a marine fish belonging to the family Pomacentridae, the clownfishes and damsselfishes.



Clearfin lionfish

Pterois radiata

The clearfin lionfish (*Pterois radiata*), also called the tailbar lionfish, radiata lionfish, fireworks fish or radial firefish, is a carnivorous, ray-finned fish with venomous spines belonging to the family Scorpaenidae, the scorpionfishes and...



Clearnose skate

Raja eglanteria

The clearnose skate (*Rostroraja eglanteria*) is a species of cartilaginous fish in the family Rajidae. *R. eglanteria* is also known by other common names such as the brier skate and summer skate. Clearnose skates are easily identified by the...



Cleftbelly trevally

Atropus atropus

The cleftbelly trevally (*Atropus atropus*), also known as the cleftbelly kingfish, Kuweh trevally or thin crevalle, is a species of tropical marine fish of the jack family, Carangidae. The species inhabits coastal waters throughout the Indo-West...



Cleisthenes pinetorum

Cleisthenes pinetorum

Cleisthenes pinetorum is a flatfish of the family Pleuronectidae. It is a demersal fish that lives on sublittoral sand and mud bottoms at depths of between 50 and 200 metres (160 and 660 ft). Its native habitat is the temperate waters of the...



Climbing galaxias

Galaxias brevipinnis

The climbing galaxias or kōaro (Galaxias brevipinnis) is a fish of the family Galaxiidae found in Australia, New Zealand, and nearby islands. The name climbing galaxias is used in Australia, and koaro or kōaro in New Zealand. Further vernacular...



Clipper barb

Enteromius callipterus

The clipper barb (Enteromius callipterus) or Congo barb is a freshwater and brackish tropical fish belonging to the minnow family (Cyprinidae). Its native habitat is from Côte d'Ivoire through the Chad Basin to Nigeria and Cameroon. It was...



Cloudy catshark

Scyliorhinus torazame

The cloudy catshark (Scyliorhinus torazame) is a common species of catshark, belonging to the family Scyliorhinidae. It is a bottom-dweller that inhabits rocky reefs in the northwestern Pacific Ocean, from the shore to a depth of 320 m (1,050...



Clown barb

Barbodes everetti

Barbodes everetti, the Clown barb or Everett's barb is a species of cyprinid fish native to Borneo and Sumatra. It inhabits clear streams in forested foothills and can also be found in forest floor puddles as shallow as 15 centimetres (5.9 in) or...



Clown coris

Coris aygula

The clown coris (Coris aygula), also known as the clown wrasse, false clownwrasse, humphead wrasse, hump-headed wrasse, red-blotched rainbowfish or twinspace wrasse, is a species of wrasse native to the Indian Ocean and the western Pacific Ocean.



Clown featherback

Chitala ornata

The clown featherback (Chitala ornata), also known as the clown knifefish and spotted knifefish, is a nocturnal species of tropical fish with a long, knife-like body. This knifefish is native to freshwater habitats in Cambodia, Laos, Thailand,...



Clown killi

Epiplatys annulatus

The clown killi (Epiplatys annulatus) or banded panchax, is a species of fish in the family Nothobranchiidae, an African rivuline, native to fresh water habitats in Guinea, Liberia and Sierra Leone in West Africa. This species was described as...



Clown loach

Chromobotia macracanthus

The clown loach (Chromobotia macracanthus), or tiger botia, is a tropical freshwater fish belonging to the botiid loach family. It is the sole member of the genus Chromobotia. It originates in inland waters in Indonesia on the islands of Sumatra...



Clown triggerfish

Balistoides conspicillum

The clown triggerfish (Balistoides conspicillum), also known as the bigspotted triggerfish, is a demersal marine fish belonging to the family Balistidae, or commonly called triggerfish.



Cnidoglanis macrocephalus

Cnidoglanis macrocephalus

Cnidoglanis macrocephalus is a species of catfish (order Siluriformes) of the family Plotosidae, and is the only species of the genus Cnidoglanis. It is commonly known as the cobbler, estuary cobbler, deteira, estuary catfish, South Australian...



Coastal cutthroat trout

Oncorhynchus clarkii clarkii

The coastal cutthroat trout (Oncorhynchus clarkii clarkii), also known as the sea-run cutthroat trout, blue-back trout or harvest trout, is one of the several subspecies of cutthroat trout found in Western North America. The coastal cutthroat...



Coastal trevally

Carangoides coeruleopinnatus

The coastal trevally (Carangoides coeruleopinnatus), also known as the onion trevally, Japanese trevally or bluefin kingfish, is a species of inshore marine fish in the jack family Carangidae. The species is distributed throughout the tropical...



Coastrange sculpin

Cottus aleuticus

The Coastrange sculpin (Cottus aleuticus) is a freshwater sculpin of the genus Cottus. They are commonly found near the ocean in western North America, namely Canada and the United States. It is also known as the Aleutian sculpin.



Cobia

Rachycentron canadum

The cobia (Rachycentron canadum) (, KOH-bee-ə) is a species of carangiform marine fish, the only extant representative of the genus Rachycentron and the family Rachycentridae. Its other common names include black kingfish, black salmon, ling,....



Coelotilapia joka

Coelotilapia joka

Coelotilapia joka is a vulnerable species of cichlid fish from rivers in Sierra Leone and Liberia in West Africa. This relatively small tilapia formerly was included in the genus Tilapia, but in 2013 it was moved to its own genus Coelotilapia and...



Coffinfish

Chaunax endeavouri

The coffinfish or furry coffinfish (Chaunax endeavouri) is a species of sea toad of the family Chaunacidae. It is found in salty temperate waters of southwestern Pacific, off east coast of Australia. The coffinfish was first discovered around...



Coho salmon

Oncorhynchus kisutch

The coho salmon (Oncorhynchus kisutch; Karuk: achvuun) is a species of anadromous fish in the salmon family and one of the five Pacific salmon species. Coho salmon are also known as silver salmon or "silvers". The scientific species name is based...



Coilia nasus

Coilia nasus

Coilia nasus, also known as ungeo and the Japanese grenadier anchovy or Chinese tapertail anchovy is a species of ray-finned fish from the family Engraulidae (anchovies). It grows to 41 cm (16 in) total length; it is a relatively large species...



Colomesus asellus

Colomesus asellus

Colomesus asellus, the Amazon puffer, asellus puffer, South American freshwater puffer, Peruvian puffer, or Brazilian puffer is a species of pufferfish confined to the Amazon, Essequibo and Orinoco basins in tropical South America. It is a...



Colomesus psittacus

Colomesus psittacus

Colomesus psittacus, the Banded puffer, parrot puffer or South American estuarine puffer, is a species of pufferfish found all along the Western Atlantic coastline of South America from the Gulf of Paria down to the mouth of the Amazon River in...



Colorado pikeminnow

Ptychocheilus lucius

The Colorado pikeminnow (Ptychocheilus lucius, formerly squawfish) is the largest cyprinid fish of North America and one of the largest in the world, with reports of individuals up to 6 ft (1.8 m) long and weighing over 100 pounds (45 kg). Native...



Colorado River cutthroat trout

Oncorhynchus clarkii pleuriticus

The Colorado River cutthroat trout (Oncorhynchus clarkii pleuriticus) is a subspecies of cutthroat trout native only to the Green and Colorado River basins, which are west of the Continental Divide.

Cutthroat trout found in other river basins...



Columbia River redband trout

Oncorhynchus mykiss gairdneri

The Columbia River redband trout, the inland redband trout or the interior redband trout (Oncorhynchus mykiss gairdneri) is one of three redband trout subspecies of the rainbow trout in the family Salmonidae. It is native in the Columbia River...



Comber

Serranus cabrilla

The comber (; Serranus cabrilla) is a species of marine ray-finned fish from the family Serranidae, the sea basses. It is widely distributed in the eastern North and South Atlantic Oceans and into the southwestern Indian Ocean. It is caught for...



Comet

Calloplelesios altivelis

The comet or marine betta (Calloplelesios altivelis) is a species of reef-associated tropical marine fish in the longfin family Plesiopidae, most commonly found between 3 and 50 m deep. It is native to the Indo-Pacific Ocean. It can reach a...



Comet grouper

Epinephelus morrhua

The comet grouper (*Epinephelus morrhua*), also known as the comet cod or dot-dash grouper, is a species of marine ray-finned fish, a grouper from the subfamily Epinephelinae which is part of the family Serranidae, which also includes the anthias...



Commerson's frogfish

Antennarius commerson

Commerson's frogfish or the giant frogfish, Antennarius commerson, is a marine fish belonging to the family Antennariidae.



Common barbel

Barbus barbus

The common barbel (*Barbus barbus*) is a species of freshwater fish belonging to the family Cyprinidae. It shares the common name 'barbel' with its many relatives in the genus Barbus, of which it is the type species. In Great Britain it is usually...



Common bleak

Alburnus alburnus

The common bleak (*Alburnus alburnus*) is a small freshwater coarse fish of the cyprinid family.



Common bluestripe snapper

Lutjanus kasmira

The common bluestripe snapper (*Lutjanus kasmira*), bluestripe snapper, bluebanded snapper, bluestripe sea perch, fourline snapper, blue-line snapper or moonlighter, is a species of snapper belonging to the family Lutjanidae. It is native to the...



Common bream

Abramis brama

The common bream (Abramis brama), also known as the freshwater bream, bream, bronze bream, carp bream or sweaty bream, is a European species of freshwater fish in the family Cyprinidae. It is now considered to be the only species in the genus...



Common bully

Gobiomorphus cotidianus

The common bully (Gobiomorphus cotidianus), or toitoi (Māori), is a fish endemic to New Zealand, and is present throughout the country. There are three other bully species that can be confused with common bullies. There are few characteristics of...



Common dab

Limanda limanda

The common dab (Limanda limanda) is an edible flatfish of the family Pleuronectidae. It is a demersal fish native to shallow seas around Northern Europe, in particular the North Sea, where it lives on sandy bottoms down to depths of about 100...



Common dace

Leuciscus leuciscus

The common dace (Leuciscus leuciscus) is a species of freshwater and brackish water ray-finned fish from the family Cyprinidae which is native to Europe but which has been introduced to other parts of the world. It is a quarry species for coarse...



Common dragonet

Callionymus lyra

The common dragonet (*Callionymus lyra*) is a species of dragonet which is widely distributed in the eastern North Atlantic where it is common near Europe from Norway and Iceland southwards. It is a demersal species that occurs over sand bottoms....



Common eagle ray

Myliobatis aquila

The common eagle ray or bullray (*Myliobatis aquila*) is a species of fish in the family Myliobatidae. It inhabits the eastern Atlantic Ocean (North Sea to South Africa), the Mediterranean Sea and the southwestern Indian Ocean.



Common galaxias

Galaxias maculatus

The common galaxias (*Galaxias maculatus*) or inanga (from the Māori *īnanga*) is a very widespread Southern Hemisphere fish in the family Galaxiidae. It is a slim, narrow fish with a forked tail and a mottled, spotty pattern, typically about 10 cm...



Common goby

Pomatoschistus microps

The common goby (*Pomatoschistus microps*) is a species of ray-finned fish native to fresh and brackish waters along the Atlantic and Baltic Sea coasts of Europe and northern Africa, with a range stretching from Norway to Morocco and Mauritania. It...



Common guitarfish

Rhinobatos rhinobatos

The common guitarfish (*Rhinobatos rhinobatos*) is a species of cartilaginous fish in the family Rhinobatidae. It is native to the eastern Atlantic Ocean and the Mediterranean Sea. It is a bottom-dwelling fish feeding on crustaceans, other...



Common hatchetfish

Gasteropelecus sternicla

The common hatchetfish or river hatchetfish (Gasteropelecus sternicla) is a tropical fish belonging to the freshwater hatchetfish family (Gasteropelecidae). Originating in South America in the Peruvian and middle Amazon, the Guianas and...



Common ling

Molva molva

The common ling (Molva molva), also known as the white ling or simply the ling, is a large member of the family Lotidae, a group of cod-like fishes. It resembles the related rocklings, but it is much larger and has a single barbel. This species...



Common logperch

Percina caprodes

The common logperch (Percina caprodes), sometimes simply known as the logperch, is a species of freshwater ray-finned fish, a darter from the subfamily Etheostomatinae, part of the family Percidae, which also contains the perches, ruffes and...



Common minnow

Phoxinus phoxinus

The Eurasian minnow, minnow, or common minnow (Phoxinus phoxinus) is a small species of freshwater fish in the carpfamily Cyprinidae. It is the type species of genus Phoxinus. It is ubiquitous throughout much of Eurasia, from Britain and Spain to...



Common mudskipper

Periophthalmus kalolo

The common mudskipper (Periophthalmus kalolo) is a species of mudskipper native to marine and brackish waters of the Indo-Pacific from eastern Africa to Samoa. This species can be found in mangrove forests where it spends most of its time out of...



Common nase

Chondrostoma nasus

The common nase (Chondrostoma nasus) is a European potamodromous cyprinid fish. It is often simply called the nase, but that can refer to any species of its genus Chondrostoma. Another name is sneep.



Common pandora

Pagellus erythrinus

The common pandora (Pagellus erythrinus), also known as king of the breams, pandora red pandora or Spanish seabream, is a species of marine ray-finned fish belonging to the family Sparidae, which includes the seabreams and porgies. This fish is...



Common remora

Remora remora

The common remora (Remora remora) is a pelagic marine fish belonging to family Echeneidae. The dorsal fin, which has 22 to 26 soft rays, acts as a suction cup, creating a vacuum to allow it to attach to larger marine animals, such as whales,....



Common roach

Rutilus rutilus

The roach, or rutilus roach (*Rutilus rutilus*), also known as the common roach, is a fresh- and brackish-water fish of the family Cyprinidae, native to most of Europe and western Asia. Fish called roach can be any species of the genera *Rutilus*,...



Common rudd

Scardinius erythrophthalmus

The common rudd (*Scardinius erythrophthalmus*) is a benthopelagic freshwater fish, widely spread in Europe and middle Asia, around the basins of the North, Baltic, Black, Caspian and Aral seas.



Common seadragon

Phyllopteryx taeniolatus

The common seadragon or weedy seadragon (*Phyllopteryx taeniolatus*) is a marine fish related to the seahorses. Adult common seadragons are a reddish colour, with yellow and purple markings; they have small leaf-like appendages that resemble kelp...



Common shiner

Luxilus cornutus

The common shiner (*Luxilus cornutus*) is a freshwater fish of the family Cyprinidae, found in North America. It ranges in length between 4 and 6 inches (10 and 15 cm), although they can reach lengths of up to 8 inches (20 cm).



Common silver-biddy

Gerres oyena

The common silver-biddy (*Gerres oyena*), also known as the blacktip silver biddy, Darnley Island silverbelly, longtail silverbiddy, oceanic silver biddy, shining silver-belly or slender silver belly, is a species of mojarra native to marine and...



Common skate

Dipturus batis

The common skate (Dipturus batis), also known as the blue skate, is the largest skate in the world, attaining a length of up to 2.85 m (9 ft 4 in). Historically, it was one of the most abundant skates in the northeast Atlantic Ocean and the...



Common smooth-hound

Mustelus mustelus

Sharks portalThe common smooth-hound (Mustelus mustelus) is a houndshark of the family Triakidae. It is found in the eastern Atlantic Ocean from the British Isles to South Africa, and in the Mediterranean Sea, Madeira, and the Canary Islands at...



Common snook

Centropomus undecimalis

The common snook (Centropomus undecimalis) is a species of marine fish in the family Centropomidae of the order Perciformes. The common snook is also known as the sergeant fish or robalo. It was originally assigned to the sciaenid genus Sciaena;...



Common snowtrout

Schizothorax richardsonii

The common snowtrout or snowtrout (Schizothorax richardsonii), also known as "Asala", or "Asela" by the locals, is a species belong to the class of ray-finned fishes, family Cyprinidae, order Cypriniformes in the genus Schizothorax. The genus...



Common sole

Solea solea

The common sole, Dover sole, or black sole (Solea solea) is a species of flatfish in the family Soleidae. It is one of the largest fish in the Solea genus. It lives on the sandy or muddy seabed of the northern Atlantic and the Mediterranean Sea...



Common stingray

Dasyatis pastinaca

The common stingray (Dasyatis pastinaca) is a species of stingray in the family Dasyatidae, found in the northeastern Atlantic Ocean and the Mediterranean and Black Seas. It typically inhabits sandy or muddy habitats in coastal waters shallower...



Common thresher

Alopias vulpinus

The common thresher (Alopias vulpinus), also known as Atlantic thresher, is the largest species of thresher shark, family Alopidae, reaching some 6 m (20 ft) in length. About half of its length consists of the elongated upper lobe of its caudal...



Common torpedo

Torpedo torpedo

The common torpedo (Torpedo torpedo), also known as ocellate torpedo or eyed electric ray, is a species of electric ray in the family Torpedinidae. It is found in the Mediterranean Sea and the eastern Atlantic Ocean from the Bay of Biscay to...



Conger erebennus

Conger erebennus

Conger erebennus is an eel in the family Congridae (conger/garden eels). It was described by David Starr Jordan and John Otterbein Snyder in 1901, originally under the genus Leptocephalus. It is a marine, temperate water-dwelling eel which is...



Congo pufferfish

Tetraodon miurus

The Congo puffer or potato puffer (Tetraodon miurus) is a freshwater pufferfish found in areas of the Congo River in Africa, including rapids. The Congo puffer populations in the wild are still healthy in suitable habitats and it is considered...



Congo tetra

Phenacogrammus interruptus

The Congo tetra (Phenacogrammus interruptus) is a species of fish in the African tetra family, found in the central Congo River Basin in Africa. It is commonly kept in aquaria.



Congoli

Pseudaphritis urvillii

The congoli (Pseudaphritis urvillii), also known as the freshwater flathead, marble fish, marbled flathead, sand trout, sanding, sandy, sandy whiting or tupong, is a species of marine ray-finned fish and it is the only species of fish in the...



Congrogadus subducens

Congrogadus subducens

The carpet eel-blenny (Congrogadus subducens), also known as the green wolf eel or green wolf eel blenny, is a relatively large species of dottyback found in coastal parts of the Indo-West Pacific, including coral reefs, among rocks, seagrass...



Convict cichlid

Amatitlania nigrofasciata

The convict cichlid (Amatitlania nigrofasciata) is a fish species from the family Cichlidae, native to Central America, also known as the zebra cichlid. Convict cichlids are popular aquarium fish and have also been the subject of numerous studies...



Convict julie

Julidochromis regani

The convict julie (Julidochromis regani) is a cichlid species in the subfamily Pseudocrenilabrinae family endemic to Lake Tanganyika. Hence it is found in Burundi, the Democratic Republic of the Congo, Tanzania, and Zambia. The fish is named...



Cookeolus japonicus

Cookeolus japonicus

Cookeolus japonicus is a species of fish in the family Priacanthidae, the bigeyes and catalufas. It is the only extant species of Cookeolus, except for C. spinolacrymatus, an extinct Late Pliocene fish known from a fossil specimen collected in...



Cookiecutter shark

Isistius brasiliensis

The cookiecutter shark (Isistius brasiliensis), also called the cigar shark, is a species of small squaliform shark in the family Dalatiidae. This shark occurs in warm, oceanic waters worldwide, particularly near islands, and has been recorded as...



Copadichromis azureus

Copadichromis azureus

Copadichromis azureus is a species of haplochromine cichlid. It is endemic to Lake Malawi where it is found in the country of Malawi.



Copadichromis borleyi

Copadichromis borleyi

Copadichromis borleyi is a species of haplochromine cichlid fish endemic to Lake Malawi in East Africa. The species is popular in the fishkeeping hobby where it is frequently kept in aquariums. The species has numerous common names, including...



Copella arnoldi

Copella arnoldi

Copella arnoldi, commonly known as the splash tetra or the splashing tetra, is a species of tropical freshwater fish belonging to the family Lebiasinidae. It is native to South America. The fish is named in honor of German aquarist Johann Paul...



Copper redhorse

Moxostoma hubbsi

The copper redhorse (Moxostoma hubbsi) is a North American species of freshwater fish in the family Catostomidae. It is found only in Canada. Its extremely small range, which is restricted to a few rivers in the lowlands of southwestern Quebec, has...



Copper rockfish

Sebastes caurinus

The copper rockfish (Sebastes caurinus), also known as the copper seaperch, is a species of marine ray-finned fish belonging to the subfamily Sebastinae, the rockfishes, part of the family Scorpaenidae. It is found in the eastern Pacific.



Copper shark

Carcharhinus brachyurus

The copper shark (Carcharhinus brachyurus), bronze whaler, or narrowtooth shark is a species of requiem shark found mostly in temperate latitudes. It is distributed in a number of separate populations in the northeastern and southwestern...



Copperband butterflyfish

Chelmon rostratus

The copperband butterflyfish (Chelmon rostratus), also known as the beaked coral fish, is found in reefs in both the Pacific and Indian Oceans. This butterflyfish is one of the three species that make up the genus Chelmon and all have long beaks.



Coral catshark

Atelomycterus marmoratus

The coral catshark (Atelomycterus marmoratus) is a species of catshark in the family Scyliorhinidae. It is common on shallow coral reefs across the Indo-West Pacific, from Pakistan to New Guinea. Reaching up to 70 cm (28 in) in length, the coral...



Coral grouper

Cephalopholis miniata

Cephalopholis miniata, also known as the coral grouper, coral hind, coral rock cod, coral cod, coral trout, round-tailed trout or vermillion seabass is a species of marine ray-finned fish, a grouper from the subfamily Epinephelinae which is in...



Coral hawkfish

Cirrhitichthys oxycephalus

The coral hawkfish (Cirrhitichthys oxycephalus), the pixy hawkfish or sharp-headed hawkfish, is a species of marine ray-finned fish, a hawkfish belonging to the family Cirrhitidae. It is native to tropical reefs of the Indian Ocean and the...



Coregonus albula

Coregonus albula

Coregonus albula, known as the vendace or as the European cisco, is a species of freshwater whitefish in the family Salmonidae. It is found in lakes in northern Europe, especially Finland, Latvia, Lithuania, Sweden, Russia and Estonia, and in...



Coregonus artedi

Coregonus artedi

Coregonus artedi, commonly known as the cisco, is a North American species of freshwater whitefish in the family Salmonidae. The number of species and definition of species limits in North American ciscoes is a matter of debate. Accordingly,...



Coregonus baicalensis

Coregonus baicalensis

Coregonus baicalensis is a species of freshwater whitefish in the family Salmonidae. It is endemic to the Russian Federation's Lake Baikal where it is found on the bottom demersal. The maximum length recorded for this species is 60 centimetres...



Coregonus bavaricus

Coregonus bavaricus

Coregonus bavaricus, the Ammersee kilch, is a species of freshwater whitefish endemic to Lake Ammersee in the German state of Upper Bavaria. A small, silver-colored fish, it typically lives between 60–85 m (197–279 ft) deep, though shallower in...



Coregonus hoyi

Coregonus hoyi

Coregonus hoyi, also known as the bloater, is a species or form of freshwater whitefish in the family Salmonidae. It is a silvery-coloured herring-like fish, 25.5 centimetres (10.0 in) long. It is found in most of the Great Lakes and in Lake...



Coregonus lavaretus

Coregonus lavaretus

Coregonus lavaretus is a species of freshwater whitefish, in the family Salmonidae. It is the type species of its genus Coregonus. There are widely different concepts about the delimitation of the species Coregonus lavaretus and about the number...



Coregonus maraena

Coregonus maraena

Coregonus maraena, referred to in English as the maraene, maraena whitefish, vendace, cisco, lake herring, lake whitefish or the whitefish, is a whitefish of the family Salmonidae that occurs in the Baltic Sea basin - in the sea itself and the...



Coregonus sardinella

Coregonus sardinella

Coregonus sardinella, known as the least cisco or the sardine cisco, is a fresh- and brackishwater salmonid fish that inhabits rivers, estuaries and coastal waters of the marginal seas of the Arctic Basin, as well as some large lakes of those...



Coregonus trybomi

Coregonus trybomi

Coregonus trybomi is a freshwater whitefish in the family Salmonidae. It is a spring-spawning type of cisco, which probably has evolved from sympatric vendace (Coregonus albula) independently in a number of Swedish lakes. Only one of those...



Coregonus vandesius

Coregonus vandesius

Coregonus vandesius, the vendace, is a freshwater whitefish found in the United Kingdom. Population surveys since the 1960s have revealed a steady decline and the fish is no longer present in some of its previous haunts but is still present in...



Coregonus widegreni

Coregonus widegreni

Coregonus widegreni, also called the Valaam whitefish, is a putative species of freshwater whitefish, a part of the common whitefish complex (*Coregonus lavaretus sensu lato*) from Northern Europe. It is a demersal form of freshwater whitefish that...



Coreoleuciscus splendidus

Coreoleuciscus splendidus

Coreoleuciscus splendidus, also known as the swiri or Korean splendid dace, is a species of fish from the cyprinid family endemic to rivers of the Korean peninsula. It was first scientifically described by Tamezo Mori in 1935, who noted in 1936...



Coris gaimard

Coris gaimard

Coris gaimard, the yellowtail wrasse or African coris, among other vernacular names, is a species of wrasse native to the tropical waters of the central Indian Ocean and the western Pacific Ocean, from Christmas Islands and Cocos Keeling Islands...



Corkwing wrasse

Symphodus melops

The corkwing wrasse (Symphodus melops) is a species of wrasse native to the eastern Atlantic Ocean from Norway to Morocco and out to the Azores, as well as being found in the Mediterranean Sea and the Adriatic Sea. This species can be found in...



Cornish jack

Mormyrops anguilloides

The Cornish jack, Mormyrops anguilloides, is a species of weakly electric fish in the family Mormyridae, native to quiet waters in much of Sub-Saharan Africa. The largest species in its family, the Cornish jack is a nocturnal group hunter of...



Cortez rainbow wrasse

Thalassoma lucasanum

The Cortez rainbow wrasse (*Thalassoma lucasanum*) is a species of wrasse native to the eastern Pacific Ocean from Baja California to Peru, as well as around the Galapagos Islands. It is a reef inhabitant, occurring in small schools from the...



Corydoras atropersonatus

Corydoras atropersonatus

Corydoras atropersonatus is a tropical freshwater fish belonging to the subfamily Corydoradinae of the family Callichthyidae. It originates in inland waters in South America, and is found in the upper Amazon River basin in Ecuador and Peru.The...



Corydoras haraldschultzi

Corydoras haraldschultzi

Corydoras haraldschultzi is a tropical freshwater fish belonging to the Corydoradinae sub-family of the family Callichthyidae. It is found in Brazil and Bolivia. The fish will grow to 7 centimeters. It lives in a tropical climate in water with a...



Corydoras julii

Corydoras julii

Corydoras julii (also known as the julii cory or leopard catfish) is a small freshwater catfish native to eastern Brazil. It is often confused with Corydoras trilineatus, the three stripe corydoras. Corydoras julii are small, peaceful shoaling...



Corydoras leopardus

Corydoras leopardus

Corydoras leopardus, the leopard cory, is a catfish very similar to, and often confused with, the three line cory, Corydoras trilineatus. The most obvious differences are that C. leopardus has a longer, more pointed snout than C. trilineatus, and...



Corydoras loxozonus

Corydoras loxozonus

Corydoras loxozonus is a tropical freshwater fish belonging to the Corydoradinae sub-family of the family Callichthyidae. It originates in inland waters in South America, and is found in the Meta River basin in Colombia. In the system of...



Corydoras melini

Corydoras melini

Corydoras melini, the bandit corydoras or false bandit catfish, is a tropical freshwater fish belonging to the subfamily Corydoradinae of the family Callichthyidae. It originates in inland waters in South America, and is found in the upper Rio...



Corydoras nanus

Corydoras nanus

Corydoras nanus is a tropical freshwater fish belonging to the Corydoradinae sub-family of the family Callichthyidae. It originates in inland waters in South America, and is found in the Suriname and Maroni River basins in Suriname and the...



Corydoras paleatus

Corydoras paleatus

Corydoras paleatus is a species of catfish (order Siluriformes) of the family Callichthyidae. Its common names include blue leopard corydoras, mottled corydoras, and peppered catfish. It originates from the lower Paraná River basin and coastal...



Corydoras panda

Corydoras panda

Corydoras panda is a species of catfish belonging to the genus Corydoras, of the family Callichthyidae, and is a native member of the riverine fauna of South America. It is found in Peru and Ecuador, most notably in the Huánuco region, where it...



Corydoras punctatus

Corydoras punctatus

Corydoras punctatus (spotted catfish) is a species of catfish found in the Suriname river basin and Iracoubo river basin in tropical Suriname and French Guiana. The species is rarely seen. It is hard to tell the sex of the fish until they are an...



Coryphaenoides rupestris

Coryphaenoides rupestris

Coryphaenoides rupestris is a species of marine ray-finned fish in the family Macrouridae. Its common names include the rock grenadier, the roundnose grenadier and the roundhead rat-tail. In France it is known as grenadier de roche and in Spain...



Corythoichthys flavofasciatus

Corythoichthys flavofasciatus

Corythoichthys flavofasciatus, known commonly as the network pipefish, reticulate pipefish and yellow-banded pipefish, is a species of marine fish in the family Syngnathidae.



Costello tetra

Hemigrammus hyanuary

The Costello tetra is a species of characin from the Amazon basin and is found in Brazil and Peru. The specific name comes from Lake Hyanuary in Brazil. Other common names used in English are January tetra and green neon. The name "green neon" is...



Cottus perifretum

Cottus perifretum

Cottus perifretum, the bullhead or miller's thumb, is a species of freshwater ray-finned fish belonging to the family Cottidae, the typical sculpins. It is found on both sides of the English Channel, native to Great Britain. It also inhabits...



Cownose ray

Rhinoptera bonasus

The cownose ray (Rhinoptera bonasus) is a species of Batoidea found throughout a large part of the western Atlantic and Caribbean, from New England to southern Brazil (the East Atlantic populations are now generally considered a separate species,...



Cowtail stingray

Pastinachus sephen

The cowtail stingray (Pastinachus sephen) is a species of stingray in the family Dasyatidae, widespread in the Indo-Pacific region and occasionally entering freshwater habitats. Other common names include banana-tail ray, drab stingray, fantail...



Crazy fish

Butis butis

Butis butis, the crazy fish, duckbill sleeper, or upside-down sleeper, is a species of sleeper goby that are native to brackish and freshwater coastal habitats of the Indian Ocean and the western Pacific Ocean from the African coast to the...



Creek chubsucker

Erimyzon oblongus

The creek chubsucker (Erimyzon oblongus) is a freshwater fish of the sucker family (Catostomidae).



Creek whaler

Carcharhinus fitzroyensis

The creek whaler (Carcharhinus fitzroyensis) is a common species of requiem shark, and part of the family Carcharhinidae, endemic to northern Australia. It frequents shallow waters close to shore, including estuaries. This small, stocky shark...



Creole wrasse

Clepticus parrae

The creole wrasse (Clepticus parrae) is a species of wrasse native to the western Atlantic Ocean.



Crested bullhead shark

Heterodontus galeatus

The crested bullhead shark (Heterodontus galeatus) is an uncommon species of bullhead shark, in the family Heterodontidae. It lives off the coast of eastern Australia from the coast to a depth of 93 m (305 ft). This shark can be distinguished...



Crevalle jack

Caranx hippos

The crevalle jack (Caranx hippos), also known as the common jack, black-tailed trevally, couvalli jack, black cavalli, jack crevale, or yellow cavalli is a common species of large marine fish classified within the jack family, Carangidae. The...



Croaking gourami

Trichopsis vittata

The croaking gourami (*Trichopsis vittata*) is a species of small freshwater labyrinth fish of the gourami family. They are native to still waters in Southeast Asia and are distributed worldwide via the aquarium trade. Croaking gouramis are capable...



Crocodile shark

Pseudocarcharias kamoharai

The crocodile shark (Pseudocarcharias kamoharai) is a species of mackerel shark and the only extant member of the family Pseudocarchariidae. A specialized inhabitant of the mesopelagic zone, the crocodile shark can be found worldwide in tropical...



Crocodile snake eel

Brachysomophis crocodilinus

The crocodile snake eel (Brachysomophis crocodilinus) is an eel in the family Ophichthidae (worm/snake eels). It was described by Edward Turner Bennett in 1833. It is a tropical, marine eel which is known from the Indo-Pacific, including East...



Crossocheilus langei

Crossocheilus langei

Crossocheilus langei, also known as the red algae eater (but sold sometimes sold as "Siamese algae eater"), is a species of ray-finned fish in the genus Crossocheilus. It is native to Malaysia and Sumatra. This bottom-dwelling tropical fish is...



Crossocheilus reticulatus

Crossocheilus reticulatus

Crossocheilus reticulatus (silver flying fox) is a freshwater fish in the family Cyprinidae from Southeast Asia. It grows to 17 cm (6.7 in) standard length.



Crucian carp

Carassius carassius

The crucian carp (Carassius carassius) is a medium-sized member of the common carp family Cyprinidae. It occurs widely in northern European regions. Its name derives from the Low German karusse or karutze, possibly from Medieval Latin coracinus...



Crystal darter

Crystallaria asprella

The crystal darter (Crystallaria asprella) is a species of freshwater ray-finned fish, a darter from the subfamily Etheostomatinae, part of the family Percidae, which also contains the perches, ruffes and pikeperches. This small North American...



Ctenochaetus binotatus

Ctenochaetus binotatus

Ctenochaetus binotatus, the twospot bristletooth or twospot surgeonfish, is a species of marine ray-finned fish belonging to the family Acanthuridae, the surgeonfishes unicornfishes and tangs. This species has a wide Indo-Pacific distribution.



Ctenochaetus cyanocheilus

Ctenochaetus cyanocheilus

Ctenochaetus cyanocheilus, the bluelip bristletooth, yelloweye bristletooth, goldring bristletooth, Indo-Pacific yellow tang or short-tail bristle-tooth, is a species of marine ray-finned fish belonging to the family Acanthuridae which includes...



Ctenochaetus hawaiiensis

Ctenochaetus hawaiiensis

Ctenochaetus hawaiiensis thechevron tang, black surgeonfish, Hawaiian bristletooth, Hawaiian kole or Hawaiian surgeonfish is a species of marine ray-finned fish belonging to the family Acanthuridae which includes the surgeonfishes, unicornfishes...



Ctenochaetus striatus

Ctenochaetus striatus

Ctenochaetus striatus, the striated surgeonfish, lined bristletooth, fine-lined bristletooth or striped bristletooth, is a species of marine ray-finned fish belonging to the family Acanthuridae, the surgeonfishes, unicoornfishes and tangs This...



Ctenochaetus strigosus

Ctenochaetus strigosus

Ctenochaetus strigosus, the kole tang, spotted bristletooth, spotted surgeonfish, goldring bristletooth, goldring surgeonfish, yelloweye tang or yellow-eyed surgeonfish, is a species of marine ray-finned fish belonging to family Acanthuridae...



Ctenochaetus tominiensis

Ctenochaetus tominiensis

Ctenochaetus tominiensis, the Tomini surgeonfish, Tomini bristletooth, yellowtip bristletooth or orangetip bristletooth, is a species of marine ray-finned fish belonging to the family Acanthuridae which includes the surgeonfishes, unicornfishes...



Ctenochaetus truncatus

Ctenochaetus truncatus

Ctenochaetus truncatus, the Indian gold-ring bristletooth, squaretail bristletooth, yelloweye bristletooth, truncate bristletooth or spotted yellow eye tang, is a species of marine ray-finned fish belonging to the family Acanthuridae which...



Ctenolucius hujeta

Ctenolucius hujeta

Ctenolucius hujeta, also known as silver gar, rocket gar, hujeta gar and freshwater barracuda, is a slender, medium-sized freshwater fish found in Venezuela and Colombia, where it is known as "agujeta"; Spanish for "shoelace".Large specimens can...



Ctenopoma multispine

Ctenopoma multispine

Ctenopoma multispine, known as the many-spined ctenopoma or climbing perch, is an African freshwater fish. It is up to 14 cm (5.5 in) long and occurs in the Lufira River, Lualaba River, Upemba River, Luapula-Moero in the Democratic Republic of...



Cuban dogfish

Squalus cubensis

The Cuban dogfish (Squalus cubensis) is a dogfish, a member of the family Squalidae in the order Squaliformes.



Cuban gar

Atractosteus tristoechus

The Cuban gar (Atractosteus tristoechus), also known as the manjuarí, is a fish in the family Lepisosteidae. It is a tropical, freshwater species, although it also inhabits brackish water. It is found in rivers and lakes of western Cuba and the...



Cubera snapper

Lutjanus cyanopterus

The cubera snapper (*Lutjanus cyanopterus*), also known as the Cuban snapper, is a species of marine ray-finned fish, a snapper belonging to the family Lutjanidae. It is native to the western Atlantic Ocean. It is a commercially important species...



Cuckoo wrasse

Labrus mixtus

The cuckoo wrasse (*Labrus mixtus*) is a species of wrasse native to the eastern Atlantic Ocean from Norway to Senegal, including the Azores and Madeira. It is also found in the Mediterranean Sea. They occur in weedy, rocky areas mostly between 40...



Cui-ui

Chasmistes cujus

The cui-ui (*Chasmistes cujus*) is a large sucker fish endemic to Pyramid Lake and, prior to its desiccation in the 20th century, Winnemucca Lake in northwestern Nevada. It feeds primarily on zooplankton and possibly on nanoplankton (such as algae...



Culter alburnus

Culter alburnus

Culter alburnus, the topmouth culter, is an East Asian species of freshwater ray-finned fish in the cyprinid genus *Culter*.



Cunner

Tautoglabrus adspersus

The cunner (*Tautoglabrus adspersus*), also known as the blue perch, bergall, chogset, choggie, conner or sea perch, is a species of wrasse native to the northwestern Atlantic, where it is found from the Gulf of St. Lawrence and Newfoundland to...



Cusk

Brosme brosme

The cusk or tusk (Brosme brosme) is a North Atlantic cod-like fish in the ling family Lotidae. It is the only species in the genus Brosme. Its other common names include European cusk, and brosmius.



Cutthroat trout

Oncorhynchus clarkii

The cutthroat trout (Oncorhynchus clarkii) is a fish species of the family Salmonidae native to cold-water tributaries of the Pacific Ocean, Rocky Mountains, and Great Basin in North America. As a member of the genus Oncorhynchus, it is one of...



Cyclopterus

Cyclopterus lumpus

Cyclopterus is a genus of marine ray-finned fish belonging to the family Cyclopteridae, the lumpsuckers or lumpfish. Its only species is Cyclopterus lumpus, the lumpsucker or lumpfish. It is found in the North Atlantic and adjacent parts of the...



Cynoglossus cynoglossus

Cynoglossus cynoglossus

Cynoglossus cynoglossus, commonly known as the Bengal tonguesole, sometimes called the Gangetic tongue sole, is a species of tonguefish. It is commonly found in the Indian Ocean, particularly in brackish water and muddy or sandy bottoms off the...



Cynoglossus lingua

Cynoglossus lingua

Cynoglossus lingua, commonly known as the long tongue sole or textile tongue sole is a species of fish found along the coast of the Philippines, Thailand and the Malay peninsula, Myanmar, the Bay of Bengal and as far west as the Red Sea.



Cynoscion

Cynoscion regalis

Cynoscion is a genus of marine ray-finned fishes belonging to the family, Sciaenidae, the drums and croakers. These fishes are found off the coasts of North and South America in the western Atlantic and eastern Pacific Oceans. Many fishes in this...



Cynoscion acoupa

Cynoscion acoupa

Cynoscion acoupa, the acoupa weakfish, blacktail basher or grey snapper, is a species of marine ray-finned fish belonging to the family Sciaenidae, the drums and croakers. This fish is found in the western Atlantic.



Cynoscion arenarius

Cynoscion arenarius

Cynoscion arenarius, sand seatrout, sand weakfish or white trout, is a species of marine ray-finned fish belonging to the family Sciaenidae, the drums and croakers. It is found in the western Atlantic Ocean. It is closely related to the common...



Cynoscion nebulosus

Cynoscion nebulosus

Spotted seatrout (Cynoscion nebulosus), also known as speckled trout, is a common estuarine fish found in the southern United States along coasts of Gulf of Mexico and the coastal Atlantic Ocean from Maryland to Florida. While most of these fish...



Cynotilapia afra

Cynotilapia afra

Cynotilapia afra, the afra cichlid or dogtooth cichlid, is a small species of cichlid fish from Lake Malawi in East Africa, where found in rocky habitats. The genus name roughly translates as dogtooth cichlid (hence its common name) which...



Cyphotilapia frontosa

Cyphotilapia frontosa

Cyphotilapia frontosa, also called the front cichlid and frontosa cichlid, is an east African species of fish endemic to Lake Tanganyika. The genus name is a combination of the Ancient Greek "cypho-", meaning "curved", and tilapia, which means...



Cyphotilapia gibberosa

Cyphotilapia gibberosa

Cyphotilapia gibberosa is a species of fish in the cichlid family, one of two species in the genus Cyphotilapia. Native to Lake Tanganyika in East Africa, it was described in 2003 nearly 100 years after its congener, C. frontosa. This species is...



Cyprichromis leptosoma

Cyprichromis leptosoma

Cyprichromis leptosoma is a mouthbrooding species of fish in the family Cichlidae. It is endemic to Lake Tanganyika in Zambia and Tanzania. It seems quite common within its range and faces no particular threats, so the International Union for...



Cyprinella spiloptera

Cyprinella spiloptera

The spotfin shiner (Cyprinella spiloptera) is a species of ray-finned fish in the family Cyprinidae. It is a small sized freshwater fish found abundantly in many watercourses of North America.



Cyprinodon julimes

Cyprinodon julimes

The Julimes pupfish (Cyprinodon julimes) (Spanish: Cachorrito de Julimes) is a species of killifish in the family Cyprinodontidae. This pupfish is endemic to El Pandeño, a hot spring in Julimes, Chihuahua, Mexico, and it is adapted to life in...



Cyprinodon longidorsalis

Cyprinodon longidorsalis

Cyprinodon longidorsalis, the cachorrito de charco palmar or La Palma pupfish, is a species of fish in the family Cyprinodontidae. It was endemic to the Ojo de Agua la Presa in southwestern Nuevo Leon state in Mexico, but became extinct in the...



Cyprinodon nevadensis

Cyprinodon nevadensis

Cyprinodon nevadensis is a species of pupfish in the genus Cyprinodon. The species is also known as the Amargosa pupfish, but that name may also refer to one subspecies, Cyprinodon nevadensis amargosae. All six subspecies are or were endemic to...



Cyprinus carpio carpio

Cyprinus carpio carpio

Cyprinus carpio carpio is a subspecies of the common carp that is commonly found in Europe. They are native to much of Europe (notably the Danube and Volga Rivers) and can also be found in the Caucasus and Central Asia. Mitochondrial DNA analysis...



Cyprinus rubrofuscus

Cyprinus rubrofuscus

Cyprinus rubrofuscus, the Amur carp, is a species of cyprinid fish, and is the wild form of the well-known koi. It is widespread in the fresh waters of eastern Asia, native to China, Korea, Russia, Vietnam and Laos from the Amur to Red River...



Cyrtocara

Cyrtocara moorii

Cyrtocara moorii, commonly known as the hump-head, is a species of haplochromine cichlid endemic to Lake Malawi in east Africa where they prefer areas with sandy substrates. It can grow to a length of 20 centimetres (7.9 in) TL. The species is...



Dabry's sturgeon

Acipenser dabryanus

Dabry's sturgeon (Acipenser dabryanus), also known as the Yangtze sturgeon, Changjiang sturgeon and river sturgeon, is a species of fish in the sturgeon family, Acipenseridae. It is endemic to China and today restricted to the Yangtze River...



Dacodraco

Dacodraco hunteri

Dacodraco is a monotypic genus of marine ray-finned fish belonging to the family Channichthyidae, the crocodile icefishes, its only member is Dacodraco hunteri. This species is found in the Southern Ocean.



Dadio

Laubuka dadiburjori

The dadio (Laubuka dadiburjori) is a cyprinid fish, is rarely seen in the aquarist hobby but is not entirely unlike the Danio nigrofasciatus in appearance. Laubuka dadiburjori is a gold/silver fish with a blue line, it has two colour morphs, one...



Daggernose shark

Isogomphodon oxyrhynchus

The daggernose shark (Isogomphodon oxyrhynchus) is a little-known species of requiem shark, in the family Carcharhinidae, and the only extant member of its genus. It inhabits shallow tropical waters off northeastern South America, from Trinidad...



Daggertooth pike conger

Muraenesox cinereus

The daggertooth pike conger (Muraenesox cinereus) also known as the darkfin pike eel in Australia, to distinguish it from the related pike-eel (Muraenesox bagio), is a species of eel in the pike conger family, Muraenesocidae. They primarily live...



Daisy stingray

Fontitrygon margarita

The daisy stingray, Fontitrygon margarita, is a little-known species of stingray in the family
Dasyatidae, found in shallow waters along the coast of West Africa. This species typically grows to 60
cm (24 in) across and has a rounded pectoral fin...



Damba

Paretroplus damii

The damba (Paretroplus damii) is a species of cichlid.



Danio annulosus

Danio annulosus

Danio annulosus is a species of Chain danio fish in the family Cyprinidae in Danio genus. The species is named and described jointly by Sven O Kullander, M.D Mizanur Rahman, Michael Noren and Abdur Rob Mollah in 2015 in association with the...



Danio dangila

Danio dangila

Danio dangila, the moustached danio, is a freshwater fish, and is the largest of the true Danio species at up to (6 in) long. Its name is from its particularly long barbels. It is sometimes kept in aquariums, where its relatively passive nature...



Danio erythromicron

Danio erythromicron

Danio erythromicron, often known as emerald dwarf danio and emerald dwarf rasbora, is a species of cyprinid fish which is endemic to Inle Lake in Myanmar.



Danio kyathit

Danio kyathit

Danio kyathit is also call ocelot danio,is a small, schooling species of fish in the family Cyprinidae. It is endemic to the upper reaches of Irrawaddy River near Myitkyina in northern Myanmar. Described in 1998, it is closely related to the...



Danio margaritatus

Danio margaritatus

Danio margaritatus, the celestial pearl danio, often referred to in the aquarium trade as galaxy rasbora or Microrasbora sp. 'Galaxy', is a small cyprinid from Myanmar and Northern Thailand (In Salween basin) . It has so far been found only in a...



Danio tinwini

Danio tinwini

Danio tinwini, commonly called gold-ring danio, is a newly discovered species of Danio from Myanmar. It is also referred to as Danio sp. "TW02". It is a tiny gold fish whose body and fins are covered with blue spots. It has also been referred to...



Danionella translucida

Danionella translucida

Danionella translucida is an extremely small (1.1 cm long) species of cyprinid fish endemic to Myanmar. When described, it was considered to be one of the smallest fish. It was collected from the roots of floating aquatic plants in a...



Danube bleak

Alburnus chalcoides

The Danube bleak or Caspian shemaya (Alburnus chalcoides) is a species of freshwater fish in the family Cyprinidae. It is found in Iran, Ukraine, Georgia, Armenia, Slovakia, Moldova, Greece, Czechia, Azerbaijan, Turkiye, Afghanistan, Austria,...



Dara

Parakuhlia macrophthalmus

The dara (Parakuhlia macrophthalmus) is a species of marine ray-finned fish, a grunt belonging to the family Haemulidae. It is native to the Atlantic coast of Africa. It is the only species in the monospecific genus Parakuhlia.



Dark shyshark

Haploblepharus pictus

The dark shyshark or pretty happy (Haploblepharus pictus) is a species of catshark, belonging to the family Scyliorhinidae, endemic to the temperate waters off southern Namibia and western South Africa. It is benthic in nature and inhabits...



Dark sleeper

Odontobutis obscura

The dark sleeper, Odontobutis obscura, is a species of freshwater sleeper native to China, Japan, and Korea. This species can reach up to 25 cm (9.8 in) in length. It is a commercially important species. This species is also variously called...



Dascyllus albisella

Dascyllus albisella

Dascyllus albisella commonly known as the Hawaiian dascyllus, Hawaiian domino, Domino damselfish, or white-spotted damsel is a marine fish found in the Eastern Central Pacific.



Dascyllus melanurus

Dascyllus melanurus

Dascyllus melanurus, known commonly as the four stripe damselfish, blacktail dascyllus, humbug damselfish, blacktail damselfish, and blacktail humbug, is a species of fish in the family Pomacentridae. It is native to the western Pacific Ocean. It...



Dascyllus reticulatus

Dascyllus reticulatus

Dascyllus reticulatus, known commonly as the reticulate dascyllus or two-stripe damselfish among other vernacular names, is a species of marine fish in the family Pomacentridae. Reticulate dascyllus is widespread throughout the tropical waters of...



Dash-and-dot goatfish

Parupeneus barberinus

The dash-and-dot goatfish (Parupeneus barberinus) is a species of goatfish native to the Indian and Pacific oceans.



Dawkinsia assimilis

Dawkinsia assimilis

Dawkinsia assimilis is a species of ray-finned fish in the genus Dawkinsia. It is endemic to the southern Western Ghats especially to the Southwest Indian states of Karnataka and Kerala. They are known as Mascara Barb. Filament barbs are a group...



Dawkinsia filamentosa

Dawkinsia filamentosa

Dawkinsia filamentosa, the filament barb, or poovali paral is a species of barb. Young fish have barely any color and black spots. They start having more color at three months old. The fish is a swift swimmer. Males are larger than females and...



Dawn tetra

Aphyocharax paraguayensis

The dawn tetra, panda tetra, or Paraguay tetra (Aphyocharax paraguayensis) comes from the Rio Paraguay basin in Paraguay and Brazil. It is a schooling fish; suitable temperatures for the Paraguay tetra are 24 to 27 °C. This species is omnivorous...



Death Valley pupfish

Cyprinodon salinus

The Death Valley pupfish (Cyprinodon salinus), also known as Salt Creek pupfish, is a small species of fish in the family Cyprinodontidae found only in Death Valley National Park, California, United States. There are two recognized subspecies: C....



Decapterus russelli

Decapterus russelli

Decapterus russelli, also known as the Indian scad, northern mackerel scad, round scad, Russell's mackerel scad, slender scad, or three lined grunter, is a species of ray-finned fish in the family Carangidae. It naturally occurs in the Indian and...



Deepsea lizardfish

Bathysaurus ferox

The deepsea lizardfish, Bathysaurus ferox, is an aulopiform of the family Bathysauridae, found in tropical and subtropical seas across the world. The deepsea lizardfish should not be confused with the true or "typical" lizardfishes of the related...



Deepsea skate

Bathyraja abyssicola

The deepsea skate (Bathyraja abyssicola) is a species of softnose skate, in the family Arhynchobatidae, found in deep water from 362 to 2,906 m, usually on the continental slope. They are distributed from off northern Baja California around...



Deepwater sculpin

Myoxocephalus thompsonii

The deepwater sculpin (Myoxocephalus thompsonii) is a species of freshwater fish in the family Cottidae of order Scorpaeniformes. It is a glacial relict, native to a limited number of deep, cold lakes in Canada and the United States.The deepwater...



Deepwater stingray

Plesiobatis daviesi

The deepwater stingray or giant stingaree (Plesiobatis daviesi) is a species of stingray and the sole member of the family Plesiobatidae. It is widely distributed in the Indo-Pacific, typically over fine sediments on the upper continental slope...



Delminichthys adspersus

Delminichthys adspersus

Delminichthys adspersus (or spotted minnow; Croatian: imotska gaovica) is a species of cyprinid fish.It is found in Bosnia and Herzegovina and Croatia. Its natural habitats are rivers, intermittent rivers, freshwater lakes, and inland karsts.It...



Delta smelt

Hypomesus transpacificus

The delta smelt (*Hypomesus transpacificus*) is an endangered slender-bodied smelt, about 5 to 7 cm (2.0 to 2.8 in) long, in the family Osmeridae. Endemic to the upper Sacramento-San Joaquin Estuary of California, it mainly inhabits the...



Dendrochirus biocellatus

Dendrochirus biocellatus

Dendrochirus biocellatus, the twospot turkeyfish, twinspot lionfish, twoeyed lionfish or ocellated lionfish, is a species of marine ray-finned fish belonging to the family Scorpaenidae, the scorpionfishes and lionfishes. This species is...



Dendrochirus brachypterus

Dendrochirus brachypterus

Dendrochirus brachypterus, the dwarf lionfish, short-finned turkeyfish, shortspine rockcod or shortspine scorpionfish, is a species of marine ray-finned fish belonging to the family Scorpaenidae, the scorpionfishes and lionfishes. It is found in...



Dendrochirus zebra

Dendrochirus zebra

Dendrochirus zebra, known commonly as the zebra turkeyfish or zebra lionfish among other vernacular names, is a species of marine fish in the family Scorpaenidae. The zebra turkeyfish is widespread throughout the tropical waters of the Indo-West...



Denise's pygmy seahorse

Hippocampus denise

Hippocampus denise, also known as Denise's pygmy seahorse or the yellow pygmy seahorse, is a seahorse of the family Syngnathidae native to the western Pacific.



Denison barb

Sahyadria denisonii

The Denison barb, Denison's barb, Miss Kerala, red-line torpedo barb, or roseline shark (*Sahyadria denisonii*) is an endangered species of cyprinid fish endemic to the fast-flowing hill streams and rivers of the Western Ghats in India. It is...



Denticle herring

Denticeps clupeoides

The denticle herring (Denticeps clupeoides) is a small, up to 15 cm (5.9 in) long, species of ray-finned fish found only in the rivers of Benin, Nigeria, and western Cameroon. It is related to the herrings, but notable for its large anal fin and...



Desert pupfish

Cyprinodon macularius

The desert pupfish (Cyprinodon macularius) is a rare species of bony fish in the family Cyprinodontidae. It is a small fish, typically less than 7.62 cm (3 in) in length. Males are generally larger than females, and have bright-blue coloration,...



Desert sucker

Catostomus clarkii

The desert sucker or Gila Mountain sucker (Catostomus clarkii), is a freshwater species of ray-finned fish in the sucker family, endemic to the Great Basin and the Colorado River Basin in the United States. It inhabits rapids and fast-flowing...



Desmodema polystictum

Desmodema polystictum

Desmodema polystictum, also called the deal fish, polka-dot ribbonfish, or spotted ribbonfish, is a fish in the family Trachipteridae. It is found near New Zealand, the northwestern Atlantic Ocean, and South Africa. The species became more known...



Desmopuntius rhomboocellatus

Desmopuntius rhomboocellatus

Desmopuntius rhomboocellatus, the snakeskin barb, is a species of cyprinid fish endemic to Borneo where it is found in western and central Kalimantan. This species reaches a length of 8.8 centimetres (3.5 in) TL. The specific rhomboocellatus...



Devil fish

Mobula mobular

The devil fish or giant devil ray (Mobula mobular) is a species of ray in the family Mobulidae. It is currently listed as endangered, mostly due to bycatch mortality in unrelated fisheries.



Devils Hole pupfish

Cyprinodon diabolis

The Devils Hole pupfish (Cyprinodon diabolis) is a critically endangered species of the family Cyprinodontidae (pupfishes) found only in Devils Hole, a water-filled cavern in the US state of Nevada. It was first described as a species in 1930 and...



Diamond darter

Crystallaria cincotta

The diamond darter (Crystallaria cincotta) is a species of freshwater ray-finned fish, a darter from the subfamily Etheostominae, part of the family Percidae, which also contains the perches, ruffes and pikeperches. It is named for the...



Diamond lizardfish

Synodus synodus

The diamond lizardfish (Synodus synodus) is a species of lizardfish that lives in tropical and subtropical waters of the Atlantic Ocean.



Diamond stingray

Dasyatis dipterura

The diamond stingray (Hypanus dipterurus) is a species of stingray in the family Dasyatidae. It is found in the coastal waters of the eastern Pacific Ocean from southern California to northern Chile, and around the Galápagos and Hawaiian Islands....



Diamond tetra

Moenkhausia pittieri

The diamond tetra (Moenkhausia pittieri) is a small freshwater fish of the characin family (family Characidae) of order Characiformes. It is found in and around Lake Valencia in Venezuela, South America.



Diana's hogfish

Bodianus diana

Diana's hogfish, *Bodianus diana*, is a species of wrasse native to the Indian Ocean from the African coast to the Nicobars and the Cocos-Keeling Islands. Reports of its presence in the western Pacific Ocean are erroneous. It occurs on the seaward...



Dichotomyctere nigroviridis

Dichotomyctere nigroviridis

Dichotomyctere nigroviridis (syn. *Tetraodon nigroviridis*) or the green spotted puffer is a species of pufferfish. It is found across South and Southeast Asia in coastal freshwater, but survives the longest in brackish to saltwater, and brackish...



Dichotomyctere ocellatus

Dichotomyctere ocellatus

Dichotomyctere ocellatus (syn. Tetraodon biocellatus), commonly the figure 8 puffer or eyespot puffer, is a pufferfish found in freshwater in Southeast Asia. It is known from the lower reaches of the Mekong (Cambodia), the Peninsular Malaysia as...



Dichotomyctere sabahensis

Dichotomyctere sabahensis

Dichotomyctere sabahensis is a species of pufferfish endemic to the southeast Asian island of Borneo. This species grows to a standard length of 10.8 cm (4.3 in).It is also called the giant spotted puffer or the Saba puffer. The juvenile looks...



Dicrossus filamentosus

Dicrossus filamentosus

Dicrossus filamentosus is a species of dwarf cichlid fish. It occurs in the Rio Negro and Orinoco basins of Colombia, Venezuela, and Brazil.These fish lays eggs on leaves or directly on the substrate. The female takes care of the fry alone, since...



Diplodus cervinus

Diplodus cervinus

Diplodus cervinus, the zebra sea bream, is a species of marine ray-finned fish belonging to the family Sparidae, which includes the seabreams and porgies. This species is found in the temperate north eastern Atlantic Ocean and the Mediterranean Sea.



Diplodus sargus

Diplodus sargus

Diplodus sargus, the white seabream or sargo is a species of marine ray-finned fish belonging to the family Sparidae, which includes the seabreams and porgies. This fish is found in the eastern Atlantic Ocean and in the Mediterranean Sea. It is a...



Diplodus vulgaris

Diplodus vulgaris

Diplodus vulgaris, the common two-banded sea bream, is a species of marine ray-finned fish belonging to the family Sparidae, which includes the seabreams and porgies. This species is found in the northeastern Atlantic Ocean and the Mediterranean....



Diploprion bifasciatum

Diploprion bifasciatum

Diploprion bifasciatum, the barred soapfish, also known as the doublebanded soapfish, two banded grouper, two-banded sea perch, two-banded soapfish, yellow emperor or yellow striped grouper, is a species of marine ray-finned fish, related to the...



Dog snapper

Lutjanus jocu

The dog snapper (Lutjanus jocu), also known as the dogtooth snapper, pargue or snuggetooth snapper, is a species of marine ray-finned fish, a snapper belonging to the family Lutjanidae. It is native to the Atlantic Ocean. It is a commercially...



Dogtooth tuna

Gymnosarda unicolor

The dogtooth tuna (Gymnosarda unicolor), also known as white tuna, is a species of pelagic marine fish which belongs to the family Scombridae.



Dollar sunfish

Lepomis marginatus

The dollar sunfish (Lepomis marginatus) is a species of freshwater fish in the sunfish family (family Centrarchidae) of order Perciformes. It is categorized as a warm water pan-fish. Early settlers said that this species of sunfish resembled a...



Dolly Varden trout

Salvelinus malma

The Dolly Varden trout (Salvelinus malma) is a species of salmonid ray-finned fish native to cold-water tributaries of the Pacific Ocean in Asia and North America. Despite the name "trout" (which typically refers to freshwater species from the...



Donets ruffe

Gymnocephalus acerina

The Donets ruffe (Gymnocephalus acerina) is a species of perch native to eastern Europe where it occurs in the basins of the Black Sea and the Sea of Azov. They inhabit freshwater lakes and rivers usually over hard compacted sand. They are...



Dorab wolf-herring

Chirocentrus dorab

The dorab wolf-herring (Chirocentrus dorab) is a fish species from the genus Chirocentrus of the family Chirocentridae. It is a coastal fish, silvery below and bright blue above. It is found in both marine and brackish or estuarine waters,...



Dormitator latifrons

Dormitator latifrons

Dormitator latifrons, the Pacific fat sleeper, is a species of fish in the family Eleotridae found on the Pacific coast of the Americas from around Palos Verdes, California, to Peru, where it can be found in stagnant or sluggish fresh or brackish...



Dormitator maculatus

Dormitator maculatus

The fat sleeper (Dormitator maculatus) is a species of fish belonging to the family Eleotridae, known for their flat heads; they are generally found in fresh water, usually found in lakes, ponds and rivers.



Doryrhamphus excisus

Doryrhamphus excisus

Doryrhamphus excisus is a species of flagtail pipefish from the genus Doryrhamphus. Its common names include blue-striped pipefish and blue-and-orange cleaner pipefish. The fish is found throughout much of the Indo-Pacific and tropical East Pacific.



Double-lined mackerel

Grammatorcynus bilineatus

The double-lined mackerel (Grammatorcynus bilineatus), is a species of Spanish mackerel (tribe Scomberomorini) in the family Scombridae. This species is sometimes also called the scad mackerel.



Doubleband surgeonfish

Acanthurus tennenti

The doubleband surgeonfish or lieutenant tang (*Acanthurus tennenti*), is a marine ray-finned fish in the family Acanthuridae. It is found in the tropical and sub-tropical Indo-Pacific region and grows to a maximum length of 31 cm (12 in).



Doublespotted queenfish

Scomberoides lysan

The doublespotted queenfish (*Scomberoides lysan*) is a tropical game fish in family Carangidae (jacks). It is associated with reefs and ranges widely throughout the Indian and Pacific Oceans. Other common names for this fish are giant dart,...



Dracula fish

Danionella dracula

The dracula fish (Danionella dracula) is a species of tropical danionin fish from the cyprinid family. It is a freshwater fish endemic to Myanmar. A close relative is Danio rerio, the zebrafish of aquariums. It is named dracula after its unusual...



Draughtsboard shark

Cephaloscyllium isabellum

The draughtsboard shark (Cephaloscyllium isabellum) is a species of catshark, and part of the family Scyliorhinidae, so named for its "checkerboard" colour pattern of dark blotches. It is endemic to New Zealand, where it is also known as the...



Drepane punctata

Drepane punctata

Drepane punctata, the butterfish, concertinafish, jetto, peppercorn, sickle-fish, silver moonfish, spotted batfish, spotted sicklefish, sicklefish or spotted spadefish, is a species of ray-finned fish belonging to the family Drepaneidae, the...



Dumb gulper shark

Centrophorus harrissoni

Sharks portalThe dumb gulper shark (Centrophorus harrissoni) is a rare and endangered deepwater dogfish, found along the east coast of Australia and in isolated spots north and west of New Zealand. It is also known as the dumb shark, Harrison's...



Dusky morwong

Dactylophora nigricans

The dusky morwong (*Dactylophora nigricans*) is a species of marine ray-finned fish, traditionally regarded as belonging to the family Cheilodactylidae, the members of which are commonly known as morwongs. It is native to the western and southern...



Dusky shark

Carcharhinus obscurus

The dusky shark (*Carcharhinus obscurus*) is a species of requiem shark, in the family Carcharhinidae, occurring in tropical and warm-temperate continental seas worldwide. A generalist apex predator, the dusky shark can be found from the coast to...



Dusky smooth-hound

Mustelus canis

The dusky smooth-hound (*Mustelus canis*), also called the smooth dogfish or the dog shark, is a species of houndshark in the family Triakidae. This shark is an olive grey or brown in color, and may have shades of yellow or grayish white. Females...



Dusky spinefoot

Siganus luridus

The dusky spinefoot (*Siganus luridus*), also known the squaretail rabbitfish, is a species of marine ray-finned fish, a rabbitfish belonging to the family Siganidae. It is native to the western Indian Ocean which has spread to the Mediterranean Sea...



Dusky tripletooth goby

Tridentiger obscurus

The dusky tripletooth goby (*Tridentiger obscurus*) is a species of goby native to marine, fresh and brackish waters along the coasts of Korea and Japan. This species can reach a length of 14 centimetres (5.5 in) TL. This species is of minor...



Dwarf black stingray

Hemitrygon parvonigra

The dwarf black stingray (Hemitrygon parvonigra) is a little-known species of stingray in the family Dasyatidae, found off northwestern Australia and perhaps throughout Southeast Asia at depths of 60–185 m (197–607 ft). Growing to a width of 51...



Dwarf corydoras

Corydoras hastatus

The dwarf corydoras (Corydoras hastatus), dwarf catfish, tail spot pygmy catfish, or micro catfish is a tropical freshwater fish belonging to the subfamily Corydoradinae of the family Callichthyidae. It originates in inland waters in South...



Dwarf gourami

Trichogaster lalius

The dwarf gourami (Trichogaster lalius) is a species of gourami native to South Asia.



Dwarf gulper shark

Centrophorus atromarginatus

Sharks portalThe dwarf gulper shark (Centrophorus atromarginatus) is a dogfish of the family Centrophoridae found in the Indo-West Pacific oceans, from the Gulf of Aden, Japan, Taiwan, and northern Papua New Guinea.As a Squaliform, Centrophorus...



Dwarf hawkfish

Cirrhichthys falco

The dwarf hawkfish, (Cirrhitichthys falco), is a small species of hawkfish found on tropical reefs in the Indo-Pacific region from the Maldives to the Caroline Islands and Samoa. It can sometimes be found in the aquarium trade.



Dwarf lanternshark

Etmopterus perryi

The dwarf lanternshark (Etmopterus perryi) is a species of dogfish shark in the family Etmopteridae and is the smallest shark in the world, reaching a maximum known length of 20 cm (8 in). It is known to be present only on the upper continental...



Dwarf loach

Ambastaia sidthimunki

The dwarf loach, ladderback loach, pygmy loach, chain loach or chain botia (Ambastaia sidthimunki) is a freshwater fish belonging to the family Botiidae. Formerly included in the genus Yasuhikotakia, it is frequently seen in the aquarium trade,...



Dwarf pufferfish

Carinotetraodon travancoricus

The dwarf pufferfish (Carinotetraodon travancoricus), also known as the Malabar pufferfish, pygmy pufferfish, or pea pufferfish, is a small freshwater pufferfish endemic to Kerala and southern Karnataka in Southwest India. They are popular in...



Dwarf pygmy goby

Pandaka pygmaea

The dwarf pygmy goby or Philippine goby (Pandaka pygmaea) is a tropical species of fish in the subfamily Gobionellinae from brackish water and mangrove areas in Southeast Asia. It is one of the smallest fish species in the world. Males reach...



Dwarf rainbowfish

Melanotaenia praecox

The dwarf rainbowfish (Melanotaenia praecox) is a species of rainbowfish in the subfamily Melanotaeniinae. It is known under a number of common names including diamond rainbowfish, neon rainbowfish, Praecox rainbowfish, dwarf neon rainbowfish,...



Dwarf rasbora

Boraras maculatus

The dwarf rasbora (Boraras maculatus) is a species of ray-finned fish in the genus Boraras, native to freshwater habitats of southeast Asia. It grows to be about 10-20mm (0.4-0.8 inches) long at adulthood.



Dwarf sawfish

Pristis clavata

The dwarf sawfish or Queensland sawfish, Pristis clavata, is a sawfish of the family Pristidae, found in tropical Australia. This endangered species is the smallest species in its family.



Dwarf seahorse

Hippocampus zosterae

The dwarf seahorse (*Hippocampus zosterae*) is a species of seahorse found in the subtidal aquatic beds of the Bahamas and parts of the United States. It is threatened by habitat loss. According to Guinness World Records, it is the slowest-moving...



Dwarf sturgeon

Pseudoscaphirhynchus hermanni

The dwarf sturgeon, little shovelnose sturgeon, or small Amu-Darya shovelnose sturgeon (Pseudoscaphirhynchus hermanni) is a species of fish in the family Acipenseridae. It is found in Turkmenistan, Uzbekistan and probably in Tajikistan.



Dysommia rugosa

Dysommia rugosa

Dysommia rugosa is an eel in the family Synphobranchidae (cutthroat eels). It was described by Isaac Ginsburg in 1951. It is a marine, deep water-dwelling eel which is known from the western Atlantic and eastern central Pacific Ocean. It dwells...



Eagle Lake trout

Oncorhynchus mykiss aquilarum

Eagle Lake trout, Oncorhynchus mykiss aquilarum, is a subspecies of rainbow trout endemic to Eagle Lake, in Lassen County, California. It is a type of trout known for its ability to withstand high alkalinity. Its unique adaptations to the harsh...



East Atlantic peacock wrasse

Symphodus tinca

The East Atlantic peacock wrasse (Symphodus tinca) is a species of wrasse native to the eastern Atlantic Ocean from Spain to Morocco and in the coastal waters of the Mediterranean Sea and the Black Sea. This species lives around rocks surrounded...



Eastern blacknose dace

Rhinichthys atratulus

Eastern blacknose dace (Rhinichthys atratulus) is a species of ray-finned fish in the genus Rhinichthys. Its name originates from the Old French word "dars" which is the nominative form of the word "dart" in reference to their swimming pattern....



Eastern freshwater cod

Maccullochella ikei

The eastern freshwater cod (Maccullochella ikei), also known as the eastern cod or Clarence River cod, is a large predatory freshwater fish of the genus Maccullochella and the family Percichthyidae, that occur in the coastal Clarence River system...



Eastern mosquitofish

Gambusia holbrooki

The eastern mosquitofish (Gambusia holbrooki) is a species of freshwater fish, closely related to the western mosquitofish, Gambusia affinis. It is a member of the family Poeciliidae of order Cyprinodontiformes. The eastern mosquitofish is native...



Eastern mudminnow

Umbra pygmaea

The eastern mudminnow (Umbra pygmaea) is a species of freshwater fish belonging to the family Umbridae. It is native to eastern North America, but has been introduced to Europe. It feeds primarily on insect larvae and other small aquatic...



Eastern nannygai

Centroberyx affinis

The eastern nannygai (Centroberyx affinis), also known as the redfish, bight redfish, red snapper, golden snapper or koarea, is an alfoncino of the genus Centroberyx. It is found around Australia and New Zealand at depths between 10 and 450...



Eastern rainbowfish

Melanotaenia splendida splendida

Melanotaenia splendida splendida, also known as the eastern rainbowfish, is a subspecies of fish in the family Melanotaeniidae endemic to Australia. The Eastern rainbowfish is widespread on the mainland, and is found in river systems east of the...



Eastern sand darter

Ammocrypta pellucida

The eastern sand darter (Ammocrypta pellucida) is a species of freshwater ray-finned fish, a darter from the subfamily Etheostomatinae, part of the family Percidae, which also contains the perches, ruffes and pikeperches. The eastern sand darter...



Echeneis neucratoides

Echeneis neucratoides

The whitefin sharksucker (Echeneis neucratoides) or short-disk sharksucker, is a species of remora native to subtropical waters of the western Atlantic Ocean, Gulf of Mexico and Caribbean Sea. This fish can reach a length of 75 centimetres (30...



Ecsenius bicolor

Ecsenius bicolor

Ecsenius bicolor, commonly known as the flame tail blenny or bicolor blenny, is a blenny from the Indo-Pacific. It frequently makes its way into the aquarium trade. It grows to a size of 11 centimetres (4.3 in) in length.



Ecsenius stigmatura

Ecsenius stigmatura

Ecsenius stigmatura, commonly known as the tail-spot blenny, is a blenny from the Western Pacific. It occasionally makes its way into the aquarium trade. A coppery-coloured fish with a distinct blackish spot at the base of the tail. It has some...



Ecsenius yaeyamaensis

Ecsenius yaeyamaensis

Ecsenius yaeyamaensis, known commonly as the Yaeyama blenny in Guam and Micronesia, and also known as the Yaeyama clown blenny in Micronesia, or the Pale-spotted combtooth-blenny in Indonesia is a species of combtooth blenny in the genus...



Eel catfish

Channallabes apus

The eel catfish (Channallabes apus) is an airbreathing catfish found in the muddy swamps of the tropics of Central Africa. It grows up to 32.7 cm in total length (12.9 in), and is notable for its ability to propel itself out of the water to catch...



Eel-tailed catfish

Tandanus tandanus

The eel-tailed catfish, Tandanus tandanus, is a species of catfish (order Siluriformes) of the family Plotosidae. This fish is also known as dewfish, freshwater catfish, jewfish, and tandan. This species is a freshwater fish native to the...



Egyptian mouthbrooder

Pseudocrenilabrus multicolor

The Egyptian mouthbrooder (Pseudocrenilabrus multicolor) is a species of cichlid. This small mouthbrooder reaches about 8 cm (3.1 in) in length, and is found in rivers, lakes, and other freshwater habitats in Eastern Africa from Egypt and as far...



Egyptian sole

Solea aegyptiaca

The Egyptian sole (Solea aegyptiaca) is a species of flatfish in the true sole family, Soleidae. It lives on the sandy or muddy seabed of the Mediterranean Sea, and is now colonising the Red Sea. It often semi-immerses itself in the substrate....



Elacatinus evelynae

Elacatinus evelynae

Elacatinus evelynae, commonly known as the sharknose goby, Caribbean cleaner goby, or Caribbean cleaning goby, is a species of goby native to the Western Atlantic Ocean from the Bahamas and the Lesser Antilles to the northern coast of South...



Elassoma evergladei

Elassoma evergladei

Elassoma evergladei, or the Everglades pygmy sunfish, is a species of fish from the genus Elassoma (the pygmy sunfishes) that is endemic to North America.



Electric blue hap

Sciaenochromis ahli

The electric blue hap (Sciaenochromis ahli) is a species of cichlid fish endemic to Lake Malawi. It prefers to live in caves and crevices in rocky substrates. This species can reach a length of 20 centimetres (7.9 in) TL. It can also be found in...



Electrophorus electricus

Electrophorus electricus

Electrophorus electricus is the best-known species of electric eel. It is a South American electric fish. Until the discovery of two additional species in 2019, the genus was classified as the monotypic, with this species the only one in the...



Elegant corydoras

Corydoras elegans

The elegant corydoras or elegant catfish (Corydoras elegans) is a tropical freshwater fish belonging to the subfamily Corydoradinae of the family Callichthyidae. It originates in inland waters in South America, and is found in the Upper Amazon...



Elegant firefish

Nemateleotris decora

Nemateleotris decora, the elegant firefish or purple firefish, is a species of dartfish native to tropical waters in the Indo-West Pacific, from Mauritius to Samoa, north to Ryukyu Islands, and south to New Caledonia, from depths ranging between...



Eleotris sandwicensis

Eleotris sandwicensis

Eleotris sandwicensis, the Sandwich Island sleeper, Hawaiian sleeper or oopu, is a species of fish in the family Eleotridae endemic to the Hawaiian Islands, where it can be found in marine, fresh, and brackish waters around the coast. Due to this...



Elephant ear gourami

Osphronemus exodon

The elephant ear gourami (Osphronemus exodon) is a large species of gourami native to the Mekong river basin in Thailand, Laos, and Cambodia.



Eleutheronema tetradactylum

Eleutheronema tetradactylum

Eleutheronema tetradactylum, the fourfinger threadfin, known as ranwas in Pakistan, is a species of marine ray-finned fish, a threadfin from the family Polynemidae which occurs in the Indian and western Pacific Ocean.



Elongate glassy perchlet

Chanda nama

The elongate glassy perchlet (Chanda nama) is a species of freshwater fish in the Asiatic glassfish family Ambassidae of order Perciformes, the only species in the genus Chanda. It is native to an area of south Asia from Pakistan to Burma, in the...



Elopichthys bambusa

Elopichthys bambusa

Elopichthys bambusa, the yellowcheek or kanyu (Chinese: 鲢鱼; pinyin: gǎnyú), is a large cyprinid fish that is found in freshwater habitats in eastern Asia. It ranges from the Amur River in Russia, through China to the Red and Lam Rivers in...



Elops hawaiiensis

Elops hawaiiensis

The Hawaiian ladyfish (Elops hawaiiensis), also known as the Hawaiian tenpounder or banana fish, is a species of ray-finned fish in the family Elopidae. It is sometimes referred to as the giant herring, though it is not closely related to the true...



Elops machnata

Elops machnata

Elops machnata, the tenpounder, is a species of ray-finned fish in the family Elopidae in the order Elopiformes (tarpons and tenpounders). This species is found in coastal regions of the Indian Ocean.



Elops saurus

Elops saurus

The ladyfish or tenpounder (Elops saurus) is a species of fish in the genus Elops, the only genus in the monotypic family Elopidae.



Ember parrotfish

Scarus rubroviolaceus

The ember parrotfish (Scarus rubroviolaceus) is a species of marine ray-finned fish, a parrotfish, in the family Scaridae. It is native to the Indian and Pacific Oceans. It is also known as the bicolor parrotfish and the redlip parrotfish.



Ember tetra

Hyphessobrycon amandae

The ember tetra (Hyphessobrycon amandae) is a freshwater fish of the characin family (family Characidae) of order Characiformes. It is native to the Araguaia River basin of Brazil and was discovered in 1987 and named in honor of the fish explorer...



Embiotoca jacksoni

Embiotoca jacksoni

Embiotoca jacksoni, commonly known as the black surfperch, is a species of surfperches native to shallow coastal areas of the eastern Pacific. Other common names of the species include black perch and butterlips. They are usually a dark reddish...



Embiotoca lateralis

Embiotoca lateralis

Embiotoca lateralis, commonly known as the striped surfperch or striped seaperch, is a species of surfperch native to the north-eastern Pacific Ocean.



Emerald catfish

Corydoras splendens

The emerald catfish (Corydoras splendens) is a tropical freshwater fish belonging to the Corydoradinae sub-family of the family Callichthyidae native to the Amazon Basin in South America. It has traditionally been known as Brochis splendens. The...



Emerald rockcod

Trematomus bernacchii

The emerald rockcod (Trematomus bernacchii), also known as the emerald notothen is a species of marine ray-finned fish belonging to the family Nototheniidae, the notothen or cod icefishes. It is native to the Southern Ocean where it is a...



Emerald shiner

Notropis atherinoides

The emerald shiner (Notropis atherinoides) is one of hundreds of small, silvery, slender fish species known as shiners. The identifying characteristic of the emerald shiner is the silvery emerald color on its sides. It can grow to 3.5 inches in...



Emmelichthys nitidus

Emmelichthys nitidus

Emmelichthys nitidus, the Cape bonnetmouth, bonnetmouth, redbait, pearl fish, picarel, red baitfish, red herring or Southern rover, is a species of marine ray-finned fish belonging to the family Emmelichthyidae, the rovers and bonnetmouths. This...



Emperor angelfish

Pomacanthus imperator

The emperor angelfish (Pomacanthus imperator) is a species of marine angelfish. It is a reef-associated fish, native to the Indian and Pacific Oceans, from the Red Sea to Hawaii and the Austral Islands. This species is generally associated with...



Enchelycore schismatorhynchus

Enchelycore schismatorhynchus

Enchelycore schismatorhynchus is a moray eel found in coral reefs in the Pacific and Indian Oceans. It was first named by Bleeker in 1853, and is commonly known as the white-margined moray, brown moray eel, or the funnel-nostril moray.



English sole

Parophrys vetulus

The English sole (Parophrys vetulus) is a flatfish of the family Pleuronectidae. It is a demersal fish that lives on sandy and muddy bottoms in estuaries and near shore areas, at depths of up to 550 metres (1,800 ft). It reaches up to 57...



Enneacampus ansorgii

Enneacampus ansorgii

Enneacampus ansorgii, the African freshwater pipefish, is a pipefish in the family Syngnathidae (pipefishes and seahorses). It is widely distributed in coastal rivers and streams of Western Africa, being found in both slow and fast flowing water...



Epaulette shark

Hemiscyllium ocellatum

The epaulette shark (, Hemiscyllium ocellatum) is a species of longtailed carpet shark of the family Hemiscylliidae, found in shallow, tropical waters off Australia and New Guinea (and possibly elsewhere). The common name of this shark comes from...



Epigonus telescopus

Epigonus telescopus

Epigonus telescopus, the black cardinal fish, is a species of deepwater cardinalfish found in most temperate oceans worldwide, at depths of between 75 and 1,200 metres (246 and 3,937 ft) though mostly between 300 and 800 metres (980 and 2,620...



Epinephelus andersoni

Epinephelus andersoni

Epinephelus andersoni, the catface grouper, brown-spotted grouper, catface rockcod or brown spotted rockcod, is a species of marine ray-finned fish, a grouper from the subfamily Epinephelinae which is part of the family Serranidae, which also...



Epinephelus bruneus

Epinephelus bruneus

Epinephelus bruneus, commonly known as the longtooth grouper or the kelp grouper, is a species of marine ray-finned fish, a grouper from the subfamily Epinephelinae which is part of the family Serranidae, which also includes the anthias and sea...



Epinephelus corallicola

Epinephelus corallicola

Epinephelus corallicola, also known as the black-dotted cod, coral cod, coral rockcod or duskyfin grouper, is a species of marine ray-finned fish, a grouper from the subfamily Epinephelinae which is part of the family Serranidae, which also...



Epinephelus flavocaeruleus

Epinephelus flavocaeruleus

Epinephelus flavocaeruleus, commonly called blue-and-yellow grouper, is a species of marine ray-finned fish, a grouper from the subfamily Epinephelinae which is part of the family Serranidae, which also includes the anthias and sea basses. It is...



Epinephelus marginatus

Epinephelus marginatus

Epinephelus marginatus (.mw-parser-output.IPA-label-small{font-size:85%}.mw-parser-output.references.IPA-label-small,.mw-parser-output.infobox.IPA-label-small,.mw-parser-output.navbox.IPA-label-small{font-size:100%}Latin pronunciation:), the...



Epinephelus sexfasciatus

Epinephelus sexfasciatus

Epinephelus sexfasciatus, the sixbar grouper, sixbar rockcod or six-banded rockcod, is a species of marine ray-finned fish, a grouper from the subfamily Epinephelinae which is part of the family Serranidae, which also includes the anthias and sea...



Epinephelus summana

Epinephelus summana

Epinephelus summana, the summan grouper, is a species of marine ray-finned fish, a grouper from the subfamily Epinephelinae which is part of the family Serranidae, which also includes the anthias and sea basses. It comes from the Eastern Indian...



Eritrean butterflyfish

Chaetodon paucifasciatus

The Eritrean butterflyfish or crown butterflyfish (Chaetodon paucifasciatus) is a species of marine ray-finned fish, a butterflyfish belonging to the family Chaetodontidae. It is essentially just known from the Red Sea and the Gulf of Aden, but...



Erythrinus erythrinus

Erythrinus erythrinus

Erythrinus erythrinus, the red wolf fish, is a relatively small species of trahira (family Erythrinidae) from freshwater habitats in South America.



Escolar

Lepidocybium flavobrunneum

The escolar, *Lepidocybium flavobrunneum*, a species of fish in the family Gempylidae, is found in deep (200–885 metres, or 656–2,904 ft) tropical and temperate waters around the world. It is also known as snake mackerel, walu walu (Hawaiian,...



Escualosa thoracata

Escualosa thoracata

The white sardine (Escualosa thoracata) is a species of fish in the family Clupeidae. It was described by Achille Valenciennes in 1847. It is a tropical fish of the Indo-Pacific distributed from Thailand to Indonesia and Australia. Other common...



Esox cisalpinus

Esox cisalpinus

Esox cisalpinus, the southern pike or cisalpine pike, is a species of freshwater fish known from central and northern Italy, southeastern France and Switzerland, and it might also occur in the western Balkans. It has traditionally been considered...



Estuary perch

Macquaria colonorum

The estuary perch (Macquaria colonorum) is a species of temperate perch endemic to south-eastern Australia, where it prefers brackish waters such as lower tidal reaches of coastal lakes, rivers, and streams.



Estuary stingray

Hemitrygon fluviorum

The estuary stingray (Hemitrygon fluviorum), also called the estuary stingaree or brown stingray, is a species of stingray in the family Dasyatidae. Endemic to eastern Australia, it typically inhabits shallow, mangrove-lined tidal rivers,....



Etheostoma exile

Etheostoma exile

Etheostoma exile, the Iowa darter, is a species of freshwater ray-finned fish, a darter from the subfamily Etheostomatinae, part of the family Percidae, which also contains the perches, ruffes and pikeperches. It is, along with about 17 other...



Etheostoma obama

Etheostoma obama

Etheostoma obama, the spangled darter, is a species of freshwater ray-finned fish, a darter from the subfamily Etheostomatinae, part of the family Percidae, which also contains the perches, ruffes and pikeperches. It is endemic to the eastern...



Etheostoma osburni

Etheostoma osburni

Etheostoma osburni, the candy darter or finescale saddled darter, is a species of fish in the family Percidae, a member of the group known as darters. This species is endemic to the eastern United States where it is known only from the Kanawha...



Etheostoma zonale

Etheostoma zonale

Etheostoma zonale, the banded darter, is a species of freshwater ray-finned fish, a darter from the subfamily Etheostomatinae, part of the family Percidae, which also contains the perches, ruffes and pikeperches. It is endemic to the eastern...



Etrumeus sadina

Etrumeus sadina

Etrumeus sadina, the red-eye round herring, is a species of fish belonging to the family Dussumieriidae, a family notable for their rounded bellies.



Eulachon

Thaleichthys pacificus

The eulachon (*Thaleichthys pacificus*), also spelled oolichan, ooligan, hooligan), or the candlefish, is a small anadromous species of smelt that spawns in some of the major river systems along the Pacific coast of North America from northern...



Eumecichthys

Eumecichthys fiski

Eumecichthys fiski, the unicorn crestfish or unicornfish, is a very rare, little-known species of crestfish in the family Lophotidae, and the only member of the genus Eumecichthys. It likely has a worldwide distribution, having been first...



Eurasian carp

Cyprinus carpio

The Eurasian carp or European carp (Cyprinus carpio), widely known as the common carp, is a widespread freshwater fish of eutrophic waters in lakes and large rivers in Europe and Asia. The native wild populations are considered vulnerable to...



European anchovy

Engraulis encrasicolus

The European anchovy (Engraulis encrasicolus) is a forage fish somewhat related to the herring. It is a type of anchovy; anchovies are placed in the family Engraulidae. It lives off the coasts of Europe and Africa, including in the Mediterranean...



European bitterling

Rhodeus amarus

The European bitterling (Rhodeus amarus) is a temperate freshwater fish belonging to the subfamily Acheilognathinae of the family Cyprinidae. It originates in Europe, ranging from the Rhone River basin in France to the Neva River in Russia. It...



European bullhead

Cottus gobio

The European bullhead (*Cottus gobio*) is a freshwater fish that is widely distributed in Europe, mainly in rivers. It is a member of the family Cottidae, a type of sculpin. It is also known as the miller's thumb, freshwater sculpin, common...



European conger

Conger conger

The European conger (*Conger conger*) is a species of conger of the family Congridae. It is the heaviest eel in the world and native to the northeast Atlantic, including the Mediterranean Sea.



European eel

Anguilla anguilla

The European eel (*Anguilla anguilla*) is a species of eel. They are critically endangered due to overfishing by fisheries on coasts for human consumption and parasites.



European flounder

Platichthys flesus

The European flounder (*Platichthys flesus*) is a flatfish of European coastal waters from the White Sea in the north to the Mediterranean and the Black Sea in the south. It has been introduced into the United States and Canada accidentally through...



European mudminnow

Umbra krameri

The European mudminnow (*Umbra krameri*) is a species of fish in the Umbridae family found in Austria, Bosnia and Herzegovina, Bulgaria, Croatia, the Czech Republic, Hungary, Moldova, Romania, Serbia, Montenegro, Slovakia, Slovenia, and Ukraine.



European perch

Perca fluviatilis

The European perch (Perca fluviatilis), also known as the common perch, redfin perch, big-scaled redfin, English perch, Euro perch, Eurasian perch, Eurasian river perch, Hatch, poor man's rockfish or in Anglophone parts of Europe, simply the...



European plaice

Pleuronectes platessa

The European plaice (Pleuronectes platessa), commonly referred to as simply plaice, is a species of marine flatfish in the genus Pleuronectes of the family Pleuronectidae.



European sea sturgeon

Acipenser sturio

The European sea sturgeon (Acipenser sturio), also known as the Atlantic sturgeon or common sturgeon, is a species of sturgeon native to Europe. It was formerly abundant, being found in coastal habitats all over Europe. It is anadromous and...



European seabass

Dicentrarchus labrax

The European seabass (Dicentrarchus labrax), also known as the European bass, sea bass, common bass, white bass, capemouth, white salmon, sea perch, white mullet, sea dace or Loup de Mer, is a primarily ocean-going fish native to the waters off...



European smelt

Osmerus eperlanus

The smelt or European smelt (Osmerus eperlanus) is a species of fish in the family Osmeridae.



European sprat

Sprattus sprattus

The European sprat (Sprattus sprattus), also known as bristling, brisling, garvie, garvock, Russian sardine, russlet, skipper or whitebait, is a species of small marine fish in the herring family Clupeidae. Found in European waters, it has silver...



Euthynnus affinis

Euthynnus affinis

Euthynnus affinis, the mackerel tuna, little tuna, wavyback skipjack tuna, kawakawa, or tongkol komo is a species of ray-finned bony fish in the family Scombridae, or mackerel family. It belongs to the tribe Thunnini, better known as the tunas....



Euthynnus lineatus

Euthynnus lineatus

Euthynnus lineatus, the black skipjack tuna or black skipjack, is a species of ray-finned bony fish in the family Scombridae. It belongs to the tribe Thunnini, better known as the tunas.



Eutropiichthys vacha

Eutropiichthys vacha

Eutropiichthys vacha is a species of schilbid catfish native to India, Nepal, Bhutan, and Bangladesh. It can reach a length of 34 cm, and a mass of 1.35 kg. Eutropiichthys vacha is a climate sensitive species that may be selected as a target...



Exquisite wrasse

Cirrhilabrus exquisitus

The exquisite wrasse (Cirrhilabrus exquisitus) is a species of ray-finned fish from the family Labridae, the wrasses, which is native to reefs in the Indo-West Pacific region. It can be found in the aquarium trade.



Exyrias akihito

Exyrias akihito

Exyrias akihito is a species of marine goby. Since the recent description of this species it has been found to have a wide distribution in the western Pacific including the Great Barrier Reef, New Guinea, Indonesia, the Philippines and the...



Eyed flounder

Bothus ocellatus

The eyed flounder (*Bothus ocellatus*) is a species of fish in the family Bothidae (lefteye flounders). The species is found on or near the sandy seabed in relatively shallow waters in the western Atlantic Ocean, the Caribbean Sea and the Gulf of...



Eyespot rasbora

Brevibora dorsiocellata

The eyespot rasbora (Brevibora dorsiocellata) is a small fish belonging to the Family Cyprinidae, subfamily Danioninae, which is known by the common names of ocellated rasbora, hi-spot rasbora, and eye-spot rasbora, an allusion to the marking...



Faceless cusk

Typhlonus nasus

The faceless cusk (Typhlonus nasus) is a genus of cusk-eel found in the Indian and Pacific Oceans at depths from 3,935 to 5,100 m (12,910 to 16,732 ft). This species grows to 46.5 cm (18.3 in) in standard length. The fish is named after its...



Fahaka pufferfish

Tetraodon lineatus

The fahaka pufferfish (from Arabic: فِهْقَة), also known as the Nile puffer, globe fish, lineatus puffer (Tetraodon lineatus), is a tropical freshwater pufferfish found in the upper Nile, Chad, Senegal, Gambia, Geba, Volta and Turkana basins in...



Fallfish

Semotilus corporalis

The fallfish (Semotilus corporalis) is a North American freshwater fish, a chub in the family Cyprinidae. The fallfish is the largest minnow species native to Eastern North America.



False catshark

Pseudotriakis microdon

The false catshark or sofa shark (*Pseudotriakis microdon*) is a species of ground shark in the family Pseudotriakidae, and the sole member of its genus. It has a worldwide distribution, and has most commonly been recorded close to the bottom over...



False cleanerfish

Aspidontus taeniatus

The false cleanerfish (*Aspidontus taeniatus*) is a species of combtooth blenny, a mimic that copies both the dance and appearance of *Labroides dimidiatus* (the bluestreak cleaner wrasse), a similarly colored species of cleaner wrasse. It likely...



False osman

Schizopygopsis stoliczkai

The false osman (*Schizopygopsis stoliczkai*) is an Asian cyprinid freshwater fish living in the highlands of Afghanistan, southwestern China, Iran, northern India, Kyrgyzstan, Pakistan and Tajikistan. Young individuals and small adults live in...



False scad

Caranx rhonchus

The false scad (*Caranx rhonchus*), also known as the spotfin scad, ten-finned horse mackerel and yellow horse mackerel, is a species of medium-sized marine fish classified in the jack family Carangidae. The false scad is distributed throughout the...



False scorpionfish

Centrogenys vaigiensis

The false scorpionfish (*Centrogenys vaigiensis*), also known as prettyfins, is a species of perciform fish, the only species in genus *Centrogenys*, which in turn is the only genus in the family Centrogenyidae. They are pale grey or brown and...



False trevally

Lactarius lactarius

The false trevally (Lactarius lactarius) is a species of fish in the family Lactariidae, currently the sole member of the family.



Fangtooth moray

Enchelycore anatina

The fangtooth moray (Enchelycore anatina) sometimes also known as tiger moray or bird-eye conger is a moray eel of the family Muraenidae found in warmer parts of the eastern Atlantic Ocean, including the Canary Islands, Madeira and various other...



Fantail darter

Etheostoma flabellare

The fantail darter (Etheostoma flabellare) is a species of freshwater ray-finned fish, a darter from the subfamily Etheostomatinae, part of the family Percidae, which also contains the perches, ruffes and pikeperches. It is widely distributed...



Farlowella acus

Farlowella acus

Farlowella acus is the type species of the armored catfish genus Farlowella. Members of this genus are commonly known as twig catfish or whiptail catfish. The specific name of the species, acus, means “pointed” or a “needle” or “spine”.



Fathead minnow

Pimephales promelas

Fathead minnow (Pimephales promelas), also known as fathead or tuffy, is a species of temperate freshwater fish belonging to the genus Pimephales of the cyprinid family. The natural geographic range extends throughout much of North America, from...



Featherfin squeaker

Synodontis eupterus

The featherfin squeaker (Synodontis eupterus) or featherfin synodontis is a species of Synodontis catfish. This species is found in the basins of the White Nile, Volta and Niger Rivers and the Chad Basin. Featherfin squeakers are called such due...



Ferox trout

Salmo ferox

Ferox trout (Salmo ferox) is a variety of trout found in oligotrophic lakes/lochs of Ireland, England, Scotland and Wales. Ferox trout is a traditional name for large, piscivorous trout, which in Scotland feed largely on Arctic char. It has been...



Fimbriated moray

Gymnothorax fimbriatus

The fimbriated moray (Gymnothorax fimbriatus), also known as dark-spotted moray or spot-face moray, is a moray eel of the family Muraenidae.



Finescale dace

Chrosomus neogaeus

The finescale dace (Chrosomus neogaeus) is a species of freshwater fish in the family Cyprinidae of order Cypriniformes. It is native to the northern portions of Minnesota, with relatively smaller populations in northern New York, Vermont, New...



Finetooth shark

Carcharhinus isodon

The finetooth shark (Carcharhinus isodon) is a species of requiem shark, in the family Carcharhinidae, found in the western Atlantic Ocean, from North Carolina to Brazil. It forms large schools in shallow, coastal waters, and migrates seasonally...



Fire eel

Mastacembelus erythrotaenia

The fire eel (Mastacembelus erythrotaenia) is a relatively large species of spiny eel. This omnivorous freshwater fish is native to in Southeast Asia but is also found in the aquarium trade. Although it has declined locally (especially in parts...



Firemouth cichlid

Thorichthys meeki

The firemouth cichlid (Thorichthys meeki) is a species of cichlid fish native to Central America. They occur in rivers of the Yucatán Peninsula, Mexico, south through Belize and into northern Guatemala. Their natural habitat is typically shallow...



Fistularia corneta

Fistularia corneta

Fistularia corneta, commonly known as the Pacific cornetfish or the deepwater cornetfish, is a marine fish in the family Fistulariidae. It is endemic to the eastern Pacific Ocean, being found from California to Peru, including many offshore...



Fistularia tabacaria

Fistularia tabacaria

Fistularia tabacaria, the cornetfish, blue-spotted cornetfish, tobacco trumpetfish or unarmed trumpetfish, is a species of cornetfish found along the Atlantic coasts of the Americas and in the central Atlantic off West Africa and the Macaronesian...



Fiveband barb

Desmopuntius pentazona

The fiveband barb (Desmopuntius pentazona) is a species of cyprinid freshwater fish from Southeast Asia. This species is restricted to blackwater streams and peat swamps in northwestern Borneo and possibly Peninsular Malaysia, but it has often...



Fivebeard rockling

Ciliata mustela

The fivebeard rockling (Ciliata mustela) is a coastal ray-finned fish of the family Lotidae. the lings and rocklings. It is found in the eastern Atlantic Ocean. It is not a fish of any commercial importance.



Flagfish

Jordanella floridae

The flagfish (Jordanella floridae), also known as the American flagfish or Florida flagfish, is a species of pupfish, a type of killifish from the family Cyprinodontidae which is endemic to Florida. It is found in the aquarium trade. Its common...



Flagtail catfish

Dianema urostriatum

The flagtail catfish or stripedtail catfish, *Dianema urostriatum*, is a tropical freshwater fish belonging to the Callichthyinae sub-family of the family Callichthyidae. It originates in the Amazon Basin in South America. The fish will grow in...



Flame angelfish

Centropyge loricula

The flame angelfish (*Centropyge loricula*) is a marine angelfish of the family Pomacanthidae found in tropical waters of the Pacific Ocean. Other common names include flame angel, flaming angelfish and Japanese pygmy angelfish.



Flame hawkfish

Neocirrhites armatus

The flame hawkfish (*Neocirrhites armatus*) is a species of marine ray-finned fish, a hawkfish belonging to the family Cirrhitidae. It is native to tropical reefs of the Pacific Ocean at depths of from 1 to 10 metres (3.3 to 32.8 ft). This species...



Flame tetra

Hyphessobrycon flammeus

The flame tetra (*Hyphessobrycon flammeus*), also known as the red tetra or Rio tetra, is a small freshwater fish of the characin family Characidae. This tetra was first introduced as aquarium fish in 1920 by C. Bruening, Hamburg, Germany, and...



Flameback angelfish

Centropyge aurantonotus

The flameback angelfish (*Centropyge aurantonotus*), also known as the flameback pygmy angelfish, Brazilian flameback angelfish, Caribbean flameback angelfish or fireball angelfish, is a species of marine ray-finned fish, a marine angelfish...



Flapnose houndshark

Scylliogaleus quecketti

Sharks portalThe flapnose houndshark (*Scylliogaleus quecketti*) is a houndshark of the family Triakidae, and the only member of the genus *Scylliogaleus*. It is found in the waters off subtropical South Africa, in the western Indian Ocean between...



Flat needlefish

Ablennes hians

The flat needlefish (*Ablennes hians*), or barred longtom, the only known member of the genus *Ablennes*, is a marine fish of the family Belonidae. Flat needlefish are considered gamefish, frequently caught with the help of artificial lights, but are...



Flathead catfish

Pylodictis olivaris

The flathead catfish (*Pylodictis olivaris*), also called by several common names including mudcat or shovelhead cat, is a large species of North American freshwater catfish in the family Ictaluridae. It is the only species of the genus *Pylodictis*...



Flathead chub

Platygobio gracilis

The flathead chub (Platygobio gracilis) is a species of fish in the carp family, Cyprinidae. It is the only member of the monotypic genus Platygobio. It is native to North America, where it is distributed throughout central Canada and the central...



Flathead grey mullet

Mugil cephalus

The flathead grey mullet (Mugil cephalus) is an important food fish species in the mullet family Mugilidae. It is found in coastal tropical and subtropical waters worldwide. Its length is typically 30 to 75 centimetres (12 to 30 in). It is known...



Flavescent peacock

Aulonocara stuartgranti

The flavescent peacock (Aulonocara stuartgranti), also known as Grant's peacock, is a species of haplochromine cichlid. Its common name refers to its "flavescent" (yellowish) colour. It is endemic to Lake Malawi where found in the countries of in...



Flier

Centrarchus macropterus

The flier (Centrarchus macropterus) is a species of freshwater ray-finned fish, a sunfish from the family Centrarchidae which is endemic to the southern United States of America. It is the only species in the monospecific genus Centrarchus. A...



Florida gar

Lepisosteus platyrhincus

The Florida gar (*Lepisosteus platyrhincus*) is a species of gar found in the US from the Savannah River and Ochlockonee River watersheds of Georgia and throughout peninsular Florida. Florida gar can reach a length over 3 ft (91 cm). The young feed...



Florida pompano

Trachinotus carolinus

The Florida pompano (*Trachinotus carolinus*) is a species of marine fish in the *Trachinotus* (pompano) genus of the family Carangidae. It has a compressed body and short snout; coloration varies from blue-greenish silver on the dorsal areas and...



Flying fox

Epalzeorhynchus kalopterus

The flying fox (*Epalzeorhynchus kalopterus*) is a Southeast Asian species of freshwater fish in the family Cyprinidae. It is commonly kept in the aquarium trade. Among other foods, it is known to eat green algae. It is sometimes mistakenly...



Flying gurnard

Dactylopterus volitans

The flying gurnard (*Dactylopterus volitans*), also known as the helmet gurnard, is a bottom-dwelling fish of tropical to warm temperate waters on both sides of the Atlantic. On the American side, it is found as far north as Massachusetts...



Forcipiger longirostris

Forcipiger longirostris

Forcipiger longirostris, commonly known as the longnose butterflyfish or big longnose butterflyfish, is a species of butterflyfish found on coral reefs throughout the tropical waters of the Indo-Pacific. Even with its distinctive, point-like long...



Forest snakehead

Channa lucius

The forest snakehead (Channa lucius) is a species of snakehead, a fish of the family Channidae. Its range includes most of Southeast Asia and parts of southern China. It lives in forest streams and can reach 40 cm (16 in) in length. The forest...



Forktail blue-eye

Pseudomugil furcatus

The forktail blue-eye (Pseudomugil furcatus) is a diminutive species of rainbowfish in the subfamily Pseudomugilinae. It is endemic to Papua New Guinea, south-east of Popondetta, where it is found in rainforest streams. It used to be placed in the...



Fort Maguire aulonocara

Aulonocara hansbaenschi

The Fort Maguire aulonocara (Aulonocara hansbaenschi), also known as Aulonocara 'Fort Maguire' in the aquarium fish trade, is a species of haplochromine cichlid endemic to Lake Malawi. It is found in Malawi and Mozambique. Its natural habitat is...



Fossorochromis rostratus

Fossorochromis rostratus

Fossorochromis rostratus is a species of cichlid fish that is endemic to Lake Malawi in East Africa. The only species in its genus, it is a relatively large (up to 24.4 centimetres (9.6 in) in total length), sexually dimorphic cichlid. It feeds on...



Fountain darter

Etheostoma fonticola

The fountain darter (Etheostoma fonticola) is a species of freshwater ray-finned fish, a darter from the subfamily Etheostominae, part of the family Percidae, which also contains the perches, ruffes and pikeperches. It is found in the...



Fourbeard rockling

Enchelyopus cimbrius

The fourbeard rockling or four-bearded rockling (Enchelyopus cimbrius) is a species of lotid fish found in the northern Atlantic Ocean. This species grows to 41 cm (16 in) in total length. It is of minor importance in commercial fisheries.



Foureye butterflyfish

Chaetodon capistratus

The foureye butterflyfish (Chaetodon capistratus) is a butterflyfish (family Chaetodontidae). It is alternatively called the four-eyed butterflyfish. This species is found in the Western Atlantic from Massachusetts, USA and Bermuda to the West...



Fourhorn sculpin

Myoxocephalus quadricornis

The fourhorn sculpin (Myoxocephalus quadricornis) is a species of ray-finned fish belonging to the family Cottidae, the typical sculpins. This species has a Holarctic distribution and can be found in marine, brackish and fresh waters.



Foxface rabbitfish

Siganus vulpinus

The foxface rabbitfish (Siganus vulpinus), also known as the foxface, black-face rabbitfish, common foxface or badger fish is a species of marine ray-finned fish, a rabbitfish belonging to the family Siganidae. It is found in the Indian and...



Foxfish

Bodianus frenchii

The foxfish, Bodianus frenchii, is a species of wrasse native to the temperate marine waters in southwestern Western Australia to eastern South Australia, and from southern Queensland to eastern Tasmania, at depths between 10 and 40 m. A gap in...



Freckled madtom

Noturus nocturnus

The freckled madtom (Noturus nocturnus) a species of freshwater madtom (catfish) found in the eastern United States. The freckled madtom is a benthic feeder mostly of invertebrates and usually inhabits waters that are medium to large in size,...



French angelfish

Pomacanthus paru

The French angelfish (Pomacanthus paru) is a species of marine ray-finned fish, a marine angelfish belonging to the family Pomacanthidae. It occurs in the Western Atlantic Ocean.



Freshwater blenny

Salaria fluviatilis

The freshwater blenny (Salariopsis fluviatilis) is a species of fish in the family Blenniidae. It is found in African rivers and brooks in Algeria and Morocco flowing to the Mediterranean Sea. In Europe it is widespread in the freshwaters of...



Freshwater butterflyfish

Pantodon buchholzi

The freshwater butterflyfish or African butterflyfish, Pantodon buchholzi, is the only extant species in the family Pantodontidae within the order Osteoglossiformes. It is not closely related to saltwater butterflyfishes.



Freshwater cobbler

Tandanus bostocki

The freshwater cobbler, Tandanus bostocki, is a species of catfish (order Siluriformes) of the family Plotosidae. The freshwater cobbler originates from coastal drainages of southwestern Australia, from the Frankland River to the Moore River. The...



Freshwater drum

Aplodinotus grunniens

The freshwater drum, Aplodinotus grunniens, is a fish endemic to North and Central America. It is the only species in the genus Aplodinotus, and is a member of the family Sciaenidae. It is the only North American member of the group that inhabits...



Freshwater mullet

Pseudomyxus capensis

The freshwater mullet (Pseudomyxus capensis) is a species of fish in the family Mugilidae. It is endemic to South Africa.



Freshwater whipray

Urogymnus dalyensis

The freshwater whipray (Urogymnus dalyensis) is a little-known species of stingray in the family Dasyatidae, found in a number of large rivers and associated estuaries in northern Australia. Until recently, this species was regarded as a regional...



Frigate tuna

Auxis thazard

The frigate tuna, frigate mackerel or alagaduwa (Auxis thazard) is a species of tuna, in the family Scombridae, found around the world in tropical oceans. The eastern Pacific population is now regarded as a separate species by some authorities....



Frilled shark

Chlamydoselachus anguineus

The frilled shark (Chlamydoselachus anguineus), also known as the lizard shark, is one of the two extant species of shark in the family Chlamydoselachidae (the other is the southern African frilled shark (Chlamydoselachus africana)). The frilled...



Frillfin goby

Bathygobius soporator

The frillfin goby (Bathygobius soporator) is a species of marine fish in the genus Bathygobius.



Fringed-lipped peninsula carp

Labeo fimbriatus

The fringed-lipped peninsula carp (*Labeo fimbriatus*) is a cyprinid fish in genus *Labeo* from Pakistan, India, Nepal, Bangladesh and Myanmar. It is listed as Least Concern in the global IUCN Red List of threatened species.



Fringefin lanternshark

Etmopterus schultzi

Sharks portalThe fringefin lanternshark (*Etmopterus schultzi*) is a shark of the family Etmopteridae found in the western central Atlantic from Texas to Florida, northern Gulf of Mexico, and Mexico. It is endemic to this area. It is a deep water...



Frog shark

Somniosus longus

The frog shark (Somniosus longus) is a very rare species of squaliform shark mainly found in deep water in the Pacific Ocean. It is in the sleeper shark family Somniosidae with the Greenland shark.



Gafftopsail catfish

Bagre marinus

The gafftopsail catfish (Bagre marinus) is a species of marine catfish found in the waters of the western central Atlantic Ocean, as well as the Gulf of Mexico and the Caribbean Sea. It has long, venomous spines which can cause painful wounds. It...



Gagata gagata

Gagata gagata

Gagata gagata is a species of sisorid catfish found in Bangladesh, Myanmar and India. This species grows to a length of 30.5 centimetres (12.0 in) TL. It is native to Asia, found in India, Bangladesh and Burma.



Galapagos bullhead shark

Heterodontus quoyi

Sharks portalThe Galapagos bullhead shark, Heterodontus quoyi, is a bullhead shark of the family Heterodontidae found in the tropical eastern Pacific Ocean between latitudes 0° to 10°S, at depths between 3 and 40 m. It can reach a length of 1.07...



Galapagos shark

Carcharhinus galapagensis

The Galapagos shark (Carcharhinus galapagensis) is a species of requiem shark, in the family Carcharhinidae, found worldwide. It favors clear reef environments around oceanic islands, where it is often the most abundant shark species. A large...



Galaxias olidus

Galaxias olidus

Galaxias olidus, the mountain galaxias, is a species of freshwater galaxiid fish widely found in southeastern Australia.



Galeichthys feliceps

Galeichthys feliceps

Galeichthys feliceps, the white barbel, sea barbel, white baggar or white sea catfish, is a species of sea catfish found in coastal waters and estuaries over muddy bottoms at depths of from 1 – 120 metres where they gather in large shoals. They...



Galjoen

Dichistius capensis

The galjoen, black bream, or blackfish (*Dichistius capensis*) is a species of marine fish found only along the coast of South Africa. Galjoen is the national fish of South Africa.



Ganges shark

Glyphis gangeticus

The Ganges shark (Glyphis gangeticus) is a critically endangered species of requiem shark found in the Ganges River (Padma River) and the Brahmaputra River of India and Bangladesh. It is often confused with the more common bull shark...



Garfish

Belone belone

The garfish (Belone belone), also known as the garpike or sea needle, is a pelagic, oceanodromous needlefish found in brackish and marine waters of the Atlantic Ocean and the Mediterranean, Caribbean, Black, and Baltic Seas.



Garibaldi

Hypsypops rubicundus

The Garibaldi, known historically also as the Catalina goldfish and marine goldfish and now commonly as the Garibaldi damselfish (Hypsypops rubicundus) is a species of bright orange fish in the damselfish family. It occurs in the subtropical...



Garra barreimiae

Garra barreimiae

Garra barreimiae, the Oman garra, is a species of ray-finned fish in the family Cyprinidae. It is found in the mountains of northern Oman and in the United Arab Emirates. Most populations inhabit wadis, streams, pools and springs, but one...



Garra flavatra

Garra flavatra

Garra flavatra can also be called Panda Garra, is a species of cyprinid fish in the genus Garra.



Garra gotyla

Garra gotyla

Garra gotyla is a species of ray-finned fish in the cyprinid family from freshwater in South Asia. There are two subspecies, which are widely regarded as separate species by some authorities (e.g., the IUCN): Garra gotyla gotyla (Gray, 1830) –...



Gastromyzon punctulatus

Gastromyzon punctulatus

Gastromyzon punctulatus, the hillstream loach, is a species of ray-finned fish in the genus Gastromyzon.



Genicanthus bellus

Genicanthus bellus

Genicanthus bellus, the ornate angelfish, bellus angelfish or bellus lyretail angelfish, is a species of marine ray-finned fish, a marine angelfish belonging to the family Pomacanthidae. It is found in the Indo-Pacific region.



Genicanthus personatus

Genicanthus personatus

Genicanthus personatus the masked angelfish, is a species of marine ray-finned fish, a marine angelfish belonging to the family Pomacanthidae. It is endemic to Hawaii.



Genidens barbatus

Genidens barbatus

Genidens barbatus, the white sea catfish or marine catfish, is a species of catfish in the family Ariidae. It was described by Bernard Germain de Lacépède in 1803, originally under the genus Pimelodus. It is an oceanodromous species that is found...



Geometric moray

Gymnothorax griseus

The geometric moray (Gymnothorax griseus) is a moray eel of the family Muraenidae found throughout the western Indian Ocean at depths to 40 m. Its length is up to 65 cm. It is parasitized by Ichthyoxenus puhi, a species of isopod. The fish is...



Geophagus altifrons

Geophagus altifrons

Geophagus altifrons is a freshwater eartheater cichlid fish native to the Amazon River Basin, Brazil. Introduced to Singapore, it thrives in slow-moving streams and reservoirs. Its ecological impact on its adopted habitat is still unknown. A member...



Gerres erythrourus

Gerres erythrourus

Gerres erythrourus the deep-bodied mojarra, also known as the short silverbidy or short silverbelly, is a species of ray-finned fish from the family Gerreidae, a mojarra. It is native to marine and brackish waters of coastal waters of the Indian...



Gerres filamentosus

Gerres filamentosus

Gerres filamentosus, the whipfin silver-biddy, flagfin mojarra or threadfin silver belly, is a fish native to the coastline of Africa and Madagascar east to Japan, Australia and New Caledonia.



Gerres subfasciatus

Gerres subfasciatus

Gerres subfasciatus, the common silver belly roach, ovate silver bidy., common silverbidy or Southern silver bidy, is a species of mojarra native to Indian and Pacific coastal waters of Australia.



Ghost catshark

Apristurus manis

Sharks portalThe ghost catshark (*Apristurus manis*) is a catshark of the family Scyliorhinidae found on the continental slopes in the northwest Atlantic off Massachusetts, the northeast Atlantic from the Porcupine Bank west of Ireland and the...



Giant African threadfin

Polydactylus quadrifilis

The giant African threadfin (*Polydactylus quadrifilis*) is a species of ray-finned fish from the threadfin family Polynemidae. It is found in the eastern Atlantic Ocean off the west coast of Africa.



Giant barb

Catlocarpio siamensis

The giant barb, Siamese giant carp, or simply Siamese carp (*Catlocarpio siamensis*; Khmer: ត្រីគីលីរាំង, *trei kól reäng*; Thai: กระโห้, RTGS:...



Giant catfish

Netuma thalassina

The giant catfish (*Netuma thalassina*), also known as the giant sea catfish, giant salmon catfish, giant marine-catfish, or the khagga, is a species of catfish in the family Ariidae. It was described by Eduard Rüppell in 1837, originally under the...



Giant cichlid

Boulengerochromis microlepis

The giant cichlid (*Boulengerochromis microlepis*), also known as the emperor cichlid, is a species of fish in the family Cichlidae, endemic to Lake Tanganyika in Africa. It is the only member of its genus *Boulengerochromis* and tribe...



Giant danio

Devario aequipinnatus

The giant danio (*Devario aequipinnatus*) is a tropical fish belonging to the minnow family Cyprinidae. Originating in Sri Lanka, Nepal, and the west coast of India, this species grows to a maximum length of 4-6 inches (10-15 cm), making it one of...



Giant freshwater stingray

Urogymnus polylepis

The giant freshwater stingray (*Urogymnus polylepis*, also widely known by the junior synonym *Himantura chaophraya*) is a species of stingray in the family Dasyatidae. It is found in large rivers and estuaries in Southeast Asia and Borneo, though...



Giant goby

Gobius cobitis

The giant goby (*Gobius cobitis*) is a species of goby native to coastal marine and brackish waters of the eastern Atlantic, the Mediterranean Sea and the Black Sea at depths of from 10 to 35 metres (33 to 115 ft). This species is of minor...



Giant gourami

Osphronemus goramy

The giant gourami (*Osphronemus goramy*) is a species of large gourami native to freshwater habitats in Southeast Asia. It has also been introduced elsewhere. The species is commercially important as a food fish and is also farmed. It can be found...



Giant grouper

Epinephelus lanceolatus

The giant grouper (Epinephelus lanceolatus), also known as the Queensland groper (grouper), brindle grouper or mottled-brown sea bass, is a species of marine ray-finned fish, a grouper from the subfamily Epinephelinae which is part of the family...



Giant hawkfish

Cirrhitus rivulatus

The giant hawkfish (Cirrhitus rivulatus), also known as the hieroglyphic hawkfish, is a species of marine ray-finned fish, a hawkfish belonging to the family Cirrhitidae. It is a marine fish and the largest of the hawkfish family with maximum...



Giant kelpfish

Heterostichus rostratus

The giant kelpfish (Heterostichus rostratus) is a species of clinid native to the west coast of North America, where it is found from California to southern Baja California. It inhabits rocky areas with kelp and other large seaweeds. Its diet...



Giant kōkopu

Galaxias argenteus

The giant kōkopu (Galaxias argenteus) is a threatened species of ray-finned fish in the genus Galaxias, found only in New Zealand. It can reach up to 58 cm (23 in) in length and 2.7 kg (6.0 lb) in weight, making it the largest species in the...



Giant moray

Gymnothorax javanicus

The giant moray (Gymnothorax javanicus) is a species of moray eel and a species of marine fish in the family Muraenidae. In terms of body mass, it is the largest moray eel; however, the slender giant moray is the largest in terms of body length.



Giant mottled eel

Anguilla marmorata

The giant mottled eel (Anguilla marmorata), also known as the marbled eel, is a species of tropical anguillid eel that is found in the Indo-Pacific and adjacent freshwater habitats.



Giant mudskipper

Periophthalmodon schlosseri

The giant mudskipper (Periophthalmodon schlosseri) is a species of mudskipper native to the tropical shores of the eastern Indian Ocean and the western Pacific Ocean where it occurs in marine, brackish and fresh waters. It is most frequently...



Giant oarfish

Regalecus glesne

The giant oarfish (Regalecus glesne) is a species of oarfish of the family Regalecidae. It is an oceanodromous species with a worldwide distribution, excluding polar regions. Other common names include Pacific oarfish, king of herrings,....



Giant oceanic manta ray

Mobula birostris

The giant oceanic manta ray, giant manta ray, or oceanic manta ray (Mobula birostris) is a species of ray in the family Mobulidae, and the largest type of ray in the world. It is circumglobal and is typically found in tropical and subtropical...



Giant pangasius

Pangasius sanitwongsei

The giant pangasius, paroon shark, pangasid-catfish or Chao Phraya giant catfish (Pangasius sanitwongsei) is a species of freshwater fish in the shark catfish family (Pangasiidae) of order Siluriformes, found in the Chao Phraya and Mekong basins...



Giant red tail gourami

Osphronemus laticlavius

The giant red tail gourami (Osphronemus laticlavius) is a large species of gourami belonging to the family Osphronemidae. This endangered fish is native to the Southeast Asian island of Borneo, where only known from the Kinabatangan and Segama...



Giant salmon carp

Aptosyax grypus

The giant salmon carp (Aptosyax grypus), also termed the Mekong giant salmon carp, is a species of freshwater fish in the family Cyprinidae and the single species in the monotypic genus Aptosyax. It is endemic to the middle reaches of the...



Giant sea bass

Stereolepis gigas

The giant sea bass (Stereolepis gigas) is a fish native to the North Pacific Ocean. Although commonly referred to as a giant sea bass, black sea bass or giant black sea bass, it is actually a wreckfish in the family Polyprionidae rather than in...



Giant stargazer

Kathetostoma giganteum

The giant stargazer (Kathetostoma giganteum) is a stargazer of the family Uranoscopidae, found on the continental shelf around New Zealand and endemic to that area. It is commonly called monkfish, but this should not be confused with the Northern...



Giant sunfish

Mola alexandrini

The giant sunfish or bumphead sunfish (Mola alexandrini), (also known as the Ramsay's sunfish, southern sunfish, southern ocean sunfish, short sunfish or bump-head sunfish in various parts of the world), is a fish belonging to the family...



Giant trevally

Caranx ignobilis

The giant trevally (Caranx ignobilis), also known as the lowly trevally, barrier trevally, ronin jack, giant kingfish, or ulua, is a species of large marine fish classified in the jack family, Carangidae. The giant trevally is distributed...



Gibbonsia montereyensis

Gibbonsia montereyensis

Gibbonsia montereyensis, the crevice kelpfish, is a species of clinid found along the Pacific coast of North America from British Columbia, Canada, to Baja California, Mexico where it prefers areas close to the shore amongst algae. This species...



Gila chub

Gila intermedia

The Gila chub (Gila intermedia) is a species of ray-finned fish in the family Cyprinidae. It is found in Mexico and the United States. The Gila chub is closely related to the roundtail chub. This species is commonly found in association with the...



Gila topminnow

Poeciliopsis occidentalis

The Gila topminnow or charalito (Poeciliopsis occidentalis) is a species of fish in the family Poeciliidae. It is found in Mexico and the United States.



Gila trout

Oncorhynchus gilae

The Gila trout (Oncorhynchus gilae) is a species of salmonid, related to the rainbow trout, native to the Southwest United States. Prior to 2006 the Gila trout was federally listed as endangered. In July 2006, after much work by the Game and Fish...



Gilded catfish

Zungaro zungaro

The gilded catfish or jau (Zungaro zungaro) is a South American catfish (order Siluriformes) of the family Pimelodidae. It is also known as manguruyu or black manguruyu.



Gilded triggerfish

Xanthichthys auromarginatus

The gilded triggerfish or blue-throated triggerfish, Xanthichthys auromarginatus, is a spotted gray triggerfish. Males of the species have blue cheeks and yellow-bordered, white fins. It is widely, but locally, distributed at islands in the...



Gillaroo

Salmo stomachicus

The gillaroo (Salmo stomachicus}; historically included in Salmo trutta) is a species of trout which eats primarily snails and is only proven to inhabit Lough Melvin in Ireland. The name gillaroo is derived from the Irish for "red fellow" (giolla...



Ginbuna

Carassius langsdorfii

The ginbuna (Carassius langsdorfii), sometimes referred to as silver crucian carp or Japanese silver crucian carp, is a species of freshwater fish in the carp family (family Cyprinidae). It is native to lakes and rivers in Japan.



Giraffe catfish

Auchenoglanis occidentalis

The giraffe catfish, Auchenoglanis occidentalis, is an African catfish. It eats plants off the floor of lakes and streams.



Girella cyanea

Girella cyanea

Girella cyanea, also known as the blue drummer or Australian bluefish, is a species of sea chub native to inshore waters, around 6 metres (20 ft) depth, from Australia to New Zealand and the Kermadec Islands. Sightings were first recorded in 1881...



Girella nigricans

Girella nigricans

Girella nigricans, commonly known as the opaleye or rudderfish, is a species of sea chub found in the Eastern Pacific, from California to southern Baja California. A rarely documented isolated population also exists in the Gulf of California,...



Giuris margaritacea

Giuris margaritacea

Giuris margaritacea, the snakehead gudgeon, Aporos sleeper, or ornate sleeper, is a species of fish in the family Eleotridae found in marine, brackish, and fresh waters from Madagascar to Melanesia. This species grows to a total length of 29 cm...



Glaucosoma hebraicum

Glaucosoma hebraicum

Glaucosoma hebraicum, the West Australian dhufish, Westralian jewfish, or West Australian pearl perch, is a species of fish in the family Glaucosomatidae, the pearl perches. It is endemic to the waters around Western Australia from Shark Bay,...



Glossamia aprion

Glossamia aprion

Glossamia aprion is a species of freshwater ray-finned fish from the family Apogonidae, the cardinalfishes, from northern Australia and New Guinea. In Australia, it is commonly known as the mouth almighty.



Glowlight danio

Danio choprae

The glowlight danio (Danio choprae) is a small, schooling fish closely related to the popular zebrafish Danio rerio. This should not be confused with the GloFish, a trademarked brand of fluorescent zebrafish that appear to glow in the dark under...



Glowlight rasbora

Trigonostigma hengeli

The glowlight rasbora (Trigonostigma hengeli) is a species of cyprinid fish in the genus Trigonostigma.



Gnathochromis permaxillaris

Gnathochromis permaxillaris

Gnathochromis permaxillaris is a species of African fish in the family Cichlidae. It is endemic to relatively deep waters in Lake Tanganyika in the countries of Burundi, the Democratic Republic of the Congo, Tanzania, and Zambia. It is widespread...



Gobiesox strumosus

Gobiesox strumosus

The skilletfish (*Gobiesox strumosus*) is a member of the family Gobiesocidae, composed of clingfishes. These are mainly marine, tropical and temperate fishes characterized by a powerful sucking disk on the underside of the forebody.



Gobio gobio

Gobio gobio

Gobio gobio, or the gudgeon, is a species of fish in the family Cyprinidae. This small fish is widely distributed in fresh-water streams and lakes across central and temperate Eurasia. The gudgeon inhabits various fresh-water habitats with sandy...



Gobiocypris rarus

Gobiocypris rarus

Gobiocypris rarus is a freshwater species of cyprinid fish endemic to China where it is only found near Jiuxiang in Sichuan province. It is the only species in its genus.



Gobiodon citrinus

Gobiodon citrinus

Gobiodon citrinus, the poison goby, is a species of goby native to the Indian Ocean from the Red Sea and the coast of Africa to the western Pacific Ocean to Japan, Samoa and the Great Barrier Reef. They are reef dwellers being found at depths of...



Gobiodon histrio

Gobiodon histrio

Gobiodon histrio, the Broad-barred goby, is a species of goby native to the Indian Ocean from the Red Sea to the western Pacific Ocean to southern Japan, Samoa and the Great Barrier Reef. This species is a reef dweller, being found at depths of...



Gobiosoma bosc

Gobiosoma bosc

Gobiosoma bosc (naked goby) is a fish named for its lack of scales. This is a true goby; it is part of the genus Gobiosoma.



Gobius tropicus

Gobius tropicus

Gobius tropicus is a species of fish currently classified in the family Gobiidae. It is native the Atlantic waters around Ascension Island. The actual taxonomic position of this species is uncertain and it is suspected that it is not even a goby.



Goblin shark

Mitsukurina owstoni

The goblin shark (Mitsukurina owstoni) is a rare species of deep-sea shark. Sometimes called a "living fossil", it is the only extant representative of the family Mitsukurinidae, a lineage some 125 million years old. This pink-skinned animal has...



Gold barb

Barbodes semifasciolatus

Barbodes semifasciolatus, the Chinese barb, is a species of cyprinid fish native to the Red River basin in southeast Asia where they occur in fresh waters at depths of 5 metres (16 ft) or less. The captive variant popularly known as the gold barb...



Gold tetra

Hemigrammus rodwayi

The gold tetra (Hemigrammus rodwayi) is a species of small freshwater fish in the family Characidae native to lowland South America, where they are abundant in coastal floodplains. Their name comes from a powdery golden tint on their body that is...



Goldband fusilier

Pterocaesio chrysozona

The gold-band fusilier (*Pterocaesio chrysozona*) also known as the yellow-band fusilier or black-tipped fusilier, is a species of marine ray-finned fish, a fusilier belonging to the family Caesionidae. It is widespread around reefs in the...



Golden barb

Pethia gelius

The golden barb or golden dwarf barb (*Pethia gelius*) is a species of cyprinid fish native to inland waters in Asia, and is found in Pakistan, India, and Bangladesh. It has also been introduced to waters in Colombia. It natively inhabits rivers,...



Golden grey mullet

Chelon aurata

The golden grey mullet (*Chelon aurata*) is a fish in the family Mugilidae.



Golden grouper

Saloptia powelli

The golden grouper (Saloptia powelli), also known as the pink grouper or Powell's grouper, is a species of marine ray-finned fish, a grouper from the subfamily Epinephelinae which is part of the family Serranidae, which also includes the anthias...



Golden mandarin fish

Siniperca scherzeri

The golden mandarin fish (Siniperca scherzeri), also known as the leopard mandarin fish, is a species of freshwater ray-finned fish, an Asian perch from the family Sinipercidae, which is native to eastern Asia (Korea, China, and Vietnam). This...



Golden perch

Macquaria ambigua

The golden perch (Macquaria ambigua) is a medium-sized, yellow or gold-coloured species of Australian freshwater fish found primarily in the Murray-Darling River system, though a subspecies is found in the Lake Eyre-Cooper Creek system, and...



Golden redhorse

Moxostoma erythrurum

The golden redhorse (Moxostoma erythrurum) is a species of freshwater fish endemic to Ontario and Manitoba in Canada and the Midwestern, southern, and eastern United States. It lives in calm, often silty or sandy waters in streams, small to large...



Golden shiner

Notemigonus crysoleucas

The golden shiner (*Notemigonus crysoleucas*) is a cyprinid fish native to eastern North America. It is the sole member of its genus. Much used as a bait fish, it is probably the most widely pond-cultured fish in the United States. It can be found...



Golden skiffia

Skiffia francesae

Skiffia francesae, the golden skiffia or tiro dorado, is a species of splitfin endemic to the Rio Teuchitlán, a tributary of Río Ameca in western Mexico. It is extinct in the wild, but has been maintained in aquaria and the aquarium hobbyist trade.



Golden topminnow

Fundulus chrysotus

The golden topminnow (Fundulus chrysotus) is a fish of the genus Fundulus and is a United States native fish mostly distributed throughout the southeast, ranging from Kentucky and Ohio south into Florida. Although it has such a wide distribution...



Golden trevally

Gnathanodon speciosus

The golden trevally (Gnathanodon speciosus), also known as the golden kingfish, banded trevally or king trevally, is a species of large marine fish classified in the jack and horse mackerel family Carangidae, and the only member of the...



Golden trout

Oncorhynchus mykiss aguabonita

The California golden trout (Oncorhynchus aguabonita or Oncorhynchus mykiss aguabonita), is a species of trout native to California. The golden trout is normally found in the Golden Trout Creek (tributary to the Kern River), Volcano Creek...



Goldeneye cichlid

Nannacara anomala

The goldeneye cichlid (*Nannacara anomala*) is a species of cichlid found in fresh water from the Aruka River in Guyana, to the Maroni River in Suriname. It is often found on flooded savannas near the coast. The male grows to a length of about 5.6...



Goldenstriped soapfish

Grammistes sexlineatus

The goldenstriped soapfish (*Grammistes sexlineatus*), also known as the lined soapfish, golden-striped bass, radio fish, sixline soapfish, six-lined perch or white-lined rock cod, is a species of marine ray-finned fish, related to the groupers and...



Goldeye

Hiodon alosoides

The goldeye (*Hiodon alosoides*) is a freshwater fish found in Canada and the northern United States. It is one of only two extant species in the family Hiodontidae, the other species being *Hiodon tergisus*. The species name *alosoides* means...



Goldfish

Carassius auratus

The goldfish (*Carassius auratus*) is a freshwater fish in the family Cyprinidae of order Cypriniformes. It is commonly kept as a pet in indoor aquariums, and is one of the most popular aquarium fish. Goldfish released into the wild have become an...



Goldsinny wrasse

Ctenolabrus rupestris

The goldsinny wrasse (*Ctenolabrus rupestris*) is a species of ray-finned fish, a wrasse from the family Labridae which is native to the eastern Atlantic Ocean and associated seas. This species is the only known member of its genus.



Goldstripe sardinella

Sardinella gibbosa

The goldstripe sardinella (Sardinella gibbosa) is a species of fish of the family Clupeidae. It is native to shallow tropical waters of the western Indo-Pacific, living at depths down to 70 m, and being associated with coral reefs. It grows up to...



Goldtail angelfish

Pomacanthus chrysurus

The goldtail angelfish (Pomacanthus chrysurus), also known as the earspot angelfish, is a species of marine ray-finned fish, a marine angelfish belonging to the family Pomacanthidae. It is found in the western Indian Ocean.



Gomphosus caeruleus

Gomphosus caeruleus

Gomphosus caeruleus, the green birdmouth wrasse, is a species of wrasse belonging to the family Labridae. It can be found in the aquarium trade.



Gomphosus varius

Gomphosus varius

The bird wrasse, Gomphosus varius, is a species of the wrasse family.



Gonorynchus gonorynchus

Gonorynchus gonorynchus

Gonorynchus gonorynchus is a species of fish in the family Gonorynchidae, found on temperate continental shelves worldwide. Common names for this fish include mousefish, ratfish, sandfish, and sand eel.



Gopher rockfish

Sebastes carnatus

The gopher rockfish (Sebastes carnatus), also known as the gopher sea perch, is a species of marine ray-finned fish belonging to the subfamily Sebastinae, the rockfishes, part of the family Scorpaenidae. It is found in the eastern Pacific,...



Gorgasia preclara

Gorgasia preclara

Gorgasia preclara, the splendid garden eel or orange-barred garden eel, is a species of marine garden eel.



Graceful shark

Carcharhinus amblyrhynchoides

The graceful shark or Queensland shark (Carcharhinus amblyrhynchoides) is a species of requiem shark, in the family Carcharhinidae, found in the tropical Indo-Pacific, from the Gulf of Aden to northern Australia. It is a midwater species that has...



Gramma brasiliensis

Gramma brasiliensis

Gramma brasiliensis is a species of fish endemic to the Atlantic coast of Brazil where it is a reef inhabitant. It prefers areas with rocks or coral. This species can reach a length of 6.6 centimetres (2.6 in) SL. It can also be found in the...



Grammatostomias flagellibarba

Grammatostomias flagellibarba

Grammatostomias flagellibarba is a species of barbeled dragonfish. They live at depths of up to 1,500 m (5,000 ft) below the surface and usually measure up to 15.2 cm (6 in) in length.



Grass carp

Ctenopharyngodon idella

The grass carp (Ctenopharyngodon idella) is a species of large herbivorous freshwater fish in the family Cyprinidae, native to the Pacific Far East, with a native range stretching from northern Vietnam to the Amur River on the Sino-Russian...



Grass goby

Zosterisessor ophiocephalus

The grass goby (Zosterisessor ophiocephalus) is a species of goby native to the Mediterranean Sea, the Sea of Azov and the Black Sea. It is the only known member of its genus.



Grass puffer

Takifugu niphobles

The grass puffer (Takifugu niphobles), or Kusa-fugu (Japanese: 草河豚), is a species of fish in the pufferfish family (Tetraodontidae). This common to abundant species is found in the northwest Pacific Ocean in China (including Hong Kong and...



Gray angelfish

Pomacanthus arcuatus

The gray angelfish (*Pomacanthus arcuatus*), also written as grey angelfish and known in Jamaica as the pot cover, is a species of marine ray-finned fish belonging to the marine angelfish family, Pomacanthidae. It is found in the Western Atlantic...



Gray smooth-hound

Mustelus californicus

Sharks portalThe gray smooth-hound (*Mustelus californicus*) is a houndshark of the family Triakidae. It is found on the continental shelves of the subtropical eastern Pacific, from northern California to the Gulf of California, between latitudes...



Graysby

Cephalopholis cruentata

The graysby (*Cephalopholis cruentata*) is a species of marine ray-finned fish, a grouper from the subfamily Epinephelinae which is in the family Serranidae which also includes the anthias and sea basses. It is found in the western Atlantic. It is...



Greasy grouper

Epinephelus tauvina

The greasy grouper (*Epinephelus tauvina*), also known as the Arabian grouper or greasy rockcod, is an Indo-Pacific fish species of economic importance belonging to the family Serranidae.



Great barracuda

Sphyraena barracuda

Sphyraena barracuda, commonly known as the great barracuda, is a species of barracuda: large, predatory ray-finned fish found in subtropical oceans around the world.



Great Basin redband trout

Oncorhynchus mykiss newberrii

The Great Basin redband trout (Oncorhynchus mykiss newberrii) is one of three redband trout subspecies of the rainbow trout in the western United States. The Great Basin redband trout is native to drainages in south central Oregon east of the...



Great hammerhead

Sphyrna mokarran

The great hammerhead (Sphyrna mokarran) is the largest species of hammerhead shark, belonging to the family Sphyrnidae, attaining an average length of 4.6 m (15 ft) and reaching a maximum length of 6.2 m (20 ft). It is found in tropical and warm...



Great northern tilefish

Lopholatilus chamaeleonticeps

The great northern tilefish (Lopholatilus chamaeleonticeps) or golden tile, is the largest species in the family Malacanthidae (tilefishes). It grows to a length of between 38 and 44 inches (970 and 1,120 mm). The great northern tilefish is a...



Great sand eel

Hyperoplus lanceolatus

The great sand eel (Hyperoplus lanceolatus) is the greater species of sand eel. The maximum size is 35 centimetres (14 in).



Great seahorse

Hippocampus kelloggi

The great seahorse (Hippocampus kelloggi), also known as Kellogg's seahorse is a species of fish in the family Syngnathidae. It is one of the largest of the 54 species of seahorse.



Great white shark

Carcharodon carcharias

The great white shark (Carcharodon carcharias), also known as the white shark, white pointer, or simply great white, is a species of large mackerel shark which can be found in the coastal surface waters of all the major oceans. It is the only...



Greater amberjack

Seriola dumerili

The greater amberjack (Seriola dumerili), also known as the allied kingfish, great amberfish, greater yellowtail, jenny lind, Sea donkey, purplish amberjack, reef donkey, rock salmon, sailors choice, yellowtail, and yellow trevally, is a species...



Greater argentine

Argentina silus

The greater argentine (Argentina silus ; synonymous with Salmo silus), also known as the Atlantic argentine, great silver smelt, herring smelt or simply smelt, is a northern Atlantic herring smelt and can be found at depths from 140 to 1,440...



Greater pipefish

Syngnathus acus

The greater pipefish (Syngnathus acus) is a pipefish of the family Syngnathidae. It is a seawater fish and the type species of the genus Syngnathus.



Greater redhorse

Moxostoma valenciennesi

The greater redhorse (Moxostoma valenciennesi) is a species of freshwater fish found in northeastern North America. It is the largest member of the genus Moxostoma, with a maximum length of 80 cm (31.5 in). The greater redhorse is long lived with...



Greater weever

Trachinus draco

The greater weever (Trachinus draco, Linnaeus 1758) is a benthic and demersal venomous marine fish of the family Trachinidae. The greater weever is widely distributed along the eastern Atlantic coastline from Norway to Morocco, extending to the...



Green chromide

Eetroplus suratensis

The green chromide (Eetroplus suratensis) is a species of cichlid fish that is native to fresh and brackish water habitats in some parts in India such as Kerala, Goa, Chilika Lake in Odisha and Sri Lanka. The species was first described by Marcus...



Green gold catfish

Corydoras melanotaenia

The green gold catfish (Corydoras melanotaenia) is a tropical freshwater fish belonging to the Corydoradinae subfamily of the family Callichthyidae. It originates in inland waters in South America, and is found in the Meta River basin in...



Green humphead parrotfish

Bolbometopon muricatum

The green humphead parrotfish (*Bolbometopon muricatum*) is the largest species of parrotfish, growing to lengths of 1.5 m (4.9 ft) and weighing up to 75 kg (165 lb). It is found on reefs in the Indian and Pacific Oceans, from the Red Sea in the...



Green jack

Caranx caballus

The green jack (Caranx caballus), also known as the horse jack, is an abundant species of coastal marine fish in the jack family Carangidae. The species is distributed in the eastern Pacific Ocean along the American coastline from Santa Cruz...



Green jobfish

Aprion virescens

The green jobfish (Aprion virescens), also known as the gray jobfish, gray snapper, or slender snapper, and in Hawaiian as uku, is a species of marine ray-finned fish, a snapper belonging to the family Lutjanidae. It is found in the Indo-Pacific...



Green lanternshark

Etmopterus virens

The green lanternshark (Etmopterus virens) is a species of dogfish shark in the family Etmopteridae, found in the western central Atlantic Ocean. This species usually occurs on the upper continental slope below a depth of 350 m (1,150 ft)....



Green moray

Gymnothorax funebris

The green moray (Gymnothorax funebris) is a moray eel of the family Muraenidae, found in the western Atlantic Ocean from Long Island, New York, Bermuda, and the northern Gulf of Mexico to Brazil, at depths down to 40 metres (130 ft). Its length...



Green neon tetra

Paracheirodon simulans

The green neon tetra (Paracheirodon simulans) is a freshwater fish of the family Characidae of order Characiformes. It is native to the upper Orinoco and Negro Rivers in South America. This fish is similar in appearance to the closely related and...



Green pufferfish

Dichotomyctere fluviatilis

The green pufferfish or Ceylon pufferfish (Dichotomyctere fluviatilis, syn. Tetraodon fluviatilis) is a species of pufferfish found in South and Southeast Asia. Its habitat includes rivers, estuaries, lakes and flood plains. It lives in fresh to...



Green sturgeon

Acipenser medirostris

The green sturgeon (Acipenser medirostris) is a species of sturgeon native to the northern Pacific Ocean, from China and Russia to Canada and the United States.



Green sunfish

Lepomis cyanellus

The green sunfish (Lepomis cyanellus) is a species of aggressive freshwater fish in the sunfish family (Centrarchidae) of order Perciformes. A panfish popular with anglers, the green sunfish is also kept as an aquarium fish by hobbyists, they...



Green swordtail

Xiphophorus hellerii

The green swordtail (Xiphophorus hellerii) is a species of freshwater/brackish fish in family Poeciliidae of order Cyprinodontiformes. A live-bearer, it is closely related to the southern platyfish or 'platy' (X. maculatus) and can crossbreed...



Green terror

Andinoacara rivulatus

The green terror (Andinoacara rivulatus, syn. Aequidens rivulatus) is a colorful freshwater fish in the cichlid family. The fish originates from the Pacific side of South America from the Tumbes River in Peru to the Esmeraldas River in Ecuador...



Greenback cutthroat trout

Oncorhynchus clarkii stomias

The greenback cutthroat trout (Oncorhynchus clarkii stomias) is the easternmost subspecies of cutthroat trout. The greenback cutthroat, once widespread in the Arkansas and South Platte River drainages of Eastern Colorado and Southeast Wyoming,....



Greenback horse mackerel

Trachurus declivis

The greenback horse mackerel or greenback scad (Trachurus declivis) is a species of jack in the family Carangidae, found around western and southern Australia, and around New Zealand, from the surface to depths of 460 m. Its length is up to 64...



Greenbone

Odax pullus

Odax pullus, known by the names greenbone, butterfish or its Māori language name rarī, is a species of ray-finned fish, a weed whiting from the family Odacidae, which is found around New Zealand. It is of minor importance to local commercial...



Greeneye spurdog

Squalus chloroculus

Sharks portalThe greeneye spurdog (Squalus chloroculus) is a species of dogfish described in 2007. It is a member of the family Squalidae, found off the coast of southeastern Australia. The length of the longest specimen measured is 85.6 cm...



Greenland cod

Gadus ogac

The Greenland cod (*Gadus ogac*), commonly known also as ogac, is a species of ray-finned fish in the cod family, Gadidae. Genetic analysis has shown that it may be the same species as the Pacific cod (*Gadus macrocephalus*). It is a bottom-dwelling...



Greenland halibut

Reinhardtius hippoglossoides

The Greenland halibut or Greenland turbot (*Reinhardtius hippoglossoides*) belongs to the family Pleuronectidae (the right-eye flounders), and is the only species of the genus Reinhardtius. It is a predatory fish that mostly ranges at depths...



Greenland shark

Somniosus microcephalus

The Greenland shark (*Somniosus microcephalus*), also known as the gurry shark or grey shark, is a large shark of the family Somniosidae ("sleeper sharks"), closely related to the Pacific and southern sleeper sharks. The Greenland shark is a...



Greenside darter

Etheostoma blennioides

The greenside darter (*Etheostoma blennioides*) is a species of freshwater ray-finned fish, a darter from the subfamily Etheostomatinae, part of the family Percidae, which also contains the perches, ruffes and pikeperches. It inhabits swift riffles...



Grey bamboo shark

Chiloscyllium griseum

The grey bamboo shark, *Chiloscyllium griseum*, is a species of carpet shark in the family Hemiscylliidae, found in the Indo-West Pacific Oceans from the Arabian Sea to Pakistan, India, Malaysia, Thailand, Indonesia, China, Japan, the Philippines,...



Grey gurnard

Eutrigla gurnardus

The grey gurnard (Eutrigla gurnardus) is a species of ray-finned fish from the family Triglidae, the gurnards and sea robins. It is native to the eastern Atlantic Ocean, the Mediterranean Sea, and the Black Sea. It is caught as a food fish and is...



Grey reef shark

Carcharhinus amblyrhynchos

The grey reef shark or gray reef shark (Carcharhinus amblyrhynchos, sometimes misspelled amblyrhynchus or amblyrhinchos) is a species of requiem shark, in the family Carcharhinidae. One of the most common reef sharks in the Indo-Pacific, it is...



Grey sharpnose shark

Rhizoprionodon oligolinx

Sharks portalThe grey sharpnose shark, Rhizoprionodon oligolinx, is a requiem shark of the family Carcharhinidae. It is found in the tropical waters of the Indo-West Pacific Oceans, between latitudes 30° N and 18° S, from the surface to a depth...



Grey triggerfish

Balistes capriscus

The grey triggerfish (Balistes capriscus), or gray triggerfish, is a species of ray-finned fish in the triggerfish family. The species is native to shallow parts of the western Atlantic from Nova Scotia to Argentina and also the eastern Atlantic,...



Greyface moray eel

Gymnothorax thyrsoideus

The greyface moray eel, also called the freckled moray, slender moray, or white-eyed moray. Gymnothorax thyrsoideus, is a species of marine fish in the family Muraenidae.



Grunt sculpin

Rhamphocottus richardsonii

The grunt sculpin or grunt-fish (Rhamphocottus richardsonii) is a small fish mainly found in the eastern Pacific Ocean. The grunt sculpin generally remains close to shore and is often found in empty giant barnacle shells. The common name comes...



Guachanche barracuda

Sphyraena guachancho

The Guachanche barracuda (*Sphyraena guachancho*) is an ocean-going species of game fish in the barracuda family, Sphyraenidae. It was described by the French zoologist Georges Cuvier in 1829. The description was part of the second edition of Le...



Guadalupe bass

Micropterus treculii

The Guadalupe bass (*Micropterus treculii*) is a rare species of fish endemic to the U.S. state of Texas, where it also is the official state fish. It is restricted to creeks and rivers (including the Guadalupe River, hence the name Guadalupe...



Guinean bichir

Polypterus ansorgii

The Guinean bichir (*Polypterus ansorgii*) is a ray-finned fish from rivers and other freshwater habitats in Western Africa, ranging from Guinea-Bissau to Nigeria. It reaches a maximum length of 72 cm (28 in), is greenish-brown to black in color,...



Gulf butterfish

Peprilus burti

The Gulf butterfish (*Peprilus burti*) is a fish species of the family Stromateidae found in the Gulf of Mexico and off the coast of the Eastern United States.



Gulf Coast pygmy sunfish

Elassoma gilberti

The Gulf Coast pygmy sunfish, *Elassoma gilberti*, is a species of pygmy sunfish endemic to Florida, United States. This species can reach 2.5 centimetres (0.98 in) in standard length.



Gulf flounder

Paralichthys albigutta

The Gulf flounder (Paralichthys albigutta) is a species of saltwater flounder.



Gulf grouper

Mycteroperca jordani

The Gulf grouper (Mycteroperca jordani) is a species of marine ray-finned fish, a grouper from the subfamily Epinephelinae which is part of the family Serranidae, which also includes the anthias and sea basses. It is endemic to Mexico.



Gulf killifish

Fundulus grandis

The Gulf killifish (Fundulus grandis) is one of the largest members of the genus Fundulus; it is capable of growing up to 7 inches (18 cm) in length, whereas the majority of other Fundulus reach a maximum length of 4 inches (10 cm). Therefore, F....



Gulf pipefish

Syngnathus scovelli

The Gulf pipefish (Syngnathus scovelli) is a species of pipefish in the member of the taxonomic family Sygnathidae. Syngnathus scovelli is native to the region of south Florida, United States, the Atlantic Ocean, etc. S. scovelli is similar to...



Gulf sturgeon

Acipenser oxyrinchus desotoi

The Gulf sturgeon (Acipenser oxyrinchus desotoi) is a subspecies of sturgeon that lives in the Gulf of Mexico and some rivers draining into it. The Gulf sturgeon was first recognized as a separate subspecies in 1955. The nominate subspecies is...



Gulf toadfish

Opsanus beta

The Gulf toadfish (Opsanus beta) is a species of toadfish found in the Gulf of Mexico.



Gulf torpedo

Torpedo sinuspersici

The Gulf torpedo (Torpedo sinuspersici) or variable electric ray, is a species of electric ray in the family Torpedinidae. It is found in the Indian Ocean, but may represent a species flock of several local endemic species. It is distinguishable...



Gulper shark

Centrophorus granulosus

The gulper shark (Centrophorus granulosus) is a long and slender dogfish usually about three feet in length generally found in deep, murky waters all around the world. It is a light grayish brown, paler ventrally, with a long snout and large...



Gummy shark

Mustelus antarcticus

The gummy shark (*Mustelus antarcticus*), also known as the Australian smooth hound, flake, sweet william or smooth dog-shark, is a species of ground shark in the genus *Mustelus* of the family *Triakidae*. These small to medium-sized bottom-dwelling...



Guppy

Poecilia reticulata

The guppy (*Poecilia reticulata*), also known as millionfish or the rainbow fish, is one of the world's most widely distributed tropical fish and one of the most popular freshwater aquarium fish species. It is a member of the family *Poeciliidae*...



Gwyniad

Coregonus pennantii

The gwyniad (*Coregonus pennantii*) is a freshwater whitefish native to Bala Lake (Welsh: Llyn Tegid) in northern Wales. The population is threatened by deteriorating water quality and by the ruffe, a fish introduced to the lake in the 1980s and now...



Gymnammodytes cicerelus

Gymnammodytes cicerelus

Gymnammodytes cicerelus, also known as Mediterranean sand eel, sonso in Catalan, and barrinaire or enfú in Menorca, is a fish in the family *Ammodytidae*. It is the only species of this family in the Mediterranean Sea. It is a species from the...



Gymnarchus

Gymnarchus niloticus

Gymnarchus niloticus – commonly known as the aba, aba aba, frankfish, freshwater rat-tail, poisson-cheval, or African knifefish – is an electric fish, and the only species in the genus *Gymnarchus* and the family *Gymnarchidae* within the order...



Gymnocypris przewalskii

Gymnocypris przewalskii

Gymnocypris przewalskii (common name: Przewalskii's naked carp; in Chinese: 青海湖裸鲤; lit. 'Qinghai Lake naked carp') is a species of cyprinid that is endemic to the Lake Qinghai basin in China, where it is the dominant fish species (the other...



Gymnothorax miliaris

Gymnothorax miliaris

Gymnothorax miliaris, the goldentail moray, bastard eel, or conger moray, is a species of marine fish in the family Muraenidae.



Gymnothorax nudivomer

Gymnothorax nudivomer

Gymnothorax nudivomer, the starry moray or yellowmouth moray, is a species of marine fish in the family Muraenidae.



Gymnothorax pictus

Gymnothorax pictus

Gymnothorax pictus, the painted moray, paintspotted moray or peppered moray, is a moray eel. The Chamorro name of the eel is títugi.



Gymnothorax polyuranodon

Gymnothorax polyuranodon

Gymnothorax polyuranodon, commonly known as the freshwater moray, is a species of moray eel that is native to the Indo-Pacific region, including Sri Lanka, the Philippines, Indonesia, Papua New Guinea, the northern coastline of Australia, and...



Gymnothorax rueppelliae

Gymnothorax rueppelliae

Gymnothorax rueppelliae, the banded moray, banded reef-eel, Rüppell's moray, Rüppell's moray eel, black barred eel, yellow-headed moray eel or yellow-headed moray, is a moray eel found in tropical coral reefs. Gymnothorax rueppelliae is a pale...



Gyrinocheilus aymonieri

Gyrinocheilus aymonieri

Gyrinocheilus aymonieri is a freshwater fish native to large parts of Southeast Asia. It is of interest as a local food source and for the aquarium trade. Its common names include honey sucker, sucking loach and Chinese algae eater.G. aymonieri...



Gyrinomimus grahami

Gyrinomimus grahami

Gyrinomimus grahami is a flabby whalefish of the genus Gyrinomimus, found in all the southern oceans.



Haddock

Melanogrammus aeglefinus

The haddock (Melanogrammus aeglefinus) is a saltwater ray-finned fish from the family Gadidae, the true cods. It is the only species in the monotypic genus Melanogrammus. It is found in the North Atlantic Ocean and associated seas, where it is an...



Haemulon flavolineatum

Haemulon flavolineatum

Haemulon flavolineatum, the French grunt, banana grunt, gold laced grunt, open-mouthed grunt, redmouth grunt, or yellow grunt, is a species of marine ray-finned fish, a grunt belonging to the family Haemulidae. It is native to the western...



Haemulon plumierii

Haemulon plumierii

Haemulon plumierii, the white grunt or common grunt, is a species of ray-finned fish in the family Haemulidae native to the western Atlantic Ocean. It grows to a length of about 30 cm (12 in) and is a silvery-cream color, with narrow yellow and...



Halfmoon

Medialuna californiensis

The halfmoon (*Medialuna californiensis*), also known as the blue perch, is a species of marine ray-finned fish, a sea chub from the subfamily Scorpidinae, part of the family Kyphosidae. It is native to the coasts of the eastern Pacific Ocean off...



Halichoeres chrysus

Halichoeres chrysus

Halichoeres chrysus, commonly called the canary wrasse, golden wrasse or yellow wrasse, is a fish species in the wrasse family native to central Indo-Pacific area.



Halichoeres maculipinna

Halichoeres maculipinna

Halichoeres maculipinna, the clown wrasse, is a species of tropical fish that lives throughout the Caribbean Sea and adjacent parts of the western Atlantic Ocean. It is a carnivorous, multi-colored wrasse that is common throughout its range.



Haliichthys taeniophorus

Haliichthys taeniophorus

The ribboned pipefish (Haliichthys taeniophorus), ribboned pipehorse or ribboned seadragon, is a species of pipefish found along the coast of northern Australia (Shark Bay to Torres Strait) and New Guinea (both West New Guinea and Papua New...



Halloween darter

Percina crypta

The Halloween darter (*Percina crypta*) is a small freshwater fish native to North America. It is found in Georgia and Alabama in the drainage basin of the Apalachicola River, specifically in the Flint River system and the Chattahoochee River...



Halosaurus ovenii

Halosaurus ovenii

Halosaurus ovenii, also called the Oven's halosaur, is a deep-sea fish in the family Halosauridae. It is found in the eastern Atlantic where it is widely distributed in tropical and subtropical waters along both coasts. In the Mediterranean, H....



Hammerjaw

Omosudis lowii

The hammerjaw, Omosudis lowii, is a small deep-sea aulopiform fish, found worldwide in tropical and temperate waters to 4,000 m (13,000 ft) depth. It is the only representative of its family, Omosudidae (from the Greek omo, "shoulder", and Latin...



Hampala macrolepidota

Hampala macrolepidota

The hampala barb (Hampala macrolepidota) is a relatively large southeast Asian species of cyprinid from the Mekong and Chao Phraya basins, as well as Peninsular Malaysia and the Greater Sundas (Borneo, Java and Sumatra). It prefers running rivers...



Haplochromis latifasciatus

Haplochromis latifasciatus

Formally Haplochromis latifasciatus, Now Astatotilapia Latifasciata is a species of cichlid that is endemic to Uganda where restricted to the Lake Kyoga system, including Lake Bisina (a medium-sized lake to the east of Lake Kyoga itself) and Lake...



Haplochromis nyererei

Haplochromis nyererei

Haplochromis nyererei is a species of cichlid endemic to Lake Victoria in Africa. This species can reach a length of 7.7 centimetres (3.0 in) SL. The specific name honours Julius Nyerere (1922-1999) who was President of Tanzania from 1961 to...



Haplochromis obliquidens

Haplochromis obliquidens

Haplochromis obliquidens is an African species of cichlid found in Lake Victoria and the adjacent reaches of the Nile. This species can reach a standard length of 8.9 cm (3.5 in). Another species sometimes seen in the aquarium trade, Haplochromis...



Haplochromis vonlinnei

Haplochromis vonlinnei

Haplochromis vonlinnei is a species of cichlid endemic to Lake Victoria. It is greyish in color with a distinct mid-lateral band, and a rather slender shape. It feeds mainly on smaller fish. This species can reach a length of 15.9 centimetres...



Hara jerdoni

Hara jerdoni

Hara jerdoni, common name in English Sylhet Hara; also known as Asian Stone Catfish, or Dwarf Anchor Catfish, is a species of South Asian river catfish native to northeastern India and Bangladesh. This species grows to a length of 4 centimetres...



Hardhead catfish

Ariopsis felis

The hardhead catfish (Ariopsis felis) is a species of sea catfish from the northwest Atlantic and Gulf of Mexico, and similar to the gafftopsail catfish (Bagre marinus). It is one of four species in the genus Ariopsis. The common name, hardhead...



Hardnose shark

Carcharhinus macloti

The hardnose shark (Carcharhinus macloti) is a species of requiem shark, in the family Carcharhinidae, so named because of the heavily calcified cartilages in its snout. A small bronze-coloured shark reaching a length of 1.1 m (3.6 ft), it has a...



Hardyhead silverside

Atherinomorus lacunosus

The hardyhead silverside (Atherinomorus lacunosus), also known as the broad-banded hardyhead, broad-banded silverside, Capricorn hardyhead, pitted hardyhead, robust hardyhead, robust silverside, slender hardyhead and wide-banded hardyhead...



Harlequin fish

Othos dentex

The harlequin fish (Othos dentex) is a species of marine ray-finned fish in the subfamily Anthiinae, part of the family Serranidae, the groupers and sea basses. It is the only member of the genus Othos. It is also known commonly as the Chinese...



Harlequin rasbora

Trigonostigma heteromorpha

The harlequin rasbora (Trigonostigma heteromorpha) is a small fish in the family Cyprinidae. The species became an instant favorite among aquarists after its introduction in the early 1900s and is the best known and most widely kept species among...



Harlequin tuskfish

Choerodon fasciatus

The harlequin tuskfish, *Choerodon fasciatus*, is a species of wrasse native to the western Pacific Ocean. It occasionally makes its way into the aquarium trade.



Hawaiian cleaner wrasse

Labroides phthiophagus

The Hawaiian cleaner wrasse or golden cleaner wrasse (*Labroides phthiophagus*), is a species of wrasse (genus *Labroides*) found in the waters surrounding the Hawaiian Islands. The fish is endemic to Hawaii. These cleaner fish inhabit coral reefs,...



Hawaiian freshwater goby

Lentipes concolor

The Hawaiian freshwater goby, or 'O'opu (Lentipes concolor) ('o'opu 'alamo'o or 'o'opu hi'u koleis), is a species of goby endemic to Hawaii, where it occurs in mountain streams. Males of this species can reach a standard length of 7 cm (2.8 in),...



Hawaiian garden eel

Gorgasia hawaiiensis

The Hawaiian garden eel (Gorgasia hawaiiensis) is an eel in the family Congridae (conger/garden eels). It was described by John Ernest Randall and James Robert Chess in 1980. It is a marine, tropical eel which is known from the Hawaiian...



Hawaiian grouper

Hyporthodus quernus

The Hawaiian grouper (Hyporthodus quernus), also known as the Hawaiian black grouper, Seale's grouper or Hapu'upu'u, is a species of marine ray-finned fish, a grouper from the subfamily Epinephelinae which is part of the family Serranidae, which...



Hawkfish anthias

Serranocirrhitus latus

The hawkfish anthias (Serranocirrhitus latus), also known as the swallowtail basslet, coral perch, hawk anthias, fathead anthias, or sunburst anthias, is a species of marine ray-finned fish, an anthias from the subfamily Anthiinae part of the...



Head-and-taillight tetra

Hemigrammus ocellifer

Hemigrammus ocellifer is a South American freshwater fish, also described by the synonyms Hemibrycon ocellifer and Tetragonopterus ocellifer. Common names include beacon fish, beacon tetra, head-and-tail light tetra and head-and-taillight tetra....



Hector's lanternfish

Lampanyctodes hectoris

Hector's lanternfish (Lampanyctodes hectoris) is a lanternfish in the family Myctophidae, the only species in the genus Lampanyctodes. It is named after James Hector. It is a widespread marine fish, known from shallow tropical waters in the...



Hedgehog seahorse

Hippocampus spinosissimus

The hedgehog seahorse (Hippocampus spinosissimus) is a species of fish of the family Syngnathidae. It inhabits coastal waters from India and Sri Lanka to Taiwan and northern Australia. It is threatened by overfishing, as both targeted catch and...



Helicolenus dactylopterus

Helicolenus dactylopterus

Helicolenus dactylopterus, blackbelly rosefish, bluemouth rockfish, and bluemouth seaperch, is a species of marine ray-finned fish belonging to the subfamily Sebastinae which is classified within the family Scorpaenidae. This Atlantic species is...



Helicolenus percoides

Helicolenus percoides

Helicolenus percoides, the reef ocean perch, coral cod, coral perch, Jock Stewart, kuriarki, ocean perch, red gurnard perch, red gurnard scorpionfish, red ocean perch, red perch, red rock perch, scarpee or sea perch, is a species of marine...



Hemibagrus nemurus

Hemibagrus nemurus

Hemibagrus nemurus is a species of catfishes in the family Bagridae. After a major review by Ng and Kottelat (2013), its distribution is believed to be confined (endemic) to Java.:233 It is found in Sumatra in the Bukit Tigapuluh National Park...



Hemibagrus punctatus

Hemibagrus punctatus

Hemibagrus punctatus, or Nilgiri mystus, is a species of fish in the family Bagridae. It was first described by Jerdon in 1849. It is endemic to east-flowing rivers in the Western Ghats of India. Of these, the species is likely only found in the...



Hemibagrus wyckii

Hemibagrus wyckii

Hemibagrus wyckii is a species of catfish (order Siluriformes) of the family Bagridae. It is occasionally called the crystal-eyed catfish or black devil catfish.



Hemibagrus wyckioides

Hemibagrus wyckioides

Hemibagrus wyckioides, the Asian redtail catfish, is a species of catfish (order Siluriformes) of the family Bagridae.



Hemichromis fasciatus

Hemichromis fasciatus

Hemichromis fasciatus, also called the banded jewelfish and five-spot cichlid, is a species of fish in the cichlid family. It is distributed throughout West Africa. It can also be found in the Nile Basin, Lake Chad, and the upper Zambezi. It can...



Hemichromis frempongi

Hemichromis frempongi

Hemichromis frempongi is a species of fish in the cichlid family. It is endemic to Lake Bosumtwi in Ghana. Its validity as a species is questionable, and some believe it should be included in the widespread H. fasciatus.



Hemichromis lifalili

Hemichromis lifalili

Hemichromis lifalili, common name blood-red jewel cichlid, is a species of fish in the family Cichlidae.



Hemigrammocyppris rasborella

Hemigrammocyppris rasborella

Hemigrammocyppris rasborella, the golden venus chub, is a species of cyprinid fish endemic to the islands of Honshu, Shikoku and Kyushu in Japan. It is found widely in lowland habitats, including ditches and ponds. It is listed as endangered on...



Hemigrammus erythrozonus

Hemigrammus erythrozonus

Hemigrammus erythrozonus, commonly known as the glowlight tetra, is a small tropical fish from the Essequibo River, Guyana, South America. It is silver in colour and a bright iridescent orange to red stripe extends from the snout to the base of...



Hemiramphus far

Hemiramphus far

Hemiramphus far, the halfbeak, black-barred halfbeak, black-barred garfish, barred halfbeak, barred garfish or spotted halfbeak, is a schooling marine fish belonging to the family Hemiramphidae, the halfbeaks. It has an Indo-Pacific distribution...



Hemiscyllium galei

Hemiscyllium galei

Hemiscyllium galei, the Cenderawasih epaulette shark, is a species of bamboo shark in the family Hemiscylliidae. Together with *H. henryi*, it was only scientifically described in 2008 by Gerald R. Allen and Mark V. Erdmann. At present, *H. galei* is...



Hemiscyllium halmahera

Hemiscyllium halmahera

Hemiscyllium halmahera, or the Halmahera epaulette shark, is a species of bamboo shark from Indonesia. This species is described from two specimens collected near Ternate island in 2013, off the coast of larger Halmahera island. This species is...



Hemiscyllium henryi

Hemiscyllium henryi

The Triton epaulette shark (*Hemiscyllium henryi*) is a species of bamboo shark in the genus *Hemiscyllium*, that is composed of nine morphologically similar, yet distinct, sharks that are geographically restricted to New Guinea and northern...



Hemitripteris americanus

Hemitripteris americanus

Hemitripteris americanus, the sea raven, is a species of marine ray-finned fish belonging to the subfamily Hemitripterinae of the family Agonidae. The sea raven is found along the Atlantic coast of North America.



Hemitripteris villosus

Hemitripteris villosus

Hemitripteris villosus, the sea raven, is a species of marine ray-finned fish belonging to the subfamily Hemitripterinae of the family Agonidae. The sea raven is found in the northwestern Pacific Ocean off the coast of Russia and Japan.



Hephaestus carbo

Hephaestus carbo

Hephaestus carbo, the coal grunter or black grunter, is a species of freshwater ray-finned fish, a grunter from the family Terapontidae. It is endemic to rivers in northern Australia.



Hepsetus cuvieri

Hepsetus cuvieri

Hepsetus cuvieri, sometimes known as the African pike or Kafue pike characin, is a predatory freshwater fish found in southern Africa. This species was described in 1861 by the French naturalist Francis de Laporte de Castelnau. The fish is named...



Hepsetus odoe

Hepsetus odoe

Hepsetus odoe, the African pike characin, is a predatory freshwater characin belonging to the family Hepsetidae. It was formerly considered that there was a single species of Hepsetus pike characin but recent studies have led to the species being...



Herichthys carpintis

Herichthys carpintis

Herichthys carpintis, the lowland cichlid, pearl scale cichlid, is a species of cichlid.



Heros efasciatus

Heros efasciatus

Heros efasciatus is a species of tropical freshwater cichlid native to the Amazon basin in South America. It is commonly found in the aquarium trade, often in its gold colored variation rather than the wild-type olive green one.



Heros liberifer

Heros liberifer

Heros liberifer is a species of tropical cichlid native to lentic habitats, especially flooded forests, in the upper and middle Orinoco basin in South America. Up until 2014 the liberifer was incorrectly believed to be the previously defined H....



Heros notatus

Heros notatus

Heros notatus is a species of tropical freshwater cichlid native to Rio Negro and Essequibo River in South America.



Herring scad

Alepes vari

The herring scad (Alepes vari) (also known as the duskyfin crevalle and trevally scad), is a common species of tropical marine fish in the jack family Carangidae. The species inhabits the surface waters of coastal regions throughout the Indo-West...



Heterandria formosa

Heterandria formosa

Heterandria formosa (known as the least killifish, dwarf topminnow, mosqu or midget livebearer) is a species of livebearing fish within the family Poeciliidae. This is the same family that includes familiar aquarium fishes such as guppies and...



Heterobranchus bidorsalis

Heterobranchus bidorsalis

Heterobranchus bidorsalis, the African catfish or eel-like fattyfin catfish, is an airbreathing catfish found in Africa. It is closely related to the vundu catfish, which is well-known among fishermen.



Heteropneustes fossilis

Heteropneustes fossilis

Heteropneustes fossilis, also known as Asian stinging catfish or fossil cat, is a species of airsac catfish found in India, Bangladesh, Pakistan, Nepal, Sri Lanka, Thailand, Myanmar, and Bhutan. It has also been introduced to the Tigris River...



Heteropneustes microps

Heteropneustes microps

Heteropneustes microps is a species of airsac catfish possibly endemic to Sri Lanka, though records from India have been made. This species grows to a total length of 15.0 cm (5.9 in). This fish is a component of local commercial fisheries, and...



Heteropriacanthus

Heteropriacanthus cruentatus

Heteropriacanthus, the glasseyes or glass bigeyes, are a genus of the bigeye family found in all tropical seas around the world. It occasionally makes its way into the aquarium trade. It grows to a size of 50.7 cm (20.0 in) in total length. The...



Heterotilapia buttikoferi

Heterotilapia buttikoferi

Heterotilapia buttikoferi, also known as the zebra tilapia, is a West African species of cichlid.



Hickory shad

Alosa mediocris

The hickory shad (Alosa mediocris), fall herring, mottowacca, freshwater taylor or bonejack is a member of the herring family Clupeidae, ranging along the East Coast of the United States from Florida to the Gulf of Maine. It is an anadromous fish...



Highfin carpsucker

Carpiodes velifer

The highfin carpsucker (Carpiodes velifer) is a freshwater fish found in the southeastern and midwestern United States. Its usual habitat is medium to large-size rivers where it is mostly found 1 to 3 metres (3 to 10 ft) beneath the surface. It...



Himantolophus groenlandicus

Himantolophus groenlandicus

Himantolophus groenlandicus, the Atlantic footballfish or Atlantic football-fish, is an anglerfish found primarily in mesopelagic depths of the ocean. Despite its name, this species might not be restricted to the Atlantic Ocean, with its range...



Hingemouth

Phractolaemus ansorgii

The hingemouth (Phractolaemus ansorgii) is a small freshwater fish that is found only in west central Africa, the sole member of the subfamily Phractolaeminae of the family Kneriidae. The mouth can extend like a small trunk, thus the name, and has...



Hiodon tergisus

Hiodon tergisus

Hiodon tergisus, the mooneye, is a freshwater fish that is widespread across eastern North America.



Hippocampus bargibanti

Hippocampus bargibanti

Hippocampus bargibanti, also known as Bargibant's seahorse or the pygmy seahorse, is a seahorse of the family Syngnathidae found in the central Indo-Pacific area. It is tiny, usually less than 2 centimetres (0.79 in) in size and lives exclusively...



Hippocampus haema

Hippocampus haema

Hippocampus haema, the Korean seahorse, is a seahorse of the family Syngnathidae native to the northern Pacific Ocean (Korea Strait, Sea of Japan, northeastern coast of Honshu), and it usually lives in Sargassum and weeds on shallow soft bottom...



Hippocampus kuda

Hippocampus kuda

Hippocampus kuda is a species of seahorse, also known as the common seahorse, estuary seahorse, yellow seahorse or spotted seahorse. The common name sea pony has been used for populations formerly treated as the separate species Hippocampus...



Hippocampus paradoxus

Hippocampus paradoxus

The paradoxical seahorse (Hippocampus paradoxus) is a small seahorse in the genus Hippocampus. The only known specimen was captured in 1995 and remained unnoticed in a museum until 2006.



Hippocampus patagonicus

Hippocampus patagonicus

The Patagonian seahorse (Hippocampus patagonicus) is a species of marine fish of the family Syngnathidae. It inhabits coastal waters from northeastern Brazil to Chubut, Argentina. It generally is found at shallow depths attached to natural or...



Hippocampus pontohi

Hippocampus pontohi

Hippocampus pontohi, also known as Pontoh's pygmy seahorse or the weedy pygmy seahorse, is a seahorse of the family Syngnathidae native to the central Indo-pacific. Named after Hentje Pontoh, the Indonesian dive guide from Bunaken (Manado) who...



Hippocampus waleananus

Hippocampus waleananus

Hippocampus waleananus, the Walea pygmy seahorse, is a species of seahorse endemic to the Togian Islands in Indonesia, and is associated with specific soft corals. The species was described in 2009 from a single specimen which was found close to...



Hipposcarus harid

Hipposcarus harid

Hipposcarus harid, the Longnose parrotfish or Candelamoa parrotfish, is a species of marine ray-finned fish, a parrotfish from the family Scaridae found on coral reefs of Indian Ocean and the Red Sea.



Hitch

Lavinia exilicauda

The hitch (Lavinia exilicauda) is a cyprinid fish endemic to central California, and was once very common. The common name may derive from a Pomoan word for this species. It is the only species in the monospecific genus Lavinia.



Hog-nosed catfish

Corydoras multiradiatus

The hog-nosed catfish (Corydoras multiradiatus) is a tropical freshwater fish belonging to the Corydoradinae sub-family of the family Callichthyidae. It is native to South America, and is found in the western Amazon basin in Ecuador and Peru....



Hogchoker

Trinectes maculatus

The hogchoker (Trinectes maculatus) is a small species of flatfish found along the Atlantic and Gulf coasts of North and South America, ranging from Massachusetts to Venezuela. They prefer brackish water, and are abundant in many bays and...



Hogfish

Lachnolaimus maximus

The hogfish (Lachnolaimus maximus), also known as boquinete, doncella de pluma or pez perro in Mexico is a species of wrasse native to the Western Atlantic Ocean, living in a range from Nova Scotia, Canada, to northern South America, including...



Holacanthus africanus

Holacanthus africanus

Holacanthus africanus, the Guinean angelfish or West African angelfish, is species of marine ray-finned fish, a marine angelfish belonging to the family Pomacanthidae. It is found in the warmer sections of the eastern Atlantic Ocean off the coast...



Holacanthus bermudensis

Holacanthus bermudensis

Holacanthus bermudensis, also known as the blue angelfish or Bermuda blue angelfish, is a species of ray-finned fish, a marine angelfish belonging to the family Pomacanthidae. It occurs in the western Atlantic Ocean.



Holacanthus passer

Holacanthus passer

Holacanthus passer (king angelfish or passer angelfish) is a large marine angelfish of the family Pomacanthidae. Its large size and bright colors make it a popular aquarium fish, even though it can be difficult to keep.



Holcomycteronus profundissimus

Holcomycteronus profundissimus

Holcomycteronus profundissimus, also known by its synonym Grimaldichthys profundissimus, (Danish: Grimaldis slangekvabbe; Finnish: Grimaldikala; Italian: Pesce di Grimaldi) is a species of deep-sea fish in the cusk-eel family. The fish has an...



Holocentrus adscensionis

Holocentrus adscensionis

Holocentrus adscensionis is a squirrelfish of the family Holocentridae found in the Atlantic Ocean. Its range extends from North Carolina, USA to Brazil and throughout the Caribbean Sea in the Western Atlantic and from Gabon to Ascension Island...



Honey blue-eye

Pseudomugil mellis

The honey blue-eye (Pseudomugil mellis) is an endangered species of fish in the subfamily Pseudomugilinae. It is endemic to southeastern Queensland, Australia, where it is found in mildly acidic, often tannin-stained, ponds and streams in wallum...



Honey gourami

Trichogaster chuna

The honey gourami (Trichogaster chuna) is a species of gourami native to India and Bangladesh.



Honeycomb cowfish

Acanthostracion polygonius

The honeycomb cowfish (Acanthostracion polygonius) belongs to family Ostraciidae. It is also known simply as the trunkfish or cowfish.



Honeycomb grouper

Epinephelus merra

The honeycomb grouper (Epinephelus merra), also known as black-spotted rock-cod, common birdwire rockcod, dwarf spotted rockcod, dwarf-spotted grouper, honeycomb cod, wire-netted reefcod or wire-netting cod, is a species of marine ray-finned...



Honeycomb whipray

Himantura undulata

The honeycomb whipray (Himantura undulata) is a species of stingray in the family Dasyatidae, found widely in the shallow coastal waters of the Indo-Pacific from India to the Malay Archipelago. This large species grows to 1.3 m (4.3 ft) across...



Hong Kong grouper

Epinephelus akaara

The Hong Kong grouper (Epinephelus akaara) is a species of marine ray-finned fish, a grouper from the subfamily Epinephelinae which is part of the family Serranidae, which also includes the anthias and sea basses. It is found in eastern and...



Hooktooth shark

Chaenogaleus macrostoma

The hooktooth shark (Chaenogaleus macrostoma), is a weasel shark of the family Hemigaleidae, the only extant member of the genus Chaenogaleus, but there is an extinct species, Chaenogaleus affinis. The hooktooth shark is found in the tropical...



Hoosier cavefish

Amblyopsis hoosieri

The Hoosier cavefish (Amblyopsis hoosieri) is a subterranean species of blind fish from southern Indiana in the United States. Described in 2014, A. hoosieri was the first new species of amblyopsid cavefish to be discovered in 40 years.



Hoplias aimara

Hoplias aimara

Hoplias aimara, also known as anjumara, traíra, trahira, manjuma, anjoemara and wolf fish, is a species of freshwater fish found in the rivers of South America. In Amazonia, the native populations are concerned by high levels of mercury...



Hoplias curupira

Hoplias curupira

Hoplias curupira, also known as the black wolf-fish, has a wide distribution in the Amazon basin but was described as recently as 2009.



Hoplias malabaricus

Hoplias malabaricus

Hoplias malabaricus, also known as the wolf fish, tiger fish, guabine or trahira, is a predatory Central and South American freshwater ray-finned fish of the characiform family Erythrinidae.



Hoplosternum littorale

Hoplosternum littorale

Hoplosternum littorale is a species of catfish (order Siluriformes) belonging to the Callichthyinae subfamily of the family Callichthyidae. It is known as tamuatá in Brazil, atipa in French Guiana, hassa in Guyana, kwi kwi (or kwie kwie) in...



Horabagrus brachysoma

Horabagrus brachysoma

Horabagrus brachysoma or the sun catfish is a species of catfish endemic to rivers in the Western Ghats of India. It is known as Günther's catfish or yellow catfish. It is also known as Manjakoori in its native range. It is also known by a host...



Horaglanis krishnai

Horaglanis krishnai

Horaglanis krishnai, the Indian blind catfish, is a species of airbreathing catfish endemic to India, mainly in wells and underground water channels around Kottayam, Kerala. It lacks pigmentation and eyes, like other cavefish, and reaches about...



Horn shark

Heterodontus francisci

Sharks portalThe horn shark (*Heterodontus francisci*) is a species of bullhead shark, in the family Heterodontidae. It is endemic to the coastal waters off the western coast of North America, from California to the Gulf of California. Young...



Hornyhead chub

Nocomis biguttatus

The hornyhead chub (*Nocomis biguttatus*) is a small species of minnow in the family Cyprinidae of order Cypriniformes. It mainly inhabits small rivers and streams of the northern central USA, up into Canada. The adults inhabit faster, rocky pools...



Horse-eye jack

Caranx latus

The horse-eye jack (*Caranx latus*), also known as the big-eye jack, is a game fish and minor commercial fish in the family Carangidae. Its appearance is similar to that of the crevalle jack, although the horse-eye jack's head is not as blunt. The...



Houndfish

Tylosurus crocodilus

The houndfish (*Tylosurus crocodilus*) is a game fish of the family Belonidae. It is the largest member of its family, growing up to 5 feet (1.5 m) in length and 10 pounds (4.5 kg) in weight. It is also often called the crocodile needlefish.



Huchen

Hucho hucho

The huchen (*Hucho hucho*) (, from German), also known as Danube salmon or redfish (German: Rotfisch), is a large species of freshwater fish in the family Salmonidae native to the Danube basin in Central and Eastern Europe. It is the type species...



Hucho bleekeri

Hucho bleekeri

Hucho bleekeri, the Sichuan taimen, is a species of freshwater fish in the salmon family (Salmonidae). endemic to the Yangtze basin in China. Their typical habitat includes mountain streams and small rivers, typically being found in the catchment...



Hucho ishikawae

Hucho ishikawae

Hucho ishikawae, the Korean taimen, is a species of salmonid fish found in the border region between North Korea and China, including the Am-nok or Yalu, Dok-ro, Weon-ju and Jang-jin Rivers. Monitoring of the species has been made very difficult...



Hucho taimen

Hucho taimen

Siberian taimen (Hucho taimen), also known as the common taimen (Russian: Обыкновенный таймень, romanized: Obyknovénnyĭ taǐmén'), Siberian giant trout or Siberian salmon, is a species of salmon-like ray-finned fish from the genus Hucho in the...



Humboldt cutthroat trout

Oncorhynchus clarkii humboldtensis

The Humboldt cutthroat trout (Oncorhynchus clarkii humboldtensis) is a subspecies of cutthroat trout, a North American fish in the family Salmonidae. It is one of the several subspecies of cutthroat trout. It was formally scientifically named in...



Humpback anglerfish

Melanocetus johnsonii

The humpback anglerfish (Melanocetus johnsonii) is a species of black seadevils in the family of Melanocetidae, which means "black whale" in Greek. The species is named after James Yate Johnson, the English naturalist who discovered the first...



Humpback chub

Gila cypha

The humpback chub (Gila cypha) is a federally protected fish that lived originally in fast waters of the Colorado River system in the United States. This species takes its name from the prominent hump between the head and dorsal fin, which is...



Humpback grouper

Cromileptes altivelis

The humpback grouper (Cromileptes altivelis), panther grouper, or (in Australia) barramundi cod is a species of marine ray-finned fish, a grouper from the subfamily Epinephelinae which is in the family Serranidae which also includes the anthias...



Humpback mahseer

Hypselobarbus mussullah

The humpback mahseer (*Hypselobarbus mussullah*) is a species of freshwater ray-finned fish from the Indian endemic genus *Hypselobarbus* in the carp and minnow family Cyprinidae.



Humpback red snapper

Lutjanus gibbus

The humpback red snapper (*Lutjanus gibbus*), the paddletail, paddletail snapper or hunchback snapper, is a species of marine ray-finned fish, a snapper belonging to the family Lutjanidae. It has a wide Indo-West Pacific distribution. It is a...



Humpback whitefish

Coregonus pidschian

The humpback whitefish (*Coregonus pidschian*), also referred to as the bottom whitefish, the Arctic whitefish or the pidschian, is a species of freshwater whitefish with a northern distribution. It is one of the members in the broader common...



Humphead wrasse

Cheilinus undulatus

The humphead wrasse (*Cheilinus undulatus*) is a large species of wrasse mainly found on coral reefs in the Indo-Pacific region. It is also known as the Māori wrasse, Napoleon wrasse, Napoleon fish, so mei 蘇眉 (Cantonese), mameng (Filipino), and...



Hydrocynus brevis

Hydrocynus brevis

Hydrocynus brevis, also known as the tigerfish, Nile tigerfish or Sahelian tigerfish, is a predatory freshwater fish distributed throughout Africa.



Hydrocynus forskahlii

Hydrocynus forskahlii

Hydrocynus forskahlii, the elongate tigerfish, is a species of predatory characin from the family Alestidae which is found in northern and western Africa.



Hydrocynus goliath

Hydrocynus goliath

Hydrocynus goliath, also known as the goliath tigerfish, giant tigerfish, or mbenga, is a very large African predatory freshwater fish of the family Alestidae.



Hydrocynus tanzaniae

Hydrocynus tanzaniae

Hydrocynus tanzaniae, the blue tigerfish, is a large African predatory freshwater fish.



Hydrocynus vittatus

Hydrocynus vittatus

Hydrocynus vittatus, the African tigerfish, tiervis or ngwesh is a predatory freshwater fish distributed throughout much of Africa. This fish is generally a piscivore but it has been observed leaping out of the water and catching barn swallows in...



Hydrolycus armatus

Hydrolycus armatus

Hydrolycus armatus is a species of dogtooth characin found in freshwater of tropical South America. It is sometimes known as the payara, a name it shares with the related H. scomberoides. This predatory fish occasionally makes its way into the...



Hydrolycus tatauaia

Hydrolycus tatauaia

Hydrolycus tatauaia is a species of dogtooth characin found in the Amazon, Orinoco and Essequibo basins in tropical South America. Adults mainly occur in deep and/or fast-flowing rivers. It is migratory, moving upstream to breed in...



Hypancistrus contradens

Hypancistrus contradens

Hypancistrus contradens is a species of armored catfish endemic to Venezuela where it occurs in the Orinoco River.



Hyperoglyphe antarctica

Hyperoglyphe antarctica

Hyperoglyphe antarctica, the Antarctic butterfish, bluenose warehou, deepsea trevally, blue eye trevalla, blue-eye cod, bluenose sea bass, or deep sea trevalla, is a medusafish of the family Centrolophidae found in all the southern oceans, at...



Hyperopisus

Hyperopisus bebe

The Bebe mormyrid (Hyperopisus bebe) is a species of elephantfish in the family Mormyridae. It is the only species in its genus and is divided into two subspecies. This species is known from many rivers in the northern half of Africa, ranging...



Hyphessobrycon bentosi

Hyphessobrycon bentosi

Hyphessobrycon bentosi, the Bentos tetra, ornate tetra or candy cane tetra, is a species of characin fish found in sluggish tributaries at the Amazon Basin in Brazil and Peru. Occasionally, it makes its way into the aquarium trade. It has often...



Hyphessobrycon columbianus

Hyphessobrycon columbianus

Hyphessobrycon columbianus (Colombian tetra or blue-red Colombian tetra) is a species of Tropical freshwater fish of the characin family. The Colombian tetra is native to the Acandí River near Acandí) in northwestern Colombia. The Colombian...



Hyphessobrycon sweglesi

Hyphessobrycon sweglesi

Hyphessobrycon sweglesi is a species of tetra that lives in the Orinoco River drainage basin in South America. The species' scientific name used to be Megalamphodus sweglisi and the species' common name is red phantom tetra. It grows up to 4 cm...



Hypomesus japonicus

Hypomesus japonicus

Hypomesus japonicus, the Japanese smelt, is a coastal fish species of the northwestern Pacific Ocean, ranging from the Korean Peninsula and northern Japan to the Kuril Islands and Peter the Great Bay.



Hypomesus nipponensis

Hypomesus nipponensis

Hypomesus nipponensis (Japanese smelt, in Japanese: wakasagi) is a commercial food fish native to the lakes and estuaries of northern Honshu and Hokkaido, Japan, Korea, and Sakhalin, Khabarovsk Krai, and Primorsky Krai, Russia. It has been...



Hypomesus pretiosus

Hypomesus pretiosus

Hypomesus pretiosus, or surf smelt, is a marine smelt with a range from Prince William Sound, Alaska to Long Beach, California, although its population declines south of San Francisco. The surf smelt grows to be about 10 inches in southern...



Hypoplectrus gemma

Hypoplectrus gemma

Hypoplectrus gemma, the blue hamlet, is a species of marine ray-finned fish, a sea bass from the subfamily Serraninae which is part of the family Serranidae, which also includes the groupers and anthias. It is found in the western Atlantic Ocean...



Hypoplectrus indigo

Hypoplectrus indigo

Hypoplectrus indigo, the indigo hamlet, is a species of marine ray-finned fish, a sea bass from the subfamily Serraninae which is part of the family Serranidae, which also includes the groupers and anthias. It is found in the central western...



Hypoplectrus nigricans

Hypoplectrus nigricans

Hypoplectrus nigricans, the black hamlet, is a species of marine ray-finned fish, a sea bass from the subfamily Serraninae which is part of the family Serranidae, which also includes the groupers and anthias. It is native to shallow parts of the...



Hyporhamphus ihi

Hyporhamphus ihi

Hyporhamphus ihi, the known as the garfish, piper or by its Māori name takeke, is a halfbeak found all around New Zealand in shallow inshore waters.



Hyporthodus nigrinus

Hyporthodus nigrinus

Hyporthodus nigrinus, the Warsaw grouper, is a species of marine ray-finned fish, a grouper from the subfamily Epinephelinae which is part of the family Serranidae, which also includes the anthias and sea basses. It is found in the Western...



Hypostomus luteus

Hypostomus luteus

Hypostomus luteus, also known in the pet trade as the golden sailfin pleco, is an armored catfish in the genus Hypostomus of the family Loricariidae. The species experiences a notable three-phase shift in its coloration as it ages. It grows to a...



Hypostomus plecostomus

Hypostomus plecostomus

Hypostomus plecostomus, also known as the suckermouth catfish or the common pleco, is a tropical freshwater fish belonging to the armored catfish family (Loricariidae), named for the longitudinal rows of armor-like scutes that cover the upper...



Hypostomus punctatus

Hypostomus punctatus

Hypostomus punctatus, the suckermouthed catfish, is a tropical fish belonging to the armored suckermouth catfish family, Loricariidae. Hypostomus punctatus is a freshwater fish native to South America, in the coastal drainages of southeastern...



Hypseleotris compressa

Hypseleotris compressa

Hypseleotris compressa, the empire gudgeon, is a species of Gobiiform fish in the family Eleotridae endemic to Australia and south-central New Guinea.



Hypseleotris galii

Hypseleotris galii

Hypseleotris galii, the firetail gudgeon, is a species of gudgeon native to eastern Australia, where it is found in freshwater streams. Gale's carp-gudgeon is an alternate common name.



Hypselobarbus carnaticus

Hypselobarbus carnaticus

Hypselobarbus carnaticus, also known as the Carnatic carp, is a species of cyprinid fish from the Western Ghats in India where it inhabits riffles and larger pools in rapidly flowing rivers and streams. It prefers to shelter underneath boulders...



Hypsophrys nicaraguensis

Hypsophrys nicaraguensis

Hypsophrys nicaraguensis, the moga, is a species of cichlid native to the Atlantic slope of Central America, from Nicaragua to Costa Rica. The species is a popular aquarium fish and is traded under a variety of common names that include nickie,...



Ice goby

Leucopsarion petersii

The ice goby (Leucopsarion petersii) is a species of goby in the subfamily, Gobionellinae, and the only member of the monotypic genus Leucopsarion. It is native to the northwestern Pacific Ocean, where it occurs in China, Japan, and Korea. Its...



Ictalurus pricei

Ictalurus pricei

Ictalurus pricei, the Yaqui catfish, is a species of North American freshwater catfish native to Mexico and Arizona.



Ide

Leuciscus idus

The ide (Leuciscus idus), or orfe, is a freshwater fish of the family Cyprinidae found in larger rivers, ponds, and lakes across Northern Europe and Asia. It has been introduced outside its native range into Europe, North America, and New...



Idiacanthus atlanticus

Idiacanthus atlanticus

Idiacanthus atlanticus, the black dragonfish, is a barbeled dragonfish of the family Stomiidae, found circumglobally in southern subtropical and temperate oceans between latitudes 25°S and 60°S, at depths down to 2,000 metres (6,600 ft). The...



Ilish

Tenualosa ilisha

The ilish (Tenuulosa ilisha) (Bengali: ইলিশ, romanized: ilíś), also known as the ilishi, hilsa, hilsa herring or hilsa shad, is a species of fish related to the herring, in the family Clupeidae. It is a very popular and sought-after food fish in...



Ilisha africana

Ilisha africana

Ilisha africana, called the West African ilisha, is a species of longfin herring native to the coasts, lagoons and estuaries of western Africa from Senegal to Angola. It prefers shallow waters, and is unlikely to be found below 25 m. Some...



Ilisha elongata

Ilisha elongata

The elongate ilisha (Ilisha elongata), also known as the Chinese herring (simplified Chinese: 勒鱼; traditional Chinese: 勒魚; pinyin: lèyú, or simply 鰵; 鰵; lè) or slender shad (although not a true herring or shad), is a species of longfin herring...



Indian anchovy

Stolephorus indicus

The Indian anchovy (Stolephorus indicus), also known as Hardenberg's anchovy, is a species of oceanodromous ray-finned fish in the family Engraulidae. It is known as handalla (හැන්දලා) in Sri Lanka, where it is widely sold at most markets and...



Indian flying barb

Esomus danrica

The Indian flying barb (Esomus danrica), historically flying barb, is one of the species known in the group flying barbs owing to their extremely long barbels. It was discovered as long ago as 1822 by Hamilton. However, it is rarely seen in...



Indian lizardfish

Synodus indicus

The Indian lizardfish (Synodus indicus) is a species of lizardfish that lives mainly in the Indo-West Pacific Ocean.



Indian mackerel

Rastrelliger kanagurta

The Indian mackerel (Rastrelliger kanagurta) is a species of mackerel in the scombrid family (family Scombridae) of order Perciformes. It is commonly found in the Indian and WestPacific oceans, and their surrounding seas. It is an important food...



Indian mud moray eel

Gymnothorax tile

The Indian mud moray eel, (Gymnothorax tile) is a moray eel found in the western Pacific and Indian Oceans. It was first named by Hamilton in 1822, and is also commonly known as the freshwater moray or freshwater snowflake eel.



Indian Ocean oriental sweetlips

Plectorhinchus vittatus

The Indian Ocean oriental sweetlips (*Plectorhinchus vittatus*), also known as the oriental sweetlips or oriental blubberlips, is a species of marine ray-finned fish, a sweetlips belonging to the subfamily Plectorhinchinae, one of two subfamilies...



Indian threadfin

Leptomelanosoma indicum

The Indian threadfin (Leptomelanosoma indicum) is a species of marine ray-finned fish from the family Polynemidae, the threadfins. It is a coastal species from south-east Asia which has been recorded in Papua New Guinea.



Indian threadfish

Alectis indica

The Indian threadfish (Alectis indica), also known as the Indian threadfin, diamond trevally, mirror fish or plumed trevally, is a large species of coastal marine fish of the jack family, Carangidae. The species is widespread in the waters of the...



Indian triggerfish

Melichthys indicus

The Indian triggerfish (Melichthys indicus), also known as the black-finned triggerfish, has a brown body and black fins with white lines at the base of the dorsal and anal fins. It is found across the Indian Ocean. They can grow up to 10 in (25...



Indo-Pacific blue marlin

Makaira mazara

The Indo-Pacific blue marlin (Makaira mazara) is a species of marlin belonging to the family Istiophoridae.



Indo-Pacific king mackerel

Scomberomorus guttatus

Indo-Pacific king mackerel (*Scomberomorus guttatus*), also known as the spotted seer fish or spotted Spanish mackerel, is a sea fish among the mackerel variety of fishes. It is found in around the Indian Ocean and adjoining seas. It is a popular...



Indo-Pacific sailfish

Istiophorus platypterus[1]

The Indo-Pacific sailfish (*Istiophorus platypterus*) is a sailfish native to the Indian and Pacific Oceans and is naturalized in the Atlantic where it has entered the Mediterranean Sea via the Suez Canal as a Lessepsian migrant. It is dark blue on...



Indo-Pacific sergeant

Abudefduf vaigiensis

The Indo-Pacific sergeant (*Abudefduf vaigiensis*) is a species of damselfish in the family Pomacentridae. It may also be known as the Sergeant major although this name is usually reserved for the closely related species *Abudefduf saxatilis*. The...



Indo-Pacific tarpon

Megalops cyprinoides

The Indo-Pacific tarpon (*Megalops cyprinoides*), also known as the oxeye herring or simply herring due to its superficial resemblance to the true herrings, of which it is not a member, is the smaller of the two species of tarpon and lives in...



Indonesian speckled carpetshark

Hemiscyllium freycineti

The Indonesian speckled carpetshark, Hemiscyllium freycineti, is a species of bamboo shark in the family Hemiscylliidae. It is found in the shallow ocean around the Raja Ampat Islands in West Papua, Indonesia, but was formerly believed to be more...



Indonesian wobbegong

Orectolobus leptolineatus

The Indonesian wobbegong (Orectolobus leptolineatus) is a species of carpet shark in the family Orectolobidae, that can grow up to 120 cm in length. It lives in the Western Pacific on the ocean floor, and is harmless to humans.



Iniistius pavo

Iniistius pavo

Iniistius pavo, the peacock wrasse or blue razorfish, is a species of marine ray-finned fish from the family Labridae, the wrasses, which has a wide Indo-Pacific distribution.



Inimicus didactylus

Inimicus didactylus

Inimicus didactylus, also known as sea goblin, demon stinger or devil stinger, is a Western Pacific member of the Inimicus genus of venomous fishes, closely related to the true stonefishes. It can reach a body length of 25 cm (10 in) and is...



Inimicus filamentosus

Inimicus filamentosus

Inimicus filamentosus, also known as the filament-finned stinger, barred ghoul, two-stick stingfish, or devil scorpionfish, is a member of the Inimicus genus of venomous fishes. It is a member of the Synanceiidae (devilfishes, goblinfishes, and...



Inland silverside

Menidia beryllina

The inland silverside (Menidia beryllina) is a neotropical silverside native to eastern North America, and introduced into California. It is a fish of estuaries and freshwater environments. Inland silversides are quite elongate even for...



Inshore lizardfish

Synodus foetens

The inshore lizardfish (Synodus foetens) is a member of the family Synodontidae found in the western Atlantic.



Iodotropheus sprengerae

Iodotropheus sprengerae

Iodotropheus sprengerae, the rusty cichlid, lavender mbuna or lavender cichlid, is a species of cichlid endemic to the Boadzulu and Chinyankwazi Islands and Chinyamwezi Island regions of Lake Malawi. It prefers a temperature range of 24-26...



Ipnops meadi

Ipnops meadi

Ipnops meadi, also known as the grideye fish, is a highly specialized species of Placodithyran abyssal fish found in the Clarion-Clipperton Zone of the Indo-Pacific Ocean. The species was named after Giles W. Mead of the Harvard Museum of...



Iridescent shark

Pangasianodon hypophthalmus

The iridescent shark, iridescent shark catfish (Pangasianodon hypophthalmus) is a species of shark catfish (family Pangasiidae) native to the rivers of Southeast Asia. Despite its name, it is not a shark. It is found in the Mekong basin as well...



Island mackerel

Rastrelliger faughni

The island mackerel (Rastrelliger faughni) is a species of true mackerel in the scombrid family (Scombridae) from the Indo-Pacific. Their maximum reported length is 20 cm, and the maximum reported weight is 0.75 kg. While the FAO report no...



Island trevally

Carangoides orthogrammus

The island trevally, island jack, thicklip trevally or false bluefin trevally (Carangoides orthogrammus) is a widespread species of offshore marine fish classified in the jack family Carangidae. The island trevally is common through the tropical...



Izak catshark

Holohalaelurus regani

The Izak catshark or simply Izak (Holohalaelurus regani) is a species of catshark, belonging to the family Scyliorhinidae, common off the coasts of South Africa and southern Namibia. It typically inhabits the outer continental shelf at depths of...



Jack Dempsey

Rocio octofasciata

The Jack Dempsey (Rocio octofasciata) is a species of cichlid fish that is native to freshwater habitats from southern Mexico to Honduras, but also introduced elsewhere. Its common name refers to its aggressive nature and strong facial features....



Jack silverside

Atherinopsis californiensis

Atherinopsis californiensis, the jack silverside or jacksmelt, is a species of neotropical silverside native to the Pacific coast of North America from Oregon, United States to southern Baja California, Mexico. This species grows to 45 cm (18 in)...



Jaggedhead gurnard

Gargariscus prionocephalus

The jaggedhead gurnard (Gargariscus prionocephalus) is a species of marine ray-finned fish belonging to the family Peristediidae, the armoured gurnards or armored searobins. This species is found in the eastern Indian Ocean and the western...



Jaguar catshark

Bythaelurus giddingsi

The jaguar catshark (Bythaelurus giddingsi), also known as the Galápagos catshark, is a species of catshark from the Galápagos Islands. The species was first described in 2012. This catshark is about a foot long when mature, and it is colored...



Japanese amberjack

Seriola quinqueradiata

The Japanese amberjack or yellowtail, Seriola quinqueradiata, is a species of jack fish in the family Carangidae, native to the northwest Pacific Ocean. It is known as shiyu (鰺魚) in China, bang'eo (방어) in Korea, and buri (鰺) or hamachi (鰺) in...



Japanese anchovy

Engraulis japonicus

The Japanese anchovy (*Engraulis japonicus*) is a schooling fish of the family Engraulidae. It is common in the Pacific Ocean south from the Sea of Okhotsk, widespread in the Sea of Japan, Yellow Sea, and East China Sea, and near the coasts of...



Japanese angelfish

Centropyge interrupta

The Japanese angelfish (*Centropyge interrupta*) or Japanese pygmy angelfish, is a species of marine ray-finned fish, a marine angelfish belonging to the family Pomacanthidae. It is found in the western Pacific Ocean.



Japanese angelshark

Squatina japonica

The Japanese angelshark (*Squatina japonica*) is a species of angelshark, family Squatinidae, found in the northwestern Pacific Ocean off China, Japan, and Korea. It is a bottom-dwelling shark found in sandy habitats down to 300 m (980 ft) deep....



Japanese bullhead shark

Heterodontus japonicus

The Japanese bullhead shark (*Heterodontus japonicus*) is a species of bullhead shark in the family Heterodontidae found in the northwestern Pacific Ocean off the coasts of Japan, Korea, and China. This benthic shark occurs at depths of 6–37 m...



Japanese catshark

Apristurus japonicus

Sharks portalThe Japanese catshark (Apristurus japonicus) is a catshark of the family Scyliorhinidae, found in the northwest Pacific off Chiba Prefecture, Honshū, Japan, between 36 and 34°N. This shark has a relatively slender body, with the...



Japanese eel

Anguilla japonica

The Japanese eel (*Anguilla japonica*; nihon unagi (日本鰻)) is a species of anguillid eel found in Japan, Korea, Taiwan, China, and Vietnam, as well as the northern Philippines. There are three main species under the *Anguilla* genus, and all three...



Japanese fluvial sculpin

Cottus pollux

The Japanese fluvial sculpin (*Cottus pollux*) is a species of freshwater ray-finned fish belonging to the family Cottidae, the typical sculpins. It is endemic to Japan, where it inhabits mountain streams in Honshu, Shikoku, and Kyushu. It reaches...



Japanese gissu

Pterothrissus gissu

Pterothrissus gissu, also known as the Japanese gissu, is a species of ray-finned fish in the family Albulidae. The Japanese gissu is a rare fish that is distributed in deep water off northwest Pacific Ocean. This fish is known to pass through a...



Japanese jack mackerel

Trachurus japonicus

The Japanese jack mackerel (*Trachurus japonicus*), also known as the Japanese horse mackerel or Japanese scad, is a species named after its resemblance to mackerel but which is in the family Carangidae, the jacks, pompanos, trevallies and scads....



Japanese lates

Lates japonicus

The Japanese lates (*Lates japonicus*), also known as the akame (from the Japanese 赤目, アカメ, literally "red eye") or Japanese barramundi, is a species of fish found in Japan. A bottom-dweller restricted to estuarine habitats and large rivers in the...



Japanese rice fish

Oryzias latipes

The Japanese rice fish (*Oryzias latipes*), also known as the medaka, is a member of genus *Oryzias* (ricefish), the only genus in the subfamily Oryziinae. This small (up to about 3.6 cm or 1.4 in) native of Japan is a denizen of rice paddies....



Japanese roughshark

Oxynotus japonicus

Sharks portalThe Japanese roughshark (*Oxynotus japonicus*) is a rare species of shark in the family Oxynotidae, known only from a handful of specimens recovered from Suruga Bay and the Enshunada Sea off Japan. It is a benthic species that occurs...



Japanese sandfish

Arctoscopus japonicus

The Japanese sandfish (*Arctoscopus japonicus*), also known as the sailfin sandfish, is a species of fish of the Percomorpha (perch-like) clade in the order Trachiniformes, being one of the two genera in the family Trichodontidae, the sandfishes....



Japanese sawshark

Pristiophorus japonicus

Sharks portalThe Japanese sawshark (Pristiophorus japonicus) is a species of sawshark in the family Pristiophoridae. This shark has a long, narrow rostrum. Its first dorsal fin originates behind the tips of the pectoral fins, and its caudal fin...



Japanese sea bass

Lateolabrax japonicus

The Japanese sea bass (Lateolabrax japonicus) is a species of catadromous marine ray-finned fish from the Asian sea bass family Lateolabracidae which is found in the Western Pacific.



Japanese seahorse

Hippocampus mohnikei

The Japanese seahorse (in Japanese, kitano-umi-uma and sangotatsu) or lemur-tail seahorse (Hippocampus mohnikei) is a species of fish in the family Syngnathidae. The Japanese seahorse reaches a maximum length of 8.0 cm, is usually dark brown and...



Japanese snapper

Paracaesio caerulea

The Japanese snapper (Paracaesio caerulea) is a species of marine ray-finned fish, a snapper belonging to the family Lutjanidae. It is native to the Western Pacific Ocean.



Japanese Spanish mackerel

Scomberomorus niphonius

The Japanese Spanish mackerel (Scomberomorus niphonius), also known as the Japanese seer fish, is a species of true mackerel in the scombrid family (Scombridae). Their maximum reported length is 100 cm, and the maximum reported weight is 10.57 kg.



Japanese sturgeon

Acipenser schrenckii

The Japanese sturgeon or Amur sturgeon (Acipenser schrenckii) is a species of fish in the family Acipenseridae found in the Amur River basin in China and Russia. Claims of its presence in the Sea of Japan need confirmation. The species has 11–16...



Japanese topeshark

Hemitriakis japonica

The Japanese topeshark (Hemitriakis japonica) is a species of houndshark, in the family Triakidae. It can reach a length of up to 1.1 m. It is found in the subtropical northwest Pacific from China, Taiwan, Korea, and Japan, between latitudes 40°...



Japanese white crucian carp

Carassius cuvieri

The Japanese white crucian carp, also known as Japanese carp, white crucian carp, or gengoro-buna (Carassius cuvieri), is a species of freshwater fish in the carp family (family Cyprinidae). It is found in Japan and, as an introduced species, in...



Japanese whiting

Sillago japonica

The Japanese whiting (*Sillago japonica*), also known as the Japanese sillago or Shiro-gisu, is a common species of coastal marine fish belonging to the smelt-whiting family, Sillaginidae. As suggested by its name, the Japanese whiting was first...



Japanese wobbegong

Orectolobus japonicus

The Japanese wobbegong (*Orectolobus japonicus*) is a carpet shark in the family Orectolobidae of the wobbegong family, found in the tropical western Pacific Ocean from Japan and Korea to Vietnam and the Philippines, between latitudes 43 and 6°N....



Java barb

Barbonymus gonionotus

The Java barb (*Barbonymus gonionotus*; Thai: ตะเพียน Ta-phian; Lao: Pa keng; Khmer: ត្រីម្លិះ Trey Chpin; Indonesian: Tawes; Vietnamese: Mè Vinh), more commonly known as silver barb in aquaculture, is a species of ray-finned fish in the genus...



Java loach

Pangio oblonga

The Java loach (*Pangio oblonga*) is a species of tropical freshwater fish, an unbanded kuhli loach, native to the sandy streams of Southeast Asia. Its alternative common names include the black kuhli loach, chocolate kuhli loach and cinnamon...



Javaen barb

Systemus rubripinnis

The Javaen barb (Systemus rubripinnis), is a species of cyprinid fish native to Southeast Asia. It can reach a length of 25 centimetres (9.8 in) SL and is of minor importance to local commercial fisheries.



Jenkins' whipray

Pateobatis jenkinsii

The Jenkins' whipray (Pateobatis jenkinsii) is a species of stingray in the family Dasyatidae, with a wide distribution in the Indo-Pacific region from South Africa to the Malay Archipelago to northern Australia. This large species grows to 1.5 m...



John Dory

Zeus faber

John Dory, St Pierre or Peter's fish, refers to fish of the genus Zeus, especially Zeus faber, of widespread distribution. It is an edible demersal coastal marine fish with a laterally compressed olive-yellow body which has a large dark spot, and...



Johnny darter

Etheostoma nigrum

The johnny darter (Etheostoma nigrum) is a species of freshwater ray-finned fish, a darter from the subfamily Etheostomatinae, part of the family Percidae, which also contains the perches, ruffes and pikeperches. It is native to shallow waters...



Johnrandallia

Johnrandallia nigrirostris

The blacknosed butterflyfish or barberfish (*Johnrandallia nigrirostris*) (from the Spanish names, El Barbero or Mariposa Barbero, "the barber" or "butterfly barber"), is a species of fish in the family Chaetodontidae, the butterfly fishes. It is...



Jolthead porgy

Calamus bajonado

The jolthead porgy (*Calamus bajonado*) is an ocean-going species of fish in the family Sparidae. In Bermuda, it is known as the blue bone porgy, in the United States, it is also known by the Spanish name bojanado, in Jamaica, it is one of the...



Julidochromis marlieri

Julidochromis marlieri

Julidochromis marlieri is a species of cichlid endemic to Lake Tanganyika where it is only known from the northwestern portion preferring rocky shorelines in deep waters. In the aquarium trade, it is commonly known as Marlier's Julie, Spotted...



Jullien's golden carp

Probarbus jullieni

The Jullien's golden carp (*Probarbus jullieni*) is a species of endangered freshwater ray-finned fish in the family Cyprinidae found in Southeast Asian river basins. Its existence is being threatened by various economic activities, such as...



June sucker

Chasmistes liorus

The June sucker (*Chasmistes liorus*) is an endangered species of fish endemic to Utah Lake and the Provo River in the U.S. state of Utah. It is named after the month in which it spawns. It is a gray or brownish fish with a paler belly, growing up...



Kaluga

Huso dauricus

The kaluga (Huso dauricus), also known as the river beluga, is a large predatory sturgeon found in the Amur River basin. With a maximum size of at least 1,000 kg (2,205 lb) and 5.6 m (18.6 ft), the kaluga is one of the biggest of the sturgeon...



Kamchatka flounder

Atheresthes evermanni

The Kamchatka flounder, *Atheresthes evermanni*, is a flatfish of the family Pleuronectidae. It is a demersal fish that lives at depths of between 20 metres (66 ft) and 1,200 metres (3,900 ft). Its native habitat is the temperate waters of the...



Kamchatkan rainbow trout

Oncorhynchus mykiss mykiss

The Kamchatkan rainbow trout (*Oncorhynchus mykiss mykiss*) is a subspecies of the rainbow trout, which is a fish in the family Salmonidae. It is native to Russian Far East. Its main range is on the Kamchatka Peninsula, and it has also been...



Kamloops rainbow trout

Oncorhynchus mykiss kamloops

The Kamloops rainbow trout or Kamloops trout (*Oncorhynchus mykiss kamloops*) is a local variety of the rainbow trout, a fish in the family Salmonidae. From its native range in British Columbia, Canada, it has been transferred to several other...



Kampango

Bagrus meridionalis

The kampango or kampoyo (*Bagrus meridionalis*) is a critically endangered species of large and predatory bagrid catfish that is endemic to Lake Malawi, Lake Malombe and the upper Shire River in Africa. It prefers areas near rocks in water...



Kanadi kingfish

Scomberomorus plurilineatus

The Kanadi kingfish (Scomberomorus plurilineatus) is a species of ray-finned bony fish in the family Scombridae, the mackerel family. Also known as the Kanadi seerfish, queen mackerel, or spotted mackerel, it is found in subtropical waters of the...



Kaup's arrowtooth eel

Synaphobranchus kaupii

The Kaup's arrowtooth eel (Synaphobranchus kaupii, also known as the Kaup's cut-throat eel, the Gray's cutthroat, the Longnosed eel, the Northern cutthroat eel, or the Slatjaw cutthroat eel) is an eel in the family Synaphobranchidae (cutthroat...



Keeltail needlefish

Platybelone argalus

The keeltail needlefish (Platybelone argalus), sometimes called the keeled needlefish, is a tropical fish of the family Belonidae. It was described by the French naturalist Charles Alexandre Lesueur in 1821.



Kelp greenling

Hexagrammos decagrammus

The kelp greenling (Hexagrammos decagrammus) is a species of marine ray-finned fish belonging to the family Hexagrammidae, the greenlings. It occurs in the eastern Pacific Ocean



Kern River rainbow trout

Oncorhynchus mykiss gilberti

The Kern River rainbow trout (*Oncorhynchus mykiss gilberti*) is a localized subspecies of the rainbow trout (*Oncorhynchus mykiss*), a variety of fish in the family Salmonidae. It is found in a short section of the main stem of the Kern River and...



Keyhole cichlid

Cleithracara maronii

The keyhole cichlid (*Cleithracara maronii*) is a cichlid fish endemic to tropical South America, occurring in the lower Orinoco Basin in Venezuela and river basins in The Guianas. It is the only species in the genus *Cleithracara*. The species is...



Kidako moray

Gymnothorax kidako

The Kidako moray (*Gymnothorax kidako*) is a species of marine fish in the family Muraenidae. It inhabits coral reefs or lagoons and could be found in tropical and subtropical seas near Taiwan, Japan, and Australia. The species is diurnal, which...



Killarney shad

Alosa killarnensis

The Killarney shad (Alosa killarnensis), also called the goureen, is a freshwater fish in the family Clupeidae, endemic to a single lake in Ireland, Lough Leane in County Kerry. Research has shown that it is a landlocked subspecies of the...



King dory

Cyttus traversi

The king dory or lookdown dory (Cyttus traversi) is a Dory, in the genus Cyttus, found around South Africa, southern Australia, and New Zealand, over the continental shelf at depths of between 200 and 800 m. Its length is between 20 and 40 cm.



King George whiting

Sillaginodes punctatus

The King George whiting (Sillaginodes punctatus), also known as the spotted whiting or spotted sillago, is a coastal marine fish of the smelt-whitings family Sillaginidae. The King George whiting is endemic to Australia, inhabiting the south...



King mackerel

Scomberomorus cavalla

The king mackerel (Scomberomorus cavalla) surmayi or kingfish, is a migratory species of mackerel of the western Atlantic Ocean and Gulf of Mexico. It is an important species to both the commercial and recreational fishing industries.



King threadfin

Polydactylus macrochir

The king threadfin (Polydactylus macrochir), also known as the blind salmon, blink tassel-fish, burnett salmon, gold threadfin, king salmon, kingfish, Sheridan threadfin, triped tassel fish, or threadfin salmon, is a species of marine ray-finned...



King-of-the-salmon

Trachipterus altivelis

King-of-the-salmon, Trachipterus altivelis, is a species of ribbonfish in the family Trachipteridae. Its common name comes from the legends of the Makah people west of the Strait of Juan de Fuca, which hold that this fish leads the salmon...



Kingklip

Genypterus capensis

Genypterus capensis, commonly known as kingklip, is a species of cusk eel occurring along the Southern African coast from Walvis Bay in Namibia to Algoa Bay in South Africa. It is closely related to Genypterus blacodes from New Zealand. The...



Kirikuchi char

Salvelinus leucomaenis japonicus

The kirikuchi char, Salvelinus leucomaenis japonicus, is a freshwater fish in the family Salmonidae. It is endemic to the Kii Peninsula of central Honshu in Japan. It is the southernmost population of the char genus Salvelinus and is considered a...



Kissing gourami

Helostoma temminckii

Kissing gouramis, also known as kissing fish or kissers (*Helostoma temminckii*), are medium-sized tropical freshwater fish comprising the monotypic labyrinth fish family Helostomatidae (from the Greek *elos*, stoma). These fish originate from...



Kitefin shark

Dalatias licha

The kitefin shark or seal shark (*Dalatias licha*) is a species of squaliform shark in the family Dalatiidae, and the type species in its genus. It is found sporadically around the world, usually close to the sea floor at depths of 200–600 m...



Kiyi

Coregonus kiyi

The kiyi (Coregonus kiyi) is a species of freshwater whitefish, a deepwater cisco, endemic to the Great Lakes of North America. It previously inhabited Lake Superior, Lake Michigan, Lake Huron and Lake Ontario, but is now believed to persist only...



Klunzinger's wrasse

Thalassoma rueppellii

Klunzinger's wrasse (Thalassoma rueppellii), also known as Rüppell's wrasse, is a species of ray-finned fish, a wrasse from the family Labridae which is endemic to the Red Sea. It inhabits the margins and seaward slopes of reefs at depths from 1...



Knifetooth sawfish

Anoxypristis cuspidata

The narrow sawfish (Anoxypristis cuspidata), also known as the pointed sawfish or knifetooth sawfish, is a species of sawfish in the family Pristidae, part of the Batoidea, a superorder of cartilaginous fish that include the rays and skates....



Knobsnout parrotfish

Scarus ovifrons

The knobsnout parrotfish (Scarus ovifrons) is a species of marine ray-finned fish, a parrotfish, in the family Scaridae which is famous for its characteristic blue color. It lives in reefs and coral reefs. It grows to a maximum length of about 90...



Knysna seahorse

Hippocampus capensis

The Knysna /'naɪznə/ seahorse or Cape seahorse (*Hippocampus capensis*) is a species of fish in the family Syngnathidae. It is endemic to the south coast of South Africa, where it has been found in only three brackish water habitats: the estuary of...



Kōhaku

Cyprinus rubrofasciatus var. "Kōhaku"

Kōhaku (紅白 (kōhaku, "red and white")) is a variety of ornamental koi (carp). The Kōhaku has a white body, with red markings across the body. It is considered one of the 'Big Three' varieties of Koi, along with the Sanke, and Showa. The Kōhaku...



Koheru

Decapterus koheru

The koheru (Decapterus koheru) is a ray-finned fish of the genus Decapterus, part of the family Carangidae. It is endemic to New Zealand where it occurs from Manawatāwhi / Three Kings Islands to southern North Island. This is a schooling, pelagic...



Koi

Cyprinus rubrofusus var. "koi"

Koi (鯉, English: /'kɔɪ/, Japanese:) or more specifically nishikigoi (錦鯉, Japanese:, literally "brocaded carp") are colored varieties of the common carp (Cyprinus carpio) that are kept for decorative purposes in outdoor koi ponds or water...



Kokanee salmon

Oncorhynchus nerka

The kokanee salmon (Oncorhynchus nerka), also known as the kokanee trout, little redfish, silver trout, kikanning, Kennerly's salmon, Kennerly's trout, or Walla, is the non-anadromous form of the sockeye salmon (meaning that they do not migrate...



Korean mackerel

Scomberomorus koreanus

The Korean mackerel (Scomberomorus koreanus) also known as the Korean seerfish, is a ray-finned bony fish in the family Scombridae, better known as the mackerel family. Within that family, this fish is a member of the tribe Scomberomorini, the...



Korean sandlance

Hypoptychus dybowskii

The Korean sandlance (Hypoptychus dybowskii) is a species of marine ray-finned fish belonging to the family Hypoptychidae. The Korean sandlance is the only species in this monotypic family and genus and is found in the northwestern Pacific Ocean.



Korean skate

Hongoeo koreana

The Korean skate (Hongoeo koreana) is a skate in the family Rajidae, and is the sole member of the genus Hongoeo. It was previously classified in the genus Raja as Raja koreana until discoveries identified this species as a member of a distinct...



Krabi mouth-brooding betta

Betta simplex

The Krabi mouth-brooding betta, Betta simplex, is a species of gourami endemic to Krabi Province, Thailand. It inhabits karst springs and wetlands, where it lives amongst vegetation close to the banks. This species can reach a length of 8 cm (3...



Kryptopterus bicirrhis

Kryptopterus bicirrhis

Kryptopterus bicirrhis, often called the glass catfish, is an Asian glass catfish species of the genus Kryptopterus. Until 1989, the concept of K. bicirrhis included its smaller relative Kryptopterus vitreolus ("ghost catfish"; often confused...



Kryptopterus kryptopterus

Kryptopterus kryptopterus

Kryptopterus kryptopterus (blue sheatfish) is a species of catfish, the type species of the genus Kryptopterus. It can be distinguished from all its congeners, with the exception of the newly split Kryptopterus geminus, by the almost flat dorsal...



Kryptopterus minor

Kryptopterus minor

Kryptopterus minor is a small species of Asian glass catfish from the Kapuas River basin in Borneo, Indonesia. Until 2013, the ghost catfish (K. vitreolus) was included in K. minor. The true K. minor is rarely (if ever) seen in the aquarium...



Kryptopterus vitreolus

Kryptopterus vitreolus

Kryptopterus vitreolus, known in the aquarium trade traditionally as the glass catfish and also as the ghost catfish or phantom catfish, is a small species of Asian glass catfish. It is commonly seen in the freshwater aquarium trade, but its...



Kuhl's maskray

Neotrygon kuhlii

Kuhl's maskray (Neotrygon kuhlii), also known as the blue-spotted stingray, blue-spotted maskray, or Kuhl's stingray, is a species of stingray of the family Dasyatidae. It was recently changed from Dasyatis kuhlii in 2008 after morphological and...



Kuhli loach

Pangio kuhlii

The true Kuhli loach (Pangio kuhlii), occasionally referred to as eel loach, is a small eel-like freshwater fish belonging to the loach family (Cobitidae). They originate from the island of Java in Indonesia. This snake-like creature is very...



Kuhlia marginata

Kuhlia marginata

Kuhlia marginata, the dark-margined flagtail, spotted flagtail, silver flagtail, orange-finned flagtail, northern jungle perch or mountain bass, is a species of diadromous ray-finned fish, a flagtail from the family Kuhliidae. It is found in...



Kuhlia mugil

Kuhlia mugil

Kuhlia mugil, the barred flagtail, the fiveband flagtail or the five-bar flagtail, is a species of marine ray-finned fish, a flagtail belonging to the family Kuhliidae. It is found in the Indo-Pacific region.



Kuhlia rupestris

Kuhlia rupestris

Kuhlia rupestris, the rock flagtail, jungle perch, mountain trout, buffalo bream, dusky-finned bulleye, rockmountain bass or spotted flagtail, is a species of ray-finned fish, a flagtail, from the family Kuhliidae. It is a catadromous species...



Kuhlia sandvicensis

Kuhlia sandvicensis

Kuhlia sandvicensis, the reticulated flagtail, zebra-headed flagtail or Hawaiian flagtail, is a species of ray-finned fish, a flagtail from the family Kuhliidae which is found in the central Pacific Ocean. It is popular as a game fish and can...



Kuhlia xenura

Kuhlia xenura

Kuhlia xenura, the strange-tailed flagtail or Hawaii flagtail, is a species of ray-finned fish, a flagtail from the family Kuhliidae which is endemic to Hawaii, where it occurs in fresh, brackish, and marine waters. It can be found in tide pools,...



Kuria labeo

Labeo gonius

The Kuria labeo (Labeo gonius) (Assamese: কুঁটি, romanized: kurhi, কুৰিহা kuriha; Bengali: ঘনিয়া, romanized: ghonia, গৈন্না gôinna) is a species of fish in the carp family, Cyprinidae. It is native to Pakistan, India, Bangladesh, and Burma, and...



Kurtus gulliveri

Kurtus gulliveri

Kurtus gulliveri, the nurseryfish, is a species of fish in the family Kurtidae native to fresh and brackish waters in southern New Guinea and northern Australia. They can be found in estuaries, mangrove swamps, nipa swamps and slow-flowing rivers...



Kyphosus sectatrix

Kyphosus sectatrix

Kyphosus sectatrix, the Bermuda chub, Pacific drummer, beaked chub, grey drummer, Pacific chub or white chub, is a species of marine ray-finned fish, a sea chub from the family Kyphosidae. This species is found in tropical and subtropical coastal...



Labeo bata

Labeo bata

Labeo bata (Bengali: বাটা, romanized: bātā) is a fish in genus Labeo. It is commonly known as bata, and is a native fish to India and Bangladesh.



Labeo boga

Labeo boga

Labeo boga is fish in genus Labeo. It is in the family Cyprinidae and the Actinopterygii class which is widespread in India, Nepal, Bangladesh, Pakistan and Myanmar.



Labeo coubie

Labeo coubie

Labeo coubie, the African carp, is a cyprinid fish, widespread in Africa, where it occurs within the drainage basin of the Nile (Blue, White, Lake Albert) and in the Chad, Niger-Benue, Volta, Senegal and Gambia Rivers, as well as the Cross River...



Labeo dyocheilus

Labeo dyocheilus

Labeo dyocheilus is fish in genus Labeo from southern Asia.



Labeobarbus bynni

Labeobarbus bynni

Labeobarbus bynni, the Niger barb, is an African species of cyprinid freshwater fish. It has often been placed in the genus Barbus, but is now usually placed in Labeobarbus. This is a relatively large barb, up to 82 cm (32 in) in total length. It...



Labidochromis caeruleus

Labidochromis caeruleus

Labidochromis caeruleus is a species of cichlid endemic to the central western coastal region of Lake Malawi in East Africa. It is also known as lemon yellow lab, the blue streak hap, the electric yellow or yellow prince, depending on the colour...



Labrisomus nuchipinnis

Labrisomus nuchipinnis

Labrisomus nuchipinnis, the hairy blenny, is a species of labrisomid blenny native to the Atlantic Ocean from the coast of the Americas to the African coast. This species prefers areas that give them crevices and holes to shelter in such as areas...



Labroides bicolor

Labroides bicolor

Labroides bicolor is a species of wrasse endemic to the Indian Ocean and the Pacific Ocean. It is known by various names including bicolor cleanerfish, bicolor(ed) cleaner wrasse, cleaner wrasse, two-color cleaner wrasse and yellow diesel wrasse.



Labrus viridis

Labrus viridis

Labrus viridis (the green wrasse) is a species of wrasse native to the eastern Atlantic Ocean from Portugal to Morocco, as well as through the Mediterranean Sea to the Black Sea. This species is found around rocky reefs amongst the rocks or in...



Laced moray

Gymnothorax favagineus

The laced moray (Gymnothorax favagineus), also known as the leopard moray, leopard moray eel, tessellate moray or honeycomb moray, is a species of marine fish in the family Muraenidae.



Lactophrys trigonus

Lactophrys trigonus

Lactophrys trigonus, commonly called the buffalo trunkfish or trunkfish, is a species of boxfish native to the Western Atlantic.



Laetacara curviceps

Laetacara curviceps

Laetacara curviceps, the Flag acara, Dwarf flag cichlid, is a species of cichlid that lives in slow-moving rivers and streams as well as ponds and lakes of the Amazon Basin. This species can reach a total length of 10 cm (4 in).



Lagocephalus scleratus

Lagocephalus scleratus

Lagocephalus scleratus (Gmelin, 1789), commonly known as the silver-cheeked toadfish, or Sennin-fugu (Japanese: 仙人河豚), is an extremely poisonous marine bony fish in the family Tetraodontidae (puffer fishes).



Lagocephalus spadiceus

Lagocephalus spadiceus

Lagocephalus spadiceus, also known as the half-smooth golden pufferfish, is a species of fish in the family Tetraodontidae. It is a common fish in the Red Sea, as well as the Indian Ocean, but can be found also in the Mediterranean, where it...



Lagodon

Lagodon rhomboides

Lagodon is a monospecific genus of marine ray-finned fish belonging to the family Sparidae, which includes the seabreams and porgies. The only species in the genus is Lagodon rhomboides, the pinfish, red porgy, bream, pin perch, sand perch,...



Lagoon triggerfish

Rhinecanthus aculeatus

The lagoon triggerfish (*Rhinecanthus aculeatus*), also known as the blackbar triggerfish, the Picasso triggerfish, or the Picassofish, is a triggerfish, up to 30 cm in length, found on reefs in the Indo-Pacific region. This species has been studied...



Lahontan cutthroat trout

Oncorhynchus clarkii henshawi

Lahontan cutthroat trout (Oncorhynchus clarkii henshawi) is the largest subspecies of cutthroat trout, and the state fish of Nevada. It is one of three subspecies of cutthroat trout that are listed as federally threatened.



Lake Chala tilapia

Oreochromis hunteri

The Lake Chala tilapia (Oreochromis hunteri) is a species of cichlid fish that is endemic to Lake Chala, a small crater lake on the border of Kenya and Rombo District of Kilimanjaro Region in Tanzania. It mostly lives in relatively deep water, at...



Lake chub

Couesius plumbeus

The lake chub (Couesius plumbeus) is a freshwater cyprinid fish found in Canada and in parts of the United States. Of all North American minnows, it is the one with the northernmost distribution. Its genus, Couesius is considered monotypic today....



Lake chubsucker

Erimyzon sucetta

The lake chubsucker (Erimyzon sucetta) is a species of freshwater fish endemic to North America, found in the Great Lakes and the Mississippi River basin, as far north as Ontario, Canada, extending south to the Gulf of Mexico. It is mainly found...



Lake Eacham rainbowfish

Melanotaenia eachamensis

The Lake Eacham rainbowfish (Melanotaenia eachamensis) is a species of rainbowfish in the subfamily Melanotaeniidae which was thought to be endemic to Yidyam (Lake Eacham), Queensland, Australia but has proven to have a wider range.



Lake Kutubu rainbowfish

Melanotaenia lacustris

The Lake Kutubu rainbowfish (*Melanotaenia lacustris*) is a species of fish in the family Melanotaeniidae. It is also known as turquoise rainbowfish. It is one of 13 fish endemic to Lake Kutubu, a lake found within the Kikori River system in Papua...



Lake Malawi sardine

Engraulicypris sardella

The Lake Malawi sardine, lake sardine, or usipa (Engraulicypris sardella), is an African species of freshwater fish in the family Cyprinidae. It is endemic to Lake Malawi and its outlet, the (upper) Shire River; it is found in Malawi, Mozambique,...



Lake minnow

Rhynchocypris percunurus

The lake minnow or swamp minnow (Rhynchocypris percunurus) is a Eurasian species of small freshwater cyprinid fish. It has a wide but disjunct distribution including parts of Europe (Belarus, Czech Republic, Germany, Lithuania, Poland, Russia, and...



Lake salmon

Opsaridium microlepis

The lake salmon or mpasa (Opsaridium microlepis) is an African species of freshwater fish, endemic to Lake Malawi, in the family Cyprinidae found in Malawi, Mozambique, and Tanzania. Its natural habitats are rivers and freshwater lakes.



Lake sturgeon

Acipenser fulvescens

The lake sturgeon (Acipenser fulvescens), also known as the rock sturgeon, is a North American temperate freshwater fish, one of about 25 species of sturgeon. Like other sturgeons, this species is a bottom feeder and has a partly cartilaginous...



Lake Tanganyika sardine

Limnothrissa miodon

The Lake Tanganyika sardine (Limnothrissa miodon) is a species of freshwater fish in the family Dorosomatidae which was endemic to Lake Tanganyika but which has now been introduced to other lakes in Africa as a food source. It is monotypic within...



Lake trout

Salvelinus namaycush

The lake trout (Salvelinus namaycush) is a freshwater char living mainly in lakes in northern North America. Other names for it include mackinaw, namaycush, lake char (or charr), touladi, togue, and grey trout. In Lake Superior, it can also be...



Lake Turkana barb

Enteromius turkanae

The Lake Turkana barb (Enteromius turkanae) is a species of ray-finned fish in the family Cyprinidae. It is endemic to Lake Turkana in Kenya. It is not considered a threatened species by the IUCN.



Lampris guttatus

Lampris guttatus

Lampris guttatus, commonly known as the opah, cravo, moonfish, kingfish, and Jerusalem haddock, is a large, colorful, deep-bodied pelagic lampriform fish belonging to the family Lampridae, which comprises the genus Lampris. It is a pelagic fish...



Lampris immaculatus

Lampris immaculatus

Lampris immaculatus, commonly known as the southern opah or southern moonfish, is a species of fish native to the Southern Ocean. The species is found commonly in New Zealand waters. They are caught commercially using long-line fishery in New...



Lamprologus callipterus

Lamprologus callipterus

Lamprologus callipterus is a species of cichlid endemic to Lake Tanganyika where it very actively moves about in search of crustaceans and other invertebrates. Males of this species can reach a length of 12.4 centimetres (4.9 in) TL while the...



Lamprologus lethops

Lamprologus lethops

Lamprologus lethops is a species of cichlid fish from areas with fast current in the Congo River in Central Africa, where it is believed to live in depths as great as 160 m (520 ft) or 200 m (660 ft) below the surface. It reaches up to about 12.5...



Lamprologus ocellatus

Lamprologus ocellatus

Lamprologus ocellatus is a species of shell dwelling cichlid endemic to Lake Tanganyika. It is a popular aquarium fish due to its small size, appearance, and intelligence. Juveniles measure about 6 millimetres (0.24 in) at hatching, while adults...



Lane snapper

Lutjanus synagris

The lane snapper (Lutjanus synagris), the Mexican snapper, redbtail snapper or spot snapper, is a species of marine ray-finned fish, a snapper belonging to the family Lutjanidae. It is native to the western Atlantic Ocean.



Large kelpfish

Chironemus marmoratus

The large kelpfish (Chironemus marmoratus), the Eastern kelpfish, hiwihwi, surgefish or kelpfish, is a species of marine ray-finned fish, a kelpfish belonging to the family Chironemidae. It is found in southern Australia, and off the North...



Large-eyed rabbitfish

Hydrolagus mirabilis

The large-eyed rabbitfish (*Hydrolagus mirabilis*) is a species of fish in the family Chimaeridae. It is found in several areas of the Atlantic Ocean and within the Mediterranean Sea.



Largehead hairtail

Trichiurus lepturus

The largehead hairtail (*Trichiurus lepturus*) or beltfish is a member of the cutlassfish family, Trichiuridae. This common to abundant species is found in tropical and temperate oceans throughout the world. The taxonomy is not fully resolved, and...



Largemouth bass

Micropterus salmoides

The largemouth bass (*Micropterus salmoides*) is a carnivorous freshwater ray-finned fish in the Centrarchidae (sunfish) family, native to the eastern and central United States, southeastern Canada and northern Mexico. It is known by a variety of...



Largemouth yellowfish

Labeobarbus kimberleyensis

The largemouth yellowfish or Vaal-Orange largemouth yellowfish (*Labeobarbus kimberleyensis*) is a ray-finned fish species in the family Cyprinidae. This large freshwater barb is found in southern Africa. It has long been placed in *Barbus*, the...



Largescale silver carp

Hypophthalmichthys harmandi

The largescale silver carp (Hypophthalmichthys harmandi) is a freshwater fish in the carp family Cyprinidae. It is native to Hainan and Vietnam.



Largescale sucker

Catostomus macrocheilus

The largescale sucker (Catostomus macrocheilus) is a species of sucker, a type of freshwater fish, found in western North America.



Largetooth cookiecutter shark

Isistius plutodus

The largetooth cookiecutter shark (Isistius plutodus) is a rare species of squaliform shark in the family Dalatiidae, reported from depths of 60–200 m (200–660 ft) at scattered locations in the Atlantic and Pacific Oceans. As its common name...



Largetooth sawfish

Pristis pristis

The largetooth sawfish (Pristis pristis, syn. P. microdon and P. perotteti) is a species of sawfish in the family Pristidae. It is found worldwide in tropical and subtropical coastal regions, but also enters freshwater. It has declined...



Larimichthys crocea

Larimichthys crocea

Larimichthys crocea, commonly called the large yellow croaker, yellow croaker or croceine croaker, is a species of marine ray-finned fish belonging to the family Sciaenidae, the drums and croakers. This species is found in the Western Pacific...



Larimichthys polyactis

Larimichthys polyactis

Larimichthys polyactis, called the redlip croaker, small yellow croaker, little yellow croaker or yellow corvina, is a species of croaker native to the western Pacific, generally in temperate waters such as the East China Sea and the Yellow Sea.



Lasiognathus amphirhamphus

Lasiognathus amphirhamphus

Lasiognathus amphirhamphus is a species of wolftap angler found in the Madeira Abyssal Plain in the east-central Atlantic Ocean where it occurs at a depth of from 1,200 to 1,305 metres (3,937 to 4,281 ft). The females of this species grow to a...



Lasiognathus dinema

Lasiognathus dinema

Lasiognathus dinema is a species of wolftap angler found in the deep waters of the northern Gulf of Mexico. It is found at depths of around 3,280 to 4,900 feet (1,000 to 1,500 meter). This species is part of the Angler Fish species. This species...



Latris lineata

Latris lineata

Latris lineata, the striped trumpeter, common trumpeter, copper moki, Hobart-town trumpeter, kokikohi, real trumpeter, Tasmanian striped trumpeter or Tasmanian trumpeter, is a species of marine ray-finned fish belonging to the family Latridae,...



Latticed butterflyfish

Chaetodon rafflesii

The latticed butterflyfish (Chaetodon rafflesii), also known as Raffles' coralfish, is a species of marine ray-finned fish, a butterflyfish belonging to the family Chaetodontidae. It is found in the Indo-Pacific region.



Leafscale gulper shark

Centrophorus squamosus

The leafscale gulper shark (Centrophorus squamosus) is a dogfish of the family Centrophoridae. C. squamosus is reported to have a lifespan of approximately 70 years, based on otolith ring counts. It was the first described species in the genus...



Leafy seadragon

Phycodurus eques

The leafy seadragon (Phycodurus eques) or Glauert's seadragon, is a marine fish. It is the only member of the genus Phycodurus in the family Syngnathidae, which includes seadragons, pipefish, and seahorses. It is found along the southern and...



Leaping blenny

Alticus saliens

The leaping blenny (*Alticus saliens*), also known as the jumping blenny, is a species of combtooth blenny (family Blenniidae) in the genus *Alticus*. There is some uncertainty as to whether it was first described by J.R. Forster in 1788 or B.G.E....



Leaping mullet

Chelon saliens

The leaping mullet (Chelon saliens) is a species of fish in the family Mugilidae. It is found in coastal waters and estuaries in the northeast Atlantic, ranging from Morocco to France, and including the Mediterranean and Black Sea. It has been...



Least chub

Iotichthys phlegethontis

The least chub (Iotichthys phlegethontis) is a species of ray-finned fish in the family Cyprinidae, the only member of the genus Iotichthys. This species is found only in Utah in the United States. Its native habitat formerly included much of the...



Least puffer

Sphoeroides parvus

Least puffer, Sphoeroides parvus, is a species in the family Tetraodontidae, or pufferfishes. This species is the common bay and inshore puffer for the waters around Texas and Louisiana. It has also been found as far east as Apalachicola Bay and...



Leatherjacket fish

Oligoplites saurus

The leatherjacket fish (Oligoplites saurus), also known as leather jack, is a species of jack in the family Carangidae. Leather jack may also refer to other members of the Carangidae, such as the pilot fish. The largest are about a foot long.



Lebranche mullet

Mugil liza

The Lebranche mullet or the liza (Mugil liza) is a species of saltwater fish in the family Mugilidae. It is found in the western Atlantic Ocean and Caribbean Sea and is fished commercially.



Leerfish

Lichia amia

The leerfish or garrick (*Lichia amia*) is a species of marine fish in the family Carangidae, and is native to the Mediterranean and the coastal waters of western Africa to the coastal waters of eastern South Africa. Also recorded in the Black Sea....



Leiognathus equula

Leiognathus equulus

Leiognathus equula, the common ponyfish is a species of marine ray-finned fish, a ponyfish from the family Leiognathidae. It occurs in brackish and marine waters from East Africa to Fiji in the Indian and western Pacific Oceans, Red Sea and...



Leiopotherapon plumbeus

Leiopotherapon plumbeus

Leiopotherapon plumbeus, known commonly as the silver perch, is a species of fish in the family Terapontidae, the grunters. It is endemic to the Philippines, where it is called ayungin, bugaong, bigaong, and багаong.



Leiopotherapon unicolor

Leiopotherapon unicolor

Leiopotherapon unicolor, the spangled grunter or spangled perch is a species of ray-finned fish, a grunter from the family Terapontidae. It is endemic to Australia.



Lemon shark

Negaprion brevirostris

The lemon shark (Negaprion brevirostris) is a species of shark from the family Carcharhinidae and is classified as a Vulnerable species by the International Union for the Conservation of Nature. Lemon sharks can grow to 3.4 metres (11 ft) in...



Lemon sole

Microstomus kitt

The lemon sole (Microstomus kitt) is a flatfish of the family Pleuronectidae. It is native to shallow seas around Northern Europe, where it lives on stony bottoms down to depths of about 1,400 metres (4,600 ft). It grows up to 65 centimetres (26...



Lemon tetra

Hyphessobrycon pulchripinnis

The lemon tetra (Hyphessobrycon pulchripinnis) is a species of tropical freshwater fish which originates from South America, belonging to the family Characidae. It is a small tetra growing to 5 cm (2 in) in length. The species is a long...



Lemonpeel angelfish

Centropyge flavissima

The lemonpeel angelfish (Centropyge flavissima), also known as the yellow angelfish, is a species of marine ray-finned fish, a marine angelfish belonging to the family Pomacanthidae. It is found in the Indo-Pacific region.



Leopard bush fish

Ctenopoma acutirostre

The leopard bush fish (Ctenopoma acutirostre), also known as leopard bushfish, spotted ctenopoma, leopard ctenopoma, spotted climbing perch, spotted leaf fish, spotted cichlid or spotted bushfish, is a freshwater fish. It is a member of the...



Leopard catshark

Poroderma pantherinum

The leopard catshark (Poroderma pantherinum) is a species of catshark, and part of the family Scyliorhinidae, endemic to the coastal waters of South Africa. Abundant in inshore waters under 20 m (66 ft) deep, this bottom-dweller favors rocky...



Leopard coral grouper

Plectropomus leopardus

The leopard coral grouper (Plectropomus leopardus), also known as the common coral trout, leopard coral trout, blue-dotted coral grouper or spotted coral grouper, is a species of marine ray-finned fish, a grouper from the subfamily Epinephelinae...



Leopard eel

Myrichthys pardalis

The leopard eel (Myrichthys pardalis) is an eel in the worm or snake eels family, Ophichthidae. It was described by Achille Valenciennes in 1839, originally under the genus Ophisurus. Male leopard eels can reach a maximum total length of 64.8 cm...



Leopard flounder

Bothus pantherinus

The leopard flounder (Bothus pantherinus) or panther flounder, is a flatfish found in the Pacific and Indian Oceans.



Leopard moray eel

Enchelycore pardalis

The leopard moray eel, tiger moray eel or dragon moray (Enchelycore pardalis), is a species of marine fish in the family Muraenidae. The leopard moray eel is widespread throughout the Indo-Pacific oceans from Réunion to the Hawaiian, Line and...



Leopard round stingray

Urobatis pardalis

The leopard round stingray (Urobatis pardalis) or Central American round stingray and Costa Rican round stingray is a small species of round ray found in shallow waters off the coast from Costa Rica to Colombia in the eastern Pacific Ocean.



Leopard shark

Triakis semifasciata

The leopard shark (*Triakis semifasciata*) is a species of houndshark, in the family Triakidae. It is found along the Pacific coast of North America, from the U.S. state of Oregon to Mazatlán in Mexico. Typically measuring 1.2–1.5 m (3.9–4.9 ft)...



Leopard torpedo

Torpedo panthera

Torpedo panthera also known as the leopard torpedo is a species of fish in the family Torpedinidae. It is found in Djibouti, Egypt, Eritrea, India, Iran, Oman, Pakistan, Saudi Arabia, Somalia, Sudan, and Yemen. Its natural habitat is open seas.



Leopard whipray

Himantura leoparda

The leopard whipray (*Himantura leoparda*) is a little-known species of stingray in the family Dasyatidae, found in the Indian and Pacific Oceans from the Andaman Sea to the Coral Triangle. It is found close to shore at depths shallower than 70 m...



Lepidocephalichthys guntea

Lepidocephalichthys guntea

Lepidocephalichthys guntea, also known as guntea loach, scavenger loach or peppered loach, is a species of cobitid loach native to southern and south-eastern Asia. This species reaches a length of 15 centimetres (5.9 in) TL. It uses its...



Lepidogalaxias

Lepidogalaxias salamandroides

Lepidogalaxias salamandroides is a species of small freshwater fish of Western Australia. It is the only member of the family Lepidogalaxiidae and genus Lepidogalaxias. Common names for this fish include salamanderfish and Shannon mudminnow....



Lepidorhombus boscii

Lepidorhombus boscii

The four-spot megrim (Lepidorhombus boscii) is a species of flatfish in the family Scophthalmidae. It is found a depths between 7 and 800 m (23 and 2,625 ft) in the northeast Atlantic and Mediterranean. It can be separated from its close...



Leporinus fasciatus

Leporinus fasciatus

Leporinus fasciatus, commonly known as the banded leporinus or the black-banded leporinus, is a species of characin in the family Anostomidae. L. fasciatus is native to the Amazon Basin in South America, but has been introduced into the US states...



Leptobarbus hoevenii

Leptobarbus hoevenii

Hoven's carp (Leptobarbus hoevenii; Jelawat in Malay), also known as the mad barb or sultan fish, is a species of fish in the barb family. Named in honor of Bleeker's fellow Dutchman, "le célèbre professeur de zoologie" Jan van der Hoeven...



Leptobotia elongata

Leptobotia elongata

Leptobotia elongata, the imperial flower loach, elongate loach or royal clown loach, is a species of botiid fish found in flowing water in the upper and middle Yangtze basin in China. It is the largest species in the family, reaching up to 50 cm...



Lepturacanthus savala

Lepturacanthus savala

The Savalai hairtail, *Lepturacanthus savala*, also known as small-head hairtail, is a species of cutlassfish native to the Indian Ocean and into the west Pacific Ocean as far east as Australia. They inhabit deep waters at depths from 100 m (330...



Lesser devil ray

Mobula hypostoma

The lesser devil ray (*Mobula hypostoma*) is a species of devil ray in the family Mobulidae.



Lesser electric ray

Narcine bancroftii

The lesser electric ray (*Narcine bancroftii*), also known as the Brazilian electric ray, small electric ray, spotted torpedo ray, torpedofish or trembler, is a species of numbfish in the family Narcinidae found on the western coastal fringes of...



Lesser sand eel

Ammodytes tobianus

The lesser sand eel or sand lance (*Ammodytes tobianus*) is a species of fish in the sand lance family Ammodytidae. It is an elongated cylindrical fish which may be up to 20 centimetres (7.9 in) long.



Lesser spiny eel

Macrognathus aculeatus

The lesser spiny eel, *Macrognathus aculeatus*, is a Southeast Asian tropical freshwater fish belonging to the family Mastacembelidae. They are uncommonly found in aquaria.



Lesser weever

Echiichthys vipera

The lesser weever (Echiichthys vipera) is a venomous weever of the family Trachinidae, in the order Perciformes, and the class Actinopterygii. It is generally found on the sandy sea beds of the open sea, near the shore. Lesser weevers may sting...



Lethrinus atkinsoni

Lethrinus atkinsoni

Lethrinus atkinsoni, the Pacific yellowtail emperor,, Atkinson's emperor, reticulated emperor, tricky snapper, Tuamotu emperor, yellow morwong or yellow-tailed emperor is a species of marine ray-finned fish benomging to the family Lethrinidae,...



Lethrinus harak

Lethrinus harak

Lethrinus harak, the thumbprint emperor blackblotch emperor or blackspot emperor, is a species of marine ray-finned fish belonging to the Family Lethrinidae. the emperors and emperor breams. This species is found in the Indo-Pacific region.



Lethrinus lentjan

Lethrinus lentjan

Lethrinus lentjan, the pink ear emperor, redspot emperor, purple ear emperor or purple-headed emperor, is a species of marine ray-finned fish belonging to the family Lethrinidae, the emperors and emperor breams. This fish is found in the...



Lethrinus mahsena

Lethrinus mahsena

Lethrinus mahsena, common names the sky emperor, mahsena emperor, and cutthroat emperor, is a species of emperor fish. It grows to 65 cm (26 in) in length, but is commonly found at between 35 and 45 cm (14 and 18 in). This fish may be yellow to...



Lethrinus miniatus

Lethrinus miniatus

Lethrinus miniatus, the trumpet emperor, redthroat emperor, sweetlip emperor, Sweetlip Swoose, island snapper, yellowmouth perch, yellowmouth snapper, lipper or nannygal, is a species of marine ray-finned fish belonging to the family Lethrinidae,...



Lethrinus nebulosus

Lethrinus nebulosus

Lethrinus nebulosus, the spangled emperor, green snapper, morwong, north-west snapper, sand bream, sand snapper, sixteen-pounder, sharie, sheri and yellow sweetlip, is a species of marine ray-finned fish belonging to the family Lethrinidae, the...



Lethrinus olivaceus

Lethrinus olivaceus

Lethrinus olivaceus, common name longface emperor or long-nosed emperor, is a species of marine ray-finned fish belonging to the family Lethrinidae, the emperors or emperor breams. This species has a wide Indo-Pacific distribution.



Leucaspius delineatus

Leucaspius delineatus

Leucaspius delineatus, known as the sunbleak, belica or moderlieschen is a species of freshwater fish in the family Cyprinidae. It is currently the only species included in genus Leucaspius, whereas formerly others were included, which now have...



Leuresthes tenuis

Leuresthes tenuis

Leuresthes tenuis, the California grunion, is a species of ray-finned fish native to the Pacific coast of North America from Monterey Bay in California to Baja California. This species grows to 19 cm (7 in) in total length and is of minor...



Liberty molly

Poecilia salvatoris

The liberty molly (Poecilia salvatoris) is a fish species from El Salvador. It is one of the short fin molly species in the Poecilia sphenops complex, and had been often regarded as synonymous with P. sphenops until recently. Males and females are...



Limia melanonotata

Limia melanonotata

Limia melanonotata, the blackbanded limia, is a toothcarp in the family Poeciliidae. It is endemic to the island of Hispaniola (in both Haiti and the Dominican Republic).



Lined butterflyfish

Chaetodon lineolatus

The lined butterflyfish (Chaetodon lineolatus) is a species of marine ray-finned fish, a butterflyfish belonging to the family Chaetodontidae, one of the largest species in the genus Chaetodon. It has a wide range from the Red Sea to South Africa...



Lined seahorse

Hippocampus erectus

The lined seahorse (Hippocampus erectus), northern seahorse or spotted seahorse, is a species of fish that belongs to the family Syngnathidae. H. erectus is a diurnal species with an approximate length of 15 cm (5.9 inches) and lifespan of one to...



Lingcod

Ophiodon elongatus

The lingcod or ling cod (Ophiodon elongatus), also known as the buffalo cod or cultus cod, is a fish of the greenling family Hexagrammidae. It is the only extant member of the genus Ophiodon. A slightly larger, extinct species, Ophiodon...



Linophryne arborifera

Linophryne arborifera

Linophryne arborifera, or illuminated netdevil, is an anglerfish of the family Linophrynidae, found in all tropical and subtropical oceans at depths below 1,000 m (3,300 ft) in the Bathyal zone. Its length is up to 77 mm (3 inches). The female is...



Lionhead cichlid

Steatocranus casuarius

The lionhead cichlid (Steatocranus casuarius), also known as African blockhead, buffalohead, humphead cichlid, lionhead or lumphead is a species of rheophilic cichlid native to Pool Malebo and the Congo River. It uses caves for spawning. This...



Liopropoma carmabi

Liopropoma carmabi

Liopropoma carmabi, the candy basslet, is a species of fish in the family Serranidae.



Liparis fabricii

Liparis fabricii

Liparis fabricii, commonly known as the gelatinous seasnail or gelatinous snailfish, is a benthopelagic species of snailfish from the Arctic Ocean. It has a tadpole-like body with a maximum length of about 20 cm (7.9 in). It is brown to black in...



Liparis liparis

Liparis liparis

Liparis liparis, the common seasnail, striped seasnail or seasnail, is a small species of marine ray-finned fish belonging to the family Liparidae, the snailfishes, in the order Scorpaeniformes, the scorpionfishes and flatheads. It is found in...



Liparis marmoratus

Liparis marmoratus

Liparis marmoratus, or the festive snailfish, is a marine ray-finned fish from the genus Liparis. It was first described by Schmidt in 1950.



Lipophrys pholis

Lipophrys pholis

Lipophrys pholis, commonly known as shanny, also known as the smooth blenny or common blenny, is a species of combtooth blenny. It matures at two years of age. Distributed in the Eastern Atlantic from the southern Norway to Morocco and Madeira,...



Lithodoras

Lithodoras dorsalis

Lithodoras dorsalis, the rock-bacu, is the only species in the genus Lithodoras of the catfish (order Siluriformes) family Doradidae. This species originates from the Amazon basin in Brazil and estuaries near Cayenne, French Guiana. These fish...



Little dragonfish

Eurypegasus draconis

The little dragonfish or short dragonfish (Eurypegasus draconis) is a species of marine fish in the family Pegasidae. It is widespread throughout the tropical waters of the Indo-Pacific, including the Red Sea. The little dragonfish can grow up to...



Little Kern golden trout

Oncorhynchus mykiss whitei

The Little Kern golden trout (Oncorhynchus mykiss whitei) is a brightly colored subspecies of rainbow trout native to the main stem and tributaries of the Little Kern River in Tulare County, California. Together with the California golden trout...



Little skate

Leucoraja erinacea

The little skate (Leucoraja erinacea) is a species of skate in the family Rajidae, found from Nova Scotia to North Carolina on sand or gravel habitats. They are one of the dominant members of the demersal fish community in the northwestern...



Little tunny

Euthynnus alletteratus

The little tunny (*Euthynnus alletteratus*), also known as the false albacore, little tuna, bonita, or erroneously as the blue bonito, is a species of tuna in the family Scombridae. It can be found in the Atlantic Ocean and the Mediterranean and...



Live sharksucker

Echeneis naucrates

The live sharksucker or slender sharksucker (*Echeneis naucrates*) is a species of marine fish in the family Echeneidae, the remoras.



Loach minnow

Rhinichthys cobitis

The loach minnow (Rhinichthys cobitis) is a species of freshwater fish. It is a member of the carp family (family Cyprinidae) of order Cypriniformes. It occurs in streams and small rivers throughout the Gila River and San Pedro River systems in...



Lobed river mullet

Cestraeus goldiei

The lobed river mullet (Cestraeus goldiei), also known as ludong or banak, is a freshwater mullet.

While it is claimed to be endemic to Cagayan River and tributaries extending through the watersheds of Cagayan Valley and the Santa-Abra River...



Lollipop catshark

Cephalurus cephalus

The lollipop catshark (Cephalurus cephalus) is a little-known species of deep sea catshark, belonging to the family Scyliorhinidae, and the only described member of its genus. A diminutive, bottom-dwelling shark of the outer continental shelf and...



Long-snouted seahorse

Hippocampus guttulatus

Hippocampus guttulatus, commonly known as the long-snouted seahorse and in Great Britain as the spiny seahorse, is a marine fish belonging to the family Syngnathidae, native from the northeast Atlantic, including the Mediterranean.



Longbill spearfish

Tetrapturus pfluegeri

The longbill spearfish (Tetrapturus pfluegeri) is a species of marlin native to the Atlantic Ocean where it is found above the thermocline in open waters between 40°N and 35°S. This species can reach a length of 254 centimetres (100 in) FL and...



Longcomb sawfish

Pristis zijsron

The longcomb sawfish, narrowsnout sawfish or green sawfish (Pristis zijsron) is a species of sawfish in the family Pristidae, found in tropical and subtropical waters of the Indo-West Pacific. It has declined drastically and is now considered a...



Longear sunfish

Lepomis megalotis

The longear sunfish (Lepomis megalotis) is a freshwater fish in the sunfish family, Centrarchidae, of order Perciformes. It is native to the area of eastern North America stretching from the Great Lakes down to northeastern Mexico. The longear...



Longfin crevalle jack

Caranx fischeri

The longfin crevalle jack (Caranx fischeri) is a recently described species of large marine fish classified in the jack family, Carangidae. It inhabits the subtropical waters of the east Atlantic Ocean, ranging along the African coast from...



Longfin grouper

Epinephelus quoyanus

The longfin grouper (Epinephelus quoyanus), also known as the longfin rockcod, bar-breasted rock-cod, Gilbert's rock-cod, honeycomb rockcod, spotted groper or wirenet cod, is a species of marine ray-finned fish, a grouper from the subfamily...



Longfin mako shark

Isurus paucus

The longfin mako shark (Isurus paucus) is a species of mackerel shark in the family Lamnidae, with a probable worldwide distribution in temperate and tropical waters. An uncommon species, it is typically lumped together under the name "mako" with...



Longfin smelt

Spirinchus thaleichthys

The longfin smelt (Spirinchus thaleichthys) is a smelt that is found in several estuaries and lakes along the northern Pacific coast of North America. Its most distinctive characteristic is the long pectoral fins that reach nearly to the base of...



Longfin trevally

Carangoides armatus

The longfin trevally (*Carangoides armatus*), also known as the longfin kingfish, longfin cavalla or armed trevally, is a species of inshore marine fish in the jack family, Carangidae. The species is common in tropical to subtropical waters of the...



Longfin yellowtail

Seriola rivoliana

The longfin yellowtail (*Seriola rivoliana*), also known as the almaco or silvercoat jack, deep-water, falcate, European or highfin amberjack, rock salmon, longfin or yellow kingfish, is a game fish of the family Carangidae; they are in the same...



Longhorn cowfish

Lactoria cornuta

The longhorn cowfish (*Lactoria cornuta*), also called the horned boxfish, is a species of boxfish from the family Ostraciidae, recognizable by its long horns that protrude from the front of its head, rather like those of a cow or bull. They are a...



Longhorn sculpin

Myoxocephalus octodecemspinosus

The longhorn sculpin (*Myoxocephalus octodecemspinosus*) is a species of marine ray-finned fish belonging to the family Cottidae, the typical sculpins. This species is found in the Northwest Atlantic Ocean. It is a predatory and scavenging fish that...



Longjaw mudsucker

Gillichthys mirabilis

The longjaw mudsucker *Gillichthys mirabilis* is a goby (Gobiidae) of the Pacific Ocean coast of California and Baja California, noted for its extremely large mouth and ability to survive out of water for short periods. As the common name suggests,...



Longlure frogfish

Antennarius multiocellatus

The longlure frogfish (Antennarius multiocellatus) is a species of frogfish in the genus Antennarius found in the western Atlantic from Bermuda and the Bahamas, south along the coasts of Central and South America, to Brazil around warm shallow...



Longnose dace

Rhinichthys cataractae

The longnose dace (Rhinichthys cataractae) is a freshwater minnow native to North America. Rhinichthys means snout fish (reference to the long snout) and cataractae means of the cataract (first taken from Niagara Falls). Longnose dace are small,...



Longnose gar

Lepisosteus osseus

The longnose gar (Lepisosteus osseus), also known as longnose garpike or billy gar, is a ray-finned fish in the family Lepisosteidae. The genus may have been present in North America for about 100 million years. References are made to gars being...



Longnose hawkfish

Oxycirrhites typus

The longnose hawkfish (Oxycirrhites typus) is a species of marine ray-finned fish, a hawkfish belonging to the family Cirrhitidae. It is found on tropical reefs of the Indian Ocean and the Pacific Ocean, where it can be found at depths around 10...



Longnose sawshark

Pristiophorus cirratus

The longnose sawshark (Pristiophorus cirratus) or common sawshark, is a sawshark of the family Pristiophoridae.



Longnose stingray

Dasyatis guttata

The longnose stingray (Hypanus guttatus) is a species of stingray in the family Dasyatidae, native to the western Atlantic Ocean from the southern Gulf of Mexico to Brazil. Found in coastal waters no deeper than 36 m (118 ft), this demersal...



Longnose sucker

Catostomus catostomus

The longnose sucker (Catostomus catostomus) is a species of cypriniform freshwater fish in the family Catostomidae. It is native to North America from the northern United States to the top of the continent. It is also found in Russia in rivers of...



Longnose trevally

Carangoides chrysophrys

The longnose trevally (Carangoides chrysophrys), also known as the tea-leaf trevally, club-nosed trevally, grunting trevally or dusky trevally, is a species of inshore marine fish in the jack family, Carangidae. The species is distributed...



Longnose velvet dogfish

Centroselachus crepidater

The longnose velvet dogfish (*Centroselachus crepidater*) is a sleeper shark of the family Somniosidae, found circumglobally in southern hemisphere subtropical seas, at depths of between 230 and 1,500 m. It reaches a length of 130 cm. It has a diet...



Longrakered trevally

Ulua mentalis

The longrakered trevally (Ulua mentalis), also known as the cale cale trevally and heavyjawed kingfish, is a species of marine fish in the jack and horse mackerel family Carangidae. The longrakered trevally is distributed throughout the tropical...



Longsnout dogfish

Deania quadrispinosa

Sharks portalThe longsnout dogfish (Deania quadrispinosa) is a little-known deepwater dogfish, found in the Atlantic and Indian Oceans from Namibia to Mozambique and in the South Pacific off southern Australia and New Zealand.The longsnout...



Longspine snipefish

Macroramphosus scolopax

The longspine snipefish, bellowfish, common bellowsfish, snipe-fish, snipefish, spine trumpet fish, or trumpetfish, Macroramphosus scolopax, is a snipefish of the genus Macroramphosus. It is also known as the slender snipefish off the South...



Longspine squirrelfish

Holocentrus rufus

The longspine squirrelfish (Holocentrus rufus) is a silvery red, sea fish with orange-gold body stripes. One of about 150 species of squirrelfish, their most distinguishing characteristics are their large eyes and the long third spine of the anal...



Longspined bullhead

Taurulus bubalis

The long-spined bullhead (Taurulus bubalis), also known as the long-spined sea-scorpion, and the long-spined scorpion fish is a coastal fish of the sculpin family Cottidae, inhabiting marine waters of Europe.



Longtail stingray

Dasyatis longa

The longtail stingray (Hypanus longus, often misspelled longa), is a species of stingray in the family Dasyatidae, found in the eastern Pacific Ocean from Baja California to Colombia. It inhabits sandy habitats down to a depth of 90 m (300 ft)....



Lookdown

Selene vomer

The lookdown (Selene vomer) is a species of game fish in the family Carangidae. It was first described in 1758 by Carl Linnaeus, in the 10th edition of Systema Naturae.



Lophiosilurus alexandri

Lophiosilurus alexandri

Lophiosilurus alexandri is a species of catfish (order Siluriformes) of the family Pseudopimelodidae, and the only species of the monotypic genus Lophiosilurus.



Lophius americanus

Lophius americanus

Lophius americanus is a goosefish in the family Lophiidae, also called all-mouth, American anglerfish, bellows-fish, devil-fish, headfish, molligut, satchel-mouth, or wide-gape. It is native to the eastern coast of North America.



Lophius budegassa

Lophius budegassa

Lophius budegassa, the blackbellied angler, is a species of anglerfish in the family Lophiidae. It was described by Maximilian Spinola in 1807. The fish is found in a depth range of 300–1,000 metres (1,000–3,300 ft) in the eastern Ionian Sea...



Lophius piscatorius

Lophius piscatorius

Lophius piscatorius, commonly known as the angler, European angler or common monkfish, is a monkfish in the family Lophiidae. It is found in coastal waters of the northeast Atlantic, from the Barents Sea to the Strait of Gibraltar, the...



Lost River sucker

Deltistes luxatus

The Lost River sucker (Deltistes luxatus) is a species of ray-finned fish in the family Catostomidae. It is the only living member of the genus Deltistes. It is found only in California and Oregon. Its population is much reduced from historical...



Lost shark

Carcharhinus obsolerus

The lost shark (Carcharhinus obsoletus), previously known as the false smalltail shark, is a possibly extinct species of requiem shark (family Carcharhinidae). It is known only from the Western Central Pacific Ocean, in the southern South China Sea.



Louisiana pancake batfish

Halieutichthys intermedius

The Louisiana pancake batfish, Halieutichthys intermedius, belongs to the batfish family Ogcocephalidae. It is native to the Gulf of Mexico, and was discovered in 2010. The known range of the species lies within the area of the Gulf of Mexico...



Louvar

Luvarus imperialis

The louvar or luvar (Luvarus imperialis) is a species of marine ray-finned fish, it is the only extant species in the genus Luvarus and family Luvaridae. This taxon is classified within the suborder Acanthuroidei, which includes the surgeonfish,...



Lucania goodei

Lucania goodei

Lucania goodei, the bluefin killifish, is a small species of fish in the topminnow family Fundulidae. It is native to the southeastern United States, but has been introduced to California, Texas and North Carolina. Other common names for the fish...



Luciocephalus pulcher

Luciocephalus pulcher

Luciocephalus pulcher, the pikehead, giant pikehead or crocodile pikehead, is a species of gourami native to the Malaya Peninsula, Sumatra, and Borneo. It is a yellowish-brown fish with dark brown longitudinal bands and stripes, and can reach a...



Luna lionfish

Pterois lunulata

The Luna lionfish (Pterois lunulata), the dragon's beard fish or Japanese lionfish, is a species of marine ray-finned fish belonging to the family Scorpaenidae, which consists of scorpionfishes and lionfishes. It is found in the western Pacific...



Lusitanian toadfish

Halobatrachus didactylus

The Lusitanian toadfish (Halobatrachus didactylus) is a species of toadfish found along the Atlantic and Mediterranean Coasts of western Europe and western Africa, from the Bay of Biscay to Ghana, with an isolated and old northern record from the...



Lutjanus agennes

Lutjanus agennes

Lutjanus agennes, the African red snapper, Guinean snapper or African cubera snapper, is a species of marine ray-finned fish belonging to the family Lutjanidae, the snappers. It is native to the coastal Atlantic waters of Africa.



Lutjanus bohar

Lutjanus bohar

Lutjanus bohar, the two-spot red snapper, the red bass, twinspot snapper or bohar snapper, is a species of marine ray-finned fish, a snapper belonging to the family Lutjanidae. It has a wide Indo-Pacific distribution.



Lutjanus carponotatus

Lutjanus carponotatus

Lutjanus carponotatus, the Spanish flag snapper, stripey snapper, dusky-striped sea-perch, gold-banded sea perch, gold-stripe sea-perch, striped seaperch or stripey seaperch, is a species marine ray-finned fish, a snapper belonging to the family...



Lutjanus decussatus

Lutjanus decussatus

Lutjanus decussatus, the checkered snapper, checkered seaperch or cross-hatched snapper, is a species marine ray-finned fish belonging to the family Lutjanidae. It is native to the western Pacific and Indian Oceans.



Lutjanus erythropterus

Lutjanus erythropterus

Lutjanus erythropterus, the crimson snapper, crimson seaperch, high-brow sea-perch, Longman's sea perch, red bream, saddle-tailed perch, small-mouth nannygai or smallmouth sea perch is a species of marine ray-finned fish, a snapper belonging to...



Lutjanus fulviflamma

Lutjanus fulviflamma

Lutjanus fulviflamma, the dory snapper, blackspot snapper, black-spot sea perch, finger-mark bream, long-spot snapper, Moses perch or red bream, is a species of marine ray-finned fish belonging to the family Lutjanidae, the snappers. It has a...



Lutjanus fulvus

Lutjanus fulvus

Lutjanus fulvus, the blacktail snapper, flametail snapper, redmargined seaperch, Waigeu snapper or yellowmargined sea perch, is a species of marine ray-finned fish, a snapper belonging to the family Lutjanidae. It is native to the Indo-West...



Lutjanus goldiei

Lutjanus goldiei

Lutjanus goldiei, the Papuan black snapper, Papuan black bass, New Guinea bass or Niugini black bass, is a species of freshwater and brackish water ray-finned fish, a snapper in the family Lutjanidae. It is found in the western Pacific Ocean.



Lutjanus johnii

Lutjanus johnii

Lutjanus johnii, John's snapper, the golden snapper, big-scaled bream, fingermark bream, fingerbanger, fingermark seaperch, John's sea-perch, or spotted-scale sea perch, is a species of marine ray-finned fish, a snapper belonging to the family...



Lutjanus malabaricus

Lutjanus malabaricus

Lutjanus malabaricus, the Malabar blood snapper, saddletail snapper, large-mouthed nannygai, large-mouthed sea-perch, Malabar snapper, nannygai, red bass, red bream, red emperor, red Jew, red snapper, saddletail seaperch, scarlet emperor or...



Lutjanus monostigma

Lutjanus monostigma

Lutjanus monostigma, the one-spot snapper, onspot seaperch or Moses snapper, is a species of marine ray-finned fish, a snapper belonging to the family Lutjanidae. It is found in the Indo-Pacific region.



Lutjanus rivulatus

Lutjanus rivulatus

Lutjanus rivulatus, the blubberlip snapper, Maori snapper, blue-spotted seaperch, Maori bream, Maori seaperch, multi-coloured snapper, scribbled snapper, speckled snapper or yellowfin snapper, is a species of marine ray-finned fish, a snapper...



Lutjanus russellii

Lutjanus russellii

Lutjanus russellii, Russell's snapper, Moses snapper, fingermark bream, Moses seaperch or Russell's sea-perch, is a species of marine ray-finned fish, a snapper belonging to the family Lutjanidae. It is native to the western Pacific Ocean.



Lutjanus sebae

Lutjanus sebae

Lutjanus sebae, also known as red emperor, emperor red snapper, emperor snapper, government bream, king snapper, queenfish or red kelp, is a species of marine ray-finned fish, a snapper belonging to the family Lutjanidae. It is native to the...



Lutjanus vitta

Lutjanus vitta

Lutjanus vitta, the brownstripe red snapper, brownstripe snapper, broadband seaperch, brownstripe seaperch, one-band sea-perch, one-lined snapper or striped seaperch, is a species of marine ray-finned fish, a snapper belonging to the family...



Lythrypnus dalli

Lythrypnus dalli

Lythrypnus dalli, commonly known as the blue-banded goby or Catalina goby, is a species of goby. It is native to the eastern Pacific where it is found from Monterey Bay, California to northern Peru, including the Gulf of California. It can be...



Mackerel icefish

Champscephalus gunnari

The mackerel icefish (Champscephalus gunnari) is a benthopelagic species of fish found in the Southern Ocean and the southernmost waters of the Atlantic Ocean. They are mainly to be found near Heard and McDonald Islands, Îles Kerguelen and...



Mackerel scad

Decapterus macarellus

The mackerel scad (Decapterus macarellus), or speedo, is a species of fish of the family, Carangidae. While it can be considered gamefish, it is usually used as bait. It is popular for consumption in Hawai'i, the Philippines and the U.A.E. In...



Macquarie perch

Macquaria australasica

The Macquarie perch (Macquaria australasica) is an Australian native freshwater fish of the Murray-Darling river system. It is a member of the family Percichthyidae and is closely related to the golden perch (Macquaria ambigua).The Macquarie...



Macrognathus pancalus

Macrognathus pancalus

Macrognathus pancalus, the barred spiny eel or Indian spiny eel, or পাঁকাল in bengali is a small freshwater fish in southern Asia. It usually is found in slow and shallow rivers. Males are more slender and often smaller than the females.



Macrognathus siamensis

Macrognathus siamensis

The peacock eel or spotfin spiny eel (Macrognathus siamensis) is a spiny eel found in freshwater habitats throughout Southeast Asia. They are commercially important as food and aquarium fish.



Macropharyngodon meleagris

Macropharyngodon meleagris

Macropharyngodon meleagris, the black-spotted wrasse, Eastern leopard wrasse or reticulated wrasse, is a species of ray-finned fish from the family Labridae, the wrasses. This species is native to the eastern Indian Ocean and the Pacific Ocean....



Macrotocinclus

Macrotocinclus affinis

The golden otocinclus (Macrotocinclus affinis, formerly Otocinclus affinis) is one of the smallest known suckermouth catfish, often called a 'dwarf oto'. Endemic to Southeast Brazil, this herbivorous, rheophilic, bottom-feeder only grows to...



Macrourus berglax

Macrourus berglax

Macrourus berglax, also known as the roughhead grenadier or onion-eye grenadier, is a species of marine ray-finned fish in the family Macrouridae. It is a deep-water fish found in the Atlantic Ocean.



Madeira rockfish

Scorpaena maderensis

The Madeira rockfish (*Scorpaena maderensis*) is a species of scorpionfish (*Scorpaenidae*) in the genus *Scorpaena*, found in the coastal waters of the eastern Atlantic Ocean and the Mediterranean Sea. This species reaches a length of around 14...



Magnificent rabbitfish

Siganus magnificus

The magnificent rabbitfish (*Siganus magnificus*), also known as the magnificent foxface or the Andaman foxface, is a species of marine ray-finned fish, a rabbitfish belonging to the family Siganidae. It is from the eastern Indian Ocean. It...



Magpie perch

Cheilodactylus nigripes

The magpie perch (*Cheilodactylus (Pseudogoniistius) nigripes*), magpie morwong or black-striped morwong, is a species of marine ray-finned fish, traditionally regarded as belonging to the family Cheilodactylidae, the members of which are commonly...



Mahi-mahi

Coryphaena hippurus

The mahi-mahi or common dolphinfish (*Coryphaena hippurus*) is a surface-dwelling ray-finned fish found in off-shore temperate, tropical, and subtropical waters worldwide. Also widely called dorado (not to be confused with *Salminus brasiliensis*, a...



Malabar danio

Devario malabaricus

The Malabar danio (*Devario malabaricus*) is a tropical fish belonging to the minnow family (*Cyprinidae*). Originating in Sri Lanka and the west coast of India, the fish has been circulated throughout the world through the aquarium fish trade. It...



Malabar grouper

Epinephelus malabaricus

The Malabar grouper (Epinephelus malabaricus) also known as blackspot rockcod, estuary rockcod, giant rock cod, greasy grouper, Malabar rockcod, Morgan's cod or speckled grouper, is a species of marine ray-finned fish, a grouper from the...



Malabar snakehead

Channa diplogramma

The Malabar snakehead (Channa diplogramma) is a vulnerable species of snakehead from fresh water in the Western Ghats in India. Until 2011, its scientific name was usually considered to be a synonym of C. micropeltes, the giant snakehead.



Malabar trevally

Carangoides malabaricus

The Malabar trevally (Carangoides malabaricus), also known as the Malabar jack, Malabar kingfish or nakedshield kingfish, is a species of large inshore marine fish of the jack family, Carangidae. It is distributed throughout the Indian and west...



Malacosteus australis

Malacosteus australis

Malacosteus australis, the southern stoplight loosejaw, is a species of barbeled dragonfish. This species is mainly distinguished from Malacosteus niger by a smaller postorbital photophore in both sexes and lower numbers of lateral photophores....



Malacosteus niger

Malacosteus niger

Malacosteus niger, commonly known as the black dragon fish, is a species of deep-sea fish. Some additional common names for this species include: northern stoplight loosejaw, lightless loosejaw, black loosejaw, and black hinged-head. It belongs...



Malapterurus electricus

Malapterurus electricus

Malapterurus electricus is a thickset fish with six mouth barbels and a single fin on its back, just anterior to the rounded tail fin. It is brownish or grayish, irregularly spotted with black, and attains a length and weight of about 1.2 metres...



Malawi eyebiter

Dimidiochromis compressiceps

The Malawi eyebiter (Dimidiochromis compressiceps) is a species of fish in the family Cichlidae. This predatory cichlid is endemic to Lake Malawi in East Africa.



Malawi squeaker

Synodontis njassae

The Malawi squeaker (*Synodontis njassae*) is a species of upside-down catfish endemic to Lake Malawi. This species grows to a length of 19.2 centimetres (7.6 in) TL. This species is a minor component of local commercial fisheries and can also be...



Maltese skate

Leucoraja melitensis

The Maltese skate (*Leucoraja melitensis*), or Maltese ray, is a species of fish in the family Rajidae. It is a rare endemic species from the Mediterranean Sea found in the coastal waters of Algeria, Italy, Malta, Tunisia, Greece and Turkey. Its...



Maluti redfin

Pseudobarbus quathlambae

The Maluti redfin (Pseudobarbus quathlambae) is a ray-finned fish species in the family Cyprinidae. It is colloquially called the Maluti minnow, but it is not a true minnow. It is Lesotho's only endemic freshwater fish, and threatened by the...



Man-of-war fish

Nomeus gronovii

Nomeus gronovii, the man-of-war fish or bluebottle fish, is a species of fish in the family Nomeidae, the driftfish. It is a distinct species characterized by an elongated body, large eyes, and blackish-blue stripes, growing up to 39 cm long....



Mandarin dogfish

Cirrhigaleus barbifer

Sharks portalThe mandarin dogfish (Cirrhigaleus barbifer) is a dogfish, a member of the family Squalidae in the order Squaliformes. It is found at depths of 140–650 metres (460–2,130 ft) off southern Japan, Taiwan, and Indonesia (Bali and...



Mangar

Luciobarbus esocinus

The Mangar or pike barbel, (Luciobarbus esocinus) is a large species of ray-finned fish in the genus Luciobarbus, native to the Tigris–Euphrates river system in Iran, Iraq, Syria and Turkey. The species is highly prized as a food fish, but it has...



Mango tilapia

Sarotherodon galilaeus

The mango tilapia (Sarotherodon galilaeus) is a species of fish from the cichlid family that is native to fresh and brackish waters in Africa and the Levant. Other common names include Galilaea tilapia, Galilean comb, Galilee St. Peter's fish,...



Mangrove red snapper

Lutjanus argentimaculatus

The mangrove red snapper (Lutjanus argentimaculatus), also known as mangrove jack, grey snapper, creek red bream, Stuart evader, dog bream, purple sea perch, red bream, red perch, red reef bream, river roman, or rock barramundi, is a species of...



Mangrove rivulus

Kryptolebias marmoratus

The mangrove rivulus or mangrove killifish, Kryptolebias marmoratus (syn. Rivulus marmoratus), is a species of killifish in the family Rivulidae. It lives in brackish and marine waters (less frequently in fresh water) along the coasts of Florida,...



Mangrove snapper

Lutjanus griseus

The mangrove snapper or gray snapper (*Lutjanus griseus*) is a species of snapper native to the western Atlantic Ocean from Massachusetts to Brazil, the Gulf of Mexico, Bermuda, and the Caribbean Sea. The species can be found in a wide variety of...



Mangrove whipray

Urogymnus granulatus

The mangrove whipray (*Urogymnus granulatus*) or whitetail stingray, is a species of stingray in the family Dasyatidae. It is widely distributed in the Indo-Pacific region from the Red Sea to northern Australia and Micronesia. A benthic inhabitant...



Map puffer

Arothron mappa

The map puffer (*Arothron mappa*), also known as the map pufferfish, scribbled pufferfish, or Kesho-fugu, is a demersal marine fish belonging to the family Tetraodontidae. The map puffer is typically found in tropical and subtropical waters from...



Marbled electric ray

Torpedo marmorata

The marbled electric ray (*Torpedo marmorata*) is a species of electric ray in the family Torpedinidae found in the coastal waters of the eastern Atlantic Ocean from the North Sea to South Africa. This benthic fish inhabits rocky reefs, seagrass...



Marbled grouper

Dermatolepis inermis

The marbled grouper (*Dermatolepis inermis*), donkey fish, mutton hamlet, rockhind and sicklefish grouper, is a species of marine ray-finned fish, a grouper from the subfamily Epinephelinae which is part of the family Serranidae, which also...



Marbled hatchetfish

Carnegiella strigata

The marbled hatchetfish (Carnegiella strigata) is a small, normally 3.5 cm (1.4 in) in length, freshwater ray-finned fish native to South America. Hatchet shaped, it presents a gold line extending from its eye to its caudal fin while the area...



Marbled parrotfish

Leptoscarus vaigiensis

The marbled parrotfish (Leptoscarus vaigiensis), also known as the seagrass parrotfish, is a species of marine ray-finned fish, a parrotfish from the family Scaridae and is the only known member of the genus Leptoscarus. It has a wide...



Marbled rockcod

Notothenia rossii

The marbled rockcod (Notothenia rossii) is a species of marine ray-finned fish, belonging to the family Nototheniidae, the nototheniids or cod icefishes. It is native to the Southern Ocean, where it can be found at depths from 5 to 350 m (16 to...



Marbled swordtail

Xiphophorus meyeri

The marbled swordtail (Xiphophorus meyeri) is a species of freshwater fish in the family Poeciliidae. It was endemic to the Rio Salado system (itself a part of the Rio Grande basin) in Coahuila, northeastern Mexico. It was restricted to springs....



Marbled whipray

Fluvitrygon oxyrhyncha

The marbled whipray (Fluvitrygon oxyrhynchus) is a little-known species of stingray in the family Dasyatidae, native to several freshwater rivers in Southeast Asia. This species has an oval pectoral fin disc with an elongated, pointed snout and a...



Maroon clownfish

Premnas biaculeatus

Premnas biaculeatus, commonly known as spine-cheeked anemonefish or the maroon clownfish, is a species of anemonefish found in the Indo-Pacific from western Indonesia to Taiwan and the Great Barrier Reef. They can grow up to be about 17 cm (6.7...



Mary River cod

Maccullochella mariensis

The Mary River cod (Maccullochella mariensis) is a species of temperate perch native to the coastal Mary River system of southern Queensland, Australia. Mary River cod are one of Australia's most endangered freshwater fishes and are notable for...



Masked puffer

Arothron diadematus

The masked puffer (Arothron diadematus) is a pufferfish in the family Tetraodontidae.



Mastacembelus ellipsifer

Mastacembelus ellipsifer

Mastacembelus ellipsifer is a species of spiny eel that is endemic to Lake Tanganyika in Africa and sometimes kept in aquariums. Although sometimes called the Tanganyikan spiny eel, it is only one of fifteen spiny eel species in the Tanganyikan...



Mayaheros urophthalmus

Mayaheros urophthalmus

Mayaheros urophthalmus, also known as the Mayan cichlid or Mexican mojarra is a species of cichlid.



Maylandia callainos

Maylandia callainos

Maylandia callainos (sometimes referred to as cobalt zebra, cobalt blue mbuna or cobalt blue zebra cichlid) is a species of cichlid endemic to Lake Malawi where they only occurred naturally in Nkhata Bay though it has now been introduced to other...



Maylandia estherae

Maylandia estherae

Maylandia estherae (the Red Zebra mbuna, Red Zebra Cichlid, or Esther Grant's Zebra) is a haplochromine cichlid. It is a rock dwelling fish or mbuna from Lake Malawi. This fish, like most cichlids from Lake Malawi, is a mouthbrooder - females...



Maylandia lombardoi

Maylandia lombardoi

Maylandia lombardoi, is a 13 centimetres (5.1 in) long freshwater fish from the family Cichlidae. This species is popular in the aquarium hobby where it is sold under a variety of common names including: lombardoi mbuna, kenya mbuna or kennyi...



Mbu pufferfish

Tetraodon mbu

The Mbu pufferfish, also known as Mbuna pufferfish, giant pufferfish, or giant freshwater pufferfish (Tetraodon mbu), is a carnivorous freshwater pufferfish originating from the middle and lower sections of the Congo River in Africa, as well as...



McCloud River redband trout

Oncorhynchus mykiss stonei

The McCloud River redband trout (Oncorhynchus mykiss stonei) is one of three redband trout subspecies of the rainbow trout in the family Salmonidae. The trout is native in small tributaries of the McCloud River and Pit River which are tributaries...



Mediterranean horse mackerel

Trachurus mediterraneus

The Mediterranean horse mackerel (Trachurus mediterraneus), also known as the Black Sea horse mackerel, horse mackerel, Mediterranean scad, common scad, or simply scad, is a species of mackerel in the family Carangidae found in the eastern...



Mediterranean killifish

Aphanius fasciatus

The Mediterranean killifish, Mediterranean banded killifish or South European toothcarp (Aphanius fasciatus) is a species of fish in the family Cyprinodontidae. It is found in Albania, Algeria, Bosnia and Herzegovina, Croatia, Cyprus, Egypt,...



Mediterranean moray

Muraena helena

The Mediterranean moray (Muraena helena), also known as Roman eel, is a species of fish in the moray eel family. It has a long eel-like body and is found in the eastern Atlantic Ocean and Mediterranean Sea. Its bite can be dangerous to humans.



Mediterranean rainbow wrasse

Coris julis

The Mediterranean rainbow wrasse (Coris julis) is a small, colourful fish in the family Labridae. It can be found in the Mediterranean Sea and in the northeast Atlantic Ocean from Sweden to Senegal (though it is a rare wanderer to the southern...



Mediterranean sand smelt

Atherina hepsetus

The Mediterranean sand smelt, Atherina hepsetus, is a species of fish in the family Atherinidae.



Mediterranean scaldfish

Arnoglossus laterna

The Mediterranean scaldfish (Arnoglossus laterna), also known as the scaldfish, is a species of benthic left eyed flatfish belonging to the family Bothidae. It is found in the eastern Atlantic of Europe and Africa as well as the Mediterranean,...



Mediterranean spearfish

Tetrapturus belone

The Mediterranean spearfish (Tetrapturus belone) is a species of marlin native to the Mediterranean Sea where it is particularly common around Italy, although there is a probable record of one caught off Madeira. It is an open-water fish, being...



Megalancistrus parananus

Megalancistrus parananus

Megalancistrus parananus is a species of armored catfish native to Argentina, Brazil, Paraguay and Uruguay where it is found in the Paraguay, Paraná, and Uruguay River basins. This species grows to a length of 58.8 centimetres (23.1 in) SL.



Megaleporinus obtusidens

Megaleporinus obtusidens

Megaleporinus obtusidens is a South American species of headstander that inhabits the basins of the Paraná River and the Río de la Plata, the Uruguay River, and the São Francisco River (including minor rivers and tributaries like the Bermejo,...



Megalodoras uranoscopus

Megalodoras uranoscopus

The giant talking catfish or giant raphael catfish (Megalodoras uranoscopus) is a species of thorny catfish that is native to the Amazon Basin in Bolivia, Brazil, Colombia, Ecuador and Peru. This species grows to a length of 60 centimetres (24...



Megamouth shark

Megachasma pelagios

The megamouth shark (Megachasma pelagios) is a species of deepwater shark. Rarely seen by humans, it measures around 5.2 m (17 ft) long and is the smallest of the three extant filter-feeding sharks alongside the relatively larger whale shark and...



Megrim

Lepidorhombus whiffiagonis

The megrim, megrim sole, whiff, or Cornish sole (Lepidorhombus whiffiagonis) is a species of left-eyed flatfish in the family Scophthalmidae. It is found in the northeast Atlantic and Mediterranean Sea between 100 and 700 m (330 and 2,300 ft)...



Meiacanthus grammistes

Meiacanthus grammistes

Meiacanthus grammistes, the striped blenny, also called the striped fang blenny, grammistes blenny, line-spot harptail blenny or striped poison-fang blenny, is a species of combtooth blenny from the western Pacific Ocean. The fish stays in the...



Mekong freshwater stingray

Hemitygon laosensis

The Mekong freshwater stingray, Hemitygon laosensis, is a species of stingray in the family Dasyatidae, restricted to the Mekong and Chao Phraya Rivers in Laos and Thailand; the occurrence in Chao Phraya is considered an introduction. Measuring...



Mekong giant catfish

Pangasianodon gigas

The Mekong giant catfish (Pangasianodon gigas; Thai: ปลาบึก, RTGS: pla...



Melanocetus eustalus

Melanocetus eustalus

Melanocetus eustalus is a deep sea anglerfish in the family Melanocetidae, found off the Pacific coast of Mexico at depths down to about 1,675 m (5,500 ft).



Melanocetus murrayi

Melanocetus murrayi

Melanocetus murrayi, commonly known as Murray's abyssal anglerfish, is a deep sea anglerfish in the family Melanocetidae, found in tropical to temperate parts of the world's oceans at depths down to over 2,000 m (6,600 ft). Its length is up to...



Melanocetus niger

Melanocetus niger

Melanocetus niger is a species of black seadevil, a type of anglerfish. They have big mouths, with plenty of long teeth, and numerous fin rays. The fish is bathypelagic and has been found at depths ranging from 1,200 to 2,000 metres (3,900 to...



Melanochromis auratus

Melanochromis auratus

Melanochromis auratus, the auratus cichlid, is a freshwater fish of the cichlid family. It is also known as golden mbuna and Malawi golden cichlid. It is endemic to the southern region of Lake Malawi, particularly from Jalo Reef southward along...



Melanotaenia duboulayi

Melanotaenia duboulayi

Melanotaenia duboulayi, the crimson-spotted rainbowfish, less commonly known as the Duboulay's rainbowfish, is a species of freshwater fish endemic to coastal eastern Australia, although M. duboulayi has also been kept as an aquarium fish since...



Melon barb

Haludaria fasciata

The melon barb (Haludaria fasciata) is a common species of cyprinid fish that is endemic to rivers in Goa, Karnataka, Kerala and Tamil Nadu in the Western Ghats of South India. They live in a tropical climate in water that typically has a pH of...



Melon butterflyfish

Chaetodon trifasciatus

The melon butterflyfish (*Chaetodon trifasciatus*) or the Indian redfin butterflyfish, is a species of marine ray-finned fish, a butterflyfish belonging to the family Chaetodontidae. It is found in the Indian Ocean from East Africa to Western Java....



Mene maculata

Mene maculata

Mene maculata, the moonfish, is the only extant member of the genus Mene and of the family Menidae. The body is highly compressed laterally and very deep vertically. The ventral profile is steep, with a sharp ventral edge. The caudal (tail) fin...



Menticirrhus americanus

Menticirrhus americanus

Menticirrhus americanus, the southern kingfish, southern kingcroaker, king whiting, Carolina whiting, sea mullet, roundhead, or whiting, is a species of marine fish in the family Sciaenidae. It lives in shallow coastal waters on the western...



Menticirrhus saxatilis

Menticirrhus saxatilis

Menticirrhus saxatilis, the northern kingfish or northern kingcroaker, is a species of marine fish in the family Sciaenidae (commonly known as the "drum" or "croaker" family). It lives in the shallow coastal waters of the western Atlantic Ocean...



Merlangius

Merlangius merlangus

Merlangius merlangus, commonly known as whiting or merling, is an important food fish in the eastern North Atlantic Ocean and the northern Mediterranean, western Baltic, and Black Sea. In Anglophonic countries outside the Whiting's natural range,...



Merluccius australis

Merluccius australis

Merluccius australis, the southern hake, is a species of fish from the family Merlucciidae, the true hakes. It is found in the southern Pacific and Atlantic Oceans with two disjunct populations, one around southern South America and the other in...



Merluccius capensis

Merluccius capensis

Merluccius capensis (shallow-water Cape hake or South African hake) is a ray-finned fish in the genus Merluccius, found in the south-eastern Atlantic Ocean, along the coast of South Africa. It is a long, lean fish with a large head, similar in...



Merluccius gayi

Merluccius gayi

Merluccius gayi is a merluccid hake of the genus Merluccius, with two subspecies, the South Pacific hake or Chilean common hake (M. g. gayi) and the Peruvian hake (M. g. peruanus), found in the south-western Pacific Ocean, along the coast of...



Merluccius merluccius

Merluccius merluccius

Merluccius merluccius, the European hake, is a merluccid hake of the genus Merluccius. Other vernacular names include Cornish salmon and herring hake. It is a predatory species which was often netted alongside one of its favoured prey, the...



Merluccius paradoxus

Merluccius paradoxus

Merluccius paradoxus, the deep-water Cape hake, is a merluccid hake of the genus Merluccius, found in the south-eastern Atlantic Ocean, along the coast of Southern Africa, south of Angola. Its range extends in decreasing abundance around the...



Merluccius patagonicus

Merluccius patagonicus

Merluccius patagonicus, the Patagonian hake, is a species of fish from the family Merlucciidae from the western South Atlantic which was described in 2003. However some authorities consider the distinguishing features of M. patagonicus to be...



Mesonauta festivus

Mesonauta festivus

Mesonauta festivus, the flag cichlid, is a species of cichlid native to the Paraná, Paraguay, Madre de Dios, Guaporé, Mamoré, Jamari and Tapajós river basins in Brazil, Peru, Paraguay and Bolivia. It can reach a standard length of 12.1 cm (4.8...



Messmate pipefish

Corythoichthys haematopterus

The messmate pipefish (Corythoichthys haematopterus) is a species of marine fish in the family Syngnathidae. It is widespread throughout the tropical waters of the Indo-Pacific region, from the eastern coast of Africa to the Vanuatu Islands. This...



Metynnis argenteus

Metynnis argenteus

Metynnis argenteus is a species of serrasalmid fish endemic to the Tapajós River Basin in Brazil. It is one of the species known in the aquarium trade as the "silver dollar". M. argenteus is generally considered the archetypal silver dollar...



Metynnis hypsauchen

Metynnis hypsauchen

Metynnis hypsauchen is a species of serrasalmid found in Amazon and Paraguay river basins, as well as rivers of the Guiana Shield. It is one of the fish known as the "silver dollar" in the aquarium trade. It has frequently been confused with the...



Metynnis lippincottianus

Metynnis lippincottianus

Metynnis lippincottianus, commonly known as the spotted silver dollar, is a species of serrasalmid native to the Brazilian Amazon and multiple rivers of French Guiana. An average *M. lippincottianus* can grow to a length of 13 cm. The fish is named...



Metynnis maculatus

Metynnis maculatus

Metynnis maculatus, the spotted metynnis, is a species of serrasalmid fish found in Argentina and Brazil, and possibly in Bolivia, Paraguay, and Peru. It is one of the species of fish known in the aquarium trade as the "silver dollar".



Mexican barred snapper

Hoplopagrus guentherii

The Mexican barred snapper (Hoplopagrus guentherii), also known as the barred pargo, is a species of marine ray-finned fish, a snapper belonging to the family Lutjanidae. It is native to the eastern Pacific Ocean. It is the only species in its...



Mexican blind brotula

Typhliasina pearsei

The Mexican blind brotula (Typhliasina pearsei) is a species of viviparous brotula endemic to Mexico, where it is found in sinkholes and caves. It is known as sak kay in Mayan and dama blanca ciega in Spanish. This cavefish grows to a standard...



Mexican golden trout

Oncorhynchus chrysogaster

The Mexican golden trout (*Oncorhynchus chrysogaster*) is a species of fish in the family Salmonidae. The species is endemic to high-elevation headwaters of the Fuerte River, Sinaloa River, and Culiacán River drainages in the Sierra Madre...



Mexican hornshark

Heterodontus mexicanus

The Mexican hornshark (Heterodontus mexicanus) is a bullhead shark of the family Heterodontidae. This shark is grey-brown in color, with black spots scattered on the fins and body. It has a cylindrical trunk, conical head, and small spiracles...



Mexican tetra

Astyanax mexicanus

The Mexican tetra (Astyanax mexicanus), also known as the blind cave fish, blind cave characin, and blind cave tetra, is a freshwater fish of the family Characidae of the order Characiformes. The type species of its genus, it is native to the...



Microctenopoma ansorgii

Microctenopoma ansorgii

Microctenopoma ansorgii is a small freshwater fish, known in the aquarium trade as the ornate ctenopoma, orange ctenopoma, ornate climbing perch, pretty ctenopoma, or rainbow ctenopoma. It is related to the more familiar spotted climbing perch...



Microdevario kubotai

Microdevario kubotai

Microdevario kubotai is a species of cyprinid found in southeast Asian rivers and streams. It belongs to the genus Microdevario, which contains small danionins. The type locality is in Ranong Province, Peninsular Thailand. It is also known from...



Microgadus tomcod

Microgadus tomcod

Microgadus tomcod, also commonly known as frostfish, Atlantic tomcod or winter cod, is a type of cod found in North American coastal waters from the Gulf of St. Lawrence, St. Lawrence River and northern Newfoundland, south to Virginia. The fishing...



Microspathodon chrysurus

Microspathodon chrysurus

The yellowtail damselfish (Microspathodon chrysurus) is a species of damselfish native to tropical areas such as the Caribbean coast of Panama. Damselfish are abundant in coral reef environments. The International Union for Conservation of Nature...



Microstomus pacificus

Microstomus pacificus

The Pacific Dover sole (Microstomus pacificus), also called the slime sole or slippery sole, is a Pacific flatfish of the flounder family which ranges from Baja California to the Bering Sea. It takes its name from a resemblance to the common sole...



Midas blenny

Ecsenius midas

Ecsenius midas, known commonly as the Midas blenny, Persian blenny, lyretail blenny or golden blenny, is a species of marine fish in the family Blenniidae. The Midas blenny is widespread throughout the tropical waters of the Indo-Pacific area from...



Midnight parrotfish

Scarus coelestinus

The midnight parrotfish (Scarus coelestinus) is a species of parrotfish that inhabits coral reefs mainly in the Caribbean, Bahamas, and Florida. The typical size is between 30 and 60 cm, but it can grow to almost 1 m. It has been observed as far...



Mikrogeophagus altispinosus

Mikrogeophagus altispinosus

Mikrogeophagus altispinosus is a species of fish endemic to the southern Amazon River basin in Brazil and Bolivia. The species is part of the family Cichlidae and subfamily Geophaginae. It is a popular aquarium fish, traded under the common names...



Milk shark

Rhizoprionodon acutus

The milk shark (Rhizoprionodon acutus) is a species of requiem shark, and part of the family Carcharhinidae, whose common name comes from an Indian belief that consumption of its meat promotes lactation. The largest and most widely distributed...



Milkfish

Chanos chanos

The milkfish (Chanos chanos) is the sole living species in the family Chanidae. However, there are at least five extinct genera from the Cretaceous. The repeating scientific name (tautonym) is from Greek khanos (χᾶνος ‘mouth’). The species has many...



Millet butterflyfish

Chaetodon miliaris

The millet butterflyfish (Chaetodon miliaris) is a species of butterflyfish in the family Chaetodontidae. Other common names include the lemon butterflyfish and the millet-seed butterflyfish. It is endemic to the Hawaiian Islands and the Johnston...



Mimic shiner

Notropis volucellus

The mimic shiner (Notropis volucellus) is a species of North American cyprinid freshwater fish in the genus Notropis. The genus Notropis is commonly known as the eastern shiners. It is native to areas of the Hudson Bay drainage, Great Lakes...



Mimoblennius cirrosus

Mimoblennius cirrosus

Mimoblennius cirrosus, the fringed blenny, is a species of combtooth blenny found in coral reefs in the western Indian Ocean. This species grows to a length of 5.4 centimetres (2.1 in) TL.



Mirogrex terraesanctae

Acanthobrama terraesanctae

Mirogrex terraesanctae, the Kinneret bream or Kinneret bleak, is a species of freshwater fish in the family Cyprinidae. It is known from two lakes: Lake Tiberias (Sea of Galilee, Lake Kinneret), Israel, and Lake Muzayrib, Syria. This is a small...



Mirror dory

Zenopsis nebulosa

The mirror dory (Zenopsis nebulosa) is a dory of the family Zeidae, found in the southern Pacific Ocean at depths of between 30 and 800 m. Its length is up to 70 cm. The mirror dory has a high first dorsal fin containing 9 spiny rays, and about 27...



Misgurnus fossilis

Misgurnus fossilis

The weatherfish (Misgurnus fossilis) is a species of true loach that has a wide range in Europe and some parts of Asia. It is an omnivorous scavenger bottom feeder, using its sensitive barbels to find edible items. The diet mostly consists of...



Moapa dace

Moapa coriacea

The moapa dace (Moapa coriacea) is a rare cyprinid fish of southern Nevada, United States, found only in the upper parts of the Muddy (formerly Moapa) River, and in the warm springs that give rise to the river. It is the only species of the...



Mobula eregoodootenkee

Mobula eregoodootenkee

Mobula eregoodootenkee, the pygmy devil ray or longhorned mobula, is a species of ray in the family Mobulidae. It is endemic to the Indian Ocean and central-west Pacific Ocean. It ranges from South Africa in the west to the Philippines in the...



Mobula kuhlii

Mobula kuhlii

Mobula kuhlii, the shortfin devil ray, is a species of ray in the family Mobulidae. It is endemic to the Indian Ocean and central-west Pacific Ocean. It ranges from South Africa, Mozambique, Tanzania and the Seychelles in the west to the...



Mobula munkiana

Mobula munkiana

Mobula munkiana, commonly known as the manta de monk, Munk's devil ray, pygmy devil ray, smoothtail mobula or Munk's pygmy devil ray is a species of ray in the family Mobulidae. It is found in tropical parts of the eastern Pacific Ocean, ranging...



Moggel

Labeo umbratus

Moggel (Labeo umbratus) is a freshwater African fish in genus Labeo. It occurs within the drainage basin of the Orange River. The species has been recorded in the, Olifants River in Mpumalanga and Limpopo, introduced there by anglers. This species...



Mogurnda adspersa

Mogurnda adspersa

Mogurnda adspersa (commonly known as the southern purple-spotted gudgeon) is a species of endangered gudgeon that is endemic to south-eastern mainland Australia (the states of South Australia, New South Wales and Victoria and Queensland). The...



Mogurnda mogurnda

Mogurnda mogurnda

Mogurnda mogurnda, commonly known as the northern trout gudgeon or northern purple-spotted gudgeon is a freshwater fish native to northern Australia and New Guinea.



Mohave tui chub

Siphateles bicolor mohavensis

The Mohave tui chub (Siphaletes bicolor mohavensis) is a subspecies of the tui chub endemic to the Mojave River. The Mohave tui chub was listed as endangered under the Endangered Species Act in 1970, then added to California's list of endangered...



Mola carplet

Amblypharyngodon mola

The mola carplet (Amblypharyngodon mola) is a species of carplet in the family Cyprinidae. It is found in Afghanistan, Pakistan, India, Bangladesh, and Myanmar, although IUCN considers its presence certain only in Pakistan, India, and Bangladesh....



Mola tecta

Mola tecta

Mola tecta, the hoodwinker sunfish, belongs to the family Molidae and genus Mola. It is closely related to the more widely known ocean sunfish (Mola mola). The Latin word "tecta" means hidden. The word "hidden" was adopted for the name because...



Mollisquama mississippiensis

Mollisquama mississippiensis

Mollisquama mississippiensis or the American pocket shark is a species of pocket shark native to the Gulf of Mexico. It is the second species of pocket shark to be described.



Molly Miller

Scartella cristata

Molly Miller (née Carter; born October 19, 1986) is an American women's college basketball coach at Grand Canyon University. As of 2023–24, she is leading the Antelopes women's basketball team in her fourth season after taking over in 2020–21....



Monkey goby

Neogobius fluviatilis

The monkey goby (Neogobius fluviatilis) is a species of goby native to the basins of the Black Sea and the Sea of Azov.



Monocentris japonica

Monocentris japonica

Monocentris japonica, the Japanese pineapplefish, is a pinecone fish of the family Monocentridae, found in the tropical Indo-West Pacific Oceans, at depths between 2 and 100 m and can be found on both rocky and coral reefs. The fish is nocturnal...



Monodactylus argenteus

Monodactylus argenteus

Monodactylus argenteus is a species of fish in the family Monodactylidae, the moonyfishes. Its common names include silver moonyfish, or silver moony, butter bream, and diamondfish. It is native to the western Pacific and Indian Oceans, including...



Monodactylus sebae

Monodactylus sebae

Monodactylus sebae, the African moony, is a species of moonyfish native to fresh, brackish and marine waters from the eastern Atlantic, ranging from the Canary Islands down to Angola. It inhabits mangrove swamps and estuaries and can occasionally...



Monognathus ahlstromi

Monognathus ahlstromi

Monognathus ahlstromi, the paddletail onejaw, is an ocean-dwelling eel found in the North Pacific Ocean off of the coast of the United States. It is found up to a depth of 2,000 m (6,600 ft). It does not provide parental care. Little information...



Monotaxis grandoculis

Monotaxis grandoculis

Monotaxis grandoculis, the humpnose big-eye bream, bigeye barenose, bigeye bream, bigeye emperor, grand-eyed porgy fish, humpnose sea-bream, large-eye bream, mu and roundtooth large-eye bream, is a species of marine ray-finned fish belonging to...



Montana Arctic grayling

Thymallus arcticus montanus

The Montana Arctic grayling (Thymallus arcticus montanus) is a North American freshwater fish in the salmon family Salmonidae. The Montana Arctic grayling, native to the upper Missouri River basin in Montana and Wyoming, is a disjunct population...



Montecristo cichlid

Oscura heterospila

The Montecristo cichlid (Oscura heterospila) is a species of freshwater fish from the Atlantic slope of southern Mexico and Guatemala. This cichlid occurs in lagoons, creeks and rivers with slight to moderate current in the Grijalva–Usumacinta,...



Monterrey platyfish

Xiphophorus couchianus

The Monterrey platyfish (Xiphophorus couchianus) is a species of freshwater fish in family Poeciliidae. A live-bearer, it was native to a very small section of the San Juan River system (itself part of the Rio Grande basin) in the vicinity of the...



Moon wrasse

Thalassoma lunare

The moon wrasse (Thalassoma lunare) also known as the crescent wrasse or lyretail wrasse, is a species of wrasse native to the Indian Ocean and the western Pacific Ocean. It is an inhabitant of coral reefs and surrounding areas at depths from 1...



Moonlight gourami

Trichopodus microlepis

The moonlight gourami (Trichopodus microlepis), also known as the moonbeam gourami, is a labyrinth fish of the family Osphronemidae native to Indochina. This peaceful species is a popular aquarium fish.



Moorish idol

Zanclus cornutus

The Moorish idol (Zanclus cornutus) is a species of marine ray-finned fish belonging to the family Zanclidae. It is the only member of the monospecific genus Zanclus and the only extant species within the Zanclidae. This species is found on reefs...



Mora moro

Mora moro

Mora moro, the common mora, is a deep-sea fish, the only species in the genus Mora. It is found worldwide in temperate seas, at depths of between 300 and 2,500 m. Its length is up to about 80 cm. Other names in English include goodly-eyed cod,...



Mormyrus longirostris

Mormyrus longirostris

Mormyrus longirostris, commonly referred as the eastern bottle-nosed mormyrid, is a medium-sized ray-finned fish species belonging to the family Mormyridae. It was originally described by Wilhelm Peters in Monatsberichte der Akad. Wiss. Berlin,...



Mosquitofish

Gambusia affinis

The western mosquitofish (Gambusia affinis) is a North American freshwater poeciliid fish, also known commonly, if ambiguously, as simply mosquitofish or by its generic name, Gambusia, or by the common name gambezi. Its sister species, the...



Mottled grouper

Mycteroperca rubra

The mottled grouper (Mycteroperca rubra) is a species of marine ray-finned fish, a grouper from the subfamily Epinephelinae which is part of the family Serranidae, which also includes the anthias and sea basses. It is found in the eastern...



Mottled sculpin

Cottus bairdii

The mottled sculpin (Cottus bairdii) is a species of freshwater ray-finned fish belonging to the family Cottidae, the typical sculpins. This species has a wide but scattered North American distribution. As the name suggests, its coloration is a...



Mottled skate

Beringraja pulchra

The mottled skate (Beringraja pulchra) is a species of skate in the family Rajidae. An inhabitant of shallow coastal waters, it is found in the northwestern Pacific Ocean off Korea, Japan, and China. This species grows to 1.12 m (3.7 ft) long and...



Mountain mullet

Dajaus monticola

The mountain mullet (Dajaus monticola) is a freshwater fish of the family Mugilidae. It can be found in North and South America, from North Carolina, Florida, Louisiana and Texas in the United States to Colombia and Venezuela, including the West...



Mountain sucker

Catostomus platyrhynchus

The mountain sucker (Catostomus platyrhynchus) is a sucker found throughout western North America, on both sides of the Rocky Mountains, including the upper Missouri River, Columbia River, Sacramento River, and Colorado River. It is not limited...



Mountain whitefish

Prosopium williamsoni

The mountain whitefish (Prosopium williamsoni) is one of the most widely distributed salmonid fish of western North America. It is found from the Mackenzie River drainage in Northwest Territories, Canada through western Canada and the...



Mouse catshark

Galeus murinus

The mouse catshark (*Galeus murinus*) is a species of catshark and part of the family Scyliorhinidae. It is common in the northeastern Atlantic Ocean from Iceland to Western Sahara. There is much taxonomic confusion regarding this species in...



Mozambique tilapia

Oreochromis mossambicus

The Mozambique tilapia (*Oreochromis mossambicus*) is an oreochromine cichlid fish native to southeastern Africa. Dull colored, the Mozambique tilapia often lives up to a decade in its native habitats. It is a popular fish for aquaculture. Due to...



Mrigal carp

Cirrhinus cirrhosus

The mrigal carp (*Cirrhinus cirrhosus*), (Bengali: মৃগেল, romanized: mrigél) also known as the white carp, is a species of ray-finned fish in the carp family. Native to streams and rivers in India, the only surviving wild population is in the...



Mud sunfish

Acantharchus pomotis

The mud sunfish (*Acantharchus pomotis*) is a freshwater ray-finned fish, a sunfish from the family Centrarchidae, which widely distributed in the fresh waters along the Atlantic coast of North America, ranging from New York to Alabama. It is the...



Mueller's pearlside

Maurolicus muelleri

Maurolicus muelleri, commonly referred to as Mueller's pearlside, Mueller's bristle-mouth fish (not to be confused with the Gonostomatidae), or the silvery lightfish (not to be confused with the Phosichthyidae), is a marine hatchetfish in the...



Muksun

Coregonus muksun

The muksun (Coregonus muksun) is a type of whitefish widespread in the Siberian Arctic waters. It is mostly found in the freshened areas of the Kara and Laptev Seas and up the major rivers, as well as in Lake Taymyr. Like all whitefishes, it is a...



Mullus barbatus

Mullus barbatus

Mullus barbatus (red mullet) is a species of goatfish found in the Mediterranean Sea, Sea of Marmara, the Black Sea and the eastern North Atlantic Ocean, where its range extends from Scandinavia to Senegal. They are fished, mostly by trawling....



Mummichog

Fundulus heteroclitus

The mummichog (Fundulus heteroclitus) is a small killifish found along the Atlantic coast of the United States and Canada. Also known as Atlantic killifish, mummies, gudgeons, and mud minnows, these fish inhabit brackish and coastal waters...



Muraena augusti

Muraena augusti

Muraena augusti is a moray eel found on the eastern Atlantic ocean on waters surrounding the archipelagos of the Azores, Madeira, the Canary Islands and Cape Verde (collectively known as Macaronesia). It was described by Johann Jakob Kaup in...



Muraena lentiginosa

Muraena lentiginosa

Muraena lentiginosa is a moray eel from the Eastern Pacific. It occasionally makes its way into the aquarium trade. It grows to a size of 61 centimetres (2 ft 0 in) in length.



Murray cod

Maccullochella peelii

The Murray cod (Maccullochella peelii) is a large Australian predatory freshwater fish of the genus Maccullochella in the family Percichthyidae. Although the species is called a cod in the vernacular, it is not related to the Northern Hemisphere...



Murray hardyhead

Craterocephalus fluviatilis

The Murray hardyhead (Craterocephalus fluviatilis) is a species of fish in the family Atherinidae endemic to inland parts of southeastern Australia. The fish is an omnivore, feeding on small crustaceans, aquatic insects and algae.



Murray River rainbowfish

Melanotaenia fluviatilis

The Murray River rainbowfish (Melanotaenia fluviatilis), known less commonly as the Australian rainbowfish, is a species of freshwater fish endemic to southeastern Australia. The southernmost species of all rainbowfishes, these fish are very...



Muskellunge

Esox masquinongy

The muskellunge (Esox masquinongy), often shortened to muskie, musky, ski, or lunge, is a species of large freshwater predatory fish native to North America. It is the largest member of the pike family, Esocidae.



Mutton snapper

Lutjanus analis

The mutton snapper (Lutjanus analis) is a species of marine ray-finned fish, a snapper belonging to the family Lutjanidae. It is found in the Western Atlantic Ocean.



Myanmar arowana

Scleropages inscriptus

The Batik Arowana or Myanmar arowana(Scleropages inscriptus), is a large, bony-tongued fish native to Myanmar. The species is found in the Tanintharyi River basin on the Indian Ocean coast of peninsular Myanmar.This species is differentiated from...



Mycteroperca bonaci

Mycteroperca bonaci

Mycteroperca bonaci, the black grouper, black rockfish or marbled rockfish, is a species of marine ray-finned fish, a grouper from the subfamily Epinephelinae which is part of the family Serranidae, which also includes the anthias and sea basses....



Mycteroperca fusca

Mycteroperca fusca

Mycteroperca fusca, the Island grouper or comb grouper, is a species of grouper from the family Serranidae which is endemic to the Macaronesian Islands in the eastern Atlantic. It is classified as Vulnerable in The IUCN Red List of Endangered...



Mycteroperca interstitialis

Mycteroperca interstitialis

The yellowmouth grouper (Mycteroperca interstitialis), also known as the crossband rockfish, grey maddock, hamlet, harlequin rockfish, princess rockfish, rockfish, salmon grouper, salmon rock fish or scamp, is a species of marine ray-finned fish....



Mycteroperca microlepis

Mycteroperca microlepis

The gag grouper (*Mycteroperca microlepis*), also known as velvet rockfish, the gag, or charcoal belly, is a species of marine ray-finned fish, a grouper from the subfamily Epinephelinae which is part of the family Serranidae, which also includes...



Mycteroperca rosacea

Mycteroperca rosacea

Mycteroperca rosacea, the leopard grouper or golden grouper, is a species of marine ray-finned fish, a grouper from the subfamily Epinephelinae which is part of the family Serranidae, which also includes the anthias and sea basses. It is found in...



Mycteroperca xenarcha

Mycteroperca xenarcha

The broomtail grouper or mangrove grouper (*Mycteroperca xenarcha*) is a species of ray-finned fish from the family Serranidae which is found in the eastern Pacific along the western coast of the Americas from California to Peru.



Myctophum nitidulum

Myctophum nitidulum

Myctophum nitidulum, common name pearly lanternfish, is a species of deep sea fish in the family Myctophidae, the "lanternfish".



Myctophum punctatum

Myctophum punctatum

Myctophum punctatum is a species of mesopelagic fish in the family Myctophidae. Its common name is spotted lanternfish, sometimes spelled spotted lanternfish. It is found in the Northern Atlantic and in the Mediterranean at depths down to 1000m....



Mylocheilus caurinus

Mylocheilus caurinus

Mylocheilus caurinus, the peamouth, peamouth chub, redmouth sucker or northwestern dace, is a species of freshwater ray-finned fish from the family Cyprinidae, the carps and minnows, that is found in western North America. It is the only species...



Mylopharodon conocephalus

Mylopharodon conocephalus

Mylopharodon conocephalus, known as the hardhead, is a freshwater ray-finned fish from the family Cyprinidae, the carps and minnows, which is endemic to California. It is the sole member of the monotypic genus Mylopharodon.



Myloplus schomburgkii

Myloplus schomburgkii

Myloplus schomburgkii, also known as the Disk tetra, Disk pacu, Black-ear pacu, Black-band myleus or Black-barred myleus is a species of serrasalmid with a black bar on its side. This species is found in the middle and lower Amazon River basin,...



Myoxocephalus aeneus

Myoxocephalus aeneus

Myoxocephalus aeneus, commonly known as the grubby, or little sculpin, is a species of marine ray-finned fish belonging to the family Cottidae, the typical sculpins. This species is found in the northwestern Atlantic Ocean.



Myoxocephalus scorpius

Myoxocephalus scorpius

Myoxocephalus scorpius, typically known as the shorthorn sculpin or bull-rout, is a species of fish in the family Cottidae. It is a demersal species of the Northern Atlantic and adjacent subarctic and Arctic seas. The species has many English...



Myripristis jacobus

Myripristis jacobus

Myripristis jacobus, the blackbar soldierfish, is a soldierfish from the Western Atlantic. It occasionally makes its way into the aquarium trade. It grows to a size of 25 cm in length. Myripristis jacobus can be located in a marine environment...



Myripristis murdjan

Myripristis murdjan

Myripristis murdjan is a species of soldierfish found in the Indo-Pacific.



Mystus ankutta

Mystus ankutta

Mystus ankutta, also known as the Sri Lanka dwarf catfish or yellow dwarf catfish, is a species of catfish of the family Bagridae that is endemic to Sri Lanka. In the wild it is found in freshwater bodies from Kelani river to Nilwala river in Sri...



Mystus cavasius

Mystus cavasius

Mystus cavasius, the Gangetic mystus, is a species of catfish of the family Bagridae. In the wild it is found in Indian Subcontinent countries such as, India, Bangladesh, Pakistan, Sri Lanka, Nepal, and Myanmar. Reports of this species from the...



Mystus gulio

Mystus gulio

Mystus gulio, the Long Whiskers Catfish, is a species of catfish of the family Bagridae. The generic name is probably derived from the Latin "mystax", meaning "moustache", due to the long barbels. It is found in India, Sri Lanka, Pakistan, Nepal...



Mystus leucophasis

Mystus leucophasis

Mystus leucophasis is an Asian species of upside-down catfish belonging to the family Bagridae. It is known for its unusual behaviour, as it primarily swims in an inverted position and never upright. Mystus leucophasis originate in the Sittang and...



Mystus nigriceps

Mystus nigriceps

Mystus nigriceps, the two-spot catfish, is a species of catfish of the family Bagridae. The species was available in aquarium circles without a scientific name until a connection was made to a description of Mystus micracanthus by the...



Mystus vittatus

Mystus vittatus

Mystus vittatus, the striped dwarf catfish, is a species of catfish of the family Bagridae. It is found in brackish water systems with marginal vegetation in lakes and swamps with a mud substrate of Asian countries Pakistan, India, Sri Lanka,...



Nandus nandus

Nandus nandus

Nandus nandus, commonly known as the Gangetic leaffish, is a species of Asian leaffish native to South Asia and Indochina. They are common in slow-moving or stagnant bodies of water, including ponds, lakes, ditches, and flooded fields. Other...



Nannostomus anduzei

Nannostomus anduzei

Nannostomus anduzei (common name: Anduzi's pencilfish) is a freshwater species of fish belonging to the genus Nannostomus in the characin family Lebiasinidae. It is native to Venezuela and northern Brazil, particularly the upper Orinoco and Rio...



Nannostomus beckfordi

Nannostomus beckfordi

Nannostomus beckfordi, (from the Greek: nanos = small, and the Latin stomus = relating to the mouth; beckfordi = in honor of British naturalist, F.J.B. Beckford), commonly known as the golden pencil fish or Beckford's pencil fish, is a freshwater...



Nannostomus bifasciatus

Nannostomus bifasciatus

Nannostomus bifasciatus, (from the Greek: nanos = small, and the Latin stomus = relating to the mouth; from the Latin: bifasciatus = two bands), is a freshwater species of fish belonging to the characin family Lebiasinidae. It is commonly known...



Nannostomus eques

Nannostomus eques

Nannostomus eques, (from the Greek: nanos = small, and the Latin stomus = relating to the mouth; from the Latin: eques = horseman), commonly known as the diptail pencilfish or brown pencilfish, is a freshwater species of fish belonging to the...



Nannostomus espei

Nannostomus espei

Nannostomus espei, (from the Greek: nanos = small, and the Latin stomus = relating to the mouth; espei = in honor of tropical fish importer/exporter, Heinrich Espe), commonly known as Espe's pencilfish or barred pencilfish, is a freshwater...



Nannostomus marginatus

Nannostomus marginatus

Nannostomus marginatus, (from the Greek: nanos = small, and the Latin stomus = relating to the mouth; from the Latin: marginatus = marginated), commonly known as the dwarf pencilfish, is a freshwater species of fish belonging to the genus...



Nannostomus mortenthaleri

Nannostomus mortenthaleri

Nannostomus mortenthaleri, commonly known as the coral red pencilfish, is a freshwater species of fish belonging to the characin family Lebiasinidae. It is one of the most colourful of the genus, being suffused with a bright coral red colouration...



Nannostomus trifasciatus

Nannostomus trifasciatus

Nannostomus trifasciatus, (from the Greek: nanos = small, and the Latin stomus = relating to the mouth; from the Latin: trifasciatus = three bands), commonly known as the three-lined or three-stripe pencilfish, is a freshwater species of fish...



Nannostomus unifasciatus

Nannostomus unifasciatus

Nannostomus unifasciatus, (from the Greek: nanos = small, and the Latin stomus = relating to the mouth; from the Latin: unifasciatus = one band), commonly known as the one-lined pencilfish, is a freshwater species of fish belonging to the genus...



Narcine brasiliensis

Narcine brasiliensis

Narcine brasiliensis, the Brazilian electric ray or lesser numbfish, is a species of electric ray in the family Narcinidae. It inhabits coastal waters of the Southwest Atlantic from Southern Brazil through Uruguay to northern Argentina.



Narcine entemedor

Narcine entemedor

Narcine entemedor, the giant electric ray or Cortez electric ray, is a species of numbfish, family Narcinidae, native to the eastern Pacific Ocean from the Gulf of California to Panama. It is found in shallow water on sandy bottoms and sometimes...



Narrow-barred Spanish mackerel

Scomberomorus commerson

The narrow-barred Spanish mackerel (Scomberomorus commerson) is a mackerel of the family Scombridae found in a wide-ranging area in Southeast Asia, but as far west as the east coast of Africa and from the Middle East and along the northern...



Narrow-lined puffer

Arothron manilensis

The narrow-lined pufferfish (Arothron manilensis) is a demersal marine fish belonging to the family Tetraodontidae.



Narrowmouthed catshark

Schroederichthys bivius

The narrowmouthed catshark (Schroederichthys bivius) is a catshark of the family Scyliorhinidae, found from central Chile around the Straits of Magellan, to Argentina between latitudes 23° S and 56° S, at depths down to about 180 m (600 ft) in...



Narrownose chimaera

Harriotta raleighana

The narrownose chimaera (Harriotta raleighana) is a longnose chimaera of the family Rhinochimaeridae, the longnose chimaeras, consisting of eight species belonging three genera. This species is found in temperate seas worldwide, at depths between...



Naso brevirostris

Naso brevirostris

Naso brevirostris, also known as the short-nosed unicornfish, spotted unicornfish, brown unicornfish, lined unicornfish, longnose surgeonfish, palefin unicornfish, paletail unicornfish, shorthorned unicornfish or shortsnouted unicornfish, is a...



Naso elegans

Naso elegans

Naso elegans, the elegant unicornfish, the blonde naso tang, Indian orange-spine unicorn, lipstick surgeonfish, lipstick tang, orangespine unicornfish or smoothheaded unicornfish, is a species of marine ray-finned fish belonging to the family...



Naso lituratus

Naso lituratus

Naso lituratus, the clown unicornfish, orangespined unicornfish, black-finned unicornfish, Pacific orangespined unicornfish, blackfinned unicornfish or stripefaced unicornfish, is a species of marine ray-finned fish belonging to the family...



Naso vlamingii

Naso vlamingii

Naso vlamingii, the bignose unicornfish, scibbled unicornfish, Vlaming's unicornfish, and zebra unicornfish, is a species of marine ray-finned fish belonging to the family Acanthuridae, the surgeonfishes, unicornfishes and tangs. This species is...



Nassau grouper

Epinephelus striatus

The Nassau grouper (Epinephelus striatus) is one of the large number of perciform fishes in the family Serranidae commonly referred to as groupers. It is the most important of the groupers for commercial fishery in the West Indies, but has been...



Natal shyshark

Haploblepharus kistnasamyi

The Natal shyshark, eastern shyshark or happy chappie (Haploblepharus kistnasamyi) is a species of catshark, belonging to the family Scyliorhinidae. It was once regarded as the "Natal" form of the puffadder shyshark (H. edwardsii). This shark is...



Navaga

Eleginus nawaga

The navaga (*Eleginus nawaga*) is a relatively small species of fish in the cod family Gadidae. It inhabits the European arctic and subarctic waters of the Barents, White, and Kara seas, from the Kola Bay to the Ob River estuary. Navaga fish usually...



Necklace carpetshark

Parascyllium variolatum

Sharks portalThe necklace carpetshark (*Parascyllium variolatum*), also known as the varied carpetshark, is a carpetshark of the family Parascylliidae endemic to the waters off Australia's southern coast between latitudes 37°S and 41°S. It is...



Nelma

Stenodus nelma

Stenodus nelma, known alternatively as the nelma, sheefish, siifish, inconnu or connie, is a commercial species of freshwater whitefish in the family Salmonidae. It is widespread in the Arctic rivers from the Kola Peninsula (White Sea basin)...



Nemadactylus macropterus

Nemadactylus macropterus

Nemadactylus macropterus, the tarakihi, jackass morwong or deep sea perch, is a species of marine ray-finned fish, traditionally regarded as belonging to the family Cheilodactylidae, the members of which are commonly known as morwongs. It is...



Nemateleotris helfrichi

Nemateleotris helfrichi

Nemateleotris helfrichi, Helfrich's Dartfish, is a species of dartfish native to the Pacific Ocean.



Nemateleotris magnifica

Nemateleotris magnifica

Nemateleotris magnifica, known by a variety of common names including fire goby, magnificent fire fish, fire dartfish, or red fire goby, is a species of dartfish native to coral reefs of the Indian and Pacific oceans.



Nematobrycon palmeri

Nematobrycon palmeri

Nematobrycon palmeri, commonly known as the emperor tetra, is a species of characid fish found in the Atrato and San Juan river basins in western Colombia. It was first imported in the aquarium trade to the United States in 1960 and has since...



Nemipterus japonicus

Nemipterus japonicus

Nemipterus japonicus, the Japanese threadfin bream, is a species of marine ray-finned fish belonging to the family Nemipteridae, the threadfin and whiptail breams. This species is found in the Indo Pacific region and is an important food fish.



Nemipterus randalli

Nemipterus randalli

Nemipterus randalli, or Randall's threadfin bream, is a species of ray-finned fish from the family Nemipteridae, the threadfin breams, which is native to the western Indian Ocean and the Red Sea, it has invaded the eastern Mediterranean by...



Nemipterus virgatus

Nemipterus virgatus

Nemipterus virgatus, the golden threadfin bream or yellowlipped threadfin bream, is a species of marine ray-finned fish belonging to the family Nemipteridae, the threadfin breams. This species is found in the Western Pacific Ocean.



Neoarius berneyi

Neoarius berneyi

Neoarius berneyi, the highfin catfish, Berney's catfish, Berney's shark catfish, or the lesser salmon catfish, is a freshwater sea catfish that is commonly kept in aquariums. The origin of the name Neoarius berneyi is Greek, with the genus name...



Neoglyphidodon melas

Neoglyphidodon melas

Neoglyphidodon melas, also known as the bowtie damselfish, black damsel, bluefin or royal damsel, is a species of damselfish found in the Indo-West Pacific. It often makes its way into the aquarium trade. It grows to a size of 18 cm (7.1 in) in...



Neolamprologus brichardi

Neolamprologus brichardi

Neolamprologus brichardi is a species of cichlid endemic to the alkaline waters of Lake Tanganyika in East Africa. It is a popular aquarium fish kept in the fishkeeping hobby, where it is known under a variety of common names including Princess...



Neolamprologus caudopunctatus

Neolamprologus caudopunctatus

Neolamprologus caudopunctatus is a species of cichlid endemic to Lake Tanganyika, where it occurs along the Zambian shores. It has a silvery-beige colored body, accented only by a distinctive, goldish-orange dorsal fin and blue eyes. Its tail fin...



Neolamprologus leleupi

Neolamprologus leleupi

Neolamprologus leleupi (lemon cichlid) is a species of cichlid endemic to Lake Tanganyika where it occurs throughout the lake. It is a recess-dweller, inhabiting cracks and crevices. It feeds on invertebrates living in the rich biocover of the...



Neolamprologus multifasciatus

Neolamprologus multifasciatus

Neolamprologus multifasciatus is one of the small shell-dwelling cichlids endemic to Lake Tanganyika. The male reaches 5 cm (2 in) in length, and the female only 2.5 cm (1 in) in the aquarium. In the wild, they reach only 3 centimetres (1.2 in)...



Neolamprologus pulcher

Neolamprologus pulcher

Neolamprologus pulcher is a species of cichlid endemic to Lake Tanganyika where it prefers locations with plenty of sedimentation. The common names for N. pulcher include daffodil cichlid, fairy cichlid, princess of Zambia and lyretail cichlid....



Neolamprologus similis

Neolamprologus similis

Neolamprologus similis is a shell-dwelling cichlid endemic to Lake Tanganyika in Africa, where it is only known along the shores of the Democratic Republic of the Congo and Tanzania. N. similis is copper-coloured with vertical white stripes...



Neon tetra

Paracheirodon innesi

The neon tetra (Paracheirodon innesi) is a Freshwater fish of the characin family (family Characidae) of order Characiformes. The type species of its genus, it is native to blackwater and clearwater streams in the Amazon basin of South America....



Neopomacentrus cyanomos

Neopomacentrus cyanomos

Neopomacentrus cyanomos, commonly known as the regal demoiselle, is a fish native to the Indo-Pacific region from the Persian Gulf, the Red Sea and the coastal waters of east Africa east to the Philippines, north to southern Japan, south to...



Neosalanx tangkahkeii

Neosalanx tangkahkeii

Neosalanx tangkahkeii, the Chinese icefish or short-snout icefish, is a species of icefish endemic to fresh and brackish waters in China. Despite its common name it is not the only icefish in China; the majority of the species in this family are...



Nervous shark

Carcharhinus cautus

The nervous shark (Carcharhinus cautus) is a species of requiem shark, and part of the family Carcharhinidae, so named because of its timid behavior in regard to humans. It is common in shallow, coastal waters off northern Australia, Papua New...



New Holland seahorse

Hippocampus whitei

Hippocampus whitei, commonly known as White's seahorse, New Holland seahorse, or Sydney seahorse, is a species of marine fish of the family Syngnathidae. It is thought to be endemic to the Southwest Pacific, from Sydney, New South Wales and...



New Zealand eagle ray

Myliobatis tenuicaudatus

The New Zealand eagle ray or Australian eagle ray (Myliobatis tenuicaudatus) is an eagle ray of the family Myliobatidae, found in bays, estuaries, and near rocky reefs around New Zealand and southern Australia on the inner continental shelf. It...



New Zealand longfin eel

Anguilla dieffenbachii

The New Zealand longfin eel (Anguilla dieffenbachii), also known as ōrea, is a species of freshwater eel that is endemic to New Zealand. It is the largest freshwater eel in New Zealand and the only endemic species – the other eels found in New...



New Zealand sand flounder

Rhombosolea plebeia

The New Zealand sand flounder (*Rhombosolea plebeia*) is a righteye flounder of the genus *Rhombosolea*, found around New Zealand in shallow waters down to depths of 100 m.



New Zealand smelt

Retropinna retropinna

The New Zealand smelt (*Retropinna retropinna*), also known as the New Zealand common smelt, New Zealand cucumber fish, or silveries is a smelt of the family Retropinnidae, found only in New Zealand at shallow depths in estuaries and rivers. Their...



New Zealand sprat

Sprattus muelleri

The New Zealand sprat (*Sprattus muelleri*) is a herring-like, marine fish in the family Clupeidae found in the subtropical southwest Pacific Ocean endemic to New Zealand. It belongs to a genus *Sprattus* of small oily fish, usually known by their...



New Zealand turbot

Colistium nudipinnis

The New Zealand turbot, Colistium nudipinnis, is a righteye flounder of the subfamily Rhombosoleinae in the family Pleuronectidae, found around New Zealand in shallow enclosed waters.



Niger stingray

Fontitrygon garouaensis

The Niger stingray or smooth freshwater stingray (Fontitrygon garouaensis) is a species of stingray in the family Dasyatidae, native to rivers in Nigeria and Cameroon. Attaining a width of 40 cm (16 in), this species can be distinguished by its...



Night shark

Carcharhinus signatus

The night shark (Carcharhinus signatus) is a species of requiem shark, in the family Carcharhinidae, found in the temperate and tropical waters of the Atlantic Ocean. An inhabitant of the outer continental shelf and upper continental slope, this...



Nightfish

Bostockia porosa

The nightfish (Bostockia porosa) is a species of freshwater ray-finned fish, a temperate perch from the family Percichthyidae which is endemic to southwestern Australia.



Nile carp

Labeo niloticus

The Nile carp (Labeo niloticus) is a fish species in the genus Labeo. It feeds primary on plankton, and is distributed along the entire Nile valley. It is generally believed to be the fish that swallowed the phallus of the Egyptian god Osiris in...



Nile perch

Lates niloticus

The Nile perch (Lates niloticus), also known as the African snook, Goliath perch, African barramundi, Goliath barramundi, Giant lates or the Victoria perch, is a species of freshwater fish in family Latidae of order Perciformes. It is widespread...



Nile tilapia

Oreochromis niloticus

The Nile tilapia (Oreochromis niloticus) is a species of tilapia, a cichlid fish native to parts of Africa and the Levant, particularly Israel and Lebanon. Numerous introduced populations exist outside its natural range. It is also commercially...



Nimbochromis fuscotaeniatus

Nimbochromis fuscotaeniatus

Nimbochromis fuscotaeniatus (spothead hap, fuscotaeniatus hap) is a species of cichlid endemic to Lake Malawi and Lake Malombe. Males of this species can reach a length of 25 centimetres (9.8 in) TL while the females grow to 20 centimetres (7.9...



Nimbochromis livingstonii

Nimbochromis livingstonii

Nimbochromis livingstonii, Livingston's cichlid or (locally) kalingono, is a freshwater mouthbrooding cichlid native to Lake Malawi, an African Rift Lake. It is also found in the upper Shire River and Lake Malombe. They are found in inshore areas...



Nimbochromis polystigma

Nimbochromis polystigma

Nimbochromis polystigma is a species of cichlid endemic to Lake Malawi, in the Great Rift Valley system of Africa. This species prefers areas with rock/sand substrate or areas with plentiful vegetation. It can grow to a length of 23 centimetres...



Nimbochromis venustus

Nimbochromis venustus

Nimbochromis venustus, commonly called venustus hap or giraffe hap, is a Haplochromine cichlid endemic to Lake Malawi in Africa. It prefers the deeper regions of the lake (6 to 23 metres (20 to 75 ft)) where it hunts smaller juvenile cichlids...



Ninespine stickleback

Pungitius pungitius

The ninespine stickleback (Pungitius pungitius), also called the ten-spined stickleback, is a freshwater species of fish in the family Gasterosteidae that inhabits temperate waters. It is widely but locally distributed throughout Eurasia and...



Ningu

Labeo victorinus

The ningu (Labeo victorinus) is a species of ray-finned fish in the family Cyprinidae. It is found in the Lake Victoria basin in Burundi, Kenya, Tanzania, and Uganda. Its natural habitats are rivers, swamps, freshwater lakes, freshwater marshes,...



Ninja lanternshark

Etmopterus benchleyi

The ninja lanternshark (Etmopterus benchleyi) is a lanternshark of the family Etmopteridae found in the eastern Pacific Ocean from Nicaragua, south to Panama and Costa Rica. The depth range of collections is from 836 to 1443 m along the...



Nkhomo-benga peacock

Aulonocara baenschi

The nkomo-benga peacock (Aulonocara baenschi), also known as the new yellow regal peacock, is a species of haplochromine cichlid which is endemic to Lake Malawi. This species is threatened by capture for the aquarium trade.



North Pacific hake

Merluccius productus

The North Pacific hake, Pacific hake, Pacific whiting, or jack salmon (Merluccius productus) is a ray-finned fish in the genus Merluccius, found in the northeast Pacific Ocean from northern Vancouver Island to the northern part of the Gulf of...



Northern cavefish

Amblyopsis spelaea

The northern cavefish or northern blindfish (Amblyopsis spelaea) is found in caves through Kentucky and southern Indiana. The International Union for Conservation of Nature lists the species as near threatened. The life cycle of northern cavefish...



Northern clingfish

Gobiesox maeandricus

The northern clingfish (Gobiesox maeandricus) is a species of saltwater fish. It is a member of the family Gobiesocidae of order Gobiesociformes. It is native to the Pacific coast of North America from Revillagigedo Island and Baja California...



Northern hogsucker

Hypentelium nigricans

The northern hogsucker (*Hypentelium nigricans*) is a freshwater ray-finned fish belonging to the family Catostomidae, the suckers. It is native to the United States and Canada where it is found in streams and rivers. It prefers clear, fast-flowing...



Northern madtom

Noturus stigmosus

The northern madtom (Noturus stigmosus) is a freshwater fish.



Northern pike

Esox lucius

The northern pike (Esox lucius) is a species of carnivorous fish of the genus Esox (pikes). They are commonly found in moderately salty and fresh waters of the Northern Hemisphere (i.e. holarctic in distribution). They are known simply as a pike...



Northern pikeminnow

Ptychocheilus oregonensis

The Northern pikeminnow, Columbia River dace or colloquially Squawfish (Ptychocheilus oregonensis) is a large member of the minnow family, Leuciscidae. This predatory freshwater fish is native to northwestern North America, ranging from the Nass...



Northern pipefish

Syngnathus fuscus

The northern pipefish (Syngnathus fuscus) is a northwest Atlantic species of fish belonging to the family Syngnathidae.



Northern puffer

Sphoeroides maculatus

The northern puffer, Sphoeroides spheroides, is a species in the family Tetraodontidae, or pufferfishes, found along the Atlantic coast of North America. Unlike many other pufferfish species, the flesh of the northern puffer is not poisonous....



Northern red snapper

Lutjanus campechanus

The northern red snapper (Lutjanus campechanus) is a species of marine ray-finned fish, a snapper belonging to the family Lutjanidae. It is native to the western Atlantic Ocean, the Caribbean Sea, and the Gulf of Mexico, where it inhabits...



Northern redbelly dace

Chrosomus eos

The northern redbelly dace (Chrosomus eos) is a fresh water cyprinid fish, generally found in lakes and small streams in the northeastern United States and eastern Canada. Ranging from 1–3 inches, it is one of forty-four species from the minnow...



Northern river shark

Glyphis garricki

The northern river shark or New Guinea river shark (Glyphis garricki) is a species of requiem shark, in the family Carcharhinidae, found in scattered tidal rivers and associated coastal waters in northern Australia and in Papua New Guinea. This...



Northern rock sole

Lepidopsetta polyxystra

The northern rock sole (Lepidopsetta polyxystra) is a flatfish of the family Pleuronectidae. It is a demersal fish that lives on sand, mud and gravel bottoms at depths of up to 700 metres (2,300 ft), though it is most commonly found between 19...



Northern sennet

Sphyraena borealis

The northern sennet, Sphyraena borealis, is an ocean-going species of fish in the barracuda family, Sphyraenidae. It was described by the American zoologist James Ellsworth De Kay in 1842. De Kay's description was part of several volumes he...



Northern snakehead

Channa argus

The northern snakehead (Channa argus) is a species of snakehead fish native to temperate East Asia, in China, Russia, North Korea, and South Korea. Their natural range goes from the Amur River watershed in Siberia and Manchuria down to Hainan. It...



Northern studfish

Fundulus catenatus

The northern studfish (Fundulus catenatus) is the largest of the killifish and is native to the southcentral United States.



Northern sunfish

Lepomis peltastes

The northern sunfish (Lepomis peltastes) is a species of North American freshwater fish in the sunfish family (Centrarchidae) of order Perciformes. Lepomis peltastes was traditionally recognized as a subspecies of longear sunfish (Lepomis...



Northern tidewater goby

Eucyclogobius newberryi

Eucyclogobius newberryi, the northern tidewater goby, is a species of goby native to lagoons of streams, marshes, and creeks along the coast of California, United States. The northern tidewater goby is one of six native goby species of...



Northern whiting

Sillago sihama

The northern whiting (Sillago sihama), also known as the silver whiting and sand smelt, is a marine fish, the most widespread and abundant member of the smelt-whiting family Sillaginidae. The northern whiting was the first species of sillaginid...



Northern wolffish

Anarhichas denticulatus

The northern wolffish (Anarhichas denticulatus), also known as the blue sea cat or jelly cat, is a species of marine ray-finned fish belonging to the family Anarhichadidae, the wolf fishes. This species is found in the North Atlantic Ocean and...



Nothobranchius furzeri

Nothobranchius furzeri

Nothobranchius furzeri, the turquoise killifish, is a species of killifish from the family Nothobranchiidae native to Africa where it is only known from Zimbabwe and Mozambique. This annual killifish inhabits ephemeral pools in semi-arid areas...



Nothobranchius rachovii

Nothobranchius rachovii

Nothobranchius rachovii, the bluefin notho, is a species of freshwater annual killifish from Mozambique. It can grow up to 6 cm (2.4"). It is popular among killifish enthusiasts, who raise them from eggs in aquaria.



Notolabrus fucicola

Notolabrus fucicola

Notolabrus fucicola, the banded parrotfish, blue wrasse, kelpie, New Zealand banded wrasse, purple parrotfish, saddled wrasse, Southern purple wrasse, Southern wrasse, winter bream or yellow-saddled wrasse, is a species of wrasse native to the...



Notoliparis kermadecensis

Notoliparis kermadecensis

Notoliparis kermadecensis (from Greek: noton, back, and liparos, fat) is a species of snailfish (Liparidae) that lives in the deep sea. Endemic to the Kermadec Trench in the Southwest Pacific, it is hadobenthic with a depth range between 6,474...



Notothenia angustata

Notothenia angustata

Notothenia angustata, the Maori chief or black cod, is a species of marine ray-finned fish, belonging to the family Nototheniidae, the notothen or cod icefishes. It is native to the Southern Ocean



Notothenia microlepidota

Notothenia microlepidota

Notothenia microlepidota, the black cod or small-scaled cod, is a species of marine ray-finned fish, belonging to the family Nototheniidae, the notothen or cod icefishes. It is native to the Pacific waters around New Zealand and Macquarie...



Noturus flavus

Noturus flavus

Noturus flavus, the stonecat, is a North American freshwater catfish of the family Ictaluridae. The common name is due to its habit of hiding near or under stones in fast-moving water.



Noturus insignis

Noturus insignis

Noturus insignis (margined madtom) is a small species of North American catfish belonging to the family Ictaluridae.



Novaculichthys

Novaculichthys taeniourus

Novaculichthys taeniourus, also known as the rockmover wrasse, carpet wrasse, dragon wrasse, bar-cheeked wrasse, olive-scribbled wrasse or reindeer wrasse, is a species of wrasse mainly found in coral reefs and lagoons in the Indo-Pacific region....



Nurse shark

Ginglymostoma cirratum

The nurse shark (*Ginglymostoma cirratum*) is an elasmobranch fish in the family Ginglymostomatidae. The conservation status of the nurse shark is globally assessed as Vulnerable in the IUCN List of Threatened Species. They are considered to be a...



Nursehound

Scyliorhinus stellaris

The nursehound (*Scyliorhinus stellaris*), also known as the large-spotted dogfish, greater spotted dogfish or bull huss, is a species of catshark, belonging to the family Scyliorhinidae, found in the northeastern Atlantic Ocean. It is generally...



Obscure snakehead

Parachanna obscura

The obscure snakehead, *Parachanna obscura*, is a medium-sized carnivorous fish that has an elongated shape tapered on both ends and is covered in medium circular scales (cycloid). The head, resembling a snake, is long and depressed anteriorly and...



Obtuse barracuda

Sphyraena obtusata

The obtuse barracuda, *Sphyraena obtusata*, a barracuda of the family Sphyraenidae, is found in tropical oceans around the world. Its length is up to 55 cm.



Ocean pout

Zoarces americanus

The ocean pout (*Zoarces americanus*) is an eelpout in the family Zoarcidae. It is found in the Northwest Atlantic Ocean, off the coast of New England and eastern Canada. The fish has antifreeze proteins in its blood, giving it the ability to...



Ocean sunfish

Mola mola

The ocean sunfish or common mola (Mola mola) is one of the largest bony fish in the world. It was misidentified as the heaviest bony fish, which was actually a different species, Mola alexandrini. Adults typically weigh between 247 and 1,000 kg...



Ocean surgeon

Acanthurus bahianus

Ocean surgeon may refer to: Acanthurus bahianus, a fish found off the coast of Brazil. Acanthurus tractus, a fish found in the western Atlantic Ocean, Florida, the Bahamas, the Caribbean Sea and the Gulf of Mexico.



Ocean whitefish

Caulolatilus princeps

The ocean whitefish (Caulolatilus princeps), also known as the ocean tilefish, is a species of marine ray-finned fish, a tilefish belonging to the family Malacanthidae. It is native to the eastern Pacific Ocean.



Oceanic puffer

Lagocephalus lagocephalus

The oceanic puffer, sci-name: Lagocephalus lagocephalus (meaning "rabbit head"), is a pufferfish of the family Tetraodontidae, found in all tropical and subtropical oceans, at depths of between 10 and 475 m. Though indigenous to the Pacific,...



Oceanic whitetip shark

Carcharhinus longimanus

The oceanic whitetip shark (Carcharhinus longimanus) is a large pelagic requiem shark inhabiting tropical and warm temperate seas. It has a stocky body with long, white-tipped, rounded fins. The species is typically solitary, though they may...



Ocellaris clownfish

Amphiprion ocellaris

The ocellaris clownfish (Amphiprion ocellaris), also known as the false percula clownfish or common clownfish, is a marine fish belonging to the family Pomacentridae, which includes clownfishes and damselfishes. Amphiprion ocellaris are found in...



Ocellate river stingray

Potamotrygon motoro

The ocellate river stingray (Potamotrygon motoro), also known as the peacock-eye stingray or black river stingray, is a species of freshwater stingray in the family Potamotrygonidae. It was the first species to be described in the family and is...



Ocellate spot skate

Okamejei kenojei

The ocellate spot skate (Okamejei kenojei), also known as the spiny rasp skate or swarthy skate, is a species of skate in the family Rajidae and is commonly found in the north-western Pacific Ocean. O. kenojei is a bottom-feeding carnivore that...



Ocellated dragonet

Neosynchiropus ocellatus

The ocellated dragonet or scooter dragonet (*Neosynchiropus ocellatus*) is a species of tropical marine fish in the family Callionymidae. It is native to the southwest Pacific Ocean from southern Japan to the Marquesan Islands.



Ocellated frogfish

Fowlerichthys ocellatus

The ocellated frogfish, Fowlerichthys ocellatus, is a frogfish of the family Antennariidae, found in the western Atlantic including the eastern portion of the Gulf of Mexico, and the Caribbean Sea, to depths of 150 m. Its grows to a length of 38...



Odessa barb

Pethia padamya

The Odessa barb (Pethia padamya) is a species of cyprinid fish known from Central Myanmar, where it is known to occur in an artificial pond above the Anisakan Falls and also from the lower Chindwin River. For many years it has been known to the...



Odontamblyopus lacepedii

Odontamblyopus lacepedii

Odontamblyopus lacepedii, also known as warasubo, is a species of eel goby found in muddy-bottomed coastal waters in China, Korea and Japan. This species excavates elaborate vertical burrows up to 90 centimetres (35 in) long in the sea bed. This...



Odontesthes regia

Odontesthes regia

Odontesthes regia is a fish belonging to the family Atherinopsidae, commonly referred to in English as Chilean silverside, Peruvian silverside or sea silverside, and in Spanish as pejerrey or pejerrey de mar. This is an epipelagic species...



Ogcocephalus corniger

Ogcocephalus corniger

Ogcocephalus corniger, the longnose batfish, is a species of batfish found at depths between 29 and 230 m (95 and 755 ft) in the Atlantic Ocean, ranging from North Carolina to the Gulf of Mexico and the Bahamas. Like other members of the family...



Ogcocephalus parvus

Ogcocephalus parvus

Ogcocephalus parvus, the roughback batfish, is a species of batfish found along the coast of the western Atlantic Ocean from North Carolina in the United States to Brazil. This species grows to a length of 10 centimetres (3.9 in) TL.



Ogcocephalus porrectus

Ogcocephalus porrectus

Ogcocephalus porrectus, the rosy-lipped batfish, is endemic to Cocos Island off the Pacific coast of Costa Rica. Though members of Ogcocephalidae occur in tropical, warm waters in both the Western Atlantic and Eastern Pacific. Rosy-lipped batfish...



Ogcocephalus radiatus

Ogcocephalus radiatus

Ogcocephalus radiatus or the polka-dot batfish is an arrow-shaped fish in the family Ogcocephalidae with an elongated thin tail. It is dorso-ventrally flattened with round pectoral fins that sit flat on the bottom of the sea floor. It uses its...



Ogcocephalus vespertilio

Ogcocephalus vespertilio

Ogcocephalus vespertilio, the Brazilian batfish or seadevil, is a species of batfish. Its distribution includes the western Atlantic, from the Antilles to Brazil, more specifically on the coast of Brazil, occurring from the Amazon River to La...



Ohrid trout

Salmo letnica

Ohrid trout or the Lake Ohrid brown trout (Salmo letnica) is an endemic species of trout in Lake Ohrid and in its tributaries and outlet, the Black Drin river, in North Macedonia and Albania. Locally, the fish is known as охридска пастрмка...



Oilfish

Ruvettus pretiosus

Oilfish (Ruvettus pretiosus) are a species of snake mackerel with a cosmopolitan distribution in tropical and temperate oceans. They can be found at depths from 100 to 800 metres (330 to 2,620 ft), but most often between 200 and 400 metres (660...



Okefenokee pygmy sunfish

Elassoma okefenokee

The Okefenokee pygmy sunfish, Elassoma okefenokee, is a species of pygmy sunfish found in southeastern United States, where it prefers waters with dense vegetation growth in the Altamaha drainage in southern Georgia south to Lake Okeechobee,...



Old wife

Enoplosus armatus

Enoplosus armatus, commonly referred to as the old wife (plural: old wives), is a species of perciform fish endemic to the temperate coastal waters of Australia. It is the only modern species in the family Enoplosidae. It has a deep and compressed...



Olive barb

Systemus sarana

The olive barb (Systemus sarana) (Malayalam: കുറ്റുവ/മുണ്ടത്തി) is a species of cyprinid fish native to Afghanistan, India, Nepal, Bangladesh, Pakistan, Bhutan, Sri Lanka, Thailand and Myanmar. This species can reach a length of 42 centimetres (17...



Olive flounder

Paralichthys olivaceus

The olive flounder (Paralichthys olivaceus), bastard halibut or Korean halibut, is a temperate marine species of large-tooth flounder native to the North-western Pacific Ocean. It is often referred to as the Korean flatfish (광어) or Japanese...



Olympic mudminnow

Novumbra hubbsi

The Olympic mudminnow (*Novumbra hubbsi*) is a fish native to the western lowlands of Washington: the Chehalis River basin, Deschutes River basin, and some Olympic Peninsula basins. It grows to 8 cm (about 3 in) in length, and is Washington's only...



Oman bullhead shark

Heterodontus omanensis

Sharks portalThe Oman bullhead shark, *Heterodontus omanensis*, is a bullhead shark of the family Heterodontidae found in the tropical western Indian Ocean around central Oman, from the surface to a depth of 72 m (236 ft) on the continental shelf....



Ompok bimaculatus

Ompok bimaculatus

Ompok bimaculatus, known as butter catfish, is a species of sheatfishes native to Asian countries such as Bangladesh, India, Pakistan, and Sri Lanka, but recently identified in Myanmar. It also indentified in Mekong Basin of Vietnam Known in...



Omul

Coregonus migratorius

The omul, *Coregonus migratorius*, also known as Baikal omul (Russian: байкальский омуль), is a whitefish species of the salmon family endemic to Lake Baikal in Siberia, Russia. It is considered a delicacy and is the object of one of the largest...



Oncorhynchus kawamurae

Oncorhynchus kawamurae

Black kokanee (*Oncorhynchus kawamurae*), or kunimasu (国鱒, クニマス, "national trout") in Japanese, is a species of landlocked Pacific trout in Japan that was thought to be extinct for 70 years before being rediscovered in 2010. The species was...



Oncorhynchus masou

Oncorhynchus masou

The masu salmon (*Oncorhynchus masou*), also known as masu (Japanese: マス, lit. 'trout') or cherry trout (桜鱒 サクラマス, sakura masu) in Japan, is a species of salmonid belonging to the genus *Oncorhynchus*, found in the North Pacific along Northeast/East...



Oncorhynchus masou formosanus

Oncorhynchus masou formosanus

Oncorhynchus masou formosanus, commonly known as the Formosan salmon, Taiwanese trout (Japanese: タイワンマス, romanized: taiwan masu), Tsugitaka trout (次高鱒, tsugitaka masu), Lishan trout (Chinese: 梨山鱒, after its native Lishan area in Heping District,...



Oncorhynchus masou macrostomus

Oncorhynchus masou macrostomus

The amago or the red-spotted masu salmon (*Oncorhynchus masou macrostomus*) is a salmonid fish endemic to western Japan, and a subspecies of the more widespread Northwest Pacific masu salmon or cherry salmon (*Oncorhynchus masou*). It is...



Ophidion barbatum

Ophidion barbatum

Snake blenny (*Ophidion barbatum*) is a fish species in the family Ophidiidae. It is widespread in the eastern Atlantic from southern England to Senegal in West Africa, and the northern Mediterranean. It is a marine subtropical demersal fish, up to...



Ophioblennius atlanticus

Ophioblennius atlanticus

Ophioblennius atlanticus, also known as the redlip blenny and the horseface blenny, is a species of combtooth blenny, family Blenniidae, found primarily in the western central Atlantic ocean. Redlip blennies can be found in coral crests and...



Ophisternon bengalense

Ophisternon bengalense

Ophisternon bengalense the Bengal eel, Bengal mudeel or onegill eel, is a species of fish in the family Synbranchidae. It is endemic to freshwater and brackish water rivers and swamps in Oceania and South Asia. It is normally 100 cm in maximum...



Orange chromide

Pseudetroplus maculatus

The orange chromide (Pseudetroplus maculatus ; more commonly Etroplus maculatus) is a species of cichlid fish that is endemic to freshwater and brackish streams, lagoons and estuaries in southern India and Sri Lanka. It is also known as pallathi...



Orange clownfish

Amphiprion percula

The orange clownfish (Amphiprion percula) also known as percula clownfish and clown anemonefish, is widely known as a popular aquarium fish. Like other clownfishes (also known as anemonefishes), it often lives in association with sea anemones. A....



Orange River mudfish

Labeo capensis

Orange River mudfish (Labeo capensis) is a species of fish in genus Labeo. It inhabits the Orange River system of southern Africa.



Orange roughy

Hoplostethus atlanticus

The orange roughy (Hoplostethus atlanticus), also known as the red roughy, slimehead and deep sea perch, is a relatively large deep-sea fish belonging to the slimehead family (Trachichthyidae). The UK Marine Conservation Society has categorized...



Orange skunk clownfish

Amphiprion sandaracinos

Amphiprion sandaracinos, also known as the orange skunk clownfish or orange anemonefish, is a species of anemonefish that is distinguished by its broad white stripe along the dorsal ridge. Like all anemonefishes it forms a symbiotic mutualism...



Orange spotted filefish

Oxymonacanthus longirostris

The orange spotted filefish or harlequin filefish, Oxymonacanthus longirostris, is a filefish in the family Monacanthidae found on coral reefs in the Indo-Pacific Oceans. The orangespotted filefish is a different species and refers to...



Orange-dotted tuskfish

Choerodon anchorago

The orange-dotted tuskfish (Choerodon anchorago) is a species of wrasse native to the Indian Ocean from Sri Lanka eastward to French Polynesia in the western Pacific. Its range extends north to the Ryukyus and south to New Caledonia. It inhabits...



Orange-fin anemonefish

Amphiprion chrysopterus

The orange-fin anemonefish (*Amphiprion chrysopterus*) is a marine fish belonging to the family Pomacentridae, the clownfishes and damselfishes, found in the Western Pacific north of the Great Barrier Reef from the surface to 20 m, to include the...



Orange-lined triggerfish

Balistapus undulatus

The orange-lined triggerfish (*Balistapus undulatus*) is a demersal triggerfish. Although *Balistapus* is a monotypic genus, it is closely related to the genus *Balistoides*.



Orange-spotted grouper

Epinephelus coioides

The orange-spotted grouper (*Epinephelus coioides*), also known as the brown-spotted rockcod, estuary cod, estuary rockcod, goldspotted rockcod, greasy cod, North-west groper, orange spotted cod or blue-and-yellow grouper, is a species of marine...



Orange-spotted snakehead

Channa aurantimaculata

The orange-spotted snakehead (*Channa aurantimaculata*) is a species of snakehead fish. Its body is of brownish colour intermixed with vertical orange stripes. Males have taller dorsal fins with more intense coloration, and narrower heads. It is...



Orange-spotted spinefoot

Siganus guttatus

The orange-spotted spinefoot (*Siganus guttatus*), also known as the deepbody spinefoot, gold-saddle rabbitfish, golden rabbitfish, golden-spotted spinefoot, goldlined spinefoot or yellowblotch spinefoot, is a species of marine ray-finned fish, a...



Orange-spotted trevally

Carangoides bajad

The orange-spotted trevally, Carangoides bajad (also known as the gold-spotted trevally) is a species of inshore marine fish in the jack family, Carangidae. The species is fairly common in tropical to subtropical waters of the Indo-Pacific,...



Orangeback angelfish

Centropyge acanthops

The orangeback angelfish (Centropyge acanthops), also known as the flameback angelfish, African pygmy angelfish and in South Africa as the Jumping Bean or Bean, is a species of ray-finned fish, a marine angelfish belonging to the family...



Orangefin labeo

Labeo calbasu

The orangefin labeo (Labeo calbasu) is a fish of the carp family Cyprinidae, found commonly in rivers and freshwater lakes in and around South Asia and South-East Asia. Native to Bangladesh and India.



Orangespotted sunfish

Lepomis humilis

The orangespotted sunfish (Lepomis humilis) is a North American species of freshwater fish in the sunfish family (Centrarchidae) of order Perciformes. These fish are widely distributed across the middle and eastern United States, from the Rocky...



Orangethroat darter

Etheostoma spectabile

The orangethroat darter (Etheostoma spectabile) is a species of freshwater ray-finned fish, a darter from the subfamily Etheostomatinae, part of the family Percidae, which also contains the perches, ruffes and pikeperches. It is endemic to the...



Orectolobus hutchinsi

Orectolobus hutchinsi

Orectolobus hutchinsi, the western wobbegong, is a species of carpet shark in the family Orectolobidae. The western wobbegong shark is a moderate sized marine shark found off the coast of Western Australia. Its scientific name is Orectolobus...



Orectolobus reticulatus

Orectolobus reticulatus

Orectolobus reticulatus, the network wobbegong, is a recently described species of carpet shark found in relatively shallow waters off Kimberley and Darwin in north-western Australia. With a known maximum length of only 52.3 centimetres (20.6...



Oreochromis amphimelas

Oreochromis amphimelas

Oreochromis amphimelas is a species of tilapia cichlid endemic to north–central Tanzania, where it is found in Lake Manyara and a number of other saline lakes with closed basins. Maximum recorded size is 28 cm (11 in) in standard length.



Oreochromis andersonii

Oreochromis andersonii

Oreochromis andersonii, the three-spotted tilapia, threespot tilapia, or threespot bream, is a species of cichlid native to Africa, where it is found in rivers and swamps in the southern half of the continent. This species reaches a length of 61...



Oreochromis aureus

Oreochromis aureus

The blue tilapia (*Oreochromis aureus*) is a species of tilapia, a fish in the family Cichlidae. Native to Northern and Western Africa, and the Middle East, through introductions it is now also established elsewhere, including parts of the United...



Oreochromis esculentus

Oreochromis esculentus

Oreochromis esculentus, the Singida tilapia or Graham's tilapia, is a species of cichlid endemic to the Lake Victoria basin, including some of its satellite lakes such as Kyoga, in Tanzania, Uganda, and Kenya. Its common name refers to Lake...



Oreochromis jipe

Oreochromis jipe

Oreochromis jipe, the Jipe tilapia, is a critically endangered species of cichlid fish native to Kenya and Tanzania, where it is restricted to Lake Jipe and the Pangani River. The population in the Pangani River (including Ruva River) shows some...



Oreochromis leucostictus

Oreochromis leucostictus

Oreochromis leucostictus (the blue-spotted tilapia) is a species of cichlid native to Albertine Rift Valley lakes and associated rivers in DR Congo and Uganda. It has now been introduced widely elsewhere East Africa, and is believed to have...



Oreochromis lidole

Oreochromis lidole

Oreochromis lidole is a species of freshwater fish in the family Cichlidae. This tilapia is native to Malawi, Mozambique and Tanzania, where it is found in Lake Malawi, Lake Malombe, the Shire River and perhaps some crater lakes further north. It...



Oreochromis macrochir

Oreochromis macrochir

Oreochromis macrochir (longfin tilapia, greenhead tilapia, or greenhead bream) is a species of cichlid native to the Zambezi Basin, Lake Mweru, and Lake Bangweulu. It has been used extensively for stocking ponds and dams in other parts of...



Oreochromis tanganicae

Oreochromis tanganicae

Oreochromis tanganicae, the Tanganyika tilapia, is a species of cichlid endemic to Lake Tanganyika and the mouths of its larger affluent rivers. This species can reach a length of 42 centimetres (17 in) SL.



Oreochromis urolepis hornorum

Oreochromis urolepis hornorum

The Wami tilapia is a tilapiine cichlid that grows to over 20 cm in length and is considered a useful food fish in Tanzania and the island of Zanzibar, where it may have been introduced by man. It is tolerant of brackish water and grows well in...



Oreochromis variabilis

Oreochromis variabilis

Oreochromis variabilis, the Victoria tilapia, is a species of African cichlid native to Lake Victoria and its tributaries, Lake Kyoga, Lake Kwanja, and Lake Bisina (Salisbury), as well as being found in the Victoria Nile above Murchison Falls....



Oriental flying gurnard

Dactyloptena orientalis

Dactyloptena orientalis, known commonly as the Oriental flying gurnard or purple flying gurnard among other vernacular names, is a species of marine fish in the family Dactylopteridae. Their name is derived from the French word 'gurnard' meaning...



Ornate butterflyfish

Chaetodon ornatissimus

The ornate butterflyfish (Chaetodon ornatissimus), or clown butterflyfish is a species of marine ray-finned fish, a butterflyfish in the family Chaetodontidae. The ornate butterflyfish is a close relative of the mailed butterflyfish (C....



Ornate rainbowfish

Rhadinocentrus ornatus

The ornate rainbowfish (Rhadinocentrus ornatus) is a species of rainbowfish endemic to an area in eastern Australia, where it is native to coastal regions and sandy offshore islands in southern Queensland and northern New South Wales. It is the...



Ornate wobbegong

Orectolobus ornatus

The ornate wobbegong (*Orectolobus ornatus*) is a species of carpet shark that lives in Australia and possibly other countries in the Western Pacific Ocean. It is coloured golden brown, yellow-green and blueish-grey, and it grows to maximum 120...



Ornate wrasse

Thalassoma pavo

The ornate wrasse (*Thalassoma pavo*) is a species of wrasse native to the rocky coasts of the eastern Atlantic Ocean and the Mediterranean Sea. This species is of minor importance to local commercial fisheries, is also popular as a game fish, and...



Orthopristis chrysoptera

Orthopristis chrysoptera

Orthopristis chrysoptera, the pigfish, hogfish, piggy perch, redmouth grunt or sailor's choice, is a species of marine ray-finned fish, a grunt belonging to the family Haemulidae. It is found in the western Atlantic Ocean. This name derives from...



Oryzias dancena

Oryzias dancena

Oryzias dancena or the Indian ricefish are a freshwater–brackish fish species native to the India, Bangladesh, Sri Lanka, Myanmar and Thailand. Their maximum length is only 3.1 centimetres (1.2 in). They are normally found in brackish habitats...



Oryzias javanicus

Oryzias javanicus

Oryzias javanicus, the Javanese ricefish, is a small species of fish in the family Adrianichthyidae. It lives in Peninsular Thailand, Malaysia, Singapore, and Indonesia (Java, Sumatra, Borneo, Sulawesi, Bali and Lombok), where it can be seen in...



Oscar

Astronotus ocellatus

The oscar (Astronotus ocellatus) is a species of fish from the cichlid family known under a variety of common names, including tiger oscar, velvet cichlid, and marble cichlid. In tropical South America, where the species naturally resides, A....



Osteobrama belangeri

Osteobrama belangeri

Osteobrama belangeri (Meitei: Pengba) is a species of ray-finned fish in the genus Osteobrama. It was found in the Indian state of Manipur, but has been extirpated there and as of 2022 is found in the wild only in Myanmar. Farmed populations in...



Osteobrama cotio

Osteobrama cotio

Osteobrama cotio is a species of ray-finned fish in the genus Osteobrama. This species had three subspecies named O. cotio cotio, O. cotio cunma and O. cotio peninsularis but these are now considered separate species. This species is found in the...



Osteochilus longidorsalis

Osteochilus longidorsalis

Osteochilus longidorsalis is a species of fish in the family Cyprinidae. Its common names are hiffin carp and long finned barb. It is endemic to the Western Ghats (India) and only found in Chalakudy River and Periyar River in Kerala. Locally it is...



Osteochilus vittatus

Osteochilus vittatus

Osteochilus vittatus is a species of cyprinid fish from Southeast Asia. Its common name is bonylip barb, Hasselt's bony-lipped barb, hard-lipped barb, or silver sharkminnow. It grows to 32 cm (13 in) SL.



Ostracion solorensis

Ostracion solorensis

Ostracion solorensis is a species of boxfish native to the western Pacific Ocean. Its common name is reticulate boxfish. It grows to 12 centimeters in length. It is sometimes kept as an aquarium fish.



Otjikoto tilapia

Tilapia guinasana

The Otjikoto tilapia (*Tilapia guinasana*) is a critically endangered species of cichlid fish endemic to Namibia where it was originally only found in Lake Guinas. This very small sinkhole lake contains quite clear water that generally ranges...



Otocinclus arnoldi

Otocinclus arnoldi

Otocinclus arnoldi is a species of armoured catfish native to South America, where it occurs in the lower río Paraná drainage, in the lower and middle rio Uruguai and in the río de La Plata. This species reaches a maximum length of 4.8...



Otocinclus batmani

Otocinclus batmani

Otocinclus batmani is a species of armored catfish (Loricariidae) native to South America. It can grow up to 37 millimetres (1.5 in) (SL).It is named for the comic-book hero Batman, who had a bat shape for a symbol, and has the pigmented mark on...



Oval butterflyfish

Chaetodon lunulatus

The oval butterflyfish (Chaetodon lunulatus), red-finned butterflyfish or redfin butterflyfish, is a species of marine ray-finned fish, a butterflyfish belonging to family Chaetodontidae. It is found in the Pacific Ocean from Eastern Indonesia to...



Owens pupfish

Cyprinodon radiosus

The Owens pupfish (Cyprinodon radiosus) is a rare species of fish in the family Cyprinodontidae, the pupfish. It is endemic to California in the United States, where it is limited to the Owens Valley. It is a federally listed endangered species...



Oxydoras niger

Oxydoras niger

Oxydoras niger, the rip saw catfish or cuiu cuiu, is a species of thorny catfish native to the Amazon, Essequibo and São Francisco basins in Bolivia, Brazil, Colombia, Ecuador, Guyana, Peru and Venezuela. This species grows to a length of 100...



Oxyleotris marmorata

Oxyleotris marmorata

The marble goby (*Oxyeleotris marmorata*) is a widely distributed species of fish in the family Butidae native to fresh and brackish waters of the Mekong and Chao Praya basins, as well as rivers and other water bodies in Cambodia (where it is...



Oxyjulis

Oxyjulis californica

Oxyjulis californica is a species of wrasse native to the eastern Pacific Ocean along the coasts of California and Baja California. Its distribution extends from Salt Point in Sonoma County, California, to southern central Baja California, near...



Oxynoemacheilus ercisanus

Oxynoemacheilus ercisanus

Oxynoemacheilus ercisanus, the Van loach, is a species of stone loach endemic to the Lake Van basin in Eastern Turkey.



Oyster toadfish

Opsanus tau

The oyster toadfish, *Opsanus tau*, also known as the oyster toad, ugly toad, oyster cracker, oyster catcher, and bar dog, is a Northeast Atlantic species of fish of the family Batrachoididae. The maximum length of this toadfish is 43.2 cm (17 in),...



Ozark bass

Ambloplites constellatus

The Ozark bass (*Ambloplites constellatus*) is a species of freshwater fish in the sunfish family (Centrarchidae) of order Perciformes. It is native only to the White River, Sac River, James River, and Pomme de Terre River drainages of Missouri and...



Ozark cavefish

Amblyopsis rosae

The Ozark cavefish, Amblyopsis rosae, is a small subterranean freshwater fish endemic to the United States. It has been listed as a threatened species in the US since 1984; the IUCN currently lists the species as Near Threatened, though it was...



Pacific angelshark

Squatina californica

The Pacific angelshark (Squatina californica) is a species of angelshark, family Squatinidae, found in the eastern Pacific Ocean from Alaska to the Gulf of California, and from Ecuador to Chile, although those in the Gulf of California and...



Pacific barracuda

Sphyraena argentea

Sphyraena argentea (also known as the Pacific barracuda, California barracuda or the silver barracuda) is a predatory species of marine barracuda fish of the family Sphyraenidae. They are found in the northeast Pacific Ocean, from Cabo San Lucas,...



Pacific blue-eye

Pseudomugil signifer

The Pacific blue-eye (Pseudomugil signifer) is a species of fish in the subfamily Pseudomugilinae native to eastern Australia. Described by Austrian naturalist Rudolf Kner in 1866, it comprises two subspecies that have been regarded as separate...



Pacific bluefin tuna

Thunnus orientalis

The Pacific bluefin tuna (Thunnus orientalis) is a predatory species of tuna found widely in the northern Pacific Ocean, but it is migratory and also recorded as a visitor to the south Pacific. In the past it was often included in T. thynnus, the...



Pacific bonito

Sarda lineolata

Pacific bonito, Sarda lineolata, is a marine species of bonito that is a game fighter but not highly thought of as a food fish. While it has been considered to be a subspecies of Sarda chiliensis, recent treatments recognize it as a species, S....



Pacific cod

Gadus macrocephalus

The Pacific cod (Gadus macrocephalus) is a species of ray-finned fish in the family Gadidae. It is a bottom-dwelling fish found in the northern Pacific Ocean, mainly on the continental shelf and upper slopes, to depths of about 900 m (3,000 ft)....



Pacific crevalle jack

Caranx caninus

The Pacific crevalle jack (Caranx caninus) is a species of large marine fish classified in the jack family Carangidae. The species is distributed through the tropical waters of the eastern Pacific Ocean from California in the north to Peru in the...



Pacific electric ray

Tetronarce californica

Tetronarce californica also known as the Pacific electric ray is a species of electric ray in the family Torpedinidae, endemic to the coastal waters of the northeastern Pacific Ocean from Baja California to British Columbia. It generally inhabits...



Pacific footballfish

Himantolophus sagamius

The Pacific footballfish (Himantolophus sagamius) is a species found in the Pacific. It has a wide range, extending from the coasts of Honshu (Gulf of Sagami) and Hokkaido islands through the Kuril-Kamchatka trough, in the northwest Pacific, to...



Pacific goliath grouper

Epinephelus quinquefasciatus

The Pacific goliath grouper (Epinephelus quinquefasciatus), also known as the Pacific itajara grouper, is a species of marine ray-finned fish, a grouper from the subfamily Epinephelinae which is part of the family Serranidae, which also includes...



Pacific halibut

Hippoglossus stenolepis

Hippoglossus stenolepis, the Pacific halibut, is a species of righteye flounder. This very large species of flatfish is native to the North Pacific and is fished by commercial fisheries, sport fishers, and subsistence fishers.



Pacific herring

Clupea pallasii

The Pacific herring (Clupea pallasii) is a species of the herring family associated with the Pacific Ocean environment of North America and northeast Asia. It is a silvery fish with unspined fins and a deeply forked caudal fin. The distribution...



Pacific jack mackerel

Trachurus symmetricus

The Pacific jack mackerel (Trachurus symmetricus), also known as the Californian jack mackerel or simply jack mackerel, is an abundant species of pelagic marine fish in the jack family, Carangidae. It is distributed along the western coast of...



Pacific ladyfish

Elops affinis

The Pacific ladyfish (Elops affinis), also known as the Pacific tenpounder and machete, is a species of ray-finned fish in the genus Elops, the only genus in the monotypic family Elopidae. The Pacific ladyfish can be found throughout the...



Pacific leaping blenny

Alticus arnoldorum

The Pacific leaping blenny (Alticus arnoldorum), also known as the leaping rockskipper, is a species of combtooth blenny (family Blenniidae) in the genus Alticus. The blennies are oviparous, and form distinct pairs when mating. Males can reach a...



Pacific nurse shark

Ginglymostoma unami

Sharks portalGinglymostoma unami, also known as the Pacific nurse shark is a nurse shark of the family Ginglymostomatidae. It is found in southeastern coast of Baja California, Mexico to Costa Rica including Gulf of California. It is 2.8 meters...



Pacific ocean perch

Sebastes alutus

The Pacific ocean perch (Sebastes alutus), also known as the Pacific rockfish, rose fish, red bream or red perch, is a fish whose range spans across the North Pacific : from southern California around the Pacific rim to northern Honshū, Japan,...



Pacific rudderfish

Psenopsis anomala

The Pacific rudderfish (Psenopsis anomala) is a marine fish also known by such names as Japanese butterfish, melon seed, wart perch, ibodai (Japanese name, イボダイ) or simply but ambiguously as butterfish. This fish, which can grow to 30 cm (12 in)...



Pacific sand sole

Psettichthys melanostictus

The Pacific sand sole (Psettichthys melanostictus), also known as simply sand sole, is a flatfish species inhabiting the northeastern Pacific waters where it lives on sandy bottoms. The only species in the genus, Psettichthys, it ranges from the...



Pacific sanddab

Citharichthys sordidus

The Pacific sanddab (Citharichthys sordidus) is a species of flatfish. It is by far the most common sanddab, and it shares its habitat with the longfin sanddab (C. xanthostigma) and the speckled sanddab (C. stigmaeus). It is a medium-sized...



Pacific sandfish

Trichodon trichodon

The Pacific sandfish (Trichodon trichodon) is a species of sandfish native to the Pacific coast of North America, from San Francisco Bay to Unimak Island in the Aleutian Islands. Like its cousin the sailfin or Japanese sandfish (Arctoscopus...



Pacific saury

Cololabis saira

The Pacific saury (*Cololabis saira*) is species of fish in the family Scomberesocidae. Saury is a seafood in several East Asian cuisines and is also known by the name mackerel pike.



Pacific seahorse

Hippocampus ingens

The Pacific seahorse, also known as the giant seahorse, (*Hippocampus ingens*) is a species of fish in the family Syngnathidae. Their genus name (*Hippocampus*) is derived from the Greek word hippos, which means "horse" and campus, which means "sea...



Pacific sharpnose shark

Rhizoprionodon longurio

The Pacific sharpnose shark (Rhizoprionodon longurio) often migrates along the Pacific coast of Mexico. They are most commonly caught in the artisanal fishery of Mazatlán. When a large sample of Pacific sharpnose sharks were observed scientists...



Pacific sierra

Scomberomorus sierra

The Pacific sierra (Scomberomorus sierra) also known as the Mexican sierra, is a ray-finned bony fish in the family Scombridae, better known as the mackerel family. More specifically, this fish is a member of the tribe Scomberomorini, the Spanish...



Pacific sleeper shark

Somniosus pacificus

The Pacific sleeper shark (Somniosus pacificus) is a sleeper shark of the family Somniosidae, found in the North Pacific on continental shelves and slopes in Arctic and temperate waters between latitudes 70°N and 22°N and in at least two places...



Pacific smalltail shark

Carcharhinus cerdale

The Pacific smalltail shark (Carcharhinus cerdale) is a species of requiem shark, in the family Carcharhinidae. It was described in 1898, but later mistakenly merged with Carcharhinus porosus. The mistake was corrected in 2011. It is relatively...



Pacific spadenose shark

Scoliodon macrorhynchos

The Pacific spadenose shark (Scoliodon macrorhynchos) is a species of requiem shark, in the family Carcharhinidae. It was once regarded as conspecific to the spadenose shark (S. laticaudus).



Pacific spiny dogfish

Squalus suckleyi

The Pacific spiny dogfish (Squalus suckleyi) is a common species of the Squalidae (dogfish) family of sharks and are among the most abundant species of sharks in the world. This species is closely related to Squalus acanthias and for many years...



Pacific staghorn sculpin

Leptocottus armatus

The Pacific staghorn sculpin (Leptocottus armatus) is a species of marine ray-finned fish belonging to the family Cottidae, the typical sculpins. This species is found in the eastern Pacific Ocean. It is the only species in the monospecific genus...



Pacific thread herring

Opisthonema libertate

The Pacific thread herring or deep-bodied Pacific thread herring (Opisthonema libertate) is a herring-like fish in the family Clupeidae. It is found in the Eastern Pacific. It can grow to 30 cm (12 in) total length.



Pacific viperfish

Chauliodus macouni

The Pacific viperfish, Chauliodus macouni, is a predatory deep-sea fish found in the North Pacific. It is reported as being either mesopelagic or bathypelagic, with diel vertical migration to shallower waters. The Pacific viperfish is one of the...



Pacific white skate

Bathyraja spinosissima

The Pacific white skate (Bathyraja spinosissima) is a species of skate in the family Arhynchobatidae. It is one of the deepest-living of all skates, occurring at a depth of 800 to 2,906m (2,600 and 9,500 feet) on the continental slope. It is...



Pacuma toadfish

Batrachoides surinamensis

The Pacuma toadfish (Batrachoides surinamensis) is a species of toadfish found in the Caribbean Sea and Atlantic Ocean along the coasts of Central and South America from Honduras to Brazil. It is the largest toadfish, reaching a length up to 57...



Paedocypris progenetica

Paedocypris progenetica

Paedocypris progenetica is a species of tiny cyprinid fish endemic to the Indonesian islands of Sumatra and Bintan where it is found in peat swamps and blackwater streams. It was discovered by Singaporean ichthyologist Heok Hui Tan. He has...



Pagrus major

Pagrus major

Pagrus major, the red seabream, red pargo, red porgy or silver seabream, is a species of marine ray-finned fish in the family Sparidae, which includes the seabreams and porgies. This species is found in the Western Pacific Ocean. The fish has...



Painted comber

Serranus scriba

The painted comber (Serranus scriba) is a species of marine ray-finned fish, a sea bass from the subfamily Serraninae, classified as part of the family Serranidae which includes the groupers and anthias. It is found in the eastern Atlantic Ocean,...



Painted frogfish

Antennarius pictus

The painted frogfish or spotted frogfish, Antennarius pictus, is a marine fish belonging to the family Antennariidae.



Painted greenling

Oxylebius pictus

The painted greenling (*Oxylebius pictus*) is a species of marine ray-finned fish belonging to the family Zaniolepididae, which includes this species and the combfishes. It is endemic to the northeast Pacific Ocean. It is the only species in the...



Painted sweetlips

Diagramma pictum

The painted sweetlips (*Diagramma pictum*), also known as the Australian slatey, blackall, bluey, grey sweetlips, moke, morwong, mother-in-law fish, painted blubber-lips, slate bream, slate sweetlips, smokey bream, thicklip or yellowdot sweetlips...



Paiute cutthroat trout

Oncorhynchus clarkii seleniris

Paiute cutthroat trout (Oncorhynchus clarkii seleniris) is one of fourteen subspecies of cutthroat trout. Paiute Cutthroat are native only to Silver King Creek, a headwater tributary of the Carson River in the Sierra Nevada, in California. This...



Paiute sculpin

Cottus beldingii

The Paiute sculpin (Cottus beldingii) is a species of fish in the family Cottidae. It is found in the United States, inhabiting the Columbia River drainage from Idaho, western Wyoming, and northeastern Nevada to western Washington and Oregon, and...



Pale ghost shark

Hydrolagus bemisi

The pale ghost shark (*Hydrolagus bemisi*) is a shortnose chimaera of the family Chimaeridae. It is endemic to New Zealand waters.



Pale-edged stingray

Telatrygon zugei

The pale-edged stingray or sharpnose stingray (Telatrygon zugei) is a species of stingray in the family Dasyatidae, found in the Indian and Pacific Oceans from India to the western Malay Archipelago and southern Japan. This bottom-dwelling ray is...



Pallid sturgeon

Scaphirhynchus albus

The pallid sturgeon (Scaphirhynchus albus) is an endangered species of ray-finned fish, endemic to the waters of the Missouri and lower Mississippi river basins of the United States. It may have even reached the St. Croix River before...



Pampus argenteus

Pampus argenteus

Pampus argenteus, the silver pomfret or white pomfret (or pompano to avoid confusion with true pomfrets of the genus Bramidae), is a species of butterfish that lives in the Indo-West Pacific, spanning the coastal waters of the Middle East...



Panaqolus maccus

Panaqolus maccus

Panaqolus maccus, commonly called the clown panaque, clown plecostomus, clown pleco, or ringlet pleco, is a dwarf loricariid. By numbering systems such as the L-number system, this fish may also be known as L104, L162, or LDA22.



Panaque armbrusteri

Panaque armbrusteri

Panaque armbrusteri is a species of fish in the South American armoured catfish family Loricariidae. This species is distributed throughout the Tapajós river, a large tributary of the Amazon River, with similar populations found in the Xingu,...



Panaque nigrolineatus

Panaque nigrolineatus

Panaque nigrolineatus, the royal panaque, royal plec, or royal pleco, is an herbivorous freshwater armored catfish native to Brazil, Colombia and Venezuela where it occurs in the Orinoco and Amazon basins. It is known for being one of the few...



Panaque suttonorum

Panaque suttonorum

Panaque suttonorum, commonly known as the blue-eye panaque, is a species of freshwater fish from the South American catfish family Loricariidae. It is known from the Maracaibo Basin in Venezuela. In the wild, the species feeds on algae and...



Pancake batfish

Halieutichthys aculeatus

The pancake batfish (*Halieutichthys aculeatus*) belongs to the batfish family Ogcocephalidae. Their distribution includes western Atlantic, North Carolina, northern Gulf of Mexico to northern South America. They inhabit a subtropical, sandy....



Panga

Pterogymnus lanarius

Panga or panga seabream is the common South African name for *Pterogymnus lanarius*, a small, ocean-dwelling fish, native to the southeast Atlantic Ocean and southwest Indian Ocean. Alternatively called "torpedo scads", they are cold-blooded with...



Pangasius conchophilus

Pangasius conchophilus

Pangasius conchophilus is a species of shark catfish. It is a freshwater, benthopelagic, potamodromous and tropical fish, measuring up to 120 centimetres (3.9 ft) long. It is found in the Mekong, Bangpakong, and Chao Phraya basins.



Pangasius kinabatanganensis

Pangasius kinabatanganensis

Pangasius kinabatanganensis is a species of shark catfish. It is a freshwater, benthopelagic and tropical fish, measuring up to 23.8 centimetres (0.78 ft) long. It is found in the Kinabatangan basin, in northeastern Borneo which is in the state...



Pangasius myanmar

Pangasius myanmar

Pangasius myanmar is a species of shark catfish. It is a freshwater, benthopelagic, tropical fish, measuring up to 120 centimetres (3.9 ft) long. It is found from Irrawaddy to Salween and in Rangoon.



Pangasius pangasius

Pangasius pangasius

Pangasius pangasius, the Pangas catfish, is a species of shark catfish native to fresh and brackish waters of Bangladesh, India, Myanmar, and Pakistan. It has also been introduced to Cambodia and Vietnam. This species grows to a standard length...



Pangio semicincta

Pangio semicincta

Pangio semicincta, the half-banded kuhli loach, is a species of cobitid loach in the genus Pangio found in the Malay Peninsula, Borneo, and Sumatra. It is a very popular fish in the aquarium trade, however, it is often confused for and mislabeled...



Panther danio

Danio aesculapii

The panther danio (Danio aesculapii) is a species of danio endemic to Myanmar. Before its description it was referred to as Danio species TW03. The panther is a smallish danio which on first glance looks a little dull, but displays a variety of...



Pao abei

Pao abei

Pao abei is a species of freshwater pufferfish from the Mekong, Chao Phraya and Mae Klong river basins in Southeast Asia. It is named after Japanese ichthyologist Tokiharu Abe. They are molluscivores, using the beak-like teeth to break open the...



Pao baileyi

Pao baileyi

Pao baileyi, the hairy pufferfish, is a species of pufferfish usually found in the rocky habitats, including rapids, of the Mekong mainstream and its larger tropical freshwater tributaries.



Pao cochinchinensis

Pao cochinchinensis

Pao cochinchinensis is a species of freshwater pufferfish native to the basins of the Mekong and Chao Phraya Rivers. This species grows to a length of 7 centimetres (2.8 in) SL.It is known for its reputation to attack people by slicing off bits...



Pao leiurus

Pao leiurus

Pao leiurus is a species of pufferfish found in Southeast Asia from Thailand to Indonesia and in particular the Mekong basin. It is occasionally imported into the aquarium trade but is reported to be aggressive and snappish.



Pao palembangensis

Pao palembangensis

Pao palembangensis (formerly Tetraodon palembangensis), is a species of freshwater pufferfish native to Thailand, Laos, Malaysia and Indonesia in Southeast Asia. Its commonly called the humpback puffer or dragon puffer. It is bred for aquaria as...



Pao suvattii

Pao suvattii

Pao suvattii, or Arrowhead puffer, is a species of pufferfish. It is a medium-sized pufferfish, reaching 11.5 cm (4.5 in) SL. It is also known as a Pignose Puffer or a Mekong Puffer. It is locally common in the Lower Mekong basin, and is...



Papyrocranus afer

Papyrocranus afer

The reticulated knifefish (Papyrocranus afer) is a species of freshwater fish in the family Notopteridae, the featherbacks, found in tropical West Africa. It reaches a maximum length of 80 cm (31 in) and a reported weight of 1.3 kg (2.9 lb).



Paracanthocobitis urophthalma

Paracanthocobitis urophthalma

Acanthocobitis (Paracanthocobitis) urophthalma, also known as the banded mountain zipper loach, is a species of ray-finned fish in the genus, or subgenus, Paracanthocobitis. This species is endemic to southwestern Sri Lanka.



Paracanthurus

Paracanthurus hepatus

Paracanthurus hepatus is a species of Indo-Pacific surgeonfish. A popular fish in marine aquaria, it is the only member of the genus Paracanthurus. A number of common names are attributed to the species, including regal tang, palette surgeonfish,...



Parachaetodon

Parachaetodon ocellatus

Parachaetodon is a monotypic genus of butterflyfishes, the only species being the sixspine butterflyfish (Parachaetodon ocellatus), which is also known as the ocellate butterflyfish or eyespot butterflyfish. is a species of butterflyfish native...



Parachanna africana

Parachanna africana

Parachanna africana, the African snakehead or Niger snakehead, is a species of fish from west-central Africa. Little is published on its biology. Limited primarily to coastal sections of rivers, it is thought to be a nest-building, thrust...



Parachromis dovii

Parachromis dovii

Parachromis dovii, the guapote, rainbow bass, or wolf cichlid, is a species of cichlid native to Central America where it occurs in lakes, rivers and streams in Honduras, Nicaragua and Costa Rica. It is one of the largest cichlids, reaching up to...



Parachromis friedrichsthalii

Parachromis friedrichsthalii

Parachromis friedrichsthalii, the Yellowjacket cichlid, is a species of cichlid native to Central America where it is found along the Atlantic Slope in Mexico, Belize, Honduras and Guatemala. This species grows to a length of 28 centimetres (11...



Parachromis managuensis

Parachromis managuensis

Parachromis managuensis is a large species of cichlid native to freshwater habitats in Central America, where it is found from Honduras to Costa Rica. The binomial name refers to Lake Managua in Nicaragua, from which the holotype was obtained. It...



Parachromis motaguensis

Parachromis motaguensis

Parachromis motaguensis, the False yellowjacket cichlid, is a species of cichlid native to Guatemala and Honduras in Central America. This species grows to a length of 30 centimetres (12 in) TL. This species is farmed.



Paradise fish

Macropodus opercularis

The fork tailed paradisefish (often just called paradise fish, paradise-fish, paradisefish), or paradise gourami (Macropodus opercularis) is a species of gourami found in most types of fresh water in East Asia, ranging from the Korean Peninsula...



Parajulis

Parajulis poecilopterus

Parajulis poecilopterus is a species of wrasse native to the northwestern Pacific Ocean along the coast of Asia. It is an inshore species, being found in areas with a substrate of pebbles. This species grows to 34 cm (13 in) in total length. This...



Paralabrax clathratus

Paralabrax clathratus

Paralabrax clathratus, the kelp bass, bull bass or calico bass, is a species of marine ray-finned fish, a sea bass from the subfamily Serraninae, classified as part of the family Serranidae which includes the groupers and anthias. It is found in...



Paralichthys lethostigma

Paralichthys lethostigma

Paralichthys lethostigma, the southern flounder, is a species of large-tooth flounder native to the East Coast of the United States and the northern Gulf of Mexico. It is a popular sport fish and is the largest and most commercially valuable...



Parambassis ranga

Parambassis ranga

Parambassis ranga, commonly known as the Indian glassy fish, Indian glassy perch, or Indian X-ray fish, is a species of freshwater fish in the Asiatic glassfish family Ambassidae of order Perciformes. It is native to an area of South Asia from...



Paranthias colonus

Paranthias colonus

Paranthias colonus (the Pacific creolefish) is a species of grouper found in the Eastern Pacific Ocean. They are typically found in small aggregations well above reefs, but will retreat to the reef at the approach of danger. They form large...



Paraplesiops bleekeri

Paraplesiops bleekeri

Paraplesiops bleekeri, commonly known as the eastern blue devil, blue-tipped long-fin or Bleeker's blue devil fish, is a species of fish in the family Plesiopidae. This colourful, secretive fish is endemic to Australia, where it is a protected...



Paratilapia polleni

Paratilapia polleni

Paratilapia polleni is a medium-sized cichlid endemic to Madagascar. It is also a popular fish for display at public aquaria. It is sometimes referred to by the common names starry-night cichlid, polleni cichlid, and black diamond cichlid, while...



Pardachirus marmoratus

Pardachirus marmoratus

Pardachirus marmoratus, the finless sole, speckled sole or Red Sea Moses sole, is a species of flatfish native to the western Indian Ocean.



Paretroplus menarambo

Paretroplus menarambo

Paretroplus menarambo (pinstripe menarambo or pinstripe damba) is a species of cichlid fish. It is threatened by invasive species and over-fishing. It is part of a captive breeding program by public institutions like London Zoo and Bolton Museum...



Pareutropius debauwi

Pareutropius debauwi

Pareutropius debauwi is a species that belong to the Order Actinopterygii, ray-finned fishes, and can be placed in the Schilbeidae family, Schilbid catfishes. This family includes 9 genera and 46 species. The common name for Pareutropius debauwi...



Paristiopterus labiosus

Paristiopterus labiosus

Paristiopterus labiosus, the giant boarfish, is a species of armorhead native to the coastal waters of southern Australia, and New Zealand. It occurs over the continental shelf at depths from 20 to 200 m (66 to 656 ft). This species can reach a...



Parore

Girella tricuspidata

The parore (Girella tricuspidata) also known as luderick, black bream, black snapper or blackfish is a species of marine ray-finned fish, a sea chub from the family Kyphosidae which is found in the southwestern Pacific Ocean off Australia and New...



Parosphromenus harveyi

Parosphromenus harveyi

Parosphromenus harveyi is a species of gourami endemic to Malaysia, where it is only known from Selangor. The most extreme freshwater habitats in Peninsular Malaysia is the peat swamp forest that consist of dark-coloured and highly acidic waters....



Parupeneus cyclostomus

Parupeneus cyclostomus

Parupeneus cyclostomus, the gold-saddle goatfish, blue goatfish or yellowsaddle goatfish, is a species of goatfish native to the Indo-Pacific. It is a commercially important species, as well as being sought out as a game fish, though it has been...



Parupeneus forsskali

Parupeneus forsskali

Parupeneus forsskali, common name Red Sea goatfish, is a species of goatfish belonging to the family Mullidae.



Parupeneus heptacanthus

Parupeneus heptacanthus

Parupeneus heptacanthus, commonly known as cinnabar goatfish, and sold in UK as "Red Mullet", is a marine fish native to the western Pacific and Indian Oceans.



Parupeneus multifasciatus

Parupeneus multifasciatus

Parupeneus multifasciatus, the manybar goatfish, is a species of goatfish native to the eastern Indian Ocean Pacific Ocean where it is found from Christmas Island to the Hawaiian Islands and from southern Japan to Lord Howe Island. Its Hawaiian...



Patagonian toothfish

Dissostichus eleginoides

The Patagonian toothfish (Dissostichus eleginoides), also known as Chilean sea bass, mero, icefish, and Antarctic cod, is a species of notothen found in cold waters (1–4 °C or 34–39 °F) between depths of 45 and 3,850 m (150 and 12,630 ft) in the...



Payara

Hydrolycus scomberoides

The payara, Hydrolycus scomberoides, is a species of dogtooth tetra. This predatory fish is found in the Amazon Basin in tropical South America. It was the first of four species to be described in the genus Hydrolycus.



Peaceful betta

Betta imbellis

The peaceful betta or crescent betta (Betta imbellis) is a species of gourami native to Southeast Asia. Betta imbellis has a pair of suprabranchial chambers that each house a labyrinth organ, a complex bony structure lined with thin, highly...



Peacock blenny

Salaria pavo

Salaria pavo, the peacock blenny, is a species of combtooth blenny found in the eastern Atlantic coast from France to Morocco; also in the Mediterranean Sea, Black Sea and the eastern Adriatic Sea. This species has colonised the northern Red Sea...



Peacock flounder

Bothus mancus

The peacock flounder (Bothus mancus), also known as the flowery flounder, is a species of fish in the family Bothidae (lefteye flounders). The species is found widely in relatively shallow waters in the Indo-Pacific, also ranging into warmer...



Peacock gudgeon

Tateurndina ocellicauda

The peacock gudgeon or peacock goby (*Tateurndina ocellicauda*) is a tropical freshwater species of fish in the family Eleotridae that is endemic to the eastern part of Papua New Guinea. It can be found in schools hovering over the substrate in...



Pearl cichlid

Geophagus brasiliensis

The pearl cichlid (*Geophagus brasiliensis*) is a species of fish in the cichlid family. It is native to southeastern Brazil (north to Bahia), Paraguay, Uruguay and far northeastern Argentina, where it is found in rivers, lakes, and nearby slightly...



Pearl danio

Danio albolineatus

The pearl danio (*Danio albolineatus*) is a tropical fish belonging to the minnow family Cyprinidae. Originating in Sumatra, Myanmar, and Thailand and Vietnam (Phu Quoc Island), this fish is sometimes found in aquariums by fish-keeping hobbyists....



Pearl gourami

Trichopodus leerii

The pearl gourami (*Trichopodus leerii*) is a species of gourami native to Southeast Asia.



Pearl stingray

Fontitrygon margaritella

The pearl stingray (Fontitrygon margaritella) is a little-known species of stingray in the family Dasyatidae, found in shallow coastal waters from Mauritania to Angola, though fossils have been found in Portugal. Growing to 30 cm (12 in) across....



Pearlscale butterflyfish

Chaetodon xanthurus

The pearlscale butterflyfish (Chaetodon xanthurus), also known as yellow-tailed butterflyfish, crosshatch butterflyfish or Philippines chevron butterflyfish is a species of marine ray-finned fish, a butterflyfish belonging to the family...



Pearly razorfish

Xyrichtys novacula

The pearly razorfish (Xyrichtys novacula), commonly known as cleaver wrasse, is a species of wrasse. It is of minor importance to local commercial fisheries and is popular as a game fish. It can also be found in the aquarium trade.



Peckoltia pankimpuju

Peckoltia pankimpuju

Peckoltia pankimpuju is a species of armored catfish from the family Loricariidae, native to the Marañón River in the upper Amazon basin of Peru. It is commonly called the coal pleco, Peruvian lyre-tail, and L350 under the L-number code. It...



Peckoltia vittata

Peckoltia vittata

Peckoltia vittata is a species of catfish belonging to the subfamily Hypostominae of the family Loricariidae.



Pedder galaxias

Galaxias pedderensis

The Pedder galaxias (Galaxias pedderensis) is an Australian freshwater fish. It is considered to be extinct in the wild since 2005 by the EPBC Act, and was originally found only in Lake Pedder in Tasmania.



Pelagic stingray

Pteroplatytrygon violacea

The pelagic stingray (Pteroplatytrygon violacea) is a species of stingray in the family Dasyatidae, and the sole member of its genus. It is characterized by the wedge-like shape of its pectoral fin disc, which is much wider than long, as well as...



Pelagic thresher

Alopias pelagicus

The pelagic thresher (Alopias pelagicus) is a species of thresher shark, family Alopiidae; this group of sharks is characterized by the greatly elongated upper lobes of their caudal fins. The pelagic thresher occurs in the tropical and...



Pelates quadrilineatus

Pelates quadrilineatus

Pelates quadrilineatus, also known as the trumpeter perch or fourlined terapon, is a species of ray-finned fish in the family Terapontidae, the grunters. It occurs in the western Indo-Pacific region, and also in the eastern Mediterranean Sea,...



Pelecus cultratus

Pelecus cultratus

Pelecus cultratus, commonly known as the ziege, sichel, sabre carp or sabrefish, is a cyprinid fish species from Eastern Europe and adjacent Asian regions, the only one in its genus, inhabiting the lower reaches of rivers and brackish waters in...



Peled

Coregonus peled

The peled (Coregonus peled), also called the northern whitefish, is a species of freshwater whitefish in the family Salmonidae. It is found in northern Europe and Asia. The peled is related to ciscoes of the Coregonus sardinella complex.



Pelican eel

Eurypharynx pelecyanoides

The pelican eel (Eurypharynx pelecyanoides) is a deep-sea eel. It is the only known member of the genus Eurypharynx and the family Eurypharyngidae. It belongs to the "saccopharyngiforms", members of which were historically placed in their own...



Pelvicachromis pulcher

Pelvicachromis pulcher

Pelvicachromis pulcher is a freshwater fish of the cichlid family, endemic to Nigeria and Cameroon. It is popular amongst aquarium hobbyists, and is most commonly sold under the name kribensis, although it has other common names, including...



Pelvicachromis sacrimontis

Pelvicachromis sacrimontis

Pelvicachromis sacrimontis is a freshwater fish of the cichlid family known only from a small area of southeastern Nigeria. Currently Fishbase considers this binomial to be a junior synonym of P. pulcher and, it was also known as Pelvicachromis...



Pelvicachromis taeniatus

Pelvicachromis taeniatus

Pelvicachromis taeniatus is a species of cichlid from Benin and Nigeria that is occasionally kept as an aquarium fish. It is native to the soft-water rivers. This species can reach a length of 7.1 centimetres (2.8 in) SL. It is known to exist in...



Penang betta

Betta pugnax

The Penang betta (Betta pugnax) is a species of gourami native to Southeast Asia and common in swiftly flowing forest streams of the Malay Peninsula, Thailand, Sumatra, and the Riau Islands. In addition to its native range, the species has been...



Pennant coralfish

Heniochus acuminatus

The pennant coralfish (Heniochus acuminatus), also known as the longfin bannerfish, reef bannerfish or coachman, is a species of fish of the family Chaetodontidae, native to the Indo-Pacific area.



Peppermint angelfish

Centropyge boylei

The peppermint angelfish (Centropyge boylei) is a relatively small species of marine ray-finned fish, a marine angelfish belonging to the family Pomacanthidae. It is found in the Central area of the South Pacific Ocean, and is known to be native...



Peprilus paru

Peprilus paru

Peprilus paru, (harvestfish or American harvestfish; syn. Peprilus alepidotus), also occasionally known by a few local names as star butter fish or sometimes even simply as butterfish, is a marine, benthopelagic, circular-shaped and deep-bodied...



Percina shumardi

Percina shumardi

Percina shumardi (river darter) is a benthic species of freshwater ray-finned fish, a darter from the subfamily Etheostominae, part of the family Percidae, which also contains the perches, ruffes and pikeperches. It is native to North America....



Percopsis omiscomaycus

Percopsis omiscomaycus

Percopsis omiscomaycus, also known as the trout-perch, the grounder or the sand minnow, is one of two species in the family Percopsidae. Its name comes from the Greek root words perc, meaning perch and opsi meaning appearance. The species name...



Periophthalmus gracilis

Periophthalmus gracilis

The slender mudskipper (Periophthalmus gracilis) is a species of mudskippers native to marine and brackish waters of the eastern Indian Ocean and the western Pacific Ocean where it is an inhabitant of the intertidal zone, capable of spending time...



Perissodus microlepis

Perissodus microlepis

Perissodus microlepis is a species of cichlid endemic to Lake Tanganyika. This species reaches a length of 11 centimetres (4.3 in) TL. This species can also be found in the aquarium trade. It is a scale-eating 'parasite' on other fish species. It...



Permit

Trachinotus falcatus

The permit (Trachinotus falcatus) is a game fish of the western Atlantic Ocean belonging to the family Carangidae. Adults feed on crabs, shrimp, and smaller fish.



Persian sturgeon

Acipenser persicus

The Persian sturgeon (Acipenser persicus) is a species of fish in the family Acipenseridae. It is found in the Caspian Sea and to a lesser extent the Black Sea and ascends certain rivers to spawn, mainly the Volga, Kura, Araks and Ural Rivers. It...



Peruvian anchoveta

Engraulis ringens

The Peruvian anchoveta (Engraulis ringens) is a species of fish of the anchovy family, Engraulidae, from the Southeast Pacific Ocean. It is one of the most commercially important fish species in the world, with annual harvests varying between...



Peruvian Pacific sardine

Sardinops sagax sagax

The Peruvian Pacific sardine (scientific name Sardinops sagax sagax) is a subspecies of the South American pilchard found in Peru. Related subspecies and species of international importance include Sardinops sagax caeruleus (USA), Sardinops sagax...



Petenia splendida

Petenia splendida

Petenia splendida, the bay snook, is a species of cichlid from Guatemala and northern Central America. It is important to local commercial fisheries.



Peters's elephantnose fish

Gnathonemus petersii

Peters's elephant-nose fish (Gnathonemus petersii) is an African freshwater elephantfish in the genus Gnathonemus. Other names in English include elephantnose fish, long-nosed elephant fish, and Ubangi mormyrid, after the Ubangi River. The Latin...



Pethia cumingii

Pethia cumingii

Pethia cumingii, known as the Cuming's barb or the two spot barb (though that name can also apply to the Ticto barb, *P. ticto*), is a species of cyprinid fish endemic to Sri Lanka. Its stocks have declined in recent decades, and in 1996 it was...



Pethia reval

Pethia reval

Pethia reval, the red-finned barb, is a species of cyprinid fish endemic to Sri Lanka. This species can reach a length of 3.4 centimetres (1.3 in) SL. Congener, Pethia cumingii, also a Sri Lankan endemic, has long been presumed to be dichromatic...



Pethia stoliczkana

Pethia stoliczkana

Pethia stoliczkana is a fresh water tropical cyprinid fish native to the upper Mekong, Salween, Irrawaddy, Meklong and upper Charo Phraya basins in the countries of Nepal, India, Pakistan, Myanmar, Bangladesh, Laos, Thailand, China and Sri Lanka.



Petitella bleheri

Hemigrammus bleheri

Petitella bleheri is a species of characin found in Amazon Basin in Brazil and Peru. One of three species called rummy-nose tetra, but is also called firehead tetra according to FishBase. Reaching a standard length of 2 in (5 cm) it is a common...



Petrale sole

Eopsetta jordani

The Petrale sole (Eopsetta jordani) is an edible flatfish of the family Pleuronectidae. It is a demersal fish that lives on sandy bottoms, usually in deep water, down to depths of about 550 metres (1,800 ft). Males can grow to 53 centimetres (21...



Petrocephalus bane

Petrocephalus bane

Petrocephalus bane is a species of elephantfish native to Africa where it occurs in the basins of the Nile, Bénoué and Volta rivers and the Chad Basin. It prefers relatively still waters of lakes, irrigation canals and lagoons. This species can...



Pholidichthys leucotaenia

Pholidichthys leucotaenia

Pholidichthys leucotaenia, commonly known as the convict blenny/goby or the engineer blenny/goby, is a marine fish from the west-central Pacific Ocean. Despite its common names, it is neither a blenny nor a goby, but is in fact one of two species...



Photoblepharon palpebratum

Photoblepharon palpebratum

Photoblepharon palpebratum (Syn. P. palpebratus), the eyelight fish or one-fin flashlightfish, is a species of saltwater anomalopid fish of the order Beryciformes. It is native to the western and central Pacific Ocean where it is found along...



Photoblepharon steinitzi

Photoblepharon steinitzi

The flashlight fish (Photoblepharon steinitzi) is a species of anomalopid fish of the order Beryciformes. It is native to the western Indian Ocean and the Red Sea where it is found near coral reefs. It is a nocturnal predator, hiding in crevices...



Photocorynus

Photocorynus spiniceps

Photocorynus spiniceps is a species of anglerfish in the family Linophrynidae. It is in the monotypic genus Photocorynus. Photocorynus spiniceps originates from the Philippines. The known mature male individuals are 6.2–7.3 millimeters (0.25-0.3...



Phreatichthys andruzzii

Phreatichthys andruzzii

Phreatichthys andruzzii, a species of cyprinid fish, is the only species of the genus Phreatichthys, and is endemic to Somalia. This cave-adapted fish is whitish (not pigmented) and blind. It has no scales. It is considered to have evolved in the...



Phreatobius cisternarum

Phreatobius cisternarum

Phreatobius cisternarum is a species of catfish in the genus Phreatobius. This Brazilian fish is one of the few fish species that live underground in phreatic habitats. It has proved problematic in its classification.



Phreatobius sanguijuela

Phreatobius sanguijuela

Phreatobius sanguijuela is a South American species of heptapterid catfish that lives in underground waters. This species was first discovered in a hand-dug well located within the Parag  a River drainage, a tributary of the Guapor   (It  nez) River...



Phycis phycis

Phycis phycis

Phycis phycis, the forkbeard, is a species of phycid hakes in the family Phycidae.



Piaractus brachypomus

Piaractus brachypomus

Piaractus brachypomus, the pirapitinga, is a large species of pacu, a close relative of piranhas and silver dollars, in the serrasalmid family. It is native to the Amazon basin in tropical South America, but it formerly included populations in...



Piaractus mesopotamicus

Piaractus mesopotamicus

Piaractus mesopotamicus, the small-scaled pacu, Paran   River pacu or simply pacu (a name shared with other species), is a South American ray-finned fish that is native to the Paraguay-Paran   River basin, but it has been introduced by aquaculture...



Pickhandle barracuda

Sphyr  na jello

The pickhandle barracuda (Sphyraena jello) is a species of barracuda found throughout the Indo-Pacific region. Its common name derives from the dark marks along its sides, which look like the thick ends of pickaxe handles. Sea anglers sometimes...



Pictichromis diadema

Pictichromis diadema

Pictichromis diadema, the bicolor dottyback, diadem dottyback, or purple-top dottyback is a fish species of the dottyback family. It is a popular aquarium fish species.



Pictichromis paccagnellorum

Pictichromis paccagnellae

The royal dottyback (Pictichromis paccagnellorum), also called the bicolor dottyback, gramma dotty or false gramma is a fish commonly kept in marine aquariums. The front of the fish is bright purple and the posterior is yellow. In an aquarium, it...



Pictichromis porphyrea

Pictichromis porphyrea

Pictichromis porphyrea, the magenta dottyback, is a fish in the dottyback family from the western Pacific. It can be found from the Philippines to Samoa, extending north to the Ryukyu Islands and south to the Moluccas and Admiralty Islands. It...



Picturesque dragonet

Synchiropus picturatus

The picturesque dragonet (Synchiropus picturatus) is a brightly colored member of the dragonet family native to the Indo-West Pacific: Philippines, eastern Indonesia and northwest Australia. It occasionally makes its way into the aquarium trade,...



Piebald madtom

Noturus gladiator

The piebald madtom (Noturus gladiator) is a species of madtom catfish that is found in seven river systems in the Tennessee and Mississippi coastal tributaries. The most common systems N. gladiator is found in are, the Hatchie River, Obion River,...



Pigeye shark

Carcharhinus amboinensis

The pigeye shark or Java shark (Carcharhinus amboinensis) is an uncommon species of requiem shark, in the family Carcharhinidae, found in the warm coastal waters of the eastern Atlantic and western Indo-Pacific. It prefers shallow, murky...



Pigo

Rutilus pigus

The pigo (Rutilus pigus) is a species of freshwater fish in the roach genus Rutilus of the family Cyprinidae. The fish is native to northern Italy and Switzerland. It inhabits subalpine lakes of the northern Adriatic basin from the Livenza to the...



Pike topminnow

Belonesox belizanus

The pike topminnow (Belonesox belizanus), more commonly known as pike killifish and sometimes referred to as topminnow, is a species of poeciliid found from Mexico to Nicaragua. It has also been introduced to Florida, USA. It is the only known...



Pilot fish

Naucrates ductor

The pilot fish (*Naukrates ductor*) is a carnivorous fish of the trevally, or jackfish family, Carangidae. It is widely distributed and lives in warm or tropical open seas.



Pimelodus pictus

Pimelodus pictus

Pimelodus pictus, also known as the pictus cat or pictus catfish, is a small (11.0 centimetres (4.3 in TL) member of the catfish family Pimelodidae, native to the Amazon and Orinoco river basins and commonly kept as a pet in freshwater...



Pimelodus pohli

Pimelodus pohli

Pimelodus pohli is a species of catfish belonging to the family Pimelodidae. It is only found in the São Francisco River, Brazil, and its tributaries. Smaller, younger members of this species (under 10.0 centimetres (3.9 in) standard length) have...



Pindani

Pseudotropheus socolofi

The pindani (Pseudotropheus socolofi) is a species of cichlid endemic to Lake Malawi preferring areas with sandy substrates and nearby rocks where the males establish their territories. This species can reach a length of 6.7 centimetres (2.6 in)...



Pinhead pearlfish

Encheliophis boraborensis

The pinhead pearlfish, Encheliophis boraborensis, is a species of slender, ray-finned fish in the family Carapidae found in the tropical Indo-Pacific Ocean; it normally lives inside the body cavity of a sea cucumber such as the pineapple sea...



Pink cusk-eel

Genypterus blacodes

The pink cusk-eel, Genypterus blacodes, is a demersal species of cusk-eel found in the oceans around southern Australia, Chile, Brazil, and around New Zealand except the east coast of Northland, in depths of 22 to 1,000 metres (70 to 3,280 feet;...



Pink lanternshark

Etmopterus dianthus

Sharks portalThe pink lanternshark (*Etmopterus dianthus*) is a shark of the family Etmopteridae found around Australia and New Caledonia, at depths of between 110 and 880 m. Its length is up to 41 cm.Reproduction is ovoviviparous.



Pink maomao

Caprodon longimanus

The pink maomao (*Caprodon longimanus*), also known as the longfin perch, is a species of marine ray-finned fish, a member of the subfamily Anthiinae which is part of the grouper and sea bass family Serranidae. It is found in the southern Pacific...



Pink salmon

Oncorhynchus gorbuscha

Pink salmon or humpback salmon (*Oncorhynchus gorbuscha*) is a species of euryhaline ray-finned fish in the family Salmonidae. It is the type species of the genus *Oncorhynchus* (Pacific salmon), and is the smallest and most abundant of the seven...



Pink skunk clownfish

Amphiprion perideraion

The pink skunk clownfish (*Amphiprion perideraion*), also known as the pink anemonefish, is a species of anemonefish that is widespread from northern Australia through the Malay Archipelago and Melanesia. Like all anemonefishes, it forms a...



Pink whipray

Pateobatis fai

The pink whipray (Pateobatis fai) is a species of stingray in the family Dasyatidae, with a wide but ill-defined distribution in the tropical Indo-Pacific from southern Africa to Polynesia. It is a bottom dweller that generally inhabits shallow...



Pinktail triggerfish

Melichthys vidua

The pinktail triggerfish (Melichthys vidua) is a species of triggerfish from reefs in the Indo-Pacific. It is very dark green (often appears almost black) with translucent whitish-pink dorsal and tail fins. They can grow up to 40 centimetres (16...



Piper gurnard

Trigla lyra

The piper gurnard (Trigla lyra), also known as the piper or the lyre gurnard, is a species of marine, demersal ray-finned fish from the family Triglidae, the gurnards and sea robins. It is found in the eastern Atlantic Ocean and the Mediterranean...



Pirate perch

Aphredoderus sayanus

The pirate perch (Aphredoderus sayanus) is a freshwater fish that commonly inhabits coastal waters along the east coast of the United States and the backwater areas of the Mississippi Valley. This species is often found towards the bottom of...



Placidochromis phenochilus

Placidochromis phenochilus

Placidochromis phenochilus is a species of cichlid endemic to the northern parts of Lake Malawi. This species can reach a length of 15.7 centimetres (6.2 in) TL.



Plains topminnow

Fundulus sciadicus

The plains topminnow (Fundulus sciadicus) is a species of freshwater topminnow found in North America. The fish has a small range within the United States of America which consists of two major populations.



Plate fish

Bothus lunatus

The plate fish (Bothus lunatus) is a flounder in the genus Bothus, found in the warmer parts of the Atlantic including the Caribbean. Its typical habitat is sandy plains near coral reefs and it is able to change its colouring to make it...



Platycephalus fuscus

Platycephalus fuscus

Platycephalus fuscus, the dusky flathead or black flathead, (literally translating from Ancient Greek as "flat-head dusky") is a large predatory fish and the largest member of the family Platycephalidae. Dusky flathead are a largely estuarine...



Platydoras costatus

Platydoras costatus

Platydoras costatus, the Raphael catfish, is a species of thorny catfish native to rivers in Brasil, Suriname and French Guiana. It was long confused with other species in the genus Platydoras, especially the more widespread "true" striped...



Plectorhinchus caeruleonothus

Plectorhinchus caeruleonothus

Plectorhinchus caeruleonothus, the blue bastard, is a species of ray-finned fish belonging to the subfamily Plectorhinchinae, the sweetlips, one of two subfamilies in the family Haemulidae, the grunts. It was discovered in Australia in 2015.



Plectorhinchus chaetodonoides

Plectorhinchus chaetodonoides

Plectorhinchus chaetodonoides, the harlequin sweetlips, clown sweetlips, spotted sweetlips or many-spotted sweetlips, is a species of marine ray-finned fish, a sweetlips belonging to the subfamily Plectorhinchinae, one of two subfamilies in the...



Plectorhinchus gaterinus

Plectorhinchus gaterinus

Plectorhinchus gaterinus, commonly known as the blackspotted rubberlip or African grunt, is a species of marine ray-finned fish, a sweetlips belonging to the subfamily Plectorhinchinae, one of two subfamilies in the family Haemulidae, the grunts....



Plectorhinchus gibbosus

Plectorhinchus gibbosus

Plectorhinchus gibbosus, commonly known as the Harry hotlips, black sweetlips, brown sweetlips, dusky sweetlips, gibbous sweetlips, hairy hotlips or humpback sweetlips, is a species of marine ray-finned fish, a sweetlips belonging to the...



Plectorhinchus lineatus

Plectorhinchus lineatus

Plectorhinchus lineatus, also known as the yellowbanded sweetlips, oblique-banded sweetlips, diagonal-banded sweetlips, Goldman's sweetlips, lined blubber-lips, lined sweetlips, many-lined sweetlips, striped sweetlips and yellowband sweetlips, is...



Plectropomus laevis

Plectropomus laevis

Plectropomus laevis, known commonly as the black-saddled coral grouper, cluespotted coral trout, blacksaddled coral trout, blue-spot trout, Chinese footballer, footballer cod, footballer coral trout, oceanic coral trout or tiger trout, is a...



Plotosus canius

Plotosus canius

Plotosus canius is a species of eeltail catfish, in the genus Plotosus, native to the Indian Ocean, the western Pacific Ocean and New Guinea.



Plotosus lineatus

Plotosus lineatus

Plotosus lineatus, common name striped eel catfish, is a species of eeltail catfishes belonging to the family Plotosidae.The fish is known in Israel as "Nasrallah fish" after the Lebanese cleric and political figure Hassan Nasrallah.



Pocket shark

Mollisquama parini

Sharks portalThe pocket shark (*Mollisquama parini*) is a species of kitefin shark in the family Dalatiidae. The species is found in deep water off Chile in the southeastern Pacific Ocean. It was the only member of the genus *Mollisquama*, until...



Poecilia mexicana

Poecilia mexicana

Poecilia mexicana, commonly known as the shortfin molly or Atlantic molly, is a species of poeciliid fish native to fresh and brackish water in Mexico and Guatemala. One population is found in caves and known as the cave molly.



Poecilia sphenops

Poecilia sphenops

Poecilia sphenops is a species of fish, of the genus *Poecilia*, known under the common name molly; to distinguish it from its congeners, it is sometimes called short-finned molly or common molly. They inhabit fresh water streams, coastal brackish,...



Poecilia velifera

Poecilia velifera

Poecilia velifera, known as the Yucatan molly and also as the giant sailfin molly amongst aquarists, is a large livebearer that lives in coastal waters of the Yucatan peninsula. These live-bearer (*Poeciliidae*) fish are particularly well known for...



Poecilia wingei

Poecilia wingei

Poecilia wingei, known to aquarists as Endlers or Endler's livebearer, in the genus Poecilia, is a small fish native to the Paria Peninsula in Venezuela. They are prolific breeders and often hybridize with guppies. These very colorful hybrids are...



Pointy-nosed blue chimaera

Hydrolagus trolli

The pointy-nosed blue chimaera (Hydrolagus trolli), also known as the pointy-nosed blue ratfish, Ray Troll's chimaera or abyssal ghostshark, is a species of deep-sea fish in the family Chimaeridae.



Pollachius pollachius

Pollachius pollachius

Pollachius pollachius is a species of marine fish in the family Gadidae. FAO uses the English name pollack for this species, whereas in American English it is known as European pollock. Other vernacular names include lythe, and in the Isle of...



Pollachius virens

Pollachius virens

The saithe (or) (Pollachius virens) is a species of marine fish in the genus Pollachius. Together with P. pollachius, it is generally referred to in the United States as pollock. Other names include the Boston blue (separate from bluefish),...



Pollan

Coregonus pollan

The pollan or Irish pollan (Coregonus pollan or Coregonus autumnalis) is a freshwater whitefish known only from five Irish lakes, Lough Neagh, Lower Lough Erne, Lough Ree, Lough Derg and Lough Allen. The pollan faces competition from introduced...



Polydactylus sexfilis

Polydactylus sexfilis

Polydactylus sexfilis, the six-finger threadfin or yellowthread threadfin, is a species of marine ray-finned fish, a threadfin from the family Polynemidae which is found in the Indian and Pacific Oceans.



Polypterus bichir

Polypterus bichir

Polypterus bichir, the Nile bichir, is a fish which lives in the Nile and some of its tributaries in Africa. It is a dark grayish color on the top, with a dark vertical marking and bands on the flank. This marking is more prominent on juveniles,...



Polypterus congicus

Polypterus congicus

Polypterus congicus, the Congo bichir, is a species of bichir with a maximum recorded size of 970 millimetres (38 in). The colour also varies from yellowish brown to grey, darker in the top, paler in the ventral area. It has a pattern of around 8...



Polypterus endlicheri

Polypterus endlicheri

Polypterus endlicheri, the saddled bichir, is one of the largest species of the Polypterus genus of freshwater fish. Named in honor of botanist Stephan Endlicher (1804–1849), who apparently discovered the species in the fish collection at the...



Polypterus makelembembe

Polypterus makelembembe

Polypterus makelembembe is a species of the fish genus Polypterus, found in the central basin of the Congo River. It was once considered a morph of the closely related Polypterus retropinnis, but was given species status in 2006 with a...



Polypterus ornatipinnis

Polypterus ornatipinnis

Polypterus ornatipinnis, the ornate bichir, is a bony fish found in Lake Tanganyika and the Congo River basin in Central and East Africa.



Polypterus palmas

Polypterus palmas

Polypterus palmas, also called the shortfin or marbled bichir, is a fish in the family Polypteridae found in freshwater environments throughout West Africa.



Polypterus polli

Polypterus polli

Polypterus polli, Poll's bichir, is a species of bichir from the Malebo Pool and the lower and central basins of the Congo River. It was named in honor of Belgian ichthyologist Max Poll.



Polypterus senegalus

Polypterus senegalus

Polypterus senegalus, the Senegal bichir, gray bichir or Cuvier's bichir, and sometimes called the "dinosaur eel" (a misnomer, as the creature is neither an eel nor a dinosaur), "dinosaur bichir", or "dragon fish" is in the pet trade due to its...



Polypterus teugelsi

Polypterus teugelsi

Polypterus teugelsi (Teugelsi bichir) is a species of carnivorous, nocturnal bichir (a group of primitive ray-finned fishes) that lives in the Cross River drainage basin in the country of Cameroon. P. teugelsi was described in 2004 by Ralf Britz.



Polypterus weeksii

Polypterus weeksii

Polypterus weeksii, the mottled bichir, is a fish in the family Polypteridae found in the central basin of the Congo River. It grows to about 54 cm in head-to-tail length. Named in honor of John Henry Weeks (1861-1924), Baptist missionary....



Pomacanthus asfur

Pomacanthus asfur

Pomacanthus asfur, the Arabian angelfish, is a species of marine ray-finned fish, a marine angelfish belonging to the family Pomacanthidae. It is found in the Western Indian Ocean.



Pomacanthus maculosus

Pomacanthus maculosus

Pomacanthus maculosus, the yellowbar angelfish, half-moon angelfish, yellow-marked angelfish, yellowband angelfish or yellow-blotched angelfish, is a species of marine ray-finned fish, a marine angelfish belonging to the family Pomacanthidae. It...



Pomacanthus navarchus

Pomacanthus navarchus

Pomacanthus navarchus, the blue-girdled angelfish or majestic angelfish, is a marine angelfish from the Indo-Pacific region. It occasionally makes its way into the aquarium trade. P. navarchus is one of the smallest of the larger species of...



Pomacanthus semicirculatus

Pomacanthus semicirculatus

Pomacanthus semicirculatus, also known as the semicircled angelfish, Koran angelfish, blue angelfish, zebra angelfish or half-circled angelfish, is a species of ray-finned fish, a marine angelfish, in the family Pomacanthidae. It is found in the...



Pomacanthus xanthometopon

Pomacanthus xanthometopon

Pomacanthus xanthometopon is a marine ray-finned fish, a marine angelfish belonging to the family Pomacanthidae found in shallow parts of the Indo-Pacific. It is commonly known as the blueface angelfish or the yellowface angelfish because of its...



Pomacanthus zonipectus

Pomacanthus zonipectus

Pomacanthus zonipectus, the Cortez angelfish, is a species of marine ray-finned fish, a marine angelfish belonging to the family Pomacanthidae. It is from the Eastern Pacific. It occasionally makes its way into the aquarium trade.



Pomacentrus coelestis

Pomacentrus coelestis

Pomacentrus coelestis, the neon damselfish, is a species of damselfish in the family Pomacentridae. It is found in the Indo-Pacific. It can grow to a maximum size of 9 centimetres (3.5 in) in length. It occasionally makes its way into the...



Pomacentrus moluccensis

Pomacentrus moluccensis

Pomacentrus moluccensis, the lemon damselfish, is a species of bony fish in the family Pomacentridae, from the Western Pacific Ocean. It occasionally makes its way into the aquarium trade. It grows to a size of 9 cm (4 in) in length.



Pomacentrus pavo

Pomacentrus pavo

Pomacentrus pavo, sapphire damsel, peacock damsel or blue damsel, is a damselfish from the Indo-Pacific. It occasionally makes its way into the aquarium trade. It grows to a size of 8.5 cm (3.3 in) in length.



Pomadasys commersonnii

Pomadasys commersonnii

Pomadasys commersonnii, the smallspotted grunter, is a species of ray-finned fish, a grunt belonging to the family Haemulidae. It is native to the brackish and marine waters of the western Indian Ocean.



Pomadasys kaakan

Pomadasys kaakan

Pomadasys kaakan, the javelin grunter or barred javelin is a species of marine ray-finned fish, a grunt belonging to the family Haemulidae. It is native to the Indian and Pacific Oceans, from Africa to Australia.



Pomadasys maculatus

Pomadasys maculatus

Pomadasys maculatus, commonly known as the saddle grunt, blotched grunt or blotched javelin, is a species of marine ray-finned fish, a grunt belonging to the family Haemulidae. It is native to the Indo-West Pacific region.



Pompano dolphinfish

Coryphaena equiselis

The pompano dolphinfish (Coryphaena equiselis) is a species of surface-dwelling ray-finned fish found in tropical and subtropical waters. They are one of only two members of the family Coryphaenidae, the other being the mahi-mahi or common...



Pond loach

Misgurnus anguillicaudatus

The pond loach (Misgurnus anguillicaudatus), also known as the Dojo loach, oriental weatherloach or oriental weatherfish, is a freshwater fish in the loach family Cobitidae. They are native to East Asia, but are also popular as an aquarium fish...



Pond smelt

Hypomesus olidus

The pond smelt (Hypomesus olidus) is a fresh and brackish water species of smelt. It is found in the East Asia (eastern Siberia, northeast China, Korea, Hokkaido) and the northwestern North America (Alaska, northwestern Canada). It can grow to 20...



Pondicherry shark

Carcharhinus hemiodon

The Pondicherry shark (Carcharhinus hemiodon) is an extremely rare species of requiem shark, in the family Carcharhinidae. A small and stocky gray shark, it grows not much longer than 1 m (3.3 ft) and has a fairly long, pointed snout. This...



Pontic shad

Alosa immaculata

The Pontic shad (Alosa immaculata, previously Alosa pontica), also referred to as the Black Sea shad or Kerch shad, is a species of clupeid fish in the genus Alosa, native to the Black Sea and Sea of Azov basins.



Ponticola kessleri

Ponticola kessleri

Ponticola kessleri, the bighead goby or Kessler's goby, is a species of goby native to Eurasia. The bighead goby is a Ponto-Caspian relict species. It inhabits the fresh and oligohaline waters, with mineralisation from 0-0.5‰ up to 1.5-3.0‰.



Pool barb

Puntius sophore

The pool barb, spotfin swamp barb, or stigma barb (Puntius sophore) is a tropical freshwater and brackish fish belonging to the Puntius genus in the family Cyprinidae. It is native to inland waters in Asia and is found in India, Nepal,...



Poor cod

Trisopterus minutus

The poor cod (Trisopterus minutus) is a temperate marine fish belonging to the cod family (Gadidae). It is red brown in colour and has a pronounced chin barbel. It may grow up to a length of 40 cm. It is usually found in small shoals at depths...



Popeye catalufa

Pristigenys serrula

The popeye catalufa (Pristigenys serrula), also known as the bigeye soldierfish, is a species of marine ray-finned fish in the family Priacanthidae, the bigeyes. This fish has an overall dusky orange to red colour with white markings. The dorsal...



Porae

Nemadactylus douglasii

The porae (Nemadactylus douglasii), the grey morwong, blue morwong, butterfish, Douglas' morwong, Eastern blue morwong, great perch, queen snapper, rubberlip morwong or silver morwong, is a species of marine ray-finned fish, traditionally...



Porbeagle

Lamna nasus

The porbeagle or porbeagle shark (Lamna nasus) is a species of mackerel shark in the family Lamnidae, distributed widely in the cold and temperate marine waters of the North Atlantic and Southern Hemisphere. In the North Pacific, its ecological...



Porcupine ray

Urogymnus asperrimus

The porcupine ray (Urogymnus asperrimus) is a rare species of stingray in the family Dasyatidae. This bottom-dweller is found throughout the tropical Indo-Pacific, as well as off West Africa. It favors sand, coral rubble, and seagrass habitats in...



Porcupine river stingray

Potamotrygon histrix

The porcupine river stingray (Potamotrygon histrix, sometimes incorrectly modified to Potamotrygon hystrix) is a species of river stingray in the family Potamotrygonidae, the type of the Potamotrygon genus. It is found in the basins of the Paraná...



Porichthys notatus

Porichthys notatus

Porichthys notatus is a species of batrachoid toadfish. It is a member of the midshipman genus, Porichthys, and is known by the common name plainfin midshipman. It is native to the eastern Pacific Ocean, where its distribution extends along the...



Port Jackson shark

Heterodontus portusjacksoni

The Port Jackson shark (*Heterodontus portusjacksoni*) is a nocturnal, oviparous (egg laying) type of bullhead shark of the family Heterodontidae, found in the coastal region of southern Australia, including the waters off Port Jackson. It has a...



Portuguese dogfish

Centroscymnus coelolepis

The Portuguese dogfish (*Centroscymnus coelolepis*) or Portuguese shark, is a species of sleeper shark of the family Somniosidae. This globally distributed species has been reported down to a depth of 3,675 m (12,057 ft), making it the...



Potamotrygon falkneri

Potamotrygon falkneri

Potamotrygon falkneri, the largespot river stingray or reticulated freshwater stingray, is a species of freshwater stingray in the family Potamotrygonidae from tropical and subtropical South America. It is sometimes seen in the aquarium trade,...



Potamotrygon rex

Potamotrygon rex

Potamotrygon rex, the great river stingray, is a species of freshwater stingray belonging to the family Potamotrygonidae, native to South America in the Neotropical region. P. rex, along with some other Potamotrygon species form a sub-species...



Potamotrygon tigrina

Potamotrygon tigrina

Potamotrygon tigrina, also known as the tiger river stingray, is a species of freshwater ray in the family Potamotrygonidae. This endangered species is endemic to black- and whitewater rivers in the upper Amazon basin in northeastern Peru. It is...



Potato grouper

Epinephelus tukula

The potato grouper (Epinephelus tukula), also called the potato cod or potato bass, is a species of marine ray-finned fish, a grouper from the subfamily Epinephelinae which is part of the family Serranidae, which also includes the anthias and sea...



Potosi pupfish

Cyprinodon alvarezii

Potosi pupfish (Cyprinodon alvarezii) is a species of fish in the family Cyprinodontidae. It is endemic to Mexico, but is now extinct in the wild, only surviving in captivity.



Powan

Coregonus clupeoides

The powan (Coregonus clupeoides) is a kind of freshwater whitefish endemic to two lochs in Scotland, Loch Lomond and Loch Eck. It has been successfully introduced in two other sites, Loch Sloy and the Carron Valley Reservoir. Powan populations are...



Predatory carp

Chanodichthys erythropterus

The predatory carp (Chanodichthys erythropterus), also known as the redfin culter or skygazer, is a species of ray-finned fish in the genus Chanodichthys. This East Asian freshwater cyprinid ranges from the Amur River south to Taiwan and the Red...



Priacanthus blochii

Priacanthus blochii

Priacanthus blochii, the paeony bulleye, is a species of marine ray finned fish, a bigeye in the family Priacanthidae. Other common names for this species include Bloch's bigeye, blotched bigeye, glass-eye bigeye, shortfin bigeye, silver big-eye...



Priacanthus hamrur

Priacanthus hamrur

Priacanthus hamrur, the lunar-tailed bigeye, goggle eye, or moontail bullseye, is a species of marine fishes belonging to the family Priacanthidae.



Priacanthus meeki

Priacanthus meeki

Priacanthus meeki is a species of marine ray finned fish, a bigeye in the family Priacanthidae. It is a red fish found in the Hawaiian and Midway Islands. It grows to a size of 33 cm in length. Common names are Hawaiian bigeye in English and ula...



Priacanthus tayenus

Priacanthus tayenus

Priacanthus tayenus, commonly known as purple-spotted bigeye, is a species of marine ray finned fish, a bigeye in the family Priacanthidae. It is native to the Indian and Pacific Oceans.



Prickly anglerfish

Himantolophus appeli

The prickly anglerfish (Himantolophus appeli) is a footballfish of the family Himantolophidae, found around the world in the southern oceans (apart from eastern Pacific), in deep water. Its length is up to 40 cm (16 in). It is a mesopelagic...



Prickly dogfish

Oxynotus bruniensis

The prickly dogfish (Oxynotus bruniensis) is a poorly known species of dogfish shark in the family Oxynotidae, inhabiting temperate Australian and New Zealand waters. Reaching a length of 75 cm (30 in), this brown to gray shark has a very thick...



Prickly sculpin

Cottus asper

The prickly sculpin (Cottus asper) is a species of ray-finned fish belonging to the family Cottidae, the typical sculpins. It is native to the river drainages of the Pacific Slope of North America from Seward, Alaska south to the Ventura River of...



Prickly shark

Echinorhinus cookei

The prickly shark (Echinorhinus cookei) is one of the two species of sharks in the family Echinorhinidae (the other one is the bramble shark), found in the Pacific Ocean over continental and insular shelves and slopes, and in submarine canyons....



Prietella phreatophila

Prietella phreatophila

The Mexican blindcat (Prietella phreatophila), in Spanish bagre de muzquiz, is a species of North American freshwater catfish (family Ictaluridae). Until recently, it was believed to be endemic to Coahuila in the Rio Bravo drainage in northern...



Princess parrotfish

Scarus taeniopterus

The princess parrotfish (Scarus taeniopterus) is a species of marine ray-finned fish, a parrotfish, in the family Scaridae. It is typically 20 to 25 centimetres (7.9 to 9.8 in) long, found in the Caribbean, South Florida, the Bahamas, and...



Prionotus carolinus

Prionotus carolinus

Prionotus carolinus, the northern sea robin, is a species of marine ray-finned fish belonging to the family Triglidae, the sea robins. This fish is found in the western Atlantic Ocean.



Prionotus evolans

Prionotus evolans

Prionotus evolans, the striped searobin, is a species of marine ray-finned fish belonging to the family Triglidae, the sea robins. This fish is found in the western Atlantic Ocean.



Prionurus punctatus

Prionurus punctatus

Prionurus punctatus, the yellowtail surgeonfish, is a species of marine ray-finned fish belonging to the family Acanthuridae, the surgeonfishes, unicornfishes and tangs. This fish is found in the eastern central Pacific Ocean.



Pristella maxillaris

Pristella maxillaris

Pristella maxillaris is a species of characin in the genus Pristella, and is commonly known as the X-ray fish or X-ray tetra because of its translucent body. It is a widely distributed and adaptable fish, found in the Amazon and Orinoco basins,...



Pristigenys alta

Pristigenys alta

Pristigenys alta, the toro or short bigeye, is a species of fish in the family Priacanthidae. Some anglers refer to this fish as "toro snapper", but it is not a snapper, and only distantly related to the fish of the snapper family.



Pristiophorus nancyae

Pristiophorus nancyae

Sharks portalPristiophorus nancyae, the African dwarf sawshark or dwarf sawshark, is a sawshark of the family Pristiophoridae. The species was discovered in 2011 when a specimen was caught off the coast of Mozambique at a depth of 1,600 ft (490 m).



Pristipomoides filamentosus

Pristipomoides filamentosus

Pristipomoides filamentosus, also known as the crimson jobfish, rosy snapper, bluespot jobfish, crimson snapper, king emperor, king snapper or rosy jobfish, is a species of ray-finned fish, a snapper belonging to the family Lutjanidae. It is...



Pristipomoides multidens

Pristipomoides multidens

Pristipomoides multidens, the goldbanded jobfish or goldbanded snapper, is a species of ray-finned fish, a snapper belonging to the family Lutjanidae. It is found in the Indian and Pacific Oceans.



Pristolepis fasciata

Pristolepis fasciata

Pristolepis fasciatus, commonly known as the Malayan leaffish, is a fish in the family Pristolepididae. It lives in the Mekong and Chao Phraya basins in Thailand, Cambodia, Vietnam, Myanmar, the Malay Peninsula, Sumatra, and Borneo, possibly also...



Prochilodus lineatus

Prochilodus lineatus

Prochilodus lineatus, the streaked prochilod, is a species of ray-finned fish in the family Prochilodontidae. It is native to the Paraná—Paraguay and Paraíba do Sul river basins in South America. It performs long breeding migrations and supports...



Prochilodus magdalenae

Prochilodus magdalenae

Prochilodus magdalenae is a tropical prochilodontid freshwater fish from Colombia. It is found in the Atrato, Sinú, Cauca and Magdalena Rivers. It has been measured to reach 36 cm (14 in) in standard length and 905 g (1.995 lb) in weight. They...



Protanguilla

Protanguilla palau

Protanguilla palau is a species of eel, the only species in the genus Protanguilla (first eel), which is in turn the only genus in its family, Protanguillidae. Individuals were found swimming in March 2010 in a deep underwater cave in a fringing...



Protomelas taeniolatus

Protomelas taeniolatus

Protomelas taeniolatus (often sold under various trade names: Haplochromis red empress, red empress or spindle hap) is a Haplochromine cichlid endemic to Lake Malawi in Eastern Africa. The fish is popular in the aquarium hobby due to the bright...



Prowfish

Zaprora silenus

The prowfish (Zaprora silenus) is a species of scorpaeniform marine fish found in the northern Pacific Ocean. It is the only extant member of the family, Zaproridae. Another species, Araeosteus rothi, is known from Late Miocene marine strata in...



Prussian carp

Carassius gibelio

The Prussian carp, silver Prussian carp or Gibel carp (Carassius gibelio) is a member of the family Cyprinidae, which includes many other fish, such as the common carp, goldfish, and the smaller minnows. It is a medium-sized cyprinid, and does...



Pseudacanthicus leopardus

Pseudacanthicus leopardus

Pseudacanthicus leopardus is a species of armored catfish endemic to Guyana where it occurs in the Rupununi River basin. This species grows to a length of 15 centimetres (5.9 in) TL.



Pseudacanthicus spinosus

Pseudacanthicus spinosus

Pseudacanthicus spinosus is a species of armored catfish endemic to Brazil where it occurs in the Amazon Basin. This species grows to a length of 26 centimetres (10 in) SL.



Pseudanthias bartlettorum

Pseudanthias bartlettorum

Pseudanthias bartlettorum, Bartlett's anthias is a species of marine ray-finned fish in the subfamily Anthiinae of the family Serranidae, the groupers and sea basses. It occurs in the Pacific Ocean. This fish is sometimes kept in aquaria.



Pseudanthias bicolor

Pseudanthias bicolor

Pseudanthias bicolor, the bicolor anthias or yellowback basslet is a species of marine ray-finned fish from the subfamily Anthiinae of the family Serranidae, the groupers and sea basses. It is from the Indo-Pacific Ocean. It occasionally makes...



Pseudanthias hypselosoma

Pseudanthias hypselosoma

Pseudanthias hypselosoma is a species of marine ray-finned fish, a member of the genus Pseudanthias which is part of the subfamily Anthiinae, which in turn is part of the family Serranidae, the groupers and sea basses. It is an Indo-Pacific...



Pseudanthias pleurotaenia

Pseudanthias pleurotaenia

Pseudanthias pleurotaenia also known as the squarespot anthias, pink square anthias, mirror basslet or squarespot fairy basslet is a species of marine ray-finned fish in the subfamily Anthiinae of the family Serranidae, the groupers and sea...



Pseudobalistes fuscus

Pseudobalistes fuscus

Pseudobalistes fuscus (common names: blue triggerfish, rippled triggerfish, yellow-spotted triggerfish and blue-and-gold triggerfish) is a fish belonging to the family Balistidae.



Pseudochaenichthys

Pseudochaenichthys georgianus

Pseudochaenichthys is a monotypic genus of marine ray-finned fish belonging to the family Channichthyidae, the crocodile icefishes. Its only member is Pseudochaenichthys georgianus, the South Georgia icefish, which is found in the Southern Ocean.



Pseudochromis aldabraensis

Pseudochromis aldabraensis

Pseudochromis aldabraensis, the orange dottyback, neon dottyback or Arabian dottyback, is a species of ray-finned fish from the family Pseudochromidae, the dottybacks. It is commonly kept in marine aquariums.



Pseudochromis fridmani

Pseudochromis fridmani

Pseudochromis fridmani is a species of fish in the family Pseudochromidae, the dottybacks. Its common name is orchid dottyback. It is endemic to the Red Sea. This fish is up to 6.3 centimeters long. It lives in tropical marine waters up to 60...



Pseudocyttus

Pseudocyttus maculatus

Pseudocyttus maculatus, the smooth oreo or smooth dory, is an oreo, the only species in the genus Pseudocyttus. It is found in all southern oceans at depths of between 400 and 1,500 metres (1,300 and 4,900 ft). Its length is up to 60 centimetres...



Pseudohemiodon apithanos

Pseudohemiodon apithanos

Pseudohemiodon apithanos is a species of armored catfish endemic to Ecuador where it occurs in the San Miguel River basin. (Kasaset) This species grows to a length of 14.5 centimetres (5.7 in) SL.



Pseudoliparis amblystomopsis

Pseudoliparis amblystomopsis

Pseudoliparis amblystomopsis, or the hadal snailfish, is a species of snailfish from the hadal zone of the Northwest Pacific Ocean, including the Kuril–Kamchatka and Japan Trenches. In October 2008, a team from British and Japanese institutes...



Pseudoliparis swirei

Pseudoliparis swirei

Pseudoliparis swirei, the Mariana snailfish or Mariana hadal snailfish, is a species of snailfish found at hadal depths in the Mariana Trench in the western Pacific Ocean. It is known from a depth range of 6,198–8,076 m (20,335–26,496 ft),...



Pseudoplatystoma corruscans

Pseudoplatystoma corruscans

Pseudoplatystoma corruscans, the spotted sorubim, is a species of long-whiskered catfish native to the São Francisco and Paraná—Paraguay River basins in South America. This species grows to a length of 166 centimetres (65 in) TL.



Pseudoplatystoma fasciatum

Pseudoplatystoma fasciatum

Pseudoplatystoma fasciatum or barred sorubim or barred catfish is a species of long-whiskered catfish native to the Suriname, Corantijn and Essequibo. The nocturnal predator feeds mainly on other fish and crabs. Females reach a more notable size....



Pseudoplatystoma tigrinum

Pseudoplatystoma tigrinum

Pseudoplatystoma tigrinum, the tiger sorubim, tiger shovelnose or caparari is a species of long-whiskered catfish native to the Amazon Basin in South America. The largest individuals of this species grows to 130 cm (51 in) in total length....



Pseudorhombus arsius

Pseudorhombus arsius

Pseudorhombus arsius, the largetooth flounder, is a species of left-eyed flatfish, that is the dark side with the eyes on the adult fish is the left side of the fish's body, from the family Paralichthyidae. As Rhombus polyspiros it was named as...



Pseudorinelepis

Pseudorinelepis genibarbis

Pseudorinelepis genibarbis is a species of armored catfish native to Brazil, Peru, and Ecuador where it is found in the Amazon basin.



Pseudosphromenus cupanus

Pseudosphromenus cupanus

Pseudosphromenus cupanus, also known as the spiketail paradisefish is a species of medium-small gouramies native to southern India and Sri Lanka. This species is often found in shallow, slow-moving or stagnant water, such as ditches and paddy...



Pseudotropheus crabro

Pseudotropheus crabro

Pseudotropheus crabro, the Bumblebee Cichlid or Hornet Cichlid, is a species of cichlid endemic to Lake Malawi where it is found in different habitats but most frequently in large caves or in the vicinity of large boulders. This species can reach...



Pseudotropheus cyaneorhabdos

Pseudotropheus cyaneorhabdos

Pseudotropheus cyaneorhabdos is a species of cichlid endemic to Lake Malawi where it is only known from around Likoma Island where it prefers rocky substrates at depths of from 5 to 10 metres (16 to 33 ft). This species can grow to a length of...



Pseudotropheus demasoni

Pseudotropheus demasoni

Pseudotropheus demasoni is a species of cichlid endemic to Lake Malawi where it is only known from the Pombo Rocks in Tanzanian waters. This species can potentially reach a maximum length of 10 centimetres (3.9 in) SL. It is now commonly found in...



Pseudotropheus johannii

Pseudotropheus johannii

Pseudotropheus johannii or the bluegray mbuna is an African freshwater fish from the family Cichlidae.



Pseudotropheus saulosi

Pseudotropheus saulosi

Pseudotropheus saulosi is a species of cichlid endemic to Lake Malawi in East Africa, where it lives in areas with rocky substrates. It is classified as a dwarf-mbuna and was first described by Ad Konings in 1990, who gave it the specific name...



Psilorhynchus sucatio

Psilorhynchus sucatio

Psilorhynchus sucatio, the river stone carp or sucatio minnow, is freshwater ray-finned fish a species of torrent minnow. The specific name is a latinization of one of the local names for this species, sukati. It is a widely distributed species...



Psychedelic frogfish

Histiophryne psychedelica

The psychedelic frogfish (Histiophryne psychedelica) is a yellow-brown or peach colored frogfish named for its pink and white stripes arranged in a fingerprint pattern. The fish is from waters near Ambon Island and Bali, Indonesia.



Psychrolutes marcidus

Psychrolutes marcidus

Psychrolutes marcidus, the smooth-head blobfish, also known simply as blobfish, is a deep-sea fish of the family Psychrolutidae. It inhabits the deep waters off the coasts of mainland Australia and Tasmania, as well as the waters of New...



Psychrolutes microporos

Psychrolutes microporos

Psychrolutes microporos is a species of deepwater marine fish in the family Psychrolutidae, commonly known as a blobfish or fathead. It is found in the abyssal zone in waters around Australia and New Zealand. A photograph of an individual taken...



Psychrolutes occidentalis

Psychrolutes occidentalis

Psychrolutes occidentalis, also known as the Western Australian sculpin or western blobfish, is a species of deep-sea fish of the family Psychrolutidae. It is a bathydemersal fish found along the continental slope to the west of Australia.



Ptereleotris evides

Ptereleotris evides

Ptereleotris evides, the Blackfin dartfish or scissortail goby, is a species of dartfish native to the Indian Ocean and the western Pacific Ocean. It is a reef inhabitant and can be found at depths of from 2 to 15 metres (6.6 to 49.2 ft). This...



Pterodoras granulosus

Pterodoras granulosus

The granulated catfish (Pterodoras granulosus) is a species of thorny catfish found in the Paraná and Amazon basin as well as the coastal drainages of Suriname and Guyana. This species is commercially caught for human consumption as well as being...



Pteroidichthys amboinensis

Pteroidichthys amboinensis

Pteroidichthys amboinensis, the Ambon scorpionfish or Godfrey's scorpionfish, is a species of marine ray-finned fish belonging to the family Scorpaenidae, the scorpionfishes. It is found in the Indian and Pacific oceans.



Pterois antennata

Pterois antennata

Pterois antennata, the spotfin lionfish, banded lionfish, broadbarred lionfish, broadbarred firefish, raggedfinned firefish, raggedfinned scorpionfish or roughscaled lionfish, is a species of marine ray-finned fish belonging to the family...



Pterois miles

Pterois miles

Pterois miles, the devil firefish or common lionfish, is a species of ray-finned fish native to the western Indo-Pacific region. It is frequently confused with its close relative, the red lionfish (Pterois volitans). The scientific name is from...



Pterois mombasae

Pterois mombasae

Pterois mombasae, the African lionfish, deepwater firefish or frillfin turkeyfish, is a species of marine ray-finned fish belonging to the family Scorpaenidae, the scorpionfishes and lionfishes. It is found in the tropical Indian Ocean, typically...



Pterois russelii

Pterois russelii

Pterois russelii, the largetail turkeyfish, plaintail firefish, plaintail turkeyfish, Russell's firefish, Russell's lionfish, spotless butterfly-cod or the spotless firefish, is a species of ray-finned fish with venomous spines belonging to the...



Pterois sphex

Pterois sphex

Pterois sphex, the Hawaiian turkeyfish or Hawaiian lionfish is a species of ray-finned fish with venomous spines belonging to the family Scorpaenidae, the scorpionfishes and lionfishes. It is found in the eastern Central Pacific, specifically in...



Pterophyllum altum

Pterophyllum altum

Pterophyllum altum, also referred to as the altum angelfish, deep angelfish, or Orinoco angelfish, occurs strictly in the Orinoco River Basin and the Upper Rio Negro watershed in Southern Venezuela, Southeastern Colombia and extreme Northern Brazil.



Pterophyllum leopoldi

Pterophyllum leopoldi

Pterophyllum leopoldi, also referred to as the teardrop angelfish, Leopold's angelfish dwarf angelfish, or roman-nosed angelfish, is an angelfish species native to the Amazon River (between Manacapuru and Santarém), Essequibo River and Rupununi...



Pterophyllum scalare

Pterophyllum scalare

Pterophyllum scalare, most commonly referred to as angelfish or freshwater angelfish, is the most common species of Pterophyllum kept in captivity. It is native to the Amazon Basin in Peru, Colombia, and Brazil. Particularly to the Ucayali river...



Pterygoplichthys gibbiceps

Pterygoplichthys gibbiceps

Pterygoplichthys gibbiceps is a species of armored catfish native to Brazil, Ecuador, Peru and Venezuela where it is found in the Orinoco and Amazon basins. Pterygoplichthys gibbiceps shows all the characteristic features of its genus—a large...



Pterygoplichthys joselimaianus

Pterygoplichthys joselimaianus

Pterygoplichthys joselimaianus is a species of armored catfish endemic to Brazil where it occurs in the Tocantins River basin. It grows to a length of 30.5 centimetres (12.0 in) SL.



Pterygoplichthys multiradiatus

Pterygoplichthys multiradiatus

Pterygoplichthys multiradiatus is one of several tropical fish commonly known as Orinoco sailfin catfish, plecostomus (or plecos). It belongs to the armored catfish family (Loricariidae). Named for its sail-like dorsal fin, the part of its...



Pterygoplichthys pardalis

Pterygoplichthys pardalis

Pterygoplichthys pardalis, the Amazon sailfin catfish, is a freshwater tropical fish in the armored catfish family (Loricariidae). It is one of a number of species commonly referred to as the common pleco or "leopard pleco" by aquarists. It can be...



Ptychochromis insolitus

Ptychochromis insolitus

Ptychochromis insolitus, also known as the Mangarahara cichlid or joba mena, is a species of cichlid endemic to certain river systems in northern Madagascar. This critically endangered fish is threatened by habitat loss and competition from...



Puddingwife wrasse

Halichoeres radiatus

The puddingwife wrasse, *Halichoeres radiatus*, is a species of wrasse native to the western Atlantic Ocean from North Carolina to Bermuda, through the West Indies and Gulf of Mexico, to offshore islands of Brazil, being absent from Brazilian...



Puffadder shyshark

Haploblepharus edwardsii

The puffadder shyshark (*Haploblepharus edwardsii*), also known as the Happy Eddie, is a species of catshark, belonging to the family Scyliorhinidae, endemic to the temperate waters off the coast of South Africa. This common shark is found on or...



Pumpkinseed

Lepomis gibbosus

The pumpkinseed (*Lepomis gibbosus*), also referred to as pond perch, common sunfish, punkie, sunfish, sunny, and kivver, is a small/medium-sized North American freshwater fish of the genus *Lepomis* (true sunfishes), from family Centrarchidae...



Pungtungia herzi

Pungtungia herzi

Pungtungia herzi is a species of cyprinid fish found in Japan and the Korean Peninsula. Named in honor of German entomologist Alfred Otto Herz (1856–1905), who collected the type specimen. It has been observed to deposit its eggs into nests of...



Puntigrus anchisporus

Puntigrus anchisporus

Puntigrus anchisporus is a species of cyprinid fish endemic to Borneo. This species can grow to a length of 6.6 centimetres (2.6 in) TL.



Puntigrus partipentazona

Puntigrus partipentazona

Puntigrus partipentazona, the Dwarf Tiger Barb, is a species of cyprinid fish native to Southeast Asia where it is found in the Mekong, Mae Klong, and Chao Phraya basins of Thailand, the Malay Peninsula, and coastal streams of southeast Thailand...



Pycnochromis iomelas

Pycnochromis iomelas

Pycnochromis iomelas , also known as the half and half chromis, is a species of reef dwelling fish in the family Pomacentridae. It is occasionally seen for sale in the aquarium trade.



Pygmy corydoras

Corydoras pygmaeus

Corydoras pygmaeus, or the pygmy corydoras or pygmy catfish is a tropical and freshwater fish belonging to the subfamily Corydoradinae of the family Callichthyidae. It originates in tropical inland waters in South America, and is found in the...



Pygmy gourami

Trichopsis pumila

The pygmy gourami (Trichopsis pumila), also known as the sparkling gourami, is a freshwater species of gourami native to Southeast Asia.



Pygmy ribbontail catshark

Eridacnis radcliffei

The pygmy ribbontail catshark (Eridacnis radcliffei) is a species of finback catshark, family Proscylliidae, distributed patchily in the western Indo-Pacific from Tanzania to the Philippines. It occurs around the edges of continental and insular...



Pygmy shark

Euprotomicros bispinatus

Sharks portalThe pygmy shark (Euprotomicros bispinatus), the second-smallest of all the shark species after the dwarf lanternshark, is a squaliform shark of the family Dalatiidae, the only member of the genus Euprotomicros. Their lengths are up...



Pygmy whitefish

Prosopium coulterii

The pygmy whitefish (Prosopium coulterii) is a freshwater whitefish of the genus Prosopium in the family Salmonidae. Found in the mountain streams and lakes in western North America, it also has isolated populations in Lake Superior and in...



Pygocentrus cariba

Pygocentrus cariba

Pygocentrus cariba or black spot piranha is a species of piranha native to the Orinoco River basin lowlands and the Llanos region in Venezuela and Colombia. This species can reach a total length of 27.9 centimetres (11.0 in). It is popular as a...



Pygocentrus palometa

Pygocentrus palometa

Pygocentrus palometa is a species of piranha native to South America, where restricted to the Orinoco Basin. Although recognized by FishBase, the scientific name may be a nomen dubium."Palometa" is a general common name used in South America for...



Pygocentrus piraya

Pygocentrus piraya

Pygocentrus piraya, often called the piraya piranha or San Francisco piranha, and sometimes sold as the man-eating piranha, is a large, aggressive piranha from the São Francisco River basin in Brazil.



Pyjama shark

Poroderma africanum

The pyjama shark or striped catshark (Poroderma africanum) is a species of catshark, and part of the family Scyliorhinidae, endemic to the coastal waters of South Africa. This abundant, bottom-dwelling species can be found from the intertidal...



Pyramid butterflyfish

Hemitaurichthys polylepis

The pyramid butterflyfish (Hemitaurichthys polylepis) is a species of marine ray-finned fish, a butterflyfish belonging to the family Chaetodontidae, native from central Indo-Pacific.



Queen angelfish

Holacanthus ciliaris

The queen angelfish (Holacanthus ciliaris), also known as the blue angelfish, golden angelfish, or yellow angelfish, is a species of marine angelfish found in the western Atlantic Ocean. It is a benthic (ocean floor) warm-water species that lives...



Queen coris

Coris formosa

The queen coris (Coris formosa), also known as the sand wrasse, Queen coris wrasse, Formosan coris, Formosa coris wrasse, Formosa wrasse, Indian Ocean wrasse or Indian sand wrasse, is a species of wrasse native to the Indian Ocean from the Red...



Queen parrotfish

Scarus vetula

The queen parrotfish (Scarus vetula) is a species of marine ray-finned fish, a parrotfish, in the family Scaridae. It is found on reefs in the tropical West Atlantic Ocean and the Caribbean Sea. Other common names include blownose, blue chub,...



Queen snapper

Etelis oculatus

The queen snapper (Etelis oculatus), also known as the night snapper or brim snapper, is a species of ray-finned fish, a snapper belonging to the family Lutjanidae. It is native to the western Atlantic Ocean, and is the only species in the genus...



Queenfish

Seriphus politus

The queenfish (Seriphus politus) is a species of fish in the family Sciaenidae, the drums and croakers. It is native to the eastern Pacific Ocean, where it occurs along the North American coastline from Oregon to Baja California; it has been...



Queensland school mackerel

Scomberomorus queenslandicus

The Queensland school mackerel (Scomberomorus queenslandicus) is a species of fish in the family Scombridae. It is found in tropical waters of the western Pacific, largely confined to inshore coastal waters of southern Papua New Guinea and...



Quillback

Carpiodes cyprinus

The quillback (Carpiodes cyprinus), also known as the quillback carpsucker, is a type of freshwater fish of the sucker family widely distributed throughout North America. It is deeper-bodied than most suckers, leading to a carplike appearance....



Quillback rockfish

Sebastes maliger

The quillback rockfish (Sebastes maliger), also known as the quillback seaperch, is a species of marine ray-finned fish belonging to the subfamily Sebastinae, the rockfishes, part of the family Scorpaenidae. This species primarily dwells in salt...



Quillfish

Ptilichthys goodei

The quillfish, (Ptilichthys goodei), is a species of marine ray-finned fish, it is the only species in the genus Ptilichthys and family Ptilichthyidae. This fish occurs in the northern North Pacific Ocean.



Rabbit fish

Chimaera monstrosa

Chimaera monstrosa, also known as the rabbit fish or rat fish, is a northeast Atlantic and Mediterranean species of cartilaginous fish in the family Chimaeridae. The rabbit fish is known for its characteristically large head and small, tapering...



Raccoon butterflyfish

Chaetodon lunula

The raccoon butterflyfish (Chaetodon lunula), also known as the crescent-masked butterflyfish, lunule butterflyfish, halfmoon butterflyfish, moon butterflyfish, raccoon butterfly, raccoon, raccoon coralfish, and redstriped butterflyfish, is a...



Racer goby

Babka gymnotrachelus

The racer goby (Babka gymnotrachelus) is a species of goby native to fresh, sometimes brackish, waters, of the Black Sea basin. It is a Ponto-Caspian relict species. The species is placed a monotypic genus, Babka, which was once considered a...



Ragfish

Icosteus aenigmaticus

The ragfish (Icosteus aenigmaticus) is a ray-finned fish of the northern Pacific Ocean; although a perciform, its skeleton is mostly cartilage, and the larvae have pelvic fins that disappear as they mature. It is the sole member of the family...



Rainbow cichlid

Herotilapia multispinosa

Herotilapia multispinosa (previously: Archocentrus multispinosus) also known as the rainbow cichlid is a Central American freshwater fish of the cichlid family. It is found on the Atlantic slope of Honduras, Nicaragua, and Costa Rica from Patuca...



Rainbow darter

Etheostoma caeruleum

The rainbow darter (Etheostoma caeruleum) is a small species of freshwater ray-finned fish, a darter from the subfamily Etheostomatinae, part of the family Percidae, which also contains the perches, ruffes and pikeperches. It is native to North...



Rainbow parrotfish

Scarus guacamaia

The rainbow parrotfish (Scarus guacamaia) is a species of fish in the family Scaridae.



Rainbow runner

Elagatis bipinnulata

The rainbow runner (*Elagatis bipinnulata*), also known as the rainbow yellowtail, Spanish jack and Hawaiian salmon, is a common species of pelagic marine fish of the jack family, Carangidae. The species is widespread throughout the tropical and...



Rainbow sardine

Dussumieria acuta

The rainbow sardine (*Dussumieria acuta*), also known as common sprat, dwarf round herring, rainbow herring, and sharpnosed sprat, is a bony fish important to aquaculture and commercial fisheries.



Rainbow shark

Epalzeorhynchus frenatum

The rainbow shark (*Epalzeorhynchus frenatum*) is a species of Southeast Asian freshwater fish from the family Cyprinidae. It is also known as the ruby shark, red-fin shark, red-finned shark, rainbow sharkminnow, green fringelip labeo, whitefin...



Rainbow shiner

Notropis chrosomus

The rainbow shiner (*Notropis chrosomus*) is a North American species of ray-finned fish in the genus *Notropis*. The rainbow shiner has a length of 5 to 8 centimeters. It has translucent color from pink to golden with a silver-black stripe along its...



Rainbow smelt

Osmerus mordax

The rainbow smelt (*Osmerus mordax*) is a North American species of fish of the family Osmeridae. Walleye, trout, and other larger fish prey on these smelt. The rainbow smelt prefer juvenile ciscoes, zooplankton such as calanoid copepods...



Rainbow trout

Oncorhynchus mykiss

The rainbow trout (Oncorhynchus mykiss) is a species of trout native to cold-water tributaries of the Pacific Ocean in Asia and North America. The steelhead (sometimes called steelhead trout) is an anadromous (sea-run) form of the coastal rainbow...



Rainwater killifish

Lucania parva

The rainwater killifish (Lucania parva) is a small silvery fish with yellow flashes and diamond shaped scales that is widespread from Cape Cod, Massachusetts, through to Tampico, Mexico. It is commonly found in large numbers in fresh to brackish...



Raitt's sand eel

Ammodytes marinus

Raitt's sand eel (Ammodytes marinus), also known as the lesser sand eel, is a small semi-pelagic ray-finned fish found in the North Atlantic Ocean. The Raitt's sand eel is member of the family Ammodytidae which includes all 31 species of sand...



Raja radula

Raja radula

The rough ray (Raja radula) is a Mediterranean ray species of the Rajidae family described by Delaroche in 1809.



Ram cichlid

Mikrogeophagus ramirezi

The ram cichlid, Mikrogeophagus ramirezi, is a species of freshwater fish endemic to the Orinoco River basin, in the savannahs of Venezuela and Colombia in South America. The species has been examined in studies on fish behaviour and is a popular...



Randall's prawn goby

Amblyeleotris randalli

Amblyeleotris randalli, Randall's prawn goby, is a marine benthic species of goby native to tropical reefs of the central Indo-Pacific. This species can also be found in the aquarium trade.



Raniceps raninus

Raniceps raninus

Raniceps raninus, the tadpole fish, is a species of Gadidae fish native to the northeast Atlantic Ocean around the coasts of France, Ireland, and the United Kingdom and the North Sea. This species grows to a total length of 27.5 cm (10.8 in). It...



Rasbora sarawakensis

Rasbora sarawakensis

Rasbora sarawakensis, also known as the blue line rasbora or Sarawak rasbora, is a species of ray-finned fish in the genus Rasbora which is endemic to the island of Borneo.



Rasbora trilineata

Rasbora trilineata

Rasbora trilineata is a species of ray-finned fish in the genus Rasbora. Common names include scissortail rasbora and three-lined rasbora. It comes from Southeast Asia. Its length is up to 15 cm (6").



Razorback sucker

Xyrauchen texanus

The razorback sucker (Xyrauchen texanus) is a suckerfish found in rivers and lakes in the southwestern United States and formerly northwestern Mexico. It can grow to 91 cm (3 ft) in length and is recognisable by the keel between its head and...



Reba carp

Cirrhinus reba

The Reba carp (Malayalam: കാവേരിക്കുഴി) (Cirrhinus reba) is a species of ray-finned fish in the genus Cirrhinus. This freshwater edible fish is found in large streams, rivers, tanks, lakes, reservoirs. It is native to Bangladesh, India, Nepal,...



Red codling

Pseudophycis bachus

The red codling or hoka (*Pseudophycis bachus*) is a morid cod of the genus *Pseudophycis*, restricted to New Zealand, from the surface to 700 m. A closely related species is found in Australia. Its length is up to 90 cm. *P. bachus* is a food source...



Red cornetfish

Fistularia petimba

The red cornetfish (*Fistularia petimba*), also known as the rough flutemouth, is a cornetfish of the family *Fistulariidae*, found in subtropical and tropical oceans worldwide, at depths between 10 m (33 ft) and 200 metres (660 ft). They are up to 2...



Red dragonet

Foetorepus altivelis

The red dragonet (Foetorepus altivelis) is a species of dragonet native to the Indian Ocean and the western Pacific Ocean where it occurs at depths of from 70 to 600 metres (230 to 1,970 ft). Males of this species reaches a length of 17...



Red drum

Sciaenops ocellatus

The red drum (Sciaenops ocellatus), also known as redfish, channel bass, puppy drum, spottail bass, or simply red, is a game fish found in the Atlantic Ocean from Massachusetts to Florida and in the Gulf of Mexico from Florida to northern Mexico....



Red dwarf rasbora

Microrasbora rubescens

Red dwarf rasbora (Microrasbora rubescens) is a species of cyprinid found endemic to Lake Inle in Shan State in Myanmar. It belongs to the genus Microrasbora, which contains two small species of danionins.



Red garra

Garra rufa

The red garra (Garra rufa), also known as the doctor fish or nibble fish, is a species of cyprinid that is native to a wide range of freshwater habitats in subtropical parts of Western Asia. This small fish typically is up to about 14 centimeters...



Red grouper

Epinephelus morio

The red grouper (Epinephelus morio) is a species of marine ray-finned fish, a grouper from the subfamily Epinephelinae which is part of the family Serranidae, which also includes the anthias and sea basses. It is found in the western Atlantic Ocean.



Red gurnard

Chelidonichthys cuculus

The red gurnard (Chelidonichthys cuculus), also known as the East Atlantic red gurnard or soldier, is a benthic species of ray-finned fish belonging to the family Triglidae, the gurnards and sea robins. This fish is found in the eastern Atlantic...



Red hake

Urophycis chuss

The red hake or squirrel hake fish, *Urophycis chuss*, is a species of phycid hake. It is found in the Atlantic Ocean at depths between 10 and 500 m. It grows to about 30 in (75 cm) and 7 lb (3.2 kg). Red hake are edible, and are sought out by...



Red hind

Epinephelus guttatus

The red hind (*Epinephelus guttatus*), also known as the koon or lucky grouper in Caribbean vernacular, is a species of marine ray-finned fish, a grouper from the subfamily Epinephelinae which is part of the family Serranidae, which also includes...



Red Irish lord

Hemilepidotus hemilepidotus

The red Irish lord (*Hemilepidotus hemilepidotus*) is a species of marine ray-finned fish in the family Agonidae. It is found in the northern Pacific Ocean, from Russia to Alaska and as far south as Monterey Bay. It is a distinctly red fish, with...



Red lionfish

Pterois volitans

The red lionfish (*Pterois volitans*) is a venomous coral reef fish in the family Scorpaenidae, order Scorpaeniformes. It is mainly native to the Indo-Pacific region, but has become an invasive species in the Caribbean Sea, as well as along the...



Red moki

Cheilodactylus spectabilis

The red moki (*Cheilodactylus spectabilis*) also known as the banded morwong, brown-banded morwong, carp or nanua is a species of marine ray-finned fish, traditionally regarded as belonging to the family Cheilodactylidae, commonly referred to as...



Red morwong

Cheilodactylus fuscus

The red morwong (Cheilodactylus (Morwong) fuscus), also known as the sea carp, is a species of marine ray finned fish traditionally regarded as belonging to the family Cheilodactylidae, the members of which are commonly known as morwongs. It is...



Red neon blue-eye

Pseudomugil luminatus

The red neon blue-eye (Pseudomugil luminatus) is a species of fish in the subfamily Pseudomugilinae. It has only been recorded from swamps in the vicinity of Timika in Papua, Indonesia.



Red pigfish

Bodianus unimaculatus

The red pigfish (Bodianus unimaculatus), also known as the red hogfish or eastern pigfish, is a species of wrasse native to the southwestern Pacific Ocean from eastern Australia to New Zealand, including Lord Howe Island, Norfolk Island, and the...



Red rainbowfish

Glossolepis incisus

The red rainbowfish or salmon-red rainbowfish (Glossolepis incisa) is a species of rainbowfish from Lake Sentani in Irian Jaya, Indonesia. Belonging to the family Melanotaeniidae, in the subfamily Melanotaeniinae, the Australian rainbowfishes. It...



Red Sea bannerfish

Heniochus intermedius

The Red Sea bannerfish (Heniochus intermedius) is a species of marine ray-finned fish, a butterflyfish from the family Chaetodontidae. It is found in the western Indian Ocean. It has been recorded as an introduced species off Florida and as a...



Red Sea clownfish

Amphiprion bicinctus

The Red Sea Clownfish (Amphiprion bicinctus, meaning "both sawlike with two stripes"), commonly known as the Red Sea or two-banded anemonefish is a marine fish belonging to the family Pomacentridae, the clownfishes and damselfishes. Like other...



Red shiner

Cyprinella lutrensis

The red shiner or red-horse minnow (Cyprinella lutrensis) is a North American species of freshwater fish in the family Cyprinidae. They are deep-bodied and laterally compressed, and can grow to about three inches in length. For most of the year,...



Red steenbras

Petrus rupestris

The red steenbras (Petrus rupestris) is a species of fish in the family Sparidae and the only known member of the monospecific genus Petrus. This species is endemic to South Africa. The species' population has been depleted by overfishing in...



Red stingray

Hemitygon akajei

The red stingray (Hemitygon akajei) is a species of stingray in the family Dasyatidae, found in the northwestern Pacific Ocean off Japan, Korea, and China, and possibly elsewhere. It primarily inhabits shallow, sandy habitats close to shore, and...



Red tailed tinfoil

Barbonymus altus

The red tailed tinfoil or red tailed tinfoil barb (Barbonymus altus) is a species of freshwater cyprinid fish from South-East Asia. It lives in the Mekong and Chao Phraya river basins.



Red-bellied piranha

Pygocentrus nattereri

The red-bellied piranha, also known as the red piranha (Pygocentrus nattereri), is a type of piranha native to South America, found in the Amazon, Paraguay, Paraná and Essequibo basins, as well as coastal rivers of northeastern Brazil. This fish...



Red-breasted wrasse

Cheilinus fasciatus

The red-breasted wrasse (Cheilinus fasciatus) is a species of wrasse native to the Indian Ocean and the western Pacific Ocean.



Red-finned blue-eye

Scaturiginichthys vermeilipinnis

The red-finned blue-eye (Scaturiginichthys vermeilipinnis) is a tiny, critically endangered species of fish in the family Pseudomugilidae. It is the only species in its genus. The species was first recorded in 1990. It is endemic to central...



Red-lipped batfish

Ogcocephalus darwini

The red-lipped batfish or Galápagos batfish (Ogcocephalus darwini) is a fish of unusual morphology found around the Galápagos Islands and off Peru at depths of 3 to 76 m (10 to 249 ft). Red-lipped batfish are closely related to rosy-lipped...



Red-tailed black shark

Epalzeorhynchos bicolor

The red-tailed black shark (Epalzeorhynchos bicolor; syn. Labeo bicolor), also known as the redtail shark and redtail sharkminnow, is a species of freshwater fish in the carp family, Cyprinidae. Despite its name, it is more closely related to...



Redbanded rockfish

Sebastes babcocki

The redbanded rockfish (Sebastes babcocki), also known as the bandit, barber pole, flag rockfish, Spanish flag, Hollywood, convict, and canary, is a species of marine ray-finned fish belonging to the subfamily Sebastinae, the rockfishes, part of...



Redbelly tilapia

Coptodon zillii

The redbelly tilapia (*Coptodon zillii*, syn. *Tilapia zillii*), also known as the Zille's redbreast tilapia or St. Peter's fish (a name also used for other tilapia in Israel), is a species of fish in the cichlid family. This fish is found widely in...



Redbreast sunfish

Lepomis auritus

The redbreast sunfish (*Lepomis auritus*) is a species of freshwater fish in the sunfish family (family Centrarchidae) of the order Perciformes. The type species of its genus, it is native to the river systems of eastern Canada and the United...



Redbreast tilapia

Coptodon rendalli

The redbreast tilapia (*Coptodon rendalli*) is a species of fish in the family Cichlidae. It is found widely in the southern half of Africa. Its natural habitats are freshwater lakes and freshwater marshes. It is known as the redbreast kurper in...



Redear sunfish

Lepomis microlophus

The redear sunfish (*Lepomis microlophus*), also known as the shellcracker, Georgia bream, cherry gill, chinquapin, improved bream, rouge ear sunfish and sun perch) is a freshwater fish in the family Centrarchidae and is native to the southeastern...



Redeye bass

Micropterus coosae

The redeye bass, redeye, or Coosa bass (*Micropterus coosae*) is a species of freshwater fish in the sunfish family (Centrarchidae) native to the Coosa River system of Georgia, Alabama. The waters it is normally found in are cool streams and rivers...



Redeye piranha

Serrasalmus rhombeus

The redeye piranha (Serrasalmus rhombeus), also known as the black piranha, white piranha, spotted piranha or yellow piranha, is a species of freshwater ray-finned fish, a piranha from the family Serrasalminae. It is found in northern South...



Redeye tetra

Moenkhausia sanctaefilomenae

The redeye tetra (Moenkhausia sanctaefilomenae), is a species of tetra from the São Francisco, upper Paraná, Paraguay and Uruguay river basins in eastern and central South America. This freshwater fish is commonly kept in aquariums and bred in...



Redfin bully

Gobiomorphus hutereaui

The redfin bully (Gobiomorphus hutereaui) is a species of freshwater fish in the family Eleotridae endemic to New Zealand. Being amphidromous, it spends part of its life cycle at sea. Males have distinctive bright red patterns and stripes on their...



Redfin pickerel

Esox americanus americanus

The redfin pickerel (Esox americanus americanus) is a subspecies of freshwater fish belonging to the pike family (Esocidae) of the order Esociformes. Not to be confused with its close relatives, the grass pickerel and the chain pickerel, this...



Redfin shiner

Lythrurus umbratilis

The redfin shiner (Lythrurus umbratilis) is a freshwater fish in the family Cyprinidae. The redfin shiner is most commonly found in the Ohio and Mississippi River basins, as well as in drainages of the Great Lakes. The diet of the redfin shiner...



Redhook myleus

Myloplus rubripinnis

The redhook myleus, Myloplus rubripinnis, is a medium to large omnivorous fish of the family Serrasalminidae from South America, where found in the Amazon and Orinoco basins, as well as rivers of the Guiana Shield. It is peaceful and schools often,...



Redhump eartheater

Geophagus steindachneri

The redhump eartheater ('Geophagus steindachneri') is a species of eartheater cichlid from freshwater habitats in northwestern South America.



Redmouth grouper

Aethaloperca rogaa

The redmouth grouper (*Aethaloperca rogaa*), also known as the red-flushed rock cod is a species of ray-finned fish, a grouper from the subfamily Epinephelinae which is part of the family Serranidae, which also includes the anthias and sea basses....



Redside barb

Puntius bimaculatus

The redside barb or two-spot barb (*Puntius bimaculatus*) is a species of ray-finned fish in the genus *Puntius*. It is found in India and Sri Lanka. It was identified and classified by Pieter Bleeker in 1863.



Redside dace

Clinostomus elongatus

The redside dace (*Clinostomus elongatus*) is a species of ray-finned fish in the family Cyprinidae, found in the United States and Canada. It is unique among minnows, being the only species to routinely feed on flying insects by leaping from...



Redside shiner

Richardsonius balteatus

The redside shiner (*Richardsonius balteatus*) is a species of cyprinid fish found in the western United States and Canada.



Redspotted sunfish

Lepomis miniatus

The redspotted sunfish (*Lepomis miniatus*), also known as a stumpknocker, is a species of freshwater ray-finned fish, a sunfish from the family Centrarchidae which is native to the United States. The redspotted sunfish was previously considered to...



Redtail butterflyfish

Chaetodon collare

The red-tailed butterflyfish (Chaetodon collare), also known as the brown butterflyfish, Pakistani butterflyfish or redbill butterflyfish, is a species of marine ray-finned fish, a butterflyfish belonging to the family Chaetodontidae. It can be...



Redtail catfish

Phractocephalus hemiliopterus

The redbill catfish, Phractocephalus hemiliopterus, is a pimelodid (long-whiskered) catfish. In Venezuela, it is known as cajaró, in Guyana, it is known as a banana catfish, and in Brazil, it is known as pirarara, stemming from the Tupi language...



Redtail notho

Nothobranchius guentheri

The redbtail notho (Nothobranchius guentheri) is a species of killifish in the family Nothobranchiidae. It is endemic to Zanzibar. Its natural habitats are intermittent rivers and intermittent freshwater marshes. The redbtail notho eats mosquito...



Redtail splitfin

Xenotoca eiseni

The redbtail splitfin or redbtail goodeid (Xenotoca eiseni) is a species of goodeid fish from the family Goodeidae and subfamily Goodeinae. Like other members of Goodeinae, the redbtail splitfin is native to Mexico and a livebearer. However, the...



Redtail surfperch

Amphistichus rhodoterus

The redbtail surfperch (Amphistichus rhodoterus) is a marine fish that inhabits the near-shore and estuarine waters of North American Pacific coasts.



Redtoothed triggerfish

Odonus niger

The Redtoothed triggerfish (Odonus niger) is a triggerfish of the tropical Indo-Pacific area, and the sole member of its genus. Some other common names include blue triggerfish, redfang triggerfish, redtoothed filefish, and Niger triggerfish.



Reedfish

Erpetoichthys calabaricus

The reedfish, ropefish (more commonly used in the United States), or snakefish, Erpetoichthys calabaricus, is a species of fish in the family Polypteridae alongside the bichirs. It is the only member of the genus Erpetoichthys. It is native to...



Reef manta ray

Mobula alfredi

The reef manta ray (Mobula alfredi) is a species of ray in the family Mobulidae, one of the largest rays in the world. Among generally recognized species, it is the second-largest species of ray, only surpassed by the giant oceanic manta ray. The...



Reef triggerfish

Rhinecanthus rectangulus

The reef triggerfish (*Rhinecanthus rectangulus*), also known as the rectangular triggerfish, wedgetail triggerfish or by its Hawaiian name humuhumunukunukuāpua‘a...



Reptilian snake eel

Brachysomophis henshawi

The reptilian snake eel (*Brachysomophis henshawi*), also known as Henshaw's snake eel, the Hawaiian crocodile eel or the crocodile snake eel, is an eel in the family Ophichthidae (worm/snake eels). It was described by David Starr Jordan and John...



Reticulate whipray

Himantura uarnak

The reticulate whipray or honeycomb stingray (*Himantura uarnak*) is a species of stingray in the family Dasyatidae. It inhabits coastal waters in the western Indian Ocean including the Red Sea, Natal and the Arabian Sea; also a Lessepsian...



Retroculus lapidifer

Retroculus lapidifer

Retroculus lapidifer is a species of cichlid native to tropical South America, where it is found in the rivers of the southeastern Amazon basin in Brazil. This fish was first described in 1855 by the French naturalist Francis de Laporte de...



Rex sole

Glyptocephalus zachirus

The rex sole (*Glyptocephalus zachirus*) is a flatfish of the family Pleuronectidae. Locally, it may also be known as a witch or threadfin sole (due to the filamentous pelvic fin on the eyed surface). It is a demersal fish that lives in temperate...



Rhabdosargus holubi

Rhabdosargus holubi

Rhabdosargus holubi, the Cape stumpnose, is a species of fish in the seabream family, Sparidae. It is native to southern Africa, where it can be found mainly along the eastern coast of South Africa. This fish is usually around 15 centimeters long,...



Rhabdosargus sarba

Rhabdosargus sarba

Rhabdosargus sarba, also known as the goldlined seabream, silver bream, tarwhine, or yellowfin bream, is a species of fish in the seabream family, Sparidae. This species occurs Red Sea and the Persian Gulf to Eastern Cape, South Africa, eastwards...



Rhinecanthus assasi

Rhinecanthus assasi

Rhinecanthus assasi, commonly known as the Assasi triggerfish or Arabian picassofish, is a species of fish in the family Balistidae, the triggerfishes.



Rhinochimaera pacifica

Rhinochimaera pacifica

Rhinochimaera pacifica, commonly known as the Pacific spookfish, knifenose chimaera, narrownose chimaera, Pacific long-nosed chimaera, or Pinocchiofish, is a species of chimaera in the family Rhinochimaeridae. It lives in various parts of the...



Rhinogobiops

Rhinogobiops nicholsii

Rhinogobiops is a genus of true gobies in the family Gobiidae. It is monotypic, being represented by the single species, Rhinogobiops nicholsii, also known as the blackeye goby, bluespot goby, and crested goby. They are common inhabitants of...



Rhinogobius flumineus

Rhinogobius flumineus

Rhinogobius flumineus, commonly known as the lizard goby or kawa-yoshinobori, is a species of goby endemic to Japan where it is found in the mid- to upper reaches of fast-flowing rivers. This species can reach a length of 7 centimetres (2.8 in) TL.



Rhinogobius giurinus

Rhinogobius giurinus

Rhinogobius giurinus is a species of goby native to eastern Asia where it inhabits marine, brackish and fresh waters of rivers and estuaries. This species can reach a length of 8 centimetres (3.1 in) TL. It is of importance to local peoples as a...



Rhinopias eschmeyeri

Rhinopias eschmeyeri

Rhinopias eschmeyeri or Eschmeyer's scorpionfish or paddle-flap scorpionfish, is a species of marine ray-finned fish belonging to the family Scorpaenidae, the scorpionfishes. This species is found in the Indo-West Pacific. It grows to an average...



Rhinopias frondosa

Rhinopias frondosa

Rhinopias frondosa, the weedy scorpionfish or the weed fish, is a species of marine ray-finned fish belonging to the family Scorpaenidae, the scorpionfishes. This species is found in the Indo-West Pacific. It is a rare but highly desirable fish...



Rhizosomichthys totae

Rhizosomichthys totae

Rhizosomichthys totae (sometimes known as Greasefish in English) is a species of catfish (order Siluriformes) of the family Trichomycteridae, and the only species of the genus Rhizosomichthys. This fish grew to about 13.8 centimetres (5.4 in) and...



Ribbon eel

Rhinomuraena quaesita

The ribbon eel (Rhinomuraena quaesita), also known as the leaf-nosed moray eel or bernis eel, is a species of moray eel, the only member of the genus Rhinomuraena. What is now known as R. quaesita also includes the former R. amboinensis. R....



Ribbon sawtail fish

Idiacanthus fasciola

The ribbon sawtail fish, Idiacanthus fasciola, is a barbeled dragonfish of the family Stomiidae, found around the world at depths over 500 m between latitudes 40° N and 54° S. Length is up to 35 cm in general for the female, and 15 cm for the male.



Ribboned sweetlips

Plectorhinchus polytaenia

Plectorhinchus polytaenia, the ribboned sweetlips, also known as Tesone di mare or yellow-ribbon sweetlips, is a species of marine ray-finned fish, a sweetlips belonging to the subfamily Plectorhinchinae, one of two subfamilies in the family...



Rice-paddy eel

Pisodonophis boro

The rice-paddy eel (*Pisodonophis boro*; also known commonly as the Bengal's snake-eel, the estuary snake eel, or the snake eel) is an eel in the family Ophichthidae (worm-snake eels). It was described by Francis Buchanan-Hamilton in 1822,...



Ridge scaled rattail

Macrourus carinatus

The ridge scaled rattail or ridge-scaled grenadier, *Macrourus carinatus*, is a species of deep-water fish in the family Macrouridae. It has southern circumglobal distribution in temperate to subantarctic waters (34°S–65°S) and is found in the...



Ring-tailed cardinalfish

Ostorhinchus aureus

The ring-tailed cardinalfish (*Ostorhinchus aureus*) is a widespread fish species in the family Apogonidae found in the Red Sea and off East Africa to Papua New Guinea, north to Japan, and south to Australia.



Rio Grande cutthroat trout

Oncorhynchus clarkii virginalis

The Rio Grande cutthroat trout (*Oncorhynchus clarkii virginalis*), a member of the family Salmonidae, is found in northern New Mexico and southern Colorado in tributaries of the Rio Grande. It is one of 14 subspecies of cutthroat trout native to the...



Rio Grande silvery minnow

Hybognathus amarus

The Rio Grande silvery minnow or Rio Grande minnow (Hybognathus amarus) is a small herbivorous North American fish. It is one of the seven North American members of the genus Hybognathus, in the cyprinid family. The Rio Grande silvery minnow is...



Rita rita

Rita rita

Rita rita (Common names: rita (English), রিটা (Bengali)) is a species of bagrid catfish that is found across southern Asia. It has been recorded in Afghanistan, Bangladesh, India, Myanmar, Nepal and Pakistan. It is one of the giants of its genus....



River blackfish

Gadopsis marmoratus

The river blackfish (Gadopsis marmoratus) is a freshwater fish endemic to the temperate waters of south-eastern Australia. It is found from southern Queensland through to central Victoria, including in the Murray-Darling river system. It is also...



River carpsucker

Carpionodes carpio

The river carpsucker (Carpionodes carpio) is a freshwater fish found in the inland United States and northern Mexico. This species has a slightly arched back and is somewhat stout and compressed. While the fins are usually opaque, in older fish...



River chub

Nocomis micropogon

The river chub (*Nocomis micropogon*) is a minnow in the family Cyprinidae. It is one of the most common fishes in North American streams.



River redhorse

Moxostoma carinatum

The river redhorse (*Moxostoma carinatum*) is a species of freshwater fish endemic to the eastern half of the United States and southeastern Canada. They can range from 10 to 30 inches in length and can potentially reach a weight of more than 10...



Roanoke bass

Ambloplites cavifrons

The Roanoke bass (Ambloplites cavifrons) is a species of freshwater fish in the sunfish family (Centrarchidae) of order Perciformes. This species is endemic to rivers in the eastern United States of America, where it is native only to the states...



Roanoke logperch

Percina rex

The Roanoke logperch (Percina rex) is a species of freshwater ray-finned fish, a darter from the subfamily Etheostomatinae, part of the family Percidae, which also contains the perches, ruffes and pikeperches. It is found in the Roanoke and...



Robust ghost pipefish

Solenostomus cyanopterus

The robust ghost pipefish (Solenostomus cyanopterus), also known as the blue-finned ghost pipefish, Racek's ghost pipefish, robust-snouted ghost pipefish, or the squaretail ghost-pipefish, is a species of false pipefish belonging to the family...



Robust redhorse

Moxostoma robustum

The robust redhorse (Moxostoma robustum) is a redhorse-type fish that lives in the freshwater streams of the eastern United States. It was previously called the smallfin redhorse.



Rock bass

Ambloplites rupestris

The rock bass (*Ambloplites rupestris*), also known as the rock perch, goggle-eye, red eye, and black perch, is a freshwater fish native to east-central North America. This red-eyed fish is a species of freshwater fish in the sunfish family...



Rock beauty

Holacanthus tricolor

The rock beauty (*Holacanthus tricolor*), also known as corn sugar, coshubba, rock beast, catalineta, and yellow nanny, is a species of marine ray-finned fish, a marine angelfish belonging to the family Pomacanthidae. It is found in the western...



Rock cod

Lotella rhacina

The rock cod (*Lotella rhacina*) is a temperate fish found off the coasts of southeastern Australia, Tasmania, the Great Australian Bight and northwards up the southwestern Australia coasts. They are also found around the coasts of New Zealand and...



Rock goby

Gobius paganellus

The rock goby (*Gobius paganellus*) is a small coastal goby of eastern Atlantic waters, from Scotland to Senegal. It is also reported from the Mediterranean and Black Seas, and is an anti-Lessepsian migrant in the Gulf of Eilat and Red Sea. There...



Rock greenling

Hexagrammos lagocephalus

The rock greenling (*Hexagrammos lagocephalus*) is a species of marine ray-finned fish belonging to the family Hexagrammidae, the greenlings. It is sometimes known as fringed greenling and erroneously as the red rock trout.



Rock sole

Lepidopsetta bilineata

The rock sole (Lepidopsetta bilineata), also known as the Pacific rock sole or Southern rock sole is a flatfish of the family Pleuronectidae. It is a demersal fish that lives on sand and gravel bottoms at depths of up to 575 metres (1,886 ft),...



Rohu

Labeo rohita

The rohu, rui, ruhi or roho labeo (Labeo rohita) is a species of fish of the carp family, found in rivers in South Asia. It is a large omnivore and extensively used in aquaculture.



Romanichthys

Romanichthys valsanicola

Romanichthys is a genus of ray-finned fish, one of two genera in the tribe Romanichthyini, which along with the tribe Luciopercini, forms the subfamily Luciopercinae of the family Percidae, alongside the perches, ruffes, and darters. The genus...



Roosterfish

Nematistius pectoralis

The roosterfish, Nematistius pectoralis, is a game fish found in the warmer waters of the East Pacific from Baja California to Peru. It is the only species in the genus Nematistius and the family Nematistiidae. It is distinguished by its "rooster..."



Rose danio

Danio roseus

The rose danio (Danio roseus) is a tropical fish of the family Cyprinidae. At first glance, this species resembles the pearl danio, but lacks the orange stripe on the side. It has a rosy colouration along the lower part of the fish and lower fins...



Rosy barb

Pethia conchonius

The rosy barb (Pethia conchonius) is a subtropical freshwater cyprinid fish found in southern Asia from Afghanistan to Bangladesh.



Rosy bitterling

Rhodeus ocellatus

The rosy bitterling or Tairiku baratanago (Rhodeus ocellatus) is a small freshwater fish belonging to the family Cyprinidae (carp), native to East Asia from the Amur River basin to the Pearl River basin. Females are about 4–5 cm (1.6–2.0 in) long...



Rosy tetra

Hyphessobrycon rosaceus

The rosy tetra (Hyphessobrycon rosaceus) is a small species of Characin from the South American countries of Guyana and Suriname. It is popular in the aquarium trade.



Roughback whipray

Fluvitrygon kittipongi

The roughback whipray (Fluvitrygon kittipongi) is a rare species of freshwater stingray in the family Dasyatidae, found over sandy bottoms in the Mae Klong and Chao Phraya Rivers of Thailand. Growing no more than 29 cm (11 in) across, this small...



Rougheye rockfish

Sebastes aleutianus

The rougheye rockfish (Sebastes aleutianus) is a rockfish of the genus Sebastes. It is also known as the blackthroat rockfish, rougheye seaperch, blacktip seaperch, longlife seaperch or the blacktip rockfish and grows to a maximum of about 97 cm...



Roughskin dogfish

Centroscyrnus owstonii

Sharks portalThe roughskin dogfish (Centroscyrnus owstonii) is a sleeper shark of the family Somniosidae, found around the world on continental shelves in tropical, subtropical and temperate seas, at depths of between 100 and 1,500 m. It reaches...



Roughtail stingray

Bathytoshia centroura

The roughtail stingray (Bathytoshia centroura) is a species of stingray in the family Dasyatidae, with separate populations in coastal waters of the northwestern and southwestern Atlantic Ocean. This bottom-dwelling species typically inhabits...



Round fantail stingray

Taeniurops grabatus

The round fantail stingray (Taeniurops grabatus) or round stingray, is a poorly known species of stingray in the family Dasyatidae. It inhabits sandy, muddy, or rocky coastal habitats in the eastern Atlantic Ocean and the southern Mediterranean...



Round goby

Neogobius melanostomus

The round goby (Neogobius melanostomus) is a euryhaline bottom-dwelling species of fish of the family Gobiidae. It is native to Central Eurasia, including the Black Sea and the Caspian Sea. Round gobies have established large non-native...



Round ribbontail ray

Taeniurops meyeni

The round ribbontail ray or blotched fantail ray, (Taeniurops meyeni) is a species of stingray in the family Dasyatidae, found throughout the nearshore waters of the tropical Indo-Pacific, as well as off islands in the eastern Pacific. It is a...



Round sardinella

Sardinella aurita

The round sardinella (Sardinella aurita) is a species of ray-finned fish in the genus Sardinella found in both sides of the Atlantic Ocean and the Mediterranean Sea. S. aurita went through a large boom in catch population around 1990. However, its...



Round scad

Decapterus punctatus

The round scad (Decapterus punctatus) (or cigar minnow) is a species of fish in the Carangidae. It was described in 1829 by the French naturalist and zoologist, Georges Cuvier. Although the round scad is considered a good food fish, it is mostly...



Round stingray

Urobatis halleri

The round stingray (Urobatis halleri) or Haller's round ray and Little round stingray is a species of round ray, family Urotrygonidae, found in the coastal waters of the tropical and subtropical parts of the northeastern Pacific Ocean. It is a...



Round whitefish

Prosopium cylindraceum

The round whitefish (Prosopium cylindraceum) is a freshwater species of fish that is found in lakes from Alaska to New England, including the Great Lakes. It has an olive-brown back with light silvery sides and underside and its length is...



Round-tailed paradise fish

Macropodus ocellatus

The round-tailed paradise fish (Macropodus ocellatus) is a species of gourami native to eastern Asia, where it is found in Korea and China. It is also found in Japan but it is believed that they were introduced from Korea in the 1910s. It is also...



Roundscale spearfish

Tetrapturus georgii

The roundscale spearfish (Tetrapturus georgii) is an Istiophoridae species of marlin living in the epipelagic zone of the Atlantic Ocean. It has long been misidentified as white marlin but can be differentiated thanks to their scale shapes that...



Roundtail chub

Gila robusta

The roundtail chub (Gila robusta) is a cyprinid fish in the genus Gila, of southwestern North America. It is native to the Colorado River drainage basin, including the Gila River and other tributaries, and in several other rivers. It is part of...



Roving coral grouper

Plectropomus pessuliferus

The roving coral grouper (Plectropomus pessuliferus), also known as the spotted coral grouper, is a species of marine ray-finned fish, a grouper from the subfamily Epinephelinae which is part of the family Serranidae, which also includes the...



Royal angelfish

Pygoplites diacanthus

The royal angelfish (Pygoplites diacanthus), or regal angelfish, is a species of marine ray-finned fish, a marine angelfish belonging to the family Pomacanthidae, and the monotypic genus Pygoplites. It is found in tropical Indo-Pacific oceans. It...



Royal gramma

Gramma loreto

The royal gramma (Gramma loreto), also known as the fairy basslet, is a species of fish in the family Grammatidae native to reef environments of the tropical western Atlantic Ocean. They are commonly kept in aquariums.



Royal knifefish

Chitala blanci

The royal knifefish or Indochina featherback, Chitala blanci, is a species of ray-finned fish in the family Notopteridae found in the Mekong basin in Cambodia, Laos, Thailand and Vietnam.



Royal tetra

Inpaichthys kerri

The royal tetra (Inpaichthys kerri) is a species of characin endemic to Brazil, where it is found in tributaries of the Aripuanã River. It is the sole member of its genus. The fish was named in honor of Warwick Estevam Kerr (1922-2018), a...



Ruby seadragon

Phyllopteryx dewysea

The ruby seadragon (Phyllopteryx dewysea) is a marine fish in the family Syngnathidae, which also includes seahorses. It inhabits the coast of Western Australia. The species was first described in 2015, making it only the third known species of...



Ruby tetra

Axelrodia riesei

The ruby tetra (Axelrodia riesei) is a species of freshwater fish in the family Characidae. It is found in the Río Meta, Colombia, South America. In nature, it has a natural red colour, but this is somewhat lessened in captivity. Ruby tetras grow...



Rubyfish

Plagiogeneion rubiginosum

The rubyfish (Plagiogeneion rubiginosum), also known as the cosmopolitan rubyfish, red ruby or ruby rover, is a species of marine ray-finned fish, belonging to the family Emmelichthyidae, the rovers, bonnetmouths and rubyfishes. This species is...



Ruffe

Gymnocephalus cernua

The ruffe (Gymnocephalus cernua), also known as the Eurasian ruffe or pope, is a freshwater fish found in temperate regions of Europe and northern Asia. It has been introduced into the Great Lakes of North America, reportedly with unfortunate...



Rummy-nose tetra

Hemigrammus rhodostomus

The rummy-nose tetra (Petitella rhodostoma) is a species of tropical freshwater characin fish originating in South America, popular among fishkeepers as a tropical aquarium fish. One of many small tetras belonging to the same genus, it is on...



Russian sturgeon

Acipenser gueldenstaedtii

The Russian sturgeon (Acipenser gueldenstaedtii), also known as the diamond sturgeon or Danube sturgeon, is a species of fish in the family Acipenseridae. It is found in Azerbaijan, Bulgaria, Georgia, Iran, Kazakhstan, Romania, Russia, Turkey....



Rusty parrotfish

Scarus ferrugineus

The rusty parrotfish (*Scarus ferrugineus*) is a species of marine ray-finned fish, a parrotfish belonging to the family Scaridae. It is associated with reefs in the north western Indian Ocean and the Red Sea.



Rutilus frisii

Rutilus frisii

Do not confuse it with another fish called the kutum, *Rutilus kutum**Rutilus frisii*, called the vyrezub, Black Sea roach, or kutum, is a species of fish in the family Cyprinidae, native to the basins of the Black Sea, Sea of Azov, and Sea of...



Rutilus heckelii

Rutilus heckelii

Rutilus heckelii is a species of roach, a genus in the family Cyprinidae. This species occurs in freshened areas of Black and Azov Seas, entering the Don, Kuban, Dniepr, Dniestr and rarely Danube drainages. Newer research, however, suggests that...



Sablefish

Anoplopoma fimbria

The sablefish (Anoplopoma fimbria) is one of two members of the fish family Anoplopomatidae and the only species in the genus Anoplopoma. In English, common names for it include sable (US), butterfish (US), black cod (US, UK, Canada), blue cod...



Saccopharynx ampullaceus

Saccopharynx ampullaceus

Saccopharynx ampullaceus, referred to as the gulper eel, gulper, taillight gulper or pelican-fish, is an ocean-dwelling eel found in the North Atlantic Ocean. They are found up to a depth of 3,000 m (9,800 ft). These fish are rarely observed, so...



Sacramento blackfish

Orthodon microlepidotus

The Sacramento blackfish (Orthodon microlepidotus) is a species of freshwater fish in central California. A cyprinid, the blackfish is the sole member of its genus.



Sacramento perch

Archoplites interruptus

The Sacramento perch (Archoplites interruptus) is an endangered sunfish (family Centrarchidae) native to the Sacramento–San Joaquin River Delta, Pajaro, and Salinas River areas in California, but widely introduced throughout the western United...



Sacramento pikeminnow

Ptychocheilus grandis

The Sacramento pikeminnow (Ptychocheilus grandis), formerly known as the Sacramento squawfish, is a large cyprinid fish of California, United States. It is native to the Los Angeles River, Sacramento-San Joaquin, Pajaro-Salinas, Russian River,...



Sacramento splittail

Pogonichthys macrolepidotus

The splittail (Pogonichthys macrolepidotus), also called Sacramento splittail, is a cyprinid fish native to the low-elevation waters of the Central Valley in California. It was first described by William O. Ayres in 1854. It is the sole living...



Saddle butterflyfish

Chaetodon ephippium

The saddle butterflyfish (Chaetodon ephippium) is a species of marine ray-finned fish, a butterflyfish belonging to the family Chaetodontidae. It is found in the Indian and Pacific Oceans from Sri Lanka and the Cocos-Keeling Islands to the...



Saddleback clownfish

Amphiprion polymnus

Amphiprion polymnus, also known as the saddleback clownfish or yellowfin anemonefish, is a black and white species of anemonefish with a distinctive saddle. Like all anemonefishes it forms a symbiotic mutualism with sea anemones and is unaffected...



Saddletail grouper

Epinephelus daemeli

The saddletail grouper (Epinephelus daemeli), also known as black cod or black rock-cod in Australia, and as saddle-tailed grouper or spotted black grouper in New Zealand, is a large marine fish of the family Serranidae. It is found off the...



Saffron cod

Eleginus gracilis

The saffron cod (Eleginus gracilis) is a commercially harvested fish closely related to true cods (genus Gadus). It is dark grey-green to brown, with spots on its sides and pale towards the belly. It may grow to 55 cm and weigh up to 1.3 kg.



Sahyadria chalakkudiensis

Sahyadria chalakkudiensis

Sahyadria chalakkudiensis is a species of cyprinid fish endemic to the Chalakkudy River, Kerala, India in the Western Ghats where it can be found in well-vegetated upper reaches of rivers. This species can reach a length of 12.5 centimetres (4.9...



Sailback houndshark

Gogolia filewoodi

Sharks portalThe sailback houndshark (Gogolia filewoodi) is a houndshark of the family Triakidae, and the only member of the genus Gogolia. It is found in the deep waters of continental shelf off northern Papua New Guinea. Only one specimen has...



Sailfin flying fish

Parexocoetus brachypterus

The sailfin flying fish (Parexocoetus brachypterus) is a member of the flying fish family (Exocoetidae). As is typical of other members of its family, this species has the ability to jump out of the water and glide on hypertrophied fins in order...



Sailfin molly

Poecilia latipinna

The sailfin molly (Poecilia latipinna) is a livebearer fish typically found in both freshwater and brackish waterways along the East Coast of the United States, from North Carolina south to Florida, and around the Gulf of Mexico to Texas, and...



Sailfin roughshark

Oxynotus paradoxus

The sailfin roughshark (Oxynotus paradoxus) is a species of dogfish shark in the family Oxynotidae, found in the eastern North Atlantic from Scotland to Senegal between latitudes 61°N and 11°N, at depths of between 265 and 720 m (869 and 2,362...



Sailfin snapper

Symphoricthys spilurus

The sailfin snapper (*Symphorichthys spilurus*), blue-lined sea bream or blue-lined sea perch is a species of marine ray-finned fish, a snapper belonging to the family Lutjanidae. It is native to the Indo-Pacific region. It is of minor importance...



Sailfin tang

Zebrasoma velifer

The sailfin tang (*Zebrasoma velifer*), the Pacific sailfin tang, purple sailfinned tang or sailfin surgeonfish, is a marine ray-finned fish belonging to the family Acanthuridae which includes the surgeonfishes, unicornfishes and tangs. This fish...



Sakhalin sturgeon

Acipenser mikadoi

The Sakhalin sturgeon (*Acipenser mikadoi*) is a species of fish in the family Acipenseridae. It is found in Japan and Russia.



Salangichthys

Salangichthys microdon

Salangichthys microdon, the Japanese icefish, is a species of icefish found in Japan, Korea and the Russian Far East. With the recent removal of *S. ishikawae* to the genus *Neosalangichthys* this species is the only remaining member of the genus...



Salarias fasciatus

Salarias fasciatus

Salarias fasciatus, commonly known as the jewelled blenny or lawnmower blenny is a benthic, neritic, marine fish species endemic to Australasia. Despite being known as the lawnmower blenny, due to its propensity to consume algae growth in...



Salminus brasiliensis

Salminus brasiliensis

Salminus brasiliensis (dourado, dorado, golden dorado, river tiger or jaw characin) is a large, predatory characiform freshwater fish found in central and east-central South America. Despite having *Salminus* in its name, the dorado is not related...



Salmo aphelios

Salmo aphelios

Salmo aphelios is a species of trout, a salmonid fish endemic to Lake Ohrid in North Macedonia and Albania in the Balkans. Salmo aphelios is one of four different forms of the Ohrid trout complex within the single lake, along with S. balcanicus,...



Salmo balcanicus

Salmo balcanicus

Salmo balcanicus is a type of trout, a fish in the family Salmonidae. It is endemic to Lake Ohrid and its outlet in North Macedonia and Albania in the Balkans. Salmo balcanicus is one of four different forms of the Ohrid trout complex which is...



Salmo carpio

Salmo carpio

Salmo carpio, also known as the carpione (carpione del Garda or Lake Garda carpione) is a salmonid fish endemic to Lake Garda in Italy. It has been introduced to a number of other lakes in Italy and elsewhere but unsuccessfully in all cases. The...



Salmo cettii

Salmo cettii

Salmo cettii, or the Mediterranean trout, is a species of trout, a freshwater fish in the family Salmonidae. It lives in the Mediterranean region in Corsica, Sardinia, Sicily, and on the Italian mainland in the Magra drainage and further south....



Salmo ciscaucasicus

Salmo ciscaucasicus

Salmo ciscaucasicus, the Caspian salmon or Terek trout, is a salmonid fish endemic to the Caspian Sea and its inflowing rivers. It was described in 1967 originally as a subspecies of Salmo trutta. S. ciscaucasicus lives on the western shore of...



Salmo dentex

Salmo dentex

Salmo dentex is a variety of trout, a freshwater fish in the family Salmonidae, found in the western Balkans. Until recently the identity, biological distinctness and species status of the dentex trout were not properly clarified, but genetic...



Salmo ezenami

Salmo ezenami

Salmo ezenami is a critically endangered freshwater salmonid fish, endemic to Lake Kezenoi-Am (Lake Eizenam) in Northern Caucasus. The only natural habitat where this species occurs is in Lake Kezenoi-Am in Chechnya at the border of Dagestan. This...



Salmo macedonicus

Salmo macedonicus

Salmo macedonicus is a species of fish in the family Salmonidae. It is found only in rivers in the upper Vardar river basin in North Macedonia. It is threatened by habitat loss. The taxonomic status of this fish is controversial.



Salmo macrostigma

Salmo macrostigma

Salmo macrostigma is a species of freshwater trout endemic to Algeria in northwest Africa. It can reach a length of 60 centimetres (24 in) TL. The name Salmo trutta macrostigma (or Salmo macrostigma) has previously been used of many populations...



Salmo marmoratus

Salmo marmoratus

Salmo marmoratus, the marble trout, is a species of freshwater fish in the family Salmonidae. It is characterized by a distinctive marbled color pattern and high growth capacity. The marble trout is found in only a handful of drainages and rivers...



Salmo obtusirostris

Salmo obtusirostris

Salmo obtusirostris, also known as the Adriatic trout, Adriatic salmon, and softmouth trout, is a species of salmonid fish endemic to the rivers of Western Balkans in southeastern Europe. The scientific name has changed several times through...



Salmo obtusirostris salonitana

Salmo obtusirostris salonitana

Salmo obtusirostris salonitana (also termed by the common names soft-muzzled trout or Solin salmon and the scientific synonym Salmothymus obtusirostris salonitana) is an endemic trout subspecies found in the Jadro River in the vicinity of Solin...



Salmo ohridanus

Salmo ohridanus

Salmo ohridanus, also known by the local name as the belvica in North Macedonia or belushka in Albania, is a species of freshwater salmonid fish, endemic to Lake Ohrid in Albania and North Macedonia. Salmo ohridanus is a relatively small fish,...



Salmo peristericus

Salmo peristericus

Salmo peristericus, or the Prespa trout is a variety of trout, a freshwater fish in the family Salmonidae. It is endemic to the Lake Prespa watershed at the border area of Greece and North Macedonia. Four populations are known: one in the Agios...



Salmo platycephalus

Salmo platycephalus

Salmo platycephalus, known as the flathead trout, Ala balik or the Turkish trout, is a type of trout, a fish in the family Salmonidae. It is endemic to southeastern Turkey. It is known only from one population, which occupies three streams,...



Salmo trutta fario

Salmo trutta fario

Salmo trutta fario, sometimes called the river trout, is a river-dwelling freshwater predatory fish from the genus Salmo of the family Salmonidae. It is one of the three main subspecies or morphs of the brown trout (Salmo trutta), besides sea...



Salmon shark

Lamna ditropis

The salmon shark (Lamna ditropis) is a species of mackerel shark found in the northern Pacific ocean. As an apex predator, the salmon shark feeds on salmon, squid, sablefish, and herring. It is known for its ability to maintain stomach...



Salt and pepper catfish

Corydoras habrosus

The salt and pepper catfish (Corydoras habrosus) is a tropical freshwater fish belonging to the Corydoradinae sub-family of the family Callichthyidae. It originates in inland waters in South America, and is found in the Upper Orinoco River basin...



Salvelinus colii

Salvelinus colii

Salvelinus colii, also called Cole's char, Enniskillen char or Trevelyan's char, is a cold-water species of char fish in the family Salmonidae. Salvelinus colii is currently located in Ireland, in several lakes draining westward, in County Clare,...



Salvelinus elgyticus

Salvelinus elgyticus

Salvelinus elgyticus is a species of fish in the salmon family, Salmonidae. It is a member of genus Salvelinus, the chars. It is known commonly as the small-mouth char (Russian: малоротая палия). It is endemic to Lake El'gygytgyn in eastern...



Salvelinus evasus

Salvelinus evasus

Salvelinus evasus, is a vulnerable deepwater char or trout living in the Ammersee lake in Bavaria, Southern Germany. This fish lives in the great depths of the lake, below 80 m. It can reach up to a foot in length, about 30 cm. Salvelinus evasus...



Salvelinus grayi

Salvelinus grayi

Salvelinus grayi, also called Gray's char, Lough Melvin char or freshwater herring, is a species of lacustrine char in the family Salmonidae. It is only found in Lough Melvin, Ireland; numbers of fish are declining and the species is considered...



Salvelinus inframundus

Salvelinus inframundus

Salvelinus inframundus, also known as Orkney charr is a cold-water fish in the family Salmonidae which is endemic to Scotland.



Salvelinus killinensis

Salvelinus killinensis

Salvelinus killinensis, also known as Haddy charr is a variety of charr found in certain lakes in Scotland. Salvelinus killinensis lives in Loch Killin (Inverness-shire), after which it is named. It is also found in Loch Doine (Trossachs) and...



Salvelinus leucomaenis

Salvelinus leucomaenis

Salvelinus leucomaenis, the whitespotted char, is an East Asian trout in the genus Salvelinus, called iwana in Japanese and kundzha (кунджа) in Russian. Both landlocked and ocean-run forms occur. The landlocked form typically grows up to 35 cm (1...



Salvelinus obtusus

Salvelinus obtusus

Salvelinus obtusus, commonly called the blunt-nosed Irish charr or blunt-snouted Irish char, is a species of lacustrine char fish in the family Salmonidae, found in the Lakes of Killarney, Ireland.



Salvelinus profundus

Salvelinus profundus

Salvelinus profundus (German: Tiefseesaibling 'Deepwater char') is a deepwater char species found only in deep areas of Lake Constance. This fish can reach 24 cm (9.4 in) in length and has a blunt snout with the mouth in subinferior position. Its...



Salvelinus umbla

Salvelinus umbla

Salvelinus umbla, also known as lake char, is a species of char found in certain lakes of the region of the Alps in Europe. This char species usually inhabits the deeper waters of the lakes, feeding on crustaceans, insects and benthic fauna....



Salvin's cichlid

Trichromis salvini

The Salvin's cichlid (*Trichromis salvini*), also known as the yellow-belly cichlid or tricolored cichlid, is a species of the family Cichlidae. It is found in rivers of the Atlantic slope of southern Mexico, Belize, and Guatemala.



Samson fish

Seriola hippos

The samson fish (Seriola hippos) is a jack of the genus Seriola. It is found in the Indo-Pacific Oceans to eastern Northland in Australia. It is not found anywhere in New Zealand. Its length is between 80 and 150 cm.



Sand devil

Squatina dumeril

The sand devil or Atlantic angel shark (Squatina dumeril) is a species of angelshark, family Squatinidae, native to the northwestern Atlantic Ocean. It occurs off the eastern United States, in the northern Gulf of Mexico, and possibly in parts of...



Sand goby

Pomatoschistus minutus

The sand goby (Pomatoschistus minutus), also known as a polewig or pollybait, is a species of ray-finned fish native to marine and brackish European waters from the Baltic Sea through the Mediterranean Sea and into the Black Sea where it occurs...



Sand grey mullet

Myxus elongatus

The sand grey mullet (Myxus elongatus) is a species of mullet found in coastal marine waters of Australia.



Sand shiner

Notropis stramineus

The sand shiner (*Notropis stramineus*) is a widespread North American species of freshwater fish in the family Cyprinidae. Sand shiners live in open clear water streams with sandy bottoms where they feed in schools on aquatic and terrestrial...



Sand smelt

Atherina presbyter

The sand smelt (*Atherina presbyter*) is a species of marine fish of the family Atherinidae, common in the northeastern Atlantic from the Danish straits, where it is rare, and Scotland to the Canary Islands and the western Mediterranean Sea. Sand...



Sand sole

Pegusa lascaris

The sand sole (*Pegusa lascaris*) is a fish species in the family Soleidae. It is a marine, subtropical, demersal fish up to 40 centimetres (16 in) long. Widespread in the northeastern and southeastern Atlantic, to the Gulf of Guinea in the south...



Sand tiger shark

Carcharias taurus

The sand tiger shark (*Carcharias taurus*), gray nurse shark, spotted ragged-tooth shark or blue-nurse sand tiger, is a species of shark that inhabits subtropical and temperate waters worldwide. It inhabits the continental shelf, from sandy...



Sand whiting

Sillago ciliata

The sand whiting (*Sillago ciliata*), also known as the summer whiting, yellowfin whiting or blue-nose whiting, is a common species of coastal marine fish of the family Sillaginidae, the smelt-whittings. It is a slender, slightly compressed fish...



Sandbar Shark

Carcharhinus plumbeus

The Sandbar shark (Carcharhinus plumbeus) is a species of requiem shark that are native to the Atlantic Ocean and the Indo-Pacific. It shouldn't be confused with the similarly named Sand tiger shark (Carcharias taurus) and can be distinguished by...



Sander marinus

Sander marinus

Sander marinus, the estuarine perch, also called sea pikeperch or sea zander, is a species of ray-finned fish in the family Percidae which includes the perches, ruffes and darters. It is found in eastern Europe in the Black Sea and the Caspian...



Santa Ana sucker

Catostomus santaanae

The Santa Ana sucker (Catostomus santaanae) is a freshwater ray-finned fish, endemic to California. It is closely related to the mountain sucker and has dark grey upper parts and silvery underparts. It grows to a maximum length of 25 cm (10 in),...



Saratoga Springs pupfish

Cyprinodon nevadensis nevadensis

The Saratoga Springs pupfish (Cyprinodon nevadensis nevadensis) is a subspecies of the Amargosa pupfish (Cyprinodon nevadensis) of the family Cyprinodontidae. The native population is endemic to Saratoga Springs, a small wetland in Death Valley...



Sarcastic fringehead

Neoclinus blanchardi

The sarcastic fringehead (Neoclinus blanchardi) is a small but hardy saltwater fish that possesses a large mouth and aggressive territorial behavior, for which it has been given its common name. They can be up to 30 centimetres (12 in) long....



Sarda chiliensis

Sarda chiliensis

Sarda chiliensis, the eastern Pacific bonito, is a marine species of bonito. It ranges from Ecuador to Chile. Sarda lineolata, which ranges from Alaska to Mexico was formerly considered a subspecies, as Sarda chiliensis lineolata, but this...



Sardinella albella

Sardinella albella

The white sardinella (Sardinella albella), also known as deep-bodied sardine, perforated-scale sardine or short-bodied sardine, is a species of ray-finned fish in the genus Sardinella. It is an important food fish, which can be feed as dried,....



Sardinella tawilis

Sardinella tawilis

Sardinella tawilis (the freshwater sardinella, freshwater herring, bombon sardine or freshwater sardine) is a freshwater sardine found exclusively in the Philippines. It is the only member of the genus Sardinella known to exist entirely in fresh...



Sardinops

Sardinops sagax

Sardinops is a monotypic genus of sardines of the family Clupeidae. The only member of the genus is Sardinops sagax. It is found in the Indo-Pacific and East Pacific oceans. Its length is up to 40 cm (16 in). It has numerous common or vernacular...



Sargassum fish

Histrio histrio

The sargassum fish, anglerfish, or frog fish (Histrio histrio) is a frogfish of the family Antennariidae, the only species in its genus. It lives among Sargassum seaweed which floats in subtropical oceans. The scientific name comes from the Latin...



Sargocentron diadema

Sargocentron diadema

Sargocentron diadema, known commonly as the crowned squirrelfish, is a species of marine ray-finned fish belonging to the family Holocentridae, which is the only family in the order Holocentriformes. Squirrelfish in general are large, active,...



Sargocentron rubrum

Sargocentron rubrum

Sargocentron rubrum, also known as redcoat, is a member of the family Holocentridae of the order Beryciformes. Squirrelfish in general are large, active, nocturnal fish which are usually red in color. It is found in the wide Indo-Pacific, from the...



Sargocentron spiniferum

Sargocentron spiniferum

Sargocentron spiniferum, common name sabre squirrelfish, giant squirrelfish and spiny squirrelfish, is a large Indo-Pacific species of squirrelfish belonging to the family Holocentridae.



Sargocentron xantherythrum

Sargocentron xantherythrum

Sargocentron xantherythrum, commonly known as Hawaiian squirrelfish or striped squirrelfish, is a member of the squirrelfish family. It is known from the Hawaiian Islands and Johnston Atoll. It is occasionally available in the aquarium trade. It...



Satanoperca jurupari

Satanoperca jurupari

Satanoperca jurupari, the demon eartheater, is a species of cichlid endemic to the Amazon basin in South America. It can reach a length of 18.5 centimetres (7.3 in) SL.



Satomi's pygmy seahorse

Hippocampus satomiae

Satomi's pygmy seahorse (Hippocampus satomiae) is the smallest known seahorse in the world with an average length of 13.8 millimetres (0.54 in) and an approximate height of 11.5 millimetres (0.45 in). This member of the family Syngnathidae is...



Saucereye porgy

Calamus calamus

The saucereye porgy (Calamus calamus) is an ocean-going species of fish in the family Sparidae. In Bermuda, they are also known as the goat's head porgy. In Jamaica, they are known as the Porgi grunt and the sugareye porgy. They may also be known...



Sauger

Sander canadensis

The sauger (Sander canadensis) is a freshwater perciform fish of the family Percidae that resembles its close relative, the walleye. The species is a member of the largest vertebrate order, the Perciformes. It is the most migratory percid species...



Saury

Cololabis adocetus

The saury (Cololabis adocetus) is a species of fish that is a member of the family Scomberesocidae, or the saury family. It is widespread in the Eastern Pacific in the surface waters, typically remaining in the top 50 centimeters of the water...



Sawback angelshark

Squatina aculeata

The sawback angelshark (Squatina aculeata) is an angelshark of the family Squatinidae. It is one of the rarest species of sharks known to date, and one of the three species of angelsharks that inhabit the Mediterranean. The Sawback angelshark lives...



Sawbwa barb

Sawbwa resplendens

The sawbwa barb (*Sawbwa resplendens*), also known as the Burmese rammy nose, Asian rummynose or rummynose rasbora, is an endangered species of cyprinid fish in the monotypic genus *Sawbwa*. The species is endemic to Inle Lake in Myanmar (Burma). It...



Scale-eye plaice

Acanthopsetta nadeshnyi

The scale-eye plaice (Acanthopsetta nadeshnyi) is a flatfish of the family Pleuronectidae. It is a demersal fish that lives at depths from between 18 metres (59 ft) to 900 metres (3,000 ft). It can reach 46 centimetres (18 in) in length and can...



Scaled sardine

Harengula jaguana

The scaled sardine, Harengula jaguana, is a herring-like fish in the family Clupeidae. It is native to coastal waters of the western Atlantic Ocean, from the Gulf of Mexico (where it is known as the pilchard or whitebait) down to Brazil where it...



Scalloped bonnethead

Sphyrna corona

The scalloped bonnethead (Sphyrna corona) is a rare, little-known species of hammerhead shark in the family Sphyrnidae. Its other common names include the mallethead shark and the crown shark. It is found in tropical and subtropical waters in the...



Scalloped hammerhead

Sphyrna lewini

The scalloped hammerhead (Sphyrna lewini) is a species of hammerhead shark in the family Sphyrnidae. It was originally known as Zygaena lewini. The Greek word sphyrna translates into "hammer" in English, referring to the shape of this shark's...



Scaly yellowfish

Labeobarbus natalensis

The scaly yellowfish, or KwaZulu-Natal yellowfish, (Labeobarbus natalensis) is a species of freshwater fish in the family Cyprinidae. This fish is found in the Tugela River System in South Africa, as well as in the Umzimkulu, Umfolozi and the...



Scamp grouper

Mycteroperca phenax

The scamp grouper (Mycteroperca phenax), also known as scamp, is a species of marine ray-finned fish, a grouper from the subfamily Epinephelinae which is part of the family Serranidae, which also includes the anthias and sea basses. It is found...



Scarlet badis

Dario dario

The scarlet badis (Dario dario) is a tropical freshwater fish and one of the smallest known percoid fish species. It is a micropredator, feeding on small aquatic crustaceans, worms, insect larvae and other zooplankton. It is sold under a variety...



Scarus frenatus

Scarus frenatus

Scarus frenatus is a species of parrotfish. Common names include bridled parrotfish, sixband or six-banded parrotfish or vermiculate parrotfish.



Scarus ghobban

Scarus ghobban

Scarus ghobban, also known as the blue-barred parrotfish, blue trim parrotfish, cream parrotfish, globe-headed parrotfish, green blotched parrotfish, yellow scale parrotfish, and bluechin parrotfish, is a species of marine ray-finned fish in the...



Scarus niger

Scarus niger

Scarus niger, common names the swarthy parrotfish, dusky parrotfish, and black parrotfish, is a species of parrotfish. It is in the phylum Chordata, class Actinopterygii, and family Scaridae. Like other members of its family, it has...



Scarus prasiognathos

Scarus prasiognathos

Scarus prasiognathos, the Singapore parrotfish, is a species of marine ray-finned fish, a parrotfish, in the family Scaridae. It is native to the eastern Indian and western Pacific Oceans, where it lives in coral reefs. This species was first...



Scarus quoyi

Scarus quoyi

Scarus quoyi, commonly known as Quoy's or greenblotch parrotfish, is a species of marine ray-finned fish, a parrotfish, in the family Scaridae. It is native to the western Pacific Ocean, where it lives in coral reefs. The species can grow to a...



Scatophagus argus

Scatophagus argus

Scatophagus argus, the spotted scat, butterfish, mia mia, spotted butterfish or tiger scat, is a species of fish in the scat family Scatophagidae. It occurs in two basic color morphs which are called green scat and ruby or red scat. This fish is...



Schelly

Coregonus stigmaticus

The schelly (Coregonus stigmaticus) is a living fresh water fish of the salmon family, endemic to four lakes in the Lake District, England. Its taxonomy is disputed with some recognizing it as a distinct species and others as a variant of the...



Schilbe intermedius

Schilbe intermedius

Schilbe intermedius or the silver butter catfish is a widespread species of African catfish. It seems closely related to Schilbe uranoscopus and these two species are sympatric over part of their ranges.



Schindleria brevipinguis

Schindleria brevipinguis

Schindleria brevipinguis is a species of marine fish in family Gobiidae of Perciformes. Known as the stout infantfish, it is native to Australia's Great Barrier Reef and to Osprey Reef in the Coral Sea.



Schindleria praematura

Schindleria praematura

Schindleria praematura, Schindler's fish is a species of neotenic goby which was formerly placed in the monogeneric family Schindleriidae but which is currently classified within the Gobiidae. It is associated with reefs and has an Indo-Pacific...



Schistura oedipus

Schistura oedipus

Schistura oedipus is a species of troglobitic stone loach endemic to caves in the Pang Mapha karst formation in Mae Hongson Province in Thailand. It is only found in very fast flowing cave streams where it feeds on micro-organism and organic...



Schizothorax nepalensis

Schizothorax nepalensis

Schizothorax nepalensis is a cyprinid fish species of the genus Schizothorax. It was first collected in 1979 in the alpine fresh water Rara Lake located in Nepal's Rara National Park.



Schizothorax plagiostomus

Schizothorax plagiostomus

Schizothorax plagiostomus is a species of ray-finned fish in the genus Schizothorax. Schizothorax plagiostomus locally known as khont, snow trout, snow carp, snow barbel and Swati fish has an elongated sub cylindrical body with short, blunt and...



School shark

Galeorhinus galeus

The school shark (Galeorhinus galeus) is a houndshark of the family Triakidae, and the only member of the genus Galeorhinus. Common names also include tope, tope shark, snapper shark, and soupfin shark. It is found worldwide in temperate seas at...



Schooling bannerfish

Heniochus diphreutes

The schooling bannerfish (Heniochus diphreutes), also known as the false moorish idol, is a marine ray-finned fish, a butterflyfish from the family Chaetodontidae. It is native to the Indo-Pacific area.



Schoolmaster snapper

Lutjanus apodus

The schoolmaster snapper (Lutjanus apodus), is a species of marine ray-finned fish, a snapper belonging to the family Lutjanidae. It is found in the western Atlantic Ocean. Like other snapper species, it is a popular food fish.



Sciades parkeri

Sciades parkeri

Sciades parkeri, the gillbacker sea catfish, is a species of sea catfish found in coastal rivers in Brazil, French Guiana, Guyana, Suriname, and Venezuela. While usually no more than 90 cm in length, it can reach lengths of up to 190 cm and weigh...



Sciaenochromis fryeri

Sciaenochromis fryeri

Sciaenochromis fryeri is a species of cichlid endemic to Lake Malawi where it is found along the rocky coastal areas at depths of 10 to 40 metres (33 to 131 ft). It can reach a length of 11.5 centimetres (4.5 in) SL. The specific name honours...



Scissortail sergeant

Abudefduf sexfasciatus

The scissortail sergeant or striptailed damselfish (Abudefduf sexfasciatus) is a large damselfish. It earns its name from the black-striped tail and sides, which are reminiscent of the insignia of a military Sergeant, being similar to those of...



Scleropages jardinii

Scleropages jardinii

Scleropages jardinii, the Gulf saratoga, Australian bonytongue, Pearl arowana or northern saratoga, is a freshwater bony fish native to Australia and New Guinea, one of two species of fishes sometimes known as Australian arowana, the other being...



Scombrops boops

Scombrops boops

Scombrops boops, the gnomefish, is a species of marine ray-finned fish, a gnomefish from the family Scombridae. It is found in the Indo-West Pacific and marginally in the south eastern Atlantic. It occurs off South Africa and Mozambique and...



Scoophead

Sphyrna media

The scoophead (Sphyrna media) is a little-known species of hammerhead shark, part of the family Sphyrnidae. It inhabits the tropical waters of the western Atlantic Ocean, from Panama to southern Brazil, and in the eastern Pacific Ocean from the...



Scophthalmus maeoticus

Scophthalmus maeoticus

Scophthalmus maeoticus (Black Sea turbot or kalkan) is a fish species in the family Scophthalmidae. It is widespread in the Black Sea. It is sometimes treated as a subspecies of the turbot, Scophthalmus maximus, which is common in the...



Scorpaena azorica

Scorpaena azorica

Scorpaena azorica, the Azores scorpionfish, is a species of marine ray-finned fish belonging to the family Scorpaenidae, the scorpionfishes. This fish occurs in the Northeastern Atlantic region in European waters. This fish is found in demersal...



Scorpaena cardinalis

Scorpaena cardinalis

Scorpaena cardinalis, the eastern red scorpionfish, grandfather hapuku, cardinal scorpionfish, Cook's scorpionfish, Cook's rockcod, Kermadec scorpionfish, Northern scorpionfish, red scorpion fish, red scorpion-cod or Sandy-bay cod, is a species...



Scorpaena guttata

Scorpaena guttata

Scorpaena guttata is a species of fish in the scorpionfish family known by the common name California scorpionfish. It is native to the eastern Pacific Ocean, where it can be found along the coast of California and Baja California. Its...



Scorpaena papillosa

Scorpaena papillosa

Scorpaena papillosa, the red rock cod, Southern red scorpionfish, chained scorpionfish, common red gurnard, Southern red scorpioncod, Southern rockcod or dwarf scorpionfish, is a species of venomous marine ray-finned fish belonging to the family...



Scorpaena scrofa

Scorpaena scrofa

Scorpaena scrofa, the red scorpionfish, bigscale scorpionfish, large-scaled scorpion fish, or rascasse is a venomous marine species of ray-finned fish in the family Scorpaenidae, the scorpionfishes. It is found in the Mediterranean Sea, in the...



Scorpaenopsis diabolus

Scorpaenopsis diabolus

Scorpaenopsis diabolus, the false stonefish, false scorpionfish or the devil scorpionfish, is a species of venomous marine ray-finned fish belonging to the family Scorpaenidae, the scorpionfishes. It has venomous spines and lives in the tropical...



Scorpaenopsis oxycephalus

Scorpaenopsis oxycephala

Scorpaenopsis oxycephalus, the tasseled scorpionfish, or small-scaled scorpionfish, is a species of venomous marine ray-finned fish belonging to the family Scorpaenidae, the scorpionfishes. It has a wide distribution in the Indian and Pacific...



Scortum barcoo

Scortum barcoo

Scortum barcoo is a species of fish in the family Terapontidae, known by the common names Barcoo grunter and jade perch. It is endemic to Australia, where it can be found in certain major rivers, including the Barcoo River. It is reared in...



Scrapermouth mbuna

Labeotropheus trewavasae

The scrapermouth mbuna (Labeotropheus trewavasae) is a species of cichlid endemic to Lake Malawi where it prefers areas with rocky substrates. This species can reach a length of 11.7 centimetres (4.6 in) TL. This species can also be found in the...



Scrawled butterflyfish

Chaetodon meyeri

The scrawled butterflyfish (Chaetodon meyeri), also known as Meyer's butterflyfish or the maypole butterflyfish, is a species of marine ray-finned fish, a butterflyfish belonging to the family Chaetodontidae. It is found in the Indian and Pacific...



Scrawled cowfish

Acanthostracion quadricornis

The scrawled cowfish (Acanthostracion quadricornis) is a species of boxfish native to the western tropical and equatorial Atlantic, as well as the Gulf of Mexico. They range in size from 8–15 inches (20–38 cm), with a maximum length of 18 inches...



Scup

Stenotomus chrysops

The scup (Stenotomus chrysops) is a fish that occurs primarily in the Atlantic from Massachusetts to South Carolina. Along with many other fish of the family Sparidae, it is also commonly known as porgy. Scup grow as large as 18 in (460 mm) and...



Scuticaria tigrina

Scuticaria tigrina

Scuticaria tigrina is a moray eel found in coral reefs in the Pacific and Indian Oceans. It is commonly known as the tiger reef-eel, tiger snake moray, tiger moray eel, tiger moray, tiger eel, spotted eel, or the spotted snake moray.



Sea goldie

Pseudanthias squamipinnis

The sea goldie (Pseudanthias squamipinnis), also known as the orange basslet, lyretail coralfish, onestripe goldie, lyretail anthias, lyretail fairy basslet, orange fairy basslet, orange seaperch, scalefin basslet, scalefin Fairy basslet and...



Sebae clownfish

Amphiprion sebae

Amphiprion sebae, also known as the sebae clownfish, is an anemonefish found in the northern Indian Ocean, from Java to the Arabian Peninsula. Like all anemonefish it is usually found living in association with sea anemones. While the common name...



Sebastes atrovirens

Sebastes atrovirens

Sebastes atrovirens, the kelp rockfish, is a species of marine ray-finned fish belonging to the subfamily Sebastinae, the rockfishes, part of the family Scorpaenidae. It is native to the Pacific Ocean along the coast of California in the United...



Sebastes capensis

Sebastes capensis

Sebastes capensis, the false jacobever or Cape redfish, is a species of marine ray-finned fish belonging to the subfamily Sebastinae, the rockfishes, part of the family Scorpaenidae. It is found in the South Atlantic Ocean and may also occur off...



Sebastes chrysomelas

Sebastes chrysomelas

Sebastes chrysomelas, commonly known as the black-and-yellow rockfish, is a marine fish species of the family Sebastidae. It is found in rocky areas in the Pacific off California and Baja California. Although it is similar in appearance to...



Sebastes inermis

Sebastes inermis

Sebastes inermis, the Japanese red seaperch or dark-banded rockfish, is a species of marine ray-finned fish belonging to the subfamily Sebastinae, the rockfishes, part of the family Scorpaenidae. It is found in the northwestern Pacific Ocean....



Sebastes mentella

Sebastes mentella

Sebastes mentella, the beaked redfish, deepwater redfin, ocean perch, Atlantic redfish, Norway haddock, red perch, golden redfish, or hemdurgan, is a species of marine ray-finned fish belonging to the subfamily Sebastinae, the rockfishes, part of...



Sebastes miniatus

Sebastes miniatus

Sebastes miniatus, the vermilion rockfish, vermilion seaperch, red snapper, red rock cod, and rasher, is a species of marine ray-finned fish belonging to the subfamily Sebastinae, the rockfishes, part of the family Scorpaenidae. It is native to...



Sebastes norvegicus

Sebastes norvegicus

Sebastes norvegicus, the rose fish, rock fish, ocean perch, Atlantic redfish, Norway haddock, golden redfish, pinkbelly rosefish or bergylt, is a species of marine ray-finned fish belonging to the subfamily Sebastinae, the rockfishes, part of the...



Sebastes schlegelii

Sebastes schlegelii

Sebastes schlegelii, also known as the Korean rockfish, northern black seaperch, and black rockfish, is a predatory species of marine ray-finned fish belonging to the subfamily Sebastinae, the rockfishes, part of the family Scorpaenidae It is...



Sebastiscus marmoratus

Sebastiscus marmoratus

Sebastiscus marmoratus, the sea ruffe, false kelpfish or dusky stingfish, is a species of marine ray-finned fish belonging to the subfamily Sebastinae, the rockfishes, part of the family Scorpaenidae. It is found in the Western Pacific from...



Securicula

Securicula gora

Securicula gora is a species of cyprinid fish found in Bangladesh, India, Pakistan and possibly in Nepal. It is the only species in its genus.



Selar boops

Selar boops

Selar boops, the oxeye scad, is a species of ray-finned fish in the family Carangidae, the scads, jacks and trevallies. It is found in the Indo-Pacific. It is an important species for fisheries in some parts of its range.



Selene setapinnis

Selene setapinnis

Selene setapinnis, the Atlantic moonfish, is a West Atlantic fish belonging to the family Carangidae.



Semotilus atromaculatus

Semotilus atromaculatus

Semotilus atromaculatus, known as the creek chub or the common creek chub, is a small minnow, a freshwater fish found in the eastern US and Canada. Differing in size and color depending on origin of development, the creek chub can usually be...



Sergeant major

Abudefduf saxatilis

The sergeant major or p ntano (Abudefduf saxatilis) is a species of damselfish. It grows to a maximum length of about 22.9 centimetres (9.0 in).



Seriola carpenteri

Seriola carpenteri

Seriola carpenteri is a species of bony fish commonly known as the Guinean amberjack (French: Sériole guinéenne, Spanish: Medregal de Guinea), which feeds on squids and fishes. It attains a size of at least 48 cm (18.9 in) fork length, and...



Seriola dorsalis

Seriola dorsalis

Seriola dorsalis, the California yellowtail is a species of ray-finned fish of the family Carangidae. This species is also known by several alternate names, such as amberjack, forktail, mossback, white salmon and yellowtail tunis or tuna or by...



Serpae tetra

Hyphessobrycon eques

The Serpae tetra (Hyphessobrycon eques), also known as the Red Minor tetra, Jewel tetra or Callistus tetra, is a species of tetra, a tropical freshwater fish of the characin family (family Characidae) of order Characiformes. It is native to the...



Serpent eel

Ophisurus serpens

The serpent eel or sand snake-eel (Ophisurus serpens) is an eel found in the eastern Atlantic Ocean, western Mediterranean Sea, western Indian Ocean, western Pacific Ocean and the north-east and west coast of North Island in New Zealand. It is an...



Serra Spanish mackerel

Scomberomorus brasiliensis

The Serra Spanish mackerel (Scomberomorus brasiliensis) is a species of fish in the family Scombridae. Specimens have been recorded at up to 125 cm in length, and weighing up to 6.71 kg. It is found in the western Atlantic, along the Caribbean...



Serranus tortugarum

Serranus tortugarum

Serranus tortugarum, the chalk bass, is a species of marine ray-finned fish, a sea bass from the subfamily Serraninae, classified as part of the family Serranidae which includes the groupers and anthias. It is found in the western Atlantic Ocean....



Serrasalmus elongatus

Serrasalmus elongatus

Serrasalmus elongatus is a slender and elongated shaped piranha from the S. humeralis group. It is commonly known as elongated piranha or pike piranha in reference to its shape. In its native range it is one of the smaller piranha species locally...



Serrasalmus geryi

Serrasalmus geryi

Serrasalmus geryi, known as the violet line piranha or Gery's piranha, is a species of piranha of the family Serrasalminae. It is found in the lower Tocantins and Araguaia Rivers of Brazil. The species is notable for being the one of only the...



Setipinna taty

Setipinna taty

The scaly hairfin anchovy (Setipinna taty) is a species of ray-finned fish in the family Engraulidae. It is found in coastal waters and estuaries in the tropical western Indo-Pacific region.



Sevan trout

Salmo ischchan

The Sevan trout (Salmo ischchan) is an endemic fish species of Lake Sevan in Armenia, known as ishkhan...



Shabout

Arabibarbus grypus

The shabout (Arabibarbus grypus) is a species of cyprinid fish also called in English, Persian or Arabic by the alternate common names shirbot and variations shabut, shabboot or shabbout, and in local languages by several other common names. It...



Shadow bass

Ambloplites ariommus

The shadow bass (Ambloplites ariommus) is a species of freshwater fish in the sunfish family (Centrarchidae) of order Perciformes. It is endemic to southeastern United States of America.



Shaggy frogfish

Antennarius hispidus

The shaggy frogfish (Antennarius hispidus), is a marine fish in the family Antennariidae.



Shark mackerel

Grammatorcynus bicarinatus

The shark mackerel (Grammatorcynus bicarinatus) is a species of Spanish mackerel (tribe Scomberomorini) in the scombrid family (Scombridae). Their maximum reported length is 112 cm (44 in), and the maximum reported weight is 13.5 kg (30 lb).This...



Sharpbelly

Hemiculter leucisculus

The sharpbelly or wild carp, sharpbelly, or common sawbelly (Hemiculter leucisculus), is a tropical freshwater and brackish water fish belonging to the Cultrinae subfamily of the family Cyprinidae. It originates in large streams and reservoirs...



Sharpfin barracuda

Sphyraena acutipinnis

The sharpfin barracuda (Sphyraena acutipinnis) is a schooling species of barracuda that inhabits lagoons, bays and seaward reefs. It is nocturnally active. It grows to 80 cm (31 in) total length, although it is commonly somewhat smaller. The...



Sharpnose sevengill shark

Heptranchias perlo

The sharpnose sevengill shark (Heptranchias perlo), also known as one-finned shark, perlon shark, sevengill cow shark, sharpsnouted sevengill or slender sevengill, is a species of shark in the family Hexanchidae, and the only living species in...



Sharpsnout stingray

Fontitrygon geijskesi

The sharpsnout stingray or wingfin stingray (Fontitrygon geijskesi) is a species of stingray in the family Dasyatidae, found from off Venezuela to northern Brazil. It inhabits shallow, brackish water, shifting towards the coast in the dry season...



Sharptooth houndshark

Triakis megalopterus

The sharptooth houndshark, or spotted gully shark (Triakis megalopterus) is a species of houndshark in the family Triakidae found in shallow inshore waters from southern Angola to South Africa. Favoring sandy areas near rocky reefs and gullies....



Sheepshead minnow

Cyprinodon variegatus

The sheepshead minnow (Cyprinodon variegatus), also known as sheepshead pupfish, is a species of ray-finned fish in the family Cyprinodontidae, the pupfishes. It is found in salt marsh and estuary environments and is native to the eastern coasts...



Sheepshead porgy

Calamus penna

The sheepshead porgy (Calamus penna), also known as the littlemouth porgy or speckled porgy, is a species of marine ray-finned fish belonging to the family Sparidae, the seabreams and porgies. This species is found in the Western Atlantic Ocean.



Shiner perch

Cymatogaster aggregata

The shiner perch (Cymatogaster aggregata) is a common surfperch found in estuaries, lagoons, and coastal streams along the Pacific coast from Alaska to Baja California. It is the sole member of its genus. The shiner perch is also known as...



Shishamo

Spirinchus lanceolatus

Shishamo (柳葉魚, literally "Willow Leaf Fish"), or Spirinchus lanceolatus, is an anadromous fish (smelt) native to Hokkaido, Japan.



Shoal bass

Micropterus cataractae

The shoal bass (Micropterus cataractae) is a species of freshwater fish in the sunfish family (Centrarchidae) of order Perciformes. One of the black basses, it is native to waters in Florida and Georgia. It is also occasionally found in rivers...



Shore rockling

Gaidropsarus mediterraneus

The shore rockling (Gaidropsarus mediterraneus) is a mottled brown, small, elongated fish. This eel-like fish has three barbels on its head, with the second dorsal fin and the anal fin running the length of most of its body. These fins may be...



Short mackerel

Rastrelliger brachysoma

The short mackerel or shortbodied mackerel (Rastrelliger brachysoma) is a species of mackerel in the family Scombridae. Its habitat is the shallow waters of Southeast Asia and Melanesia, feeding mainly on small zooplankton. It is of major...



Short-finned eel

Anguilla australis

The short-finned eel (Anguilla australis), also known as the shortfin eel, is one of the 15 species of eel in the family Anguillidae. It is native to the lakes, dams and coastal rivers of south-eastern Australia, New Zealand, and much of the...



Short-snouted seahorse

Hippocampus hippocampus

The short-snouted seahorse (*Hippocampus hippocampus*) is a species of seahorse in the family Syngnathidae. It was endemic to the Mediterranean Sea and parts of the North Atlantic, particularly around Italy and the Canary Islands. In 2007, colonies...



Short-tail nurse shark

Pseudoginglymostoma brevicaudatum

The short-tail nurse shark (*Pseudoginglymostoma brevicaudatum*) is a nurse shark in the family Ginglymostomatidae, the only member of the genus *Pseudoginglymostoma*. It is found in the tropical western Indian Ocean between latitudes 0° and 27° S,...



Short-tail stingray

Bathytoshia brevicaudata

The short-tail stingray or smooth stingray (*Bathytoshia brevicaudata*) is a common species of stingray in the family Dasyatidae. It occurs off southern Africa, typically offshore at a depth of 180–480 m (590–1,570 ft), and off southern Australia...



Short-tailed river stingray

Potamotrygon brachyura

The short-tailed river stingray (Potamotrygon brachyura) is a species of river stingray (family Potamotrygonidae) native to the Río de la Plata Basin in South America. It is sometimes known as the giant freshwater stingray, but this name is...



Shortbill spearfish

Tetrapturus angustirostris

The shortbill spearfish (Tetrapturus angustirostris), sometimes called the short-nosed spearfish, is a species of marlin native to the Indian and Pacific Oceans, with occasional records from the Atlantic Ocean. This species occurs in open waters...



Shortfin mako shark

Isurus oxyrinchus

The shortfin mako (; Māori: /'ma:ko/; Isurus oxyrinchus), also known as the shortfin mako shark, blue pointer, or bonito shark, is a large mackerel shark. It is commonly referred to as the mako shark, as is the longfin mako shark (Isurus paucus)....



Shorthead redhorse

Moxostoma macrolepidotum

The shorthead redhorse (Moxostoma macrolepidotum) is a wide-ranging species in North America. The shorthead redhorse is native to central and eastern North America. However, its range has expanded to include areas like the Hudson estuary and...



Shortjaw bonefish

Albula glossodonta

Albula glossodonta (roundjaw bonefish, shortjaw bonefish, Indo-Pacific bonefish, sharpjaw bonefish, or smallmouth bonefish) is a type of marine fish found in the Pacific Ocean. They grow up to 70 cm. Shortjaw bonefish are important to food...



Shortjaw cisco

Coregonus zenithicus

The shortjaw cisco (Coregonus zenithicus) is a North-American freshwater whitefish in the salmon family. Adult fish range to about 30 cm (12 in) in length and are silver, tinged with green above and paler below. One of the members of the broader...



Shortjaw kōkopu

Galaxias postvectis

The shortjaw kōkopu (Galaxias postvectis) is a galaxiid fish, endemic to New Zealand. They are large, scaleless fish which inhabit stable pools in cascading, bouldery streams with forest cover. Shortjaw kōkopu are amphidromous, with the fry going...



Shortnose cisco

Coregonus reighardi

The shortnose cisco (Coregonus reighardi) is a North American freshwater whitefish in the salmon family Salmonidae. One of the members of the broader Coregonus artedii species complex of ciscoes, it is native to the Great Lakes of Canada and the...



Shortnose gar

Lepisosteus platostomus

The shortnose gar (Lepisosteus platostomus) is a primitive freshwater fish of the family Lepisosteidae. It is native to the United States where its range includes the Mississippi and Missouri River basins, ranging from Montana to the west and the...



Shortnose sawshark

Pristiophorus nudipinnis

Sharks portalThe shortnose sawshark, (Pristiophorus nudipinnis) or southern sawshark, is a sawshark of the family Pristiophoridae. The body of this shark is slightly compressed, and its rostrum is narrow. The two dorsal fins are almost identical...



Shortnose sturgeon

Acipenser brevirostrum

The shortnose sturgeon (Acipenser brevirostrum) is a small and endangered species of North American sturgeon. The earliest remains of the species are from the Late Cretaceous Period, over 70 million years ago. Shortnose sturgeons are long-lived...



Shortnose sucker

Chasmistes brevirostris

The shortnose sucker (Chasmistes brevirostris) is a rare species of fish in the family Catostomidae, the suckers. This fish is native to southern Oregon and northern California in the United States. It is a federally listed endangered species of...



Shortraker rockfish

Sebastes borealis

The shortraker rockfish (Sebastes borealis) is an offshore, demersal species of marine ray-finned fish belonging to the subfamily Sebastinae, the rockfishes, part of the family Scorpaenidae. It is found in the northern Pacific Ocean.



Shortspine spurdog

Squalus mitsukurii

The shortspine spurdog (Squalus mitsukurii) is a dogfish, a member of the family Squalidae, found on continental shelves off Japan in temperate waters, from the surface to 950 m. Its length is up to 75 cm.



Shortspine thornyhead

Sebastolobus alascanus

The shortspine thornyhead (Sebastolobus alascanus), also known as the channel rockcod, shortspine channel rockfish or spinycheek rockfish, is a species of marine ray-finned fish belonging to the subfamily Sebastinae, the rockfishes, part of the...



Shoshone pupfish

Cyprinodon nevadensis shoshone

The Shoshone pupfish (Cyprinodon nevadensis shoshone) is a subspecies of Cyprinodon nevadensis from California in the United States. It is characterized by large scales and a "slab-sided," narrow, slender body, with the arch of the ventral...



Shotted halibut

Eopsetta grigorjewi

The shotted halibut (Eopsetta grigorjewi) is a flatfish of the family Pleuronectidae. It is a demersal fish that lives on sandy mud bottoms in the sublittoral zone at depths of between 60 metres (200 ft) and 1,325 metres (4,347 ft). It can reach...



Shovelnose guitarfish

Rhinobatos productus

The shovelnose guitarfish, Pseudobatos productus, is a ray in the family Rhinobatidae. It becomes mature at an estimated seven to eight years old. Males are between 90–100 cm (35–39 in) long, while females are around 99 cm (39 in) at that age. The...



Shovelnose sturgeon

Scaphirhynchus platyrhynchus

The shovelnose sturgeon (Scaphirhynchus platyrhynchus) is the smallest species of freshwater sturgeon native to North America. It is often called hackleback, sand sturgeon, or switchtail. Switchtail refers to the long filament found on the upper...



Shrimp eel

Ophichthus gomesii

The shrimp eel (Ophichthus gomesii) is an eel in the family Ophichthidae (worm/snake eels). It was described by Francis de Laporte de Castelnau in 1855. It is a common inshore species of eel usually found in the shallow Gulf of Mexico and the...



Shrimp scad

Alepes djedaba

The shrimp scad (*Alepes djedaba*) (also known as the slender yellowtail kingfish), is a species of widespread tropical marine fish of the jack family, Carangidae. The shrimp scad is widely distributed in the tropical and subtropical western Indian...



Shuttles hoppfish

Periophthalmus modestus

The shuttles hoppfish or shuttles mudskipper (*Periophthalmus modestus*) is a species of mudskippers native to fresh, marine and brackish waters of the northwestern Pacific Ocean from Vietnam to Korea and Japan. This species occurs in muddy...



Siamese algae-eater

Crossocheilus oblongus

The Siamese algae-eater (*Crossocheilus oblongus*) is a species of freshwater fish in the carp family, Cyprinidae. This bottom-dwelling tropical fish is found in mainland Southeast Asia, including the Chao Phraya and Mekong basins as well as the...



Siamese fighting fish

Betta splendens

The Siamese fighting fish (*Betta splendens*), commonly known as the betta, is a freshwater fish native to Southeast Asia, namely Cambodia, Laos, Myanmar, Malaysia, Indonesia, Thailand, and Vietnam. It is one of 73 species of the genus *Betta*, but...



Siamese mud carp

Henicorhynchus siamensis

The Siamese mud carp (*Henicorhynchus siamensis*) is a species of freshwater cyprinid fish, a variety of Asian carp native to the Mekong and Chao Phraya Rivers in Southeast Asia, especially in Cambodia, Laos and Thailand. It is very common in...



Siamese tigerfish

Datnioides pulcher

The Siamese tigerfish (Datnioides pulcher), also known as the Siamese tiger perch, is a species of ray-finned fish belonging to the family Lobotidae, the tripletails and tiger perches. This fish is endemic to Indochina and is assessed as...



Siberian sturgeon

Acipenser baerii

The Siberian sturgeon (Acipenser baerii) is a species of sturgeon in the family Acipenseridae. It is most present in all of the major Siberian river basins that drain northward into the Kara, Laptev and East Siberian Seas, including the Ob,...



Sicklefin lemon shark

Negaprion acutidens

The sicklefin lemon shark (Negaprion acutidens) or sharptooth lemon shark is a species of requiem shark belonging to the family Carcharhinidae, widely distributed in the tropical waters of the Indo-Pacific. It is closely related to the...



Sicklefin weasel shark

Hemigaleus microstoma

The sicklefin weasel shark (Hemigaleus microstoma) is an uncommon species of ground shark in the family Hemigaleidae. It is native to southern India, southern China, and parts of Southeast Asia, living in shallow waters down to a depth of 170 m...



Sicyases sanguineus

Sicyases sanguineus

Sicyases sanguineus is a species of amphibious marine clingfish in the family Gobiesocidae. It lives in the Southeast Pacific along the entire coast of Chile and southern Peru. Locally, it is known as pejesapo (literally, frog-fish). It inhabits...



Sicydium plumieri

Sicydium plumieri

Sicydium plumieri (Spanish vernacular: Olivo, Ceti; English vernacular: Sirajo Goby) is a freshwater species of the goby native to the Antilles from Cuba to Trinidad and Tobago, though not recorded from all islands. This species can reach a...



Sicyopterus stimpsoni

Sicyopterus stimpsoni

Sicyopterus stimpsoni, commonly known as the Nopoli rockclimbing goby, oopu nopili, or Stimpson's goby, is a species of amphidromous goby endemic to Hawaii. This species can reach a length of 19.8 centimetres (7.8 in) SL.



Sicyopus cebuensis

Sicyopus cebuensis

Sicyopus cebuensis is a species of fish that was first documented by Chen and Shao in 1998, who documented just a single adult pair located in the Uling brook within the upper Naga River basin on the island of Cebu, Cebu province, the...



Siganus doliatus

Siganus doliatus

Siganus doliatus, commonly known as the barred spinefoot, scribbled rabbitfish, pencil-streaked rabbitfish, barred Spanish mackerel, blue-lined rabbitfish or two-barred rabbitfish, is a species of marine ray-finned fish, a rabbitfish belonging to...



Siganus rivulatus

Siganus rivulatus

Siganus rivulatus, the marbled spinefoot, rivulated rabbitfish or surf parrotfish, is a gregarious, largely herbivorous ray-finned fish of the family Siganidae. Its natural range encompasses the western Indian Ocean and the Red Sea from where it...



Siganus sutor

Siganus sutor

Siganus sutor, the shoemaker spinefoot rabbitfish,whitespotted rabbitfish or African whitespotted rabbitfish, is a species of marine ray-finned fish, a rabbitfish belonging to the family Siganidae. It lives in Indo-Pacific coral reefs. It is...



Siganus virgatus

Siganus virgatus

Siganus virgatus, the barhead spinefoot, doublebar rabbitfish or doublebar spinefoot is a species of marine ray-finned fish, a rabbitfish belonging to the family Siganidae. It is found in the Indo-Pacific region.



Silky Shark

Carcharhinus falciformis

The Silky shark (Carcharhinus falciformis) is a species of requiem shark named for the smooth texture of its skin. It is one of the most abundant sharks in the pelagic zone and is a swift, inquisitive, and persistent hunter.



Silurus biwaensis

Silurus biwaensis

Silurus biwaensis, the giant Lake Biwa catfish, is a large predatory catfish species endemic to Lake Biwa in Japan.



Silurus soldatovi

Silurus soldatovi

Silurus soldatovi, Soldatov's catfish or northern sheatfish, is a large catfish species from the Far East.



Silver arowana

Osteoglossum bicirrhosum

The silver arowana (Osteoglossum bicirrhosum) is a South American freshwater bony fish of the family Osteoglossidae. Silver arowanas are sometimes kept in aquariums, but they are predatory and require a very large tank. The generic name...



Silver carp

Hypophthalmichthys molitrix

The silver carp (*Hypophthalmichthys molitrix*) is a species of freshwater cyprinid fish, a variety of Asian carp native to China and eastern Siberia, from the Amur River drainage in the north to the Xi Jiang River drainage in the south. Although a...



Silver chub

Macrhybopsis storeriana

The silver chub (Macrhybopsis storeriana) is a species of freshwater fish of the family Cyprinidae found in North America.



Silver cyprinid

Rastrineobola argentea

The silver cyprinid (Rastrineobola argentea) also known as the Lake Victoria sardine, mukene, and omena (native language), dagaa (Swahili) is a species of pelagic, freshwater ray-finned fish in the carp family, Cyprinidae from East Africa. It is...



Silver drummer

Kyphosus sydneyanus

The silver drummer (Kyphosus sydneyanus), also known as the buff bream, buffalo bream, buffs, common buffalo bream, drummer bream, Southern silver drummer or Sydney drummer, is a species of marine ray-finned fish, a sea chub from the family...



Silver eel

Ariosoma mellissii

The silver eel (Ariosoma mellissii), also known as the Melliss's conger, is an eel in the family Congridae (conger/garden eels). It was described by Albert Günther in 1870. It is a rare tropical, marine eel which is known solely from St. Helena,...



Silver gemfish

Rexea solandri

Rexea solandri, the silver gemfish, is a species of snake mackerel found in the southwestern Pacific Ocean around Australia and New Zealand with reports of possible records from Madagascar and Japan. Gemfish appear as infrequent, but regular...



Silver hake

Merluccius bilinearis

The silver hake, Atlantic hake, or New England hake (Merluccius bilinearis) is a merluccid hake of the genus Merluccius, found in the northwest Atlantic Ocean. It is highly predatory and typically feeds on fish and crustaceans.



Silver pearlfish

Encheliophis homei

The silver pearlfish, Encheliophis homei, is a species of eel-like fish in the family Carapidae. This pearlfish lives inside the coelom of sea cucumbers such as Bohadschia argus, Thelenota ananas , and Stichopus chloronotus. It is native to...



Silver rasbora

Rasbora argyrotaenia

The silver rasbora (*Rasbora argyrotaenia*) is a species of ray-finned fish in the genus *Rasbora*. It occurs in the Mekong, Chao Phraya and Mae Klong basins, Malay Peninsula as well as Borneo, Java and Sumatra.



Silver redhorse

Moxostoma anisurum

The silver redhorse (*Moxostoma anisurum*: *Moxostoma*= mouth to suck; *anisurum* = unequal tail) is a species of freshwater fish endemic to Canada and the United States. It is the longest-lived redhorse known (a group of 25 extant species), with a...



Silver scabbardfish

Lepidopus caudatus

The silver scabbardfish (*Lepidopus caudatus*), also known as the frostfish or beltfish is a benthopelagic cutlassfish of the family *Trichiuridae* found throughout the temperate seas of the world. It grows to over 2 metres (6 ft 7 in) in length.



Silver spinyfin

Diretmus argenteus

The silver spinyfin (*Diretmus argenteus*) is a spinyfin of the genus *Diretmus*, found around the world except the Mediterranean, at depths down to 2,000 m. It belongs to the monotypic genus *Diretmus*. Their length is between 30 and 40 cm. The silver...



Silver sweep

Scorpiis lineolata

The silver sweep (*Scorpiis lineolata*), also known as the false pompano, sweep, trumps or windawindawi, is a species of marine ray-finned fish from the subfamily *Scorpidinae* of the sea chub family *Kyphosidae*. It is native to the southwestern...



Silver warehou

Seriolella punctata

The silver warehou, *Seriolella punctata*, is a medusafish of the family Centrolophidae found in the southern Indian and southern Pacific oceans, at depths of between 100 and 650 m. Its length is up to about 65 cm.



Silver-banded whiting

Sillago argentifasciata

The silver-banded whiting (Sillago argentifasciata) is a narrowly distributed species of inshore marine fish of the smelt whiting family Sillaginidae that inhabits the coastline of a single province within the Philippines. The species' taxonomic...



Silverjaw minnow

Ericymba buccata

Silverjaw minnows (Ericymba buccata) are part of the family Cyprinidae, which includes carps and minnows. With over 300 known species, there are more species of minnows native to North America's fresh waters than any other fish. Minnows can be...



Silvertip shark

Carcharhinus albimarginatus

The silvertip shark (Carcharhinus albimarginatus) is a large species of requiem shark, in the family Carcharhinidae, with a fragmented distribution throughout the tropical Indian and Pacific Oceans. It is often encountered around offshore islands...



Silvertip tetra

Hasemania nana

The silvertip tetra (Hasemania nana) is a species of characid freshwater fish omnivores native to streams and creeks in the São Francisco basin in Brazil, but frequently kept in aquariums. Silvertip tetras are slightly more aggressive than other...



Sinarapan

Mistichthys luzonensis

The sinarapan or tabyos (*Mistichthys luzonensis*) is a species of fish in the goby subfamily, Gobionellinae, and the only member of the monotypic genus *Mistichthys*. It is endemic to the Philippines, where it occurs along the Bicol River and in...



Siniperca chuatsi

Siniperca chuatsi

Siniperca chuatsi, the mandarin fish (Chinese: 鱈魚,), is a species of freshwater ray-finned fish from the family Sinipercidae, the Oriental perches. It is the type species of the genus *Siniperca* (Chinese perches).



Six-banded tiger barb

Desmopuntius hexazona

The six-banded tiger barb (*Desmopuntius hexazona*) is a Southeast Asian species of cyprinid fish native to blackwater streams, peat swamps and other freshwater habitats with little movement in the Malay Peninsula, Sumatra and Borneo. Although...



Six-line wrasse

Pseudocheilinus hexataenia

The six-line wrasse (*Pseudocheilinus hexataenia*) is a species of marine ray-finned fish from the family Labridae which has a wide Indo-Pacific distribution. This species is associated with coral reefs and can be found in the aquarium trade.



Sixbar angelfish

Pomacanthus sexstriatus

The sixbar angelfish (*Pomacanthus sexstriatus*), also known as the six banded angelfish, is a species of marine ray-finned fish, a marine angelfish belonging to the family Pomacanthidae. It is found in Indo-Pacific region.



Sixgill sawshark

Pliotrema warreni

The sixgill sawshark, Pliotrema warreni is a sawshark of the family Pristiophoridae. Presence of 6 pairs of gill slits highlights this genus among sharks; outside Hexanchiformes order, Pliotrema is the only shark with more than 5 gill slits....



Sixgill stingray

Hexatrygon bickelli

The sixgill stingray (Hexatrygon bickelli) is a species of stingray and the only extant member of the family Hexatrygonidae. Although several species of sixgill stingrays have been described historically, they may represent variations in a...



Skilfish

Erilepis zonifer

The skilfish (Erilepis zonifer) is a species of ray-finned fish, one of two species belonging to the family Anoplopomatidae and the only species in the genus Erilepis. It is also known as black cod, though they have no relation to cod, family...



Skipjack shad

Alosa chrysochloris

The skipjack herring (Alosa chrysochloris) is a North American, migratory, fresh- and brackish water fish species in the herring family Clupeidae. The name skipjack shad comes from the fact that it is commonly seen leaping out of the water while...



Skipjack tuna

Katsuwonus pelamis

The skipjack tuna (Katsuwonus pelamis) is a medium-sized perciform fish in the tuna family, Scombridae, and is the only member of the genus Katsuwonus. It is otherwise known as katsuo, arctic bonito, mushmouth, oceanic bonito, striped tuna or...



Skunk loach

Yasuhikotakia morleti

The skunk loach, skunk botia or Hora's loach (*Yasuhikotakia morleti*; syn. *Botia morleti*, *Botia horae*) is a species of botiid loach found in the Mekong River basin in Indochina, as well as the Chao Phraya and Mae Klong basins in Thailand. The...



Sleek lates

Lates stappersii

The sleek lates (Lates stappersii) is a species of lates perch endemic to Lake Tanganyika. It is locally known as Mikebuka (Tanzania) or Mukeke (Democratic Republic of the Congo and Burundi). This species can reach a length of 45 centimetres (18...



Sleek unicornfish

Naso hexacanthus

The sleek unicornfish (Naso hexacanthus), also known as the blue-tail unicorn, Thorpe's unicornfish, blacktongue unicornfish or ‘opelu kala, is a species of marine ray-finned fish belonging to the family Acanthuridae, the surgeonfishes,...



Sleepy cod

Oxyeleotris lineolata

The sleepy cod (Oxyeleotris lineolata) is a medium-sized fish in the family Butidae, native to tropical fresh waters of northern Australia and questionably from New Guinea. It is a member of the order Perciformes, thus is unrelated to the true...



Slender bamboo shark

Chiloscyllium indicum

The slender bamboo shark, Chiloscyllium indicum, is a bamboo shark in the family Hemiscylliidae found in the Indo-West Pacific Oceans between latitudes 40° N and 10° S, and longitude 65° E and 160° E. It is harmless to humans.



Slender giant moray

Strophidon sathete

The slender giant moray or Gangetic moray, Strophidon sathete, is the longest member of the family of moray eels. It is in the genus Strophidon. The longest recorded specimen was caught in 1927, on the Maroochy River in Queensland; it measured...



Slender grouper

Anyperodon leucogrammicus

The slender grouper (Anyperodon leucogrammicus) is a species of marine ray-finned fish, a grouper from the subfamily Epinephelinae which is part of the family Serranidae, which also includes the anthias and sea basses. It is the only species in...



Slender rainbow sardine

Dussumieria elopsoides

The slender rainbow sardine (*Dussumieria elopoides*) is a small, subtropical, salt water fish of the Indian Ocean and the South China Sea which was first described by Bleeker in 1849. *Dussumieria hasselti* and *Dussumieria productissima* are...



Slender rasbora

Rasbora daniconius

The black-line rasbora or slender rasbora (*Rasbora daniconius*) (called darka at Rangpur), is a species of ray-finned fish in the genus cyprinid family. It is found in rivers of South and Southeast Asia, ranging from Sri Lanka and the Indus basin...



Slender seahorse

Hippocampus reidi

The slender seahorse or longsnout seahorse (*Hippocampus reidi*) is a species of fish in the family Syngnathidae that usually inhabits subtropical regions.



Slender smooth-hound

Gollum attenuatus

The slender smooth-hound or gollumshark (*Gollum attenuatus*) is a species of ground shark in the family Pseudotriakidae. It is endemic to the waters around New Zealand, where it is usually found close to the bottom over the continental slope at...



Slender snipe eel

Nemichthys scolopaceus

The slender snipe eel (*Nemichthys scolopaceus*), also known as the deep sea duck, is a fish that can weigh only a few ounces, yet reach 5 feet or 1.5 m in length. Features include a bird-like beak with curving tips, covered with tiny hooked teeth,...



Slender suckerfish

Phtheichthys lineatus

The slender suckerfish or lousefish (Phtheichthys lineatus) is a rare species of remora found around the world in tropical and subtropical seas, in areas like the Atlantic, Pacific, and Indian Ocean, from depths from 1 to 100 meters deep.The...



Slender sunfish

Ranzania laevis

The slender sunfish (Ranzania laevis) is a mola of the family Molidae, the only extant member of the genus Ranzania, found globally in tropical and temperate seas. Its length is up to 1 m (3.3 ft). Several stranding and mass stranding events have...



Slender tuna

Allothunnus fallai

The slender tuna, *Allothunnus fallai*, is a species of tuna, the only species in the genus *Allothunnus*, found around the world in the southern oceans between latitudes 20° and 50° South, although there are two records of probable vagrants, one in...



Slender white sardine

Escualosa elongata

The slender white sardine (Escualosa elongata) is a species of sardine (family Clupeidae) in the genus Escualosa. It was described by Thosaporn Wongratana in 1983. It is a tropical fish which was discovered at a Sunday market in Bangkok,....



Slimy sculpin

Cottus cognatus

The slimy sculpin (Cottus cognatus) is a freshwater species of fish belonging to the family Cottidae, which is the largest sculpin family. They usually inhabit cold rocky streams or lakes across North America, ranging from the Great Lakes,....



Sling-jaw wrasse

Epibulus insidiator

The slingjaw wrasse (Epibulus insidiator) is a species of wrasse from the family Labridae which is native to the tropical waters of the Indo-Pacific where it occurs around coral reefs. This species is of minor importance to local commercial...



Slippery dick

Halichoeres bivittatus

The slippery dick (Halichoeres bivittatus) is a species of wrasse native to shallow, tropical waters of the western Atlantic Ocean.



Sliteye shark

Loxodon macrorhinus

The sliteye shark (Loxodon macrorhinus) is a species of requiem shark, in the family Carcharhinidae, and the only member of its genus. It is found in the tropical waters of the Indo-West Pacific between latitudes 34° N and 30° S, from depths of 7...



Sloane's viperfish

Chauliodus sloani

Sloane's viperfish, Chauliodus sloani, is a predatory mesopelagic dragonfish found in waters across the world. The species was first described by German scientists Marcus Elieser Bloch and Johann Gottlob Schneider in their 1801 book Systema...



Small red scorpionfish

Scorpaena notata

Small red scorpionfish (Scorpaena notata) is a venomous Scorpionfish, common in marine subtropical waters. It is widespread in the Eastern Atlantic from the Bay of Biscay to Senegal, Madeira, Azores and the Canary Islands, including the...



Small snakehead

Channa asiatica

The small snakehead (Channa asiatica) is a species of snakehead. It is one of four species of the genus Channa native to China. It also can be found in Taiwan and southern Japan, to which it migrated (or was introduced). It is a medium-sized...



Small-scale whiting

Sillago parvisquamis

The small-scale whiting (*Sillago parvisquamis*), also known as the blue whiting, is a species of inshore marine fish of the smelt-whiting family Sillaginidae. The small-scale whiting is very similar in body shape and colour to other species in the...



Small-spotted catshark

Scyliorhinus canicula

The small-spotted catshark (*Scyliorhinus canicula*), also known as the sandy dogfish, lesser-spotted dogfish, rough-hound or morgay (in Scotland and Cornwall), is a catshark of the family Scyliorhinidae. It is found on the continental shelves and...



Smalleye hammerhead

Sphyrna tudes

The smalleye hammerhead (Sphyrna tudes), also called the golden hammerhead or curry shark, is a small species of hammerhead shark in the family Sphyrnidae. This species was historically common in the shallow coastal waters of the western Atlantic...



Smalleye stingray

Megatrygon microps

The smalleye stingray (Megatrygon microps) is a large species of stingray in the family Dasyatidae, measuring up to 2.2 m (7.2 ft) across. Rare but widely distributed, it is found in the Indo-Pacific from Mozambique to India to northern...



Smallmouth bass

Micropterus dolomieu

The smallmouth bass (Micropterus dolomieu) is a species of freshwater fish in the sunfish family (Centrarchidae) of the order Perciformes. It is the type species of its genus Micropterus (black basses), and is a popular game fish sought by...



Smallmouth buffalo

Ictiobus bubalus

The smallmouth buffalo (Ictiobus bubalus, from the Greek for "bull-fish" and "buffalo") is one of the longest-lived fishes, capable of living more than a century. They are a catostomid fish species native to the major tributaries and surrounding...



Smallmouth yellowfish

Labeobarbus aeneus

The smallmouth yellowfish (Labeobarbus aeneus) is a species of ray-finned fish in the genus Labeobarbus. It has become an invasive species in rivers of the Eastern Cape, South Africa, such as the Mbhashe River.



Smallscale archerfish

Toxotes microlepis

The smallscale archerfish (Toxotes microlepis) is a perciform fish of genus Toxotes. As its name suggests, the scales of the smallscale archerfish are smaller than those of other archerfish. They reach a maximum length of 15 centimetres (5.9 in)....



Smallspine spookfish

Harriotta haeckeli

The smallspine spookfish (Harriotta haeckeli) is a species of fish in the family Rhinochimaeridae with a rather disjunct population. Its natural habitat is open seas.



Smallspotted dart

Trachinotus baillonii

The smallspotted dart (Trachinotus baillonii) is an Indo-Pacific species of pompano in the family Carangidae.



Smalltail shark

Carcharhinus porosus

The smalltail shark (Carcharhinus porosus) is a species of requiem shark, and part of the family Carcharhinidae. It is found in the western Atlantic Ocean, from the northern Gulf of Mexico to southern Brazil. It inhabits shallow waters close to...



Smalltooth sand tiger

Odontaspis ferox

The smalltooth sand tiger or bumpytail ragged-tooth (Odontaspis ferox) is a species of mackerel shark in the family Odontaspidae, with a patchy but worldwide distribution in tropical and warm temperate waters. They usually inhabit deepwater...



Smalltooth sawfish

Pristis pectinata

The smalltooth sawfish (Pristis pectinata) is a species of sawfish in the family Pristidae. It is found in shallow tropical and subtropical waters in coastal and estuarine parts of the Atlantic. Reports from elsewhere are now believed to be...



Smooth butterfly ray

Gymnura micrura

The smooth butterfly ray (Gymnura micrura) is a species of cartilaginous fish in the family Gymnuridae. It is a member of the order Myliobatiformes, which contains 10 total families. Its natural habitats are shallow seas, subtidal aquatic beds,...



Smooth hammerhead

Sphyrna zygaena

The smooth hammerhead (*Sphyrna zygaena*) is a species of hammerhead shark, and part of the family Sphyrnidae. This species is named "smooth hammerhead" because of the distinctive shape of the head, which is flattened and laterally extended into a...



Smooth lanternshark

Etmopterus pusillus

The smooth lanternshark or slender lanternshark (*Etmopterus pusillus*) is a species of dogfish shark in the family Etmopteridae, found widely in the Atlantic and Pacific Oceans. It inhabits benthic environments at a depth of 274–1,000 m (899–3,281...



Smooth toadfish

Tetractenos glaber

The smooth toadfish (*Tetractenos glaber*) is a species of fish in the pufferfish family Tetraodontidae. It is native to shallow coastal and estuarine waters of southeastern Australia, where it is widespread and abundant. French naturalist...



Smooth trunkfish

Lactophrys triqueter

Lactophrys triqueter also known as the smooth trunkfish, is a species of boxfish found on and near reefs in the Caribbean Sea, Gulf of Mexico and subtropical parts of the western Atlantic Ocean.



Smoothback angelshark

Squatina oculata

Sharks portalThe smoothback angelshark (*Squatina oculata*) is an angelshark of the family Squatinidae found in the eastern Atlantic and Mediterranean.



Smoothtooth blacktip shark

Carcharhinus leiodon

The smoothtooth blacktip shark (Carcharhinus leiodon) is a species of requiem shark in the family Carcharhinidae. It is known only from the type specimen caught from the Gulf of Aden, off eastern Yemen, and a handful of additional specimens...



Snaggletooth shark

Hemipristis elongata

The snaggletooth shark, or fossil shark (Hemipristis elongata), is a species of weasel shark in the family Hemigaleidae, and the only extant member of the genus Hemipristis. It is found in the Indo-West Pacific, including the Red Sea, from...



Snail darter

Percina tanasi

The snail darter (Percina tanasi) is a small species of freshwater ray-finned fish, a darter from the subfamily Etheostomatinae, part of the family Percidae, which also contains the perches, ruffes and pikeperches. It is found in East Tennessee...



Snake mackerel

Gempylus serpens

The snake mackerel (Gempylus serpens) is the sole species of fish in the monotypic genus Gempylus, belonging to the family Gempylidae (which is also referred to generally as "snake mackerels"). It is found worldwide in tropical and subtropical...



Snake pipefish

Entelurus aequoreus

The snake pipefish (*Entelurus aequoreus*) is a species of pipefish, from the family Syngnathidae, native to the northeastern Atlantic Ocean where they are generally found amongst algae close in to shore. It is the largest species of pipefish...



Snake River fine-spotted cutthroat trout

Oncorhynchus clarkii behnkei

The Snake River fine-spotted cutthroat trout is a form of the cutthroat trout (*Oncorhynchus clarkii*) that is considered either as a separate subspecies *O. c. behnkei*, or as a variety of the Yellowstone cutthroat trout (*O. c. bouvieri*). The fish...



Snakeskin gourami

Trichopodus pectoralis

The snakeskin gourami (Trichopodus pectoralis) is a species of gourami native to Southeast Asia. Prior to the merging of Belontiidae to the family Osphronemidae, the snakeskin gourami was regarded as the largest member of its family. It is still...



Snowflake moray

Echidna nebulosa

The snowflake moray (Echidna nebulosa), also known as the clouded moray among many vernacular names, is a species of marine eel of the family Muraenidae. It has blunt teeth ideal for its diet of crustaceans, a trait it shares with the zebra moray...



Snowy grouper

Hyporthodus niveatus

The snowy grouper (Hyporthodus niveatus) is a species of marine ray-finned fish, a grouper from the subfamily Epinephelinae which is part of the family Serranidae, which also includes the anthias and sea basses. It is found in the western...



Snubnosed eel

Simenchelys parasitica

The snubnosed eel, Simenchelys parasitica, also known as the pug-nosed eel, slime eel, or snub-nose parasitic eel, is a species of deep-sea eel and the only member of its genus. Some authors classify it as the sole member of the subfamily...



So-iuy mullet

Planiliza haematocheila

The so-iuy mullet (Planiliza haematocheilus), also known as the haarder, redlip mullet or so-iny mullet, is a species of fish in the family Mugilidae. Previously the species was included in the genus Mugil (as Mugil soiuy), but is now considered a...



Sohal surgeonfish

Acanthurus sohal

The sohal surgeonfish (Acanthurus sohal) or sohal tang, is a species of marine ray-finned fish belonging to the family Acanthuridae, which includes the surgeonfishes, unicornfishes and tangs. This fish is found in the northwestern Indian Ocean.



Solea senegalensis

Solea senegalensis

Solea senegalensis, the Senegalese sole, is a species of flatfish from the family of the true soles, the Soleidae, from the eastern Atlantic and the Mediterranean Sea.



Solenostomus paradoxus

Solenostomus paradoxus

The ornate ghost pipefish or harlequin ghost pipefish, Solenostomus paradoxus, is a false pipefish of the family Solenostomidae. The species' name comes from the Greek paradoxos, referring to this fish's unusual external features. Ornate ghost...



Sonora sucker

Catostomus insignis

The Sonora Sucker (Gila Sucker), *Catostomus insignis*, is a medium-sized catostomid fish with 16 other species in the genus scattered throughout North America. This species is remarkably similar in appearance to the Yaqui Sucker (*C. bernardini*).



Sonoyta pupfish

Cyprinodon eremus

The Sonoyta pupfish or Quitobaquito pupfish (Cyprinodon eremus) is an endangered species of pupfish from Sonora in Mexico and Arizona in the United States.Originally, it was considered to be a subspecies of the desert pupfish (C. macularius), but...



Sooty grunter

Hephaestus fuliginosus

The sooty grunter (Hephaestus fuliginosus), also known by the name black bream, blubberlips, Northern grunter or purple grunter, is a species of freshwater ray-finned fish, a grunter from the family Terapontidae. It inhabits coastal and inland...



South African mullet

Chelon richardsonii

The South African mullet (Chelon richardsonii), also called a harder mullet or simply harder, is a species of mullet. It is found in South African coastal waters from Walvis Bay (Namibia) to KwaZulu-Natal, and grows to a maximum length of 40.5...



Southern African anchovy

Engraulis capensis

The southern African anchovy (Engraulis capensis) is a species of anchovy which occurs in the southeast Atlantic Ocean near Namibia and South Africa.



Southern African frilled shark

Chlamydoselachus africana

The southern African frilled shark (Chlamydoselachus africana) is a species of shark in the family Chlamydoselachidae, described in 2009. It is found in the deep waters off southern Angola to southern Namibia. This species is difficult to...



Southern blue devil

Paraplesiops meleagris

The southern blue devil (Paraplesiops meleagris) is a species of fish in the longfin family Plesiopidae endemic to southern Australia. It is a close relative of the eastern blue devil (Paraplesiops bleekeri), which lives in the coastal waters of...



Southern blue whiting

Micromesistius australis

The southern blue whiting (Micromesistius australis) is a codfish of the genus Micromesistius, found in the southern oceans with temperatures between 3 and 7 °C, at depths of 50 to 900 m. Its length is commonly between 30 and 60 cm, with a...



Southern bluefin tuna

Thunnus maccoyii

The southern bluefin tuna (Thunnus maccoyii) is a tuna of the family Scombridae found in open southern Hemisphere waters of all the world's oceans mainly between 30°S and 50°S, to nearly 60°S. At up to 2.5 metres (8 ft 2 in) and weighing up to...



Southern conger

Conger verreauxi

The southern conger (Conger verreauxi) is a conger of the family Congridae, found in the eastern Indian Ocean and south-western Pacific Ocean, including southern Australia and New Zealand, at depths down to 100 m in broken rocky reef areas....



Southern eagle ray

Myliobatis goodei

The Southern eagle ray (Myliobatis goodei), sometimes known as the Southern eagle fish or the rockfish, is a ray species in the family Myliobatidae. It lives in waters just off of the Atlantic coast, from the tip of Florida down to Argentina,...



Southern platyfish

Xiphophorus maculatus

The southern platyfish, common platy, or moonfish (Xiphophorus maculatus) is a species of freshwater fish in family Poeciliidae of order Cyprinodontiformes. A live-bearer, it is closely related to the green swordtail (X. hellerii) and can...



Southern puffer

Sphoeroides nephelus

The southern puffer (Sphoeroides nephelus) is a species of pufferfish in the family Tetraodontidae. It is found in the Caribbean Sea, and possibly further south on the coast of Brazil.



Southern pygmy perch

Nannoperca australis

The southern pygmy perch (Nannoperca australis), also known as the Tasmanian pygmy perch, is a species of freshwater ray-finned fish, a temperate perch from the family Percichthyidae which is native to south-eastern Australia and Tasmania.



Southern redbelly dace

Chrosomus erythrogaster

The southern redbelly dace (*Chrosomus erythrogaster*), is a North American species of temperate freshwater fish of the family Cyprinidae. The natural geographic range extends from Western New York to Minnesota, and south to Oklahoma, Arkansas, and...



Southern saratoga

Scleropages leichardti

The southern saratoga (*Scleropages leichardti*), also known as the spotted bonytongue, spotted saratoga, or simply saratoga, is a freshwater bony fish native to Australia. It belongs to the subfamily Osteoglossinae, or arowanas, a primitive group...



Southern school whiting

Sillago bassensis

The southern school whiting (*Sillago bassensis*), also known as the silver whiting or trawl whiting, is a common species of coastal marine fish of the smelt-whiting family that inhabits the south and south-west coasts of Australia. Its distribution...



Southern sennet

Sphyraena picudilla

The southern sennet (Sphyraena picudilla) is an ocean-going species of game fish in the barracuda family, Sphyraenidae. It was described by the Cuban zoologist Felipe Poey. The description was part of a two-volume work, which Poey published in...



Southern sleeper shark

Somniosus antarcticus

Sharks portalThe southern sleeper shark, or Whitley's sleeper shark (Somniosus antarcticus), is a deepwater benthopelagic sleeper shark of the family Somniosidae found in the southern and subantarctic extremes of the Atlantic, Indian and Pacific...



Southern stargazer

Astroscopus y-graecum

The southern stargazer (Astroscopus y-graecum) is a species of marine fish in the family Uranoscopidae and genus Astroscopus. They are native to the United States.



Southern stingray

Hypanus americanus

The southern stingray (Hypanus americanus) is a whiptail stingray found in tropical and subtropical waters of the Western Atlantic Ocean from New Jersey to southern Brazil. It has a flat, diamond-shaped disc, with a mud brown, olive, and grey...



Spadenose shark

Scoliodon laticaudus

The spadenose shark (Scoliodon laticaudus) is a species of requiem shark in the family Carcharhinidae. It is common in the tropical Indian and western Pacific Oceans, where it forms large schools in shallow water. A small shark reaching a length...



Spanish hogfish

Bodianus rufus

The Spanish hogfish, Bodianus rufus, is a species of wrasse native to the western Atlantic Ocean, where it can be found from North Carolina and Bermuda through the Caribbean and the Gulf of Mexico to southern Brazil. It inhabits coral or rock...



Spanish toothcarp

Aphanius iberus

The Spanish toothcarp (Aphanius iberus), also known as the Spanish pupfish or Iberian killifish, is a small, endemic species of fish in the family Cyprinodontidae. Its risk of extinction is one of the greatest of any Iberian vertebrate. Its...



Spanner barb

Barbodes lateristriga

Barbodes lateristriga, the spanner barb or T-barb, is a species of cyprinid fish native to the Malay Peninsula and the island of Borneo. It can reach a length of 18 centimetres (7.1 in) TL. This species can also be found in the aquarium trade.



Sparisoma cretense

Sparisoma cretense

The Mediterranean parrotfish (Sparisoma cretense) is a species of parrotfish found at depths up to 50 m (160 ft) along rocky shores in the Mediterranean and the eastern Atlantic, from Portugal south to Senegal. It is generally common, but...



Spearfish remora

Remora brachyptera

The spearfish remora (Remora brachyptera) is a species of remora with a worldwide distribution in tropical and subtropical seas. Remoras attach themselves to other fish with a sucker on the head and this fish is almost exclusively found living on...



Speartooth shark

Glyphis glyphis

The speartooth shark (*Glyphis glyphis*) is a rare species of river shark, belonging to the family Carcharhinidae. It inhabits coastal marine waters and tidal reaches of large tropical rivers in northern Australia and New Guinea. Despite being a...



Speckled dace

Rhinichthys osculus

The speckled dace (*Rhinichthys osculus*), also known as the spotted dace and the carpita pinta, is a member of the minnow family. It is found in temperate freshwater in North America, from Sonora, Mexico to British Columbia, Canada. Canada is the...



Speckled hind

Epinephelus drummondhayi

The speckled hind (*Epinephelus drummondhayi*), also known as the calico grouper, kitty mitchell or strawberry grouper, is a species of marine ray-finned fish, a grouper from the subfamily Epinephelinae which is part of the family Serranidae, which...



Speckled longfin eel

Anguilla reinhardtii

The speckled longfin eel, Australian long-finned eel or marbled eel (*Anguilla reinhardtii*) is one of 15 species of eel in the family Anguillidae. It has a long snake-like cylindrical body with its dorsal, tail and anal fins joined to form one...



Spectrunculus grandis

Spectrunculus grandis

Spectrunculus grandis is a species of Rhizopharyngia ray-finned fish in the cusk-eel family known by the common names pudgy cusk-eel and giant cusk-eel. It is one of two species in the formerly monotypic genus *Spectrunculus*, the other species, *S....*



Sperata aor

Sperata aor

Sperata aor, the long-whiskered catfish (Bengali: আড় ar, Assamese: আঁৰি ãri), is a species of catfish of the family Bagridae found in southern Asia in the nations of India, Pakistan, Nepal, Bangladesh and Myanmar. It grows to a length of 180...



Sperata seenghala

Sperata seenghala

Sperata seenghala, the Giant river-catfish, is a species of bagrid catfish. It is known locally as Guizza, Guizza ayer, Auri, Ari, Pogal, Singhara and Seenghala, among other names. It is found in southern Asia in the countries of Afghanistan,...



Sphaeramia nematoptera

Sphaeramia nematoptera

The pajama cardinalfish, spotted cardinalfish, coral cardinalfish, or polkadot cardinalfish (Sphaeramia nematoptera) is a species of fish belonging to the family Apogonidae. It is a popular aquarium fish. It grows to a total length around 8.5 cm...



Sphaeramia orbicularis

Sphaeramia orbicularis

The orbiculate cardinalfish, chubby cardinal, or polka-dot cardinalfish (Sphaeramia orbicularis) is a species of Apogonidae (cardinalfishes). It grows to about 10 cm total length, and has a thin, dark vertical 'waistband' with scattered dark...



Sphoeroides annulatus

Sphoeroides annulatus

Sphoeroides annulatus (bullseye puffer) is a species in the family Tetraodontidae, or pufferfishes. It is found in the eastern Pacific Ocean from California, USA to Pisco, Peru and the Galápagos Islands.



Sphyraena chrysotaenia

Sphyraena chrysotaenia

Sphyraena chrysotaenia, the yellowstripe barracuda, is a species of predatory, ray finned fish from the family Sphyraenidae which is found in the Indo-West Pacific region. It has entered the Mediterranean Sea from the Red Sea through the Suez...



Sphyraena helleri

Sphyraena helleri

Sphyraena helleri, the Heller's barracuda, is a schooling species of barracuda in the family Sphyraenidae. The species is named in honor of zoologist Edmund Heller



Sphyraena sphyraena

Sphyraena sphyraena

Sphyraena sphyraena, also known as the European barracuda or Mediterranean barracuda, is a ray-finned predatory fish of the Mediterranean basin and the warmer waters of the Atlantic Ocean.



Sphyraena viridensis

Sphyraena viridensis

Sphyraena viridensis, the yellowmouth barracuda or yellow barracuda is a predatory ray-finned fish from the family Sphyraenidae, the barracudas. It is found in the warmer waters of the eastern Atlantic Ocean and the Mediterranean. It is often...



Spined loach

Cobitis taenia

The spined loach (Cobitis taenia) is a common freshwater fish in Europe. It is sometimes known as spotted weather loach, not to be confused with the "typical" weather loaches of the genus Misgurnus. This is the type species of the spiny loach...



Spined pygmy shark

Squaliolus laticaudus

The spined pygmy shark (Squaliolus laticaudus) is a species of squaliform shark in the family Dalatiidae found widely in all oceans. Growing no larger than roughly 28 cm (11 in), it is one of the smallest sharks alive, with this record beaten by...



Spinetail mobula

Mobula japanica

The spinetail devil ray (Mobula japanica), also known as the spinetail mobula ray or Japanese mobula ray, is a species of pelagic marine fish which belongs to the family Mobulidae. It is found throughout the tropical and sub-tropical waters of...



Spinner shark

Carcharhinus brevipinna

The spinner shark (Carcharhinus brevipinna) is a type of requiem shark, in the family Carcharhinidae, named for the spinning leaps it makes as a part of its feeding strategy. This species occurs in tropical and warm temperate waters worldwide,...



Spiny butterfly ray

Gymnura altavela

The spiny butterfly ray or giant butterfly ray (Gymnura altavela) is a species of butterfly ray, family Gymnuridae, native to the shallow coastal waters of the Atlantic Ocean. A large ray that can measure over 2 m (6 ft 7 in) across, it may be...



Spiny chromis

Acanthochromis polyacanthus

The spiny chromis (Acanthochromis polyacanthus) is a species of damselfish from the western Pacific. It is the only member of the genus Acanthochromis.



Spiny Dogfish

Squalus acanthias

The Spiny dogfish (Squalus acanthias) is one of the best-known representatives of the shark family. It is distinguished by two spines (one anterior to each dorsal fin) and no anal fin. It lives in shallow waters and further offshore in most...



Spiny icefish

Chaenodraco wilsoni

Chaenodraco wilsoni, the spiny icefish, is a species of marine ray-finned fish belonging to the family Channichthyidae, the crocodile icefishes. It is native to the Southern Ocean. This species is the only known member of its genus. It is of...



Spiny red gurnard

Chelidonichthys spinosus

The spiny red gurnard (*Chelidonichthys spinosus*) is a species of marine ray-finned fish belonging to the family Triglidae, the gurnards and sea robins. This species is found in the northwestern Pacific Ocean where they occur at depths of from 25...



Spiny seahorse

Hippocampus histrix

The spiny seahorse (*Hippocampus histrix*), also referred to as the thorny seahorse, is a small marine fish in the family Syngnathidae, native to the Indo-Pacific area. It is classified as a Vulnerable species by the IUCN.



Splendid alfonsino

Beryx splendens

The splendid alfonsino (*Beryx splendens*) is an alfonsino of the genus *Beryx*, found around the world at depths between 25 and 1,250 metres (82 and 4,101 ft), usually between 400 and 600 metres (1,300 and 2,000 ft). Although its most common size is...



Splendid toadfish

Sanopus splendidus

The splendid toadfish (*Sanopus splendidus*) also called the coral toadfish and the Cozumel splendid toadfish is a species of toadfish once believed to be entirely endemic to the island of Cozumel but have been found on the reefs of Honduras all...



Spoonhead sculpin

Cottus ricei

The spoonhead sculpin (Cottus ricei) is a species of freshwater ray-finned fish belonging to the family Cottidae, the typical sculpins. This species is widespread in northeastern North America.



Spot

Leiostomus xanthurus

The spot (Leiostomus xanthurus), also known commonly as the spot croaker, Norfolk spot and the Virginia spot, is a species of small short-lived saltwater fish in the family Sciaenidae. The species inhabits estuary and coastal waters from...



Spot-tail shark

Carcharhinus sorrah

The spot-tail shark, or sorrah shark (Carcharhinus sorrah), is a species of requiem shark, in the family Carcharhinidae, found in the tropical Indo-West Pacific Ocean between latitudes 31°N and 31°S from the surface to a depth around 72 m (236...



Spotbanded scat

Selenotoca multifasciata

The spotbanded scat,(Selenotoca multifasciata), also known as the striped scat, banded scat, barred scat, butterfish, John Dory, Johnny Dory, old maid, Southern butter-fish or striped butterfish, is a species of ray-finned fish, belonging to the...



Spotfin betta

Betta macrostoma

The spotfin betta or Brunei beauty (Betta macrostoma) is a species of labyrinth fish endemic to the island of Borneo. It generally inhabits pools of very slow-moving water along rivers and streams in very shallow waters of 30 cm (12 in) or less....



Spotfin butterflyfish

Chaetodon ocellatus

The spotfin butterflyfish (Chaetodon ocellatus) is species of marine ray-finned fish, a butterflyfish from the family Chaetodontidae. It is found in the western Atlantic Ocean, in the Gulf of Mexico and most commonly found in the Caribbean...



Spotfin croaker

Roncador stearnsii

The spotfin croaker (Roncador stearnsii) is a species of marine ray-finned fish belonging o the family Sciaenidae, the drums and croakers. This species occurs from Mazatlán, Mexico, to Point Conception, California, including the Gulf of...



Spotfin hogfish

Bodianus pulchellus

The spotfin hogfish or Cuban hogfish (Bodianus pulchellus) is a species of wrasse native to the Atlantic Ocean, where it is mainly found from North Carolina, United States, through the Caribbean to Brazil. It has also been recorded from São Tomé...



Spotless smooth-hound

Mustelus griseus

Sharks portalThe spotless smooth-hound (*Mustelus griseus*) is a species of houndshark, in the family Triakidae, found on the continental shelves of the northwest Pacific, between latitudes 40° N and 11° N, from the surface to a depth of 300 m. It...



Spottail pinfish

Diplodus holbrookii

The spottail pinfish (*Diplodus holbrookii*) is an ocean-going species of fish in the family Sparidae. It is also known as the spottail seabream or spottail pinkish porgy. Along with other members of their family, Spottail pinfish are occasionally...



Spottail shiner

Notropis hudsonius

The spottail shiner or spottail minnow (*Notropis hudsonius*) is a small- to medium-sized freshwater minnow. It can be found as far north as Canada and as far south as the Chattahoochee River in Georgia. These shiners live in lakes, rivers, and...



Spotted barb

Barbodes binotatus

Barbodes binotatus, commonly known as the spotted barb or common barb, is a tropical species of cyprinid fish endemic to Java, Indonesia.



Spotted bass

Micropterus punctulatus

The spotted bass (*Micropterus punctulatus*), also called spotty, or spots in various fishing communities, is a species of North American freshwater fish belonging to the sunfish family (Centrarchidae) of the order Perciformes. It is noted for the...



Spotted betta

Betta picta

The spotted betta or Java fighting fish (Betta picta) is a species of gourami endemic to Indonesia where it inhabits relatively cool, 22–25 °C (72–75 °F), highland streams of Java. In 1967, D. S. Johnson reported the species to be common in...



Spotted blue-eye

Pseudomugil gertrudae

The spotted blue-eye (Pseudomugil gertrudae) is a species of fish in the family Pseudomugilidae. It is native to Papua New Guinea and northern Australia. It adapts readily to captivity and can be kept in a small freshwater aquarium.



Spotted boxfish

Ostracion meleagris

The spotted or white-spotted boxfish (Ostracion meleagris), is a species of boxfish found in the Indian and Pacific Oceans. It is found on reefs at depths of from 1 to 30 metres (3.3 to 98.4 feet). This species grows to a length of 25 centimetres...



Spotted danio

Danio nigrofasciatus

The spotted danio or dwarf danio (Danio nigrofasciatus) is a tropical fish belonging to the minnow family (Cyprinidae). Originating in northern Myanmar, this fish is sometimes found in community tanks by fish-keeping hobbyists. It grows to a...



Spotted drum

Equetus punctatus

The spotted drum or spotted ribbonfish (Eques punctatus), is a species of marine ray-finned fish belonging to the family Sciaenidae, the drums and croakers. This species is found in the western Atlantic Ocean.



Spotted eagle ray

Aetobatus narinari

The spotted eagle ray (Aetobatus narinari) is a cartilaginous fish of the eagle ray family, Aetobatidae. As traditionally recognized, it is found globally in tropical regions, including the Atlantic, Pacific and Indian Oceans. Recent authorities...



Spotted estuary smooth-hound

Mustelus lenticulatus

Sharks portalThe spotted estuary smooth-hound or rig (*Mustelus lenticulatus*) is a houndshark of the family Triakidae, found on the continental shelves and in estuaries around New Zealand. It is closely related to the gummy shark (*Mustelus...*



Spotted fanfish

Pteraclis velifera

The spotted fanfish (*Pteraclis velifera*) is a species of pomfret found in the Indian and western Pacific oceans from South Africa to New Zealand. It is found at depths to 500 metres (1,600 ft). It reaches a length of 50 centimetres (20 in) TL.



Spotted galaxias

Galaxias truttaceus

Spotted galaxias (*Galaxias truttaceus*) is a largish, primarily-freshwater galaxias species found in southern Australia. Spotted galaxias are perhaps the most beautiful of the Australian galaxias species. They are a somewhat tubular, deep-bodied...



Spotted gar

Lepisosteus oculatus

The spotted gar (*Lepisosteus oculatus*) is a freshwater fish native to North America that has an abundance of dark spots on its head, fins, and dart-like body. Spotted gar have an elongated mouth with many needle-like teeth to catch other fish and...



Spotted garden eel

Heteroconger hassi

The spotted garden eel (*Heteroconger hassi*) is a heteroconger belonging to the family Congridae. It is native to the Indo-Pacific.



Spotted halibut

Verasper variegatus

The spotted halibut (*Verasper variegatus*) is a flatfish of the family Pleuronectidae. It is a demersal fish that lives on sandy, muddy bottoms in the sublittoral coastal zone at depths of up to 100 metres (330 ft). It can reach 60 centimetres (24...



Spotted handfish

Brachionichthys hirsutus

The spotted handfish (*Brachionichthys hirsutus*) is a rare Australian fish in the handfish family, *Brachionichthyidae*, classified as critically endangered on the IUCN Red List 2020. It has a highly restricted range, being found only in the estuary...



Spotted houndshark

Triakis maculata

Sharks portalThe spotted houndshark, *Triakis maculata*, is a houndshark of the family *Triakidae* found in tropical waters in the eastern Pacific Ocean off the coast of South America. It usually grows to a length around 180 centimetres (71 in). The...



Spotted moray

Gymnothorax moringa

The spotted moray (*Gymnothorax moringa*) is a medium to large moray eel. Other common names include conger, spotted eel, red moray, speckled moray, white cong, white jawed moray, white-chinned moray and white-jawed moray eel. Spotted eels have a...



Spotted moray eel

Gymnothorax isingteena

The spotted moray eel (*Gymnothorax isingteena*) is a moray eel found in coral reefs in the Pacific and Indian Oceans. It was first named by John Richardson in 1845.



Spotted prawn goby

Amblyeleotris guttata

Amblyeleotris guttata, the spotted prawn-goby is a species of goby native to reefs of the Western Pacific Ocean, that includes the Philippines to Tonga, north to the Ryukyu Islands, south to Australia. This species can reach a length of 11...



Spotted ratfish

Hydrolagus colliei

The spotted ratfish (*Hydrolagus coliei*) is a chimaera found in the north-eastern Pacific Ocean. Often seen by divers at night in the Pacific Northwest, this cartilaginous fish gets its characteristic name from a pointed rat-like tail. The...



Spotted ray

Raja montagui

The spotted ray or spotted skate (*Raja montagui*) is a species of skate in the family Rajidae.



Spotted sand bass

Paralabrax maculatofasciatus

The spotted bay bass (Paralabrax maculatofasciatus) is a species of marine ray-finned fish, a sea bass from the subfamily Serraninae, classified as part of the family Serranidae which includes the groupers and anthias. It is found in the central...



Spotted seabass

Dicentrarchus punctatus

The spotted seabass (Dicentrarchus punctatus) is a species of ray-finned fish belonging to the family Moronidae, the temperate basses. This species is found in the marine and brackish waters of the coastal eastern Atlantic Ocean from the English...



Spotted sucker

Minytrema melanops

The spotted sucker (Minytrema melanops) is a species of sucker (fish) that is native to eastern North America. The spotted sucker inhabits deep pools of small to medium rivers over clay, sand or gravel. They are occasionally found in creeks and...



Spotted sunfish

Lepomis punctatus

The spotted sunfish (Lepomis punctatus), also known as a stumpknocker, is a member of the freshwater sunfish family Centrarchidae and order perciformes. The redspotted sunfish, redear sunfish and pumpkinseed sunfish are its closest...



Spotted tilapia

Pelmatolapia mariae

The spotted tilapia (Pelmatolapia mariae), also known as the spotted mangrove cichlid or black mangrove cichlid, is a species of fish of the cichlid family. It is native to fresh and brackish water in West and Central Africa, but has been...



Spotted trunkfish

Lactophrys bicaudalis

The spotted trunkfish (Lactophrys bicaudalis) is a member of the family Ostraciidae. It lives in reefs throughout the Caribbean, as well as the southwestern Atlantic Ocean. It gets its name from the black spots on its whitish or yellow-golden...



Spotted weever

Trachinus araneus

The spotted weever, Trachinus araneus, is a fish of the family Trachinidae, order Perciformes, and class Actinopterygii. It is up to 45 cm long, brown and yellow on the head and back, paler below with darker spots along the sides. The shape of its...



Spotted wobbegong

Orectolobus maculatus

The spotted wobbegong (*Orectolobus maculatus*) is a carpet shark in the family Orectolobidae, endemic to Australia. It is a large, robust species, typically reaching 150–180 centimetres (59–71 in) in length. Coloured green, yellow, or brown, it...



Spotty

Notolabrus celidotus

The spotty or paketi, *Notolabrus celidotus*, is a species of wrasse endemic to the waters around New Zealand, including Stewart Island and the surrounding areas. It can be found on reefs at depths from 22 to 145 m (72 to 476 ft), though most...



Spotwing flying gurnard

Dactyloptena macracantha

The spotwing flying gurnard (Dactyloptena macracantha) is an unusual looking fish because of its huge pectoral fins. The fish has dark spots and wavy lines on the fins. It has a dull head and a grey or sometimes brown body that is covered with...



Spring cavefish

Forbesichthys agassizii

The spring cavefish (Forbesichthys agassizii) is the only member of the genus Forbesichthys and is one of seven species in the family Amblyopsidae. This species is listed as state endangered in Missouri, but it is considered to be of least...



Spring pygmy sunfish

Elassoma alabamae

The spring pygmy sunfish, *Elassoma alabamae*, is a species of pygmy sunfish endemic to springs in northern Alabama. It was historically known to occur in springs in North Alabama along the Tennessee River in Limestone and Lauderdale counties. The...



Squalius alburnoides

Squalius alburnoides

Squalius alburnoides is a species of fish in the family Cyprinidae. It is found in Portugal and Spain. Its natural habitats are rivers and intermittent rivers. It may be threatened by habitat loss. This species is a highly peculiar fish in regard...



Squalius cephalus

Squalius cephalus

The common chub (Squalius cephalus), also known as the European chub or simply chub, is a species of European freshwater ray-finned fish in the carp family Cyprinidae, that frequents both slow and moderate rivers, as well as canals, lakes and...



Squalius malacitanus

Squalius malacitanus

Squalius malacitanus, commonly known as the Málaga chub, is a species of freshwater fish in the carp family Cyprinidae. It was first isolated from the Guadalmina River in Málaga, hence its name. It is considered a vulnerable species. S....



Squalus clarkae

Squalus clarkae

Squalus clarkae, also known as Genie's dogfish, is a species of shark from the Gulf of Mexico and western Atlantic. It was described in 2018 and named in honor of ichthyologist Eugenie Clark. It was previously believed to be a part of Squalus...



Squatina caillieti

Squatina caillieti

The Philippines angelshark (*Squatina caillieti*) is a species of angelshark, family Squatinidae, known only from a 33 cm (13 in) long immature female caught in the Philippines, where it is the only known representative of its family. It has a...



Squatina squatina

Squatina squatina

Squatina squatina, the angelshark or monkfish, is a species of shark in the family Squatinidae (known generally also as angel sharks), that were once widespread in the coastal waters of the northeastern Atlantic Ocean. Well-adapted for...



Star pearlfish

Carapus mourlani

The star pearlfish, Carapus mourlani, is a species of slender, ray-finned fish in the family Carapidae. It normally lives inside a starfish or a sea cucumber.



Starry flounder

Platichthys stellatus

The starry flounder (Platichthys stellatus), also known as the grindstone, emery wheel and long-nosed flounder, is a common flatfish found around the margins of the North Pacific. The distinctive features of the starry flounder include the...



Starry rockfish

Sebastes constellatus

The starry rockfish (Sebastes constellatus), also known as the spotted corsair, spotted rockfish, chinafish, and red rock cod, is a species of marine ray-finned fish belonging to the subfamily Sebastinae, the rockfishes, part of the family...



Starry smooth-hound

Mustelus asterias

Sharks portal The starry smooth-hound (Mustelus asterias) is a houndshark of the family Triakidae. It is found on the continental shelves of the northeast Atlantic, between latitudes 61 and 16° N, from the surface to a depth of 200 m (660 ft).



Starry sturgeon

Acipenser stellatus

The starry sturgeon (*Acipenser stellatus*) also known as stellate sturgeon or sevruga (Drakul, Persian: اوزون برون, and Turkish: Uzun Burun, lit. 'long nosed'), is a species of sturgeon. It is native to the Black, Azov, Caspian and Aegean sea...



Starry triggerfish

Abalistes stellaris

The starry triggerfish (Abalistes stellaris), or flat-tailed triggerfish, is a tropical, harmless, oviparous bottom dweller, characterized by some white spots along the spinal dark band. The tail is dorsoventral and looks very thin, when looked...



Starry weever

Trachinus radiatus

The starry weever, or streaked weever, Trachinus radiatus, is a fish of the family Trachinidae widespread in the eastern Atlantic from Gibraltar to the Gulf of Guinea, and probably further south; it is also known from the Mediterranean Sea. A...



Stegastes partitus

Stegastes partitus

Stegastes partitus or the bicolor damselfish is a species of bony fish in the family Pomacentridae found near the sea bed on shallow rocky and coral reefs in the Caribbean Sea, the Gulf of Mexico and off the coasts of Florida.



Stegastes planifrons

Stegastes planifrons

Stegastes planifrons (threespot damselfish) is a damselfish from the Western Atlantic. It occasionally makes its way into the aquarium trade.



Stegastes variabilis

Stegastes variabilis

Stegastes variabilis, commonly known as the Brazilian cocoa damselfish, is a species of damselfish in the family Pomacentridae, found on coral reefs in the western Atlantic Ocean and off Brazil. They are often solitary fish.



Stenogobius hawaiiensis

Stenogobius hawaiiensis

Stenogobius hawaiiensis, the Naniha goby, is a species of goby endemic to the Hawaiian Islands where it can be found in marine, brackish and fresh waters. This species is not as adept a climber as are many other Hawaiian gobies and thus is not...



Stephanolepis diaspros

Stephanolepis diaspros

Stephanolepis diaspros, commonly known as the reticulated filefish or the reticulated leatherjacket, is a species of bony fish, a ray-finned fish in the family Monacanthidae. Its natural range is the western Indian Ocean but it is also one of the...



Sterba's corydoras

Corydoras sterbai

Sterba's corydoras (*Corydoras sterbai*) is a member of the South American *Corydoras* genus of freshwater aquarium catfish and one of the most popular species of *Corydoras* due to its attractive markings. The fish is native to the Guaporé River...



Sterlet

Acipenser ruthenus

The sterlet (*Acipenser ruthenus*) is a relatively small species of sturgeon from Eurasia native to large rivers that flow into the Black Sea, Azov Sea, and Caspian Sea, as well as rivers in Siberia as far east as Yenisei. Populations migrating...



Stigmatogobius sadanundio

Stigmatogobius sadanundio

Stigmatogobius sadanundio is a species of goby native to south Asia from India to Indonesia including Sri Lanka and the Andaman Islands. It can be found in mostly fresh waters (occasionally in brackish waters) of estuaries and the tidal zones of...



Stolephorus commersonii

Stolephorus commersonii

Commerson's anchovy (Stolephorus commersonii), also known as Devis's anchovy, long-jawed anchovy, Teri anchovy, is a species of anadromous ray-finned fish in the family Engraulidae. It is known as haalmassa in Sri Lanka, where it is widely used...



Stomias boa boa

Stomias boa boa

Stomias boa boa, also called the scaly dragonfish or boa dragonfish, is a subspecies of abyssal barbeled dragonfish of the family Stomiidae. It is found at great depths worldwide in tropical to temperate oceans but is absent from the northern...



Stone loach

Barbatula barbatula

The stone loach (Barbatula barbatula) is a European species of fresh water ray-finned fish in the family Nemacheilidae. It is one of nineteen species in the genus Barbatula. Stone loaches live amongst the gravel and stones of fast flowing water...



Stone moroko

Pseudorasbora parva

The stone moroko (Pseudorasbora parva), also known as the topmouth gudgeon, is a fish belonging to the Cyprinid family, native to Asia, but introduced and now considered an invasive species in Europe and North America. The fish's size is rarely...



Stone triggerfish

Pseudobalistes naufragium

The stone triggerfish (Pseudobalistes naufragium) is the largest species of triggerfish.



Stonogobiops yasha

Stonogobiops yasha

Stonogobiops yasha, the Orange-striped shrimpgoby, is a species of goby native to the Western Pacific Ocean where it occurs at depths of from 15 to 40 metres (49 to 131 ft). It inhabits sandy areas along the outer slopes of reefs where it lives...



Stoplight parrotfish

Sparisoma viride

The stoplight parrotfish (Sparisoma viride) is a species of marine ray-finned fish, a parrotfish from the family Scaridae, inhabiting coral reefs in Florida, Caribbean Sea, Gulf of Mexico, Bermuda and as far south as Brazil. It mainly feeds on...



Straightnose pipefish

Nerophis ophidion

The straightnose pipefish (Nerophis ophidion) is a species of pipefish which lives in brackish water in the northeastern Atlantic, the Baltic, Mediterranean and Black Sea.



Streaked Spanish mackerel

Scomberomorus lineolatus

The streaked Spanish mackerel (Scomberomorus lineolatus) is found mainly in and around India, especially along the Maharashtra and Gujarat coasts. The peak season for fishing this fish is from October to December. It is also known by other names,...



Streaked spinefoot

Siganus javus

The streaked spinefoot (Siganus javus), also known as the Java rabbitfish, blue-spotted spinefoot, blue-spotted trevally, Java spinefoot or white-spotted rabbit-fish, is a species of marine ray-finned fish, a rabbitfish belonging to the family...



Streamlined spinefoot

Siganus argenteus

The streamlined spinefoot (Siganus argenteus), also known as the forktail rabbitfish, schooling rabbitfish or silver spinefoot, is a species of marine ray-finned fish, a rabbitfish belonging to the family Siganidae. It is found in the...



Streber

Zingel streber

The streber (Zingel streber), also known as the Danube streber, is a species of freshwater ray-finned fish in the family Percidae. It is found in strongly flowing waters in the Danube and Dniester drainages of Austria, Bosnia and Herzegovina,...



Striated frogfish

Antennarius striatus

The striated frogfish or hairy frogfish (Antennarius striatus) is a marine fish belonging to the family Antennariidae.



Striped bass

Morone saxatilis

The striped bass (Morone saxatilis), also called the Atlantic striped bass, striper, linesider, rock, or rockfish, is an anadromous perciform fish of the family Moronidae found primarily along the Atlantic coast of North America. It has also been...



Striped beakfish

Oplegnathus fasciatus

The striped beakfish (Oplegnathus fasciatus) is a species of marine ray-finned fish, a knifejaw from the family Oplegnathidae. It is native to the north-western Pacific Ocean, though a smattering of records exist from other localities in the...



Striped bonito

Sarda orientalis

The striped bonito (Sarda orientalis) is a species of marine perciform fish. They have been recorded at lengths of 102 centimetres (40 in), though they are commonly no longer than 55 centimetres (22 in). Distributed through the Indo-Pacific and...



Striped burrfish

Chilomycterus schoepfi

The striped burrfish or simply burrfish (Chilomycterus schoepfi) is a member of the porcupinefish family Diodontidae.



Striped killifish

Fundulus majalis

The striped killifish (*Fundulus majalis*), also called the striped mummichog, is a North American species of fundulid killifish. It lives in salt and brackish waters in shallow coastal regions from New Hampshire to Florida, and in the northern...



Striped marlin

Kajikia audax

The striped marlin (*Tetrapturus audax*, also *Kajikia audax*) is a species of marlin found globally in tropical to temperate oceans not far from the surface. It is a desirable commercial and game fish, although conservation measures are in place to...



Striped panchax

Aplocheilichthys lineatus

The striped panchax (*Aplocheilichthys lineatus*) is a species of killifish, of the genus *Aplocheilichthys*. An aquarium variant of this species with a more yellowish coloration is known as golden wonder killifish. The striped panchax inhabits fresh and...



Striped Raphael catfish

Platydoras armatulus

The striped Raphael catfish (*Platydoras armatulus*) is a catfish of the family Doradidae. It may also be called Southern striped Raphael, talking catfish, chocolate doradid, chocolate catfish or thorny catfish. It is native to the Amazon,...



Striped red mullet

Mullus surmuletus

The striped red mullet or surmullet (*Mullus surmuletus*) is a species of goatfish found in the Mediterranean Sea, eastern North Atlantic Ocean, and the Black Sea. They can be found in water as shallow as 5 metres (16 ft) or as deep as 409 metres...



Striped shiner

Luxilus chrysocephalus

The striped shiner (Luxilus chrysocephalus) is a member of the family Cyprinidae. It a silvery fish with three to four dorsolateral stripes and dark crescents on the sides. The scales and sensory pores on the snout can be outlined in black. Fins...



Stripey

Microcanthus strigatus

The stripey (Microcanthus strigatus), also known as the footballer or convict fish, is a species of ray-finned fish, a sea chub from the subfamily Microcanthinae which is part of the family Kyphosidae. It is native to the Pacific Ocean where it...



Strongylura marina

Strongylura marina

The Atlantic needlefish (Strongylura marina) is a common demersal needlefish species common in marinas and other areas with minimal currents. Its extremely long jaw and body set this fish apart from other predators. Atlantic needlefish are found...



Sturisomatichthys aureum

Sturisomatichthys aureum

Sturisomatichthys aureum, also known by several common names, including whiptail (English), Gylden Størmalle (Danish), Helikoptermal (Swedish), and Stoerwels (German), is a species of armored catfish endemic to Colombia. It grows to a length of...



Sturisomatchthys panamense

Sturisomatchthys panamense

Sturisomatchthys panamense is a species of armored catfish native to Caribbean-slope rivers of Colombia, as well as Pacific-slope rivers of Ecuador and Panama. This species grows to a length of 26 centimetres (10 in) SL.



Stylephorus

Stylephorus chordatus

Stylephorus chordatus, the tube-eye or thread-tail, is a deep-sea fish, the only fish in the genus Stylephorus and family Stylephoridae. It is found in deep subtropical and tropical oceans around the world, living at depths during the day and...



Sulfurhead aulonocara

Aulonocara maylandi

The sulfurhead aulonocara (Aulonocara maylandi) or sulfurhead peacock is a species of haplochromine cichlid which is endemic to Lake Malawi in the country of Malawi and has only been observed on West Reef (Luwala Reef) and Eccles Reef (Chimwalani...



Sulphur molly

Poecilia sulphuraria

The sulphur molly (Poecilia sulphuraria), locally known as molly del Teapa, is an endangered species of fish in the family Poeciliidae. It is endemic to Mexico, specifically to the Baños del Azufre (Grijalva River basin) near Teapa, Tabasco. The...



Summer flounder

Paralichthys dentatus

The summer flounder or fluke (Paralichthys dentatus) is a marine flatfish that is found in the Atlantic Ocean off the East Coast of the United States and Canada. It is especially abundant in waters from North Carolina to Massachusetts.



Sunburst butterflyfish

Chaetodon kleinii

The sunburst butterflyfish (Chaetodon kleinii), also known as the black-lipped butterflyfish, "blacklip butterflyfish" or Klein's butterflyfish, is a species of marine ray-finned fish, a butterflyfish belonging to the family Chaetodontidae. This...



Sundadanio axelrodi

Sundadanio axelrodi

Sundadanio axelrodi is a danionin in the family Cyprinidae, with origin from Borneo, Bangka, and Sumatra. Originally described as a rasbora, Rasbora axelrodi, this tiny species was later deemed to be more closely related to the danios but not...



Super VC-10 hap

Placidochromis milomo

The Super VC-10 hap (Placidochromis milomo) is a species of cichlid endemic to Lake Malawi along the Malawian shores of the lake. It prefers areas with rocky substrates. This species can reach a length of 18.7 centimetres (7.4 in) SL. It can also...



Surge wrasse

Thalassoma purpurum

The surge wrasse (Thalassoma purpurum), also known as the green-blocked wrasse, purple wrasse or red and green wrasse, is a species of wrasse native to the southeast Atlantic Ocean through the Indian and Pacific Oceans, where it inhabits reefs...



Suwannee bass

Micropterus notius

The Suwannee bass (Micropterus notius) is a species of freshwater fish in the sunfish family (Centrarchidae) of order Perciformes. One of the black basses, This species is native to just two river systems in Florida and Georgia, although it has...



Swamp barb

Puntius chola

The swamp barb (Puntius chola), also known as chola barb, is a species of tropical freshwater fish belonging to the subfamily Cyprininae of the family Cyprinidae. It originates in inland waters in Asia, and is found in Pakistan, India, Nepal,...



Swamp darter

Etheostoma fusiforme

The swamp darter (*Etheostoma fusiforme*) is a species of freshwater ray-finned fish, a darter from the subfamily Etheostomatinae, part of the family Percidae, which also contains the perches, ruffes and pikeperches. It is endemic to the Eastern...



Swell shark

Cephaloscyllium ventriosum

The swell shark (*Cephaloscyllium ventriosum*) is a catshark in the family Scyliorhinidae. It is found in the tropical and subtropical eastern Pacific Ocean from between central California to southern Mexico, with an additional population off the...



Swordfish

Xiphias gladius

Swordfish (*Xiphias gladius*), also known as broadbills in some countries, are large, highly migratory predatory fish characterized by a long, flat, pointed bill. They are a popular sport fish of the billfish category, though elusive. Swordfish are...



Symphurus thermophilus

Symphurus thermophilus

Symphurus thermophilus is a species of tonguefish notable for being the only flatfish known to be an obligate inhabitant of hydrothermal vents. It is known from several widely dispersed locations in the western Pacific Ocean and occurs in great...



Symphysodon aequifasciatus

Symphysodon aequifasciatus

Symphysodon aequifasciatus, the blue discus or brown discus, is a species of cichlid native to rivers of the eastern and central Amazon Basin downriver from the Purus Arch. This discus is found in black-, clear- and whitewater, but its preference...



Symphysodon discus

Symphysodon discus

Symphysodon discus, the red discus or Heckel discus, is a species of cichlid native to the Amazon Basin, where found in the lower Rio Negro, upper Uatumã, Nhamundá, Trombetas and Abacaxis. This species is essentially restricted to blackwater...



Symphysodon tarzoo

Symphysodon tarzoo

Symphysodon tarzoo, the green discus, is a species of cichlid native to rivers of the western Amazon Basin upriver from the Purus arch, although it occasionally occurs downstream. An introduced population in the Nanay River is based on stock from...



Synanceia horrida

Synanceia horrida

Synanceia horrida, the estuarine stonefish, hollow-cheek stonefish, horrid stonefish, rough stonefish or true stonefish, is a species of venomous, marine ray-finned fish, a stonefish belonging to the subfamily Synanceiinae which is classified as...



Synanceia nana

Synanceia nana

Synanceia nana, the Red Sea stonefish or dwarf scorpionfish, is a species of venomous, marine ray-finned fish, a stonefish belonging to the subfamily Synanceiinae which is classified as being within the family Scorpaenidae, the scorpionfishes and...



Synanceia verrucosa

Synanceia verrucosa

Synanceia verrucosa, the reef stonefish or simply stonefish, is a species of venomous, marine ray-finned fish, belonging to the subfamily Synanceiinae which is classified as being within the family Scorpaenidae, the scorpionfishes and their...



Synbranchus marmoratus

Synbranchus marmoratus

Synbranchus marmoratus, the marbled swamp eel, neotropical swamp eel, marmorated swamp eel, mottled swamp eel, zange, or muçum is a species of swamp eel native to Central and South America, including the island of Trinidad and Grenada.



Synchiropus splendidus

Synchiropus splendidus

Synchiropus splendidus, the mandarinfish or mandarin dragonet, is a small, brightly colored member of the dragonet family, which is popular in the saltwater aquarium trade. The mandarinfish is native to the Pacific, ranging approximately from the...



Syncrossus berdmorei

Syncrossus berdmorei

Syncrossus berdmorei is a freshwater fish in the loach family Botiidae. It is native to streams and rivers in Thailand, Burma, and nearby parts of northeastern India, where typically found over a soft bottom near boulders and submerged trees....



Syncrossus helodes

Syncrossus helodes

Syncrossus helodes, commonly known as banded loach, tiger botia or lesser katy loach, is a freshwater fish in the loach family Botiidae. It is native to rivers in Vietnam, Thailand, Laos and Cambodia. S. helodes resembles S. hymenophysa from...



Syncrossus hymenophysa

Syncrossus hymenophysa

Syncrossus hymenophysa, commonly known as the tiger loach, tiger botia or green tiger loach, is a species of freshwater fish in the loach family Botiidae. It is native to fast mountain streams and large rivers, over soft substrates and often near...



Synodontis angelicus

Synodontis angelicus

Synodontis angelicus is a species of upside-down catfish commonly named polkadot squeaker, black clown catfish, whitespotted squeaker, pearl squeaker, or angel squeaker. This species is native to the Congo Basin in the Democratic Republic of the...



Synodontis batensoda

Synodontis batensoda

Synodontis batensoda, the upside-down catfish, is a species of mochokid upside-down catfish. It is unevenly distributed in inland waters across Africa from Senegal to Ethiopia, and is also known as a squeaker or giant upside-down catfish. It was...



Synodontis decorus

Synodontis decorus

Synodontis decorus is a species of upside-down catfish. Common names include clown catfish, clown synodontis, clown syno, clown squeaker, and barredtail squeaker.



Synodontis grandiops

Synodontis grandiops

Synodontis grandiops is a species of upside-down catfish endemic to the Democratic Republic of the Congo, Burundi, and Tanzania, where it is only known from Lake Tanganyika. It was first described by Jeremy John Wright and Lawrence M. Page in...



Synodontis lucipinnis

Synodontis lucipinnis

Synodontis lucipinnis is a species of upside-down catfish endemic to Zambia, where it is only known from the Musende Rocks area (Mpulungu) of Lake Tanganyika. It was first described by Jeremy John Wright and Lawrence M. Page in 2006. The species...



Synodontis membranaceus

Synodontis membranaceus

Synodontis membranaceus, known as the moustache catfish, is a species of upside-down catfish that is native to northern Africa. It was first described by French naturalist Geoffroy Saint-Hilaire in 1809 as Pimelodus membranaceus, from specimens...



Synodontis multipunctatus

Synodontis multipunctatus

Synodontis multipunctatus, also known as the cuckoo catfish, cuckoo squeaker, or multipunk, is a small catfish from Lake Tanganyika, one of the lakes in the Great Rift Valley system in Africa. It is a brood parasite upon mouthbrooding cichlids....



Synodontis nigrita

Synodontis nigrita

Synodontis nigrita, known as the false upside down catfish, is a species of upside-down catfish that occurs widely in northern Africa. It was first described by French zoologist Achille Valenciennes in 1840. The type specimen is in the Muséum...



Synodontis nigriventris

Synodontis nigriventris

Synodontis nigriventris, the blotched upside-down catfish, is a species of upside-down catfish native to the Congo Basin of Cameroon, the Democratic Republic of the Congo and the Republic of the Congo.



Synodontis petricola

Synodontis petricola

Synodontis petricola, known as the cuckoo catfish, or the pygmy leopard catfish, is a species of upside-down catfish endemic to Burundi, Zambia, the Democratic Republic of the Congo, and Tanzania where it is only known from Lake Tanganyika. It...



Synodontis schall

Synodontis schall

Synodontis schall, the Wahrindi, is a species of upside-down catfish widespread in northern Africa. This species is in the largest genus of the family Mochokidae. This species grows to a length of 49.0 centimetres (19.3 in) TL.



Synodus intermedius

Synodus intermedius

Synodus intermedius, the common sand diver, is a species of fish in the lizardfish family, the Synodontidae, a basal ray-finned fish in the class Actinopterygii. Sand divers inhabit subtropical marine ecosystems, (37-17°N), including sandy-...



Syr Darya sturgeon

Pseudoscaphirhynchus fedtschenkoi

The Syr Darya sturgeon (Pseudoscaphirhynchus fedtschenkoi), or Syr Darya shovelnose sturgeon, is a species of fish in the family Acipenseridae. It is found in Kazakhstan, Tajikistan and Uzbekistan, where it is endemic to the Syr Darya River and,...



Tachysurus fulvidraco

Tachysurus fulvidraco

Tachysurus fulvidraco, the yellowhead catfish or Korean bullhead, is a species of bagrid catfish found in eastern Asia from Siberia to China, Korea, Vietnam, and Laos, where it can be found in lakes and river channels. It can reach a maximum...



Tadpole madtom

Noturus gyrinus

The tadpole madtom (*Noturus gyrinus*) is a species of fish in the family Ictaluridae. It is native to Canada and the United States.



Taenianotus

Taenianotus triacanthus

Taenianotus is a monotypic genus of marine ray-finned fish belonging to the family Scorpaenidae, the scorpionfishes. Its only species is Taenianotus triacanthus, the leaf scorpionfish, paperfish, paper scorpionfish, sailfin leaffish or threespine...



Taenioides cirratus

Taenioides cirratus

Taenioides cirratus, known as the Bearded worm goby, is a species of worm goby native to the Indian Ocean and the western Pacific Ocean from islands offshore of eastern Africa to New Caledonia and from Japan to Australia.



Tahoe sucker

Catostomus tahoensis

The Tahoe sucker (Catostomus tahoensis) is a freshwater cypriniform fish inhabiting the Great Basin region of the Western United States.



Tail-spot wrasse

Halichoeres melanurus

The tail-spot wrasse, Halichoeres melanurus, is a species of wrasse in the western Pacific from Japan to Samoa and Tonga and south to the Great Barrier Reef. This species is found along rocky shores or on coral reefs at depths from 1 to 15 m (3.3...



Taillight shark

Euprotomicroides zantedeschia

The taillight shark (Euprotomicroides zantedeschia) is a little-known species of shark in the family Dalatiidae and the only member of its genus. It is known from only four specimens collected from deep oceanic waters in the southern Atlantic...



Takifugu chinensis

Takifugu chinensis

Takifugu chinensis, the Chinese puffer, is a species of fish in the family Tetraodontidae (pufferfish) that is found in coastal waters of China, Japan and Korea. This critically endangered pufferfish has drastically declined because of...



Takifugu rubripes

Takifugu rubripes

Takifugu rubripes, commonly known as the Japanese puffer, Tiger puffer, or torafugu (Japanese: 虎河豚), is a pufferfish in the genus Takifugu. It is distinguished by a very small genome that has been fully sequenced because of its use as a model...



Tambaqui

Colossoma macropomum

The tambaqui (Colossoma macropomum) is a large species of freshwater fish in the family Serrasalminae. It is native to tropical South America, but kept in aquaculture and introduced elsewhere. It is also known by the names black pacu,...



Tanganyika killifish

Lamprichthys tanganicanus

The Tanganyika killifish (Lamprichthys tanganicanus) is a species of poeciliid endemic to Lake Tanganyika, where it forms large schools, mainly close to rocky shores but also pelagically off shore. This species grows to a length of 15 centimetres...



Tanganyika lates

Lates angustifrons

The Tanganyika lates (*Lates angustifrons*) is a species of lates perch endemic to Lake Tanganyika. It is a widespread predator on other fishes. This species can reach a length of 200 centimetres (79 in) SL and the greatest recorded weight is 100...



Tanichthys micagemmae

Tanichthys micagemmae

Tanichthys micagemmae is a species of freshwater ray-finned fish. It is a member of the carp family (family Cyprinidae) of order Cypriniformes. It was only discovered in 2001, in a tributary of the Bến Hải River in Quảng Bình Province, Vietnam....



Tank goby

Glossogobius giuris

Glossogobius giuris, the tank goby, is a species of goby native to fresh, marine and brackish waters from the Red Sea and East Africa through South Asia and the Indian Ocean to China, Australia and the islands of the Pacific Ocean. This species...



Taractichthys steindachneri

Taractichthys steindachneri

Taractichthys steindachneri, the sickle pomfret, is a species of marine ray-finned fish, a pomfret of the family Bramidae. It is found in the Indian and Pacific Oceans.



Tasmanian mudfish

Neochanna cleaveri

The Tasmanian mudfish, Neochanna cleaveri, is a small Australian amphidromous fish in the galaxiid family, of the order Osmeriformes.



Tasmanian whitebait

Lovettia sealii

The Tasmanian whitebait (Lovettia sealli), also known as the Australian whitebait or Derwent whitebait, is a semi-anadromous osmeriform fish of the family Galaxiidae, found only in Tasmania and southern Victoria, Australia.



Tasselled wobbegong

Eucrossorhinus dasypogon

The tasselled wobbegong (Eucrossorhinus dasypogon) is a species of carpet shark in the family Orectolobidae and the only member of its genus. It inhabits shallow coral reefs off northern Australia, New Guinea, and adjacent islands. Reaching 1.8 m...



Tautog

Tautoga onitis

The tautog (Tautoga onitis), also known as the blackfish, is a species of wrasse native to the western Atlantic Ocean from Nova Scotia to South Carolina. This species inhabits hard substrate habitats in inshore waters at depths from 1 to 75 m (5...



Tawny nurse shark

Nebrius ferrugineus

The tawny nurse shark (Nebrius ferrugineus) is a species of carpet shark in the family Ginglymostomatidae, and the only extant member of the genus Nebrius. It is found widely along coastlines in the Indo-Pacific, preferring reefs, sandy flats, and...



Teardrop butterflyfish

Chaetodon unimaculatus

The teardrop butterflyfish (*Chaetodon unimaculatus*) is a species of marine ray-finned fish, a butterflyfish belonging to the (family Chaetodontidae. It is found in the Indo-Pacific region.



Tench

Tinca tinca

The tench or doctor fish (*Tinca tinca*) is a fresh- and brackish-water fish of the order Cypriniformes found throughout Eurasia from Western Europe including the British Isles east into Asia as far as the Ob and Yenisei Rivers. It is also found in...



Tephrinectes

Tephrinectes sinensis

Tephrinectes sinensis, the Chinese brill, is a species of flatfish in the large-tooth flounder family, Paralichthyidae. It is the only member of its genus Tephrinectes. Like the rest of the large-tooth flounders, it has both eyes on the left side...



Terapon jarbua

Terapon jarbua

Terapon jarbua, the jarbua terapon, crescent grunter, crescent banded grunter, crescent perch, spiky trumpeter, thornfish or tiger perch, is a species of ray-finned fish, a grunter of the family Terapontidae. It occurs in the Indo-Pacific, it is...



Terapon puta

Terapon puta

Terapon puta, the spiny-checked grunter, three-lined grunter, small-scaled banded grunter, small-scaled terapon, squeaking perch or two-lined grunter, is a species of fish from the Indo-Pacific region, it is a member of the grunter family...



Terapon theraps

Terapon theraps

Terapon theraps, the largescaled terapon, banded grunter, banded trumpeter, flagtail grunter, flagtail trumpeter, large-scaled grunter-perch, northern grunter and spiky trumpeter, is a species of marine ray-finned fish, a grunter from the family...



Tessellated darter

Etheostoma olmstedii

The tessellated darter (*Etheostoma olmstedi*) is a species of freshwater ray-finned fish, a darter from the subfamily Etheostomatinae, part of the family Percidae, which also contains the perches, ruffes and pikeperches. It is native to Canada and...



Tete sea catfish

Ariopsis seemanni

The Tete sea catfish (*Ariopsis seemanni*) or Colombian shark catfish is a species of sea catfish in the family Ariidae, native to Pacific-draining rivers and estuaries in Central and South America.



Tetraodon duboisi

Tetraodon duboisi

Tetraodon duboisi is a species of pufferfish endemic to the Democratic Republic of the Congo where it is known only from Stanley Pool. Some sources speculate that they are critically endangered in the wild, but that theory is difficult to verify...



Tetraodon schoutedeni

Tetraodon schoutedeni

Tetraodon schoutedeni is a species of pufferfish native to Africa's Congo Basin. This species grows to a length of 9 cm (3.5 in). This species is named in honour of the Belgian zoologist Henri Schouteden.



Texas cichlid

Herichthys cyanoguttatus

The Texas cichlid (Herichthys cyanoguttatus), also known as the Rio Grande cichlid, is a freshwater fish of the cichlid family, and the only cichlid species that is native to the United States. It is found in the lower Rio Grande drainage in...



Texas shiner

Notropis amabilis

The Texas shiner (Notropis amabilis) is a species of ray-finned fish in the genus Notropis. It is found in the Colorado River to Rio Grande drainage from Texas and northeastern Mexico and the Rio Salado and Rio San Juan systems in Mexico to the...



Thalassoma bifasciatum

Thalassoma bifasciatum

Thalassoma bifasciatum, the bluehead, bluehead wrasse or blue-headed wrasse, is a species of marine ray-finned fish, a wrasse from the family Labridae. It is native to the coral reefs of the tropical waters of the western Atlantic Ocean....



Thaumatichthys axeli

Thaumatichthys axeli

Prince Axel's wonderfish (Thaumatichthys axeli) is a bottom-dwelling deep-sea anglerfish of the family Thaumatichthyidae. Thaumatichthys axeli lives at a depth of around 3,600 meters (in the abyssal zone), deeper than any other member of the...



Thaumatichthys binghami

Thaumatichthys binghami

Thaumatichthys binghami is a species of wolftap angler known from the western central Atlantic Ocean, where it occurs at a depth of 2,532 metres (8,307 ft). This species is similar to T. axeli, except that its esca bears 2–3 pairs of lateral...



Thaumatichthys pagidostomus

Thaumatichthys pagidostomus

Thaumatichthys pagidostomus is a species of wonderfish found in the Indian and Atlantic Oceans at a depth of around 1,392 metres (4,567 ft). This species grows to a length of 8.5 centimetres (3.3 in) TL. This species has relatively longer...



Thayeria boehlkei

Thayeria boehlkei

Thayeria boehlkei is a species of characin fish endemic to the Amazon river basin and Araguaia river, in Peru and Brazil respectively. The species is popular with aquarium hobbyists where it is traded under a variety of common names including...



Thermarces cerberus

Thermarces cerberus

Thermarces cerberus is a species of ray-finned fish in the family Zoarcidae. This fish, commonly known as the pink vent fish, is associated with hydrothermal vents and cold seeps at bathypelagic depths in the East Pacific.



Thick-lipped gourami

Trichogaster labiosa

The thick-lipped gourami (Trichogaster labiosa) is a species of gourami native to Southeast Asia, and is a popular aquarium fish.



Thicklip grey mullet

Chelon labrosus

The thicklip grey mullet, Chelon labrosus, is a coastal fish of the family Mugilidae. It typically is about 32 cm (13 in) long, with 75 cm (30 in) being the maximum recorded. It is named after its thick upper lip and silvery-grey appearance. It is...



Thinlip mullet

Chelon ramada

The thinlip mullet (Chelon ramada) is a species of fish in the family Mugilidae. It is found in shallow European waters and is a migratory species.



Thornback cowfish

Lactoria fornasini

The thornback cowfish (Lactoria fornasini), is a poisonous species of boxfish found throughout the tropical Indo-Pacific from East Africa to the Bass Islands (French Polynesia). It can grow to a maximum length of 23 cm (9 in). It is an uncommon...



Thornback ray

Raja clavata

The thornback ray (Raja clavata), or thornback skate, is a species of ray fish in the family Rajidae.



Thorntail stingray

Dasyatis thetidis

The thorntail stingray, black stingray, or longtail stingray (Dasyatis thetidis) is a species of stingray in the family Dasyatidae. It is found off southern Africa, Australia, and New Zealand from the intertidal zone to a depth of 440 m (1,440...



Thorny skate

Amblyraja radiata

The thorny skate (Amblyraja radiata) is a species of fish in the family Rajidae. This bottom-living skate lives in the North and south-eastern Atlantic Ocean in depths ranging from 20 to 1,000 m (66–3,281 ft) and water temperatures from –1 to 14...



Threadfin acara

Acarichthys heckelii

The threadfin acara (Acarichthys heckelii), also known as Heckel's thread-finned acara, is a South American species of cichlid fish. It is the only member of the genus Acarichthys and is native to rivers in the Amazon and Essequibo basins in...



Threadfin butterflyfish

Chaetodon auriga

The threadfin butterflyfish (Chaetodon auriga) is a species of marine ray-finned fish belonging to the family Chaetodontidae. It has a wide Indo-Pacific distribution.



Threadfin rainbowfish

Iriatherina werneri

The threadfin rainbowfish or featherfin rainbowfish (Iriatherina werneri) is a rainbowfish, the only species in the genus Iriatherina. It is characterized by long beautiful fins, and is among the most attractive of the rainbowfishes. It is native...



Threadfin shad

Dorosoma petenense

The threadfin shad (Dorosoma petenense) is a small pelagic freshwater forage fish common in lakes, large streams and reservoirs of the Southeastern United States. Like the American gizzard shad, the threadfin shad has an elongated dorsal fin, but...



Three spot cichlid

Cichlasoma trimaculatum

The three spot cichlid (Amphilophus trimaculatum), also known as the trimac or red-eyed cichlid, is a species of cichlid from Mexico and Central America, from the subfamily Cichlasomatinae. It is rarely found as an aquarium fish. Although still...



Three spot gourami

Trichopodus trichopterus

The three spot gourami (Trichopodus trichopterus), also known as the opaline gourami, blue gourami, and gold gourami, is a species of fish native to southeastern Asia, but also introduced elsewhere. This gourami gets its name from the two spots...



Three-bearded rockling

Gaidropsarus vulgaris

The three-bearded rockling (*Gaidropsarus vulgaris*) is found in European waters from the central Norwegian coast and the Faroe Islands, through the North Sea, and around the British Isles to the region around the western Mediterranean. They can...



Three-spined stickleback

Gasterosteus aculeatus

The three-spined stickleback (*Gasterosteus aculeatus*) is a fish native to most inland and coastal waters north of 30°N. It has long been a subject of scientific study for many reasons. It shows great morphological variation throughout its range....



Threespot dascyllus

Dascyllus trimaculatus

The threespot dascyllus (*Dascyllus trimaculatus*), also known as the domino damsel or simply domino, is a species of damselfish from the family Pomacentridae. It is native to the Indo-Pacific from the Red Sea and East Africa, to the Pitcairn...



Threestripe corydoras

Corydoras trilineatus

The three stripe corydoras (*Corydoras trilineatus*), leopard catfish, false julii corydoras, or three line catfish is a tropical freshwater fish belonging to the subfamily Corydoradinae of the family Callichthyidae. It originates in inland waters...



Threetooth puffer

Triodon macropterus

Triodon macropterus (common name the threetooth puffer and the black-spot keeled pufferfish) is a tetraodontiform fish, the only living species in the genus *Triodon* and family Triodontidae. Other members of the family are known from fossils...



Thunnus tonggol

Thunnus tonggol

Thunnus tonggol is a species of tuna of tropical Indo-West Pacific waters. It is commonly known as the longtail tuna or northern bluefin tuna. The usage of the latter name, mainly in Australia to distinguish it from the southern bluefin tuna,...



Thymallus thymallus

Thymallus thymallus

Thymallus thymallus, the grayling or European grayling, is a species of freshwater fish in the salmon family Salmonidae. It is the only species of the genus Thymallus (the graylings) native to Europe, where it is widespread from the United...



Thymallus yaluensis

Thymallus yaluensis

Thymallus yaluensis, also known as Yalu grayling, is a putative species of freshwater fish, a grayling in the salmon family Salmonidae. It is endemic to the upper Yalu River in Korea, on the Chinese border. Thymallus yaluensis is a small fish, the...



Thyrsites

Thyrsites atun

Thyrsites atun (Euphrasén, 1791), known as the snoek in South Africa and as the barracouta in Australasia, is a long, thin species of snake mackerel found in the seas of the Southern Hemisphere, and a popular food fish in South Africa,...



Ticto barb

Pethia ticto

The ticto barb or twospot barb (Pethia ticto) is a species of subtropical freshwater fish belonging to the family Cyprinidae. It is a native of the upper Mekong, Salween, Irrawaddy, Meklong and upper Charo Phraya basins in the countries of Nepal,...



Tidepool sculpin

Oligocottus maculosus

The tidepool sculpin (Oligocottus maculosus) is a fish species in the sculpin family Cottidae that ranges from the Bering Sea to southern California. Individuals reach up to 8 cm (3 in) in length and are common in tidepools.



Tiger barb

Puntigrus tetrazona

The tiger barb or Sumatra barb (Puntigrus tetrazona), is a species of tropical cyprinid fish. The natural geographic range reportedly extends throughout the Malay Peninsula, Sumatra and Borneo in Indonesia, with unsubstantiated sightings reported...



Tiger grouper

Mycteroperca tigris

The tiger grouper (Mycteroperca tigris) is a species of marine ray-finned fish, a grouper from the subfamily Epinephelinae which is part of the family Serranidae, which also includes the anthias and sea basses. It is found in the warmer waters of...



Tiger rockfish

Sebastes nigrocinctus

The tiger rockfish (Sebastes nigrocinctus), also called tiger seaperch, banded rockfish and black-banded rockfish, is a species of marine ray-finned fish belonging to the subfamily Sebastinae, the rockfishes, part of the family Scorpaenidae. It...



Tiger Shark

Galeocerdo cuvier

The Tiger shark (Galeocerdo cuvier) is a large species of ground shark, the largest order of sharks, with over 270 species. It is one of the most solitary, nocturnal hunters among sharks. It is also notable for having the widest food variety of...



Tiger tail seahorse

Hippocampus comes

The tiger tail seahorse (Hippocampus comes) is a species of fish in the family Syngnathidae. The species was first described by Theodore Cantor in 1850. It is found in India, Indonesia, Malaysia, the Philippines, Singapore, Thailand and Vietnam....



Tilapia busumana

Tilapia busumana

Tilapia busumana is a species of cichlid native to Lake Bosumtwi, the Bia River Basin, the Pra River Basin and the Tano River Basin in Ghana and Côte d'Ivoire. This species can reach a length of 18 centimetres (7.1 in) SL. It can also be found in...



Tilapia sparrmanii

Tilapia sparrmanii

Tilapia sparrmanii, the banded tilapia, or vlei kurper, is a widespread and adaptable cichlid fish that is found in warmer freshwater habitats of southern Africa. They prefer water with ample plant cover, and occur naturally as far north as DR...



Tinfoil barb

Barbonymus schwanenfeldii

The tinfoil barb (Barbonymus schwanenfeldii) is a tropical Southeast Asian freshwater fish of the family Cyprinidae. This species was originally described as Barbus schwanenfeldii by Pieter Bleeker in 1853, and has also been placed in the genera...



Titan triggerfish

Balistoides viridescens

The titan triggerfish, giant triggerfish or moustache triggerfish (Balistoides viridescens) is a large species of triggerfish found in lagoons and at reefs to depths of 50 m (160 ft) in most of the Indo-Pacific, though it is absent from Hawaii....



Toadstool groper

Trachypoma macracanthus

The toadstool groper (Trachypoma macracanthus), also known as the Pacific rockcod, Pacific perch, Strawberry cod and whitespotted sea bass is a sea bass of the subfamily Anthiinae of the family Serranidae, is found in the southern Pacific Ocean....



Toba betta

Betta rubra

The Toba betta (Betta rubra) is a species of gourami endemic to Sumatra, Indonesia where it is an inhabitant of peat swamps. The species grows to a length of 5 centimetres (2.0 in) TL, and it can also be found in the aquarium trade.



Tokyo bitterling

Tanakia tanago

The Tokyo bitterling (Tanakia tanago) is a temperate freshwater fish of the carp family (Cyprinidae). Taxonomically, it belongs to the subfamily Acheilognathinae. The species was first described as Rhodeus tanago by Shigeo Tanaka in 1909. It is...



Tomato clownfish

Amphiprion frenatus

The tomato clownfish (Amphiprion frenatus) is a species of marine fish in the family Pomacentridae, the clownfishes and damselfishes. It is native to the waters of the Western Pacific, from the Japan to Indonesia. Other common names include...



Tompot blenny

Parablennius gattorugine

The tompot blenny (Parablennius gattorugine) is a species of combtooth blenny from the north eastern Atlantic Ocean which is found in shallow, coastal waters off western Europe, the Mediterranean and North Africa.



Toothless blindcat

Trogloglanis pattersoni

The toothless blindcat (*Trogloglanis pattersoni*) is a species of North American freshwater catfish endemic to Texas in the United States.



Topsmelt silverside

Atherinops affinis

The topsmelt silverside (*Atherinops affinis*), also known as the topsmelt, is a species of Neotropical silverside native to the eastern Pacific Ocean.



Tor douronensis

Tor douronensis

Tor douronensis, also known as Labeobarbus douronensis, is a species of ray-finned fish of the family Cyprinidae in the genus Tor. This Asian fresh water river carp can be discovered in southern Thailand, east to Vietnam and south to Indonesia....



Tor khudree

Tor banna 5 baris murah

Tor khudree, the Deccan mahseer, Khudree mahseer, or black mahseer, is a freshwater fish of the carp family found in major rivers and reservoirs of India and Sri Lanka. Found throughout India, following large-scale introductions of...



Tor putitora

Tor putitora

Tor putitora, the Putitor mahseer, Himalayan mahseer, or golden mahseer, is an endangered species of cyprinid fish that is found in rapid streams, riverine pools, and lakes in the Himalayan region. Its native range is within the basins of the...



Tor remadevii

Tor remadevii

Tor remadevii, the orange-finned mahseer, also known as the hump-backed mahseer, is a Critically Endangered species of freshwater fish endemic to the Western Ghats of India. It is restricted to the Kaveri river basin. It can be distinguished from...



Tor tambroides

Tor tambroides

Tor tambroides, known as empurau in Malay, is a species of mahseer native to Southeast Asia.



Tor tor

Tor tor

Tor tor, commonly known as the tor mahseer or tor barb, is a species of cyprinid fish found in fast-flowing rivers and streams with rocky bottoms in India, Bangladesh, Bhutan, Nepal, and Pakistan. It is a commercially important food and game...



Torpedo scad

Megalaspis cordyla

The torpedo scad (Megalaspis cordyla), also known as the hardtail scad, finny scad, finletted mackerel scad or cordyla scad, is a species of moderately large marine fish classified in the jack and horse mackerel family, Carangidae. The torpedo...



Torquigener albomaculosus

Torquigener albomaculosus

Torquigener albomaculosus, or the white-spotted pufferfish, is the 20th discovered species of the genus Torquigener. The species was discovered in the ocean waters around the Ryukyu Islands in Japan off the south coast of Amami Ōshima Island....



Torquigener flavimaculosus

Torquigener flavimaculosus

Torquigener flavimaculosus is a pufferfish of the family Tetraodontidae, capable of very rapid body inflation when threatened. Native to the Western Indian Ocean, it is also found in the Indo Pacific from the Persian Gulf to the Seychelles....



Torrentfish

Cheimarrichthys fosteri

The torrentfish (Cheimarrichthys fosteri), or panoko (Māori), is an amphidromous freshwater fish that is endemic to New Zealand. Torrentfish are well adapted to life in shallow, fast-flowing riffles and rapids. They grow to a maximum of 20 cm...



Tortonese's stingray

Dasyatis tortonesei

Tortonese's stingray (Dasyatis tortonesei) is a species of stingray of the family Dasyatidae. It occurs in the eastern Atlantic from Morocco to Mauritania, and is also found in the Mediterranean Sea.



Tosanoides obama

Tosanoides obama

Tosanoides obama, Obama's basslet, is a coral reef fish species from the subfamily Anthiinae part of the family Serranidae, the groupers and sea basses. It was discovered in Papahānaumokuākea Marine National Monument, Hawaii. Tosanoides obama was...



Totoaba

Totoaba macdonaldi

The totoaba or totuava (Totoaba macdonaldi) is a species of marine fish, a very large member of the drum family Sciaenidae that is endemic to the Gulf of California in Mexico. It is the only species in the genus Totoaba. Formerly abundant and...



Toxotes blythii

Toxotes blythii

Toxotes blythii, the clouded archerfish or zebra archerfish, is a perciform fish of genus Toxotes. It is found in rivers and estuaries in Myanmar, ranging from the lower Irrawaddy to the Tenasserim Division, including lower Sittaung and Salween....



Toxotes chatareus

Toxotes chatareus

Toxotes chatareus, sometimes known by the common names common archerfish, seven-spot archerfish or largescale archerfish, is a species of perciform fish in the archerfish genus Toxotes. They are usually no larger than 20 centimetres (7.9 in) but...



Trachinotus goodei

Trachinotus goodei

Trachinotus goodei, the palometa, is an ocean-going game fish of the family Carangidae. Other common names include banner pompano, camade fish, cobbler, gafftopsail, great pompano, joefish, longfin pompano, old wife, sand mackerel, streamers...



Trachinotus ovatus

Trachinotus ovatus

Trachinotus ovatus, the pompano which is also known as the derbio or silverfish, is a species of ray-finned fish in the family Carangidae, the jacks. It has large, strong fins. It is common in the Mediterranean Sea, and in the Atlantic Ocean from...



Trachipterus arcticus

Trachipterus arcticus

Trachipterus arcticus is a species of ribbonfish found predominantly in the North Atlantic Ocean, with one report from the Mediterranean Sea. They are rarely encountered by humans due to their deep-sea habitat and the fact that they are of no...



Trachipterus trachipterus

Trachipterus trachipterus

Trachipterus trachipterus, is a ribbonfish of the family Trachipteridae, found in tropical and subtropical seas worldwide. Its length is up to 3 m.



Trachurus lathami

Trachurus lathami

Trachurus lathami is a species of fish in the family Carangidae and the genus Trachurus, the jack mackerels. Common names include rough scad and horse mackerel in English, as well as chinchard frappeur (French), chicharro garretón (Spanish),...



Transparent goby

Aphia minuta

Aphia minuta, the transparent goby, is a species of the goby native to the northeastern Atlantic Ocean where it can be found from Trondheim, Norway to Morocco. It is also found in the Mediterranean, Black Sea and the Sea of Azov. It is a pelagic...



Treefish

Sebastes serriceps

The treefish (Sebastes serriceps) is a species of marine ray-finned fish belonging to the subfamily Sebastinae, the rockfishes, part of the family Scorpaenidae. It is native to the eastern Pacific Ocean.



Trichogaster fasciata

Trichogaster fasciata

Trichogaster fasciata, the banded gourami or striped gourami or Colisa or Kholche, is a tropical labyrinth perch found in some Asian countries like Bangladesh, Eastern India, Northeastern India, Nepal, Upper Myanmar, China and Pakistan.It is also...



Trichomycterus rivulatus

Trichomycterus rivulatus

Trichomycterus rivulatus is a species of pencil catfish that is native to high-altitude Andean streams and lakes (including Junin, Poopó and Titicaca) in southern Peru, western Bolivia and northern Chile. It is the largest species in the genus...



Trigonostigma espei

Trigonostigma espei

The espei rasbora, or lambchop rasbora (Trigonostigma espei) is a species of ray-finned fish in the genus Trigonostigma. It is named after the dark band that appears like a lamb chop along its body. The species is predominantly found in Thailand...



Trigonostigma somphongsi

Trigonostigma somphongsi

Trigonostigma somphongsi is a species of ray-finned fish in the genus Trigonostigma. It is endemic to Thailand. It is threatened by habitat loss.



Trimmatom nanus

Trimmatom nanus

Trimmatom nanus, the midget dwarfgoby, is a species of marine goby native to the Indian Ocean and the western Pacific Ocean. It can mainly be found on oceanic drop-offs at depths of from 20 to 30 metres (66 to 98 ft) though it can occasionally be...



Tripletail wrasse

Cheilinus trilobatus

The tripletail wrasse (Cheilinus trilobatus) is a species of marine ray-finned fish from the family Labridae, the wrasses.



Triplewart seadevil

Cryptopsaras couesii

The triplewart seadevil (Cryptopsaras couesii) is a sea devil of the family Ceratiidae and the order Lophiiformes. This species is the only member of its genus. Noted for its extreme sexual dimorphism, the triplewart seadevil's length ranges...



Trisopterus esmarkii

Trisopterus esmarkii

Trisopterus esmarkii, the Norway pout, is a species of fish in the cod family. It is found in the Barents Sea, North Sea, Baltic Sea, off the coasts of Norway, Iceland, the British Isles and elsewhere in the northeast Atlantic Ocean. It prefers...



Trisopterus luscus

Trisopterus luscus

Trisopterus luscus (.mw-parser-output.IPA-label-small{font-size:85%}.mw-parser-output.references.IPA-label-small,.mw-parser-output.infobox.IPA-label-small,.mw-parser-output.navbox.IPA-label-small{font-size:100%}Latin pronunciation: ; bib, pout...



Tristramella simonis

Tristramella simonis

Tristramella simonis, the short jaw tristramella, is a vulnerable species of cichlid fish from the Jordan River system, including Lake Tiberias (Kinneret), in Israel and Syria, with introduced populations in the Nahr al-Kabir and Orontes basins...



Tropheus duboisi

Tropheus duboisi

Tropheus duboisi, the white spotted cichlid, is a species of cichlid endemic to Lake Tanganyika. It can reach a length of 12 cm (4.7 in).



Tropheus moorii

Tropheus moorii

Tropheus moorii (blunthead cichlid) is a species of cichlid endemic to Lake Tanganyika in Africa. Over 40 different color morphs of this species are dispersed throughout the lake, ranging from dark green to flame red and yellow. They mostly feed...



Tropical gar

Atractosteus tropicus

The tropical gar (Atractosteus tropicus) is a species of fish from Central America, where it is found in the Pacific and Atlantic drainages from southern Mexico to Costa Rica. In Central America it is known as gaspar and in Mexico it is known as...



Trout barb

Raiamas bola

The trout barb, or Indian trout (Raiamas bola) is a carp of the family Cyprinidae, which occurs in freshwaters around the Bay of Bengal.



Trout cod

Maccullochella macquariensis

The trout cod (Maccullochella macquariensis) or bluenose cod, is a large predatory freshwater fish of the genus Maccullochella and the family Percichthyidae, closely related to the Murray cod. It was originally widespread in the south-east corner...



Trumpeter whiting

Sillago maculata

The trumpeter whiting (Sillago maculata), also known as the winter whiting or diver whiting, is a common species of coastal marine fish of the smelt-whiting family, Sillaginidae. The trumpeter whiting is endemic to Australia, inhabiting the...



Trypauchen vagina

Trypauchen vagina

Trypauchen vagina, commonly known as the burrowing goby, is a species of eel goby found in the Indo-Pacific region. It has an elongated body about 20 to 22 cm (7.9 to 8.7 in) in length. It is reddish-pink in color and possesses distinctive...



Tub gurnard

Chelidonichthys lucerna

The tub gurnard (Chelidonichthys lucerna), also known as the sapphirine gurnard, tube-fish, tubfish or yellow gurnard, is a species of marine ray-finned fish belonging to the family Triglidae, the gurnards and sea robins. It is found in the...



Tui chub

Siphateles bicolor

The Tui chub (Siphateles bicolor) is a cyprinid fish native to western North America. Widespread in many areas, it is a highly adaptable fish that has historically been a staple food source for native peoples.



Tule perch

Hysterocarpus traskii

The tule perch (Hysterocarpus traskii) is a surfperch (Embiotocidae) native to the rivers and estuaries of central California. It is the sole member of its genus, and the only freshwater surfperch. The tule perch is small, at most 15 centimetres...



Turbot

Scophthalmus maximus

The turbot (Scophthalmus maximus) is a relatively large species of flatfish in the family Scophthalmidae. It is a demersal fish native to marine or brackish waters of the Northeast Atlantic, Baltic Sea and the Mediterranean Sea. It is an...



Turkey moray

Gymnothorax meleagris

The turkey moray (Gymnothorax meleagris), also known as the guineafowl moray or as the white-mouth/whitemouth moray is a species of marine fish in the family Muraenidae.



Twait shad

Alosa fallax

The twait shad or twaite shad (Alosa fallax) is a species of fish in the family Clupeidae. It is found in the eastern Atlantic Ocean and the Mediterranean Sea and is an anadromous fish which lives in the sea but migrates into fresh water to...



Twin-spot goby

Signigobius biocellatus

The Twin-spot Goby, or Crab-eyed Goby, (Signigobius biocellatus) is a species of goby native to the Western Pacific Ocean where they can be found in areas of sand, silt in lagoons, or coastal bays with nearby cover such as rubble, coral, or leaf...



Two-spined blackfish

Gadopsis bispinosus

The two-spined blackfish (*Gadopsis bispinosus*) is a species of temperate perch endemic to Australia. It is found in the cool, clear, strong-flowing, cobble bottomed, sub-alpine rivers and streams (ranging from small to large) in the southeast...



Two-spotted goby

Gobiusculus flavescens

The two-spotted goby (*Pomatoschistus flavescens*) is a species of goby native to marine and brackish waters of the eastern Atlantic Ocean where it can be found from the Faeroes and Norway to the northwestern coast of Spain. It has also been...



Twospined angelfish

Centropyge bispinosa

The twospined angelfish (*Centropyge bispinosa*), also known as the dusky angelfish, or coral beauty, is a species of marine ray-finned fish, a marine angelfish belonging to the family Pomacanthidae. They are found in the Indo-Pacific.



Tylosurus choram

Tylosurus choram

Tylosurus choram, the Red Sea houndfish, is a species of needlefish from the family Belonidae. A marine fish bluish in color with a long slender body, and a pointed long toothed beak, found in most temperate, warm seas, and sometimes rivers, it...



Typhleotris madagascariensis

Typhleotris madagascariensis

Typhleotris madagascariensis is a species of fish in the family Milyeringidae that is endemic to Madagascar, where it is only known from underground waters in the southwestern portion of the island. This cavefish is blind and lacks pigmentation,...



Typhlichthys subterraneus

Typhlichthys subterraneus

Typhlichthys subterraneus, the southern cavefish, is a species of cavefish in the family Amblyopsidae endemic to karst regions of the eastern United States.



Tyrannochromis macrostoma

Tyrannochromis macrostoma

Tyrannochromis macrostoma, or big-mouth hap, is a species of cichlid endemic to Lake Malawi where it prefers rocky shallows. This species can reach a length of 30 centimetres (12 in) TL. It can also be found in the aquarium trade.



Uaru amphiacanthoides

Uaru amphiacanthoides

Uaru amphiacanthoides, the uaru or triangle cichlid, is a species of cichlid native to South America where it occurs in clear streams of the Amazon Basin. This species can reach a length of 25 centimetres (9.8 in) SL. It is also of importance as...



Ukrainian stickleback

Pungitius platygaster

The Ukrainian stickleback (Pungitius platygaster) also known as the Caspian ninespine stickleback, southern ninespine stickleback, and Aral ninespine stickleback, is a species of fish in the family Gasterosteidae. It is found in Afghanistan,...



Umbrina cirrosa

Umbrina cirrosa

Umbrina cirrosa, the shi drum, is a species of marine fish from the warmer waters of the eastern Atlantic Ocean, Mediterranean Sea and the Black Sea. It is a commercially important species which is trawled for and farmed in aquaculture, as well...



Undulate ray

Raja undulata

The undulate ray (Raja undulata) is a species of ray and cartilaginous fish found in the Mediterranean and East Atlantic from southern Ireland and England to the Gulf of Guinea. It is found in areas with mud or sand, and may occur as deep as 200...



Undulated moray

Gymnothorax undulatus

The undulated moray (Gymnothorax undulatus) is a moray eel of the family Muraenidae, found in the Indo-Pacific and east-central Pacific Ocean at depths down to 30 m. Their length is up to 1.5 m.



Unicorn leatherjacket

Aluterus monoceros

The unicorn leatherjacket (*Aluterus monoceros*) is a filefish of the family Monacanthidae, found around the world in subtropical oceans between latitudes 43° N and 35° S, at depths down to 50 m. Its length is up to 76 cm.



Upeneichthys lineatus

Upeneichthys lineatus

Upeneichthys lineatus, also known as the blue-striped mullet, blue-lined goatfish, blue-striped goatfish, blue-spotted goatfish and blue striped red mullet, is a species of marine ray-finned fish, a goatfish from the family Mullidae. It is native...



Upeneus moluccensis

Upeneus moluccensis

Upeneus moluccensis, the goldband goatfish, golden-banded goatfish or Moluccan goatfish, is a species of Indo-Pacific goatfish from the red mullet and goatfish family, the Mullidae. It is widespread in the warmer waters of the Indian and Pacific...



Utah chub

Gila atraria

The Utah chub (Gila atraria) is a cyprinid fish native to western North America, where it is abundant in the upper Snake River and throughout the Lake Bonneville basin. This chub generally follows the cyprinid body plan. Its dorsal fin lies...



Utah sucker

Catostomus ardens

The Utah sucker (Catostomus ardens) is a species of freshwater fish in the family Catostomidae found in the upper Snake River and the Lake Bonneville areas of western North America where it lives in a wide range of habitats. It is a large sucker...



Vadigo

Campogramma glaycos

The vadigo, Campogramma glaycos (also known as the big-toothed pompano, zippered pompano, lexa and lexola), is a species of medium-sized coastal marine fish in the jack family, Carangidae. The species is distributed throughout the eastern...



Vagabond butterflyfish

Chaetodon vagabundus

The vagabond butterflyfish (Chaetodon vagabundus), also known as the crisscross butterflyfish, is a species of marine ray-finned fish, a butterflyfish belonging to the family Chaetodontidae. It is found in the Indo-Pacific region.



Valencia letourneuxi

Valencia letourneuxi

Valencia letourneuxi, the Corfu toothcarp, is a species of fish in the family Valenciidae. It is found in Albania and Greece. Its natural habitats are rivers, freshwater marshes, freshwater spring, and coastal saline lagoons. It is threatened by...



Valenciennea puellaris

Valenciennea puellaris

Valenciennea puellaris, the Orange-spotted sleeper-goby, Orange-dashed goby, or Maiden goby, Diamond Watchman goby, is a species of goby native to the Indian Ocean and the western Pacific Ocean. It inhabits lagoons and outer reefs where it occurs...



Valenciennea strigata

Valenciennea strigata

Valenciennea strigata is a species of fish in the family Gobiidae, the gobies. Its common names include the blueband goby, golden-head sleeper goby, and pennant glider. It is native to the Indian Ocean and the western Pacific Ocean where it can...



Valentin's sharpnose puffer

Canthigaster valentini

Valentin's sharpnose puffer (Canthigaster valentini), also known as the saddled puffer or black saddled toby, is a demersal marine fish belonging to the family Tetraodontidae. The saddled puffer is a small sized fish which grows up to 11 cm. It...



Variabilichromis moorii

Variabilichromis moorii

Variabilichromis moorii has no common name and is a species of freshwater cichlid endemic to Lake Tanganyika in eastern Africa. It is a small ovate bodied fish named for an early collector of fish from the lake, John Edmund Sharrock Moore...



Variable platyfish

Xiphophorus variatus

The variable platyfish (Xiphophorus variatus), also known as variatus platy or variegated platy, is a species of freshwater fish in family Poeciliidae of order Cyprinodontiformes. A livebearer, it is native to southern Tamaulipas and northern...



Variegated lizardfish

Synodus variegatus

The variegated lizardfish (Synodus variegatus) is a lizardfish of the family Synodontidae found in the western Pacific and Indian Oceans, at depths from 4 to 90 m. It can reach a maximum length of 40 cm.



Velvet belly lanternshark

Etmopterus spinax

The velvet belly lanternshark (or simply velvet belly) (Etmopterus spinax) is a species of dogfish shark in the family Etmopteridae. One of the most common deepwater sharks in the northeastern Atlantic Ocean, the velvet belly is found from...



Velvet dogfish

Zameus squamulosus

Sharks portalThe velvet dogfish (Zameus squamulosus), the only member of the genus Zameus, is a small sleeper shark of the family Somniosidae, found around the world between latitudes 64°N and 48°S, from the surface to 2,000 m.



Velvet whalefish

Barbourisia rufa

The velvet whalefish (Barbourisia rufa) is a deep-sea whalefish, the sole known member of its family Barbourisiidae. It is found throughout the tropical and temperate parts of the world's oceans, mainly in the Pacific near Japan and New Zealand,...



Vermilion snapper

Rhomboplites aurorubens

The vermilion snapper (Rhomboplites aurorubens), the clubhead snapper, night snapper, Brian's red fish or beeliner is a species of marine ray-finned fish, a snapper belonging to the family Lutjanidae. It is native to the western Atlantic Ocean.



Vieja melanurus

Vieja melanurus

Vieja melanurus, the quetzal cichlid, redhead cichlid or firehead cichlid, is a species of cichlid that is native to the Lake Petén system, the Grijalva–Usumacinta River basin and other Atlantic river drainages in southern Mexico, Belize and...



Vimba vimba

Vimba vimba

Vimba vimba, called also the vimba bream, vimba, zante, or zarte, is a European fish species in the family Cyprinidae. It largely lives in the sea but makes an annual migration up-river each year to breed.



Violet goby

Gobioides broussonnetii

The violet goby (*Gobioides broussonnetii*) is a species of goby native to marine, fresh and brackish waters near the Atlantic coast of North and South America from South Carolina in the United States of America, to northern Brazil. It prefers...



Viper dogfish

Trigonognathus kabeyai

The viper dogfish or viper shark (Trigonognathus kabeyai) is a rare species of dogfish shark in the family Etmopteridae, and the only extant member of its genus. It has been found in the Pacific Ocean off southern Japan, the Bonin Islands,...



Viper moray

Enchelynassa formosa

The viper moray (Enchelynassa formosa or Enchelynassa canina) is a species of saltwater eel, the only member of the genus Enchelynassa of the family Muraenidae (Moray eels). It is found in the Indo-Pacific oceans. It can grow up to 250 cm.



Viviparous eelpout

Zoarces viviparus

The viviparous eelpout (Zoarces viviparus), also known as the, viviparous blenny and European eelpout is species of marine ray-finned fish belonging to the family Zoarcidae, the eelpouts. It is notable for being ovoviviparous and gives birth to...



Volga pikeperch

Sander volgensis

The Volga pikeperch, or Volga zander (Sander volgensis), is a species of fish in the perch family Percidae. It is found in Austria, Azerbaijan, Bosnia and Herzegovina, Bulgaria, Croatia, Germany, Hungary, Moldova, Romania, Russia, Serbia,...



Vundu

Heterobranchus longifilis

The vundu (*Heterobranchus longifilis*) is a species of large airbreathing catfish found widely in rivers and other freshwater habitats of sub-Saharan Africa, as well as the Nile. It is also called the sampa, cur, lenda, or certa.



Wahoo

Acanthocybium solandri

Wahoo (*Acanthocybium solandri*) is a scombrid fish found worldwide in tropical and subtropical seas. It is best known to sports fishermen, as its speed and high-quality flesh makes it a prized and valued game fish. In Hawaii, the wahoo is known as...



Walking catfish

Clarias batrachus

The walking catfish (*Clarias batrachus*) is a species of freshwater airbreathing catfish native to Southeast Asia. It is named for its ability to "walk" and wiggle across dry land, to find food or suitable environments. While it does not truly...



Wallago attu

Wallago attu

Wallago attu is a freshwater catfish of the family Siluridae, native to South and Southeast Asia. W. attu is found in large rivers and lakes in two geographically disconnected regions (disjunct distribution), with one population living over much...



Wallagonia leerii

Wallagonia leerii

Wallagonia leerii, also known as the Great Tapah, helicopter catfish, and formerly the striped wallago catfish is a species of catfish native to Southeast Asia. Its habitat ranges from the river drainages of Thailand through the Malayan peninsula...



Wallagonia micropogon

Wallagonia micropogon

Wallagonia micropogon is a species of catfish in the genus Wallagonia. This species was recently discovered and is found in the Mekong River drainage area between southern Vietnam and northern Laos as well as Chao Phraya River in Thailand. It is...



Walleye

Sander vitreus

The walleye (Sander vitreus, synonym Stizostedion vitreum), also called the yellow pike or yellow pikeperch or yellow pickerel, is a freshwater perciform fish native to most of Canada and to the Northern United States. It is a North American...



Warmouth

Lepomis gulosus

The warmouth (Lepomis gulosus) is a freshwater fish of the sunfish family (Centrarchidae) that is found throughout the eastern United States. Other local names include molly, redeye, goggle-eye, red-eyed bream, and strawberry perch.



Warpaint shiner

Luxilus coccogenis

The warpaint shiner (Luxilus coccogenis) is a species of freshwater fish found in North America. It is common in the upper Tennessee River basin as well as in the Savannah River, the Santee River, and the New River in North Carolina. Adults have...



Warty frogfish

Antennarius maculatus

The warty frogfish or clown frogfish (*Antennarius maculatus*) is a marine fish belonging to the family Antennariidae.



Warty oreo

Allocyttus verrucosus

The warty oreo (*Allocyttus verrucosus*) is an oreo found in all southern oceans at depths of between 300 and 1,600 m. Its length is up to 42 cm. The warty oreo resembles the spiky oreo. However, there are differences: the body, which has a shape...



Watercress darter

Etheostoma nuchale

The watercress darter (Etheostoma nuchale) is a species of freshwater ray-finned fish, a darter from the subfamily Etheostomatinae, part of the family Percidae, which also contains the perches, ruffes and pikeperches. It is endemic to the eastern...



Waterfall climbing cave fish

Cryptotora thamicola

The waterfall climbing cave fish (Cryptotora thamicola), also known as the cave angel fish, is a species of troglotic hillstream loach endemic to Thailand. It reaches a length of 2.8 centimetres (1.1 in) SL. This fish is known for its fins,...



Wedge sole

Dicologlossa cuneata

The wedge sole (Dicologlossa cuneata), is a flatfish of the family Soleidae. It is a bottom dwelling predatory fish inhabiting both sandy and muddy soils at depths between 10 and 450 m (33 and 1,476 ft) in the East Atlantic and Mediterranean Sea....



Wels catfish

Silurus glanis

The wels catfish (or ; Silurus glanis), also called sheatfish or just wels, is a large species of catfish native to wide areas of central, southern, and eastern Europe, in the basins of the Baltic, Black and Caspian Seas. It has been introduced...



West African Spanish mackerel

Scomberomorus tritor

The West African Spanish mackerel (Scomberomorus tritor) is a species of fish in the family Scombridae. Specimens have been recorded at up to 100 cm in length, and weighing up to 6 kg. Coloration is bluish-green on the back fading to silvery on...



Western blacknose dace

Rhinichthys obtusus

Western blacknose dace (Rhinichthys obtusus) is a common species of ray-finned fish belonging to the family Cyprinidae (minnow or carp) and the genus Rhinichthys. Western blacknose dace have tan to dark brown backs, lighter sides, and cream...



Western blue devil

Paraplesiops sinclairi

The western blue devil (Paraplesiops sinclairi) is a species of fish in the longfin family Plesiopidae endemic to Western Australia. The fish is found in rocky reef habitats in the coastal, inshore waters of southwestern Western Australia, from...



Western carp gudgeon

Hypseleotris klunzingeri

The western carp gudgeon (Hypseleotris klunzingeri) is one of several carp gudgeon species. Carp gudgeons are very small perciform fish (similar in size, shape and colour) found in the Australian Murray-Darling River system, mainly in lowland...



Western pygmy perch

Nannoperca vittata

The western pygmy perch (Nannoperca vittata) is a species of temperate perch endemic to southwestern Australia.



Western tubenose goby

Proterorhinus semilunaris

The western tubenose goby (Proterorhinus semilunaris) is a species of goby native to fresh waters of the Black Sea and Aegean Sea basins. It has recently spread as an invasive species to Central and Western Europe and to North America. Previously...



Westslope cutthroat trout

Oncorhynchus clarkii lewisi

The westslope cutthroat trout (Oncorhynchus clarkii lewisi), also known as the black-spotted trout, common cutthroat trout and red-throated trout is a subspecies of the cutthroat trout (Oncorhynchus clarkii) and is a freshwater fish in the salmon...



Whale Shark

Rhincodon typus

The Whale shark (Rhincodon typus) is a slow-moving, filter-feeding shark and the largest known extant fish species. It holds many records for size in the animal kingdom, most notably being by far the most massive living non-cetacean animal.



Whalesucker

Remora australis

The whalesucker (Remora australis) is a species of remora in the family Echeneidae, so named because it attaches itself exclusively to cetaceans. It is found worldwide in tropical and warm waters; in the Gulf of Mexico and western Atlantic Ocean,...



Whiptail catfish

Dasyloricaria filamentosa

Whiptail catfish (Dasyloricaria filamentosa) is a species of armored catfish endemic to Colombia where it is found in the Magdalena River basin and is suspected to also occur in the Catatumbo River. This species grows to a length of 26...



Whiskery shark

Furgaleus macki

The whiskery shark (Furgaleus macki) is a species of houndshark in the family Triakidae, and the only member of its genus. This common shark inhabits the Australian continental shelf from Western Australia to the Bass Strait, to a depth of 220 m...



White Amur bream

Parabramis pekinensis

The white Amur bream (Parabramis pekinensis) is a species of cyprinid freshwater fish, of the monotypic genus Parabramis. It is native to eastern Asia, where found from the Amur River basin in Russia south to Ningbo and Shanghai in China. It is...



White bass

Morone chrysops

The white bass, silver bass, or sand bass (Morone chrysops) is a freshwater fish of the temperate bass family Moronidae. commonly around 12-15 inches long. The species' main color is silver-white to pale green. Its back is dark, with white sides...



White bullhead

Ameiurus catus

The white bullhead (Ameiurus catus), also known as the white catfish, is a member of the family Ictaluridae of the order Siluriformes.



White Cloud Mountain minnow

Tanichthys albonubes

The White Cloud Mountain minnow (Tanichthys albonubes) is a hardy species of freshwater fish and coldwater fish often kept in an aquarium. The species is a member of the carp family (family Cyprinidae) of the order Cypriniformes, native to China....



White crappie

Pomoxis annularis

The white crappie (*Pomoxis annularis*) is a freshwater fish found in North America, one of the two species of crappies. Alternate common names for the species include goldring, silver perch, white perch and sac-a-lait. USS Goldring is named for...



White croaker

Genyonemus lineatus

White croaker ('*Genyonemus lineatus*') is a species of croaker occurring in the Eastern Pacific. White croakers have been taken from Magdalena Bay, Baja California, to Vancouver Island, British Columbia, but are not abundant north of San...



White grouper

Epinephelus aeneus

The white grouper (*Epinephelus aeneus*) is a species of marine ray-finned fish, a grouper from the subfamily Epinephelinae which is part of the family Serranidae, which also includes the anthias and sea basses. The white grouper is found in the...



White hake

Urophycis tenuis

The white hake or mud hake (*Urophycis tenuis*) is a phycid hake. It is found in the deeper waters in the northwest Atlantic Ocean.



White marlin

Kajikia albida

The white marlin (*Kajikia albida*), also known as Atlantic white marlin, marlin, skilligalee, is a species of billfish that lives in the epipelagic zone of the tropical and subtropical Atlantic Ocean. They are found between the latitudes of 45° N...



White mullet

Mugil curema

The white mullet or silver mullet (Mugil curema) is a tropical and subtropical marine fish of the family Mugilidae. It is commonly about 30 cm (12 in) long. Note that silver mullet is also a common name used for fantail mullet (Mugil trichodon).



White perch

Morone americana

The white perch (Morone americana) is not a true perch but is a fish of the temperate bass family, Moronidae, notable as a food and game fish in eastern North America. In some places it is referred to as "Silver Bass". The common name "white...



White ribbon eel

Pseudechidna brummeri

The white ribbon eel, Pseudechidna brummeri, is a species of saltwater eels, the only member of the genus Pseudechidna of the Muraenidae (Moray eel) family. It is found in the Indo-Pacific oceans from the western Indian Ocean to Samoa, and north...



White River springfish

Crenichthys baileyi

The White River springfish (Crenichthys baileyi) is a species of fish in the family Goodeidae, the splitfins. It is a rare species of the Great Basin of western United States, where it is endemic to isolated warm springs in the White River...



White Sands pupfish

Cyprinodon tularosa

The White Sands pupfish (Cyprinodon tularosa) is a species of pupfish in the family Cyprinodontidae. It is endemic to the Tularosa Basin in southern New Mexico, in the Western United States. They are only found in two springs and two small...



White Sea herring

Clupea pallasii marisalbi

The White Sea herring, *Clupea pallasii marisalbi*, is a subspecies of the Pacific herring, *Clupea pallasii*, in the genus *Clupea* of the family *Clupeidae*.



White seabass

Atractoscion nobilis

White seabass or white weakfish, *Atractoscion nobilis*, is a species of croaker occurring from Magdalena Bay, Baja California, to Juneau, Alaska. They usually travel in schools over deep rocky bottoms (0–122 m) and in and out of kelp beds.



White sturgeon

Acipenser transmontanus

White sturgeon (*Acipenser transmontanus*) is a species of sturgeon in the family *Acipenseridae* of the order *Acipenseriformes*. They are an anadromous (migratory) fish species ranging in the Eastern Pacific; from the Gulf of Alaska to Monterey,...



White sucker

Catostomus commersonii

The white sucker (*Catostomus commersonii*) is a species of freshwater cypriniform fish inhabiting the upper Midwest and Northeast in North America, but it is also found as far south as Georgia and as far west as New Mexico. The fish is commonly...



White trevally

Pseudocaranx dentex

The white trevally (*Pseudocaranx dentex*), also known as striped jack,) is a jack of the family Carangidae widespread in tropical and warm temperate areas between 40°N and 47°S, in the Atlantic, Mediterranean, Indian, and Pacific Oceans. It has a...



White-edge freshwater whipray

Fluvitrygon signifer

The white-edge freshwater whipray (Fluvitrygon signifer) is an extremely rare species of stingray in the family Dasyatidae, native to four river systems in Southeast Asia. Measuring up to 60 cm (24 in) across, this ray has an oval pectoral fin...



White-eye bream

Ballerus sapa

White-eye bream (Ballerus sapa) is a fish species of the family Cyprinidae. It is widespread in many large rivers in Europe and Asia in drainages of the Black Sea, Caspian Sea, and Aral Sea. It was introduced to several rivers in northern Russia,...



White-spotted puffer

Arothron hispidus

The white-spotted puffer fish (Arothron hispidus) is a medium to large-sized puffer fish, it can reach 50 cm length. It is light grey in color, or greyish or yellowish, and clearly covered with more or less regular white points, that become...



Whitecheek shark

Carcharhinus dussumieri

Sharks portalThe whitecheek shark or widemouth blackspot shark (Carcharhinus dussumieri) is a requiem shark of the family Carcharhinidae, found in the Indo-West Pacific Ocean between latitudes 34°N and 25°S. It can reach a length of 1 m. It...



Whitefin trevally

Carangoides equula

The whitefin trevally (*Carangoides equula*), also known as the horse trevally, is a species of deep water offshore fish in the jack family Carangidae. The species inhabits the tropical to temperate waters of the Indo-Pacific and central Pacific....



Whitemargin stargazer

Uranoscopus sulphureus

Whitemargin stargazer (*Uranoscopus sulphureus*) is a fish of family Uranoscopidae, widespread in the Indopacific: Red Sea, Indonesia, Fiji, Samoa, and Tonga. Marine reef-associated fish, up to 45.0 cm maximal length.



Whitemargin unicornfish

Naso annulatus

The whitemargin unicornfish (*Naso annulatus*), also known as the ringtailed unicornfish or short-horned unicorn-fish, is a tropical fish found throughout the Indo-Pacific. It can reach a length of 100 cm, making it one of the largest members of...



Whitenose shark

Nasolamia velox

The whitenose shark (*Nasolamia velox*) is a species of shark of the family Carcharhinidae. The only member of the genus *Nasolamia*, it is found in the tropical waters of the eastern Pacific Ocean between latitudes 31° N and 18° S, between depths of...



Whitesaddled catshark

Scyliorhinus hesperius

The whitesaddled catshark (Scyliorhinus hesperius) is a species of catshark, and part of the family Scyliorhinidae. It is found on the upper continental slope of the western central Atlantic Ocean, off the coasts of Honduras, Panama and Colombia,...



Whitespot ghost shark

Hydrolagus albus

The whitespot ghost shark (Hydrolagus albus) is a chimaera species in the family Chimaeridae, which lives in parts of the Galápagos Islands in the southeast Pacific Ocean. It lives in waters with steep slopes and boulders and grows to a total...



Whitespotted bamboo shark

Chiloscyllium plagiosum

The whitespotted bamboo shark (Chiloscyllium plagiosum) is a species of carpet shark with an adult size that approaches one metre in length. This small, mostly nocturnal species is harmless to humans. The whitespotted bamboo shark is occasionally...



Whitespotted bullhead shark

Heterodontus ramalheira

Sharks portalThe whitespotted bullhead shark, Heterodontus ramalheira, is a bullhead shark of the family Heterodontidae found in the western and northern Indian Ocean between latitudes 22°N to 26°S, at depths between 40 and 305 m. It can grow up...



Whitespotted conger

Conger myriaster

The whitespotted conger (Conger myriaster) (Japanese: 真穴子; ma-anago) is a marine conger eel, widespread in the Northwest Pacific near the coasts of Japan, Korean Peninsula, and the East China Sea. C. myriaster inhabits shallow sea bottom sand and...



Whitespotted grouper

Epinephelus coeruleopunctatus

The whitespotted grouper (Epinephelus coeruleopunctatus), also known as the rankin cod, ocellated rockcod, small-spotted cod, white-spotted reef-cod or whitespotted rockcod, is a species of marine ray-finned fish, a grouper from the subfamily...



Whitespotted whipray

Maculabatis gerrardi

The whitespotted whipray or sharpnose stingray (Maculabatis gerrardi) is a species of stingray in the family Dasyatidae. It is found in coastal regions including estuaries, in the Indo-Pacific, and has also been recorded in the Ganges River. It...



Whitetail dascyllus

Dascyllus aruanus

Dascyllus aruanus, known commonly as the whitetail dascyllus or humbug damselfish among other vernacular names, is a species of marine fish in the family Pomacentridae.



Whitetip Reef Shark

Triaenodon obesus

The Whitetip reef shark (Triaenodon obesus) is a species of small requiem shark, meaning it doesn't have to keep swimming to breathe. It is one of the most common sharks found on Indo-Pacific coral reefs, typically inhabiting the bottom or near...



Whitetip weasel shark

Paragaleus leucomatus

Sharks portalThe whitetip weasel shark (*Paragaleus leucomatus*) is a weasel shark of the family Hemigaleidae. Only one specimen, caught off Kosi Bay, South Africa, has been caught. That specimen was 96 cm long. In 2020, a fuzzy image believed...



Widemouth blindcat

Satan eurystomus

The widemouth blindcat (Satan eurystomus) is a species of North American freshwater catfish endemic to Texas in the United States. It is the only species in the genus Satan.



Widow rockfish

Sebastes entomelas

The widow rockfish (Sebastes entomelas), or brown bomber, is a species of marine ray-finned fish belonging to the subfamily Sebastinae, the rockfishes, part of the family Scorpaenidae. It is found in the northeastern Pacific Ocean.



William's mbuna

Maylandia greshakei

William's mbuna (Maylandia greshakei), also known as ice blue zebra mbuna or ice blue zebra or the Pseudotropheus ice blue among the aquarium enthusiasts, is a species of cichlid fish endemic to Lake Malawi where it is only found at Makokola in...



Windowpane flounder

Scophthalmus aquosus

The windowpane flounder (Scophthalmus aquosus) is a species of fish in the family Scophthalmidae. It is widespread at a depth of 5–73 meters in the western Atlantic from the Gulf of St. Lawrence in Canada to Florida in the United States. An...



Winghead shark

Eusphyra blochii

The winghead shark (*Eusphyra blochii*) is a species of hammerhead shark, and part of the family Sphyrnidae. Reaching a length of 1.9 m (6.2 ft), this small brown to gray shark has a slender body with a tall, sickle-shaped first dorsal fin. Its...



Winter flounder

Pseudopleuronectes americanus

The winter flounder (*Pseudopleuronectes americanus*), also known as the black back, is a right-eyed ("dextral") flatfish of the family Pleuronectidae. It is native to coastal waters of the western north Atlantic coast, from Labrador, Canada to...



Witch

Glyptocephalus cynoglossus

The witch (*Glyptocephalus cynoglossus*), known in English by a variety of other common names including the witch flounder, pole flounder, craig fluke, Torbay sole, and grey sole, is a species of flatfish from the family Pleuronectidae. It occurs...



Witch

Arnoglossus scapha

The witch, Arnoglossus scapha, is a lefteye flounder of the family Bothidae, found around China and New Zealand, in waters less than 400 m in depth. Their length is from 20 to 40 cm.



Worm pearlfish

Encheliophis vermicularis[1]

The worm pearlfish (Encheliophis vermicularis) is an eel-like fish in the family Carapidae.



Wrestling halfbeak

Dermogenys pusilla

The wrestling halfbeak (Dermogenys pusilla) also known as Malayan halfbeak is a species of viviparous halfbeak native to the fresh and brackish waters of rivers and coastal regions in South-East Asia, in Singapore, Thailand, Indonesia, Thailand,...



Wrymouth

Cryptacanthodes maculatus

The wrymouth (Cryptacanthodes maculatus) sometimes called ghostfish or "potato-head fish", is a slim, eel-like creature belongs to the wrymouth family Cryptacanthodidae. It outgrows the blennies, its relatives, and may reach a length of 97...



Wuchang bream

Megalobrama amblycephala

The Wuchang bream (Megalobrama amblycephala; Chinese: 武昌魚; pinyin: Wǔchāng yú) is a species of cyprinid fish native to bodies of water throughout the Yangtze basin, China, including Liangzi Lake. It is an important object of fish farming, and in...



Xanthichthys mento

Xanthichthys mento

Xanthichthys mento, the redbtail triggerfish, blue-throat triggerfish, or crosshatch triggerfish, is a species of triggerfish from the Pacific. It inhabits outer-reef areas at depths of 6–131 m (20–430 ft), and feeds on zooplankton. Xanthichthys...



Xenentodon cancila

Xenentodon cancila

Xenentodon cancila, the freshwater garfish, is a species of needlefish found in freshwater and brackish habitats in South and Southeast Asia.



Xenobrama microlepis

Xenobrama microlepis

Xenobrama microlepis, the golden pomfret, is a species of pomfret, a type of fish, found in the Subantarctic Pacific Ocean. This species is known to grow to a length of 49.4 centimetres (19.4 in) SL. This species is the only known member of the...



Xingu River ray

Potamotrygon leopoldi

The Xingu River ray, white-blotched river stingray, or polka-dot stingray (Potamotrygon leopoldi) is a species of freshwater fish in the family Potamotrygonidae. It is endemic to the Xingu River basin in Brazil and as such prefers clear waters...



Xiphophorus montezumae

Xiphophorus montezumae

Xiphophorus montezumae, the Montezuma swordtail, is a livebearing freshwater fish of the order Cyprinodontiformes, family Poeciliidae, and genus Xiphophorus. It is in the same genus as the common platy and the swordtail. Xiphophorus means...



Xiphophorus nezahualcoyotl

Xiphophorus nezahualcoyotl

Xiphophorus nezahualcoyotl, the mountain swordtail, is a live bearing fish in the family Poeciliidae. It is endemic to the northwestern Pánuco River basin in Mexico. The specific name of this fish refers to the poet, philosopher and emperor of...



Yaoshania pachychilus

Yaoshania pachychilus

Yaoshania pachychilus, the panda loach, is a species of gastromyzontid loach endemic to mountain streams in Jinxiu County, Guangxi in China. This species grows to a length of 5.8 centimetres (2.3 in) SL. This species is monotypic, but it was...



Yasuhikotakia modesta

Yasuhikotakia modesta

Yasuhikotakia modesta (blue botia) is a tropical freshwater fish of the family Botiidae. It is native to large rivers in Thailand, Laos, Cambodia, and Vietnam. The blue botia is a widely available fish in the aquarium trade and can be purchased...



Yellow bass

Morone mississippiensis

The yellow bass (Morone mississippiensis) is a member of the family Moronidae. This species is a deep bodied fish that possesses five to seven dark stripes laterally along the sides, the lowest few of these are often broken or disrupted anterior...



Yellow boxfish

Ostracion cubicum

The yellow boxfish (Ostracion cubicum) is a species of boxfish found in reefs throughout the Pacific Ocean and Indian Ocean as well as the southeastern Atlantic Ocean. Recorded occasionally since 2011 in the Levantine waters of the Mediterranean...



Yellow bullhead

Ameiurus natalis

The yellow bullhead (*Ameiurus natalis*) is a species of bullhead catfish, a ray-finned fish that lacks scales.



Yellow clown goby

Gobiodon okinawae

The yellow clown goby, *Gobiodon okinawae*, also known as the Okinawa goby or yellow coral goby, is a member of the goby family native to the western Pacific from southern Japan to the southern reaches of the Great Barrier Reef. As the name...



Yellow goatfish

Mulloidichthys martinicus

The yellow goatfish (*Mulloidichthys martinicus*), also known as yellowsaddle, is a species of goatfish native to the Atlantic Ocean around the coasts of Africa and the Americas. This species can reach a total length of 39.4 cm (15.5 in), but most...



Yellow grouper

Epinephelus awoara

The yellow grouper (*Epinephelus awoara*), also known as the banded grouper, is a species of marine ray-finned fish, a grouper from the subfamily *Epinephelinae* which is part of the family *Serranidae*, which also includes the anthias and sea basses....



Yellow jack

Carangoides bartholomaei

The yellow jack (*Carangoides bartholomaei*), also known as coolihoo, is a species of marine fish in the jack family, *Carangidae*. It is one of only two representatives of its genus present in the Atlantic Ocean, inhabiting waters off the east coast...



Yellow longnose butterflyfish

Forcipiger flavissimus

The yellow longnose butterflyfish or forceps butterflyfish (Forcipiger flavissimus) is a species of marine fish in the family Chaetodontidae. The yellow longnose butterflyfish is widespread throughout the tropical waters of the Indo-Pacific area...



Yellow moray

Gymnothorax prasinus

The yellow moray (Gymnothorax prasinus) is a moray eel of the genus Gymnothorax, found in southern Australia and between North Cape and the Māhia Peninsula on the North Island of New Zealand.



Yellow perch

Perca flavescens

The yellow perch (Perca flavescens), commonly referred to as perch, striped perch or preacher is a freshwater perciform fish native to much of North America. The yellow perch was described in 1814 by Samuel Latham Mitchill from New York. It is...



Yellow prawn-goby

Cryptocentrus cinctus

The yellow prawn-goby (Cryptocentrus cinctus) is a species of goby native to the Western Pacific, where it can be found at depths of from 1 to 25 metres (3.3 to 82.0 ft) in coastal bays and lagoons. This species is symbiotic with alpheid shrimps....



Yellow rasbora

Rasbora lateristriata

The yellow rasbora (Rasbora lateristriata) is a species of ray-finned fish in the genus Rasbora from Southeast Asia. It is a primarily freshwater fish originally from Java island in Indonesia. It is known as the Wader pari fish in the Indonesian...



Yellow stingray

Urobatis jamaicensis

The yellow stingray (Urobatis jamaicensis) is a species of stingray in the family Urotrygonidae, found in the tropical western Atlantic Ocean from North Carolina to Trinidad. This bottom-dwelling species inhabits sandy, muddy, or seagrass bottoms...



Yellow tang

Zebrasoma flavescens

The yellow tang (Zebrasoma flavescens), also known as the lemon sailfin, yellow sailfin tang or somber surgeonfish, is a species of marine ray-finned fish belonging to the family Acanthuridae which includes the surgeonfishes, unicornfishes and...



Yellow-edged lyretail

Variola louti

The yellow-edged lyretail (Variola louti) also known as the yellowedge coronation trout, fairy cod, lunar tail rock cod, lunartailed cod, lyre-tail cod or moontail seabass,, is a species of marine ray-finned fish, a grouper from the subfamily...



Yellow-edged moray

Gymnothorax flavimarginatus

The yellow-edged moray, also known as yellow-margin(ed) moray, leopard moray, and speckled moray, (Gymnothorax flavimarginatus) is a moray eel of the family Muraenidae, found in the Indo-Pacific Oceans at depths to 150 m (500 ft).



Yellow-eye mullet

Aldrichetta forsteri

Yellow-eye mullet (Aldrichetta forsteri), also known as Coorong mullet (after the Coorong area of South Australia), conmuri, estuary mullet, Forster's mullet, freshwater mullet, pilch, pilchard, Victor Harbor mullet, yelloweye, yellow-eyed mullet...



Yellow-tailed African tetra

Alestopetersius caudalis

The yellow-tailed African tetra (Alestopetersius caudalis) is a freshwater fish that lives in the Congo River basin. It is found in both the Democratic Republic of the Congo and the Republic of the Congo (via an unsubstantiated report). It is an...



Yellowback fusilier

Caesio xanthonota

The yellowback fusilier (Caesio xanthonota) is a pelagic marine ray-finned fish, a fusilier belonging to the family Caesionidae. It is native to the tropical Indo-Pacific, being found in shallow water from the East African coast to Indonesia.



Yellowbelly flounder

Rhombosolea leporina

The yellowbelly flounder (Rhombosolea leporina) is a flatfish of the genus Rhombosolea, found around New Zealand. A different species from the genus Rhombosolea is found in Australia and also goes by the name yellow-belly flounder. The Māori...



Yelloweye rockfish

Sebastes ruberrimus

The yelloweye rockfish (Sebastes ruberrimus) is a species of marine ray-finned fish belonging to the subfamily Sebastinae, the rockfishes, part of the family Scorpaenidae. and one of the biggest members of the genus Sebastes. Its name derives...



Yellowfin barbel

Luciobarbus xanthopterus

The yellowfin barbel (Luciobarbus xanthopterus) is a species of cyprinid fish endemic to the Tigris-Euphrates River System.



Yellowfin croaker

Umbrina roncadore

The yellowfin croaker (Umbrina roncadore) is a species of croaker occurring from the Gulf of California, Mexico, to Point Conception, California. They frequent bays, channels, harbors and other nearshore waters over sandy bottoms. These croakers...



Yellowfin goatfish

Mulloidichthys vanicolensis

The yellowfin goatfish (Mulloidichthys vanicolensis) is a species of goatfish native to the Indian Ocean and the Pacific Ocean.



Yellowfin grouper

Mycteroperca venenosa

The yellowfin grouper (Mycteroperca venenosa) is a species of marine ray-finned fish, a grouper from the subfamily Epinephelinae which is part of the family Serranidae, which also includes the anthias and sea basses. It is found in the warmer...



Yellowfin sole

Limanda aspera

The yellowfin sole (Limanda aspera) is a flatfish of the family Pleuronectidae. It is a demersal fish that lives on soft, sandy bottoms at depths of up to 700 metres (2,300 ft), though it is most commonly found at depths of around 91 metres (299...



Yellowfin tuna

Thunnus albacares

The yellowfin tuna (*Thunnus albacares*) is a species of tuna found in pelagic waters of tropical and subtropical oceans worldwide. Yellowfin is often marketed as ahi, from the Hawaiian 'ahi, a name also used there for the closely related bigeye...



Yellowfin whiting

Sillago schomburgkii

The yellowfin whiting (Sillago schomburgkii), also known as the western sand whiting or fine-scaled whiting, is a species of inshore marine fish in the smelt-whiting family Sillaginidae. The species is endemic to the eastern Indian Ocean, ranging...



Yellowhead jawfish

Opistognathus aurifrons

The yellowhead jawfish (Opistognathus aurifrons) is a species of jawfish native to coral reefs in the Caribbean Sea. It is found at depths of from 3 to 40 metres (9.8 to 131.2 ft). The head and upper body are a light, but brilliant, yellow color...



Yellowhead wrasse

Halichoeres garnoti

The yellowhead wrasse (Halichoeres garnoti) is a fish species belonging to wrasse family native to shallow tropical waters in the Caribbean Sea and western Atlantic Ocean.



Yellowmargin triggerfish

Pseudobalistes flavimarginatus

The yellowmargin triggerfish, pineapple trigger, yellowface triggerfish or yellow-face triggerfish (Pseudobalistes flavimarginatus) is a marine fish in the family Balistidae. It is found in coastal tropical waters and reefs of the Indo-Pacific...



Yellowspotted trevally

Carangoides fulvoguttatus

The yellowspotted trevally (Carangoides fulvoguttatus), also known as the yellowspotted kingfish, goldspotted trevally, tarrum, or yellowspot, is a widespread species of large inshore marine fish in the jack family Carangidae. The yellowspotted...



Yellowstone cutthroat trout

Oncorhynchus clarkii bouvieri

The Yellowstone cutthroat trout (Oncorhynchus clarkii bouvieri) is a subspecies of the cutthroat trout (Oncorhynchus clarkii). It is a freshwater fish in the salmon family (family Salmonidae). Native only to a few U.S. states, their original...



Yellowstripe scad

Selaroides leptolepis

The yellowstripe scad (Selaroides leptolepis), also known as the yellowstripe trevally, yellow-banded trevally, smooth-tailed trevally, slender-scaled trevally and slender trevally, is a species of small inshore fish in the jack and horse...



Yellowtail amberjack

Seriola lalandi

The yellowtail amberjack, yellowtail kingfish, hiramasa or great amberjack (Seriola lalandi) is a large fish found in the Southern Ocean. Although previously thought to be found in all oceans and seas, recent genetic analysis restricts S. lalandi...



Yellowtail barracuda

Sphyraena flavicauda

The yellowtail barracuda (Sphyraena flavicauda) is one of the smaller species of barracuda of the family Sphyraenidae, which can be found in Indo-West Pacific oceans. It has also invaded the Mediterranean through the Suez Canal from the Red Sea,...



Yellowtail flounder

Limanda ferruginea

The yellowtail flounder (Limanda ferruginea), also known as the rusty dab, is a species of flatfish in the family Pleuronectidae (righteye flounders). Reaching 56 cm (22 in) in length, it has reddish brown upperparts, pale underparts and yellow...



Yellowtail horse mackerel

Trachurus novaezelandiae

The yellowtail horse mackerel (Trachurus novaezelandiae), also known as the yellowtail scad, is a jack in the family Carangidae found around Australia and New Zealand at depths to 500 m. Its length is up to 50 centimetres (20 in). The yellowtail...



Yellowtail rockfish

Sebastes flavidus

The yellowtail rockfish (Sebastes flavidus), or yellowtail seaperch, is a species of marine ray-finned fish belonging to the subfamily Sebastinae, the rockfishes, part of the family Scorpaenidae. This species lives mainly off the coast of western...



Yellowtail scad

Atule mate

The yellowtail scad (Atule mate) (also known as the northern yellowtail scad, one-finlet scad, deep trevally and omaka), is an abundant species of small inshore marine fish of the jack family, Carangidae. The species is widespread in the...



Yellowtail snapper

Ocyurus chrysurus

The yellowtail snapper (Ocyurus chrysurus) is an abundant species of snapper native to the western Atlantic Ocean including the Gulf of Mexico and the Caribbean Sea. Although they have been found as far north as Massachusetts, their normal range...



Yoyo loach

Botia almorhae

The yoyo loach, Almora loach or Pakistani loach (Botia almorhae) is a freshwater fish belonging to the loach family Botiidae. It originates in the slow-running and still waters of the Ganges basin in northern India and possibly Nepal. Despite the...



Zacco platypus

Zacco platypus

The pale chub (Zacco platypus), also known as pale bleak or fresh-water sprat, is one of the most extensively distributed Asiatic cyprinids and is found in Korea, Japan, Taiwan, and most of Southeast China. This species of fish is native to...



Zander

Sander lucioperca

The zander (Sander lucioperca), sander or pikeperch, is a species of ray-finned fish from the family Percidae, which also includes perch, ruffe and darter. It is found in freshwater and brackish habitats in western Eurasia. It is a popular game...



Zappa confluentus

Zappa confluentus

Zappa confluentus, the New Guinea slender mudskipper, is a mudskipper endemic to New Guinea, where it is only known from the lower parts of the Fly, Ramu and Bintuni Rivers. It is found on mudflats adjacent to turbid rivers. This species can reach...



Zebra bullhead shark

Heterodontus zebra

Sharks portalThe zebra bullhead shark (*Heterodontus zebra*) is a bullhead shark of the family Heterodontidae found in the central Indo-Pacific between latitudes 40°N and 20°S, from Japan and Korea to Australia. It is typically found at relatively...



Zebra loach

Botia striata

The zebra loach (*Botia striata*) is a freshwater loach native to rivers and streams in the Western Ghats of India. The maximum size is about 9 cm (3.5 in). It lives in tropical climate with temperature range of 21–26 °C (70–79 °F), and prefers...



Zebra mbuna

Maylandia zebra

The zebra mbuna (*Maylandia zebra*) is a species of cichlid endemic to Lake Malawi in Africa. This species can reach a length of 11.3 cm (4.4 in). It feeds on aufwuchs, a surface layer of mostly algal material that grows on rocks. This cichlid is a...



Zebra moray

Gymnomuraena zebra

The zebra moray (*Gymnomuraena zebra*) is a species of marine fish in the family Muraenidae. It is the only member of the genus *Gymnomuraena*, though it sometimes has been included in *Echidna* instead.



Zebra oto

Otocinclus cocama

The zebra oto or tiger oto (Otocinclus cocama) is a fish of the genus Otocinclus of the family Loricariidae that originates from Peru.



Zebra pleco

Hypancistrus zebra

The zebra pleco (Hypancistrus zebra) is a species of catfish endemic to a small section of the Rio Xingu in Brazil. It occurs in the big bend area of the river, downstream from the town of Altamira, Para. It was first described in 1991. It gets...



Zebra seahorse

Hippocampus zebra

The zebra seahorse (Hippocampus zebra) is a species of fish in the family Syngnathidae. It is endemic to northern Australia.



Zebra Shark

Stegostoma tigrinum

Zebra sharks (Stegostoma tigrinum) are slow-swimming, bottom-dwelling sharks. They are called so because, as juveniles, these sharks have dark bodies with yellowish stripes. As they mature, the pattern changes to small dark spots on a grayish-tan...



Zebrafish

Danio rerio

The zebrafish (Danio rerio) is a freshwater fish belonging to the minnow family (Cyprinidae) of the order Cypriniformes. Native to India and South Asia, it is a popular aquarium fish, frequently sold under the trade name zebra danio (and thus...



Zebrasoma desjardinii

Zebrasoma desjardinii

Zebrasoma desjardinii, the Red Sea sailfin tang, Desjardin's sailfin tang, Indian sailfin tang or Indian sailfin surgeonfish, is a species of marine ray-finned fish belonging to the family Acanthuridae, the surgeonfishes, unicornfishes and tangs....



Zebrasoma gemmatum

Zebrasoma gemmatum

Zebrasoma gemmatum, the gem tang, jewelled tang, spotted tang or Mauritian tang, is a species of marine ray-finned fish belonging to the family Acanthuridae which includes the surgeonfishes, unicornfishes and tangs. This species is found in the...



Zebrasoma rostratum

Zebrasoma rostratum

Zebrasoma rostratum, the longnose surgeonfish, longnose tang or black tang, is a species of marine ray-finned fish belonging to the family Acanthuridae, which includes the surgeonfishes, unicornfishes and tangs. This fish is found in the western...



Zebrasoma scopas

Zebrasoma scopas

Zebrasoma scopas, the brown tang, twotone tang, scopas tang or brush-tail tang, is a species of marine ray-finned fish belonging to the family Acanthuridae which includes the surgeonfishes, unicornfishes and tangs. The brown tang is found throughout...



Zebrasoma xanthurum

Zebrasoma xanthurum

Zebrasoma xanthurum, the purple tang or yellowtail tang, is a species of marine ray-finned fish belonging to the family Acanthuridae which includes the surgeonfishes, unicornfishes and tangs. This species is endemic to the north western Indian...



Zeugopterus punctatus

Zeugopterus punctatus

Zeugopterus punctatus, the common topknot, is a species of left eyed flatfish in the family Scophthalmidae, from the eastern Atlantic Ocean.



Zig-zag eel

Mastacembelus armatus

The zig-zag eel (*Mastacembelus armatus*), also known as the tire-track eel, tire-track spiny eel or marbled spiny eel, is a species of ray-finned, spiny eels belonging to the genus *Mastacembelus* (Scopoli, 1777) of the family Mastacembelidae, and is...



Zingel asper

Zingel asper

Zingel asper, also known as the Rhone streber, is a species of freshwater ray-finned fish in the family Percidae that is at a high risk of becoming extinct due to habitat destruction and pollution. It is endemic to the rivers Durance and Beaume in...



Zingel zingel

Zingel zingel

Zingel zingel, the zingel, is a species of freshwater ray-finned fish in the family Percidae. It is found in fast-flowing streams in southeastern Europe. It is the type species of the genus Zingel.



Zona

Bedotia geayi

The red-tailed silverside, or zona (*Bedotia geayi*) is a species of Madagascar rainbowfish endemic to the Mananjary River drainage in Madagascar. It is threatened by habitat loss and introduced species. It has often been confused with the related B...



Zoogoneticus tequila

Zoogoneticus tequila

Zoogoneticus tequila, Tequila splitfin or simply Tequila fish, is a species of goodeid fish (family Goodeidae) from Mexico. The specific epithet, tequila, derives from the Tequila Volcano, which looms near the type locality.



Zu cristatus

Zu cristatus

Zu cristatus, the scalloped ribbonfish, is a ribbonfish of the family Trachipteridae found circumglobally in all oceans at tropical latitudes, at depths down to 90 m. Its length is up to 118 cm. In the past, the binomial name Trachipterus iris was...